k-mer spectra and allelic coverage analyses reveal pervasive polymorphic duplications in *Ostrea edulis*

Lila Colston-Nepali[1,2]*, Sylvie Lapègue[1], and Nicolas Bierne[2]*

Affiliations:

[1] **MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Montpellier, France**

[2] **ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France**

* corresponding authors: lila.colston-nepali@umontpellier.fr ; nicolas.bierne@umontpllier.fr

ABSTRACT

Marine bivalves are known for their high genetic diversity and potentially high genetic load, as well as their structurally complex genomes. The European flat oyster, *Ostrea edulis*, is no exception. In this study we performed both *k*-mer based and reference-based analyses on short-read data with high coverage (~70-160X) from five individuals. Up to one-third of heterozygous SNPs showed allelic coverage fractions that deviated from expectation. Despite no evidence of recent whole-genome duplication, we detected significant signals for genomic regions of increased ploidy, and 15% and 25% of *k*-mer pairs distant by a single nucleotide displayed triploid-like and tetraploid-like profiles respectively. In a diploid species, this indicates the presence of heterozygote and homozygote duplications. These results were confirmed by coverage-based genotype inference, which showed that deviant SNPs are located in polymorphic duplications. While duplications affected genic and non-genic regions similarly, their genomic distribution varied across chromosomes, with a notable enrichment in non-metacentric chromosomes. The latter being prone to somatic aneuploidy, we suggest that aneuploidy could act as an indirect driver, with duplications potentially buffering the expression of recessive deleterious mutations in aneuploid cells. Finally, we consider how such widespread structural variation may complicate population genomic analyses in *O. edulis* and other marine bivalves.

INTRODUCTION

Marine bivalves often exhibit very high levels of genetic diversity, making them ideal models for population genetics since the advent of allozyme markers (Ward, Woodwark, and Skibinski 1994; Bazin, Glémin, and Galtier 2006). Early studies have revealed a number of surprising observations, including heterozygote deficiencies relative to Hardy-Weinberg equilibrium (Zouros and Foltz 1984), segregation distortions in the offspring of mating crosses (Bierne et al. 1998; Launey and Hedgecock 2001), and heterozygosity-fitness correlations (David 1998). These have led to interpretations based on profuse adaptive variation (Mitton 2000; Koehn 1983), unusual demographics such as chaotic genetic patchiness (Eldon et al. 2016; David et al. 1997) and sweepstakes reproductive variance (Hedgecock and Pudovkin 2011), and high rates of somatic aneuploidy (Thiriot-Quiévreux 1986; Leitão et al. 2001), but also technical issues such as null alleles (Foltz 1986). Although the very large size of bivalve populations could explain both the high polymorphism (Harrang et al. 2013; Takeuchi 2017) leading *de facto* to technical and cytogenetical problems, and a high genetic load (Plough 2016), not all issues have been fully resolved. The transition to DNA sequence analysis has confirmed high nucleotide diversity both at silent and amino-acid mutations in marine bivalves (Sauvage et al. 2007; Harrang et al. 2013; Romiguier et al. 2014). However, recent access to structural variation in genome sequence studies has revived the idea of potentially unique bivalve genetics underpinned by widespread presence-absence (Gerdol et al. 2020) and copy number (Modak et al. 2021; Qi, Li, and Zhang 2021) variation, which have been suggested to be adaptive (i.e. maintained by selection). As for allozyme and SNP markers, though, high population size and genetic load, possibly associated with sweepstakes reproduction, might also explain this observation similarly well.

Highly polymorphic species usually have a compact genome (Lynch and Walsh 2007), though this is not always the case, such as in marine bivalves. A high proportion of their genomes are made of repetitive sequences (Šatović and Plohl 2021). This includes pearl oysters (*Pinctada fucata*, Takeuchi et al. 2012; Takeuchi et al. 2022), Pacific oysters (*Magallana gigas*, Zhang et al. 2012), Mediterranean mussels (*Mytilus galloprovincialis*, Murgarella et al. 2016), Manila clams (*Ruditapes philippinarum*, Mun et al. 2017), deep sea and shallow-water mussels (*Bathymodiolus platifrons* and *Modiolus philippinarum*, Sun et al. 2017), scallops (*Patinopecten yessoensis*, Wang et al. 2017), and golden mussels (*Limnoperna fortunei*, Uliano-Silva et al. 2018). Although repetitive sequences include highly repeated elements such as mobile elements, satellite DNA, or tandem repeats (Šatović and Plohl 2021), segmental duplications (SDs) have been found to make up 26% of the Pacific oyster genome - the highest recorded in published genomes (Qi, Li, and Zhang 2021). Studies of resequenced genomes have also revealed presence-absence variation (PAV) throughout the Mediterranean mussel genome, with approximately 25% of genes subject to PAV (Gerdol et al. 2020), but also persistent copy number variable regions (CNVRs) consisting of over 43% of the Pacific oyster genome (Qi, Li, and Zhang 2021), and duplication variation contributing to over 16.5% of the eastern oyster (*Crassostrea virginica*) genome (Modak et al. 2021).

However, population genomics data have been predominantly obtained with low coverage
genome sequencing in order to analyze many samples within constrained project budgets. These
studies focus on SNPs and ignore structural variation, even though some of these SNPs are likely
to be found in these parts of the genome. Downstream analyses rely on a number of assumptions
about genome assemblies, although next-generation sequencing methods have inherent technical
risks of error that may violate these assumptions and lead to misinterpretation of the data. For
this reason complex filtering strategies are used to improve the accuracy and reliability of
genome-wide SNP datasets (O'Leary et al. 2018; Hemstrom et al. 2024). Very few studies,
though, consider the effect of genome complexities on next-generation sequencing. Dallaire et al.
(2023) described the effects of rediploidization and repetitive elements on low-coverage whole
genome sequencing in a salmonid fish, which they hypothesize causes collapsed assembled
genomic regions. As a result SNPs may be called that do not conform to expected patterns of
heterozygosity and allelic coverage fractions, which they refer to as 'deviant SNPs'. For
example, deviant SNPs deviated from the expected allelic coverage fraction of 1/2, with a
hotspot of SNPs demonstrating an allelic coverage fraction of 1/4 (Dallaire et al. 2023). These
deviant SNPs have a significant effect on downstream analyses and population genetic
inferences, and most filtering approaches will fail to remove them (Dallaire et al. 2023).
Consideration then, of the presence of these deviant SNPs in non-model species with complex
genomes, such as bivalves, represents a potential blind spot in population genomic analyses.

The European flat oyster (*Ostrea edulis*) is a marine bivalve from the family Ostreidae, or the
true oysters. *O. edulis* has highly variable reproductive success, both in the wild and in
experimental conditions, and a low Ne/N ratio (Hedgecock et al. 2007; Lallias et al. 2010). *O.*
edulis has been found to have high genetic load causing segregation distortions in pair crosses
(Bierne et al. 1998), and up to a third of nonsynonymous mutations have been estimated to be
mildly deleterious in this species (Harrang et al. 2013). The genome size of *O. edulis* is also
larger than other sequenced oyster species (Boutet et al. 2022; Gundappa et al. 2022; Li et al.
2023), and contains comparable repeated elements as *M. gigas*, *C. virginica*, and *Saccostrea*
glomerata (Boutet et al. 2022; Li et al. 2023). Synteny analyses indicate that *O. edulis* has not
experienced a recent whole genome duplication event since divergence from Crassostreinae (Li
et al. 2023), although such an event may have occurred at the origin of all the Ostreidae.

In this study we used *k*-mer and allelic coverage analyses on five very high coverage re-
sequenced *O. edulis* genomes (70-160X) in order to infer the proportion of SNPs that deviate
from the expectation of diploid genotypes in single copy orthologs. Our *k*-mer analyses showed
that ~15% and ~25% of our SNPs are triploid-like (3X coverage) and tetraploid-like (4X
coverage) respectively. Allelic coverage analyses also identified SNPs that did not adhere to
diploid expectation. Investigating how these are shared between individuals, we obtained
evidence that they are located in polymorphic duplications.

METHODS

1. Sampling and sequencing

Five *O. edulis* samples with high coverage short-read sequencing were used: four newly sequenced samples from Galicia, Spain (sequencing depth ranging from ~70-160X coverage), and one from Brittany, France, using publicly available reads (SRA accession: SRR1723031, ~70X coverage, named ROSC in this study). Three of the Galician samples (2C4M, 6C9M, 8C17M) came from a culture raft in Cambados, Ría de Arousa, and one sample (4C6M) came from a natural bed in Ría de Ferrol. DNA was extracted from mantle tissue for all four using Machery-Nagel's NucleoSpin Tissue extraction kit. The four DNA samples were sent to the MGX sequencing platform, which produced the TruSeq DNA PCR-Free libraries, followed by sequencing on Illumina Novaseq 6000 on an S4 line.

2. Reference-free k-mer analyses

As a first step, reference-free *k*-mer based analyses were used. *k*-mer approaches allow users to evaluate large, and potentially complex, genomic datasets computationally efficiently and without the need for a reference genome (Jenike et al. 2025). *k*-mer frequency histograms were computed using the program FastK (https://github.com/thegenemyers/FASTK) on the raw reads, using a *k*-mer count of 21, and then two different tools were used to analyse the *k*-mer spectrum. First, GenomeScope 2.0 (Ranallo-Benavidez, Jaron and Schatz 2020) was used to analyze the coverage of the *k*-mers, and fit to a model of 2 x p negative binomial distributions (p = ploidy) representing the unique and multi-copy heterozygous and homozygous *k*-mers. A ploidy of 2, or diploid, was used. GenomeScope 2.0 was also used to estimate genome size and sequencing error in the dataset.

Next, the program Smudgeplot (v. 0.3.0) (Ranallo-Benavidez, Jaron and Schatz 2020) was used to find *k*-mer pairs that differ by exactly one nucleotide, representing either heterozygous alleles, or complex genome structures (i.e. duplications, paralogs, etc.). Total coverage of the *k*-mer pairs is plotted against the minor *k*-mer coverage to reveal hotspots that represent the haplotype structures present, and their frequency, in the genome. Input parameters for error margins and coverage ranges were set based on GenomeScope 2.0 output. Unlike GenomeScope 2.0, ploidy is not needed *a priori* to run Smudgeplot.

In order to investigate and compare genome-wide duplication events in other ostreids, GenomeScope 2.0 and Smudgeplot analyses were also performed on publicly available short-read datasets of three oyster species: the lamellated oyster *O. denselamellosa* (SRA accession: SRR19238449), the Olympia oyster *O. lurida* (SRA accessions: SRR3306651-SRR3306656), and the Pacific oyster *M. gigas* (SRA accession: SRR7474375). To compare the duplications in the flat oyster to a species with known PAV (Gerdol et al. 2020), *k*-mer analyses were also used

203   on the Mediterranean mussel *M. galloprovincialis* (SRA accession: SRR17154976). Further
204   analyses were implemented on two other *Mytilus* species: the common mussel *M. edulis* (SRA
205   accession: SRR28968509), and the thick-shelled mussel *M. coruscus* (SRA accession:
206   SRR10502233). The SRA toolkit (https://github.com/ncbi/sra-tools) was used to download and
207   extract FASTQ files. For *O. lurida*, six sequencing runs of the same individual were pooled.
208

3.   *Reference-based analyses*

3.1 Alignment to reference genome and SNP calling

Sequence data was trimmed using the program fastp (Chen et al. 2018) to filter for quality and
remove both polyG tails and Illumina adapters. Samples were aligned to an *O. edulis* reference
genome (Li et al. 2023, GenBank assembly: GCA_032173915.1) using the program BWA-MEM
(Li and Durbin 2010). Mapped reads were then filtered using the program Sambamba (Tarasov
et al. 2015) to remove reads that did not pair properly, and secondary alignments. SNPs were
called using bcftools mpileup and bcftools call (Danecek et al. 2021), with maximum coverage
increased to 2000, resulting in 20,868,611 sites.

3.2 Dosage based inference of genotypes

Output VCF files were filtered to include only biallelic SNPs, and to remove both indels and all
missing data using vcftools (Danecek et al. 2011) and bcftools (Danecek et al. 2021). SNPs
mapping to unplaced scaffolds were removed in R (R Core Team 2023), with 15,106,345
remaining sites for downstream analyses. A custom python script was used to extract the
reference (A) and alternative (B) allele counts, total allele count, and calculate the allele
coverage fraction (site frequency) per SNP (defined as alternative reads / total reads) for each of
the 5 individuals. The output CSV files were then analyzed in R (R Core Team 2023). Firstly,
allelic coverage fraction distributions for heterozygous sites were assessed using histograms of
alternative allele frequency for each individual. Histograms of site coverage were then used to
estimate the coverage per individual, and determine the expected relative depth for 2X (diploid),
3X (triploid-like), and 4X (tetraploid-like) sites. Higher relative depth values (i.e. 5X and above)
that correspond to the presence of alleles with more than two repeated copies were ignored, as
they represent a small proportion of SNPs. The mode of the distribution was used as an estimate
for the 2X relative depth. This estimate was then used alongside the density plots to estimate the
expected 1X, 3X and 4X relative depth for each individual. Density plots were visualized with
ggplot2 (Wickham 2016) to investigate the relationship between the relative depth and the allelic
coverage fraction. Gene dosage clusters were identified and circumscribed for 12 potential
genotypes at biallelic sites: AA, AB, BB, AAA, AAB, BBA, BBB, AAAA, AAAB, AABB,
BBBA, BBBB (see Supplementary Fig. 1). These dosage-inferred genotypes (DI-genotypes),
were identified for a subset of SNPs for which we have good confidence of the inferred

genotype, and were then used to further investigate variation between individuals, chromosomes,
and functional categories (e.g. genic vs. non-genic).
245
246   *3.2.1 Variation between individuals*
247
248 To determine if the same SNPs were categorised into the same DI-genotype in different
249 individuals, SNP lists were created for each of the 12 DI-genotypes for each individual.
250 Comparisons were made for each individual (i.e. the percentage of AA SNPs in individual X that
251 are present in each of the 12 categories in individual Y), leading to a total of 20 repetitions for
252 144 comparisons.
253
254   *3.2.2 Location and density of different genomic regions*
255
256 The 12 DI-genotypes were combined to represent the three types of dosage 2X, 3X and 4X (i.e.
257 AA, AB, BB were pooled as 2X etc.). To see if the 3X and 4X SNPs are present in specific
258 regions in the genome, the tool CMplot (Yin et al. 2021) was used to visualize SNP-density. The
259 proportions of 2X, 3X and 4X SNPs were also calculated per chromosome for each individual.
260
3.2.3 Functional annotation of SNPs
The program SnpEff (Cingolani et al. 2012) was used to annotate the SNPs. A new database for
the flat oyster was built using annotation, coding sequence (CDS), and protein sequence files for
the same assembly that was used as a reference genome (Li et al. 2023). Annotation categories
were simplified into four main groups- intergenic, intronic, synonymous and nonsynonymous
variants (see Supplementary Table 1 for details). The proportion of 2X, 3X, and 4X SNPs was
calculated for each of these annotation categories.

268
269 **RESULTS AND DISCUSSION**
270
271 ***k*-mer analyses reveals a high proportion of non-diploid *k*-mers in *O. edulis* and other**
272 **bivalve genomes**
273
The *k*-mer coverage distributions obtained with Genomescope 2.0 showed that a significant
proportion of *k*-mers were classified outside of the standard diploid coverage expectation in the
genomes of all five individuals (Fig. 1A and Supplementary Figures 2A-5A), with signals for
both 3X and 4X *k*-mers present. To further investigate this signal, we conducted Smudgeplot
analyses that use the co-distribution of total coverage and the minor *k*-mer coverage of closely
related *k*-mer pairs to determine how many copies of each allele exist across a genome.
Smudgeplot allows the delineation of distinct dosage clusters that would be otherwise difficult to
determine using coverage alone (even when coverage is high, as is the case here). The AAB
cluster with a 3X coverage and an allelic coverage fraction of 1/3 is not expected to be found in a

diploid genome, except in cases of heterozygous duplication (where one chromosome has a 1-
copy allele and the other has a 2-copy allele). Smudgeplot analyses indicated strong evidence for
non-diploid *k*-mer pairs, with 14-16% with a 3X (AAB) signal, and 21-26% with a 4X (AAAB
or AABB) signal (Fig. 1B and Supplementary Figures 2B-5B).
287
288

289
290   **Figure 1.** A) GenomeScope profile for sample 6C9M with dotted lines indicating 1X, 2X, 3X and 4X *k*-
291   mer coverages and B) corresponding Smudgeplot analysis for the same individual, using an error
292   threshold of 46 and a coverage range of 50-110.
293
Given that Smudgeplot has rarely been used in bivalves, we reanalysed several public NGS
datasets in both ostreids (*Ostrea denselamellosa*, *O. lurida*, and *Magalllana gigas*), and mytilids
(Mytilus galloprovincialis, *M. edulis*, and *M. coruscus*) in order to compare patterns of non-
diploid genomic regions. We observed that 3X and 4X clusters were found in every genome, but
O. edulis had the strongest signal for 3X and 4X regions in both Genomescope 2.0 and
Smudgeplot analyses out of the species re-analysed in this study (Supplementary Figures 6-11).
Further, *O. edulis* is the only species in this study that has a higher proportion of 4X than 3X *k*-
mer pairs. Although *M. gigas* was described as having a very high proportion of both CNVRs
and SDs (Qi, Li, and Zhang 2021), it was the species with the weakest signals amongst the
ostreids analysed here. Surprisingly, a Smudgeplot analysis performed in the brooding Chilean
oyster, *O. chilensis*, indicated no *k*-mer pairs deviating from diploid expectation (see Piccoli
2024). The signal of non-diploid *k*-mer pairs in *Mytilus* mussels suggests that this may be a
general feature of bivalve genomes, and that the presence or absence of one of the two copies of
a duplication could be a potential driver of the presence-absence signal detected in mussels
(Gerdol et al. 2020).
309

Reference-free *k*-mer analyses thus provide a first indication (with the high frequency of AAB,
AAAB, and AABB clusters) of the presence of heterozygous duplications in *O. edulis* and other
bivalve genomes. However, a reference mapping approach is needed in order to better
understand the variation between individuals, how 3X and 4X SNPs cluster in the genome, how
called SNPs are affected by polymorphic duplications, and to investigate the distribution of these
duplications between chromosomes and between functional categories (i.e. coding vs. non-
coding).
317
318 **Reference mapping analysis confirms polymorphic duplications in *O. edulis***
319
First we investigated the fraction of allelic coverage of heterozygous SNPs and found a
surprisingly large proportion (approximately 30%) of heterozygous SNPs with a coverage
fraction between 0.2 and 0.4 (Fig. 2 and Supplementary Figure 12). Although routine inspection
of allelic coverage should be included in best practice for NGS data analysis (Jaron et al. 2022,
Bierne 2022, Ament-Velásquez 2016), it is unfortunately not reported on sufficiently in the
literature. Nevertheless, it is evident that such a significant deviation from the expected
distribution is highly unusual.
327

328
329
330 **Figure 2.** Histogram demonstrating the distribution of the allelic coverage fraction for called
331 heterozygous sites in individual 6C9M.
332
333 In order to better delineate 3X and 4X SNPs, we adopted a similar approach to Smudgeplot by
334 investigating the co-distribution of total coverage and allelic coverage fraction of heterozygote
335 calls. SNPs adhering to the expectations for heterozygous diploid sites, AB sites (i.e. 2X relative
336 depth and 0.5 allelic coverage fraction), had the strongest density (Fig. 3) as expected.
337

338
Figure 3. Density plot for sample 6C9M showing allelic coverage fraction on the x-axis and coverage on
the y-axis. Marked on the right hand side of the plot is relative depth, the coverage of the SNP divided by
the genome average coverage. Areas of increased density shown in brighter yellow. Locations on plot
corresponding to specific dosage inferred genotypes marked.

However, corroborating the Smudgeplot results, strong density was also seen in many of the
other dosage-inferred clusters corresponding to 3X and 4X genotypes as well as outside these
regions (including at relative depth below 2X, as expected when one allele is less efficiently
mapped with lower quality scores). This confirms that there are numerous heterozygous and
homozygous duplications distributed throughout the species genome. Most SNPs that adhere to
diploid expectations (2X SNPs), were also identified as regular diploid SNPs in other
individuals. On average, over 95% of SNPs identified as AB in one individual were classified as
AA (50.3%), AB (29.7%), or BB (15.7%) SNPs in other individuals (Fig. 4A and Supplementary
Table 2).

353
354
355

356
357 **Figure 4.** Boxplots showing the percentage of shared sites between SNPs categorized into **A** AB DI-
358 genotypes, **B** AAB DI-genotypes, and **C** AABB DI-genotypes in one individual, and the 12 DI-genotypes
359 in other individuals.
360
The AAB SNPs in one individual were primarily classified as AA (42%) or AAB (18.5%) in
other individuals (Figure 4B). This is what we would expect when a 2-copy allele with a
mutation in one copy segregates at low frequency (the duplication is mainly found at the
heterozygous state). In this case individuals homozygous for the two-copy allele are rare, which
could explain why only 5% of these SNPs are categorized as AABB SNPs in other individuals.
The rest were classified as AAA (12.4%), AB (8.9%), AAAA (4.2%), AAAB (3.7%), and BB
(2.5%) SNPs. Finally, if we look at AABB SNPs in one individual, the majority are classified as
AA (24.5%), AAB (21.7%) or AABB (20.8%) SNPs in other individuals (Fig. 4C and
Supplementary Table 2). SNPs switch between DI-genotypes in different individuals, suggesting

that these duplications are not fixed. This result is in better accordance with the expectation for
polymorphic duplications (where the 2-copy allele has a mutation in one copy). Given that the
focal individual is a homozygote for the duplication (AABB), we expect the frequency of the 2-
copy allele to be higher than when the focal individual is a heterozygote for the duplication
(AAB), and that we should mostly observe the three expected genotypes (AA, AAB, AABB).
The rest were mainly classified as AB (8.3%), AAA (5.9%) and AAAB (5%).
376

The other DI-genotypes observed can be explained if we consider the possible mutational paths
taken to obtain the polymorphism observed in this study (Figure 5A and 5B). First, we must have
a duplication creating the 2-copy allele. The two copies are initially identical, and until a
nucleotide mutation occurs, there is no SNP. A nucleotide mutation can then occur in one copy
of the 2-copy allele (Figure 5A), creating a SNP at this position. This results in three alleles
segregating in the population: the 1-copy allele carrying the ancestral allele of the SNP (one dose
of A), the 2-copy allele carrying the ancestral allele of the SNP on both copies (two doses of A),
and the 2-copy allele with the derived allele on one copy and the ancestral allele on the other
(one dose of A plus one dose of B). Here allele A refers to the ancestral allele, and allele B to the
derived allele. However, in the dataset A will refer to the allele carried by the reference genome,
and B will refer to the alternative allele. In this scenario, we expect not only AA, AAB, and
AABB DI-genotypes, but also AAA, AAAB, and AAAA DI-genotypes, which correspond to the
presence of the unmutated 2-copy allele, contributing two doses of the ancestral allele in a DI-
genotype. Alternatively, following duplication a mutation can occur on the 1-copy allele instead,
creating an alternative mutational path (Fig. 5B). This results in three different alleles
segregating in the population: the 1-copy allele carrying the ancestral allele of the SNP (one dose
of A), the 2-copy allele carrying the ancestral allele of the SNP on both copies (two doses of A),
and the 1-copy allele carrying the derived allele of the SNP (one dose of B). In this scenario, we
expect the DI-genotypes AA, AB and BB corresponding to the nucleotide polymorphism in the
1-copy allele, but also the AAA, AAB, and AAAA genotypes corresponding to the presence of
the 2-copy allele with the ancestral SNP allele on both copies, contributing two doses of A in the
DI genotype.
399

Figure 5. Two potential mutational paths leading to the polymorphic duplications seen in *O. edulis*. Each pentagon represents a portion of the genome, and the vertical lines in the pentagons represent the variant position. Stars represent mutations. Plots to the right of the mutational paths demonstrate the relative depth and allelic coverage fraction of the resulting genotypes. In mutational path **A**, a duplication event occurs, leading to a 2-copy allele. Then a mutation occurs in one of the two copies of the 2-copy allele, resulting in an ancestral variant (A) and a derived variant (B) at the focal SNP. The result is three types of allele present: a 1-copy allele with the ancestral variant (A), a 2-copy allele with the same variant at the SNP (AA), and a 2-copy allele with two different variants (AB). In mutational path **B**, a duplication event occurs, leading to a 2-copy allele. Then a mutation occurs in the 1-copy allele. The result is three types of allele present in the population: a 1-copy allele with the ancestral variant (A), a 2-copy allele with the same variant at the SNP (AA) and a 1-copy allele with a mutation (B).

These two scenarios can explain the diversity of DI-genotypes observed, and their frequencies given the genotype of the focal individual. When the alternative allele corresponds to the ancestral allele, we expect all the symmetrical DI-genotypes that we observe in the data. Of course, there are many other potential scenarios, such as when the polymorphic duplication is due to a deletion of one copy of a fixed duplication. There may also be more complex

demographic scenarios than a single population model, such as secondary contact between a
lineage that has a fixed duplication, and has accumulated some divergence between the two
copies (AABB), and a lineage that has kept the ancestral 1-copy allele (AA). This scenario
would generate the polymorphisms we observe the most- AA, AAB, AABB. Further, we note
that the duplication is not necessarily a tandem repeat, and there may be recombination between
the two copies, generating recombined genotypes. However, these more complex scenarios
mostly tend to remove intermediate DI-genotypes of the mutational path, and we believe that our
results are well explained simply by widespread polymorphic duplication.
427
428 **Duplications are not randomly distributed between chromosomes, but affect the functional**
429 **categories of SNPs equally.**
430
431 The density of 3X and 4X SNPs in 100kb windows varied throughout the genome (Fig. 6,
432 Supplementary Figures 13-16), allowing us to identify windows of non-diploid SNPs
433 corresponding to collapsed duplicated regions.
434

435
436
437 **Figure 6.** SNP-density plots for A) 3X SNPs and B) 4X SNPs in sample 6C9M. Density of SNPs per
438 100kb window shown for all 10 chromosomes. Position on the chromosome indicated on the x-axis.
439
Surprisingly, some chromosomes were found to have a higher density of non-diploid SNP
clusters. Chromosomes 6, 7 and 8 appeared to have the highest density for all individuals, and
this was verified by the average count for each chromosome (Fig. 7A). At this stage we do not
have a clear explanation for this result, and further work will be needed to better understand the
pattern. However, there are some different characteristics between the first five and last five
chromosomes that may indicate potential drivers of this result. The first five chromosomes each
have an evident putative centromere at their centre, revealed by a valley of nucleotide diversity
(Supplementary Figure 17). Conversely, there are no valleys of nucleotide diversity in the central
part of the last five chromosomes. Further, only chromosomes 8 and 10 have a reduction in
diversity towards their ends, perhaps because these are low-recombining regions as often found

in telomeric regions (*O. edulis* does not have acrocentric chromosomes so this can not explain
these regions).
452

453
454
Figure 7. A Percentage of SNPs classified as 2X, 3X and 4X in all 10 chromosomes, averaged across all
456 five individuals and **B** percentage of intergenic, intronic, nonsynonymous and synonymous variants
457 classified as 2X, 3X and 4X SNPs, averaged across all five individuals.
458

Variation in chromosomal morphology has been noted in both *O. edulis* and its closely related
sister species *O. angasi* (Thiriot-Quiévreux 1984 ; Li and Havenhand 1997), as well as in other
ostreids (Leitão et al. 2002). In both species, the first five chromosomes are metacentric, which
conforms to the position of putative centromeres that we identify using valleys of nucleotide
diversity in these chromosomes. The last five shorter chromosomes consist of submetacentric
and subtelocentric pairs. This conforms to the lack of valleys of nucleotide diversity in the centre
of these chromosomes. Duplications are often produced by problems during recombination, or
mistakes during cellular division (Ohta 2000, Zhang 2003), and these issues may be more likely
to occur in chromosomes prone to cohesion defects and mispairing. Under this hypothesis, the
higher density of polymorphic duplications in chromosomes 6, 7 and 8 would be caused by a
higher rate of duplication mutation, owing to chromosome pairing and problems with
recombination. It is not clear whether this explanation is plausible, but it would mean that
duplication alleles would behave mostly neutrally and mutation rate defines their density, or that
the genetic load of deleterious duplicated or deleted copy alleles would be greater on these three
chromosomes. The proportion of 4X to 3X SNPs is higher in these 3 chromosomes (0.44, 0.36
and 0.42 for chromosomes 6, 7 and 8 respectively, compared with an average of 0.24 for the
other chromosomes), suggesting, as explained above, that the 2-copy alleles segregate at higher
frequency in these chromosomes.

477
Another explanation could be linked to somatic aneuploidy, which is known to occur frequently
in bivalves (Thiriot-Quievreux 2002), including *O. edulis* (Thiriot-Quiévreux 1986).
Chromosome loss and gain are well known to be nonrandom and influenced by centromere size
and cohesion defects (Sheltzer and Amon 2011). In marine bivalves, aneuploidy is biased
towards chromosome loss, but affects only a small proportion of cells in a sample (Thiriot-
Quiévreux, Pogson, and Zouros 1992), and affects specific chromosomes only (Bouilly et al.
2005; Thiriot-Quiévreux, Pogson, and Zouros 1992; Li & Havenhand 1997; Leitão et al. 2001;
Khatir et al. 2022). In *O. edulis*, chromosome loss is always found at chromosomes 6 to 10
(Thiriot-Quiévreux 1986), in accordance with these chromosomes being more prone to instability
during cell divisions. We do not think that somatic aneuploidy directly explains our results, as it
only affects a fraction of cells in a sample, which would create dosages intermediate to our
defined categories, while clusters are well-defined in our *k*-mer and reference based analyses. It
may still explain the uncategorized SNPs though, which are found at significantly higher rates in
these chromosomes (χ^2 test significant at p < 2.2e-16 for all individuals. See Supplementary
Statistics). We suggest that somatic chromosome loss exposes recessive deleterious mutations to
negative selection, and that extra copies producing compensation for this loss, including dosage
compensation, can buffer the genetic load exposed by aneuploidy (Kondrashov 2012; Katju and
Bergthorsson 2013; Basilicata and Valsecchi 2021). We investigated whether 2X, 3X, and 4X
SNPs are found in similar proportions between functional categories. Intergenic and
nonsynonymous sites had a slightly lower, although significant (χ^2 test significant at p < 2.2e-16
for all individuals. See Supplementary Statistics), proportion of 2X SNPs than intronic and
synonymous sites (Fig. 7B). This result suggests that coding regions are similarly affected by
duplications than non-coding regions, and that functional sites may even be slightly more
affected than nonfunctional sites. This concurs with the observations made in *C. virginica*
(Modak et al. 2021), and *M. galloprovincialis* (Gerdol et al. 2020).
503
Finally, similar patterns occurring in *M. gigas* further support our hypothesis concerning dosage
compensation in chromosomes prone to aneuploidy. Qi, Li, and Zhang (2021) observe a higher
rate of segmental duplications in chromosomes 1, 4, 8 and 10, and a higher rate of CNVRs in
chromosomes 1, 3 and 4. Synteny analysis performed by Li et al. (2023) indicates that these
chromosomes correspond to the last 5 chromosomes in *O. edulis* (see Supplementary Table 3).
Bouilly et al. (2005) investigated somatic aneuploidy in *M. gigas*, and found certain
chromosomes to be more prone to chromosome loss. The study predates the reference genome
for *M. gigas* and it is not possible to definitively determine which chromosome numbers
correspond to the chromosomes described by Qi, Li, and Zhang (2021). However, Bouilly et al.
(2005) numbered the chromosomes by size, and therefore it is very likely that their chromosome
1 corresponds to chromosome 1 as numbered by Qi, Li, and Zhang (2021), as this chromosome
is significantly larger than the other chromosomes. Chromosome 1, with a high rate of segmental
duplications and CNVRs (Qi, Li, and Zhang 2021), was identified as experiencing the highest
rates of somatic aneuploidy (Bouilly et al. 2005).

518
Implications for the population genetics of bivalves
520
Though the high proportion of non-diploid regions in *O. edulis*, demonstrated by both the *k*-mer
analyses and the reference mapping approach in this study, is in-line with previous results in
marine bivalves (e.g. Modak et al. 2021, Qi, Li, and Zhang 2021), the skewed distribution of
individual allelic coverage fraction (~30% of SNPs between 0.2 and 0.4), is still surprising. This
result raises two fundamental questions. Firstly, how can low or medium coverage data be
analysed for the purposes of population structure, demographic reconstruction, and selection
scan studies? Secondly, is there, as proposed regularly for decades, something potentially unique
about the biology, ecology, or genetics of marine bivalves that can explain their unusual
polymorphism, including rampant polymorphic duplications?
530
The availability of five high coverage genomes gave us the opportunity to take an unusual
approach to the study of structural variation and its impact on SNP calling. Usually, only one
individual used as a reference genome is deeply sequenced, while the resequenced genomes have
insufficient coverage to undertake the dosage-based genotyping that we used in this study. Short
indels or CNVs can be called with short-read data, but calling large duplications requires pair-
mate, linked-read, or long-read sequencing, methods that are still rarely used in population
genomics, especially in bivalves. Further, even with these methods, if the duplicated copy is too
long for the long DNA molecules to pass through, and if the two copies are too similar to resolve
duplication by assembly, a recent long duplication can easily be missed. With these issues in
mind, dosage-based inference with high coverage provides valuable information. However,
filtering of medium or low coverage datasets on the basis of coverage of a single individual alone
can not provide the necessary information. The distributions of 2X, 3X, and 4X coverage, and
1/2, 1/3, and 1/4 allelic coverage fraction overlap too much with insufficient sequencing depth.
In addition, the duplications causing the inflated coverage and skewed allelic coverage fractions
occur in different regions in different individuals, making it difficult to filter specific SNPs. The
analysis of the coverage of a number of individuals may help alleviate the issue. If there are
enough sufficiently covered heterozygote and homozygote duplications in the sample then the
excess coverage and skewed allelic coverage fraction can be detected on a sample-wide scale.
This is the information that the programs ngsParalog (Linderoth 2018) or paramask (Tjeng et al.
2025) use to filter SNPs considered 'deviant' in multicopy regions. These methods also use
heterozygote frequencies in the sample, assuming Hardy-Weinberg proportions for the former,
and fitting an unknown inbreeding coefficient for the latter. These programs also rely on the
expectation that deviant SNPs cluster together in collapsed duplicated regions of the genome.
The programs aim to mainly target 4X coverage and 1/4 allelic coverage fraction, with the
assumption that the duplications are fixed. Therefore, these methods would need to be modified
in order to better account for polymorphic duplications. These methods are likely to remove
many problematic SNPs, and filtering with them is highly recommended in bivalves, and indeed
any non-model species. However, they are unlikely to remove variants in rare duplicated alleles,

or SNPs that do not depart much from Hardy-Weinberg proportions. In the case of *O. edulis*,
however, we can advise the removal of the three most affected chromosomes (i.e. 6, 7, 8) for
specific analyses such as population structure or demographic inference.
562

The explanation for why *O. edulis*, and perhaps bivalves in general, have such a high rate of
polymorphic duplication can not be solved here. Pangenome studies will be needed to better
understand the mechanistic constraints and evolutionary forces that shape these patterns. The
putative whole genome duplication ancestral to Ostreidae (>50MYA), or ancestral to *Mytilus*
mussels (Corrochano-Fraile et al. 2022) might have helped initiate the dynamic of gain and loss
of gene copies, but cannot reasonably explain our result. Divergence between the two copies
must be low, because the duplication mutation must be recent to be still polymorphic, or because
the merging of duplicates in the reference genome would not occur with a strong divergence
between copies. In fact, the dosage-based genotypes observed suggest that the reference genome
most often has the duplication merged into a single copy, or has the 1-copy allele at the
polymorphic duplication identified. This is expected given that most duplications were found to
be polymorphic and should follow a classic allelic frequency spectrum (one given individual has
the 1-copy allele at most of the polymorphic duplications). Follow-up studies should attempt to
obtain the frequency of the 2-copy allele in different populations. This will allow testing for
deviation from neutral expectations. We propose that the high burden of polymorphic
duplications could be a simple consequence of the global high genetic diversity of marine
bivalves with large population sizes. Indeed, a similar proportion of polymorphism in coding and
non-coding regions is usually taken as evidence for neutral or nearly-neutral theory. However,
adaptive or maladaptive forces may also be at play. The possible role of presence-absence and
duplication variation in the evolutionary success, phenotypic variation, and local adaptation of
marine bivalves has been emphasized by others (e.g. Gerdol et al. 2020; Modak et al. 2021). We
suggest instead that our observations are in accordance with the high genetic load of these
abundant and highly fertile organisms, which may be further enhanced by their sweepstakes
reproduction (Hedgecock and Pudovkin 2011; Harrang et al. 2013; Plough 2016). Just as a high
rate of segregating deleterious non-synonymous mutations have been suspected (Harrang et al.
2013), the same explanation could be proposed for mildly deleterious copy-number variants. The
population genetics of accurately called duplication variants is needed to provide an answer to
this question.

Acknowledgements

The authors are very grateful to Antonio Villalba for carrying out the sampling of the four Galician flat oyster samples used in this study. They also express their thanks to Xavier Dallaire, Pierre-Alexandre Gagnaire, Céline Reisser, and Christelle Fraïsse for valued discussion and insight.

Funding

- This work was supported by the scientific management of Ifremer and the Occitanie region as part of Lila Colston-Nepali's doctoral contract.
- Data used in this work were partly produced through the GenSeq technical facilities of the "Institut des Sciences de l'Evolution de Montpellier" with the support of LabEx CeMEB, an ANR "Investissements d'avenir" program (ANR-10-LABX-04-01).

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. The authors declare the following non-financial conflict of interest: Nicolas Bierne is a recommender for PCI.

Data, scripts, code and supplementary information availability

The four newly sequenced *O. edulis* whole genomic read datasets are deposited in DATAREF, the Ifremer centralized and secure storage through data archiving. A DOI has been obtained and they have been submitted to the ENA.

Supplementary materials are available on bioRxiv:
https://www.biorxiv.org/content/10.1101/2025.06.24.661118v1.supplementary-material

LITERATURE CITED

Ament-Velásquez SL, Figuet E, Ballenghien M, Zattara EE, Norenburg JL, Fernández-Álvarez FA, Bierne J, Bierne N, Galtier N. 2016. Population genomics of sexual and asexual lineages in fissiparous ribbon worms (Lineus, Nemertea): hybridization, polyploidy and the Meselson effect. *Molecular Ecology* 25:3356–3369. DOI: 10.1111/mec.13717.

Basilicata MF, Valsecchi CIK. 2021. The good, the bad, and the ugly: Evolutionary and pathological aspects of gene dosage alterations. *PLOS Genetics* 17:e1009906. DOI:

10.1371/journal.pgen.1009906.

Bazin E, Glémin S, Galtier N. 2006. Population size does not influence mitochondrial genetic

diversity in animals. *Science* 312:570–572. DOI: 10.1126/science.1122033.

Bierne N. 2022. Pressing NGS data through the mill of *k*-mer spectra and allelic coverage ratios

in order to scan reproductive modes in non-model species. *Peer Community in Evolutionary*

Biology 1:100142. DOI: 10.24072/pci.evolbiol.100142.

Bierne N, Launey S, Naciri-Graven Y, Bonhomme F. 1998. Early effect of inbreeding as revealed

by microsatellite analyses on *Ostrea edulis* larvae. *Genetics* 148:1893–1906. DOI:

10.1093/genetics/148.4.1893.

Bouilly K, Leitão A, Chaves R, Guedes-Pinto H, Boudry P, Lapègue S. 2005. Endonuclease

banding reveals that atrazine-induced aneuploidy resembles spontaneous chromosome loss in

Crassostrea gigas. *Genome* 48:177–180. DOI: 10.1139/g04-087.

Boutet I, Alves Monteiro HJ, Baudry L, Takeuchi T, Bonnivard E, Billoud B, Farhat S, Gonzales-

Araya R, Salaun B, Andersen AC, Toullec J, Lallier FH, Flot J, Guiglielmoni N, Guo X, Li C,

Allam B, Pales-Espinosa E, Hemmer-Hansen J, Moreau P, Marbouty M, Koszul R, Tanguy A.

2022. Chromosomal assembly of the flat oyster (*Ostrea edulis* L.) genome as a new genetic

resource for aquaculture. *Evolutionary Applications* 15:1730–1748. DOI: 10.1111/eva.13462.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics (Oxford, England) 34:i884–i890. DOI: 10.1093/bioinformatics/bty560.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Ruden DM, Lu X. 2012.

A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff:

SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly* 6:80-92. DOI:

10.4161/fly.19695.

Corrochano-Fraile A, Davie A, Carboni S, Bekaert M. 2022. Evidence of multiple genome

duplication events in Mytilus evolution. *BMC Genomics* 23:340. DOI: 10.1186/s12864-022-

08575-9.

661 Dallaire X, Bouchard R, Hénault P, Ulmo-Diaz G, Normandeau E, Mérot C, Bernatchez L,

662 Moore J-S. 2023. Widespread deviant patterns of heterozygosity in whole-genome sequencing

663 due to autopolyploidy, repeated elements, and duplication. *Genome Biology and Evolution*

664 15:evad229 DOI: 10.1101/2023.07.27.550877.

665 Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,

666 Marth GT, Sherry ST, McVean G, Durbin R, 1000 Genomes Project Analysis Group. 2011. The

667 variant call format and VCFtools. *Bioinformatics (Oxford, England)* 27:2156–2158. DOI:

668 10.1093/bioinformatics/btr330.

669 Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,

670 McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and BCFtools. *GigaScience*

671 10:giab008. DOI: 10.1093/gigascience/giab008.

672 David P. 1998. Heterozygosity–fitness correlations: new perspectives on old problems. *Heredity*

673 80:531–537. DOI: 10.1046/j.1365-2540.1998.00393.x.

674 David P, Berthou P, Noel P, Jarne P. 1997. Patchy recruitment patterns in marine invertebrates: a

675 spatial test of the density-dependent hypothesis in the bivalve Spisula ovalis. *Oecologia* 111:331–

676 340. DOI: 10.1007/s004420050243.

677 Eldon B, Riquet F, Yearsley J, Jollivet D, Broquet T. 2016. Current hypotheses to explain genetic

678 chaos under the sea. *Current Zoology* 62:551–566. DOI: 10.1093/cz/zow094.

679 Foltz DW. 1986. Null alleles as a possible cause of heterozygote deficiencies in the oyster

680 *Crassostrea virginica* and other bivalves. *Evolution* 40:869–870. DOI: 10.1111/j.1558-

681 5646.1986.tb00549.x.

682 Gerdol M, Moreira R, Cruz F, Gómez-Garrido J, Vlasova A, Rosani U, Venier P, Naranjo-Ortiz

683 MA, Murgarella M, Greco S, Balseiro P, Corvelo A, Frias L, Gut M, Gabaldón T, Pallavicini A,

684 Canchaya C, Novoa B, Alioto TS, Posada D, Figueras A. 2020. Massive gene presence-absence

685 variation shapes an open pan-genome in the Mediterranean mussel. *Genome Biology* 21:275.

686 DOI: 10.1186/s13059-020-02180-3.

Gundappa MK, Peñaloza C, Regan T, Boutet I, Tanguy A, Houston RD, Bean TP, Macqueen DJ.

2022. Chromosome-level reference genome for European flat oyster (*Ostrea edulis* L.).

Evolutionary Applications 15:1713–1729. DOI: 10.1111/eva.13460.

Harrang E, Lapègue S, Morga B, Bierne N. 2013. A high load of non-neutral amino-acid

polymorphisms explains high protein diversity despite moderate effective population size in a

marine bivalve with sweepstakes reproduction. *G3 Genes|Genomes|Genetics* 3:333–341. DOI:

10.1534/g3.112.005181.

Hedgecock D, Launey S, Pudovkin AI, Naciri Y, Lapègue S, Bonhomme F. 2007. Small effective

number of parents (Nb) inferred for a naturally spawned cohort of juvenile European flat oysters

Ostrea edulis. *Marine Biology* 150:1173–1182. DOI: 10.1007/s00227-006-0441-y.

Hedgecock D, Pudovkin AI. 2011. Sweepstakes reproductive success in highly fecund marine

fish and shellfish: A Review and Commentary. *Bulletin of Marine Science* 87:971–1002. DOI:

10.5343/bms.2010.1051.

Hemstrom W, Grummer JA, Luikart G, Christie MR. 2024. Next-generation data filtering in the

genomics era. *Nature Reviews Genetics*:1–18. DOI: 10.1038/s41576-024-00738-6.

Jaron KS, Hodson CN, Ellers J, Baird SJE, Ross L. 2022. Genomic evidence of paternal genome

elimination in the globular springtail Allacma fusca. *Genetics* 222:iyac117. DOI:

10.1093/genetics/iyac117.

Jenike KM, Campos-Domínguez L, Boddé M, Cerca J, Hodson CN, Schatz MC, Jaron KS. 2025.

k-mer approaches for biodiversity genomics. *Genome Research*. DOI: 10.1101/gr.279452.124.

Katju V, Bergthorsson U. 2013. Copy-number changes in evolution: rates, fitness effects and

adaptive significance. *Frontiers in Genetics* 4. DOI: 10.3389/fgene.2013.00273.

Khatir Z, Hizam Z, Lyons B, Leitão A. 2022. Aneuploidy in the pearl oyster *Pinctada radiata*

(Leach, 1814): evidence for nonrandom chromosome loss and gain in marine bivalves.

Malacologia 65(1-2), 71-77. DOI: 10.4002/040.065.0105.

Koehn RK. 1983. 9 - Biochemical Genetics and Adaptation in Molluscs. In: Hochachka PW ed.

The Mollusca. San Diego: Academic Press, 305–330. DOI: 10.1016/B978-0-12-751402-4.50016-

9.

Kondrashov FA. 2012. Gene duplication as a mechanism of genomic adaptation to a changing

environment. *Proceedings of the Royal Society B: Biological Sciences* 279:5048–5057. DOI:

10.1098/rspb.2012.1108.

Lallias D, Taris N, Boudry P, Bonhomme F, Lapègue S. 2010. Variance in the reproductive

success of flat oyster *Ostrea edulis* L. assessed by parentage analyses in natural and experimental

conditions. *Genetics Research* 92:175–187. DOI: 10.1017/S0016672310000248.

Launey S, Hedgecock D. 2001. High Genetic Load in the Pacific Oyster *Crassostrea gigas*.

Genetics 159:255–265. DOI: 10.1093/genetics/159.1.255.

Leitão A, Boudry P, Thiriot-Quiévreux C. 2001. Evidence of differential chromosome loss in

aneuploid karyotypes of the Pacific oyster, *Crassostrea gigas*. *Genome*, 44: 735-737.

Leitao A, Chaves R, Santos S, Boudry P, Guedes Pinto H, Thiriot Quievreux C, Santos S, Boudry

P, Guedes Pinto H, Thiriot Quievreux C. 2002. Cytogenetic study of *Ostrea conchaphila*

(Mollusca : Bivalvia) and comparative karyological analysis within Ostreinae. *Journal of Shellfish*

Research 21:685–690.

Li X, Bai Y, Dong Z, Xu C, Liu S, Yu H, Kong L, Li Q. 2023. Chromosome-level genome

assembly of the European flat oyster (*Ostrea edulis*) provides insights into its evolution and

adaptation. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*

45:101045. DOI: 10.1016/j.cbd.2022.101045.

Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform.

Bioinformatics (Oxford, England) 26:589–595. DOI: 10.1093/bioinformatics/btp698.

Li XX, Havenhand JN. 1997. Karyotype, nucleolus organiser regions and constitutive

heterochromatin in *Ostrea angasi* (Molluscae: Bivalvia): evidence of taxonomic relationships

within the Ostreidae. *Marine Biology* 127:443–448. DOI: 10.1007/s002270050031.

Linderoth T. 2018. Identifying population histories, adaptive genes and genetic duplication from

population-scale next generation sequencing. Berkeley: University of California.

Lynch M, Walsh B. 2007. The origins of genome architecture. Sinauer Associated, Sunderland,

Massachusetts.

Mitton JB. 2000. Selection in natural populations. Oxford University Press.

Modak TH, Literman R, Puritz JB, Johnson KM, Roberts EM, Proestou D, Guo X, Gomez-

Chiarri M, Schwartz RS. 2021. Extensive genome-wide duplications in the eastern oyster

(*Crassostrea virginica*). *Philosophical Transactions of the Royal Society B: Biological Sciences*

376:20200164. DOI: 10.1098/rstb.2020.0164.

Mun S, Kim Y-J, Markkandan K, Shin W, Oh S, Woo J, Yoo J, An H, Han K. 2017. The whole-

genome and transcriptome of the Manila clam (*Ruditapes philippinarum*). *Genome Biology and*

Evolution 9:1487–1498. DOI: 10.1093/gbe/evx096.

Murgarella M, Puiu D, Novoa B, Figueras A, Posada D, Canchaya C. 2016. A First insight into

the genome of the filter-feeder mussel *Mytilus galloprovincialis*. *PLOS ONE* 11:e0151561. DOI:

10.1371/journal.pone.0151561.

Ohta T. 2000. Evolution of gene families. *Gene* 259:45–52. DOI: 10.1016/S0378-

1119(00)00428-5.

O'Leary SJ, Puritz JB, Willis SC, Hollenbeck CM, Portnoy DS. 2018. These aren't the loci you're

looking for: Principles of effective SNP filtering for molecular ecologists. *Molecular Ecology*

27:3193–3206. DOI: 10.1111/mec.14792.

Piccoli GAR. 2024. *Ostrea chilensis* genomic analysis and population diversity: widespread-

hemizygosity genomics. Victoria University of Wellington.

Plough LV. 2016. Genetic load in marine animals: a review. *Current Zoology* 62:567–579. DOI:

10.1093/cz/zow096.

Qi H, Li L, Zhang G. 2021. Construction of a chromosome-level genome and variation map for

the Pacific oyster *Crassostrea gigas*. *Molecular Ecology Resources* 21:1670–1685. DOI:

10.1111/1755-0998.13368.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. Vienna, Austria:

R Foundation for Statistical Computing. https://www.R-project.org/

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for

reference-free profiling of polyploid genomes. *Nature Communications* 11:1432. DOI:

10.1038/s41467-020-14998-3.

Romiguier J, Gayral P, Ballenghien M, Bernard A, Cahais V, Chenuil A, Chiari Y, Dernat R,

Duret L, Faivre N, Loire E, Lourenco JM, Nabholz B, Roux C, Tsagkogeorga G, Weber A a.-T.,

Weinert LA, Belkhir K, Bierne N, Glémin S, Galtier N. 2014. Comparative population genomics

in animals uncovers the determinants of genetic diversity. *Nature* 515:261–263. DOI:

10.1038/nature13685.

Šatović Vukšić E, Plohl M. 2021. Exploring satellite DNAs: specificities of bivalve mollusks

genomes. *Progress in molecular and subcellular biology* 60:57–83. DOI: 10.1007/978-3-030-

74889-0_3.

Sauvage C, Bierne N, Lapègue S, Boudry P. 2007. Single nucleotide polymorphisms and their

relationship to codon usage bias in the Pacific oyster *Crassostrea gigas*. *Gene* 406:13–22. DOI:

10.1016/j.gene.2007.05.011.

Sheltzer JM, Amon A. 2011. The aneuploidy paradox: costs and benefits of an incorrect

karyotype. *Trends in Genetics* 27:446–453. DOI: 10.1016/j.tig.2011.07.003.

Sun J, Zhang Y, Xu T, Zhang Y, Mu H, Zhang Y, Lan Y, Fields CJ, Hui JHL, Zhang W, Li R,

Nong W, Cheung FKM, Qiu J-W, Qian P-Y. 2017. Adaptation to deep-sea chemosynthetic

environments as revealed by mussel genomes. *Nature Ecology & Evolution* 1:1–7. DOI:

10.1038/s41559-017-0121.

Takeuchi T. 2017. Molluscan genomics: implications for biology and aquaculture. *Current*

Molecular Biology Reports 3:297–305. DOI: 10.1007/s40610-017-0077-3.

Takeuchi T, Kawashima T, Koyanagi R, Gyoja F, Tanaka M, Ikuta T, Shoguchi E, Fujiwara M,

Shinzato C, Hisata K, Fujie M, Usami T, Nagai K, Maeyama K, Okamoto K, Aoki H, Ishikawa

T, Masaoka T, Fujiwara A, Endo K, Endo H, Nagasawa H, Kinoshita S, Asakawa S, Watabe S,

Satoh N. 2012. Draft genome of the pearl oyster *Pinctada fucat*a: a platform for understanding

bivalve biology. *DNA Research* 19:117–130. DOI: 10.1093/dnares/dss005.

Takeuchi T, Suzuki Y, Watabe S, Nagai K, Masaoka T, Fujie M, Kawamitsu M, Satoh N, Myers

EW. 2022. A high-quality, haplotype-phased genome reconstruction reveals unexpected

haplotype diversity in a pearl oyster. *DNA Research* 29:dsac035. DOI: 10.1093/dnares/dsac035.

Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast processing of NGS

alignment formats. *Bioinformatics (Oxford, England)* 31:2032–2034. DOI:

10.1093/bioinformatics/btv098.

Thiriot-Quiévreux C. 1984. (Bivalvia). *Cahiers de Biologie Marine* 25:407-418.

Thiriot-Quiévreux C. 1986. Etude de l'aneuploïdie dans différents naissains d'Ostreidae

(Bivalvia). *Genetica* 70:225–231. DOI: 10.1007/BF00122190.

Thiriot-Quievreux C. 2002. Review of the literature on bivalve cytogenetics in the last ten years.

Cahiers de Biologie Marine 43:17-26.

Thiriot-Quiévreux C, Pogson GH, Zouros E. 1992. Genetics of growth rate variation in bivalves:

aneuploidy and heterozygosity effects in a *Crassostrea gigas* family. *Genome* 35:39–45. DOI:

10.1139/g92-007.

Tjeng B, Arimond M, Grindeland HB, Libera AD, Fulgione A. 2024. ParaMask, a new method to

identify multicopy genomic regions, corrects major biases in whole-genome sequencing data.

DOI: 10.21203/rs.3.rs-4660628/v1.

Uliano-Silva M, Dondero F, Dan Otto T, Costa I, Lima NCB, Americo JA, Mazzoni CJ,

Prosdocimi F, Rebelo M de F. 2018. A hybrid-hierarchical genome assembly strategy to sequence

the invasive golden mussel, *Limnoperna fortunei*. *GigaScience* 7:gix128. DOI:

10.1093/gigascience/gix128.

Wang S, Zhang J, Jiao W, Li J, Xun X, Sun Y, Guo X, Huan P, Dong B, Zhang L, Hu X, Sun X,

Wang J, Zhao C, Wang Y, Wang D, Huang X, Wang R, Lv J, Li Y, Zhang Z, Liu B, Lu W, Hui

Y, Liang J, Zhou Z, Hou R, Li X, Liu Y, Li H, Ning X, Lin Y, Zhao L, Xing Q, Dou J, Li Y, Mao

J, Guo H, Dou H, Li T, Mu C, Jiang W, Fu Q, Fu X, Miao Y, Liu J, Yu Q, Li R, Liao H, Li X,

Kong Y, Jiang Z, Chourrout D, Li R, Bao Z. 2017. Scallop genome provides insights into

evolution of bilaterian karyotype and development. *Nature Ecology & Evolution* 1:1–12. DOI:

10.1038/s41559-017-0120.

Ward RD, Woodwark M; Skibinski DOF. 1994. A comparison of genetic diversity levels in

marine, freshwater and anadromous fish. *Journal of Fisheries Biology* 44:213-232. DOI:

10.1111/j.1095-8649.1994.tb01200.x

Wickham H. 2016. ggplot2: elegant graphics for data analysis. Springer-Verlag New York. ISBN

978-3-319-24277-4. https://ggplot2.tidyverse.org.

Yin L, Zhang H, Tang Z, Xu J, Yin D, Zhang Z, Yuan X, Zhu M, Zhao S, Li X, Liu X. 2021.

rMVP: A Memory-Efficient, Visualization-Enhanced, and Parallel-Accelerated Tool for Genome-

Wide Association Study. *Genomics, Proteomics & Bioinformatics* 19:619–628. DOI:

10.1016/j.gpb.2020.10.007.

Zhang J. 2003. Evolution by gene duplication: an update. *Trends in Ecology & Evolution* 18:292–

298. DOI: 10.1016/S0169-5347(03)00033-8.

Zhang G, Fang X, Guo X, Li L, Luo R, Xu F, Yang P, Zhang L, Wang X, Qi H, Xiong Z, Que H,

Xie Y, Holland PWH, Paps J, Zhu Y, Wu F, Chen Y, Wang J, Peng C, Meng J, Yang L, Liu J,

Wen B, Zhang N, Huang Z, Zhu Q, Feng Y, Mount A, Hedgecock D, Xu Z, Liu Y, Domazet-

Lošo T, Du Y, Sun X, Zhang S, Liu B, Cheng P, Jiang X, Li J, Fan D, Wang W, Fu W, Wang T,

Wang B, Zhang J, Peng Z, Li Y, Li N, Wang J, Chen M, He Y, Tan F, Song X, Zheng Q, Huang

R, Yang H, Du X, Chen L, Yang M, Gaffney PM, Wang S, Luo L, She Z, Ming Y, Huang W,

Zhang S, Huang B, Zhang Y, Qu T, Ni P, Miao G, Wang J, Wang Q, Steinberg CEW, Wang H,

Li N, Qian L, Zhang G, Li Y, Yang H, Liu X, Wang J, Yin Y, Wang J. 2012. The oyster genome

reveals stress adaptation and complexity of shell formation. *Nature* 490:49–54. DOI:

10.1038/nature11413.

843 Zouros E and Foltz DW. 1984. Possible explanations of heterozygote deficiency in bivalve

844 molluscs. *Malacologia* 25:583-591.

845