Genomic analysis identifies *Campylobacter concisus* genomospecies 2 as a novel species

and proposes the name *Campylobacter oralis*

Fang Liu[1#], Siying Chen[1#], Joanna M Biazik[2], Christopher Luk[1], Anjaneyaswamy Ravipati[2],

Ruiting Lan[1], Stephen M Riordan[3], Li Zhang[1]*

1. School of Biotechnology and Biomolecular Sciences, University of New South Wales,

Sydney, NSW 2052, Australia

2. Mark Wainwright Analytical Centre, University of New South Wales, Sydney, NSW 2052,

Australia

3. Gastrointestinal and Liver Unit, Prince of Wales Hospital, University of New South

Wales, Sydney, NSW 2031, Australia

18   * Correspondence author: L.Zhang@unsw.edu.au

19   # Co-first authors

ABSTRACT

Campylobacter concisus comprises two distinct genomospecies (GS) based on core genome

analysis. Here we conducted further analysis and demonstrated that *C. concisus* GS2 strains

belong to a novel *Campylobacter* species.

A total of 245 *C. concisus* genomes including 85 GS1 and 160 GS2 strains were analysed.

DNA-DNA hybridization (DDH) values between strains of GS1 and GS2 *C. concisus* were

from 42.8% to 66%. Average Nucleotide Identity (ANI) between GS1 and GS2 *C. concisus*

strains were from 87.8% to 89.7%. The average GC contents of GS1 and GS2 *C. concisus*

strains were 37.3% and 39.2% respectively. GS1 and GS2 specific genes have been identified.

GS1 and GS2 *C. concisus* strains can also be differentiated based on core genome, 23S rRNA

polymorphisms and protein profiles. Their morphology appears different under certain

conditions.

In conclusion, ANI and DDH values clearly support that *C. concisus* GS2 as a novel

Campylobacter species, for which we propose the name *Campylobacter oralis*. The

identification of *C. concisus* and *C. oralis* specific genes, along with differences in their protein

profiles, morphology, core genome and 23S rRNA phylogeny, further support this

classification.

INTRODUCTION

Campylobacter concisus was first isolated by Tanner et al. from human dental specimen in 1981, with strain ATCC 33237 as the type strain [1]. Later studies isolated *C. concisus* from fecal samples and biopsies of patients with gastritis, inflammatory bowel disease (IBD) which includes Crohn's disease and ulcerative colitis, and other diseases [2-8]. In 2016, Chung et al. conducted a comparative genome analysis of 36 *C. concisus* strains and reported that *C. concisus* contained two GS based on the core genome and identified GS specific genes [9]. Based on the core genome, the *C. concisus* reference strain ATCC33237 is a GS1 strain [9]. Kirk et al. in 2018 examined 113 *C. concisus* genomes and reported that the genomes of GS2 *C. concisus* strains contained higher numbers of genes compared to genomes of GS1 *C. concisus* strains [10]. In 2020, Liu et al. analysed 16 complete sequenced *C. concisus* genomes and reported that the genomes of GS2 *C. concisus* strains had larger sizes and higher GC contents than the genomes of GS1 *C. concisus* strains [11]. In 2021, Cornelius et al. analysed 190 *C. concisus* genomes, also confirmed the existence of GS1 and GS2 *C. concisus* [12].

We hypothesize that GS2 *C. concisus* is a novel species and conducted further analysis of 245 *C. concisus* genomes and also investigated methods that may phenotypically differentiate *C. concisus* GS1 and GS2 strains. Our data support that the GS2 *C. concisus* is a novel *Campylobacter* species, and we propose the name *Campylobacter oralis*. We also examined the presence of *csep1* gene, a molecular marker that is associated with active Crohn's disease and found that one of these markers, *csep1* gene, was only present in *C. oralis*, not in *C. concisus*.

MATERIALS AND METHODS

Analysis of genomes of *C. concisus* GS1 and GS2 strains

The genomes of 245 *C. concisus* strains, including 85 GS1 and 160 GS2 strains, were obtained

from National Center for Biotechnology Information (NCBI) genome database for analysis

(Supplementary Table 1).

Phylogenetic analysis based on core genome, 23S and 16S rRNA genes

The core genome of the 245 *C. concisus* strains was generated by using Roary (v3.12.0) without

splitting paralogs [13]. The phylogenetic tree of core genome was generated by using FastTree

(v2.1.10) based on the approximate-maximum-likelihood method [14]. The sequences of 16S

and 23S rRNA gene were used to generate phylogenetic trees based on the maximum-

likelihood method (100 bootstrap) by using MEGAcc (v10.1.6), and *Campylobacter jejuni*

NCTC113651 was used as an outgroup [15].

DNA-DNA hybridization (DDH), average nucleotide identity (ANI) and GC content

In silico DDH and ANI analyses were performed on all 245 *C. concisus* genomes. DDH values

were   calculated   on   Genome-to-Genome   Distance   Calculator   (GGDC   2.1

https://ggdc.dsmz.de). ANI values were generated using fastANI (v1.3) [16, 17]. The GC

contents were determined using SeqKit [18].

Identification of genomospecies-specific genes

The results on the presence and absence of genes among 245 *C. concisus* strains obtained from

Roary were further analysed through Scoary (v1.6.16), GS-specific genes were then identified

[19]. Redundant genes were excluded until mutually exclusive GS-specific genes were yielded.

The presence of each of the candidate GS specific genes in 245 *C. concisus* strains was

confirmed by blasting each gene against whole genomes and annotated genes using nucleotide

BLAST, and annotated proteins using protein BLAST. In this study, GS specific genes are

defined as genes present in all GS1 strains but absent in all GS2 strains, *vice versa*, as described

previously [9].

Matrix assisted laser desorption ionization-time of flight mass spectrometry (MALDI-

TOF MS) analysis

A total of 20 *C. concisus* strains (10 GS1 strains and 10 GS2 strains) were used for this analysis

(Supplementary Table 1). The complete genomes of these strains were sequenced previously

[11]. These strains were cultured on horse blood agar (HBA) plates under anaerobic conditions

supplemented with 5% H_2 for 48 hours as described previously [20].

Bacterial proteins were prepared using ethanol-formic acid extraction procedure. Briefly, a

loopful bacteria of each strain was suspended in 300 µL of distilled water, and 900 µL of

ethanol was added. After mixing thoroughly, the bacterial cell suspension was centrifuged at

13,000 × g for 2 mins and the supernatant was discarded. The residual ethanol was removed

by repeating centrifugation. The pellet was dried at room temperature for 5 mins and suspended

in 5 to 50 µL of formic acid-water (70:30 [vol/vol]), and an equal volume of ethanol was added.

Following centrifugation at 13,000 × g for 2 mins, 1 µL of supernatant was spotted on ground

steel MALDI target plate and allowed to dry at room temperature prior to overlaying 1 µL of

matrix solution.

MALDI-TOF MS analysis was performed using the linear positive ion mode on Ultraflex TOF-

TOF mass spectrometer (Bruker Daltonik GmbH, Bremen, Germany). External calibrations

were performed using The Bruker Bacterial Test Standard (Bruker Daltanics) according to the

manufacturer's instructions. The raw spectra data were exported to Biotyper Version 4.0 (Build

19) for bacterial identification and FlexAnalysis Version 3.4 for further downstream data

analysis.

Scanning electron microscope (SEM) imaging of GS1 and GS2 *C. concisus* strains

Six *C. concisus* strains were randomly chosen for SEM examination, including three GS1

strains including P26UCO-S2, P10CDO-S2, and P3UCO1; and three GS2 strains including

BEO2, P15UCO-S2, and P2CDO4.

C. concisus strains were cultured on HBA plates for 24 hours, followed by cultivation in heart

infusion broth containing 5% fetal bovine serum for 12 hours. Bacterial cells were collected

and fixed overnight at 4 °C in 2.5% glutaraldehyde in 0.2 M sodium phosphate buffer. Fixed

samples were dehydrated using a graded series of ethanol washes (30%, 50%, 70%, 80%, 90%

and 2 × 100%) for 5 mins in each solution change. Samples were then air-dried using

hexamethyldisilazane evaporative dehydration. Samples were then mounted onto SEM stubs,

platinum coated and viewed using a FEI Nova NanoSEM 230 (Oregon, USA) operating at 5kV.

Examination of the presence of *csep1* gene, plasmids pSma1 and pICON in GS1 and GS2

***C. concisus* strains**

The *C. concisus csep1* gene, plasmids pSma1 and pICON were previously found to be

associated with IBD, including both CD and UC [11, 21]. Their presence in the genomes of 85

GS1 and 160 GS2 strains, as well as in other *Campylobacter* species was examined by NCBI

nucleotide BLAST [22].

Growth characters of GS2 *C. concisus* strains and catalase, oxidase and urease tests

Growth of 10 GS2 *C. concisus* strains with complete sequenced genomes (Supplementary

Table 1) under different conditions, including incubation temperature, atmospheric conditions,

resistance to bile were examined. Gram staining was performed. Catalase, oxidase and urease

tests were also performed.

RESULTS

***C. concisus* GS1 and GS2 strains can be separated based on core genome and 23S rRNA**

gene

The core genome of 245 *C. concisus* strains contained 675 genes, as determined by Roary

default setting (the gene is present in more than or equal to 99% of strains). The 85 GS1 *C.*

concisus strains and 160 GS2 *C. concisus* strains were clearly separated based on the sequences

of core genome (Figure 1). The GS2 *C. concisus* strains contained two clusters, most of the

strains belonged to cluster 1. The 23S rRNA gene also consistently separated GS1 and GS2 *C.*

concisus strains (Figure 1). However, the sequences of 16S rRNA genes were unable to

differentiate GS1 and GS2 *C. concisus* (Supplementary Figure 1).

The average GC content of the 85 GS1 *C. concisus* strains was 37.3%, ranging from 34.7 to

38.2%. The average GC content of the 160 GS2 *C. concisus* strains was 39.2%, ranging from

36.4 to 40.3%.

The DDH and ANI values between GS1 and GS2 *C. concisus* strains were below the

recommended cut off value for defining a bacterial species

The DDH values between each of the 85 GS1 *C. concisus* strains and each of the 160 GS2 *C.*

concisus strains were all below 70%. The average DDH value between GS1 and GS2 *C.*

concisus strains was 54.9%, ranging from 42.8 to 66% (Table 1).

The average ANI value between each of the 85 GS1 *C. concisus* strains and each of the 160

GS2 *C. concisus* strains was 88.7%, ranging from 87.8 to 89.7% (Table 1).

The DDH values between all GS1 *C. concisus* strains and the *C. concisus* type strain

ATCC 33237

The DDH values between the *C. concisus* type strain ATCC 33237 and the 84 remaining GS1

strains were all ≥69% (Table 1). The average DDH values between the 84 GS1 *C. concisus*

strains and the *C. concisus* type strain ATCC 33237 was 82.9%, ranging from 74.5 to 99.2 %

(Table 1).

The ANI values between the 84 GS1 *C. concisus* strains and the *C. concisus* type strain ATCC

33237 ranged from 93.83 to 99.99% (Table 1).

The DDH values between all GS2 *C. concisus* strains and the *C. concisus* type strain

ATCC 33237 were below 70%

The DDH values between the 160 GS2 strains and the *C. concisus* type strain ATCC 33237

were all below 70%, ranging from 50.5 to 63.7 %. The ANI values between the 160 GS2 *C.*

concisus strains and the *C. concisus* type strain ATCC 33237 ranged from 88.41 to 89.15%

(Table 1).

Both GS1 and GS2 *C. concisus* strains carry GS-specific genes

Both GS1 and GS2 *C. concisus* strains contained their respective specific genes. Four GS1 and

three GS2 *C. concisus* specific genes were identified (Table 2).

MALDI-TOF MS profiles were different between GS1 and GS2 *C. concisus* strains

On MALDI-TOF MS analysis, protein peaks specific to GS1 or GS2 *C. concisus* strains were

identified. Two protein peaks, at m/z 7401 and 7557, were found to be present only in GS1 *C.*

concisus strains, with the prevalence being 80 and 60% respectively (Table 3). Four protein

peaks with m/z 6924, 7424, 8045 and 10359, were present only in GS2 *C. concisus* strains,

with the prevalence being 90%, 90%, 50% and 50% respectively (Table 3). Combination of

m/z 6924 with any of the other three protein peaks can 100% differentiate GS2 from GS1

strains.

GS1 and GS2 *C. concisus* strains showed difference in morphology

Electron microscopy examination showed that strains of GS1 and GS2 *C. concisus* differ in

morphology. GS1 *C. concisus* strains were straight to curved rods and the shapes of GS2 *C.*

concisus strains were spiral to curved (Figure 2).

The Crohn's disease associated *csep1* gene and plasmids were predominately present in

GS2 *C. concisus* strains

The Crohn's disease associated *csep1* gene was found in the chromosome of 38.75% (62/160)

GS2 *C. concisus* strains and none of the GS1 strains. The Crohn's disease associated pICON

plasmid was found in two GS1 strains (2/85, 2.4%), four GS2 strains (4/160, 2.5%) and one

strain of *Campylobacter showae* (B32_SW) [23]. The severe ulcerative colitis associated

pSma1 plasmid was found in three GS1 strains (3/85, 3.5%) and 13 GS2 strains (13/160, 8.1%)

(Supplementary Table 1).

DISCUSSION

We previously reported that *C. concisus* consisted of GS1 and GS2 strains based on the

difference in the core genome, which was confirmed by other researchers [9-12]. Our findings

in this study show that GS2 *C. concisus* is a novel *Campylobacter* species, we propose naming

it *C. oralis,* whereas GS1 *C. concisus* solely remains as *C. concisus*.

The recommended cut off DDH and ANI values for defining bacterial species are 70% DDH

and 95% ANI [17, 24-26]. Both the DDH and ANI values between *C. oralis* and *C. concisus*

strains were clearly below these criteria (Table 1). Phenotypically, *C. oralis* has a spiral to

curved rod shape and *C. concisus* has a straight or curved rod shape under the culture conditions

that we have used (Figure 2). MALDI-TOF MS analysis identified *C. oralis* and *C. concisus*

specific protein peaks. Our study and the study by On et al. consistently identified four *C. oralis*

protein peaks with the m/z 6924, 7424, 8045 and 10359 (Table 3) [27]. Combination of these

protein peaks can 100% differentiate *C. oralis* from *C. concisus*. Based on our genome analysis,

C. oralis and *C. concisus* specific genes were not related to common metabolic pathways,

making biochemical differentiation difficult. PCR targeting 23S rRNA gene can also be used

as a quick method for differentiating *C. oralis* from *C. concisus* and the sequences of PCR

primers were reported in previous studies [28, 29]. The 16S rRNA gene cannot differentiate *C.*

oralis from *C. concisus* (Supplementary Figure 1).

We and other researchers previously reported the association between *C. concisus* and IBD [2-

5]. In these earlier studies, the identification of *C. concisus* was based on 16S rRNA gene, thus

both *C. oralis* and *C. concisus* were detected. Recently, we also reported the associations

between *csep1* gene, pICON and pSma1 plasmids with IBD [11, 21]. In this study, we found

that Crohn's disease associated *csep1* gene was present in 38.75% of *C. oralis* strains and none

of the *C. concisus* strains. The plasmid pICON is rare, we found it in only seven strains of three

Campylobacter species (four *C. oralis* strains, two *C. concisus* strains and one *C. showae*

strain), interestingly all these strains were isolated from patients with Crohn's disease [10, 21,

23]. The severe ulcerative colitis associated pSma1 plasmid was found in less than 5% of *C.*

concisus strains and 10% of *C. oralis* strains. These suggest that *C. oralis* may play a more

important role in the development of Crohn's disease than *C. concisus.*

In conclusion, ANI and DDH values clearly support that *C. concisus* GS2 as a novel

Campylobacter species, for which we propose the name *Campylobacter oralis.* The

identification of *C. concisus* and *C. oralis* specific genes, along with differences in their protein

profiles, morphology, core genome and 23S rRNA phylogeny, further support this

classification.

Description of *C. oralis*

C. oralis bacterial cells are Gram negative, with a curved to spiral shape, $0.5 - 0.8 \times 4 - 8$ μm,

having rounded ends, motile with darting motility by means of a single polar flagellum (Figure

2). Colonies are circular, with convex elevation, translucent, smooth margins, no pigmentation,

and no observable haemolysis on HBA plates. The growth and biochemical characteristics of

C. oralis are listed in Table 4.

P15UCO-S2 strain is the type strain of *C. oralis*. This strain was isolated from saliva of a patient

with ulcerative colitis and its complete genome has been sequenced [11]. The DDH values

between the 159 *C. oralis* strains and the *C. oralis* type strain P15UCO-S2 ranged from 66.2

to 98.7%. The ANI values between the 159 *C. oralis* strains and the type strain P15UCO-S2

ranged from 93.6 to 98.7%. Based on these analyses, we recommend 66.2% DDH and 93.6%

ANI with the type strain P15UCO-S2 as the cut off values for identifying *C. oralis*. These

values are able to exclude *C. concisus* strains, the DDH and ANI values between the 85 *C.*

concisus strains and the *C. oralis* type strain P15UCO-S2 were $53.4 - 64.9\%$ and $88.2 - 89.3\%$

respectively. The *C. oralis* type strain has been deposited in the DSMZ under the number DSM

115209.

C. oralis was most closely related to *C. concisus* (Supplementary Table 2). *C. oralis* and *C.*

concisus can be differentiated based on the core genome sequence, 23S rRNA gene, DDH

value, ANI value and species-specific genes. *C. oralis* and *C. concisus* show different

morphology under electron microscope under certain culture conditions. PCR targeting 23S

rRNA gene can be used to differentiate *C. oralis* and *C. concisus* in clinical laboratories.

MALDI-TOF MS is a method that can be potentially used to differentiate *C. oralis* and *C.*

concisus.

TABLES

Table 1 DDH and ANI values obtained from analyses of GS1 and GS2 *C. concisus* genomes

		DDH	ANI
85 GS1 vs 160 GS2 strains	Max	66	89.69
	Min	42.8	87.83
	Mean	54.9	88.69
84 GS1 strains versus ATCC 33237	Max	99.2	99.56
	Min	74.5	94.01
	Mean	82.9	94.74
160 GS2 strains versus ATCC 33237	Max	63.7	89.15
	Min	50.5	88.41
	Mean	57.4	88.73

GS: genomospecies. ATCC 33237: *C. concisus* type strain described by Tanner et al [1]

Table 2 GS1 and GS2 *C. concisus* specific genes

GS	Gene	Annotation	Gene locus	Protein length (AA)
1	*tehA*	Tellurite resistance protein TehA	CVT06_06385	330
	abgT	p-aminobenzoyl-glutamate transport protein	CVT06_04695	508
	pstS	Phosphate-binding protein PstS	CVT06_05910	346
2	*hcp*	Hydroxylamine reductase	CCC13826_1540	550
	aqpZ	Aquaporin Z	CCC13826_1636	231

GS: genomospecies. AA: amino acid. GS specific genes: present in all the strains from one GS and absent in all the strains from the other GS.

282 **Table 3 MALDI-TOF MS markers differentiating GS1 and GS2 *C. concisus***

Strain ID	GS	Average m/z					
		6924.389	7401.524	7424.504	7557.485	8045.274	10359.59
P10CDO-S2	1		7392.891				
P26UCO-S2	1		7409.682		7558.543		
H25O-S1	1						
P3UCO1	1		7411.752		7556.284		
P19CDO-S1	1				7558.897		
H17O-S1	1		7373.999				
H28O-S1	1		7406.009				
H1O1	1		7404.886		7555.046		
P27CDO-S2	1		7403.849		7557.604		
P28CDO-S1	1		7409.12		7558.533		
P1CDO2	2	6924.038		7426.252		8044.699	
P1CDO3	2	6926.038					10359.56
P15UCO-S2	2	6924.789		7425.441			
P2CDO4	2	6923.513		7424.174		8044.692	10359.5
H14O-S1	2			7424.693		8045.149	10358.19
P16UCO-S2	2	6923.745		7426.193		8045.902	10360.75
H19O-S1	2	6924.265		7422.043			10359.98
P20CDO-S2	2	6921.875		7424.294			
13826	2	6926.661		7423.399		8045.93	
P13UCO-S1	2	6924.579		7424.045			

283 GS: genomospecies

284   **Table 4 The growth and biochemical characters of _C. oralis_**

Conditions	Growth
37 ºC	Growth
42 ºC	Growth
Microaerobic conditions (generated using Oxford gas-pak)	No growth
Anaerobic conditions (generated using Oxford gas-pak)	Slow growth
Microaerobic conditions with 5 -10% hydrogen	Growth
Anaerobic conditions with 5 -10% hydrogen	Growth
Aerobic conditions containing 10% CO2	No growth
HBA plates containing 1% ox bile	Growth
HBA plates containing 2% ox bile	Growth
MacConkey agar	Growth
HBA plates containing 0.4% Sodium fumarate	Increased growth
HBA plates containing 1% NaCl	Growth
HBA plates containing 2% NaCl	Growth
Catalase	Negative
Oxidase	Positive
Urease	Negative

285   _C. oralis_ and _C. concisus_ displayed same growth and biochemical characters when same
286   parameters were used.

287   **FIGURES**

Figure 1

A   **B**

Figure 1 Phylogenetic trees of GS1 and GS2 *C. concisus* based on sequences of *C. concisus*

core genome and the 23S rRNA gene. *C. concisus* core genome and 23S rRNA were able to

291   separate GS1 and GS2 strains. **A:** Core genome was generated by Roary (v3.12.0) and the

292   approximate maximum-likelihood tree was generated by FastTree (v2.1.10). **B:** 23S rRNA

293   maximum-likelihood tree (100 bootstrap) was constructed with MEGAXcc. Strains P21CDO-

294   S2 and H25O-S1 were removed for the 23S rRNA phylogenetic tree due to incomplete 23S

295   rRNA sequences.

Figure 2

Figure 2 Scanning electron microscope images showing the different morphologies of GS1
and GS2 *C. concisus.* **A:** *C. concisus* strains were cultured on HBA plates for 24 hours
followed by cultivation in heart infusion broth with 5% fetal bovine serum for 12 hours.

300   Bacterial cells were then subjected to scanning electron microscopic examination. Top row:

301   GS1 strains including P26UCO-S2, P10CDO-S2 and P3UCO1. Bottom row: GS2 strains

302   including BEO2, P15UCO-S2 and P2CDO4. Magnification: 30000×, scale bars represent 2

303   μm. **B:** Showing the single polar flagellum of *C. oralis*. Magnification:500000×. GS:

304   genomospecies.

AUTHOR CONTRIBUTION:

Fang Liu: Conducted analysis of *csep1* gene and plasmids. Performed confirmation analysis on GS specific genes.

Siying Chen: Performed genome analysis, most of the experiments and prepared the type strain for submission to ATCC.

Joanna M Biazik: Generated SEM pictures for Figure 2A using bacterial strains cultured by Siying Chen.

Christopher Luk: Conducted catalase, oxidase and urease tests.

Anjaneyaswamy Ravipati: Generated data for Table 3 using bacterial strains cultured by Siying Chen.

Ruiting Lan: Provided critical discussion during this project.

Li Zhang and Stephen M Riordan: Conceived the project and Li Zhang supervised the project.

Li Zhang, Fang Liu and Siying Chen played a major role in writing the manuscript. All authors have read the manuscript and provided feedback.

ACKNOWLEDGEMENT:

Picture in Figure 2B was generated by Vikneswari Mahendran during her study for a Masters by Research degree in Associate Professor Li Zhang group at the University of New South Wales in 2011.

FUNDING SUPPORT:

This work is supported by a Faculty Research Grant awarded to LZ from the University of New South Wales.

CONFLICTS OF INTEREST

330   The authors declare that there are no conflicts of interest

REFERENCE:

1. Tanner, A.C.R., et al., *Wolinella gen. nov., WoZinella succinogenes (Vibrio succinogenes Wolin et al.) comb. nov., and description of Bacteroides gracilis sp. nov., Wolinella recta sp. nov., Campylobacter concisus sp. nov., and Eikenella corrodens from Humans with Periodontal Disease.* Int J Syst Bacteriol, 1981(ll): p. 432-445.

2. Zhang, L., et al., *Detection and isolation of Campylobacter species other than C. jejuni from children with Crohn's disease.* J Clin Microbiol, 2009. **47**(2): p. 453-455.

3. Mukhopadhya, I., et al., *Detection of Campylobacter concisus and other Campylobacter species in colonic biopsies from adults with ulcerative colitis.* PLOS ONE, 2011. **6**(6): p. e21490-e21490.

4. Mahendran, V., et al., *Prevalence of Campylobacter species in adult Crohn's disease and the preferential colonization sites of Campylobacter species in the human intestine.* PLOS ONE, 2011. **6**(9): p. e25417.

5. Kirk, K.F., et al., *Optimized cultivation of Campylobacter concisus from gut mucosal biopsies in inflammatory bowel disease.* Gut Pathog, 2016. **8**: p. 27.

6. Lastovica, A.J. and E. Roux, *Efficient isolation of Campylobacteria from stools.* J Clin Microbiol, 2000. **38**(7): p. 2798-2800.

7. Macfarlane, S., et al., *Microbial colonization of the upper gastrointestinal tract in patients with Barrett's esophagus.* Clin Infect Dis, 2007. **45**(1): p. 29-38.

8. Zhang, L., et al., *Isolation and detection of Campylobacter concisus from saliva of healthy individuals and patients with inflammatory bowel disease.* J Clin Microbiol, 2010. **48**(8): p. 2965-7.

9. Chung, H.K., et al., *Genome analysis of Campylobacter concisus strains from patients with inflammatory bowel disease and gastroenteritis provides new insights into pathogenicity.* Sci Rep, 2016. **6**: p. 38442.

10.   Kirk, K.F., et al., *Molecular epidemiology and comparative genomics of Campylobacter concisus strains from saliva, faeces and gut mucosal biopsies in inflammatory bowel disease.* Sci Rep, 2018. **8**(1): p. 1902.

11.   Liu, F., et al., *Analysis of complete Campylobacter concisus genomes identifies genomospecies features, secretion systems and novel plasmids and their association with severe ulcerative colitis.* Microb Genom, 2020. **6**(11): p. mgen000457.

12.   Cornelius, A.J., et al., *Genetic characterisation of Campylobacter concisus: Strategies for improved genomospecies discrimination.* Syst Appl Microbiol, 2021. **44**(3): p. 126187.

13.   Page, A.J., et al., *Roary: rapid large-scale prokaryote pan genome analysis.* Bioinformatics, 2015. **31**(22): p. 3691-3693.

14.   Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree: computing large minimum evolution trees with profiles instead of a distance matrix.* Mol Biol Evol, 2009. **26**(7): p. 1641-1650.

15.   Kumar, S., et al., *MEGA-CC: computing core of molecular evolutionary genetics analysis program for automated and iterative data analysis.* Bioinformatics, 2012. **28**(20): p. 2685-2686.

16.   Meier-Kolthoff, J.P., et al., *Genome sequence-based species delimitation with confidence intervals and improved distance functions.* BMC Bioinformatics, 2013. **14**(1): p. 60.

17.   Jain, C., et al., *High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.* Nat Commun, 2018. **9**(1): p. 5114.

18.   Shen, W., et al., *SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file manipulation.* PLOS ONE, 2016. **11**(10): p. e0163962.

19.   Brynildsrud, O., et al., *Rapid scoring of genes in microbial pan-genome-wide*
association studies with Scoary. Genome Biol, 2016. **17**(1): p. 1-9.

20.   Lee, H., et al., *Examination of the anaerobic growth of Campylobacter concisus strains.*
Int J Microbiol, 2014. **2014**: p. 476047.

21.   Liu, F., et al., *Genomic analysis of oral Campylobacter concisus strains identified a*
potential bacterial molecular marker associated with active Crohn's disease. Emerg
Microbes Infect, 2018. **7**(1): p. 64.

22.   Johnson, M., et al., *NCBI BLAST: a better web interface.* Nucleic Acids Res, 2008.
36(suppl 2): p. W5-W9.

23.   Hsu, T., et al., *Comparative genomics and genome biology of Campylobacter showae.*
Emerg Microbes Infect, 2019. **8**(1): p. 827-840.

24.   Wayne, L.G., *International Committee on Systematic Bacteriology: Announcement of*
the Report of the Ad Hoc Committee on Reconciliation of Approaches to Bacterial
Systematics. Syst Appl Microbiol, 1988. **10**(2): p. 99-100.

25.   Goris, J., et al., *DNA-DNA hybridization values and their relationship to whole-genome*
sequence similarities. Int J Syst Evol Microbiol, 2007. **57**(1): p. 81-91.

26.   Richter, M. and R. Rossello-Mora, *Shifting the Genomic Gold Standard for the*
Prokaryotic Species Definition. Proc Natl Acad Sci U S A, 2009. **106**(45): p. 19126-
19131.

27.   On, S.L., et al., *Markers for discriminating Campylobacter concisus genomospecies*
using MALDI-TOF analysis. CRMS, 2021. **2**: p. 100019.

28.   Bastyns, K., et al., *Specific detection of Campylobacter concisus by PCR amplification*
of 23S rDNA areas. Mol Cell Probes, 1995. **9**(4): p. 247-50.

29.   Wang, Y., et al., *Campylobacter concisus genomospecies 2 is better adapted to the human gastrointestinal tract as compared with Campylobacter concisus genomospecies 1*. Front Physiol, 2017. **8**: p. 543.