Mycobacterial Metallophosphatase MmpE Acts as a Nucleomodulin to Regulate Host Gene Expression and Promote Intracellular Survival

Liu Chen[1,2], Baojie Duan[1], Qiang Jiang[1], Yifan Wang[1], Yingyu Chen[1,3], Lei Zhang[1,3]* and Aizhen Guo[1,2,3]*

[1]   National Key Laboratory of Agricultural Microbiology, College of Veterinary Medicine, Huazhong Agricultural University, Wuhan 430070, China

[2]   Hubei Hongshan Laboratory, Huazhong Agricultural University, Wuhan 430070, China

[3]   National Professional Laboratory for Animal Tuberculosis, Ministry of Agriculture and Rural Affairs, Huazhong Agricultural University, Wuhan 430070, China

*Address correspondence to: Lei Zhang, zhanglei2023@mail.hzau.edu.cn; Aizhen Guo, aizhen@mail.hzau.edu.cn

Abstract

Mycobacterium tuberculosis, the causative agent of tuberculosis, remains a major global health challenge. Nucleomodulins, bacterial effectors that target the host cell nuclei, are increasingly recognized as key virulence factors, but their roles in mycobacterial pathogenesis remain incompletely elucidated. Here, we characterize a hypothetical protein Rv2577 (designated MmpE) not only as a Fe^{3+}/Zn^{2+}-dependent metallophosphatase but also as a critical nucleomodulin involved in immune evasion and intracellular persistence. MmpE utilizes two nuclear localization signals, RRR[20–22] and RRK[460–462], to enter the host cell nucleus, where it binds to the promoter region of the vitamin D receptor (VDR) gene, thereby inhibiting host inflammatory gene expression. Additionally, MmpE represses the PI3K–Akt–mTOR signaling pathway, which arrests phagosome maturation. These actions collectively facilitate immune suppression and promote mycobacterial survival in macrophages and in mice. Our findings identify MmpE as a conserved nucleomodulin in mycobacteria and reveal a novel mechanism of MmpE-mediated intracellular survival.

Keywords Mycobacterium; Nucleomodulin; Metallophosphatase; Immune evasion; Intracellular survival.

Introduction

Mycobacterium tuberculosis (Mtb), the etiological agent of tuberculosis, has posed a persistent global health burden for centuries. Despite significant advances in elucidating the molecular mechanisms of its pathogenesis, current research has largely focused on effector proteins that modulate host cell membrane and cytoplasmic processes (Nisa *et al.,* 2022; Nasiri and Venketaraman, 2025). In contrast, investigations into effector proteins that target the host cell nucleus, the central hub for gene regulation and cellular control, remain comparatively limited.

Nucleomodulins are a class of bacterial effector proteins that can translocate into the host nucleus, where they modulate nuclear processes to promote pathogen survival (Chai *et al.,* 2020; Bierne and Cossart, 2021). These proteins affect host gene

expression by directly interacting with chromatin or by mimicking transcription factors,

chromatin modifiers, or other nuclear regulators (Li *et al.,* 2013; Rolando *et al.,* 2013;

Mitra *et al.,* 2018; Hanford *et al.,* 2021). For instance, LntA from *Listeria*

monocytogenes directly binds the chromatin repressor BAHD1, disrupting its function

and derepressing interferon-stimulated genes in a type III interferon-dependent

manner, ultimately promoting bacterial persistence (Lebreton *et al.,* 2011; Lebreton *et*

al., 2014). Similarly, Ank1 and Ank6 from *Orientia tsutsugamushi* localize to the host

nucleus and facilitate the export of the NF-κB p65 subunit via exportin 1, thereby

suppressing the transcription of pro-inflammatory genes (Evans *et al.,* 2018; Steiert

and Weber, 2025). Although bacterial nucleomodulins utilize diverse strategies to

manipulate host nuclear functions, the identity and functional characterization of *M.*

tuberculosis nucleomodulins remain poorly defined.

Nucleomodulins employ various strategies to enter access to the host cell nucleus,

including passive diffusion, hijacking host proteins that contain nuclear localization

signals (NLSs), or using of their own NLSs to interact with the nuclear pore complex

(Hanford *et al.,* 2021). Among these mechanisms, NLSs play a particularly important

role. Based on their amino acid composition, NLSs are broadly classified into classical

(cNLS), non-classical (ncNLS), and other types (Lange *et al.,* 2008; Lu *et al.,* 2021).

Classical NLSs are further divided into monopartite (MP) and bipartite (BP) forms.

Monopartite NLSs consist of a single cluster of 4–8 basic residues, primarily lysine (K)

or arginine (R), with a typical consensus motif of K-K/R-X-K/R (Fontes *et al.,* 2000;

Kosugi *et al.,* 2009; Cautain *et al.,* 2015). Nuclear localization enables these effectors

to manipulate host gene expression, disrupt immune signaling, and reprogram cellular

processes to promote pathogen survival. For example, EspF from enteropathogenic

Escherichia coli contains an N-terminal NLS that directs it to the nucleolus, where it

interferes with ribosome biogenesis (Dean *et al.,* 2010; Singh *et al.,* 2018). NUE, a

SET domain-containing nucleomodulin secreted by *Chlamydia trachomatis*, also

possesses an NLS that mediates its nuclear localization, where it functions as a

histone lysine methyltransferase (HKMTase) targeting host histones H2B, H3, and H4

(Pennini *et al.,* 2010; Jermy, 2010).

Beyond their nuclear localization capabilities, some nucleomodulins directly affect host

immunity through intrinsic enzymatic activities (Agarwal *et al.,* 2012). For example, the

nucleomodulin PtpA promotes Mtb survival within macrophages by dephosphorylating

host cytoplasmic proteins (Wang *et al.,* 2017). NleC from *E. coli*, a Zn-dependent

metalloprotease, suppresses host immune responses by cleaving the p65 subunit of

NF-κB in the cytoplasm, thereby inhibiting its nuclear translocation, and can enter the

nucleus to reduce nuclear p65 levels, blocking NF-κB-dependent transcription of pro-

inflammatory genes (Baruch *et al.,* 2011; Hodgson *et al.,* 2015). Among these

enzymatically active nucleomodulins, purple acid phosphatases (PAPs) represent a

distinct and functionally significant, yet underexplored, subclass within the

metallophosphatase superfamily. PAPs are characterized by a conserved β-α-β-α-β

fold and five signature motifs (DxG, GDXXY, GNH[D/E], VXXH, GHXH), which

coordinate seven metal-ligating residues at a binuclear metal center (Rodriguez *et al.,*

2014; Bhadouria *et al.,* 2017; Bhadouria and Giri, 2022). These enzymes have been

extensively studied in plants and mammals, where they are involved in key biological

processes such as phosphorus metabolism and the generation of reactive oxygen

species (Comba *et al.,* 2013; Feder *et al.,* 2020; Liu *et al.,* 2025). In contrast, only a

few bacterial PAPs have been identified, such as BcPAP from *Burkholderia*

cenocepacia, and their functions remain largely unknown (Yeung *et al.,* 2009). Notably,

in Mtb, MmpE is the only identified PAP protein (Schenk *et al.,* 2000), but its specific

contribution to infection and pathogenesis remain elusive.

In this study, we characterized MmpE as a conserved nucleomodulin in mycobacteria

and demonstrated its nuclear translocation mediated by two NLSs, RRR[20–22] and

RRK[460–462]. Upon entering the nuclear, MmpE binds to the promoter region of the VDR

gene, a key regulator of antimicrobial responses, resulting in transcriptional repression

of pro-inflammatory genes. In addition, MmpE inhibits phagosome maturation by

suppressing the expression of genes in the PI3K–Akt–mTOR signaling pathway. These

actions collectively facilitate increased intracellular survival in both macrophage and

mouse. Beyond its nuclear function, MmpE functions as a Fe^{3+}/Zn^{2+}

metallophosphatase, with enzymatic activity reliant on the conserved residues His^{348}

and Asp^{359}, and partially contributes to intracellular survival. Taken together, our

findings identify MmpE as a bifunctional effector that integrates immune modulation

and enzymatic catalysis, underscoring its role in mycobacterial pathogenesis and its

potential as a therapeutic target.

Results

NLSs are required for the nuclear translocation of MmpE

In the prior study, we systematically identified some conserved nucleomodulins in

mycobacterium. Among them, the hypothetical protein Rv2577 (MmpE) was identified

as a candidate exhibiting strong nuclear localization in mammalian cells (Chen *et al.,*

2025). Based on these findings, we transfected HEK293T cells with a C-terminal

EGFP-tagged MmpE (MmpE-EGFP) and monitored its transcriptional level and

nuclear localization at different time points. As a result, a time-dependent increase in

the MmpE-EGFP content was observed in the host nucleus (Figure 1A, upper panel;

Figure S1A). Nuclear–cytoplasmic fractionation assays further confirmed these results,

showing a progressive enrichment of MmpE in the nuclear compartment over the

course of transfection (Figure 1A, bottom panel). Collectively, these results indicate

that MmpE possesses intrinsic nuclear translocation ability.

MmpE contains two putative NLSs, N-terminus ($RRR^{20–22}$, NLS1) and C-terminus

($RRK^{460-462}$, NLS2) (Figure 1B). To evaluate the structural contribution of these motifs,

we utilized AlphaFold to model the impact of NLS truncations on the overall

conformation of MmpE. The predicted structures indicated that deletion of either NLS1,

NLS2, or both did not obviously alter the the protein fold (Figure S1B). Based on this

analysis, we generated a series of NLS-deleted mutants, including $MmpE^{\Delta NLS1}$,

$MmpE^{\Delta NLS2}$, and $MmpE^{\Delta NLS1-2}$, to assess the functional relevance of each motif in

nuclear localization. Subcellular localization of wild-type MmpE-EGFP and NLS

deletion mutants was examined in HEK293T cells by fluorescence microscopy at 36

hpt (hours post-transfection). Deletion of either NLS1 or NLS2 alone significantly reduced the nuclear accumulation of MmpE-EGFP, with the effect of NLS2 deletion being more pronounced than that of NLS1. Simultaneous deletion of both NLS motifs (MmpE$^{\Delta NLS1\text{-}2}$) completely abolished nuclear localization, resulting in exclusive cytoplasmic distribution (Figure 1C and D). These findings were further validated by immunoblotting of nuclear and cytoplasmic fractions. Under the similar transfection conditions, wild-type MmpE-EGFP was detected in both compartments, whereas MmpE$^{\Delta NLS1}$ and MmpE$^{\Delta NLS2}$ exhibited substantially reduced nuclear abundance. By contrast, MmpE$^{\Delta NLS1\text{-}2}$ was exclusively localized to the cytoplasmic fraction, suggesting that both NLS motifs are required for efficient nuclear import (Figure 1E). Taken together, our findings identify MmpE as a nucleomodulin, with nuclear import mediated by the N- and C-terminal NLS motifs.

Deletion of NLSs does not alter the phosphatase activity of MmpE

Bioinformatic analyses identified MmpE as a Fe^{3+}/Zn^{2+}-metallophosphatase containing a purple acid phosphatase (PAP) motif (Figure 2A; Figure S2A). Multiple sequence alignment and phylogenetic reconstruction revealed that MmpE is highly conserved among mycobacterial species and harbors canonical motifs characteristic of metallophosphatases, including GD, GNHX, and GHXH (Figure 2B, Figure S2B). These motifs are typically associated with enzymes that catalyze phosphate hydrolysis via a binuclear metal ion-dependent mechanism, wherein residues such as Asp, His, and Asn coordinate essential metal cofactors, commonly Fe^{3+}, Mn^{2+}, or Zn^{2+}, at the active site.

To identify key amino acid residues of MmpE involved in metal coordination, AlphaFold was used to predict the structure of MmpE, followed by the molecular docking using PyMOL. As shown in Figure 2C, H^{348} and N^{359} were identified as candidate residues likely involved in metal ion coordination within the predicted active site. Subsequently, recombinant MBP-tagged MmpE deleting the Tat signal peptide (MmpE$^{\Delta Tat}$) was subjected to *in vitro* phosphatase assays using *p*-nitrophenyl phosphate (*p*-NPP) as the substrate. Reactions were supplemented with varying concentrations of Fe^{3+} and

Zn^{2+} to evaluate their individual effects on MmpE enzymatic activity. The results

demonstrated that, at equivalent concentrations, Fe^{3+} significantly enhanced MmpE

activity compared to Zn^{2+}, with maximal enzymatic activity was observed at 50 μM Fe^{3+}

(Figure 2D). Next, a double mutant ($MmpE^{\Delta Tat}-H^{348}AN^{359}A$) was generated via site-

directed mutagenesis to validate the functional importance of these residues. As

shown in Figure 2E, the addition of 50 μM Fe^{3+} significantly increased the *in vitro*

phosphatase activity of $MmpE^{\Delta Tat}$ compared to reactions without metal ions. However,

the enzymatic activity of the $MmpE^{\Delta Tat}-H^{348}AN^{359}A$ in the presence of 50 μM Fe^{3+} was

markedly reduced and nearly indistinguishable from that of $MmpE^{\Delta Tat}$ under metal-free

conditions. These results confirm that residues H^{348} and N^{359} are essential for Fe^{3+}

binding and catalytic activity.

Given that MmpE relies on NLSs to enter the host nucleus, we examined whether

deletion of NLS affects its phosphatase activity. Compared to wild-type MmpE, the

NLS-deleted mutant showed no significant change in enzymatic activity (Figure 2F),

which is consistent with the structural comparison results (Figure S1B). Collectively,

these results suggest that MmpE is a highly conserved Fe^{3+}/Zn^{2+}-dependent

metallophosphatase in mycobacteria, and its enzymatic activity is independent of NLS.

The nuclear translocation and phosphatase activity of MmpE are essential for *M.*

***bovis* BCG survival in macrophage cells**

To investigate the role of MmpE during infection, we first generated recombinant *M.*

bovis BCG strains expressing C-terminally Flag-tagged Ag85B (positive control) and

MmpE. Immunoblot analysis confirmed high levels of Ag85B-2×Flag and MmpE-

2×Flag in bacterial lysates, and both proteins were also detected in the culture

supernatants (Figure S3A), indicating that MmpE is indeed a secreted protein.

Furthermore, a series of recombinant BCG strains were constructed, including wild-

type (WT), *mmpE*-deleted mutant (ΔMmpE) (Figure S3B), ΔMmpE complemented with

wild-type *mmpE* (Comp-MmpE), and ΔMmpE complemented with NLS-deleted mutant

(Comp-$MmpE^{\Delta NLS1}$, Comp-$MmpE^{\Delta NLS2}$, and Comp-$MmpE^{\Delta NLS1-2}$). Deletion of NLS1

may reduce or abolish MmpE secretion (Stanley *et al.,* 2000; Buchanan *et al.,* 2001),

while deletion of NLS2 impaired its nuclear localization. All strains exhibited similar

growth rate under the normal culture conditions (Figure S3C). Subsequently, these

strains were used to infect THP-1 macrophages. As shown in Figure 3A and B, the

bacterial load of the ΔMmpE strain exhibited significantly reduced relative to the WT

strain, and bacterial clearance was accelerated over time. The phenotype of ΔMmpE

was rescued by complementation with wild-type gene *mmpE*. Meanwhile, the load of

the Comp-MmpE$^{\Delta NLS2}$ strain displayed significantly lower compared to Comp-MmpE in

both THP-1 and RAW264.7 macrophages, indicating that nuclear localization mediated

by NLS2 is critical for MmpE function. Notably, the Comp-MmpE$^{\Delta NLS1}$ and Comp-

MmpE$^{\Delta NLS1-2}$ strains demonstrated obvious higher load than those of the ΔMmpE strain,

yet lower than those of the Comp-MmpE strain (Figure 3C and D). These results

suggest that MmpE contributes to virulence not only through nuclear modulation but

also through additional mechanisms.

Next, we investigated the role of the Fe^{3+}/Zn^{2+}-dependent metallophosphatase of

MmpE on intracellular survival. As shown in Figure S3D, a double mutant strain (Comp-

MmpE-H^{348}AN^{359}A) was constructed and used to infect macrophage cells. The results

showed that this mutant exhibited significantly reduced intracellular load compared

with the Comp-MmpE strain (Figure 3E), indicating that the phosphatase activity also

plays an essential role in MmpE-mediated bacterial survival. To reveal the underlying

mechanism, cytokine expression assays were conducted in infected macrophages.

Compared with the WT and Comp-MmpE strains, cells infected with the ΔMmpE and

Comp-MmpE$^{\Delta NLS1-2}$ strains showed markedly elevated levels of pro-inflammatory

cytokines, including *IL1A*, *IL1B*, and *IL6* (Figure 3F).

Collectively, these results suggest that MmpE functions as a bifunctional effector,

possessing both nuclear import capability and phosphatase activity, promoting immune

evasion and enhancing intracellular survival during infection.

MmpE modulates host transcription network involved in inflammation response

and lysosomal maturation

To investigate the role of the nucleomodulin MmpE on host gene expression, we

performed RNA-seq assays on THP-1 macrophages infected with either wild-type

(BCG/wt) or MmpE-deleted (Ko*mmpE*) strains at 20 hpi (hours post-infection). As a

result, a total of 175 differentially expressed genes (DEGs) were identified, with 142

genes upregulated and 33 genes downregulated in Ko*mmpE*-infected macrophages

compared to those infected with BCG/wt (Figure 4A, Table S1). These DEGs were

significantly enriched in immune-related biological processes, particularly those

associated with apoptosis and cell death, as well as molecular functions related to

immune activation and cellular stress responses, such as pyrophosphatase activity

and nucleoside-triphosphatase activity (Figure 4B). Further analysis using the Kyoto

Encyclopedia of Genes and Genomes (KEGG) revealed that the DEGs were primarily

involved in immune and inflammatory signaling pathways, including cytokine–cytokine

receptor interaction, TNF signaling, Toll-like receptor signaling, chemokine signaling,

NF-κB signaling, JAK–STAT signaling, and the PI3K−Akt signaling pathway.

Additionally, several DEGs were enriched in pathways related to cell survival and

apoptosis. Another group of DEGs was associated with host defense mechanisms, as

evidenced by their enrichment in the lysosome pathway (Figure 4C). Together, these

data highlight MmpE as a critical modulator of host transcriptional responses that

facilitate mycobacterial immune evasion and intracellular survival.

To gain further insight into the functional relevance of the 175 differentially expressed

genes, we performed protein–protein interaction (PPI) network analysis, focusing on

protein-coding genes, which accounted for 38% of the total (Figure S4A). A tightly

connected network centered around key immune regulatory, including IL23A*,* IL12B*,*

CSF2*,* CD69*,* IDO1, and CEACAM1, which are critically involved in inflammatory

cytokine production (Figure 4D). Consistently, these genes expression was

upregulation in Ko*mmpE*-infected macrophages (Figure 4E). Given that inflammatory

signaling and lysosomal function are both essential for macrophage-mediated host

defense against mycobacterial infection, we further examined the expression of

representative inflammatory cytokines and key lysosomal maturation markers under

the same experimental conditions, such as *TFEB*, *LAMP1*, *LAMP2*, and several V-ATPase subunit genes (*ATP6V0A1*, *ATP6V0C*, *ATP6V1A*, *ATP6V1B2*, and *ATP6V1E1*). Consistent with transcriptomic findings, the Ko*mmpE* strain induced significantly higher expression of both inflammatory cytokines and lysosomal maturation-related genes compared to the WT strain (Figure 4F and G). Both the Comp-MmpE$^{\Delta NLS2}$ and Comp-MmpE$^{\Delta NLS1-2}$ strains showed significantly higher expression of these cytokine genes compared to the Comp-MmpE strain (Figure S4B and C). Taken together, these results suggest that MmpE manipulates host transcriptional networks governing inflammation and lysosomal function, thereby subverting antimicrobial immunity.

MmpE suppresses the expression of *VDR* and inhibits activation of the PI3K–Akt–mTOR signaling pathway

To elucidate the mechanism by which MmpE modulates host gene expression, we performed ChIP-seq in HEK293T cells transfected with EGFP-MmpE. 2,903 potential MmpE-binding sites (Table S2) were identified, with 18.49% (537) in intergenic regions and 39.44% (1145) in intragenic regions (Figure 5A). Chromosomal distribution revealed significant enrichment on chromosomes 1 and 2 (Figure 5B). Notably, 99.7% (2894) of these binding sites were associated with protein-coding regions. Among these, 1013 sites were located within 3 kb of transcription start sites (TSS), including promoter (≤1 kb, 63.18%), promoter (1–2 kb, 22.9%), and promoter (2–3 kb, 13.92%) (Figure 5C and D), suggesting that MmpE primarily targets promoter regions to regulate transcription. PPI analysis of the 1013 potential target genes revealed significant enrichment in protein kinase-related genes, such as *PRKCB*, *PLCG2* and *PIK3CB*, involved in regulating immune activation and inflammatory responses via the PI3K–Akt and mTOR signaling pathways, suggesting that MmpE may influence cell immune responses through a coordinated network of signaling pathways (Figure S5A and B). In parallel, we examined the expression of key components in this signaling axis in THP-1 macrophages infected with either BCG/wt or the Ko*mmpE* strains. Notably, Ko*mmpE* infection led to a significant upregulation of genes associated with

the pathway, including *PIK3CA*, *PIK3CB*, *PIK3R1*, *mTOR*, *AKT1–3*, and *FOXO1/3*

(Figure S5C). These genes encode critical regulators of cell survival and immune

activation within the PI3K–Akt–mTOR axis.

To identify the direct target gene of MmpE in macrophages, we focused on analysis of

the promoter regions (≤1 kb upstream of the transcription start site) of significantly

upregulated genes with | $Log_2FC > 1$| and *P* < 0.05. Four transcription factors were

identified, including CREBBP, VDR, GREB1, and FOXP4 (Figure 5E). Integrative

analysis using HOMER *de novo* motif prediction revealed that only the promoter region

of *VDR*, a key anti-inflammatory gene (Aggeletopoulou *et al.,* 2022), harbored a

potential MmpE-binding motif (Figure 5F). These findings suggest that MmpE may

directly bind to the *VDR* promoter. To validate the hypothesis, we examined *VDR*

expression in macrophages infected with recombinant strains. Compared to the

BCG/wt, *VDR* expression was significantly downregulated in THP-1 cells infected with

the Ko*mmpE* strain (Figure S5D). To further test whether MmpE directly binds the *VDR*

promoter, we cloned the *VDR* promoter region (−100 bp to +50 bp) and performed

chromatin immunoprecipitation (ChIP) assays in HEK293T cells. While both *VDR* and

GAPDH (control) sequences were present in input samples, *VDR* promoter DNA was

detected exclusively in ChIP samples from EGFP-MmpE transfected cells (Figure 5G

and H). Furthermore, the DNA-binding capacity of MmpE was unaffected by its

phosphatase activity (Figure S5E and F). To confirm these results, we performed

electrophoretic mobility shift assays (EMSA) using the same *VDR* promoter fragment.

As shown in Figure 5I and 5J, the shift bands corresponding to MmpE and *VDR*

promoter complex were observed, and a corresponding increase was observed in the

percentage of protein/DNA complexes. Collectively, these results provide strong

evidence that MmpE directly interacts with the *VDR* promoter both in vivo and *in vitro*

to repress host inflammatory signaling pathways and facilitating immune evasion

during mycobacterial infection.

To further characterize the transcriptional programs regulated by MmpE, we performed

an integrative analysis of CUT&Tag and RNA-seq datasets. This analysis identified

298	298 DEGs with direct or proximal MmpE binding sites (Table S3; Figure S6A and B).

Notably, genes such as *RHEX*, *PLXDC1*, and *GAS6* exhibited significant expression

changes and were enriched in pathways associated with cell differentiation and small

GTPase-mediated signaling (Figure S6C and D). These pathways play critical roles in

shaping immune cell function, trafficking, and inflammatory responses during host–

pathogen interactions (Li *et al.,* 2023; Ma *et al.,* 2024). Taken together, these data

establish MmpE as a nucleomodulin that directly alters host transcriptional networks

involved in immune regulation and cellular defense, thereby supporting mycobacterial

survival within host cells.

Nuclear translocation of MmpE is essential for mycobacterial survival in mice

Next, C57BL/6 mice were intranasally infected with *M. bovis* BCG strains, including

wild-type (WT), ΔMmpE, Comp-MmpE, and Comp-MmpE$^{\Delta NLS1-2}$, and systematically

assessed bacterial persistence, pulmonary inflammation, and cytokine dynamics. Our

results show that mice infected with the ΔMmpE strain exhibited significantly lower

bacterial loads in the infected mice lung compared to those infected with the WT or

Comp-MmpE strains. Remarkably, mice infected with the Comp-MmpE$^{\Delta NLS1-2}$ strain

showed significantly lower than those in the Comp-MmpE strains but higher than those

in the ΔMmpE strains (Figure 6A). A similar trend was observed in splenic bacterial

colonization (Figure S7). Hematoxylin and eosin (H&E) staining revealed extensive

inflammatory infiltrates in the lungs of mice infected with WT or Comp-MmpE, whereas

mice infected with ΔMmpE or Comp-MmpE$^{\Delta NLS1-2}$ exhibited markedly reduced

inflammation (Figure 6B). Mice infected with ΔMmpE and Comp-MmpE$^{\Delta NLS1-2}$ strains

exhibited significantly elevated levels of *Il1α*, *Il1β*, and *Il6* compared to those infected

with WT or Comp-MmpE strains, despite lower bacterial burdens (Figure 6C and E).

Collectively, these results demonstrate that MmpE indeed contributes to *M. bovis* BCG

survival *in vivo*, with its nuclear translocation being essential for mycobacterial

intracellular survival.

Discussion

MmpE Functions as a Bifunctional Protein with Nuclear Localization and Phosphatase Activity. In this study, we identified two independent NLSs within the MmpE protein, RRR[20–22] (NLS1) and RRK[460–462] (NLS2), which together form a bipartite NLS motif that is rarely observed in bacterial effector proteins. Loss of a single NLS significantly reduced the nuclear accumulation of MmpE, with NLS2 deletion causing a more pronounced effect than that of NLS1. Simultaneous deletion of both motifs completely abolished nuclear localization (Figure 1C-E). These findings indicate that NLS1 and NLS2 function in a synergistic manner, with NLS2 likely playing a dominant role in nuclear import. Further sequence analysis revealed that NLS1 (RRR[20–22]) is embedded within the conserved twin-arginine translocation (Tat) signal motif (S/TRR-X-FLK), a well-characterized determinant of Tat-dependent secretion (Gimenez *et al.,* 2018). The adjacent twin-arginine residues are critical for recognition by the Tat translocase, and mutations in this motif have been shown to significantly impair or abolish secretion (Ulfig *et al.,* 2017). Accordingly, the Comp-MmpE$^{\Delta NLS1}$ strain likely exhibits impaired secretion, and the Comp-MmpE$^{\Delta NLS2}$ strain is deficient in nuclear localization. Consistent with this, both the Comp-MmpE$^{\Delta NLS1}$ and Comp-MmpE$^{\Delta NLS2}$ strains exhibited lower intracellular survival than the Comp-MmpE strain and higher than that of the ΔMmpE strain (Figure 3A-D), suggesting that the nuclear localization ability of MmpE indeed contributes to bacterial survival. Interestingly, MmpE also functions as a Fe^{3+}/Zn^{2+}-dependent metallophosphatase, with residues H^{348} and N^{359} being essential for its catalytic activity (Figure 2D and E). The mutant strain (Comp-MmpE-$H^{348}AN^{359}A$) showed markedly reduced intracellular survival Compared to Comp-MmpE (Figure 3E), indicating that the phosphatase activity of MmpE is also required for intracellular survival. This explains why the intracellular survival of the Comp-MmpE$^{\Delta NLS1-2}$ strain is significantly lower than that of the Comp-MmpE strain, yet obviously higher than that of the ΔMmpE strain (Figure 3C and D, Figure 6A).

Taken together, these findings suggest that MmpE is a dual-function effector protein that promotes intracellular survival during infection through the coordinated action of its nuclear localization signal and phosphatase activity.

MmpE's direct targeting of the host gene *VDR* represents a unique immune evasion strategy. VDR is a critical nuclear receptor that controls the expression of numerous genes involved in host immune defense, including antimicrobial peptides and cytokines (Usategui-Martín *et al.,* 2022). For example, Mtb suppresses the expression of the host gene *CYP27B1* to reduce active vitamin D synthesis, thereby limiting VDR activation and downstream antimicrobial responses (Liu *et al.,* 2006). Similarly, *Salmonella* downregulates *VDR* through miRNA-mediated mechanisms, weakening mucosal immunity (Chen *et al.,* 2014). In this study, we found that MmpE directly binds to the promoter of the *VDR* gene (Figure 5F–J) and attenuates the expression of VDR-regulated inflammatory genes (Figure 3F; Figure 6C-E), ultimately promoting bacterial intracellular survival (Figure 6A). This mode of action represents a novel strategy employed by bacterial pathogens to evade host defenses through the direct manipulation of host gene promoters by nucleomodulins. Although previous studies have reported that bacterial nucleomodulins interfere with the host transcriptional machinery, these effects have typically occurred indirectly, such as through epigenetic modifications or inhibition of signaling pathways (Jose *et al.,* 2016; Singh *et al.,* 2023). Direct promoter binding by a nucleomodulin, however, remains relatively uncommon and represents a distinct mechanism of host gene regulation.

MmpE regulates the expression of host genes involved in phagosome maturation. As an intracellular pathogen, Mtb employs multiple strategies to evade phagolysosomal degradation, a process essential for bacterial clearance (Chandra *et al.,* 2022; Zhang *et al.,* 2023; Singh and Nagaraja, 2025). Improtantly, approximately 70% of phagosomes harboring Mtb fail to fuse with lysosomes, in part due to the activity of secreted phosphatases such as PtpA, PknG, and SapM, which disrupt vesicular trafficking (Wong *et al.,* 2011; Ge *et al.,* 2022; Zhang *et al.,* 2024; Alsayed and Gunosewoyo, 2024). MmpE has also been shown to contribute to the arrest of phagosome maturation in human cells, although the underlying mechanisms remain unclear (Forrellad *et al.,* 2020). In this study, we identify a potential mechanism by which MmpE inhibits phagosome maturation. Deletion of *mmpE* led to significant

upregulation of lysosome-associated genes, including *TFEB*, *LAMP1*, *LAMP2*, and V-

ATPase subunit genes (Figure 4C and G). Among these, TFEB is a master

transcriptional regulator of lysosomal biogenesis and function (Jeong *et al.,* 2021;

Wang *et al.,* 2024). Furthermore, ChIP-seq analysis revealed that MmpE directly binds

to promoter regions of host genes within the PI3K–Akt–mTOR signaling pathway

(Figure S5A), a key upstream regulator of TFEB activity (Fang *et al.,* 2021; Gounis *et*

al., 2025). Collectively, these results indicate that MmpE may act as a transcriptional

regulator that targets the PI3K–Akt–mTOR–TFEB axis, thereby inhibiting lysosomal

maturation.

MmpE is a Fe^{3+}/Zn^{2+}-dependent metallophosphatase that is essential for

mycobacterial intracellular survival. Previous studies have shown that

phosphatases play critical roles in host–pathogen interactions by modulating

phosphorylation-dependent signaling pathways (Sajid *et al.,* 2015; Bonne Køhler *et al.,*

2020). In bacterial pathogens, these enzymes frequently contribute to immune evasion

and intracellular survival. For example, the tyrosine phosphatase YopH from *Yersinia*

enterocolitica disrupts phagocytic signaling in host macrophages, thereby impairing

oxidative burst responses and promoting bacterial persistence (Davicino *et al.,* 2017;

Shaban *et al.,* 2020). Similarly, PpzA, a phosphatase from *Aspergillus fumigatus*, has

been implicated in fungal adaptation to immune pressure under iron-replete conditions

(Manfiolli *et al.,* 2017; Kurucz *et al.,* 2018). In this study, we found that MmpE contains

a predicted purple acid phosphatase (PAP)-like domain and exhibits phosphatase

activity that requires Fe^{3+} and Zn^{2+} ions as cofactors (Figure 2A and D). Through site-

directed mutagenesis, we identified His^{348} and Asn^{359} as critical residues for Fe^{3+}

coordination. Substitution of these residues ($H^{348}AN^{359}A$) significantly impaired

phosphatase function in the presence of Fe^{3+}, leading to a marked reduction in

bacterial intracellular load (Figure 2E; Figure 3E). However, the precise molecular

mechanism by which MmpE interacts with Fe^{3+} remains to be elucidated in future

studies.

In conclusion, our study identifies MmpE as a novel bifunctional virulence factor of

mycobacteria, combining Fe^{3+}/Zn^{2+}-dependent phosphatase activity with nuclear translocation capability. MmpE translocates into the host nucleus and directly binds to the promoters of some key host genes, such as *VDR*, thereby inhibiting the expression of the downstream targets implicated with the inflammation responses and lysosomal maturation, thereby promoting mycobacterial survival in macrophages and in mice (Figure 7). These findings reveal a sophisticated mechanism by which bacterial pathogen manipulates host gene transcription and signaling pathway to promote intracellular persistence.

Materials and methods

Key resources used in this study

Reagent or resource	Source	Identifier
Antibodies		
Anti-β-actin	Abbkine	Cat#ABL1010
Anti-Histone H3	Abbkine	Cat#ABL1070
Goat anti-rabbit IgG	Abbkine	Cat#A21020
Goat anti-mouse IgG	Abbkine	Cat#A25012
Anti-GFP	Proteintech	Cat#66002-1-Ig
Anti-Flag	Proteintech	Cat#66008-4-Ig
Cell and Mouse		
HEK293T	ATCC	Cat#CRL-3216
RAW264.7	ATCC	Cat#TIB-71
THP-1	ATCC	Cat#TIB-202
Mouse (C57BL/6)	Chang-sheng Bio (Liaoning, China)	
The plasmids used in this study are listed in Table S4		
The bacterial strains used in this study are listed in Table S5		
The primers used in this study are listed in Table S6		
Chemicals, Peptides, and Recombinant Proteins		
DMEM	Gibco	Cat#11965092
RPMI 1640	Gibco	Cat#C11875500BT
Opti-MEM™	Gibco	Cat#31985070
FBS	Gibco	Cat#A5256701
Sodium pyruvate	Gibco	Cat#11360070
L-glutamine	Gibco	Cat#A2916801

Key Resources Used in This Study (Continued)

Chemicals, Peptides, and Recombinant Proteins

HEPES	Gibco	Cat#15630080
2-Mercaptoethanol	Gibco	Cat#21985023
penicillin-streptomycin antibiotics	Gibco	Cat#15140122
PMA	Sigma-Aldrich	Cat#P8139
p-NPP	Sigma-Aldrich	Cat#4264-83-9
Formaldehyde	Sigma-Aldrich	Cat#252549
OADC	MilliporeSigma	Cat#M0678
DAPI	Beyotime	Cat# P0126
Triton-X-100	Beyotime	Cat#P0096
RIPA buffer	Beyotime	Cat#P0038
NP-40	Beyotime	Cat#P0013F
protease inhibitor cocktail	Beyotime	Cat#AR1182
HieffTrans® Liposomal Transfection Reagent	YEASEN	Cat#40802ES03
protein A/G magnetic beads	MedChemExpress	Cat#HY-K0202

Critical commercial assays

TRIpure Reagent	Aidlab	Cat#RN0101
EASYspin RNA Mini Kit	Aidlab	Cat#RN0702
EndoFreePlasmidMiniKit	Aidlab	Cat#PL0401
cDNA Reverse Transcription Kit	Vazyme	Cat#R333-01
2 × ChamQ Universal SYBR qPCR Master Mix	Vazyme	Cat#Q711-03
Uniclone One Step Seamless Cloning Kit	Genesand	Cat#SC612
Clarity™ Western ECL Substrate	BIO-RAD	Cat#1705060
Nuclear and Cytoplasmic Protein Extraction kit	Beyotime	Cat#P0027

Software and algorithms

jvenn	https://bioinfogp.cnb.csic.es/tools/venny/index.html
AlphaFold v2.2.0	https://github.com/deepmind/alphafold
iTOL	https://itol.embl.de/
ESPript 3.0	https://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi
GraphPad Prism 8	https://www.graphpad.com/
UCSF Chimera	http://www.cgl.ucsf.edu/chimera/
SignalP 5.0	https://services.healthtech.dtu.dk/services/SignalP-5.0/

Key Resources Used in This Study (Continued)	
Software and algorithms	
MEGA 12.0	https://www.megasoftware.net/
STRING 12.0	https://cn.string-db.org/
Cytoscape 3.10.3	https://cytoscape.org/
CaseViewer version 2.0	https://www.3dhistech.com/solutions/caseviewer/

Bacterial strains and culture conditions

The bacterial strains used in this study are detailed in Table S5. *E. coli* DH5α and BL21 were cultured in Luria-Bertani (LB) medium under standard conditions. *M. bovis BCG* strains were grown in Middlebrook 7H9 broth supplemented with 10% (v/v) OADC (oleic acid, albumin, dextrose, catalase), 0.05% (v/v) Tween-80, and 2% (v/v) glycerol. Antibiotics were added as required at the following final concentrations: 50 μg/mL hygromycin (for mycobacteria) and 50 μg/mL kanamycin (for both mycobacteria and *E. coli*).

Cell culture

THP-1 cells were cultured in Roswell Park memorial institute 1640 (RPMI 1640) medium supplemented with 10% (v/v) heat-inactivated fetal bovine serum (FBS), 1 mM sodium pyruvate, 2 mM L-glutamine, 10 mM HEPES buffer (pH 7.2–7.5), and 50 μM 2-mercaptoethanol. For differentiation into macrophages, THP-1 cells were treated with 200 nM phorbol 12-myristate 13-acetate (PMA) for 24 h, followed by an additional 48 h incubation in fresh PMA-free medium to allow complete differentiation prior to experimental use.

HEK293T and RAW264.7 monocytes were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (v/v) FBS and 50 μg/mL penicillin-streptomycin. All cells were maintained at 37 °C in a humidified atmosphere with 5% CO_2.

Recombinant expression and secretion of MmpE

Recombinant *M. bovis* BCG strains expressing C-terminally 2×Flag-tagged Ag85B (positive control) and MmpE were constructed to assess protein expression and secretion under standard *in vitro* culture conditions. The strains were cultured in

200 mL of Middlebrook 7H9 broth supplemented with 10% (v/v) OADC, 2% (v/v)

glycerol, and 0.05% Tween-80 at 37 °C until mid-log phase (OD_{600} = 0.6). Cultures

were centrifuged at 4,000 × g for 60 min to separate bacterial pellets from supernatants.

Pellets were washed and lysed by bead beating in phosphate-buffered saline (PBS)

supplemented with protease inhibitors. Culture supernatants were filtered through

0.22 μm sterile filters to remove residual bacteria and concentrated to a final volume

of 2 mL using Amicon Ultra centrifugal filters (10 kDa molecular weight cutoff). Both

bacterial lysates and concentrated supernatants were analyzed by SDS–PAGE

followed by immunoblotting using anti-Flag antibodies to detect the expression and

secretion of 2×Flag-tagged proteins.

Phylogenetic and sequence analysis

To investigate the evolutionary conservation and sequence features of MmpE

homologs, phylogenetic reconstruction was performed using MEGA version 12.0 and

iTOL platform with a neighbor-joining algorithm. Parameters included pairwise deletion

for gap treatment, a poisson substitution model (selected for its suitability in analyzing

closely related bacterial sequences), and 1,000 bootstrap replicates to assess node

support. Only nodes with bootstrap values ≥50% were retained, and no additional tree

pruning was applied.

The analysis combined metallophosphatase MmpE homologs from 11 *Mycobacterium*

strains, including *M. tuberculosis* (Mtu), *M. bovis* BCG (Mbb), *M. orygis* (Mory), *M.*

kubicae (Mku), *M. paraterrae* (Mpaa), *M. farcinogenes* (Mfg), *M. mucogenicum* (Mmuc),

M. vicinigordonae (Mgor), *M. lentiflavum* (Mlw), *M. avium* (Mav), *M. manitobense*

(Mman). Genomic sequences were retrieved from the NCBI RefSeq database, filtered

for completeness, and aligned using Clustal Omega v1.2.4 with default parameters

(gap opening penalty = 10, gap extension = 0.2, and iterative refinement enabled).

Sequence conservation was visualized using ESPript 3.0, with residues exhibiting ≥90%

identity across all homologs highlighted in red.

Cell transfection and confocal microscopy

HEK293T cells were seeded at 70% confluency in poly-L-lysine-coated 35-mm glass-

bottom dishes and maintained in DMEM supplemented with 10% (v/v) FBS at 37 °C under 5% CO_2. Transfection was performed using the Hieff Trans™ Liposomal Transfection Reagent according to the manufacturer's protocol. Briefly, 0.5 µg of plasmid DNA and 1.5 µL of transfection reagent were diluted in 50 µL of Opti-MEM, incubated at room temperature for 20 minutes, and then added dropwise to each well of 35-mm six-chamber glass-bottom dishes.

At 24–48 hpt, cells were washed with PBS, fixed in 4% paraformaldehyde for 15 min, and permeabilized with 0.1% Triton X-100 for 10 min. Nuclei were counterstained with 1 µg/mL DAPI (4′,6-diamidino-2-phenylindole) for 10 min. Confocal imaging was performed using an Olympus FV1000 Confocal Laser Scanning Microscope equipped with a 100×/1.40 NA oil immersion objective. EGFP and DAPI signals were detected with 488 nm and 405 nm lasers, respectively. Spectral unmixing was carried out using the Olympus FV10-ASW software (v4.2) to minimize signal cross-talk. Z-stack images were acquired in sequential mode with a 0.5 µm step size across all samples to ensure imaging consistency.

Immunoblot analysis

For immunoblotting, the cells were lysed in radioimmunoprecipitation assay (RIPA) buffer supplemented with 1.5% (v/v) protease inhibitor cocktail. Protein concentrations were quantified using the bicinchoninic acid (BCA) assay. Equal amounts of protein (20–40 µg) were resolved by SDS-PAGE and transferred onto PVDF membranes. Membranes were blocked for 1 h at room temperature with 5% (w/v) non-fat milk in TBST (25 mM Tris, 150 mM NaCl, 2 mM KCl, and 0.2% Tween-20, pH 7.4) and incubated overnight at 4 °C with primary antibodies. After three washes with TBST (10 min each), membranes were incubated with horseradish peroxidase (HRP)-conjugated secondary antibodies for 1 hour at room temperature. Protein bands were visualized using an enhanced chemiluminescence (ECL) substrate and imaged using the ChemiDoc™ MP Imaging System (Bio-Rad). β-actin, GAPDH, or histone H3 was used as an internal control for normalization. Antibody details are provided in the Key resources table.

For subcellular fractionation, cytoplasmic and nuclear proteins were isolated using a

Nuclear and Cytoplasmic Protein Extraction Kit (manufacturer's protocol). Briefly, after

treatment, cells were washed and harvested in cold PBS. Cytoplasmic and nuclear

fractions were separated according to the manufacturer's instructions. Protein

concentrations were measured using the BCA assay, and equal amounts of protein

were analyzed by immunoblotting as described above.

AlphaFold Predicted Structural Modeling of MmpE with NLS Truncations

The structural models of MmpE with NLS truncations were generated using AlphaFold

version 2.2.0. The full-length MmpE sequence was used as the input to predict the

reference structure. To assess the impact of NLS deletions, modified sequences were

prepared by removing the residues corresponding to NLS1 (RRR^{20-22}), NLS2 (RRK^{460-}

^{462}), or both motifs. Structural prediction was performed for each sequence using the

default AlphaFold parameters, which include the application of the pre-trained model

and multiple sequence alignment (MSA) generation using the UniRef90 and MGnify

databases. The confidence of the predictions was evaluated using the predicted local

distance difference test (pLDDT) scores provided by AlphaFold. The resulting models

were visualized and analyzed using UCSF ChimeraX to compare the overall protein

conformations and assess structural changes due to NLS deletions.

Prediction of metal ion-binding residues in MmpE

The structure of MmpE was predicted using AlphaFold version 2.2.0 to investigate its

potential metal ion dependence and identify key residues involved in metal

coordination. The predicted structure was analyzed using PyMOL to identify residues

within the putative active site. Residues were selected based on their spatial

arrangement in the active site, and their proximity to regions likely involved in metal

ion binding was annotated. Visualization and annotation of the predicted active site

were performed using PyMOL to determine the potential roles of these residues in

metal ion coordination.

Protein Expression and Purification

N-terminal maltose-binding protein (MBP)-tagged and C-terminal His_6-tagged wild-

type MmpE, as well as mutant variants lacking the Tat signal peptide were expressed in *E. coli* BL21 (DE3). Protein expression was performed in 1 L of LB medium. Cultures were grown at 37 °C until the optical density at 600 nm (OD_{600}) reached approximately 0.8, after which protein expression was induced with 0.2 mM isopropyl-β-D-thiogalactopyranoside (IPTG) at 18 °C for 24 h. Following induction, cells were harvested by centrifugation and lysed by sonication in ice-cold lysis buffer containing 150 mM NaCl, 30 mM Tris-HCl (pH 7.5), and 1 mM phenylmethylsulfonyl fluoride (PMSF). The lysate was clarified by centrifugation, and the supernatant was subjected to affinity purification using an MBP affinity column.

Phosphatase activity assay

Phosphatase activity was measured using *p*-nitrophenyl phosphate (*p*-NPP) as the substrate in a final reaction volume of 200 μL. Reactions were initiated by incubating 0–80 μg of purified protein with *p*-NPP at 37 °C for 60 min. The release of *p*-nitrophenol (*p*-NP), the dephosphorylated product, was quantified by measuring absorbance at 405 nm using a microplate spectrophotometer.

Kinetic parameters were determined under initial velocity (V_0) conditions using substrate concentrations ranging from 0.1 to 10 mM *p*-NPP. The kinetic constants Km and Vmax were calculated by fitting the data to the Michaelis-Menten equation using nonlinear regression analysis in GraphPad Prism 8.0. The catalytic turnover rate (Kcat) was calculated using the equation Kcat = Vmax/E_0, where Vmax was expressed as ηM *p*-NP min^{-1} and E_0 represents the active enzyme concentration in ηM. Control reactions were carried out in the absence of either the substrate or the enzyme to measure background absorbance. All assays were performed in triplicate to ensure the reproducibility of results.

Quantitative real-time PCR

THP-1 cells were infected with the *M. bovis* BCG strain and harvested 20 h post-infection for total RNA isolation. RNA was extracted using 1 mL of TRIzol® reagent following the manufacturer's protocol. To eliminate genomic DNA contamination, RNA samples were treated with DNase I for 30 min at 37 °C, followed by heat inactivation

at 65 °C for 10 min. RNA concentration and purity were assessed using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific).

For cDNA synthesis, 1 μg of total RNA was reverse transcribed using the HiScript® III cDNA Reverse Transcription Kit with random hexamers primers, according to the manufacturer's protocol. qRT-PCR was performed using ChamQ SYBR® Green RT-PCR Master Mix. Relative mRNA expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method, normalizing Ct values to the housekeeping gene *HPRT*. All qRT-PCR experiments were conducted in triplicate, with at least three independent biological replicates and two technical replicates per condition. Primer sequences are listed in Table S6.

RNA-seq analysis

THP-1 cells were seeded at a density of 1×10^6 cells well in 6-well plates and differentiated into macrophages by treatment with 100 nM PMA for 48 h. After differentiation, cells were washed with PBS and infected with *M. bovis* BCG WT or *KommpE* strains at a multiplicity of infection (MOI) of 10:1. Following 20 h of infection, total RNA was extracted using TRIzol® reagent according to the manufacturer's protocol.

High-quality sequencing reads were then aligned to the human reference genome (GRCh38) using HISAT2 (v2.2.1). Transcript quantification was performed using FeatureCounts (v2.0.3), and differential gene expression analysis was conducted using DESeq2 (v1.38.3) in R. Genes with an adjusted *P*-value < 0.05 and an absolute |$\log_2$ fold change ≥ 1| were considered significantly differentially expressed. RNA-seq data were visualized using GraphPad Prism version 8.0, as well as volcano plots, heatmaps, and other graphical outputs generated via the SRplot platform (Tang *et al.,* 2023).

CUT&Tag analysis

HEK293T cells were cultured in 10 cm^2 flat dish at a density of 1×10^7 cells per well and transfected with 10 μg of the MmpE-pEGFP expression plasmid using Hieff

Trans™ Liposomal Transfection Reagent, following the manufacturer's protocol. Cells transfected with the empty vector pEGFP-N1 served as controls. After 36 h of transfection, cells were harvested for CUT&Tag analysis.

For the CUT&Tag, cells were cross-linked with 1% formaldehyde at room temperature for 10 minutes, and the reaction was quenched with 0.125 M glycine. Cells were collected by centrifugation and permeabilized using ice-cold wash buffer containing 0.05% digitonin. Primary antibody incubation was performed overnight at 4 °C using mouse anti-GFP antibody (1:500 dilution), followed by a 2-hour incubation at room temperature with a Protein A/G-conjugated secondary antibody (1:100 dilution). The cells were then incubated at 37 °C for 1 hour with a Protein A–Tn5 transposase complex (1:100 dilution) preloaded with Illumina sequencing adapters, enabling site-specific cleavage and adapter insertion near MmpE binding sites.

The reaction was terminated using lysis buffer containing 10% SDS, followed by proteinase K digestion. DNA was purified and amplified by PCR (12 cycles) using the NEBNext Ultra II DNA Library Preparation Kit. Fragment size distribution (150–300 bp) was confirmed using the Agilent 2100 Bioanalyzer.

Sequencing was performed on the Illumina NovaSeq 6000 platform, producing 150 bp paired-end reads. Raw reads were quality-checked using FastQC and aligned to the human reference genome (hg38). Peak calling was conducted using MACS2 (v2.2.7) with the parameters --nomodel --qvalue 0.05, identifying 2,903 high-confidence MmpE binding sites with >9-fold enrichment. Genomic annotation, including promoter analysis and gene ontology (GO) enrichment, was performed using HOMER (v4.11). *De novo* motif discovery was also conducted with HOMER, and motifs were statistically validated using TOMTOM with a significance threshold of $P < 0.001$.

Chromatin Immunoprecipitation (ChIP) assays

ChIP assays were performed to validate the direct binding of MmpE to the promoter regions of target genes. HEK293T cells were transfected with either an empty control vector, EGFP, or MmpE–EGFP expression plasmids. Cross-linking, chromatin isolation,

fragmentation, and immunoprecipitation were carried out following the same procedure

described for the CUT&Tag experiments.

DNA was purified from both input and immunoprecipitated (IP) samples and subjected

to PCR using primers specific to the promoter regions of the target genes (−100 bp to

+50 bp relative to the transcription start site). *GAPDH* promoter primers were used as

a negative control. PCR products were analyzed by agarose gel electrophoresis to

assess the presence of enriched promoter fragments.

For quantitative analysis, ChIP–qPCR was performed to measure the enrichment of

specific promoter regions in IP samples. All reactions were carried out in triplicate.

Relative enrichment was calculated using the $2^{-\Delta Ct}$ method, normalized to input DNA.

Electrophoretic mobility shift assays (EMSA)

DNA substrates were amplified by PCR from the genomic DNA of *Mycobacterium*

tuberculosis H37Rv. The sequences of the oligonucleotide fragments used in the assay

are listed in Table S6. EMSA was carried out using a modified protocol as previously

described (Li *et al.,* 2020). Each 20 µL reaction contained PCR-amplified DNA

fragments, purified MmpE protein at varying concentrations, and binding buffer

composed of 50 mM Tris–HCl (pH 7.5), 50 mM NaCl, 50 mM $FeCl_3$, 0.05 mg/mL bovine

serum albumin (BSA), and 10% (v/v) glycerol. Reaction mixtures were incubated on

ice for 30 minutes to allow protein–DNA complex formation. Samples were then

resolved on a 4.8% native polyacrylamide gel using a running buffer containing 15 mM

Tris-HCl (pH 7.5) and 0.1 M glycine at 80 V. Following electrophoresis, gels were

stained with ethidium bromide and visualized using a Gel Imaging Analysis System

(JIAPENG, China).

Colony-Forming Unit (CFU) assay

For bacterial preparation, mid-log phase cultures of *M. bovis* BCG (OD_{600} = 0.8) were

harvested, washed with PBS, and resuspended in fresh medium. THP-1 cells were

seeded in 24-well plates at a density of 5×10^5 cells per well and infected at a MOI of

1:10. After 4 hours of infection, extracellular bacteria were removed by treatment with

penicillin–streptomycin (100 µg/mL) for 2 hours. Cells were then washed three times with PBS and maintained in antibiotic-free medium until harvest. To count intracellular bacteria, macrophages were lysed with 0.25% SDS for 10 minutes at room temperature. Lysates were serially diluted in PBS, and 100 µL aliquots of each dilution were plated on Middlebrook 7H10 agar supplemented with 10% OADC and kanamycin (50 µg/mL). Plates were incubated at 37 °C for 14 days. CFUs were enumerated, and final counts were calculated by applying the corresponding dilution factors. *M. bovis* BCG strains used in this study are listed in Table S5.

Animal experiments

Specific pathogen-free (SPF) female C57BL/6 mice (6–8 weeks old, 16–18 g) were obtained from Chang-sheng Bio (Liaoning, China). All animals were maintained under SPF conditions in individually ventilated cages with controlled temperature and humidity, on a 12-hour light/dark cycle. Mice had ad libitum access to sterilized food and water.

Mice infection

Mice were infected as described previously (Chen *et al.,* 2022), with minor modifications. Mid-log phase *M. bovis* BCG cultures were washed twice in PBS containing 0.05% Tween 80 and briefly sonicated to disrupt bacterial clumps. Female C57BL/6 mice were anesthetized with isoflurane and intranasally inoculated with 1×10^7 CFUs of *M. bovis* BCG in 40 µL of PBS. To confirm the inoculation dose, six mice were euthanized at 48 hours post-infection. Lungs were harvested, homogenized in 1 mL of PBS using a tissue homogenizer, and serial dilutions of the homogenates were plated on Middlebrook 7H10 agar supplemented with 10% OADC. Plates were incubated at 37 °C with 5% CO_2 for 14 days, after which CFUs were enumerated.

Histological analysis

Lung tissues from *M. bovis* BCG-infected mice were fixed in 4% phosphate-buffered formalin at room temperature for 24 h, followed by paraffin embedding. Paraffin-embedded tissue samples were sectioned into 2–3 µm thick slices using a microtome. Sections were deparaffinized, rehydrated through a graded ethanol series, and stained

with hematoxylin and eosin (H&E). Stained tissue sections were examined using an Olympus BX53 light microscope. For digital analysis and annotation, scanned slides were visualized using CaseViewer software (version 2.0; 3DHISTECH).

Ethics approval

All animal experiments were conducted in accordance with the protocols approved by the Animal Ethics Committee of the Huazhong Agricultural University.

Statistical analyses

Data are presented as mean ± standard deviation (SD). Statistical comparisons between two groups were performed using two-tailed unpaired Student's t-tests, following verification of normal distribution. A P-value less than 0.05 was considered statistically significant. Significance levels are indicated in figures as follows: $P > 0.05$ (ns, not significant), $P < 0.05$ (*), $P < 0.01$ (**), and $P < 0.001$ (***). All statistical analyses were performed using GraphPad Prism (version 8.0). Data were obtained from at least three independent biological replicates, each including two or more technical replicates.

Acknowledgments

This work was supported by the Fundamental Research Funds for the Central Universities (#2662023DKQD001), Talent Start-up Funds of Huazhong Agricultural University (#11042310008), National Natural Science Foundation of China (#32473123), China Agriculture Research System of MOF and MARA (#CARS-37), and National Key Research and Development Program of China (#2021YFD1800403).

Author contributions

Conceptualization: L. C., A. G., and L. Z.; Methodology: L. C.; Validation: L. C., B. D., and Q. J.; Formal analysis: L. C., Y. W.; Investigation: L. C., B. D., Q. J., and Y. W.; Resources: A. G., L. Z., and Y. C.; Data curation: L. C., Q. J., and Y. W.; Writing - original draft: L. C., L. Z., and A. G.; Writing - review & editing: L. C., L. Z., and A. G.; Visualization: L. C., Q. J., Y. C., L. Z., and A. G.; Supervision: A. G., and L. Z.; Project administration: A. G., and L. Z.; Funding acquisition: A. G., L. Z., and Y. C.

Disclosure and competing interest statement

The authors declare no competing interests.

References

Agarwal S, Agarwal S, Jin H, Pancholi P, Pancholi V (2012) **Serine/threonine phosphatase (SP-STP), secreted from Streptococcus pyogenes, is a pro-apoptotic protein** *J Biol Chem* **287**:9147-67.

Aggeletopoulou I, Thomopoulos K, Mouzaki A, Triantos C (2022) **Vitamin D-VDR novel anti-inflammatory molecules-new insights into their effects on liver diseases** *Int J Mol Sci* **23**:8465.

Alsayed SSR, Gunosewoyo H (2024) **Combating Tuberculosis via restoring the host immune capacity by targeting Mtb kinases and phosphatases** *Int J Mol Sci* **25**:12481.

Baruch K, Gur-Arie L, Nadler C, Koby S, Yerushalmi G, Ben-Neriah Y, Yogev O, Shaulian E, Guttman C, Zarivach R, Rosenshine I (2011) **Metalloprotease type III effectors that specifically cleave JNK and NF-κB** *EMBO J* **30**:221-31.

Bhadouria J, Giri J (2022) **Purple acid phosphatases: roles in phosphate utilization and new emerging functions** *Plant Cell Rep* **41**:33-51.

Bhadouria J, Singh AP, Mehra P, Verma L, Srivastawa R, Parida SK, Giri J (2017) **Identification of purple acid phosphatases in chickpea and potential roles of CaPAP7 in seed phytate accumulation** *Sci Rep* **7**:11012.

Bierne H, Cossart P (2021) **When bacteria target the nucleus: the emerging family of nucleomodulins** *Cell Microbiol* **14**:622-33.

Bonne Køhler J, Jers C, Senissar M, Shi L, Derouiche A, Mijakovic I (2020) **Importance of protein Ser/Thr/Tyr phosphorylation for bacterial pathogenesis** *FEBS Lett* **594**:2339-2369.

Buchanan G, Sargent F, Berks BC, Palmer T (2001) **A genetic screen for**
suppressors of *Escherichia coli* Tat signal peptide mutations establishes a
critical role for the second arginine within the twin-arginine motif *Arch Microbiol*
177:107-12.

Cautain B, Hill R, de Pedro N, Link W (2015) **Components and regulation of nuclear**
transport processes *FEBS J* **282**:445-62.

Chai Q, Wang L, Liu CH, Ge B (2020) **New insights into the evasion of host innate**
immunity by *Mycobacterium tuberculosis* *Cell Mol Immunol* **17**:901-913.

Chandra P, Grigsby SJ, Philips JA (2022) **Immune evasion and provocation by**
Mycobacterium tuberculosis *Nat Rev Microbiol* **20**:750-766.

Chen L, Duan B, Chen P, Jiang Q, Wang Y, Lu L, Chen Y, Hu C, Zhang L, Guo A (2025)
A conserved mycobacterial nucleomodulin hijacks the host COMPASS complex
to reprogram pro-inflammatory transcription and promote intracellular survival
bioRxiv https://doi.org/10.1101/2025.05.21.655295

Chen L, Li X, Xu P, He ZG (2022) **A novel zinc exporter CtpG enhances resistance**
to zinc toxicity and survival in *Mycobacterium bovis* *Microbiol Spectr*
10:e0145621.

Chen Y, Du J, Zhang Z, Liu T, Shi Y, Ge X, Li YC (2014) **MicroRNA-346 mediates**
tumor necrosis factor α-induced downregulation of gut epithelial vitamin D
receptor in inflammatory bowel diseases *Inflamm Bowel Dis* **20**:1910-8.

Comba S, Menendez-Bravo S, Arabolaza A, Gramajo H (2013) **Identification and**
physiological characterization of phosphatidic acid phosphatase enzymes
involved in triacylglycerol biosynthesis in *Streptomyces coelicolor* *Microb Cell*
Fact **12**:9.

Davicino RC, Méndez-Huergo SP, Eliçabe RJ, Stupirski JC, Autenrieth I, Di Genaro
MS, Rabinovich GA (2017) **Galectin-1-driven tolerogenic programs**

aggravate *Yersinia enterocolitica* infection by repressing antibacterial Immunity
J Immunol **199**:1382-1392.

Dean P, Scott JA, Knox AA, Quitard S, Watkins NJ, Kenny B (2010) **The**
enteropathogenic *E. coli* effector EspF targets and disrupts the nucleolus by a
process regulated by mitochondrial dysfunction *PLoS Pathog* **6**:e1000961.

Evans SM, Rodino KG, Adcox HE, Carlyon JA (2018) ***Orientia tsutsugamushi* uses**
two Ank effectors to modulate NF-κB p65 nuclear transport and inhibit NF-κB
transcriptional activation *PLoS Pathog* **14**:e1007023.

Fang S, Wan X, Zou X, Sun S, Hao X, Liang C, Zhang Z, Zhang F, Sun B, Li H, Yu B
(2021) **Arsenic trioxide induces macrophage autophagy and atheroprotection by**
regulating ROS-dependent TFEB nuclear translocation and AKT/mTOR pathway
Cell Death Dis **12**:88.

Feder D, McGeary RP, Mitić N, Lonhienne T, Furtado A, Schulz BL, Henry RJ, Schmidt
S, Guddat LW, Schenk G (2020) **Structural elements that modulate the substrate**
specificity of plant purple acid phosphatases: Avenues for improved
phosphorus acquisition in crops *Plant Sci* **294**:110445.

Fontes MR, Teh T, Kobe B (2000) **Structural basis of recognition of monopartite**
and bipartite nuclear localization sequences by mammalian importin-alpha *J Mol*
Biol **297**:1183-94.

Forrellad MA, Blanco FC, Marrero Diaz de Villegas R, Vázquez CL, Yaneff A, García
EA, Gutierrez MG, Durán R, Villarino A, Bigi F (2020) **Rv2577 of *Mycobacterium***
***tuberculosis* Is a virulence factor with dual phosphatase and phosphodiesterase**
functions *Front Microbiol* **11**:570794.

Ge P, Lei Z, Yu Y, Lu Z, Qiang L, Chai Q, Zhang Y, Zhao D, Li B, Pang Y, Liu CH, Wang
J (2022) ***M. tuberculosis* PknG manipulates host autophagy flux to promote**
pathogen intracellular survival *Autophagy* **18**:576-594.

Gimenez MR, Chandra G, Van Overvelt P, Voulhoux R, Bleves S, Ize B (2018)
Genome wide identification and experimental validation of *Pseudomonas*
***aeruginosa* Tat substrates** *Sci Rep* **8**:11950.

Gounis M, Hamidi H, Ivaska J (2025) **mTORC1 shutdown unleashes TFEB to drive**
triple-negative breast cancer invasion *Dev Cell* **60**:979-981.

Hanford HE, Von Dwingelo J, Abu Kwaik Y (2021) **Bacterial nucleomodulins: a**
coevolutionary adaptation to the eukaryotic command center *PLoS Pathog*
17:e1009184.

Hodgson A, Wier EM, Fu K, Sun X, Yu H, Zheng W, Sham HP, Johnson K, Bailey S,
Vallance BA, Wan F (2015) **Metalloprotease NleC suppresses host NF-**
κB/inflammatory responses by cleaving p65 and interfering with the p65/RPS3
interaction *PLoS Pathog* **11**:e1004705.

Jeong SJ, Stitham J, Evans TD, Zhang X, Rodriguez-Velez A, Yeh YS, Tao J,
Takabatake K, Epelman S, Lodhi IJ, Schilling JD, DeBosch BJ, Diwan A, Razani B
(2021) **Trehalose causes low-grade lysosomal stress to activate TFEB and the**
autophagy-lysosome biogenesis response. Autophagy **17**:3740-3752.

Jermy A (2010) **Chlamydia gets a NUE look** *Nat Rev Microbiol*. **8**:614.

Jose L, Ramachandran R, Bhagavat R, Gomez RL, Chandran A, Raghunandanan S,
Omkumar RV, Chandra N, Mundayoor S, Kumar RA (2016) **Hypothetical protein**
Rv3423.1 of *Mycobacterium tuberculosis* is a histone acetyltransferase *FEBS J*
283:265-81.

Kosugi S, Hasebe M, Matsumura N, Takashima H, Miyamoto-Sato E, Tomita M,
Yanagawa H (2009) **Six classes of nuclear localization signal specific to different**
binding grooves of importin alpha *J Biol Chem* **284**:478-485.

Kurucz V, Krüger T, Antal K, Dietl AM, Haas H, Pócsi I, Kniemeyer O, Emri T (2018)
Additional oxidative stress reroutes the global response of *Aspergillus*
***fumigatus* to iron depletion** *BMC Genomics* **19**:357.

Lange A, Mills RE, Devine SE, Corbett AH (2008) **A PY-NLS nuclear targeting signal**

is required for nuclear localization and function of the *Saccharomyces*

cerevisiae **mRNA-binding protein Hrp1** *J Biol Chem* **283**:12926-34.

Lebreton A, Job V, Ragon M, Le Monnier A, Dessen A, Cossart P, Bierne H (2014)

Structural basis for the inhibition of the chromatin repressor BAHD1 by the

bacterial nucleomodulin LntA *mBio* **5**: e00775-13.

Lebreton A, Lakisic G, Job V, Fritsch L, Tham TN, Camejo A, Matteï PJ, Regnault B,

Nahori MA, Cabanes D, Gautreau A, Ait-Si-Ali S, Dessen A, Cossart P, Bierne H (2011)

A bacterial protein targets the BAHD1 chromatin complex to stimulate type III

interferon response *Science* **331**:1319-21.

Li T, Lu Q, Wang G, Xu H, Huang H, Cai T, Kan B, Ge J, Shao F (2013) **SET-domain**

bacterial effectors target heterochromatin protein 1 to activate host rDNA

transcription *EMBO Rep* **14**:733-40.

Li X, Chen L, Liao J, Hui J, Li W, He ZG (2020) **A novel stress-inducible CmtR-ESX3-**

Zn^{2+} regulatory pathway essential for survival of *Mycobacterium bovis* **under**

oxidative stress *J Biol Chem* **295**:17083-17099.

Li X, Zhang M, Zhou G, Xie Z, Wang Y, Han J, Li L, Wu Q, Zhang S (2023) **Role of**

Rho GTPases in inflammatory bowel disease *Cell Death Discov* **9**:24.

Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, Ochoa MT, Schauber J, Wu K,

Meinken C, Kamen DL, Wagner M, Bals R, Steinmeyer A, Zügel U, Gallo RL,

Eisenberg D, Hewison M, Hollis BW, Adams JS, Bloom BR, Modlin RL (2006) **Toll-like**

receptor triggering of a vitamin D-mediated human antimicrobial response

Science **311**:1770-3.

Liu SF, Yi ZC, Huang ZQ, Yuan ZD, Yang YC, Zhao Y, He QY, Yang WD, Li HY, Lin

CSK, Wang X (2025) **Enhanced biodegradation of glyphosate by** *Chlorella*

sorokiniana **engineered with exogenous purple acid phosphatase** *Water Res*

268:122737.

Lu J, Wu T, Zhang B, Liu S, Song W, Qiao J, Ruan H (2021) **Types of nuclear**
localization signals and mechanisms of protein import into the nucleus *Cell*
Commun Signal **19**:60.

Ma S, Ming Y, Wu J, Cui G (2024) **Cellular metabolism regulates the differentiation**
and function of T-cell subsets *Cell Mol Immunol* **21**:419-435.

Manfiolli AO, de Castro PA, Dos Reis TF, Dolan S, Doyle S, Jones G, Riaño Pachón
DM, Ulaş M, Noble LM, Mattern DJ, Brakhage AA, Valiante V, Silva-Rocha R, Bayram
O, Goldman GH (2017) *Aspergillus fumigatus* **protein phosphatase PpzA is**
involved in iron assimilation, secondary metabolite production, and virulence
Cell Microbiol **19**.

Mitra S, Dunphy PS, Das S, Zhu B, Luo T, McBride JW (2018) *Ehrlichia chaffeensis*
TRP120 effector targets and recruits host polycomb group proteins for
degradation to promote intracellular infection *Infect Immun* **86**:e00845-17.

Nasiri MJ, Venketaraman V (2025) **Advances in host-pathogen interactions in**
Tuberculosis: emerging strategies for therapeutic intervention *Int J Mol Sci*
26:1621.

Nisa A, Kipper FC, Panigrahy D, Tiwari S, Kupz A, Subbian S (2022) **Different**
modalities of host cell death and their impact on *Mycobacterium tuberculosis*
infection *Am J Physiol Cell Physiol* **323**:C1444-C1474.

Pennini ME, Perrinet S, Dautry-Varsat A, Subtil A (2010) **Histone methylation by NUE,**
a novel nuclear effector of the intracellular pathogen *Chlamydia trachomatis*
PLoS Pathog **6**:e1000995.

Rodriguez F, Lillington J, Johnson S, Timmel CR, Lea SM, Berks BC (2014) **Crystal**
structure of the *Bacillus subtilis* **phosphodiesterase PhoD reveals an iron and**
calcium-containing active site *J Biol Chem* **289**:30889-99.

Rolando M, Sanulli S, Rusniok C, Gomez-Valero L, Bertholet C, Sahr T, Margueron R,
Buchrieser C (2013) *Legionella pneumophila* **effector RomA uniquely modifies**

host chromatin to repress gene expression and promote intracellular bacterial replication *Cell Host Microbe* **13**:395-405.

Sajid A, Arora G, Singhal A, Kalia VC, Singh Y (2015) **Protein phosphatases of pathogenic bacteria: role in physiology and virulence** *Annu Rev Microbiol* **69**:527-47.

Schenk G, Guddat LW, Ge Y, Carrington LE, Hume DA, Hamilton S, de Jersey J (2000) **Identification of mammalian-like purple acid phosphatases in a wide range of plants** *Gene* **250**:117-25.

Shaban L, Nguyen GT, Mecsas-Faxon BD, Swanson KD, Tan S, Mecsas J (2020) *Yersinia pseudotuberculosis* **YopH targets SKAP2-dependent and independent signaling pathways to block neutrophil antimicrobial mechanisms during infection** *PLoS Pathog* **16**:e1008576.

Singh AP, Sharma S, Pagarware K, Siraji RA, Ansari I, Mandal A, Walling P, Aijaz S (2018) **Enteropathogenic *E. coli* effectors EspF and Map independently disrupt tight junctions through distinct mechanisms involving transcriptional and post-transcriptional regulation** *Sci Rep* **8**:3719.

Singh PR, Dadireddy V, Udupa S, Kalladi SM, Shee S, Khosla S, Rajmani RS, Singh A, Ramakumar S, Nagaraja V (2023) **The *Mycobacterium tuberculosis* methyltransferase Rv2067c manipulates host epigenetic programming to promote its own survival** *Nat Commun* **14**:8497.

Singh PR, Nagaraja V (2025) **Epigenetic maneuvering: an emerging strategy for mycobacterial intracellular survival** *Trends Microbiol* **33**:354-369.

Stanley NR, Palmer T, Berks BC (2000) **The twin arginine consensus motif of Tat signal peptides is involved in Sec-independent protein targeting in *Escherichia coli*** *J Biol Chem* **275**:11591-6.

Steiert B, Weber MM (2025) **Nuclear warfare: pathogen manipulation of the nuclear pore complex and nuclear functions** *mBio* **16**:e0194024.

Tang D, Chen M, Huang X, Zhang G, Zeng L, Zhang G, Wu S, Wang Y (2023) **SRplot:**

A free online platform for data visualization and graphing *PLoS One* **18**:e0294236.

Ulfig A, Fröbel J, Lausberg F, Blümmel AS, Heide AK, Müller M, Freudl R (2017) **The**

h-region of twin-arginine signal peptides supports productive binding of

bacterial Tat precursor proteins to the TatBC receptor complex *J Biol Chem*

292:10865-10882.

Usategui-Martín R, De Luis-Román DA, Fernández-Gómez JM, Ruiz-Mambrilla M,

Pérez-Castrillón JL (2022) **Vitamin D receptor (VDR) gene polymorphisms modify**

the response to vitamin d supplementation: a systematic review and meta-

analysis *Nutrients* **14**:360.

Wang B, Martini-Stoica H, Qi C, Lu TC, Wang S, Xiong W, Qi Y, Xu Y, Sardiello M, Li

H, Zheng H (2024) **TFEB-vacuolar ATPase signaling regulates lysosomal function**

and microglial activation in tauopathy *Nat Neurosci* **27**:48-62.

Wang J, Ge P, Qiang L, Tian F, Zhao D, Chai Q, Zhu M, Zhou R, Meng G, Iwakura Y,

Gao GF, Liu CH (2017) **The mycobacterial phosphatase PtpA regulates the**

expression of host genes and promotes cell proliferation *Nat Commun* **8**:244.

Wong D, Bach H, Sun J, Hmama Z, Av-Gay Y (2011) *Mycobacterium tuberculosis*

protein tyrosine phosphatase (PtpA) excludes host vacuolar-H$^+$-ATPase to

inhibit phagosome acidification *Proc Natl Acad Sci U S A* **108**:19371-6.

Yeung SL, Cheng C, Lui TK, Tsang JS, Chan WT, Lim BL (2009) **Purple acid**

phosphatase-like sequences in prokaryotic genomes and the characterization of

an atypical purple alkaline phosphatase from *Burkholderia cenocepacia* **J2315**

Gene **440**:1-8.

Zhang QA, Ma S, Li P, Xie J (2023) **The dynamics of** *Mycobacterium tuberculosis*

phagosome and the fate of infection *Cell Signal* **108**:110715.

Zhang W, Dong C, Xiong S (2024) **Mycobacterial SapM hampers host autophagy initiation for intracellular bacillary survival via dephosphorylating Raptor** *iScience* **27**:109671.

Legends

Figure 1. NLSs are required for the nuclear translocation of MmpE.

(A) Expression and nuclear localization of *MmpE*. (upper) qRT-PCR analysis of *mmpE*

mRNA in HEK293T cells over 48 h post-transfection. (bottom) Western blot analysis of

nuclear fractions showing time-dependent nuclear accumulation of MmpE-EGFP.

Histone H3 and β-actin served as nuclear and cytoplasmic markers, respectively.

(B) Domain architecture of MmpE. The schematic representation of MmpE shows its

annotated functional domains, including a Tat signal peptide (1-54 amino acids, with a

twin-arginine translocation motif) and two nuclear localization signals (NLS1: 20-22

amino acids; NLS2: 460-462 amino acids). The structure of the MmpE protein was

predicted using AlphaFold 2.2.0, where the position of NLS1 is highlighted in red and

NLS2 is highlighted in green.

(C) Subcellular localization of EGFP-tagged wild-type and NLS-deleted MmpE. (left)

Schematic representation of EGFP-tagged constructs, including wild-type MmpE and

its mutants (MmpE$^{\Delta NLS1}$, MmpE$^{\Delta NLS2}$, and MmpE$^{\Delta NLS1-2}$). (right) Confocal microscopy

images of HEK293T cells transfected with the indicated constructs for 36 hpt. EGFP

fluorescence (green) and nuclear staining with DAPI (blue) were visualized using an

FLUOVIEW software (v5.0). Scale bar, 10 μm. Images were acquired with a ×100 oil

immersion objective (NA = 1.4).

(**D**) Nuclear EGFP intensity of wild-type and mutant constructs in **(C)**. Data are shown

as mean ± SD (n = 12 cells).

(**E**) Western blot analysis of nuclear and cytoplasmic fractions from HEK293T cells

transfected with wild-type and mutant MmpE-EGFP confirmed their nuclear

localization. MmpE-EGFP was detected using an anti-GFP antibody, and histone H3

and β-actin served as nuclear and cytoplasmic markers, respectively.

All experiments were performed in three biologically independent replicates. Statistical

significance was assessed by two-tailed unpaired Student's *t*-tests, with *$P < 0.05$

considered statistically significant.

Figure 2. Deletion of NLSs does not alter the phosphatase activity of MmpE.

(**A**) Domain architecture of MmpE. Schematic representation of MmpE with the annotated functional domains, including a Tat signal peptide (1–41 aa, twin-arginine translocation motif), a purple acid phosphatase domain (68–149 aa), a calcineurin-like phosphoesterase domain (211–389 aa) and two nuclear localization signals (NLS1:

20-22 aa; NLS2: 460–462 aa).

(**B**) Phylogenetic and structural conservation of MmpE. Neighbor-joining phylogenetic

tree of MmpE homologs across Mycobacterium species (1,000 bootstrap replicates;

values ≥50% shown). Species abbreviations: *M. tuberculosis* (Mtu), *M. bovis* BCG

(Mbb), *M. orygis* (Mory), *M. kubicae* (Mku), *M. paraterrae* (Mpaa), *M. farcinogenes*

(Mfg), *M. mucogenicum* (Mmuc), *M. vicinigordonae* (Mgor), *M. lentiflavum* (Mlw), *M.*

avium (Mav), *M. manitobense* (Mman).

(C) Prediction of metal ion-binding residues in MmpE. Structural modeling and

visualization of MmpE were performed using PyMOL. Conserved residues located in

the putative metal binding pocket are shown as sticks and colored according to atom

type. The predicted metal coordination site is highlighted, and key residues potentially

involved in metal ion binding are labeled. Surface representation is included to illustrate

the spatial accessibility of the binding pocket.

(D-F) Phosphatase activity assays of MmpE$^{\Delta Tat}$ under various conditions. Phosphatase

activity of MmpE$^{\Delta Tat}$ (lacking the N-terminal Tat signal peptide) was measured using *p*-

nitrophenyl phosphate (*p*-NPP) as the substrate in the presence of increasing

concentrations (0–500 µM) of Fe^{3+} and Zn^{2+} (**D**). Phosphatase activity of increasing

concentrations of MmpE$^{\Delta Tat}$ and the mutant MmpE$^{\Delta Tat}$-H^{348}AN^{359}A was measured in the

absence or presence of 50 µM Fe^{3+} (**E**). Phosphatase activity of increasing

concentrations of MmpE$^{\Delta Tat}$ and the double-NLS deletion mutant MmpE$^{\Delta Tat/\Delta NLS1-2}$ was

measured under standard conditions (**F**).

Data represent mean ± SD of three biologically independent experiments, analyzed

using two-tailed unpaired Student's *t*-tests (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$).

Figure 3. The nuclear translocation and phosphatase activity of MmpE are essential for *M. bovis* BCG survival in macrophage cells.

(A–B) Intracellular survival of *M. bovis* BCG strains in human THP-1 macrophages (**A**) and murine RAW264.7 macrophages (**B**). Strains include wild-type BCG (WT), MmpE deletion mutant (ΔMmpE), wild-type MmpE complemented strain (Comp-MmpE), and NLS2-deleted complement strain (Comp-MmpE^{ΔNLS2}).

(C–D) Intracellular survival of *M. bovis* BCG strains in THP-1 (**C**) and RAW264.7 (**D**) macrophages. Strains include ΔMmpE, Comp-MmpE, and NLS-deficient complement strains (Comp-MmpE^{ΔNLS1} and Comp-MmpE$^{ΔNLS1-2}$).

(E) Intracellular survival of metal ion binding site mutants in THP-1 cells. Strains include Comp-MmpE and a phosphatase-deficient mutant (Comp-MmpE-H^{348}AN^{359}H).

997   **(F)** Inflammatory cytokine expression in infected THP-1 cells. mRNA levels of *IL-1α*,

998   *IL-1β*, and *IL-6* were quantified by qRT-PCR 20 hpi with the indicated BCG strains.

999   Data represent mean ± SD from three independent biological replicates. Statistical

1000   analysis was performed using two-tailed unpaired Student's *t*-tests (*$P < 0.05$, **$P <$

1001 0.01, ***$P < 0.001$).

1002

Figure 4. MmpE modulates host transcription network involved in inflammation response and lysosomal maturation.

(**A**) Volcano plot of differentially expressed genes (DEGs). DEGs in MmpE deleted strain (Ko*mmpE*) compared to wild-type BCG (BCG/wt) were visualized in a volcano

plot. DEGs were defined as genes with $|\log_2(\text{fold change})| \geq 1$ and $P < 0.05$. The *x*-axis represents $\log_2$(fold change), and the *y*-axis shows $-\log_{10}$(P-value). Genes with significant upregulation and downregulation are highlighted.

(B) GO enrichment analysis of DEGs in Ko*mmpE*-infected THP-1 cells versus BCG/wt. The circular plot shows enriched biological process and molecular function terms. Outer rings represent GO terms, with red and green dots indicating upregulated and downregulated DEGs, respectively. The inner ring color reflects z-scores (blue: low; pink: high). The table lists selected GO terms with their IDs and descriptions.

(C) KEGG enrichment analysis of DEGs in KommpE-infected THP-1 cells versus BCG/wt. The bar plot displays selected significantly enriched pathways. Bar length represents the number of DEGs associated with each pathway, and color indicates statistical significance based on $-\log_{10}(P\text{-value})$.

(D) Interaction network of DEGs related to immune regulation. Network construction was performed using STRING v12.0 and visualized with Cytoscape.

(**E**) Heatmap of immune-related DEGs in Ko*mmpE*-infected and BCG/wt-infected THP-1 cells. $\log_2$fold change values are shown across three biological replicates. Red and blue indicate upregulation and downregulation, respectively.

(**F-G**) Quantitative RT-PCR analysis of gene expression in infected THP-1 cells. THP-1 cells were infected with BCG/wt or Ko*mmpE* strains for 20 hpi. mRNA levels of cytokine genes (**F**); mRNA levels of genes involved in lysosomal acidification and biogenesis (**G**).

Data are presented as mean ±SD of three biologically independent experiments, analyzed using two-tailed unpaired Student's *t*-tests (*P < 0.05, **P < 0.01, and ***P < 0.001).

1031

Figure 5. MmpE suppresses the expression of *VDR* and inhibits activation of the

PI3K–Akt–mTOR signaling pathway.

(**A**) Genomic distribution of potential MmpE-binding regions in HEK293T cells.

(**B**) Distribution of the chromosomal location of MmpE-specific ChIP-seq signals (fold-enrichment > 9).

(**C**) Biotype distribution of potential MmpE-binding regions in HEK293T cells.

(**D**) The number and position of the potential MmpE-binding sites along 20 kb from the nearest TSS of the protein-coding genes.

(**E**) Heatmap of transcription factors identified by MmpE ChIP-seq. Colors represent expression changes, with red indicating upregulation and blue indicating downregulation.

(**F**) MmpE-binding motifs identified by *de novo* motif analysis of ChIP-seq peak sequences using HOMER. Motif enrichment *P*-values were calculated using TOMTOM. "% of targets" indicates the proportion of MmpE-bound regions containing each motif. Letter height reflects the frequency of each nucleotide at the corresponding position in the consensus motif.

(**G-H**) ChIP-PCR and qPCR analyses of MmpE binding to the VDR promoter region. Chromatin was immunoprecipitated from HEK293T cells transfected with control vector (CK), EGFP, or MmpE–EGFP constructs. PCR was performed using primers targeting the *GAPDH* region (negative control) and the *VDR* promoter region, and products were analyzed by agarose gel electrophoresis (**G**). ChIP enrichment for *VDR* and *GAPDH* was quantified in HEK293T cells using qPCR, with enrichment calculated by the $2^{-\Delta Ct}$ method(**H**).

(**I-J**) EMSA analysis showing the binding of MmpE to the human *VDR* promoter. The formation of DNA–protein complexes was observed with increasing concentrations of MmpE protein. Arrows indicate the positions of free DNA and DNA–protein complexes on the native polyacrylamide gel (**I**). Quantitative analysis of DNA–protein complexes in each lane (**I**) was performed by densitometry (**J**).

Data are presented as mean ±SD of three biologically independent experiments. Statistical significance was determined by a two-tailed unpaired Student's *t*-test (***$P < 0.001$).

1063

Figure 6. Nuclear translocation of MmpE is essential for mycobacterial survival in mice.

(**A**) Bacterial burden in lungs of infected mice. Specific pathogen-free (SPF) C57BL/6 mice (n = 6 per group) were intratracheally infected with $1.0 × 10^7$ colony-forming units (CFU) of *M. bovis* BCG strains, including wild-type (WT), MmpE deleted (ΔMmpE), MmpE-complemented (Comp-MmpE), or MmpE-complemented with NLS-deleted variants (Comp-MmpE$^{ΔNLS1-2}$). Bacterial titers in lung homogenates were quantified by CFU assays at 0,14,28, and 56 days post-infection.

(**B**) Histopathology of infected lung tissues. Hematoxylin and eosin (H&E)-stained lung sections from mice infected as in (**A**) show granulomatous inflammation, Scale bars: 200 μm.

(**C-D**) Pro-inflammatory cytokine expression in the spleen of infected mice. qRT-PCR

analysis of cytokine mRNA levels of *Il1α* (**C**), *Il1β* (**D**)and *Il6* (**E**) in spleen tissues from

infected mice (n = 6/group) at 2 to 28 days post-infection.

Data are presented as mean ± SD from six biologically independent experiments.

Statistical significance was determined by a two-tailed unpaired Student's *t*-test (*P <

0.05, **P < 0.01, ***P < 0.001).

Figure 7. Schematic diagram of nucleomodulin MmpE-mediated immune suppression and mycobacterial survival.

Upon mycobacterial infection, MmpE translocates into the host cell nucleus via its

NLS2 motif (RRK[460−462]), where it binds to the promoter region of the *VDR* gene,

repressing transcription of downstream inflammatory genes. Simultaneously, MmpE

inhibits the PI3K–Akt–mTOR signaling pathway, leading to impaired phagosome

maturation. These combined actions facilitate immune evasion and promote enhanced

mycobacterial survival within macrophages and in mice.

 Supplementary Figure Legend

 Supplementary Figure 1. Identification of MmpE as a nucleomodulin in

 Mycobacterium.

 (**A**) MmpE is a mycobacterial nucleomodulin. Subnuclear localization of MmpE-EGFP.

 Confocal microscopy was used to assess the nuclear distribution of MmpE-EGFP at

 various time points post-transfection. Nuclei were stained with DAPI (blue); MmpE-

 EGFP is shown in green. Scale bar: 10 μm.

(**B**) AlphaFold-predicted structural models of MmpE with NLS truncations. Structural

models of MmpE were generated using AlphaFold 2.2.0 to assess the impact of NLS

deletions on overall protein conformation. (upper) Full-length MmpE structure shown

as a reference. (middle) MmpE structure with NLS deleted. (botten) Predicted

structures of MmpE with individual deletions of NLS1, NLS2, or both motifs.

Confidence scores (pLDDT) were used to evaluate prediction reliability, and UCSF

ChimeraX was used to visualize and compare structural changes.

A

Protein Chain	Description	Z Value	RMSD Value
6py9-C	$Fe^{(3+)}$-$Zn^{(2+)}$ purple acid phosphatase	36.5	2.3
6py9-D	$Fe^{(3+)}$-$Zn^{(2+)}$ purple acid phosphatase	36.5	2.3
6py9-A	$Fe^{(3+)}$-$Zn^{(2+)}$ purple acid phosphatase	36.4	2.3
4kkz-C	purple acid phosphatase	36.3	2.3
4kkz-A	purple acid phosphatase	36.3	2.3
6g46-D	$Fe^{(3+)}$-$Zn^{(2+)}$ purple acid phosphatase	36.3	2.3
2qfp-B	purple acid phosphatase	36.3	2.3

Z value>10 and RMSD value<4 indicate that the comparison
result is extremely reliable

1108

Supplementary Figure 2. Identification of MmpE as a conserved Fe^{3+}/Zn^{2+}-metallophosphatase in Mycobacteria.

(A) Sequence and structure-based analyses of MmpE. Sequence and structure-based analyses were performed using UCSF Chimera to investigate the potential function and structural features of MmpE. The structural comparison results, with a *Z* value >10 and RMSD value <4, indicate high confidence in the predicted structural models.

(B) Clustal Omega sequence alignment of MmpE from various mycobacterial species,

highlighting conserved residues (≥90% identity, shown in red). Blue stars indicate the

predicted NLSs, and red triangles mark characteristic sequences found in

metallophosphatases. Species abbreviations: *M. tuberculosis* (Mtu), *M. bovis* BCG

(Mbb), *M. orygis* (Mory), *M. kubicae* (Mku), *M. paraterrae* (Mpaa), *M. farcinogenes*

(Mfg), *M. mucogenicum* (Mmuc), *M. vicinigordonae* (Mgor), *M. lentiflavum* (Mlw), *M.*

avium (Mav), *M. manitobense* (Mman).

1122

1123 **Supplementary Figure 3. The nuclear translocation and phosphatase activity of**

1124 **MmpE are essential for *M. bovis* BCG survival in macrophage cells.**

(**A**) Detection of mycobacterial MmpE secretion ability. C-terminally 2×Flag-tagged Ag85B (positive control) and MmpE were expressed in *M. bovis* BCG strains. Immunoblot analysis of bacterial lysates and culture supernatants showing the expression and secretion of Flag-tagged Ag85B and MmpE.

(**B**) Construction and validation of the MmpE-deleted strain of *M. bovis* BCG. (left) Schematic diagram of the homologous recombination strategy used to delete *mmpE* from the *M. bovis* BCG genome. (right) Wild-type and mutant strains were used as templates to amplify the *mmpE* gene (600 bp upstream–600 bp downstream) by PCR. Lanes 1 and 3: wild-type genomic DNA; lanes 2 and 4: Δ*mmp*E genomic DNA.

(**C-D**) Growth curve analysis of BCG strains. Growth of *M. bovis* BCG strains, including wild-type BCG (WT), MmpE-deleted (ΔMmpE), MmpE-complemented (Comp-MmpE), and NLS-deleted complemented (Comp-MmpE$^{\Delta NLS1-2}$) strains(**C**), or phosphatase activity site mutation complement (Comp-MmpE-H^{348}AN^{359}H) (**D**), was measured in 7H9 medium.

Data are presented as mean ± SD from three biologically independent experiments. Statistical significance was determined by a two-tailed unpaired Student's *t*-test.

1141

Supplementary Figure 4. MmpE modulates host transcription network involved in inflammation response and lysosomal maturation

(**A**) Biotype distribution of potential MmpE-regulating DEGs genes in THP-1 cells.

(**B-C**) Quantitative RT-PCR analysis of gene expression in infected THP-1 cells. THP-1 cells were infected with complemented strain (Comp-MmpE) and NLS-deleted complemented (Comp-MmpE$^{\Delta NLS2}$, Comp-MmpE$^{\Delta NLS1-2}$) strains for 20 h. mRNA levels of cytokine genes (**B**); mRNA levels of genes involved in lysosomal acidification and

1149 biogenesis (**C**).

1150 Data are presented as mean ± SD from three biologically independent experiments.

1151 Statistical significance was determined by a two-tailed unpaired Student's *t*-test (*P <

1152 0.05, **P < 0.01, ***P < 0.001).

1153

Supplementary Figure 5. MmpE suppresses the expression of *VDR* and inhibits

activation of the PI3K–Akt–mTOR signaling pathway.

(A) Interaction network of DEGs associated with MmpE-binding sites. The network was

constructed using STRING v12.0 and visualized with Cytoscape, highlighting

interactions between genes potentially regulated by MmpE.

(B) KEGG pathway enrichment analysis of MmpE-binding sites identified in EGFP-

MmpE-transfected HEK293T cells revealed significant enrichment in multiple signaling

pathways, including aldosterone-regulated sodium reabsorption (hsa04925), mTOR

signaling (hsa04150), axon guidance (hsa04360), serotonergic synapse (hsa04726),

circadian entrainment (hsa04713), inflammatory mediator regulation of TRP channels

(hsa04750), cAMP signaling (hsa04024), calcium signaling (hsa04020), glutamatergic

synapse (hsa04724), and cGMP–PKG signaling (hsa04022). The bar plot shows

selected significantly enriched pathways. Bar length represents the number of MmpE-

binding sites associated with each pathway, and the color gradient indicates statistical

significance, represented as $-\log_{10}(P\text{-value})$.

(**C-D**) Quantitative RT-PCR analysis of gene expression in infected THP-1 cells. THP-

1 cells were infected with wild-type BCG (BCG/wt), MmpE-deleted (*KommpE*) strains

for 20 h. mRNA levels of PI3K-Akt pathway(**C**); mRNA levels of *VDR* gene (**D**).

(E-F) ChIP-PCR and qPCR analyses of *VDR* promoter binding by MmpE and the

phosphatase-deficient mutant MmpE-H^{348}AN^{359}A in HEK293T cells. Chromatin was

immunoprecipitated from cells transfected with EGFP control, MmpE-EGFP, or MmpE-

H^{348}AN^{359}A-EGFP constructs. PCR was performed using primers targeting the *VDR*

promoter and *GAPDH* (negative control), and products were analyzed by agarose gel

electrophoresis (**E**). ChIP enrichment for the *VDR* and *GAPDH* promoter regions was

quantified by qPCR (**F**), with enrichment calculated using the $2^{-\Delta Ct}$ method.

Data are presented as mean ± SD from three biologically independent experiments.

Statistical significance was determined by a two-tailed unpaired Student's *t*-test (***$P <$

0.001).

Supplementary Figure 6. MmpE modulates the transcription of immune-associated genes.

(**A**) Association between differentially expressed genes and differential peak-related

genes. Venn diagram showing the overlap between differential peak-related genes

identified by CUT&Tag sequencing and DEGs identified by RNA-seq analysis.

(B) Four-quadrant scatter plot comparing CUT&Tag-seq and RNA-seq results. The

$\log_2$(fold change) of peak-related genes from CUT&Tag-seq is plotted against the

$\log_2$(fold change) of FPKM values from RNA-seq analysis. The scatter plot

demonstrates the correlation between chromatin binding changes and gene

expression levels.

(C) GO enrichment analysis of DEGs revealed significant overrepresentation of

Biological Process (BP) terms in macrophages infected with Ko*mmpE* compared to

those infected with wild-type BCG (BCG/wt).

(D) Chord plot showing significantly altered genes associated with GO Biological

Process pathways in Ko*mmpE*-infected macrophages compared to BCG/wt infection.

Supplementary Figure 7. MmpE promotes mycobacterial colonization in the spleen of mice.

Bacterial colonization in the spleen. CFU assays were performed on splenic

homogenates from mice infected as described in Figure 6A. Bacterial loads (CFU/g

tissue) were quantified at 2, 14, 21, 28, and 56 days post-infection. Data are presented

as mean ± SD (n = 3). Statistical significance was determined using a two-tailed

unpaired Student's *t*-test (*P* < 0.05, **P* < 0.01, and ***P* < 0.001).