3   # Expectations and uncertainty shape pain perception during
4   learning.

6   **Mégane Lacombe-Thibault[a,b], Michel-Pierre Coll[a,b*]**
7
8
9   [a] School of Psychology, Université Laval, Québec, Québec, Canada.
10
11   [b] Centre interdisciplinaire de recherche en réadaptation et intégration sociale, Québec, Québec, Canada.
12
13
* Correspondence:
michel-pierre.coll@psy.ulaval.ca
École de psychologie,
Pavillon Félix-Antoine-Savard,
2325, rue des Bibliothèques
Université Laval, Québec (Québec) G1V 0A6
20
21
22
23

24   **Keywords:** pain | learning | uncertainty | computational modeling

30   **Acknowledgements**: MLT was supported by scholarships from the *Fonds de Recherche Québec - Santé*
31   (https://frq.gouv.qc.ca/en/) and the Canadian Institutes of Health Research (https://cihr-irsc.gc.ca/). MPC and
32   this work was supported by a career grant from the *Fonds de Recherche Québec - Santé* (grant #309041,
33   https://frq.gouv.qc.ca/en/). The funders did not play any role in the study design, data collection and analysis,
34   decision to publish or preparation of the manuscript.
35
36

37   **Conflicts of interest.** The authors declare that they have no conflict of interest.
38
39   **Data and code availability.** All raw data and code necessary to reproduce all analyses and figures in this
40   paper will be openly shared upon publication.
41
42

Abstract

Pain perception is modulated by expectations and learning processes, but the influence of uncertainty in this relationship is not well established. We aimed to examine the relationship between uncertainty, pain learning and perception using hierarchical Bayesian modeling. In an aversive learning task, fifty participants learned contingencies between auditory cues and painful stimulations under changing levels of uncertainty to create periods of stability and volatility. Model-free analysis of our data suggested unexpected trials resulted in reduced accuracy and greater response times. In unexpected trials, high pain perception was reduced, while low pain perception was increased, in line with documented effects of expectations on pain perception. Computational model fitting revealed participants' learning was best described by a two-level hierarchical gaussian filter model, suggesting participants adapted their beliefs at multiple levels during the task. Uncertainty influenced pain perception in opposite patterns for high and low pain stimulations: high pain perception was greater under high levels of uncertainty, while there was a non-significant trend for low pain perception to be reduced. Analyses of individual differences suggested depressive symptoms were associated with a reduced learning rate throughout the task. These results shed light on processes involved in pain learning in changing environments. They also suggest a possible relationship between learning alterations and psychological traits commonly found in chronic pain, such as depressive symptoms.

1. Introduction

Pain is a fundamental biological signal that serves crucial protective and learning functions [1,2]. While its immediate role in alerting us to potential tissue damage is well understood, pain's complex relationship with learning and uncertainty remains less clear. In volatile environments, where conditions change unpredictably, the ability to learn from and adapt to pain signals becomes particularly important. However, how uncertainty shapes this learning process and influences pain perception itself remains poorly understood [3,4].

Recent research suggests a fundamental link between pain perception and its learning function. Pain signals appear to carry both sensory information and learning signals in the form of prediction errors, which help update future expectations of pain [5–7]. This dual role of pain aligns with both sensory inference and reinforcement learning frameworks [8]. From a sensory inference perspective, pain perception emerges from the integration of incoming nociceptive signals with prior expectations, allowing for flexible updating of pain predictions [9,10]. Simultaneously, reinforcement learning mechanisms use these pain experiences to modify behaviour, with prediction errors serving as teaching signals that guide future actions and expectations [6,11]. Importantly, the degree to which pain carries new information contributes to modulate its perception by either increasing or decreasing pain depending on the context [3,12]. While these findings suggest that pain perception is not merely a passive sensory experience but rather an active process that combines incoming sensory information with learned expectations, how uncertainty influences this integration of learning and perception remains poorly understood.

Previous studies have mainly examined how uncertainty influences pain perception by experimentally manipulating uncertainty through various approaches. Some studies varied the probability of receiving painful

stimulation [13,14], while others manipulated the predictability of pain intensity or timing [4,15–18]. These

manipulations have yielded mixed results: some research suggests that uncertainty consistently increases

pain perception [13,14,16], while others indicate that uncertainty interacts with expectations to influence pain

experience [15,17,18]. However, these studies typically employed static manipulations of uncertainty, where

participants were explicitly informed about probabilities or experienced fixed levels of unpredictability. Few

studies have examined how people learn about and adapt to changing levels of uncertainty in pain contexts,

or how dynamically learned uncertainty affects pain perception. An exception is the work of Taylor and

collaborators [12], who found that learned uncertainty was associated with increased pain; however, their

study employed a task with limited changes in contingencies and thus with limited uncertainty. As a result,

previous research offers limited insight into the relationship between pain and uncertainty, particularly since

pain experiences frequently occur in environments where uncertainty fluctuates over time, necessitating

continuous learning and adaptation [8,10].

Hierarchical Bayesian models provide a powerful framework for understanding how uncertainty influences

pain learning. Unlike traditional associative learning models that use fixed learning rates, these models can

capture multiple forms of uncertainty that evolve dynamically over time. The Hierarchical Gaussian Filter

(HGF), for instance, represents uncertainty across different levels: from immediate uncertainty about sensory

outcomes to higher-order uncertainty about environmental volatility [19,20]. This hierarchical structure allows

modeling of different types of uncertainty in pain contexts—from uncertainty about immediate pain intensity

to uncertainty about how pain-cue relationships change over time [21,22]. These models have successfully

captured learning under uncertainty in various domains and shown, for example, that learned uncertainty is

related to stress responses [22] and illusory pain perception [23]. In the latter study, the authors found that

greater uncertainty about the cue-outcome associations amplified the perception of illusory pain. However,

the influence of various forms of uncertainty on pain perception in response to noxious stimuli has been

limited.

Here, we examined how uncertainty influences pain learning and perception using a hierarchical Bayesian

modeling approach. We hypothesized that uncertainty would increase pain perception given that pain signals

carry greater informational value when contingencies are volatile and require continuous updating of

predictions. In a task where participants learned associations between auditory cues and painful stimulations

under varying levels of uncertainty, we found that estimation uncertainty—reflecting uncertainty about

stimulus-outcome relationship—differentially affected pain ratings depending on stimulus intensity, consistent

with a sensory inference account of pain perception. We additionally explored the influence of individual

differences in learning about pain and found that depression was associated with slower updates of pain-cue

associations.

2. Methods

2.1. Participants

We aimed to recruit 50 participants based on feasibility and to obtain a sample size similar to previous studies

using a similar task [24,25]. This sample size provides adequate power (1-β = 0.80) to detect a moderate

relationship of |r| = 0.37 with a significance threshold of α = 0.05, two-tailed, for our model-free analysis. For

Bayesian model comparison, the sample size of 50 participants each completing 192 trials of the learning

task was considered sufficient based on previous studies using similar tasks. Fifty-two participants with no

history of neurological, psychiatric or pain disorders took part in the experiment. Two participants were

excluded for misidentifying a high number of stimulations (accuracies of 73.9% and 77.6%, z scores of -4.11

and -3.44 relative to the group average of 97%). The analyses were performed on the remaining 50

participants (28 females, mean age=24.38 years; SD=4.27). Written informed consent was obtained for all

participants, and this experiment was approved by the ethics committee of the *Centre intégré universitaire*

de santé et de services sociaux de la Capitale-Nationale (project #2022-2452) and participants received $20

Canadian dollars for their participation.

2.2. Electrical stimulation

Painful electrical stimuli were administered using a biphasic constant current stimulator (DS8R, Digitimer Ltd,

United Kingdom) combined with a train delay generator (DG2A, Digitimer Ltd, United Kingdom) to deliver, at

each trial, 5 monophasic stimulations of 2 ms each presented at 200 Hz. Electrical stimuli were delivered

through two 8mm Ag/AgCl cup electrodes filled with conductive gel and placed 2 centimetres apart on the

back of the participants' non-dominant hand. Prior to electrode placement, the hand was cleaned with alcohol

swabs.

2.3. Procedure

The experimental session lasted approximately 80 minutes and was divided into 3 phases: 1) Calibration, 2)

Aversive learning task and 3) Self-reported questionnaires.

2.3.1. Calibration

Pain levels for the experiment were determined using a staircase method [26]. The stimulation intensity was

increased from 1 mA in 1 mA increments until participants indicated they had reached their pain tolerance

threshold. Participants were informed a stimulation similar to their pain tolerance threshold could be used

during the task, where they would receive it repeatedly. They were asked to keep this in mind when selecting

their maximal stimulation during the calibration process. After each stimulation, participants rated the intensity

of the painful stimulation on a visual analog scale (VAS). The VAS was labelled from "no pain" to "worst

imaginable pain". Participants were also asked after each stimulation if they were willing to receive a more

intense stimulation.

This ramping procedure was repeated 3 times to ensure participants were familiar with the electrical

stimulations and rating scale. Using the results from the third ramp, the intensity at which participants first

reported perceiving pain (VAS rating > 0) was defined as the pain detection threshold, while the maximum

tolerable intensity was defined as the pain tolerance threshold, which was not exceeded during the rest of

the experiment.

The second phase of the calibration consisted of 14 equidistant electrical stimulations between the pain

detection and tolerance thresholds presented in a random order. Participants rated the intensity of each

stimulation using the same VAS as in the first part of the calibration. This process aimed to avoid expectation

effects on pain perception.

The intensities corresponding to 40% (low pain) and 90% (high pain) of the maximum pain rating were

determined by fitting various polynomial functions to the relationship between each stimulation intensity and

the corresponding VAS rating obtained during the second phase of the calibration. The function that best

represented this relationship was used to identify the low and high painful stimulations used for the rest of

the experiment. On average, the intensity of the low pain stimulation was 4.78 mA (SD=2.57, range: 1.5 mA

to 13 mA) and the intensity of the high pain stimulation was 8.93 mA (SD=4.02, range: 4 mA to 21 mA) with

an average difference of 4.21 mA (SD=1.89, range:1.25 mA to 10.5 mA) between the two stimulations. For

one participant, high and low pain stimulation intensities were reduced after the first block of the task, as the

calibrated intensities were not well tolerated. Reduced intensities are reported in the descriptive statistics

above.

2.3.2. Aversive learning task

The experimental task was based on a previously used learning paradigm in the context of volatility [25,27]

and lasted approximately 20 minutes. It included 192 trials organized into 4 blocks with a brief pause following

each block. The task began with 4 practice trials so participants could familiarize themselves with the task.

In each trial (Figure 1a), a fixation cross appeared on the screen for a random duration between 1 and 2

seconds, followed by a low (330 Hz) or a high (660 Hz) tone presented for 300 ms. The sounds were

presented at a clearly audible intensity and were easily distinguishable by participants, as verbally confirmed

by participants prior to the task. Two hundred ms after the end of the tone, a low or high intensity painful

electrical stimulus was delivered for 30 ms. With their dominant hand, participants then had to indicate as

quickly as possible within the two-second time limit whether they received the low or high intensity stimulation

by pressing one of two keys on a response box. At the end of each trial, participants had up to 3 seconds to

rate the intensity of the painful stimulus on the same VAS as used in the calibration. VAS position was

converted to a 0 to 100 score for analyses. If participants exceeded the allowed time at each step of the task,

the trial ended and was not considered in analyses.

Before the task, participants were informed that the tone frequency was associated with the probability of

receiving low or high intensity painful stimulations, with these probabilities varying throughout the task. The

tone-pain association could be strong (probability=0.84), weak (probability=0.16) or null (probability=0.5) and

was adjusted every 8, 12, 24, 36 or 48 trials to create periods of stability or volatility (Figure 1b). These exact

probabilities and their pattern of change were not disclosed to participants.

The order of stimuli presentation was adjusted so that in each block, both tones appeared as frequently and

were presented in a random order. Throughout the task, the probability of one outcome (e.g., low pain) given

a cue (e.g., low tone) was the same as the probability of the other outcome (e.g., high pain) given the other

cue (e.g., high tone). These manipulations ensured that the probability of receiving low or high pain before

the presentation of the sound was 50% on any given trial [24,27]. This ensured a similar distribution of high

and low pain stimuli across blocks.

2.3.3. Self-reported questionnaires

After the experiment, participants completed the French versions of the Pain Catastrophizing Scale (PCS;

[28,29]), the State-Trait Anxiety Inventory (STAI; [30,31]), and the Beck Depression Inventory-II (BDI-II;

[32,33]). These questionnaire data were used in exploratory analyses to examine how psychological traits

might influence aversive learning under uncertainty.

2.4. Computational modeling

To model participants' learning throughout the task, we compared several computational models: a Rescorla-

Wagner (RW) model, a Pearce-Hall (PH) model and the Hierarchical Gaussian Filter (HGF). For the HGF,

we tested four variants that differed in their complexity: models with either two or three levels, each

implemented with or without a perceptual uncertainty parameter.

For learning models, the pain outcome at each trial was contingency coded so that high pain following low

tone and low pain following low tones were coded 1 and high pain following high tone and low pain following

high tones were coded 0. These binary outcomes served as observed sensory inputs in the models. Using

these outcomes, we assessed each model's ability to predict participants' log response times using a linear

response model. We used the default implementation of each learning model in the TAPAS toolbox and we

thus refer readers to the open code (https://github.com/translationalneuromodeling/tapas, version 7.1.2) for

the update equations of each model. The required priors for the HGF models were adjusted from the default

values defined in the TAPAS toolbox based on the priors used in a previous study using a similar task [25]

(see Supplementary Table S1 for all priors for the HGF models). Trials with mistakes or response times below

200 ms were ignored in the response model fitting procedure (average number of trials included = 186.22,

SD=7.38, range = 154-192).

The models were fitted to the data using the HGF module of the TAPAS toolbox and model fits at the group

level were compared using the log model evidence (LME) within a Bayesian model comparison procedure,

as implemented in the Variational Bayesian Analysis toolbox (VBA; https://mbb-team.github.io/VBA-toolbox

14). Below, we briefly describe each model and the response models linking the trial-wise estimates derived

from these models to the log of the response time at each trial.

2.4.1. Rescorla-Wagner model

The RW model uses prediction errors to drive associative learning: larger discrepancies between actual and

expected outcomes produce greater changes in associative strength [34]. At each trial (t), the associative

strength (V) between auditory cues and pain outcomes is updated based on a learning rate parameter (α)

and the prediction error (δ)—the difference between the actual outcome (λ) and expected outcomes. The

model assumes that response times are predicted by the trial-specific absolute prediction error ($|\delta|$), with five

free parameters: the learning rate (α), initial associative strength (V_0) and the response model coefficients

and noise (β_0, β_1, ζ).

(1) $V_{(t+1)} = V_{(t)} + \alpha * \delta_{(t)}$

(2) $\delta_{(t)} = \lambda_{(t)} - V_{(t)}$

(3) $\log(RT) = \beta_0 + \beta_1 * |\delta| + \zeta$

2.4.3. Pearce-Hall model

The Pearce-Hall model has previously been used in uncertainty studies [12,35] and combines associative

learning with dynamic attention. The model assumes that response times were influenced by attention (α) at

each trial (t). The attention parameter is updated based on the absolute prediction error from the previous

trial, while associative strength (V) follows the Rescorla-Wagner learning rule. This model included five free

parameters: initial attention (α_0), attention learning rate (η), three response coefficients (β_0, β_1), plus

Gaussian noise (ζ), as specified in the linear response model below:

(4) $V_{(t+1)} = V_{(t)} + \eta * \alpha(t) * \delta(t)$

Where

(5) $\delta(t) = \lambda(t) - V(t)$

(6) $\alpha(t) = |\delta(t)|$

(7) $\log(RT) = \beta_0 + \beta_1 * \alpha + \zeta$

2.4.3. Hierarchical gaussian filter

The Hierarchical Gaussian Filter model (HGF; Figure 2a-b) assumes that response times were influenced by

uncertainty estimates across two (HGF2) or three (HGF3) hierarchical levels. The model tracks belief updates

about stimulus transitions at two or three coupled levels: stimulus outcomes (x_1), probability of stimulus

transitions (x_2), and, for the three-level version, volatility of these probabilities (x_3). The beliefs (μ) at each

level above the first evolve as Gaussian random walks, where ω_2 determines the step size of probability

updates at the second level, and in the HGF3, ω_3 determines the step size of volatility updates at the third

level. Each level is updated based on precision-weighted (ψ) prediction errors (ε, δ) from the level below.

The equations can be found in previous descriptions of the model [19,20].

For both HGF2 and HGF3, we tested versions that took into account perceptual uncertainty in pain processing

(HGF2pu and HGF3pu). In these versions, rather than treating outcomes as binary events, the model

represents them as continuous values drawn from a mixture of Gaussian distributions. The perceptual

uncertainty parameter α represents the variance of these distributions centered at 0 and 1, with higher values

indicating greater difficulty in discriminating between high and low pain intensities [19].

The HGF linear response model incorporated uncertainty estimates from each hierarchical level transformed

to the outcome level [24,25]: outcome uncertainty (unc_1), estimation uncertainty (unc_2), and, for the HGF3,

phasic volatility (unc_3). The number of free parameters varied across HGF variants: five to eight parameters

depending on model complexity. For the learning component, these included step size parameters (2nd level

tonic volatility; ω_2, and for HGF3, 3rd level volatility ω_3) and, where applicable, perceptual uncertainty (α).

The response model included coefficients for each uncertainty term ($\beta_{0\text{-}2}$ for HGF2, $\beta_{0\text{-}3}$ for HGF3) and

Gaussian noise (ζ):

(8) HGF2: $\log(RT) = \beta_0 + \beta_1 * unc_1 + \beta_2 * unc_2 + \zeta$

(9) HGF3: $\log(RT) = \beta_0 + \beta_1 * unc_1 + \beta_2 * unc_2 + \beta_3 * unc_3 + \zeta$

2.5. Statistical analyses

2.5.1. Model-free analyses

We first confirmed the influence of the variations in contingencies throughout the task on behavioural responses independently of the computational modeling procedure. To this end, we used repeated-measures analyses of variance (ANOVA) to test the effect of Trial type (expected [84% probability], unexpected [16% probability] or neutral [50% probability]) and Pain intensity (high or low) and their interaction on response times (RTs), proportion of errors and pain intensity ratings. The Greenhouse-Geisser correction was used in case of violation of the sphericity assumption and post-hoc pairwise comparisons were conducted. The level of significance was set at $p<0.05$, bilateral, for all analyses.

2.5.2 Model-based analyses

We used linear mixed-effects models to examine how trial-by-trial pain ratings were influenced by computational estimates from the winning model, the HGF2: the absolute prediction error at the first level (δ_1), outcome uncertainty ($\hat{\sigma_1}$) and estimation uncertainty ($\hat{\sigma_2}$). For each computational estimate, we constructed a separate model that included the estimate, stimulation intensity (high vs. low), and their interaction as fixed effects. All models included random intercepts and random slopes for participants for all fixed effects, accounting for individual differences in baseline ratings and in the strength of relationships between predictors and pain ratings. Estimation uncertainty and outcome uncertainty were respectively log transformed and squared to take into account their positive and negatively skewed distributions and favour model convergence.

2.5.3. Individual differences in learning, pain catastrophizing, depressive and anxiety symptoms

321 The individual differences in the PCS, BDI and STAI-Y1 and STAI-Y2 questionnaires total scores were

322 examined by correlating these values with the parameters of the winning learning model. Pearson r

323 correlations were conducted with the data from 49 participants only, as one participant had missing data on

324 the questionnaires due to a technical issue. The Holm-Bonferroni method was used to correct alpha tests.

325
Figure 1. (a) Schematic representation of an experimental trial. **(b)** Tone and pain level contingencies throughout the experimental
task showing phases in which the high or low pain level could be expected (Exp., 0.84), Neutral (Neut. 0.5) or Unexpected (Unexp.,
0.16). **(c)** Proportion of errors in identifying the pain levels, **(d)** response time **(e)** pain rating and **(f)** pain rating centered on each
pain level mean as a function of pain level and contingencies. All error bars show the 95% confidence interval of the mean.

3. Results

3.1. Model-free analyses

We confirmed that Trial type influenced both participants' accuracy [$F_{(1.644, 80.542)}$=7.423, p=0.002,

η_p^2=0.030; Figure 1c] and response times [$F_{(1.908, 93.508)}$=8.272, p<0.001, η_p^2=0.008; Figure 1d]. As

expected, participants made more errors and were slower in unexpected trials compared to both expected

[accuracy: t(49)=3.101, p=0.003, Hedge's g=0.414; RT: t(49)=-3.955, p<0.001, g=-0.236] and neutral trials

[accuracy: t(49)=2.921, p=0.005, g=0.420; RT: (t(49)=-2.137, p=0.038, g=-0.128]. Participants also made

more errors [($F_{(1,49)}$=5.027, p=0.030, η_p^2=0.016, *g*=-0.324)] and were slower [$F_{(1,49)}$=12.287, p<0.001,

$\eta_p^2=0.015$)] in high pain trials compared to low pain trials. There was no interaction between Trial type and

Pain intensity on accuracy [$F(1.627,79.697)=0.248$, $p=0.735$, $\eta_p^2<0.001$] or RTs [$F(1.677,82.169)=2.474$,

$p=0.100$, $\eta_p^2=0.002$].

We further confirmed that pain intensity ratings differed between the two administered intensities

($F(1,49)=315.539$, $p<0.001$, $\eta_p^2=0.707$; Figure 1e). Pain intensity and Trial type showed an interaction

($F(1.575,77.176)=8.259$, $p=0.001$, $\eta_p^2=0.001$): participants rated high pain stimulations lower during

unexpected trials compared to both expected ($t(49)=2.084$, $p=0.042$, $g=0.056$) and neutral trials ($t(49)=2.400$,

$p=0.020$, $g=0.073$), while ratings for low pain trials increased in unexpected conditions compared to both

expected ($t(49)=-2.151$, $p=0.036$, $g=-0.071$) and neutral trials ($t(49)=-2.616$, $p=0.012$, $g=-0.111$; Figure 1f).

349
3.2 Computational model fitting and validation

Bayesian model comparison identified the two-level HGF without perceptual uncertainty (HGF2) as the best

predictor of participants' log response times (Figure 2c; protected exceedance probability [PEP] = 0.99; model

frequency = 0.540).

To validate this model, we conducted two analyses. First, we performed parameter recovery by simulating

ten response time datasets for each participant using their fitted HGF2 parameters (500 datasets total) and

generating response times by adding random gaussian noise to the linear response model. We then fitted

the model to these simulated data and compared the recovered parameters, averaged across the ten

datasets per participant, to the original simulation parameters. This analysis revealed high Spearman ρ

correlations between simulated and recovered parameters (Figure 2d; parameter relationships illustrated in

Supplementary Figure S1c-g).

Second, we assessed model recovery by fitting all models to data simulated from each model. The HGF2

was correctly identified as the winning model for 95% of datasets simulated using its parameters with no

evidence for better fit from alternative models. However, the HGF2 was not specific in recovering itself and

showed high recovery for datasets generated using other models (Supplementary Figure S1b), further

suggesting that the HGF2 was a superior fit to our task.

3.3. Model-based analyses

Analysis of the relationship between the HGF2 model's trial-wise estimates and pain ratings revealed a

significant interaction between estimation uncertainty and pain intensity ($\beta = 8.194$, SE = 2.191, $p < 0.001$;

Figure 3a). Decomposition of this interaction revealed that estimation uncertainty was positively associated

with pain ratings in high-intensity trials ($\beta = 5.669$, SE = 1.891, $p = 0.003$), while showing a negative but not

significant relationship with pain ratings in low-intensity trials ($\beta = -2.702$, SE = 1.385, $p = 0.05$). Similarly,

there was a significant interaction between absolute prediction errors and pain ratings ($\beta = -2.994$, SE =

1.437, $p = 0.037$; Figure 3b). The decomposition of this interaction revealed that absolute prediction errors

were negatively associated with pain ratings in high pain trials (B = -2.264, SE = 1.122, $p = 0.044$) but showed

no significant relationship in low pain trials (B = 0.731, SE = 0.664, $p = 0.271$). Outcome uncertainty did not

significantly affect pain ratings ($\beta = -3.540$, SE = 10.952, $p = 0.747$), and its interaction with pain intensity

was also non-significant ($\beta = -1.221$, SE = 24.563, $p = 0.960$).

Figure 2. (a) Structure of the HGF2 model. **(b)** Example of absolute prediction error, irreducible and estimation uncertainty across trials. **(c)** Results of the Bayesian model comparison. **(d)** HGF2 model parameter recovery.

3.4. Individual differences in learning

High depressive symptoms were associated with lower tonic volatility for tone-pain associations (ω_2; r(47) = -0.41, p = 0.015; Figure 3f). Pain catastrophizing scores (r(47) = -0.29, p = 0.123; Figure 3c), state anxiety (r(47) = -0.20, p = 0.170; Figure 3d) and trait anxiety (r(47) = -0.29, p = 0.123; Figure 3e) showed non-significant negative correlations with tonic volatility.

Figure 3. Model-based analyses results of linear mixed-effects models. (a) Pain ratings by estimation uncertainty where lighter lines represent each participant's individual slope and darker bold lines represent the average slope (red for high pain intensity and blue for low pain intensity). (b) Pain ratings by absolute prediction errors. (c-f) Individual differences in PCS, STAI state, STAI trait and BDI scores and the tonic volatility parameter (ω_2). The reported p-values were corrected with the Holm-Bonferroni method.

Discussion

Pain perception is fundamentally shaped by expectations and learning. Using a dynamic learning task, we asked participants to learn associations between auditory cues and painful electrical stimulations of two different intensities. The task was designed to generate periods of high and low uncertainty by frequently changing the cue-pain contingencies throughout the experiment. Results of computational modeling revealed that under high uncertainty, high-intensity pain was perceived as more painful, while low-intensity pain was perceived as less painful. These findings extend previous research on expectation and uncertainty effects in

pain perception [4,12,18,36] by showing how learned uncertainty in a changing environment modulates pain

processing in an intensity-dependent manner—amplifying the perception of high pain while attenuating the

perception of low pain.

Model-free analyses showed expected results, as more errors and longer response times were found in

unexpected trials. Our results also suggest an opposite effect of expectations on pain perception for high and

low intensity stimuli, where high pain perception decreased and low pain perception increased in unexpected

trials. These findings are in line with Bayesian inference theories where perception is biased toward

expectations [37], as found in similar studies [18]. Although these analyses provide insight on our findings,

they fail to explain the underlying mechanisms responsible for the effect of unexpectedness. Computational

modeling allows for a more in-depth explanation of the dynamic relationships between expectations and

uncertainty and how they influence perception to optimize learning.

Our computational modeling approach revealed that learning about pain in volatile environments is best

captured by hierarchical learning processes. The two-level Hierarchical Gaussian Filter (HGF2; [19,20])

outperformed other models in predicting participants' response times, suggesting that participants

continuously updated beliefs at multiple levels. Similarly to a recent study [23], the three-level HGF3, which

additionally models estimates of environmental uncertainty, did not outperform the simpler two-level HGF

and showed poor parameter recovery performance in comparison to the HGF2. This discrepancy, when

compared to several other studies using a similar task [24,25], could be attributed to the specific contingency

structure of our task, which included fewer trials and contingency switches as well as a repetitive contingency

switching pattern compared to a random one, therefore not requiring estimations of volatility to perform

adequately.

Using the HGF2 to estimate participants' subjective perception of environmental uncertainty, we found that

its relationship with pain perception depended on stimulus intensity: under higher estimation uncertainty,

high-intensity pain was perceived as more intense, while there was a trending negative relationship for low-

intensity pain, suggesting it was perceived as less intense, though this effect did not reach statistical

significance. This pattern aligns with Bayesian frameworks of perception and suggests that under uncertainty,

the pain system shifts toward relying more on incoming sensory information [37]. It could also be the case

that the differential impact of uncertainty on pain perception depending on stimulus intensity reflects

asymmetric cost functions for pain perception and serves to optimize pain's learning and control function [6].

When facing high pain stimuli under uncertainty, the biological cost of underestimating the threat outweighs

the cost of overestimating, leading to elevated pain ratings as a protective learning mechanism for avoiding

significant threats. Conversely, for low pain stimuli, the system favors underestimation when uncertain, as

this minimizes costs associated with unnecessary defensive responses, resulting in lower pain ratings. This

could optimize learning by filtering out noisier signals of unthreatening pain. Our results could also explain

the previously observed mixed effects of uncertainty on pain perception [13,14,16–18], as studies using

different pain intensities might find apparently contradictory effects of uncertainty depending on where their

stimuli fell relative to participants' average expectations. Here, our dynamic modulation of uncertainty through

changing contingencies provides a more ecologically valid approach compared to previous studies that used

static manipulations of uncertainty or fixed probabilities.

Additionally, we found absolute prediction errors, as estimated by the HGF2, influenced pain perception

differently according to stimulus intensity: high pain was rated lower as prediction errors increased, while

there was evidence for low pain to be rated higher, although this effect did not reach statistical significance.

These results align with learning theories of pain suggesting that perception is dynamically adjusted in

response to prediction errors to optimize learning [6]. The greater malleability of high pain compared to low

pain could reflect an asymmetric cost of enhancing the salience of prediction errors. Less threatening signals

of pain could carry less informational value for harm avoidance, thus requiring less adjustment in response

to prediction errors. In contrast, more salient signals of high pain could be more flexible in response to

prediction errors and serve a greater learning purpose for threat avoidance.

Individual differences in self-reported measures of depression, anxiety and pain catastrophizing were

examined in relation to learning parameters estimated by the HGF2 model. We found a non-significant

negative trend between anxiety and pain catastrophizing and ω_2—a parameter representing the step size of

the learning rate for updating cue-outcome associations over time [19]—while participants reporting more

symptoms of depression showed significantly reduced ω_2 values. A lower ω_2 indicates reduced flexibility in

updating beliefs about how pain-cue relationships change. This finding has particular relevance for chronic

pain populations, where elevated catastrophizing, depression and anxiety are common [38] and represent

important risk factors for chronic pain development [39]. Our results suggest that these traits may be linked

with impaired ability to update pain expectations in changing environments, potentially contributing to

maladaptive pain processing. It should be noted that the three measures were moderately to highly correlated

(r = 0.5 to 0.72), suggesting that they tap into a common trait related to learning. Nevertheless, the

relationships between these traits and learning may help explain why some individuals develop chronic pain

conditions while others recover from similar injuries and thus provide a computational hypothesis for the long

theorized link between chronic pain and learning mechanisms [40,41] that should be tested in future studies.

Several limitations warrant consideration. First, while we used a similar design to previous studies [42,43],

our task employed a single contingency structure. Our findings should be tested for generalization across

different types of contingency patterns and learning environments. Second, while our findings suggest a link

between psychological characteristics and learning mechanisms, these relationships were relatively weak

and observed in a healthy sample with few individuals reporting high anxiety, depression or catastrophizing.

Testing these relationships in clinical populations will be crucial to establish their relevance to chronic pain.

Finally, while we apply our results to pain perception, similar influences of uncertainty on perception might

be observed in other phenomena, and psychological traits might be linked to general learning mechanisms

rather than being specific to pain learning.

In conclusion, our study demonstrates that pain perception is dynamically modulated by learned uncertainty,

with distinct effects depending on pain intensity, and is influenced by individual differences in psychological

characteristics. The successful application of hierarchical Bayesian models to capture these effects provides

a promising framework for understanding pain learning in complex, changing environments. Our finding that

symptoms of depression are associated with reduced learning efficiency could be particularly relevant for

understanding how individuals with chronic pain learn and deal with uncertainty in their daily lives.

References

481   **References**

1. Melzack R, Casey KL. Sensory, motivational, and central control determinants of pain: A new conceptual model. Skin Senses. 1968;1: 423–443.

2. Fields HL. How expectations influence pain. Pain. 2018;159: S3–S10. doi:10.1097/j.pain.0000000000001272

3. Coll M-P, Walden Z, Bourgoin P-A, Taylor V, Rainville P, Robert M, et al. Pain reflects the informational value of nociceptive inputs. Pain. 2024;165: e115–e125. doi:10.1097/j.pain.0000000000003254

4. Zaman J, Van Oudenhove L, Vlaeyen JWS. Uncertainty in a context of pain: disliked but also more painful? Pain. 2021;162: 995–998. doi:10.1097/j.pain.0000000000002106

5. Roy M, Shohamy D, Daw N, Jepma M, Wimmer GE, Wager TD. Representation of aversive prediction errors in the human periaqueductal gray. Nat Neurosci. 2014;17: 1607–1612. doi:10.1038/nn.3832

6. Seymour B. Pain: A Precision Signal for Reinforcement Learning and Control. Neuron. 2019;101: 1029–1041. doi:10.1016/j.neuron.2019.01.055

7. Seymour B, Dolan RJ. Emotion, Motivation, and Pain. 6. ed. In: McMahon SB, Wall PD, Melzack R, McMahon SB, editors. Wall and Melzack's textbook of pain. 6. ed. Philadelphia, Pa: Elsevier, Saunders; 2013.

8. Seymour B, Mancini F. Hierarchical models of pain: Inference, information-seeking, and adaptive control. NeuroImage. 2020;222: 117212. doi:10.1016/j.neuroimage.2020.117212

9. Büchel C, Geuter S, Sprenger C, Eippert F. Placebo Analgesia: A Predictive Coding Perspective. Neuron. 2014;81: 1223–1239. doi:10.1016/j.neuron.2014.02.042

10. Tabor A, Burr C. Bayesian Learning Models of Pain: A Call to Action. Curr Opin Behav Sci. 2019;26: 54–61. doi:10.1016/j.cobeha.2018.10.006

503   11.   Seymour B, O'Doherty JP, Dayan P, Koltzenburg M, Jones AK, Dolan RJ, et al. Temporal difference

504   models   describe   higher-order   learning   in   humans.   Nature.   2004;429:   664–667.

505   doi:10.1038/nature02581

506   12.   Taylor VA, Chang L, Rainville P, Roy M. Learned expectations and uncertainty facilitate pain during

507   classical conditioning. Pain. 2017;158: 1528–1537. doi:10.1097/j.pain.0000000000000948

508   13.   Ploghaus A, Narain C, Beckmann CF, Clare S, Bantick S, Wise R, et al. Exacerbation of Pain by Anxiety

509   Is   Associated   with   Activity   in   a   Hippocampal   Network.   J   Neurosci.   2001;21:   9896–9903.

510   doi:10.1523/JNEUROSCI.21-24-09896.2001

511   14.   Rhudy JL, Meagher MW. Fear and anxiety: divergent effects on human pain thresholds. Pain. 2000;84:

512   65–75. doi:10.1016/S0304-3959(99)00183-9

513   15.   Pavy F, Zaman J, Von Leupoldt A, Torta DM. Expectations underlie the effects of unpredictable pain:

514   a   behavioral   and   electroencephalogram   study.   Pain.   2024;165:   596–607.

515   doi:10.1097/j.pain.0000000000003046

516   16.   Yoshida W, Seymour B, Koltzenburg M, Dolan RJ. Uncertainty Increases Pain: Evidence for a Novel

517   Mechanism of Pain Modulation Involving the Periaqueductal Gray. J Neurosci. 2013;33: 5638–5646.

518   doi:10.1523/JNEUROSCI.4984-12.2013

519   17.   Zaman J, Vanpaemel W, Aelbrecht C, Tuerlinckx F, Vlaeyen JWS. Biased pain reports through

520   vicarious information: A computational approach to investigate the role of uncertainty. Cognition.

521   2017;169: 54–60. doi:10.1016/j.cognition.2017.07.009

522   18.   Brown CA, Seymour B, Boyle Y, El-Deredy W, Jones AKP. Modulation of pain ratings by expectation

523   and uncertainty: Behavioral characteristics and anticipatory neural correlates. Pain. 2008;135: 240–

524   250. doi:10.1016/j.pain.2007.05.022

525   19.   Mathys C, Daunizeau J, Friston KJ, Stephan KE. A bayesian foundation for individual learning under

526   uncertainty. Front Hum Neurosci. 2011;5: 39. doi:10.3389/fnhum.2011.00039

20.   Mathys CD, Lomakina EI, Daunizeau J, Iglesias S, Brodersen KH, Friston KJ, et al. Uncertainty in

perception and the Hierarchical Gaussian Filter. Front Hum Neurosci. 2014;8.

doi:10.3389/fnhum.2014.00825

21.   Anchisi D, Zanon M. A Bayesian Perspective on Sensory and Cognitive Integration in Pain Perception

and Placebo Analgesia. Daunizeau J, editor. PLOS ONE. 2015;10: e0117270.

doi:10.1371/journal.pone.0117270

22.   de Berker AO, Rutledge RB, Mathys C, Marshall L, Cross GF, Dolan RJ, et al. Computations of

uncertainty mediate acute stress responses in humans. Nat Commun. 2016;7.

doi:10.1038/ncomms10996

23.   Ehmsen JF, Nikolova N, Christensen DE, Banellis L, Böhme RA, Brændholt M, et al. Thermosensory

predictive coding underpins an illusion of pain. Sci Adv. 2025;11: eadq0261.

doi:10.1126/sciadv.adq0261

24.   Iglesias S, Mathys C, Brodersen KH, Kasper L, Piccirelli M, den Ouden HEM, et al. Hierarchical

Prediction Errors in Midbrain and Basal Forebrain during Sensory Learning. Neuron. 2013;80: 519–

530. doi:10.1016/j.neuron.2013.09.009

25.   Lawson RP, Mathys C, Rees G. Adults with autism overestimate the volatility of the sensory

environment. Nat Neurosci. 2017;20: 1293–1299. doi:10.1038/nn.4615

26.   Vlaev I, Seymour B, Dolan RJ, Chater N. The Price of Pain and the Value of Suffering. Psychol Sci.

2009;20: 309–317. doi:10.1111/j.1467-9280.2009.02304.x

27.   den Ouden HEM, Daunizeau J, Roiser J, Friston KJ, Stephan KE. Striatal Prediction Error Modulates

Cortical Coupling. J Neurosci. 2010;30: 3210–3219. doi:10.1523/JNEUROSCI.4458-09.2010

28.   Sullivan MJL, Bishop SR, Pivik J. The Pain Catastrophizing Scale: Development and validation. Psychol

Assess. 1995;7: 524–532. doi:10.1037/1040-3590.7.4.524

29.   French DJ, Noël M, Vigneau F, French JA, Cyr CP, Evans RT. L'Échelle de dramatisation face à la

551   douleur PCS-CF: Adaptation canadienne en langue française de l'échelle «Pain Catastrophizing

552   Scale». Can J Behav Sci Rev Can Sci Comport. 2005;37: 181–192. doi:10.1037/h0087255

553   30.   Spielberger CD. State-Trait Anxiety Inventory for Adults. 1983. doi:10.1037/t06496-000

554   31.   Gauthier J, Bouchard S. Adaptation canadienne-française de la forme révisée du State–Trait Anxiety

555   Inventory de Spielberger. Rev Can Sci Comport. 1993;25: 559–578. doi:10.1037/h0078881

556   32.   Beck AT. An Inventory for Measuring Depression. Arch Gen Psychiatry. 1961;4: 561.

557   doi:10.1001/archpsyc.1961.01710120031004

558   33.   Bourque P, Beaudette D. Étude psychométrique du questionnaire de dépression de Beck auprès d'un

559   échantillon d'étudiants universitaires francophones. Can J Behav Sci Rev Can Sci Comport. 1982;14:

560   211–218. doi:10.1037/h0081254

561   34.   Rescorla RA, Wagner A. A theory of Pavlovian conditioning: Variations in the effectiveness of

562   reinforcement and nonreinforcement. In: Black A, Prokasy W, editors. Classical Conditioning II: Current

563   Research and Theory. New York: Appleton- Century-Crofts; 1972. pp. 64–99.

564   35.   Taylor VA, Roy M, Chang L, Gill L-N, Mueller C, Rainville P. Reduced Fear-Conditioned Pain

565   Modulation in Experienced Meditators: A Preliminary Study. Psychosom Med. 2018;80: 799–806.

566   doi:10.1097/PSY.0000000000000634

567   36.   Koyama T, McHaffie JG, Laurienti PJ, Coghill RC. The subjective experience of pain: Where

568   expectations become reality. Proc Natl Acad Sci. 2005;102: 12950–12955.

569   doi:10.1073/pnas.0408576102

570   37.   Hoskin R, Berzuini C, Acosta-Kane D, El-Deredy W, Guo H, Talmi D. Sensitivity to pain expectations:

571   A Bayesian model of individual differences. Cognition. 2019;182: 127–139.

572   doi:10.1016/j.cognition.2018.08.022

573   38.   Brouwer B, Waardenburg S, Jacobs C, Overdijk M, Leue C, Köke A, et al. Biopsychosocial baseline

574   values of 15 000 patients suffering from chronic pain: Dutch DataPain study. Reg Anesth Pain Med.

575   2020;45: 774–782. doi:10.1136/rapm-2020-101476

576   39.   Mills SEE, Nicolson KP, Smith BH. Chronic pain: a review of its epidemiology and associated factors

577   in population-based studies. Br J Anaesth. 2019;123: e273–283. doi:10.1016/j.bja.2019.03.023

578   40.   Apkarian AV, Baliki MN, Geha PY. Towards a theory of chronic pain. Prog Neurobiol. 2009;87: 81–97.

579   doi:10.1016/j.pneurobio.2008.09.018

580   41.   Zaman J, Vlaeyen JWS, Van Oudenhove L, Wiech K, Van Diest I. Associative fear learning and

581   perceptual discrimination: A perceptual pathway in the development of chronic pain. Neurosci Biobehav

582   Rev. 2015;51: 118–125. doi:10.1016/j.neubiorev.2015.01.009

583   42.   Atlas LY, Dildine TC, Palacios-Barrios EE, Yu Q, Reynolds RC, Banker LA, et al. Instructions and

584   experiential learning have similar impacts on pain and pain-related brain responses but produce

585   dissociations in value-based reversal learning. eLife. 2022;11: e73353. doi:10.7554/eLife.73353

586   43.   Horing B, Büchel C. The human insula processes both modality-independent and pain-selective

587   learning   signals.   Summerfield   C,   editor.   PLOS   Biol.   2022;20:   e3001540.

588   doi:10.1371/journal.pbio.3001540

Supplementary Materials

S1.1. Priors for the HGF models

Table S1. Priors for the different HGF perceptual and response models used during model fitting.

Parameter name	Prior mean	Prior variance
Perceptual model		
Tonic volatility (level 2: ω_2)	-5	16
Tonic volatility (level 3: ω_3)	-6	16
Perceptual uncertainty (α)	log(0.5)	1
Response model		
Baseline RT (β_0)	log(760)	4
RT predictors (β)	0	4
Decision noise (ζ)	log(log(20))	log(2)

S1.2. Model recovery

Model recovery performance for other models was also compared and suggests the RW model (proportion = 44%) had the best recovery performance while the HGF3pu model (proportion = 28%) had moderate recovery performance (Figure S1). All other models showed poor model recovery performance. Analyses of model confusion suggest the HGF2 model was consistently mistaken with the HGF2pu (proportion = 100%), HGF3 (proportion = 78%), HGF3pu (proportion = 64%), RW (proportion = 56%) and PH models (proportion = 90%; Figure S1a). Protected exceedance probability (PEP) analyses suggested the HGF2 model clearly outperformed itself as well as the HGF2pu, HGF3, HGF3pu, RW and PH models (PEPs = 1.00; Figure S1b).

608
609 **Figure S1.** (a) Proportion and (b) protected exceedance probability of simulated and recovered models. (c-g) Spearman ρ
610 correlations between simulated and mean recovered parameters of winning models (HGF-3pu).
611

S1.3. Full results for statistical analyses of model-free results

Proportion of errors

3 (trial type) x 2 (stimulus intensity) ANOVA

	Statistics
Trial type	$F(1.644,80.542)=7.423$, $p=0.002$, $\eta_p^2=0.030$
Stimulus intensity	$F(1,49)=5.027$, $p=0.030$, $\eta_p^2=0.016$
Trial type x stimulus intensity	$F(1.627,79.697)=0.248$, $p=0.735$, $\eta_p^2=0.001$

Pain ratings

3 (trial type) x 2 (stimulus intensity) ANOVA

	Statistics
Trial type	$F(1.696,83.078)=0.632$, $p=0.509$, $\eta_p^2<0.001$
Stimulus intensity	$F(1,49)=315.539$, $p<0.001$, $\eta_p^2=0.707$
Trial type x stimulus intensity	$F(1.575,77.176)=8.259$, $p=0.001$, $\eta_p^2=0.001$

Response times

3 (trial type) x 2 (stimulus intensity) ANOVA

	Statistics
Trial type	$F(1.908,93.508)=8.272$, $p<0.001$, $\eta_p^2=0.008$

Stimulus intensity	$F(1,49)=12.287$, $p<0.001$, $\eta_p^2=0.015$				
Trial type x stimulus intensity	$F(1.677,82.169)=2.474$, $p=0.100$, $\eta_p^2=0.002$				

Proportion of errors

Comparison		T	df	p	BF10	Hedge's g
Trial type	Expected v. Neutral	0.055	49	0.956	0.154	0.007
	Expected v. Unexpected	3.101	49	0.003	10.152	0.414
	Neutral v. Unexpected	2.921	49	0.005	6.557	0.420
Stimulus intensity	Low v. High	-2.242	49	0.030	1.505	-0.324
Stimulus intensity (high) v. trial type	Expected v. Neutral	0.469	49	0.641	0.171	0.069
	Expected v. Unexpected	2.609	49	0.012	3.209	0.347
	Neutral v. Unexpected	1.846	49	0.071	0.739	0.291
Stimulus intensity (low) v. trial type	Expected v. Neutral	-0.473	49	0.638	0.171	-0.083
	Expected v. Unexpected	2.164	49	0.035	1.296	0.333
	Neutral v. Unexpected	2.259	49	0.028	1.554	0.401

Pain ratings

Comparison		T	df	p	BF10	Hedge's g
Trial type	Expected v. Neutral	-0.106	49	0.916	0.155	-0.002
	Expected v. Unexpected	0.960	49	0.342	0.238	0.023
	Neutral v. Unexpected	0.858	49	0.395	0.218	0.025
Stimulus intensity	Low v. High	17.763	49	<0.001	7.379e19	3.072

Stimulus intensity (high) v. trial type	Expected v. Neutral	-1.100	49	0.277	0.272	-0.019
	Expected v. Unexpected	2.084	49	0.042	1.118	0.056
	Neutral v. Unexpected	2.400	49	0.020	2.061	0.073
Stimulus intensity (low) v. trial type	Expected v. Neutral	1.057	49	0.296	0.260	0.041
	Expected v. Unexpected	-2.151	49	0.036	1.265	-0.071
	Neutral v. Unexpected	-2.616	49	0.012	3.262	-0.111

Response times

Comparison		T	df	p	BF10	Hedge's g
Trial type	Expected v. Neutral	-1.975	49	0.054	0.919	-0.101
	Expected v. Unexpected	-3.955	49	<0.001	100.812	-0.236
	Neutral v. Unexpected	-2.137	49	0.038	1.232	-0.128
Stimulus intensity	Low v. High	3.505	49	<0.001	28.855	0.253
Stimulus intensity (high) v. trial type	Expected v. Neutral	-2.094	49	0.041	1.138	-0.131
	Expected v. Unexpected	-2.266	49	0.028	1.577	-0.161
	Neutral v. Unexpected	-0.367	49	0.716	0.164	-0.027
Stimulus intensity (low) v. trial type	Expected v. Neutral	-0.999	49	0.323	0.246	-0.057
	Expected v. Unexpected	-4.203	49	<0.001	208.023	-0.299
	Neutral v. Unexpected	-2.886	49	0.006	6.029	-0.227