Expectations and uncertainty shape pain perception during learning.

6   **Mégane Lacombe-Thibault[a,b], Michel-Pierre Coll[a,b]***

7
8
9   [a] School of Psychology, Université Laval, 2325 Rue des Bibliothèques, G1V 0A6, Québec, Québec, Canada.

10
11 [b] Centre interdisciplinaire de recherche en réadaptation et intégration sociale, 525 Boulevard Wilfrid-Hamel,
12   G1M 2S8, Québec, Québec, Canada.

13
14
* Correspondence:
Michel-Pierre Coll
michel-pierre.coll@psy.ulaval.ca
École de psychologie,
Pavillon Félix-Antoine-Savard,
2325, rue des Bibliothèques
Université Laval, Québec (Québec) G1V 0A6

22
23
24

Abstract

Pain perception is modulated by expectations and learning processes, but the influence of uncertainty in this relationship is not well established. We aimed to examine the relationship between uncertainty, pain learning and perception using hierarchical Bayesian modeling. In an aversive learning task, fifty participants learned contingencies between auditory cues and painful stimulations under changing levels of uncertainty to create periods of stability and volatility. Model-free analysis of our data suggested unexpected trials resulted in reduced accuracy and greater response times. In unexpected trials, high pain perception was reduced, while low pain perception was increased, in line with documented effects of expectations on pain perception. Computational model fitting revealed participants' learning was best described by a two-level hierarchical gaussian filter model, suggesting participants adapted their beliefs at multiple levels during the task. Uncertainty influenced pain perception in opposite patterns for high and low pain stimulations: high pain perception was greater under high levels of uncertainty, while there was a non-significant trend for low pain perception to be reduced. Analyses of individual differences suggested depressive symptoms were associated with a reduced learning rate throughout the task. These results shed light on processes involved in pain learning in changing environments. They also suggest a possible relationship between learning alterations and psychological traits commonly found in chronic pain, such as depressive symptoms.

Perspective: This article explores the influence of varying levels of uncertainty on pain perception and learning. Findings reveal that uncertainty modulates pain perception differently depending on pain intensity and that pain learning is influenced by psychological traits. These results contribute to our understanding of pain modulation in dynamic environments.

Keywords: pain | learning | expectations | uncertainty | computational modeling

Introduction

Pain is a fundamental biological signal that serves crucial protective and learning functions[1,2]. While its immediate role in alerting us to potential tissue damage is well understood, pain's complex relationship with learning and uncertainty remains less clear. In volatile environments, where conditions change unpredictably, learning from pain signals becomes particularly important. However, how uncertainty shapes this process and influences pain perception remains poorly understood[3,4].

Recent research suggests a fundamental link between pain perception and its learning function. Pain signals appear to carry sensory information and learning signals as prediction errors, which help update future expectations of pain[5–7]. This dual role of pain aligns with both sensory inference and reinforcement learning frameworks[8]. From a sensory inference perspective, pain perception integrates incoming nociceptive signals with prior expectations, allowing for flexible updating of pain predictions[9,10]. Simultaneously, reinforcement learning mechanisms use pain to modify behaviour, with prediction errors serving as teaching signals that guide future actions and expectations[6,11]. The degree to which pain carries new information modulates its perception by increasing or decreasing pain depending on the context[3,12]. While these findings suggest that pain perception is not merely a passive sensory experience but rather an active process combining sensory information with learned expectations, how uncertainty influences this integration remains poorly understood.

Previous studies have examined how uncertainty influences pain perception by experimentally manipulating uncertainty through various approaches. Some varied the probability of receiving pain[13,14], while others manipulated the predictability of pain intensity or timing[4,15–18]. These manipulations have yielded mixed results: some research suggests that uncertainty increases pain perception[13,14,16], while others indicate that uncertainty interacts with expectations to influence pain experience[15,17,18]. However, these studies typically employed static manipulations of uncertainty, where participants were explicitly informed about probabilities

or experienced fixed levels of unpredictability. Few studies have examined how people learn about changing

levels of uncertainty in pain contexts, or how it affects pain perception. An exception is the work of Taylor

and collaborators[12], who found that learned uncertainty was associated with increased pain; however, they

employed a task with limited changes in contingencies and thus, limited uncertainty. Consequently, previous

research offers limited insight into the relationship between pain and uncertainty, particularly since pain

experiences frequently occur in environments where uncertainty fluctuates, necessitating continuous

learning[8,10].

Hierarchical Bayesian models provide a powerful framework for understanding how uncertainty influences

pain learning. Unlike traditional associative learning models with fixed learning rates, these models capture

multiple forms of uncertainty that evolve dynamically. The Hierarchical Gaussian Filter (HGF), for instance,

represents uncertainty from immediate uncertainty about sensory outcomes to higher-order uncertainty about

environmental volatility[19,20]. This hierarchical structure allows modeling of different types of uncertainty in

pain contexts—from uncertainty about immediate pain intensity to uncertainty about how pain-cue

relationships change over time[21,22]. These models have successfully captured learning under uncertainty in

various domains and shown, for example, that learned uncertainty is related to stress responses[22] and illusory

pain perception[23]. However, the influence of various forms of uncertainty on pain perception in response to

noxious stimuli has been limited.

Using hierarchical Bayesian modeling, we integrated principles from the sensory inference and reinforcement

learning frameworks to assess how pain perception is influenced by both sensory prediction errors and

uncertainty. We hypothesized that uncertainty would increase pain perception, as pain signals carry greater

informational value in volatile contexts requiring continuous updating of predictions. In a task where

participants learned associations between auditory cues and painful stimulations under varying levels of

uncertainty, we found that estimation uncertainty—reflecting uncertainty about stimulus-outcome

relationships—differentially affected pain ratings depending on stimulus intensity. We additionally found that

depression was associated with slower updates of pain-cue associations.

103   **Methods**

Participants

We aimed to recruit 50 participants based on feasibility and to obtain a sample size similar to previous studies

using a similar task[24,25]. This sample size provides adequate power ($1-\beta = 0.80$) to detect a moderate

relationship of $|r| = 0.37$ with a significance threshold of $\alpha = 0.05$, two-tailed, for our model-free analysis. For

Bayesian model comparison, the sample size of 50 participants each completing 192 trials of the learning

task was considered sufficient based on previous studies using similar tasks. Fifty-two participants with no

history of neurological, psychiatric or pain disorders took part in the experiment. Two participants were

excluded for misidentifying a high number of stimulations (accuracies of 73.9% and 77.6%, z scores of -4.11

and -3.44 relative to the group average of 97%). The analyses were performed on the remaining 50

participants which included 28 participants who self-identified as cisgender women and 22 as cisgender men

(mean age=24.38 years; SD=4.27). Forty-three participants identified as White (86%), 2 as Asian (4%), 2 as

Latino/a American (4%), 1 as Black (2%), 1 as Middle Eastern (2%) and 1 as North African (2%). To

encourage diversity and inclusion in our sample, we employed a broad recruitment strategy that included

emails sent to the entire university community as well as posters in various locations across campus and

within the research center. No members of the public or patients were involved in this study's design or

implementation.

Written informed consent was obtained for all participants, and this experiment was approved by the ethics

committee of the *Centre intégré universitaire de santé et de services sociaux de la Capitale-Nationale* (project

#2022-2452) and participants received $20 Canadian dollars for their participation.

Electrical stimulation

Painful electrical stimuli were administered using a biphasic constant current stimulator (DS8R, Digitimer Ltd,

United Kingdom) combined with a train delay generator (DG2A, Digitimer Ltd, United Kingdom) to deliver, at

each trial, 5 monophasic stimulations of 2 ms each presented at 200 Hz. Electrical stimuli were delivered

through two 8mm Ag/AgCl cup electrodes filled with conductive gel and placed 2 centimetres apart on the

back of the participants' non-dominant hand. Prior to electrode placement, the hand was cleaned with alcohol

swabs.

Procedure

133 The experimental session lasted approximately 80 minutes and was divided into 3 phases: 1) Calibration, 2)

134 Aversive learning task and 3) Self-reported questionnaires.

Calibration

Pain levels for the experiment were determined using a staircase method[26]. The stimulation intensity was

increased from 1 mA in 1 mA increments until participants indicated they had reached their pain tolerance

threshold. Participants were informed a stimulation similar to their pain tolerance threshold could be used

during the task, where they would receive it repeatedly. They were asked to keep this in mind when selecting

their maximal stimulation during the calibration process. After each stimulation, participants rated the intensity

of the painful stimulation on a visual analog scale (VAS). The VAS was labelled from "no pain" to "worst

imaginable pain". Participants were also asked after each stimulation if they were willing to receive a more

intense stimulation.

This ramping procedure was repeated 3 times to ensure participants were familiar with the electrical

stimulations and rating scale. Using the results from the third ramp, the intensity at which participants first

reported perceiving pain (VAS rating > 0) was defined as the pain detection threshold, while the maximum

tolerable intensity was defined as the pain tolerance threshold, which was not exceeded during the rest of

the experiment.

The second phase of the calibration consisted of 14 equidistant electrical stimulations between the pain

detection and tolerance thresholds presented in a random order. Participants rated the intensity of each

stimulation using the same VAS as in the first part of the calibration. This process aimed to avoid expectation

effects on pain perception.

The intensities corresponding to 40% (low pain) and 90% (high pain) of the maximum pain rating were

determined by fitting various polynomial functions to the relationship between each stimulation intensity and

the corresponding VAS rating obtained during the second phase of the calibration. The function that best

represented this relationship was used to identify the low and high painful stimulations used for the rest of

the experiment. On average, the intensity of the low pain stimulation was 4.78 mA (SD=2.57, range: 1.5 mA

to 13 mA) and the intensity of the high pain stimulation was 8.93 mA (SD=4.02, range: 4 mA to 21 mA) with

an average difference of 4.21 mA (SD=1.89, range:1.25 mA to 10.5 mA) between the two stimulations. For

one participant, high and low pain stimulation intensities were reduced after the first block of the task, as the

calibrated intensities were not well tolerated. Reduced intensities are reported in the descriptive statistics

above.

Aversive learning task

The experimental task was based on a previously used learning paradigm in the context of volatility[25,27] and

lasted approximately 20 minutes. It included 192 trials organized into 4 blocks with a brief pause following

each block. The task began with 4 practice trials so participants could familiarize themselves with the task.

In each trial (Figure 1a), a fixation cross appeared on the screen for a random duration between 1 and 2

seconds, followed by a low (330 Hz) or a high (660 Hz) tone presented for 300 ms. The sounds were

presented at a clearly audible intensity and were easily distinguishable by participants, as verbally confirmed

by participants prior to the task. Two hundred ms after the end of the tone, a low or high intensity painful

electrical stimulus was delivered for 30 ms. With their dominant hand, participants then had to indicate as

quickly as possible within the two-second time limit whether they received the low or high intensity stimulation

by pressing one of two keys on a response box. At the end of each trial, participants had up to 3 seconds to

rate the intensity of the painful stimulus on the same VAS as used in the calibration. VAS position was

converted to a 0 to 100 score for analyses. If participants exceeded the allowed time at each step of the task,

the trial ended and was not considered in analyses.

Although participants were not informed of the exact task structure and pattern of tone-pain association

changes shown in Figure 1b, they were aware that the tone frequency was associated with the probability of

receiving low or high intensity painful stimulations. The tone-pain probability structure was predetermined

and changed every 8, 12, 24, 36 or 48 trials (blocks), creating periods of stability and volatility (Figure 1b).

Within each block, trials were pseudo-randomly presented to match the block's length and probability

structure. During each block, the probability of receiving high or low pain after a given tone could be strong

(probability=0.84), while the pain-intensity probability for the other tone was weak (probability=0.16), or both

probabilities could be null (probability=0.5). For model free analyses, we used the same labels as previous

studies[25,27] and we label these outcomes as expected (P(pain level|tone) = 0.84), unexpected (P(pain

level|tone) = 0.16) and neutral (P(pain level|tone) = 0.5) to reflect the outcome probabilities defined by the

task structure.

The order of stimuli presentation was adjusted so that in each block, both tones appeared as frequently and

were presented in a random order. Throughout the task, the probability of one outcome (e.g., low pain) given

a cue (e.g., low tone) was the same as the probability of the other outcome (e.g., high pain) given the other

cue (e.g., high tone). These manipulations ensured that the probability of receiving low or high pain before

the presentation of the sound was 50% on any given trial[24,27]. This ensured a similar distribution of high and

low pain stimuli across blocks.

Self-reported questionnaires

After the experiment, participants completed the French versions of the Pain Catastrophizing Scale

(PCS[28,29]), the State-Trait Anxiety Inventory (STAI[30,31]), and the Beck Depression Inventory-II (BDI-II[32]).

These questionnaire data were used in exploratory analyses to examine how psychological traits might

influence aversive learning under uncertainty.

Computational modeling

To model participants' learning throughout the task, we compared several computational models: a Rescorla-

Wagner (RW) model, a Pearce-Hall (PH) model and the Hierarchical Gaussian Filter (HGF). For the HGF,

we tested four variants that differed in their complexity: models with either two or three levels, each

implemented with or without a perceptual uncertainty parameter.

For the learning models, pain outcomes at each trial were coded in contingency space to account for the

tone-pain associations, as recommended in the HGF toolbox[19,20] and previous studies using a similar

task[25,33]. Specifically, high or low pain following low tones were coded 1 and high or low pain following high

tones were coded 0, therefore reflecting the contingency between cue (tones) and outcome (pain intensity).

These binary outcomes served as observed sensory inputs in the models, capturing both the cue presented

and the resulting sensory outcome. Using these outcomes as observed sensory inputs, we assessed each

model's ability to predict participants' log response times using a linear response model. We used the default

implementation of each learning model in the TAPAS toolbox[34] and we thus refer readers to the open code

(https://github.com/translationalneuromodeling/tapas, version 7.1.2) for the update equations of each model.

The required priors for the HGF models were adjusted from the default values defined in the TAPAS toolbox

based on the priors used in a previous study using a similar task[25] (see Supplementary Table S1 for all priors

for the HGF models). Trials with mistakes or response times below 200 ms were ignored in the response

model fitting procedure (average number of trials included = 186.22, SD=7.38, range = 154-192).

The models were fitted to the data using the HGF module of the TAPAS toolbox[34]. Model fits at the group

level were compared within a Bayesian model comparison procedure, as implemented in the Variational

Bayesian Analysis toolbox (VBA; https://mbb-team.github.io/VBA-toolbox). Specifically, model fit was

quantified using the log model evidence (LME) for each participant, which represents the negative surprise

of the data under a given model[19] and offers a trade-off between model fit and complexity[25]. Model selection

was done by performing group-level estimations of the frequency of a given model being the best across

participants and the protected exceedance probability of each model, representing the probability that a given

model is more likely than others[35]. The model with the highest exceedance probability and most frequently

identified as the best model was considered the winning model.

Below, we briefly describe each model and the response models linking the trial-wise estimates derived from

these models to the log of the response time at each trial.

Rescorla-Wagner model

The RW model uses prediction errors to drive associative learning: larger discrepancies between actual and

expected outcomes produce greater changes in associative strength[36]. At each trial (t), the associative

strength (V) between auditory cues and pain outcomes is updated based on a learning rate parameter (α)

and the prediction error (δ)—the difference between the actual outcome (λ) and expected outcomes. The

model assumes that response times are predicted by the trial-specific absolute prediction error ($|\delta|$), with five

free parameters: the learning rate (α), initial associative strength (V_0) and the response model coefficients

and noise (β_0, β_1, ζ).

(1) $V_{(t+1)} = V_{(t)} + \alpha * \delta_{(t)}$

(2) $\delta_{(t)} = \lambda_{(t)} - V_{(t)}$

(3) $\log(RT) = \beta_0 + \beta_1 * |\delta| + \zeta$

Pearce-Hall model

The Pearce-Hall model has previously been used in uncertainty studies[12,37] and combines associative

learning with dynamic attention. The model assumes that response times were influenced by attention (α) at

each trial (t). The attention parameter is updated based on the absolute prediction error from the previous

trial, while associative strength (V) follows the Rescorla-Wagner learning rule. This model included five free

parameters: initial attention (α_0), attention learning rate (η), three response coefficients (β_0, β_1), plus

Gaussian noise (ζ), as specified in the linear response model below:

(4) $V_{(t+1)} = V_{(t)} + \eta * \alpha(t) * \delta(t)$

Where

(5) $\delta(t) = \lambda(t) - V(t)$

(6) $\alpha(t) = |\delta(t)|$

(7) $\log(RT) = \beta_0 + \beta_1 * \alpha + \zeta$

Hierarchical gaussian filter

The Hierarchical Gaussian Filter model (HGF; Figure 2a-b) assumes that response times were influenced by

uncertainty estimates across two (HGF2) or three (HGF3) hierarchical levels. The model tracks belief updates

about stimulus transitions at two or three coupled levels: stimulus outcomes (x_1), probability of stimulus

transitions (x_2), and, for the three-level version, volatility of these probabilities (x_3). The beliefs (μ) at each

level above the first evolve as Gaussian random walks, where ω_2 determines the step size of probability

updates at the second level, and in the HGF3, ω_3 determines the step size of volatility updates at the third

level. Each level is updated based on precision-weighted (ψ) prediction errors (ε, δ) from the level below.

The equations can be found in previous descriptions of the model[19,20].

For both HGF2 and HGF3, we tested versions that took into account perceptual uncertainty in pain processing

(HGF2pu and HGF3pu). In these versions, rather than treating outcomes as binary events, the model

represents them as continuous values drawn from a mixture of Gaussian distributions. The perceptual

uncertainty parameter α represents the variance of these distributions centered at 0 and 1, with higher values

indicating greater difficulty in discriminating between high and low pain intensities[19].

The HGF linear response model incorporated uncertainty estimates from each hierarchical level transformed

to the outcome level[24,25]: outcome uncertainty (unc_1), estimation uncertainty (unc_2), and, for the HGF3,

phasic volatility (unc_3). The number of free parameters varied across HGF variants: five to eight parameters

depending on model complexity. For the learning component, these included step size parameters (2nd level

tonic volatility; ω_2, and for HGF3, 3rd level volatility ω_3) and, where applicable, perceptual uncertainty (α).

The response model included coefficients for each uncertainty term (β_{0-2} for HGF2, β_{0-3} for HGF3) and

Gaussian noise (ζ):

(8) HGF2: $\log(RT) = \beta_0 + \beta_1 * unc_1 + \beta_2 * unc_2 + \zeta$

(9) HGF3: $\log(RT) = \beta_0 + \beta_1 * unc_1 + \beta_2 * unc_2 + \beta_3 * unc_3 + \zeta$

Statistical analyses

Model-free analyses

We first confirmed the influence of the variations in contingencies throughout the task on behavioural

responses independently of the computational modeling procedure. To this end, we used repeated-measures

analyses of variance (ANOVA) to test the effect of Trial type (expected [84% probability], unexpected [16%

probability] or neutral [50% probability]) and Pain intensity (high or low) and their interaction on response

times (RTs), proportion of errors and pain intensity ratings. The Greenhouse-Geisser correction was used in

case of violation of the sphericity assumption and post-hoc pairwise comparisons were conducted. The level

of significance was set at $p < 0.05$, bilateral, for all analyses.

Model-based analyses

We used linear mixed-effects models to examine how trial-by-trial pain ratings were influenced by

computational estimates from the winning model, the HGF2: the absolute prediction error at the first level

(\delta_1), outcome uncertainty (\hat{\sigma_1}) and estimation uncertainty (\hat{\sigma_2}). For each

computational estimate, we constructed a separate model that included the estimate, stimulation intensity

(high vs. low), and their interaction as fixed effects. All models included random intercepts and random slopes

for participants for all fixed effects, accounting for individual differences in baseline ratings and in the strength

of relationships between predictors and pain ratings. Estimation uncertainty and outcome uncertainty were

respectively log transformed and squared to take into account their positive and negatively skewed

distributions and favour model convergence.

Individual differences in learning, pain catastrophizing, depressive and anxiety symptoms

The individual differences in the PCS, BDI and STAI-Y1 and STAI-Y2 questionnaires total scores were

examined by correlating these values with the parameters of the winning learning model. Pearson r

correlations were conducted with the data from 49 participants only, as one participant had missing data on

the questionnaires due to a technical issue. The Holm-Bonferroni method was used to correct alpha tests.

Figure 1. (a) Schematic representation of an experimental trial. (b) Tone and pain level contingencies

throughout the experimental task showing phases in which the high or low pain level could be expected (Exp.,

0.84), Neutral (Neut. 0.5) or Unexpected (Unexp., 0.16). (c) Proportion of errors in identifying the pain levels,

(d) response time (e) pain rating and (f) pain rating centered on each pain level mean as a function of pain

level and contingencies. All error bars show the 95% confidence interval of the mean.

327
328   **Results**

Model-free analyses

We confirmed that Trial type influenced both participants' accuracy [$F_{(1.644,80.542)}$=7.423, p=0.002,

η_p^2=0.030; Figure 1c] and response times [$F_{(1.908,93.508)}$=8.272, p<0.001, η_p^2=0.008; Figure 1d]. As

expected, participants made more errors and were slower in unexpected trials compared to both expected

[accuracy: $t_{(49)}$=3.101, p=0.003, Hedge's g=0.414; RT: $t_{(49)}$=-3.955, p<0.001, g=-0.236] and neutral trials

[accuracy: $t_{(49)}$=2.921, p=0.005, g=0.420; RT: ($t_{(49)}$=-2.137, p=0.038, g=-0.128]. Participants also made

more errors [($F_{(1,49)}$=5.027, p=0.030, η_p^2=0.016, *g*=-0.324)] and were slower [$F_{(1,49)}$=12.287, p<0.001,

η_p^2=0.015)] in high pain trials compared to low pain trials. There was no interaction between Trial type and

Pain intensity on accuracy [$F_{(1.627,79.697)}$=0.248, p=0.735, η_p^2<0.001] or RTs [$F_{(1.677,82.169)}$=2.474,

p=0.100, η_p^2=0.002].

We further confirmed that pain intensity ratings differed between the two administered intensities

($F_{(1,49)}$=315.539, p<0.001, η_p^2=0.707; Figure 1e). Pain intensity and Trial type showed an interaction

($F_{(1.575,77.176)}$=8.259, p=0.001, η_p^2=0.001): participants rated high pain stimulations lower during

unexpected trials compared to both expected ($t_{(49)}$=2.084, p=0.042, g=0.056) and neutral trials ($t_{(49)}$=2.400,

p=0.020, g=0.073), while ratings for low pain trials increased in unexpected conditions compared to both

expected (t(49)=-2.151, p=0.036, g=-0.071) and neutral trials (t(49)=-2.616, p=0.012, g=-0.111; Figure 1f).

346
Computational model fitting and validation

Bayesian model comparison identified the two-level HGF without perceptual uncertainty (HGF2) as the best

predictor of participants' log response times (Figure 2c; protected exceedance probability [PEP] = 0.99; model

frequency = 0.540).

To validate this model, we conducted two analyses. First, we performed parameter recovery by simulating

ten response time datasets for each participant using their fitted HGF2 parameters (500 datasets total) and

generating response times by adding random gaussian noise to the linear response model. We then fitted

the model to these simulated data and compared the recovered parameters, averaged across the ten

datasets per participant, to the original simulation parameters. This analysis revealed high Spearman ρ

correlations between simulated and recovered parameters (Figure 2d; parameter relationships illustrated in

Supplementary Figure S1c-g).

Second, we assessed model recovery by fitting all models to data simulated from each model. The HGF2

was correctly identified as the winning model for 95% of datasets simulated using its parameters with no

evidence for better fit from alternative models. However, the HGF2 was not specific in recovering itself and

showed high recovery for datasets generated using other models (Supplementary Figure S1b), further

suggesting that the HGF2 was a superior fit to our task.

Model-based analyses

Analysis of the relationship between the HGF2 model's trial-wise estimates and pain ratings revealed a

significant interaction between estimation uncertainty and pain intensity (β = 8.194, SE = 2.191, p < 0.001;

Figure 3a). Decomposition of this interaction revealed that estimation uncertainty was positively associated

with pain ratings in high-intensity trials (β = 5.669, SE = 1.891, p = 0.003), while showing a negative but not

significant relationship with pain ratings in low-intensity trials (β = -2.702, SE = 1.385, p = 0.05). Similarly,

there was a significant interaction between absolute prediction errors and pain ratings (β = -2.994, SE =

1.437, p = 0.037; Figure 3b). The decomposition of this interaction revealed that absolute prediction errors

were negatively associated with pain ratings in high pain trials (B = -2.264, SE = 1.122, p = 0.044) but showed

no significant relationship in low pain trials (B = 0.731, SE = 0.664, p = 0.271). Outcome uncertainty did not

significantly affect pain ratings (β = -3.540, SE = 10.952, p = 0.747), and its interaction with pain intensity

was also non-significant (β = -1.221, SE = 24.563, p = 0.960).

Figure 2. (a) Structure of the HGF2 model. (b) Example of absolute prediction error, irreducible and estimation

uncertainty across trials. (c) Results of the Bayesian model comparison. (d) HGF2 model parameter recovery.

Individual differences in learning

High depressive symptoms were associated with lower tonic volatility for tone-pain associations (ω_2; r(47) =

-0.41, p = 0.015; Figure 3c). Pain catastrophizing scores (r(47) = -0.29, p = 0.123; Figure 3d), state anxiety

(r(47) = -0.20, p = 0.170; Figure 3e) and trait anxiety (r(47) = -0.29, p = 0.123; Figure 3f) showed non-

significant negative correlations with tonic volatility.

Figure 3. Model-based analyses results of linear mixed-effects models. (a) Pain ratings by estimation uncertainty where lighter lines represent each participant's individual slope and darker bold lines represent the average slope (red for high pain intensity and blue for low pain intensity). (b) Pain ratings by absolute prediction errors. (c-f) Individual differences in BDI, PCS, STAI state and STAI trait scores and the tonic volatility parameter (ω_2). The reported p-values were corrected with the Holm-Bonferroni method.

Discussion

Pain perception is fundamentally shaped by expectations and learning. Using a dynamic learning task, we asked participants to learn associations between auditory cues and painful electrical stimulations of two different intensities. The task was designed to generate periods of high and low uncertainty by frequently changing the cue-pain contingencies throughout the experiment. Results of computational modeling revealed

that under high uncertainty, high-intensity pain was perceived as more painful, while low-intensity pain was

perceived as less painful. These findings extend previous research on expectation and uncertainty effects in

pain perception[4,12,18,38] by showing how learned uncertainty in a changing environment modulates pain

processing in an intensity-dependent manner—amplifying the perception of high pain while attenuating the

perception of low pain.

Model-free analyses showed expected results, as more errors and longer response times were found in

unexpected trials. Our results also suggest an opposite effect of expectations on pain perception for high and

low intensity stimuli, where high pain perception decreased and low pain perception increased in unexpected

trials. These findings are in line with Bayesian inference theories where perception is biased toward

expectations[39], as found in similar studies[18]. Although these analyses provide insight on our findings, they

fail to explain the underlying mechanisms responsible for the effect of unexpectedness. Computational

modeling allows for a more in-depth explanation of the dynamic relationships between expectations and

uncertainty and how they influence perception to optimize learning.

Our computational modeling approach revealed that learning about pain in volatile environments is best

captured by hierarchical learning processes. The two-level Hierarchical Gaussian Filter (HGF2[19,20])

outperformed other models in predicting participants' response times, suggesting that participants

continuously updated beliefs at multiple levels. Similarly to a recent study[23], the three-level HGF3, which

additionally models estimates of environmental uncertainty, did not outperform the simpler two-level HGF

and showed poor parameter recovery performance in comparison to the HGF2. This discrepancy, when

compared to several other studies using a similar task[24,25], could be attributed to the specific contingency

structure of our task, which included fewer trials and contingency switches as well as a repetitive contingency

switching pattern compared to a random one, therefore not requiring estimations of volatility to perform

adequately.

Using the HGF2 to estimate participants' subjective perception of environmental uncertainty, we found that

its relationship with pain perception depended on stimulus intensity: under higher estimation uncertainty,

high-intensity pain was perceived as more intense, while there was a trending negative relationship for low-

intensity pain, suggesting it was perceived as less intense, though this effect did not reach statistical

significance. This pattern aligns with Bayesian frameworks of perception and suggests that under uncertainty,

the pain system shifts toward relying more on incoming sensory information[39]. It could also be the case that

the differential impact of uncertainty on pain perception depending on stimulus intensity reflects asymmetric

cost functions for pain perception and serves to optimize pain's learning and control function[6]. When facing

high pain stimuli under uncertainty, the biological cost of underestimating the threat outweighs the cost of

overestimating, leading to elevated pain ratings as a protective learning mechanism for avoiding significant

threats. Conversely, for low pain stimuli, the system favors underestimation when uncertain, as this minimizes

costs associated with unnecessary defensive responses, resulting in lower pain ratings. This could optimize

learning by filtering out noisier signals of unthreatening pain. Our results could also explain the previously

observed mixed effects of uncertainty on pain perception[13,14,16–18], as studies using different pain intensities

might find apparently contradictory effects of uncertainty depending on where their stimuli fell relative to

participants' average expectations. Here, our dynamic modulation of uncertainty through changing

contingencies provides a more ecologically valid approach compared to previous studies that used static

manipulations of uncertainty or fixed probabilities.

Additionally, we found absolute prediction errors, as estimated by the HGF2, influenced pain perception

differently according to stimulus intensity: high pain was rated lower as prediction errors increased, while

there was evidence for low pain to be rated higher, although this effect did not reach statistical significance.

These results align with learning theories of pain suggesting that perception is dynamically adjusted in

response to prediction errors to optimize learning[6]. The greater malleability of high pain compared to low pain

could reflect an asymmetric cost of enhancing the salience of prediction errors. Less threatening signals of

pain could carry less informational value for harm avoidance, thus requiring less adjustment in response to

prediction errors. In contrast, more salient signals of high pain could be more flexible in response to prediction

errors and serve a greater learning purpose for threat avoidance.

Individual differences in self-reported measures of depression, anxiety and pain catastrophizing were

examined in relation to learning parameters estimated by the HGF2 model. We found a non-significant

negative trend between anxiety and pain catastrophizing and ω_2—a parameter representing the step size of

the learning rate for updating cue-outcome associations over time[19]—while participants reporting more

symptoms of depression showed significantly reduced ω_2 values. A lower ω_2 indicates reduced flexibility in

updating beliefs about how pain-cue relationships change. This finding has particular relevance for chronic

pain populations, where elevated catastrophizing, depression and anxiety are common[40] and represent

important risk factors for chronic pain development[41]. Our results suggest that these traits may be linked with

impaired ability to update pain expectations in changing environments, potentially contributing to maladaptive

pain processing. It should be noted that the three measures were moderately to highly correlated (r = 0.5 to

0.72), suggesting that they tap into a common trait related to learning. Nevertheless, the relationships

between these traits and learning may help explain why some individuals develop chronic pain conditions

while others recover from similar injuries and thus provide a computational hypothesis for the long-theorized

link between chronic pain and learning mechanisms[42,43] that should be tested in future studies.

Several limitations warrant consideration. First, while we used a similar design to previous studies[44,45], our

task employed a single contingency structure. Our findings should be tested for generalization across

different types of contingency patterns and learning environments. Second, while our findings suggest a link

between psychological characteristics and learning mechanisms, these relationships were relatively weak

and observed in a healthy sample with few individuals reporting high anxiety, depression or catastrophizing.

Testing these relationships in clinical populations will be crucial to establish their relevance to chronic pain.

Finally, while we apply our results to pain perception, similar influences of uncertainty on perception might

be observed in other phenomena, and psychological traits might be linked to general learning mechanisms

rather than being specific to pain learning.

In conclusion, our study demonstrates that pain perception is dynamically modulated by learned uncertainty,

with distinct effects depending on pain intensity, and is influenced by individual differences in psychological

characteristics. The successful application of hierarchical Bayesian models to capture these effects provides

a promising framework for understanding pain learning in complex, changing environments. Our finding that

symptoms of depression are associated with reduced learning efficiency could be particularly relevant for

understanding how individuals with chronic pain learn and deal with uncertainty in their daily lives.

Disclosures

MLT was supported by scholarships from the *Fonds de Recherche Québec – Santé* (2023-2024, 2024-2028)

and the Canadian Institutes of Health Research (2023, 2025-2028). MPC and this work was supported by a

career grant from the *Fonds de Recherche Québec - Santé* (grant #309041, 2021-2025). The authors declare

that they have no conflict of interest. All raw data and code necessary to reproduce all analyses and figures

are available at https://github.com/meganel-t/painuncertainty.

References

1.   Melzack R, Casey KL. Sensory, motivational, and central control determinants of pain: A new

conceptual model. The *skin Senses*. 1968;1:423–443.

2.   Fields HL. How expectations influence pain. *Pain*. 2018;159(1):S3–S10.

doi:10.1097/j.pain.0000000000001272

3.   Coll MP, Walden Z, Bourgoin PA et al. Pain reflects the informational value of nociceptive inputs. *Pain*.

2024;165(10):e115–e125. doi:10.1097/j.pain.0000000000003254

4.   Zaman J, Van Oudenhove L, Vlaeyen JWS. Uncertainty in a context of pain: disliked but also more

painful? *Pain*. 2021;162(4):995–998. doi:10.1097/j.pain.0000000000002106

5.   Roy M, Shohamy D, Daw N, Jepma M, Wimmer GE, Wager TD. Representation of aversive prediction

errors in the human periaqueductal gray. *Nat Neurosci*. 2014;17(11):1607–1612. doi:10.1038/nn.3832

6.   Seymour B. Pain: A Precision Signal for Reinforcement Learning and Control. *Neuron*.

2019;101(6):1029–1041. doi:10.1016/j.neuron.2019.01.055

7.   Seymour B, Dolan RJ. Emotion, Motivation, and Pain. In: McMahon SB, Wall PD, Melzack R, McMahon

SB, eds. *Wall and Melzack's textbook of pain*. 6. ed. Elsevier, Saunders; 2013.

8.   Seymour B, Mancini F. Hierarchical models of pain: Inference, information-seeking, and adaptive

control. *NeuroImage*. 2020;222:117212. doi:10.1016/j.neuroimage.2020.117212

9.   Büchel C, Geuter S, Sprenger C, Eippert F. Placebo Analgesia: A Predictive Coding Perspective.

Neuron. 2014;81(6):1223–1239. doi:10.1016/j.neuron.2014.02.042

10.   Tabor A, Burr C. Bayesian Learning Models of Pain: A Call to Action. *Curr Opin Behav Sci*. 2019;26:54–

61. doi:10.1016/j.cobeha.2018.10.006

11.   Seymour B, O'Doherty JP, Dayan P, et al. Temporal difference models describe higher-order learning

in humans. *Nature*. 2004;429(6992):664–667. doi:10.1038/nature02581

12.   Taylor VA, Chang L, Rainville P, Roy M. Learned expectations and uncertainty facilitate pain during

classical conditioning. *Pain*. 2017;158(8):1528–1537. doi:10.1097/j.pain.0000000000000948

13.   Ploghaus A, Narain C, Beckmann CF, et al. Exacerbation of Pain by Anxiety Is Associated with Activity

in a Hippocampal Network. *J Neurosci*. 2001;21(24):9896–9903. doi:10.1523/JNEUROSCI.21-24-

09896.2001

14.   Rhudy JL, Meagher MW. Fear and anxiety: divergent effects on human pain thresholds. *Pain*.

2000;84(1):65–75. doi:10.1016/S0304-3959(99)00183-9

15.   Pavy F, Zaman J, Von Leupoldt A, Torta DM. Expectations underlie the effects of unpredictable pain:

a behavioral and electroencephalogram study. *Pain*. 2024;165(3):596–607.

doi:10.1097/j.pain.0000000000003046

16.   Yoshida W, Seymour B, Koltzenburg M, Dolan RJ. Uncertainty Increases Pain: Evidence for a Novel

Mechanism of Pain Modulation Involving the Periaqueductal Gray. *J Neurosci*. 2013;33(1):5638–5646.

doi:10.1523/JNEUROSCI.4984-12.2013

17.   Zaman J, Vanpaemel W, Aelbrecht C, Tuerlinckx F, Vlaeyen JWS. Biased pain reports through

vicarious information: A computational approach to investigate the role of uncertainty. *Cognition*.

2017;169:54–60. doi:10.1016/j.cognition.2017.07.009

18.   Brown CA, Seymour B, Boyle Y, El-Deredy W, Jones AKP. Modulation of pain ratings by expectation

and uncertainty: Behavioral characteristics and anticipatory neural correlates. *Pain*. 2008;135(3):240–

250. doi:10.1016/j.pain.2007.05.022

19.   Mathys C, Daunizeau J, Friston KJ, Stephan KE. A bayesian foundation for individual learning under

uncertainty. *Front Hum Neurosci*. 2011;5:39. doi:10.3389/fnhum.2011.00039

20.   Mathys CD, Lomakina EI, Daunizeau J, et al. Uncertainty in perception and the Hierarchical Gaussian

Filter. *Front Hum Neurosci*. 2014;8. doi:10.3389/fnhum.2014.00825

21. Anchisi D, Zanon M. A Bayesian Perspective on Sensory and Cognitive Integration in Pain Perception

and Placebo Analgesia. *PLoS ONE*. 2015;10(2):e0117270. doi:10.1371/journal.pone.0117270

22. de Berker AO, Rutledge RB, Mathys C, et al. Computations of uncertainty mediate acute stress

responses in humans. *Nat Commun*. 2016;7(1). doi:10.1038/ncomms10996

23. Ehmsen JF, Nikolova N, Christensen DE, et al. Thermosensory predictive coding underpins an illusion

of pain. *Sci Adv*. 2025;11(11):eadq0261. doi:10.1126/sciadv.adq0261

24. Iglesias S, Mathys C, Brodersen KH, et al. Hierarchical Prediction Errors in Midbrain and Basal

Forebrain during Sensory Learning. *Neuron*. 2013;80(2):519–530. doi:10.1016/j.neuron.2013.09.009

25. Lawson RP, Mathys C, Rees G. Adults with autism overestimate the volatility of the sensory

environment. *Nat Neurosci.* 2017;20(9): 1293–1299. doi:10.1038/nn.4615

26. Vlaev I, Seymour B, Dolan RJ, Chater N. The Price of Pain and the Value of Suffering. *Psychol Sci*.

2009;20(3):309–317. doi:10.1111/j.1467-9280.2009.02304.x

27. den Ouden HEM, Daunizeau J, Roiser J, Friston KJ, Stephan KE. Striatal Prediction Error Modulates

Cortical Coupling. *J Neurosci.* 2010;30(9): 3210–3219. doi:10.1523/JNEUROSCI.4458-09.2010

28. Sullivan MJL, Bishop SR, Pivik J. The Pain Catastrophizing Scale: Development and validation. *Psychol

Assess*. 1995;7(4):524–532. doi:10.1037/1040-3590.7.4.524

29. French DJ, Noël M, Vigneau F, French JA, Cyr CP, Evans RT. L'Échelle de dramatisation face à la

douleur PCS-CF: Adaptation canadienne en langue française de l'échelle «Pain Catastrophizing

Scale». *Can J Behav Sci Rev Can Sci Comport*. 2005;37(3):181–192. doi:10.1037/h0087255

30. Spielberger CD. State-Trait Anxiety Inventory for Adults. Published online 1983. doi:10.1037/t06496-

000

31. Gauthier J, Bouchard S. Adaptation canadienne-française de la forme révisée du State–Trait Anxiety

Inventory de Spielberger. *Rev Can Sci Comport*. 1993;25(4):559–578. doi:10.1037/h0078881

32.   Beck AT. An Inventory for Measuring Depression. *Arch Gen Psychiatry*. 1961;4(6):561.

doi:10.1001/archpsyc.1961.01710120031004

33.   Lawson RP, Bisby J, Nord CL, Burgess N, Rees G. The Computational, Pharmacological, and

Physiological Determinants of Sensory Learning under Uncertainty. *Current Biology*. 2021;31(1):163-

172.e4.

34.   Frässle S, Aponte EA, Bollmann S, et al. TAPAS: An Open-Source Software Package for Translational

Neuromodeling and Computational Psychiatry. *Front Psychiatry*. 2021;12:680811.

35.   Rigoux L, Stephan KE, Friston KJ, Daunizeau J. Bayesian model selection for group studies —

Revisited. *NeuroImage*. 2014;84:971-85.

36.   Rescorla RA, Wagner AR. A theory of Pavlovian conditioning: Variations in the effectiveness of

reinforcement and nonreinforcement. In: Black A, Prokasy W, eds. *Classical Conditioning II: Current*

Research and Theory. Appleton-Century-Crofts; 1972:64–99.

37.   Taylor VA, Roy M, Chang L, Gill L-N, Mueller C, Rainville P. Reduced Fear-Conditioned Pain

Modulation in Experienced Meditators: A Preliminary Study. *Psychosom* Med. 2018;80(9):799–806.

doi:10.1097/PSY.0000000000000634

38.   Koyama T, McHaffie JG, Laurienti PJ, Coghill RC. The subjective experience of pain: Where

expectations become reality. *Proc Natl Acad Sci.* 2005;102(36):12950–12955.

doi:10.1073/pnas.0408576102

39.   Hoskin R, Berzuini C, Acosta-Kane D, El-Deredy W, Guo H, Talmi D. Sensitivity to pain expectations:

A Bayesian model of individual differences. *Cognition*. 2019;182:127–139.

doi:10.1016/j.cognition.2018.08.022

40.   Brouwer B, Waardenburg S, Jacobs C, et al. Biopsychosocial baseline values of 15 000 patients

suffering from chronic pain: Dutch DataPain study. *Reg Anesth Pain Med*. 2020;45(10):774–782.

doi:10.1136/rapm-2020-101476

41.   Mills SEE, Nicolson KP, Smith BH. Chronic pain: a review of its epidemiology and associated factors

580   in population-based studies. *Br J Anaesth*. 2019;123(2):e273–e283. doi:10.1016/j.bja.2019.03.023

581   42.   Apkarian AV, Baliki MN, Geha PY. Towards a theory of chronic pain. *Prog Neurobiol*. 2009;87(2):81–

582   97. doi:10.1016/j.pneurobio.2008.09.018

583   43.   Zaman J, Vlaeyen JWS, Van Oudenhove L, Wiech K, Van Diest I. Associative fear learning and

584   perceptual discrimination: A perceptual pathway in the development of chronic pain. *Neurosci Biobehav*

585   *Rev*. 2015;51:118–125. doi:10.1016/j.neubiorev.2015.01.009

586   44.   Atlas LY, Dildine TC, Palacios-Barrios EE, et al. Instructions and experiential learning have similar

587   impacts on pain and pain-related brain responses but produce dissociations in value-based reversal

588   learning. *eLife*. 2022;11:e73353. doi:10.7554/eLife.73353

589   45.   Horing B, Büchel C. The human insula processes both modality-independent and pain-selective

590   learning signals. *PLoS Biol*. 2022;20(5):e3001540. doi:10.1371/journal.pbio.3001540