PmaI-dependent pH gradients initiate inter- and intraspecies germling fusion in filamentous
fungi

Rudolph A. Y.[1], Rothermund, N.[2], Vasilevska, N.[1], Fleißner A.[2], Nordzieke D.E.[1,*]

[1]University of Göttingen, GZMB, Institute of Microbiology and Genetics, Genetics of Eukaryotic
Microorganisms, Grisebachstr. 8, 37077 Göttingen, Germany

[2]University of Braunschweig, Institute of Technology, Institute of Genetics, Spielmannstraße
7, 38106 Braunschweig, Germany

*Correspondence: dnordzi@gwdg.de

Keywords: *Colletotrichum graminicola*, *Neurospora crassa*, *Botrytis cinerea*, Pmal ATPase,
fungal GPCR, MAPK pathway, CAT fusion

<u>Abstract</u>

Somatic cell fusion among spore germlings is a complex process in filamentous fungi that
enables the formation of a syncytial unit and facilitates the exchange of genetic material apart
from sexual reproduction. In this study, we investigated the molecular mechanisms underlying
germling fusion in the maize pathogen Colletotrichum graminicola. Based on secretomes
harvested during the germling fusion process of fusion-competent oval and fusion-deficient
falcate conidia, we identified pH as a communication signal. We showed that the plasma
membrane localized ATPase PmaI is a probable source for pH gradient formation. Our results
also suggest that the a-pheromone receptor CgSte3 mediates sensing of acidic pH gradients,
activating downstream the Cell Wall Integrity Mitogen Activated Protein Kinase pathway. We
propose a new model for germling interactions, in which pH serves as an interaction initiation
signal, resulting in the activation of the cell-dialogue. Our findings have implications for our
understanding of fungal communication and pathogenicity, and highlight the importance of pH
as a signal molecule in germling fusion.

<u>Introduction</u>

Somatic cell fusion is a complex process that enables filamentous fungi to merge genetically
identical or non-identical cells, resulting in the formation of a syncytial unit. This process is
essential for the efficient development and growth of fungal colonies, and is mediated by a
variety of signaling pathways and molecular mechanisms (Bastiaans *et al.*, 2015, Richard *et*
al., 2012, Herzog *et al.*, 2015). In *Neurospora crassa*, a model organism for studying fungal
biology, somatic cell fusion is initiated by the germination of conidia, which undergo mutual
attraction and fusion to form a syncytial unit (Read *et al.*, 2012). This process is regulated by
a complex interplay of signaling molecules and molecular components, including transcription
factors, the striatin-interacting phosphatase and kinase (STRIPAK) complex, NADPH oxidase
complexes, and mitogen-activated protein kinase (MAPK) pathways (Fischer and Glass,
2019).

The current understanding of germling fusion in fungi suggests that it is a three-step process:
(I) the formation of conidial anastomosis tubes (CATs), (II) bidirectional chemotropic growth of

CATs towards each other (CAT homing), and (III) cell fusion at the CAT contact side (Glass *et*
al., 2004, Kurian *et al.*, 2018). Central to the reciprocal recognition of interaction partners are
the two proteins SO (soft) and MAK-2, which are part of the cell wall integrity (CWI) and
pheromone Mitogen Activated Protein Kinase (MAPK) pathways, respectively (Fleissner *et al.*,
2009, Serrano *et al.*, 2018). During the tropic growth of the interacting cells, both proteins are
recruited to the plasma membrane of the growing cell tips in an alternating manner. The current
hypothesis proposes that SO localization to the CAT tip is associated with the release of a
previously unknown communication signal, which is recognized by MAK-2 in the interaction
partner, forming the so-called 'cell-dialogue' (Fleissner et al., 2009). Recent studies have
challenged the long-held assumption that germling fusion occurs only among germlings of the
same species, revealing that interspecies interactions and true fusion can occur between
different fungal species (Haj Hammadeh *et al.*, 2022, Mehta and Baghela, 2021). For example,
germlings of *N. crassa* and *Botrytis cinerea* interact with each other albeit the fusion process
is not completed (Haj Hammadeh et al., 2022). Additionally, *Colletotrichum gloeosporioides*
and *Colletotrichum siamense* fuse and form viable progeny, with the genome of the resulting
progeny undergoing genetic recombination (Mehta and Baghela, 2021).

Colletotrichum graminicola is a hemibiotrophic maize pathogen responsible for anthracnose
disease, causing high economic damage with corn loss of 10-20% of harvest annually
(Belisário *et al.*, 2022). For successful disease development, *C. graminicola* generates two
different asexual spore types, falcate conidia in acervuli on infected maize leaves, and oval
conidia inside parenchyma cells nearby the plant vascular system (Belisário et al., 2022,
Panaccione *et al.*, 1989). Those spore types differ significantly in their vegetative development
and infection-relevant processes, making falcate conidia more efficient in leaf infection and
enable oval conidia to sense maize root-exudated signals, followed by chemotropic growth
towards the host and root infection (Nordzieke *et al.*, 2019, Schunke *et al.*, 2020, Rudolph *et*
al., 2024a, Chaky *et al.*, 2001). One prominent difference between both spore types is the
ability to form a network by vegetative fusions: oval conidia form fusion bridges at high rates
under starvation conditions, whereas falcate conidia stay dormant in the same experimental
setups. However, close proximity of oval and falcate conidia can release dormancy in falcate
conidia and result in fusion of both spore types (Nordzieke et al., 2019). From these results we
conclude that the germling interaction signal is secreted by oval conidia but is absent from
falcate conidia. Intriguingly, oval conidia fusion takes place also on leaves. There, fusion bridge
formation is correlated to the generation of hyphopodia for leaf penetration, although both
processes do not directly depend on each other (Nordzieke et al., 2019, Nordzieke, 2022).

In this study, we aimed to identify signals responsible for reciprocal germling attraction during
the fusion process in *C. graminicola*. To identify these signal(s), a methodical setup was
established and verified, enabling us to collect secreted molecules during the fusion process.
Comparing the germling fusion secretome of oval and falcate conidia, we observed difference
in secretome pH. In the following, we verified that distinct pHs are able to attract *C. graminicola*
germlings and medium pH influences CAT fusion frequency. We further explored that the a-
pheromone receptor CgSte3 and the CWI pathway mediate pH sensing. Applying an inhibitor
of the plasma membrane standing ATPase PmaI, we provided evidence that pH signal
formation is dependent on ATPase activity and links germination and fusion in *C. graminicola*.
From those findings, we propose a new model for germling interactions, in which pH serves as
an interaction initiation signal, resulting in the activation of the cell-dialogue. Since PmaI-
dependent pH gradients are formed in all fungi in the course of germination, we further
explored intraspecies interaction of *C. graminicola* with *N. crassa* and *B. cinerea*. All together,
we identified that pH gradients serve as signal during reciprocal germling attraction prior to
fusion in *C. graminicola*. Since the fusion rate of other filamentous fungi does depend on pH
as well, we propose a general function as interaction initiation signal.

Material and methods

Strains, growth conditions, and collection of spores

The *C. graminicola* (Ces.) G.W. Wilson (teleomorph *Glomerella graminicola* D. J. Politis) wild-type strain M2 (also referred to as M1.001) was provided by H. Deising (Martin-Luther University Halle-Wittenberg) and used in this study (Forgey *et al.*, 1978, O'Connell *et al.*, 2012). Falcate and oval conidia of *C. graminicola* were generated and collected as described previously (Rudolph *et al.*, 2024b). *N. crassa* matA (strain FGSC2489 provided by Fungal Genetics Stock Center) and *B. cinerea* (strain B05.10; Quidde *et al.*, 1999) were grown on Vogel´s minimal medium (Vogel´s MM), supplemented with 2% sucrose (Vogel, 1956). All strains used in this study are listed in Supplementary Table 1.

Generation and analysis of germling secretome during conidial interaction

In our standard protocol for the quantification of germling fusions, we spread 50 µl of spore solution (c = 5 x 10^7 ml^{-1}) on water agar (1% serva agar, 1% agarose, 25 mM NaNO$_3$) and incubate the plates for 17 h at 23°C. For oval conidia preparations, we see a high percentage of germling interacting and fusing after that time. However, falcate conidia do not germinate nor fuse under these conditions (Nordzieke et al., 2019). To enable the identification of signals generated during the germling interaction and fusion process, we aimed to collect the secretome of oval and falcate in this experimental setup and to compare their chemical properties. Therefore, we spread oval and falcate conidia of *C. graminicola* CgM2 on cellophane (SERVA cellophane sheets II (140 x 133), SERVA Electrophoresis GmBH, Heidelberg, Germany; 50 µl of spore solution with c = 5 x 10^7 ml^{-1}) topping water agar and let the liquid evaporate (clean bench). Afterwards, the cellophane sheets were transferred to petri dishes filled with a single layer of glass beads (Glass Beads Ø 2.85 – 3.45 mm, Roth, Karlsruhe, Germany) and 10 ml of a 25 mM NaNO$_3$ solution. After 17 h incubation at 23°C, the liquid was collected in glass ware and frozen at -80°C. To control the formation of germling fusion networks, microscopic pictures were taken, verifying the formation of fusion bridges for oval conidia and their absence for falcate conidia (Figure 1). pH of secretome solutions was measured using pH strips (20RO5121, Roth, Karlsruhe, Germany) of freshly harvested or 10 x concentrated secretome.

Germling fusion and germination assays in *C. graminicola*

To determine the impact of concentrated germling fusion secretome on the fusion and germination ability of oval and falcate conidia, 10 µl of different concentrated spores solutions (c = 10^6 ml^{-1}; 5 x 10^5 ml^{-1}; 10^5 ml^{-1}) were dropped on water agar (1% serva agar, 1% agarose, 25 mM NaNO$_3$) and dried under a laminar flow. The dried spore drops were topped with 20 µl of 10 x concentrated secretome generated from oval or falcate conidia germlings during fusion conditions. After drying of the secretome drops, the plates were inoculated for 17 h at 23°C. At least 100 touching germlings and conidia were accessed for fusion bridge formation and 100 conidia for germination in at least three biological replicates.

For the determination of pH-dependent germling fusion and germination frequencies, 50 µl of oval conidia (c = 5 x 10^7 ml^{-1}) were spread on water agar (1% serva agar, 1% agarose, 25 mM NaNO$_3$) adjusted to different pH values after autoclaving using pH strips (pH-Fix 4.5-10.0, Roth, Karlsruhe, Germany). For buffered conditions, the water agar was prepared with citrate phosphate buffer to reach a stable pH of 6.5. Microscopic evaluation was performed after 17

h of cultivation at 23°C, counting at least 100 touching conidia or germlings and accessing their
fusion state. Germination rates were determined by evaluating at least 100 conidia. Both
experiments were performed in at least three biological replicates.

Concentration-dependent germling fusion assays were performed by dropping 10 µl of oval
conidia spore solution (10^6 m^{-1}; 7.5 x 10^5 m^{-1}; 5 x 10^5 m^{-1}; 2.5 x 10^5 m^{-1}; 10^5 m^{-1}; 7.5 x 10^4 m^{-1};
5 x 10^4 m^{-1}) of *C. graminicola* wildtype (CgM2), *Cgste3* deletion and complementation strains
on water agar (1% serva agar, 1% agarose, 25 mM NaNO$_3$), dried under laminar flow, and
incubated for 17 h at 23°C. For higher concentrations, at least 100 touching conidia or
germlings were microscopically analyzed for the presence of fusions, for lower concentrations
the maximal amount of conidia and germlings present in the 10 µl drop were evaluated. For
each concentration, the experiments were performed at least three times.

To determine the effect of Pmal inhibitor Demethoxycurcumin (DMCM) (Demethoxycurcumin
≥ 98% HPLC, Sigma-Aldrich, St. Louis, USA) on germling fusion and germinate rate of oval
conidia, DMCM was dissolved in DMSO. Working solutions were prepared just before use by
further diluting with distilled water. As control, diluted DMSO concentration was used
equalizing 15 µM of DMCM. 20 µl of different DMCM concentrations or the control were applied
on dried 10 µl drops of oval conidia (c = 10^6 ml^{-1}) on water agar (1% serva agar, 1% agarose,
25 mM NaNO$_3$) and dried as well under a laminar flow. Microscopic evaluation was performed
after 17 h of cultivation at 23°C, counting at least 100 touching conidia or germlings and
accessing their fusion state. Germination rates were determined by evaluating at least 100
conidia. Both experiments were performed in at least three biological replicates.

Chemotropic assay

Chemotropic growth is defined as directed growth along a gradient of a given chemical
stimulus, which can be positive (attractant signal) or negative (repellant signal). To display
positive and negative chemotropic growth responses, the 'chemotropic index' is used as a
standard for result depiction, indicating the difference of a preferred growth direction compared
to a situation, in which neither attraction no repulsion occurs (Turrà *et al.*, 2015). In our lab, we
developed a 3D printed device to easily monitor directed growth responses (Schunke et al.,
2020). Using this device, we determined growth direction of fungal germlings derived from *C.*
graminicola oval conidia of CgM2 (wildtype), Δ*Cgste3*, Δ*Cgso*, and the corresponding
complementation strains after incubation for 6 h at 23°C. As attractant molecules, we evaluated
10 x concentrated secretome of oval and falcate conidia generated during the germling fusion
process and potassium phosphate buffer adjusted to different pH values (pH 6, 7, ,7.5, 8). All
experiments were performed in at least two biological replicates.

Microscopy

Light (differential interference contrast (DIC)) was performed with the Axiolmager M1
microscope (Zeiss, Jena, Germany). Images were captured with the Photometrix coolSNAP
HQ camera (Roper Scienific, Photometrics, Tucson, AZ, USA) and processed with the ZEISS
ZEN software (version 2.3, Zeiss). Alternatively, images were taken with the Axio Observer
microscope (Zeiss, Jena, Germany). Images were captured with sCMOS pco.panda 4.2 M
(Excelitas PCO GmbH) and processed with the software 4-D MICROSCOPE FLMV 1.1 and 4-
D MICROSCOPE Observer Z.1 (Schnabel et al., 1997).

Inter- and intraspecies interaction studies

The interspecies interaction experiments were performed on Vogel´s MM (20 ml of 50x Vogel's
trace elements solution, 20 g Sucrose, 15 g Agar-Agar; Vogel, 1956). The interactions
experiment with *C. graminicola* and *B. cinerea* was carried out at 23°C with an incubation time
of 17 h. First, fungal spore solutions were adjusted to distinct concentrations (*B. cinerea*: c =
3.33×10^7 ml^{-1}; *C. graminicola*: c = 5×10^7 ml^{-1}). 300 µl of *B. cinerea* spore solutions were
plated on Vogel's MM medium and 50 µl of *C. graminicola* oval conidia added. Using a pipette
tip, the spores from both species were mixed and spread onto the medium plate. For the
interspecies interaction experiment between *C. graminicola* and *N. crassa*, 50 µl of a
C. graminicola oval conidia spore solution (c = 5×10^7 x ml^{-1}) was first distributed on Vogel´s
MM medium using glass beads and incubated for 14 h at room temperature. After that time,
300µl of *N. crassa* spore solution (c = 10^7 ml^{-1}) was added on those pre-incubated plates, dried
and further incubated at 28°C for 3.5 - 4 h. To observe the cell dialog during interspecies
interactions with *C. graminicola*, the MAK-2-GFP of *N. crassa* (Fleissner et al., 2009) was used.
The interspecies interaction rate was calculated based on several screen fields in which spores
of both species were present. In total, ≥ 76 interactions were calculated in three biological
replicates.

pH-dependent fusion and germination rates of *C. graminicola*, *B. cinerea*, and *N. crassa* was
performed on Vogels' MM adjusted to distinct pH without buffering the medium. For *C.*
graminicola, 50 µl of oval conidia spore solution (c = 5×10^7 x ml^{-1}) was spread on Vogels' MM
using glass beads. For *B. cinerea* and *N. crassa*, 300 µl of c = 10^7 ml^{-1} spore solutions of were
added on Vogels' MM and spread using pipette tips. After incubation (*C. graminicola*: 17 h,
23°C, *B. cinerea*: 17 h, 23°C, *N. crassa*: 3 h, 30°C), at least 100 contacting germlings or conidia
were accessed for formed fusion bridges. In the same experiments, germination rates were
determined from at least 100 conidia. Both fusion and germination experiments were
conducted in at least 3 biological replicates.

Localization of MAK-2-GFP was monitored during *N. crassa* – *N. crassa* and *N. crassa* – *C.*
graminicola interactions, using the same conditions described above. As readout served
fluorescence microscopy.

Statistics

For statistical analysis in this study, the T-test for unequal variances was used for all
experiments (Ruxton, 2006).

AI statement

The AI chatbot 'Chat AI' provided by the GWDG at the Georg-August University Göttingen was
used to improve the text of this manuscript regarding English grammar and typing.

<u>Results</u>

Secreted molecules secreted from oval conidia during fusion induce chemotropic growth of *C.*
graminicola germlings

While numerous proteins and second messengers are necessary for CAT fusion (Fu *et al.*,
2011, Fischer and Glass, 2019), signals resulting in bidirectional attraction of germlings are
unknown. To explore the potential of CAT signal identification, we established an experimental

design enabling to harvest secreted signals during fusion bridge formation. We spread equal
amounts of fusion competent oval or fusion-incompetent falcate conidia on cellophane sheets
topping water agar. After drying the spore solution, we transferred the cellophane to a new
petri dish filled with a single layer of glass beads and 25 mM $NaNO_3$ solution. After 17 h of
incubation, the germination and fusion state of the cultures were evaluated, showing that oval
conidia germinated in high percentages and undergo fusions eagerly. In contrast, falcate
conidia did not germinate or form fusion bridges (Figure 1, A). Since those results are in line
with previous observations (Nordzieke et al., 2019), glass beads and liquid were separated by
filtration and the flow through analyzed.

Figure 1 CAT secretome of *C. graminicola* oval conidia elicits chemotropic growth and shapes fusion
and germination of oval and falcate conidia. (A) Schematic representation of fusion secretome
generation. 50 µl of oval and falcate conidia spore solutions (c = 5 x 10^7 ml^{-1}) of the *C. graminicola*
wildtype were spread on cellophane topping water agar. After evaporation of the liquid, cellophane
sheets were transferred to petri dishes filled with a single layer of glass beads and 10 ml of a 25 mM
$NaNO_3$ solution. After 17 h incubation at 23°C, the fusion secretome was collected and the cellophane
evaluated for the presence or absence of fusions. (B) Chemotropic index of *C. graminicola* germlings
derived from oval conidia to 10 x concentrated fusion secretome harvested from oval (Sec_{oc}) or falcate
(Sec_{fc}) conidia incubations. 20 µl of oval conidia (c = 5 x 10^5 ml^{-1}) were applied in a 3D printed device
between two wells containing 40 µl solvent control and attractant. Microscopic evaluation of fungal tip
growth was performed after incubation for 6 h, 23°C, n = 8 biological replicates, statistical significance
was calculated with a two-tailed t-test (*, p < 0.05). (C) Germling fusion rates in dependence of spore
concentration and the supplementation with 10 x concentrated Sec_{oc} and Sec_{fc}. 10 µl of oval or falcate
conidia were dropped in different amounts (10^4, 5 x 10^3, 10^3) on water agar plates agar (1% serva agar,
1% agarose, 25 mM $NaNO_3$). After drying, the drops were topped with 20 µl of 10 x concentrated Sec_{oc}
and Sec_{fc} and the liquid dried again. After 17 h incubation at 23°C, at least 100 contacting germlings
and conidia were microscopically examined for the formation of fusion bridges, n ≥ 3, ns = not significant.

In a first experiment, we tested whether the fusion secretomes of oval and falcate conidia are
able to attract *C. graminicola* germlings using a 3D printed device developed in our group
(Schunke et al., 2020). Indeed, we found that 10x concentrated fusion secretome from oval
conidia (Sec_{oc}) induce a strong chemotropic response, whereas the 10x concentrated fusion

secretome of falcate conidia (Sec_{fc}) showed minor attraction potential (Figure 1, B), indicating
a successful harvest of interaction signal(s). In a second approach, we supplemented oval and
falcate conidia with 10x Sec_{oc} and Sec_{fc} and accessed their germination and fusion rates after
17 h of incubation. Since germling fusion is a spore concentration-dependent process, we
tested the effect of the secretomes on optimal (10^4) and suboptimal (10^3) spore concentrations
(Figure 1, C, Supplementary Figure S1). For oval conidia, we observed a spore-density
dependent effect of the secretomes: whereas in optimal experimental conditions, no effect of
secretome addition was visible, 10x Sec_{oc} did increase the fusion frequency of oval conidia
applied in suboptimal concentrations (Figure 1, C). Regarding falcate conidia, the addition of
concentrated Sec_{oc} or Sec_{fc} did not induce fusions; however, in experimental conditions with
10^3 applied spores, germination of this spore type was induced (Supplementary Figure S1).
Together, these first experiments indicate that isolation of a CAT fusion signal was successful
with the experimental approach chosen and point to a possible interconnection between
germination and fusion.

Distinct pHs attract *C. graminicola* germlings and induce germling fusion

Next, we aimed on secretome characterization. When we monitored the pH of the Sec_{oc} and
Sec_{fc} and compared it with control conditions, we observed a more basic pH for 10x Sec_{oc}
(Figure 2, A). We thus aimed to explore, whether pH can act as attractant for *C. graminicola*
and influence fusion frequency. As depicted in Figure 2, B, pH of 7.5 strongly attracts
germlings, whereas higher or lower pH values do not, indicating the sensing is specific to
distinct pH gradients. Monitoring of CAT fusion events on water agar plates with adjusted pH
values revealed that in comparison to not adapted conditions, pH values of 5 and >7 are
inhibiting fusion bridge formation. In contrast, medium adjusted to pH 6.5 induces fusion
frequency. From these results, we were wondering whether pH could be a mere factor of
wellbeing for *C. graminicola* germlings, therefore enabling increased number of fusion events.
To test this, we generated buffered water agar medium with a pH adjusted to 6.5, which
provides optimal fusion condition in not buffered conditions. Intriguingly, no fusions were
detected in those experiments, underpinning the necessity of active pH change by fungal
germlings or conidia for fusion bridge formation (Figure 2, C). In the same experiments, we
also monitored germination. In contrast to network formation by fusion, the germination
process is only affected under buffered conditions (Supplementary Figure 2).

The a-pheromone receptor CgSte3 and the MAPK scaffold CgSo mediate pH sensing

In recent work of our group and others, fungal pheromone receptors were shown to perceive
defense molecules secreted from plant roots like peroxidases and diterpenoids besides fungal
pheromones (Rudolph et al., 2024a, Rudolph et al., 2024b, Turrà et al., 2015, Sridhar *et al.*,
2020, Sharma *et al.*, 2022, Ramaswe *et al.*, 2024, Vitale *et al.*, 2019). Similarly, the Cell Wall
Integrity Mitogen Activated Protein Kinase pathway (CWI) was shown to mediate the
extracellular signals and transform them into chemotropic growth responses (Turrà et al., 2015,
Sridhar et al., 2020, Sharma et al., 2022). To test whether this receptor-MAPK module is also
involved in pH sensing, deletion mutants of genes encoding for the a-pheromone receptor
Cgste3 and the CWI scaffold *Cgso* were examined for their ability to sense and re-direct growth
to pH 7.5 (Figure 3). In contrast to wild-type strain CgM2, which show a strong growth re-
direction, both mutant strains fail to grow towards a gradient of pH 7.5. For both mutants, this
effect was restored by reintroducing the corresponding gene (Figure 3, A).

Figure 2 Distinct pH values attract *C. graminicola* germlings and impact their ability to undergo fusions.
(A) pH of freshly harvested not-concentrated and concentrated fusion secretome of oval (Sec$_{oc}$) and
falcate (Sec$_{fc}$) was examined using pH strips, n = 4. (B) Chemotropic index of *C. graminicola* wild-type
germlings to potassium phosphate buffer adjusted to different pH values (pH 6; 7;7.5; 8). 20 µl of oval
conidia (c = 5 x 10^5 ml^{-1}) were applied in a 3D printed device between two wells containing 40 µl solvent
control and attractant. Microscopic evaluation of fungal tip growth was performed after incubation for 6
h, 23°C for least two biological replicates, statistical significance was calculated with a two-tailed t-test
(*, p < 0.05). (C) Germling fusion rate of *C. graminicola* wild-type germlings spread on water agar (1%
serva agar, 1% agarose, 25 mM NaNO$_3$) adjusted to different medium pH with and without buffering. 50
µl of oval conidia (c = 5 x 10^7 ml^{-1}) were spread on the corresponding medium plates. After 17 h of
incubation at 23°C, at least 100 contacting germlings and conidia were microscopically examined for
fusion bridge formation, n ≥ 4. Statistical significance was calculated with a two-tailed t-test
corresponding to control conditions (c) *, p < 0.05.

In an earlier report, we investigated the impact of *Cgso* deletion on developmental processes
in *C. graminicola*. Besides the absence of germling and hyphal fusion, this mutant shows a
strong conidiation phenotype, provoked by the impaired nutrition of asexual fruiting bodies, the
acervuli (Nordzieke, 2022). Therefore, Δ*Cgso* appear dark and generate increased levels of
aerial mycelia in contrast to wildtype. In contrast, the a-pheromone receptor mutant Δ*Cgste3*
shows normal level of conidiation, indicating that this strain is able to undergo vegetative
fusions. Indeed, microscopic investigations showed normal fusion frequencies of Δ*Cgste3*
compared to wildtype, when both strains are plated in optimal spore concentrations (Figure 3,
B and C). Interestingly, a decrease in spore concentration revealed a significant reduction of
Δ*Cgste3* fusion rate compared to wild-type CgM2, which was reversed in a *Cgste3*
complementing strain. At the same time, Δ*Cgste3* germination rates don't differ from the
wildtype (Figure S3). From these results, we concluded that pH might act as long-distance
signaling molecule, able to guided conidia or germlings towards each other, but dispensable
when germlings are in close proximitiy.

Next, we aimed to identify the source of pH gradient formation. In filamentous fungi and plants,
growth is strictly correlated to the activity of a plasma membrane located ATPase, PmaI. This

ATPase pumps protons into the extracellular space, creating pH gradients used for the uptake
of diverse molecules (Portillo, 2000, Palmgren, 2001, Li *et al.*, 2022, Zhgun *et al.*, 2020). Since
pmaI is an essential gene in many organisms (Harris *et al.*, 1994), we decided to use a
Demethoxycurcumin (DMCM), a described inhibitor of PmaI (Dao *et al.*, 2016), to test a
probable impact of PmaI-dependent pH gradients for germling fusion. Intriguingly, we see a
concentration-dependent inhibition of germling fusion compared to control conditions (Figure
3 D). At the same time, germination is inhibited by DMCM, further indicating that germination
and fusion are interconnected processes.

Indications for pH as a conserved fungal fusion initiation signal

At first, germling fusion was considered as a highly species-specific process, providing only
germling interaction and fusion of individuals of the same species or communication group
(Heller *et al.*, 2016, Serrano et al., 2018). However, this perspective is changing. For
filamentous fungi like *N. crassa* and *B. cinerea*, intraspecies germling interaction and even the
induction of the SO-MAK-2-dependent cell dialog was shown (Haj Hammadeh et al., 2022).
However, those interactions never result in cross-species progeny. In *N. crassa – B. cinerea*
interactions, the fusion process is abolished upon contact of both interaction partners. Instead
of cell wall degradation to allow fusion, a cell wall thickens (Haj Hammadeh et al., 2022).
Additionally, there are reports about true fusions in between closely related species. The plant
pathogenic fungi *Colletotrichum gloeosporioides* and *Colletotrichum siamense* form progeny
after fusion, which show signs of genetic recombination of both parental species, making
intraspecies fusion a driver of evolution and host adaptation (Mehta and Baghela, 2021, Haj
Hammadeh et al., 2022). A prerequisite for intraspecies fusion is the presence of common or
very similar interaction signals, which could guide individuals of two different species towards
each other. Since PmaI-dependent pH gradients, we were interested in testing, whether
C. graminicola germlings are able to interact with other fungal species.

In a first approach, we plated either fusion competent spores of *C. graminicola* and *B. cinerea*
or *C. graminicola* and *N. crassa* on Vogel's minimal medium, providing conditions for fusion
bridge formation in all three species. Since those fungi grow with different speed and have
distinct temperature optima, specific incubation times had to be chosen. For *C. graminicola –*
B. cinerea interactions, conidia were spread in different concentrations on Vogel's MM and
incubated at 23°C. Since *N. crassa* conidia germinate and fuse much faster than *C.*
graminicola, the latter was pre-incubated for 14 h. After that time-interval, *N. crassa* conidia
were added and both fungi incubated further at 28°C. Both described experimental setups
resulted in interspecies and intraspecies interactions, whereas the proportion of species-
species interactions was least (Figure 4, A - D).

The hallmark of germling interaction in *N. crassa* is the oscillatory cell-dialogue, in which SO
and MAK-2 show an alternate recruiting to the plasma membrane of interacting partners. At
the end of this dialogue, MAK-2 accumulates at the contact point. Intriguingly, in intraspecies
interactions with *N. crassa*, this late accumulation of MAK-2 was shown, indicating an intact
cell dialogue (Fleissner et al., 2009, Haj Hammadeh et al., 2022). To test whether this is true
for *N. crassa – C. graminicola* interactions as well, a *N. crassa* wildtype strain expressing MAK-
2-GFP was used in interaction studies with *C. graminicola*. Likewise as described for *N. crassa*
– N. crassa and *N. crassa – B. cinerea* interactions, MAK-2-GFP accumulates at the contact
point with *C. graminicola* germlings (Supplementary Figure 4).

Figure 3 Sensing of acidic pH is mediated by the a-pheromone receptor CgSte3 and the CWI MAPK scaffold protein CgSo. (A) Chemotropic index of *C. graminicola* germlings of wildtype (CgM2), *Cgste3* and *Cgso* deletion strains and corresponding complementations to potassium phosphate buffer adjusted to pH 7.5. 20 µl of oval conidia (c = 5 x 10⁵ ml⁻¹) were applied in a 3D printed device between two wells containing 40 µl solvent control and attractant. Microscopic evaluation of fungal tip growth was performed after incubation for 6 h, 23°C for least two biological replicates, statistical significance was calculated with a two-tailed t-test to CgM2 (*, p < 0.05). (B) Microscopic evaluation of germling fusions after spreading of 50 µl *C. graminicola* wildtype (CgM2) or Δ*Cgste3* oval conidia spore solutions (c = 5 x 10⁷ ml⁻¹) on water agar (1% serva agar, 1% agarose, 25 mM NaNO₃) and incubation for 17 h, 23°C, scale = 10 µm, fusions are indicated with white asterisk. (C) Spore concentration-dependent germling fusion rates of *C. graminicola* wildtype, Δ*Cgste3* and Δ*Cgste3* complementation strain Δ*Cgste3*::*Cgste3*. 10 µl of oval conidia spore solutions were dropped on water agar (1% serva agar, 1% agarose, 25 mM NaNO₃) to reach spore amounts of 10⁴ - 5 x 10². Microscopic evaluation after 17 h of incubation at 23°C was performed for at least 100 contacting germlings and conidia in higher spore concentration. In lower spore concentrations, the full area was examined for fusion bridge formation, n ≥ 2, statistical significance was calculated with a two-tailed t-test to CgM2 values of the corresponding spore concentration (*, p < 0.05). (D) Germling fusion and germination rates of *C. graminicola* wildtype in dependence of treatment with the PmaI inhibitor Demethoxycurcumin (DMCM). 20 µl of different DMCM

concentrations or the control were applied on dried 10 µl drops of oval conidia (c = 10^6 ml^{-1}) on water
agar (1% serva agar, 1% agarose, 25 mM NaNO$_3$) and dried prior to incubation for 17 h at 23°C. Since
DMCM was dissolved in DMSO, diluted DMSO equalizing 15 µM DMCM solutions were used as control.
Germling fusion was accessed for at least 100 contacting conidia or germlings. Germination rates were
determined by evaluating at least 100 conidian, n = 3, statistical significance was calculated with a two-
tailed t-test to control conditions (-), *, $p < 0.05$.

Together these results indicate, that indeed all three fungi share a common interaction signal,
but indicate also species-specific signals, explaining the higher proportion of self-interactions
observed. To test whether the shard interaction signal might be pH, conidia of all three species
were grown on Vogel's MM adjusted to different pH without buffering the medium. As seen for
C. graminicola on water agar, interaction frequency of all three species is pH dependent,
although to different extents: *C. graminicola* shows the most pH-stable interaction pattern,
reduced only significantly at basic pHs (Figure 4 E). The interaction frequency of *B. cinerea*
already dropped at pH 7, and in *N. crassa*, pH of 6.5 already reduced interaction rates
significantly. A similar pattern can be seen for the germination of conidia on Vogel's MM.
Germination rates of *C. graminicola* oval conidia is hardly affected also at high pH and the
germination rates of *B. cinerea* are stable until pH 9. However, *N. crassa* conidia are highly
sensitive to pH changes, showing already significant reduction on medium pH of 6.5
(Supplementary Figure 5).

Discussion

The perception of extracellular signals is important for fungi to find nutrients, hosts, toxic
environments, or germlings and spores of the same species (Turrà *et al.*, 2016). In this study,
we identified pH as a communication signal outgoing from secretomes harvested during the
germling fusion. Already in earlier studies, pH was identified as a fusion-modulation factor:
over a broad pH range fusions *Fusarium oxysporum* take place, however, when the medium
is buffered, only pH of 6.4 still allows fusion to occur (Kurian et al., 2018). From those results,
it was concluded that an active modulation of pH by the fungal germlings are required for
germling fusion. In our experiments, we see a similar influence on pH of fusion frequency of
C. graminicola: whereas not-buffered media allow fusions over a broad pH spectrum, buffering
abolishes fusion bridge formation. Analysis of chemotropic growth further emphasized the role
of distinct pHs (7.5) on germling attraction and thus their signaling nature (Figure 2). Likewise,
acidic pH was shown to induce chemotropic growth in germlings of *Aspergillus nidulans*
(Yamamoto *et al.*, 2024). We further identified the plasma membrane localized ATPase PmaI
as probable source for pH gradient formation using the specific PmaI-inhibitor
Demthoxycurcumin (Dao et al., 2016). In fungi and plants, PmaI ATPase-formed pH gradient
drive the uptake of nutrients and its activity is directly linked to cell growth (Serrano, 1984,
Divon and Fluhr, 2007, Yamamoto et al., 2024, Palmgren, 2001, Haruta *et al.*, 2010,
Morsomme and Boutry, 2000). Additionally to the provision of an extracellular proton gradient,
PmaI activity results in a lower cytosolic pH, which is involved in the regulation of MAPK
pathways and thus affects development and pathogenicity in *F. oxysporum* (Mariscal *et al.*,
2022). Intriguingly, PmaI was also shown to be crucial for the sensing of acidic pH in *A.
nidulans*, however it is unclear how the determining mechanism works (Yamamoto et al.,
2024).

Figure 4 Working model for the pH-dependent induction of the cell dialogue. Pmal-depenent pH gradients (H⁺) form immediately when a conidium starts to germinate. Those pH gradients are sensed by surrounding germlings or conidia via the a-pheromone receptor Ste3, resulting in the activation of the Cell Wall Integrity (CWI) MAPK pathway and induction of growth redirection towards the interaction partner. From work in other fungi, we propose that CWI activation by Ste3 induces the relocalization of SO to the tip of the interacting germling, followed by the release of a SO-dependent, so far unknown communication signal (blue circles).

pH is a crucial environmental factor for fungi and its fluctuations influence diverse cellular processes like cell wall remodeling, nutrient availability, and enzyme activity (Li *et al.*, 2021).

Many plant and human pathogenic fungi modulate the extracellular pH to explore their surroundings, enable successful host infection and maintain cellular homeostasis (Masachis *et al.*, 2016, Fernandes *et al.*, 2017, Kane, 2016, Alkan *et al.*, 2013, Vylkova, 2017, Kesten *et al.*, 2019). Traditionally, plant-interacting fungi were thought to be strictly alkalizing or acidifying their surroundings. However, in the last decade it was shown, that induced changes in ambient pH are flexible and depend on carbon availability (Bi *et al.*, 2016). Sensing of extracellular, alkaline pH is primarily mediated by the pal/RIM pathway in fungi, with PalH as the central pH sensor (Selvig and Alspaugh, 2011). Together with MAPK modules, the pal/RIM pathway is central for the sensing and the response to extracellular pH levels, enabling to maintain cellular integrity also in the face of stress and pathogenic host interaction (Lara-Martínez *et al.*, 2025). PalH is a 7 transmembrane receptor resembling classical G-protein coupled receptors (GPCR), interacting downstream with an arrestin for signal transduction (Lucena-Agell *et al.*, 2016). As transcription factor downstream of PalH, PacC is described, acting as activator of gene expression at alkaline pH and as a repressor of genes expressed at acidic pH (Lara-Martínez et al., 2025). In contrast, the sensor for acidic environmental pH is unknown in fungi (Yamamoto et al., 2024). In this study, we provide evidence that the a-pheromone receptor CgSte3 mediates sensing of acidic pH gradients, activating downstream the CWI pathway (Figure 3). Fungal pheromone receptors are GPCRs first described in the yeast *Saccharomyces cerevisiae*, in which they mediate chemotropic attraction of mating partners during sexual fusion (Hagen and Sprague Jr, 1984, Nakayama *et al.*, 1985, Jenness *et al.*, 1986). In the last years, new roles for fungal pheromone receptors were identified. This includes the spore density dependent regulation of germination in *F. oxysporum*, and the sensing of plant defense molecules like class III peroxidases and diterpenoids in different host-pathogen systems (Turrà et al., 2015, Vitale et al., 2019, Sharma et al., 2022, Rudolph et al., 2024a, Rudolph et al., 2024b). Downstream of diterpenoid and peroxidase sensing, in those studies the CWI was shown to be responsible for consequent growth redirection (Turrà et al., 2015, Sharma et al., 2022, Rudolph et al., 2024b), indicating a conserved link between external signal sensing by pheromone receptors and CWI activation. So far, only one study investigated how the two pheromone receptors interact upon peroxidase sensing. They results point to a scenario, in which peroxidase directly or a peroxidase-generated ligand stimulates heterodimer formation in between Ste2 and Ste3 of *Fusarium graminearum* (Sharma et al., 2022). Intriguingly, *C. graminicola* is the only species, in which the α-pheromone receptor Ste2 was lost during evolution (Wilson *et al.*, 2021) and therefore the recognition of external signals is dependent only on the a-pheromone receptor Ste3. How the perception in this one-receptor system functions on the molecular level, is still to be explored. Another open question is how those different attractant molecules are sensed by the same receptor. Since the chemical properties of the perceived signals are such diverse, a direct ligand-receptor interaction seems unlikely. For mammalian cells it was shown that GPCRs and growth factor receptor tyrosine kinases (RTKs) can activate each other in a process termed 'transactivation', without the required of ligand binding for the secondary receptor (Wang, 2016, Delcourt *et al.*, 2007). In such a scenario, the a-pheromone receptor CgSte3 could be associated with other, more specific receptors, which serve as sensors for diterpenoids, peroxidases or pH specifically. Additionally, modifications of amino acids can induce GPCR activation (Wang *et al.*, 2018). In mammalian cells, a current study shows that protonation of histidine can induce GPCR activation (Guo *et al.*, 2025). As a third possibility, pH could influence the binding-capacity of another ligand by CgSte3, resulting in receptor activation (Sun *et al.*, 2025). Which of these possibilities or also other probable mechanisms are truly mediating CgSte3 receptor activation, has to be investigated in future studies.

Together, our results about *C. graminicola* CAT fusion point to a scenario, in which Pmal-depenent pH gradients are formed as soon as a conidium starts to germinate. Those pH gradients are sensed by surrounding germlings or conidia via the a-pheromone receptor

CgSte3, resulting in the activation of CWI and induction of growth redirection towards the
interaction partner. From work in other fungi, especially *N. crassa*, we know the oscillatory
recruitment of SO and MAK-2 during germling interaction is crucial for fusion bridge formation
(Fleissner et al., 2009, Serrano et al., 2018). Since CgSo is a scaffold of CWI, pH sensing
might additionally induce CgSo localization to the plasma membrane, which could be
associated with the secretion of a second interaction signal, still to be discovered (Figure 5).
This model could also explain why germling fusions are absent in falcate conidia: those spores
have a pronounced dormant phase, mediate by secondary metabolites like mycosporines and
other factors (Leite and Nicholson, 1992, Vasselli *et al.*, 2022, Chaky et al., 2001). Since
germination of falcate conidia does not occur under starvation conditions, PmaI is inactive and
no pH gradients build up, making an induction of the fusion process impossible. However,
when oval conidia are close by, they can provide the pH fusion-initiation signal, which is
perceived by falcate conidia and subsequently induce dormancy breaking, germling
interaction, and fusion.

Figure 5 Interaction of *C. graminicola* with *N. crassa* and *B. cinerea*. (A-D) Inter- and Intraspecies
interactions of *C. graminicola* and *B. cinerea* or *N. crassa* on Vogels minimal medium (MM). (A, C) The
interactions experiment with *C. graminicola* and *B. cinerea* was carried out at 23°C with an incubation
time of 17 h. First, *C. graminiocla* oval conidia (50 µl of c = 5 x 10^7 ml^{-1}) and 300 µl *B. cinerea* spore
solution (c= 3.33 x 10^7 ml^{-1}) were spread together on Vogels' MM. (B, D) To access *C. graminicola* and
N. crassa germling interactions, 50 µl of a *C. graminicola* oval conidia spore solution (c = 5x10^7 x ml^{-1})
was distributed on incubation medium and incubated for 14 h at 23°C. After that time, 300µl of *N. crassa*
spore solution (c = 10^7 ml^{-1}) was added on those pre-incubated plates and further incubated at 28°C for
3.5 - 4 h. (A-B) Microscopic images of intra- and interspecies interaction, interactions are marked with a
black asterisk, scale = 10 µm. (C-D) Inter- and intraspecies interaction rates of ≥ 76 germling interactions
of *C. graminicola* and *B. cinerea* (C) and *C. graminicola* with *N. crassa* (D), n = 3. (E) Germling interaction
rates of *C. graminicola*, *B. cinerea* and *N. crassa* on Vogel's minimal medium adjusted to different pH.
Species-specific conidia amounts (*C. graminicola*: 50 µl of 5 x 10^7 ml^{-1}; *B. cinerea* and *N. crassa*: 300
µl of c = 10^7 ml^{-1}) were spread on Vogels' MM and incubated at 23°C for 17 h (*C. graminicola*, *B. cinerea*)
or 28°C for 3,5 - 4 h (*N. crassa*), n ≥ 3, statistical significance was calculated with a two-tailed t-test to
control conditions (c), *, p < 0.05.

The ability of germlings to fuse in an early stage of colony formation, fastens radial growth, by
sharing of nutrients or cellular components (Roca *et al.*, 2005, Richard et al., 2012). Recently,
studies have shown that germling communication mediating bidirectional chemotropic growth

prior to fusion is not species-specific (Haj Hammadeh et al., 2022). Closely related species of
Colletotrichum and *Aspergillus* are even able to fuse and exchange genetic material, resulting
in progeny with recombinant genomes (Ishitani *et al.*, 1956, Mehta and Baghela, 2021). In this
study, we show that the maize pathogen *C. graminicola* communicates with the saprophyte *N.*
crassa and the pathogen *B. cinerea* with 8 and 6 % of interactions respectively. The three
species differ strongly in their preferred habitat. *N. crassa* is a saprophytic fungus primarily
living on burned vegetation in humid subtropical and tropical regions (Turner *et al.*, 2001). *B.*
cinerea, the grey mold-causing fungus, has a high host range including several vegetables like
tomato, lettuce, and beans, or fruit crops such as strawberries and grapes (Chen *et al.*, 2023).
Even though *B. cinerea* cause lesions on *Zea mays* under artificial conditions, cereals are not
host plants of the pathogen (Bi *et al.*, 2023), thus it is unlikely that those fungi meet in the
environment. In line, we never observed the outgrowth of new fungal hyphae from the contact
point of intraspecies interactions, as it was reported for the interaction of *C. siamense* and *C.*
gloeosporioides (Mehta and Baghela, 2021). Intriguingly, the two pathogenic species *B.*
cinerea and *C. graminicola* showed more stable fusion and germination frequencies as the
saprophytic fungus *N. crassa*. This is in line with previous observations, that especially plant-
interacting species are able to tolerate wide ranges of ambient pH (Felix *et al.*, 1993).

In conclusion, our study has provided new insights into the molecular mechanisms underlying
germling fusion in the maize pathogen *C. graminicola*. We have identified pH as a
communication signal outgoing from secretomes harvested during the germling fusion, and
have shown that the plasma membrane localized ATPase PmaI is a probable source for pH
gradient formation. Our results also suggest that the a-pheromone receptor CgSte3 mediates
sensing of acidic pH gradients, activating downstream the CWI MAPK pathway. These findings
are consistent with previous studies that have shown that pH is a crucial environmental factor
for fungi and its fluctuations influence diverse cellular processes. Our results also suggest that
the maize pathogen *C. graminicola* communicates with the saprophyte *N. crassa* and the
pathogen *B. cinerea*, although these three species differ strongly in their preferred habitat.
Overall, our study contributes to a better understanding of the molecular mechanisms
underlying germling fusion in fungi and indicate pH as fusion initiation signal across species
borders.

Acknowledgments

We thank Gabriele Beyer and Gertrud Stahlhut for excellent technical assistance and Lilith
Elisabeth Gück and Simone Lewandowski for help with experimental work. This work was
funded by the Deutsche Forschungsgemeinschaft (Bonn-Bad Godesberg). Grant was provided
to D.E.N. (project NO 1230/3-1 (447175909). This work was partly supported by the Göttingen
Graduate Center for Neurosciences, Biophysics, and Molecular Biosciences at the Georg-
August-Universität Göttingen.

References

Alkan, N., Espeso, E. A. and Prusky, D. (2013) "Virulence regulation of phytopathogenic fungi
by pH." *Antioxidants & Redox Signaling,* 19**,** 1012-1025.
Bastiaans, E., Debets, A. J. and Aanen, D. K. (2015) "Experimental demonstration of the
benefits of somatic fusion and the consequences for allorecognition." *Evolution,* 69**,**
1091-1099.
Belisário, R., Robertson, A. E. and Vaillancourt, L. J. (2022) "Maize anthracnose stalk rot in
the genomic era." *Plant Disease,* 106**,** 2281-2298.

Bi, F., Barad, S., Ment, D., et al. (2016) "Carbon regulation of environmental pH by secreted small molecules that modulate pathogenicity in phytopathogenic fungi." *Molecular plant pathology,* 17**,** 1178-1195.

Bi, K., Liang, Y., Mengiste, T. and Sharon, A. (2023) "Killing softly: a roadmap of *Botrytis cinerea* pathogenicity." *Trends in plant science,* 28**,** 211-222.

Chaky, J., Anderson, K., Moss, M. and Vaillancourt, L. (2001) "Surface hydrophobicity and surface rigidity induce spore germination in *Colletotrichum graminicola.*" *Phytopathology,* 91**,** 558-564.

Chen, T., Zhang, Z., Chen, Y., Li, B. and Tian, S. (2023) "*Botrytis cinerea.*" *Current Biology,* 33**,** R460-R462.

Dao, T. T., Sehgal, P., Tung, T. T., et al. (2016) "Demethoxycurcumin is a potent inhibitor of p-type ATPases from diverse kingdoms of life." *PLoS One,* 11**,** e0163260.

Delcourt, N., Bockaert, J. and Marin, P. (2007) "GPCR-jacking: from a new route in RTK signalling to a new concept in GPCR activation." *Trends in pharmacological sciences,* 28**,** 602-607.

Divon, H. H. and Fluhr, R. (2007) "Nutrition acquisition strategies during fungal infection of plants." *FEMS microbiology letters,* 266**,** 65-74.

Felix, G., Regenass, M. and Boller, T. (1993) "Specific perception of subnanomolar concentrations of chitin fragments by tomato cells: induction of extracellular alkalinization, changes in protein phosphorylation, and establishment of a refractory state." *The Plant Journal,* 4**,** 307-316.

Fernandes, T. R., Segorbe, D., Prusky, D. and Di Pietro, A. (2017) "How alkalinization drives fungal pathogenicity." *PLoS pathogens,* 13**,** e1006621.

Fischer, M. S. and Glass, N. L. (2019) "Communicate and fuse: how filamentous fungi establish and maintain an interconnected mycelial network." *Frontiers in microbiology,* 10**,** 619.

Fleissner, A., Leeder, A. C., Roca, M. G., Read, N. D. and Glass, N. L. (2009) "Oscillatory recruitment of signaling proteins to cell tips promotes coordinated behavior during cell fusion." *Proceedings of the National Academy of Sciences,* 106**,** 19387-19392.

Forgey, W., Blanco, M. and Loegering, W. (1978) "Differences in pathological capabilities and host specificity of *Colletotrichum graminicola* on *Zea mays.*" *Plant Disease Reporter,* 62**,** 573-576.

Fu, C., Iyer, P., Herkal, A., Abdullah, J., Stout, A. and Free, S. J. (2011) "Identification and characterization of genes required for cell-to-cell fusion in *Neurospora crassa.*" *Eukaryotic cell,* 10**,** 1100-1109.

Glass, N. L., Rasmussen, C., Roca, M. G. and Read, N. D. (2004) "Hyphal homing, fusion and mycelial interconnectedness." *Trends in microbiology,* 12**,** 135-141.

Guo, L., Zhu, K., Zhong, Y.-N., et al. (2025) "Structural basis and biased signaling of proton sensation by GPCRs mediated by extracellular histidine rearrangement." *Molecular Cell,* 85**,** 1658-1673. e1657.

Hagen, D. C. and Sprague Jr, G. F. (1984) "Induction of the yeast α-specific STE3 gene by the peptide pheromone a-factor." *Journal of molecular biology,* 178**,** 835-852.

Haj Hammadeh, H., Serrano, A., Wernet, V., et al. (2022) "A dialogue-like cell communication mechanism is conserved in filamentous ascomycete fungi and mediates interspecies interactions." *Proceedings of the National Academy of Sciences,* 119**,** e2112518119.

Harris, S. L., Na, S., Zhu, X., et al. (1994) "Dominant lethal mutations in the plasma membrane H (+)-ATPase gene of *Saccharomyces cerevisiae.*" *Proceedings of the National Academy of Sciences,* 91**,** 10531-10535.

Haruta, M., Burch, H. L., Nelson, R. B., et al. (2010) "Molecular characterization of mutant *Arabidopsis* plants with reduced plasma membrane proton pump activity." *Journal of Biological Chemistry,* 285**,** 17918-17929.

Heller, J., Zhao, J., Rosenfield, G., Kowbel, D. J., Gladieux, P. and Glass, N. L. (2016) "Characterization of greenbeard genes involved in long-distance kind discrimination in a microbial eukaryote." *PLoS biology,* 14**,** e1002431.

Herzog, S., Schumann, M. R. and Fleißner, A. (2015) "Cell fusion in *Neurospora crassa.*" *Current opinion in microbiology,* 28**,** 53-59.

Ishitani, C., Ikeda, Y. and Sakaguchi, K.-I. (1956) "Hereditary variation and genetic
recombination in koji-molds (*Aspergillus oryzae* and *Asp. sojae*) VI. Genetic
recombination in heterozygous diploids." *The Journal of General and Applied*
Microbiology, 2**,** 401-430.
Jenness, D. D., Burkholder, A. C. and Hartwell, L. H. (1986) "Binding of α-factor pheromone to
Saccharomyces cerevisiae a cells: dissociation constant and number of binding sites."
Molecular and cellular biology, 6**,** 318-320.
Kane, P. M. (2016) Proton transport and pH control in fungi. In: *Yeast Membrane Transport.*
Ramos, J., Sychrová, H. and Kschischo, M., eds.: Springer, pp. 33-68.
Kesten, C., Gámez-Arjona, F. M., Menna, A., et al. (2019) "Pathogen-induced pH changes
regulate the growth-defense balance in plants." *The EMBO Journal,* 38**,** e101822.
Kurian, S. M., Di Pietro, A. and Read, N. D. (2018) "Live-cell imaging of conidial anastomosis
tube fusion during colony initiation in *Fusarium oxysporum*." *PLoS One,* 13**,** e0195634.
Lara-Martínez, D., Tristán-Flores, F. E., Cervantes-Montelongo, J. A. and Silva-Martínez, G.
A. (2025) "Fungal stress responses and the importance of GPCRs." *Journal of Fungi,*
11**,** 213.
Leite, B. and Nicholson, R. L. (1992) "Mycosporine-alanine: A self-inhibitor of germination from
the conidial mucilage of *Colletotrichum graminicola*." *Experimental mycology,* 16**,** 76-
86.
Li, B., Chen, Y. and Tian, S. (2021) "Function of pH-dependent transcription factor PacC in
regulating development, pathogenicity, and mycotoxin biosynthesis of phytopathogenic
fungi." *The FEBS Journal,* 289**,** 1723-1730.
Li, J., Yang, S., Li, D., Peng, L., Fan, G. and Pan, S. (2022) "The plasma membrane H^+-ATPase
is critical for cell growth and pathogenicity in *Penicillium digitatum*." *Applied*
Microbiology and Biotechnology, 106**,** 5123-5136.
Lucena-Agell, D., Hervás-Aguilar, A., Múnera-Huertas, T., et al. (2016) "Mutational analysis of
the *Aspergillus* ambient pH receptor PalH underscores its potential as a target for
antifungal compounds." *Molecular Microbiology,* 101**,** 982-1002.
Mariscal, M., Miguel-Rojas, C., Hera, C., Fernandes, T. R. and Di Pietro, A. (2022) "*Fusarium*
oxysporum casein kinase 1, a negative regulator of the plasma membrane H^+-ATPase
Pma1, is required for development and pathogenicity." *Journal of Fungi,* 8**,** 1300.
Masachis, S., Segorbe, D., Turrà, D., et al. (2016) "A fungal pathogen secretes plant
alkalinizing peptides to increase infection." *Nature Microbiology,* 1**,** 1-9.
Mehta, N. and Baghela, A. (2021) "Quorum sensing-mediated inter-specific conidial
anastomosis tube fusion between *Colletotrichum gloeosporioides* and *C. siamense*."
IMA fungus, 12**,** 1-17.
Morsomme, P. and Boutry, M. (2000) "The plant plasma membrane H^+-ATPase: structure,
function and regulation." *Biochimica et Biophysica Acta (BBA)-Biomembranes,* 1465**,**
1-16.
Nakayama, N., Miyajima, A. and Arai, K. (1985) "Nucleotide sequences of STE2 and STE3,
cell type-specific sterile genes from *Saccharomyces cerevisiae*." *The EMBO journal,* 4**,**
2643.
Nordzieke, D. E. (2022) "Hyphal fusions enable efficient nutrient distribution in *Colletotrichum*
graminicola conidiation and symptom development on maize." *Microorganisms,* 10**,**
1146.
Nordzieke, D. E., Sanken, A., Antelo, L., Raschke, A., Deising, H. B. and Pöggeler, S. (2019)
"Specialized infection strategies of falcate and oval conidia of *Colletotrichum*
graminicola." *Fungal Genetics and Biology,* 133**,** 103276.
O'Connell, R. J., Thon, M. R., Hacquard, S., et al. (2012) "Lifestyle transitions in plant
pathogenic *Colletotrichum* fungi deciphered by genome and transcriptome analyses."
Nature genetics, 44**,** 1060-1065.
Palmgren, M. G. (2001) "Plant plasma membrane H^+-ATPases: powerhouses for nutrient
uptake." *Annual review of plant biology,* 52**,** 817-845.
Panaccione, D. G., Vaillancourt, L. J. and Hanau, R. M. (1989) "Conidial dimorphism in
Colletotrichum graminicola." *Mycologia,* 81**,** 876-883.

Portillo, F. (2000) "Regulation of plasma membrane H^+-ATPase in fungi and plants." *Biochimica et Biophysica Acta (BBA)-Reviews on Biomembranes,* 1469, 31-42.

Quidde, T., Büttner, P. and Tudzynski, P. (1999) "Evidence for three different specific saponin-detoxifying activities in *Botrytis cinerea* and cloning and functional analysis of a gene coding for a putative avenacinase." *European Journal of Plant Pathology,* 105, 273-283.

Ramaswe, J. B., Steenkamp, E. T., De Vos, L., Fru, F. F., Adegeye, O. O. and Wingfield, B. D. (2024) "Sex pheromone receptor Ste2 orchestrates chemotropic growth towards pine root extracts in the pitch canker pathogen *Fusarium circinatum.*" *Pathogens,* 13, 425.

Read, N. D., Goryachev, A. B. and Lichius, A. (2012) "The mechanistic basis of self-fusion between conidial anastomosis tubes during fungal colony initiation." *Fungal Biology Reviews,* 26, 1-11.

Richard, F., Glass, N. and Pringle, A. (2012) "Cooperation among germinating spores facilitates the growth of the fungus, *Neurospora crassa.*" *Biology Letters,* 8, 419-422.

Roca, G. M., Read, N. D. and Wheals, A. E. (2005) "Conidial anastomosis tubes in filamentous fungi." *FEMS microbiology letters,* 249, 191-198.

Rudolph, A., Schunke, C., Sasse, C., et al. (2024a) "Maize diterpenoid sensing via Ste3 a-pheromone receptor and rapid germination of *Colletotrichum graminicola* oval conidia facilitating root infection." *bioRxiv,* 2024.2004. 2005.588234.

Rudolph, A. Y., Schunke, C. and Nordzieke, D. E. (2024b) "Conserved perception of host and non-host signals via the a-pheromone receptor Ste3 in *Colletotrichum graminicola.*" *Frontiers in Fungal Biology,* 5, 1454633.

Ruxton, G. D. (2006) "The unequal variance t-test is an underused alternative to Student's t-test and the Mann–Whitney U test." *Behavioral Ecology,* 17, 688-690.

Schunke, C., Pöggeler, S. and Nordzieke, D. E. (2020) "A 3D printed device for easy and reliable quantification of fungal chemotropic growth." *Frontiers in Microbiology,* 11, 584525.

Selvig, K. and Alspaugh, J. A. (2011) "pH response pathways in fungi: adapting to host-derived and environmental signals." *Mycobiology,* 39, 249-256.

Serrano, A., Illgen, J., Brandt, U., et al. (2018) "Spatio-temporal MAPK dynamics mediate cell behavior coordination during fungal somatic cell fusion." *Journal of cell science,* 131, jcs213462.

Serrano, R. (1984) "Plasma membrane ATPase of fungi and plants as a novel type of proton pump." *Current topics in cellular regulation,* 23, 87-126.

Sharma, T., Sridhar, P. S., Blackman, C., et al. (2022) "*Fusarium graminearum* Ste3 G-protein coupled receptor: A mediator of hyphal chemotropism and pathogenesis." *Msphere,* 7, e00456-00422.

Sridhar, P. S., Trofimova, D., Subramaniam, R., et al. (2020) "Ste2 receptor-mediated chemotropism of *Fusarium graminearum* contributes to its pathogenicity against wheat." *Scientific Reports,* 10, 10770.

Sun, L., Su, T., Fisher, A. C., et al. (2025) "Dynamic protonation of ligand sites in molecular catalysts enhances electrochemical CO_2 reduction." *Science Advances,* 11, eadu6915.

Turner, B. C., Perkins, D. D. and Fairfield, A. (2001) "*Neurospora* from natural populations: a global study." *Fungal Genetics and Biology,* 32, 67-92.

Turrà, D., El Ghalid, M., Rossi, F. and Di Pietro, A. (2015) "Fungal pathogen uses sex pheromone receptor for chemotropic sensing of host plant signals." *Nature,* 527, 521-524.

Turrà, D., Nordzieke, D., Vitale, S., El Ghalid, M. and Di Pietro, A. (2016) Hyphal chemotropism in fungal pathogenicity. In: *Seminars in cell & developmental biology.* Elsevier, pp. 69-75.

Vasselli, J. G., Hancock, H., Bedsole, C. O., Kainer, E., Chappell, T. M. and Shaw, B. D. (2022) "The conidial coin toss: A polarized conidial adhesive in *Colletotrichum graminicola.*" *Fungal Genetics and Biology,* 163, 103747.

Vitale, S., Di Pietro, A. and Turrà, D. (2019) "Autocrine pheromone signalling regulates
community behaviour in the fungal pathogen *Fusarium oxysporum*." *Nature*
microbiology, 4**,** 1443-1449.
Vogel, H. J. (1956) "A convenient growth medium for *Neurospora* (Medium N)." *Microbial*
Genet. Bull., 13**,** 42-43.
Vylkova, S. (2017) "Environmental pH modulation by pathogenic fungi as a strategy to conquer
the host." *PLoS pathogens,* 13**,** e1006149.
Wang, W., Qiao, Y. and Li, Z. (2018) "New insights into modes of GPCR activation." *Trends in*
pharmacological sciences, 39**,** 367-386.
Wang, Z. (2016) "Transactivation of epidermal growth factor receptor by G protein-coupled
receptors: recent progress, challenges and future research." *International journal of*
molecular sciences, 17**,** 95.
Wilson, A. M., Lelwala, R. V., Taylor, P. W., Wingfield, M. J. and Wingfield, B. D. (2021)
"Unique patterns of mating pheromone presence and absence could result in the
ambiguous sexual behaviors of *Colletotrichum* species." *G3,* 11**,** jkab187.
Yamamoto, R., Miki, H., Itani, A. and Takeshita, N. (2024) "Hyphae of the fungus *Aspergillus*
nidulans demonstrate chemotropism to nutrients and pH." *Plos Biology,* 22**,** e3002726.
Zhgun, A., Dumina, M., Valiakhmetov, A. and Eldarov, M. (2020) "The critical role of plasma
membrane H$^+$-ATPase activity in cephalosporin C biosynthesis of *Acremonium*
chrysogenum." *PLoS One,* 15**,** e0238452.