Title: Protein kinase CK2α' as a dual modulator of immune signaling and synaptic

dysfunction in Tauopathy

Authors: Angel White[1], Peter Gavrilyuk[1], Persephone Gu[1], Rafael Falcon Moya[1], Reid

Thurston[1], Amal Fickak[1], Nicholas B Rozema[1], Prarthana Keshavaram[1], Scott Vermilyea[1], Riley

Schlichte[1], Joyce Meints[1], Ying Zhang[1], Alfonso Araque[1], Michael Lee[1], Rocio Gomez-Pastor[1].

Affiliation: [1]University of Minnesota, School of Medicine, Department of Neuroscience,

Minneapolis, MN, USA. [2]Minnesota Supercomputing Institute, University of Minnesota,

Minneapolis, MN, USA.

One Sentence summary: This study highlights CK2α' as a key node at the intersection of tau

pathology, synaptic dysfunction, and neuroimmune signaling.

Abstract:

Tauopathies are a group of neurodegenerative diseases characterized by tau accumulation,

neuroinflammation, and synaptic dysfunction, yet effective treatments remain elusive. Protein

Kinase CK2 has been previously associated with different aspects of tau pathology but genetic

evidence for the contribution of CK2 to tauopathy remained lacking. Here, we show CK2α', one

of the two catalytic subunits of CK2, as a novel regulator of tau-mediated neurodegeneration.

We found that CK2α' expression is elevated in postmortem brains of dementia patients and in

the hippocampus of PS19 tauopathy mice, especially in neurons and microglia. Using genetic

haploinsufficiency in PS19 mice, we demonstrated that reduced CK2α' levels significantly

decrease phosphorylated tau and total tau burden in the hippocampus and cortex. CK2α'

depletion also attenuated microglial activation, pro-inflammatory cytokine production, and

microglia synaptic engulfment, enhanced synaptic gene expression, synaptic density, and LTP.

Importantly, CK2α' depletion rescued cognitive deficits assessed in the Barnes maze. These

effects appear to be mediated through both neuronal and glial functions and may involve CK2α'-

dependent modulation of tau-associated phosphorylation and neuroinflammatory and immune

signaling pathways.

Main text:

Background

Tauopathies encompass a broad range of neurodegenerative conditions, most notably

Alzheimer's disease (AD) and related dementias (ADRD), for which no cure currently exists (*1-*

3). Tauopathies are typically marked by progressive neuronal loss, synaptic and cognitive

dysfunction, and widespread inflammation in both the central nervous system and peripheral

tissues(*4-8*). A deeper understanding of the molecular mechanisms underlying tauopathies and

their downstream consequences is essential for developing effective disease-modifying

therapies. One key area of research is the role neuroinflammation plays in the progression of

tauopathies. These disorders are characterized by chronic, excessive inflammatory responses

to abnormal tau proteins (*5, 6, 9*). This persistent inflammation can exacerbate tau pathology by

promoting sustained neuronal damage and dysfunction (*9-12*). However, despite ongoing

research, the mechanisms that connect neuroinflammation and tau pathology are still poorly

understood.

Protein kinase CK2 (formerly known as Casein Kinase II), is a serine/threonine kinase that has

emerged as an important regulator of neuroinflammation and protein homeostasis in several

neurodegenerative diseases (*13-15*), and it has previously been connected to tau pathology in

AD (*16-19*). However, its specific role in the inflammatory pathways associated with tauopathies

remains poorly defined. Interestingly, CK2 has demonstrated potential as a therapeutic target in

various neurodegenerative contexts (*14, 15*). In AD, CK2 has been implicated in processes such

as synaptic plasticity (*20, 21*), amyloid precursor protein (APP) processing (*22-24*), and tau

accumulation (*18, 19*), highlighting its relevance for further investigation.

CK2 is a tetrameric enzyme composed of 2 regulatory β subunits (CK2β) and 2 catalytic

subunits α (CKα) or α' (CK2α') (*13*). CK2α and CK2α' are highly similar in structure but display

differential expression and substrate specificity (*25-31*). CK2α is expressed throughout the body

at relatively equal levels and has hundreds of substrates, while CK2α' expression is more

restricted to testes and brain and has very few validated substrates (*25-31*). Another important

difference between these two catalytic subunits is the embryonic lethality of CK2α knock-out

mice, while complete deletion of CK2α' does not have any major defects other than male sterility

(*30, 31*). Recent studies show that CK2α' is selectively upregulated in HD, and partial genetic

reduction of CK2α' in HD mouse models improves synaptic function, reduces protein

aggregation, and alleviates behavioral deficits, indicating its key role in disease pathology (*14,*

15).

CK2 was one of the first protein kinases identified to be altered in an AD brain (*32, 33*), but its

role in AD has been somewhat controversial due to discrepancies in the ability to detect

consistent alterations in different affected tissues. Early studies using pan-CK2 antibodies (non-

selective for CK2α'/CK2α subunits) showed increased levels of total CK2 in the hippocampus

and frontal cortex of AD mouse models and patients with AD, especially in glial cells (*34*) but

others have reported opposite results (*32, 35, 36*). In those studies where levels of CK2 were

elevated, the authors associated the increase in CK2 immunoreactivity with the activation of

neuroinflammatory processes and AD pathology (*34*).

Although genetic studies have not yet clarified whether CK2α and CK2α' differentially contribute

to AD pathology, multiple lines of evidence implicate CK2 in Tau phosphorylation and tau-

mediated dysfunction. Kinase inhibitor screens identified CK2 as a major driver of okadaic-acid–

induced pTau in N2a cells (*17*), and CK2-mediated phosphorylation of the PP2A inhibitor SET

increases pTau levels in neurons exposed to Aβ or overexpressing hTau (*16*). Moreover, mice

overexpressing CK2 exhibit elevated pTau, impaired synaptic plasticity, and cognitive deficits

reminiscent of tauopathies (*16*). Despite this, most studies have treated CK2 as a single

holoenzyme, overlooking known differences in substrate specificity between CK2α and CK2α'.

Determining whether these catalytic subunits play distinct roles in tauopathy is essential for

developing targeted therapies, especially given the current lack of subunit-selective CK2

inhibitors and the opportunity this presents for designing more precise and effective treatments.

Due to previous findings specifically connecting the dysregulation of CK2α' with various

neurodegenerative diseases and reports linking this subunit to the phosphorylation of Tau in

different contexts, we explored the role of CK2α' in various tauopathy models. We confirmed a

specific upregulation at the RNA and protein levels of CK2α' in affected tissues of AD and FTD

patients as well as a mouse model of tauopathy (PS19). CK2α' upregulation was specifically

observed in both neurons and microglia. Importantly, genetic depletion of CK2α' in both cell and

mouse models of tauopathy (expressing Tau-P301L or Tau-P301S mutations) resulted in

decreased pTau. Furthermore, PS19 mice haploinsufficient for CK2α' showed attenuated

microglial activation, pro-inflammatory cytokine production, and microglia synaptic engulfment,

enhanced synaptic gene expression, synaptic density, and LTP. Importantly, CK2α' depletion

rescued cognitive deficits assessed in the Barnes maze. Overall, our data showed a large role

of the CK2α' catalytic subunit specifically in tau pathology, largely connected to the modulation

of microglia and synaptic functions. These studies highlight CK2α' as an attractive therapeutic

target for the treatment of AD/ADRD.

Results

Levels of Protein kinase CK2α' are increased in the brains of dementia patients and

mouse models of tauopathy

There are two different genes (*Csnk2a1* and *Csnk2a2*) that code for two different CK2 catalytic

subunits CK2α and CK2α', respectively. These two catalytic subunits share a high percent of

homology and they can arrange in various combinations (α-α, α-α' or α'-α') depending on their

expression and tissue availability although the canonical arrangement in the brain is expected to

be α-α' (**Fig. 1A, B**) (*13, 25-31*). Analyzing existing RNA-seq data from the Allen Brain Institute:

Aging, Dementia, and Traumatic Brain Injury (TBI) Study (*37*) we found a specific increase in

the expression of *CSNK2a2* (CK2α'), but not in *CSNK2a1* (CK2α), in the parietal cortex (PCx) of

patients with dementia compared to non-dementia controls (**Fig. 1C-E**). We also found a trend

towards increased CK2α' in the temporal cortex (TCx) and hippocampus (HIP) although data did

not reach statistical significance (**Fig. 1C**) (*37*). Importantly, we also observed a significant

increase in CK2α' protein levels, but not CK2α, in the frontal cortex of FTD patients compared to

age and sex-match controls (**Fig.1F left panel, Fig. 1 G, H, File S1**). Immunoblotting for CK2α'

in CK2α' knock-out mice confirmed antibody specificity (**Fig. 1F, right panel**). Enhanced levels

of CK2α' was also associated with the presence of tau pathology, as previously assessed by

AT8 immunostaining and reported in the sample metadata provided by the NeuroBioBank (**Fig.**

1I, File S1). The majority of FTD samples grouped in the top left quadrant representing high

levels of CK2α' and the presence of tau pathology, while most control samples grouped in the

bottom right quadrant representing low levels of CK2α' and absence of tau pathology. For those

two control samples (#1 and #6) in which tau pathology was observed, the AT8 phospho-tau

(pTau) signal and distribution differed from the canonical tau pathology associated with

dementia and individuals lacked classical cognitive decline (**File S1**).

To examine whether CK2α' was also elevated in mouse models of tauopathy we utilized the

PS19 mouse model, which expresses human tau with a P301S mutation associated with familial

forms of FTD and other tauopathies (*38*). Pathology and neuronal loss in PS19 mice are seen

largely in the hippocampus, although it does spread to other brain regions including the

neocortex, entorhinal cortex and amygdala (*38, 39*). CK2α', but not CK2α, was found elevated

in the hippocampus of symptomatic (10-12 months) PS19 (*38, 40, 41*) mice by both

immunoblotting and RNA in situ hybridization (**Fig. 1J-R**). CK2α' was elevated in all regions of

the hippocampus specially in the CA1 and DG regions (**Fig. 1M-O**), and signal colocalized with

NeuN staining, pointing to enhanced expression in neurons (**Fig. 1P**). Enhanced expression of

CK2α' was also observed outside the neuronal layer in microglia (Iba1+) (**Fig. 1Q, R**). Single

cell RNA-Seq studies available in The Alzheimer's Cell Atlas (TACA) (*42-46*) confirmed

increased *CSNK2a2* expression in patients with AD in both microglia and neurons in the

occipital cortex (OL) and entorhinal cortex (EC) respectively (**Fig. S1**). However, studies in

prefrontal cortex (PFC) and superior frontal gyrus (SFG) within the frontal lobe revealed

enhanced expression of *CSNK2a2* in cells other than neurons and microglia, suggesting a cell-

specific upregulation of *CSNK2a2* that seems to be brain region-dependent. Overall, these

results demonstrate a specific upregulation of CK2α' in brain affected regions of patients with

dementia and in hippocampal neurons and microglia of mouse models of tauopathy.

Genetic depletion of Protein kinase CK2α' impacts tau pathology *in vitro* and *in vivo*

Previous studies have shown that CK2 holoenzyme regulates tau phosphorylation in various cell

and mouse models of AD (*16, 47*), but whether CK2α' subunit specifically contributes to this

phenomenon is unknown. We first explored the impact of silencing CK2α or CK2α' in Neuro2a

(N2a) cells expressing a pathological tau variant (Tau-P301L) in Tau phosphorylation. N2a cells

were treated with either scrambled RNA (ScrRNA) or silencing RNA for CK2α (siCK2α) or CK2α'

(siCK2α') and transfected with either Tau-P301L-EGFP (*48*) or EGFP empty vector (control)

(**Fig. 2A**). The AT8 phospho-tau (pTau) antibody was then used to examine the impact of

silencing CK2α and CK2α' on tau pathology (**Fig. 2B**). First, we confirmed the efficacy of the

siRNAs decreasing the expression of CK2α or CK2α' without altering the other subunits (**Fig.**

2C-F) and the increased expression of Tau in N2A transfected cells (**Fig. 2G**). Importantly,

siCK2α' reduced AT8 levels in Tau-P301L transfected cells (p=0.0130) but not siCK2α

(p=0.9722) (**Fig. 2B, H**). This data demonstrates an impact of the CK2α' subunit specifically on

pTau accumulation *in vitro*.

We then generated a PS19 mouse haploinsufficient for CK2α' (PS19;CK2α'$^{(+/-)}$) (**Fig. S2A-C**) to

assess the impact of depleting CK2α' on tau pathology *in vivo* in both prodromal (~7 months)

and at a late symptomatic age (~10-12 months) (*38*) (**Fig. 3A-B**). In the prodromal age no

significant differences were observed in the accumulation of AT8 in PS19 and PS19;CK2α'$^{(+/-)}$ in

all three examined regions of the hippocampus (CA1, CA3, and DG) and overlaying cortex (**Fig.**

3C-E, Fig. S2D-F). As expected, in the late symptomatic groups we observed a significant

increase in AT8+ area in both the PS19 and PS19;CK2α'$^{(+/-)}$ mice in the CA1, CA3, and DG (**Fig.**

3C-E) and cortex (**Fig. S2D-F)** relative to all other groups. However, PS19;CK2α'$^{(+/-)}$ mice

showed a significant decrease in AT8 signal compared to PS19 in the CA1, DG, and cortex

(CA1; p=0.0075, DG; p=0.0186 and Cortex; p<0.0001) (**Fig. 3C,E, Fig. S2D-F**) suggesting an

overall decrease in pTau upon depletion of CK2α'. Similar results were obtained using the

AT100 phospho-Tau antibody (**Fig. S2G-J**). Decreased pTau in PS19;CK2α'$^{(+/-)}$ was also

confirmed by immunoblotting from whole hippocampal samples at 9 months (**Fig. 3F-I**).

Tau pathology has been previously reported to follow a characteristic progressive pattern of

deposition in the hippocampus that can be categorized into 4 pathology types (*49-52*) (**Fig. 3J**).

These pathology types have been shown to correlate with hippocampal volume (*50, 52*) and

offer a more holistic pathological assessment. We observed differential distributions of patterns

between PS19 and PS19;CK2α'$^{(+/-)}$ mice with a more dramatic alteration in late symptomatic

stages where nearly all PS19 mice presented a type 4 pathology in contrast to the

PS19;CK2α'$^{(+/-)}$ that presented a variation in types with almost 50% of mice presenting type 2

pathology (prodromal p=0.11, and symptomatic p=0.0587) (**Fig. 3K**). The observed modulation

of tau pathology underscored a critical role for CK2α' in the modulation of tau phosphorylation

and associated pathology.

RNA-Seq analyses revealed CK2α' alters the expression of genes related to immune

response and synaptic function

To determine whether the reduction in Tau pathology translates into wider changes in cellular

signaling and gene regulation, we next analyzed RNA-seq profiles from CK2α'-deficient mice

(**Fig. 4**). We focused on the prodromal phase to identify pathways involved in disease onset that

may be missed if only later stages of the disease are studied where significant neuronal loss is

present. Gene expression analyses in the PS19 vs WT mice identified 477 significant DGEs

(**Fig. 4A**). Gene ontology pathway analysis revealed that many of the DGEs belonging to the top

eight dysregulated pathways corresponded to immune and inflammatory functions (**Fig. 4B**), as

previously reported (*53, 54*) . Other important GO pathways were associated with synaptic

dysfunction where the top 5 corresponded to dysregulation in synapse assembly, regulation of

synapse organization, regulation of synapse structure or activity, post synapse organization and

synaptic pruning (**Fig. 4B**). Synaptic pruning was upregulated in PS19 vs WT, whereas the other

synaptic pathways related to organization and regulation of activity were largely downregulated

(**Fig. 4C**).

Direct comparison of PS19 vs PS19;CK2α'$^{(+/-)}$ mice yield a small set of 26 significant DGEs (**Fig.**

4D). Several of these DGEs were related to Apoptotic/phagocytic and Immune functions (*Cept1,*

Ube2n, Oxct1, Exoc3, Adgrb1, Sh3glb2). Most of the DGEs between PS19 and PS19;CK2α'$^{(+/-)}$

were up-regulated in PS19 mice compared to WT but reduced expression in PS19;CK2α'$^{(+/-)}$.

One example Adgrdb1, an adhesion-GPCR, strongly associated with the engulfment and

pruning of synapses (*55, 56*), was significantly downregulated in PS19;CK2α'$^{(+/-)}$ mice (**Fig. 4D**).

We then assessed whether the manipulation of CK2α' had a broader effect on modules of genes

with similar biological functions despite not reaching the strict >Log2 criteria. Weighted Gene

Co-Expression Network Analysis (WGNCA) revealed 34 gene modules with 8 modules (red,

lightcyan, royalblue, skyblue, tan, lightyellow, turquoise, and black) showing statistical

significance detected via a Kruskal-Wallis test among our groups (p<0.05) (**Fig. 4E, File S2**).

Among the 8 significant gene modules the red, tan, and black modules showed an expression

pattern in the PS19;CK2α'$^{(+/-)}$ mice that more closely resembled the expression pattern of WT

mice showing an overall rescue in expression compared to PS19 mice (**Fig. 4F**). We focused

our analyses on the tan module for being the most significant gene module among the three

rescued modules (**Fig. 4E**). The tan module composed on 215 genes (Kruskal-wallis p=0.016)

(**File S2**) was represented by pathways associated with synaptic function and assembly, with

top pathways being vesicle-mediated transport in synapse, synaptic vesicle cycle, and

regulation of synapse organization (**Fig. 4G**). The overall expression of genes associated with

these pathways presented a depletion in the PS19 mice that was ameliorated in the

PS19;CK2α'$^{(+/-)}$ group (**Fig. 4F**). Overall, these results suggested an amelioration in synapse

pruning and an improvement in synaptic function and organization when deleting CK2α' in PS19

mice.

PS19 mice lacking CK2α' ameliorated microglia reactivity, neuroinflammation and

phagocytic activity

pTau is a potent driver of microglia reactivity (*9, 38, 57-60*) which in turn contributes to

exacerbate Tau-mediated pathology (*53, 61-65*). Having established that CK2α' is elevated in

PS19 microglia, and that deletion CK2α' decreased phospho-Tau accumulation and resulted in

the downregulation of genes associated with apoptosis/phagocytosis and synaptic pruning, we

investigated whether these phenotypes were associated with changes in microglia reactivity. We

first examined the number of Iba1+ microglia in the hippocampus of mice as an indicator of

microgliosis (**Fig. 5A, B, Fig. S3A-E**). Iba1+ cells significantly increased in PS19 mice

compared to WT, especially in symptomatic mice. Importantly, PS19;CK2α'$^{(+/-)}$ significantly

reduced the number of Iba1+ microglia in the CA1 and CA3 (**Fig. 5A, B, Fig. S3A-E**).

Microgliosis was also reduced in the cortex of PS19;CK2α'$^{(+/-)}$ mice compared to PS19 (**Fig.**

S3F-H). We also analyzed the number of GFAP+ cells in the hippocampus of PS19 mice in both

age groups (**Fig. S4**), as increased astrogliosis is also reported in PS19 mice (*38*). However, no

significant differences were found between PS19 and PS19;CK2α'$^{(+/-)}$ mice (**Fig. S4**) suggesting

that CK2α' mediated changes in glial cells are specific to microglia.

We then performed morphological assessments of microglia using measurements

corresponding to ramification and correlating with microglia state (**Fig. 5A, C-E**). It has been

previously reported that microglia adopt an amoeboid morphology in PS19 mice, characterized

by the loss of microglia processes, and that is associated with a more reactive and phagocytic

state (*9, 66*). Both the PS19 and PS19;CK2α'$^{(+/-)}$ mice showed a decrease in the average branch

length of microglia in the CA1 (**Fig. 5C**). However, PS19;CK2α'$^{(+/-)}$ mice showed a significant

increase in the number of branches and number of endpoints compared to the PS19 mice (**Fig.**

5D,E) indicating an improvement in the morphological state of microglia.

Since the morphological state of microglia is strongly related to its reactive state and

neuroinflammation we first assessed the expression of a variety of cytokines and other

inflammatory molecules associated with microglia reactivity by using a cytokine array panel with

hippocampal extracts from symptomatic PS19 and PS19;CK2α'$^{(+/-)}$ (**Fig. 5F, G**). Importantly, we

confirmed that several cytokines were significantly decreased ($p<0.05$) in PS19;CK2α'$^{(+/-)}$ mice

relative to PS19 including, MIP-1α, IL17, IL7, IL4, and I309 with several others showing trending

decreases ($0.05<p<0.1$) including TREM1 and TNFα (**Fig. 5F, G**). We also investigated whether

changes in overall cytokine levels and microglia reactivity upon CK2α' depletion related to an

altered phagocytic state. For this we assessed the levels of CD68, a lysosomal/endosomal

marker strongly associated with phagocytic function of microglia and normalized CD68 levels to

the number of Iba1+ cells observed in each animal and time point (**Fig. 6A-E, Fig. S3I**). We

observed increased levels of CD68 in all regions of the hippocampus in the PS19 mice at both

prodromal and symptomatic stages (**Fig. 6C-E**)**.** PS19;CK2α'$^{(+/-)}$ mice also presented increased

CD68 levels compared to WT but displayed significantly less CD68 in the CA1 and a trending

decrease in the CA3 and DG at the prodromal stage compared to PS19 mice (**Fig. 6C-E**).

Importantly, symptomatic PS19;CK2α'$^{(+/-)}$ mice displayed significantly less CD68 in all

hippocampal regions compared to PS19 mice (**Fig. 6C-E**). Expression of another phagocytic

marker Clec7a was also significantly reduced in PS19;CK2α'$^{(+/-)}$ compared to PS19 mice

supporting the impact of CK2α' deletion in the amelioration of microglia phagocytosis (**Fig. 6F-**

H).

Aberrant microglia-mediated synaptic pruning has been extensively reported in PS19 mice and

contributes to synaptic dysfunction and neuronal loss (*53, 61-65*). Considering the changes

observed in CD68 and Clec7a markers of microglia phagocytosis and transcriptional changes

related to apoptosis and synaptic pruning upon deletion of CK2α' we assessed whether

PS19;CK2α'$^{(+/-)}$ microglia decreased synaptic engulfment. To quantify synapse

phagocytosis/engulfment we co-stained for the postsynaptic marker PSD-95 and Iba1 (*67, 68*)

and examined the CA1 stratum radiatum of symptomatic mice (**Fig. 6I**). PS19 mice exhibited a

significant increase in PSD-95+ puncta within Iba1+ cells relative to control mice (**Fig. 6I, J**)**,**

demonstrating increased phagocytosis in PS19 microglia. Importantly, PS19;CK2α'$^{(+/-)}$ showed a

significant reduction in PSD-95 engulfment (**Fig. 6I, J**) indicating an amelioration in the

phagocytic activity of microglia upon deletion of CK2α'.

CK2α' happloinsufficiency improved hippocampal synaptic density and synaptic function in PS19 mice

We then assessed whether the observed changes in synaptic gene expression and synaptic

engulfment by microglia upon deletion of CK2α' were translated into enhanced synaptic density

and neuronal activity in PS19 mice. We first conducted NeuN immunostaining analyses in the

hippocampus of PS19 and PS19;CK2α'$^{(+/-)}$ mice to determine if depletion of CK2α' could impact

the overall neuronal loss characteristic of PS19 mice (*38, 69-71*). Significant depletion of NeuN+

cells and hippocampal atrophy was only seen in PS19 groups at a symptomatic age (**Fig. 7A, B,**

Fig. S5A-C). Importantly, PS19;CK2α'$^{(+/-)}$ mice revealed a significant increase in the number of

NeuN+ cells compared to PS19 in the CA1 (**Fig. 7A, B, Fig. S5A-C**).

We then evaluated if total synapse density was also positively altered in the CA1. We conducted

immunostaining and colocalization analyses for the vesicular glutamate transporter 1 (VGlut1)

(pre-synaptic marker of excitatory synapses), and the postsynaptic marker PSD-95 (**Fig. 7C-I**).

As expected, PS19 mice showed a significant decrease in the number of synapses (colocalized

puncta VGlut1-PSD95) compared to WT mice (**Fig. 7F, I**). The reduction in colocalized puncta in

PS19 mice was due to a concomitant reduction in both PSD-95 and VGlut1 markers (**Fig. 7D, E,**

G, H). In PS19;CK2α'$^{(+/-)}$ mice we observed a decrease in the % loss of both PSD-95 (**Fig. 7D,**

G) and VGlut1 (**Fig. 7E, H**). This resulted in the amelioration of total synapse loss in

PS19;CK2α'$^{(+/-)}$ mice and obtained values for colocalization were no longer significant respect to

WT (**Fig. 7F, I**). These results demonstrate a partial rescue in the density of excitatory synapses

in the CA1 of PS19;CK2α'$^{(+/-)}$ mice aligning with the amelioration of hippocampal atrophy

observed in this region (**Fig. 7B**) and the reduced synapse engulfment mediated by microglia

(**Fig. 6I, J**). In line with these results, silencing CK2α' in primary cortical neuron cultures

transfected with Tau-P301L-EGFP increased spine density compared to scramble (**Fig. S6**).

To investigate functional implications of the rescue seen in PS19;CK2α'$^{(+/-)}$ hippocampal atrophy

and synaptic density we assessed hippocampal synaptic plasticity by performing extracellular

field excitatory postsynaptic potential (fEPSP) recordings in the CA1 region of acute

hippocampal slices (**Fig. 7J, K**). Theta-burst stimulation (TBS) at the Schaffer collaterals

induced short-term potentiation (STP), which resulted in a significant increase in synaptic

efficacy at CA1 synapses in both WT and CK2α'$^{(+/-)}$ mice, as evidenced by the enhanced fEPSP

amplitude (170 ±13%, n = 6; 170 ± 12%, n = 7) (**Fig. 7L, M**). In contrast, the increase in fEPSP

amplitude was significantly reduced in PS19 and PS19;CK2α'$^{(+/-)}$ mice relative to WT controls

post TBS (123 ± 4%, n = 7; 128 ± 7%,n = 6; **Fig. 7L, M**), indicating impaired STP in these

models. Furthermore, we examined long-term potentiation (LTP) at CA3-CA1 synapses, a form

of synaptic plasticity linked to learning and memory. It has been previously shown that PS19

mice present impaired LTP in the CA1-CA3 pathway (*38, 72, 73*). First, we observed significant

differences in input-output curves at CA3-CA1 synapses between PS19 and PS19;CK2α'$^{(+/-)}$

mice when compared to WT, which suggests that excitatory synaptic input properties were

altered in these models (**Fig. 7N**). TBS at the Schaffer collaterals induced robust LTP in both

WT (173 ± 12%, n = 6) and CK2α'$^{(+/-)}$ (164 ± 9%, n = 7; **Fig. 7L, O**). However, in the PS19

mouse model, the magnitude of LTP was significantly diminished compared to WT (128 ± 9%, n

= 7; vs. WT p=0.007) (**Fig. 7L, O**), suggesting a disruption in the molecular mechanisms

underlying LTP. Importantly, in the PS19;CK2α'$^{(+/-)}$ mice, LTP was partially restored (150 ± 6%, n

= 6; vs. WT p=0.199) (**Fig. 7L, O**), pointing to a potential compensatory mechanism that partially

mitigates the LTP deficits observed in the PS19 mice. Importantly, paired-pulse facilitation (PPF)

was unaltered across all experimental groups, suggesting that the observed changes in synaptic

plasticity were most likely due to postsynaptic mechanisms dependent on the hippocampus

(**Fig. S7**).

CK2α' haploinsufficiency improved PS19 behavior on Barnes Maze

Finally, we assessed whether the overall amelioration in Tau-pathology, microgliosis and synaptic dysfunction mediated by CK2α' had an impact in cognitive behaviors. Mice were trained on the Barnes maze task for 5 consecutive days on the location of a target hole (training) and on the 6th day a probe trial was conducted with the target hole removed (**Fig. 8A**). During the training trials the primary latency or the time to first locate the escape hole was recorded (**Fig. 8B-C**). All mice in the prodromal group demonstrated an improvement on time needed to locate the escape hole from day 1 vs day 5 (WT p=0.0272; PS19 p=0.0.0067; CK2α'$^{(+/-)}$ p=0.0002; PS19;CK2α'$^{(+/-)}$ p<0.0001) with no significant difference between the genotypes (**Fig. 8B**). All mice in the symptomatic group also showed a capability of acquiring the task demonstrating a significant improvement from day 1 vs day 5 (WT p<0.0001; PS19 p=0.0109; CK2α'$^{(+/-)}$ p<0.0001; PS19;CK2α'$^{(+/-)}$ p=0.0037). However, PS19 and PS19;CK2α'$^{(+/-)}$ showed a significant impairment compared to WT, starting on day 3 for PS19 (p=0.0009) and day 4 for PS19;CK2α'$^{(+/-)}$ (p=0.0306) (**Fig. 8C**). The training data suggested that both symptomatic PS19 and PS19;CK2α'$^{(+/-)}$ mice showed similar impaired learning on the Barnes Maze compared to WT mice, although deficits in PS19;CK2α'$^{(+/-)}$ were attenuated.

Following the training phase of the Barnes maze we conducted a probe phase in which the escape hole was covered and the time spent exploring the escape hole quadrant was recorded. At the prodromal age there was no significant difference in time spent in the escape hole quadrant for any genotype (**Fig. 8D**). At the symptomatic age, the PS19 mice showed a significant impairment in the time spent in the goal quadrant compared to WT (p<0.0001) (**Fig. 8D**). Importantly, the PS19;CK2α'$^{(+/-)}$ mice spent significantly more time than the PS19 mice in the goal quadrant with no significant difference from WT (vs. PS19 p=0.0113 vs. WT p=0.2335) demonstrating a rescue in spatial memory for the PS19;CK2α'$^{(+/-)}$ on the probe phase of the Barnes maze (**Fig. 8D**).

The PS19 mouse model is known to present motor deficits due to hindlimb paralysis and

weakness (*38*), which could confound Barnes Maze results if PS19 mice perform more poorly

due to less distance traveled. Prodromal mice displayed no difference in distance traveled on

Barnes Maze (**Fig. S8A**). While the symptomatic PS19 mice traveled less distance in the

Barnes Maze compared to WT mice, they did not show a significant difference in distance

traveled compared to PS19;CK2$\alpha'^{(+/-)}$ mice (**Fig. S8B**). In addition, in open-field testing we found

no differences between groups at the prodromal time point (**Fig. S8C**). On the contrary,

symptomatic mice showed significant hyperactivity in the open field task, demonstrating the

ability of PS19 mice to move (**Fig. S8D**). Indeed, no significant differences were observed in the

open field between PS19 and PS19;CK2$\alpha'^{(+/-)}$ mice (**Fig. S8D**). This data supports the

improvement in cognitive abilities seen in the PS19;CK2$\alpha'^{(+/-)}$ mice are not due to changes in

motor function but rather are related to an improvement in memory.

Mice can use different cognitive strategies to navigate the Barnes Maze and locate the escape

hole (*74*). As the mouse learns the task they show improvement in the strategy used to locate

the escape hole (*74*) (**Fig. 8E**). Assessing these cognitive strategies is important to provide a

deeper insight into how memory and spatial learning are being utilized. All animals in the

prodromal group demonstrated a similar pattern in spatial strategies over the training phase

(**Fig. 8F**) and showed a predominance in the corrected and direct strategies in the probe phase

(**Fig. 8G**). Symptomatic PS19 and PS19;CK2$\alpha'^{(+/-)}$ mice both showed impaired spatial strategies

over the training phase, by day 5 both genotypes largely displayed either failing, random or

serial strategies (**Fig. 8H**). However, during the shortened probe phase, PS19 displayed a

worsened overall strategy, with predominant failing, serial, and long correction strategies

compared to PS19;CK2$\alpha'^{(+/-)}$ mice that displayed more serial, corrected and direct strategies

(**Fig. 8I**). Overall, the PS19;CK2$\alpha'^{(+/-)}$ mice displayed improvement in the probe phase of the

Barnes Maze in both the time spent in the escape quadrant and strategy used to locate the

target, indicating improvements in spatial memory.

Discussion

Tauopathies are a heterogeneous group of neurodegenerative disorders for which effective

treatments are still not available, in part due to an incomplete understanding of the underlying

mechanisms. In this study, we present a novel target, CK2α', a catalytic subunit of CK2, as a

potential upstream regulator of tau mediated neurodegeneration, acting at the intersection of

neuroimmune signaling and synaptic dysfunction.

Here, we report that CK2α' expression is abnormally elevated in the brains of dementia patients

and in the hippocampus of PS19 mice, with preferential upregulation observed in hippocampal

neurons and microglia. These findings are supported by multiple single-cell RNA-sequencing

(snRNA-seq) studies in human AD, which consistently show increased CK2α' expression in

excitatory neurons (*44-46*). Additionally, the study by Gerrits et al. (*43*) in the occipital lobe (OL)

has reported preferential CK2α' expression in AD-associated microglia. Other scRNA-seq

studies have detected elevated CK2α' expression in various brain cell types, with patterns

varying by dataset and brain region (*44-46*). On the other hand, CK2α seems to be increased in

cortical astrocytes of the 5x FAD and PS19 models (*75*). This collective evidence suggests that

CK2α' dysregulation in AD/ADRD may be both cell-type- and region-specific. Consistent with

our findings, a concurrently submitted study has reported that CK2α' is elevated in AD models

and contributes to APP-regulated glial inflammation (*76*), independently supporting and

strengthening our conclusions.

Increased CK2 levels have been associated with tau pathology (*16, 17, 47, 77*), microglia state

(*78, 79*) and general inflammation (*80-86*). However, many of these studies were conducted in

cell models (*16, 17, 34, 47*) and genetic evidence for the specific contributions of the individual

CK2 subunits to tau pathology remained largely unexplored. Our studies *in vitro* using N2A cells

transfected with Tau-P301L showed that silencing CK2α', but not CK2α, decreased pTau

accumulation. Furthermore, our studies *in vivo* reducing the levels of CK2α' in the PS19 model

also influenced tau pathology by decreasing the accumulation of pTau and tau burden in the

hippocampus and cortex of PS19;CK2α'$^{(+/-)}$ mice. In support to our data, previous studies *in vitro*

showed that overexpression of CK2α' subunit decreased the activity of I2PP2A/SET, a cellular

protein that functions as an endogenous inhibitor of protein phosphatase 2A (PP2A), increasing

pTau. In contrast overexpression of CK2α increased the activity of I2PPA2a/SET (*77*). Although

early studies demonstrated that Tau is a physiological substrate of CK2, at least in specific

contexts such as neurogenesis (*87*), the findings by Perez et al. (*77*) suggest that the effects we

observed after manipulating CK2α' on pTau accumulation could be mediated through

intermediary molecules that modulate Tau phosphorylation.

Haploinsufficiency of CK2α' also increased the expression of synaptic genes, synaptic density,

and synaptic function in the PS19 mice. Along these lines, haploinsufficiency of CK2α' in a

mouse model of HD also showed improved synaptic density and function (*15, 88*), although how

exactly CK2α' influences these processes is still unknown. A previous study showed that CK2α'

can influence the expression of synaptic genes, especially those related to glutamatergic

excitatory signaling (*15*). A more recent study showed that pharmacological inhibition of CK2

mitigated AD tau pathology by preventing the NMDA receptor subunit NR2B synaptic

mislocalization (*47*). NR2B mediates long-term depression (LTD), and LTD specifically induces

the phosphorylation of tau. These studies support our findings and strengthen the relationship

between CK2α', tau phosphorylation, and the regulation of neuronal function in both AD and

other neurodegenerative diseases. However, since CK2α' is found elevated in both neurons and

microglia, it is still unknown whether the effects mediated by CK2α' in pTau accumulation and

synaptic function are cell or non-cell autonomous, warranting further investigation.

CK2 has been identified as a mediator of microglia reactivity and inflammation in different

models (*78, 79*) and it has been well-established as a mediator of inflammation in a variety of

contexts including SARS-CoV infection, cancers, bacterial infections, intestinal inflammation and

renal failure (*80-85*). This is largely mediated via CK2's involvement in regulating several large

inflammatory factors such as NF-κB, STAT1, and EGR-1 (*86*). It is known that chronic activation

of NF-kB increases the accumulation of pTau (*9*). On the other hand, pathological tau has also

been shown to induce inflammation and NF-kB activation by interacting directly with

inflammatory receptors such as toll-like receptors 2 (TLR2) (*10, 11*) creating a vicious cycle of

worsening inflammation and tau pathology (*9-12*). Our data demonstrated that CK2α'

haploinsufficiency significantly reduced the number of Iba1+ cells in the hippocampus, reduced

the levels of the microglia reactive and phagocytic markers CD68 and Clec7a, ameliorated

microglia morphological changes associated with pathology, reduced the levels of pro-

inflammatory cytokines in PS19 mice and reduced microglia-mediated synaptic engulfment. In

support to our findings, previous studies in human microglia derived from hiPSCs treated with

CK2 inhibitors also reduced cytokines production (*79*). Therefore, all together, these data

support a specific key role of CK2α' in mediating neuroinflammation via activation of microglia.

Loss of synaptic density and neuronal dysfunction in AD models has been previously connected

to increased phagocytic microglia (*39, 63, 89, 90*). Our results demonstrated a decrease in

microglia phagocytosis of the PSD-95 synaptic marker in PS19;CK2α'$^{(+/-)}$ mice suggesting that

CK2α' is influencing the phagocytosis of synapses by microglia, but the mechanisms behind this

are yet to be elucidated. CK2α' could facilitate these processes by regulating the transcription of

genes involved in phagocytosis and synaptic engulfment and/or by phosphorylating and

activating specific proteins related to these processes. Deletion of CK2α' in PS19 mice resulted

in the down-regulation of several genes related to apoptosis/phagocytosis and synapse

engulfment such as *Cept1, Ube2n, Oxct1, Exoc3, Adgrb1* and *Sh3glb2 (55, 56, 91-95)* which

can contribute to the phenotypes observed in PS19;CK2α'$^{(+/-)}$ mice. It is also possible that CK2α'

phosphorylates elements of the complement component such as C9 and C1r, known CK2

substrates (*96, 97*) and influence microglia-mediated synaptic pruning.

In conclusion, our findings identified CK2α' as a central and previously underappreciated

regulator of tauopathy pathogenesis, acting at the crossroads of neuroimmune signaling,

synaptic function, and tau phosphorylation. By demonstrating that CK2α' haploinsufficiency

alleviates tau pathology, reduces neuroinflammation, and improves synaptic integrity and

function in PS19 mice, we provide compelling genetic evidence that CK2α' contributes to

multiple pathological hallmarks of tau-driven neurodegeneration. Importantly, these effects

appear to occur through complex, context-dependent mechanisms that may differ across cell

types and disease stages. The observation that CK2α' influences both neuronal and microglial

compartments underscore the need to dissect its cell-autonomous versus non-cell-autonomous

roles. Given the growing interest in CK2 as a therapeutic target and the limitations of current

inhibitors that lack subunit specificity, the development of CK2α'-selective modulators (*76, 98*)

may represent a promising avenue for disease-modifying therapies in tauopathies and related

neurodegenerative conditions. Future studies elucidating the molecular and cellular pathways

downstream of CK2α' will be critical for translating these findings into clinical applications.

Materials and Methods

Study design

In this study we report the upregulation of CK2α' in AD human samples and mouse models and

investigated the role of CK2α' in tauopathy. We hypothesized that CK2α' contributes to tau-

associated pathology and that reducing its levels would improve outcomes in tauopathy models.

Data was collected from AD/ADRD post-mortem human tissue and AD mouse and cell models.

Methods used in this study include western blotting, immunostaining (tissue and cells), bulk

RNA-Seq, electrophysiology and behavioral assays. Publicly available human genomic datasets

were also analyzed, as described.

For mouse experiments, animals were assigned to cohorts at birth based on genotype and aged

to predetermined time points; no additional randomization was performed. Sample sizes for all

experiments were chosen based on previous literature and our prior experience with similar

experiments; no formal power calculations were performed, exact sample sizes are detailed in

figure legends. Equal numbers of male and female mice were used. Data is presented as

mouse averages unless otherwise detailed in the figure legend.

Experimenters were not blinded during behavioral testing, however all image analysis was

performed on blinded samples to reduce experimenter bias.

Cell culture: Culturing, transfections, and immunocytochemistry

Neuro-2a cells (N2a; ATCC CCL-131) were cultured in DMEM with 10% FBS and

penicillin/streptomycin at 37 °C and 5% CO_2. For silencing experiments, cells were plated in 12-

well plates or Matrigel-coated coverslips. Cells were transfected with siRNA targeting CK2α or

CK2α′ (Qiagen) or a negative control using DharmaFECT 1 (25 nM). After 24 h, cells were

transfected with pEGFP-C1 or Tau-P301L-EGFP using jetOptimus and processed 24 h later. For

immunocytochemistry, cells were fixed, blocked in 5% NGS/TBST, incubated with AT8 primary

antibody overnight, followed by Alexa-fluor secondary antibodies, and mounted with ProLong

Gold for imaging (see Supplementary Methods for primary neuron culturing details;(*99*)).

Mouse Lines

For this study we used the B6;C3-Tg(Prnp-MPAT-P301S)PS19Vle/J (or PS19) (*38*) mouse line

(Jackson #008169 maintained with purchased B6C3H breeders; obtained from Dr. Karen Ashe

at University of Minnesota), and CK2α'$^{(+/-)}$ mice, originally obtained from Dr. Seldin at Boston

University (Taconic biosciences TF3062) (*31*) and maintained on the C57Bl/6J background for

several generations. We crossed PS19 and CK2α'$^{(+/-)}$ mice and this cross generated 4

genotypes of interest WT (PS19$^{0/0}$;CK2α'$^{(+/+)}$), CK2α'$^{(+/-)}$ (PS19$^{0/0}$;CK2α'$^{(+/-)}$), PS19

(PS19$^{Tg/0}$;CK2α'$^{(+/+)}$), and PS19;CK2α'$^{(+/-)}$ (PS19$^{Tg/0}$;CK2α'$^{(+/-)}$) mice. For all experiments

littermate WT and CK2α'$^{(+/-)}$ controls were used. Two time points were used in experiments; pre-

symptomatic (6-7 months) and symptomatic (10-12 months). All mice were housed under

standard SPF conditions. All animal care and sacrifice procedures were approved by the

University of Minnesota Institutional Animal Care and Use Committee (IACUC) in compliance

with the National Institutes of Health guidelines for the care and use of laboratory animals under

the approved animal protocol 2307-A41243.

Human postmortem tissue

Postmortem human frontal cortex (Brodmann's area 46) FTD patients and controls subjects was

provided by NeuroBioBank of National Institutes of Health. A total of 6 female (3 control/3FTD)

and 8 male (4 control/4 FTD) samples were used, and samples were age and sex matched.

Ages ranged from 60-77 with the average age of all samples being 70. More detailed

information on specific samples can be found in **file S1**.

Immunoblotting

Human and mouse protein was extracted as previously described (*14*). Tissue was

homogenized in 25 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Triton

X-100, incubated on ice 15-30 min, then SDS was added to 2% and samples were heated at

100°C for 5 min. Lysates were centrifuged at 4°C, 13,000 g for 20 min and supernatants

collected. Protein concentration determined by BCA assay (Pierce).

Protein samples were separated on 4–20% SDS Criterion TGX Stain-Free gels (BioRad) and

transferred to a 0.2 μm nitrocellulose membrane in Tris–Glycine Buffer (25 nM Tris-Base,

200 mM Glycine) at 25 V for 30 min (Trans Turbo Transfer system). Membranes were

blocked in 5% milk in TBST (0.5% Tween-20) for 1 hour , incubated overnight at 4 °C in

primary antibody diluted in 2.5% milk in TBST, then incubated in HRP-conjugated secondary

antibody (Amersham 1:5000) for 1 hour. Bands were visualized with SuperSignal

Chemiluminiscent substrate (Thermo Scientific) using a GE ImageQuant Las4000 imager.

Primary antibodies: CK2α' (1:2000; Rabbit, Novus NB100-379), CK2α (1:1000; Rabbit, Abcam

ab76040), Clec7a (1:500; Rat invivoGen mabg-mdect), At8 (1:1000; Mouse, Invitrogen

MN1020), At100 (1:1000; Mouse, Invitrogen MN1060), Tau-5 (1:1000; Mouse, Abcam ab80579),

and GAPDH (1:5000; Mouse, Santa cruz sc-365062).(See supplementary Methods for cytokine

immunoblotting details)

Immunohistochemistry

Immunohistochemistry experiments were conducted as previously described (*14, 100, 101*)

using fluorescent or 3,3'-diaminobenzidine (DAB) detection. Mice were anesthetized (Avertin,

250 mg/kg) and perfused intracardially with TBS (25 mM Tris-base, 135 mM Nacl, 3 mM KCl,

pH 7.6) supplemented with 7.5 mM heparin. Brains were fixed with 4% PFA at 4°C for 4–5 days,

cryoprotected with 30% sucrose for 4–5 days and embedded in a 2:1 mixture of 30% sucrose in

TBS:OCT (Tissue-Tek). Coronal sections (16 μm) were stored in a 50% glycerol-50% at -20ºC.

Three hippocampal sections were analyzed per mouse.

For immunofluorescent staining, sections were blocked in 5% normal goat serum (NGS) in

TBST for 1 h at room temperature. Primary antibodies were incubated overnight at 4°C in TBST

containing 5% NGS. Secondary Alexa-fluorophore-conjugated antibodies (Invitrogen) were

added (1:200 in TBST with 5% NGS) for 1 h at room temperature. Slides were mounted in

ProLong Gold Antifade with DAPI (Invitrogen) and subsequently imaged. Primary antibodies

used and dilutions are as follows: GFAP (1:2000; Chicken Millipore Sigma AB5541), Iba1

(1:500; Rabbit Fujifilm Wako 019-19741), Iba1 (1:500, Goat Fujifilm Wako 011-27991), NeuN

(1:1000; Mouse Millipore MAB377), Clec7a (1:1000;Rat inviogen mabg-mdect), PSD95 (1:500;

Rabbit Invitrogen 51-6900) and Vglut1 (1:500; Guinea pig Millipore AB5905). Imaging of

immunofluorescent slices was conducted using a confocal microscope (Stellaris, Lecia). For

NeuN, Iba1, and GFAP imaging 20x tiles were acquired with a 16um z-dimension and 1um step

size. For PSD95 and VGlut1 images were acquired as described for synapse analysis.

For DAB staining antigen retrieval was performed using either Tris-EDTA pH=9.0 (Biolegend

422703) or Rodent decloaker (Biocare medical RD913) at 80ºC for 30 minutes. Sections were

blocked at room temperature in 10% NGS/TBST for 1 hour, then incubated in primary antibodies

overnight at 4C in 5%NGS/TBST. Sections were incubated in biotin-conjugated secondaries

(Jackson Immuno Research Labs Biotin-SP (long spacer) AffiniPure) in 5%NGS/TBST for 1

hour, followed by 3% hydrogen peroxide for 20 minutes. Sections were incubated in tertiary

antibodies (VECTASTAIN Elite ABC, HRPS kit, Vector laboratories) in 5%NGS/TBST for 1 hour

per manufacturer instructions. DAB chromogen diluted in DAB substrate buffer (Biolegend) was

applied for sufficient staining. Sections were counterstained with 0.1% cresly violet, dehydrated,

cleared in ethanol/xylene and mounted in Permount mounting medium (Fisher scientific).

Primary antibodies used and dilutions are as follows: pTau (Ser202, Thr205) AT8 (1:1000;

Mouse, Invitrogen Mn1020), pTau (Ser214, Thr212) AT100 (1:5000; Mouse Invitrogen MN1060)

and CD68 [RM1031] (1:2500; Rabbit, Abcam ab303565). DAB images were captured at 10x

(Echo Revolve) and at 40x (EasyScan One, Motic)(See Supplementary Methods for Tau

pathology type assessment;(*49-52*)). All analyses were conducted in either Qupath (*102, 103*) or

ImageJ (FIJI) (*104*) (See Supplementary Methods for Image Analysis details, including

microglia morphology analysis (*105, 106*))

In situ hybridization

RNAscope was conducted using a custom Csnk2a2 probe (ACDbio protocol). Slides were then

blocked and incubated overnight with anti-IBA1. The next day, Alexa-fluorophore secondary

antibodies were applied, and slides were mounted with ProLong Gold Antifade containing DAPI

for imaging.(See Supplementary Methods for details).

Synapse analysis

Synapse analysis was conducted as previously described (*14, 88, 101, 107, 108*).

Immunohistochemical staining was conducted as described above with Vglut1 and PSD95, with

blocking increased to 20% NGS+TBST for 2 hours and primary and secondary antibodies

diluted in 10% NGS+TBST. Fluorescent images from the molecular layer of the CA1 were taken

on a confocal microscope (Stellaris, Lecia) at 63x with z-step dimension of 0.34 μm with 15

steps were generated. Lecia stellaris software lightning processing was applied to images post-

acquisition and edited images were used for analysis. Maximum projections of three sections

per slice were generated. Puncta analyses were conducted blinded using the PunctaAnalyzer

Plugin (Durham,NC,USA) in FIJI as previously described (*14, 88, 101, 107, 108*). Data is

represented as slice averages with three slices per animal.

Electrophysiology

Mice were anesthetized with isoflurane (2%) and decapitated for slice preparation. Brains were

rapidly removed into (sucrose 189 mM, glucose 10 mM, NaHCO3 26 mM, KCl 3 mM, MgSO4 5

mM, CaCl2 0.1 mM, NaH2PO4 1.25 mM). Transverse hippocampal slices (350 μm) were cut on

a vibratome (LEICA VT1000S) and incubated at room temperature in oxygenated artificial

cerebrospinal fluid (aCSF; NaCl 124 mM, KCl 5 mM, NaH2PO4 1.25 mM, MgSO4 2 mM,

NaHCO3 26 mM, CaCl2 2 mM, glucose 10 mM; gassed with 95% O2, 5% CO2; pH=7.4) for 45

min-1 h. Experiments were performed at 30–34°C with continuous perfusion.

Field excitatory postsynaptic potentials (fEPSPs) were recorded in the CA1 region of the

hippocampus. fEPSPs in were evoked with a stimulating electrode placed on the Schaffer

collateral (0.33 Hz) using brief current pulses (200 µs, 0.1–0.2 mA). Extracellular recording

electrodes were filled with aCSF. Stimulation was adjusted to obtain a fEPSP peak amplitude of

approximately 1 mV during control conditions. After a stable fEPSP baseline period of 10 min,

LTP was induced by a Theta-burst stimulation (TBS) consisting of a series of 10 bursts of 5

stimuli (100 Hz within the burst, 200 ms interburst interval), which was repeated 4 times (5 s

apart). Data were filtered at 3 kHz and acquired at 10 kHz using pCLAMP 10.2 software

(Molecular Devices, RRID: SCR_011323).

A stimulus-response curve (0.05-0.4 mV, mean of five fEPSPs at each stimulation strength) was

compiled for the different mice used. For paired-pulse ratio experiments, two fEPSPs were

evoked 40 ms apart for 0.5 min at baseline frequency (6 times) at the beginning of the baseline

recording. The PPR was expressed as the amplitude of the second fEPSP divided by the

amplitude of the first fEPSP.

Behavior

Behavioral testing was performed with the support from the Universtiy of Minnesota Mouse

Behavior Core. Mice were handled daily for $\geq$ 1 week and habituated to testing room $\geq$ 30 min

before tasks. All tasks were recorded using ANYmaze software (Stoelting Co., Wood Dale,

Illinois) and overhead cameras.

Barnes maze was conducted on a 20 hole Barnes Maze (San Diego Instruments) under 450-

500lux illumination with black and white visual cues around the room, as described (*74, 109,*

110). Mice were placed in the center under a cover with lights off, lights were turned on and the

cover removed within 30 seconds. Training consisted of four 3-minute trials per mouse for 5

days, with 20-30 minutes intertrial intervals. The maze was cleaned with 70% ethanol between

mice and rotated between trials. Primary latency, or the time to first explore the escape hole,

was recorded. Trials ended when mice entered the escape box or after 3 minutes, at which point

the mouse was gently guided to the escape box. During the probe trial, the escape hole was

covered and time spent in each maze zone was recorded. Spatial strategy was determined by a

blinded investigator.

Open field was conducted in 40cmx40cm boxes under 150-200lux illumination. Mice were

placed in boxes and behavior was recorded for 1 hour. Boxes were thoroughly cleaned between

mice with 70% ethanol.

RNA-Seq Analysis

RNA-Sequencing was conducted by Novogene (Sacramento, CA). Gene expression analysis

was carried out using the CHURP pipeline (https://doi-

org.ezp2.lib.umn.edu/10.1145/3332186.3333156) using n=4-7 mice/genotype with

female(F)/male(M) ratios as follows; 4 WT (2F/2M), 5 CK2$\alpha^{'(+/-)}$ (2F/3M), PS19 (3F/4M) (3 high

pathology and 4 low pathology), and 4 PS19;CK2$\alpha^{'(+/-)}$ (3F/1M). Differential gene expression was

determined with DESeq2 using default setting (*111*) (v1.46.0). Genes with a FDR <=0.05 were

considered significant. Outliers' identification was performed using Cook's distance (DESeq2).

Driver factors of gene expression variance (genotype and/or sex) were evaluated using R

(v4.4.2) package variancePartition (1.36.3). Pathway and clustering analysis were completed

gProfiler2 (*112*) (v0.2.3) and clusterProfiler (v4.14.6). Data visualization was done using various

R graphic packages, including ggplot2, ggraph, and DESeq2 visualization functions. The RNA-

seq data set generated in this manuscript has been deposited at GEO (accession number

GSE298505). The reviewer token to access the GEO deposited data is **mlwxmuactlibryl**.(See

Supplementary methods for RNA extraction and qPCR details).

Statistical analyses and data representation

For electrophysiology experiments data were analyzed using Clampfit 10.2 software (pCLAMP,

Molecular Devices, RRID: SCR_011323). Data are presented as mean ± SEM. To estimate

changes in synaptic efficacy, STP was quantified by comparing the mean fEPSP amplitude

during 1 minute after the application of the TBS. LTP was quantified as for STP but 40 minutes

after the protocol. Graphs were obtained using SigmaPlot 14.0. Before applying any statistical

comparison, the data were subjected to Shapiro- Wilk normality and equal variance tests. For

any comparisons between two groups, two- paired Student's t-test was used. For multiple

comparisons to the same control, One- way ANOVA and Holm-Sidak test was used. P-values

less than 0.05 were considered statistically significant.

For all other experiments GraphPad Prism (GraphPad, San Diego, CA, USA) was used to

create graphs and conduct statical testing. For all other experiments data is represented as

mean $\pm$ SEM. Statistical analyses were performed using paired Student's t-test, unpaired

student's t-test, one-way or two-way ANOVA with post-hoc analysis, as indicated in each figure

legend. A p-value <0.05 was considered significant.

List of supplemental materials

Supplemental materials and methods

Fig. S1 to S8

Data File S1-S2

Funding

This work was supported by R01NS110694 (Alzheimer's disease supplement) (RGP) and R03

AG087281 (RGP), R01AG077743 (MLK) and R01NS108686 (MLK), R01MH119355 (AA)

and R01DA048822 (AA) and Department of Defense (W911NF2110328) (AA).

Authors contributions

RGP conceived and directed the project, helped with data analysis and interpretation, figures

preparation and manuscript writing. AW and RT conducted in vitro experiments using N2A cells.

AW conducted immunoblotting with human samples, generated all mice used in the study,

conducted immunostaining for pTau, CD68, and cytokine analyses, performed all behavioral

studies, and contributed to preparing all figures and writing the manuscript. PK helped with

behavioral data analysis. PG conducted RNA in situ hybridization for CK2a' and co-staining with

NeuN and Iba1, performed microglia morphology analyses, synapse engulfment by microglia,

and GFAP and Iba1 staining. PGU assisted with Clec7a immunoblotting. NBR, PG and AF

conducted immunostaining for PSD95/VGlut1 and synapse density analyses. SV and RS helped

validate siCK2a' in neurons. JM helped with execution of DAB immunostaining. MKL contributed

to experimental supervision and data interpretation. AW and YZ conducted RNAseq

experiments and data analyses. RFM conducted electrophysiological studies in the CA1 and AA

helped with experimental supervision and data interpretation of such data.

Competing interests

The authors declare no competing interests

Data and materials availability

RNA-Seq data generated in this study is available in the Gene Expression Omnibus repository

found at https://www.ncbi.nlm.nih.gov/geo/ accession number GSE298505. The

reviewer token to access the GEO deposited data is **mlwxmuactlibryl**.

Allan Brain Atlas Aging, Dementia and TBI dataset that supported conclusions of this article is

publicly available at https://aging.brain-map.org/overview/home (*37*).

Human brain tissue was received from the NIH NeuroBioBank repository and is available

publicly for request at https://neurobiobank.nih.gov/.

scRNA-Seq study data from human samples that contributed to conclusions of this article can

be found in The Alzheimer's Cell Atlas (TACA) https://taca.lerner.ccf.org/ (*42-46*).

Figure legends

Figure 1. CK2α' is increased in tissues of patients with AD/ADRD and in PS19 mice. **(A, B)**

Representative diagram of CK2 holoenzyme, including regulatory and catalytic subunits and

differential distribution of expression of catalytic subunits of CK2α and CK2α' in the human body.

A and B were created in BioRender (White, A. (2025) eqznm6m https://BioRender.com/). **(C)**

Averaged expression of CSNK2A2 in non-cognitively impaired (NCI) and dementia samples

from the Allen Brain Institute: Aging, Dementia and TBI study in the parietal cortex (PCx),

Temporal cortex (TCx), and Hippocampus (HIP) are shown (patients with TBI were excluded

from the analyses). **(D, E)** Quantification of the expression of CSNK2A2 and CSNK2A1 in PCx

from NCI and dementia from the Allen Brain Institute: Aging, Dementia and TBI study (n=20-

27/group). Significance determined by unpaired t-test, $*p<0.05$. **(F)** Immunoblot for CK2α' and

CK2α in protein lysates from frontal cortex of age and sex matched healthy controls (C) and

patients with FTD (left panel). Arrow indicates CK2α' specific band, asterisk indicates unspecific

band. (Right panel) Immunoblot for CK2α' in WT and CK2α' [(-/-)] mice showing antibody-band

specificity. **(G, H)** CK2α' (G) and CK2α (H) protein levels were normalized to GAPDH and shown

relative to age/sex matched control (n=7/group). Statistical analyses were conducted using

unpaired t-test with welch's correction p=0.0377. Data are shown as mean ± SEM. (**I**)

Relationship between CK2α' levels from (**F**) and the presence of Tau pathology reported by the

NIH biobank. Horizontal line represents the group CK2α' intensity average normalized by

GAPDH. (**J**) Immunoblot for CK2α' and CK2α in the hippocampus of symptomatic WT and PS19

mice. (**K, L**) CK2α' (K) and CK2α (L) protein levels were analyzed and normalized to GAPDH

(n=3 mice/genotype). **(M)** Csnk2a2 in situ hybridization images from symptomatic WT and PS19

hippocampus. Panels show 10x images of hippocampal cell layers with Csnk2a2 RNA probe

and DAPI, scale bar=50 μm. (**N, O**) Quantification of Csnk2a2 puncta in the CA1 (p=0.0163)

(**N**), and DG (p=0.0003) (**O**) quantified within the cell layer identified by DAPI, (n=3-4

mice/genotype). **(P)** Csnk2a2 in situ hybridization and NeuN immuno-fluorescence images from

symptomatic WT and PS19 CA1. Scale bar= 20 μm. **(Q)** Csnk2a2 RNA in situ hybridization and

Iba1 immunofluorescence in symptomatic WT and PS19 hippocampus. Top: representative

image of single Iba1+ cell in the Stratum Radiatum (CA1). Scale bar is 20 μm. Bottom: 3D

rendering of cells Iba1+ cells containing Csnk2a2 puncta showing puncta reside within the

microglia cell. **(R)** Quantification of the number of Csnk2a2+ puncta/Iba1+ soma (p<0.0001,n=3-

4 mice/group, 8-9 cells/mouse, every data point is a cell). All Data shown are mean ± SEM.

Statistical analyses were conducted using unpaired t-test and displayed with compact letter

display.

Figure 2. Silencing CK2α' reduces pTau levels in N2a cells expressing Tau-P301L. (A)

Schematic of experimental design for N2a cell transfection and immunostaining. **(B)**

Representative images of AT8 immunostaining and EGFP fluorescence in N2a cells transfected

with either EGFP or Tau-P301L-EGFP and treated with ScrRNA, siCK2α, or siCK2α'. Scale bar=

100 μm. **(C-E)** RT-qPCR analysis of mRNA expression levels of CK2α' **(C)** (Scr-EGFP vs.

siCK2α'-EGFP p=0.0003, Scr-EGFP vs. siCK2α'-Tau-P301L p=0.0003, Scr-Tau-P301L vs.

siCK2α'-EGFP p=0.0002, Scr-Tau-P301L vs. siCK2α'-Tau-P301L p=0.0002, siCK2α-EGFP vs.

siCK2α'-EGFP p=0.0049, siCK2α-EGFP vs. siCK2α'-Tau-P301L p=0.0056, siCk2α-Tau-P301L

vs. siCK2α'-EGFP p=0.0059, siCk2α-Tau-P301L vs. siCK2α'-Tau-P301L p=0.0067), CK2α **(D)**

(Scr-EGFP vs. siCK2α-EGFP p=0.0008, Scr-EGFP vs. siCk2α-Tau-P301L p=0.0013, Scr-Tau-

P301L vs. siCK2α-EGFP p=0.0161, Scr Tau-P301L vs. siCk2α Tau-P301L p=0.0262, siCK2α

EGFP vs. siCK2α' EGFP p=0.004, siCK2α-EGFP vs. siCK2α'-Tau-P301L p=0.0144, siCk2α-Tau-

P301L vs. siCK2α'-EGFP p=0.0065, siCk2α-Tau-P301L vs. siCK2α'-Tau-P301L p=0.0235), and

CK2β **(E)** in transfected N2a cells. Expression levels were normalized to GAPDH and presented

relative to ScrRNA-EGFP condition (n=3 independent replicas). **(F)** Immunoblot of CK2α', CK2α

and CK2β in N2A cells transfected with EGFP or Tau-P301L-EGFP and treated with siCK2α' or

siCK2α. (**G**) RT-qPCR analysis of mRNA expression levels of huTau expression in N2a cells

transfected with EGFP or Tau-P301L-EGFP transfected N2a cells. Expression levels were

normalized to GAPDH and presented relative to EGFP condition (p=0.0464, n=6/condition). **(H)**

Quantification of AT8+ cells normalized to the number of EGFP+ cells, presented relative to

ScrRNA-Tau-P301L condition. (n=4/condition; Scr-Tau-P301L vs. siCK2α'-Tau-P301L p=0.013).

All data shown as mean ± SEM. Statistical analyses were conducted using one-way ANOVA

with tukey's multiple and displayed with compact letter display.

Figure 3. CK2α' haploinsufficiency reduced tau pathology in PS19 mice. (A-B)

Representative hippocampal sections immunostained for pTau (AT8;Ser202/Thr205). Scale

bar=500 μm. **(C-E)** Quantification of AT8+ immunoreactivity (% area) in the CA1 (C), CA3 (D)

and DG (E) regions in prodromal and symptomatic cohorts (n=6-9 mice/genotype). Data

represented as mean ± SEM. Statistical analyses were conducted using two-way ANOVA with

tukey's posthoc analysis and displayed with compact letter display. (**CA1 significant**

comparisons: prodromal: WT vs. symptomatic: PS19 p<0.0001, prodromal: WT vs.

symptomatic: PS19;CK2α'$^{(+/-)}$ p<0.0001, prodromal: PS19 vs. symptomatic: PS19 p<0.0001,

prodromal: PS19 vs. symptomatic: PS19;CK2α'$^{(+/-)}$ p<0.0001, prodromal: CK2α'$^{(+/-)}$ vs.

symptomatic: PS19 p<0.0001, prodromal: CK2α'$^{(+/-)}$ vs. symptomatic: PS19;CK2α'$^{(+/-)}$ p<0.0001,

prodromal: PS19;CK2α'$^{(+/-)}$ vs. symptomatic: PS19 p<0.0001, prodromal: PS19;CK2α'$^{(+/-)}$ vs.

symptomatic: PS19;CK2α'$^{(+/-)}$ p<0.0001, symptomatic: WT vs. symptomatic: PS19 p<0.0001 ,

symptomatic: WT vs. symptomatic: PS19;CK2α'$^{(+/-)}$ p<0.0001, symptomatic: PS19 vs.

symptomatic: CK2α'$^{(+/-)}$ p<0.0001, symptomatic: PS19 vs. symptomatic: PS19;CK2α'$^{(+/-)}$

p=0.0075, symptomatic: CK2α'$^{(+/-)}$ vs. symptomatic: PS19;CK2α'$^{(+/-)}$ p<0.0001; **CA3 significant**

comparisons: all significant comparisons p<0.0001; **DG significant comparisons:** prodromal:

WT vs. symptomatic: PS19 p<0.0001, prodromal: WT vs. symptomatic: PS19;CK2α'$^{(+/-}$p<0.0001,

prodromal: PS19 vs. symptomatic: PS19 p=<0.0001, prodromal: PS19 vs. symptomatic:

PS19;CK2α'$^{(+/-)}$ p<0.0001, prodromal: CK2α'$^{(+/-)}$ vs. symptomatic: PS19 p<0.0001, prodromal:

CK2α'$^{(+/-)}$ vs. symptomatic:PS19;CK2α'$^{(+/-)}$ p<0.0001, prodromal: PS19;CK2α'$^{(+/-)}$ vs.

symptomatic: PS19 p<0.0001, prodromal :PS19 CK2α'$^{(+/-)}$ vs. symptomatic: PS19;CK2α'$^{(+/-)}$

p<0.0001, symptomatic: WT vs. symptomatic: PS19 p<0.0001, symptomatic: WT vs.

symptomatic: PS19;CK2α'$^{(+/-)}$ p<0.0001, symptomatic: PS19 vs. symptomatic: CK2α'$^{(+/-)}$

p<0.0001, symptomatic: PS19 vs. symptomatic :PS19;CK2α'$^{(+/-)}$ p=0.0186, symptomatic:

CK2α'$^{(+/-)}$ vs. symptomatic:PS19;CK2α'$^{(+/-)}$ p<0.0001). **(F)** Immunoblot for AT8 (pS202/T205),

AT100 (pT212/pS214), and Tau-5 (Total Tau) in hippocampal samples obtained from 9 month

mice. **(G-I)** Quantification of Tau-5 (G), AT8/Tau-5 (H) and AT100/Tau-5 (I) from images shown in

F (n=3-4 mice/genotype). All data are shown as mean $\pm$ SEM. **(J)** Representative pTau

pathology types. **(K)** Proportion of each pTau pathology type in prodromal and symptomatic

cohorts (n=7-10 mice/genotype). Fisher's exact t-test 7m p=0.11 & 12m p=0.0587.

Figure 4. CK2α' haploinsufficiency reduced transcriptional alterations in

apoptotic/phagocytic and synaptic-function related genes in PS19 mice. (A) Volcano plot

of DGEs in PS19 vs WT comparison (n=3-4 mice/genotype). **(B)** Gene-ontology showing the top

biological pathways for DGEs in PS19 vs WT. **(C)** Gene network for top 5 synaptic pathways

impacted in PS19 vs WT. Red dots indicate genes upregulated in PS19 mice. Blue dots indicate

genes down-regulated in PS19 mice. **(D)** Heat map displaying DGEs identified between PS19

and PS19;CK2α'$^{(+/-)}$ mice (n=3-4 mice/genotype). Red boxes indicate genes with

apoptotic/phagocytic and immune-related functions. *q>0.05. **(E)** WGCNA identified modules,

dotted line indicates significance as determined by the Kruskal wallis test between WT, CK2α'[(+/-]

), PS19, and PS19;CK2α'[(+/-)] mice (n=3-5 mice/genotype). **(F)** Heatmap of modules detected

from WGCNA indicating differences across genotypes. * indicates p<0.05 determined by

Kruskal-wallis test. **(G)** Gene ontology for top biological pathways for MEtan module.

Figure 5. CK2α' haploinsufficiency impacts Iba1+ microglia morphology and cytokine

production in PS19 mice. (A) Iba1+ microglia immunostaining and microglia skeleton

representation from the CA1 of symptomatic animals used for microglia morphology analysis.

(B) Quantification of the number of Iba1+ positive cells in the CA1 (n=3-6 mice/genotype; WT

vs. PS19 p<0.0001, WT vs. CK2α'[(+/-)] p=0.9963, WT vs. PS19; CK2α'[(+/-)] p<0.0001, PS19 vs.

CK2α'[(+/-)] p<0.0001, PS19 vs. PS19;CK2α'[(+/-)] p=0.0313, CK2α'[(+/-)] vs. PS19;CK2α'[(+/-)] p<0.0001).

(C) Quantification of microglia branch length (WT vs. PS19 p=0.4402, WT vs. CK2α'[(+/-)]

p=0.1371, WT vs. PS19; CK2α'[(+/-)] p=0.041, PS19 vs. CK2α'[(+/-)] p=0.0184, PS19 vs.

PS19;CK2α'[(+/-)] p=0.927, CK2α'[(+/-)] vs. PS19;CK2α'[(+/-)] p=0.0004). **(D)** Number of branches per

microglia (WT vs. PS19 p=0.0008, WT vs. CK2α'[(+/-)] p=0.2116, WT vs. PS19;CK2α'[(+/-)]

p=0.9908, PS19 vs. CK2α'[(+/-)] p=0.3063, PS19 vs. PS19;CK2α'[(+/-)] p=0.0114, CK2α'[(+/-)] vs. PS19;

CK2α'[(+/-)] p=0.0757). **(E)** Number of endpoints per microglia (WT vs. PS19 p=0.0003, WT vs.

CK2α'[(+/-)] p=0.3541, WT vs. PS19;CK2α'[(+/-)] p=0.5111, PS19 vs. CK2α'[(+/-)] p=0.0556, PS19 vs.

PS19;CK2α'[(+/-)] p=0.0004, CK2α'[(+/-)] vs. PS19;CK2α'[(+/-)] p=0.0179) (n=27 cells (3 mice/genotype,

3 slices/mouse, 3 cells/slice). **(F)** Representative cytokine arrays from PS19 and PS19;CK2α'[(+/-)]

at 9 months from hippocampal protein extracts. Cytokines that showed significant differences

are circled in red. **(G)** Selected quantifications of cytokines showing significance or trending

significance. Quantifications are represented relative to WT (n=4 mice/genotype;Trem1

p=0.0624, Tnfα p=0.1009, IL10 p=0.0948, IL3 p=0.0952, IFNy p=0.0972, GM-CSF p=0.0816, G-

CSF p=0.0562, C5 p=0.0522, Mip 1α p=0.0482, IL17 p=0.0386, IL7 p=0.0481, IL4 p=0.046,

I309 p=0.039). Data are shown as mean $\pm$ SEM. Statistical analyses were conducted using

ANOVA with Geisser-greenhouse correction and Tukey's post-hoc analysis in **B-E** and unpaired

t-test in **G.** Significance displayed with compact letter display or #p$\leq$0.1, *p<0.05.

Figure 6. CK2α' haploinsufficiency decreases microglia reactivity and synaptic

engulfment. (A) CD68 immunostaining with cresyl violet counterstain in the hippocampus of

symptomatic mice. Scale bar=150 μm. (**B**) zoomed images from A in the CA1 and DG. **(C-E)**

Quantifications of % area CD68+ normalized by Iba1+ cell count in prodromal and symptomatic

cohorts in CA1, CA3 and DG (n=3-6 mice/genotype). Comparisons shown relative to cohort age

group (**CA1:** *Prodromal:* WT vs. PS19 p<0.0001, PS19 vs. CK2α'$^{(+/-)}$ p<0.0001, PS19 vs.

PS19;CK2α'$^{(+/-)}$ p=0.0003, *Symptomatic:* WT vs. PS19 p<0.0001, WT vs. PS19;CK2α'$^{(+/-)}$

p=0.0036, PS19 vs. CK2α'$^{(+/-)}$ p<0.0001, PS19 vs. PS19;CK2α'$^{(+/-)}$ p=0.0042, CK2a'(+/-) vs.

PS19;CK2α'$^{(+/-)}$ p=0.0045 **CA3:** *Prodromal:* WT vs. PS19 p=0.0328, WT vs. PS19;CK2α'$^{(+/-)}$

p=0.1739, PS19 vs. CK2α'$^{(+/-)}$ p=0.0193, PS19 vs. PS19; CK2α'$^{(+/-)}$ p=0.6895, CK2α'(+/-) vs.

PS19;CK2α'$^{(+/-)}$ p=0.1045 *Symptomatic:* WT vs. PS19 p<0.0001, WT vs. PS19;CK2α'$^{(+/-)}$

p=0.0003, PS19 vs. CK2α'$^{(+/-)}$ p<0.0001, PS19 vs. PS19;CK2α'$^{(+/-)}$ p=0.0404, CK2α'$^{(+/-)}$ vs. PS19;

CK2α'$^{(+/-)}$ p=0.0003 **DG:** *Prodromal:* WT vs. PS19 p=0.0176, WT vs. PS19; CK2α'$^{(+/-)}$ p=0.9281,

PS19 vs. CK2α'$^{(+/-)}$ p=0.0091, PS19 vs. PS19;CK2α'$^{(+/-)}$ p=0.0536, CK2α'$^{(+/-)}$ vs. PS19;CK2α'$^{(+/-)}$

p=0.7762 *Symptomatic:* WT vs. PS19 p<0.0001, WT vs. PS19;CK2α'$^{(+/-)}$ p=0.0018, PS19 vs.

CK2α'$^{(+/-)}$ p<0.0001, PS19 vs. PS19;CK2α'$^{(+/-)}$ p=0.011, CK2α'$^{(+/-)}$ vs. PS19; CK2α'$^{(+/-)}$ p=0.002. **(F,**

G) Immunoblotting and quantification of protein levels for Clec7a in the hippocampus of

symptomatic animals (n=3 mice/genotype). (**H**) Representative immunofluorescence of Clec7a

in the hippocampus of symptomatic animals. **(I)** Iba1 and PSD95 immunostaining from the CA1

of symptomatic animals. Top row: whole microglia including processes, scale bar= 5μm. Middle

row: higher magnification view of the soma (white dotted box in top row), scale bar= 2µm.

Bottom row: 3D reconstruction of Iba1 and PSD-95 signal generated from Imaris Software,

scale bar= 2µm. **(J)** Quantification of the number of PSD-95 puncta/Iba1+ soma (n= 16-18 cell,

3 mice/genotype, 2-3 slices/mouse, 22 cells/slice; WT vs. PS19 p<0.0001, WT vs. PS19;

CK2α'$^{(+/-)}$ p=0.0854, PS19 vs. CK2α'$^{(+/-)}$ p<0.0001, PS19 vs. PS19; CK2α'$^{(+/-)}$ p=0.0025, CK2α'$^{(+/-)}$

vs. PS19;CK2α'$^{(+/-)}$ p=0.0208). Data are show as mean $\pm$ SEM. Statistical analyses were

conducted using one-way ANOVA with Tukey's post-hoc displayed with compact letter display.

Figure 7. CK2α' haploinsufficiency partially rescued hippocampal synapse loss and

improved LTP PS19 mice. (A) Representative NeuN staining in the CA1 of symptomatic mice.

Scale bar=200 µm. **(B)** Quantification of NeuN+ cells (cells/mm^2) in the CA1 of prodromal and

symptomatic cohorts (n= 3-6 mice/genotype; 2-3 slices/mouse; 1 data point= 1 slice, slices from

same animal share a shape). Comparisons shown relative to age group (Significant

comparisons: *Symptomatic:* WT vs. PS19 p<0.0001, WT vs. PS19;CK2α'$^{(+/-)}$ p=0.0002, PS19 vs.

CK2α'$^{(+/-)}$ p<0.0001, CK2α'$^{(+/-)}$ vs. PS19;CK2α'$^{(+/-)}$ p=0.0001). **(C)** Vglut1 and PSD95

immunostaining in the CA1 of symptomatic mice. White arrow heads indicate colocalization.

Scale bar= 2.5 µm. **(D-F)** Quantification of PSD95 puncta (WT vs. PS19 p=0.0445, PS19 vs.

CK2α'$^{(+/-)}$ p=0.0008) (**D**), Vglut1 puncta (WT vs. PS19 p=0.0003, WT vs. PS19;CK2α'$^{(+/-)}$

p=0.0212, PS19 vs. CK2α'$^{(+/-)}$ p<0.0001) (**E**) and Vglut1/PSD95 colocalized puncta (WT vs.

PS19 p=0.0017, PS19 vs. CK2α'$^{(+/-)}$ p=0.0004) (**F**) relative to WT. (n=3 mice/genotype (3

slices/mouse, 3 images/slice, 1 data point=slice average). **(G-I)** Analyses from (D-F)

represented as a % decrease from WT (G: WT vs. PS19 p=0.0445, PS19 vs. CK2α'$^{(+/-)}$

p=0.0008 H: WT vs. PS19 p=0.0003, WT vs. PS19;CK2α'$^{(+/-)}$ p=0.0212, PS19 vs. CK2α'$^{(+/-)}$

p<0.0001, I: WT vs. PS19 p=0.0017, PS19 vs. CK2α'$^{(+/-)}$ p=0.0004) **(J)** Experimental set-up: R,

recording electrode; S stimulating electrode. **(K)** Representative traces before (1) and after (2)

Theta-burst stimulation (TBS). **(L)** Course temporal of TBS application inducing LTP in

pyramidal neurons of CA1 (n=6-7 mice/genotype). **(M)** Short-term potentiation (STP) results,

significance reported relative to WT (WT vs. PS19 p=0.014, WT vs. PS19;CK2$\alpha'^{(+/-)}$ p=0.007).

(N) Input-output curves. Averaged fEPSP amplitude, (n=3mice/genotype). **(O)** Long-term

potentiation (LTP) results, significance reported relative to WT (WT vs. PS19 p=0.007). All data

are shown as mean ± SEM. Statistics: one way-ANOVA with Tukey's (**B**), repeated measures

ANOVA with Geisser-greenhouse correction and Tukey's (**D-I**), or one-way ANOVA with Holm-

Sidak (**M**-**O**). Significance displayed as compact letter display.

Figure 8. Depletion of CK2α' improved spatial memory of PS19 mice on the Barnes maze.

(A) Schematic of the Barnes Maze divided into 4 quadrants (Target, Opposite, +1 and -1) and

experimental set up. **(B-C)** Primary latency (s), or time to first explore escape hole at prodromal

time point. Statistical analyses were conducted using two-way ANOVA with Bonferroni's multiple

comparison's. Prodromal; genotype p=0.0573, training day p<0.0001 and interaction p=0.9690.

Symptomatic: genotype p<0.0001, training day p<0.0001 and interaction p=0.7018. Differences

for each training day were conducted using Bonferroni's multiple comparisons and * show

significance relative to WT. **(D)** Time spent in goal quadrant of Barnes maze during probe trial at

prodromal and symptomatic time points. n=10-12 mice/genotype, 2 outliers were removed from

PS19 group using ROUT (Q=1%). **(E)** Representative search strategy traces based on their

navigation patterns on the maze. **(F,H)** Spatial strategies used to locate the escape hole across

training days for each genotype at prodromal and symptomatic stages, respectively. **G,I)** Spatial

strategies used to locate the previously learned escape target hole during probe trial. Data are

shown as mean ± SEM. Statistical analyses were conducted using Two-way ANOVA with

Bonferroni's multiple comparisons (B-C) or Two-Way ANOVA with Tukey's post-hoc analysis (D).

Significance displayed with either *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 or compact letter

display.

References

1.　B. Kovacech, R. Skrabana, M. Novak, Transition of tau protein from disordered to

misordered in Alzheimer's disease. *Neurodegener Dis* **7**, 24-27 (2010).

2.　V. M. Lee, M. Goedert, J. Q. Trojanowski, Neurodegenerative tauopathies. *Annu Rev*

Neurosci **24**, 1121-1159 (2001).

3.　K. A. Josephs *et al.*, Neuropathological background of phenotypical variability in

frontotemporal dementia. *Acta Neuropathol* **122**, 137-153 (2011).

4.　H. C. Tai *et al.*, The synaptic accumulation of hyperphosphorylated tau oligomers in

Alzheimer disease is associated with dysfunction of the ubiquitin-proteasome system.

Am J Pathol **181**, 1426-1435 (2012).

5.　M. Dani *et al.*, Microglial activation correlates in vivo with both tau and amyloid in

Alzheimer's disease. *Brain* **141**, 2740-2754 (2018).

6.　S. H. Lee, E. J. Bae, S. J. Park, S. J. Lee, Microglia-driven inflammation induces

progressive tauopathies and synucleinopathies. *Exp Mol Med*, (2025).

7.　A. Bejanin *et al.*, Tau pathology and neurodegeneration contribute to cognitive

impairment in Alzheimer's disease. *Brain* **140**, 3286-3300 (2017).

8.　C. Langworth-Green *et al.*, Chronic effects of inflammation on tauopathies. *Lancet*

Neurol **22**, 430-442 (2023).

9.　C. Wang *et al.*, Microglial NF-κB drives tau spreading and toxicity in a mouse model of

tauopathy. *Nat Commun* **13**, 1969 (2022).

10.   Y. Kim, S. H. Ryu, J. Hyun, Y. S. Cho, Y. K. Jung, TLR2 immunotherapy suppresses

904   neuroinflammation, tau spread, and memory loss in rTg4510 mice. *Brain Behav Immun*

905   **121**, 291-302 (2024).

906   11.   D. Dutta *et al.*, Tau fibrils induce glial inflammation and neuropathology via TLR2 in

907   Alzheimer's disease-related mouse models. *J Clin Invest* **133**, (2023).

908   12.   K. Bhaskar *et al.*, Regulation of tau pathology by the microglial fractalkine receptor.

909   *Neuron* **68**, 19-31 (2010).

910   13.   D. W. Litchfield, Protein kinase CK2: structure, regulation and role in cellular decisions of

911   life and death. *Biochem J* **369**, 1-15 (2003).

912   14.   R. Gomez-Pastor *et al.*, Abnormal degradation of the neuronal stress-protective

913   transcription factor HSF1 in Huntington's disease. *Nat Commun* **8**, 14405 (2017).

914   15.   D. Yu *et al.*, CK2 alpha prime and alpha-synuclein pathogenic functional interaction

915   mediates synaptic dysregulation in huntington's disease. *Acta Neuropathol Commun* **10**,

916   83 (2022).

917   16.   Q. Zhang *et al.*, CK2 Phosphorylating I 2 PP2A /SET Mediates Tau Pathology and

918   Cognitive Impairment. *Front Mol Neurosci* **11**, 146 (2018).

919   17.   H. Yadikar *et al.*, Screening of tau protein kinase inhibitors in a tauopathy-relevant cell-

920   based model of tau hyperphosphorylation and oligomerization. *PLoS One* **15**, e0224952

921   (2020).

922   18.   L. Baum *et al.*, Casein kinase II is associated with neurofibrillary tangles but is not an

923   intrinsic component of paired helical filaments. *Brain Res* **573**, 126-132 (1992).

924   19.   A. C. Lim, S. Y. Tiu, Q. Li, R. Z. Qi, Direct regulation of microtubule dynamics by protein

925   kinase CK2. *J Biol Chem* **279**, 4433-4439 (2004).

926   20.   R. Kimura, N. Matsuki, Protein kinase CK2 modulates synaptic plasticity by modification

927   of synaptic NMDA receptors in the hippocampus. *J Physiol* **586**, 3195-3206 (2008).

928   21.   H. J. Chung, Y. H. Huang, L. F. Lau, R. L. Huganir, Regulation of the NMDA receptor

929   complex and trafficking by activity-dependent phosphorylation of the NR2B subunit PDZ

930   ligand. *J Neurosci* **24**, 10248-10259 (2004).

931   22.   S. C. Lenzken *et al.*, Recruitment of casein kinase 2 is involved in AbetaPP processing

932   following cholinergic stimulation. *J Alzheimers Dis* **20**, 1133-1141 (2010).

933   23.   M. Raftery *et al.*, Phosphorylation of apolipoprotein-E at an atypical protein kinase CK2

934   PSD/E site in vitro. *Biochemistry* **44**, 7346-7353 (2005).

935   24.   J. Walter, A. Schindzielorz, B. Hartung, C. Haass, Phosphorylation of the beta-amyloid

936   precursor protein at the cell surface by ectocasein kinases 1 and 2. *J Biol Chem* **275**,

937   23523-23529 (2000).

938   25.   Y. Bian *et al.*, Global screening of CK2 kinase substrates by an integrated

939   phosphoproteomics workflow. *Sci Rep* **3**, 3460 (2013).

940   26.   S. F. Rusin, M. E. Adamo, A. N. Kettenbach, Identification of Candidate Casein Kinase 2

941   Substrates in Mitosis by Quantitative Phosphoproteomics. *Front Cell Dev Biol* **5**, 97

942   (2017).

943   27.   B. Guerra, S. Siemer, B. Boldyreff, O. G. Issinger, Protein kinase CK2: evidence for a

944   protein kinase CK2beta subunit fraction, devoid of the catalytic CK2alpha subunit, in

945   mouse brain and testicles. *FEBS Lett* **462**, 353-357 (1999).

946   28.   M. Montenarh, C. Götz, Protein Kinase CK2α', More than a Backup of CK2α. *Cells* **12**,

947   (2023).

948   29.   I. Ceglia, M. Flajolet, H. Rebholz, Predominance of CK2α over CK2α' in the mammalian

949   brain. *Mol Cell Biochem* **356**, 169-175 (2011).

950   30.   D. Y. Lou *et al.*, The alpha catalytic subunit of protein kinase CK2 is required for mouse

951   embryonic development. *Mol Cell Biol* **28**, 131-139 (2008).

952   31.   X. Xu, P. A. Toselli, L. D. Russell, D. C. Seldin, Globozoospermia in mice lacking the

953   casein kinase II alpha' catalytic subunit. *Nat Genet* **23**, 118-121 (1999).

32.   D. S. Iimoto, E. Masliah, R. DeTeresa, R. D. Terry, T. Saitoh, Aberrant casein kinase II in

Alzheimer's disease. *Brain Res* **507**, 273-280 (1990).

33.   J. E. Allende, C. C. Allende, Protein kinases. 4. Protein kinase CK2: an enzyme with

multiple substrates and a puzzling regulation. *FASEB J* **9**, 313-323 (1995).

34.   A. F. Rosenberger *et al.*, Increased occurrence of protein kinase CK2 in astrocytes in

Alzheimer's disease pathology. *J Neuroinflammation* **13**, 4 (2016).

35.   M. V. Aksenova, G. S. Burbaeva, K. V. Kandror, D. V. Kapkov, A. S. Stepanov, The

decreased level of casein kinase 2 in brain cortex of schizophrenic and Alzheimer's

disease patients. *FEBS Lett* **279**, 55-57 (1991).

36.   T. Saitoh, D. Iimoto, Aberrant protein phosphorylation and cytoarchitecture in Alzheimer's

disease. *Prog Clin Biol Res* **317**, 769-780 (1989).

37.   J. A. Miller *et al.*, Neuropathological and transcriptomic characteristics of the aged brain.

Elife **6**, (2017).

38.   Y. Yoshiyama *et al.*, Synapse loss and microglial activation precede tangles in a P301S

tauopathy mouse model. *Neuron* **53**, 337-351 (2007).

39.   C. K. Walker *et al.*, Dendritic Spine Remodeling and Synaptic Tau Levels in PS19

Tauopathy Mice. *Neuroscience* **455**, 195-211 (2021).

40.   J. P. Anderson *et al.*, Phosphorylation of Ser-129 is the dominant pathological

modification of alpha-synuclein in familial and sporadic Lewy body disease. *J Biol Chem*

281, 29739-29752 (2006).

41.   H. Takeuchi *et al.*, P301S mutant human tau transgenic mice manifest early symptoms

of human tauopathies with dementia and altered sensorimotor gating. *PLoS One* **6**,

e21050 (2011).

42.   Y. Zhou *et al.*, The Alzheimer's Cell Atlas (TACA): A single-cell molecular map for

translational therapeutics accelerator in Alzheimer's disease. *Alzheimers Dement (N Y)*

8, e12350 (2022).

43.   E. Gerrits *et al.*, Distinct amyloid-β and tau-associated microglia profiles in Alzheimer's

disease. *Acta Neuropathol* **141**, 681-696 (2021).

44.   M. Otero-Garcia *et al.*, Molecular signatures underlying neurofibrillary tangle

susceptibility in Alzheimer's disease. *Neuron* **110**, 2929-2948.e2928 (2022).

45.   K. Leng *et al.*, Molecular characterization of selectively vulnerable neurons in

Alzheimer's disease. *Nat Neurosci* **24**, 276-287 (2021).

46.   S. F. Lau, H. Cao, A. K. Y. Fu, N. Y. Ip, Single-nucleus transcriptome analysis reveals

dysregulation of angiogenic endothelial cells and neuroprotective glia in Alzheimer's

disease. *Proc Natl Acad Sci U S A* **117**, 25800-25809 (2020).

47.   C. A. Marshall *et al.*, Inhibition of CK2 mitigates Alzheimer's tau pathology by preventing

NR2B synaptic mislocalization. *Acta Neuropathol Commun* **10**, 30 (2022).

48.   B. R. Hoover *et al.*, Tau mislocalization to dendritic spines mediates synaptic dysfunction

independently of neurodegeneration. *Neuron* **68**, 1067-1081 (2010).

49.   Y. Shi *et al.*, Microglia drive APOE-dependent neurodegeneration in a tauopathy mouse

model. *J Exp Med* **216**, 2546-2561 (2019).

50.   Y. Shi *et al.*, ApoE4 markedly exacerbates tau-mediated neurodegeneration in a mouse

model of tauopathy. *Nature* **549**, 523-527 (2017).

51.   H. Martini-Stoica *et al.*, TFEB enhances astroglial uptake of extracellular tau species and

reduces tau spreading. *J Exp Med* **215**, 2355-2377 (2018).

52.   H. Takahashi *et al.*, Reduced progranulin increases tau and α-synuclein inclusions and

alters mouse tauopathy phenotypes via glucocerebrosidase. *Nat Commun* **15**, 1434

(2024).

53.   A. Litvinchuk *et al.*, Complement C3aR Inactivation Attenuates Tau Pathology and

Reverses an Immune Network Deregulated in Tauopathy Models and Alzheimer's

Disease. *Neuron* **100**, 1337-1353.e1335 (2018).

54.   J. Kim *et al.*, Cerebral transcriptome analysis reveals age-dependent progression of

1006   neuroinflammation in P301S mutant tau transgenic male mice. *Brain Behav Immun* **80**,

1007   344-357 (2019).

1008   55.   F. H. Shiu *et al.*, ADGRB1 contributes to astrocyte-mediated phagocytosis of excitatory

1009   synapses. *Exp Neurol* **395**, 115451 (2025).

1010   56.   T. Lala, R. A. Hall, Adhesion G protein-coupled receptors: structure, signaling,

1011   physiology, and pathophysiology. *Physiol Rev* **102**, 1587-1624 (2022).

1012   57.   X. Chen *et al.*, Microglia-mediated T cell infiltration drives neurodegeneration in

1013   tauopathy. *Nature* **615**, 668-677 (2023).

1014   58.   K. F. Odfalk, K. F. Bieniek, S. C. Hopp, Microglia: Friend and foe in tauopathy. *Prog*

1015   *Neurobiol* **216**, 102306 (2022).

1016   59.   C. Gao, J. Jiang, Y. Tan, S. Chen, Microglia in neurodegenerative diseases: mechanism

1017   and potential therapeutic targets. *Signal Transduct Target Ther* **8**, 359 (2023).

1018   60.   S. Fixemer *et al.*, Microglia phenotypes are associated with subregional patterns of

1019   concomitant tau, amyloid-β and α-synuclein pathologies in the hippocampus of patients

1020   with Alzheimer's disease and dementia with Lewy bodies. *Acta Neuropathol Commun*

1021   **10**, 36 (2022).

1022   61.   R. Hassan-Abdi, A. Brenet, M. Bennis, C. Yanicostas, N. Soussi-Yanicostas, Neurons

1023   Expressing Pathological Tau Protein Trigger Dramatic Changes in Microglial Morphology

1024   and Dynamics. *Front Neurosci* **13**, 1199 (2019).

1025   62.   S. Hong *et al.*, Complement and microglia mediate early synapse loss in Alzheimer

1026   mouse models. *Science* **352**, 712-716 (2016).

1027   63.   B. Dejanovic *et al.*, Changes in the Synaptic Proteome in Tauopathy and Rescue of Tau-

1028   Induced Synapse Loss by C1q Antibodies. *Neuron* **100**, 1322-1336.e1327 (2018).

1029   64.   H. Lui *et al.*, Progranulin Deficiency Promotes Circuit-Specific Synaptic Pruning by

1030   Microglia via Complement Activation. *Cell* **165**, 921-935 (2016).

65.   K. Pampuscenko *et al.*, Extracellular tau stimulates phagocytosis of living neurons by
activated microglia via Toll-like 4 receptor-NLRP3 inflammasome-caspase-1 signalling
axis. *Sci Rep* **13**, 10813 (2023).

66.   J. Leyh *et al.*, Classification of Microglial Morphological Phenotypes Using Machine
Learning. *Front Cell Neurosci* **15**, 701673 (2021).

67.   S. Parhizkar *et al.*, Lemborexant ameliorates tau-mediated sleep loss and
neurodegeneration in males in a mouse model of tauopathy. *Nat Neurosci*, (2025).

68.   E. R. Roy *et al.*, Concerted type I interferon signaling in microglia and neural cells
promotes memory impairment associated with amyloid β plaques. *Immunity* **55**, 879-
894.e876 (2022).

69.   Y. B. Choi *et al.*, Erythropoietin-derived peptide treatment reduced neurological deficit
and neuropathological changes in a mouse model of tauopathy. *Alzheimers Res Ther*
13, 32 (2021).

70.   S. L. DeVos *et al.*, Tau reduction prevents neuronal loss and reverses pathological tau
deposition and seeding in mice with tauopathy. *Sci Transl Med* **9**, (2017).

71.   J. Wagner *et al.*, Reducing tau aggregates with anle138b delays disease progression in
a mouse model of tauopathies. *Acta Neuropathol* **130**, 619-631 (2015).

72.   N. Zheng *et al.*, Electrophysiology-based screening identifies neuronal HtrA serine
peptidase 2 (HTRA2) as a synaptic plasticity regulator participating in tauopathy. *Transl*
Psychiatry **15**, 5 (2025).

73.   R. El Fatimy *et al.*, MicroRNA-132 provides neuroprotection for tauopathies via multiple
signaling pathways. *Acta Neuropathol* **136**, 537-555 (2018).

74.   T. Illouz *et al.*, Unbiased classification of spatial strategies in the Barnes maze.
Bioinformatics **32**, 3314-3320 (2016).

75.   K. Chen *et al.*, Selective removal of astrocytic PERK protects against glymphatic
impairment and decreases toxic aggregation of β-amyloid and tau. *Neuron*, (2025).

76.   I. I. N. Da Silva *et al.*, CK2 inhibition suppresses glial inflammation in the brain. *bioRxiv*,

1058   (2025).

1059   77.   M. Pérez, J. Avila, The expression of casein kinase 2alpha' and phosphatase 2A activity.

1060   *Biochim Biophys Acta* **1449**, 150-156 (1999).

1061   78.   J. Liu, J. Tian, R. Xie, L. Chen, CK2 inhibitor DMAT ameliorates spinal cord injury by

1062   increasing autophagy and inducing anti-inflammatory microglial polarization. *Neurosci*

1063   *Lett* **805**, 137222 (2023).

1064   79.   S. Mishra, C. Kinoshita, A. D. Axtman, J. E. Young, Evaluation of a Selective Chemical

1065   Probe Validates That CK2 Mediates Neuroinflammation in a Human Induced Pluripotent

1066   Stem Cell-Derived Microglial Model. *Front Mol Neurosci* **15**, 824956 (2022).

1067   80.   I. Y. Chen *et al.*, Upregulation of the chemokine (C-C motif) ligand 2 via a severe acute

1068   respiratory syndrome coronavirus spike-ACE2 signaling pathway. *J Virol* **84**, 7703-7712

1069   (2010).

1070   81.   S. R. Larson *et al.*, Myeloid Cell CK2 Regulates Inflammation and Resistance to

1071   Bacterial Infection. *Front Immunol* **11**, 590266 (2020).

1072   82.   D. Drygin *et al.*, Protein kinase CK2 modulates IL-6 expression in inflammatory breast

1073   cancer. *Biochem Biophys Res Commun* **415**, 163-167 (2011).

1074   83.   K. Parhar, J. Morse, B. Salh, The role of protein kinase CK2 in intestinal epithelial cell

1075   inflammatory signaling. *Int J Colorectal Dis* **22**, 601-609 (2007).

1076   84.   M. Yamada *et al.*, Inhibition of protein kinase CK2 prevents the progression of

1077   glomerulonephritis. *Proc Natl Acad Sci U S A* **102**, 7736-7741 (2005).

1078   85.   J. Huang *et al.*, Sphingosine kinase 1 mediates diabetic renal fibrosis via NF-κB

1079   signaling pathway: involvement of CK2α. *Oncotarget* **8**, 88988-89004 (2017).

1080   86.   N. N. Singh, D. P. Ramji, Protein kinase CK2, an important regulator of the inflammatory

1081   response? *J Mol Med (Berl)* **86**, 887-897 (2008).

87. J. Avila, J. J. Lucas, M. Perez, F. Hernandez, Role of tau protein in both physiological

and pathological conditions. *Physiol Rev* **84**, 361-384 (2004).

88. N. Zarate *et al.*, Neurochemical correlates of synapse density in a Huntington's disease

mouse model. *J Neurochem* **164**, 226-241 (2023).

89. E. M. Coomans *et al.*, In vivo tau pathology is associated with synaptic loss and altered

synaptic function. *Alzheimers Res Ther* **13**, 35 (2021).

90. T. Vogels, A. N. Murgoci, T. Hromádka, Intersection of pathological tau and microglia at

the synapse. *Acta Neuropathol Commun* **7**, 109 (2019).

91. X. Liu *et al.*, Proteomic analysis of ferroptosis pathways reveals a role of CEPT1 in

suppressing ferroptosis. *Protein Cell* **15**, 686-703 (2024).

92. P. Yin *et al.*, Aged monkey brains reveal the role of ubiquitin-conjugating enzyme UBE2N

in the synaptosomal accumulation of mutant huntingtin. *Hum Mol Genet* **24**, 1350-1362

(2015).

93. J. Y. Qiu *et al.*, OXCT1 regulates hippocampal neurogenesis and alleviates cognitive

impairment via the Akt/GSK-3β/β-catenin pathway after subarachnoid hemorrhage. *Brain*

Res **1827**, 148758 (2024).

94. J. M. Serfass *et al.*, Endophilin B2 facilitates endosome maturation in response to growth

factor stimulation, autophagy induction, and influenza A virus infection. *J Biol Chem* **292**,

10097-10111 (2017).

95. Y. H. Wang *et al.*, Endophilin B2 promotes inner mitochondrial membrane degradation by

forming heterodimers with Endophilin B1 during mitophagy. *Sci Rep* **6**, 25153 (2016).

96. O. Bohana-Kashtan, L. A. Pinna, Z. Fishelson, Extracellular phosphorylation of C9 by

protein kinase CK2 regulates complement-mediated lysis. *Eur J Immunol* **35**, 1939-1948

(2005).

97.   S. Pelloux *et al.*, Identification of a cryptic protein kinase CK2 phosphorylation site in

human complement protease C1r, and its use to probe intramolecular interaction. *FEBS*

Lett **386**, 15-20 (1996).

98.   D. Mudaliar *et al.*, Discovery of a CK2α'-Biased ATP-Competitive Inhibitor from a High-

Throughput Screen of an Allosteric-Inhibitor-Like Compound Library. *ACS Chem*

Neurosci, (2024).

99.   S. C. Vermilyea *et al.*, Loss of tau expression attenuates neurodegeneration associated

with α-synucleinopathy. *Transl Neurodegener* **11**, 34 (2022).

100.   T. G. Brown *et al.*, Striatal spatial heterogeneity, clustering, and white matter association

of GFAP. *Front Cell Neurosci* **17**, 1094503 (2023).

101.   M. A. Solem *et al.*, Absence of hippocampal pathology persists in the Q175DN mouse

model of Huntington's disease despite elevated HTT aggregation. *J Huntingtons Dis* **14**,

59-84 (2025).

102.   P. Bankhead *et al.*, QuPath: Open source software for digital pathology image analysis.

Sci Rep **7**, 16878 (2017).

103.   A. C. Ruifrok, D. A. Johnston, Quantification of histochemical staining by color

deconvolution. *Anal Quant Cytol Histol* **23**, 291-299 (2001).

104.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat*

Methods **9**, 676-682 (2012).

105.   K. Young, H. Morrison, Quantifying Microglia Morphology from Photomicrographs of

Immunohistochemistry Prepared Tissue Using ImageJ. *J Vis Exp*, (2018).

106.   I. Arganda-Carreras, R. Fernández-González, A. Muñoz-Barrutia, C. Ortiz-De-Solorzano,

3D reconstruction of histological sections: Application to mammary gland tissue. *Microsc*

Res Tech **73**, 1019-1029 (2010).

107.   N. Zarate *et al.*, Heat Shock Factor 1 Directly Regulates Postsynaptic Scaffolding PSD-

95 in Aging and Huntington's Disease and Influences Striatal Synaptic Density. *Int J Mol*

Sci **22**, (2021).

108.   S. U. McKinstry *et al.*, Huntingtin is required for normal excitatory synapse development

in cortical and striatal circuits. *J Neurosci* **34**, 9455-9472 (2014).

109.   M. W. Pitts, Barnes Maze Procedure for Spatial Learning and Memory in Mice. *Bio*

Protoc **8**, (2018).

110.   C. A. Barnes, Memory deficits associated with senescence: a neurophysiological and

behavioral study in the rat. *J Comp Physiol Psychol* **93**, 74-104 (1979).

111.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

112.   U. Raudvere *et al.*, g:Profiler: a web server for functional enrichment analysis and

conversions of gene lists (2019 update). *Nucleic Acids Res* **47**, W191-W198 (2019).

1143

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8