Protein kinase CK2α' as a dual modulator of immune signaling and synaptic dysfunction
in Tauopathy

Angel White[1], Peter Gavrilyuk[1], Rafael Falcon Moya[1], Reid Thurston[1], Amal Fickak[1], Nicholas B
Rozema[1], Prarthana Keshavaram[1], Scott Vermilyea[1], Riley Schlichte[1], Joyce Meints[1], Ying
Zhang[2], Alfonso Araque[1], Michael Lee[1], Rocio Gomez-Pastor[1]

[1]University of Minnesota, School of Medicine, Department of Neuroscience.

[2]Minnesota Supercomputing Institute, University of Minnesota.

Abstract

Tauopathies are a group of neurodegenerative diseases characterized by tau accumulation,
neuroinflammation, and synaptic dysfunction, yet effective treatments remain elusive. Protein
Kinase CK2 has been previously associated with different aspects of tau pathology but genetic
evidence for the contribution of CK2 to tauopathy remained lacking. Here, we show CK2α', one
of the two catalytic subunits of CK2, as a novel regulator of tau-mediated neurodegeneration.
We found that CK2α' expression is elevated in the hippocampus of PS19 tauopathy mice and in
postmortem brains of dementia patients, particularly in neurons and microglia. Using genetic
haploinsufficiency in PS19 mice, we demonstrated that reduced CK2α' levels significantly
decrease phosphorylated tau and total tau burden in the hippocampus and cortex. CK2α'
depletion also enhanced synaptic gene expression, synaptic density, and LTP, while attenuating
microglial activation, synaptic engulfment, and pro-inflammatory cytokine levels. Importantly,
CK2α' depletion rescued cognitive deficits assessed in the Barnes maze. These effects appear
to be mediated through both neuronal and glial functions and may involve CK2α'-dependent
modulation of tau-associated phosphorylation and neuroinflammatory and immune signaling
pathways. Our findings highlight CK2α' as a key node at the intersection of tau pathology,
synaptic dysfunction, and neuroimmune signaling. Targeting CK2α' may offer a novel and
selective therapeutic strategy for modifying disease progression in tauopathies.

Introduction

Tauopathies encompass a broad range of neurodegenerative conditions, most notably
Alzheimer's disease (AD) and related dementias (ADRD), for which no cure currently exists[1-3].
These disorders are generally categorized as either primary or secondary tauopathies. Primary
tauopathies result from direct mutations in the microtubule-associated protein tau (MAPT), a
protein critical for microtubule stabilization and intracellular transport[4,5]. In contrast, secondary
tauopathies arise from pathological tau accumulation triggered by mutations in genes other than
MAPT[6]. Regardless of the cause, tauopathies are typically marked by progressive neuronal
loss, synaptic and cognitive dysfunction, and widespread inflammation in both the central
nervous system and peripheral tissues[7-11]. A deeper understanding of the molecular
mechanisms underlying tauopathies and their downstream consequences is essential for
developing effective disease-modifying therapies. One key area of research is the role

neuroinflammation plays in the progression of tauopathies. These disorders are characterized
by chronic, excessive inflammatory responses to abnormal tau proteins[8,9,12]. This persistent
inflammation can exacerbate tau pathology by promoting sustained neuronal damage and
dysfunction[12-15]. However, despite ongoing research, the mechanisms that connect
neuroinflammation and tau pathology are still poorly understood.

Protein kinase CK2 (formerly known as Casein Kinase II), is a serine/threonine kinase that has
emerged as an important regulator of neuroinflammation and protein homeostasis in several
neurodegenerative diseases[16-18], and it has previously been connected to tau pathology in AD[19-
22]. However, its specific role in the inflammatory pathways associated with tauopathies remains
poorly defined. Interestingly, CK2 has demonstrated potential as a therapeutic target in various
neurodegenerative contexts [17,18]. In AD, CK2 has been implicated in processes such as synaptic
plasticity[23,24], amyloid precursor protein (APP) processing[25-27], and tau accumulation[21,22],
highlighting its relevance for further investigation.

CK2 is a tetrameric enzyme composed of 2 regulatory β subunits (CK2β) and 2 catalytic
subunits either α (CKα) or α' (CK2α')[16]. CK2α and CK2α' are highly similar in structure but
display differential expression and substrate specificity[28-34]. CK2α is expressed throughout the
body at relatively equal levels and has hundreds of substrates, while CK2α' expression is more
restricted to the testes and the brain and has very few validated substrates[28-34]. Another
important difference between these two catalytic subunits is the embryonic lethality of CK2α
knock-out mice, while complete deletion of CK2α' does not have any major defects other than
male sterility[33,34]. Recently, studies in Huntington's disease (HD) have shown the specific
upregulation of CK2α' in cell and mouse models of HD as well as in affected brain tissues of
patients with HD[17,18]. Importantly, HD mice lacking one allele of CK2α' showed improved
synaptic function, reduced protein aggregates, and ameliorated behavioral deficits, suggesting a
key role of CK2α' in this pathology[17,18].

CK2 was one of the first protein kinases identified to be altered in an AD brain[35,36], but its role in
AD has been somewhat controversial due to discrepancies in the ability to detect consistent
alterations in different affected tissues. Early studies using pan-CK2 antibodies (non-selective
for CK2α'/CK2α subunits) showed increased levels of total CK2 in the hippocampus and frontal
cortex of AD mouse models and patients with AD, especially in glial cells [37] but others have
reported opposite results[35,38,39]. In those studies where levels of CK2 were elevated, the authors
associated the increase in CK2 immunoreactivity with the activation of neuroinflammatory
processes and AD pathology[37]. However, to date, genetic studies testing the role of CK2 (either
CK2α' or CK2α) in AD/ADRD models are lacking.

Despite the lack of genetic evidence for the potential differential contribution between CK2α' and
CK2α to AD pathology, several studies have proposed the role of CK2 in the phosphorylation of
Tau (pTau) and tau-mediated pathology. In a kinase inhibitor screening using okadaic acid-
induced pTau in Neuro-2a (N2a) cells CK2 was returned as one of the kinases whose inhibition
most impacted pTau[20]. CK2 has been shown to phosphorylate the phosphatase PP2a inhibitor
SET (I2PP2A/SET) increasing pTau in primary neuronal cultures treated with amyloid-β (Aβ) or
overexpressing human Tau (hTau)[19]. Additionally, wild-type (WT) mice overexpressing CK2 were
also shown to have increased pTau, impaired synaptic plasticity, and display cognitive deficits

like those seen in tauopathies[19]. Overall, this evidence suggests a role of CK2 in tauopathies
including pTau accumulation, synaptic plasticity, and cognitive deficits.

Previous studies assessing the role of CK2 in tauopathy have largely focused on the CK2
holoenzyme without differentiating between the different catalytic CK2α and CK2α' subunits.
Given the substantial difference in substrate specificity between the two catalytic subunits, it is
crucial to determine whether CK2α' and CK2α contribute differently to the onset and/or
progression of tauopathies. Understanding these differences is essential for designing
therapeutic strategies that are both highly effective and minimize off-target effects. The lack of
commercially available inhibitors that discriminate between the different catalytic subunits offers
an attractive window for designing and development of specific inhibitors to selectively inhibit
CK2 catalytic subunits.

Due to previous findings specifically connecting the dysregulation of CK2α' with various
neurodegenerative diseases and reports linking this subunit to the phosphorylation of Tau in
different contexts, we explored the role of CK2α' in various tauopathy models. We confirmed a
specific upregulation at the RNA and protein levels of CK2α' in affected tissues of AD and FTD
patients as well as a mouse model of tauopathy (PS19). CK2α' upregulation was specifically
observed in both neurons and microglia. Importantly, genetic depletion of CK2α' in both cell and
mouse models of tauopathy (expressing Tau-P301L or Tau-P301S mutations) resulted in
decreased pTau. Furthermore, PS19 mice haploinsufficient for CK2α' showed improved synaptic
density, synapse function, and cognition. However, the effects of CK2α' haploinsufficiency were
more profoundly noted in inflammation and immune processes and in the restoration of
microglia morphology, cytokine levels and phagocytic activity. Overall, our data showed a large
role of the CK2α' catalytic subunit specifically in tau pathology, largely connected to the
modulation of microglia functions. These studies highlight CK2α' as an attractive therapeutic
target for the treatment of AD/ADRD.

Methods

Mouse Lines

For this study we used the B6;C3-Tg(Prnp-MPAT-P301S)PS19Vle/J (or PS19)[40] mouse line
(Jackson #008169 maintained with purchased B6C3H breeders; obtained from Dr. Karen Ashe
at University of Minnesota), and CK2α'$^{(+/-)}$ mice, originally obtained from Dr. Seldin at Boston
University (Taconic biosciences TF3062)[34] and maintained on the C57Bl/6J background for
several generations. We crossed PS19 and CK2α'$^{(+/-)}$ mice and this cross generated 4
genotypes of interest WT (PS19$^{0/0}$;CK2α'$^{(+/+)}$), CK2α'$^{(+/-)}$ (PS19$^{0/0}$;CK2α'$^{(+/-)}$), PS19
(PS19$^{Tg/0}$;CK2α'$^{(+/+)}$), and PS19;CK2α'$^{(+/-)}$ (PS19$^{Tg/0}$;CK2α'$^{(+/-)}$) mice. For all experiments
littermate WT and CK2α'$^{(+/-)}$ controls were used. Two time points were used in experiments; pre-
symptomatic (6-7 months) and symptomatic (10-12 months). All mice were housed under
standard SPF conditions. All animal care and sacrifice procedures were approved by the
University of Minnesota Institutional Animal Care and Use Committee (IACUC) in compliance
with the National Institutes of Health guidelines for the care and use of laboratory animals under
the approved animal protocol 2307-A41243.

Human postmortem tissue

Postmortem human frontal cortex (Brodmann's area 46) FTD patients and controls subjects was provided by NeuroBioBank of National Institutes of Health. A total of 6 female (3 control/3FTD) and 8 male (4 control/4 FTD) samples were used, and samples were age and sex matched. Ages ranged from 60-77 with the average age of all samples being 70. More detailed information on specific samples can be found in **supplemental table 1**.

Immunoblotting

Human and mouse protein was extracted as previously described[17]. Brain tissue was homogenized with either tissue-tearor (Biospec products, Inc) or Pellet pestle motor (Kimble Kontes) in 25 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Triton X-100 and samples were incubated on ice 15-30 min and vortexed at full speed for 30 s every 5 min. Additional SDS was added to samples to bring total concentration up to 2%. Mouse protein was heated at 100°C for 5 min. Samples were centrifuged at 4°C, 13,000 g for 20 min and the supernatant was collected. Protein concentration determined using the BCA protein quantification assay (Pierce).

Immunoblotting was conducted as previously described[17]. Human and mouse protein samples were separated on 4–20% SDS Criterion TGX Stain-Free gels (BioRad) at 80 V. Proteins were transferred to a nitrocellulose membrane (BioRad 0.2 µm) in Tris–Glycine Buffer (25 nM Tris-Base, 200 mM Glycine) at 25 V for 30 min in Trans Turbo Transfer system (BioRad). The membrane was blocked with 5% non-fat dry milk in TBS containing 0.5% Tween-20 (TBST) for 1 h at room temperature. Membrane was incubated in TBST containing 2.5% milk overnight at 4 °C. The next day, membranes were incubated in secondary antibody (Amersham ECL HRP Conjugated Antibodies 1:5000) in TBST containing 2.5% milk) for 1 h at room temperature. Followed by band detection using SuperSignal Chemiluminiscent substrate (Thermo Scientific) and GE ImageQuant Las4000 mulit-mode imager. Primary antibodies and their concentrations are as follows: CK2α' (1:2000; Rabbit, Novus NB100-379) and GAPDH (1:5000; Mouse, Santa cruz sc-365062).

Cell culture: Culturing, transfections, and immunocytochemistry

Neuro-2a cells (N2a; ATCC CCL-131) were originally obtained from Dr. Veena Prahlad (University of Iowa). Cells were cultured at 37ºC 5% CO_2 in Dulbecco's modified Eagle's medium (DMEM, Genesee) supplemented with 10% fetal bovine serum (FBS), 100 U ml^{-1} penicillin/streptomycin as previously described[20,41,42]. For silencing experiments cells were simultaneously plated in 12-well plates for RNA-extraction or 24-well plates containing Matrigel matrix (Corning) coated glass coverslips for immunocytochemistry at a density of 80,000 cells/cm^2.

24 hours after plating all cells were transfected with siRNA for CK2α (Qiagen SI02007180, Rn_RGD:621663_4 FlexiTube siRNA), CK2α' (Qiagen SI00961072 Mm_Csnk2a2_4 Flexitube siRNA, Qiagen SI00961051 Mm_Csnk2a2_1 FlexiTube siRNA, Qiagen SI00961058 Mm_Csnk2a2_2 FlexiTube siRNA, and Qiagen SI00961065 Mm_Csnk2a2_3 FlexiTube siRNA pooled) or AllStars negative scrRNA control (Qiagen S103650318) using DharmaFECT 1

(Horizon discovery) at a final concentration of 25 nM in cell medium. 24 hours after siRNA
transfection cells were transfected with either pEGFP-C1 plasmid (Clontech) or prk5-Tau-
P301L-EGFP (Addgene #46908)[43] using jetOptimus transfection reagents (Polyplus). 24 hours
after transient transfection RNA was extracted as described below and cells were fixed with 4%
PFA for immunocytochemistry.

For immunocytochemistry cells were blocked in 5% NGS in TBST (0.3% triton x-100 in TBS) for
1 hour, then incubated in primary antibody diluted in overnight 5% NGS in TBST (0.3% triton x-
100 in TBS) (pTau (Ser202, Thr205) AT8 (1:500; Mouse Invitrogen Mn1020)). Secondary Alexa-
fluorophore-conjugated antibodies (Invitrogen) were added (1:200 in TBST with 5% NGS) for 1
h at room temperature. Coverslips were mounted in ProLong Gold Antifade with DAPI
(Invitrogen) and subsequently imaged.

Immunohistochemistry

Immunohistochemistry experiments were conducted as previously described[17,44,45], and stained
with either fluorescent secondaries or carried through 3,3'-diaminobenzidine (DAB) staining.
Mice were anesthetized with Avertin (250 mg/kg Tribromoethanol) and perfused intracardially
with tris-buffered saline (TBS) (25 mM Tris-base, 135 mM Nacl, 3 mM KCl, pH 7.6)
supplemented with 7.5 mM heparin. Brains were dissected, fixed with 4% PFA in TBS at 4°C for
4–5 days, cryoprotected with 30% sucrose in TBS for 4–5 days and embedded in a 2:1 mixture
of 30% sucrose in TBS:OCT (Tissue-Tek). Brains were cryo-sectioned into 16 µm-thick coronal
slices and stored in a 50% glycerol-50% TBS (25 mM Tris-base, 125 mM NaCl, 3 mM KCl, pH
7.6) solution at -20°C. For each experiment, three hippocampal containing sections were used
per mouse.

For immunofluorescent staining, sections were blocked in 5% normal goat serum (NGS) in
TBST for 1 h at room temperature. Primary antibodies were incubated overnight at 4°C in TBST
containing 5% NGS. Secondary Alexa-fluorophore-conjugated antibodies (Invitrogen) were
added (1:200 in TBST with 5% NGS) for 1 h at room temperature. Slides were mounted in
ProLong Gold Antifade with DAPI (Invitrogen) and subsequently imaged. Primary antibodies
used and dilutions are as follows: GFAP (1:2000; Chicken Millipore Sigma AB5541), Iba1
(1:500; Rabbit Fujifilm Wako 019-19741), Iba1 (1:500, Goat Fujifilm Wako 011-27991), NeuN
(1:1000; Mouse Millipore MAB377), PSD95 (1:500; Rabbit Invitrogen 51-6900) and Vglut1
(1:500; Guinea pig Millipore AB5905). Imaging of immunofluorescent slices was conducted
using a confocal microscope (Stellaris, Lecia). For NeuN, Iba1, and GFAP imaging 20x tiles
were acquired with a 16um z-dimension and 1um step size. For PSD95 and VGlut1 images
were acquired as described for synapse analysis.

For DAB staining antigen retrieval was performed using either Tris-EDTA pH=9.0 (Biolegend
422703) or Rodent decloaker (Biocare medical RD913) at 80°C for 30 minutes. Sections were
blocked at room temperature in TBST containing 10% NGS for 1 hour, then incubated in primary
antibodies overnight at 4C in TBST containing 5% NGS. Sections were incubated in biotin-
conjugated secondaries (Jackson Immuno Research Labs Biotin-SP (long spacer) AffiniPure) in
TBST containing 5% NGS for 1 hour, then blocked in 3% hydrogen peroxide for 20 minutes.
Sections were incubated in tertiary antibodies (VECTASTAIN Elite ABC, HRPS kit, Vector

laboratories) in TBST containing 5% NGS for 1 hour per manufacturer instructions. Sections
were then incubated in DAB chromogen diluted in DAB substrate buffer (Biolegend) for sufficient
staining. For counterstaining sections were incubated in 0.1% Cresyl violet for 10-12 minutes at
37ºC. Slides were then dehydrated and cleared in a series of ethanol and xylene washes before
being mounted in Permount mounting medium (Fisher scientific). Primary antibodies used and
dilutions are as follows: pTau (Ser202, Thr205) AT8 (1:1000; Mouse, Invitrogen Mn1020) and
CD68 [RM1031] (1:2500; Rabbit, Abcam ab303565). DAB images were taken at 10x on a hybrid
microscope using the brightfield settings (Echo Revolve) and at 40x on the EasyScan One slide
scanner (Motic).

in situ **hybridization**

Tissues were collected as previously described in the immunohistochemistry procedure. Slices
were mounted on superfrost plus glass slices using PBS and stored at -80ºC overnight before *in*
situ hybridization. RNAscope was performed according to the ACDbio manufacturer's protocol
using a custom probe for *Csnk2a2*. Immediately following RNAScope protocol, the slides were
blocked with 10% Normal Goat Serum (NGS) in Co-detection antibody diluent (CDD) for 1 hour.
Anti-IBA1 (Rabbit, Fujifilm Wako 019-19741) was diluted at 1:200 in 5% NGS with CDD and
slides were incubated overnight. The next day secondary Alexa-fluorophore-conjugated
antibodies (Invitrogen) were added (1:40 in CDD with 5% NGS) for 2h at room temperature.
Slides were mounted in ProLong Gold Antifade with DAPI (Invitrogen) and subsequently
imaged.

Tau pathology type

Tau pathology type analysis was conducted by 3 independent and blinded investigators. Tau
pathologies were defined based on previous publications[46-49] and based on the pattern of AT8+
staining. Briefly as described by Shi et al 2017[46] type 1; displays mossy fiber and sparse and
diffuse cell body staining in the dentate gyrus (DG), type 2; displays compact dense tangle like
cell body staining in the DG and CA3 with some sparse CA1 cell body staining, type 3; staining
is seen in the neuropil of the stratum radiatum of the CA1 staining along with staining of
dendrites from pyramidal neurons and sparse cell body staining, type 4; dense fragmented,
dotted, and grainy staining is observed all over the hippocampus. The percentage of pathology
for the cohort for each type was determined by the following calculation; ((number of mice with
pathology type)/(total number of mice in cohort)) x 100.

Image quantification

All image quantifications were conducted using either QuPath[50] or Image J (FIJI)[51] and were
blinded to the experimenter. For all quantifications images names were either blinded in FIJI
using the BlindAnalysis plugin or in Qupath using the automated image masking tool.
Quantifications were performed with at least two blinded experimenters.

Quantification of NeuN was done automatically using QuPath cell detection. Identical ROIs were
applied to each image for all 3 regions of the hippocampus. The cell detection tool was used to
automatically detect NeuN+ cells based on identical thresholding settings applied to all images.
Quantification of Iba1+ and GFAP+ cell counts was conducted as we previously described[44,45].

An average projection of 20x tiled z-stack confocal image was generated in FIJI and used for
counting. Two consistent ROIs (~3900 um^2) were drawn within the CA1, CA3, and DG for counts
and the cell counter FIJI plugin was used to count cells manually. A cell was determined to be
Iba1+ or GFAP+ based on the presence of DAPI within the stained area. Results were averaged
and normalized per mm^2 and reported as animal averages (2 ROIs/slice; 3 slices/animal).
Quantification of DAB positive percent area was quantified using either the Colordevonvolution2
plugin in FIJI[52] or in QuPath. Briefly a threshold of positive DAB staining was determined and
applied to all images to then calculate the percentage positive area.

Quantification of PSD95+ puncta within Iba1+ microglia were performed using Imaris software
(Bitplane). A confocal microscope (Stellaris, Lecia) was used to acquire images in the CA1
stratum radiatum region of the hippocampus with a 40x objective and 3x zoom factor with 0.34
size z-steps for a total of 15 steps. Lecia Stellaris software lighting processing was applied
before analysis. Images were then 3D rendered in Imaris and cropped to isolate individual Iba1+
cells (2 per slice). Iba1+ surfaces were generated and used to mask PSD95+ signal. The
masked PSD95+ signal was then used to create spots, to ensure it was located within Iba1+
signal. The number of spots within the Iba1+ surfaces per cell soma defined as the region where
Iba1+ signal colocalized with DAPI and within the whole Iba1+ cell surface were counted and
reported as a number.

Synapse analysis

Synapse analysis was conducted as previously described[17,45,53-55]. Immunohistochemical
staining was conducted as described above with Vglut1 and PSD95, with blocking increased to
20% NGS+TBST for 2 hours and primary and secondary antibodies diluted in 10% NGS+TBST.
Fluorescent images from the molecular layer of the CA1 were taken on a confocal microscope
(Stellaris, Lecia) at 63x with z-step dimension of 0.34 μm with 15 steps were generated. Lecia
stellaris software lightning processing was applied to images post-acquisition and edited images
were used for analysis. Maximum projections of three sections per slice were generated. Puncta
analyses were conducted blinded using the PunctaAnalyzer Plugin (Durham,NC,USA) in FIJI as
previously described[17,45,53-55]. Data is represented as slice averages with three slices per animal.

Electrophysiology

Mice were anesthetized with isoflurane (2%) and decapitated for slice preparation. After
decapitation, the whole brain, containing the two hippocampi, was removed into ice-cold
solution consisting of: sucrose 189 mM, glucose 10 mM, NaHCO3 26 mM, KCl 3 mM, MgSO4 5
mM, CaCl2 0.1 mM, NaH2PO4 1.25 mM. Brains were positioned on the stage of a vibratome
slicer and cut to obtain transverse hippocampal slices (350 μm thick; LEICA VT1000S). Slices
were incubated at room temperature in artificial cerebrospinal fluid (aCSF) continuously
oxygenated for 45 min-1 h before use. aCSF contained NaCl 124 mM, KCl 5 mM, NaH2PO4
1.25 mM, MgSO4 2 mM, NaHCO3 26 mM, CaCl2 2 mM, glucose 10 mM; gassed with 95% O2,
5% CO2. The pH was adjusted to 7.4 with NaOH. All experiments were carried out at 30–34°C,
during which the slices were continuously perfused with solution.
Field excitatory postsynaptic potentials (fEPSPs) were recorded in the CA1 region of the
hippocampus. fEPSPs in the CA1 region of the hippocampus were evoked with a stimulating
electrode placed on the Schaffer collateral (0.33 Hz) using brief current pulses (200 μs, 0.1–0.2

mA). Extracellular recording electrodes were filled with aCSF. Stimulation was adjusted to
obtain a fEPSP peak amplitude of approximately 1 mV during control conditions. After a stable
fEPSP baseline period of 10 min, LTP was induced by a Theta-burst stimulation (TBS)
consisting of a series of 10 bursts of 5 stimuli (100 Hz within the burst, 200 ms interburst
interval), which was repeated 4 times (5 s apart). Data were filtered at 3 kHz and acquired at 10
kHz using pCLAMP 10.2 software (Molecular Devices, RRID: SCR_011323).
A stimulus-response curve (0.05-0.4 mV, mean of five fEPSPs at each stimulation strength) was
compiled for the different mice used. For paired-pulse ratio experiments, two fEPSPs were
evoked 40 ms apart for 0.5 min at baseline frequency (6 times) at the beginning of the baseline
recording. The PPR was expressed as the amplitude of the second fEPSP divided by the
amplitude of the first fEPSP.
297
Behavior

Behavioral testing was performed with the support and guidance of the Mouse Behavior Core at
University of Minnesota. Mice were handled daily for a minimum of one week prior to all
behavioral testing. Mice were habituated to behavior room for a minimum of 30 minutes each
day prior to beginning of task. All tasks were recorded and tracked using ANYmaze software
(Stoelting Co., Wood Dale, Illinois) and overhead cameras.

Barnes maze was conducted on a 20 hole Barnes Maze (San Diego Instruments) divided into 4
quadrants and a center zone under 450-500lux light intensity, with black and white visual cues
placed around the room, as similarly described[56-58]. Mice were placed in the center under a
covering with lights off, lights were turned on and covering lifted within 30 seconds. The training
phase consisted of four 3-minute trials per mouse for 5 days, with an inter trial interval of
approximately 20-30 minutes. The maze was thoroughly cleaned with 70% between all mice
and rotated between trials to avoid olfactory cues. Mice were allowed to explore the maze and
primary latency, or the time to first explore the escape hole, was recorded. The trial either
concluded when the mouse entered the escape hole, or at 3 minutes at which point the mouse
was gently guided to the escape box. For the probe day the escape hole was covered with an
identical covering used on all other holes, mouse was placed on the maze in the same manner
as training. The time spent in each maze zone was then recorded. Spatial strategy was
determined by a blinded investigator.

Open field was conducted in 40cmx40cm boxes with a 150-200lux overhead light. Mice were
placed in boxes and behavior was recorded for 1 hour. Boxes were thoroughly cleaned between
mice with 70% ethanol.

RNA extraction and qPCR

RNA was extracted from cells and mouse striatal tissues by using the RNeasy extraction kit
(Qiagen) according to the manufacturer's instructions. cDNA was prepared using the
Superscript First Strand Synthesis System (Invitrogen). SYBR green based PCR was performed
with SYBR mix (Roche). The qPCR amplification was performed using the LightCycler 480
System (Roche). Each sample was tested in triplicate and normalized to GAPDH levels.

For qPCR the following primers were used: CK2α' **FWD-**CGACTGATTGATTGGGGTCT **REV-**
AGAATGGCTCCTTTCGGAAT; CK2α **FWD-**TCCCCATGCTGTGACAATAA **REV-**
AAGACCCTGTGTCACGAACC; CK2β **FWD-** AGTCCTCCAGACACCACCAC **REV-**
GACTGGGCTCTTGAAGTTGC; huTau **FWD-** GCTGGCCTGAAAGCTGAAGA **REV-**
CGTTTTACCATCAGCCCCT.

RNA-Seq Analysis

RNA was extracted as previously described and RNA-Sequencing was conducted by Novogene
(Sacramento, CA). Gene expression analysis was carried out using the CHURP pipeline
(https://doi-org.ezp2.lib.umn.edu/10.1145/3332186.3333156) using n=4-7 mice/genotype with
female(F)/male(M) ratios as follows; 4 WT (2F/2M), 5 CK2α'$^{(+/-)}$ (2F/3M), PS19 (3F/4M) (3 high
pathology and 4 low pathology), and 4 PS19;CK2α'$^{(+/-)}$ (3F/1M). Differential gene expression was
determined with DESeq2 using default setting[59] (v1.46.0). Genes with a FDR $<=$ 0.05 were
considered significant. Outliers' identification was performed using Cook's distance (DESeq2).
Driver factors of gene expression variance (genotype and/or sex) were evaluated using R
(v4.4.2) package variancePartition (1.36.3). Pathway and clustering analysis were completed
gProfiler2[60] (v0.2.3) and clusterProfiler (v4.14.6). Data visualization was done using various R
graphic packages, including ggplot2, ggraph, and DESeq2 visualization functions. The RNA-seq
data set generated in this manuscript has been deposited at GEO (accession number
GSE298505).

Microglia morphology

Microglia morphology measurements were adopted from previously described protocols[61].
Slices were stained with Iba1+ as described in the immunohistochemistry methods. Nine total
cells were imaged per animal across 3 slices with 3 cells per slice. Images were taken on a
confocal microscope (Stellaris, Lecia) at 40x magnification with a zoom factor of 2.5. Z-stacks
were acquired with a Z-step size of 0.3 µm, covering the entire slice. Maximum projection of z-
stacks were generated via FIJI for analysis. For image analysis, brightness and contrast were
adjusted to subtract background noise before converting the image into a binary black-and-
white version of the original. The paintbrush and despeckle functions were used to remove any
additional noise before applying the skeletonizing function. Finally, the AnalyzeSkeleton plugin[62]
was used to quantify the resulting skeleton image.

Cytokine proteome profiling

The Proteome Profiler Mouse Cytokine Array Panel (ARY006, R&D Systems) was used to
detect the levels of cytokine/chemokine as per manufacturer's instructions. Frozen hippocampi
from 9 months old PS19 and PS19;CK2α'$^{(+/-)}$ mice were homogenized in PBS containing Halt
protease inhibitor cocktail and phosphatase inhibitors (Fisher Scientific) and 1% triton X-100
(Sigma). Samples were stored at -80°C for 15 min, thawed and centrifuged at 10,000 × g for 5
min to remove cell debris. 300 µg of protein was applied to each membrane and the
manufacturer's protocol was followed. A total of n=4 mice/genotype with a female(F)/male(M)
ratio (2F/2M PS19 and 2F/2M PS19;CK2α'$^{(+/-)}$) were analyzed. Imaging was conducted as
described for immunoblotting. Spot intensity was quantified using FIJI software and a batch

correction was applied to normalize data. Membrane spots corresponding to 40 different
cytokines or chemokines were measured as outlined in manufacturer protocol.

Statistical analyses and data representation

For electrophysiology experiments data were analyzed using Clampfit 10.2 software (pCLAMP,
Molecular Devices, RRID: SCR_011323). Data are presented as mean ± SEM. To estimate
changes in synaptic efficacy, STP was quantified by comparing the mean fEPSP amplitude
during 1 minute after the application of the TBS. LTP was quantified as for STP but 40 minutes
after the protocol. Graphs were obtained using SigmaPlot 14.0. Before applying any statistical
comparison, the data were subjected to Shapiro- Wilk normality and equal variance tests. For
any comparisons between two groups, two- paired Student's t-test was used. For multiple
comparisons to the same control, one- way ANOVA and Holm-Sidak test was used. P-values
less than 0.05 were considered statistically significant.
378
For all other experiments GraphPad Prism (GraphPad, San Diego, CA, USA) was used to
create graphs and conduct statical testing. For all other experiments data is represented as
mean $\pm$ SEM. Statistical analyses were performed using paired Student's t-test, unpaired
student's t-test, one-way or two-way ANOVA with post-hoc analysis, as indicated in each figure
legend. A p-value <0.05 was considered significant.

Results

Levels of Protein kinase CK2α' are increased in the brains of dementia patients and
mouse models of tauopathy

There are two different genes (*Csnk2a1* and *Csnk2a2*) that code for two different CK2 catalytic
subunits CK2α and CK2α', respectively. These two catalytic subunits share a high percent of
homology and they can arrange in various combinations (α-α, α-α' or α'-α') depending on their
expression and tissue availability although the canonical arrangement in the brain is expected to
be α-α' (**Fig. 1A, B**)[16,28-34]. To determine whether CK2 catalytic subunits are altered in patients
with dementia we utilized the RNA-seq data from the Allen Brain Institute: Aging, Dementia, and
Traumatic Brain Injury (TBI) Study. In this dataset, three different regions were available:
parietal cortex (PCx), temporal cortex (TCx) and hippocampus (HIP). Analyzing this dataset [63]
we found a specific increase in the expression of *CSNK2a2,* encoding for CK2α', but not in
CSNK2a1, encoding for CK2α, in patients with dementia compared to non-dementia controls in
the PCx (**Fig. 1C-F**). We also found a trend towards increased CK2α' in the TCx and HIP
although data did not reach statistical significance (**Fig. 1C, G, H**). Importantly, we also
observed a significant increase in CK2α' protein levels in the frontal cortex of FTD patients
compared to age and sex-match controls (**Fig.1I, J, Supp. Table 1**). Enhanced levels of CK2α'
also associated with the presence of tau pathology previously assessed in those samples using
the AT8 antibody (**Fig. 1K, Supp. Table 1**). The majority of FTD samples grouped in the top left
quadrant representing high levels of CK2α' and the presence of tau pathology, while most
control samples grouped in the bottom right quadrant representing low levels of CK2α' and

absence of tau pathology. For those two control samples (#1 and #6) in which tau pathology
was observed, the AT8 phospho-tau (pTau) signal and distribution differed from the canonical
tau pathology associated with dementia and individuals lacked classical cognitive decline
(**Supp. Table 1**).

To examine whether CK2α' was also elevated in mouse models of tauopathy we utilized the
PS19 mouse model, which expresses human tau with a P301S mutation associated with familial
forms of FTD and other tauopathies[40]. Pathology and neuronal loss in PS19 mice are seen
largely in the hippocampus, although it does spread to other brain regions including the
neocortex, entorhinal cortex and amygdala[40,64]. We conducted RNA in situ hybridization for
CK2α' in the hippocampus of PS19 mice at a symptomatic age (10-12 months), defined as the
age at which animals present overt tau pathology and cognitive decline[40,65,66]. We observed a
significant increase in CK2α' in all regions of the hippocampus specially in the pyramidal and
granular layers, that colocalized with NeuN staining, pointing to enhanced expression in
neurons (**Fig. 2A-D**). Expression outside the neuronal layer was also observed which
associated with enhanced CK2α' levels in the microglia (Iba1+) of PS19 mice compared to WT
(**Fig. 2E-G**). Single cell RNA-Seq studies available in The Alzheimer's Cell Atlas (TACA)[67-71]
confirmed increased CSNK2a2 expression in patients with AD in both microglia and neurons in
the occipital cortex (OL) and entorhinal cortex (EC) respectively (**Supp. Fig. 1**). However,
studies in prefrontal cortex (PFC) and frontal lobes (SFG) revealed enhanced expression of
CSNK2a2 in cells other than neurons and microglia, suggesting a cell-specific upregulation of
CSNK2a2 that seems to be brain region-dependent. Overall, these results demonstrate a
specific upregulation of CK2α' in brain affected regions of patients with dementia and in
hippocampal neurons and microglia of mouse models of tauopathy.

Genetic depletion of Protein kinase CK2α' impacts tau pathology *in vitro* and *in vivo*

Previous studies have shown that CK2 holoenzyme regulates tau phosphorylation in various cell
and mouse models of AD[19,72], but whether CK2α' subunit specifically contributes to this
phenomenon is unknown. To explore whether CK2α' has any role in the regulation of tau
phosphorylation and pathology we first explored the impact of silencing CK2α or CK2α' in
Neuro2a (N2a) cells expressing a pathological tau variant (Tau-P301L). N2a cells were treated
with either scrambled RNA (ScrRNA) or silencing RNA for CK2α (siCK2α) or CK2α' (siCK2α'),
and transfected with either Tau-P301L-EGFP[43] or EGFP empty vector (control) (**Fig. 3A**). The
AT8 phospho-tau (pTau) antibody was then used to examine the impact of silencing CK2α and
CK2α' on tau pathology (**Fig. 3B**). First, we confirmed the efficacy of the siRNAs decreasing the
expression of CK2α or CK2α' without altering the other subunits (**Fig. 3C-E**) and the increased
expression of Tau in N2A transfected cells (**Fig. 3F**). As expected, the levels of AT8 increased
upon expression of Tau-P301L in N2A control cells (**Fig. 3B, G**). Importantly, AT8 levels
decreased in Tau-P301L transfected cells treated with siCK2α' (p=0.0130) but not siCK2α
relative to Tau-P301L ScrRNA condition (p=0.9722) (**Fig. 3B, G**). Our data demonstrates an
impact of the CK2α' subunit specifically on pTau accumulation *in vitro* (**Fig. 3H**).

We then generated a PS19 mouse haploinsufficient for CK2α' (PS19;CK2α'[(+/-)])(**Supp. Fig. 2A-**
F), to assess the impact of depleting CK2α' on survival, tau pathology and symptomatology *in*
vivo. We then assessed tau phosphorylation in the hippocampus and overlaying cortex of mice

at a prodromal stage (~7 moths) and at a late symptomatic age (~10-12 moths)[40] (**Fig. 4A-B,**
Supp. Fig. 3A-B). In the prodromal age no significant differences were observed in the
accumulation of AT8 in PS19 and PS19;CK2α'[(+/-)] in all three examined regions of the
hippocampus (CA1, CA3, and DG) and overlaying cortex (**Fig. 4C-E, Supp Fig. 3C**). As
expected, in the late symptomatic groups we observed a significant increase in AT8+ area in
both the PS19 and PS19;CK2α'[(+/-)] mice in the CA1, CA3, and DG and cortex relative to all other
groups (**Fig. 4C-E, Supp. Fig. 3C**). However, PS19;CK2α'[(+/-)] mice showed a significant
decrease in AT8 signal compared to PS19 in the CA1, DG, and cortex (CA1; p=0.0075, DG;
p=0.0186 and Cortex; p<0.0001) (**Fig. 4C,E, Supp. Fig. 3C**) suggesting an overall decrease in
pTau upon depletion of CK2α'.

Tau pathology has been previously reported to follow a characteristic progressive pattern of
deposition in the hippocampus that can be categorized into 4 pathology types[46-49] (**Fig. 4F**).
These pathology types have been shown to correlate with hippocampal volume[46,49] and offer a
more holistic pathological assessment. We observed differential distributions of patterns
between PS19 and PS19;CK2α'[(+/-)] mice with a more dramatic alteration in late symptomatic
stages where nearly all PS19 mice presented a type 4 pathology in contrast to the
PS19;CK2α'[(+/-)] that presented a variation in types with almost 50% of mice presenting type 2
pathology (prodromal p=0.11, and symptomatic p=0.0587) (**Fig. 4G**).

Interestingly, although CK2α' seemed to play a clear role in modifying the progression of tau
pathology in PS19 mice, no significant differences in survival were observed between PS19 and
PS19;CK2α'(+/-) animals, with both groups showing approximately 50% mortality by 50 weeks
of age (**Supp. Fig. 2G**). Survival is a relatively coarse endpoint, influenced by the dysfunction of
multiple neural and systemic pathways like the spinal cord and brain stem[40,73]. While CK2α'
depletion clearly impacted tau phosphorylation and tau burden in the hippocampus and cortex,
these effects were insufficient to alter the overall lifespan of this aggressive model. Nonetheless,
the observed modulation of tau pathology underscored a critical role for CK2α' in specific
aspects of tau-mediated neurodegeneration. We therefore decided to further explore the
molecular and cellular changes influenced by CK2α' modulation in PS19 mice.

CK2α' happloinsufficiency improved hippocampal synaptic density and synaptic function
in PS19 mice

We conducted NeuN immunostaining analyses in the hippocampus of PS19 and PS19;CK2α'[(+/-)]
mice to determine if depletion of CK2α' could impact the overall neuronal loss characteristic of
PS19 mice[40,74-76]. Significant depletion of NeuN+ cells and hippocampal atrophy was only seen
in PS19 groups at a symptomatic age (**Fig. 5A, B, Supp. Fig. 4A-C**). An ~30-50% reduction in
the number of NeuN counts is often reported at late symptomatic ages in PS19 mice[40,74-76].
However, we noticed a great variability in the number of NeuN+ cells and hippocampal area
among PS19 littermates at this disease stage that was not related to sex (**Supp. Fig. 4D-F**).
Wide variability in tau pathology and neuronal degeneration within this mouse model has been
previously reported and associated to nuances in the genetics of this model[77]. Therefore, we
categorized PS19 and PS19;CK2α'[(+/-)] animals presenting NeuN counts within 10% of the WT as

"abnormally low pathology" and as a result 3 PS19 animals and 1 PS19;CK2α'(+/-) were excluded
from subsequent analyses. Analyses in curated groups of PS19 and PS19;CK2α'(+/-) mice
revealed a significant increase in the number of NeuN+ cells in PS19;CK2α'(+/-) mice compared
to PS19 in the CA1 (**Fig. 5A, B, Supp. Fig. 4A-C**).

We then evaluated if total synapse density was also positively altered in the CA1. We conducted
immunostaining and colocalization analyses for the vesicular glutamate transporter 1 (VGlut1)
(pre-synaptic marker of excitatory synapses), and the postsynaptic density protein 95 (PSD95)
(post-synaptic marker) (**Fig. 5C-I**). As expected, PS19 mice showed a significant decrease in
the number of synapses (colocalized puncta VGlut1-PSD95) compared to WT mice (**Fig. 5F, I**).
The reduction in colocalized puncta in PS19 mice was due to a concomitant reduction in both
PSD95 and VGlut1 markers (**Fig. 5D, E, G, H**). In PS19;CK2α'(+/-) mice we found that although
there is not a statistical significance compared to PS19 mice, CK2α' haploinsufficiency
decreased the % loss of both PSD95 (**Fig. 5D, G**) and VGlut1 (**Fig. 5E, H**). This resulted in the
amelioration of total synapse loss in PS19;CK2α'(+/-) mice and obtained values for colocalization
were no longer significant respect to WT (**Fig. 5F, I**). These results demonstrate a partial rescue
in the density of excitatory synapses in the CA1 of PS19;CK2α'(+/-) mice aligning with the
amelioration of hippocampal atrophy observed in this region (**Fig. 5B**).

To investigate functional implications of the rescue seen in CA1 hippocampal atrophy and
synaptic density we assessed hippocampal synaptic plasticity by performing extracellular field
excitatory postsynaptic potential (fEPSP) recordings in the CA1 region of acute hippocampal
slices (**Fig. 5J, K**). Theta-burst stimulation (TBS) at the Schaffer collaterals induced short-term
potentiation (STP), which resulted in a significant increase in synaptic efficacy at CA1 synapses
in both WT and CK2α'(+/-) mice, as evidenced by the enhanced fEPSP amplitude (170 ±13%, n =
6; 170 ± 12%, n = 7) (**Fig. 5L, M**). In contrast, the increase in fEPSP amplitude was significantly
reduced in PS19 and PS19;CK2α'(+/-) mice relative to WT controls post TBS (123 ± 4%, n = 7;
128 ± 7%,n = 6; **Fig. 5L, M**), indicating impaired STP in these models. Furthermore, we
examined long-term potentiation (LTP) at CA3-CA1 synapses, a form of synaptic plasticity
linked to learning and memory. It has been previously shown that PS19 mice present impaired
LTP in the CA1-CA3 pathway[40,78,79]. First, we observed significant differences in input-output
curves at CA3-CA1 synapses between PS19 and PS19;CK2α'(+/-) mice when compared to WT,
which suggests that excitatory synaptic input properties were altered in these models (**Fig. 5N**).
TBS at the Schaffer collaterals induced robust LTP in both WT (173 ± 12%, n = 6) and CK2α'(+/-)
(164 ± 9%, n = 7; **Fig. 5L, O**). However, in the PS19 mouse model, the magnitude of LTP was
significantly diminished compared to WT (128 ± 9%, n = 7; vs. WT p=0.007) (**Fig. 5L, O**),
suggesting a disruption in the molecular mechanisms underlying LTP. Interestingly, in the
PS19;CK2α'(+/-) mice, LTP was partially restored (150 ± 6%, n = 6; vs. WT p=0.199) (**Fig. 5L, O**),
pointing to a potential compensatory mechanism that partially mitigates the LTP deficits
observed in the PS19 mice. Importantly, paired-pulse facilitation (PPF) was unaltered across all
experimental groups, suggesting that the observed changes in synaptic plasticity were most
likely due to postsynaptic mechanisms dependent on the hippocampus (**Supp**. **Fig. 5**).

CK2α' haploinsufficiency improved PS19 behavior on Barnes Maze

We then tested whether the partial rescue observed in PS19;CK2α'$^{(+/-)}$ mice associated to
synaptic density and functional LTP along with the decrease in Tau pathology had any impact in
cognitive behaviors. Mice were trained on the Barnes maze task for 5 consecutive days on the
location of a target hole (training) and on the 6th day a probe trial was conducted with the target
hole removed (**Fig. 6A**). During the training trials the primary latency or the time to first locate
the escape hole was recorded (**Fig. 6B-C**). All mice in the prodromal group demonstrated an
improvement on time needed to locate the escape hole from day 1 vs day 5 (WT p=0.0272;
PS19 p=0.0.0067; CK2α'$^{(+/-)}$ p=0.0002; PS19;CK2α'$^{(+/-)}$ p<0.0001) with no significant difference
between the genotypes (**Fig. 6B**). All mice in the symptomatic group also showed a capability of
acquiring the task demonstrating a significant improvement from day 1 vs day 5 (WT p<0.0001;
PS19 p=0.0109; CK2α'$^{(+/-)}$ p<0.0001; PS19;CK2α'$^{(+/-)}$ p=0.0037). However, PS19 and
PS19;CK2α'$^{(+/-)}$ showed a significant impairment compared to WT, starting on day 3 for PS19
(p=0.0009) and day 4 for PS19;CK2α'$^{(+/-)}$ (p=0.0306) (**Fig. 6C**). The training data suggested that
both symptomatic PS19 and PS19;CK2α'$^{(+/-)}$ mice showed similar impaired learning on the
Barnes Maze compared to WT mice, although deficits in PS19;CK2α'$^{(+/-)}$ were attenuated.

Following the training phase of the Barnes maze we conducted a probe phase in which the
escape hole was covered and the time spent exploring the escape hole quadrant was recorded.
At the prodromal age there was no significant difference in time spent in the escape hole
quadrant for any genotype (**Fig. 6D**). At the symptomatic age, the PS19 mice showed a
significant impairment in the time spent in the goal quadrant compared to WT (p<0.0001) (**Fig.
6D**). Importantly, the PS19;CK2α'$^{(+/-)}$ mice spent significantly more time than the PS19 mice in
the goal quadrant with no significant difference from WT (vs. PS19 p=0.0113 vs. WT p=0.2335)
demonstrating a rescue in spatial memory for the PS19;CK2α'$^{(+/-)}$ on the probe phase of the
Barnes maze (**Fig. 6D**).

The PS19 mouse model is known to present motor deficits due to hindlimb paralysis and
weakness[40], which could confound Barnes Maze results if PS19 mice perform more poorly due
to less distance traveled. Prodromal mice displayed no difference in distance traveled on
Barnes Maze (**Supp. Fig. 6A**). While the symptomatic PS19 mice traveled less distance in the
Barnes Maze compared to WT mice, they did not show a significant difference in distance
traveled compared to PS19;CK2α'$^{(+/-)}$ mice (**Supp Fig. 6B**). In addition, in open-field testing we
found no differences between groups at the prodromal time point (**Supp. Fig. 6C**). On the
contrary, symptomatic mice showed significant hyperactivity in the open field task,
demonstrating the ability of PS19 mice to move (**Supp Fig. 6D**). Indeed, no significant
differences were observed in the open field between PS19 and PS19;CK2α'$^{(+/-)}$ mice (**Supp Fig.
6D**). This data supports the improvement in cognitive abilities seen in the PS19;CK2α'$^{(+/-)}$ mice
are not due to changes in motor function but rather are related to an improvement in memory.

Mice can use different cognitive strategies to navigate the Barnes Maze and locate the escape
hole[57]. As the mouse learns the task they show improvement in the strategy used to locate the
escape hole[57] (**Fig. 6E**). Assessing these cognitive strategies is important to provide a deeper
insight into how memory and spatial learning are being utilized. All animals in the prodromal
group demonstrated a similar pattern in spatial strategies over the training phase (**Fig. 6F**) and
showed a predominance in the corrected and direct strategies in the probe phase (**Fig. 6G**).

Symptomatic PS19 and PS19;CK2α'(+/-) mice both showed impaired spatial strategies over the
training phase, by day 5 both genotypes largely displayed either failing, random or serial
strategies (**Fig. 6H**). However, during the shortened probe phase, PS19 displayed a worsened
overall strategy, with predominant failing, serial, and long correction strategies compared to
PS19;CK2α'(+/-) mice that displayed more serial, corrected and direct strategies (**Fig. 6I**).
Overall, the PS19;CK2α'(+/-) mice displayed improvement in the probe phase of the Barnes Maze
in both the time spent in the escape quadrant and strategy used to locate the target, indicating
improvements in spatial memory.

RNA-Seq analyses revealed CK2α' alters the expression of genes related to immune
response and synaptic function

To start interrogating the mechanisms by which CK2α' influences tau pathology and cognitive
decline in PS19 mice we conducted RNA-Seq analysis in the hippocampus of animals in the
prodromal phase. The reason we focused on this particular age group is because the prodromal
phase represents the initial stage of disease development that precedes overt symptoms.
During this time, sometimes subtle but significant molecular changes begin to occur, including
alterations in gene expression that precede the onset of full-blown pathology. By focusing on
this phase, we aimed to identify pathways involved in disease onset that may be missed if only
later stages of the disease are studied where significant neuronal loss is present. Our RNA-seq
analyses revealed a large variability in the expression pattern of PS19 mice that associated to
the variability observed in NeuN counts (**Supp. Fig. 4D-F**). By comparing the expression pattern
of the different animals along with the tau pathology we found two distinct expression profiles
that allowed the mice to segregate into a low (Low-PS19) and high (High-PS19) pathology
groups (**Supp. Fig. 7A**). These two different pathology groups showed differential gene
expression compared with WT from one another (**Supp. Fig. 7B-C**). With the Low-PS19 group
primarily differentially dysregulated genes (DGEs) related to developmental and organizational
pathways (**Supp. Fig. 7B**) and the High-PS19 showing DGEs related to immune processes
(**Supp. Fig. 7C**). Interestingly, the High-PS19 group was more similar in the gene expression
pattern to the PS19;CK2α'(+/-) group than the Low-PS19 group (**Supp. Fig. 7D-E**). Moreover, the
PS19;CK2α'(+/-) mice showed a more similar expression pattern among animals of the same
group. Moving forward we focused our comparisons on the High-PS19 group with
PS19;CK2α'(+/-).

Gene expression analyses in the High-PS19 cohort (referred hereafter as PS19) vs WT mice
identified 477 significant DGEs in the PS19 vs WT (**Fig. 7A**). Gene ontology pathway analysis
revealed that many of the DGEs belonging to the top dysregulated pathways corresponded to
immune and inflammatory functions (**Fig. 7B**), as previously reported[80,81]. Other important GO
pathways were associated with synaptic dysfunction where the top 5 impacted synaptic
pathways in PS19 mice corresponded to dysregulation in synapse assembly, regulation of
synapse organization, regulation of synapse structure or activity, post synapse organization and
synaptic pruning (**Fig. 7B-C**). Synaptic pruning was upregulated in PS19 vs WT, whereas the
other synaptic pathways related to organization and regulation of activity were largely
downregulated (**Fig. 7C**). These results aligned with the decreased synapse density and

impaired synaptic function seen in PS19 mice and indicated a potential early prodromal
transcriptional signature for these deficits.

Direct comparison of PS19 vs PS19;CK2α'$^{(+/-)}$ mice yield a small set of 26 significant DGEs (**Fig.**
7D). This small gene set returned no biologically connected pathways via gene ontology
analysis, but several of the DGEs were related to Apoptotic/phagocytic and
Immune/inflammatory functions (*Cept1, Ube2n, Oxct1, Exoc3, Adgrb1, Sh3glb2*, *Adgrdb1*). We
then assessed whether the manipulation of CK2α' had any overall effect on modules of genes
with similar biological functions despite not reaching the >Log2 criteria. Weighted Gene Co-
Expression Network Analysis (WGNCA) revealed 34 gene modules with 8 modules (red,
lightcyan, royalblue, skyblue, tan, lightyellow, turquoise, and black) showing statistical
significance detected via a Kruskal-Wallis test among our groups (p<0.05) (**Fig. 7E,F, Supp.**
Table 2). Interestingly several of these modules (red, tan, and black) showed a trend where the
PS19;CK2α'$^{(+/-)}$ mice more closely resembled the expression pattern of WT mice. On the
contrary, other modules showed a larger dysregulation in the PS19;CK2α'$^{(+/-)}$ compared to WT
and PS19, such as the lightcyan, lightyellow, and turquoise. Among the 8 significant gene
modules, we focused our analyses on the tan and turquoise modules for showing a different
expression pattern between PS19 and PS19;CK2α'$^{(+/-)}$ mice. The tan module composed on 215
genes (Kruskal-wallis p=0.016) (**Fig. 7F-G, Supp. Fig. 8A, Supp. Table 2**) was represented by
pathways associated with synaptic function and assembly, with top pathways being vesicle-
mediated transport in synapse, synaptic vesicle cycle, and regulation of synapse organization
(**Fig. 8G**). The overall expression of genes associated with these pathways presented a
depletion in the PS19 mice that was ameliorated in the PS19;CK2α'$^{(+/-)}$ group (**Fig. 7F**).
Changes in these pathways aligned with the improvements observed in NeuN abundance,
synapse density and function in PS19;CK2α'$^{(+/-)}$ mice (**Fig. 5**).

On the other hand, the turquoise module composed of 2765 genes (Kruskal-wallis p=0.0041)
displayed a dysregulation in the same direction in both PS19 and PS19;CK2α'$^{(+/-)}$ mice, with a
greater dysregulation in PS19;CK2α'$^{(+/-)}$ (**Fig. 7F,H, Supp. Fig. 8B, Supp. Table 2**). This module
was a large collection of genes including top significant pathways related to innate immune
response and immune response-activating signaling pathways (**Fig. 7H**). PS19;CK2α'$^{(+/-)}$ mice
showed a higher expression than PS19 of a large set genes associated with immune response
perhaps indicating an improved activation of immune systems.

To further explore DGEs specific to the PS19 and PS19;CK2α'$^{(+/-)}$ mice we focused specifically
on their comparisons to WT. We set criteria to yield two comparisons (*comparison 1*) significant
DGEs in PS19 vs WT but not PS19;CK2α'$^{(+/-)}$ vs WT and (*comparison 2*) significant DGEs in
PS19;CK2α'$^{(+/-)}$ vs WT but not PS19 vs WT. This more restrictive criteria then yielded two unique
set of genes (**Supp. Table 3**). Biological pathways associated with *comparison 1* included
regulation of neurogenesis, positive regulation of nervous system development and regulation of
synapse organization (**Supp. Fig. 8C**), similar to those pathways identified in the tan module.
Biological pathways associated with *comparison 2* included those associated with inflammatory
or immune responses (**Supp**. **Fig. 8D**). The majority of genes included in the top two biological
pathways were upregulated in PS19;CK2α'$^{(+/-)}$ mice and associated with immune response
(**Supp. Fig. 8D, E**). These results indicate there are more immune-related genes significantly

differentially expressed in the PS19;CK2α'$^{(+/-)}$ compared to WT than PS19 compared to WT

similar to what was revealed in the turquoise module (**Fig. 7H**). This was also confirmed by

gene set variation analysis (GSVA) activity scores of the dysregulated pathways within

comparison 2 showing increased activity of pathways associated with immune response relative

to WT (**Fig. 7I**). This analysis also revealed interesting alterations in apoptotic pathways. We

observed a decrease in activity in extrinsic apoptotic pathways and an increase in intrinsic

apoptotic pathways in the PS19;CK2α'$^{(+/-)}$ when comparing to PS19 mice (**Fig. 7I**). This

suggests a shift in the molecular mechanisms governing the apoptotic pathway towards a more

beneficial and selective apoptotic environment perhaps contributing to the beneficial outcomes

observed in PS19;CK2α'$^{(+/-)}$ mice. Overall, the RNA-Seq analyses revealed that CK2α'

haploinsufficiency caused alterations in networks of genes associated with important and

relevant biological pathways demonstrating the amelioration of synaptic dysfunction, the

enhanced activation of immune related pathways, and impacts on apoptotic signaling.

PS19 mice lacking CK2α' ameliorated microglia reactivity, neuroinflammation and

phagocytic activity

Considering the enhanced levels of CK2α' in microglia, the activation of pathways involved in

immune response upon deletion of CK2α', and the key role of microglia as CNS immune cells in

tauopathies and dementia[12,40,82-85], we examined the state of glial cells (astrocytes and

microglia) in the hippocampus in both the prodromal and symptomatic mice groups. First, we

observed a significant increase in the number of GFAP+ cells in the hippocampus of PS19 mice

in both age groups (**Supp. Fig. 9A-E**), as previously reported[40], but we did not find any

differences compared to PS19;CK2α'$^{(+/-)}$ mice (**Supp. Fig. 9C-E**). However, GFAP+ cells

significantly decreased in the PS19;CK2α'$^{(+/-)}$ compared to the PS19 mice in the overlaying

cortex of symptomatic mice (**Supp. Fig. 9F-H**). When looking at Iba1+ cells we observed a

modest but significant increase in Iba1+ positive cells throughout the hippocampus in the

prodromal PS19 mice compared to WT (**Supp. Fig. 10A,C-E**). In the symptomatic group, both

the PS19 and PS19;CK2α'$^{(+/-)}$ demonstrated a much greater increase in the number of Iba1+

cells compared to WT, but PS19;CK2α'$^{(+/-)}$ mice presented significantly less Iba1+ cells than

PS19 in both the CA1 (**Fig. 8A,B, Supp. Fig. 10B, C**) and CA3 (**Supp. Fig. 10B,E**). No

significant differences were observed among the PS19 gin the DG (**Supp. Fig. 10B, E**). We also

observed significantly less Iba1+ cells in the PS19;CK2α'$^{(+/-)}$ compared to PS19 mice in the

overlaying cortex of symptomatic mice (**Supp. Fig. 10F-H**).

We also examined microglia morphology using measurements corresponding to ramification

and correlating with microglia state (**Fig. 8A, C-E**). It has been previously reported that microglia

adopt an amoeboid morphology in PS19 mice, characterized by the loss of microglia processes,

and that is associated with a more reactive and phagocytic state[12,86]. We examined microglia

morphology in the CA1 in symptomatic mice as we observed the largest decrease in Iba1+ cell

counts in this hippocampal region (**Fig. 8B, Supp. 10C-E**). Both the PS19 and PS19;CK2α'$^{(+/-)}$

mice showed a decrease in the average branch length (**Fig. 8C**). However, PS19;CK2α'$^{(+/-)}$ mice

demonstrated a significant increase in the number of branches and number of endpoints

compared to the PS19 mice (**Fig. 8D,E**) indicating an improvement in the morphological state of
microglia.

Since the morphological state of microglia is strongly related to its reactive state and
neuroinflammation, we first assessed the expression of a variety of cytokines and other
inflammatory molecules associated with microglia reactivity by using a cytokine array panel with
hippocampal extracts from symptomatic PS19 and PS19;CK2α'$^{(+/-)}$ (**Fig. 8F, G**). Importantly, we
confirmed that several cytokines were significantly decreased (p<0.05) in PS19;CK2α'$^{(+/-)}$ mice
relative to PS19 including, MIP-1α, IL17, IL7, IL4, and I309 with several others showing trending
decreases (0.05<p<0.1) including TREM1 and TNFα (**Fig. 8F, G**). We also investigated whether
changes in overall cytokine levels and microglia morphology upon CK2α' depletion related to an
altered microglia reactive state. For this we assessed the levels of CD68 in the hippocampus of
mice and normalized to the number of Iba1+ cells observed in each animal and time point (**Fig.**
9A-E, Supp. Fig. 10I). We observed increased levels of CD68 in all regions of the hippocampus
in the PS19 mice at both prodromal and symptomatic stages (**Fig. 9C-E**)**.** PS19;CK2α'$^{(+/-)}$ mice
also presented increased CD68 levels compared to WT but displayed significantly less CD68 in
the CA1 and a trending decrease in the CA3 and DG at the prodromal stage compared to PS19
mice (**Fig. 9C-E**). Importantly, symptomatic PS19;CK2α'$^{(+/-)}$ mice displayed significantly less
CD68 in all hippocampal regions compared to PS19 mice (**Fig. 9C-E**).

Reactive microglia have also been implicated in synaptic and neuronal loss in tauopathy[80,87-91].
Prior studies have demonstrated that reactivity state of microglia is associated with microglial
engulfment of synapses which can be assessed through colocalization of synaptic markers such
as PSD95 within microglia[92,93]. To quantify synapse phagocytosis/engulfment we co-stained for
PSD95 and Iba1 and examined the CA1 stratum radiatum of symptomatic mice (**Fig. 9F**).
PSD95+ puncta were quantified within the Iba1+ cell soma, which is considered to be
lysosomes containing degraded material are present[94,95]. PS19 mice exhibited a significant
increase in PSD95+ puncta within Iba1+ relative to control mice (**Fig. 9F, G**)**,** demonstrating
increased phagocytosis in PS19 microglia. Importantly, PS19;CK2α'$^{(+/-)}$ showed a significant
reduction in PSD95+ engulfment (**Fig. 9F, G**) indicating an amelioration in the phagocytic
activity of microglia upon deletion of CK2α'.

Discussion

Tauopathies are a heterogeneous group of neurodegenerative disorders for which effective
treatments are still not available, in part due to an incomplete understanding of the underlying
mechanisms. In this study, we present a novel target, CK2α', a catalytic subunit of CK2, as a
potential upstream regulator of tau mediated neurodegeneration, acting at the intersection of
immune signaling and synaptic dysfunction.

Here, we report that CK2α' expression is abnormally elevated in the brains of dementia patients
and in the hippocampus of PS19 mice, with preferential upregulation observed in hippocampal
neurons and microglia. These findings are supported by multiple single-cell RNA-sequencing
(snRNA-seq) studies in human AD, which consistently show increased CK2α' expression in
excitatory neurons[69-71]. Additionally, the study by Gerrits et al.[68] in the occipital lobe (OL) has

reported preferential CK2α' expression in AD-associated microglia. Other scRNA-seq studies
have detected elevated CK2α' expression in various brain cell types, with patterns varying by
dataset and brain region[69-71]. Although these datasets lack specific information on the
hippocampus, the collective evidence suggests that CK2α' dysregulation in AD/ADRD may be
both cell-type- and region-specific.

Increased CK2 levels and/or activity have been previously reported in AD and AD models[19,37,72],
and CK2 has been associated with tau pathology[19,20,72,96], microglia state[97,98] and general
inflammation[99-105]. However, many of these studies were conducted in cell models[19,20,37,72] and
genetic evidence for the specific contributions of the individual CK2 subunits to these processes
remained largely unexplored. Our studies in vitro using N2A cells transfected with Tau-P301L
showed that silencing CK2α', but not CK2α, decreased pTau accumulation. Furthermore, our
studies in vivo reducing the levels of CK2α' in the PS19 model also influenced tau pathology by
decreasing the accumulation of pTau and tau burden in the hippocampus and cortex of
PS19:CK2α'$^{(+/-)}$ mice. In support to our data, previous studies *in vitro* by Perez et al. showed that
overexpression of CK2α' subunit decreased the activity of I2PP2A/SET, a cellular protein that
functions as an endogenous inhibitor of protein phosphatase 2A (PP2A), increasing pTau. In
contrast overexpression of CK2α increased the activity of I2PPA2a/SET[96]. Although early
studies demonstrated that Tau is a physiological substrate of CK2, at least in specific contexts
such as neurogenesis[106], the findings by Perez et al.[96] suggest that the effects we observed
after manipulating CK2α' on pTau accumulation could be mediated through intermediary
molecules that modulate Tau phosphorylation and not directly.

Haploinsufficiency of CK2α' also increased the expression of synaptic genes, synaptic density,
and synaptic function in the PS19 mice. Along these lines, haploinsufficiency of CK2α' in a
mouse model of HD also showed improved synaptic density and function[18,54], although how
exactly CK2α' influences these processes is still unknown. A previous study showed that CK2α'
can influence the expression of synaptic genes, especially those related to glutamatergic
excitatory signaling[18]. A more recent study showed that pharmacological inhibition of CK2
mitigated AD tau pathology by preventing the NMDA receptor subunit NR2B synaptic
mislocalization[72]. NR2B mediates long-term depression (LTD), and LTD specifically induces the
phosphorylation of tau. These studies support our findings and strengthen the relationship
between CK2α', tau phosphorylation, and the regulation of neuronal function in both AD and
other neurodegenerative diseases. However, since CK2α' is found elevated in both neurons and
microglia, it is still unknown whether the effects mediated by CK2α' in pTau accumulation and
synaptic function are cell or non-cell autonomous, warranting further investigation.

CK2 has been identified as a mediator of microglia reactivity and inflammation in different
models[97,98] and it has been well-established as a mediator of inflammation in a variety of
contexts including SARS-CoV infection, cancers, bacterial infections, intestinal inflammation and
renal failure [99-104]. This is largely mediated via CK2's involvement in regulating several large
inflammatory factors such as NF-κB, STAT1, and EGR-1[105]. It is known that chronic activation of
NF-kB increases the accumulation of pTau[12]. On the other hand, pathological tau has also been
shown to induce inflammation and NF-kB activation by interacting directly with inflammatory
receptors such as toll-like receptors 2 (TLR2)[13,14] creating a vicious cycle of worsening

inflammation and tau pathology[12-15]. Our data demonstrated that CK2α' haploinsufficiency
significantly reduced the number of Iba1+ cells in the hippocampus, reduced the levels of the
microglia reactive marker CD68, ameliorated microglia morphological changes associated with
pathology, reduced the levels of pro-inflammatory cytokines in PS19 mice and reduced
microglia phagocytic activity. Previous studies in human microglia derived from hiPSCs treated
with CK2 inhibitors reduced cytokines production[98]. Therefore, all together, these data support a
specific key role of CK2α' in mediating neuroinflammation via activation of microglia.

Loss of synaptic density and neuronal dysfunction has been previously connected to increased
phagocytic microglia[64,89,107,108]. The activation of the Complement Component pathway in
microglia is associated with enhanced phagocytic activity. While such activation is beneficial to
clear Tau aggregates and get rid of dysfunctional synapses, chronic activation results in
excessive pruning and an aberrant loss of synapses[80,89]. Our results demonstrated a decrease
in microglia phagocytosis of the PSD95 synaptic marker in PS19:CK2α'$^{(+/-)}$ mice suggesting that
CK2α' is influencing the phagocytosis of synapses by microglia, but the mechanisms behind this
are unclear. Complement component q1 (C1q) is often seen at synapses that accumulate pTau,
and the levels of C1q are associated with enhanced phagocytic microglia engulfment of
synapses leading to the decline of synapse density, synaptic function and cognitive decline
associated with AD[89]. In line with these previous findings, our RNA-seq data revealed the
upregulation of genes related to synaptic pruning in PS19 mice, especially those associated
with C1q. Importantly, antibodies targeting C1q have rescued synapse loss in tauopathy
models[89]. While manipulation of CK2α' did not impact the expression of C1q, it is possible that
CK2α' is impacting the complement pathway via phosphorylation of critical components, such as
Complement component (C9) and complement component (C1r), which are known targets of
CK2 [109,110]. Further studies focusing on the impact of CK2α' on the classical cascade mediated
synapse engulfment will be important to elucidate the mechanisms involved in improved
synaptic density, function, and cognition.

An intriguing finding from our study is the apparent discrepancy between the upregulation of
genes associated with immune and innate responses and the reduction of neuroinflammatory
phenotypes in microglia in PS19:CK2α'$^{(+/-)}$ mice. Typically, the transcriptional activation of innate
immune pathways is linked to pathology and neuroinflammation. In PS19 mice, such activation,
along with increased expression of genes related to microglial dysfunction and reactivity, has
been extensively reported[80,81]. Surprisingly, while CK2α' haploinsufficiency further amplified this
transcriptional immune response in PS19 mice, we observed a concomitant reduction in
neuroinflammatory markers, accompanied by improvements in neuronal function and behavior.
This apparent paradox may reflect a primed but non-pathological immune state, the
engagement of anti-inflammatory regulatory mechanisms, or a temporal or cell-type-specific
decoupling between transcriptional signatures and functional outcomes. Similar findings have
been reported in studies involving partial agonists of the tropomyosin receptor kinase B (TrkB)
and C (TrkC), where synaptic deficits were rescued in the APP model of AD despite enhanced
expression of microglial neuroinflammatory markers and immune responses typically associated
with late-stage pathology[111]. Additionally another study reported that INPP5D haploinsufficiency
in the PS19 mouse resulted in decreased tau pathology and ameliorated motor deficits[112], yet
increased inflammatory signatures seen via RNA-Seq. Collectively, these observations

underscore the nuanced role of immune signaling in the CNS and caution against interpreting
transcriptional evidence of inflammation as a direct indicator of detrimental neuroimmune
activation.

In conclusion, our findings identify CK2α' as a central and previously underappreciated regulator
of tauopathy pathogenesis, acting at the crossroads of immune signaling, synaptic function, and
tau phosphorylation. By demonstrating that CK2α' haploinsufficiency alleviates tau pathology,
reduces neuroinflammation, and improves synaptic integrity and function in PS19 mice, we
provide compelling genetic evidence that CK2α' contributes to multiple pathological hallmarks of
tau-driven neurodegeneration. Importantly, these effects appear to occur through complex,
context-dependent mechanisms that may differ across cell types and disease stages. The
observation that CK2α' influences both neuronal and microglial compartments underscore the
need to dissect its cell-autonomous versus non-cell-autonomous roles. Given the growing
interest in CK2 as a therapeutic target and the limitations of current inhibitors that lack subunit
specificity, the development of CK2α'-selective modulators[113] may represent a promising avenue
for disease-modifying therapies in tauopathies and related neurodegenerative conditions. Future
studies elucidating the molecular and cellular pathways downstream of CK2α' will be critical for
translating these findings into clinical applications.

Figure Legends

Figure 1. CK2α' is increased in tissues of patients with AD/ADRD. **(A, B)** Representative
diagram of CK2 holoenzyme, including regulatory and catalytic subunits and differential
distribution of expression of catalytic subunits of CK2α and CK2α' in the human body. A and B
were created in BioRender (White, A. (2025) eqznm6m https://BioRender.com/). **(C)** Averaged
expression of *CSNK2A2* in non-cognitively impaired (NCI) and dementia samples from the Allen
Brain Institute: Aging, Dementia and TBI study; parietal cortex (PCx), Temporal cortex (TCx),
and Hippocampus (HIP) are shown (patients with TBI were excluded from the analyses). **(E-H)**
Quantification of the expression of *CSNK2A2* and *CSNK2A1* in PCx (**E, F**), TCx (**G**), and HIP
(**H**) from NCI and dementia from the Allen Brain Institute: Aging, Dementia and TBI study (n=20-
27/group). Significance determined by unpaired t-test, *p<0.05. **(I)** Representative immunoblot
for CK2α' in protein lysates from frontal cortex of age and sex matched healthy controls (C) and
patients with FTD. Arrow indicates CK2α' specific band, asterisk indicates unspecific band. **(J)**
CK2α' protein levels were normalized to GAPDH and shown relative to age/sex matched
control, quantified using ImageJ software (n=7/group). Data are shown as mean ± SEM.
Statistical analyses were conducted using unpaired t-test with welch's correction. * p<0.05. (**K**)
Relationship between CK2α' levels from (I) and the presence of Tau pathology reported by the
NIH biobank for those samples. Horizontal line represents the group CK2α' intensity average
normalized by GAPDH.

Figure 2. *Csnk2a2* **expression is increased in both neurons and microglia in the hippocampus of PS19 mice. (A)** *Csnk2a2 in situ* hybridization images from symptomatic WT and PS19 hippocampus. Left panels show 10x images of hippocampal cell layers with *Csnk2a2* RNA probe and DAPI, scale bar=50 µm. White boxes indicate zoomed inlets shown on the right panels at 20x, scale bar=10 µm. **(B)** (B) Csnk2a2 in situ hybridization and NeuN immuno-fluorescence images from symptomatic WT and PS19 CA1. Scale bar= 20 µm. **(C-E)** Quantification of *Csnk2a2* puncta in the CA1 (**C**), CA3 (**D**) and DG (**E**) region quantified within the cell layer identified by DAPI, (n=3-4 mice/genotype). **(F)** Representative images *Csnk2a2* RNA probe and Iba1 immunofluorescence in symptomatic WT and PS19 hippocampus. **(G)** Top: representative image of single Iba1+ cell in the Stratum Radiatum (CA1) with *Csnk2a2* RNA Scope. Scale bar is 20 µm. Bottom: 3D rendering of cells Iba1+ cells containing *Csnk2a2* puncta showing puncta reside within the cell. **(H)** Quantification of the number of *Csnk2a2*+ puncta/Iba1+ soma (n=3-4 mice/group, 8-9 cells/mouse, every data point is a cell). All Data shown are mean $\pm$ SEM. Statistical analyses were conducted using unpaired t-test * $p<0.05$, *** $p<0.001$, **** $p<0.0001$.

Figure 3. Silencing CK2α' reduces pTau levels in N2a cells expressing Tau-P301L. (A) Schematic of experimental design for N2a cell transfection and immunostaining. **(B)** Representative images of AT8 immunostaining and EGFP fluorescence in N2a cells transfected with either EGFP or Tau-P301L-EGFP and treated with ScrRNA, siCK2α, or siCK2α'. Scale bar= 100 µm. **(C-E)** RT-qPCR analysis of mRNA expression levels of CK2α' **(C)**, CK2α **(D)**, and CK2β **(E)** in transfected N2a cells. Expression levels were normalized to GAPDH and presented relative to ScrRNA EGFP condition (n=3/condition). **(F)** RT-qPCR analysis of mRNA expression levels of huTau expression in N2a cells transfected with EGFP or Tau-P301L-EGFP transfected N2a cells. Expression levels were normalized to GAPDH and presented relative to EGFP condition (n=6/condition). **(G)** Quantification of AT8+ cells normalized to the number of EGFP+ cells, presented relative to ScrRNA Tau-P301L condition. (n=4/condition). All data shown as mean ± SEM. Statistical analyses were conducted using one-way ANOVA with tukey's multiple and displayed with compact letter display. **(H)** Diagram summarizing results of experiment in B. **A** and **H** were created in BioRender (White, A. (2025) https://BioRender.com/eqznm6m).

Figure 4. CK2α' haploinsufficiency reduced tau pathology in PS19 mice. (A) Representative whole-slice sections immunostained for pTau using the AT8 antibody (Ser202/Thr205) in prodromal and symptomatic mice, scale bar= 1000 µm. **(B)** Higher magnification images of hippocampus (CA1, CA3, and DG) from sections shown in (A). Scale bar=500 µm. **(C-E)** Quantification of AT8+ immunoreactivity (% area) in the CA1, CA3 and DG regions in prodromal and symptomatic cohorts (n=6-9 mice/genotype). Data represented as mean $\pm$ SEM. Statistical analyses were conducted using two-way ANOVA with tukey's posthoc analysis and displayed with compact letter display. **(E)** Representative pTau pathology types. **(F)** Proportion of each pTau pathology type in prodromal and symptomatic cohorts (n=7-10 mice/genotype). Fisher's exact t-test 7m p=0.11 & 12m p=0.0587.

Figure 5. CK2α' haploinsufficiency partially rescued hippocampal synapse loss and improved LTP in PS19 mice. (A) Representative NeuN staining in the CA1 of symptomatic mice. Scale bar=200 μm. **(B)** Quantification of NeuN+ cells (cells/mm^2) in the CA1 of prodromal and symptomatic cohorts (n= 3-6 mice/genotype; 2-3 slices/mouse; 1 data point= 1 slice, shape of data point indicates same animal slices). Data was shown as mean ± SEM. Statistical analyses were conducted using one way-ANOVA with Tukey's post-hoc analysis and displayed by compact letter display. Comparisons shown relative to cohort age group. **(C)** Vglut1 and PSD95 immunostaining in the CA1 of symptomatic mice. White arrow heads indicate colocalization of Vglut1 and PSD95. Scale bar= 2.5 μm. **(D-F)** Quantification of PSD95 puncta (**D**), Vglut1 puncta (**E**) and Vglut1/PSD95 colocalized puncta (**F**) relative to WT. (n=3 mice/genotype (3 slices/mouse, 3 images/slice, 1 data point=slice average). **(G-I)** Synaptic analyses from (D-F) represented as a % decrease from WT. Data are shown as mean ± SEM. Statistical analyses were conducted using repeated measures ANOVA with Geisser-greenhouse correction and Tukey's post-hoc analysis *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. **(J)** Scheme showing the general experimental set-up: R, recording electrode; S stimulating electrode. **(K)** Representative traces before (1) and after (2) the TBS application. **(L)** Course temporal of Theta-burst stimulation (TBS) application inducing LTP in pyramidal neurons of CA1 region from hippocampus (WT, Black n=6; CK2α'$^{(+/-)}$, Grey n=7; PS19, Red, n=7; PS19;CK2α'$^{(+/-)}$, Pink n=6). **(M)** Summary of short-term potentiation (STP) results; significance reported relative to WT. **(N)** Input-output curves. Averaged fEPSP amplitude, (WT, n=3; CK2α'$^{(+/-)}$,n=3; PS19, n=3; PS19;CK2α'$^{(+/-)}$,n=3). (*p < 0.05,**p < 0.01). All data are shown as mean ± SEM. Statistical analyses were conducted using two-sided Student's *t* test or One-way ANOVA with Holm-Sidak multiple comparisons. **(O)** Summary of long-term potentiation (LTP) results; significance reported relative to WT.

Figure 6. Depletion of CK2α' improved spatial memory of PS19 mice on the Barnes maze.
(A) Schematic of the Barnes Maze divided into 4 quadrants (Target, Opposite, +1 and -1) and experimental set up. **(B-C)** Primary latency (s), or time to first explore escape hole at prodromal time point. Statistical analyses were conducted using two-way ANOVA with Bonferroni's multiple comparison's. Prodromal; genotype p=0.0573, training day p<0.0001 and interaction p=0.9690. Symptomatic: genotype p<0.0001, training day p<0.0001 and interaction p=0.7018. Differences for each training day were conducted using Bonferroni's multiple comparisons and * show significance relative to WT. **(D)** Time spent in goal quadrant of Barnes maze during probe trial at prodromal and symptomatic time points. n=10-12 mice/genotype, 2 outliers were removed from PS19 group using ROUT (Q=1%). **(E)** Representative search strategy traces based on their navigation patterns on the maze. **(F,H)** Spatial strategies used to locate the escape hole across training days for each genotype at prodromal and symptomatic stages, respectively. **G,I)** Spatial strategies used to locate the previously learned escape target hole during probe trial. Data are shown as mean ± SEM. Statistical analyses were conducted using Two-way ANOVA with Bonferroni's multiple comparisons (B-C) or Two-Way ANOVA with Tukey's post-hoc analysis (D).

Significance displayed with either *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 or compact letter
display.

Figure 7. CK2α' haploinsufficiency has transcriptional impacts on synapses and immune
related gene pathways in PS19 mice. (A) Volcano plot of DGEs in PS19 vs WT comparison
(n=3-4 mice/genotype). **(B)** Gene-ontology top biological pathways for DGEs in PS19 vs WT.
(C) Gene network for top 5 synaptic pathways impacted in PS19 vs WT. Red dots indicate
genes upregulated in PS19 mice. Blue dots indicate genes down-regulated in PS19 mice. **(D)**
Heat map displaying average FC across three comparisons PS19 vs WT, PS19 vs PS19;
CK2α'$^{(+/-)}$ and PS19; CK2α'$^{(+/-)}$ vs WT for all significant DGEs identified via PS19 vs PS19;
CK2α'$^{(+/-)}$ comparison. Red boxes indicate genes with apoptotic/phagocytotic and or immune-
related functions. *q>0.05. **(E)** WGCNA identified modules, dotted line indicates significance as
determined by the Kruskal wallis test between WT, CK2α'$^{(+/-)}$, PS19, and PS19:CK2α'$^{(+/-)}$ mice
(n=3-5 mice/genotype). **(F)** Heatmap of modules detected from WGCNA indicating differences
across genotypes. * indicates p<0.05 determined by Kruskal-wallis test. **(G, H)** Gene ontology
for top biological pathways for *MEtan* module (**G**) and *MEturquoise* module (**H**). **(I)** Average
representation of pathway activity scores determined via GSVA analysis. Panels **G** and **H** were
created in BioRender (White, A. (2025) https://BioRender.com/eqznm6m).

Figure 8. CK2α' haploinsufficiency impacts Iba1+ microglia morphology and cytokine
production in PS19 mice. (A) Iba1+ microglia immunostaining and microglia skeleton
representation from the CA1 of symptomatic animals used for microglia morphology analysis.
(B) Quantification of the number of Iba1+ positive cells in the CA1 (n=3-6 mice/genotype). **(C)**
Quantification of microglia branch length, **(D)** the number of branches per microglia, **(E)** and
number of endpoints per microglia (n=27 cells (3 mice/genotype, 3 slices/mouse, 3 cells/slice).
(F) Representative cytokine arrays from PS19 and PS19;CK2α'$^{(+/-)}$ at 9 moths from hippocampal
protein extracts. Cytokines that showed significant differences are circled in red. **(G)** Selected
quantifications of cytokines showing significance or trending significance. Quantifications are
represented relative to WT (n=4 mice/genotype). Data are shown as mean ± SEM. Statistical
analyses were conducted using ANOVA with Geisser-greenhouse correction and Tukey's post-
hoc analysis in **B-E,** and unpaired t-test in **G.** #p<0.1, *p<0.05, **p<0.01, ***p<0.001,
****p<0.0001.

Figure 9. CK2α' haploinsufficiency decreases microglia reactivity and synaptic
engulfment. (A) CD68 immunostaining with cresyl violet counterstain in the hippocampus of
symptomatic mice. Scale bar=150 µm. (**B**) zoomed images from A in the CA1 and DG. **(C-E)**
Quantifications of % area CD68+ normalized by Iba1+ cell count in prodromal and symptomatic
cohorts in CA1, CA3 and DG (n=3-6 mice/genotype). Comparisons shown relative to cohort age
group. **(F)** Iba1 and PSD95 immunostaining from the CA1 of symptomatic animals. Top row
displays whole microglia including processes. Scale bar= 5µm. Middle row displays higher
magnification view of the soma (white dotted box in top row), scale bar= 2µm. Bottom row
displays 3D reconstruction of Iba1 and PSD95 signal generated from Imaris Software, scale

bar= 2μm. **(G)** Quantification of the number of PSD95 puncta/Iba1+ soma (n= 16-18 cell, 3
mice/genotype, 2-3 slices/mouse, 22 cells/slice). Data are show as mean $\pm$ SEM. Statistical
analyses were conducted using one-way ANOVA with Tukey's post-hoc *p<0.05, **p<0.01,
***p<0.001, ****p<0.0001.

Supplementary Figure legends

Supplementary Figure 1. *CSNK2a2* **scRNA-seq expression in human brain tissue. (A)**
Summary of *CSNK2a2* single cell expression from studies found in The Alzheimer's Cell Atlas
(TACA) in occipital lobe (OL), occipitotemporal cortex (OTC), entorhinal cortex (EC), superior
frontal gyrus (SFG), and prefrontal cortex (PFC). Absence of expression for a cell type indicates
that cell type was not identified as a cluster in the study.

Supplementary Figure 2. **Generation of a CK2α' haploinsufficient tauopathy model. (A)**
The PS19 mouse model was crossed with mice haploinsufficient for CK2α' (Illustration from
NIAID NIH BIOART Source (bioart.niaid.nih.gov/bioart/283 & bioart.niaid.nih.gov/bioart/281). **(B)**
CK2α' immunoblotting in the hippocampus of symptomatic mice. (**C, D**) Quantification of CK2α'
levels from B in CK2α'$^{(+/-)}$ mice relative to WT and PS19;CK2α'$^{(+/-)}$ mice relative to PS19 mice in
symptomatic hippocampus (n=6 mice/genotype). Arrow indicates specific CK2α' band. Asterisk
indicates an unspecific band. **(E, F)** Relative gene expression of CK2α' in CK2α'$^{(+/-)}$ mice relative
to WT and PS19;CK2α'$^{(+/-)}$ mice relative to PS19 mice in the symptomatic hippocampus from
qPCR (n=3 mice/genotype). **(G)** Survival curves up to 60 weeks for all 4 genotypes (n=64-74
mice/genotype). Data are shown as mean $\pm$ SEM. Statistical analyses were conducted using
unpaired t-test *p<0.05, **p<0.01, ***p<0.001.

1009

Supplementary Figure 3. PS19;CK2α'(+/-) mice display decreased AT8 staining in the
cortex of symptomatic mice. (A) Representative diagram of cortical region imaged for
quantification in B and C. **(B)** Representative images of AT8 staining in the cortex of
symptomatic mice. **(B)** Quantification of % area AT8+ in the cortex of prodromal and
symptomatic mice (n=3-7 mice/genotype). Data are shown as mean ± SEM. Statistical analyses
were conducted using Two-way ANOVA with Tukey's post-hoc analysis and displayed with
compact letter display. Panel A was created in BioRender (White, A. (2025)
https://BioRender.com/eqznm6m).

1018

Supplementary Figure 4. NeuN pathology and variability among symptomatic PS19 mice.
(A) NeuN immunostaining in the hippocampus of prodromal and symptomatic mice. Dotted
boxes indicate quantified subregions: CA1, CA3, and DG. Scale bar = 250 μm. **(B-D)**
Quantifications of NeuN+ cells (cells/mm^2) in the CA3 (**B**), and DG (**C**) in prodromal and
symptomatic cohorts (n= 3-6 mice/genotype;2-3 slices/mouse;1 data point=1 slice, shape of
data point indicates slices of the same animal). Data was shown as mean $\pm$ SEM. Statistical
analyses were conducted using one way-ANOVA with Tukey's post-hoc analysis and displayed

by compact letter display. Comparisons shown relative to cohort age group. **(D)** Representative
NeuN immunofluorescence in the hippocampus of symptomatic PS19 mice, demonstrating high
and low pathology within littermates. Scale bar=250 µm. **(E)** Quantification of NeuN+ cells in the
CA1, CA3 and DG of the hippocampus in high and low pathology mice (n=3 mice/genotype). **(F)**
Quantification of hippocampal area in high and low pathology mice (n=3 mice/genotype). Data
are shown as mean ± SEM. Statistical analyses were conducted using unpaired t-test. **p<0.01,
***p<0.001.

1033

1034   **Supplementary Figure 5. Paired pulse ratio of hippocampal slices shows no deficits**
1035   **between genotypes. (A)** Paired-pulse ratio (PPR) in acute hippocampal slices of (WT, Black
1036   n=5; CK2α'$^{(+/-)}$, Grey, n=5; PS19, Red, n=5; PS19; CK2α'$^{(+/-)}$, Pink n=4). (*p < 0.05,**p < 0.01).
1037   Data are shown as mean ± SEM. Statistical analyses were conducted using two-sided Student's
1038   t test. n.s. indicates no significance.

1039

Supplementary Figure 6. Symptomatic PS19 demonstrates decreased movement on
Barnes Maze, but hyperactive phenotype in open field. (A-B) Distance traveled in prodromal
and symptomatic mice during the probe phase of Barnes Maze. **(C-D)** Distance traveled by
prodromal and symptomatic mice during 1 hour in an open field arena. Data are shown as mean
± SEM (n=10-12 mice/genotype, outliers removed using ROUT (Q=1%). Statistical analyses
were conducted using One-Way ANOVA with Tukey's post-hoc analysis. *p<0.05, **p<0.01,
***p<0.001, ****p<0.0001.

1047

Supplementary Figure 7. RNA-Seq analyses in PS19 cohort show genetic variability
associated with high and low pathology. (A) Representative AT8 staining of a low and high
pathology prodromal PS19 mice scale bar= 1000 µm. **(B)** Gene ontology top related pathways
for Low pathology PS19 vs WT. **(C)** Gene ontology top related pathways for High pathology
PS19 vs WT comparison. **(D-E)** Venn diagram of the number of unique and overlapping DGEs
for 6 total comparisons of interest.

1054

Supplementary Figure 8. PS19;CK2α'(+/-) mice display increased immune-related gene
expression. (A) Eigengene Average expression for tan module. **(A)** Eigengene Average
expression for turquoise module. **(C)** Gene ontology for the top-related pathways for the DGEs
in PS19 vs WT not present in the PS19;CK2α'$^{(+/-)}$ vs WT comparison. **(D)** Gene ontology for the
top-related pathways for the DGEs in PS19;CK2α'$^{(+/-)}$ vs WT not present in the PS19 vs WT
comparison. **(E)** Gene network for the top 2 pathways returned in (D).

1061

Supplementary Figure 9. CK2α' haploinsufficency decreased GFAP+ astrocytes in the
overlaying cortex but not in the hippocampus of PS19 mice. (A-B) Immunostaining for

GFAP in the CA1, CA3 and DG of the hippocampus in prodromal (A) and symptomatic (B)
cohorts. Scale bar=100 µm. **(C-E)** Quantification of GFAP+ cells in CA1, CA3 and DG of
hippocampus. (n=3-6 mice/genotype). **(F)** Representative area corresponding to overlying
cortex is indicated with a boxed area. **(G)** Representative GFAP immunostaining in the cortex.
Scale bar= 50 µm. **(H)** Quantification of GFAP+ cells in the cortex (n=3-6/genotype). All Data
are shown as mean ± SEM. Statistical analyses were conducted using two-way ANOVA with
tukey's multiple corrections and represented with compact letter display. Panel **F** was created in
BioRender (White, A. (2025) https://BioRender.com/eqznm6m).

1072

Supplementary Figure 10. CK2α' haploinsufficency decreased Iba1+ microglia cell
number in the hippocampus and overlaying cortex of PS19 mice. (A-B) Immunostaining for
Iba1 in the CA1, CA3 and DG of the hippocampus in prodromal (**A**) and symptomatic (**B**)
cohorts. Scale bar=100 µm. **(C-E)** Quantification of Iba1+ cell counts in CA1, CA3 and DG of
hippocampus (n=3-6 mice/genotype). **(F)** Representative area corresponding to overlying cortex
is indicated with a boxed area. **(G)** Representative Iba1 immunostaining in the cortex scale
bar=50 µm. **(H)** Quantification of Iba1+ cells in the cortex (n=3-6 mice/genotype). All data are
shown as mean ± SEM. Statistical analyses were conducted using two-way ANOVA with tukey's
multiple corrections and represented with compact letter display. Panel **F** was created in
BioRender (White, A. (2025) https://BioRender.com/eqznm6m).

1083

1084 **References**

1085

1086

1.	Kovacech B, Skrabana R, Novak M. Transition of tau protein from disordered to
misordered in Alzheimer's disease. *Neurodegener Dis.* 2010;7(1-3):24-7.
doi:10.1159/000283478
2.	Lee VM, Goedert M, Trojanowski JQ. Neurodegenerative tauopathies. *Annu Rev*
Neurosci. 2001;24:1121-59. doi:10.1146/annurev.neuro.24.1.1121
3.	Josephs KA, Hodges JR, Snowden JS, et al. Neuropathological background of
phenotypical variability in frontotemporal dementia. *Acta Neuropathol.* Aug 2011;122(2):137-53.
doi:10.1007/s00401-011-0839-6
4.	Weingarten MD, Lockwood AH, Hwo SY, Kirschner MW. A protein factor essential for
microtubule assembly. *Proc Natl Acad Sci U S A.* May 1975;72(5):1858-62.
doi:10.1073/pnas.72.5.1858
5.	Strang KH, Golde TE, Giasson BI. MAPT mutations, tauopathy, and mechanisms of
neurodegeneration. *Lab Invest.* Jul 2019;99(7):912-928. doi:10.1038/s41374-019-0197-x
6.	Sexton C, Snyder H, Beher D, et al. Current directions in tau research: Highlights from
Tau 2020. *Alzheimers Dement.* May 2022;18(5):988-1007. doi:10.1002/alz.12452
7.	Tai HC, Serrano-Pozo A, Hashimoto T, Frosch MP, Spires-Jones TL, Hyman BT. The
synaptic accumulation of hyperphosphorylated tau oligomers in Alzheimer disease is associated
with dysfunction of the ubiquitin-proteasome system. *Am J Pathol.* Oct 2012;181(4):1426-35.
doi:10.1016/j.ajpath.2012.06.033

8.   Dani M, Wood M, Mizoguchi R, et al. Microglial activation correlates in vivo with both tau and amyloid in Alzheimer's disease. *Brain*. Sep 01 2018;141(9):2740-2754. doi:10.1093/brain/awy188

9.   Lee SH, Bae EJ, Park SJ, Lee SJ. Microglia-driven inflammation induces progressive tauopathies and synucleinopathies. *Exp Mol Med*. May 01 2025;doi:10.1038/s12276-025-01450-z

10.   Bejanin A, Schonhaut DR, La Joie R, et al. Tau pathology and neurodegeneration contribute to cognitive impairment in Alzheimer's disease. *Brain*. Dec 01 2017;140(12):3286-3300. doi:10.1093/brain/awx243

11.   Langworth-Green C, Patel S, Jaunmuktane Z, et al. Chronic effects of inflammation on tauopathies. *Lancet Neurol*. May 2023;22(5):430-442. doi:10.1016/S1474-4422(23)00038-8

12.   Wang C, Fan L, Khawaja RR, et al. Microglial NF-κB drives tau spreading and toxicity in a mouse model of tauopathy. *Nat Commun*. Apr 12 2022;13(1):1969. doi:10.1038/s41467-022-29552-6

13.   Kim Y, Ryu SH, Hyun J, Cho YS, Jung YK. TLR2 immunotherapy suppresses neuroinflammation, tau spread, and memory loss in rTg4510 mice. *Brain Behav Immun*. Oct 2024;121:291-302. doi:10.1016/j.bbi.2024.08.002

14.   Dutta D, Jana M, Paidi RK, et al. Tau fibrils induce glial inflammation and neuropathology via TLR2 in Alzheimer's disease-related mouse models. *J Clin Invest*. Sep 15 2023;133(18)doi:10.1172/JCI161987

15.   Bhaskar K, Konerth M, Kokiko-Cochran ON, Cardona A, Ransohoff RM, Lamb BT. Regulation of tau pathology by the microglial fractalkine receptor. *Neuron*. Oct 06 2010;68(1):19-31. doi:10.1016/j.neuron.2010.08.023

16.   Litchfield DW. Protein kinase CK2: structure, regulation and role in cellular decisions of life and death. *Biochem J*. Jan 01 2003;369(Pt 1):1-15. doi:10.1042/BJ20021469

17.   Gomez-Pastor R, Burchfiel ET, Neef DW, et al. Abnormal degradation of the neuronal stress-protective transcription factor HSF1 in Huntington's disease. *Nat Commun*. 02 13 2017;8:14405. doi:10.1038/ncomms14405

18.   Yu D, Zarate N, White A, et al. CK2 alpha prime and alpha-synuclein pathogenic functional interaction mediates synaptic dysregulation in huntington's disease. *Acta Neuropathol Commun*. Jun 03 2022;10(1):83. doi:10.1186/s40478-022-01379-8

19.   Zhang Q, Xia Y, Wang Y, et al. CK2 Phosphorylating I 2 PP2A /SET Mediates Tau Pathology and Cognitive Impairment. *Front Mol Neurosci*. 2018;11:146. doi:10.3389/fnmol.2018.00146

20.   Yadikar H, Torres I, Aiello G, et al. Screening of tau protein kinase inhibitors in a tauopathy-relevant cell-based model of tau hyperphosphorylation and oligomerization. *PLoS One*. 2020;15(7):e0224952. doi:10.1371/journal.pone.0224952

21.   Baum L, Masliah E, Iimoto DS, Hansen LA, Halliday WC, Saitoh T. Casein kinase II is associated with neurofibrillary tangles but is not an intrinsic component of paired helical filaments. *Brain Res*. Feb 21 1992;573(1):126-32. doi:10.1016/0006-8993(92)90121-o

22.   Lim AC, Tiu SY, Li Q, Qi RZ. Direct regulation of microtubule dynamics by protein kinase CK2. *J Biol Chem*. Feb 06 2004;279(6):4433-9. doi:10.1074/jbc.M310563200

23.   Chung HJ, Huang YH, Lau LF, Huganir RL. Regulation of the NMDA receptor complex and trafficking by activity-dependent phosphorylation of the NR2B subunit PDZ ligand. *J Neurosci*. Nov 10 2004;24(45):10248-59. doi:10.1523/JNEUROSCI.0546-04.2004

24.   Kimura R, Matsuki N. Protein kinase CK2 modulates synaptic plasticity by modification of synaptic NMDA receptors in the hippocampus. *J Physiol*. Jul 01 2008;586(13):3195-206. doi:10.1113/jphysiol.2008.151894

25.   Lenzken SC, Stanga S, Lanni C, De Leonardis F, Govoni S, Racchi M. Recruitment of casein kinase 2 is involved in AbetaPP processing following cholinergic stimulation. *J Alzheimers Dis*. 2010;20(4):1133-41. doi:10.3233/JAD-2010-090232

26.   Raftery M, Campbell R, Glaros EN, et al. Phosphorylation of apolipoprotein-E at an atypical protein kinase CK2 PSD/E site in vitro. *Biochemistry*. May 17 2005;44(19):7346-53. doi:10.1021/bi0504052

27.   Walter J, Schindzielorz A, Hartung B, Haass C. Phosphorylation of the beta-amyloid precursor protein at the cell surface by ectocasein kinases 1 and 2. *J Biol Chem*. Aug 04 2000;275(31):23523-9. doi:10.1074/jbc.M002850200

28.   Bian Y, Ye M, Wang C, et al. Global screening of CK2 kinase substrates by an integrated phosphoproteomics workflow. *Sci Rep*. Dec 10 2013;3:3460. doi:10.1038/srep03460

29.   Rusin SF, Adamo ME, Kettenbach AN. Identification of Candidate Casein Kinase 2 Substrates in Mitosis by Quantitative Phosphoproteomics. *Front Cell Dev Biol*. 2017;5:97. doi:10.3389/fcell.2017.00097

30.   Guerra B, Siemer S, Boldyreff B, Issinger OG. Protein kinase CK2: evidence for a protein kinase CK2beta subunit fraction, devoid of the catalytic CK2alpha subunit, in mouse brain and testicles. *FEBS Lett*. Dec 03 1999;462(3):353-7. doi:10.1016/s0014-5793(99)01553-7

31.   Montenarh M, Götz C. Protein Kinase CK2α', More than a Backup of CK2α. *Cells*. Dec 14 2023;12(24)doi:10.3390/cells12242834

32.   Ceglia I, Flajolet M, Rebholz H. Predominance of CK2α over CK2α' in the mammalian brain. *Mol Cell Biochem*. Oct 2011;356(1-2):169-75. doi:10.1007/s11010-011-0963-6

33.   Lou DY, Dominguez I, Toselli P, Landesman-Bollag E, O'Brien C, Seldin DC. The alpha catalytic subunit of protein kinase CK2 is required for mouse embryonic development. *Mol Cell Biol*. Jan 2008;28(1):131-9. doi:10.1128/MCB.01119-07

34.   Xu X, Toselli PA, Russell LD, Seldin DC. Globozoospermia in mice lacking the casein kinase II alpha' catalytic subunit. *Nat Genet*. Sep 1999;23(1):118-21. doi:10.1038/12729

35.   Iimoto DS, Masliah E, DeTeresa R, Terry RD, Saitoh T. Aberrant casein kinase II in Alzheimer's disease. *Brain Res*. Jan 22 1990;507(2):273-80. doi:10.1016/0006-8993(90)90282-g

36.   Allende JE, Allende CC. Protein kinases. 4. Protein kinase CK2: an enzyme with multiple substrates and a puzzling regulation. *FASEB J*. Mar 1995;9(5):313-23. doi:10.1096/fasebj.9.5.7896000

37.   Rosenberger AF, Morrema TH, Gerritsen WH, et al. Increased occurrence of protein kinase CK2 in astrocytes in Alzheimer's disease pathology. *J Neuroinflammation*. Jan 06 2016;13:4. doi:10.1186/s12974-015-0470-x

38.   Aksenova MV, Burbaeva GS, Kandror KV, Kapkov DV, Stepanov AS. The decreased level of casein kinase 2 in brain cortex of schizophrenic and Alzheimer's disease patients. *FEBS Lett*. Feb 11 1991;279(1):55-7. doi:10.1016/0014-5793(91)80249-3

39.   Saitoh T, Iimoto D. Aberrant protein phosphorylation and cytoarchitecture in Alzheimer's disease. *Prog Clin Biol Res*. 1989;317:769-80.

40.   Yoshiyama Y, Higuchi M, Zhang B, et al. Synapse loss and microglial activation precede tangles in a P301S tauopathy mouse model. *Neuron*. Feb 01 2007;53(3):337-51. doi:10.1016/j.neuron.2007.01.010

41.   Wauters M, Wattiez R, Ris L. Internalization of the Extracellular Full-Length Tau Inside Neuro2A and Cortical Cells Is Enhanced by Phosphorylation. *Biomolecules*. Aug 19 2016;6(3)doi:10.3390/biom6030036

42.   Namsi A, Nury T, Hamdouni H, et al. Induction of Neuronal Differentiation of Murine N2a Cells by Two Polyphenols Present in the Mediterranean Diet Mimicking Neurotrophins Activities: Resveratrol and Apigenin. *Diseases*. Jul 22 2018;6(3)doi:10.3390/diseases6030067

43.   Hoover BR, Reed MN, Su J, et al. Tau mislocalization to dendritic spines mediates synaptic dysfunction independently of neurodegeneration. *Neuron*. Dec 22 2010;68(6):1067-81. doi:10.1016/j.neuron.2010.11.030

44.   Brown TG, Thayer MN, VanTreeck JG, et al. Striatal spatial heterogeneity, clustering,
and white matter association of GFAP. *Front Cell Neurosci*. 2023;17:1094503.
doi:10.3389/fncel.2023.1094503
45.   Solem MA, G Pelzel R, Rozema NB, et al. Absence of hippocampal pathology persists in
the Q175DN mouse model of Huntington's disease despite elevated HTT aggregation. *J*
Huntingtons Dis. Feb 2025;14(1):59-84. doi:10.1177/18796397251316762
46.   Shi Y, Yamada K, Liddelow SA, et al. ApoE4 markedly exacerbates tau-mediated
neurodegeneration in a mouse model of tauopathy. *Nature*. Sep 28 2017;549(7673):523-527.
doi:10.1038/nature24016
47.   Shi Y, Manis M, Long J, et al. Microglia drive APOE-dependent neurodegeneration in a
tauopathy mouse model. *J Exp Med*. Nov 04 2019;216(11):2546-2561.
doi:10.1084/jem.20190980
48.   Martini-Stoica H, Cole AL, Swartzlander DB, et al. TFEB enhances astroglial uptake of
extracellular tau species and reduces tau spreading. *J Exp Med*. Sep 03 2018;215(9):2355-
2377. doi:10.1084/jem.20172158
49.   Takahashi H, Bhagwagar S, Nies SH, et al. Reduced progranulin increases tau and α-
synuclein inclusions and alters mouse tauopathy phenotypes via glucocerebrosidase. *Nat*
Commun. Feb 16 2024;15(1):1434. doi:10.1038/s41467-024-45692-3
50.   Bankhead P, Loughrey MB, Fernández JA, et al. QuPath: Open source software for
digital pathology image analysis. *Sci Rep*. Dec 04 2017;7(1):16878. doi:10.1038/s41598-017-
17204-5
51.   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for
biological-image analysis. *Nat Methods*. Jun 28 2012;9(7):676-82. doi:10.1038/nmeth.2019
52.   Ruifrok AC, Johnston DA. Quantification of histochemical staining by color
deconvolution. *Anal Quant Cytol Histol*. Aug 2001;23(4):291-9.
53.   Zarate N, Intihar TA, Yu D, et al. Heat Shock Factor 1 Directly Regulates Postsynaptic
Scaffolding PSD-95 in Aging and Huntington's Disease and Influences Striatal Synaptic Density.
Int J Mol Sci. Dec 04 2021;22(23)doi:10.3390/ijms222313113
54.   Zarate N, Gundry K, Yu D, et al. Neurochemical correlates of synapse density in a
Huntington's disease mouse model. *J Neurochem*. Jan 2023;164(2):226-241.
doi:10.1111/jnc.15714
55.   McKinstry SU, Karadeniz YB, Worthington AK, et al. Huntingtin is required for normal
excitatory synapse development in cortical and striatal circuits. *J Neurosci*. Jul 09
2014;34(28):9455-72. doi:10.1523/JNEUROSCI.4699-13.2014
56.   Pitts MW. Barnes Maze Procedure for Spatial Learning and Memory in Mice. *Bio Protoc*.
Mar 05 2018;8(5)doi:10.21769/bioprotoc.2744
57.   Illouz T, Madar R, Clague C, Griffioen KJ, Louzoun Y, Okun E. Unbiased classification of
spatial strategies in the Barnes maze. *Bioinformatics*. Nov 01 2016;32(21):3314-3320.
doi:10.1093/bioinformatics/btw376
58.   Barnes CA. Memory deficits associated with senescence: a neurophysiological and
behavioral study in the rat. *J Comp Physiol Psychol*. Feb 1979;93(1):74-104.
doi:10.1037/h0077579
59.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550. doi:10.1186/s13059-014-0550-8
60.   Raudvere U, Kolberg L, Kuzmin I, et al. g:Profiler: a web server for functional enrichment
analysis and conversions of gene lists (2019 update). *Nucleic Acids Res*. Jul 02
2019;47(W1):W191-W198. doi:10.1093/nar/gkz369
61.   Young K, Morrison H. Quantifying Microglia Morphology from Photomicrographs of
Immunohistochemistry Prepared Tissue Using ImageJ. *J Vis Exp*. Jun 05
2018;(136)doi:10.3791/57648

62. Arganda-Carreras I, Fernández-González R, Muñoz-Barrutia A, Ortiz-De-Solorzano C.
3D reconstruction of histological sections: Application to mammary gland tissue. *Microsc Res*
Tech. Oct 2010;73(11):1019-29. doi:10.1002/jemt.20829
63. Miller JA, Guillozet-Bongaarts A, Gibbons LE, et al. Neuropathological and
transcriptomic characteristics of the aged brain. *Elife*. Nov 09 2017;6doi:10.7554/eLife.31126
64. Walker CK, Greathouse KM, Boros BD, et al. Dendritic Spine Remodeling and Synaptic
Tau Levels in PS19 Tauopathy Mice. *Neuroscience*. Feb 10 2021;455:195-211.
doi:10.1016/j.neuroscience.2020.12.006
65. Takeuchi H, Iba M, Inoue H, et al. P301S mutant human tau transgenic mice manifest
early symptoms of human tauopathies with dementia and altered sensorimotor gating. *PLoS*
One. 2011;6(6):e21050. doi:10.1371/journal.pone.0021050
66. Zhang B, Carroll J, Trojanowski JQ, et al. The microtubule-stabilizing agent, epothilone
D, reduces axonal dysfunction, neurotoxicity, cognitive deficits, and Alzheimer-like pathology in
an interventional study with aged tau transgenic mice. *J Neurosci*. Mar 14 2012;32(11):3601-11.
doi:10.1523/JNEUROSCI.4922-11.2012
67. Zhou Y, Xu J, Hou Y, et al. The Alzheimer's Cell Atlas (TACA): A single-cell molecular
map for translational therapeutics accelerator in Alzheimer's disease. *Alzheimers Dement (N Y)*.
2022;8(1):e12350. doi:10.1002/trc2.12350
68. Gerrits E, Brouwer N, Kooistra SM, et al. Distinct amyloid-β and tau-associated microglia
profiles in Alzheimer's disease. *Acta Neuropathol*. May 2021;141(5):681-696.
doi:10.1007/s00401-021-02263-w
69. Otero-Garcia M, Mahajani SU, Wakhloo D, et al. Molecular signatures underlying
neurofibrillary tangle susceptibility in Alzheimer's disease. *Neuron*. Sep 21 2022;110(18):2929-
2948.e8. doi:10.1016/j.neuron.2022.06.021
70. Leng K, Li E, Eser R, et al. Molecular characterization of selectively vulnerable neurons
in Alzheimer's disease. *Nat Neurosci*. Feb 2021;24(2):276-287. doi:10.1038/s41593-020-00764-
7
71. Lau SF, Cao H, Fu AKY, Ip NY. Single-nucleus transcriptome analysis reveals
dysregulation of angiogenic endothelial cells and neuroprotective glia in Alzheimer's disease.
Proc Natl Acad Sci U S A. Oct 13 2020;117(41):25800-25809. doi:10.1073/pnas.2008762117
72. Marshall CA, McBride JD, Changolkar L, Riddle DM, Trojanowski JQ, Lee VM. Inhibition
of CK2 mitigates Alzheimer's tau pathology by preventing NR2B synaptic mislocalization. *Acta*
Neuropathol Commun. 03 04 2022;10(1):30. doi:10.1186/s40478-022-01331-w
73. Sahara N, Yanai R. Limitations of human tau-expressing mouse models and novel
approaches of mouse modeling for tauopathy. *Front Neurosci*. 2023;17:1149761.
doi:10.3389/fnins.2023.1149761
74. Choi YB, Dunn-Meynell AA, Marchese M, et al. Erythropoietin-derived peptide treatment
reduced neurological deficit and neuropathological changes in a mouse model of tauopathy.
Alzheimers Res Ther. Jan 27 2021;13(1):32. doi:10.1186/s13195-020-00766-4
75. DeVos SL, Miller RL, Schoch KM, et al. Tau reduction prevents neuronal loss and
reverses pathological tau deposition and seeding in mice with tauopathy. *Sci Transl Med*. Jan 25
2017;9(374)doi:10.1126/scitranslmed.aag0481
76. Wagner J, Krauss S, Shi S, et al. Reducing tau aggregates with anle138b delays
disease progression in a mouse model of tauopathies. *Acta Neuropathol*. Nov 2015;130(5):619-
31. doi:10.1007/s00401-015-1483-3
77. Crescenzi R, DeBrosse C, Nanga RP, et al. In vivo measurement of glutamate loss is
associated with synapse loss in a mouse model of tauopathy. *Neuroimage*. Nov 01
2014;101:185-92. doi:10.1016/j.neuroimage.2014.06.067
78. Zheng N, Li K, Cao J, et al. Electrophysiology-based screening identifies neuronal HtrA
serine peptidase 2 (HTRA2) as a synaptic plasticity regulator participating in tauopathy. *Transl*
Psychiatry. Jan 10 2025;15(1):5. doi:10.1038/s41398-025-03227-4

79.   El Fatimy R, Li S, Chen Z, et al. MicroRNA-132 provides neuroprotection for tauopathies
via multiple signaling pathways. *Acta Neuropathol*. Oct 2018;136(4):537-555.
doi:10.1007/s00401-018-1880-5
80.   Litvinchuk A, Wan YW, Swartzlander DB, et al. Complement C3aR Inactivation
Attenuates Tau Pathology and Reverses an Immune Network Deregulated in Tauopathy Models
and Alzheimer's Disease. *Neuron*. Dec 19 2018;100(6):1337-1353.e5.
doi:10.1016/j.neuron.2018.10.031
81.   Kim J, Selvaraji S, Kang SW, et al. Cerebral transcriptome analysis reveals age-
dependent progression of neuroinflammation in P301S mutant tau transgenic male mice. *Brain
Behav Immun*. Aug 2019;80:344-357. doi:10.1016/j.bbi.2019.04.011
82.   Chen X, Firulyova M, Manis M, et al. Microglia-mediated T cell infiltration drives
neurodegeneration in tauopathy. *Nature*. Mar 2023;615(7953):668-677. doi:10.1038/s41586-
023-05788-0
83.   Odfalk KF, Bieniek KF, Hopp SC. Microglia: Friend and foe in tauopathy. *Prog Neurobiol*.
Sep 2022;216:102306. doi:10.1016/j.pneurobio.2022.102306
84.   Gao C, Jiang J, Tan Y, Chen S. Microglia in neurodegenerative diseases: mechanism
and potential therapeutic targets. *Signal Transduct Target Ther*. Sep 22 2023;8(1):359.
doi:10.1038/s41392-023-01588-0
85.   Fixemer S, Ameli C, Hammer G, et al. Microglia phenotypes are associated with
subregional patterns of concomitant tau, amyloid-β and α-synuclein pathologies in the
hippocampus of patients with Alzheimer's disease and dementia with Lewy bodies. *Acta
Neuropathol Commun*. Mar 16 2022;10(1):36. doi:10.1186/s40478-022-01342-7
86.   Leyh J, Paeschke S, Mages B, et al. Classification of Microglial Morphological
Phenotypes Using Machine Learning. *Front Cell Neurosci*. 2021;15:701673.
doi:10.3389/fncel.2021.701673
87.   Hassan-Abdi R, Brenet A, Bennis M, Yanicostas C, Soussi-Yanicostas N. Neurons
Expressing Pathological Tau Protein Trigger Dramatic Changes in Microglial Morphology and
Dynamics. *Front Neurosci*. 2019;13:1199. doi:10.3389/fnins.2019.01199
88.   Hong S, Beja-Glasser VF, Nfonoyim BM, et al. Complement and microglia mediate early
synapse loss in Alzheimer mouse models. *Science*. May 06 2016;352(6286):712-716.
doi:10.1126/science.aad8373
89.   Dejanovic B, Huntley MA, De Mazière A, et al. Changes in the Synaptic Proteome in
Tauopathy and Rescue of Tau-Induced Synapse Loss by C1q Antibodies. *Neuron*. Dec 19
2018;100(6):1322-1336.e7. doi:10.1016/j.neuron.2018.10.014
90.   Lui H, Zhang J, Makinson SR, et al. Progranulin Deficiency Promotes Circuit-Specific
Synaptic Pruning by Microglia via Complement Activation. *Cell*. May 05 2016;165(4):921-35.
doi:10.1016/j.cell.2016.04.001
91.   Pampuscenko K, Morkuniene R, Krasauskas L, Smirnovas V, Brown GC, Borutaite V.
Extracellular tau stimulates phagocytosis of living neurons by activated microglia via Toll-like 4
receptor-NLRP3 inflammasome-caspase-1 signalling axis. *Sci Rep*. Jul 04 2023;13(1):10813.
doi:10.1038/s41598-023-37887-3
92.   Parhizkar S, Bao X, Chen W, et al. Lemborexant ameliorates tau-mediated sleep loss
and neurodegeneration in males in a mouse model of tauopathy. *Nat Neurosci*. May 27
2025;doi:10.1038/s41593-025-01966-7
93.   Roy ER, Chiu G, Li S, et al. Concerted type I interferon signaling in microglia and neural
cells promotes memory impairment associated with amyloid β plaques. *Immunity*. May 10
2022;55(5):879-894.e6. doi:10.1016/j.immuni.2022.03.018
94.   El Hajj H, Savage JC, Bisht K, et al. Ultrastructural evidence of microglial heterogeneity
in Alzheimer's disease amyloid pathology. *J Neuroinflammation*. Apr 16 2019;16(1):87.
doi:10.1186/s12974-019-1473-9

95.   Möller K, Brambach M, Villani A, Gallo E, Gilmour D, Peri F. A role for the centrosome in regulating the rate of neuronal efferocytosis by microglia in vivo. *Elife*. Nov 18 2022;11doi:10.7554/eLife.82094

96.   Pérez M, Avila J. The expression of casein kinase 2alpha' and phosphatase 2A activity. *Biochim Biophys Acta*. Mar 08 1999;1449(2):150-6. doi:10.1016/s0167-4889(99)00008-7

97.   Liu J, Tian J, Xie R, Chen L. CK2 inhibitor DMAT ameliorates spinal cord injury by increasing autophagy and inducing anti-inflammatory microglial polarization. *Neurosci Lett*. May 14 2023;805:137222. doi:10.1016/j.neulet.2023.137222

98.   Mishra S, Kinoshita C, Axtman AD, Young JE. Evaluation of a Selective Chemical Probe Validates That CK2 Mediates Neuroinflammation in a Human Induced Pluripotent Stem Cell-Derived Microglial Model. *Front Mol Neurosci*. 2022;15:824956. doi:10.3389/fnmol.2022.824956

99.   Chen IY, Chang SC, Wu HY, et al. Upregulation of the chemokine (C-C motif) ligand 2 via a severe acute respiratory syndrome coronavirus spike-ACE2 signaling pathway. *J Virol*. Aug 2010;84(15):7703-12. doi:10.1128/JVI.02560-09

100.   Larson SR, Bortell N, Illies A, Crisler WJ, Matsuda JL, Lenz LL. Myeloid Cell CK2 Regulates Inflammation and Resistance to Bacterial Infection. *Front Immunol*. 2020;11:590266. doi:10.3389/fimmu.2020.590266

101.   Drygin D, Ho CB, Omori M, et al. Protein kinase CK2 modulates IL-6 expression in inflammatory breast cancer. *Biochem Biophys Res Commun*. Nov 11 2011;415(1):163-7. doi:10.1016/j.bbrc.2011.10.046

102.   Parhar K, Morse J, Salh B. The role of protein kinase CK2 in intestinal epithelial cell inflammatory signaling. *Int J Colorectal Dis*. Jun 2007;22(6):601-9. doi:10.1007/s00384-006-0193-7

103.   Yamada M, Katsuma S, Adachi T, et al. Inhibition of protein kinase CK2 prevents the progression of glomerulonephritis. *Proc Natl Acad Sci U S A*. May 24 2005;102(21):7736-41. doi:10.1073/pnas.0409818102

104.   Huang J, Li J, Chen Z, et al. Sphingosine kinase 1 mediates diabetic renal fibrosis via NF-κB signaling pathway: involvement of CK2α. *Oncotarget*. Oct 24 2017;8(51):88988-89004. doi:10.18632/oncotarget.21640

105.   Singh NN, Ramji DP. Protein kinase CK2, an important regulator of the inflammatory response? *J Mol Med (Berl)*. Aug 2008;86(8):887-97. doi:10.1007/s00109-008-0352-0

106.   Avila J, Ulloa L, González J, Moreno F, Díaz-Nido J. Phosphorylation of microtubule-associated proteins by protein kinase CK2 in neuritogenesis. *Cell Mol Biol Res*. 1994;40(5-6):573-9.

107.   Coomans EM, Schoonhoven DN, Tuncel H, et al. In vivo tau pathology is associated with synaptic loss and altered synaptic function. *Alzheimers Res Ther*. Feb 05 2021;13(1):35. doi:10.1186/s13195-021-00772-0

108.   Vogels T, Murgoci AN, Hromádka T. Intersection of pathological tau and microglia at the synapse. *Acta Neuropathol Commun*. Jul 05 2019;7(1):109. doi:10.1186/s40478-019-0754-y

109.   Bohana-Kashtan O, Pinna LA, Fishelson Z. Extracellular phosphorylation of C9 by protein kinase CK2 regulates complement-mediated lysis. *Eur J Immunol*. Jun 2005;35(6):1939-48. doi:10.1002/eji.200425716

110.   Pelloux S, Thielens NM, Hudry-Clergeon G, Pétillot Y, Filhol O, Arlaud GJ. Identification of a cryptic protein kinase CK2 phosphorylation site in human complement protease Clr, and its use to probe intramolecular interaction. *FEBS Lett*. May 13 1996;386(1):15-20. doi:10.1016/0014-5793(96)00403-6

111.   Latif-Hernandez A, Yang T, Butler RR, et al. A TrkB and TrkC partial agonist restores deficits in synaptic function and promotes activity-dependent synaptic and microglial transcriptomic changes in a late-stage Alzheimer's mouse model. *Alzheimers Dement*. Jul 2024;20(7):4434-4460. doi:10.1002/alz.13857

112.   Soni DM, Lin PB, Lee-Gosselin A, et al. Inpp5d haplodeficiency alleviates tau pathology
in the PS19 mouse model of Tauopathy. *Alzheimers Dement*. Jul 2024;20(7):4985-4998.
doi:10.1002/alz.14078
113.   Mudaliar D, Mansky RH, White A, et al. Discovery of a CK2α'-Biased ATP-Competitive
Inhibitor from a High-Throughput Screen of an Allosteric-Inhibitor-Like Compound Library. *ACS*
Chem Neurosci. Jun 22 2024;doi:10.1021/acschemneuro.4c00062

1413

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9