TITLE: If memory serves: Multisensory male display improves female memory during

mate sampling

AUTHORS: Olivia-Rose M. Hamilton[1,2], Logan S. James[3,4], Rachel A Page[4], Michael J.

Ryan[3,4], Ryan C. Taylor[1,4], Kimberly L. Hunter[1]

Olivia-Rose Hamilton Email: hamiltonoliviarose@gmail.com

[1] Department of Biological Sciences, Salisbury University, Salisbury, MD 21801, USA.

[2] Department of Biology, Wake Forest University, Winston-Salem, NC 27109, USA.

[3] Department of Integrative Biology, University of Texas, Austin, TX 78712, USA.

[4] Smithsonian Tropical Research Institute, Balboa, Ancón, Panamá.

LAY SUMMARY:

Multisensory signals function to stimulate one or more of a receiver's senses, and in

doing so, may make signalers more memorable to receivers. Multisensory signals are

common in mating displays, including the advertisement of a male túngara frog: a call

(acoustic) and a vocal sac inflation (visual). Here, we demonstrate that female túngara

frogs were more likely to remember a multisensory male display over a unisensory

display, but only after fluctuations in the signal's intensity.

ABSTRACT:

Across many taxa, males gather in leks to perform multisensory courtship displays for

females. At a lek, changes in the sensory scene over the course of mate evaluation are

inevitable. This dynamic nature makes a female's ability to recall the location of

individual signalers an important component of female mate choice. It is hypothesized

that complex (especially multimodal) displays may improve a female's ability to

remember and thus discriminate amongst potential mates. To test this hypothesis, we

presented female túngara frogs (*Physalaemus* (=*Engystomops*) *pustulosus*) with male

calls (auditory) and robotic frogs (visual) that could later be obstructed from view via the

lowering of blinds. Specifically, we asked if the visual component of a multimodal

display improves the ability of a receiver to remember a signaler. While our initial

experiment did not find memory instantiation in female frogs, subsequent experiments

with a longer presentation time and a brief period of silence successfully instantiated

memories in females. Moreover, females continued to demonstrate significant

preferences for calls associated with the visual cue even after 25 s following the

obstruction of the visual stimulus (robotic frog). Thus, we propose that the duration of

presentation and/or the silent period impact memory capability. Silence is common in

choruses, and our data suggest that complex, multisensory stimuli may have evolved to

help females remember their preferred mate even with such fluctuations in signal

production.

ACKNOWLEDGEMENTS:

We thank Jorge López and Leslie Barría for their contribution with both field work and

data collection. We also want to thank Gregg Cohen and Luke Larter for their support

during field seasons. Paul Clements supported our team by designing the robotic units

and Christopher Maness aided with graphic design. Nana Osei-Wusu served as the blind

observer for scoring videos. Golden Frog Scuba provided compressed air cylinders that

powered the pneumatic robofrog control units. The Smithsonian Tropical Research

Institute provided logistical support and laboratory space. We are grateful to the Republic

of Panamá for hosting our team.

INTRODUCTION:

Animal communication signals constitute some of the most diverse traits observed

in nature, many of which function in mate attraction (Bradbury & Vehrencamp 2011;

Rosenthal 2017). Signaling reproductive availability can be achieved through a variety of

behaviors and sensory modalities. The calls of crickets on a warm summer evening

(Olvido & Wagner 2004), the low frequency "bellows" of a koala (Ellis et al. 2011), and

the elaborate design of a male pufferfish's nest (Kawase et al. 2013) are all produced in

efforts to procure mates. Because females often decide the outcome of a reproductively

motivated interaction, their decisions have critical fitness consequences (Andersson &

Simmons 2006). For a signaler's stimulus to be recognized, however, the nervous system

of the receiver must integrate and analyze components of the signal and assign those

components to their source (Bee & Miller 2016). Thus, the selective force of the

receiver's psychology, and the choice that is made as a result, is an important factor

driving signal evolution (Guilford & Dawkins 1991).

Often, animals communicate using more than one sensory modality (Higham &

Hebets 2013; Patricelli & Hebets 2016). During courtship displays, male butterflies, for

example, use both chemical and visual cues (Costanzo & Monteiro 2007), while wolf

spiders rely on visual and seismic signals (Kozak & Uetz 2019; McGinley et al. 2023).

Male sagebrush lizards combine chemical and visual cues to both defend their territory

and attract females (Thompson et al. 2008). Similarly, females prefer the visual display of

a wingspread paired with an acoustic song over either stimulus alone in the brown-

headed cowbird (Ronald et al. 2017). Bats, too, have been documented using multimodal

courtship displays in which males use chemical, acoustic, and visual signals (Voigt et al.

2008). Simultaneous communication channels in different modalities are often presumed

to provide more information, thereby improving a female's ability to make critical mating

decisions (Rosenthal 2017). It has therefore been predicted that the additional information

in a multimodal display should aid the female's ability to distinguish among individuals,

particularly in noisy leks (Rowe 1999). Multisensory displays may also be more

perceptible by a receiver, ultimately becoming more detectable in the environment.

Utilizing a multimodal display may therefore confer benefits to males to make

themselves more memorable when females are choosing among multiple options

(Guilford & Dawkins 1991; Rowe 1999). Indeed, some studies have provided evidence

for multicomponent displays improving a receiver's memory both within and outside the

context of mate choice (Rowe & Guilford 1999; Pardasani et al. 2021). In species with

lekking behavior, the memorability of a signaler may be even more pertinent as females

sample multiple males and make a choice within a relatively short time span (Backwell &

Passmore 1996; Schwartz et al. 2004; Pauli & Lindström 2021). There are also periodic

lapses in signal production (usually acoustic) (Greenfield 2015; Coss et al. 2020), so that

during these quiet periods, females may have to rely on memory to make a choice or

delay their decision. The "memory duration increase hypothesis" (Zhu et al. 2021) posits

that eliciting increased memorability in receivers may contribute to the evolution of

increased display complexity and multimodality.

Because females often assess multiple mates, learning and memory are important

factors for female choice (Rowe 1999; Bateson & Healy 2005; Ryan et al. 2009), even on

short time scales. For example, it is known that prior experience with conspecifics can

influence mating decisions (Schlupp et al. 1994; Hebets 2003; Coleman et al. 2004;

Cheetham et al. 2007; Bailey & Zuk 2009). Though several review papers have alluded to

the possibility of learning and memory playing a role in the evolution of complex signals

(Bateson & Healy 2005; Hebets & Papaj 2005; Bro-Jørgensen 2009; Higham & Hebets

2013; Lynch 2017; Ryan et al. 2019a; MacGillavry et al. 2023), few studies have

experimentally tested this hypothesis. There are numerous studies across taxa that have

investigated the role of spatial memory during mate searching in males (Jones et al. 2003;

Goh & Morse 2010; Foley et al. 2015), but our knowledge about the general role of

memory during mate searching in females is not as extensive (Healy & Hurly 2004).

Specifically, one aspect of female mate choice that has received little attention is the role

of multimodal cues in instantiating memory for signaler location. Females are likely to

learn and retain information about a potential mate (location, size, signaling ability, etc.)

so that they can recall it later. For instance, female house mice are able to recall the

previous location of a male pheromone for 14 days (Roberts et al. 2012). For females that

sample multiple males, especially over longer periods of time and across territories, it

may be advantageous to remember information about each signaler to make a later

choice, whether that be seconds from an initial contact or years.

The role of complex cues in memory during mate choice has primarily been

performed using anurans. Anuran communication encompasses a diversity of display

behaviors, with some species exhibiting visual, acoustic, chemical, and/or seismic signals

and cues that are attended to by females (Gerhardt & Huber 2002; Höbel & Kolodziej

2013; Starnberger et al. 2014a; Robertson & Greene 2017; Caldwell et al. 2022; James et

al. 2022). Despite this variety, acoustic signals remain the primary modality for most

anuran communication, though these too can vary in their complexity (Reichert & Höbel

2018). In one of the studies investigating memory, Akre & Ryan (2010) found that

female túngara frogs were able to recall the location of a more complex call for up to 45 s

after cessation of the call. Yet, when presented with a simpler call, this effect was no

longer exhibited, suggesting that females are able to maintain a working memory for the

location of individual male callers. However, this study focused on the unimodal,

acoustic component of the otherwise multimodal túngara frog mating display (Taylor et

al. 2008). In a more recent study, Zhu et al. (2021) investigated the role of a

multicomponent cue on the memory of the serrate-legged small treefrog (*Kurixalus*

odontotarsus) and found that the multimodal display of a vocal sac inflation (via video

playback) and an acoustic signal increased the active memory of females compared to a

unimodal display. Because multimodal cues have been suggested to improve the memory

of a receiver (Rowe 1999), these studies provide evidence that adding complexity to

displays may improve the ability of a receiver to recall the last detected location of a

preferred signaler. In the present study, we again rely on anuran communication to

further our understanding of multimodal displays and memory.

Túngara frogs are a Neotropical species, with males gathering in nocturnal

choruses to form leks (Ryan 1985). The acoustic stimulus of their mating display

involves a simple "whine", which may have up to seven additional "chucks"

accompanying the whine (Ryan 1985). Females show a strong and consistent preference

for a more complex call (whine plus chuck) (Ryan et al. 2019b). In addition to their

acoustic stimulus, males produce cues in other modalities that can attract females.

Consisting of an acoustic call, a visual cue, and a seismic component, the multimodal

display of a male túngara frog is preferred over the acoustic call alone (Taylor et al. 2008;

James et al. 2022). The male's inflating vocal sac likely evolved to maximize calling

efficiency, and its inflation secondarily became integrated into their mating display as the

visual stimulus (Pauly et al. 2006; Taylor et al. 2008). The inflation of the vocal sac

occurs synchronously with sound production during calling (Taylor et al. 2011). A

túngara frog's vocal sac is exceptionally large and, when inflated, is nearly the size of the

male itself (Dudley & Rand 1991). In complex environments like the rainforests of

Central America, the motion of a visual stimulus may be used by females to improve

their ability to localize callers (Tan & Elgar 2021).

It is known that túngara frogs will stop calling in response to a predation threat

(Dapper et al. 2011). Interruptions in the chorus are therefore common, with inter-chorus

intervals having an average duration of 25 s (Akre & Ryan 2010). Because eavesdropping

predators attend to the multimodal displays of males at a túngara frog lek (Halfwerk et al.

2014), females are vulnerable to predation while mate sampling (Bernal et al. 2007). It

may be beneficial, then, for a female to evaluate males and make a choice quickly to

mitigate predation risk. Experiments in both laboratory and natural settings seem to

corroborate this (Rand et al. 1997). Female anurans tend to assess males and choose

within a very short time period, with gray treefrogs (*Dryophytes (=Hyla) versicolor*)

assessing potential mates for only about two minutes (Schwartz et al. 2004; Feagles &

Höbel 2022). In dynamic lek environments, however, it is common for a male's display

to become interrupted or go silent during a female's assessment. In that scenario, does

she wait and restart her assessment once the chorus resumes, or does she rely on memory

to influence her choice? As Akre & Ryan (2010) demonstrated, female túngara frogs are

able to remember the location of a complex call for up to 45 s, spanning the average

inter-chorus interval shown by the same study. Túngara frogs have also been shown to

associate a visual cue with a place-learning task (Liu & Burmeister 2017; Burmeister

2022), suggesting they may depend more on landmark cues (i.e. a marker physically

linked to a "goal") than spatial cues (i.e. a marker that guides towards a "goal") when

remembering locations. It is possible, then, that the visual cue of an inflating vocal sac

aids females in remembering a male's call and, therefore, his location.

Here, we tested the hypothesis that the visual stimulus of an inflating male's vocal

sac instantiates female túngara frogs' working memory for the stimulus, compared with a

call only. In addition, we tested call playbacks with and without a silent period to better

mimic natural settings and to understand how silence influences memory instantiation.

METHODS:

General Experimental Procedures

All behavioral experiments were performed at facilities of the Smithsonian

Tropical Research Institute in Gamboa, Panamá. We collected amplexed pairs of frogs

from wild choruses between 1930 – 2300 h during the rainy season in 2021 and 2022.

After transporting them to our lab for testing, frogs were stored in total darkness for at

least one hour prior to trials to allow their eyes to dark-adapt. This dark adaption is

necessary after collecting frogs using flashlights. For each trial, we separated a female

from her mate and placed her under a visually and acoustically transparent mesh funnel in

the center of a sound chamber (ETS-Lindgren, Austin, TX, USA).

Light inside the chamber was produced via a GE brand nightlight to mimic light

conditions within the range of natural breeding conditions. The nightlight was green to

the human observer, consisting of a broad spectral peak around 510 nm. This spectrum is

within the range of what túngara frogs experience at breeding sites within forests and

along forest edges (Taylor et al. 2008). We used tape to cover most of the nightlight

surface, yielding a light-level of ~ 5.5×10^{-10} W/cm^2. Light levels in the natural

environment fluctuate widely depending on moon phase and cloud cover. Our

experimental light levels were on the lower end of what the frogs experience in nature

and within the range of the frogs' scotopic visual sensitivity (Cummings et al. 2008;

Leslie et al. 2021). The acoustic stimulus of each treatment was broadcast by a pair of

Orb Audio speakers (Sherman Oaks, CA, USA; 1.2 – 18 kHz) that were placed 80 cm

away from the release point of the female, forming an equilateral triangle (Fig. 1). For all

experiments, the speakers antiphonally broadcast identical, synthetic calls (whine-chuck).

A synthetic whine-chuck is, on average, as attractive as a natural call (Rand et al. 1992).

We used Adobe Audition v22.2 to play call files. Once the female was placed inside the

funnel, we presented the stimuli of each treatment ("presentation period") for two to three

minutes before the female was either 1) immediately released or 2) exposed to additional

testing conditions and then subsequently released. After the female was released, we

monitored and recorded her choice from outside the sound chamber using live infrared

cameras. The video files were saved for an observer (blind to the treatments) to later

rescore. We defined a choice as a female remaining within 8 cm of the base of a speaker

(the "choice zone") for at least 3 s. If a female failed to move from her initial "funnel

zone" within 4 minutes or did not make a choice within 10 minutes, we removed her

from the chamber. After a period of at least 15 minutes, the female was re-tested, and if

she failed a second time, she was set aside and not tested again.

We aimed to minimize our effect on the animals and their environment during

testing. After completion of a trial, we reunited the female with her mate and later toe-

clipped the pair, both to mark the frog (to avoid retesting the same individual) and to

obtain a genetic sample. Toes were preserved in ethanol for future analyses. If females

did not make a choice in an experiment, they were not toe-clipped. We followed

regulations from the American Society of Ichthyologists and Herpetologists regarding

toe-clipping procedures (Beaupre et al. 2004; Funk et al. 2005). Snout-vent-length and

mass were also recorded for each frog. At the end of the night, all pairs were returned to

their original collecting site and released. All frogs tested were unique individuals;

females were not retested across treatments.

Control: Robofrog Replication

Following Taylor et al. (2008), we replicated a multimodal stimulus for mate

choice experiments using speakers and robotic frogs with artificial vocal sacs (Fig. 2a)

inflated by an electromechanical pump (Fig. 2b). The robofrogs provided the visual

component of the multimodal display while the call was produced by the speaker

immediately behind the robofrog (Fig. 1). The pump was programmed to shunt air into

tubing connected to the robofrog so that a silicone vocal sac inflated at a 19 kHz tone and

deflated at a 16 kHz tone. These tones were placed at the respective beginning and end of

the synthetic call in Adobe Audition. This ensured that the robofrogs inflated in time with

the acoustic stimuli, as occurs in nature (Ryan 1985). These tones are imperceptible to

frogs (Ryan et al. 1990). We calibrated the sound-pressure level (SPL) for both speakers

at 82 dB SPL (re. 20 µPa, fast C-weighting) from the female's position within the funnel.

In this experiment, a robofrog was placed in front one of the speakers (right or left), and

its position was switched between trials. During a two-minute presentation period, both

speakers played a complex (whine-chuck) acoustic stimulus and the robofrog inflated

synchronously with the speaker behind it. As both the acoustic and visual stimuli

continued to play, the funnel was gently lifted from outside the chamber so that the

female was free to make a choice. In addition to the definition of a choice used earlier,

we also defined a choice in this experiment as the female approaching within 5 cm of the

robofrog or touching it. As the robofrog was placed in front of a speaker, it extended the

radius of the "choice zone" beyond that of a bare speaker. We note that the vocal sac

movement (rather than the mere presence of the model frog) has been shown to be the

critical aspect of the visual component, and must be closely timed with the acoustic call

to induce a preference (Taylor et al. 2008; Taylor et al. 2011).

We tested a total of 46 individual females in this treatment. Thirty-two were

tested in 2021. Data show that some climatic events may alter female túngara frog

responses across years (Cronin et al. 2024). As such, we tested an additional 14 females

in the 2022 robofrog control treatment to ensure that patterns of mate choice remained

stable across the two years. All other experimental treatments were conducted in 2022.

Memory Tests

For all subsequent treatments, we tested whether females employ their memory of

the visual stimulus of a calling male (robofrog) that is suddenly obstructed from view

when choosing between two sound sources. An opaque, black cloth served as a blind to

block the visual cue from all angles (Fig. 1). The blind wrapped around both the speaker

and the robofrog (if present) 360° to completely obscure the female's ability to see the

speaker/robofrog, irrespective of her position in the test chamber. We designed the blind

to be as tall as the speakers (13 cm) so that the females could not see anything that was

hidden behind them. The cloth was hand-sewn to a metal ring, which was then tied to

monofilament fishing line. The fishing line was accessible to the outside of the chamber

through a small port that also allowed access for speaker cables. With the blind

surrounding the speakers, the amplitude at the female's release point was 76 dB SPL; a

decrease of 6 dB from a bare speaker without the blind. This amplitude is still within the

range of what females experience in nature and has been used in previous studies in

which females continue to exhibit robust preferences (Stange et al. 2016). In addition,

both speakers were always enclosed within the blind, thus the drop in amplitude was

identical for both speakers. During the presentation period (initial playback to a female),

the blinds hung unmoving from the ceiling of the chamber. We then lowered the blinds to

obscure the robofrog. As the blinds were lowered, an observer at the live camera feed

monitored their rate and speed to ensure they were even and touched the floor of the

chamber simultaneously. The lowering of the blinds took approximately 7 s. Because the

original choice zone used for the speaker was engulfed by each blind, we defined a

choice as a female approaching within approximately 8 cm of a blind and remaining there

for at least 3 s. Despite the fact that the robofrog was occluded by the blind, we still

halted the inflation and deflation of the vocal sac whenever the blinds were lowered.

Control: Blinds

We first confirmed that females would continue to demonstrate typical phonotaxis

behavior despite the speakers being occluded by the blinds. For this control experiment

(n=13), each speaker played an identical whine-chuck call for the entirety of the

experiment. Our sample size for this control was smaller than other treatments, as we

were limited by time during our field season. We did not utilize a robofrog as we were

simply verifying that females would exhibit phonotaxis behavior and that there was no

aversion or bias for one cloth over another. After the presentation period, the blinds were

lowered, and once the blinds touched the floor, the female was immediately released

from the funnel.

Memory: Hold and/or Silence

All memory tests began with a presentation period, during which the speakers

antiphonally broadcast identical whine-chuck calls. The presentation period lasted 2 min

for the non-silent treatments and 3 min for the silent treatments. The increase in time for

the silent treatments was to ensure that a data collector could locate the monofilament

lines in a dark laboratory and be prepared to lower the blinds in a consistent manner.

Traditionally, a 3 min acclimation period has been utilized in túngara frog phonotaxis

studies (Bosch et al. 2000; Marsh et al. 2000; Baugh & Ryan 2011). One robofrog

inflated in time with the call of the speaker it was paired with and was visible to the

female during this period. We switched the robofrog between speakers across trials. After

the presentation period, the blinds were lowered and females were exposed to one of

three conditions: a silent treatment, a hold treatment, or a combined silent & hold

treatment (Fig. 3).

For the hold treatment, as soon as the blinds touched the floor, females were

retained under the funnel for 20 s but call playbacks from the speakers continued

uninterrupted (n=38). After this 20 s, the female was released and allowed to make a

choice (Fig. 3).

For the silent treatment, once the blinds touched the floor, all sound was muted

from the speakers for 5 s (n=25). After this 5 s silent period, the female was released, and

the audio (but not visual) playback began again and continued to play until the female

made a choice (Fig. 3).

For our final treatment, we combined the silent and hold conditions in sequence

(n=25). This treatment was the longest test of memory, with 25 s total time between the

lowering of the blinds and the females' release from the funnel. Females in this treatment

first received the silent period. After the 5 s silence period, the funnel remained in place

and the acoustic playback resumed. Females remained in the funnel for 20 s, as in the

hold condition, before the funnel was raised (Fig. 3). In all memory treatments, the

acoustic playback continued until the female made a choice or the 10-minute time limit

was reached.

Locomotor Behaviors

In addition to choice and time to choice (latency), we also analyzed the path

females took to reach a speaker and the number of "stops" and "turns" they made on the

way to a speaker. Using our recorded videos, we had an observer (unaware of the

hypothesis) score choice and movement behaviors. We divided the floor of the test

chamber into quadrants with the center converging on the funnel at the female's release

point (Fig. 4a). If the path the female took to reach a speaker remained within the same

quadrant as the speaker, we scored that as a "direct" path. If, at any point, the female

walked/hopped into a different quadrant than the speaker she ultimately chose, we scored

that as an "indirect" path (Fig. 4a). We defined stops as any cessation of movement for

>1 s. Turns were scored when a female changed the angle of the long-axis of her body

30° or more (Fig. 4b). Intuitively, frogs taking an indirect path are likely to exhibit higher

numbers of stops and turns (versus the direct path). Frogs taking the direct path, however,

still had sufficient space and time to express these behaviors. Locomotor behaviors have

the potential to provide additional insights into the behavior of mate searching females.

These behaviors are typically not scored in anuran phonotaxis tests (but see Feagles &

Hobel 2022; 2023).

Ethics Note:

Experimental procedures were conducted with approval from Salisbury University and

the Smithsonian Tropical Research Institute (IACUC: SU-0052; SU-0052R and STRI

2018-0411-2021; SI-21012). The Ministry of the Environment of Panamá (MiAmbiente)

approved our collection of animals and issued collecting permits for our team (ANAM:

SE/A- 39-2020). Procedures for marking animals were conducted in accordance with the

American Society of Ichthyologists and Herpetologists.

STATISTICAL ANALYSES:

We first tested the hypothesis that the proportion of choices for each treatment

were different from random chance using a binomial test. We report the mid-P-value to

smooth large jumps in p-values, as often occurs with categorical data (Agresti 2001;

Hwang & Yang 2001). Next, we used a Kruskal-Wallis test to examine how the number

of stops and turns an individual made during a trial varied depending on whether she took

a direct or indirect path to her choice. A binomial test was also used to compare the path

of females (indirect vs direct) across treatments. We used an ANOVA to compare the

mean latency to choose across treatments. To test if choice preferences were different

across years of data collection, we performed a Fisher's exact test (SISA Fisher exact

test; Uitenbroek DG 1997). All other analyses were performed in R (R Core Team 2020).

Graphs were generated in R (R Core Team 2020) using the packages: tidyverse

(Wickham et al. 2019), ggplot2 (Wickham 2016), and ggpubr (Kassambara 2023).

Figures were edited using Blender v4.2.0 and Adobe Photoshop v25.11.

RESULTS:

We tested female túngara frogs in several conditions to determine the role

memory played when choosing between two speakers, where one speaker was formerly

paired with a visual cue. Using a blind (opaque cloth), we occluded the visual stimulus

(robofrog) from the view of the female in three different memory conditions.

First, we replicated the previous finding that females ($n = 46$) preferentially

choose a speaker paired with a dynamic robofrog (multimodal display) over a bare

speaker when both speakers played the same call (whine-chuck). Because we tested frogs

in both 2021 and 2022 for this treatment, we wanted to ensure that their collection year

did not influence preference. Though we know that preferences are stable and robust

across years (Ryan et al. 2019b), a recent study demonstrated that drought can influence

females' preferences between field seasons (Cronin et al. 2024). Across the two years, the

preference ratio was 22:10 in favor of the robofrog ($p = 0.043$), and 11:3, in favor of the

robofrog ($p= 0.046$). A Fisher's exact test showed that these ratios were not significantly

different (Pearson's χ^2; $p = 0.496$), so we combined both years for the binomial test.

Overall, females expressed a significant preference for the multimodal display over the

unimodal acoustic call (binomial test: 33 robofrog:13 unimodal $p = 0.0024$; Fig. 5),

similar to what we have shown elsewhere (Taylor et al. 2008; Stange et al. 2016).

Next, we confirmed that the use of blinds had no impact on phonotaxis behavior.

Despite the decrease in call amplitude after the blinds surrounded the speakers, females

continued to demonstrate typical phonotaxis behaviors. Furthermore, there was no side

bias for one cloth over the other ($n = 13$; binomial test: 6:7 ratio; $p = 0.8953$; Fig. 5).

We ran our primary experiments testing whether females prefer a speaker where

they had previously been able to see the visual component of the robofrog. For all

memory experiments, one speaker was paired with a dynamically inflating robofrog

during the presentation period. At the end of the presentation period, we lowered the

black cloths and halted the robofrog inflation, at which point females experienced one of

three treatments: silence, hold, or a combined silence & hold.

Females in the hold treatment experienced a 2-minute presentation period and a

20 s holding period after the visual stimulus was hidden, but experienced no break in the

auditory stimulus. Data for this treatment were collected in both 2021 and 2022. The

ratios (robofrog : bare speaker) for the two years were 2:4 and 21:11. A Fisher's exact

test showed that these were not significantly different (Pearson's χ^2, $p = 0.14$). We

combined both years for the binomial test. Overall, females did not approach the

multimodal speaker significantly more often than random (binomial test: 23:15; $p =$

0.228; Fig. 5).

Females in the silent treatment ($n = 25$, all data collected in 2022) experienced a

3-minute presentation period and a 5 s silent period which began as soon as the visual

stimulus was hidden. Subsequently, the females were released as the acoustic stimuli

began playing again. Females chose the speaker that was formerly paired with the

robofrog (multimodal speaker) significantly more often than the bare speaker (binomial

test: 18:7; $p = 0.0361$; Fig. 5).

Our final group of females experienced a combined treatment ($n = 25$, all data

collected in 2022). They first experienced a 3-minute presentation period, then the 5 s

silent period, followed by a 20 s holding period where a female was held under the funnel

while calls continued to play. After the holding period, the female was released and

allowed to make a choice. Even after a total of 25 s since the robofrog was obscured,

females showed a significant preference for the speaker where they could previously

observe the robofrog model (binomial test: 18:7; $p = 0.0361$; Fig. 5).

We tested 147 individual frogs across all treatments. Six video recordings were

lost or corrupted, so locomotor behaviors were scored on the remaining 141 trials. We

categorized each trial into one of two groups: direct paths to the speaker ($n = 99$) or

indirect paths ($n = 42$). Females taking an indirect path turned significantly more often

($x\square = 11.26$ turns ± 0.97 SE) than females taking a direct path ($x\square = 5.64$ turns ± 0.33

SE) (Kruskal-Wallis H $= 27.46$; $p = <0.0001$; Fig. 6a). Similarly, females taking an

indirect path stopped significantly more often ($x\square = 20.40$ stops ± 1.67 SE) than females

taking a direct path ($x\square = 11.52$ stops ± 0.63 SE), (Kruskal-Wallis H $= 30.64$; $p =$

<0.0001; Fig. 6b). In the treatments where there was no silent period, females took a

direct path to their speaker of choice significantly more often than an indirect path

(binomial test: Robofrog Control, $p = 0.0004$; Blind + Hold, $p = 0.006$; Fig. 6c), with an

exception being the Blind Control treatment wherein stimuli were equal (binomial test: p

= 0.075; Fig. 6c). In treatments with a silent period, females did not take a direct path to

their speaker of choice significantly more often than an indirect path (binomial test: Blind

+ Silence, $p = 0.922$; Blind + Silence + Hold, $p = 0.091$; Fig. 6c).

We also investigated whether the latency of a female to make a choice was

affected by treatment type (Fig. 7). All latency data were log-transformed to meet the

assumptions of parametric tests. Treatment type did not have a significant effect on

choice latency (ANOVA: $F_{4,136} = 0.808$, $p = 0.522$). Frogs taking a direct path, however,

exhibited significantly shorter latencies ($x̄ = 46.4$ s ± 3.83 SE) than those taking an

indirect path ($x̄ = 61.45$ s ± 7.77 SE) ($t_{1,139} = 2.972$; $p = 0.004$).

DISCUSSION:

Because the accessibility of mating displays can vary both temporally and

spatially, the ability of females to recall a previously sampled mate is an important, if

underappreciated, component of mate choice. While it is to be expected that females

within a bonded pair are capable of mate recognition (Miller 1979; Clark et al. 2006;

Podraza et al. 2024), we know less about the general capability of females to recall

individual males during mate searching. When choosing among multiple potential mates,

it would be advantageous to preserve information about individuals so that females can

return to certain males and/or have that information to rely on if there is a lapse in

accessibility. For some species, this may mean preserving information for an extended

period of time (Bensch & Hasselquist 1992; Witte & Massmann 2003), while others may

only need to retain this information for less than a minute (Akre & Ryan 2010; Zhu et al.

2021). The ability to recall a single male's location becomes even more pertinent for

females attending a lek as they sample potentially dozens of other males. Amidst a

cacophony of rivals, males may increase their chance of being remembered, and therefore

chosen, simply by augmenting their display and instantiating memory in sampling

females; the noisier, brighter, more flamboyant a male is, the more memorable he may

be.

Here, to determine the effect of a multisensory display on the memory of female

túngara frogs, we asked if females could remember a "calling male" after the visual

component of the male's display – the dynamically inflating robofrog – was no longer

visually accessible. To do this, we developed a pulley system for the lowering of two

opaque black cloth blinds that visually obscured both speakers (and the robofrog) from

the females' view. After a presentation period, we gave females either a silent period of 5

s, a holding period of 20 s, or both in sequence. Once the period(s) ended, we released the

female from an acoustically and visually transparent funnel and allowed her to make a

choice.

Our results provide evidence that the addition of a visual stimulus to a male frog's

courtship call instantiates memory for the caller's location. Specifically, when females

were exposed to an additional minute of stimuli and 5 s of silence before being released,

they significantly preferred the speaker that was formerly paired with a robofrog

(multimodal speaker) over the bare speaker. However, in the hold treatment (where the

audio playback was uninterrupted), females did not exhibit a significant preference for

multimodal speaker after last being exposed to the multisensory cues 20 s before release.

Because they did seem to remember the location of the cue with exposure to 5 s of

silence, we were interested in knowing if this was simply a product of the very short, 5 s

delay since females' last exposure to the robofrog. In the combined Blind + Silence +

Hold test, females were still able to remember even after a 25 s delay (5 s of silence plus

20 s delay under the release funnel) and expressed a significant preference for the

multimodal speaker.

Here, we point out that the visual component appears to instantiate memory for a

caller's location (a significant proportion of females returned to the speaker where they

last saw the robofrog), but in all experimental treatments, females still received auditory

information about the location of the speakers. As such, the degree to which females

retained spatial memory for a caller's location is unclear. Burmeister (2022) showed that

túngara frogs were capable of learning in a two-choice spatial discrimination task, albeit

not as well as poison frogs. Regardless, the females' choices in our experiment indicate

that the visual component influenced their memory such that most females expressed

their preference for a multisensory display even when the visual component of that

display was removed.

These findings demonstrate a potential mechanism wherein more complex

displays may improve a receiver's ability to remember a signaler, as proposed by

Guilford & Dawkins (1991; Rowe 1999). This mechanism does not appear to be species

specific, as Zhu et al. (2021) demonstrated a similar effect with a species of treefrog

native to Southeast Asia. In that study, researchers were interested in learning how long a

multimodal display could instantiate memory, similar to the methods employed by Akre

& Ryan (2010) for complex calls in túngara frogs. They found that females were able to

recall a caller's location for 30 s, which was similar to the 45 s of túngara frogs (Akre &

Ryan 2010; Zhu et al. 2021). Both of those studies utilized silent periods to titrate the

length of time for which the frogs could retain their memory. However, we note that the

additional minute of presentation during treatments with a silent period (3 min) versus

those without a silent period (2 min) may also play a role in memory instantiation. It is

possible that the additional minute of presentation time influenced memory, either alone

or in conjunction with the silent period. Although a 3-min "acclimation" period has

traditionally been utilized in túngara frog phonotaxis studies (Bosch et al. 2000; Marsh et

al. 2000; Baugh & Ryan 2011), future work should aim to disentangle these potential

influences. Regardless, our results provide robust evidence that multisensory male stimuli

can instantiate memory in females for an individual signaler.

Why a silent period may play a role in memory instantiation could be due, at least

in part, to the temporal updating that female túngara frogs appear to exhibit. When male

túngara frogs continuously duet with their neighbor (or two duetting speakers in an

experiment), females seem to ignore previously available information and make rapid

assessments of their options in real-time (Baugh & Ryan 2010a). Thus, females are able

to continually modify their choice, but are therefore also sensitive to shifts in the chorus.

The sensory scene in a mating chorus is dynamic, with silence itself being a natural part

of many chorus soundscapes (Schwartz 1991; Wilson et al. 2014; Greenfield et al. 2016;

Coss et al. 2020). Male signalers commonly go silent in response to perceived predation

threats, making themselves less conspicuous (Faure & Hoy 2000), but in doing so, they

may also be unintentionally activating the female's ability to remember them. While

males are continuously calling, their location is likely discernible to females in at least

one modality. When it is already at their disposal, females do not need to retain such

spatial information until there is a lapse in the chorus, and so it is likely to be discarded in

lieu of new information (Ryan et al. 2009). The temporary cessation of sound, however,

seemed to trigger females to remember callers they previously sampled. As previous

studies have suggested, it seemed that in the absence of silence (i.e. continual calling),

females appeared to continue to update the acoustic signal information in real time and

ignore where they had previously seen the calling robofrog. Overall, there was a trend for

the non-silent treatment to choose the multimodal speaker, but the lack of silence appears

to reduce that response.

As female túngara frogs are making mate choice decisions, we commonly observe

them moving and reorienting towards a speaker each time it broadcasts a call.

Unsurprisingly, when we evaluated these locomotor behaviors, we found that females

that took an indirect path made more stops and turns. These indirect females also

displayed longer latencies prior to their choice. More notably, the number of females

taking indirect paths was significantly higher in the silent treatments than in those with

continued calling. This suggests that the task was cognitively more difficult in the silent

treatments and females were relying more on memory to make their choices. If female

túngara frogs utilize landmark cues during spatial learning, it is possible that the visual

stimulus of an inflating vocal sac, in tandem with calls, acts as a landmark cue for a

female to recall a male's location. Rather than defaulting to a random choice between two

unseen males, it seemed that females exposed to a silent period were relying on memory

to locate a potential mate whose location had been previously established.

What remains unknown is how the length of the silent period influences memory

for caller location. The silent period we employed in this study was 5 s; in nature, males

go silent for highly variable periods of time, with mean inter-chorus intervals of 25 s

(Akre & Ryan 2010; Dapper et al. 2011). Females may respond strongly to this cue and,

as a result, alter their behavior while mate sampling. When danger is sensed, instead of

completely "starting over," females may continue with their decision-making, albeit with

more caution (Bernal et al. 2007; Edomwande & Barbosa 2020; Feagles & Höbel 2022).

Previously, a female túngara frog's memory during mate evaluation had only been

tested in one modality: auditory. Akre and Ryan (2010) tested female túngara frogs'

memory for both a simpler call (whine plus one chuck) and a more complex call (whine

plus three chucks). They showed that females were only able to remember the location of

a speaker when the more complex call (whine plus three chucks) was broadcast. In the

present study, a synthetic whine and one-chuck call was used for all acoustic stimuli, and

yet, we were still able to induce memory. What both our study and Akre & Ryan (2010)

demonstrate is that in order to instantiate memory for signaler location, the display needs

to be complex in nature, whether that complexity arises from additional acoustic stimuli

(chucks) or from additional modalities (visual). What our findings also highlight is the

role that vision plays during mate choice in anurans. It has been proposed that visual

stimuli may act to attract the attention of a receiver to that signaler (Ord & Stamps 2008;

James et al 2022). In túngara frogs, not only does the visual stimulus attract the receiver's

attention, but it also acts to make the signaler more memorable. Most studies on anuran

communication focus on the importance of the acoustic component of the male's display,

but there is increasing evidence that while the acoustic signal is both necessary and

sufficient for mate attraction, the visual stimulus of the vocal sac also serves an additional

important function (Taylor & Ryan 2013; Starnberger et al. 2014b; Zhu et al. 2022).

Females themselves become more visually sensitive when in a reproductive state, which

may have evolved to aid them in mate searching and evaluation (Cummings et al. 2008;

Leslie et al. 2020). In the absence of a call, the visual component of a túngara frog male's

display does not induce mate searching behavior (Taylor et al. 2011), but here we show

that, in tandem with a call, an inflating vocal sac can induce memory for a caller's

location.

Anurans are renowned for their diversity of behavioral and life-history strategies

(Wells 2007), so it is not surprising that some species may not rely on visual cues during

communication (Li et al 2022; Augusto-Alves et al. 2024). However, we point out that Li

et al. (2022) mentioned the use of dim red light when handling frogs in their experiment.

Buchanan (1993) demonstrated that dim red light diminishes scotopic visual sensitivity in

Cope's gray treefrogs. Therefore, Li et al. (2022) may have compromised the visual

capabilities of the frogs in their experiment. We carefully calibrated our light to fall

within the range (both spectrum and intensity) of what túngara frogs experience in nature.

Additionally, all frogs remained in complete darkness before and during our experiment,

thereby preserving their visual sensitivity. While the anuran vocal sac was traditionally

thought to be related only to call production (Bucher et al. 1982; Pauly et al. 2006),

several lines of evidence point to the co-option of the vocal sac as a visual component in

the communication of a variety of species (Starnberger et al. 2014). First, several species

have been shown to preferentially respond to a vocal sac when coupled with a call

(Narins et al. 2003; Taylor et al. 2007; Taylor et al. 2008; Gomez et al. 2011; Preininger

et al. 2013; Laird et al. 2016; Zhu et al. 2021), and even when not coupled with a call

(Hirschmann et al. 2006). Second, the vocal sac varies tremendously in size and color

across species (Starnberger et al. 2014; Elias-Costa & Faivovich 2025) and in ways that

are not directly linked to sound production. In túngara frogs, the vocal sac is too large to

fully inflate unless males are floating in water; this is not true of most frogs (Halfwerk et

al. 2017). Despite its large relative size, the túngara vocal sac does not serve as a cavity

resonator (Rand & Dudley 1993) and it is too small to effectively transmit the frequencies

contained in the whine (Ryan 1985). In addition, frogs similar in size to túngaras, but

with smaller vocal sacs (e.g. *Dendropsophus ebrecatus*), produce vocalizations at roughly

the same amplitude. We have demonstrated elsewhere that female túngara frogs exhibit a

preference for the multisensory display and here again demonstrate a memory function of

this display.

Though the focus of this study was not to investigate how long a female túngara

frog is able to remember, it is important to note that the length of memory for the location

of a calling male in the present study correlates with the previous findings of Akre &

Ryan (2010). In that study, the average inter-chorus interval where males would become

silent was 25 s. In our study, the longest duration after the multimodal stimulus was last

presented was also 25 s, a combination of the previous two memory experiments'

treatment periods. When presented with a complex acoustic-only stimulus, female

túngara frogs were able to remember its location for up to 45 s (Akre & Ryan 2010); how

that duration compares to a multimodal stimulus remains unknown. In a treefrog species,

Zhu et al. (2021) demonstrated that multimodal stimuli generate a longer working

memory. Zhu et al. (2022) also showed that the presence of an inflating vocal sac plus

noise, but not either stimulus alone, restores salience to calls with missing syllables.

Future studies are needed to further compare the memory capabilities between unimodal

and multimodal stimuli and across additional taxa.

Our understanding of the active role that females play in sexual selection has

exponentially increased since the time of Darwin (Rosenthal & Ryan 2022). There are

various hypotheses to explain why females generally prefer more complex stimuli; the

receiver psychology hypothesis has provided a simple and powerful explanation for the

mechanism of female choice for multimodal displays (Rowe 1999). Interestingly, despite

being proposed more than two decades ago, the role of multimodal cues instantiating

memory in receivers has received relatively little attention. The evolutionary currency is

fitness; thus, it follows that the more complex a display the more memorable it will be

for the receiver and therefore be under positive selection. In the present study, while we

found that multimodality was important for memorability, in order for memory to be

induced, an additional minute of presentation along with a potential disturbance/predation

risk (silent period) was necessary for memory instantiation. In a lekking scenario, females

continue to make a choice despite this risk, likely so as to not lose their reproductive

investment (Bernal et al. 2007). We know that females dynamically assess males (Baugh

& Ryan 2010a,b) and will circle back to them once a "final" choice has been made (Dale

& Slavsgold 1996). An association in a female's brain between one particular male and

his location must therefore exist. Multimodal stimuli have been shown to improve both

memory and learning outside of the context of mate choice (foraging: Leonard et al.

2011; Gil-Guevara et al. 2022; aposematic signals: Speed 2000; Rowe 2002; Siddall and

Marples 2008; and song-learning: Hultsch et al. 1999; Varkevisser et al. 2022). By

utilizing a multisensory display, signalers may therefore increase their chances of being

remembered. Thus, how a signaler is chosen may be determined not by how well they

advertise quality, but simply by how memorable they are during fluctuations in a

dynamic signaling environment. Here, we used túngara frogs to better understand how

the use of multiple sensory modalities may influence a female's memory and subsequent

mate choice. Our results reflect the complex decisions females must make during mate

evaluation, especially when multiple communication modalities must be integrated and

later remembered. A receiver's cognitive ability to adapt to rapidly shifting environments

and still retain the ability to weigh attractiveness is both impressive and meaningful. The

choice a female makes has direct evolutionary consequences, and furthering our

understanding of the process of decision-making is critical for understanding the

evolution of diversity.

REFERENCES:

Agresti A. 2001. Exact inference for categorical data: recent advances and continuing controversies. Stat Med. 20:2709–2722. https://doi. org/10.1002/sim.738

Andersson M, Simmons LW. 2006. Sexual selection and mate choice. Trend Ecol & Evol. 21:296-302. https://doi.org/10.1016/j.tree.2006.03.015

Akre KL, Ryan MJ. 2010. Complexity increases working memory for mating signals. Curr Biol. 20:502-505. https://doi.org/10.1016/j.cub.2010.01.021

Augusto-Alves G, Höbel G, Toledo LF. 2024. Geographic variation in acoustic and visual cues and their potential to signal body condition in the Brazilian treefrog, *Boana albomarginata*. Behav Ecol Sociobiol. 78. https://doi.org/10.1007/s00265-024-03462-7

Backwell PRY, Passmore NI. 1996. Time constraints and multiple choice criteria in the sampling behaviour and mate choice of the fiddler crab, *Uca annulipes*. Behav Ecol Sociobiol. 38:407-416. https://doi.org/10.1007/s002650050258

Bailey NW, Zuk. 2009. Field crickets change mating preferences using remembered social information. Biol Lett. 5:449-451. https://doi.org/10.1098/rsbl.2009.0112

Bateson M, Healy SD. 2005. Comparative evaluation and its implications for mate choice. Trends Ecol Evol. 20:659-664. https://doi.org/10.1016/j.tree.2005.08.013

Baugh AT, Ryan MJ. 2010a. Mate choice in response to dynamic presentation of male

advertisement signals in túngara frogs. Anim Behav. 79:145-152.

https://doi.org/10.1016/j.anbehav.2009.10.015

Baugh AT, Ryan MJ. 2010b. Ambient light alters temporal-updating behaviour during

mate choice in a Neotropical frog. Canad J Zool. 88:448-453.

https://doi.org/10.1139/Z10-018

Baugh AT, Ryan MJ. 2011. The relative value of call embellishment in túngara frogs.

Behav Ecol Sociobiol. 65:359-367. https://doi.org/10.1007/s00265-010-1053-6

Beaupre SJ, Jacobson ER, Lillywhite HB, Zamudio K. 2004. Guidelines for use of live

amphibians and reptiles in field and laboratory research. Herpetological Animal

Care and Use Committee of the American Society of Ichthyologists and

Herpetologists.

Bee MA, Miller CT eds. 2016. Psychological mechanisms in animal communication.

Springer.

Bensch S, Hasselquist H. 1992. Evidence for active female choice in a polygynous

warbler. Anim Behav. 44:301-311. https://doi.org/10.1016/0003-3472(92)90036-

9

Bernal XE, Rand AS, Ryan MJ. 2007. Sexual differences in the behavioral response of

túngara frogs, *Physalaemus pustulosus*, to cues associated with increased

predation risk. Ethology. 113:755-763. https://doi.org/10.1111/j.1439-

0310.2007.01374.x

Bosch J, Rand AS, Ryan MJ. 2000. Signal variation and call preference for whine

frequency in the túngara frog, *Physalaemus pustulosus*. Behav Ecol Sociobiol.

49:62-66. https://doi.org/10.1007/s002650000280

Bradbury J, Vehrencamp S. 2011. Principles of animal communication. Sinauer

Associates.

Bro-Jørgensen J. 2009. Dynamics of multiple signalling systems: animal communication

in a world in flux. Trends Ecol Evol. 25:292-300.

https://doi.org/10.1016/j.tree.2009.11.003

Buchanan BW. 1993. Effects of enhanced lighting on the behaviour of nocturnal frogs.

Anim Behav. 45:893-899. https://doi.org/10.1006/anbe.1993.1109

Bucher TL, Ryan MJ, Bartholomew GW. 1982. Oxygen consumption during resting,

calling and nest building in the frog *Physalaemus pustulosus*. Physiol Zool.

55:10-22.

Burmeister SS. 2022. Ecology, cognition, and the hippocampus: a tale of two frogs. Brain

Behav Evol. 97:211-224. https://doi.org/10.1159/000522108

Caldwell M, Britt KA, Mischke LC, Collins HI. 2022. Beyond sound: bimodal acoustic

calls used in mate-choice and aggression by red-eyed treefrogs. J Exper Biol.

225:jeb244460. https://doi.org/10.1242/jeb.244460

Cheetham SA, Thom MD, Beynon RJ, Hurst JL. 2007. The effect of familiarity on mate

choice. In: Hurst JL, Beynon RJ, Roberts SC, Wyatt TD (eds). Chem Sig in Vert.

11. Springer.

Clark JA, Boersma D, Olmsted DM. 2006. Name that tune: call discrimination and

individual recognition in Magellanic penguins. Anim Behav. 72:1141-1148.

https://doi.org/10.1016/j.anbehav.2006.04.002.

Coleman SW, Patricelli GL, Borgia G. 2004. Variable female preferences drive complex

male displays. Nature. 428:742-745. https://doi.org/10.1038/nature02419.

Coss DA, Hunter KL, Taylor RC. 2020. Silence is sexy: soundscape complexity alters

mate choice in túngara frogs. Behav Ecol. 32:49-59.

https://doi.org/10.1093/beheco/araa091.

Costanzo K, Monteiro A. 2007. The use of chemical and visual cues in female mate

choice in the butterfly *Bicyclus anynana*. Proc R Soc B. 274:845-851.

https://doi.org/10.1098/rspb.2006.3729.

Cronin AD, Taylor RC, Page RA, Ryan MJ, Murphy MA, Hunter, KL. 2024. Rapid

cross-generational changes in morphology and mate choice following an extreme

climatic event. Evol Ecol. https://doi.org/10.1007/s10682-024-10324-1.

Cummings ME, Bernal XE, Reynaga R, Rand AS, Ryan MJ. 2008. Visual sensitivity to a

conspicuous male cue varies by reproductive state in *Physalaemus pustulosus*

females. J Exp Biol. 211:1203-1210. https://doi.org/10.1242/jeb.012963.

Dale S, Slagsvold T. 1996. Mate choice on multiple cues, decision rules and sampling

strategies in female pied flycatchers. Behaviour. 133:903-944.

https://doi.org/10.1163/156853996x00305.

Dapper AL, Baugh AT, Ryan MJ. 2011. The sounds of silence as an alarm cue in túngara

frogs, *Physalaemus pustulosus*. Biotropica. 43:380-385.

https://doi.org/10.1111/j.1744-7429.2010.00707.x.

Dudley R, Rand AS. 1991. Sound production and vocal sac inflation in the túngara frog,

Physalaemus pustulosus (Leptodactylidae). Copeia. 1991:460-470.

https://doi.org/10.2307/1446594.

Edomwande C, Barbosa F. 2020. The influence of predation risk on mate signaling and

mate choice in the lesser waxmoth *Achroia grisella*. Sci Reports. 10:524.

https://doi.org/10.1038/s41598-020-57481-1.

Elias-Costa AJ, Faivovich J. 2025. Evolution of vocal sacs in Anura. Bulletin Amer Mus

Nat Hist. 470:1-126. https://doi.org/10.1206/0003-0090.470.1.1.

Ellis W, Bercovitch F, FitzGibbon S, Roe P, Wimmer J, Melzer A, Wilson R. 2011.

Koala bellows and their association with the spatial dynamics of free-ranging

koalas. Behav Ecol. 22:372-377. https://doi.org/10.1093/beheco/arq216.

Faure PA & Hoy RR. 2000. The sounds of silence: cessation of singing and song pausing

are ultrasound-induced acoustic startle behaviors in the katydid *Neoconocephalus*

ensiger (Orthoptera; Tettigoniidae). J Comparative Phys A. 186:129-142.

https://doi.org/10.1007/s003590050013.

Feagles OS, Höbel G. 2022. Female gray treefrogs maintain mate choice decisions under

predation threat, but adjust movements to reduce conspicuousness during mate

approach. Behav Ecol Sociobiol. 76. https://doi.org/10.1007/s00265-021-03124-y.

Feagles OS, Höbel G. 2023. Personality affects female mate choice: frogs displaying

more consistent bold behaviors are choosier. Behav Ecol. 35: arad101.

https://doi.org/10.1093/beheco/arad101.

Foley AM, DeYoung RW, Hewitt DG, Hellickson MW, Gee KL, Wester DB, Lockwood

MA, Miller KV. 2015. Purposeful wanderings: mate search strategies of male

white-tailed deer. J Mammalogy. 96:279-286.

https://doi.org/10.1093/jmammal/gyv004.

Gerhardt HC, Huber F. 2002. Acoustic communication in insects and anurans: common

problems and diverse solutions. University of Chicago Press.

Funk WC, Donnelly MA, Lips KR. 2005. Alternative views of amphibian toe-clipping.

Nature. 433. https://doi.org/10.1038/433193c.

Gil-Guevara O, Bernal HA, Riveros AJ. 2022. Honey bees respond to multimodal stimuli

following the principle of inverse effectiveness. J Exp Biol. 225: jeb243832.

https://doi.org/10.1242/jeb.243832.

Goh MZ, Morse DH. 2010. Male mate search for female emergence sites by a parasitic

wasp. Anim Behav. 80:391-398. https://doi.org/10.1016/j.anbehav.2010.05.021.

Gomez D, Théry M, Gauthier AL, Lengagne T. 2011. Costly help of audiovisual

bimodality for female mate choice in a nocturnal anuran (*Hyla arborea*). Behav

Ecol. 22:889-898. https://doi.org/10.1093/beheco/arr039.

Greenfield MD. 2015. Signal interactions and interference in insect choruses: singing and

listening in the social environment. J Comparative Phys A. 201:143-154.

https://doi.org/10.1007/s00359-014-0938-7.

Greenfield MD, Esquer-Garrigos Y, Streiff R, Party V. 2016. Animal choruses emerge

from receiver psychology. Sci Rep. 6:34369. https://doi.org/10.1038/srep34369.

Guilford T, Dawkins MS. 1991. Receiver psychology and the evolution of animal signals.

Anim Behav. 42:1-14. https://doi.org/10.1016/S0003-3472(05)80600-1.

Halfwerk W, Jones PL, Taylor RC, Ryan MJ, Page RA. 2014. Risky ripples allow bats

and frogs to eavesdrop on a multisensory sexual display. Science. 343:413-416.

https://doi.org/10.1126/science.1244812.

Halfwerk W, Smit JAH, Loning H, Lea AM, Geipel I, Ellers J, Ryan MJ. 2017.

Environmental conditions limit attractiveness of a complex sexual signal in the

túngara frog. Nat Commun. 8:1891. https://doi.org/10.1038/s41467-017-02067-1.

Healy SD, Hurly TA. 2004. Spatial learning and memory in birds. Brain Behav Evol.

63:211-220. https://doi.org/10.1159/000076782

Hebets EA. 2003. Subadult experience influences adult mate choice in an arthropod:

Exposed female wolf spiders prefer males of a familiar phenotype. PNAS.

100:13390-13395. https://doi.org/10.1073/pnas.2333262100.

Hebets EA, Papaj DR. 2005. Complex signal function: developing a framework of

testable hypotheses. Behav Ecol Sociobiol. 57:197-214.

https://doi.org/10.1007/s00265-004-0865-7.

Higham JP, Hebets EA. 2013. An introduction to multimodal communication. Behav

783   Ecol Sociobiol. 67:1381-1388. https://doi.org/10.1007/s00265-013-1590-x.

784   Hirschmann W, Hödl W. 2006. Visual signaling in *Phrynobatrachus krefftii boulenger*,

785   1909 (Anura: Ranidae). Herpetologica. 62:18-27. https://doi.org/10.1655/04-06.1.

786   Höbel G, Kolodziej RC. 2013. Wood frogs (*Lithobates sylvaticus*) use water surface

787   waves in their reproductive behaviour. Behaviour. 150:471-483.

788   https://doi.org/10.1163/1568539X-00003062.

789   Hultsch H, Schleuss F, Todt D. 1999. Auditory-visual stimulus pairing enhances

790   perceptual learning in a songbird. Anim Behav. 58:143-149.

791   https://doi.org/10.1006/anbe.1999.1120.

792   Hwang JTG, Yang MC. 2001. An optimality theory for mid p-values in 2×2

793   contingency tables. Stat Sin. 11:807–826.

794   James LS, Baier AL, Page RA, Clements P, Hunter KL, Taylor RC, Ryan MJ. 2022.

795   Cross-modal facilitation of auditory discrimination in a frog. Biol Lett.

796   18:20220098. https://doi.org/10.1098/rsbl.2022.0098.

797   Jones CM, Braithwaite VA, Healy SD. 2003. The evolution of sex differences in spatial

798   ability. Behav Neurosci. 117:403-411. https://doi.org/10.1037/0735-

799   7044.117.3.403.

Kassambara, A. 2023. ggpubr: 'ggplot2' based publication ready plots.

Kawase H, Okata Y, Ito K. 2013. Role of huge geometric circular structures in the

reproduction of a marine pufferfish. Sci Rep. 3:2106.

https://doi.org/10.1038/srep02106.

Kozak EC, Uetz GW. 2019. Male courtship signal modality and female mate preference

in the wolf spider *Schizocosa ocreata*: results of digital multimodal playback

studies. Curr Zool. 65:705-711. https://doi.org/10.1093/cz/zoz025.

Laird KL, Clements P, Hunter KL, Taylor RC. 2016. Multimodal signaling improves

mating success in the green tree frog (*Hyla cinerea*), but may not help small

males. Behav Ecol Sociobiol. 70:1517-1525. https://doi.org/10.1007/s00265-016-

2160-9.

Leonard AS, Dornhaus A, Papaj DR. 2011. Forget-me-not: Complex floral displays,

inter-signal interactions, and pollinator cognition. Curr Zool. 57:215-224.

https://doi.org/10.1093/czoolo/57.2.215.

Leslie CE, Rosencrans RF, Walkowski W, Gordon WC, Bazan NG, Ryan MJ, Farris HE.

2020. Reproductive state modulates retinal sensitivity to light in female túngara

frogs. Front Behav Neurosci. 13. https://doi.org/10.3389/fnbeh.2019.00293.

Leslie CE, Walkowski W, Rosencrans RF, Gordon WC, Bazan NG, Ryan MJ, Farris HE.

2021. Estrogenic modulation of retinal sensitivity in reproductive female túngara

frogs. Integr Comp Biol. 61:231-239. https://doi.org/10.1093/icb/icab032.

Li H, Schrode KM, Bee MA. 2022. Vocal sacs do not function in multimodal mate

attraction under nocturnal illumination in Cope's grey treefrog. Anim Behav.

189:127-146. https://doi.org/10.1016/j.anbehav.2022.05.003.

Liu Y, Burmeister SS. 2017. Sex differences during place learning in the túngara frog.

Anim Behav. 128:61-67. https://doi.org/10.1016/j.anbehav.2017.04.002.

Lynch KS. 2017. Understanding female receiver psychology in reproductive contexts.

Integ & Comp Biol. 57:797-807. https://doi.org/10.1093/icb/icx018.

MacGillavry T, Spezie G, Fusani L. 2023. When less is more: coy display behaviours and

the temporal dynamics of animal courtship. Proc R Soc B. 290:20231684.

https://doi.org/10.1098/rspb.2023.1684.

Marsh DM, Rand AS, Ryan MJ. 2000. Effects of inter-pond distance on the breeding

ecology of túngara frogs. Oecologia. 122:505-513.

https://doi.org/10.1007/s004420050973.

McGinley RH, Starrett J, Bond JE, & Hebets EA. 2023. Light environment interacts with

visual displays in a species-specific manner in multimodal-signaling wolf spiders.

Am Nat. 201:472-490. https://doi.org/10.1086/722830.

Miller DB. 1979. The acoustic basis of mate recognition by female zebra finches

(*Taeniopygia guttata*). Anim Behav. 27:376-380. https://doi.org/10.1016/0003-

3472(79)90172-6.

Narins PM, Hödl W, Grabul DS. 2003. Bimodal signal requisite for agonistic behavior in

a dart-poison frog, *Epipedobates femoralis*. PNAS. 100:577-580.

https://doi.org/10.1073/pnas.0237165100.

Olvido AE, Wagner WE. 2004. Signal components, acoustic preference functions and

sexual selection in a cricket. Biol J Lin Soc. 83:461-472.

https://doi.org/10.1111/j.1095-8312.2004.00404.x.

Ord TJ, Stamps JA. 2008. Alert signals enhance animal communication in "noisy"

environments. PNAS. 105:18830-18835.

https://doi.org/10.1073/pnas.0807657105.

Pardasani M, Marathe SD, Purnapatre MM, Dalvi U, Abraham NM. 2021. Multimodal

learning of pheromone locations. FASEB. 35:e21836.

https://doi.org/10.1096/fj.202100167R.

Patricelli GL, & Hebets EA. 2016. New dimensions in animal communication: the case

for complexity. Curr Opinion in Behav Sci. 12:80-89.

https://doi.org/10.1016/j.cobeha.2016.09.011.

Pauli BD, Lindström K. 2021. Trade☐off between mate choice speed and decision

accuracy under mating competition in female sand gobies. J Ethol. 39:55-64.

https://doi.org/10.1007/s10164-020-00673-z.

Pauly GB, Bernal XE, Rand AS, Ryan MJ. 2006. The vocal sac increases call rate in the

túngara frog (*Physalaemus pustulosus*). Phys Biochem Zool. 79:708-719.

https://doi.org/10.1086/504613.

Podraza ME, Moss JB, Fischer EK. 2024. Evidence for individual vocal recognition in a

pair bonding poison frog, *Ranitomeya imitator*. J Exp Biol. 227:jeb246753.

https://doi.org/10.1242/jeb.246753.

Preininger D, Boeckle M, Freudmann A, Starnberger I, Sztatecsny M, Hödl W. 2013.

Multimodal signaling in the small torrent frog (*Micrixalus saxicola*) in a complex

acoustic environment. Behav Ecol Sociobiol. 67:1449-1456.

https://doi.org/10.1007/s00265-013-1489-6.

R Core Team. 2020. R: A language and environment for statistical computing. R

Foundation for Statistical Computing.

Rand AS, Ryan MJ, Wilczynski W. 1992. Signal redundancy and receiver permissiveness

in acoustic mate recognition by the túngara frog, *Physalaemus pustulosus*. Am

Zool. 32:81-90. https://doi.org/10.1093/icb/32.1.81.

Rand AS, Dudley R. 1993. Frogs in helium: the anuran vocal sac is not a cavity

resonator. Physiol Zool. 66:793-806.

https://doi.org/10.1086/physzool.66.5.30163824.

Rand AS, Bridarolli ME, Dries L, Ryan MJ. 1997. Light levels influence female choice

in túngara frogs: predation risk assessment? Copeia. 1997:447-450.

https://doi.org/10.2307/1447770.

Reichert MS, Höbel G. 2018. Phenotypic integration and the evolution of signal

repertoires: a case study of treefrog acoustic communication. Ecol Evol. 8:3410-

3429. https://doi.org/10.1002/ece3.3927.

Roberts SA, Davidson AJ, McLean L, Beynon RJ, Hurst JL. 2012. Pheromonal induction

of spatial learning in mice. Science. 338:1462-1465.

https://doi.org/10.1126/science.1225638.

Robertson JM, Greene HW. 2017. Bright colour patterns as social signals in nocturnal

frogs. Biol J Lin Soc. 121:849-857. https://doi.org/10.1093/biolinnean/blx021.

Ronald KL, Zeng R, White DJ, Fernández-Juricic E, Lucas JR. 2017. What makes a

multimodal signal attractive? A preference function approach. Behav Ecol.

28:677-687. https://doi.org/10.1093/beheco/arx015.

Rosenthal G. 2017. Mate choice: the evolution of sexual decision making from microbes

to humans. Princeton University Press.

Rosenthal G, Ryan MJ. 2022. The ascent of women: Mate choice research since Darwin.

Science. 375:eabi6308. https://doi.org/10.1126/science.abi6308.

Rowe C. 1999. Receiver psychology and the evolution of multicomponent signals. Anim

Behav. 58:921-931. https://doi.org/10.1006/anbe.1999.1242.

Rowe C, Guilford T. 1999. Novelty effects in a multimodal warning signal. Anim Behav.

57:341-346. https://doi.org/10.1006/anbe.1998.0974.

Rowe C. 2002. Sound improves visual discrimination learning in avian predators. Proc R

Soc B. 269:1353-1357. https://doi.org/10.1098/rspb.2002.2012.

Ryan MJ. 1985. The túngara frog: a study in sexual selection and communication.

University of Chicago Press.

Ryan MJ, Fox JH, Wilczynski W, Rand AS. 1990. Sexual selection for sensory

exploitation in the frog *Physalaemus pustulosus*. Nature. 343:66-67.

https://doi.org/10.1038/343066a0.

Ryan MJ, Akre K, Kirkpatrick M. 2009. Cognitive mate choice. In: Dukas R, Ratcliffe

JM (eds) Cognitive Ecology II. University of Chicago Press.

Ryan MJ, Page RA, Hunter KL, Taylor RC. 2019a. 'Crazy love': nonlinearity and

irrationality in mate choice. Anim Behav. 147:189-198.

https://doi.org/10.1016/j.anbehav.2018.04.004.

Ryan MJ, Akre KL, Baugh AT, Bernal XE, Lea AM, Leslie C, Still MB, Wylie DC,

Rand AS. 2019b. Nineteen years of consistently positive and strong female mate

choice preferences despite individual variation. Am Nat. 194:125-134.

https://doi.org/10.1086/704103.

Schlupp I, Marler C, Ryan MJ. 1994. Benefit to male sail fin mollies of mating with

heterospecific females. Science. 263:373-374.

https://doi.org/10.1126/science.8278809.

Schwartz JJ. 1991. Why stop calling? A study of unison bout singing in a Neotropical

treefrog. Anim Behav. 42:565-577. https://doi.org/10.1016/S0003-

3472(05)80240-4.

Schwartz JJ, Huth K, Hutchin T. 2004. How long do females really listen? Assessment

time for female mate choice in the grey treefrog, *Hyla versicolor*. Anim Behav.

68:533-540. https://doi.org/10.1016/j.anbehav.2003.09.016.

Siddall EC, Marples NM. 2008. Better to be bimodal: the interaction of color and odor on

learning and memory. Behav Ecol. 19:425-432.

https://doi.org/10.1093/beheco/arm155.

Speed MP. 2000. Warning signals, receiver psychology and predator memory. Anim

Behav. 60:269-278. https://doi.org/10.1006/anbe.2000.1430.

Stange N, Page RA, Ryan MJ, Taylor RC. 2016. Interactions between complex

multisensory signal components result in unexpected mate choice responses.

Anim Behav. 134:239-247. https://doi.org/10.1016/j.anbehav.2016.07.005.

Starnberger I, Preininger D, Hödl W. 2014a. From uni- to multimodality: towards an

integrative view on anuran communication. J Comp Phys A. 200:777-787.

https://doi.org/10.1007/s00359-014-0923-1.

Starnberger I, Preininger D, Hödl W. 2014b. The anuran vocal sac: a tool for multimodal

signalling. Anim Behav. 97:281-288.

https://doi.org/10.1016/j.anbehav.2014.07.027.

Tan EJ, Elgar MA. 2021. Motion: enhancing signals and concealing cues. Biol Open.

10:bio058762. https://doi.org/10.1242/bio.058762.

Taylor RC, Buchanan BW, Doherty JL. 2007. Sexual selection in the squirrel treefrog

Hyla squirella: the role of multimodal cue assessment in female choice. Anim

Behav. 74:1753-1763. https://doi.org/10.1016/j.anbehav.2007.03.010.

Taylor RC, Klein BA, Stein J, Ryan MJ. 2008. Faux frogs: multimodal signaling and the

value of robotics in animal behaviour. Anim Behav. 76:1089-1097.

https://doi.org/10.1016/j.anbehav.2008.01.031.

Taylor RC, Klein BA, Stein J, Ryan MJ. 2011. Multimodal signal variation in space and

time: how important is matching a signal with its signaler? J Exp Biol. 214:815-

820. https://doi.org/10.1242/jeb.043638.

Taylor RC, Ryan MJ. 2013. Interactions of multisensory components perceptually rescue

túngara frog mating signals. Science. 341:273-274.

https://doi.org/10.1126/science.1237113.

Thompson JT, Bissell AN, Martins EP. 2008. Inhibitory interactions between multimodal

behavioural responses may influence the evolution of complex signals. Anim

Behav. 76:113-121. https://doi.org/10.1016/j.anbehav.2007.12.015.

Uitenbroek DG. 1997. SISA binomial. http://www.quantitativeskills.com/sisa/

distributions/binomial.htm.

Varkevisser JM, Mendoza E, Simon R, Manet M, Halfwerk W, Scharff C, Riebel K.

2022. Multimodality during live tutoring is relevant for vocal learning in zebra

finches. Anim Behav. 187:263-280.

https://doi.org/10.1016/j.anbehav.2022.03.013.

Voigt CC, Behr O, Caspers B, von Helversen O, Knörnschild M, Mayer F, Nagy M.

2008. Songs, scents, and sense: Sexual selection in the greater sac-winged bat,

Saccopteryx bilineata. J Mammal. 89:1401-1410. https://doi.org/10.1644/08-

MAMM-S-060.1.

Wells KD. 2007. The ecology and behavior of amphibians. University of Chicago Press.

Wickham H. 2016. ggplot2: Elegant graphics for data analysis. Springer.

Wickham H, Averick M, Bryan J, Chang et al. 2019. Welcome to the tidyverse. J Open

Source Software. 4:1686. https://doi.org/10.21105/joss.01686.

Wilson LJ, Burrows MT, Hastie GD, Wilson B. 2014. Temporal variation and

characterization of grunt sounds produced by Atlantic cod *Gadus morhua* and

pollack *Pollachius pollachius* during the spawning season. J Fish Biol. 84:1014-

1030. https://doi.org/10.1111/jfb.12342.

Witte K, Massmann R. 2003. Female sailfin mollies, *Poecilia latipinna*, remember males

and copy of others after 1 day. Anim Behav. 65:1151-1159.

https://doi.org/10.1006/anbe.2003.2160.

Zhu B, Zhou Y, Yang Y, Deng K, Wang T, Wang J, Tang Y, Ryan MJ, Cui J. 2021.

Multisensory modalities increase working memory for mating signals in a

treefrog. J Anim Ecol. 90:1455-1465. https://doi.org/10.1111/1365-2656.13465.

Zhu B, Yang Y, Zhou Y, Deng K, Wang T, Wang J, Tang Y, Ryan MJ, Cui J. 2022.

Multisensory integration facilitates perceptual restoration of an interrupted call in

a species of frog. Behav Ecol. 33:876-883.

https://doi.org/10.1093/beheco/arac053.

FIGURE LEGENDS

Fig. 1 The internal arrangement of the sound chamber for memory phonotaxis experiments. (diagram not drawn to scale.) A female frog was placed in the center of the sound chamber in the funnel zone (black circle) and was presented with multimodal stimuli. The auditory stimulus (solid blue crescents) was broadcast by speakers while a robofrog simultaneously inflated a silicone vocal sac to serve as the visual stimulus (curved red crescents). After a presentation period, we gently lowered the blind cloths to obscure the visual stimulus from the view of the female, still held within the funnel zone.

Fig. 2 Robofrog setup. (A) Robotic, 3D-printed frog with inflated silicone vocal sac. (B) Robotic frog pneumatic controller. Compressed air drives the unit, and a solenoid valve controls the timing of inflation. During experiments, the pneumatic tube is hidden under a section of vinyl flooring that matches the floor of the test arena.

Fig. 3 Periods of stimuli for female frogs in our experiments (image not drawn to scale). All experiments first involved a presentation period of 2 or 3 min where the female was held in the funnel and speakers antiphonally broadcast attractive (whine-chuck) calls (depicted by the blue crescents). We placed a robofrog (visual stimulus, depicted by the red lines) at one speaker that had a silicone vocal sac inflating in time with the call at that speaker (multimodal speaker). Immediately after the presentation period, the blinds were lowered and the females experienced a silent period, holding period, or both. The silent period of 5 s had no stimuli (acoustic or visual) presented to the female. For the holding period, the visual stimulus was no longer presented to the female, but the acoustic stimuli were broadcast. The holding period lasted for 20 s. Immediately following one or both of these periods, the acoustic stimuli either began or continued to play and the female was released from the funnel. From here, the female was free to make a choice between speakers.

Fig. 4 (A) View of the sound chamber from above, separated into four quadrants (figure not drawn to scale). If at any time during a trial a female left the quadrant containing the speaker she chose, we considered her path to be indirect regardless of the robofrog's location. (Quadrant lines were used by a blind observer to visualize a female's path; the lines were not present during phonotaxis experiments.) Possible paths out of the funnel zone are represented by either gray arrows (indirect path) or black arrows (direct path). (B) The dorsal view of a female túngara frog. When scoring videos, the observer quantified how many stops/turns a female would make. A movement was counted as a turn if the frog oriented her body 30° or more from the long-axis of her body.

Fig. 5 Proportion of female choice across treatments. Dotted line represents a random preference (50%). Asterisks (*) above bars indicate a significant difference from the expected preference of equal probability = 0.5. Numbers inside bars represent sample sizes for each respective treatment. The dark green bars (left) denote control treatments.

Fig. 6 Female locomotor behaviors across treatments. (A) The distribution of turns made by females grouped by whether their path to choice was direct (light blue) or indirect (dark blue). (B) The distribution of stops made by females grouped by whether their path was direct (light blue) or indirect (dark blue). (C) Proportion of females making an indirect/direct choice per treatment. For treatments that did not have a silent period (i.e. Control: Robofrog Only and Blind + Hold), females were significantly more likely to take a direct path (light blue) than an indirect path (dark blue). Numbers inside the bars are the number of individuals that made a direct path or indirect path to choice. Asterisks (*) represent $P < 0.05$.

Fig. 7 Mean latency to choice across treatments using log-transformed data. The ANOVA did not report a significant difference between any of the treatments' latencies ($p = 0.964$). The dark green boxes (left) represent control treatments.

996 **FIGURES**

997 Fig. 1:

999 Fig. 2:

1000

1001   Fig. 3:

1002

1003   Fig. 4:

1004

1005

1006   Fig. 5:

1007

1008   Fig. 6:

1009

1010   Fig. 7:

1011