Title: Immune–tracheal intercellular signalling coordinates the muscle injury response in *Drosophila*

Authors: Nourhene Ammar[1], Olivier Josse[1], Aurélien Guillou[1], Jessica Perochon[2] and Hadi Boukhatmi[1]*

[1] Institut de Génétique et Développement de Rennes (IGDR), Université de Rennes, CNRS UMR6290, Rennes, France.

[2] Institute of Cancer Sciences−University of Glasgow, Wolfson Wohl Cancer Research Centre, Glasgow, UK.

* Corresponding author: hadi.boukhatmi@univ-rennes.fr

Summary

Effective tissue adaptation to damage requires precise coordination among diverse cell types. In skeletal muscle, the cellular and molecular mechanisms that orchestrate this coordination remain incompletely understood. Here, we identify adult *Drosophila* immune cells, hemocytes, as central organizers of post-injury muscle repair. Muscle damage triggers rapid recruitment of hemocytes to affected fibers. Hemocyte-derived FGF-like ligand Branchless (Bnl) activates FGF/FGFR signalling on vascular-like tracheal cells, guiding their localized expansion toward injured regions. In parallel, hemocytes deposit extracellular matrix (ECM) components, including Collagen IV, which accumulate around damaged fibers to confine Bnl distribution and facilitate tracheal recruitment. Disruption of hemocyte-mediated Bnl or ECM delivery impairs tracheal remodelling and compromises muscle recovery after injury. Together, these findings reveal an immune–muscle–vascular communication module that integrates signalling and matrix remodelling to preserve muscle function following acute damage.

Introduction

Effective tissue repair relies on precise communication and coordination across multiple cell types and tissue systems [1]. Disruption of these interactions can impair regeneration and, in severe cases, compromise organismal survival [2]. In mammals, skeletal muscle regeneration depends on a tightly orchestrated collaboration among various cell types, including macrophages, muscle stem cells (MuSCs, also known as satellite cells), and fibro/adipogenic progenitors (FAPs) [3]. Following injury, macrophages signal to MuSCs to regulate their proliferation and differentiation [4], while vascular adaptations support increased metabolic demands and structural remodelling [5]. Nonetheless, the mechanisms that coordinate cellular recruitment and tissue adaptation to muscle injury remain incompletely understood.

Drosophila frequently encounters natural wounding, requiring efficient tissue adaptation mechanisms to maintain function and ensure survival [6]. *Drosophila* has long served as a powerful model for studying myogenesis [7], and more recently, for uncovering principles of adult muscle regeneration [8-12]. While much attention has focused on the role of *Drosophila* MuSCs in regeneration, relatively little is known about how other cell types contribute to muscle adaptation to damage. In particular, the contribution of the immune blood cells-hemocytes-remains largely unexplored. Despite limited hematopoiesis in adults, hemocytes are essential for detecting and responding to bacterial infections [13]. Approximately 85% of adult hemocytes are plasmatocytes, macrophage-like cells that are distributed throughout the body [14]. Studies, including our own, have shown that they can populate the indirect flight muscles (IFMs) under normal conditions [13,15]. However, whether these immune cells engage in active intercellular signalling following muscle damage, and how they coordinate with other cell types, remains an open question.

The *Drosophila* tracheal cells -a network of air-filled tubes that functions analogously to the vertebrate vascular and respiratory systems- enwrap and penetrate muscle fibers to oxygenate them [16]. These connections are established during development through the FGF/FGFR pathway [17]. The *Drosophila* FGF-like ligand Branchless (Bnl), produced by muscle cells, directs tracheal migration by activating the Breathless (Btl)

receptor on tracheal cells. For decades, tracheal branching has been well

characterized at the morphological, cellular, and genetic levels during *Drosophila*

development [18-21]. More recently, it has been shown that the adult trachea also

undergoes dynamic remodelling to sustain adult gut regeneration [22,23], raising the

possibility that similar remodelling could contribute to muscle responses following

injury. Hemocytes closely associate with adult trachea [13] suggesting potential cross-

talk, but whether hemocytes and tracheal cells cooperate following muscle injury

remains unknown.

Here, we report that intercellular communication between muscles, hemocytes, and

tracheal cells is coordinated in responses to adult muscle injury. We show that muscle

damage triggers rapid and sustained hemocyte recruitment. Hemocyte recruitment is

required for targeted tracheal remodelling through Bnl/Btl signalling, enabling selective

recognition and support of damaged fibers. In parallel, hemocytes deposit extracellular

matrix (ECM) components, including Collagen IV, around injured fibers to spatially

confine Bnl and promote tracheal recruitment. Disrupting either tracheal recruitment or

hemocyte-mediated ECM delivery impairs muscle performance after injury. These

defects underscore the essential roles of both processes in functional recovery.

Together, these findings reveal an immune–muscle–vascular signalling module that

integrates metabolic support with ECM organization to preserve muscle function after

damage.

Results

Hemocytes are recruited to the injured adult flight muscle

Although *Drosophila* is one established model organism for studying muscle repair, it lacks molecular markers that reliably and specifically identify injured muscle fibers. In vertebrates, re-expression of embryonic Myosin Heavy Chain (MyHC-emb) in damaged fibers is a common marker of muscle injury and regeneration [24]. To assess whether Myosin Heavy Chain (Mhc) expression could similarly serve as an indicator of muscle damage in *Drosophila*, we analysed its expression following injury. Our results show that Mhc expression is drastically upregulated in injured muscles (Figure 1 A-F) and inversely correlated with the downregulation of Mef2, providing a robust molecular readout of muscle damage (Figure S1 A-B''). This upregulation persists for at least 20 days post-injury (DPI) (Figure S2 C-L). In *Drosophila*, several Mhc isoforms are generated via alternative splicing of a single *Mhc* gene [25]. To determine which isoform respond to injury, we analysed the expression of various GFP-tagged Mhc isoforms [25,26]. Only the shorter adult Mhc isoforms-K, L, and M-were upregulated in response to muscle damage (Figure S1 C-E''). To rule out the possibility that Mhc upregulation is an artifact of the injury method used, we examined its expression in *Naa30A* mutants, which exhibit developmental defects in flight muscles and spontaneous muscle detachment in eight-day-old adults, along with loss of nuclear Mef2 expression [27]. In these mutants, Mhc upregulation coincided with Mef2 downregulation (Figure S1 F-G''), further supporting Mhc as a reliable marker of muscle damage in *Drosophila*.

To study the hemocytes response to IFMs damage, we started by tracking these cells at various time points after injury using the hemocytes reporters Peroxidasin (*Pxn*) (Figure 1 A-C'') and Hemolectin (*Hml*) (Figure S1 H-I''). Co staining for these markers and Mhc revealed extensive recruitment of hemocytes to the injured muscle fibers (Figure 1 A-C'; Figure S1 H-I''). Notably, while Pxn+ hemocytes persist at the injured muscle for at least 5 DPI, Hml+ hemocytes enrichment declines over time (Figure 1 G), highlighting the heterogeneity of the adult hemocyte population, consistent with previous findings [14]. To further characterize the hemocytes recruitment dynamics, we examined the zinc finger homeodomain transcription factor Zfh1 reporter (*Enh3-Gal4 > UAS-mCD8GFP*), another hemocytes marker [15,28] (Figure 1 D-F''). Enh3+ hemocytes

exhibited recruitment kinetics similar to Pxn+ cells (Figure 1 G). Importantly, at 5 DPI

all Pxn+ cells were also Enh3+, whereas only a subset of Pxn+ cells were Hml+ (Figure

1 G' and Figure S1 H-K'''). None of the recruited Enh3+ hemocytes expressed Mef2

(Figure S1 L'-M'). To confirm that hemocytes recruitment is a response to muscle

damage, we analyzed *Naa30A* mutants. Consistently, we observed a selective

recruitment of Enh3+ hemocytes to injured muscle fibers, identified by high Mhc

expression and the loss of Mef2 nuclear expression (Figure 1 H-I'''). Altogether, these

data show that adult hemocytes are rapidly and specifically recruited to injured muscle

fibers, with distinct subpopulations displaying differential persistence over time.

Hemocytes recruitment coordinates with tracheal remodelling following muscle

injury

The selective recruitment of hemocytes to damaged fibers suggests a role in the

muscle damage response. We observed that hemocytes within the IFMs are closely

associated with tracheal cells (Figure 2 A-A'). It has previously been shown that

tracheal cells can extend new branches toward damaged regions to support adult gut

regeneration [22,23]. We therefore asked whether tracheal remodelling also occurs in

response to IFMs injury and whether it could be linked to hemocyte recruitment. Our

analysis revealed that muscle damage, whether caused by physical injury or in

Naa30A mutant animals, leads to a significant increase in tracheal coverage of the

injured fiber (Figure 2 B-D and Figure S2 A-B'). The invasion is directed toward the

injury site, with the highest branching density close to the damaged area and

progressively decreasing with distance from the injury (Figure 2 E). We further

observed that hemocytes remain spatially close to the tracheal branches, even under

injured conditions (Figure 2 C'').

To further investigate the tracheal response to injury, we focused on terminal tracheal

cells (TTCs), specialized branches of the tracheal system known for their structural

plasticity and ability to invade tissues, including muscle fibers, to deliver oxygen [16-18].

TTCs can be reliably identified by their distinctive morphology and the expression of

the transcription factor Serum Response Factor (dSRF) (Figure 2 F-F' and J-J'). We

visualized TTCs by expressing membrane-bound GFP under the control of dSRF-Gal4

(*dSRF-Gal4 > UAS-mCD8GFP*) (Figure 2 G-H'''). We observed a high density of TTCs

within injured muscle fibers, with the highest concentration near the injury site (Figure 2 H-H''' and I). Orthogonal views showed that the recruited trachea are found not only on the muscle surface but also extend branches into the injured fibers (Figure 2 G'''-H'''), consistent with the atypical muscle-tracheal organization optimized to deliver oxygen deep within the muscle, directly to sites of high metabolic demand [16,17]. Labelling TTCs nuclei with anti-dSRF confirmed the significant enrichment of dSRF+ nuclei along the injured fibers comparing to the uninjured muscles (Figure 2 K-L'' and M). The tracheal enrichment at the injured muscle results from TTCs branching and recruitment rather than proliferation, as no PH3+ cells were detected in the muscle trachea cells (Figure 2 K'''-L'''). Altogether, these data indicate that tracheal cells respond to muscle injury and target specifically the damaged fiber.

We found that tracheal enrichment around injured muscles persists throughout the repair process and is still present 20 days post-injury (Figure S2 C-K' and N), a stage at which muscle healing is expected to be complete [9,11]. A comparison of hemocyte recruitment kinetics with tracheal coverage density revealed that hemocyte recruitment precedes tracheal coverage of the injured fibers (Figure S2 M and N), raising the possibility of an interplay between the two-cell population in supporting the injured muscle fiber.

Tracheal invasion of injured muscle fibers is triggered by local hypoxia

The tracheal recruitment to injured muscle fibers (Figure 2) suggests a critical oxygen demand of the damaged muscle. If this is the case, local hypoxia within injured fibers could serve as a key cue for tracheal attraction and directional growth toward the site of injury, as seen in other physiological contexts [29]. To assess oxygen variations within the injured muscle, we first used the Synthetic HRE (Hypoxia response element) nuclear sensor (SyHREns), a hypoxia reporter that serves as a proxy for evaluating intracellular oxygen fluctuations [30]. In normal condition, the SyHREns was found to be expressed in some TTCs, identifiable by both their triangular shape and expression of dSRF (Figure 3 A-A''). Importantly, we observed a pronounced enrichment of SyHREns-positive cells surrounding injured muscle fibers, in close proximity to the recruited tracheal branches (Figure 3 B-D''). The high number of SyHREns+ cells around the injured fibers remained stable for at least 20 DPI (Figure 3 E), indicating

that local hypoxia persists in parallel with the sustained tracheal remodelling observed
over the same period.
Hypoxia-induced tracheogenesis is controlled by the transcription factor Similar (Sima)
-the *Drosophila* homolog of hypoxia-inducible factor-1α (HIF-1α)- a central regulator of
hypoxic and angiogenic responses [29]. Accordingly, silencing *sima* in TTCs severely
decreased tracheal expansion towards the injured fibers (Figure 3 F-I). Bnl is a well-
established downstream effector of hypoxic signalling, deposited locally on hypoxic
tissues to promote tracheal attraction [29]. We therefore hypothesised that if the injured
muscle undergoes hypoxia, Bnl should accumulate locally at the damaged fiber to
facilitate tracheal remodelling. Analysing the expression of an endogenously tagged
Bnl::GFP^endo allele [31] revealed marked enrichment of Bnl at injured muscle fibers
compared to adjacent intact ones (Figure 3 J-K''' and L). Altogether, these data suggest
that muscle damage induces local muscle oxygen starvation, which in turn drives
tracheal cell recruitment to the injured tissue.

Hemocytes deposit FGF-like ligand Bnl at injured muscles to mediate tracheal
recruitment

During both embryonic and post-embryonic stages, tracheal guidance is regulated in
a dose-dependent manner by local Bnl concentrations [29]. In adult flight muscle
development, muscle-tracheal connections are established through the action of Bnl,
which is secreted by muscle fibers and acts in a paracrine manner by binding to its
receptor, FGFR/Breathless (Btl), expressed on tracheal cells [17]. The observed
enrichment of Bnl in injured muscles (Figure 3 J-L) raises two key questions: Is Bnl/Btl
signaling functionally involved in directing tracheal recruitment to damaged fibers? If
so, what is the source of Bnl enrichment on the injured fibers? To begin addressing
these questions, we conditionally overexpressed Bnl in a restricted subset of adult
muscle fibers using the flip-out system combined with a regionally expressed driver
(Figure S3 A-C and Materials and Methods section). This targeted overexpression was
sufficient to recruit tracheal cells to the ectopic Bnl-expressing regions (Figure 4 A-C'),
and the extent of this recruitment depended on the duration of Bnl expression:
prolonged expression (5 days) triggered more extensive tracheal remodelling than
short-term expression (24 hours). These findings showed that the adult tracheal
system remains responsive to elevated Bnl levels in the IFMs, supporting the

hypothesis that Bnl/Btl signalling directs tracheal pathfinding in response to muscle injury.

We next sought to identify the cellular source responsible for Bnl accumulation in injured muscle fibers. Under wild-type conditions, Bnl::GFP^{endo} signal was detected in hemocytes, tracheal cells, and indirect flight muscles (IFMs) (Figure S3 D-D'). However, since Bnl is a diffusible protein, these observations alone cannot distinguish between cells that produce Bnl and those that merely take it up. To address this, we used the *Bnl-lacZ* transcriptional reporter, which revealed active *bnl* expression in hemocytes, adult IFMs, and terminal tracheal cells (TTCs), confirming that these cell types are bona fide sources of Bnl under normal conditions (Figure S3 E-F'). Given that canonical Bnl/Btl signalling operates in a paracrine manner, injured muscles and hemocytes emerge as the most plausible sources of injury-induced Bnl accumulation. To distinguish between these two possibilities, we examined *Bnl-lacZ* expression following injury. Surprisingly, we found that injured fibers lacked *Bnl-lacZ* activity, whereas neighboring uninjured fibers retained its expression (Figure 4 D-E"). These findings indicate that the injured muscle fibers themselves do not contribute to the localized Bnl accumulation after injury and suggest that the source of Bnl is extrinsic to the damaged tissue.

By contrast, we observed a marked enrichment of Bnl::GFP^{endo} signal surrounding Pxn+ hemocytes recruited to the damaged fibers, suggesting that hemocytes are responsible for depositing Bnl along injured muscles (Figure 4 F-G"). To assess the functional contribution of hemocyte-derived Bnl to tracheal invasion, we selectively silenced *bnl* in hemocytes using either the *Enh3-Gal4* or *HmlΔ-Gal4* drivers. While hemocyte recruitment to injured muscles was unaffected, *bnl* knockdown significantly reduced tracheal invasion into damaged fibers compared with controls (Figure 4 H-J" and K; Figure S3 G-I" and J). This signalling logic between the two cell types aligns with their close anatomical association under homeostatic conditions (Figure 2 A-A' and [13]).

Given the critical role of the tracheal system in supporting tissue function [21], we next asked whether hemocyte-mediated tracheal recruitment via Bnl is required to maintain muscle flight performance after injury. To test this, we performed flight assays on uninjured controls, injured flies, and injured flies in which *bnl* was specifically depleted

from hemocytes. Injured flies exhibited reduced flight capability compared with uninjured controls (Figure 4 L-L'). Importantly, flies lacking hemocyte-derived Bnl showed an even greater impairment in flight, underscoring the functional importance of this signalling in promoting tracheal remodelling and preserving muscle performance following injury (Figure 4 M).

The receptor FGFR/Btl is essential for injury-induced tracheal guided outgrowth

The finding that Bnl orchestrates tracheal morphogenesis in response to muscle damage suggested in turn that it activates its receptor Btl, on tracheal cells, thereby driving guided outgrowth and branching. To investigate this, we examined Btl expression in adult flight muscles using an endogenously tagged *Btl::cherry^endo* allele, which faithfully reports Btl expression [31]. Btl::cherry^endo was detected along tracheal branches under homeostatic conditions and within tracheal structures invading the injured muscle (Figure 5 A-B''). To complement this approach, we expressed a membrane-tethered GFP (mCD8-GFP) under the control of a *Btl-Gal4* driver (Figure 5 C-E')*.* While Btl::cherry^endo reflects consistent endogenous expression, *Btl-Gal4* drives a more heterogeneous pattern across tracheal cells (Figure 5 C-C'). Nevertheless, we observed a striking accumulation of tracheal branches labelled by *Btl-Gal4* at injured muscles, reinforcing that Btl-expressing tracheal cells are actively recruited to damaged fibers (Figure 5 D-E'). We then tested whether Btl function is required for proper tracheal pathfinding. To this end, *btl* was specifically silenced in tracheal cells using either *dSRF-Gal4* or *Btl-Gal4* drivers (Figure 5 F-I''). In both cases, tracheal invasion into injured fibers was significantly impaired compared to the control (Figure 5 J). Notably, the impairment was more pronounced with *dSRF-Gal4*, consistent with its more uniform expression in terminal tracheal cells (Figure 2 F-F') compared with *Btl-Gal4* (Figure 5 C-C'). Together, these findings show that the Bnl/Btl signalling axis plays a pivotal role in guiding tracheal remodelling after muscle damage. While Btl expression is restricted to tracheal cells, localized Bnl secretion from hemocytes at the damaged fiber refines tracheal pathfinding, ensuring precise and targeted tracheal invasion into injured muscle (Figure 5 K).

Hemocyte-derived ECM retaines Bnl ligand at injured muscles

Although Bnl is a diffusible molecule, we observed that it is specifically retained at the injured fiber (Figure 3 K'-K''), raising the question of what restricts its spatial distribution. Previous studies have shown that Collagen IV, a principal structural component of the muscle ECM, is critical for retaining diffusible signals within defined tissue regions [32]. We therefore hypothesized that ECM components may similarly act to confine Bnl to the damaged fiber.

Hemocytes are known to be major producers of ECM components during both *Drosophila* development and post-developmental stages [33,34], suggesting they could supply Collagen IV in response to adult muscle injury. To test this, we analysed Collagen IV distribution in adult flight muscles using the Vkg::GFP line (Figure 6 A-B''). Under homeostatic conditions, we observed a basal level of Vkg::GFP surrounding muscle fibers and along tracheal branches (Figure 6 A'), and we found that hemocytes express Collagen IV (Figure 6 A''). Five days after injury, Vkg::GFP was markedly enriched in damaged fibers compared with intact muscles and uninjured controls (Figure 6 A-B and A''-B''). To further assess the relationship between hemocytes and ECM remodeling, we examined Collagen IV distribution across multiple time points post-injury (1-20 DPI) and compared this to hemocyte recruitment dynamics (Figure 6 C-D'' and Figure S4 A-F). Collagen IV accumulation closely correlated with the presence and distribution of hemocytes. Notably, regions where hemocytes were evenly distributed along the injured fiber displayed stronger and more uniform Vkg::GFP signal.

To test whether hemocyte-supplied Collagen IV is necessary for Bnl retention at injured fibers, we knocked down Collagen IV specifically in hemocytes and assessed Bnl::GFP[endo] localization (Figure 6 E-F''). In control flies, Bnl signal was strongly enriched along the injured fiber; however, this enrichment was markedly reduced following Collagen IV depletion (Figure 6 E-F'' and G). These data indicate that Collagen IV is required to retain Bnl at the injury site. Together, these findings show that hemocyte-derived Collagen IV acts as a structural scaffold that immobilizes Bnl at damaged fibers (Figure 6 H). This localized Bnl/Btl signalling likely guides tracheal remodelling and contributes to the maintenance of flight ability.

Hemocyte-derived SPARC supports Collagen IV deposition, tracheal remodleling and muscle recovery

Incorporation of Collagen IV into the extracellular matrix (ECM) requires accessory molecules for proper deposition and stabilization. We therefore investigated whether hemocytes also supply such regulatory components. One key factor is SPARC, a collagen-binding glycoprotein essential for Collagen IV diffusion, stabilization, and incorporation into basement membranes [35,36]. We first examined SPARC localization in hemocytes associated with adult IFMs. Hemocytes, identified using *Enh3-Gal4* driver, were positive for SPARC (Figure 6 I-I'). Importantly, following muscle injury, SPARC levels were elevated along the injured fiber compared with controls, mirroring the distribution of Collagen IV (Figure 6 I-J'). These observations suggest that SPARC is locally produced by hemocytes in adult muscle. To further confirm the cellular origin of SPARC enrichment in injured muscles, we employed the SPARC-Gal4 driver, a Gal4 trap line inserted into the endogenous SPARC locus, to drive membrane-targeted RFP (myr-RFP) expression in flies also carrying a BM40-SPARC-GFP fosmid reporter (Figure 6 K-L'). This dual-reporter approach allowed simultaneous visualization of SPARC (via BM40-SPARC-GFP) and identification of SPARC-producing cells (via SPARC-Gal4 > myr-RFP). In uninjured muscle, BM40-SPARC-GFP and SPARC-Gal4-driven myr-RFP were co-expressed in hemocytes. Following injury, BM40-SPARC-GFP was broadly enriched along the damaged fiber, resembling endogenous SPARC protein, whereas SPARC-Gal4 activity remained restricted to hemocytes and was not detected in muscle nuclei (Figure 6 L-L'). Together, these data demonstrate that recruited hemocytes are the primary source of SPARC at injured muscle fibers. We then tested whether hemocyte-derived SPARC is required for Collagen IV enrichment and tracheal remodelling. Depletion of SPARC specifically in hemocytes was sufficient to prevent both Collagen IV accumulation and tracheal remodelling at the injured fiber (Figure 6 M-N'' and O-P). Finally, to evaluate the functional importance of SPARC in the muscle repair process, we performed flight assays and found that injured flies lacking hemocyte-derived SPARC exhibited significantly reduced flight ability compared with controls (Figure 6 Q). Altogether, these results show that hemocyte-derived SPARC is essential for ECM assembly, tracheal remodelling, and functional recovery following muscle injury. They also highlight that hemocytes act as

345 multifunctional signalling hubs, simultaneously delivering trophic, structural, and
346 guidance cues required to coordinate efficient muscle repair.

Discussion

Our study provides a compelling example of how the muscle damage response relies
on the precise coordination between multiple cell types. While *Drosophila* hemocytes
are traditionally recognized for their immunoprotective roles against pathogens and
infections [13,33,37], our findings reveal that they also play essential roles for repairing
injured muscles. We show that hemocytes both signal to the tracheal system to finely
regulate its outgrowth toward the injured muscle and directly supply key extracellular
matrix (ECM) components to the damaged fiber, two processes that support muscle
structural maintenance and recovery post injury (Figure 7).

A functional cooperation between immune cells and the respiratory system

Insect respiration has traditionally been viewed as entirely dependent on the tracheal
system, that operates independently of blood cells. However, this view has been
challenged by recent finding showing that a subtype of larval hemocytes, known as
crystal cells, can take up oxygen from tracheal cells and contribute to tissue
oxygenation [38]. Here, we reveal that in adult Drosophila, hemocytes are positioned in
close proximity to tracheal cells within muscle tissue and indirectly contribute to oxygen
delivery by guiding tracheal branches toward sites of muscle injury (Figure 7).
Following damage, hemocytes are rapidly recruited to and spread over the injured
muscle surface. They secrete Bnl that promote local tracheal outgrowth and precise
branch targeting. This finding raises an intriguing question: Why does the injured
muscle, the structure directly damaged, not itself produce Bnl? Put differently, why
delegate to extrinsic immune cells the ability to attract tracheal branches when the
demand for oxygen and repair arises intrinsically within the muscle? One possible
explanation lies in the need to prevent inappropriate or excessive activation of tracheal
growth programs. Bnl is a potent chemoattractant, and its widespread expression
within muscle fibers could trigger aberrant tracheogenesis, leading to the recruitment
of non-specific tracheal branches and disruption of the normal tracheal network
architecture. Hemocytes, by contrast, are far less numerous than muscle fiber nuclei
and, once localized along the damaged area, could secrete limited amounts of Bnl in

a spatially restricted manner. Such targeted release would ensure that tracheal

outgrowth occurs only at sites where oxygen delivery and tissue repair are most

needed.

Beyond their indirect role in oxygenation, which is mediated through the guidance of

the tracheal network, it remains to be determined whether adult hemocytes might also

directly contribute to oxygen transport, as has been described with crystal cells during

larval stages [38]. Interestingly, during development, some plasmatocytes have been

shown to transdifferentiate into crystal cells in the lymph gland, the primary

hematopoietic organ [39]. Whether a similar transdifferentiation occurs in the adult

following muscle injury remains an open question. The nature of the signals that govern

hemocyte recruitment under injury conditions remains an important avenue for future

investigation.

Do Drosophila repair or regenerate their flight muscles?

The IFMs generate the high-frequency wing beats that enable flight, a behavior

essential for predator avoidance, foraging, and mating. Given their vital function, IFMs

must be maintained in optimal condition to ensure sustained performance throughout

the animal's lifespan. This need for resilience is also underscored by the observation

that over 30% of wild-caught *Drosophila* exhibit physical injuries, frequently located in

the thorax and abdomen [6]. Although a rare population of satellite-like cells has been

identified in adult flies, the regenerative capacity of adult *Drosophila* muscles remains

debated [40]. A major limitation has been the lack of appropriate molecular markers to

monitor both muscle regeneration and muscle stem cell (MuSC) activity. In our study,

we identified a specific isoform of the Myosin Heavy Chain (Mhc) that is consistently

upregulated in damaged fibers. However, unlike in vertebrates, where MyHC-emb

levels return to baseline following regeneration, Mhc expression in *Drosophila* remains

elevated even 20 days post-injury (Figure S2). This makes Mhc a sensitive marker for

identifying injury, but less useful for assessing complete muscle recovery. Adult MuSCs

have been identified via the transcription factor Zfh1/ZEB and its reporter Enh3 [11,15],

the only presently known marker for these cells. Zfh1 is required for MuSCs

maintenance, an equivalent role of ZEB was subsequently shown in mice [41].

Interestingly, both Zfh1 and ZEB are expressed in the muscle associated immune cells

[15,41]. In our study, we found that following muscle injury, 100% of the cells recruited to

damaged fibers are both Enh3$^+$ and Pxn$^+$ (a plasmatocytes marker), while being

negative for the muscle-specific marker Mef2 (Figure S1 L-M'). This strongly suggests

that hemocytes are the predominant, if not exclusive cell population migrating to injured

muscle. These findings suggest that adult *Drosophila* flight muscles possess limited

intrinsic regenerative capacity and may instead rely on compensatory mechanisms

such as increased oxygenation through tracheal remodelling and structural

reinforcement via extracellular matrix (ECM) deposition-to cope with muscle damage.

This is consistent with the broader idea that short-lived species prioritize immediate

survival over long-term regenerative investment [42]. In such organisms, evolution favors

fast and efficient repair mechanisms that support rapid recovery and fitness over

complex, energy-intensive regeneration programs.

Functional parallels between Drosophila hemocytes and mammalian FAPs

In mammals, extracellular matrix (ECM) remodeling is essential for maintaining muscle

structure and enabling proper tissue repair. Although muscle stem cells (MuSCs) can

produce some ECM components [43], the primary contributors to ECM production and

maintenance during regeneration are fibro-adipogenic progenitors (FAPs) [44].

Drosophila lack dedicated FAPs, raising the question of how flies achieve ECM

remodelling and maintain muscle integrity. During development, hemocytes are the

primary source of ECM molecule production [33,45,46]. However, functional evidence for

their involvement in ECM remodeling in adult tissues-particularly muscle-has remained

limited. Here, we showed that adult hemocytes produce at least two key ECM

components -Collagen IV and SPARC. This production is particularly enriched around

damaged muscle fibers where hemocytes are abundant. We further showed that

hemocyte-specific ablation of ECM led to defect in muscle post injury recovery. These

findings suggest that *Drosophila* hemocytes share functional similarities with

mammalian FAPs and may compensate for their absence. This highlights an

evolutionarily flexible strategy for achieving muscle repair in organisms with fewer

specialized cell types.

436
437
438
439

Materials and Methods

Drosophila melanogater strains and genetics

All *Drosophila melanogaster* stocks were grown on standard medium at 25°C. The following stains were used: w^{118} as wild type (wt), *Pxn-RFP*[47], *HmlΔ-Gal4* (BL#30140), *Enh3-Gal4* [15], *Mhc-Iso-K, L, M (fTRG500)* and *Mhc-Iso-A, F, G (fTRG519)* [26], *weeP26* [25], *Bnl::GFP^{endo}* and *Btl::cherry^{endo}* [31], *Bnl-LacZ* (BL#11704), , *dSRF-GAL4* (BL#25753), *Btl-Gal4* (BL#78328), *UAS-Btl^{RNAi}* (BL#435544), *UAS-Bnl^{RNAi}* (BL#34572), *UAS-Bnl::GFP* [48], *Vkg::GFP* (BL#98343), *BM40-SPARC-GFP* (Sarov et al., 2016), *SME-Gal4* (VT 0576871), *UAS-H2B-RFP* [49], *UAS-mCD8GFP* (BL#5137), *UAS-myr-mRFP* (BL#7118), SyHREns [30], *UAS-Sima^{RNAi}* (BL#26207), *SPARC-Gal4* (BL#77473), *UAS-SPARC^{RNAi}* (BL#40885), *UAS-Vkg^{RNAi}* (BL#50895). *Naa30A* deletion males were obtained from crosses between *Naa30A^{Δ74}/FM0* females and either UAS-mCD8GFP; Enh3-Gal4 (Figure 1) or Mhc-Iso-K, L, M (Figure S1) males. RNAi experiments were carried out at 29°C. To achieve temporal control of gene expression in adult tissues, the tubGal80^{ts} was employed, allowing restriction of Gal4 activity to a specific time window. Fly crosses were initially maintained at 18°C, and after adult eclosion, the progenies were transferred to 29°C, where they remained until dissection. An exception was made for the Bnl knockdown experiments shown in Figure 4 J-J", which were performed without the tubGal80^{ts} transgene.

Generation of SME-FRT-stop-FRT-Gal4 fly line and flip-out induction procedure.

SME (Shavenbaby Muscle Enhancer) is one of several enhancers of the *shavenbaby* gene (VT057071-Gal4). It is active in larval muscle progenitors (AMPs) (Figure S3 A-A') and remains active in a subpopulation of localized muscle nuclei in the adult indirect flight muscles (IFMs) (Figure S3 B-B'). To generate the SME-FRT-stop-FRT-Gal4 construct, the cis-regulatory element from the original *shavenbaby* enhancer construct was used to replace the *pAct* promoter in the pAct-FRT-stop-FRT3-FRT-FRT3-Gal4 vector (Addgene plasmid #52889). The resulting construct was integrated into the attP site at cytological location 51D on chromosome II via embryo injection into vas-int; attP-51D flies. To induce localized overexpression of Bnl::GFP in adult IFM nuclei, hsflp; FRT-SME-stop-FRT-Gal4/CyO; tubGal80^{ts} females were crossed to UAS-Bnl::GFP

males. Progeny were raised at 18°C and subjected to a 2-hour heat shock at 37°C in young adulthood (Figure 4 A-C').

Immunohistochemistry and confocal microscopy

Preparation and immunofluorescence of wing discs and adult muscles were performed as previously described in [50]. The following primary antibody were used: Chicken anti-βGal (1:100, Abcam, ab9361), Chicken anti-GFP (1:300, Abcam, ab13970), Mouse anti-Mhc (1:200, DSHB, 3E8-3D3-s) , Mouse anti-dSRF (1:20, a gift from Mark A. Krasnow, Stanford university California , USA), Rabbit anti-Mef2 (1:500 , DSHB, Mef2-p), Rabbit anti-PH3 (1:100, Cell Signaling Technology, #9713), Rabbit anti-SPARC (1:500, a gift from Maurice Ringuette, University of Toronto, Canada), Rat anti-RFP (1:800, Chromotek, 5F8-20), Alexa-conjugated Phalloidin (1:200, Thermo fisher scientific life technologies).

Induction of flight muscle injury

Flight muscle injury was induced following previously described protocol [51]. Adult flies (2-3 days post-eclosion) were anesthetized and positioned laterally under a stereomicroscope. A single, localized stab wound was inflicted manually using a fine stainless-steel needle (Minutien Pins, 0.1 mm diameter; #26002–10) to minimize tissue damage and avoid extensive injury. Age-matched, uninjured flies served as controls.

Flight assay

Flight ability was assessed using a 1-m-long cylindrical flight column internally lined with paper coated with a thin layer of glue. The adhesive sheet was divided evenly into three longitudinal sections corresponding to the Top**,** Middle, and Bottom regions of the tube (Figure 4L). For each experimental condition, an equal number of flies (approximately 20–30) was released into the tube. After all flies had landed, the paper was removed, flattened, and imaged for analysis.

Image quantification was performed semi-automatically using a custom R script, Fly-Spy (available at https://github.com/Olivier-Josse/Fly-spy). Images were first converted to grayscale and digitally divided into the three predefined regions. High-

intensity spots corresponding to individual flies were identified by applying a manually adjusted threshold to ensure accurate segmentation. The number of flies detected in each region was quantified and expressed as a percentage of the total recovered flies. The distribution of flies across the three regions—reflecting high, intermediate, or low/absent flight ability—was compared between experimental conditions using a chi-square (χ^2) test. Results are presented as stacked bar histograms

Microscopy and data analysis

Samples were imaged using a Leica SP8 microscope (BIOSIT, University of Rennes) at 20×, 40×, or 63× magnification and a resolution of 1024×1024 pixels. Images were processed with ImageJ and assembled using Adobe Illustrator. To quantify fluorescence intensity, all samples were processed and analyzed in parallel under identical conditions. Confocal images were acquired using the same laser settings across all samples. Sum slice projections were generated from each confocal stack. Regions of interest (ROIs) were manually defined based on protein expression, and mean fluorescence intensities were measured within each ROI. These values were then normalized to background levels measured in comparable areas of the same sample.

Tracheal branches were visualized either by their intrinsic autofluorescence using a 405 nm UV laser [16] or by the expression of membrane-bound GFP driven by a tracheal-specific Gal4 driver (UAS-mCD8GFP). Tracheal coverage was quantified as pixel area per muscle fiber from maximum intensity Z-projections acquired at 20× magnification. Quantification was performed using the "Skeletonize" plugin in ImageJ, based on the method described by Perochon et al. (2021) [23], with minor modifications. Briefly, maximum intensity projections were generated from each confocal stack, tracheal branches were segmented by thresholding, and the resulting images were skeletonized. Skeleton pixel intensity was then measured specifically within injured muscle fibers and their corresponding contralateral control fibers. For Figure 2E, square regions of interest (ROIs) were placed at the injury site, as well as in proximal and distal regions, to assess local tracheal enrichment. Hemocyte and dSRF-positive cell numbers were quantified manually using ImageJ.

Graphs and statistical analyses were performed using GraphPad Prism. Pair comparisons were statistically tested with a two-sided Student's t-test. NS, not significant ($p>0.05$); $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ $****p< 0.0001$.

Acknowledgments

We would like to thank Jennifer Zanet, Cedric Polesello, Sougata Roy, Ruth Lehmann, Maurice Ringuette, Jean-Paul Vincent, and Mark A. Krasnow, as well as the Bloomington Drosophila Stock Center, the Vienna Drosophila Resource Center (VDRC), and the Developmental Studies Hybridoma Bank for providing *Drosophila* strains and antibodies. We are grateful to Lucas Waltzer, Laetitia Bataille, Alain Vincent, and Michèle Crozatier for their critical reading of the manuscript, and to other members of H.B's laboratory for valuable discussions. We also thank Emmanuel Gallaud for helpful input during the early phases of the project. We acknowledge Xavier Pinson from the Microscopy Rennes Imaging Center (MRic, BIOSIT, Biogenouest) for technical assistance, and Cyrille Surbled for fly stock maintenance and technical help. Work in the Boukhatmi lab was supported by an AFM-Téléthon Trampoline Grant (#23108), ATIP-Avenir (CNRS), and ANR GenepiMuSC (ANR-23-CE13-0006). N.A. was supported by an AFM-Téléthon PhD fellowship (#23846) and an FRM fellowship (FDT202404018608). Work in the Cordero lab was supported by a Sir Henry Dale Fellowship (Wellcome Trust and Royal Society, 104103/Z/14/Z), a Wellcome ISSF Returner Scheme (204820/Z/16/Z), and a Tenovus Fellowship (Tenovus/S22-28) to J.P.

Competing interest

The authors declare that they have no competing interests.

Reference

1. Sun, F., and Poss, K.D. (2023). Inter-organ communication during tissue regeneration. Development *150*. 10.1242/dev.202166.

2. Li, J., Zhang, S., and Amaya, E. (2016). The cellular and molecular mechanisms of tissue repair and regeneration as revealed by studies in Xenopus. Regeneration (Oxf) *3*, 198-208. 10.1002/reg2.69.

3. Relaix, F., Bencze, M., Borok, M.J., Der Vartanian, A., Gattazzo, F., Mademtzoglou, D., Perez-Diaz, S., Prola, A., Reyes-Fernandez, P.C., Rotini, A., and Taglietti, t. (2021). Perspectives on skeletal muscle stem cells. Nature Communications *12*, 692. 10.1038/s41467-020-20760-6.

4. Dort, J., Fabre, P., Molina, T., and Dumont, N.A. (2019). Macrophages Are Key Regulators of Stem Cells during Skeletal Muscle Regeneration and Diseases. Stem Cells Int *2019*, 4761427. 10.1155/2019/4761427.

5. Eming, S.A., Martin, P., and Tomic-Canic, M. (2014). Wound repair and regeneration: mechanisms, signaling, and translation. Sci Transl Med *6*, 265sr266. 10.1126/scitranslmed.3009337.

6. Subasi, B.S., Grabe, V., Kaltenpoth, M., Rolff, J., and Armitage, S.A.O. (2024). How frequently are insects wounded in the wild? A case study using Drosophila melanogaster. R Soc Open Sci *11*, 240256. 10.1098/rsos.240256.

7. Laurichesse, Q., and Soler, C. (2020). Muscle development : a view from adult myogenesis in Drosophila. Semin Cell Dev Biol *104*, 39-50. 10.1016/j.semcdb.2020.02.009.

8. Gunage, R.D., Dhanyasi, N., Reichert, H., and VijayRaghavan, K. (2017). Drosophila adult muscle development and regeneration. Semin Cell Dev Biol *72*, 56-66. 10.1016/j.semcdb.2017.11.017.

9. Catalani, E., Zecchini, S., Giovarelli, M., Cherubini, A., Del Quondam, S., Brunetti, K., Silvestri, F., Roux-Biejat, P., Napoli, A., Casati, S.R., et al. (2022). RACK1 is evolutionary conserved in satellite stem cell activation and adult skeletal muscle regeneration. Cell Death Discovery *8*, 459. 10.1038/s41420-022-01250-8.

10. Goh, K.Y., Lee, W.X., Choy, S.M., Priyadarshini, G.K., Chua, K., Tan, Q.H., Low, S.Y., Chin, H.S., Wong, C.S., Huang, S.Y., et al. (2024). FOXO-regulated DEAF1 controls muscle regeneration through autophagy. Autophagy *20*, 2632-2654. 10.1080/15548627.2024.2374693.

11. Chaturvedi, D., Reichert, H., Gunage, R.D., and VijayRaghavan, K. (2017). Identification and functional characterization of muscle satellite cells in Drosophila. Elife *6*. 10.7554/eLife.30107.

12. Boukhatmi, H. (2021). Drosophila, an Integrative Model to Study the Features of Muscle Stem Cells in Development and Regeneration. Cells *10*. 10.3390/cells10082112.

13. Sanchez Bosch, P., Makhijani, K., Herboso, L., Gold, K.S., Baginsky, R., Woodcock, K.J., Alexander, B., Kukar, K., Corcoran, S., Jacobs, T., et al. (2019). Adult Drosophila Lack Hematopoiesis but Rely on a Blood Cell Reservoir at the Respiratory Epithelia to Relay Infection Signals to Surrounding Tissues. Dev Cell *51*, 787-803 e785. 10.1016/j.devcel.2019.10.017.

14. Boulet, M., Renaud, Y., Lapraz, F., Benmimoun, B., Vandel, L., and Waltzer, L. (2021). Characterization of the Drosophila Adult Hematopoietic System Reveals a Rare Cell

614   Population With Differentiation and Proliferation Potential. Front Cell Dev Biol *9*,
615   739357. 10.3389/fcell.2021.739357.

616 15.   Boukhatmi, H., and Bray, S. (2018). A population of adult satellite-like cells in
617   Drosophila is maintained through a switch in RNA-isoforms. Elife *7*.
618   10.7554/eLife.35954.

619 16.   Sauerwald, J., Backer, W., Matzat, T., Schnorrer, F., and Luschnig, S. (2019). Matrix
620   metalloproteinase 1 modulates invasive behavior of tracheal branches during entry
621   into Drosophila flight muscles. Elife *8*. 10.7554/eLife.48857.

622 17.   Peterson, S.J., and Krasnow, M.A. (2015). Subcellular trafficking of FGF controls
623   tracheal invasion of Drosophila flight muscle. Cell *160*, 313-323.
624   10.1016/j.cell.2014.11.043.

625 18.   Ghabrial, A., Luschnig, S., Metzstein, M.M., and Krasnow, M.A. (2003). Branching
626   morphogenesis of the Drosophila tracheal system. Annu Rev Cell Dev Biol *19*, 623-
627   647. 10.1146/annurev.cellbio.19.031403.160043.

628 19.   Rao, P.R., Lin, L., Huang, H., Guha, A., Roy, S., and Kornberg, T.B. (2015).
629   Developmental compartments in the larval trachea of Drosophila. Elife *4*.
630   10.7554/eLife.08666.

631 20.   Ebner, A., Kiefer, F.N., Ribeiro, C., Petit, V., Nussbaumer, U., and Affolter, M. (2002).
632   Tracheal development in Drosophila melanogaster as a model system for studying the
633   development of a branched organ. Gene *287*, 55-66. 10.1016/s0378-1119(01)00895-
634   2.

635 21.   Hayashi, S., and Kondo, T. (2018). Development and Function of the Drosophila
636   Tracheal System. Genetics *209*, 367-380. 10.1534/genetics.117.300167.

637 22.   Tamamouna, V., Rahman, M.M., Petersson, M., Charalambous, I., Kux, K., Mainor, H.,
638   Bolender, V., Isbilir, B., Edgar, B.A., and Pitsouli, C. (2021). Remodelling of oxygen-
639   transporting tracheoles drives intestinal regeneration and tumorigenesis in
640   Drosophila. Nat Cell Biol *23*, 497-510. 10.1038/s41556-021-00674-1.

641 23.   Perochon, J., Yu, Y., Aughey, G.N., Medina, A.B., Southall, T.D., and Cordero, J.B.
642   (2021). Dynamic adult tracheal plasticity drives stem cell adaptation to changes in
643   intestinal homeostasis in Drosophila. Nat Cell Biol *23*, 485-496. 10.1038/s41556-021-
644   00676-z.

645 24.   Murphy, M.M., Lawson, J.A., Mathew, S.J., Hutcheson, D.A., and Kardon, G. (2011).
646   Satellite cells, connective tissue fibroblasts and their interactions are crucial for
647   muscle regeneration. Development *138*, 3625-3637. 10.1242/dev.064162.

648 25.   Orfanos, Z., and Sparrow, J.C. (2013). Myosin isoform switching during assembly of
649   the Drosophila flight muscle thick filament lattice. J Cell Sci *126*, 139-148.
650   10.1242/jcs.110361.

651 26.   Sarov, M., Barz, C., Jambor, H., Hein, M.Y., Schmied, C., Suchold, D., Stender, B.,
652   Janosch, S., K, J.V., Krishnan, R.T., et al. (2016). A genome-wide resource for the
653   analysis of protein localisation in Drosophila. Elife *5*, e12068. 10.7554/eLife.12068.

654 27.   Varland, S., Silva, R.D., Kjosas, I., Faustino, A., Bogaert, A., Billmann, M., Boukhatmi,
655   H., Kellen, B., Costanzo, M., Drazic, A., et al. (2023). N-terminal acetylation shields
656   proteins from degradation and promotes age-dependent motility and longevity. Nat
657   Commun *14*, 6774. 10.1038/s41467-023-42342-y.

658 28.   Wu, W.H., Kuo, T.H., Kao, C.W., Girardot, C., Hung, S.J., Liu, T., Furlong, E.E.M., and Liu,
659   Y.H. (2019). Expanding the mesodermal transcriptional network by genome-wide

660    identification of Zinc finger homeodomain 1 (Zfh1) targets. FEBS Lett *593*, 1698-1710.
661    10.1002/1873-3468.13443.

662 29. Jarecki, J., Johnson, E., and Krasnow, M.A. (1999). Oxygen regulation of airway
663    branching in Drosophila is mediated by branchless FGF. Cell *99*, 211-220.
664    10.1016/s0092-8674(00)81652-9.

665 30. Zhao, Y., Alexandre, C., Kelly, G., Perez-Mockus, G., and Vincent1, J.-P. (2024). Growth-
666    induced physiological hypoxia correlates with growth deceleration during normal
667    development. bioRxiv. https://doi.org/10.1101/2024.06.04.597345.

668 31. Du, L., Sohr, A., Yan, G., and Roy, S. (2018). Feedback regulation of cytoneme-
669    mediated transport shapes a tissue-specific FGF morphogen gradient. Elife *7*.
670    10.7554/eLife.38137.

671 32. Wang, X., Harris, R.E., Bayston, L.J., and Ashe, H.L. (2008). Type IV collagens regulate
672    BMP signalling in Drosophila. Nature *455*, 72-77. 10.1038/nature07214.

673 33. Gold, K.S., and Bruckner, K. (2015). Macrophages and cellular immunity in Drosophila
674    melanogaster. Semin Immunol *27*, 357-368. 10.1016/j.smim.2016.03.010.

675 34. Bunt, S., Hooley, C., Hu, N., Scahill, C., Weavers, H., and Skaer, H. (2010). Hemocyte-
676    secreted type IV collagen enhances BMP signaling to guide renal tubule
677    morphogenesis in Drosophila. Dev Cell *19*, 296-306. 10.1016/j.devcel.2010.07.019.

678 35. Bradshaw, A.D. (2009). The role of SPARC in extracellular matrix assembly. J Cell
679    Commun Signal *3*, 239-246. 10.1007/s12079-009-0062-6.

680 36. Chioran, A., Duncan, S., Catalano, A., Brown, T.J., and Ringuette, M.J. (2017). Collagen
681    IV trafficking: The inside-out and beyond story. Dev Biol *431*, 124-133.
682    10.1016/j.ydbio.2017.09.037.

683 37. Kalra, S., Bahendeka, S., Sahay, R., Ghosh, S., Md, F., Orabi, A., Ramaiya, K., Al
684    Shammari, S., Shrestha, D., Shaikh, K., et al. (2018). Consensus Recommendations on
685    Sulfonylurea and Sulfonylurea Combinations in the Management of Type 2 Diabetes
686    Mellitus - International Task Force. Indian J Endocrinol Metab *22*, 132-157.
687    10.4103/ijem.IJEM_556_17.

688 38. Shin, M., Chang, E., Lee, D., Kim, N., Cho, B., Cha, N., Koranteng, F., Song, J.J., and
689    Shim, J. (2024). Drosophila immune cells transport oxygen through PPO2 protein
690    phase transition. Nature *631*, 350-359. 10.1038/s41586-024-07583-x.

691 39. Marcetteau, J., Duarte, P., Leitao, A.B., and Sucena, E. (2025). Transdifferentiation of
692    plasmatocytes to crystal cells in the lymph gland of Drosophila melanogaster. EMBO
693    Rep *26*, 2077-2097. 10.1038/s44319-025-00366-z.

694 40. Chechenova, M., McLendon, L., Dallas, B., Stratton, H., Kiani, K., Gerberich, E.,
695    Alekseyenko, A., Tamba, N., An, S., Castillo, L., et al. (2024). Muscle degeneration in
696    aging Drosophila flies: the role of mechanical stress. Skelet Muscle *14*, 20.
697    10.1186/s13395-024-00352-4.

698 41. Siles, L., Ninfali, C., Cortes, M., Darling, D.S., and Postigo, A. (2019). ZEB1 protects
699    skeletal muscle from damage and is required for its regeneration. Nature
700    Communications *10*, 1364. 10.1038/s41467-019-08983-8.

701 42. Kirkwood, T.B. (2005). Understanding the odd science of aging. Cell *120*, 437-447.
702    10.1016/j.cell.2005.01.027.

703 43. Baghdadi, M.B., Castel, D., Machado, L., Fukada, S.I., Birk, D.E., Relaix, F., Tajbakhsh,
704    S., and Mourikis, P. (2018). Reciprocal signalling by Notch-Collagen V-CALCR retains
705    muscle stem cells in their niche. Nature *557*, 714-718. 10.1038/s41586-018-0144-9.

706   44.   Molina, T., Fabre, P., and Dumont, N.A. (2021). Fibro-adipogenic progenitors in
707   skeletal muscle homeostasis, regeneration and diseases. Open Biol *11*, 210110.
708   10.1098/rsob.210110.

709   45.   Shahab, J., Baratta, C., Scuric, B., Godt, D., Venken, K.J., and Ringuette, M.J. (2015).
710   Loss of SPARC dysregulates basal lamina assembly to disrupt larval fat body
711   homeostasis in Drosophila melanogaster. Dev Dyn *244*, 540-552.
712   10.1002/dvdy.24243.

713   46.   Mase, A., Augsburger, J., and Bruckner, K. (2021). Macrophages and Their Organ
714   Locations Shape Each Other in Development and Homeostasis - A Drosophila
715   Perspective. Front Cell Dev Biol *9*, 630272. 10.3389/fcell.2021.630272.

716   47.   Rodrigues, D., Renaud, Y., VijayRaghavan, K., Waltzer, L., and Inamdar, M.S. (2021).
717   Differential activation of JAK-STAT signaling reveals functional compartmentalization
718   in Drosophila blood progenitors. Elife *10*. 10.7554/eLife.61409.

719   48.   Lin, L. (2009). Clonal Analysis of Growth Behaviors During Drosophila Larval Tracheal
720   Development In Doctoral thesis. (University of Basel).

721   49.   Antonello, Z.A., Reiff, T., Ballesta-Illan, E., and Dominguez, M. (2015). Robust intestinal
722   homeostasis relies on cellular plasticity in enteroblasts mediated by miR-8-Escargot
723   switch. EMBO J *34*, 2025-2041. 10.15252/embj.201591517.

724   50.   Leroux, E., Ammar, N., Moinet, S., Pecot, T., and Boukhatmi, H. (2023). Studying
725   Muscle Transcriptional Dynamics at Single-molecule Scales in Drosophila. J Vis Exp.
726   10.3791/65713.

727   51.   Chakraborty, K., VijayRaghavan, K., and Gunage, R. (2018). A Method to Injure,
728   Dissect and Image Indirect Flight Muscle of Drosophila. Bio Protoc *8*, e2860.
729   10.21769/BioProtoc.2860.

730
731

Figure 1

Figure 1. Hemocytes are specifically recruited to the damaged muscle fiber. (A-F). Adult indirect flight muscles (IFMs) imaged under uninjured conditions and at 2-and 5-days post-injury (2DPI and 5DPI). Mhc (magenta) marks muscles. Hemocytes (green) are visualized using the following reporters: *Pxn-RFP* in (A-C'') and *Enh3-Gal4; UAS-mCD8GFP* in (D-F''). Dashed yellow outlines indicate injured muscle fibers showing elevated Mhc expression. **(G).** Time-course analysis of hemocytes recuitment at 1, 2, 3, 4 and 5 DPI reveals heterogeneous kinetics of response to muscle damage. Enh3-positive, Pxn-positive, and Hml-positive hemocytes are represented by light gray, black, and dark gray bars, respectively (****$p < 0.0001$, Student's t-test; n ≥ 10 heminota per condition). **(G').** Schematic representation of the proportion of recruited hemocytes expressing different reporters. **(H-I''').** Distribution of hemocytes visualized with *Enh3-Gal4; UAS-mCD8GFP* (green) in control muscles (H-H''') and *Naa30A* mutants (I-I'''). Muscles and muscle nuclei are stained with Mhc (magenta) and Mef2 (cyan) respectively. Scale bar: 200µm. Asterisks in panels H''' and I''' mark non-specific signal detected in motoneurons.

Figure 2

Figure 2. Muscle damage induces directed tracheal remodelling. (A-A'). Hemocytes (green/magenta; *Enh3-Gal4, UAS-mCD8GFP; UAS-H2B-RFP*) are associated to the muscle tracheal cells (gray, autofluorescence). Scale bars: 25 µm. **(B-C'').** Distribution of tracheal branches (gray, autofluorescence) and hemocytes (green, *Enh3-Gal4; UAS-mCD8GFP*) in control (B-B'') and at 5DPI (C-C''). Muscles are labeled with Mhc (magenta). (B'', C''). Boxed regions from (B and C), magnified 2X. Scale bars: 200 µm. **(D).** Quantification of tracheal coverage in (B) and (C) (****p < 0.0001, Student's t-test; n = 19 heminota per condition; light, dark, and intermediate shading indicate data points from three independent replicates). **(E).** Quantification of tracheal coverage in (C) at the injury site (orange box), adjacent to the injury site (green box, approximately 135 µm away from the injury site), and at distant regions (blue box, within 300 µm of the lesion border) (*p= 0.0142, Student's t-test; n=19 heminota). **(F-F').** dSRF-Gal4 marks muscle-associated terminal tracheal cells (TTCs). TTC membranes (green) and nuclei (magenta) are indicated by arrowheads and arrows, respectively (*dSRF-Gal4, UAS-mCD8GFP; UAS-H2B-RFP*). dSRF-Gal4 is inactive in primary tracheal branches (asterisks in F'). Muscles are labeled with Mhc (blue). Scale bars: 25 µm. **(G-H''').** TTCs distribution in IFMs (green, *dSRF-Gal4, UAS-mCD8GFP*) in control (G-G''') and at 5DPI (H-H'''). Muscles are labeled with Mhc (magenta).(G'',H''). 4.5X magnifications of boxed regions in (G') and (H'). Scale bar: 200 µm. (G''', H'''). Orthogonal views of (G'') and (H''). Arrow in (H''') indicates TTC enrichment at the injured muscle. **(I).** Quantification of TTCs coverage from (G) and (H) (****p < 0.0001, Student's t-test; n = 14 heminota per condition; light, dark, and intermediate shading points represent data points from three independent replicates). **(J-J').** dSRF (green) marks TTCs nuclei associated with IFMs. Scale bars: 25 µm. **(K-L''').** Distribution of TTCs nuclei (green) in IFMs under control conditions (K-K''') and at 5DPI (L-L'''). Muscles are labeled with phalloidin (blue); tracheae are visualized by autofluorescence (gray). Anti-PH3 (magenta) is used to assess cell division. (K', L'''). 2X magnification of boxed regions in (K) and (L). Scale bars: 100 µm. Arrowheads and arrows indicate dSRF expression in the muscle and TTCs nuclei, respectively. **(M).** Proportion of TTCs nuclei located within a selected muscle relative to the total number of TTCs nuclei in the imaged sample (*p = 0.036, Student's t-test; n = 17 heminota per condition; light, dark, and intermediate shading indicate data points from three independent replicates).

Figure 3

Figure 3. Local hypoxia in response to muscle damage drives tracheal branching. (A-A''). *SyHREns* reporter activity (green) is detected in a subset of TTCs marked by dSRF (magenta, arrows). Scale bars: 25 µm. **(B-D'').** Expression of the *SyHREns* reporter (green) in control IFMs (B-B''), at 5DPI (C-C'') and 20DPI (D-D''). Muscles are labeled with Mhc (magenta), and tracheal branches are visualized via autofluorescence (gray). (B'', C'' and D''). 2X magnification of the boxed regions in B', C' and D'. Scale bars: 200 µm. **(E).** Quantification of SyHREns positive cells (green) per injured muscle fiber (****p < 0.0001, Student's t-test; n > 20 heminota per condition; light, dark and intermediate shading indicate data points from three independent replicates). **(F-H'').** Sima (HIF-1α) downregulation in TTCs (*dSRF-Gal4; UAS-Sima^{RNAi}*) prevents tracheal recruitment at the injured fiber. (F'', G'' and H''). 2X magnification of the boxed regions in E', F' and G'. Scale bars: 200 µm. **(I).** Quantification of trachea coverage (****p < 0.0001, Student's t-test; n = 13 heminota per condition). **(J-K'').** Detection of Bnl (green, *Bnl::GFP^{endo}*) in control IFMs (J-J'') and at 5DPI (K-K''). Muscles are labeled with Mhc (magenta), and tracheal cells are visualized via autofluorescence (gray). (J'', K''). 4.5X magnification of the boxed regions in J and K. Scale bars: 200 µm. (J''', K'''). Orthogonal views of (J) and (K). Asterisk in K''' indicates Bnl::GFP^{endo} enrichment at the injured muscle. **(L).** Quantification of normalized Bnl::GFP^{endo} intensity level in uninjured and injured muscle fiber at 5DPI (***p<0.0001, Student t-test, n = 15 heminota per condition from three independent replicates).

Figure 4

Figure 4. Hemocytes deliver Bnl to injured muscle to promote tracheal recruitment. (A-C'). Local overexpression of UAS-Bnl::GFP (green) in adult IFMs using *SME-Gal4* driver induces enrichment of terminal tracheal cells (TTCs; dSRF, magenta) and increases tracheal branching (gray), compared with controls (A). Scale bars: 200 µm. **(D-E'').** *Bnl-LacZ* reporter expression (green) is downregulated specifically in of the injured muscle fibers overexpressing Mhc (magenta), the nuclei are marked with DAPI (blue). (D', E''). 3.5X magnification of the boxed regions in D and E. Scale bars: 200 µm. **(F-G'').** Bnl expression (green, *Bnl::GFP^endo^*) in uninjured IFMs (F-F'') and at 5DPI (G-G''). Muscles are labeled with Mhc (blue), tracheal cells visualized via autofluorescence (gray), and hemocytes detected with the *Pxn-RFP* reporter (magenta). Arrows in G'' indicate Bnl localization around hemocytes recruited to the injured muscle. Scale bars: 200 µm. **(H-J'').** Downregulating *bnl* in the hemocytes (green, *Enh3-Gal4; UAS-mCD8GFP, UAS-Bnl^RNAi^*) do not alter hemocytes recruitment but decreases tracheal invasion to the damaged fiber as shown in (J'- J'') and quantified in **(K)** (**** $p< 0.0001$, Student t-test; n = 16 heminota from two independent replicates). Scale bars: 200 µm. **(L).** Schematic of the flight-assay setup used to assess flight performance. Flies are released at the top of a graduated cylinder, and landing position across three vertical zones is recorded. **(L').** Quantification of the proportion of flies distributed across three sections under three conditions: *Control Uninjured*, *Control Injured (5DPI)* and *Enh3-Gal4; UAS-Bnl^RNAi^ Injured (5DPI)*. Bars indicate the percentage of flies landing in each of the three flight-assay zones (***$p<0.001$, * $p < 0.05$ Chi-2 test). **(M).** Schematic model of hemocyte-driven tracheal recruitment required to maintain flight capacity following muscle injury.

Figure 5

Figure 5. The FGF receptor Breathless (Btl) is expressed along tracheal branches and is required for tracheal recruitment to the damaged muscle. (A-B''). Btl::Cherry^{endo} (magenta) is detected along tracheal branches (gray) associated with indirect flight muscles (IFMs) labelled with Mhc (blue), in either uninjured A or injured muscles B. Scale bars: 50 µm. Dashed yellow outlines indicate injured muscle fibers showing elevated Mhc expression. **(C-C').** The *Btl-Gal4* driver (green and magenta; *Btl-Gal4, UAS-mCD8GFP; UASH2B-RFP*) is active in a subset of tracheal branches (arrows, C-C'). Arrowheads in C indicate the tracheal branches that do not express the *Btl-Gal4* driver. Scale bars: 25 µm. **(D-E').** Activity of the *Btl-Gal4* driver (green, *Btl-Gal4, UAS-mCD8GFP*) in control (D) and at 5DPI (E). Muscles are labelled with Mhc (magenta). Scale bars: 200 µm. **(F-I'').** Downregulating *btl* in the tracheal cells with either *Btl-Gal4* driver (*Btl-Gal4; UAS-Btl^{RNAi}*, H-H'') or *dSRF-Gal4* driver (*dSRF-Gal4, UAS- Btl^{RNAi}*, I-I'') prevents tracheal invasion to the damaged fiber. **(J).** Quantification of trachea coverage (****p <0.0001, **p=0.0053, Student's t-test; n = 7 heminota per condition). Scale bars: 200 µm. **(K).** Schematic model of hemocyte-driven Bnl activates tracheal cells recruitment via Bnl/Btl signalling pathway.

Figure 6

Figure 6. Hemocytes deposit ECM components at injury muscles to constrain Bnl to damaged muscle fibers and maintain muscle function. (A-A''). Collagen IV (green, *Vkg::GFP*) is detected at basal levels around muscle fibers (arrow, A'), tracheal branches (arrowheads, A'), and hemocytes (magenta, *Pxn-RFP*; arrow, A'') in uninjured conditions. **(B-B'').** At 5DPI, Collagen IV is strongly enriched on the injured muscle fibers, labelled with Mhc (blue). **(C-D'').** Hemocytes recruitment (magenta, *Pxn-RFP*) and Collagen IV (green, *Vkg::GFP*) distribution on the injured muscle, at 1DPI and 4DPI. Scale bars: 200 µm. **(E-F'').** Endogenous Bnl (green, *Bnl::GFP^endo*) is enriched along damaged muscle fibers (magenta, Mhc) and is markedly reduced upon hemocyte-specific depletion of Collagen IV (*Enh3-Gal4, UAS-Vkg^RNAi*). **(G).** Quantification of normalized Bnl intensity in injured fibers (****p<0.0001, Student t-test, n = 13 heminota per condition). **(H).** Schematic model of hemocyte-mediated ECM delivery required to confine Bnl to injured muscle fibers. **(I-I').** SPARC (magenta) is detected in the IFMs associated hemocytes visualized by *Enh3-Gal4* driver (green, *Enh3-Gal4, UAS-mCD8GFP*). **(J-J').** SPARC expression is increased in injured fibers compared to controls (I-I'). **(K-K').** Adult hemocytes (magenta; *SPARC-Gal4, UAS-myr-RFP*) express BM40-SPARC-GFP (green). **(L-L').** BM40-SPARC-GFP produced by adult hemocytes is broadly distributed along the injured muscle fibers. **(M-N').** Downregulation of *SPARC* in adult hemocytes (*Enh3-Gal4 > UAS-SPARC^RNAi*) reduces Collagen IV enrichment (green, *Vkg::GFP*) and impairs tracheal recruitment in injured fiber (gray, autofluorescence), as quantified in panels **(O)** and **(P)** respectively (O. ****p<0.0001, Student t-test, n = 20 heminota per condition from two independent replicates, P. ***p<0.001, Student t-test, n = 18 heminota per condition from two independent replicates). **(Q).** Quantification of the proportion of flies distributed across three sections (Top, Middle and Bottom) under three conditions: *Control Uninjured, Control Injured (5DPI)* and *Enh3-Gal4; UAS-SPARC^RNAi Injured (5DPI)* (***p<0.001, ** p < 0.01 Chi-2 test).

Figure 7

Figure 7. Hemocyte-muscle-tracheal intercellular signaling orchestrates tracheal remodelling and muscle repair after injury. (A). Schematic representation of the hemocyte-muscle-tracheal signaling module activated in response to muscle injury. **(B).** Upon muscle damage, hemocytes are rapidly recruited to the injured fiber, where they secrete the FGF ligand Branchless (Bnl), which activates its receptor Breathless (Btl) on tracheal cells. Bnl–Btl signaling drives tracheal remodeling and directional outgrowth toward the injured muscle. In parallel, hemocytes deposit extracellular matrix (ECM) components that retain Bnl and stabilize trachea-muscle attachments. Together, these coordinated cellular mechanisms promote muscle repair and preserve flight performance.

Figure S1

Supplementary data

Figure S1. Mhc upregulation marks the damaged muscle fiber. (A-B''). IFMs stained for Mhc (magenta) and Mef2 (cyan) show upregulation of Mhc and downregulation of Mef2 in the injured fiber (asterisks, B'-B'') at 5DPI, compared to the control (A'-A''). (A', B'). 2X magnification of the boxed regions in A and B. Scale bars: 200 μm. **(C-C'').** Expression of the embryonic Mhc isoform (*weeP26*, green) in uninjured (C) and injured (C'-C'') IFMs. Injury does not alter weeP26 expression (C and C''). **(D-D'').** The shorter Mhc isoforms (*Mhc-Iso-K, L, M (fTRG500)*; green) colocalize with anti-Mhc staining (magenta) and are upregulated in injured muscle (D'-D''). **(E-E'').** The longer Mhc-GFP isoforms (*Mhc-Iso-A, F, G (fTRG519)*; green) are not expressed in adult IFMs and are not upregulated in response to muscle injury (E'-E''). Scale bars: 200 μm. **(F-G'').** In *Naa30A mutant* IFMs, defective muscle attachment correlates with increased Mhc expression (magenta, G), upregulation of short Mhc-GFP isoforms (green, G'), and a loss of Mef2 (cyan, G''). **(H-I'').** Co-expression of hemocytes markers *HmlΔ-Gal4, UAS-mCD8GFP* (green) and *Pxn-RFP* (magenta) reveals two subsets of hemocytes associated with the IFMs: Pxn+Hml+ (arrows) and Pxn+Hml- (arrowheads). Nuclei are visualized with TO-PRO (blue). Data from n >10 heminota from two independent replicates. Scale bars: 200 μm. (H''-I''). 3X magnification of the boxed regions in H and I. Scale bars: 100 μm. **(J-M').** Co-expression analysis of *Enh3-Gal4; UAS-mcD8GFP* (green) and *Pxn-RFP* (magenta) shows that all Enh3+ cells express Pxn (J'-K'). No Mef2 expression (cyan) is observed in these cells, in either undamaged (L') or damaged (M', dashed lines) fibers. Scale bars: 200 μm. Yellow asterisks in panels F', G'' and L' mark non-specific signal detected in motoneurons.

Figure S2

Figure S2. Early recruitment of hemocytes initiates tracheal remodeling after muscle damage. (A-B'). In *Naa30A mutant* flies, defective muscle attachment is associated with tracheal enrichment (gray, B') and increased hemocytes accumulation (green, B') compared to controls (A-A'). **(C-K').** Time-course analysis of tracheal remodeling and hemocytes recruitment in *Enh3-Gal4; UAS-mCD8GFP* adult IFMs following localized stab injury. Tracheal branches are visualized by autofluorescence (gray), and hemocytes are labeled with GFP (green). Muscles are labeled with Mhc (magenta). Time points range from 1-hour post-injury (1HPI) to 20 days post-injury (20DPI). **(L).** Quantification of normalized Mhc intensity. **(M).** Quantification of the hemocytes number in the injured fiber shows a significant increase starting at 1HPI which remains elevated until 20DPI. **(N).** Tracheal quantification shows a marked increase in its density from 3DPI and remains sustained until 20 DPI. (****$p < 0.0001$, ***$p < 0.001$; Student's t-test; n ≥ 10 heminota per condition). Scale bars: 200 µm.

Figure S3

Figure S3. Tracheal branching requires precise control of Bnl provided by the hemocytes. (A-B'). Expression pattern of the Shavenbaby (Svb) muscle regulatory enhancer (SME) (green, *SME-Gal4; UAS-mCD8GFP*). SME-Gal4 drives GFP expression (green) in a subset of larval muscle progenitors located in the posterior region of the wing disc (A-A') and in a subset of adult muscle nuclei (green and blue (DAPI), B-B'). Scale bars: 200 µm. **(C).** Schematic representation of the FLP-OUT system used to drive UAS-Bnl::GFP expression in a subset of adult muscle nuclei, using the *Hs-FLP; FRT-SME-Stop-FRT-Gal4/CyO; tub-Gal80ts* combination. **(D-D').** Bnl (green, *Bnl::GFP^{endo}*) is detected in indirect flight muscles (IFMs, labelled with Mhc, blue; arrowheads), tracheal branches (autofluorescence, gray; asterisks), and hemocytes (*Pxn-RFP*, magenta; arrows). Scale bars: 25 µm. **(E-E').** Expression of the *Bnl-LacZ* reporter (green) is observed in terminal tracheal cells (TTCs) marked by dSRF (magenta; asterisks), and in IFMs nuclei (arrowheads). Scale bars: 25 µm. **(F-F').** Bnl-LacZ (green) is detected in IFMs associated hemocytes labelled with *Pxn-RFP* (magenta; arrows). Scale bars: 25 µm. **(G-I'').** Targeted downregulation of *bnl* in a subset of hemocytes using *HmlΔ-Gal4* driver significantly reduces tracheal enrichment at the injured fiber, as quantified in **(J).** ($*p < 0.05$, Student's t-test; n ≥ 10 heminota per condition). Scale bars: 200 µm.

Figure S4

Figure S4. Time-course analysis of hemocyte recruitment and Collagen IV deposition following muscle injury. (A-F). Hemocyte recruitment (magenta, *Pxn-RFP*) and Collagen IV distribution (green, *Vkg::GFP*) in uninjured muscles (A) and at successive time points after injury (B-F), from 1 to 20 days post injury (DPI). Scale bar: 200 µm.