Altered cognitive processes shape tactile perception in autism.

Ourania Semelidou[1,#], Mathilde Tortochot-Megne Fotso[1,*], Adinda Winderickx[1,*], Andreas Frick[1,#]

[1] *University of Bordeaux, Neurocentre Magendie, INSERM U1215, 33000, Bordeaux, France*
[*] *Equal contribution*
 # Co-corresponding authors: osemelidou@gmail.com, andreas.frick@inserm.fr

Altered sensory perception is a hallmark of autism and shapes how individuals engage with their
environment, with tactile perception playing a critical role in daily functioning and for social interactions.
While sensory alterations are thought to contribute to cognitive differences in autism, the impact of
cognition on sensory perception remains unclear. Here, we investigated how cognitive processes
modulate tactile perception in the *Fmr1*-KO genetic mouse model of autism through a translational
perceptual decision-making task. Our results revealed salience-dependent cognitive alterations that
influenced sensory performance. During training, *Fmr1*$^{-/y}$ male mice distinguishing between a high- and a
low-salience stimulus exhibited an increased choice consistency bias in low-salience trials. When tested
across a continuum of intermediate stimulus intensities, these mice demonstrated enhanced tactile
discrimination of low-salience stimuli but reduced discrimination facilitation for stimuli crossing category
boundaries. These effects were accompanied by diminished integration of sensory history and were
dissociable from the attention deficits that emerged under high cognitive load. Together, our findings
reveal that tactile perceptual alterations reflect context-dependent weighting and integration of sensory
information during decision-making rather than uniform sensory deficits or enhancements, supporting a
shift beyond traditional sensory–cognitive dichotomies.

Introduction

Autism is a neurodevelopmental condition that affects approximately 1 in 31 children (Shaw et al., 2025).
Autistic individuals display differences in social interaction and communication, repetitive behaviors,
intense or special interests, and altered sensory experience as core symptoms (American Psychiatric
Association, 2022). Sensory alterations strongly affect the way autistic individuals interact with their
environment. According to the DSM-5-TR, these are broadly expressed as "*hyper- or hyporeactivity to*
sensory input or unusual interests in sensory aspects of the environment" and are reported in 90% of
autistic individuals, spanning all sensory modalities (Robertson and Baron-Cohen, 2017).

Amongst sensory modalities, touch is the first sense to develop and plays a fundamental role in the
exploration of the environment, the definition of "self", and early social bonding (Bremner and Spence,
2017). Tactile processing alterations have a significant impact on daily functioning and are thought to
contribute to repetitive behaviors and social interaction difficulties in autism (Foss-Feig et al., 2012; Green
et al., 2018; Thye et al., 2018; He et al., 2021a; Zhai et al., 2023; Carati et al., 2024). However, clinical
studies focusing on tactile discrimination have yielded controversial results (Zetler et al., 2019), likely due
to the substantial heterogeneity of tactile processing alterations in autism (Mikkelsen et al., 2018) and
inconsistent terminology across studies (He et al., 2023). In contrast, numerous studies in the visual and
auditory domains (Plaisted et al., 1998a; O'Riordan and Plaisted, 2001; O'Riordan et al., 2001; Bonnel et
al., 2003; O'Riordan and Passetti, 2006; Samson et al., 2006; Rotschafer, 2021) have supported enhanced
sensory processing in autism, as proposed by several influential models of the condition (Plaisted, 2001;
Frith, 2003; Mottron et al., 2006).

Although clinical studies highlight the heterogeneity of tactile responses in autism, variability in
experimental protocols and stimulus types has made it difficult to draw cohesive conclusions. Tactile
discrimination of vibrotactile stimuli delivered to the glabrous skin of the fingers was reported reduced
(Puts et al., 2014; Espenhahn et al., 2023), or intact (He et al., 2021a; Asaridou et al., 2022) in autistic
individuals. Similarly, studies of sharp-dull discrimination have reported both reduced (Abu-Dahab et al.,
2013) and typical performance (Minshew et al., 1997; Minshew and Hobson, 2008), as have studies of
form discrimination (Minshew and Hobson, 2008; Demopoulos et al., 2015; Failla et al., 2017).
Translational approaches employing mouse models of autism in combination with psychometric measures
from well-controlled behavioral tasks may help clarify whether these inconsistencies reflect genuine inter-
individual variability or arise from methodological differences.

While tactile discrimination has been explored in several clinical studies, it remains underexplored in
mouse models of autism. Rodent studies leveraging innate responses have reported atypical whisker-
dependent texture discrimination (Balasco et al., 2021, 2022; Mattioni et al., 2024) and forepaw-
dependent roughness discrimination (Orefice et al., 2016). However, these studies have not employed
psychophysical methods to quantify tactile discrimination, restricting the translatability of their findings.
Implementing psychophysics in animal models might clarify previous inconsistencies between studies in
mice (Orefice et al., 2016) and humans (O'Riordan and Passetti, 2006).

Sensory alterations are thought to impact cognition in autism (Haigh, 2018). However, sensory perception
is shaped not only by the physical properties of stimuli but also by higher-order cognitive processes such
as attention and categorization—both of which are altered in autism (Allen and Courchesne, 2001; Church
et al., 2010; Gastgeb and Strauss, 2012; Cascio et al., 2015; Barbot et al., 2018; Rong et al., 2021a).
Moreover, autistic individuals demonstrate differences in how prior beliefs and expectations (priors) are

integrated with sensory input during perceptual decision-making. They tend to rely less on contextual
priors (Amoruso et al., 2019) and more heavily on prior choices (Feigin et al., 2021), and often form priors
that are imprecise or inflexible (Zaidel et al., 2015; Sapey-Triomphe et al., 2021b). Notably, variability in
the use of priors has also been linked to individual differences in sensory responsivity (Sapey-Triomphe et
al., 2021a). Thus, cognitive differences may compound or interact with primary sensory processing
alterations, contributing to the distinct perceptual experiences observed in autism.

In this study, we investigated the interplay between tactile discrimination and cognitive processes during
perceptual decision-making. To enhance translational relevance and address questions of inter-individual
heterogeneity, we developed a forepaw-based 2-Alternative Choice task for mice, using vibrotactile
stimuli analogous to those used in human psychophysical protocols. Leveraging this task, we explored
how stimuli of varying salience are discriminated in the *Fmr1$^{-/y}$* mouse model of autism and disentangled
stimulus-driven from cognitively modulated-tactile responses.

Our results revealed salience-dependent alterations in both tactile processing and cognitive modulation
in *Fmr1$^{-/y}$* mice, with a decreased influence of cognitive processes on tactile perception. During training,
Fmr1$^{-/y}$ mice exhibited heightened choice consistency bias during low-salience trials, which contributed to
reduced performance during perceptual learning. In contrast, trained *Fmr1$^{-/y}$* mice displayed enhanced
discrimination of low-salience stimuli, but a diminished facilitatory effect of categorization on across-
category discrimination. Despite their increased discrimination sensitivity at the low-salience range, *Fmr1$^{-/y}$*
mice showed attention deficits specifically for these stimuli under high cognitive load conditions. While
a strong choice consistency bias was observed during the testing phase for both WT and *Fmr1$^{-/y}$* mice,
integration of recent sensory history was selectively disrupted in *Fmr1$^{-/y}$* mice, such that past sensory
information no longer informed ongoing decisions.

These findings uncover distinct, salience-dependent cognitive alterations that shape tactile perception in
the *Fmr1$^{-/y}$* model, advancing our understanding of tactile perception in autism. Importantly, our results
indicate that the variability observed in clinical studies may arise not solely from primary sensory
alterations but also from differences in cognitive context, attentional demands, or task complexity—
offering a potential framework to reconcile prior inconsistencies in the field.

Results

***Fmr1$^{-/y}$* mice show intact learning rate and trajectory during perceptual learning.**

To investigate perceptual and cognitive alterations in a highly translational approach, we developed a
forepaw-based 2-Alternative Choice perceptual decision-making task in the *Fmr1*-KO mouse model of
autism. In this task, mice learn to discriminate between vibrotactile stimuli that differ in perceptual
salience, which we define as the prominence of a stimulus based on its physical amplitude. To ensure that
all mice could reliably detect the applied stimuli, we selected amplitudes above the perceptual thresholds
previously established for both wild-type (WT) and *Fmr1$^{-/y}$* male mice (Semelidou et al., 2024).
Accordingly, all stimuli used in the present study (40 Hz, ≥12 μm) were well above the detection threshold
(10 Hz, 4.46 μm for WT; 10Hz, 7.29 μm for *Fmr1$^{-/y}$* mice), allowing us to specifically probe discrimination,
categorization, and decision-making rather than sensory detection.

During each trial, head-fixed, water-controlled mice received a 500 ms vibrotactile stimulus to the
forepaw at 40 Hz with either high amplitude (26 μm) or low amplitude (12 μm). Stimulus salience was

defined in a relative manner: across animals, the lower-amplitude stimulus elicited consistently higher
miss rates, indicating reduced perceptual strength and attentional engagement within the suprathreshold
range. We therefore refer to the 12 µm stimulus as "low salience" and the 26 µm stimulus as "high
salience." Mice were trained to report high-salience stimuli by licking the right lick-port, and low-salience
stimuli by licking the left port within a 2-s response window following stimulus delivery. Incorrect choices
resulted in a 5-s timeout (Fig. 1A).

During early training sessions, performance was low, characterized by a low proportion of correct and a
high proportion of incorrect choices, which either remained stable or gradually improved over the course
of the session (Fig. S1A-left). Licking patterns were also inconsistent during these early sessions and poorly
time-locked to the response window (Fig. S1A-right). In contrast, late training sessions showed
consistently high correct choice rates and low incorrect choice rates (Fig. S1B-left), along with stable
licking patterns during the response window for both high- and low-salience trials (Fig. S1B-right).

Learning difficulties and altered learning trajectories have been reported in autistic individuals and
replicated in several mouse models of autism, including $Fmr1^{-/y}$ mice (Arnett et al., 2014a; Goel et al.,
2018; Mercado et al., 2020; Khachadourian et al., 2023; Shenouda et al., 2023; Mol et al., 2024). To assess
whether $Fmr1^{-/y}$ mice exhibit alterations in learning rate in our task, we compared the number of days
required to reach the predefined learning criterion (>70% correct choices for each stimulus salience across
three consecutive training days). Our results showed no significant differences between $Fmr1^{-/y}$ mice and
their WT littermates in the total number of training days to reach criterion (Fig. 1B; T test: t = 0.438, p =
0.667) or in the duration spent at each stage of the training protocol (Fig. S1C-D, see Methods; Blocks: T
test: t = 1.259, p = 0.224; Mixed: T test: t = -0.185, p = 0.855). Similarly, both genotypes required a
comparable number of days to learn the association between stimulus salience and the correct lick-port
(Fig. 1C-D; High salience: Mann-Whitney U test: U=45.5, p = 0.788; Low salience: Mann-Whitney U test: U
= 38.5, p = 0.422).

To capture potential genotype differences in the learning dynamics during training, we analyzed the
performance of all mice that reached the learning criterion during the training phase with mixed low- and
high-salience trials. Both WT and $Fmr1^{-/y}$ mice showed similar learning dynamics, with a slow, gradual
improvement across days (main effect of day: β = 0.007, SE = 0.004, z = 1.882, p = 0.060, 95% CI [0.00,
0.01]) (Fig. S2A-top). Consistent with the extended training duration required for the task, learning slopes
were shallow and comparable between genotypes (Fig. S2B; Mann-Whitney U test: U=38.0, p=0.536).
Importantly, the Genotype × Day interaction was small (β = 0.002), indicating no detectable genotype
differences in learning rate across animals.

These similar learning trajectories were accompanied by comparable baseline performance on the first
day of training across genotypes (Fig. S2A-top, S2C; genotype effect: β = −0.087, SE = 0.066, z = -1.328, p
= 0.184, p = 0.184, 95% CI [-0.22, 0.04]) and similar within-animal day-to-day performance variability (Fig.
S2A-bottom; Fig. S2D; T test: t=-0.302, p=0.767).

We next assessed whether performance differed during an intermediate stage of learning, defined for
each animal as the middle three days of its training period. Here, both genotypes again showed
comparable performance (Fig. S2E, slope; T-test: t = 0.801, p = 0.437) with similar correct-choice rates
(Fig. S2F; T-test: t=1.240, p=0.228), indicating no genotype-specific differences during intermediate
learning.

Together, these results show that *Fmr1*^{-/y} mice display an intact learning rate and trajectories in this 2-AFC
perceptual decision-making task.

Fmr1^{-/y} mice display impaired performance on low-salience trials during perceptual learning.

Given that comparable learning trajectories and training duration to reach the criterion do not exclude
the presence of performance alterations, we next evaluated additional behavioral metrics during training
with mixed high- and low-salience trials. Specifically, we analyzed overall discrimination sensitivity (d') as
well as correct and incorrect choice rates throughout training to determine whether *Fmr1*^{-/y} mice
exhibited altered task performance despite intact learning progression.

To quantify mice's sensitivity in discriminating high- versus low-salience stimuli throughout training, we
used the signal detection theory and computed the sensitivity index d'. This measure captures stimulus
discriminability by comparing normalized hit rates (rightward licks on high salience trials) and false alarm
rates (rightward licks on low salience trials). *Fmr1*^{-/y} mice showed a strong trend with a large effect size
toward reduced overall task sensitivity compared to WT littermates (Fig. 1E; T-test: t=-2.083, p=0.052,
Hedges' g=0.90), indicating diminished perceptual performance during training. These results
demonstrate that, despite acquiring the task at a comparable rate (Fig. 1B-D) and with similar learning
trajectories (Fig. S2), *Fmr1*^{-/y} mice display a trend for diminished performance during salience
discrimination training.

To complement our sensitivity (d') analysis, we also quantified correct and incorrect choice rates, allowing
us to assess both rightward and leftward lick responses. Correct choice rates were comparable between
Fmr1^{-/y} and WT mice (Fig. 1F, T-test: t=1.064, p=0.307) with similar Hit rates for both high- and low-
salience stimuli (Fig. S1E–F, High salience-T-test: t=0.721, p=0.480; Low salience-T-test: t=1.060, p=0.303).
In contrast, *Fmr1*^{-/y} mice exhibited a significantly higher rate of incorrect choices (Fig. 1G; T-test: t=-2.099,
p=0.050; Hedge's g=0.90), an effect that was specific to low-salience trials (Fig. 1H; Mann–Whitney U =
23.0, p = 0.048; Hedge's g = 0.97), while high-salience error rates were similar across genotypes (Fig. 1I;
T-test: t=-1.072; p=0.298). After outlier analysis and exclusion of one *Fmr1*^{-/y} mouse, the genotype
difference was reduced to a trend, though with a large effect size, for incorrect choices (T-test: t=-1.803;
p=0.089, Hedge's g = 0.79) and for incorrect choices during low-salience trials (Mann–Whitney U = 23.0,
p = 0.079; Hedge's g = 0.89).

These results suggest a modest reduction in task performance in *Fmr1*^{-/y} mice, primarily driven by
increased errors on low-salience trials.

Fmr1^{-/y} mice show higher choice consistency bias during training.

Perceptual learning can further reveal potential alterations in decision strategy, manifested as changes in
response bias (decision criterion c) or expectation bias (priors). To examine whether mice exhibited
stereotyped licking toward one of the two lickports—a potential indicator of response bias—we calculated
the decision criterion. This analysis revealed no significant differences between *Fmr1*^{-/y} mice and WT
littermates (Fig. 2A; Mann–Whitney U: U=65.000, p=0.254), with both groups showing a bias toward the
left lickport (WT: Wilcoxon signed-rank test W=5.000, p=0.039; *Fmr1*^{-/y:} Wilcoxon signed-rank test
W=10.000, p=0.042).

Since mice did not exhibit differences in stereotyped licking, we could rule out simple response biases and
further assess whether their decisions are influenced not only by the sensory features of stimuli but also

by previous experiences that shape expectations (i.e., priors). To investigate the impact of prior
experience on current choices, we analyzed behavioral responses on a trial-by-trial basis. We first
examined how a rewarded choice in the previous trial influenced the performance in the current trial (Fig.
2B). *Fmr1*^{-/y} mice exhibited an increased rate of incorrect responses to low-salience stimuli when these
were preceded by a rewarded high-salience stimulus (Fig. 2C; T-test: t=-2.352, p=0.030). In contrast, no
genotype difference was observed when a high-salience stimulus followed a rewarded low-salience trial
(Fig. 2D; T-test: t=0.603, p=0.554).

We next tested whether *Fmr1*^{-/y} mice were more influenced by their prior choices. To assess this, we
compared the rates of repeated prior choices in current low- and high-salience trials, controlling for the
total number of correct and incorrect responses. Our analysis revealed higher previous choice repetition
during low-salience trials in *Fmr1*^{-/y} compared to WT mice (Fig. 2E; T-test: t=-2.618, p=0.017), but a
comparable impact of prior choices during high-salience trials (Fig. 2F; T-test: t=1.215, p=0.240).

To verify that the observed differences in performance between WT and *Fmr1*^{-/y} mice reflected differential
reliance on priors rather than differences in prior strength per se, we quantified the strength of the priors
induced by high- and low-salience stimuli. Specifically, we calculated the rate of choices that were
repeated following a low- or high-salience trial, correcting for the overall rate of correct and incorrect
responses. Our results showed similar prior strength between groups, both for high-salience (Fig. S3A; T-
test: t=-1.705, p=0.105) and for low-salience stimuli (Fig. S3B; Mann–Whitney U: U=38.500, p=0.425).

Together, these results indicate an increased choice consistency bias during low-salience trials, which
contributes to performance differences during perceptual learning in *Fmr1*^{-/y} mice.

Fmr1^{-/y} mice do not show attention deficits during perceptual learning.

Stimulus salience is closely linked to attentional processes (Kerzel and Schönhammer, 2013; Forschack et
al., 2022; Bouvier et al., 2023), and attentional alterations are well-documented in autistic individuals
(Allen and Courchesne, 2001; Grubb et al., 2013; Barbot et al., 2018; Licznerski et al., 2020; Rong et al.,
2021b). Given that *Fmr1*^{-/y} mice exhibit salience-dependent performance differences during training (Fig.
1-2), we asked whether their reduced task performance might also reflect underlying attention deficits.
To assess attention, we analyzed the trials in which the animal failed to respond to the stimulus (Miss
trials) (Fig 1A). Within-genotype analysis confirmed studies linking salience and attention in humans
(Kerzel and Schönhammer, 2013), and validated that low amplitude stimuli are less salient, yielding
significantly higher Miss rates in both groups (Fig. S3C; WT: Wilcoxon signed-rank: W=0.000, p=0.004;
Fmr1^{-/y}: Paired T-test: t=-4.113, p=0.002). However, there were no genotype differences in Miss rates for
either high- or low-salience stimuli (Fig. S3D-E; Miss rate High: Mann–Whitney U: U=51.000, p=0.939; Miss
rate Low: Mann–Whitney U: U=50.000, p=1.000). These data demonstrate that low-salience stimuli are
less effective at capturing attention in both *Fmr1*^{-/y} and WT mice, and that the reduced task performance
of *Fmr1*^{-/y} mice during perceptual training cannot be attributed to an attentional deficit during low-
salience trials.

Trained _Fmr1_^{-/y} mice show enhanced discrimination of low-salience stimuli.

Altered tactile discrimination has been reported in clinical studies, though findings are inconsistent
(reviewed in (Zetler et al., 2019)), and has also been observed in mouse models of autism (Orefice et al.,
2016; Balasco et al., 2021, 2022; Mattioni et al., 2024). To assess tactile discrimination in our study, we
incorporated the original training stimuli along with 6 intermediate amplitudes, spaced 2 μm apart. Stimuli

ranging from 20 to 26 μm (20, 22, 24, 26 μm) were categorized and rewarded as high-salience stimuli,
while those from 12 to 18 μm (12, 14, 16, 18 μm) were treated as low-salience stimuli (Fig. 3A). Only
animals that successfully acquired the task during the training phase were included in the tactile
discrimination test. *Fmr1*$^{-/y}$ and WT mice showed similar performance during the last three days of
training, with no significant differences in task sensitivity (d'; T-test: t=0.800, p=0.438; Fig. S4A), rates of
correct (Fig. S4B-C; High: T-test: t=0.442, p=0.666; Low: T-test: t=0.697, p=0.498) and incorrect responses
(Fig. S4D-E; High: T-test: t=-0.680, p=0.509; Low: T-test: t=-1.069, p=0.305), decision bias (Fig. S3F;
criterion c; Mann–Whitney U: U=28.000, p=0.955), use of priors (Fig S4G-L; Fig. S4G: T-test: t=0.485,
p=0.636; S4H: T-test: t=0.155, p=0.879; S4I: T-test: t=-0.667, p=0.516; S4J: T-test: t=-0.667, p=0.516; S4K:
T-test: t=-0.072, p=0.944; S4L: T-test: t=-0.472, p=0.644), or attention (Fig. S4M-N; High: Mann–Whitney
U: U=32.500, p=0.555; Low: T-test: t=0.087, p=0.932).

To evaluate task performance, we quantified the rate of right-lick responses, corresponding to reports of
high-salience stimuli, and constructed psychometric curves, which increased with stimulus amplitude in
both genotypes. Mixed-effects modeling revealed a strong main effect of amplitude (all amplitude
coefficients p < 0.005) but no evidence for a genotype-dependent change across the psychometric
function (amplitude x genotype interaction coefficients p > 0.266), indicating comparable performance
between WT and KO animals across stimulus intensities (Fig. 3A). A genotype-by-amplitude interaction
was observed only at 18 μm amplitude (β = 0.127, SE = 0.06, z = -2.18, p = 0.030, 95% CI [-0.24, -0.01]),
where responses were slightly reduced in WT mice. However, this effect did not survive post-hoc analysis
(T-test: t = 0.568, p = 0.580) and did not generalize across amplitudes.

We then assessed the animals' ability to discriminate high-salience stimuli by comparing the proportion
of right-lick responses to 26 μm and 24 μm stimuli. Our results showed that neither WT nor *Fmr1*$^{-/y}$ mice
were able to reliably distinguish between these two amplitudes (Fig. 3B; WT: Wilcoxon signed-rank:
statistic=3.000, p=0.156; *Fmr1*$^{-/y}$: Paired t-test: t = 1.447, p = 0.221). However, both groups reliably
discriminated between 26 μm and 22 μm stimuli (Fig 3C; WT: Wilcoxon signed-rank: statistic = 0.000, p =
0.031; *Fmr1*$^{-/y}$: Paired t-test: t = -4.929, p = 0.001). These results demonstrate that high-salience stimuli
differing by 4 μm in amplitude can be discriminated by both genotypes.

We next investigated low-salience stimulus discrimination by comparing the rate of right-lick responses
(reporting high salience) to stimuli of 12 μm and 14 μm amplitude. WT mice responded similarly to both
amplitudes (Fig. 3D-left; Paired t-test: statistic = -1.684, p = 0.153), consistent with their performance on
high-salience stimuli differing by 2 μm (Fig. 3B-left). In contrast, *Fmr1*$^{-/y}$ mice exhibited enhanced
discrimination for these low-salience stimuli, with increased high-salience report rates for the 14 μm
stimuli compared to those for 12 μm (Fig. 3D-right; Paired t-test: t = -5.007, p = 0.001). Both genotypes
efficiently discriminated 12 μm from 16 μm stimuli (Fig. 3E; WT: Paired t-test: t = -3.274, p = 0.022; *Fmr1*$^{-}$
$^{/y}$: Paired t-test: t = -6.772, p < 0.001). This enhanced discrimination in *Fmr1*$^{-/y}$ mice was further supported
by low-salience reports that showed lower low-salience lick rates for 14 μm than 12 μm stimuli (Fig. S5A-
right; Paired t-test: t = 2.394, p = 0.044), whereas WT mice responded similarly (Fig. S5A-left; Paired t-test:
t = 1.589, p = 0.173), validating that improved discrimination was not driven by higher miss rates for 12
μm stimuli.

Together, these results demonstrate that *Fmr1*$^{-/y}$ mice present enhanced tactile fine-discrimination for
low-salience stimuli.

***Fmr1*$^{-/y}$ mice exhibit intact salience categorization.**

In addition to assessing stimulus discrimination, our task allowed us to evaluate whether *Fmr1*^{-/y} and WT
mice form the same categories of low- and high-salience stimuli. Using the psychometric curves of the
animals' high-salience responses (Fig. 3A), we measured the categorization bias and precision. No
differences were observed between *Fmr1*^{-/y} mice and their WT littermates regarding their categorization
bias (Fig. 3F; T-test: t=0.432, p=0.673) or precision (slope, Fig. 3G; T-test: t=-0.283, p=0.782), showing
comparable salience categorization between groups. These findings were further confirmed by similar
overall task sensitivity (d', Fig. 3H; T-test: t=0.571, p=0.578). Furthermore, *Fmr1*^{-/y} mice showed similar
rates of correct (Fig. S5B-C; High: T-test: t=-0.160, p=0.875; Low: T-test: t=1.389, p=0.188) and incorrect
choices (Fig. S5D-E; High: T-test: t=0.421, p=0.681; Low: T-test: t=-0.709, p=0.491), as well as comparable
overall decision strategy (Fig. S5F, criterion c; T-test: t=0.639, p=0.533) compared to their WT littermates.
In conclusion, these results demonstrate similar categorization of high- and low-salience stimuli in *Fmr1*⁻
^{/y} and WT mice.

***Fmr1*^{-/y} mice show reduced impact of categorization on stimulus discrimination.**

Higher-order processes such as categorization can amplify perceptual differences between stimuli
belonging to different categories (Goldstone, 1994). Alterations in this behavioral process have been
reported in autistic individuals, who, despite exhibiting intact visual categorization (in line with our
findings in the tactile domain, Fig. 3F-H), show a reduced influence of categorization on perceptual
discrimination that leads to diminished facilitation for stimuli belonging to different categories (Soulières
et al., 2007). To examine whether salience categorization differentially impacts tactile discrimination in
WT and *Fmr1*^{-/y} mice, we compared the discrimination accuracy for stimulus pairs that differ by 2 µm and
either span across different salience categories (across-category, i.e., 18 µm and 20 µm stimuli) or fall
within the same category. Discrimination accuracy was calculated as the difference in rightward choices
between each pair of stimuli.

In WT mice, discrimination accuracy was higher for across-category stimulus pairs than for both within-
category low- (Fig. 4A-left; Paired t-test, t = 4.452, p=0.007) and high-salience pairs (Fig. 4B-left; Paired t-
test, t = 5.453, p = 0.003), indicating a robust facilitatory effect of categorization. *Fmr1*^{-/y} mice, in contrast,
did not show this facilitation for low-salience stimuli, exhibiting a trend with medium effect size toward
improved discrimination for across-category versus within-category low-salience pairs (Fig. 4A-right;
Wilcoxon signed-rank: stat = 6.000, p = 0.055; Paired Hedges' g = 0.640). For high-salience stimuli,
however, *Fmr1*^{-/y} mice performed similarly to WT, showing enhanced discrimination for across-category
pairs relative to within-category high-salience comparisons (Fig. 4B-right; Paired t-test, t = 3.496, p =
0.008).

To directly compare WT and *Fmr1*^{-/y} mice, we quantified the categorical facilitation effect as the difference
in discrimination accuracy between across- and within-category (high and low-salience) stimulus pairs.
Fmr1^{-/y} mice showed a trend with a strong effect size toward reduced facilitation (T test: t = 1.81, p =
0.095; Hedge's g = 0.86), which became significant after exclusion of outliers (Fig. 4C; T test: t = 4.57, p =
0.001; Hedge's g = 1.94).

Together, these findings suggest that reduced categorization influence in *Fmr1*^{-/y} mice diminishes
discrimination facilitation for stimuli belonging to different categories, revealing a genotype-specific
alteration in how categorical information modulates tactile perception.

High cognitive load conditions reveal salience-dependent attention deficits in *Fmr1*^{-/y} mice.

Our results during perceptual learning revealed comparable, salience-dependent attention in WT and
Fmr1$^{-/y}$ mice (Fig. S3D-E). During the tactile discrimination/categorization phase of our task, animals were
exposed to a high cognitive load, as they needed to discriminate and categorize eight stimulus intensities
to obtain a water reward. To assess whether *Fmr1*$^{-/y}$ mice exhibit attentional deficits in these high
cognitive load conditions, we analyzed the rate of Miss trials as a proxy for attention.

While miss rates for high-salience stimuli did not differ between genotypes (Fig. 5A; all genotype x
amplitude interactions p > 0.296), *Fmr1*$^{-/y}$ mice showed elevated miss rates for low-salience stimuli during
categorization (Fig. 5B). A mixed-effects linear model revealed that *Fmr1*$^{-/y}$ mice had significantly higher
miss rates than WT controls at the reference amplitude of 12 μm (β = 0.067, SE = 0.024, z = 2.77, p = 0.006,
95% CI [0.019, 0.114]). The genotype differences at 14 μm (β = −0.015, SE = 0.021, z = −0.70, p = 0.483)
and 16 μm (β = −0.029, SE = 0.021, z = −1.37, p = 0.171) did not differ significantly from the effect observed
at 12 μm. However, a significant genotype × amplitude interaction was observed at the highest amplitude
(18 μm; β = −0.045, SE = 0.021, z = −2.16, p = 0.031, 95% CI [−0.087, −0.004]), indicating that genotype
differences diminished the highest stimulus intensity of the low-salience range. Post-hoc comparisons
further indicated a trend for increased misses at 12 μm with a large effect size (T-test: t = -2.8437, p =
0.058, Hedge's g = 1.23), but showed no significant differences for the other low-salience stimuli (p >
0.21). Notably, during the training period—when cognitive load was lower—miss rates were comparable
across genotypes (Fig. S3D-E).

Together, these results reveal a salience-specific attentional deficit in *Fmr1*$^{-/y}$ mice under high cognitive
load, with low-salience stimuli being particularly vulnerable.

ered *Fmr1*$^{-/y}$ mice show preserved choice perseveration but disrupted sensory history integration during
tactile categorization.

We next examined how trial history—including prior stimulus, choice, and outcome—shapes current
decisions during tactile categorization. To do so, we used generalized linear models (GLMs) with a
binomial link function to predict high-salience licks (right-lick choices) based on the current stimulus, trial
history, genotype, and their interactions. We first fit a main-effects model including the current stimulus,
previous stimulus, previous outcome, previous choice, and genotype, and then extended this model to
assess genotype-specific modulation of history effects through interactions between genotype and
previous choice, outcome, and stimulus (Fig. 6a).

As expected, the current stimulus was the dominant driver of choice in both genotypes (Fig. 6A-B; β =
1.94, SE = 0.15, z = 12.96, p < 0.001, 95% CI [1.64, 2.23]). However, previous choice also strongly influenced
behavior (Fig. 6A). Both genotypes exhibited perseveration, as psychometric curves conditioned on the
previous choice were systematically shifted following both prior right and left choices, indicating a robust
tendency to repeat prior responses (Fig. 6A-C; β = 1.16, SE = 0.10, z = 11.85, p < 0.001, 95% CI [0.97, 1.36]).
In contrast to the increased choice consistency bias observed during training (Fig. 2C-E), WT and *Fmr1*$^{-/y}$
mice exhibited comparable levels of perseveration during categorization (genotype x previous choice
interaction; β = 0.27, SE = 0.19, z = 1.47, p = 0.142, 95% CI [-0.09, 0.64]).

In contrast, previous outcome (reward or timeout) had a negligible effect on current choice in either
genotype (Fig. 6A, D; β = 0.09, SE = 0.10, z = 0.89, p = 0.373, 95% CI [-0.11, 0.28]). Interestingly, the
amplitude of the previous stimulus influenced decision-making (Fig. 6A; β = -0.12, SE = 0.04, z = -2.82, p =
0.005, 95% CI [-0.20, -0.04]) in a genotype-specific manner (Fig. 6A, E; genotype x previous stimulus
interaction, β = -0.23, SE = 0.06, z = -4.03, p < 0.001, 95% CI [-0.34, -0.12]). In WT mice, larger-amplitude

stimuli on the preceding trial slightly reduced the probability of choosing right on the subsequent trial,
whereas lower-amplitude stimuli increased it (Fig. 6E). In contrast, choices of $Fmr1^{-/y}$ mice were not
modulated by the amplitude of the previous stimulus.

Together, these results show that tactile decisions in both genotypes are dominated by current sensory
evidence and strongly biased by prior choices. While choice perseveration is preserved in $Fmr1^{-/y}$ mice,
integration of recent sensory history is selectively disrupted, revealing a genotype-specific deficit in how
past sensory information informs ongoing decision-making.

Discussion

Here, we developed a decision-making task based on psychophysics to dissociate stimulus-driven from
cognitively-modulated tactile responses in the $Fmr1^{-/y}$ mouse model of autism. Our findings reveal
salience-dependent cognitive alterations that shape sensory performance. During perceptual learning,
$Fmr1^{-/y}$ mice exhibited an increased choice consistency bias during low-salience trials, which contributed
to reduced task performance. During discrimination, $Fmr1^{-/y}$ mice displayed an enhanced tactile sensitivity
under low-salience conditions alongside decreased facilitation of across-category discrimination. Despite
this increased sensitivity, these mice showed low salience-specific attentional impairments under high
cognitive load, as well as faster updating of their world model with a decreased impact of sensory history
on their current choice. Our findings highlight the interplay between sensory and cognitive alterations in
autism, emphasizing the importance of cognitive context in interpreting sensory phenotypes, and
advocating for a shift beyond traditional sensory–cognitive dichotomies to better understand autism-
related phenotypes.

Learning alterations in autism and reliance on priors
Learning disability and intellectual disability are the second and third most common co-occurring
neurodivergencies of autism, observed in 23.5% and 21.7% of autistic individuals, respectively
(Khachadourian et al., 2023). Notably, even in the absence of these co-occurring conditions, autistic
individuals exhibit differences in learning processes and altered neural mechanisms of learning (Mercado
et al., 2020). In our study, we observed reduced performance in salience-based perceptual learning,
shaped by a higher consistency bias in choice selection in $Fmr1^{-/y}$ mice. These results align with findings in
autistic individuals showing a stronger influence of previous choices on current perceptual decisions
(Feigin et al., 2021). Moreover, decreased flexibility in perceptual decision-making is consistent with
evidence of altered auditory and visual learning in autism (Harris et al., 2015; Alispahic et al., 2022),
decreased performance under volatile conditions (Goris et al., 2021), as well as increased perseveration
and diminished sensitivity to feedback (Crawley et al., 2020). Perceptual category learning was also found
to be slower in autistic individuals, potentially due to the use of strategies that prioritize response guessing
over rule application (Bott et al., 2006; Soulières et al., 2011). Impairments in visual, tactile, and audio-
visual perceptual learning have also been shown in $Fmr1^{-/y}$ mice (Arnett et al., 2014b; Goel et al., 2018;
Mol et al., 2024).

Our findings extend this body of work by identifying a salience-dependent shift in reliance on priors in
autism, with lower-salience stimuli more strongly evoking dependence on previous choices and impacting
perceptual learning.

Increased sensory discrimination in autism
Several theories of autism, including the *weak coherence* (Frith, 2003), *enhanced perceptual functioning*
(Mottron et al., 2006), and *reduced generalization theories* (Plaisted, 2001), support the notion of a

superior perception of low-level information in autism. Indeed, a body of work has confirmed superior
pure tone discrimination (Bonnel et al., 2003; O'Riordan and Passetti, 2006) and discrimination learning
of highly confusable patterns (Plaisted et al., 1998b). Tactile discrimination in autism has been studied
using diverse experimental protocols that target different tactile features, which has often led to
contradictory results due to methodological variability and participant heterogeneity (reviewed in (Zetler
et al., 2019)). Similarly, work on roughness discrimination in rodent models (Orefice et al., 2016; Ahmadi
et al., 2023) and human studies (O'Riordan and Passetti, 2006) has shown discrepancies.

To address translational challenges between animal models and clinical research, our approach employed
vibrotactile stimuli closely aligned with those used in human studies. Consistent with findings in autistic
individuals (He et al., 2021b), $Fmr1^{-/y}$ mice showed intact amplitude discrimination for high-salience
vibrotactile stimuli. Clinical studies have typically employed stimuli with more than 10-fold higher
amplitudes compared to stimulus detection thresholds (He et al., 2021b; Asaridou et al., 2022). In
contrast, our investigation focused on suprathreshold stimuli with much lower intensities (1.5-fold higher
above threshold amplitude, 4-fold higher frequency). By employ such stimuli, our study uncovered
enhanced discrimination of low-salience stimuli in $Fmr1^{-/y}$ mice, an aspect that remains underexplored in
autistic individuals. Increased "local" processing could account for enhanced discrimination performance
(Frith, 2003). However, in our study, $Fmr1^{-/y}$ mice exhibited enhanced discrimination only under specific
stimulus salience conditions, arguing against a global increase in sensory processing precision. Future
clinical studies assessing the discrimination of lower-level stimuli are necessary to explore whether
enhanced low-salience discrimination also characterizes autistic individuals.

Intact categorization but decreased facilitation of across-category discrimination in autism
Most perceptual discrimination is influenced by categorization (Goldstone, 1994; Beauny et al., 2020;
Micher et al., 2024). As mentioned by Goldstone, "a clear distinction between sensory and cognitive
processes is not tenable" (Goldstone, 1994). The reduced generalization hypothesis of autism postulates
that, apart from superior performance on a difficult discrimination task, autistic individuals will have
inferior performance in stimulus categorization (Plaisted, 2001). However, studies on rule-based and
prototype categorization have yielded inconclusive results, reporting both intact (Klinger and Dawson,
2001; Molesworth et al., 2005), slower (Soulières et al., 2011), reduced (Froehlich et al., 2012; Gastgeb
and Strauss, 2012) and enhanced performance (Bonnel et al., 2003) in categorization tasks in autistic
individuals.

We assessed for the first time categorization and its impact on discrimination of vibrotactile stimuli in a
mouse model of autism. Our results revealed intact tactile categorization but reduced facilitation of
across-category discrimination in $Fmr1^{-/y}$ mice. Although no clinical studies to date have examined this
question in the tactile domain, our findings are consistent with previous research in the visual modality.
These studies suggest that enhanced discrimination in autistic individuals may stem from reduced
generalization (Plaisted et al., 1998b; Plaisted, 2001), and further indicate that discrimination processes
may operate with increased independence from the influence of categorization (Soulières et al., 2007).
These findings suggest a bidirectional interaction between sensory and cognitive alterations in autism,
indicating that not only atypical sensory perception can impact cognition (Haigh, 2018), but also that
cognitive differences can, in turn, shape sensory processing.

Attentional alterations in autism

Although attention deficits are not considered a general characteristic of autism (Grubb et al., 2013),
reduced attention in the presence of salient distractors (Venker et al., 2021) and weaker accuracy in
executive attention (Ridderinkhof et al., 2018) have been reported in autistic individuals. Cognitive load is
known to modulate attentional capacity (Lavie and Dalton, 2014; Murphy et al., 2016), and prior work has
shown that high cognitive load alters selective attention and distractor filtering (Head and Helton, 2014;
Lavie and Dalton, 2014; Brockhoff et al., 2022). However, less is known about attention to task-relevant
stimuli in these conditions. Our results support the view that attention deficits emerge under conditions
of high cognitive load in autism. Miss rates increased in our task when $Fmr^{1-/y}$ mice were required to
discriminate and categorize eight stimulus intensities within a single session for water reward, but not
during training sessions with only two stimulus intensities.

Notably, cognitive load appears to disproportionately affect attention to low-salience stimuli in $Fmr1^{-/y}$
mice, suggesting a disruption in late-stage attention processes, which take place after the stimulus is
perceived and are involved in top-down goal-directed behavior (Calvillo and Jackson, 2014). In contrast,
high-salience stimuli, which tend to capture attention automatically (Smallwood, 2013), may engage early
attentional processes and thus less affected by cognitive load. This context-dependent attentional
modulation aligns with previous findings in autistic individuals, showing stimulus-dependent auditory
attention deficits (Čeponiene et al., 2003). These findings suggest that attentional alterations in autism
may be conditionally expressed, particularly under tasks requiring increased cognitive resources.

Our results further dissociate attentional and tactile alterations in $Fmr1^{-/y}$ mice, as low-salience stimuli
were more accurately discriminated yet less attended to in $Fmr1^{-/y}$ mice. These findings align with studies
on linguistic processing in autistic children (Järvinen-Pasley et al., 2008; Ploog, 2010) and extend clinical
research demonstrating that tactile sensitivity alterations in autism are not linked to attentional
difficulties (He et al., 2021b).

Trial history integration during perceptual decision-making in autism
Sensory perception is shaped not only by stimulus characteristics but also by trial history. Our results show
that current stimulus intensity was the primary factor influencing the animal's choice, together with a
strong perseveration across genotypes, consistent with prior work on perceptual decision-making in mice
(Aguillon-Rodriguez et al., 2021). This effect remained intact in $Fmr1^{-/y}$ mice during tactile categorization.
Interestingly, previous work in rats and humans has revealed that stimulus history strongly influences
current perceptual choices under healthy conditions (Hachen et al., 2021). Our results confirmed these
results in control mice and further revealed that this sensory history effect was absent in $Fmr1^{-/y}$ mice.
These findings suggest poor sensory history integration and support models of autism proposing faster
world model updating, with new sensory information weighting higher than previous experience (Goris
et al., 2022).

Motivation and reward valuation during perceptual decision-making
Motivation and reward valuation are two crucial factors impacting performance during perceptual
decision-making. In this study, we minimized confounds from fatigue, satiety, or disengagement by
restricting analyses to sessions and epochs in which animals were actively performing the task. Under
these conditions, $Fmr1^{-/y}$ and WT mice showed indistinguishable task engagement, learning rates and
learning trajectories, which would not be expected if motivation, reward value, or reinforcement efficacy
were altered in our mouse model of autism.

During testing, only the sessions where the mice achieved high accuracy on well-learned training stimuli
were included in the analysis. Several additional observations further argue against altered reward

valuation as a primary driver of behavioral alterations in *Fmr1*$^{-/y}$ mice. We observed no genotype
differences in categorical response bias (criterion c), nor any effect of previous reward outcome on
subsequent choices, indicating that reward history did not differentially influence decision-making in
Fmr1$^{-/y}$ mice. In contrast, we found reduced across-category discrimination facilitation and diminished
influence of recent sensory history on choice, pointing to altered perceptual-decision processes rather
than global motivational deficits. Moreover, *Fmr1*$^{-/y}$ mice exhibited increased miss rates that emerged
selectively under high cognitive load and reduced sensory salience conditions. The salience- and context-
dependent nature of these effects, together with preserved engagement and reward sensitivity, supports
the conclusion that the observed behavioral differences reflect specific disruptions in how sensory
information is integrated and utilized during decision-making, rather than nonspecific changes in
motivation, fatigue, or stereotyped responding.

Conclusion and Future Perspectives
Numerous hypotheses have sought to provide a unified explanation for the diverse symptoms of autism.
Our findings support the view that altered cognitive processes shape perception in autism, influencing
how autistic individuals learn and engage with their environment. We demonstrate differences in choice
bias during perceptual learning and increased tactile discrimination in trained animals. These changes are
accompanied by reduced categorization influence, sensory history integration, and attention. Rather than
reflecting a global sensory deficit or enhancement, our results point to context-dependent alterations in
how sensory information is integrated, weighted, and used for decision-making.

Future work will be needed to identify the circuit mechanisms driving these effects, and to test whether
they reflect intrinsic changes in top-down cognitive networks or altered feedback between cognitive and
sensory circuits that reshape sensory representations. In addition, although the present study focused on
amplitude discrimination within the flutter range (40 Hz) to align with translational tactile paradigms,
extending this approach to a broader range of vibrotactile frequencies will be important to determine
whether sensory–cognitive alterations in autism generalize across stimuli that engage different
mechanoreceptor populations. Finally, while the present study focused on vibrotactile processing due to
its strong translational relevance, previous studies in the visual modality have reported similar cognitive
influences on perceptual decision-making, suggesting that these effects may generalize across sensory
systems. Future studies directly comparing modalities within the same experimental framework will be
important to confirm this. Elucidating how cognition modulates sensory perception may offer valuable
insights for clinical practice, help reconcile conflicting findings in the field, and inform the development of
more targeted, mechanism-based interventions.

Materials and Methods

Experimental design
To study tactile perception as well as attention and perceptual biases in autism, we developed a novel 2-
Alternative Choice task for the categorization and discrimination of flutter-range vibrotactile stimuli.
Throughout the text, we use terms that are preferred in the autistic community and are less stigmatizing
(Bottema-Beutel et al., 2021).

Ethical statement
All experimental procedures were performed in accordance with the EU directive 2010/63/EU and French
law following procedures approved by the Bordeaux Ethics Committee and Ministry for Higher Education

and Research. Mice were maintained in reversed light cycle under controlled conditions (temperature 22–
24 °C, humidity 40–60%, 12 h/12 h light/dark cycle, light on at 21:00) in a conventional animal facility with
ad libitum access to food and *ad libitum* access to water before the water restriction period. All
experiments were performed during the dark cycle, under red light.

Mice
Second-generation *Fmr1* knockout (*Fmr1$^{-/y}$*) and wild-type littermate mice 5-16 weeks old were used in
our study. Mice were maintained in a C57Bl/6 J background (Mientjes et al., 2006). Male wild-type
and *Fmr1$^{-/y}$* littermates were generated by crossing *Fmr1$^{+/-}$* females with *Fmr1$^{+/y}$* male mice from the
same production, and the resulting progeny used for our experiments was either *Fmr1$^{+/y}$* (wild type)
or *Fmr1$^{-/y}$* (KO). Mice were maintained in collective cages following weaning (2-4 litter males per cage).
Cages were balanced for genotype and supplemented with cotton nestlets and carton tubes.

The perceptual decision-making data were collected from 5 different cohorts of mice at different time-
points during their active phase of the day. Mice of both genotypes were littermates and represented in
each cohort. The number of mice is provided in the figure captions. The experimenter was blind to the
animals' genotypes throughout the experiment. The genotype of experimental animals was re-confirmed
post hoc by tail-PCR.

2-Alternative Choice task

Setup
The vibrotactile decision-making setup was positioned in an isolation cubicle to minimize interference
during the experiment. Mice were placed in a body tube and were head-fixed with their forepaws resting
on two steel bars (6 mm diameter, Thorlabs). The right bar was mounted to a Preloaded Piezo Actuator
(P-841.6, Physik Instrumente) equipped with a strain gauge feedback sensor and controlled (E-501, Physik
Instrumente) in a closed loop, as described before (Prsa et al., 2019; Semelidou et al., 2024). A 12.7 mm
stainless steel post (ThorLabs) was mounted on the actuator vertically and a 0.6 mm stainless steel rod
(ThorLabs) was clamped horizontally onto this post. The horizontal rod served as the contact bar on which
the animal rested its right forepaw. Water reward was delivered through either of the two metal feeding
needles (20G, 1,9mm tip, Agntho's AB), placed left and right of the mouse's mouth, each connected to a
lickport interface with a solenoid valve (Sanworks) equipped with a capacitive sensor
(https://github.com/poulet-lab/Bpod_CapacitivePortInterface). The perceptual decision-making setup
was controlled by Bpod (Sanworks) through scripts in Python (PyBpod,
https://pybpod.readthedocs.io/en/latest/). The lickport interface (Sanworks) was equipped with a
capacitive sensor (https://github.com/poulet-lab/Bpod_CapacitivePortInterface).

Habituation to head-fixation and water restriction
Mice (P40-P50) were handled using carton tubes and the cupping technique until they were comfortable
in the experimenter's hands, attested by eating while handled. Mice were gradually habituated to the
experimental setup and head fixation for 5 days. The third day of habituation, a water-restriction protocol
was implemented, where mice had access to liquid water in the setup and to a solid water supplement
(Hydrogel, BioServices) in their home cage. The water supplement was divided into small, individual
portions, and each mouse received its allotment after the daily training/testing session. Daily body weight
measurements were used to monitor hydration and ensure that all animals maintained stable body
weight. If necessary, supplemental water was adjusted to maintain animals within the approved weight
range. In total, the animals received 1.5-2 ml of water per day, which corresponds to 50-65% of their ad

libitum consumption, while ensuring that they did not lose more than 10% of their weight. Each mouse
received 6-8 g of Hydrogel (*ad libitum*) during the weekend. This water restriction protocol was
maintained throughout behavioral training and until the end of behavioral testing.

2-Alternative Choice task training and testing

Vibrotactile Stimuli
Stimuli were sinusoidal vibrations at 40 Hz with peak-to-peak displacements of 12 μm amplitude (low
salience) or 26 μm amplitude (high salience), well above the detection threshold of both groups (10 Hz,
4.46 μm for WT; 10Hz, 7.29 μm for *Fmr1*$^{-/y}$ mice) (Semelidou et al., 2024).
Animals were positioned in the setup to ensure stable and consistent forepaw contact with the rod
delivering the vibration. Pilot experiments with a sensor to monitor forepaw placement confirmed that
the mice did not remove their forepaws from the bar before stimulus delivery.
Analysis of trials were no response followed stimulus delivery (Miss trials) showed that the 12 μm stimulus
consistently elicited a higher proportion of missed responses compared to the 26 μm stimulus across
animals, indicating lower behavioral performance for the lower-amplitude stimulus. We therefore
referred to the 12 μm stimulus as "low salience" and the 26 μm stimulus as "high salience" to denote
relative differences in perceptual strength and attentional engagement within the suprathreshold range.

Habituated mice (8 weeks old) were trained to associate high- (26 μm) and low-salience (12 μm)
vibrotactile stimuli (pure sinusoid, 500 ms duration, 40 Hz frequency) with a water reward (8 μl) at the
right- or left-placed lickport of the setup, respectively. All trials consisted of stimulus delivery followed by
a 2 s response window during which the mice could lick to receive the reward. Inter-trial intervals were
variable (5-10 s). Training was subdivided in 4 phases: (a) automatic water delivery at the beginning of the
response window at the corresponding lickport (left for 12 μm stimuli, right for 26 μm stimuli). (b) Training
in blocks: lick-triggered water delivery during blocks of 20 trials with the same vibrotactile stimulus, of
either 12 μm or 26 μm amplitude. Licking at the wrong port resulted in 5 s timeout. (c) Training with
pseudorandomly delivered high- and low-salience trials: lick-triggered water delivery following
pseudorandom delivery of 12 μm or 26 μm amplitude stimuli. Licking at the wrong port resulted in 5 s
timeout.

All sessions consisted of 200-300 trials with a 1:1 ratio of high and low-salience stimuli. During training,
this ratio was modified when an animal showed consistent bias for one of the two lickports. Pilot
experiments with an extra sensor to monitor forepaw placement confirmed that the mice did not remove
their forepaws from the bar before stimulus delivery. To complete training in blocks, mice needed to
perform with 70% correct choices and 30% incorrect choices for both stimulus amplitudes. To complete
training with pseudorandomly delivered high- and low-salience trials, mice needed to reach the criterion
of more than 70% correct choices and less than 30% incorrect choices as an average for 3 consecutive
days. All mice that fulfilled this criterion were tested for the categorization/discrimination of vibrotactile
stimuli. During testing, stimuli (pure sinusoid, 500 ms duration, 40 Hz frequency) were delivered in a
pseudorandom manner with a 50% High-salience: 50% Low-salience ratio. Amplitudes varied on a range
between 12 μm and 26 μm. Stimuli of 20, 22, 24 and 26 μm were considered high-salience stimuli and
rewarded at the right lickport while stimuli of 12, 14, 16 and 18 μm were considered as low-salience
stimuli and rewarded at the left lickport.

600
2-Alternative choice vibrotactile task analysis
All analysis was performed with custom-made Python scripts that can be available upon request. Behavior
was quantified based on the lick events and three main outcomes were measured for each stimulus
salience: Correct choice rate (number of correct licks divided by the total number of high- or low-salience
trials), Incorrect choice rate (number of wrong licks divided by the total number of high- or low-salience

trials), and Missed stimuli rate (number of trials in which the animal did not lick, divided by the total
number of high- or low-salience trials). Sessions in which animals disengaged were analyzed only during
epochs in which the animal was actively performing the task. Training duration was calculated based on
the total number of days each animal passed in training in blocks and with pseudorandom stimulus
delivery.

For testing, only sessions with more than 70% correct choices for the training stimuli (12 μm and 26 μm)
were analyzed. Psychometric curves were fitted on the rightwards lick rate for each stimulus amplitude
using a general linear model. An average of 84 repetitions for each amplitude was used to calculate
rightward lick rates.

Based on the signal detection theory (Detection Theory,(Hautus, M.J., Macmillan, N.A., & Creelman,
2021)), sensitivity d' was calculated as:
d' = z(lick right / high amplitude stimulus trials) - z(lick right / low amplitude stimulus trials)
620

Sensitivity d' for high- and low-salience stimuli was calculated based on the Correct and Incorrect choice
rate for high- and low-salience stimuli, respectively.
The strategy of the animals was assessed through their criterion, which was calculated as:
$c = -1/2 \times [z(\textit{lick right / high amplitude stimulus trials}) + z(\textit{lick right / low amplitude stimulus trials})]$

Categorization bias was calculated as the inflection point (i.e., the midpoint parameter μ) of the fitted
logistic psychometric function and categorization precision as the slope of the curve.
Prior choice impact was calculated separately for each type of the current trial (high- or low-salience). For
low-salience trials, the prior choice impact was calculated as the rate of correct low and incorrect-low
trials, divided by the rate of all correct and all incorrect trials. Similarly, for high-salience trials, the prior
choice impact was calculated as the rate of correct high and incorrect-high trials, divided by the rate of all
correct and all incorrect trials.

Learning trajectory analysis
Learning curves were quantified from the training phase with mixed high- and low-salience trials by
calculating, for each mouse, the proportion of correct choices on each training day. For group-level
comparisons, performance was plotted across days for WT and *Fmr1*$^{-/y}$ mice, and mean learning
trajectories with 95% confidence intervals were computed for each genotype.

To statistically compare learning trajectories across genotypes, we used a linear mixed-effects model with
fixed effects for Genotype, Day, and their interaction, and a random intercept and random slope for each
animal. This model tested whether performance differed between genotypes across days and whether
the rate of learning (performance change per day) differed between WT and *Fmr1*$^{-/y}$ mice.

To quantify individual learning rates, we computed the slope of performance as a function of training day
for each animal. For each mouse, performance values (proportion of correct choices) were regressed
against each Day using a simple linear regression model:
644                  $\text{Performance}_{\text{animal}} = \beta_0 + \beta_1 * \text{Day}$

The slope (β_1) provides an estimate of the rate of improvement per training day.
Intermediate-stage slopes (middle 3 days of each mouse's training) were calculated in the same manner.

Categorization bias was calculated based on the psychometric curves, as the stimulus amplitude at the
inflection point of the sigmoid fitting curve. Categorization precision was calculated based on the slope of
the psychometric curve.

Delta discrimination accuracy was computed as the difference in the rate of high-salience (rightward) licks
between pairs of stimuli that differed by 2 μm. For analyses of low-salience and high-salience
discrimination, delta discrimination accuracy values were calculated for each within-category stimulus
pair and then averaged within the low-salience or high-salience category, respectively.

Statistics
All values are presented as mean ± s.e.m. Box plots show the median, interquartile, range, mean and
individual values. The numbers of animals used are indicated in the figure legends. Sample size was
determined based on previous studies using similar behavioral paradigms. For experiments assessing
tactile responses of trained animals, only animals that reached the training criterion were tested. For the
discrimination & categorization task, all the testing sessions where the animals showed less than 70%
correct responses to the training stimuli (12μm and 26μm) were excluded from tactile
discrimination/categorization analysis. No samples or data points were excluded from the analysis.
All statistical analysis was done using Python (SciPy, Pingouin). All datasets were tested for normality using
Shapiro–Wilk tests, and potential outliers were identified using a median absolute deviation–based
modified Z-score ($|Z| > 3.5$). When an outlier was detected, statistical analyses were performed both
including and excluding the outlier. In such cases, results from both analyses are reported when the
inclusion or exclusion of the outlier altered the statistical outcome, to ensure transparency and robustness
of the findings. For normally distributed data, two-tailed paired t-tests were used for within-subject
comparisons, and Welch's independent-samples t-tests were used for between-group comparisons;
otherwise, the Mann–Whitney U test (between groups) or the Wilcoxon signed-rank test (within subjects)
was applied. Bonferroni correction was applied where multiple comparisons were performed. Effect sizes
were calculated using Hedge's g for between-genotype comparisons and paired Hedge's g for within-
genotype comparisons, particularly when statistical analyses showed trend-level differences ($p ≈ 0.05$–
0.1), to provide a measure of the magnitude of the effect independent of sample size. Linear mixed-effects
models were used to assess statistical differences in learning trajectories, psychometric curves, and miss
rates.

Statistical modeling during categorization taking into account the trial history.
Generalized linear models (GLMs) with a binomial link function were used to quantify the effects of
current stimulus, previous trial history, genotype, and their interactions on the probability of choosing the
right port. Trials without a defined history and Miss trials were excluded. Stimulus amplitudes were z-
scored per mouse.
Two models were fit:
1. Main effects GLM – included current stimulus, previous stimulus, previous outcome, previous choice,
and genotype.
2. Interaction GLM – extended the main effects model by including genotype × previous outcome,
genotype × previous choice, and genotype × previous stimulus interactions.
Cluster-robust standard errors were calculated to account for repeated measurements within each
mouse.
Marginal effects and psychometric curves
Predicted probabilities were computed from the interaction GLM to visualize the effect of stimulus
amplitude, previous choice, and previous outcome on behavior, separately for WT and KO mice.

Data availability

The raw data of behavioral experiments generated in this study have been deposited on the figshare
database: 10.6084/m9.figshare.29459771. Source data for each figure and the Python code used for all
analyses are provided with the paper.

Figure 1. Perceptual learning performance in a forepaw-based decision-making task. For panels **B, C, D, E, F, G, H, I, J, K:** n=9 WT, 11 *Fmr1⁻/y* male mice. **A,** Left: schema showing the behavioral setup. Right: Trail protocol and behavioral outcomes depending on the type of trial and the animal's response. **B,** Total number of days spent in training for WT and *Fmr1⁻/y* mice. **C,** Total number of days spent in training until the criterion was met for high-salience stimuli for WT and *Fmr1⁻/y* mice. **D,** Total number of days spent in training until the criterion was met for low-salience stimuli for WT and *Fmr1⁻/y* mice. **E,** Sensitivity d' throughout the training period for WT and *Fmr1⁻/y* mice. **F,** Correct choice rate for both high- and low-salience trails throughout the training period for WT and *Fmr1⁻/y* mice. **G,** Incorrect choice rate for both high- and low-salience trails throughout the training period for WT and *Fmr1⁻/y* mice. **H,** Incorrect choice rate for low-salience trials throughout the training period for WT and *Fmr1⁻/y* mice. **I,** Incorrect choice rate for high-salience trails throughout the training period for WT and *Fmr1⁻/y* mice. P values were computed using two-sided t-test for panels **B, E, F, G, I,** ; Mann-Whitney test for panels **C, D, H** ; *P < 0.05 or n.s, not significant.

Figure S1. Perceptual learning duration for the different training phases and correct choice rates. n=9 WT, 11 *Fmr1*[-/y] male mice for panels **C-F**. **A,** (Left) Example trace of 100 consecutive trials from an early (mixed) training session, showing the rates of correct (top) and incorrect (bottom) choices. (Right) Licking patterns from the same session for high-salience (top) and low-salience (bottom) trials. **B,** Same as in (A), but from a late training session of the same animal. **C,** Total number of days spent in the training phase where stimulus delivery is done in blocks of high- or low-salience stimuli. **D,** Total number of days spent in training with high- and low-salience stimuli are delivered in a pseudorandom manner. **E,** Correct choice rate for high- salience trails throughout the training period for WT and *Fmr1*[-/y] mice. **F,** Correct choice rate for low- salience trails throughout the training period for WT and *Fmr1*[-/y] mice. P values were computed using two-sided t-test for panels **C, D, E, F** ; n.s, not significant.

Figure S2. Perceptual learning trajectories during training with mixed low- and high-salience trials. n=7 WT, 9 *Fmr1⁻ʸ* male mice for panels. **A,** (Top) Average learning trajectories for WT and *Fmr1⁻ʸ* mice that completed the training phase, during mixed-trial training, calculated from the correct-choice rate for each training session (day) until each mouse reached the learning criterion. Shaded areas indicate 95% confidence intervals. (Bottom) Heat map of individual performance across mixed high- and low-salience sessions, showing the correct-choice rate for each session up to each mouse's learning criterion. **B,** Slope of the learning trajectory for each mouse, calculated using a linear regression model. **C,** Performance (correct choice rate) on the first day of the mixed-trial training phase. **D,** Intra-individual variability in performance across training days, assessed as the standard deviation of the correct choice rate. **E,** Slope of the learning trajectory during the intermediate learning stage, defined as the middle 3 days of training for each animal. **F,** Performance (correct choice rate) during the intermediate learning stage. P values were computed using linear mixed-effects model for panel **A**, Mann-Whitney test for panel **B**, and two-sided t-test for panels **C, D, E, F** ; n.s, not significant.

Figure 2. Overall strategy and impact of prior choice during perceptual learning. For panels **A, C, D, E, F:** n=9 WT, 11 *Fmr1⁻/ʸ* male mice. **A,** Criterion depicting the licking strategy of the animals. **B,** Schema showing an example of how high impact of prior choice on the current trial affects the response during a high-salience trial (top) or low-salience trial (bottom). **C,** Proportion of incorrect responses in low-salience trials immediately following a correctly rewarded high-salience trial. **D,** Proportion of incorrect responses in high-salience trials immediately following a correctly rewarded low-salience trial. **E,** Proportion of correct responses in low-salience trials immediately following a correctly rewarded low-salience trial and incorrect responses in low-salience trials immediately following a correctly rewarded high-salience trial. Rates are corrected over the total number of correct and incorrect choices in low-salience trials. **F,** Proportion of correct responses in high-salience trials immediately following a correctly rewarded high-salience trial and incorrect responses in high-salience trials immediately following a correctly rewarded low-salience trial. Rates are corrected over the total number of correct and incorrect choices in high-salience trials. P values were computed using Mann-Whitney test for panel **A,;** two-sided t-test for panels **C, D, E, F,;** *P < 0.05 or n.s, not significant.

Figure S3. Prior strength and attention during perceptual learning. n=9 WT, 11 *Fmr1^-/y* male mice. **A,** Strength of the prior build for high-salience trials, calculated as the proportion of correct high-salience and incorrect low-salience responses following a correct high-salience response. Rates are corrected over the rate of overall correct high-salience and incorrect low-salience responses. **B,** Strength of the prior build for low-salience trials, calculated as the proportion of correct low-salience and incorrect high-salience responses following a correct low-salience response. Rates are corrected over the rate of overall correct low-salience and incorrect high-salience responses. **C,** Within-genotype comparisons of the proportion of missed high- and low-salience trials. **D,** Proportion of missed high-salience trials. **E,** Proportion of missed low-salience trials. P values were computed using two-sided t-test for panel **A,** ; Mann-Whitney test for panels **B, D, E** ; Wilcoxon signed-rank test for panel **C-left;** Paired t-test for panel **C-right**, ; **P < 0.01, or n.s, not significant.

Figure 3. Tactile discrimination and categorization. n=6 WT, 9 *Fmr1⁻/y* male mice. **A,** Psychometric curves for WT and *Fmr1⁻/y* mice generated based on the high-salience lick rate (rate of rightward licks) across 8 different amplitudes. Stimuli between 12–18 µm were designated as low-salience and rewarded at the left lickport, while stimuli between 20–26 µm were designated as high-salience and rewarded at the right lickport. Each amplitude was presented an average of 84 times. **B,** Comparison of high-salience (rightwards) responses for high-salience stimuli of 26 and 26 µm. **C,** Comparison of high-salience (rightwards) responses for high-salience stimuli of 26 and 22 µm. **D,** Comparison of high-salience (rightwards) responses for low-salience stimuli of 12 and 14 µm. **E,** Comparison of low-salience (leftwards) responses for low-salience stimuli of 12 and 14 µm. **F,** Categorization bias calculated based on the psychometric curves. **G**, Categorization precision computed based on the slope of the psychometric curves. **H,** Sensitivity d' of the responses in all stimulus amplitudes. P values were computed using Mixed Linear Model Regression for panel **A,**; two-sided paired t-test for panels **B-right, C-right, D, E,**; two-sided t-test for panels **F, G, H,**; Wilcoxon signed-rank test for panels **B-left, C-left,**; Bonferroni correction was applied for panels **B, C, D,** **P < 0.01, *P < 0.05, or n.s, not significant.

Figure S4. Perceptual learning performance during the last three days of training, for mice that were tested in

tactile discrimination. n=6 WT, 9 *Fmr1^{-/y}* male mice. **A,** Sensitivity d' throughout the last three days of the training period for WT

and *Fmr1^{-/y}* mice. **B,** Proportion of correct choices for high-salience trails throughout the last three days of the training period. **C,**

Proportion of correct choices for low-salience trails throughout the last three days of the training period. **D,** Proportion of incorrect

choices for high-salience trails throughout the last three days of the training period. **E,** Proportion of incorrect choices for low-salience

trails throughout the last three days of the training period. **F,** Criterion depicting the licking strategy of the animals. **G,** Proportion of

correct responses in high-salience trials immediately following a correctly rewarded high-salience trial. **H,** Proportion of correct

responses in low-salience trials immediately following a correctly rewarded low-salience trial. **I,** Proportion of incorrect responses in

high-salience trials immediately following a correctly rewarded low-salience trial. **J,** Proportion of incorrect responses in low-salience

trials immediately following a correctly rewarded high-salience trial. **K,** Proportion of correct responses in high-salience trials

immediately following a correctly rewarded high-salience trial and incorrect responses in high-salience trials immediately following a

correctly rewarded low-salience trial. Rates are corrected over the total number of correct and incorrect choices in high-salience trials.

L, Proportion of correct responses in low-salience trials immediately following a correctly rewarded low-salience trial and incorrect

responses in low-salience trials immediately following a correctly rewarded high-salience trial. Rates are corrected over the total

number of correct and incorrect choices in low-salience trials. **M,** Proportion of missed high-salience trials. **N,** Proportion of missed

low-salience trials. P values were computed using two-sided t-test for panels **A, B, C, D, E, G, H, I, J, K, L, N** ; Mann-Whitney test for

panel **F**, **M,**; n.s, not significant.

Figure S5. Sensitivity, correct, and incorrect choices during categorization of high- and low-salience stimuli. n=6 WT, 9 *Fmr1[-/y]* male mice. **A,** Sensitivity d' for high-salience stimuli during categorization. **B,** Sensitivity d' for low-salience stimuli during categorization. **C,** Correct responses for high-salience stimuli during categorization. **D,** Correct responses for low-salience stimuli during categorization. **E,** Incorrect responses for high-salience stimuli during categorization. **F,** Incorrect responses for low-salience stimuli during categorization. P values were computed using two-sided paired t-test for panel **A,**; two-sided t-test for panels **B, C, D, E, F,;** *P < 0.05, or n.s, not significant.

Figure 4. Impact of categorization on across-categories and within-category discrimination. n=6 WT, 9 *Fmr1*[-/y] male
mice. Delta discrimination accuracy, calculated as the difference in the rate of high-salience licks across stimulus pairs. **(A)** Delta
discrimination accuracy between across-category stimuli (18 μm vs 20 μm) and within-category low-salience stimulus pairs, averaged
as "low salience." **B,** Same as (**A,**) but for within-category high-salience stimulus pairs. **C,** Delta discrimination accuracy across all
stimulus pairs within the low-salience category. **D,** Same as (**C,**) but across all stimulus pairs within the high-salience category. P
values were computed using two-sided paired t-tests for panels **A-left**, **B**; Wilcoxon signed-rank test for panel **A-right**, and two-sided
t-test for panel **C,;** **P < 0.01, or n.s, not significant.

795 **Figure 5. Attention in perceptual decision-making under high cognitive load.** n=6 WT, 9 *Fmr1*[-/y] male mice. **A,** Proportion of
796 missed trials for high-salience stimuli. **B**, Proportion of missed trials for low-salience stimuli. P-values were obtained using mixed-
797 effects linear models accounting for repeated measures across stimulus amplitudes and animals for panels **A, B**,; Post-hoc two-
798 sided t-test for panel **B,-**12μm (p = 0.06); **P < 0.01, or n.s, not significant.

799
Figure 6. Trial history integration during tactile categorization. n=6 WT, 9 *Fmr1⁻/ʸ* male mice. **(A)** Generalized linear model
(GLM) coefficients from binomial regression predicting high-salience (right-lick) choices based on current stimulus amplitude,
previous stimulus amplitude, previous outcome, previous choice, genotype, and their interactions. Filled circles indicate coefficients
from the main-effects model; open circles indicate coefficients from the extended model including genotype × history interactions.
Error bars represent ±95% confidence intervals. Dark blue lines denote statistically significant effects. **(B)** Psychometric functions
showing the effect of the z-scored current stimulus amplitude for WT and *Fmr1⁻/ʸ* mice. **(C)** Psychometric curves conditioned on the
previous choice (left vs. right lick) for WT and *Fmr1⁻/ʸ* mice. **(D)** Psychometric curves conditioned on previous trial outcome (reward
vs. timeout). **(E)** Psychometric curves conditioned on the intensity of the previous stimulus (z-scored). P values were computed
using Mixed Linear Model Regression. ***P < 0.001, **P < 0.01.

Acknowledgments
We would like to thank the animal housing and genotyping facilities of INSERM U1215
Neurocenter Magendie. We thank Dr. James Poulet (Max-Delbrück-Centrum für Molekulare Medizin,
Berlin) for providing the capacitive sensors for the behavioral task. We would like to thank Dr. Melanie
Ginger for providing feedback on the manuscript. Mouse schemas were made using Biorender. This
project was funded by the Foundation for Medical Research Postdoc Fellowship (O.S.), INSERM (A.F.),
Marcel Dassault-Fondation FondaMental Award 2019 (A.F.), Simons Foundation Autism
Research Initiative (A.F.), GIS Autisme & TND (O.S.), 2024 NARSAD Young Investigator Grant (O.S.),
Fondation FondaMental (A.F.).

Author Information
Authors and Affiliations
University of Bordeaux, Neurocentre Magendie, INSERM U1215, 33000, Bordeaux, France
Ourania Semelidou, Mathilde Tortochot-Megne Fotso, Adinda Winderickx, Andreas Frick

Contributions
A.F. and O.S. conceived the project. O.S. designed the experiments. O.S., M.T-M.F. and A.W. performed
the experiments. O.S. and A.W. wrote the Python code for the behavioral experiments. O.S. analyzed
the data and wrote the necessary Python scripts. O.S. interpreted the data. O.S. prepared the figures.
O.S. wrote the manuscript, and A.F. and M.T-M.F. provided feedback.

Corresponding author
Correspondence to Andreas Frick and Ourania Semelidou.

Ethics declarations
Competing interests
The authors declare no competing interests.

Inclusion and diversity statement
We support inclusive, diverse, and equitable conduct of research. We tried to use inclusive language as
much as possible.

References

Abu-Dahab SMN, Skidmore ER, Holm MB, Rogers JC, Minshew NJ (2013) Motor and Tactile-Perceptual
Skill Differences Between Individuals with High-Functioning Autism and Typically Developing
Individuals Ages 5 – 21. J Autism Dev Disord 43:2241 Available at:
https://pmc.ncbi.nlm.nih.gov/articles/PMC3408783/ [Accessed April 16, 2025].

Aguillon-Rodriguez V et al. (2021) Standardized and reproducible measurement of decision-making in
mice. Elife 10 Available at: /pmc/articles/PMC8137147/ [Accessed November 23, 2022].

Ahmadi BBM, Afarinesh MR, Jafaripour L, Sheibani V (2023) Alteration in social interaction and tactile
discrimination of juvenile autistic-like rats following tactile stimulation and whisker deprivation.
Brain Behav 13.

Alispahic S, Pellicano E, Cutler A, Antoniou M (2022) Auditory perceptual learning in autistic adults.

Allen G, Courchesne E (2001) Attention function and dysfunction in autism. Front Biosci 6 Available at:
https://pubmed.ncbi.nlm.nih.gov/11171544/ [Accessed June 23, 2025].

American Psychiatric Association (2022) Diagnostic and statistical manual of mental disorders (5th ed.,
text rev.). American Psychiatric Association Publishing. Available at:
https://psychiatryonline.org/doi/book/10.1176/appi.books.9780890425787.

Amoruso L, Narzisi A, Pinzino M, Finisguerra A, Billeci L, Calderoni S, Fabbro F, Muratori F, Volzone A,
Urgesi C (2019) Contextual priors do not modulate action prediction in children with autism. Proc R
Soc B 286 Available at: https://royalsocietypublishing.org/doi/10.1098/rspb.2019.1319 [Accessed
April 23, 2023].

Arnett MT, Herman DH, McGee AW (2014a) Deficits in Tactile Learning in a Mouse Model of Fragile X
Syndrome. PLoS One 9 Available at: /pmc/articles/PMC4189789/ [Accessed May 9, 2024].

Arnett MT, Herman DH, McGee AW (2014b) Deficits in tactile learning in a mouse model of fragile X
syndrome. PLoS One 9 Available at: https://pubmed.ncbi.nlm.nih.gov/25296296/ [Accessed May
28, 2024].

Asaridou M, Wodka EL, Edden RAE, Mostofsky SH, Puts NAJ, He JL (2022) Could Sensory Differences Be a
Sex-Indifferent Biomarker of Autism? Early Investigation Comparing Tactile Sensitivity Between
Autistic Males and Females. J Autism Dev Disord:1–17 Available at:
https://link.springer.com/article/10.1007/s10803-022-05787-6 [Accessed December 11, 2022].

Balasco L, Pagani M, Pangrazzi L, Chelini G, Ciancone Chama AG, Shlosman E, Mattioni L, Galbusera A,
Iurilli G, Provenzano G, Gozzi A, Bozzi Y (2021) Abnormal Whisker-Dependent Behaviors and
Altered Cortico-Hippocampal Connectivity in Shank3b –/– Mice . Cereb Cortex:1–14.

Balasco L, Pagani M, Pangrazzi L, Chelini G, Viscido F, Chama AGC, Galbusera A, Provenzano G, Gozzi A,
Bozzi Y (2022) Somatosensory cortex hyperconnectivity and impaired whisker-dependent
responses in Cntnap2 -/- mice. Neurobiol Dis 169 Available at:
https://pubmed.ncbi.nlm.nih.gov/35483565/ [Accessed July 19, 2022].

Barbot A, Liu S, Kimchi R, Carrasco M (2018) Attention enhances apparent perceptual organization.
Psychon Bull Rev 25:1824 Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC5831484/
[Accessed May 20, 2025].

Beauny A, de Heering A, Muñoz Moldes S, Martin JR, de Beir A, Cleeremans A (2020) Unconscious
categorization of submillisecond complex images. PLoS One 15 Available at:
https://pubmed.ncbi.nlm.nih.gov/32785238/ [Accessed May 23, 2025].

Bonnel A, Mottron L, Peretz I, Trudel M, Gallun E, Bonnel AM (2003) Enhanced pitch sensitivity in
individuals with autism: A signal detection analysis. J Cogn Neurosci 15:226–235 Available at:
https://pubmed.ncbi.nlm.nih.gov/12676060/ [Accessed May 22, 2025].

Bott L, Brock J, Brockdorff N, Boucher J, Lamberts K (2006) Perceptual similarity in autism. Q J Exp
Psychol 59:1237–1254 Available at: https://pubmed.ncbi.nlm.nih.gov/16769623/ [Accessed April
26, 2025].

Bottema-Beutel K, Kapp SK, Lester JN, Sasson NJ, Hand BN (2021) Avoiding Ableist Language:
Suggestions for Autism Researchers. Autism adulthood challenges Manag 3:18–29 Available at:
https://pubmed.ncbi.nlm.nih.gov/36601265/ [Accessed July 17, 2024].

Bouvier B, Susini P, Marquis-Favre C, Misdariis N (2023) Revealing the stimulus-driven component of
attention through modulations of auditory salience by timbre attributes. Sci Rep 13:1–11 Available
at: https://www.nature.com/articles/s41598-023-33496-2 [Accessed April 30, 2025].

Bremner AJ, Spence C (2017) The Development of Tactile Perception. Adv Child Dev Behav 52:227–268.

Brockhoff L, Schindler S, Bruchmann M, Straube T (2022) Effects of perceptual and working memory load
on brain responses to task-irrelevant stimuli: Review and implications for future research. Neurosci
Biobehav Rev 135:104580 Available at:
https://www.sciencedirect.com/science/article/abs/pii/S0149763422000690 [Accessed May 23,
2025].

Calvillo DP, Jackson RE (2014) Animacy, perceptual load, and inattentional blindness. Psychon Bull Rev
21:670–675 Available at: https://link.springer.com/article/10.3758/s13423-013-0543-8 [Accessed
May 7, 2025].

Carati E, Angotti M, Pignataro V, Grossi E, Parmeggiani A (2024) Exploring sensory alterations and
repetitive behaviors in children with autism spectrum disorder from the perspective of artificial
neural networks. Res Dev Disabil 155 Available at: https://pubmed.ncbi.nlm.nih.gov/39577022/
[Accessed November 26, 2024].

Cascio CJ, Gu C, Schauder KB, Key AP, Yoder P (2015) Somatosensory event-related potentials and
association with tactile behavioral responsiveness patterns in children with ASD. Brain Topogr
28:895 Available at: /pmc/articles/PMC4601930/ [Accessed December 3, 2022].

Čeponiene R, Lepistö T, Shestakova A, Vanhala R, Alku P, Näätänen R, Yaguchi K (2003) Speech-sound-
selective auditory impairment in children with autism: They can perceive but do not attend. Proc
Natl Acad Sci U S A 100:5567–5572 Available at:
/doi/pdf/10.1073/pnas.0835631100?download=true [Accessed May 6, 2025].

Church B, Krauss MS, Lopata C, Toomey JA, Thomeer ML, Coutinho M V., Volker MA, Mercado E (2010)
Atypical categorization in children with high-functioning autism spectrum disorder. Psychon Bull
Rev 17:862–868.

Crawley D, Zhang L, Jones EJH, Ahmad J, Oakley B, Cáceres ASJ, Charman T, Buitelaar JK, Murphy DGM,
Chatham C, den Ouden H, Loth E (2020) Modeling flexible behavior in childhood to adulthood
shows age-dependent learning mechanisms and less optimal learning in autism in each age group.

PLoS Biol 18:e3000908 Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC7591042/
[Accessed April 26, 2025].

Demopoulos C, Brandes-Aitken AN, Desai SS, Hill SS, Antovich AD, Harris J, Marco EJ (2015) Shared and
Divergent Auditory and Tactile Processing in Children with Autism and Children with Sensory
Processing Dysfunction Relative to Typically Developing Peers. J Int Neuropsychol Soc 21:444–454
Available at: https://www.cambridge.org/core/journals/journal-of-the-international-
neuropsychological-society/article/abs/shared-and-divergent-auditory-and-tactile-processing-in-
children-with-autism-and-children-with-sensory-processing-dysfunction-relative-to-typically-
developing-peers/0F61C695B7ED89D61F5150C0978F209B [Accessed October 12, 2023].

Espenhahn S, Godfrey KJ, Kaur S, McMorris C, Murias K, Tommerdahl M, Bray S, Harris AD (2023)
Atypical Tactile Perception in Early Childhood Autism. J Autism Dev Disord 53:2891–2904 Available
at: https://pubmed.ncbi.nlm.nih.gov/35482274/ [Accessed October 12, 2023].

Failla MD, Peters BR, Karbasforoushan H, Foss-Feig JH, Schauder KB, Heflin BH, Cascio CJ (2017)
Intrainsular connectivity and somatosensory responsiveness in young children with ASD. Mol
Autism 8:1–11 Available at: https://molecularautism.biomedcentral.com/articles/10.1186/s13229-
017-0143-y [Accessed April 16, 2025].

Feigin H, Shalom-Sperber S, Zachor DA, Zaidel A (2021) Increased influence of prior choices on
perceptual decisions in autism. Elife 10.

Forschack N, Gundlach C, Hillyard S, Müller MM (2022) Attentional capture is modulated by stimulus
saliency in visual search as evidenced by event-related potentials and alpha oscillations. Attention,
Perception, Psychophys 2022 853 85:685–704 Available at:
https://link.springer.com/article/10.3758/s13414-022-02629-6 [Accessed April 30, 2025].

Foss-Feig JH, Heacock JL, Cascio CJ (2012) Tactile responsiveness patterns and their association with core
features in autism spectrum disorders. Res Autism Spectr Disord 6:337–344.

Frith U (2003) Autism: Explaining the enigma, 2nd ed. Malden: Blackwell Publishing.

Froehlich AL, Anderson JS, Bigler ED, Miller JS, Lange NT, Dubray MB, Cooperrider JR, Cariello A, Nielsen
JA, Lainhart JE (2012) Intact Prototype Formation but Impaired Generalization in Autism. Res
Autism Spectr Disord 6:921 Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC3267426/
[Accessed May 22, 2025].

Gastgeb HZ, Strauss MS (2012) Categorization in ASD: The Role of Typicality and Development. Perspect
Lang Learn Educ 19:66 Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC3372926/ [Accessed
April 23, 2025].

Goel A, Cantu DA, Guilfoyle J, Chaudhari GR, Newadkar A, Todisco B, de Alba D, Kourdougli N, Schmitt
LM, Pedapati E, Erickson CA, Portera-Cailliau C (2018) Impaired perceptual learning in a mouse
model of Fragile X syndrome is mediated by parvalbumin neuron dysfunction and is reversible. Nat
Neurosci 21:1404–1411 Available at: http://dx.doi.org/10.1038/s41593-018-0231-0.

Goldstone RL (1994) Influences of categorization on perceptual discrimination. J Exp Psychol Gen
123:178–200 Available at: https://pubmed.ncbi.nlm.nih.gov/8014612/ [Accessed May 23, 2025].

Goris J, Braem S, Van Herck S, Simoens J, Deschrijver E, Wiersema JR, Paton B, Brass M, Todd J (2022)
Reduced Primacy Bias in Autism during Early Sensory Processing. J Neurosci 42:3989–3999
Available at: https://www.jneurosci.org/content/42/19/3989 [Accessed December 11, 2022].

Goris J, Silvetti M, Verguts T, Wiersema JR, Brass M, Braem S (2021) Autistic traits are related to worse
performance in a volatile reward learning task despite adaptive learning rates. Autism 25:440–451
Available at: https://pubmed.ncbi.nlm.nih.gov/33030041/ [Accessed April 25, 2025].

Green SA, Hernandez LM, Bowman HC, Bookheimer SY, Dapretto M (2018) Sensory over-responsivity
and social cognition in ASD: Effects of aversive sensory stimuli and attentional modulation on
neural responses to social cues. Dev Cogn Neurosci 29:127–139.

Grubb MA, Behrmann M, Egan R, Minshew NJ, Carrasco M, Heeger DJ (2013) Endogenous spatial
attention: evidence for intact functioning in adults with autism. Autism Res 6:108 Available at:
https://pmc.ncbi.nlm.nih.gov/articles/PMC3661738/ [Accessed April 30, 2025].

Hachen I, Reinartz S, Brasselet R, Stroligo A, Diamond ME (2021) Dynamics of history-dependent
perceptual judgment. Nat Commun 2021 121 12:6036- Available at:
https://www.nature.com/articles/s41467-021-26104-2 [Accessed February 26, 2026].

Haigh SM (2018) Variable sensory perception in autism. Eur J Neurosci 47:602–609 Available at:
https://pubmed.ncbi.nlm.nih.gov/28474794/ [Accessed June 23, 2025].

Harris H, Israeli D, Minshew N, Bonneh Y, Heeger DJ, Behrmann M, Sagi D (2015) Perceptual learning in
autism: Over-specificity and possible remedies. Nat Neurosci 18:1574–1576 Available at:
https://pubmed.ncbi.nlm.nih.gov/26436903/ [Accessed April 25, 2025].

Hautus, M.J., Macmillan, N.A., & Creelman CD (2021) Detection Theory: A User's Guide, 3rd ed.
Routledge. Available at: https://doi.org/10.4324/9781003203636.

He JL, Williams ZJ, Harris A, Powell H, Schaaf R, Tavassoli T, Puts NAJ (2023) A working taxonomy for
describing the sensory differences of autism. Mol Autism 2023 141 14:1–17 Available at:
https://molecularautism.biomedcentral.com/articles/10.1186/s13229-022-00534-1 [Accessed July
11, 2024].

He JL, Wodka E, Tommerdahl M, Edden RAE, Mikkelsen M, Mostofsky SH, Puts NAJ (2021a) Disorder-
specific alterations of tactile sensitivity in neurodevelopmental disorders. Commun Biol 4:1–15
Available at: http://dx.doi.org/10.1038/s42003-020-01592-y.

He JL, Wodka E, Tommerdahl M, Edden RAE, Mikkelsen M, Mostofsky SH, Puts NAJ (2021b) Disorder-
specific alterations of tactile sensitivity in neurodevelopmental disorders. Commun Biol 4 Available
at: https://pubmed.ncbi.nlm.nih.gov/33483581/ [Accessed February 28, 2025].

Head J, Helton WS (2014) Sustained attention failures are primarily due to sustained cognitive load not
task monotony. Acta Psychol (Amst) 153:87–94 Available at:
https://www.sciencedirect.com/science/article/abs/pii/S0001691814002133 [Accessed May 6,
2025].

Järvinen-Pasley A, Wallace GL, Ramus F, Happé F, Heaton P (2008) Enhanced perceptual processing of
speech in autism. Dev Sci 11:109–121 Available at: https://pubmed.ncbi.nlm.nih.gov/18171373/
[Accessed June 25, 2025].

Kerzel D, Schönhammer J (2013) Salient stimuli capture attention and action. Atten Percept Psychophys
75:1633–1643 Available at: https://pubmed.ncbi.nlm.nih.gov/23918550/ [Accessed January 26,
2025].

Khachadourian V, Mahjani B, Sandin S, Kolevzon A, Buxbaum JD, Reichenberg A, Janecka M (2023)

Comorbidities in autism spectrum disorder and their etiologies. Transl Psychiatry 13:1–7 Available
at: https://www.nature.com/articles/s41398-023-02374-w [Accessed April 25, 2025].

Klinger LG, Dawson G (2001) Prototype formation in autism. Dev Psychopathol 13:111–124 Available at:
https://pubmed.ncbi.nlm.nih.gov/11346046/ [Accessed April 26, 2025].

Lavie N, Dalton P (2014) Load Theory of Attention and Cognitive Control. Oxford Handb Atten:56–75
Available at: https://academic.oup.com/edited-volume/41256/chapter/350821210 [Accessed April
30, 2025].

Licznerski P et al. (2020) ATP Synthase c-Subunit Leak Causes Aberrant Cellular Metabolism in Fragile X
Syndrome. Cell 182:1170-1185.e9 Available at: https://pubmed-ncbi-nlm-nih-
gov.proxy.insermbiblio.inist.fr/32795412/ [Accessed July 7, 2024].

Mattioni L, Barbieri A, Grigoli A, Balasco L, Bozzi Y, Provenzano G (2024) Alterations of Perineuronal Net
Expression and Abnormal Social Behavior and Whisker-dependent Texture Discrimination in Mice
Lacking the Autism Candidate Gene Engrailed 2. Neuroscience 546:63–74 Available at:
https://pubmed.ncbi.nlm.nih.gov/38537894/ [Accessed May 19, 2025].

Mercado E, Chow K, Church BA, Lopata C (2020) Perceptual category learning in autism spectrum
disorder: Truth and consequences. Neurosci Biobehav Rev 118:689–703 Available at:
https://www.sciencedirect.com/science/article/abs/pii/S0149763420305558 [Accessed April 25,
2025].

Micher N, Mazenko D, Lamy D (2024) Unconscious Processing Contaminates Objective Measures of
Conscious Perception : Evidence From the Liminal Prime Paradigm. J Cogn:1–16.

Mientjes EJ, Nieuwenhuizen I, Kirkpatrick L, Zu T, Hoogeveen-Westerveld M, Severijnen L, Rifé M,
Willemsen R, Nelson DL, Oostra BA (2006) The generation of a conditional Fmr1 knock out mouse
model to study Fmrp function in vivo. Neurobiol Dis 21:549–555 Available at:
https://pubmed.ncbi.nlm.nih.gov/16257225/ [Accessed June 25, 2025].

Mikkelsen M, Wodka EL, Mostofsky SH, Puts NAJ (2018) Autism spectrum disorder in the scope of tactile
processing. Dev Cogn Neurosci 29:140–150 Available at:
http://dx.doi.org/10.1016/j.dcn.2016.12.005.

Minshew NJ, Goldstein G, Siegel DJ (1997) Neuropsychologic functioning in autism: Profile of a complex
information processing disorder. J Int Neuropsychol Soc 3:303–316 Available at:
https://www.cambridge.org/core/journals/journal-of-the-international-neuropsychological-
society/article/abs/neuropsychologic-functioning-in-autism-profile-of-a-complex-information-
processing-disorder/581818F7F1A05B9413B7BA35EACE2543 [Accessed June 25, 2025].

Minshew NJ, Hobson JA (2008) Sensory sensitivities and performance on sensory perceptual tasks in
high-functioning individuals with autism. J Autism Dev Disord 38:1485 Available at:
https://pmc.ncbi.nlm.nih.gov/articles/PMC3077539/ [Accessed June 25, 2025].

Mol W, Post S, Lee M, Goel A (2024) Learning Impairments in Fmr1-/- mice on an Audio-Visual Temporal
Pattern Discrimination Task. bioRxiv:2024.09.25.615092 Available at:
https://www.biorxiv.org/content/10.1101/2024.09.25.615092v1 [Accessed January 22, 2025].

Molesworth CJ, Bowler DM, Hampton JA (2005) The prototype effect in recognition memory: Intact in
autism? J Child Psychol Psychiatry Allied Discip 46:661–672 Available at:
https://pubmed.ncbi.nlm.nih.gov/15877770/ [Accessed May 22, 2025].

Mottron L, Dawson M, Soulières I, Hubert B, Burack J (2006) Enhanced perceptual functioning in autism:
An update, and eight principles of autistic perception. J Autism Dev Disord 36:27–43.

Murphy G, Groeger JA, Greene CM (2016) Twenty years of load theory—Where are we now, and where
should we go next? Psychon Bull Rev 2016 235 23:1316–1340 Available at:
https://link.springer.com/article/10.3758/s13423-015-0982-5 [Accessed May 6, 2025].

O'Riordan M, Passetti F (2006) Discrimination in autism within different sensory modalities. J Autism Dev
Disord 36:665–675 Available at: https://pubmed.ncbi.nlm.nih.gov/16639532/ [Accessed December
3, 2022].

O'Riordan M, Plaisted K (2001) Enhanced discrimination in autism. Q J Exp Psychol Sect A 54:961–979
Available at: /doi/pdf/10.1080/713756000?download=true [Accessed April 24, 2025].

O'Riordan MA, Plaisted KC, Driver J, Baron-Cohen S (2001) Superior visual search in autism. J Exp Psychol
Hum Percept Perform 27:719–730 Available at: https://pubmed.ncbi.nlm.nih.gov/11424657/
[Accessed April 24, 2025].

Orefice LLL, Zimmerman ALL, Chirila AMM, Sleboda SJJ, Head JPP, Ginty DDD (2016) Peripheral
Mechanosensory Neuron Dysfunction Underlies Tactile and Behavioral Deficits in Mouse Models of
ASDs. Cell 166:299–313 Available at: http://dx.doi.org/10.1016/j.cell.2016.05.033.

Plaisted K, O'riordan M, Baron-Cohen S (1998a) Enhanced Discrimination of Novel, Highly Similar Stimuli
by Adults with Autism During a Perceptual Learning Task.

Plaisted K, O'Riordan M, Baron-Cohen S (1998b) Enhanced Visual Search for a Conjunctive Target in
Autism: A Research Note. J Child Psychol Psychiatry 39:777–783.

Plaisted KC (2001) Reduced Generalization in Autism: An Alternative to Weak Central Coherence.

Ploog BO (2010) Stimulus overselectivity four decades later: A review of the literature and its
implications for current research in autism spectrum disorder. J Autism Dev Disord 40:1332–1349
Available at: https://link.springer.com/article/10.1007/s10803-010-0990-2 [Accessed April 26,
2025].

Puts NAJ, Wodka EL, Tommerdahl M, Mostofsky SH, Edden RAE (2014) Impaired tactile processing in
children with autism spectrum disorder. J Neurophysiol 111:1803–1811 Available at:
https://pubmed.ncbi.nlm.nih.gov/24523518/ [Accessed December 3, 2022].

Ridderinkhof A, de Bruin EI, van den Driesschen S, Bögels SM (2018) Attention in Children With Autism
Spectrum Disorder and the Effects of a Mindfulness-Based Program. J Atten Disord 24:681
Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC7003152/ [Accessed May 6, 2025].

Robertson CE, Baron-Cohen S (2017) Sensory perception in autism. Nat Rev Neurosci 18:671–684.

Rong Y, Yang CJ, Jin Y, Wang Y (2021a) Prevalence of attention-deficit/hyperactivity disorder in
individuals with autism spectrum disorder: A meta-analysis. Res Autism Spectr Disord 83:101759.

Rong Y, Yang CJ, Jin Y, Wang Y (2021b) Prevalence of attention-deficit/hyperactivity disorder in
individuals with autism spectrum disorder: A meta-analysis. Res Autism Spectr Disord 83:101759
Available at: https://www.sciencedirect.com/science/article/pii/S1750946721000349 [Accessed
May 6, 2025].

Rotschafer SE (2021) Auditory Discrimination in Autism Spectrum Disorder. Front Neurosci 15:1–13.

Samson F, Mottron L, Jemel B, Belin P, Ciocca V (2006) Can spectro-temporal complexity explain the
autistic pattern of performance on auditory tasks? J Autism Dev Disord 36:65–76 Available at:
https://pubmed.ncbi.nlm.nih.gov/16382329/ [Accessed April 24, 2025].

Sapey-Triomphe LA, Temmerman J, Puts NAJ, Wagemans J (2021a) Prediction learning in adults with
autism and its molecular correlates. Mol Autism 12 Available at: https://pubmed-ncbi-nlm-nih-
gov.proxy.insermbiblio.inist.fr/34615532/ [Accessed October 12, 2023].

Sapey-Triomphe LA, Timmermans L, Wagemans J (2021b) Priors Bias Perceptual Decisions in Autism, But
Are Less Flexibly Adjusted to the Context. Autism Res 14:1134–1146 Available at: https://pubmed-
ncbi-nlm-nih-gov.proxy.insermbiblio.inist.fr/33283970/ [Accessed October 12, 2023].

Semelidou O, Gauvrit T, Vandromme C, Cornier A, Saint-Jean A, Le Feuvre Y, Ginger M, Frick A (2024)
Stimulus encoding shapes tactile perception and underlies alterations in autism.
bioRxiv:2024.08.08.607129 Available at:
http://biorxiv.org/content/early/2024/10/08/2024.08.08.607129.abstract.

Shaw KA et al. (2025) Prevalence and Early Identification of Autism Spectrum Disorder Among Children
Aged 4 and 8 Years — Autism and Developmental Disabilities Monitoring Network, 16 Sites, United
States, 2022. MMWR Surveill Summ 74:1–22 Available at:
https://www.cdc.gov/mmwr/volumes/74/ss/ss7402a1.htm [Accessed May 9, 2025].

Shenouda J, Barrett E, Davidow AL, Sidwell K, Lescott C, Halperin W, Silenzio VMB, Zahorodny W (2023)
Prevalence and Disparities in the Detection of Autism Without Intellectual Disability. Pediatrics 151
Available at: https://pubmed.ncbi.nlm.nih.gov/36700335/ [Accessed June 25, 2025].

Smallwood J (2013) Penetrating the fog of the decoupled mind: The effects of visual salience in the
sustained attention to response task. Can J Exp Psychol 67:32–40.

Soulières I, Mottron L, Giguère G, Larochelle S (2011) Category induction in autism: Slower, perhaps
different, but certainly possible. Q J Exp Psychol 64:311–327 Available at:
https://pubmed.ncbi.nlm.nih.gov/20623440/ [Accessed April 26, 2025].

Soulières I, Mottron L, Saumier D, Larochelle S (2007) Atypical categorical perception in autism:
Autonomy of discrimination? J Autism Dev Disord 37:481–490.

Thye MD, Bednarz HM, Herringshaw AJ, Sartin EB, Kana RK (2018) The impact of atypical sensory
processing on social impairments in autism spectrum disorder. Dev Cogn Neurosci 29:151–167.

Venker CE, Mathée J, Neumann D, Edwards J, Saffran J, Ellis Weismer S (2021) Competing Perceptual
Salience in a Visual Word Recognition Task Differentially Affects Children With and Without Autism
Spectrum Disorder. Autism Res 14:1147–1162 Available at: /doi/pdf/10.1002/aur.2457 [Accessed
May 6, 2025].

Zaidel A, Goin-Kochel RP, Angelaki DE (2015) Self-motion perception in autism is compromised by visual
noise but integrated optimally across multiple senses. Proc Natl Acad Sci U S A 112:6461–6466
Available at: https://www.pnas.org/doi/abs/10.1073/pnas.1506582112 [Accessed January 5,
2024].

Zetler NK, Cermak SA, Engel-Yeger B, Gal E (2019) Somatosensory discrimination in people with autism
spectrum disorder: A Scoping review. Am J Occup Ther 73:1–14.

Zhai J, Li X, Zhou Y, Fan L, Xia W, Wang X, Li Y, Hou M, Wang J, Wu L (2023) Correlation and predictive

ability of sensory characteristics and social interaction in children with autism spectrum disorder.
Front Psychiatry 14:1056051.

1118