Capsular specificity in temperate phages of *Klebsiella pneumoniae* is driven by diverse receptor-binding enzymes

Aleksandra Otwinowska[1,*], Janusz Koszucki[2,3,*], Vyshakh R. Panicker[2,3], Jade Leconte[2], Sebastian Olejniczak[1], Kathryn E. Holt[4,5], Edward J. Feil[6], Eduardo P. C. Rocha[7], Bogna Smug[2], Barbara Maciejewska[1], Zuzanna Drulis-Kawa[1†] & Rafal J. Mostowy[2,†]

1) Department of Pathogen Biology and Immunology, University of Wroclaw, Wrocław, Poland

2) Malopolska Centre of Biotechnology, Jagiellonian University, Kraków, Poland

3) Doctoral School of Exact and Natural Sciences, Jagiellonian University, Kraków, Poland

4) Department of Infection Biology, London School of Hygiene & Tropical Medicine, London, United Kingdom

5) Department of Infectious Diseases, School of Translational Medicine, Monash University, Melbourne, Victoria, Australia

6) The Milner Centre for Evolution, Department of Life Sciences, University of Bath, Bath, UK

7) Institut Pasteur, Université Paris Cité, CNRS, UMR3525, Microbial Evolutionary Genomics, Paris, France

∗ These authors contributed equally to this work.

† Correspondence: rafal.mostowy@uj.edu.pl, zuzanna.drulis-kawa@uwr.edu.pl

Abstract

Understanding host-range determinants in temperate bacteriophages is critical for elucidating phage-host co-evolution and advancing phage therapies.We analysed 3,900 *Klebsiella* genomes from diverse ecological niches to identify prophage-encoded proteins mediating capsule tropism. We applied a genome-wide association study (GWAS) correlating prophage protein clusters (from 8,105 prophages) with confidently assigned bacterial K-loci. GWAS identified high-confidence predictors for 16 out of 35 most diverse K-loci, of which 14 were receptor-binding proteins (RBPs) belonging to classical depolymerases ($n = 6$), SGNH hydrolases which deacetylate polysaccharides ($n = 6$), and structurally novel RBPs ($n = 2$). When we relaxed the filtering thresholds, we identified 26 putative depolymerases, of which 12 were deemed as strong predictions against 10 K-types. In parallel, 50 depolymerases manually found in prophages from the representative subset of 99 bacterial isolates, together with an additional 10 depolymerases based on GWAS predictions, were prepared as recombinant proteins and tested on a *Klebsiella* reference panel of 119 K-types. Most predicted prophage depolymerases (34/60) failed to yield soluble products, 6 were not active on the K-types panel, and 5/14 targeted a different K-type than their bacterial host, highlighting the unpredictability of prophages as a source of functional enzymes. A comparison of GWAS-predicted enzymes, from the manual search and from virulent phages showed that depolymerase specificity was often difficult to infer from sequence or structure alone. Our findings reveal that *Klebsiella* prophages encode structurally diverse RBPs, suggesting temperate phages may rely not only on depolymerisation but also on capsule modification—such as deacetylation—for infection. This suggests capsule diversity in *K. pneumoniae* may be substantially underestimated, with implications for phage specificity, competition and vaccine design.

Introduction

Host range is among the most critical traits of viruses, determining the variety of hosts that can be infected by a given virus[1]. Understanding and predicting viral host range is fundamental for studying viral evolution, ecology and infection dynamics[2]. In bacteriophages, host range is driven by co-evolution with bacterial hosts[3,4] and shaped by horizontal gene transfer[5,6], with broad relevance for phage therapy and microbiome engineering[7,8,9,10]. However, accurate prediction of viral host range remains challenging due to the complexity of virus-host interactions, from initial adsorption to successful evasion of host defence systems[11]. Nevertheless, recent advances in multi-omics and computational biology have been enabling statistical predictions that can be validated empirically[12]. Indeed, two recent genomic studies demonstrate that such approaches can help successfully predict the infectivity of virulent phages in two species of *Enterobacteriaceae*, *Klebsiella pneumoniae*[13] and *Escherichia coli*[14], underscoring the critical role of receptor-binding proteins (RBPs) in virus host range.

K. pneumoniae in particular serves as a great model for studying bacteria-virus interactions. It is an opportunistic bacterial pathogen characterised by the extensive diversity of its capsular polysaccharides (CPS)[15]. This diversity encompasses 77 serologically distinct CPS polysaccharides (K-types) with fully characterised biosynthetic loci (K-loci), and a at least further 86 genetically distinct K-loci of unknown polysaccharide structure[16,17,18]. Experimentally isolated virulent phages infecting *K. pneumoniae* typically produce visible halos on bacterial lawns, indicating enzymatic degradation of CPS[19]. These halos result from phage RBPs with depolymerase activity, enabling recognition, degradation, and subsequent infection[20]. Such enzymes generally exhibit highly specific activity against individual CPS types, with 58 experimentally validated depolymerases described so far, targeting 31 distinct K-types[21]. Structurally, these depolymerase-containing RBPs are typically homotrimers, featuring a characteristic right-handed parallel β-helix[22,23,24,25]. However, our understanding of the genetic, structural diversity and specificity of depolymerase-containing RBPs in *Klebsiella* phages remains far from complete.

Given that our knowledge about phage depolymerases comes predominantly from studies focused on virulent phages, their role in the biology of temperate phages remains even less understood. In *K. pneumoniae*, studies have demonstrated capsule-specific tropism in temperate phages, observing that prophages induced from bacterial genomes could infect only bacterial lineages with identical K-loci, evidenced by characteristic halos indicative of depolymerase activity[26]. Yet, attempts to computationally

identify corresponding depolymerase genes in prophages failed to show clear correlations with the observed infection patterns. Subsequent genomic analyses of nearly 4,000 *K. pneumoniae* genomes have similarly reported a surprising scarcity of identifiable depolymerase sequences in prophages [27]. A recent machine-learning-based study reported prophage-encoded depolymerase domains in approximately 20% of *Klebsiella* prophages, emphasising their potential as a novel source of host-range determinants [28]. Nevertheless, the overall rarity of depolymerase predictions raises the question of whether this scarcity results from technical limitations in annotation or fundamental biological differences between virulent and temperate phages.

To address these gaps, we aimed to identify molecular determinants of capsule specificity in temperate phages of *K. pneumoniae* using a Genome-Wide Association Study (GWAS) approach. GWAS identifies genetic variants statistically associated with phenotypes of interest, an approach already successfully adapted to bacterial genomics [29,30,31,32]. Compared to other statistical learning methods, GWAS has the advantage of not relying on pre-labelled training datasets, thereby avoiding predefined functional expectations and allowing for biologically interpretable associations between genetic variants and phenotypic traits. We applied GWAS to a diverse dataset of *K. pneumoniae* genomes from two systematically curated collections (Australia [33] and Italy [34]), representing near-random sampling from a collection of 3,900 genomes across diverse ecological and clinical niches. Within this dataset, we predicted prophages and employed GWAS to the subset of 2,527 genomes for which we could confidently predict the type of capsular polysaccharide (K-type), identifying prophage protein clusters statistically associated with 35 common K-types as candidate predictors of capsule specificity. Independently, we manually curated depolymerase-containing RBPs from prophages found in 99 representative bacterial genomes (KASPAH-REF) and performed recombinant protein preparation and enzymatic activity testing against 119 KL-type strain collection, including a set of 58 experimental depolymerases from virulent phages in the comparison [21] (see Supplementary Table S1; see also Figure 1 for a visual summary of our approach). Our results demonstrate that while prophages can be a rich source of structurally and functionally diverse depolymerases, they are also an unpredictable one. Our approach also uncovers a previously underappreciated class of receptor-binding proteins dominated by SGNH-domain enzymes, likely involved in removing capsule modifications such as acetyl groups, and pointing to capsule deacetylation as a widespread alternative mechanism of infection initiation.

1. Study population

Italy:
KLEBPAVIA
collection

Australia:
KASPAH
collection

3,900 *Klebsiella spp* isolates

2,527 *K. pneumoniae* isolates
with confident K-type predictions

2. Bioinformatics

detected prophage

bacterial genome

KL-X

SC1

KL-Y

SC2

KL-Z

SC3

3. GWAS

Predictors for KL-X

Recall

Precision

pectin-lyase

unknown
function

4. Experimental verification

A) Functional annotation

B) Recombinant protein production

C) Testing activity against 119 K-types

inactive

active

5. Comparison of predictions with experiments

GWAS prediction

Active based on manual search

Active based on GWAS prediction

Active based on literature

Figure 1. Visual summary of the study approach. (1) We assembled an ecologically diverse collection of 3,900 *Klebsiella* genomes, of which 2,527 *Klebsiella pneumoniae* isolates had high-confidence capsular (K-locus) assignments. (2) Prophage regions were predicted in each genome, and all prophage-encoded proteins were clustered and functionally annotated. (3) A bacterial GWAS (elastic-net model) was then performed on 2,527 isolates to link prophage protein clusters to 35 most diverse K-loci, yielding protein clusters as candidate predictors of capsule specificity. (4) 50 manually selected putative depolymerases from 99 high-quality reference genomes ("KASPAH-REF") were selected for recombinant protein production and enzymatic activity testing against a panel of 119 *Klebsiella* strains with diverse K-types. (5) GWAS predictors (green node) were compared to these experimental results (red node) and to previously characterized virulent-phage depolymerases (purple node); for predictors lacking any known sequence similarity, we selected representative β-helix (pectin-lyase domain) proteins for recombinant production and functional testing (blue node).

Results

Isolate selection and data

The aim of this study was threefold: (a) to identify genetic determinants of capsule tropism in temperate phages of *K. pneumoniae*, (b) to functionally characterise putative depolymerase candidates from a subset of isolates, and (c) to compare genomic predictions with experimental outcomes. To this end, we carried out a comprehensive search that is described in details in Methods and visually summarised in Figure 1. Briefly, we assembled a dataset of 3,900 *Klebsiella* genomes by combining two large and systematically curated sources. The KASPAH dataset includes 391 clinical and colonisation (gut/throat) isolates collected from hospital patients in Melbourne[35,36,33], while KLEBPAVIA comprises 3,509 isolates sampled through a One Health framework in northern Italy, spanning human, animal, environmental, and agricultural sources[34]. Together, these datasets provide a comprehensive and ecologically broad view of *Klebsiella* diversity, avoiding the clone- and outbreak-driven biases often seen in public genome databases. All isolates were short-read sequenced, and 99 representative KASPAH genomes were additionally sequenced using long reads to produce high-quality hybrid assemblies[37] (KASPAH-REF; see Figure 2A). For each genome, we identified the capsular locus (K-locus) using Kaptive[17] and assigned bacterial isolates into sequence clusters (SCs) using PopPunk[38]; the latter was done to account for population structure in the GWAS analysis, minimising confounding from recent shared ancestry and prophage co-inheritance among closely related isolates.

Prophage regions ($n = 8105$) were identified using VirSorter[39], PhiSpy[40], and CheckV[41], retaining both high-completeness (likely intact) and medium-completeness predictions to capture fragmented prophages in short-read assemblies (see Supplementary Figure S1 for a distribution of prophage completeness in most frequent K-types). To facilitate robust comparisons, we focused subsequent analyses on 2,527 genomes within the *K. pneumoniae* species complex (KPSC) that had confident K-locus predictions, and focused on 35 K-loci with sufficient diversity ($\geqslant 10$ SCs) to ensure statistical power in subsequent analyses (see Figure 2B). Those 35 K-loci each represented between 10-24 SCs and from 29 phage variants for KL47 to 112 phage variants for KL30 (median of 55), hence encompassing considerable diversity in both capsule type and prophage content. The phylogenetic tree, the genetic and prophage diversity are shown in Figure 2. The full distribution of SCs and phage variants per K-locus is shown in Supplementary Figure S2. The bacterial and prophage metadata are provided in Supplementary Tables S2 and S3, respectively.

Figure 2. The dataset. (A) Maximum-likelihood tree of 2,527 isolates with clades shaded by sequence cluster (SC) assignment. Red dots show isolates belonging to the KASPAH-REF subset ($n = 99$). Shaded areas show the three main species from the *K. pneumoniae* Species Complex with reliable K-locus predictions: *K. pneumoniae* (purple), *K. variicola* (yellow), and *K. quasipneumoniae* (green). (B) Top plot shows bacterial diversity as the number of SCs per predicted K-locus amongst KASPAH-REF isolates (red) and all other isolates (blue). Bottom plot shows phage diversity measured by the number of phage variants (similar prophages clustered using the wGRR = 0.95 threshold; see Methods) identified in the KASPAH-REF isolates (red) and using all 2,428 remaining isolates (blue). Only K-loci with at least 8 SCs are shown. Brown dashed line at 10 SCs shows the cutoff for the inclusion of the most diverse K-loci for the GWAS prediction, resulting in $n = 35$ K-loci.

GWAS predicts structurally diverse receptor-binding proteins

To perform a genome-wide association study (GWAS), we analysed prophage protein clusters as genetic predictors of capsule tropism by correlating their presence with K-locus assignments. We used an elastic-net regression model to accommodate the biological expectation that capsule tropism may be driven by a small number of key proteins, while allowing for potential diversity across variants. Population structure was controlled by including SCs as covariates, minimising confounding from lineage effects. Each protein cluster was assigned a regression coefficient (β) and a significance threshold, predictors were defined using $\beta > 0$ and a Bonferroni-corrected p-value below 0.05, and performance of these predictors was evaluated based on their ability to accurately classify K-loci using F1 score and Matthews Correlation Coefficient (MCC). Full details of the GWAS methodology are provided in Methods.

Figure 3A shows the results of GWAS for predictors filtered with F1 $\geqslant$ 0.5 and MCC $\geqslant$ 0.5. Using those thresholds, we obtained 35 protein clusters associated with 24 out of 35 K-loci, with eight K-loci (KL3, KL11, KL14, KL21, KL24, KL25, KL111 and KL127) having multiple associated predictors. Functional annotation of these 35 clusters using PHROGs[42] revealed that 23 of them encoded tail-associated proteins, of which 18 were predicted to be tail fibres—consistent with our expectation that capsule-specific associations would predominantly involve receptor-binding proteins (RBPs). We also observed a negative correlation between precision and recall across predictors (Pearson's $r = -0.58$, 95% CI: -0.76 to -0.30; see Supplementary Figure S3). While such a trade-off is expected, the presence of low-precision, high-recall predictors suggests that some candidates may target multiple K-types or represent associations unrelated to capsule interaction (more on this later).

Given that the initial GWAS results predominantly identified receptor-binding proteins (RBPs), we next focused on refining predictors based on biological expectations of polysaccharide specificity. Since most known depolymerases act on a single capsule type[21], we applied a stringent precision threshold ($\geqslant 0.8$) to prioritise highly specific predictors. Furthermore, since RBPs are often modular, with variation across N-terminal and C-terminal domains[43], we reasoned that the clustering thresholds used to define protein families could affect whether related proteins were grouped together or split into distinct predictors. To address this, we systematically repeated GWAS across six clustering thresholds, combining identity cutoffs (none, 50%, 80%) with coverage thresholds (50%, 80%), maximising chances that such potential modularity was captured. For each K-locus, we then selected the clustering level that maximised predictor performance (defined as the highest product of F1 score and MCC, with both metrics $\geqslant$ 0.5), aiming to

identify the most biologically informative protein cluster for each of them.

2   Using these criteria, we obtained predictions for 16 out of 35 K-loci (Figure 3B). We then selected a
    representative protein from best predictor (protein cluster) and generated their 3D models using Al-

4   phaFold3[44]. (Note that for 19 K-loci, there were no predictors that fulfilled our criteria; see Discussion.)
    We also mapped the locations of the major ECODs (Evolutionary Classification of Protein Domains)

6   detected in those proteins onto the 3D structures (see Figure 3B). We found that 6 of those 16 pro-
    teins (from predictors for KL3, KL14, KL28, KL38, KL62, KL127) were classical depolymerases with the

8   characteristic, parallel β-helix fold (ECOD T-level: 207.2.1, *Pectin-lyase like*). Another 6 proteins (from
    predictors for KL2, KL8, KL11, KL16, KL24 and KL55) lacked the parallel β-helix, carrying a *SGNH*

10 *hydrolase* domain (ECOD T-level: 2007.5.1), and four remaining proteins had no significant ECOD hits
    identified. Of those four, two proteins (predictors for KL47 and KL103) looked like unusual RBPs with

12 previously uncharacterised folds, while the remaining two (predictors for KL30 and KL111) did not look
    like RBPs. We used FoldSeek to search the non-RBP predictors against CATH, PDB, and AFDB and

14 found that these proteins structurally resembled known acetyltransferases[45].
    Finally, we examined how varying threshold for filtering GWAS predictors influenced the distribution of

16 ECOD domains identified among the associated protein clusters. Regardless of the filters used, filtered
    predictors were dominated by two structural folds: the *Pectin lyase-like* domain and the *SGNH hydrolase*

18 domain, along with a substantial fraction of proteins lacking functional annotation (Figure 3C). These
    results suggest that capsule tropism in temperate phages of *K. pneumoniae* involves not only classical

20 depolymerases but also other structurally diverse RBPs with distinct folds. Furthermore, we observed
    that by relaxing predictor filtering criteria — specifically, allowing lower recall while maintaining high

22 precision — additional lower-frequency protein clusters emerged, some of which may encode previously
    uncharacterised β-helix-containing depolymerases. We therefore next lowered the filtering criteria and

24 applied a series of manual curation steps to identify and putative depolymerases against as many K-loci
    as possible.

Figure 3. GWAS predicts diverse receptor-binding proteins. (A) F1 score versus Matthews Correlation Coefficient (MCC) for prophage protein clusters (PC50 clustering) identified as GWAS predictors with F1 $\geqslant$ 0.5 and MCC $\geqslant$ 0.5. Colours indicate functional annotations assigned using PHROGs. (B) Recall of the strongest predictor identified for each K-locus. For each K-locus, predictors with precision $\geqslant$ 0.8 were selected, retaining the one with the highest F1×MCC product across clustering thresholds; only those with F1 $\geqslant$ 0.5 and MCC $\geqslant$ 0.5 are shown. Shapes indicate the clustering threshold at which the strongest predictor was identified; colours reflect the top predicted ECOD domain, or grey for unannotated regions. For the 16 predictors obtained, AlphaFold3 models were generated based on representative proteins, with ECOD domains coloured based on HHpred predictions. (C) Distribution of ECOD domains across GWAS predictors across different filtering thresolds, shown as bar plots. The top row summarises predictors meeting F1 $\geqslant$ 0.5 and MCC $\geqslant$ 0.5, across varying precision thresholds (0.5–0.9). The bottom row shows predictors filtered at minimum precision $\geqslant$ 0.8, across varying F1 and MCC thresholds (0.1–0.5). ECOD domain names are abbreviated as follows: *SGNH hydrolase* (yellow; T-level: 2007.5.1), *Pectin-lyase like* (green; T-level: 207.2.1), Tail spike N-terminal (brown; Full name: *Putative tailspike protein Orf210 N-terminal domain*, T-level: 3856.1.1), Intramolecular chaperone (purple; Full name: *Intramolecular chaperone domain in virus tail spike protein*, T-level: 3240.1.1), gp11/gp12 RBD (black; Full name: *gp11/gp12 receptor-binding domain*, T-level: 877.1.1), P-loop hydrolase (pink; Full name: *P-loop containing nucleoside triphosphate hydrolases*, T-level: 2004.1.1). Grey denotes the absence of functional hits.

Prophages are an unpredictable source of active depolymerases

To comprehensively identify GWAS-predicted depolymerases containing the characteristic right-handed β-helix fold, we developed a semi-supervised curation framework combining multiple evaluation steps (full details in Supplementary Text S1). Starting from statistically significant GWAS predictors, we first assessed precision and recall for each protein cluster at both isolate and sequence-cluster (SC) levels to account for population structure. To accommodate the extensive modularity of phage receptor-binding proteins (RBPs), we then evaluated six alternative clustering thresholds to identify parameter settings that best captured biologically coherent protein families for each K-locus. Next, we applied phylogenetic mapping to visualise whether predictors consistently tracked with K-locus distribution across the bacterial tree[46]. Finally, based on combined evidence from classification metrics, clustering behaviour, domain annotation and structural modeling, predictors were conservatively classified as strong or likely. Using this approach, we identified 12 'strong' predictors for 10 K-loci (KL3, KL10, KL14, KL25, KL28, KL38, KL60, KL62, KL64, KL111) and 14 'likely' predictors for 11 K-loci (KL7, KL14, KL15, KL22, KL24, KL25, KL30, KL47, KL64, KL122, KL127), together spanning 18 of the 35 analysed K-loci (Figure 4A). Separately, we experimentally screened prophage-encoded candidate depolymerases from the KASPAH-REF dataset. Starting from 469 candidate proteins meeting initial selection criteria, we clustered these into 124 sequence groups, modelled all representatives using AlphaFold2[47], and manually evaluated the resulting structures for conserved features characteristic of depolymerases, particularly the parallel β-helix fold. This process yielded 50 proteins from 32 clusters for recombinant production (see Methods). These proteins were produced and purified, then assessed for enzymatic activity against 119 diverse *Klebsiella* strains via spot assays (Supplementary Table S4). Of those 50, 34 failed to produce soluble proteins, likely due to low expression yield or misfolding, 2 produced soluble proteins that failed to show evidence of enzymatic activity on any of the 119 tested K-types, and 14 were active on one of 11 K-types (KL23, KL28, KL32, KL38, KL46, KL52, KL60, KL62, KL64, KL127 and KL143; see Figure 4A, Supplementary Table S5 and Supplementary Text S2).

The 26 GWAS predictors (both from 'strong' and 'likely' category) were first compared to the manually curated collection of 50 depolymerases from the KASPAH-REF isolates (Supplementary Table S5) based on their protein sequence similarity. When the GWAS predictor from the 'strong' category did brea sequence similarity to any protein within our recombinant-protein set, or any of the the available enzymes with confirmed specificity described already in the literature (Supplementary Table S1), we commissioned

synthetic plasmid preparation and recombinant protein production for one representative protein from each cluster based on manual analyses to maximise chances of its successful production and enzymatic activity (see Methods).

Figure 4 shows a direct comparison of computational and experimental results per K-locus, alongside additional recombinant proteins from the literature from virulent phages. Specifically, 4A shows a table of the number of predictions per each K-locus, and Figure 4B shows a network-like comparison of the two sets of proteins, alongside 58 previously published depolymerases from virulent phages [21] (see Supplementary Table S1). We found that, out of 12 predictors in the 'strong' category, 5 were confirmed by sequence or structural similarity to the recombinant protein analysis (KL28, KL38, KL60, KL62, KL64) and 2 were confirmed by sequence or structural similarity to recombinant proteins from the literature: KL3 had distant similarity and a visibly different C-terminus, while the KL25 prediction had structurally identical receptor-binding domain (see Supplementary Text S3). Of the remaining 5 predictors, which were used to select 5 proteins for commercial production, 3 were confirmed as active enzymes (KL3, KL62 and KL111) while 2 did yield soluble proteins (KL10 and KL14) but no activity was detected against any of the 119 K-types. Additionally, one protein commercially produced based on a 'likely' predictor (KL122) was found active against isolates with the KL122 locus (Figure 4B; Supplementary Table S5).

By contrast, out of 11 K-types for which we obtained active enzymes via the manual search in the KASPAH-REF collection, five specificities (K23, K32, K46, K52, KL143) could not be compared with GWAS results as these K-types were excluded from the set of 35 loci, thus no predictors were selected. (One of them, KL23, a virulent-phage protein had already been reported and we observed structural homology of the receptor-binding domain; see Supplementary Text S3.) The enzymatic activity of the remaining six KASPAH-REF K-specific enzymes (K28, K38, K60, K62, K64 and KL127) was in line with GWAS predictions, either predictors classified as 'strong' or 'likely'. Importantly, 5 of 14 enzymes were active against a different K-type than the K-type of the prophage host (see Supplementary Figure S4).

Overall, we found that (a) of the 12 'strong' GWAS predictors, 10 were experimentally validated and 2 remained inconclusive, and (b) among the 14 active enzymes identified through manual search, only 6 corresponded to the 35 K-loci included in the GWAS analysis; all 6 were supported by GWAS predictions ('strong' or 'likely' predictors). Importantly, prophages proved to be an unpredictable source of active enzymes: many candidate genes failed to yield soluble, properly folded proteins, likely due to misfolding or expression constraints, and several enzymes displayed activity against a different K-type than that of

their host strain (see Discussion).

Figure 4. Prophages can be an unpredictable source of active depolymerases. (A) Table shows a systematic comparison of the number of bacterial SCs from the KASPAH-REF collection (red), the number of bacterial SCs from the full dataset (yellow), number of GWAS predictions in the 'strong' (dark green) 'likely' (light green) category), number of active recombinant proteins based on manual (magenta) and based on GWAS' 'strong' predictors (dark blue), and the number of experimentally verified depolymerases from virulent phages from Cheetham and colleagues[21]. (B) Sequence similarity network where each node represents a protein sequence and edges link proteins with a minimum of 50% bidirectional coverage (thin edges) or a minimum of 80% bidirectional coverage (thick edges). Colours are in accordance with panel A, with blue nodes showing additional proteins obtained via gene synthesis and recombinant protein production.

Complexity of the genotype–phenotype relationships in phage depolymerases

While the overall agreement between GWAS predictions and experimental validation was strong, closer inspection of the sequence similarity network (Figure 4B) revealed multiple genetic links between enzymes

with different K-type specificities (confirmed or predicted). To explore this complexity more systematically, we performed a detailed comparison of the protein sequence, structure and specificity relationships

across these enzymes, as described in Supplementary Text S3. This analysis revealed several key patterns that exemplify the challenges in predicting capsule specificity from sequence – or even structure – alone.

First, we observed frequent modularity, particularly variation in N-terminal domains known to be related to the virion taxonomic or morphological groups and to vary among phage depolymerases [43] (see

Supplementary Text S3). Second, we found that enzymes that shared significant sequence and structural similarity often exhibited distinct substrate specificities, indicating that proteins with related structures

of their receptor-binding domains can be functionally divergent. Figure 5 demonstrates this using the example of enzymes known or predicted to target K-types KL3, KL24 and KL28 from two connected

components of the network in Figure 4B. We found that GWAS predictors for KL3 (PC0449) and KL28 (PC0279) were both confirmed experimentally (391_03 and 1251_37), and their close comparison re-

vealed that they are genetically and structurally related to each other and to the KL24 predictions (PC0406 and PC1397; Figure 5A). Nevertheless, recombinant depolymerase against K3 (391_03) and

against K28 (1251_37) did not cross-react against K3/K24/K28 reference strains (Figure 5B) despite the fact that the K3, K24 and K28 polysaccharide structures are biochemically related (Figure 5C). Fur-

thermore, comparison to another recombinantly tested depolymerase (NCBI accession YP_03347555.1) from a virulent phage active on a K3-type capsule showed no genetic or structural similarity to any of

the other ones.

Interestingly, we found that the KL3 recombinant protein (391_03) was also active against the KL146

24 strain (see Figure 5B). We hypothesised that the dual specificity of the 391_03 protein against KL3 and KL146 reflects underlying structural similarity between the corresponding polysaccharides. Unfortu-

26 nately, the polysaccharide structure was not available for KL146 at the time of this publication, hence we compared the KL3 and KL146 loci. We found that the two loci shared two serotype-specific genes: *wcuC*

28 (glycosyl hydrolase) and *wbaZ* (glycosyl transferase), suggesting the presence of a common biochemical motif that could potentially explain the dual specificity of the KL3 prediction (see Supplementary Figure

30 S5). Intriguingly, capsular serotyping revealed a lack of KL146 antibodies cross-reactivity with K3 or any

other defined sera (Holt, personal communication).

2 In network analysis (Figure 4B), we also found examples of recombinantly tested depolymerases with the same substrate specificity that are related but genetically dissimilar. Examples include the K1-

4 specific depolymerases (NCBI accessions BAW85698.1 and YP_009302756.1, protein sequence identity of 35%), K5-specific depolymerases (APZ82848.1 and WNO47121.1, identity of 32%) or K21-specific

6 depolymerases (BAW85693.1 and YP_003347556.1, identity of 41%). Therefore, our results indicate that genetic or structural similarity is often insufficient to infer depolymerase specificity, pointing to a

8 nuanced genotype–phenotype map of these enzymes.

Figure 5. Depolymerase specificity is a complex phenotype. (A) One of the connected components from Figure 4B showing seven depolymerases: two 'strong' GWAS predictions (KL3, KL28; dark green), two 'likely' GWAS predictions (KL24; light green), one active protein from the manual search (KL28; pink), one active protein from GWAS prediction (KL3/KL146; blue), and an active protein from a virulent phage (KL3; purple). All of these proteins, except the virulent KL3 depolymerase, are genetically and structurally related, which can be seen in a systematic visualisation of pairwise alignments and 3D models obtained via AlphaFold3. (B) Spot test assay validating the predicted capsular specificity of 391_03 and 1251_37 selected proteins against reference serotypes KL3, KL24, KL28 and KL146. (C) Schematic representations of the K3, K24, and K28 capsule polysaccharides, as obtained from K-PAM[48].

K. pneumoniae prophage RBPs contain diverse enzymes

Our results so far showed that the β-helix-containing depolymerases are present in prophages, they are relatively rare, highly diverse, and not always active. Additionally, the 'pectin lyase-like' domain was not the predominant domain detected amongst the top GWAS results. Therefore, we next investigated the prevalence of different protein domains in prophage RBPs from our dataset. To this end, we analysed ECOD domain diversity across 4,598 high-completeness prophages, focusing on prophage protein clusters annotated as RBPs. For each cluster, we classified detected ECOD domains and quantified K-locus diversity via the Simpson index, which reflects the variation of K-loci associated with bacteria in which these proteins were found. We reasoned that capsule-specific RBPs should predominantly occur in genomes sharing the same K-locus.

The association between ECOD domains and K-locus diversity varied substantially across functions. Only four ECOD domains were linked to protein clusters predominantly associated with a single K-locus or few K-loci: *Phage tail fiber protein trimerization domain* (T-level: 79.1.1), *Intramolecular chaperone domain in virus tail spike protein* (T-level: 3240.1.1), *Pectin lyase-like* (T-level: 207.2.1), and *SGNH hydrolase* (T-level: 2007.5.1; see Figure 6A). Among these, only the *Phage tail fiber protein trimerization domain* and the *Pectin lyase-like* domain were predominantly found in clusters with low K-locus diversity, suggesting a stronger association with capsule specificity. In contrast, the *Intramolecular chaperone domain in virus tail spike protein* (T-level: 3240.1.1), *gp11/gp12 receptor-binding domain* (T-level: 877.1.1), and *Uncharacterized protein CV0426* (T-level: 3809.1.1) tended to occur in clusters spanning multiple K-loci, indicating broader distribution. Notably, the *SGNH hydrolase* domain exhibited the greatest variability in K-locus diversity, and on average, was less K-locus-specific than the *Pectin lyase-like* domain.

To investigate how the frequency of RBP domains varies between different K-loci, we mapped the ECOD domains from Figure 6A on the high-quality prophages found in bacterial genomes with various K-loci (see Figure 6B). Most of the RBPs detected carried the 'SGNH hydrolase' domain, while the 'Pectin lyase-like' domain was on average less prevalent, usually co-occurring with other domains (see Supplementary Figure S6). Furthermore, we found examples where while no domains were detected in prophage RBPs, adjacent proteins contained hits to 'SGNH hydrolase' (see Supplementary Figure S7). Altogether, these results highlight the structural diversity of RBPs found in *K. pneumoniae* prophages and point to the prevalence and importance of alternative capsule recognition mechanisms like SGNH hydrolases in interactions between bacteria and their temperate phages.

Figure 6. SGNH hydrolases are prevalent amongst across *K. pneumoniae* prophage RBPs. (A) Capsule-type (K-locus) diversity within RBP clusters, coloured by their dominant ECOD domain. Prophage proteins from four *Klebsiella* species were clustered at ⩾50% identity and ⩾80% coverage, then RBPs were identified by PHROG annotation (tail fiber or tail spike; probability ⩾90%, qcov ⩾ 50%, scov ⩾ 50%). ECOD domains were assigned to each RBP cluster (probability ⩾ 90%, tcov ⩾ 10%), allowing multiple domains per cluster. Each point shows a cluster with ⩾10 members from ⩾3 SCs, its Simpson index of K-locus diversity (0 = uniform, 1 = maximally diverse), and its ECOD domain. (B) Number of high-quality, nearly complete prophages encoding each ECOD domain within PHROG-defined RBPs, stratified by K-locus (completeness of ⩾90%, $n =$4,598, of which $n =$2,047 belonged to the 35 most diverse K-loci). Bars are coloured by the single ECOD domain detected (or light grey if no PHROG RBP was detected, dark grey if multiple ECOD domains co-occur). The most frequent multi-domain combinations include gp11/gp12 + SGNH hydrolase and pectin-lyase-like + Orf210 N-terminal or gp11/gp12 domains (see Supplementary Figure S6). Prophages whose RBPs lack any of the domains shown in (A) often encode adjacent proteins with other enzymatic folds (e.g., cysteine proteases, triphosphate hydrolases, SGNH hydrolases; Supplementary Figure S7).

Discussion

In this work, we have set out to uncover molecular determinants of capsular specificity in temperate phages of *K. pneumoniae*. To this end, we carried out a genome-wide association study (GWAS) on an ecologically diverse dataset of 3,900 bacterial genomes, looking for genetic predictors from their 8,105 prophage regions that correlate with the K-type predictions (K-loci). We expected to uncover a repertoire of diverse β-helix-containing depolymerases that may have escaped classical functional prediction. Instead, our analysis points to several unexpected results. First, regardless of the used classifier, GWAS did not return a result for about 30% of K-loci. Second, even amongst the best GWAS predictors, only about a half are those with the expected parallel β-helix, the second large class being proteins with the 'SGNH hydrolase' domain family, but also RBPs with previously unseen folds and non-RBP predictions like putative acetyltransferases. Third, a manual search for β-helix-containing depolymerases in a representative subset of 99 bacterial isolates and recombinant production of predicted depolymerases showed that only 14 out of 50 of them were shown to be active enzymes on a panel of bacterial isolates with 119 different K-types, while for 34 proteins expression was unsuccessful; of the 14 active ones, 5 targeted a serotype different from that of their original host (possibly due to a recent capsule switch via recombination). Furthermore, we observed strong agreement between GWAS predictions and experimental validation of enzymatic activity for receptor-binding proteins (RBPs), with only two exceptions: (i) KL14, where the protein was lacking a C-terminus, an integral part of the receptor-binding domain, and (ii) KL10, which was successfully produced as a soluble protein without enzymatic activity after both manual search and GWAS prediction. Finally, two RBPs with the SGNH hydrolase domain (KL2 and KL55) had strong GWAS predictions and were prepared as recombinant proteins but no depolymerase activity was detected in the available K-type strain collection.

Despite using an ecologically diverse dataset of bacteria and their temperate prophages, we did not detect strong GWAS hits for several important K-loci, including KL1. This absence could not be explained by insufficient data quality, as median prophage completeness was consistently near 100% (see Supplementary Figure S1), nor by capsule locus switching, which–although previously demonstrated [49,27]–is thought to occur at rates lower than phage transmission and evolution, and thus would be insufficient to fully obscure the statistical association between prophage proteins like RBPs and specific K-types. Instead, we think that the primary explanation for the weak or absent GWAS predictions is the genetic and functional

diversity of the RBPs carried by prophages. As seen by the example of K1 (see Supplementary Figure S8), prophages in bacteria with this K-type carry diverse RBPs that may not be β-helix-containing depolymerases, i.e., with the 'SGNH hydrolase' domain as well as other, unusual RBPs. The large genetic diversity of proteins with shared functional domains emphasises the importance of structural and functional characterisation of RBPs from an ecologically diverse pool of bacterial viruses.

The relatively high frequency of the 'SGNH hydrolase' domain among GWAS predictions underscores its importance as a host-range determinant in temperate phages of *K. pneumoniae*. This prevalence is also reflected by the high proportion of RBPs with this domain across all prophages in our dataset (cf., Figure 6). The 'SGNH hydrolase' domain, encompassing a diverse family of esterases and lipases, has previously been identified as an essential structural topology in RBPs of other phages[50], including those infecting *Acinetobacter baumannii*[51] and *E. coli*[52]. Specifically, Prokhorov and colleagues demonstrated that gp63.1, an RBP from an *E. coli* phage G7C, containing the SGNH domain flanked by two carbohydrate-binding modules. This SGNH domain was shown to be an esterase functioning as a deacetylase, removing an O3-acetyl group from lipopolysaccharide without degrading the O-chain[53] – a reaction essential for G7C's infectivity. Although the RBPs identified in our study are structurally different from gp63.1, all of them contain the SGNH enzymatic domain and lack the right-handed parallel β-helix, suggesting that SGNH-containing RBPs in *K. pneumoniae* phages might similarly rely on polysaccharide acetylation for their activity. Our results thus suggest that deacetylation may represent a common alternative strategy of temperate *Klebsiella* phages to recognise and "shave" polysaccharides from the acetyl groups (without depolymerisation) to initiate infection.

The idea that temperate phages of *K. pneumoniae* may often rely on deacetylation to initiate infection is consistent with the fact that O-acetylation of bacterial polysaccharides is common, for which there are multiple lines of evidence. O-acetylation has been detected in many K-types, either as part of the defined repeating-unit structure (e.g., K5, K21b, K22, K33, K54, K55, K58, K59, K82)[54,55,56,57,58,59,60,61,62] or observed in NMR spectra without precise localisation (e.g., K4, K8, K10, K11, K13, K20, K24, K30, K41, K43, K44, K49, K63, K64, K69)[63,64,65,66,67,68,69,70,71,72,73,74,75,76,77]. Likewise, K-loci in *K. pneumoniae* often feature genes annotated as acetyltransferases[17]. Incidentally, we also detected putative acetyltransferases in two of our strongest GWAS hits—despite filtering out short genes—and between 5–30% of prophages per K-locus in our dataset encode proteins with a predicted acetyltransferase function (see Supplementary Figure S9), suggesting that sugar modification may be driven – at least partially –

by incoming prophages. As the presence or absence of an acetyl group in otherwise identical or nearly-identical polysaccharide structures can be associated with different serological assignment, as in the case of K22/K37[78,56] or K30/K33[57,70], it is plausible that current serotyping and K-locus genotyping underestimate capsule diversity, as O-acetylation introduces biochemical variation that can affect phage recognition, host tropism and immune interactions beyond what is captured by these schemes.

Another interesting point is that, if the dynamic interplay between acetylation and deacetylation shapes capsule tropism of bacterial viruses, this may reflect both a direct mechanism of host recognition by temperate phages and a competitive strategy to modify host surfaces in ways that interfere with other phages, particularly those that rely on unmodified polysaccharides. Similar mechanisms have been reported across Gram-negative bacteria, where O-acetylation of surface polysaccharides alters phage susceptibility: in *E. coli* and *Salmonella typhimurium*, it modulates sensitivity to superinfecting phages[79,80]; in *Shigella flexneri*, temperate phage-mediated O-acetylation drives serotype conversion and can confer phage resistance[81,82]; and in *Pseudomonas aeruginosa*, phage D3 introduces O-acetyl modifications to prevent superinfection by related phages[83,84]. While in *K. pneumoniae* some depolymerases may tolerate acetylated substrates[85], others may not, suggesting that these subtle modifications could significantly reshape the phage infectivity landscape. Overall, these findings highlight polysaccharide acetylation and deacetylation as a potentially underappreciated layer of viral specificity and phage–phage competition (e.g., via superinfection exclusion mechanisms) in *K. pneumoniae*.

Our study also underscores a critical caveat in the growing enthusiasm for mining prophages as a source of biotechnological innovation. While prophages embedded in sequenced bacterial genomes provide a rich and accessible reservoir of genetic diversity—often more complete and better annotated than phages derived from metagenomes—our findings suggest that not all such sequences represent functional or biologically active proteins. Among the 50 candidate depolymerases we selected for recombinant production, 34 failed to produce soluble proteins. This proportion suggests that recombinant protein expression of depolymerases from prophages may be lower than this obtained from virulent phages. This would be in line with the expectation that many prophages are no longer functional: once a prophage becomes cryptic and loses the ability to enter the lytic cycle, its genes may no longer be subject to purifying selection[86]. This allows mutations to accumulate, potentially rendering key effector proteins non-functional. As a result, sequences mined from such "dead" proviruses may reflect evolutionary debris rather than viable biological tools. Therefore, while prophages are a valuable source of novel proteins, our results highlight

the importance of coupling computational discovery with careful validation and caution when interpreting the functional potential of prophage-encoded enzymes.

One important limitation of this study is that we restricted our GWAS to an ecologically diverse dataset of *Klebsiella* genomes with reliable K-locus predictions. While this strategy supports biological relevance, it limits statistical power by excluding rarer K-types and less well-characterised species. Additionally, GWAS results are inherently sensitive to various parameters—including protein choice of genetic variant, K-locus prediction or population structure correction—and we anticipate that tuning these parameters further, combined with increased sample size, would yield more associations. Another key limitation is that our experimental validation focused on depolymerase activity detectable via halo formation. As a result, we may have missed RBPs with enzymatic activity that do not produce visible halos or operates via alternative interaction mechanisms. Finally, while our findings highlight unexpected complexity in capsule recognition among temperate *Klebsiella* phages, it remains unclear to what extent these insights generalise beyond this species or to other phage lifestyles such as strictly lytic (virulent) phages.

In summary, our combined GWAS and experimental screen in a near-random sampling of *K. pneumoniae* prophages demonstrates that, although classical right-handed β-helix depolymerases do indeed mediate capsule tropism in temperate phages, they are far from the sole mechanism at play. SGNH hydrolase–driven deacetylation emerges as a likely and functionally dominant strategy for capsule recognition and modifying the capsule, revealing a previously unappreciated layer of molecular complexity. These findings uncover hidden diversity in polysaccharide modifications across *K. pneumoniae* populations, and carry direct implications for designing more effective phage therapeutics and capsule-based vaccines. More broadly, our work underscores the need to integrate population-scale genomics, structural modeling, and rigorous functional assays to fully capture the evolutionary innovations that temperate phages employ to expand and refine their host range.

Methods

Computational approach

Data

To build a representative genome dataset of *Klebsiella spp.*, we combined two collections. The first collection (KASPAH) consists of 391 *Klebsiella pneumoniae* species complex isolates (*K. pneumoniae*, *K. quasipenumoniae*, *K. variicola*), obtained from a diagnostic laboratory serving four major hospitals in Melbourne between April 2013 and March 2014[35,36,33]. These isolates were sourced from various clinical samples including pneumonia ($n = 44$), rectal swabs ($n = 82$), throat swabs ($n = 14$), urinary tract infections (UTI; $n = 193$), wound or tissue ($n = 28$), disseminated infections ($n = 25$), and other sources ($n = 5$). Sequencing was performed using short-read Illumina technology all isolates, while 99 were additionally sequenced using long-read Oxford Nanopore Technology (ONT) to generate hybrid short/long-read finished assemblies[37]. The second collection (KLEBPAVIA) encompasses 3,510 *Klebsiella spp.* isolates chosen from 6,548 samples from diagnostic laboratories in Pavia, Italy, and its surroundings between June 2017 and November 2018[34]. This dataset includes 3482 *Klebsiella spp.* isolates, with an additional 27 *K. quasipneumoniae* isolates not published in Thorpe et al. 2022, as well as a single outgroup isolate originally classified as *Klebsiella* but later identified as *Superficieibacter maynardsmithii*[87]. The full dataset consists of 3,900 isolates of *Klebsiella spp.* and a single outgroup. Importantly, the combined collection consists of diverse *Klebsiella* isolates from multiple ecological niches, hence providing a diverse overview of both capsule and prophage diversity within. All genomes were annotated with the PATRIC server between November 2022 and March 2023 and with the recipe Bacteria / Archaea and Taxonomy ID 573 (*K. pneumoniae*). The list of bacterial genomes is provided in Supplementary Table S2.

Inference of the population structure

Core genes were defined on the whole dataset (including the outgroup) using panaroo v1.3.4[88] on 'strict' mode, as genes present in at least 99% of isolates. SNPs were extracted from an alignment of 1668 core genes with snp-sites v2.5.1[89]. The maximum likelihood phylogenetic tree was inferred from those SNPs using IQ-TREE v2.1.4[90] with the following parameters: `-t PARS -ninit 2` and the number of constant sites obtained from snp-sites `-C`. The best-fitting model of sequence evolution was identified by IQTREE ModelFinder as GTR+F+I+R10. Independently, we ran Kleborate v2.4.1[91] to predict *Klebsiella* species

for all 3,900 isolates, yielding 19 different species: *K. pneumoniae, K. variicola, K. quasipneumoniae subsp. quasipneumoniae, K. quasipneumoniae subsp. similipneumoniae, K. oxytoca, K. michiganensis, K. ornithinolytica, K. aerogenes, K. pasteurii, K. grimontii, K. planticola, K. quasivariicola, K. spallanzanii, K. huaxiensis, K. Ka3, K. terrigena, K. Ko10, K. huaxiensisNEW* and *K. Kterr4*. We then used PopPunk v2.6.3 to identify bacterial genome sequence clusters (SCs) within each species, which were used as a proxy for bacterial lineages. Four of these species (*K. Ka3, K. Ko10, K. huaxiensisNEW, K. Kterr4*) are outgroups of other species (respectively of *K. aerogenes, K. huaxiensis, K. huaxiensis* and *K. quasiterrigena*) and were introducing noise in the clustering, hence were not considered for this analysis. For the remaining 3,876 genomes assigned to 15 species, we carried out the same approach as described in Thorpe et. al. 2022[34]. Briefly, for each species we fit the mixture model of PopPunk v2.6.3[38]. The number of components (k) was chosen based on the scatter plot of core and accessory distances and iteratively improved. We then ran the model refinement module of PopPunk for each species individually. However, in the case of two species (*K. spallanzanii* and *K. quasivariicola*) the refinement introduced noise in the clustering and hence was abandoned. We compared the resulting clusters to those from Thorpe et. al. with the Adjusted Rand Index (ARI) using python `sklearn.metrics.adjusted_rand_score`, for which a value of 1 indicates perfect agreement between the two clusterings and 0 indicates random labelling. We compared the resulting trees with microreact for species with ARI < 0.8, all of them appearing coherent despite the discrepancies. The resulting PopPunk clusters are henceforth referred to as bacterial sequence clusters (SCs). Code snippets to run for each species (replacing 'klebsiella' by appropriate species, adjusting K value and optionally running or not the refine command).

K-locus identification

We used Kaptive v2.0[17] implemented in Kleborate v2.4.1[91] to predict capsule polysaccharide synthesis loci (K-loci) for all isolates, which are predictive of capsular types (K-types). Species and K-loci were predicted with Kleborate for all isolates. To identify *Klebsiella* species for which K-locus prediction is reliable, we searched for the species in which 80% of isolates or more had a confident K-locus prediction (i.e., assigned 'Good', 'Very Good' or 'Perfect' by Kaptive) and which had more than 5 members. The resulting dataset consisted of 2,527 genomes of isolates belonging to four species from the *K. pneumoniae* species complex (*K. pneumoniae, K. variicola subsp. variicola, K. quasipneumoniae subsp. quasipneumoniae, K. quasipneumoniae subsp. similipneumoniae*), henceforth referred to as the GWAS dataset.

Identification of prophages

To detect prophages in bacterial genomes, we used two complementary tools: VirSorter v1.0.5[39] and PhiSpy v4.2.6[40]. VirSorter predicts clusters of viral genes based on sequence similarity to its database (`-db 1`), though it frequently over-predicts the length of prophage regions. PhiSpy, which combines heuristic methods, including protein length, transcription strand directionality or AT/GC skew, tends to predict shorter prophage regions and may capture other mobile genetic elements. As the goal was to minimise the false-negative rate of prophage prediction to consider as many likely viral proteins in the GWAS approach, we took the union of the prophage regions predicted by both tools, creating a primary detection. Manual investigation of the resulting putative viral regions indicated an occasional presence of poorly annotated receptor-binding proteins. Hence, we extended each primary detection by 2 kb upstream and downstream, forming an extended detection. Following this, we applied CheckV v1.0.1[41] to further refine our prophage predictions. CheckV not only splits merged prophages within extended regions but also trims over-predicted regions while estimating prophage completeness and confidence. To account for a potential loss of receptor-binding proteins during this trimming, we again extended the CheckV-defined prophage boundaries by 2 kb upstream and downstream. We retained only prophages with a completeness of at least 50%, confidence rated as low, medium or high, and a minimum length of 2kb. This resulted in a dataset of 8,103 prophages. The mean number of detected prophages per genome was 3.2, with a maximum of 14 prophages found in a single genome. Of those, 4,598 prophages were of high-completeness ($\geqslant 90\%$), which yields 1.8 high-quality prophage per isolate. We did not detect any prophages in 49 bacterial genomes. The prophage detection workflow is implemented in Snakemake v8.10.0 and is accessible on GitHub at `https://github.com/bioinf-mcb/mgg_prophages`.

Functional annotation

To functionally annotate predicted prophages in bacterial genomes, we predicted Open Reading Frames (ORFs) prodigal-gv (v2.11.0)[92] which were clustered MMseqs2 v13.45111[93] with sensitivity `-s 7.5`, using criteria of at least 80% coverage between query and subject sequences and a minimum of 50% amino acid identity, forming clusters (PC80) to streamline downstream analysis. Next, we aligned these protein clusters against the PHROGs database[42] with HHblits[94] using parameters: `-n 2 -cov 0 -p 0.95`. These alignments were then converted into HMM profiles and searched against the HMM ECOD70 database (version F70_20200207) using HHsearch, retaining hits with a probability of 70% or higher. Within each

PC80, we identified all unique ECOD X-levels, selecting the highest-scoring hit within each level, and reported up to five distinct ECOD T-levels per PC80 cluster based on the highest probability. Finally, we mapped predicted structural topologies onto the proteins to provide a comprehensive functional annotation. Prophage functional annotation is accessible as Snakemake v 8.10.0 workflow on GitHub repository under: `https://github.com/bioinf-mcb/mgg_annotation`.

Genome-wide association studies

To identify statistical associations between bacterial polysaccharide capsules (determined by the predicted K-locus) and protein clusters found in prophages, we conducted a genome-wide association study (GWAS) independently of the functional annotation. Protein clusters were generated by selecting proteins with a minimum length of 300 amino acids and clustering them using MMseqs. We used six different clustering levels, using combinations of identity thresholds (none, 50%, 80%) and coverage thresholds (50%, 80%), with the 50/50 parameter combination by default (PC50 clusters). Rare variants were filtered out by retaining only protein clusters found in at least three bacterial isolates from two different sequence clusters (SCs), resulting in 1,690 PC50 clusters out of an initial set of 3,745 clusters.

GWAS was run separately for each of the 35 K-loci which were found in at least 10 SCs. For each K-locus, we ran pyseer v1.3.11[95], which is an implementation of the elastic-net approach[96], which combines a ridge regression and lasso regression, with a parameter alpha which determines the amount of mixing between the two penalties, such that $\alpha = 0$ results in ridge regression and $\alpha = 1$ results in lasso regression. We used SCs as covariates in the model to control for population structure, under the assumption that capsule tropism in prophages is primarily shaped by recent horizontal acquisitions rather than deep shared ancestry; this is supported by evidence that most prophages in bacterial genomes are relatively recent acquisitions[86,27].

Given the biological nature of the polysaccharide-RBP specificity problem, we a priori expected that a given K-type would be recognised by a single enzyme family. In this situation, we would expect the associations to be determined by a small number of predictors (ideally a single protein cluster), a situation better suited for lasso-regression which eliminates most weak predictors. However, we also allowed the possibility of multiple predictors due to the potential complexities of the analysis, for example high diversity and mosaicism of RBPs resulting in multiple sequence clusters weakly correlating with the phenotype – a situation better suited for ridge-regression which allows the presence of multiple statistical

predictors. Hence, we ran pyseer in two modes, with the first being lasso-dominated ($\alpha = 0.8$) and the second being ridge-dominated ($\alpha = 0.069$). The inputs to pyseer included: (1) a binary matrix for KL types, indicating the presence or absence of the K-locus in question in all 2,527 bacterial isolates; (2) a binary matrix for the presence or absence of each of the 1,690 prophage PC50 clusters in the same isolates; and (3) a mapping of each bacterial isolate to its SC.

The model outputs a regression coefficient (β) and p-value for each protein cluster, quantifying its statistical association with the K-locus. P-values were Bonferroni-corrected (p-value of $0.05/N$, where $N = 1690$ is the number of PC50 protein clusters). While a high β value indicates strong association, visualisation of these associations across the bacterial phylogeny using Phandango v1.3.1 [46] revealed that strong statistical associations do not necessarily translate to accurate predictions, especially in imbalanced datasets where certain K-loci are relatively rare. We therefore considered all protein clusters with $\beta > 0$ and p-value below the significant threshold as predictors, and evaluated their classification performance for each K-locus separately on the full dataset using precision, recall, F1 score, and Matthews correlation coefficient (MCC):

- **Precision** measures specificity:
$$\text{Precision} = \frac{\text{TP}}{\text{TP} + \text{FP}}$$

- **Recall** measures sensitivity:
$$\text{Recall} = \frac{\text{TP}}{\text{TP} + \text{FN}}$$

where:

- **True Positive (TP)**: Protein variant present in an isolate with the corresponding K-locus.

- **False Positive (FP)**: Protein variant present in an isolate without the corresponding K-locus.

- **False Negative (FN)**: Protein variant absent from an isolate with the corresponding K-locus.

- **True Negative (TN)**: Protein variant absent from an isolate without the corresponding K-locus.

Additional metrics included:

- **F1 score** balances precision and recall:

$$\text{F1} = 2 \times \frac{\text{Precision} \times \text{Recall}}{\text{Precision} + \text{Recall}}$$

- **Matthews Correlation Coefficient (MCC)** provides a comprehensive measure of prediction quality, incorporating all four confusion matrix outcomes (TP, FP, TN, FN):

$$\mathrm{MCC} = \frac{\mathrm{TP} \times \mathrm{TN} - \mathrm{FP} \times \mathrm{FN}}{\sqrt{(\mathrm{TP} + \mathrm{FP})(\mathrm{TP} + \mathrm{FN})(\mathrm{TN} + \mathrm{FP})(\mathrm{TN} + \mathrm{FN})}}$$

Precision was particularly critical because we aimed to identify highly specific enzymes targeting distinct polysaccharides. A high precision maximises reliability of predictions, even if this occasionally reduces recall. To assess statistical uncertainty of precision and recall, we performed $n = 100$ bootstrap replicates for each K-locus, resampling bacterial isolates with replacement to recalculate TP, FP, and FN values. We derived 95% confidence intervals from the distribution quantiles of these bootstrapped estimates.

Finally, for each PC50 hit selected from the GWAS analysis, we predicted the functions of its sequence members by mapping them to their corresponding PC80 clusters, where functional annotations had previously been assigned. Since different sequence members within the same PC50 cluster could be associated with different PC80 clusters, each member might carry distinct ECOD T-level annotations (or combinations thereof, given the multi-domain nature of phage RBPs). To provide a single functional prediction for each PC50 cluster, we selected the most frequent combination of ECOD T-levels among its sequence members. Additionally, we independently checked whether any sequence member in the PC50 cluster was annotated with the *Pectin lyase-like fold* (ECOD T-level: 207.2.1) in any PC80 cluster. If this annotation was present, we labelled the PC50 cluster as a putative pectin lyase fold.

Experimental approach

Depolymerase prediction

All prophages detected in the KASPAH-REF dataset were considered if they were classified as high or medium completeness (CheckV criteria; see above). Predicted prophage proteins were clustered using MMseqs2 with parameters set to 90% bidirectional sequence coverage and 50% sequence identity. Functional annotations of protein clusters were predicted by searching representative cluster sequences against the PHROGs and UniClust30 databases using HHblits with default parameters (UniRef30 database, E-value cutoff $= 10^{-3}$, minimum sequence coverage $= 20\%$). A protein cluster was classified as a putative depolymerase candidate if it met the following criteria: (1) the average length of proteins in the cluster was at least 500 amino acids; (2) the cluster was functionally annotated as either 'tail fiber/fibre' or 'tail

spike'; (3) proteins contained predicted enzymatic domains characteristic of polysaccharide-degrading enzymes—specifically lyases (e.g., hyaluronidases, pectin/pectate lyases, alginate lyases, K5 lyases) or hydrolases (e.g., sialidases, rhamnosialidases, levanases, dextranases, or xylanases).

Domain predictions were carried out by comparing representative sequences against multiple databases, as follows: (a) BLASTp searches against the non-redundant protein sequence (nr) database, using standard parameters (expect threshold = 0.05; word size = 5; matrix = BLOSUM62; gap costs: existence = 11, extension = 1; conditional composition score matrix adjustment); (b) HMMER searches against Reference Proteomes (quick search mode; significance thresholds: sequence E-value = 0.01, hit E-value = 0.03); and (c) HHpred searches against the PDB_mmCIF70 database (HHblits MSA generation from UniRef30; maximum MSA iterations = 3; MSA generation E-value cutoff = 10^{-3}; minimum sequence identity = 0%; minimum coverage = 20%; secondary structure scoring: during alignment; alignment mode: realign with MAC [local:norealign], MAC realignment threshold = 0.3; number of target sequences = 250; minimum probability in hitlist = 20%).

Using a combination of the two approaches, a total of 469 proteins meeting all selection criteria were identified. These proteins were clustered at high coverage using MMseqs2 (80% coverage and 50% identity), obtaining 124 representative protein sequences. These sequences were then structurally modelled using AlphaFold2[47] in the monomer mode. The resulting models were screened for conserved structural features, with particular emphasis on the presence of a parallel β-helix fold characteristic of depolymerases. Structural analysis led to the identification of 50 proteins from 32 clusters, with 22 clusters yielding a single representative and 10 clusters yielding multiple representatives to capture sequence and domain-level variation within those groups (altogether 28). The complete list of these proteins is given in Supplementary Table S5, and their genomic coordinates are provided in Supplementary Table S6.

Bacterial Strains

K. pneumoniae strains used in this study (Supplementary Table S4) originate from collections of the Collection de l'Institut Pasteur (CIP), Paris, France, the National Collection of Type Cultures (NCTC), the UK Health Security Agency (UKHSA), collection of the Department of Pathogen Biology and Immunology (Wrocław, Poland), and the *Klebsiella* Acquisition Surveillance Project at Alfred Health (KASPAH) from Melbourne, Australia[35,36,33]. Bacteria were grown in Tryptone Soya Broth or Agar (TSB or TSA, Oxoid, Thermo Fisher Scientific, Waltham, MA, USA) at 37°C. The collection includes 134 strains rep-

resenting 119 distinct serotypes.

2   The *Escherichia coli* strains (Thermo Fisher Scientific, Waltham, MA, USA) used recombinant protein production: One ShotTM TOP10 for cloning and One ShotTM BL21 StarTM (DE3) for recombinant

4   protein expression. They were cultured in LB broth, Miller (Luria-Bertani) (Difco, Becton, Dickinson and Company (BD), Franklin Lakes, NJ, USA) at 37 °C supplemented with 100 µg/mL ampicillin (IBI

6   Scientific, Dubuque, IA, USA).

All bacteria were stored at -70°C in a 20% glycerol-supplemented Trypticase Soy Broth (TSB, Becton

8   Dickinson, and Company, Cockeysville, Md, USA).

Cloning, expression and purification

The genomic DNA from bacterial strains containing prophages encoding putative depolymerase was extracted using a Pure Link Genomic DNA Mini Kit (Thermo Fisher Scientific, Waltham, MA, USA)

according to the manufacturer's protocol. The genes were amplified by PCR using the specific primer pairs (Genomed, Warsaw, Poland) and Platinum SuperFi DNA polymerase (Thermo Fisher Scientific,

Waltham, MA, USA). The desired products were subsequently cloned into the Champion pET101 expression vector (Thermo Fisher Scientific, Waltham, MA, USA) with a C-terminal His tag (6x) and

propagated in *E. coli* One ShotTM TOP10 strain according to the manufacturer's protocol. Plasmid DNA was then purified and the accuracy of the clones was confirmed by sequencing (Genomed, Warsaw,

Poland). Following the transformation of the correct constructs into *E. coli* One ShotTM BL21 StarTM (DE3) strain, vector-bearing cells were cultured in 500 mL of LB broth supplemented with 100 µg/mL

ampicillin at 37 °C with agitation until reaching an optical density (OD600nm) of 0.5-0.6. The recombinant protein expression was conducted for 16 h at 18°C using isopropyl-b-D-1-thiogalactopyranoside

(IPTG; the final concentration of 1.0 mM) as an inductor of the expression. Next, the cultures were collected by centrifugation (8000 x g, 15 min, 4 °C), and the pellets were resuspended in 15 ml of lysis

buffer (500 mM NaCl, 20 mM NaH2PO4, pH 7.4). Following 3 freeze/thaw cycles and sonication, the obtained lysates were centrifuged (16000 x g, 30 min, 4 °C) and the supernatants were filtered through a

0.22 µm filter (Sarstedt AG % Co., Numbrecht, Germany). The purification of His-tagged proteins was performed independently for each protein using an FPLC system (Bio-Rad, Hercules, CA, USA) com-

bined with Bio-Scale Mini Profinity IMAC cartridges (Bio-Rad, Hercules, CA, USA). First, the column equilibration step was performed with lysis buffer (500 mM NaCl, 20 mM NaH2PO4, pH 7.4), then the

lysate was loaded and washed with 32 volumes of washing buffer. Next, the protein was eluted by using a

lysis buffer supplemented by 500 mM imidazole. After purification, protein concentration was measured spectrophotometrically (NanoPhotometer NP80, Implen GmbH, Schatzbogen, München, Germany) using a molar extinction coefficient calculated by ProtParam[97] for each protein. The quality of the purified proteins was analysed by a sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using a 12% polyacrylamide gel and Precision Plus Protein All Blue molecular weight standard (Bio-Rad, Hercules, CA, USA). In addition to employing standard cloning and expression techniques, the production of selected depolymerases (see Supplementary Table S5) were outsourced to a commercial service provider (GenScript, China). The genes encoding all the aforementioned proteins were synthesized with codon optimization for expression in *E. coli*, cloned into pET30a expression vector, and prepared by the company as highly purified protein preparations, each bearing an N-terminal 6×His tag.

Enzymatic functionality assays

Enzymatic activity was assessed using a spot assay on *K. pneumoniae* serotype collections listed in Supplementary Table S4. Bacterial cultures were grown to an optical density at 600nm $\approx$ 1.0 and 0.5 mL was spread evenly onto TSA plates. After drying, 10 µL of each recombinant enzyme was spotted on the bacterial lawn. As a negative control, 10 µL of a sample obtained from the expression and purification of an empty vector—processed using the same protocol as for the recombinant proteins was also applied. Following overnight incubation at 37 °C and an additional 24 h incubation at room temperature, plates were examined for the presence of halo zones. The minimum concentration of protein required to produce a detectable halo on the bacterial lawn (minimal halo-forming concentration; MHFC) was established using the method described previously[98].

Supplementary Material

Supplementary Tables

- S1: List of 58 active depolymerases from previously published studies on virulent phages, based on Cheetham and colleagues[21].

- S2: List of bacterial genomes used, together with the associated metadata.

- S3: List of prophage genomes detected in S1, together with the associated metadata.

- S4: List of 119 *K. pneumoniae* strains from the reference K-type collection used in this study.

- S5: List of experimentally tested, recombinantly produced proteins.

- S6: Map of proteins from Supplementary Table S3 onto prophages from Supplementary Table S2.

Supplementary Texts

- S1: Methodology and results of the relaxed GWAS approach to select β-helix-containing depolymerase predictions.

- S2: All experimental results.

- S3: Sequence and structural comparison of the GWAS predictions (Text S1) and experimental results (Text S2) based on Figure 4B.

Reproducibility

Data

All input data for this project, including genome assemblies, are provided under the following link

- https://doi.org/10.6084/m9.figshare.29181188

These data can be used as input for the data analysis pipeline which can be accessed under the following Github release:

- https://https://github.com/bioinf-mcb/mgg_gwas

The essential output of this pipeline used to generate the figures for the publication can be accessed under the following link

- `https://doi.org/10.6084/m9.figshare.29224373`

and the pipeline to generate the figures can be accessed under the following Github release:

- `https://github.com/rmostowy/klebs-gwas-figs`

Funding

This work was financed by the Polish National Agency for Academic Exchange (JK, BS, RM), the EMBO Installation Grant (JK, VRP, JL, BS, RM), the Polish National Science Centre Sonata Bis Grant (grant 2020/38/E/NZ8/00432; AO, VRP, JL, SO, BS, BM, RM). We gratefully acknowledge Polish high-performance computing infrastructure PLGrid (HPC Centers: CI TASK, ACK Cyfronet AGH) for providing computer facilities and support within computational grants no. PLG/2022/015493, PLG/2022/015858, PLG/2023/016392, PLG/2024/017293, PLG/2025/018394.

Author contributions

The following contributions can be distinguished by different authors based on the CRediT (Contributor Roles Taxonomy) system:

- **Conceptualization**: ZDK, RJM
- **Methodology**: JK, BM, ZDK, RJM
- **Software**: JK
- **Validation**: AO, JK, BS, RJM
- **Formal Analysis**: AO, JK, VRP, JL, SO, BS, RJM
- **Investigation**: AO, JK, VRP, EPCR, BS, BM, ZDK, RJM
- **Resources**: KEH, EJF, ZDK, RJM
- **Data Curation**: JK, JL
- **Writing – Original Draft**: BS, ZDK, RJM
- **Writing – Review & Editing**: AO, JK, VRP, JL, SO, KEH, EJF, EPCR, BS, BM, ZDK, RJM
- **Visualization**: AO, JK, BS, RJM

- **Supervision**: BM, ZDK, RJM
- **Project Administration**: BM, ZDK, RJM
- **Funding Acquisition**: RJM

References

[1] Strobel HM, Stuart EC, Meyer JR. A Trait-Based Approach to Predicting Viral Host-Range Evolvability. Annual Review of Virology. 2022 Sep;9(1):139–156.

[2] Tan CCS, van Dorp L, Balloux F. The evolutionary drivers and correlates of viral host jumps. Nature Ecology & Evolution. 2024 May;8(5):960–971.

[3] Meyer JR, Dobias DT, Weitz JS, Barrick JE, Quick RT, Lenski RE. Repeatability and contingency in the evolution of a key innovation in phage lambda. Science (New York, NY). 2012 Jan;335(6067):428–432.

[4] Piel D, Bruto M, Labreuche Y, Blanquart F, Goudenège D, Barcia-Cruz R, et al. Phage-host coevolution in natural populations. Nature Microbiology. 2022 Jul;7(7):1075–1086.

[5] Göller PC, Elsener T, Lorgé D, Radulovic N, Bernardi V, Naumann A, et al. Multi-species host range of staphylococcal phages isolated from wastewater. Nature Communications. 2021 Nov;12(1):6965.

[6] Kauffman KM, Chang WK, Brown JM, Hussain FA, Yang J, Polz MF, et al. Resolving the structure of phage-bacteria interactions in the context of natural diversity. Nature Communications. 2022 Jan;13(1):372.

[7] Strathdee SA, Hatfull GF, Mutalik VK, Schooley RT. Phage therapy: From biological mechanisms to future directions. Cell. 2023 Jan;186(1):17–31.

[8] Skurnik M, Alkalay-Oren S, Boon M, Clokie M, Sicheritz-Pontén T, Dąbrowska K, et al. Phage therapy. Nature Reviews Methods Primers. 2025 Feb;5(1):9. Available from: https://www.nature.com/articles/s43586-024-00377-5.

[9] Holtappels D, Alfenas-Zerbini P, Koskella B. Drivers and consequences of bacteriophage host range. FEMS microbiology reviews. 2023 Jul;47(4):fuad038.

[10] de Jonge PA, Nobrega FL, Brouns SJJ, Dutilh BE. Molecular and Evolutionary Determinants of Bacteriophage Host Range. Trends in Microbiology. 2019 Jan;27(1):51–63.

[11] Tesson F, Hervé A, Mordret E, Touchon M, d'Humières C, Cury J, et al. Systematic and quantitative view of the antiviral arsenal of prokaryotes. Nature Communications. 2022 May;13(1):2561.

[12] Lood C, Boeckaerts D, Stock M, De Baets B, Lavigne R, van Noort V, et al. Digital phagograms: predicting phage infectivity through a multilayer machine learning approach. Current Opinion in Virology. 2022 Feb;52:174–181.

[13] Boeckaerts D, Stock M, Ferriol-González C, Oteo-Iglesias J, Sanjuán R, Domingo-Calap P, et al. Prediction of Klebsiella phage-host specificity at the strain level. Nature Communications. 2024 May;15(1):4355.

[14] Gaborieau B, Vaysset H, Tesson F, Charachon I, Dib N, Bernier J, et al. Prediction of strain level phage-host interactions across the Escherichia genus using only genomic information. Nature Microbiology. 2024 Nov;9(11):2847–2861.

[15] Paczosa MK, Mecsas J. Klebsiella pneumoniae: Going on the Offense with a Strong Defense. Microbiology and molecular biology reviews: MMBR. 2016 Sep;80(3):629–661.

[16] Follador R, Heinz E, Wyres KL, Ellington MJ, Kowarik M, Holt KE, et al. The diversity of Klebsiella pneumoniae surface polysaccharides. Microbial Genomics. 2016 Aug;2(8):e000073.

[17] Lam MMC, Wick RR, Judd LM, Holt KE, Wyres KL. Kaptive 2.0: updated capsule and lipopolysaccharide locus typing for the Klebsiella pneumoniae species complex. Microbial Genomics. 2022 Mar;8(3):000800.

[18] Wyres KL, Lam MMC, Holt KE. Population genomics of Klebsiella pneumoniae. Nature Reviews Microbiology. 2020 Jun;18(6):344–359.

[19] Majkowska-Skrobek G, Łątka A, Berisio R, Maciejewska B, Squeglia F, Romano M, et al. Capsule-Targeting Depolymerase, Derived from Klebsiella KP36 Phage, as a Tool for the Development of Anti-Virulent Strategy. Viruses. 2016 Dec;8(12):324.

[20] Latka A, Maciejewska B, Majkowska-Skrobek G, Briers Y, Drulis-Kawa Z. Bacteriophage-encoded virion-associated enzymes to overcome the carbohydrate barriers during the infection process. Applied Microbiology and Biotechnology. 2017 Apr;101(8):3103–3119.

[21] Cheetham MJ, Huo Y, Stroyakovski M, Cheng L, Wan D, Dell A, et al. Specificity and diversity of Klebsiella pneumoniae phage-encoded capsule depolymerases. Essays in Biochemistry. 2024 Dec;68(5):661–677.

[22] Latka A, Leiman PG, Drulis-Kawa Z, Briers Y. Modeling the Architecture of Depolymerase-Containing Receptor Binding Proteins in Klebsiella Phages. Frontiers in Microbiology. 2019;10:2649.

[23] Squeglia F, Maciejewska B, Łątka A, Ruggiero A, Briers Y, Drulis-Kawa Z, et al. Structural and Functional Studies of a Klebsiella Phage Capsule Depolymerase Tailspike: Mechanistic Insights into Capsular Degradation. Structure (London, England: 1993). 2020 Jun;28(6):613–624.e4.

[24] Pan YJ, Lin TL, Chen CC, Tsai YT, Cheng YH, Chen YY, et al. Klebsiella Phage PhiK64-1 Encodes Multiple Depolymerases for Multiple Host Capsular Types. Journal of Virology. 2017 Mar;91(6):e02457–16.

[25] Hsieh PF, Lin HH, Lin TL, Chen YY, Wang JT. Two T7-like Bacteriophages, K5-2 and K5-4, Each Encodes Two Capsule Depolymerases: Isolation and Functional Characterization. Scientific Reports. 2017 Jul;7(1):4624.

[26] de Sousa JAM, Buffet A, Haudiquet M, Rocha EPC, Rendueles O. Modular prophage interactions driven by capsule serotype select for capsule loss under phage predation. The ISME journal. 2020 Dec;14(12):2980–2996.

[27] Haudiquet M, Buffet A, Rendueles O, Rocha EPC. Interplay between the cell envelope and mobile genetic elements shapes gene flow in populations of the nosocomial pathogen Klebsiella pneumoniae. PLoS biology. 2021 Jul;19(7):e3001276.

[28] Concha-Eloko R, Beamud B, Domingo-Calap P, Sanjuán R. Unlocking data in *Klebsiella* lysogens to predict capsular type-specificity of phage depolymerases. Bioinformatics; 2024. Available from: http://biorxiv.org/lookup/doi/10.1101/2024.07.24.604748.

[29] Power RA, Parkhill J, de Oliveira T. Microbial genome-wide association studies: lessons from human GWAS. Nature Reviews Genetics. 2017 Jan;18(1):41–50.

[30] Coll F, Phelan J, Hill-Cawthorne GA, Nair MB, Mallard K, Ali S, et al. Genome-wide analysis of multi- and extensively drug-resistant Mycobacterium tuberculosis. Nature Genetics. 2018 Feb;50(2):307–316.

[31] Lees JA, Croucher NJ, Goldblatt D, Nosten F, Parkhill J, Turner C, et al. Genome-wide identification of lineage and locus specific variation associated with pneumococcal carriage duration. eLife. 2017 Jul;6:e26255.

[32] Lees JA, Ferwerda B, Kremer PHC, Wheeler NE, Serón MV, Croucher NJ, et al. Joint sequencing of human and pathogen genomes reveals the genetics of pneumococcal meningitis. Nature Communications. 2019 May;10(1):2176.

[33] Gorrie CL, Mirčeta M, Wick RR, Judd LM, Lam MMC, Gomi R, et al. Genomic dissection of Klebsiella pneumoniae infections in hospital patients reveals insights into an opportunistic pathogen. Nature Communications. 2022 May;13(1):3017.

[34] Thorpe HA, Booton R, Kallonen T, Gibbon MJ, Couto N, Passet V, et al. A large-scale genomic snapshot of Klebsiella spp. isolates in Northern Italy reveals limited transmission between clinical and non-clinical settings. Nature Microbiology. 2022 Dec;7(12):2054–2067.

[35] Gorrie CL, Mirceta M, Wick RR, Edwards DJ, Thomson NR, Strugnell RA, et al. Gastrointestinal Carriage Is a Major Reservoir of Klebsiella pneumoniae Infection in Intensive Care Patients. Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America. 2017 Jul;65(2):208–215.

[36] Gorrie CL, Mirceta M, Wick RR, Judd LM, Wyres KL, Thomson NR, et al. Antimicrobial-Resistant Klebsiella pneumoniae Carriage and Infection in Specialized Geriatric Care Wards Linked to Acquisition in the Referring Hospital. Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America. 2018 Jul;67(2):161–170.

[37] Foster-Nyarko E, Cottingham H, Wick RR, Judd LM, Lam MMC, Wyres KL, et al. Nanopore-only assemblies for genomic surveillance of the global priority drug-resistant pathogen, Klebsiella pneumoniae. Microbial Genomics. 2023 Feb;9(2):mgen000936.

[38] Lees JA, Harris SR, Tonkin-Hill G, Gladstone RA, Lo SW, Weiser JN, et al. Fast and flexible bacterial genomic epidemiology with PopPUNK. Genome Research. 2019 Feb;29(2):304–316.

[39] Roux S, Enault F, Hurwitz BL, Sullivan MB. VirSorter: mining viral signal from microbial genomic data. PeerJ. 2015;3:e985.

[40] Akhter S, Aziz RK, Edwards RA. PhiSpy: a novel algorithm for finding prophages in bacterial genomes that combines similarity- and composition-based strategies. Nucleic Acids Research. 2012 Sep;40(16):e126–e126. Available from: https://academic.oup.com/nar/article/40/16/e126/1027055.

[41] Nayfach S, Camargo AP, Schulz F, Eloe-Fadrosh E, Roux S, Kyrpides NC. CheckV assesses the quality and completeness of metagenome-assembled viral genomes. Nature Biotechnology. 2021 May;39(5):578–585.

[42] Terzian P, Olo Ndela E, Galiez C, Lossouarn J, Pérez Bucio RE, Mom R, et al. PHROG: families of prokaryotic virus proteins clustered using remote homology. NAR genomics and bioinformatics. 2021 Sep;3(3):lqab067.

[43] Latka A, Lemire S, Grimon D, Dams D, Maciejewska B, Lu T, et al. Engineering the Modular Receptor-Binding Proteins of Klebsiella Phages Switches Their Capsule Serotype Specificity. mBio. 2021 May;12(3):e00455–21.

[44] Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature. 2024 Jun;630(8016):493–500.

[45] van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, et al. Fast and accurate protein structure search with Foldseek. Nature Biotechnology. 2024 Feb;42(2):243–246.

[46] Hadfield J, Croucher NJ, Goater RJ, Abudahab K, Aanensen DM, Harris SR. Phandango: an interactive viewer for bacterial population genomics. Bioinformatics (Oxford, England). 2018 Jan;34(2):292–293.

[47] Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein structure prediction with AlphaFold. Nature. 2021 Aug;596(7873):583–589.

[48] Patro LPP, Sudhakar KU, Rathinavelan T. K-PAM: a unified platform to distinguish Klebsiella species K- and O-antigen types, model antigen structures and identify hypervirulent strains. Scientific Reports. 2020 Oct;10(1):16732.

[49] Wyres KL, Wick RR, Judd LM, Froumine R, Tokolyi A, Gorrie CL, et al. Distinct evolutionary dynamics of horizontal gene transfer in drug resistant and virulent clones of Klebsiella pneumoniae. PLoS genetics. 2019 Apr;15(4):e1008114.

[50] Dunne M, Prokhorov NS, Loessner MJ, Leiman PG. Reprogramming bacteriophage host range: design principles and strategies for engineering receptor binding proteins. Current Opinion in Biotechnology. 2021 Apr;68:272–281.

[51] Liu S, Lei T, Tan Y, Huang X, Zhao W, Zou H, et al. Discovery, structural characteristics and evolutionary analyses of functional domains in Acinetobacter baumannii phage tail fiber/spike proteins. BMC microbiology. 2025 Feb;25(1):73.

[52] Prokhorov NS, Riccio C, Zdorovenko EL, Shneider MM, Browning C, Knirel YA, et al. Function of bacteriophage G7C esterase tailspike in host cell adsorption. Molecular Microbiology. 2017 Aug;105(3):385–398.

[53] Broeker NK, Barbirz S. Not a barrier but a key: How bacteriophages exploit host's O-antigen as an essential receptor to initiate infection. Molecular Microbiology. 2017 Aug;105(3):353–357.

[54] Van Dam JEG, Van Halbeek H, Kamerling JP, Vliegenthart JFG, Snippe H, Jansze M, et al. A bacteriophage-associated lyase acting on Klebsiella serotype K5 capsular polysaccharide. Carbohydrate Research. 1985 Oct;142(2):338–343. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621585850370`.

[55] Chowdhury TA, Jansson PE, Lindberg B, Lindquist U. Structural studies of an extracellular polysaccharide (K21b) elaborated by Klebsiella type 21b. Carbohydrate Research. 1989 Jul;190(1):145–151. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621589841540`.

[56] Parolis LAS, Parolis H, Niemann H, Stirm S. Primary structure of Klebsiella serotype K22 capsular polysaccharide: Another glycan containing 4-O-[(S-1-carboxyethyl]-d-glucuronic acid. Carbohydrate Research. 1988 Aug;179:301–314. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621588841260`.

[57] Lindberg B, Lindh F, Lönngren J, Nimmich W. Structural studies of the capsular polysaccharide of Klebsiella type 33. Carbohydrate Research. 1979 Apr;70(1):135–144. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500832772`.

[58] Dutton GGS, Merrifield EH. The capsular polysaccharide from Klebsiella serotype K54; location of the O-acyl groups, and a revised structure. Carbohydrate Research. 1982;105(2):189–203. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500849678`.

[59] Bebault GM, Dutton GGS. Structural investigation of the capsular polysaccharide of Klebsiella serotype 55. Carbohydrate Research. 1978 Jul;64:199–213. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500837015`.

[60] Dutton GGS, Savage AV. Structural investigation of the capsular polysaccharide of klebsiella serotype K58. Carbohydrate Research. 1980 Sep;84(2):297–305. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500855597`.

[61] Lindberg B, Lönngren J, Rudén U, Nimmich W. Structural studies of the capsular polysaccharide of Klebsiella type 59. Carbohydrate Research. 1975 Jun;42(1):83–93. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500841026`.

[62] Jansson PE, Lindberg B, Widmalm G, Dutton GGS, Lim AVS, Sutherland IW. Structural studies of the capsular polysaccharides from Klebsiella types 8 and 82, a reinvestigation. Carbohydrate Research. 1988 Apr;175(1):103–109. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621588801605`.

[63] Merrifield EH, Stephen AM. Structural investigation of Klebsiella K-type 4 capsular polysaccharide. Carbohydrate Research. 1981 Oct;96(1):113–120. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500847011`.

[64] Cescutti P, Ravenscroft N, Ng S, Lam Z, Dutton GGS. Structural investigation of the capsular polysaccharide produced by a novel Klebsiella serotype (SK1). Location of O-acetyl substituents using NMR and MS techniques. Carbohydrate Research. 1993 Jun;244(2):325–340. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621583850113`.

[65] Dutton GGS, Ng SK, Parolis LAS, Parolis H, Chakraborty AK. A re-investigation of the structure of the capsular polysaccharide of Klebsiella K10. Carbohydrate Research. 1989 Oct;193:147–155. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621589851146`.

[66] Thurow H, Choy YM, Frank N, Niemann H, Stirm S. The structure of Klebsiella serotype 11 capsular polysaccharide. Carbohydrate Research. 1975 May;41(1):241–255. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500870238`.

[67] Niemann H, Frank N, Stirm S. Klebsiella serotype-13 capsular polysaccharide: Primary structure and depolymerization by a bacteriophage-borne glycanase. Carbohydrate Research. 1977 Nov;59(1):165–177. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500833030`.

[68] Choy YM, Dutton GGS. Structure of the Capsular Polysaccharide of *Klebsiella* K-type 20. Canadian Journal of Chemistry. 1973 Sep;51(18):3015–3020. Available from: `http://www.nrcresearchpress.com/doi/10.1139/v73-449`.

[69] Choy YM, Dutton GGS, Zanlungo AM. The Structure of the Capsular Polysaccharide of Klebsiella K-type 24. Canadian Journal of Chemistry. 1973 Jun;51(11):1819–1825. Available from: `http://www.nrcresearchpress.com/doi/10.1139/v73-271`.

[70] Lindberg B, Lindh F, Lönngren J, Sutherland IW. Structural studies of the capsular polysaccharide of Klebsiella type 30. Carbohydrate Research. 1979 Nov;76(1):281–284. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621579800312`.

[71] Joseleau JP, Lapeyre M, Vignon M, Dutton GGS. Chemical and n.m.r.-spectroscopic investigation of the capsular polysaccharide of klebsiella serotype k41. Carbohydrate Research. 1978 Nov;67(1):197–212. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500837428`.

[72] Edebrink P, Jansson PE, Widmalm G, Nimmich W. The structure of the capsular polysaccharide from Klebsiella K43. Carbohydrate Research. 1994 Apr;257(1):107–115. Available from: `https://linkinghub.elsevier.com/retrieve/pii/000862159484111X`.

[73] Dutton GGS, Folkman TE. Structural investigation of the capsular polysaccharide of Klebsiella-serotype K44. Carbohydrate Research. 1980 Jan;78(2):305–315. Available from: `https://linkinghub.elsevier.com/retrieve/pii/0008621580900117`.

[74] Joseleau JP. Structural investigation of the capsular polysaccharide of Klebsiella serotype K 49. Carbohydrate Research. 1985 Oct;142(1):85–92. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500907353`.

[75] Joseleau JP, Marais MF. Structure of the capsular polysaccharide of Klebsiella K- type 63. Carbohydrate Research. 1979 Dec;77(1):183–190. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500838045`.

[76] Merrifield EH, Stephen AM. Structural studies on the capsular polysaccharide from klebsiella serotype k64. Carbohydrate Research. 1979 Sep;74(1):241–257. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500847801`.

[77] Hackland PL, Parolis H, Parolis LAS. A structural investigation of the capsular polysaccharide of Klebsiella K69. Carbohydrate Research. 1988 Feb;172(2):209–216. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500908553`.

[78] Lindberg B, Lindqvist B, Lönngren J, Nimmich W. Structural studies of the capsular polysaccharide of Klebsiella type 37. Carbohydrate Research. 1977 Oct;58(2):443–451. Available from: `https://linkinghub.elsevier.com/retrieve/pii/S0008621500843712`.

[79] Wollin R, Stocker BA, Lindberg AA. Lysogenic conversion of Salmonella typhimurium bacteriophages A3 and A4 consists of O-acetylation of rhamnose of the repeating unit of the O-antigenic polysaccharide chain. Journal of Bacteriology. 1987 Mar;169(3):1003–1009.

[80] Knirel YA, Prokhorov NS, Shashkov AS, Ovchinnikova OG, Zdorovenko EL, Liu B, et al. Variations in O-antigen biosynthesis and O-acetylation associated with altered phage sensitivity in Escherichia coli 4s. Journal of Bacteriology. 2015 Mar;197(5):905–912.

[81] Allison GE, Verma NK. Serotype-converting bacteriophages and O-antigen modification in Shigella flexneri. Trends in Microbiology. 2000 Jan;8(1):17–23.

[82] Teh MY, Furevi A, Widmalm G, Morona R. Influence of Shigella flexneri 2a O Antigen Acetylation on Its Bacteriophage Sf6 Receptor Activity and Bacterial Interaction with Human Cells. Journal of Bacteriology. 2020 Nov;202(24):e00363–20.

[83] Kuzio J, Kropinski AM. O-antigen conversion in Pseudomonas aeruginosa PAO1 by bacteriophage D3. Journal of Bacteriology. 1983 Jul;155(1):203–212.

[84] Kropinski AM. Sequence of the genome of the temperate, serotype-converting, Pseudomonas aeruginosa bacteriophage D3. Journal of Bacteriology. 2000 Nov;182(21):6066–6074.

[85] Ye TJ, Fung KM, Lee IM, Ko TP, Lin CY, Wong CL, et al. Klebsiella pneumoniae K2 capsular polysaccharide degradation by a bacteriophage depolymerase does not require trimer formation. mBio. 2024 Mar;15(3):e0351923.

[86] Bobay LM, Touchon M, Rocha EPC. Pervasive domestication of defective prophages by bacteria. Proceedings of the National Academy of Sciences of the United States of America. 2014 Aug;111(33):12127–12132.

[87] Batisti Biffignandi G, Gibbon MJ, Corbella M, Thorpe HA, Merla C, Castelli M, et al. Genome of Superficieibacter maynardsmithii, a novel, antibiotic susceptible representative of Enterobacteriaceae. G3 (Bethesda, Md). 2021 Feb;11(2):jkab019.

[88] Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, et al. Producing polished prokaryotic pangenomes with the Panaroo pipeline. Genome Biology. 2020 Jul;21(1):180.

[89] Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, et al. SNP-sites: rapid efficient extraction of SNPs from multi-FASTA alignments. Microbial Genomics. 2016 Apr;2(4):e000056.

[90] Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. Corrigendum to: IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution. 2020 Aug;37(8):2461.

[91] Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, Holt KE. A genomic surveillance framework and genotyping tool for Klebsiella pneumoniae and its related species complex. Nature Communications. 2021 Jul;12(1):4188.

[92] Camargo AP, Roux S, Schulz F, Babinski M, Xu Y, Hu B, et al. Identification of mobile genetic elements with geNomad. Nature Biotechnology. 2024 Aug;42(8):1303–1312.

[93] Mirdita M, Steinegger M, Söding J. MMseqs2 desktop and local web server app for fast, interactive sequence searches. Bioinformatics (Oxford, England). 2019 Aug;35(16):2856–2858.

[94] Steinegger M, Meier M, Mirdita M, Vöhringer H, Haunsberger SJ, Söding J. HH-suite3 for fast remote homology detection and deep protein annotation. BMC bioinformatics. 2019 Sep;20(1):473.

[95] Lees JA, Galardini M, Bentley SD, Weiser JN, Corander J. pyseer: a comprehensive tool for microbial pangenome-wide association studies. Bioinformatics (Oxford, England). 2018 Dec;34(24):4310–4312.

[96] Lees JA, Mai TT, Galardini M, Wheeler NE, Horsfield ST, Parkhill J, et al. Improved Prediction of Bacterial Genotype-Phenotype Associations Using Interpretable Pangenome-Spanning Regressions. mBio. 2020 Jul;11(4):e01344–20.

[97] Wilkins MR, Gasteiger E, Bairoch A, Sanchez JC, Williams KL, Appel RD, et al. Protein identification and analysis tools in the ExPASy server. Methods in Molecular Biology (Clifton, NJ). 1999;112:531–552.

[98] Maciejewska B, Squeglia F, Latka A, Privitera M, Olejniczak S, Switala P, et al. Klebsiella phage KP34gp57 capsular depolymerase structure and function: from a serendipitous finding to the design of active mini-enzymes against K. pneumoniae. mBio. 2023 Oct;14(5):e0132923.