Cryo-EM of cardiac AL-224L amyloid reveals shared features

in λ6 light chain fibril folds

Authors: Chad W. Hicks [a,#], Tatiana Prokaeva [b,c], Brian Spencer [b], Shobini
Jayaraman [a], Noorul Huda [a], Sherry Wong [b], Hui Chen [c], Vaishali Sanchorawala [b],
Francesca Lavatelli [d], Olga Gursky [a,#]

Affiliations:

[a] *Department of Pharmacology, Physiology & Biophysics, Chobanian & Avedisian School*

of Medicine, Boston University, Boston, MA, USA

[b] *Amyloidosis Center, Chobanian & Avedisian School of Medicine, Boston University,*

Boston, MA, USA

[c] *Department of Pathology and Laboratory Medicine, Chobanian & Avedisian School of*
Medicine, Boston University, Boston, MA, USA

[d] *Department of Molecular Medicine, University of Pavia, and Research Area,*
Fondazione IRCCS Policlinico San Matteo, Pavia, Italy.
16

 # Corresponding authors

ORCID ID: Chad Hicks 0000-0002-6040-6227; Olga Gursky 0000-0002-8598-4824

Keywords: Human immunoglobulin mutations; Structural polymorphism; Registry shift
with backbone flipping; Amyloid-ligand interactions; Collagen binding to amyloid; short-
chain fatty acids.

Abbreviations: AL – amyloid light chain; APR – amyloid-promoting region; CDR –
complementarity determining region; C_L – light chain constant domain; cryo-EM –
cryogenic electron microscopy; LC – light chain; LC-MS/MS - liquid chromatography
tandem mass spectrometry; V_L – light chain variable domain.

ABSTRACT

In amyloid light chain (AL) amyloidosis, aberrant monoclonal antibody light chains (LCs) deposit in vital organs causing organ damage. Each AL patient features a unique LC. Previous cryogenic electron microscopy (cryo-EM) studies revealed different amyloid structures in different AL patients. How LC mutations influence amyloid structures remains unclear. We report a cryo-EM structure of cardiac AL-224L amyloid (2.92 Å resolution) from λ6-LC family, which is overrepresented in amyloidosis. Comparison with λ6-LC structures from two other patients reveals similarities in amyloid folds. Mutation-induced structural differences in AL-224L include altered C-terminal conformation with an exposed ligand-binding surface; an enlarged hydrophilic pore with orphan density; and altered steric zipper registry with backbone flipping, which likely represent general adaptive mechanisms in amyloids. The results suggest shared features in λ6-LC amyloid folds and reveal how mutation-induced structural changes influence amyloid-ligand interactions in a patient-specific manner.

INTRODUCTION

AL amyloidosis accounts for nearly ¾ of all systemic amyloid diseases. This debilitating life-threatening systemic disorder is caused by the deposition of monoclonal immunoglobulin LCs as extracellular amyloid fibrils in vital organs, particularly heart and kidney[1]. Annually, the disease afflicts 10-12 new patients per million[2,3]. Cardiac AL amyloidosis is detrimental to patients' prognosis and is lethal if untreated. Current treatments involve chemotherapy and bone marrow transplantation to eliminate the abnormally proliferating B-cell clone that overproduces the aberrant LC protein, along with proteasome inhibitors, immunomodulatory agents, and monoclonal antibodies to eliminate the LC[2–4]. Transplantation of the afflicted organ (kidney, heart) and supportive treatments are also available to extend the patient's life and improve the quality of life. However, these treatments are not suited for all patients. New therapies are being developed, including small-molecule LC stabilizers and CRISPR-based approaches targeting the aberrant LC generation[3–5]. These efforts are complicated by the extreme variability of antibodies stemming from gene recombination and somatic hypermutations; as a result, each amyloidosis patient features a unique fibril-forming LC[6–8]. Since the pathological role of fibrils in AL and other forms of systemic amyloidoses is well established[4], understanding molecular underpinnings for amyloid fibril formation, structure and biological properties may help guide the development of much-needed amyloid-targeting therapies.

In the normal polyclonal LC repertoire, κ-LC and λ-LC proteins are found at a 2:1 ratio, yet AL amyloidosis is most frequently associated with the overexpression of λ-LCs[9]. Moreover, *IGLV6-57* gene segment encoding the variable domain (V_L) of λ6-LCs is expressed in ~2% of the polyclonal λ-LC repertoire but accounts for ~25% of AL amyloid cases[10,11]. This study is focused on the λ6-LC protein subfamily.

Native LC proteins (~25 kDa) share a common two-domain architecture consisting of the N-terminal variable (V_L) and C-terminal constant (C_L) domains joined by a short flexible linker (J-region); each domain acquires a β-sandwich immunoglobulin fold. AL deposits always contain V_L, either as a full-length LC or LC fragments[12,13]. Most LC mutations in AL patients cluster in V_L in or near three antigen-binding loops, called complementarity-determining regions (CDRs). Amyloidogenic mutations have been reported to destabilize the native structure in V_L, to perturb the protective domain-domain interactions in LC monomers or dimers, to increase protein dynamics/exposure in sensitive regions (including the interdomain linker, the amyloid-promoting regions (APRs) in V_L, and the proteolysis-prone segments in C_L), or to stabilize the amyloid structure[5,14–21]. However, oftentimes it is unclear how amyloidogenic mutations promote the misfolding of native LCs to amyloid structure. Moreover, the major triggers of misfolding are different for different LCs[22,23], suggesting different amyloidogenic pathways for different patients.

This idea is supported by the cryo-EM structural studies; 12 structures of tissue-derived amyloid fibrils formed by LCs from λ1, λ3, and λ6 sub-families are currently available[13,18,21,23–26]. The amyloid folds in these fibrils drastically differ from the native immunoglobulin fold. Importantly, fibrils from the same organ of different patients show different amyloid structures[23], while fibrils from different organs of the same patient show similar structures[24]. Since each AL patient features a unique set of mutations in the fibrillar LC, these results suggest that LC mutations are critical for determining amyloid fold.

Nevertheless, cryo-EM structures of several λ3-LCs from different patients show some
similarities, suggesting that amyloids from the same protein subfamily can be structurally
related[13,23]. Despite this recent progress, the paucity of the AL amyloid fibril structures and
their diversity prohibit using machine-learning tools such as AlphaFold[26] to predict new
amyloid structures of LCs based on their amino acid sequences.

Of the available LC fibril structures, only two proteins are from the λ6-LC subfamily:
amyloids from the heart and kidney of patient AL55 (AL-Base number AL55, PDB ID:
6HUD and 8CPE)[24,26] and amyloid from the heart of a patient AL-9ELS (PDB ID: 9ELS)[21].
Cardiac and renal AL55 amyloids have similar structures that differ from the AL-9ELS
structure. To expand the available structural repertoire of λ6-LC amyloids, we determined
a cryo-EM structure of cardiac AL-224L amyloid (PDB ID: 9OKA). This structure shares
key similarities but also shows important mutation-induced structural differences with AL55
and AL-9ELS amyloids. Interestingly, AL-224L amyloid shows an altered C-terminal
conformation with an exposed ligand-binding surface that may bind collagen; an enlarged
hydrophilic pore with orphan density; and altered steric zipper registry with backbone
flipping. These results suggest shared structural features for λ6-LC amyloids, show how
AL mutations influence amyloid structure, and reveal how this structural variability alters
amyloid-ligand interactions that define the biological properties of amyloids.

RESULTS

Analysis of AL-224L amyloid deposits and the extracted fibrils

Amyloid fibrils were extracted from the explanted heart of a previously reported 63-year-old female patient AL-224 with AL amyloidosis[28] (Table S1). Tissue analysis by Congo red staining and light microscopy indicated amyloid (Fig. S1A,B). Immunogold transmission electron microscopy showed reactivity with λ-LC antibody (Fig. S1C,D). The *IGL* gene sequence derived from the patient's bone marrow showed that AL-224L protein belonged to the λ6-LC family and was encoded by the *IGLV6-57*01, IGLJ2*02,* and *IGLC2*02* germline genes.

Liquid chromatography tandem mass spectrometry (LC-MS/MS) of fibril extracts confirmed the sequence identity of the amyloid-forming and the bone marrow-derived LCs and showed 74% peptide coverage in AL-224L residues 1-192, which encompass V_L and a part of C_L domain (Figs. 1A, S2). 2D-PAGE of fibril extracts showed two most abundant species at 15 kDa (Fig. S2A-B). LC-MS/MS of these species detected AL-224L residues 1-116 (Fig. S2C) confirming that the fibrils contained the entire V_L domain. Similar to previous reports[12,28,29], 2D western blot detected additional low-abundance species representing the full-length and fragmented LC (Fig. S2A-B).

LC-MS/MS identified additional proteins in the fibril extracts; Table S2 lists top 30 proteins. AL-224L was prevalent based on normalized spectral abundance factors. Other proteins, including amyloid signature proteins (apoE, apoA-IV) and three chains of collagen VI (ColVI), were identified with lower abundance. ColVI showed similar abundance in prior studies of cardiac λ3-AL59 fibrils that were extracted using similar protocols and contained ColVI bound to LC fibrils[13].

The native structures of LC and its V_L domain (Fig. 1B,C) differ drastically from their structure in an amyloid fibrils[13,18,21,23–26]. To obtain high-resolution structural information on AL-224L fibrils, we used cryo-EM (Fig. S3). While cryo-EM data of amyloid fibrils are typically processed using Relion[30], we performed cryo-EM data processing exclusively in CryoSPARC. Only two other investigations have determined amyloid structures using CryoSPARC[31,32]. To our knowledge, our structure of AL-224L amyloid has the highest global resolution of all similar structures processed in CryoSPARC. Methods provide detailed information of our data processing workflow to assist future investigations.

Cryo-EM micrographs showed long twisted AL-224L fibrils with either single- or double-filament morphology (Fig. 2A). 2D classification showed that ~75% of the fibril segments were single-filament with a helical crossover of approximately 700 Å, while ~25% were double-filament (Fig. 2B). We determined the cryo-EM structure of the single-filament polymorph to a resolution of 2.92 Å (Table 1 and Fig. S4A), which showed clear side-chain densities (Fig. S4B-D) and β-sheet separation (Fig. S4E). Local resolution estimation of the cryo-EM map showed a well-resolved central core at ~2.5 Å and less resolved peripheral parts at ~3.5 Å (Fig. 2C). Symmetry search analysis of the single-filament structure revealed a helical rise of 4.86 Å and a twist of 1.20°. Analysis of the corrugation in the EM density of backbone carbonyl atoms suggested a right-handed twist (Fig. S5). We were unable to determine the double-filament structure, likely due to an insufficient amount of data for this polymorph.

Cardiac λ6-LC amyloids from different patients share key structural features

Like in other patient-derived AL fibrils, the fibril core in AL-224L contains misfolded V_L
domains in a parallel in-registry cross-β-sheet conformation (Fig. 3). The β-strands are
linked via intermolecular hydrogen bonds running along the fibril z-axis while the side
chains pack via steric zippers in the x-y plane[33,34].

The AL-224L V_L domain contains 111 amino acids, while its AL55 counterpart contains 110
amino acids that differ from AL-224L in 13 point substitutions and 1 insertion (Fig. 1A).
Surprisingly, the two amyloids show similar folds consisting of a central N-terminal "snail
shell-shaped" segment[26] (NT-segment, residues 1-41 in AL-224L and 1-34 in AL55)
enclosed by a C-terminal C-shaped segment (CT-segment, residues 59-104 in AL-224L
and 66-106 in AL55) (Fig. 4A,B). These structured antiparallel segments are covalently
linked by the canonical disulfide, C22-C91. This ordered structure is flanked by two
disordered segments seen in the cryo-EM map as diffuse densities, one encompassing
the CDR2-containing linker between the NT- and CT-segments (residues 42-58 in AL-224L
and 35-65 in AL55) and the other extending C-terminally from V_L into the proximal end of
C_L (residues 105-116 in AL-224L and 107-119 in AL55) (Figs. 3B,C, 4A,B). The ordered
C-terminal tail (CT-tail, residues 95-104 in AL-224L and 95-106 in AL55) is repositioned:
the AL-224L CT-tail extends straight around the NT-segment while the AL55 CT-tail makes
two ~90° turns and folds back on itself (Fig. 4A,B).

The AL-9ELS V_L domain contains 111 amino acids[21] that differ from AL-224L in 11 point
substitutions (Fig. 1A). Like AL-224L and AL-55, the AL-9ELS amyloid structure contains
antiparallel well-ordered NT- and CT-segments (residues 1-40 and 61-112) connected by
the disordered CDR2-containing linker and by the C22-C91 disulfide (Fig. 5). The NT-
segment in the AL-9ELS amyloid is not snail-shell shaped, like in AL-224L and AL55.
Rather than occupying a sequestered central position, AL-9ELS segment 1-14 is
repositioned towards the structural periphery and rotates by ~180° to encircle residues
96-101 of the CT-segment (Fig. 5C). In contrast with AL-224L, the AL-9ELS CT-tail
(structured residues 95-112) folds back on itself.

All three λ6-LC amyloids contain a buried salt bridge, K16-D95 (E95 in AL-9ELS) (Figs.
4C,D, 5)[21,26]. This stabilizing salt bridge and the C22-C91 disulfide help maintain the
antiparallel orientation and local registry of the juxtaposed NT- and CT-segments.

V_L domain of LCs typically harbors six amyloid-promoting regions (APRs)[19]. Six APRs
were predicted by at least three different methods for AL-224L (Fig. 1A) and other λ6-LCs
using the sequence-based consensus algorithm AmylPred2[35]. Four APRs are in the
structured segments of AL224L, AL55, and AL-9ELS fibrils (Figs. 1A, 3D); one APR
overlapping CDR2 is disordered; and one APR overlapping CDR3 acquires variable
conformations in different amyloids[19,21]. The disulfide-forming residues C22 and C91 are
located in APRs in the structured regions near CDR1 and CDR3 (Fig. 1A), supporting the
importance of the disulfide-containing regions in amyloid formation[21,36].

The structured segments in AL-224L and AL55 amyloids are flattened but not planar and
are stacked in β-helix-like structures. In these structures, the rugged fibril ends expose
hydrophobic side chains I29, Y33, and Y37 from the major APR (Fig. 3E), which may help

recruit new protein molecules to the growing fibril end[26]. Residue segment N1-P14
protrudes from the plane of the CT-segment and interacts with strands across the central
cleft that are one or two molecular layers up the fibril length, n to n+1 and n to n+2 (Fig.
3D). AL-9ELS amyloid also shows cross-layer interactions involving N1-P14 segment[21]. In
all three amyloids, such cross-layer interactions sterically interlock different molecules
contributing to fibril stability.

Mutations in CDR3 alter the CT-segment conformation affecting ligand binding

Mutations in the ordered segments of λ6-LC amyloids, which include CDR1 and CDR3,
probably contribute to their structural differences (Fig. 5), while mutations in disordered
segments such as the CDR2-containing linker probably influence the amyloid formation
kinetics but not necessarily the ordered structure[22]. Amino acid sequence in CDR3 of λ6-
LCs is particularly variable[6,7]. Compared to AL-224L, AL55 residues 96-101 from CDR3
differ in three substitutions and one insertion, while CDR3 residues in AL-9ELS differ in
four substitutions, resulting in different CT-tail conformations. Importantly, in AL-224L
amyloid, the CT-tail extends along the NT-segment exposing the side chains of Y90, Q92,
and Y94 (Fig. 4A). However, in AL55 and AL-9ELS amyloids, the CT-tail folds back on the
structured CT-segment and occludes these side chains (Figs. 4B, 5B,C).

The Y90 and Y94 rings form one of the most hydrophobic solvent-exposed surfaces on
the AL-224L fibril (Fig. 3E), which is energetically unfavorable. Our cryo-EM map shows
two large orphan densities, one near Y94 and Q92 side chains and another near Y90 and
D88 side chains (Figs. 3C, 6A,B). To test the significance of these densities, we filtered
the unmasked map to a high statistical significance threshold cutoff using false-discovery
rate thresholding[37] (Supplemental Figure 1). The orphan densities were visible at a high
threshold cutoff, excluding the possibility that they were noise and suggesting bound
ligands. These ligands are absent from AL55 and AL-9ELS structures where this fibril
surface is occluded by the CT-tail[21,26] (Fig. 5B,C). These results show how high sequence
variability in CDR3 can translate into structural variability in the λ6-LC amyloid, leading to
distinct patient-specific CT-tail conformations that critically influence amyloid-ligand
binding.

AL-224L shows orphan density in a hydrophilic pore near an acidic-rich segment

Uncompensated charged segments in amyloids are highly energetically unfavorable due
to a strong repulsion within and between molecular layers[38,39]. Conflicting interatomic
forces in such segments can generate structurally frustrated regions with variable
conformations[39]. LCs feature a highly charged conserved residue segment 84-88 (EDEAD
in AL-224L and AL55, EDEAE in AL-9ELS). In the native LC structure, this segment is
stabilized by an extensive network of salt bridges and hydrogen bonds (Fig. 1C). In λ6-LC
amyloids, the acidic segment is located at the fibril surface forming a part of a variable
hydrophilic pore (Fig. 5). In AL-224L amyloid this pore is enlarged and the EDEAD
segment shows reduced local resolution indicating structural frustration (Fig. 2C).

Mutations in and near the pore probably contribute to differences in its structure. These
differences between AL-224L and AL55 include R24G, G27A, and Q82K substitutions. In
the narrow pore of AL55, the Y89 ring packs against G24 (Fig. 4B), but in AL-224L amyloid,
the Y89 side chain orients towards the C22-C91 disulfide and forms a cation-π interaction
with R24 (Fig. 4A). In AL-9ELS amyloid, R24 also interacts with Y89[21]. Notably, Y89 is
highly conserved in λ6-LCs[11]. Therefore, R24G substitution alters the conformation of the
highly conserved Y89 residue in the pore.

In AL55 and AL-9ELS amyloids the charges within the acidic-rich segment are partially
balanced by the K82-E84 salt bridge; additional charge-charge interactions stabilize the
acidic-rich segment in AL-9ELS (Fig. 5C). However, in AL-224L amyloid the K82Q
substitution eliminates the sole stabilizing salt bridge thus increasing the charge imbalance
at the acidic-reach segment (Fig. 4C), destabilizing the local backbone conformation (Fig.
2C), and contributing to the pore enlargement.

The cryo-EM map of AL-224L shows a strong orphan density in the pore near the side
chains of R24, A87, and D85 (Figs. 3C, 6A-C). This density is well above the noise level
based on the false-discovery rate map thresholding (Supplemental Fig. 1). The cryo-EM
study of AL-9ELS also reports an orphan density in the pore near R24 and Y89 side
chains[21]. In contrast, AL55 (which contains G24 instead of the bulky R24) shows no orphan
density in the pore (Fig. 6D), suggesting that this density is associated with R24 but not
G24 λ6-LC variants.

The orphan density in the hydrophilic pore of AL-224L amyloid likely represents a small
water-soluble molecule that interacts via its acidic and hydrophobic moieties with R24 and
A87, respectively. One possibility is acetate, an abundant tissue metabolite whose cardiac
uptake increases in heart failure[40], which was the cause of death of patient AL-224.
Analysis of fibril tissue extracts using an ELISA-based assay detected short-chain fatty
acids (Table S4 and Fig. S6) confirming them as potential candidates for binding amyloid
in tissues. Figure 6C illustrates how an acetate molecule can fit in the orphan density in
the pore of AL-224L amyloid.

Registry shifts and backbone flipping in amyloid helps accommodate AL mutations

Compared to AL55, the wider pore in AL-224L amyloid is facilitated by a shift in the
backbone registry of the steric zippers between the CT- and NT-segments (Fig. 4E,F).
This registry shift propagates from the EDEAD segment towards the CDR2-containing
linker encompassing a 30-residue segment 59-88. For example, in AL-224L the W36 side
chain packs between the D61 and F63 side chains, yet in AL55 the W36 side chain packs
against S69 (Fig. 4E,F). To accommodate such a 7-residue registry shift, the backbone in
segment 69-74 is flipped in AL-224L relative to AL55, inverting the side chain orientation
in this segment. Registry shifts with the backbone flipping are also seen in other structured
residue segments including 8-11, 24-29, 69-74, and 81-83 (Fig. 4E,F). As a result, most
residues pack differently in the X-Y plane of AL-224L and AL55 amyloids, yet the overall
amyloid fold is largely preserved.

The AL-9ELS structure also shows registry shifts with backbone flipping. For example,
segment 32-40, which packs antiparallel to segment 61-70 with one-residue registry shift

between AL-9ELS and AL-224L amyloids, shows an opposite side chain orientation in AL-
9ELS compared to both AL-224L and AL55 amyloids (Fig. 5). These findings suggest that
registry shifts and associated backbone flipping is a plausible general mechanism enabling
amyloids adapt to mutations while largely preserving the amyloid fold (Fig. 7).

DISCUSSION

λ6-LC amyloids have shared structural features

We determined the single-filament structure of AL-224L cardiac amyloid at 2.92 Å
resolution and compared it with the available amyloid structures of two other cardiac λ6-
LCs, AL55 (4.0 Å resolution) and AL-9ELS (3.0 Å resolution). Despite 14 amino acid
differences between AL-224L and AL55, these two amyloids have surprisingly similar
overall folds (Fig. 4), while all three amyloids share key structural features (Fig. 5). These
features include a structured fibril core comprised of antiparallel NT- and CT-segments
connected by a disordered CDR2-containing linker. In λ1-LC and λ3-LC amyloids the
CDR2-containing linker can be either ordered or disordered[13,23]. Importantly, in all LC
amyloids NT- and CT-segments are linked by the conserved internal disulfide that
maintains their antiparallel orientation providing a major constraint for amyloid formation
and structure[13,21,23,25]. Furthermore, in λ6-LC amyloids the planar part of the CT-segment
partially encircles the fibril surface, while the non-planar NT-segment forms cross-layer
interactions. Lastly, the amyloid structure adjacent to the C22-C91 disulfide contains a
variable pore; the pore is lined by mostly hydrophilic residues including the acidic-rich
segment 84-88 (Fig. 4C,D), while the Y89 side chain packs against G24/R24 in the pore
(Figs. 4A,B). Since in λ6-LCs residues 85-91 are highly conserved while residue 24 is
typically either G or R[7,11], these structural features probably extend to other λ6-LC
amyloids.

The observed structural similarities between the λ6-LC amyloids are facilitated, in part, by
the registry shifts with backbone flipping in steric zippers (Figs. 4E,F, 7). Importantly, we
were able to attribute the structural differences between amyloids to LC mutations,
providing clear insights into the role of patient-specific AL mutations in amyloid
polymorphism and ligand binding.

Germline λ6-LC is compatible with the amyloid folds in tissue-derived fibrils

The fibril-forming V_L domain in λ6-LCs is encoded by the germline gene *IGLV6-57* [10,11].
We propose that the hypothetical protein with the germline amino acid sequence can be
accommodated in the structure of AL-224L and/or other tissue-derived amyloids (Fig. 1A).
The germline-encoded residue V18 (I in AL-224L, L in AL55, and V in AL-9ELS) can be
readily accommodated instead of I18 of AL-224L in the structured NT-segment in the
hydrophobic amyloid core. The germline-encoded S44 (A in AL-224L, AL55, and AL-9ELS)
can also be accommodated in the flexible CDR2-containing segment. Furthermore, the
germline-encoded residue K82 (Q in AL-224L, K in AL55 and AL-9ELS) forms a stabilizing
salt bridge with E84 in AL55 and AL-9ELS amyloids (Figs. 4D, 5B,C)[21,26]. Lastly, germline
mutations in positions 99-101 of CDR3 are very different in AL-224L, AL55, and AL-9ELS
(Fig. 1A) and probably influence the variable CT-tail conformation at the periphery of the

amyloid structure (Fig. 5). This analysis suggests that the amyloid folds observed in AL-
224L, AL55, and AL-9ELS are compatible with the germline LC, suggesting that aspects
of these folds are probably shared by other λ6-LC amyloids.

The amyloid structure of AL-9ELS differs significantly from those of AL-224L and AL55.
The major difference involves residue segment 1-14 that forms cross-layer interactions in
all three amyloids. In AL-224L and AL55 amyloids, residues 1-14 are centered in the snail
shell-shaped core (Fig. 5A,B), yet in AL-9ELS these residues wrap around the structural
periphery (Fig. 5C)[21]. Since residues 1-14 are identical in the three λ6-LCs (Fig. 1), their
conformational differences must reflect mutations located elsewhere. The likely
candidates include residues 18 and 32 in the middle of the snail shell-shaped NT-segment
(which harbors segment 1-7 in AL-224L and AL55 amyloids) and residues 95, 96, 99 and
100 in the CT-tail (which interacts with residues 1-14 in AL-9ELS amyloid) (Fig. 5). This
comparison illustrates how the combined effects of several AL mutations define the
amyloid conformation in a mutation-free segment.

Low predicted stability of λ6-LC amyloid fibrils may promote pathology

Prior biophysical studies of the natively folded recombinant human λ6-LCs did not show
reduced thermodynamic stability of AL-224L compared to non-amyloidogenic controls but
revealed subtle differences in local stability and proteolytic processing, which were
relevant to fibrillogenesis[19]. In the current study, we assessed the amyloid fibril stability.
PDBePISA software[41] was used to calculate free energy change upon exposure of one
fibril layer to solvent for AL-224L and AL55 amyloids, as well as for the hypothetical
germline λ6-LC sequence placed in these fibril structures.

AL-224L has slightly weaker solvation energy than AL55, indicating that it is slightly less
stable than AL55 (Table S3). Structures of AL-224L or AL55 amyloids containing the
hypothetical protein with the germline λ6-LC sequence showed comparable solvation
energies as tissue-derived AL-224L and AL55, indicating comparable stabilities (Table S3).
These calculations support our hypothesis that the germline λ6-LC sequence is
compatible with amyloid structures of the tissue-derived fibrils.

Next, we computed solvation energies for all other available LC amyloid structures using
PDBePISA. AL-224L and AL55 amyloids has lower solvation energies than λ1-LC and λ3-
LC amyloids (Table S3). The Saelices team also reported lower solvation energies for
AL55 and AL-9ELS compared to λ1-LC and λ3-LC amyloids[21]. These results suggest that,
compared to other amyloids from the λ-LC family, λ6-LC fibrils are relatively unstable and
can be fragmented more easily, potentially facilitating amyloid proliferation by seeding and
enhancing amyloid cytotoxicity[42].

Role of registry shifts with backbone flipping in amyloid structures

Comparison of AL-224L, AL55, and AL-9ELS amyloid structures revealed a shift in the
steric zipper registry with the backbone flipping in several well-ordered segments (Figs.
4E,F, 5). We posit that such amyloid-specific structural changes provide a general
mechanism for adapting the fibril core to protein mutations, post-translational

modifications, and structurally frustrated regions that are often interspersed with β-strands
in amyloids[39]. Registry shifts and backbone flipping in steric zippers help explain why the
risk of amyloid formation increases for proteins containing low-complexity domains and
amino acid repeats, such as poly-Q motif in huntingtin or tandem repeats in bacterial
amyloids[43–45]. Such repeating units can facilitate formation of steric zippers differing in the
backbone registry and side chain orientation to generate energetically equivalent but
conformationally distinct fibril polymorphs of the same or closely related proteins[46].
367

A conserved acidic-rich segment is a potential gatekeeper in LC amyloid formation

Unbalanced charges in amyloids are highly energetically unfavorable due to a strong
repulsion among molecular layers of parallel in-registry cross-β-sheets[38,39]. LCs contain a
conserved residue segment 85-91[11] that overlaps the acidic-rich segment 84-88, EDEAD.
We hypothesize that this acidic segment, which forms stabilizing interactions in the native
structure (Fig. 1C) but destabilizes the local structure in amyloid (Fig. 2C), acts as a
gatekeeper protecting LCs from amyloid formation. However, AL mutations can balance
the charge on this acidic segment to promote fibrillogenesis. Examples include K83 that
forms a K83-D85 salt bridge in AL-9ELS amyloid (Fig. 5C). Ligand binding at nearby sites,
such as the internal site near A87 and the external site near D88 and Y90 in AL-224L (Fig.
6), can also stabilize the acidic-rich segment 84-88 in amyloid and thereby promote
fibrillogenesis.
380

R24 but not G24 variant is associated with ligand binding in amyloid pore

Residue 24 is a site of an AL-associated polymorphism in λ6-LCs. R24 is the major variant
found in the germline, AL-224L, and AL-9EL LCs[47]. Compared to R24, G24, the second-
major variant found in AL55, is overrepresented in amyloidosis and is considered more
amyloidogenic[11]. Stabilizing electrostatic interactions involving R24 in the native LC
structure were proposed to counteract amyloid formation[47], but the effects of R24G
substitution on the amyloid structure and stability were unknown. Cryo-EM structures of
AL-224L, AL-9ELS, and AL55 amyloids show that G24 replacement with R24 induces local
structural changes in the hydrophilic pore including the pore enlargement (Figs. 5) likely
contributing to reduced amyloidogenicity of the R24 variant. Moreover, the orphan density
seen in amyloids near R24 in AL-224L and AL-9ELS, but not near G24 in AL55, represents
a bound ligand such as acetate (Fig. 6C). Ligand binding in an internal amyloid pore
probably occurs during fibril formation and may critically influence fibrillogenesis,
suggesting the direct involvement of residue 24 in this process.
395

External ligand binding sites in AL-224L fibrils may harbor collagen triple helices

The cryo-EM map of AL-224L shows two orphan densities, one near residues Y94 and
Q92 and another near Y90 and D88 (Fig. 6A,E). Mixed hydrophilic/hydrophobic
coordinating residues and the ~18 Å center-to-center separation between these densities
suggest that they potentially represent collagen triple helices (diameter 15 Å) and/or
apolipoprotein α-helices (diameter 11-13 Å) bound laterally along the fibril spine, as

previously reported for ColVI[13] and apoE[48]. In the current study, LC-MS/MS of the fibril
tissue extracts detected three chains of ColVI as well as apoE and apoA-IV co-purified
with fibrils (Table S2). The orphan density coordinated by Q92 and Y94 appears hollow in
the x-y plane (Fig. 6A) consistent with a triple helix. To test this hypothesis, we used
pyDock software[49] for rigid body docking of a canonical collagen triple helix to the AL-224L
fibril structure. As a model we used the crystal structure of a 30-residue collagen-like
peptide containing consensus sequence repeats Pro-Hyp-Gly (PDB ID: 1CAG)[50]. The top
two docking poses show triple helices bound along the Y90 and Y94 side chain amyloid
arrays (Fig. 6E,F). These docking poses closely superimpose the two orphan densities in
the EM map of AL-224L (Fig. 6E). Therefore, these densities likely represent collagen triple
helices bound along the fibril surface. The aromatic rings of amyloid stack against the
pyrrolidine rings of collagen. We hypothesize that similar interactions mediate longitudinal
binding of ColVI to λ3-AL59 cardiac amyloid[13]; in fact, orphan densities seen in the cryo-
EM map of AL59 amyloid near the exposed arrays of Y48 and Y35[13] overlap the top
docking poses of the collagen triple helix and hence may represent bound ColVI.

Direct interactions between collagen and aromatic residues have been reported for
globular proteins[51] including proteins featuring an immunoglobulin fold[52,53] such as an
amyloidogenic LC[54]. Our docking results suggest that ring-stacking interactions also
mediate collagen binding to amyloids along the fibril spine. Such a longitudinal binding is
expected to influence mechanical and biological properties of amyloid fibrils including
amyloid nucleation, growth, fragmentation, interactions with cells including phagocytosis,
and the uptake of diagnostic dyes[13,55,56]. Future studies will identify additional details of
such amyloid-ligand interactions and their roles in vivo.

Acknowledgements The authors are grateful to Dr. Esther Bullitt, the director of the
Boston University Cryo-EM Core Facility where the data was collected, to Dr. Luca
Broggini at the University of Milan for invaluable advice on the cryo-EM structural analysis
of amyloids, and to Mei Chen and Steven Kolakowski at the Harvard Center for Mass
Spectrometry for their assistance with the proteomics analyses. We gratefully
acknowledge the patient's family for providing tissue samples for amyloid research.

Funding: This research was supported by the National Institute of General Medical
Sciences grants S10OD032253, GM135158, and GM067260**;** the Boston University
Amyloid Research Fund; Wildflower Foundation Fund; Fondo Italiano per la Scienza (FIS
2021) FIS00001548; Cariplo–Telethon grant 2022-0578; and PNRR grant ID MUR:
CN00000041.

Competing interests   The authors declare no competing interests.

Author contributions

C.W.H. prepared the cryo-EM grids, collected cryo-EM data, performed data processing,
map interpretation, model building and refinement, and prepared graphics.

T.P. initiated the study, oversaw fibril extraction, LC-MS/MS analysis and immuno-EM,
interpreted the results, prepared graphics and contributed to draft writing.

B.S. performed fibril extraction, SDS-PAGE and western blotting, prepared samples for
LC-MS/MS, contributed to immuno-EM, and prepared graphics.

S.J. performed biochemical assays, calculated solvation energies, performed molecular
docking, and contributed to data analysis and interpretation.

N.H. prepared graphics and contributed to structural interpretation.

S.W. optimized sample preparations.

H.C. performed immuno-EM of the tissues and fibrils.

V.S. contributed to the overall study administration and funding.

F.L. oversaw fibril extraction, performed LC-MS/MS analysis and interpretation, and
contributed graphics, draft writing, and funding.

O.G. administered the project, interpreted the results, wrote the first draft, and contributed
to funding. C.W.H. and O.G. wrote the paper with input from all authors.
455

Data availability The AL-224L amino acid sequence has been deposited to the AL-base,
and the gene sequence, *AL3IGLV6-57*01-IGLJ2*02-IGLC2*02* has been deposited in the
GenBank (accession number KY432418). The cryo-EM model and map data have been
deposited to the Protein Data Bank (PDB ID: 9OKA, EMDB code 70557). Raw cryo-EM
movies have been deposited to the Electron Microscopy Public Image Archive (accession
code EMPIAR-12815). The LC-MS/MS proteomics data have been deposited to the
ProteomeXchange Consortium via the PRIDE partner repository (dataset identifiers
PXD064296).

TABLES

Parameter	Data Collection
Magnification (X)	130,000
Voltage (kV)	200
Electron exposure (e⁻/Å²)	40
Dose rate (e⁻/px/s)	10.82
Defocus range (μM)	-0.8 to -2.5
Pixel size (Å)	0.900
Camera	Falcon 4i
Energy Filter slit width (eV)	10
Micrographs (#)	10,309
	Data Processing
Box size (px)	400
Segment separation distance (Å)	45
Initial uncleaned segment stack (#)	421,024
Helical rise (Å)	4.86
Helical twist (°)	1.20
Particle duplication symmetry	C1
Final segments (#)	214,100
Map resolution at 0.143 FSC cutoff (Å)	2.92
	Model Building and Validation
Non-hydrogen atoms (#)	3,335
Subunits	5
CC mask	0.75
Bond length RMSD (Å)	0.002
Bond angles RMSD (°)	0.602
Clash score	7.82
Ramachandran Outliers (%)	0
Ramachandran Allowed (%)	6.02
Ramachandran Favored (%)	93.98
Rotamer Outliers (%)	0
C-beta outliers (%)	0
MolProbity score	1.83

467
468 **Table 1 Cryo-EM statistics and validation metrics.** Table of details related the cryo-EM structure of AL-224L
469 amyloid. The details include parameters for data collection, data processing, model building, and validation.

FIGURES

Figure 1. Structural properties of λ6-LCs explored in this study.
A, Top to bottom: Amino acid sequences of the λ6 germline, AL-224L, AL55, and AL-9ELS V_L domains (consecutive residue numbering). AL55 and AL-9ELS residue differences compared to AL-224L (magenta). AL-224L, AL55, and AL-9ELS residue differences compared to germline LC (boxed). Amyloid-promoting regions in AL-224L (horizontal bars) are colored according to the consensus number predicted using AmylPred2[35]. Framework regions (FR), complementarity determining regions (CDRs), the J-region, and the proximal end of C_L domain are marked. Linear diagrams show secondary structures of λ6-LCs including β-strands (arrows), α-helices (cylinders), loops/turns (solid lines), and disordered regions (broken lines). "Native" denotes the native fold according to the x-ray crystal structure of a related λ6-LC protein, JTO (PDB ID: 6MG4); "F" denotes amyloid fibril structures of AL-224L (PDB ID: 9OKA), AL55 (PDB ID: 6HUD), and AL-9ELS (PDB ID: 9ELS) determined by cryo-EM.
B, Native structure of V_L domain (PDB ID: 6MGT). CDRs are highlighted. Spheres mark residues differentiating AL55 from AL-224L (magenta) and AL-224L from the germline LC (blue).
C, Native structure of full-length LC. The acidic-rich segment 84-88 (red). Zoomed-in view of the boxed area shows salt bridges and hydrogen bonds stabilizing the charges in this acidic segment.

A

Representative micrograph

B

Representative 2D classes

□ single-filament □ double-filament

C

488 **Figure 2. Amyloid fibril polymorphs and local resolution of the single-filament AL-224L structure.**
489 **A,** Representative cropped micrograph shows single (blue arrows) and double filaments (red arrows).
490 **B,** Representative 2D classes at a box size 1800 px (1620 Å) show an approximate crossover distance of 700
491 Å for both single and double filaments.
492 **C,** Cross-section of one molecular layer of the cryo-EM density map colored by local resolution.

Figure 3. Structural overview of the AL-224L single-filament amyloid polymorph.

A, Sharpened cryo-EM map (2.9 Å) of AL-224L amyloid (side view).

B, EM map (top view) showing well-ordered fibril core (blue, contouring threshold 0.0994) and lower-order regions (gray, threshold 0.0325). Disordered segments 42-58 and 105-116 (dashed lines).

C, Cross-section of one fibril layer superimposed on the cryo-EM map. Stick model shows ordered segments N1-P41 (NT-segment) and V59-G104 (CT-segment). CDRs are colored in green and orphan densities are labeled as 1, 2, and *.

D, A stack of molecular layers shows n to n+1 and n to n+2 interlayer contacts across the central cleft (top view). APRs in the top layer are color-coded from least amyloidogenic (cyan) to most amyloidogenic (purple), as in Fig. 1A.

E, Surface representation of five molecular layers colored according to side chain hydrophobicity (bottom view).

Figure 4. Comparison of AL-224L and AL55 fibril structures.

A, B, Cross-sectional top views show proteins' backbone in five molecular layers. Residues that differ in AL-

224L (violet) and AL55 (magenta), residues in the acidic-rich segment (red), and other residues mentioned in

the manuscript (black) are shown.

C, D, Space-filling model of one fibril layer colored according to coulombic potential.

E, F, Regions showing opposite side chain orientation (backbone flipping) in AL-224L vs. AL55 amyloids.

Figure 5. Three available λ6-LC cardiac amyloid structures.
Schematic diagram showing the polypeptide sequence and fold in the fibril core of AL-224L (PDB ID: 9OKA)
(A), AL-55 (PDB ID: 6HUD) **(B)**, and AL-9ELS (PDB ID: 9ELS) **(C)**. β-strands shown as thick lines; disordered
regions shown as dashed lines.

Figure 6. Orphan densities in the cryo-EM map of AL-224L amyloid.

A, Sharpened cryo-EM density map shows one fibril layer (top view) with orphan densities at external (1, 2) and internal (*) sites.

B, Side views show orphan densities in the maps at a threshold of 0.0781 (1, 2) or 0.0567 (*).

C, Zoomed-in view of the EM map and model of the hydrophilic pore in AL-224L shows acetate (ACE) molecule manually fitted in the extra density. Dotted lines show acetate-protein contacts within 4 Å.

D, A view similar to panel C of the amyloid pore in AL55.

E, Cryo-EM map and model of AL-224L show two top docking poses of collagen triple helix (PDB ID: 1CAG).

F, Side view of the space-filling model shows top two docking poses of collagen (yellow) to AL-224L amyloid (blue).

530 **Figure 7. Schematic diagram illustrating registry shifts and backbone flipping in a steric zipper**.
531 A steric zipper before registry shift and backbone flipping (middle). A steric zipper after registry shift (left). A
532 steric zipper after backbone flipping (right). b-strands are shown as gray arrows; side chains are shown as
533 colored bars.

References

1. Merlini, G. *et al*. Systemic immunoglobulin light chain amyloidosis. *Nat Rev Dis Primers* **4**, 38 (2018).

2. Gertz, M. A. & Dispenzieri, A. Systemic Amyloidosis Recognition, Prognosis, and Therapy: A Systematic Review. *JAMA* **324**, 79–89 (2020).

3. Sanchorawala, V. Systemic Light Chain Amyloidosis. *N Engl J Med* **390**, 2295–2307 (2024).

4. Merlini, G. AL amyloidosis: from molecular mechanisms to targeted therapies. *Hematology Am Soc Hematol Educ Program* **2017**, 1–12 (2017).

5. Morgan, G. J., Buxbaum, J. N. & Kelly, J. W. Light Chain Stabilization: A Therapeutic Approach to Ameliorate AL Amyloidosis. *Hemato* **2**, 645–659 (2021).

6. Bellotti, V., Mangione, P. & Merlini, G. Review: immunoglobulin light chain amyloidosis--the archetype of structural and pathogenic variability. *J Struct Biol* **130**, 280–289 (2000).

7. Del Pozo-Yauner, L. *et al*. Role of the mechanisms for antibody repertoire diversification in monoclonal light chain deposition disorders: when a friend becomes foe. *Front Immunol* **14**, 1203425 (2023).

8. Absmeier, R. M., Rottenaicher, G. J., Svilenov, H. L., Kazman, P. & Buchner, J. Antibodies gone bad - the molecular mechanism of light chain amyloidosis. *FEBS J* **290**, 1398–1419 (2023).

9. Haupt, C. The AL Amyloid Fibril: Looking for a Link between Fibril Formation and Structure. *Hemato* **2**, 505–514 (2021).

10. Bodi, K. *et al*. AL-Base: a visual platform analysis tool for the study of amyloidogenic immunoglobulin light chain sequences. *Amyloid* **16**, 1–8 (2009).

11. Morgan, G. J. *et al*. An updated AL-base reveals ranked enrichment of immunoglobulin light chain variable genes in AL amyloidosis. *Amyloid* **32**, 129–138 (2025).

12. Lavatelli, F. *et al*. Mass spectrometry characterization of light chain fragmentation sites in cardiac AL amyloidosis: insights into the timing of proteolysis. *J Biol Chem* **295**, 16572–16584 (2020).

13. Schulte, T. *et al*. Helical superstructures between amyloid and collagen in cardiac fibrils
from a patient with AL amyloidosis. *Nat Commun* **15**, 6359 (2024).

14. Brumshtein, B. *et al*. Inhibition by small-molecule ligands of formation of amyloid fibrils
of an immunoglobulin light chain variable domain. *Elife* **4**, e10935 (2015).

15. Blancas-Mejia, L. M. *et al*. Immunoglobulin light chain amyloid aggregation. *Chem*
Commun (Camb) **54**, 10664–10674 (2018).

16. Kazman, P. *et al*. Fatal amyloid formation in a patient's antibody light chain is caused by
a single point mutation. *Elife* **9**, e52300 (2020).

17. Peterle, D. *et al*. A Conservative Point Mutation in a Dynamic Antigen-binding Loop of
Human Immunoglobulin λ6 Light Chain Promotes Pathologic Amyloid Formation. *J Mol*
Biol **433**, 167310 (2021).

18. Radamaker, L. *et al*. Cryo-EM reveals structural breaks in a patient-derived amyloid fibril
from systemic AL amyloidosis. *Nat Commun* **12**, 875 (2021).

19. Klimtchuk, E. S. *et al*. Conformational Differences in the Light Chain Constant Domain
of Immunoglobulin G and Free Light Chain May Influence Proteolysis in AL Amyloidosis.
J Mol Biol **436**, 168837 (2024).

20. Paissoni, C. *et al*. A conformational fingerprint for amyloidogenic light chains. *Elife* **13**,
RP102002 (2025).

21. Cryo-EM reveals that cardiac IGLV6-derived AL fibrils can be polymorphic | bioRxiv.
https://www.biorxiv.org/content/10.1101/2024.12.04.626857v1.

22. Ulamec, S. M., Brockwell, D. J. & Radford, S. E. Looking Beyond the Core: The Role of
Flanking Regions in the Aggregation of Amyloidogenic Peptides and Proteins. *Front*
Neurosci **14**, 611285 (2020).

23. Karimi-Farsijani, S. *et al*. Light chain mutations contribute to defining the fibril
morphology in systemic AL amyloidosis. *Nat Commun* **15**, 5121 (2024).

24. Puri, S. *et al*. The Cryo-EM Structure of Renal Amyloid Fibril Suggests Structurally
Homogeneous Multiorgan Aggregation in AL Amyloidosis. *J Mol Biol* **435**, 168215 (2023).

25. Radamaker, L. *et al*. Cryo-EM structure of a light chain-derived amyloid fibril from a
patient with systemic AL amyloidosis. *Nat Commun* **10,** 1103 (2019).

26. Swuec, P. *et al*. Cryo-EM structure of cardiac amyloid fibrils from an immunoglobulin
light chain AL amyloidosis patient. *Nat Commun* **10**, 1269 (2019).

27. Tunyasuvunakool, K. *et al*. Highly accurate protein structure prediction for the human
proteome. *Nature* **596**, 590–596 (2021).

28. Klimtchuk, E. S. *et al*. The critical role of the constant region in thermal stability and
aggregation of amyloidogenic immunoglobulin light chain. *Biochemistry* **49**, 9848–9857
(2010).

29. Enqvist, S., Sletten, K. & Westermark, P. Fibril protein fragmentation pattern in systemic
AL-amyloidosis. *J Pathol* **219**, 473–480 (2009).

30. Scheres, S. H. W. Amyloid structure determination in RELION-3.1. *Acta Crystallogr D*
Struct Biol **76**, 94–101 (2020).

31. Hansen, K. H. *et al*. Structure of biofilm-forming functional amyloid PSMα1 from
Staphylococcus aureus. *Proc Natl Acad Sci U S A* **121**, e2406775121 (2024).

32. Bu, F., Dee, D. R. & Liu, B. Structural insight into Escherichia coli CsgA amyloid fibril
assembly. *mBio* **15**, e0041924 (2024).

33. Iadanza, M. G., Jackson, M. P., Hewitt, E. W., Ranson, N. A. & Radford, S. E. A new era for
understanding amyloid structures and disease. *Nat Rev Mol Cell Biol* **19**, 755–773
(2018).

34. Sawaya, M. R., Hughes, M. P., Rodriguez, J. A., Riek, R. & Eisenberg, D. S. The expanding
amyloid family: Structure, stability, function, and pathogenesis. *Cell* **184**, 4857–4873
(2021).

35. Tsolis, A. C., Papandreou, N. C., Iconomidou, V. A. & Hamodrakas, S. J. A consensus
method for the prediction of 'aggregation-prone' peptides in globular proteins. *PLoS*
One **8**, e54175 (2013).

36. Klimtchuk, E. S. *et al*. Role of complementarity-determining regions 1 and 3 in
pathologic amyloid formation by human immunoglobulin κ1 light chains. *Amyloid* **30**,
364–378 (2023).

37. Beckers, M., Jakobi, A. J. & Sachse, C. Thresholding of cryo-EM density maps by false
discovery rate control. *IUCrJ* **6**, 18–33 (2019).

38. Lewkowicz, E., Jayaraman, S. & Gursky, O. Protein Amyloid Cofactors: Charged Side-
Chain Arrays Meet Their Match? *Trends Biochem Sci* **46**, 626–629 (2021).

39. van der Kant, R., Louros, N., Schymkowitz, J. & Rousseau, F. Thermodynamic analysis of
amyloid fibril structures reveals a common framework for stability in amyloid
polymorphs. *Structure* **30**, 1178-1189.e3 (2022).

40. Palm, C. L., Nijholt, K. T., Bakker, B. M. & Westenbrink, B. D. Short-Chain Fatty Acids in
the Metabolism of Heart Failure - Rethinking the Fat Stigma. *Front Cardiovasc Med* **9**,
915102 (2022).

41. Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline
state. *J Mol Biol* **372**, 774–797 (2007).

42. Xue, W.-F. *et al.* Fibril fragmentation enhances amyloid cytotoxicity. *J Biol Chem* **284**,
34272–34282 (2009).

43. Lee, M., Ghosh, U., Thurber, K. R., Kato, M. & Tycko, R. Molecular structure and
interactions within amyloid-like fibrils formed by a low-complexity protein sequence
from FUS. *Nat Commun* **11**, 5735 (2020).

44. Tayeb-Fligelman, E. *et al.* Low complexity domains of the nucleocapsid protein of SARS-
CoV-2 form amyloid fibrils. *Nat Commun* **14**, 2379 (2023).

45. Nowakowska, A. W., Wojciechowski, J. W., Szulc, N. & Kotulska, M. The role of tandem
repeats in bacterial functional amyloids. *J Struct Biol* **215**, 108002 (2023).

46. Louros, N. *et al.* Local structural preferences in shaping tau amyloid polymorphism. *Nat
Commun* **15**, 1028 (2024).

47. del Pozo Yauner, L. *et al.* Influence of the germline sequence on the thermodynamic
stability and fibrillogenicity of human lambda 6 light chains. *Proteins* **72**, 684–692
(2008).

48. Lewkowicz, E., Nakamura, M. N., Rynkiewicz, M. J. & Gursky, O. Molecular modeling of
apoE in complexes with Alzheimer's amyloid-β fibrils from human brain suggests a

structural basis for apolipoprotein co-deposition with amyloids. *Cell Mol Life Sci* **80**,

376 (2023).

49. Grosdidier, S., Pons, C., Solernou, A. & Fernández-Recio, J. Prediction and scoring of

docking poses with pyDock. *Proteins* **69**, 852–858 (2007).

50. Bella, J., Eaton, M., Brodsky, B. & Berman, H. M. Crystal and molecular structure of a

collagen-like peptide at 1.9 A resolution. *Science* **266**, 75–81 (1994).

51. Li, W. *et al*. The structural basis for the collagen processing by human

P3H1/CRTAP/PPIB ternary complex. *Nat Commun* **15**, 7844 (2024).

52. Liu, Q. *et al*. The Enterococcus faecalis MSCRAMM ACE binds its ligand by the Collagen

Hug model. *J Biol Chem* **282**, 19629–19637 (2007).

53. Hoop, C. L. *et al*. Collagen I Weakly Interacts with the β-Sheets of β2-Microglobulin and

Enhances Conformational Exchange To Induce Amyloid Formation. *J Am Chem Soc*

142, 1321–1331 (2020).

54. Harris, D. L., King, E., Ramsland, P. A. & Edmundson, A. B. Binding of nascent collagen

by amyloidogenic light chains and amyloid fibrillogenesis in monolayers of human

fibrocytes. *J Mol Recognit* **13**, 198–212 (2000).

55. Jackson, J. W. *et al*. Collagen inhibits phagocytosis of amyloid in vitro and in vivo and

may act as a 'don't eat me' signal. *Amyloid* **30**, 249–260 (2023).

56. Lewkowicz, E., Jayaraman, S. & Gursky, O. Molecular basis for non-invasive diagnostics

of cardiac amyloids using bone tracers. *Biomater Sci* **12**, 4275–4282 (2024).

METHODS

Clinical data, sample collection and clinical characterization of patient AL-224

The study was conducted in accordance with the Declaration of Helsinki. Informed consent for the data and sample collection was obtained from the patient at presentation with the approval of the Institutional Review Board at Boston Medical Center. Bone marrow aspirate, clinical information, laboratory data, and post-mortem heart tissue were obtained from the sample biorepository and patient database maintained by the Boston University Amyloidosis Center.

Detailed clinical and laboratory characteristics of patient AL-224 are provided in Table S1 and in[19]. Briefly, a 63-year-old female with treatment naïve AL amyloidosis presented with cardiac and soft tissue involvements; λ LC restriction; circulating clonal IgG and λ free LC (FLC) on serum immunofixation electrophoresis; abnormal FLC κ/λ ratio; and λ FLC on urine immunofixation electrophoresis. The abdominal fat aspirate and endomyocardial and bone marrow biopsies were positive for amyloid by Congo red staining imaged using light microscopy in bright and polarized light.

The amyloid fibril typing by either immunoelectron microscopy or mass spectrometry was not available at the time of patient's evaluation. The patient passed away of cardiac arrest one month after the initial evaluation. Post-mortem evaluation demonstrated interstitial and vascular amyloid deposition in multiple organs, including the heart. The unfixed post-mortem cardiac tissue was frozen at -80 °C; formalin-fixed paraffin-embedded cardiac tissue block was made and used for further studies.

Histological and immunoelectron microscopy analyses of cardiac tissue

Post-mortem cardiac tissue was evaluated for amyloid by Congo red staining / light microscopy. Amyloid typing in cardiac tissue was performed using immunogold electron microscopy with antibodies directed against λ and κ LCs as previously described[57]. Briefly, for primary antibodies we used polyclonal rabbit anti-human antibodies against λ and κ LCs (Agilent Tech, Cat. #A019102-2 and #A019302-2). Secondary goat anti-rabbit IgG antibody conjugated to 15 nm gold particles (Ted Pella, Cat. #15727) and goat anti-mouse IgG antibody conjugated to 20 nm gold particles (Ted Pella, Cat. #15753) were used. The micrographs were collected using a JEOL JEM1020 transmission electron microscope.

Gene sequencing

The *IGL* gene was cloned and sequenced from unselected bone marrow plasma cells as previously described[58]. Once the *IGLV* gene was identified, nucleotide sequence errors introduced by FR1 primers were corrected by additional PCR amplification with 5′ primer for the *IGLV6-57* leader region and a 3′ universal *IGLC* primer. The light chain sequence, AL-224L, was deposited in GenBank (accession number KY432418).

Amyloid fibril extraction from autopsied cardiac tissue

Fibrils were extracted from the tissue as previously described[26] using overnight tissue digestion with C. histolyticum collagenase (5 mg/ml in a buffer containing 20 mM Tris, 140 mM NaCl, 2 mM $CaCl_2$, pH 8.0), followed by 15 cycles of homogenization in Tris EDTA buffer (20 mM Tris, 140 mM NaCl, 10 mM EDTA, pH 8.0) to remove soluble proteins, and by repeated homogenization of the remaining pellet in 1 ml ultrapure water. The fibril-containing supernatants from the water homogenization cycles were retained. To evaluate the protein yield, each 1 ml fraction was quantified using a Pierce BCA Protein Assay Kit (Thermo Fisher Scientific). The cryo-EM and proteomic analyses were performed using the supernatant from the third water-extraction cycle (water fraction 3), which was visibly cloudy. 2D-PAGE and western blot analyses of amyloid fibril extract 2D SDS-PAGE was performed under denaturing and reducing conditions as previously described[26]. Briefly, proteins from water fraction 3 (200 μg) solubilized in the isoelectric focusing buffer (IEF) were first separated on 11 cm strips with immobilized non-linear pH gradient 3–10 (Bio-Rad) using a Bio-Rad Protean IEF cell, and then on 8–16% polyacrylamide gradient gels (Criterion TGX gels, Bio-Rad). Gels were stained with GelCode Blue Stain Reagent (Thermo Fisher Scientific) and imaged using an ImageQuant LAS 4000 (GE). Spots of interest were excised and the proteins were identified by mass spectrometry as described below. For western blotting, proteins were transferred overnight onto a PVDF membrane (Millipore) using Criterion Blotter (Bio-Rad) and probed with a polyclonal rabbit anti-human λ LCs antibody (Fortis Life Sciences) used at a concentration of 0.125 μg/ml.

Proteomic analysis of fibril extracts by LC-MS/MS

For proteomic analysis of the extracted fibrils from water fraction 3, 25 μg of proteins were subjected to in-solution digestion with MS-grade trypsin (Thermo Fisher Scientific; enzyme:protein ratio 1:20) overnight at 37 °C. For excised protein spots #15 and #16 from

the 2D SDS-PAGE, in-gel digestion was performed as described[26]. Samples were cleaned

up with Thermo SPE C18 tips and the LC-MS/MS analysis was conducted using Q

Exactive HF-X High Resolution Orbitrap (Thermo Fisher Scientific) coupled with Ultimate

3000 nanoLC (Thermo Fisher Scientific) at Harvard Center for Mass Spectrometry.

Peptides were trapped on a trapping cartridge (300 µm x 5 mm PepMap™ Neo C18 Trap

Cartridge, Thermo Fisher Scientific) prior to separation on an analytical column (µPAC,

C18 pillar surface, 50 cm bed, Thermo Fisher Scientific). Mobile phase system consisted

of water with 1% formic acid (A) and acetonitrile with 1% formic acid (B), using a gradient

elution for the total of 110 min (1-32% B over 80 min followed by 32-45% B over 20 min),

followed by washout with up to 95% B at a flow rate of 300 nl/min. The mass spectrometer

operated in positive mode and data dependent acquisition for all analyses. A full scan

ranging from 350 to 1400 m/z was performed with a mass resolution of 12×10^4. The top

three most intensive precursor ions from each scan were used for MS2 fragmentation with

normalized collision energy of 30 at a mass resolution of 3.0×10^4. Data were processed

using the Proteome Discoverer software (Thermo Fisher Scientific) version 3.1, using a

human protein database downloaded in November 2024 from UNIPROT and augmented

with the AL-224L sequence. The following criteria were used for identification: cleavage

enzyme trypsin (semi-specific); cysteines carbamidomethylation (as a static modification);

and methionine oxidation and conversion of N-terminal glutamine into pyroglutamate (as

dynamic modifications).

Short chain fatty acid (SCFA) quantification in fibril extracts

Fibril extracts were tested for the presence of SCFA (1-6 carbons), which were quantified

using an ELISA kit (AFG Scientific, Cat. #EK700010) following manufacturer's protocols.

Homogenized and collagenase-treated cardiac tissue (220 mg) was extracted into ten 1 ml

fractions as described above. Fraction 3 was used for cryo-EM analysis and fractions 1 to

5 were used for SCFA measurements. Each fraction was diluted 1:5 in the assay buffer

and was assayed using 10 µl/well. The assay was performed in technical and biological

duplicates. SCFA concentrations were interpolated from the standard curve ($R^2 = 0.97$).

The results are reported in Figure S6.

Assay Validation: To evaluate the antibody selectivity for SCFA vs. non-esterified fatty

acids (NEFA) and potential interference from extracellular matrix components in tissue

extracts, the following validation experiments were performed. (i) Fractions 1-5 were

analyzed to assess the overall assay performance and the interference from NEFA; (ii)

Fraction 3 (Fr 3) was spiked with 12 or 24 µmol/l SCFA to test the signal recovery and

interference; (iii) 20 µmol/l SCFA was mixed with 20 µmol/l palmitic acid (a representative

NEFA) to probe antibody selectivity; (iv) combined spike tests were performed using Fr3

alone, Fr3 + 20 µmol/l SCFA, Fr3 + 20 µmol/l NEFA, and Fr3 + 20 µmol/l SCFA + 20 µmol/l

NEFA. The results, reported in Table S4, indicated minimal interference from tissue

components and no cross-reactivity with NEFAs, with high signal recovery ranging from

85 to 95%. These results confirm the specificity and robustness of the ELISA

measurements of SCFA in cardiac tissue extracts.

The endogenous SCFA concentration measured by ELISA in fraction 3 was

38.7 ± 1.2 µmol/l (or ~0.17 µmol/g tissue). Total protein concentration in fraction 3

measured by a colorimetric bicinchoninic acid assay was ~3 mg/ml. Assuming that most

of the protein in fraction 3 was the fibril-forming LC fragment 1-116, an approximate molar

ratio of LC to SCFA was 3:1. Assuming that SCFA had unit occupancy in the single-filament

structure (acetate in Figure 6C), LC to SCFA molar ratio in this structure was 1:1.

Amyloid assay using thioflavin T fluorescence

Amyloid content in tissue fractions 1-5 was assessed using thioflavin T (ThT), a diagnostic

dye that shows increased fluorescence upon binding to amyloid. Fractions 1-5 were

diluted 1:5 with Tris NaCl buffer, pH 7.92, and 10 µM ThT was added. Fluorescence

emission was measured within 5 min of ThT addition using a Tecan Infinite M1000 Pro

plate reader (excitation wavelength λ_{ex}=450 nm, emission wavelength λ_{em}=480 nm). The

measurements were performed in triplicate and the results are shown in Figure S6.

Cryo-EM sample preparation

Freshly extracted fibrils (water fraction 3) were deposited on the grids immediately upon

isolation. UltrAuFoil R 1.2/1.3 copper 300 mesh grids (Quantifoil, N1-A14nAu30-50) were

glow-discharged for 45 sec at 15 mA using a PELCO easiGlow glow discharge system to

apply a negative charge to their surface. A 3 ml sample was deposited on the grid surface

and immediately blotted for 3 sec with a blot force of 3 and plunge-frozen in liquid ethane

using a Vitrobot Mark IV apparatus (Thermo Fisher) set to 100% humidity and 4°C.

Cryo-EM data collection

A cryo-EM dataset of AL-224L amyloid was collected at the Boston University Cryo-EM Core Facility using a Glacios II at 200 kV equipped with a Falcon 4i direct electron detector, Selectris energy filter, and fringe-free imaging. A dataset of 10,309 exposures was collected using Thermo Fisher EPU in counting mode and recorded in the Electron Event Representation (EER) format using a magnification of 130,000x, pixel size of 0.90 Å, total dose of $40\,e^-Å^{-2}$, dose rate of $10.82\,e^-$ per pixel per second, defocus range of −0.8 to −2.5 µm, and energy filter slit width of 10 eV. The imaging rate was approximately 550 exposures per hour. A multishot imaging strategy was used to collect four exposures per grid hole, utilizing beam image shift to move between acquisition areas. Grid screening of the amyloid sample prior to data collection revealed fibrils more abundant near the edges of the foil holes. To maximize the number of fibrils visible in each exposure, the corner of each of the four acquisition areas was positioned partially over the gold film surface.

Cryo-EM data processing

The dataset was processed in cryoSPARC v4.5.1[59]. Exposures were imported with an EER upsampling factor of 2 then cropped by one-half back to their original resolution using patch motion correction. The contrast transfer function (CTF) correction was performed with 'patch CTF estimation'. Poor-quality micrographs were removed, yielding 7,076 high-quality micrographs. Template-free 'Filament tracer' and 'inspect picks' were used to pick filament segments with a separation distance of 45 Å from a 1000 micrograph subset of the data. Particles were extracted at a box size of 896 px and Fourier-cropped to 448 px using 'extract mics' to yield a partial uncleaned particle stack (71,140 particles). A set of 2D templates was generated with '2D classification' and 'select 2D'. The 2D templates were used with 'filament tracer' and 'inspect picks' to pick filament segments with a separation distance of 45 Å from the full 7,706 micrograph dataset. Particles were extracted at a box size of 900 px and Fourier-cropped to 450 px using 'extract mics' to yield a full uncleaned particle stack (421,024 particles). Particle cleaning was performed primarily using 2D classification. Three rounds of '2D classification' and 'select 2D' were performed to remove junk particles and double protofilaments yielding a partially cleaned particle stack (236,744 particles). An initial EM map of single protofilament AL-224L was generated using 'ab-initio reconstruction' by asking for an output of three volumes and selecting the best quality volume. An initial refinement was performed with 'helix refine' to obtain a particle stack with estimated 3D alignments. The partially cleaned, partially

aligned particle stack was extracted again at a smaller box size of 400 px. These particles

from a smaller box size were cleaned one last time using '2D classification' and 'select 2D'

to yield a cleaned particle stack (214,351 particles).

A 3D map of the cleaned particle stack was generated with 'reconstruct only' and was

used as an input for 'helix refine'. 'Helix refine' was used to generate an initial refined EM

map of the single-filament amyloid. In this first 'helix refine' job, symmetry

searching/enforcement was disabled, and the initial model was lowpass filtered to a

resolution of 8 Å. The output volume, which was used as an input volume in the next step,

showed bumps that correspond to stacked β-sheets. These β-sheet features enabled the

symmetry search utility in the next step to find the correct helical rise/twist values. 'Helix

refine' was performed again with symmetry search/enforcement enabled to produce a

Symmetry Enforced Structure. The initial model was lowpass filtered to 4.5 Å, and the twist

and rise parameters were searched over -5.2° to 3.2° and 0.8 Å to 8.8 Å, respectively.

Helical symmetry search and enforcement was set to begin at 4 Å to ensure that symmetry

averaging was applied only after the β-sheet separation along the fibril z-axis was visible.

The symmetry was calculated to have a helical rise of 4.86 Å, which is typical of amyloid

fibrils, and a twist of 1.20°. Individual particle CTF and individual particle motion was

corrected with 'reference-based motion correction'. The final EM map was generated with

a final round of 'helix refine' with symmetry search/enforcement enabled (214,100

particles, 2.92 Å, sharpening B-factor of -25.8).

Cryo-EM model building

An atomic model for AL-224L amyloid was built and refined against the sharpened EM

map. Initially, a single polyalanine chain was built into a central section across the fibril z-

axis and manually adjusted using Coot[60]. The individual residues were mutated to

correspond to the amino acid sequence of AL-224L determined by mass spectrometry.

The amino acid assignment within the EM map was determined by carefully examining

the path of the backbone, the position of the well-defined disulfide bond, C22-C91, and

the strong EM density for side chains that often corresponds to aromatic residues.

Residues 42-58 and 105-216 were not modeled because of weak EM density, suggesting

conformational heterogeneity in these segments.

The single molecular layer was duplicated to create a five-layer model. The replicated

molecules were rigid fit within the EM map above and below the original strand using the

fit-in-map function in ChimeraX[61]. The model was further refined in PHENIX[62] using

phenix.real_space_refine[63]. The β-sheet secondary structure, disulfide bond, and non-

crystallographic symmetry were manually defined in PHENIX to help guide the refinement

and impose identical conformations across all five molecular layers. The model statistics

and map-model fit were iteratively improved with multiple rounds of real-space refinements

in PHENIX and manual adjustments in Coot. To prevent the model from getting trapped in

the wrong conformation at an early stage, initial real-space refinement runs were

performed at lower map weight values, while final real space refinement runs were

performed at higher map weight values. The model was validated using the cryo-EM

Comprehensive Validation module in PHENIX running MolProbity[64].

To determine the handedness of the cryo-EM map and model of AL-224L amyloid fibrils,

we generated a highly sharpened cryo-EM map (sharpening B-factor of -200) and checked

if the orientation of the backbone carbonyl atoms matched the corrugation of the backbone

cryo-EM density. This approach was described for amyloid-like FUS fibrils[43] and is suitable

for high-resolution cryo-EM maps. In our structure of AL-224L amyloid, the right-handed

map and model had a much better map-model fit than the left-handed map and model.

Figures were generated with ChimeraX[61].

Validation of additional densities using false discovery rate (FDR) thresholding

To validate the significance of the three orphan densities seen in the cryo-EM map at the

internal and external sites, we generated a confidence map using FDR thresholding[37] with

default parameters and without incorporating local resolution and atomic model

information. We performed the map generation using the unmasked unsharpened AL-

224L cryo-EM map as the input and generated a figure viewed at a threshold of

0.99999999, equivalent to a 0.000001% false discovery rate for visible voxels

(Supplemental Fig. 1). The three densities were visible in the map indicating that their

probability of being noise was below 0.000001%.

Methods References

57. Prokaeva, T. *et al*. A novel substitution of proline (P32L) destabilises β2-microglobulin inducing hereditary systemic amyloidosis. *Amyloid* **29**, 255–262 (2022).

58. Klimtchuk, E. S., Prokaeva, T. B., Spencer, B. H., Gursky, O. & Connors, L. H. In vitro co-expression of human amyloidogenic immunoglobulin light and heavy chain proteins: a relevant cell-based model of AL amyloidosis. *Amyloid* **24**, 115–122 (2017).

59. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296 (2017).

60. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010).

61. Pettersen, E. F. *et al*. UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**, 70–82 (2021).

62. Liebschner, D. *et al*. Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861–877 (2019).

63. Afonine, P. V. *et al*. Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol* **74**, 531–544 (2018).

64. Williams, C. J. *et al*. MolProbity: More and better reference data for improved all-atom structure validation. *Protein Sci* **27**, 293–315 (2018).

903 # EXTENDED DATA

Characteristics	AL-224	Reference range
Demographic features		
Gender	F	
Age, years	63	
Age at death, years	63	
Cause of death	sudden death	
Echocardiographic features		
Interventricular septal thickness (mm)	11	
Ejection fraction, %	60	
Laboratory parameters		
Bone marrow plasma cells, %	n/a	
LC restriction	λ	
SIFE	IgG+ λ FLC	
FLC κ	9.20	3.3–19.4
FLC λ	175.00	5.7–26.3
FLC ratio	0.05	0.26–1.65
Abnormal HLC, isotype	IgG	
IgG HLC κ	1.6	3.84–12.07
IgG HLC λ	8.49	1.91–6.74
IgG HLC ratio	0.19	1.12–3.21
UIFE	λ FLC	
BNP, pg/ml	126	0-176
Troponin I, ng/ml	0.046	<0.033
Serum creatinine, mg/dl	0.91	0.7–1.3
eGFR, ml/min/1.73 m^2	71	
24-h urine protein, mg	469	
Alkaline phosphatase, U/l	127	25–100
Organ involvement		
Cardiac	+	
Soft tissue	+	
Congo red positive histology		
Fat pad aspirate	+	
Bone marrow biopsy	+	

905 **Table S1**. **Demographic, clinical and laboratory characteristics of case AL-224 with AL amyloidosis at**
906 **initial evaluation.** Abbreviations: BNP, B-type natriuretic peptide; eGFR, estimated glomerular filtration rate;
907 FLC, free light chain; HLC, heavy light chain; Ig, immunoglobulin; LC, light chain; SIFE, serum immunofixation
908 electrophoresis; UIFE, urine immunofixation electrophoresis.

Accession	Description	Coverage [%]	# PSMs	# Unique Peptides	# AAs	MW [kDa]
P12883	Myosin-7 OS=Homo sapiens OX=9606 GN=MYH7 PE=1 SV=5	76	841	133	1935	223
P13533	Myosin-6 OS=Homo sapiens OX=9606 GN=MYH6 PE=1 SV=5	50	521	3	1939	223.6
AL-224L	Amyloidogenic LC sequence AL-224L OS=Homo sapiens	74	373	40	218	23.3
Q8WZ42	Titin OS=Homo sapiens OX=9606 GN=TTN PE=1 SV=4	7	257	224	34350	3813.7
P62736	Actin, aortic smooth muscle OS=Homo sapiens OX=9606 GN=ACTA2 PE=1 SV=1	70	187	39	377	42
P12882	Myosin-1 OS=Homo sapiens OX=9606 GN=MYH1 PE=1 SV=3	25	157	4	1939	223
P35609	Alpha-actinin-2 OS=Homo sapiens OX=9606 GN=ACTN2 PE=1 SV=1	44	112	42	894	103.8
P60709	Actin, cytoplasmic 1 OS=Homo sapiens OX=9606 GN=ACTB PE=1 SV=1	47	110	5	375	41.7
P06576	ATP synthase subunit beta, mitochondrial OS=Homo sapiens OX=9606 GN=ATP5F1B PE=1 SV=3	70	99	58	529	56.5
Q14896	Myosin-binding protein C, cardiac-type OS=Homo sapiens OX=9606 GN=MYBPC3 PE=1 SV=4	38	93	64	1274	140.7
P12111	Collagen alpha-3(VI) chain OS=Homo sapiens OX=9606 GN=COL6A3 PE=1 SV=5	16	76	53	3177	343.5
P25705	ATP synthase subunit alpha, mitochondrial OS=Homo sapiens OX=9606 GN=ATP5F1A PE=1 SV=1	54	63	41	553	59.7
P17540	Creatine kinase S-type, mitochondrial OS=Homo sapiens OX=9606 GN=CKMT2 PE=1 SV=2	39	57	30	419	47.5
P04004	Vitronectin OS=Homo sapiens OX=9606 GN=VTN PE=1 SV=1	25	52	17	478	54.3
Q08043	Alpha-actinin-3 OS=Homo sapiens OX=9606 GN=ACTN3 PE=1 SV=2	14	52	3	901	103.2
P06727	Apolipoprotein A-IV OS=Homo sapiens OX=9606 GN=APOA4 PE=1 SV=4	55	45	37	396	45.3
P02511	Alpha-crystallin B chain OS=Homo sapiens OX=9606 GN=CRYAB PE=1 SV=2	65	36	23	175	20.1
Q9Y2K3	Myosin-15 OS=Homo sapiens OX=9606 GN=MYH15 PE=1 SV=6	3	32	2	1926	222
P12109	Collagen alpha-1(VI) chain OS=Homo sapiens OX=9606 GN=COL6A1 PE=1 SV=3	19	29	20	1028	108.5

P48735	Isocitrate dehydrogenase [NADP], mitochondrial OS=Homo sapiens OX=9606 GN=IDH2 PE=1 SV=2	33	28	15	452	50.9
P07437	Tubulin beta chain OS=Homo sapiens OX=9606 GN=TUBB PE=1 SV=2	34	28	3	444	49.6
P68371	Tubulin beta-4B chain OS=Homo sapiens OX=9606 GN=TUBB4B PE=1 SV=1	38	28	4	445	49.8
P17661	Desmin OS=Homo sapiens OX=9606 GN=DES PE=1 SV=3	33	27	15	470	53.5
P12814	Alpha-actinin-1 OS=Homo sapiens OX=9606 GN=ACTN1 PE=1 SV=2	14	27	6	892	103
P55809	Succinyl-CoA:3-ketoacid coenzyme A transferase 1, mitochondrial OS=Homo sapiens OX=9606 GN=OXCT1 PE=1 SV=1	28	27	19	520	56.1
P54296	Myomesin-2 OS=Homo sapiens OX=9606 GN=MYOM2 PE=1 SV=3	15	23	19	1465	164.8
P21796	Non-selective voltage-gated ion channel VDAC1 OS=Homo sapiens OX=9606 GN=VDAC1 PE=1 SV=2	57	22	16	283	30.8
P12110	Collagen alpha-2(VI) chain OS=Homo sapiens OX=9606 GN=COL6A2 PE=1 SV=4	16	21	18	1019	108.5
P45880	Voltage-dependent anion-selective channel protein 2 OS=Homo sapiens OX=9606 GN=VDAC2 PE=1 SV=2	34	21	10	294	31.5
P02649	Apolipoprotein E OS=Homo sapiens OX=9606 GN=APOE PE=1 SV=1	34	21	15	317	36.1

Table S2. Top 30 master proteins identified with at least two unique peptides of heart-tissue extracted
fibrils by LC-MS/MS.
Abbreviations: PSM, peptide-spectrum match; #AAs, number of amino acids in full-length protein; MW,
molecular weight. The proteins are ranked according to the number of PSMs. Fibril-forming protein AL-224L
(pink), collagen VI chains a1, a 2 and a 3 (green), and the amyloid signature proteins apoA-IV and apoE
(orange) are highlighted. Serum amyloid P-component was detected in low abundance (#39, not shown). The
data normalization using normalized spectral abundance factor (NSAF) values demonstrated AL-224L as the
top identified protein (data not shown). To calculate the NSAF value for each protein, the spectral counts were
divided by the protein length (PSM/#AAs) and the sum of all PSM/#AA values was divided by the number of
all identified proteins as previously reported (Neilson KA, Keighley T, Pascovici, D, Cooke B, Haynes PA.
(2013) Label-Free Quantitative Shotgun Proteomics Using Normalized Spectral Abundance Factors. Methods
Mol Biol. 2013:1002: 205-222).

LC sub-family	PDB ID	ΔG, kcal/mol
λ3	9FAA	-33.7
λ3	9FAB	-33.4
λ3	9FAC	-34.2
λ3	8R47	-26.4
λ3	9EME	-25.5
λ1	7NSL	-26.6
λ1	6IC3	-24.2
λ3	6Z1O	-21.5
λ3	6Z1I	-20.4
λ6 AL55 renal	8CPE	-23.3
λ6 AL55 cardiac	6HUD	-20.2
λ6 AL-224L cardiac	**9OKA**	**-18.0**
λ6 germline	6HUD	-18.6
λ6 germline	9OKA	-22.1

Table S3. Solvation energy calculated for the available amyloid structures of AL LCs.
ΔG values represent free energy change upon interface formation between two fibril layers, which was
estimated using Protein Interfaces, Surfaces and Assemblies (PDBePISA) server, https://www.ebi.ac.uk/msd-
srv/prot_int/cgi-bin/piserver. Bottom two rows list hypothetical values for the λ6-LC germline sequence placed
in the amyloid structure of cardiac AL55 LC (PDB: 6HUD) or cardiac AL-224L (PDB: 9OKA). Mutations to
restore the germline sequence were introduced using the Swiss-PdbViewer software, https://spdbv.unil.ch/
and the structure was energy-minimized to relieve steric clashes.

Sample	SCFA µM	Recovery µM	Recovery %
SCFA	19.97		
NEFA	0		
SCFA + NEFA	21.89	21.89	109.6
Fr3	**38.71**		
Fr3 + NEFA	36.75	36.75	95.0
Fr3 + SCFA	56.92	18.21	91.1
Fr3 + SCFA + NEFA	88.26	51.49	74.9

Table S4. Short chain fatty acid content in the amyloid-containing tissue extracts (fraction 3) measured by ELISA and assay validation.

Abbreviations: SCFA - short chain fatty acid; NEFA – non-esterified fatty acid; Fr3 – tissue-extracted fraction 3 used for cryo-EM analysis of amyloid. The experimental details are reported in Methods.

939
940
Figure S1. Histological analyses of the autopsied cardiac tissue in case AL-224.
A, Light microscopy images of Congo red-stained cardiac tissue indicate amyloid deposits. Original
magnification ×100. **B**, Areas in panel A viewed by polarized microscopy show amyloid deposits with
characteristic green birefringence. Original magnification ×100. **C, D**, Electron micrographs of autopsied
cardiac tissue show haystack-like organization of amyloid fibrils in the extracellular space. Immunogold
labeling demonstrates numerous electron dense deposits with antibody directed against immunoglobulin λ-
LC (original magnification ×20,000, **(C)** and no immunoreactivity with antibody directed against
immunoglobulin κ-LC (original magnification ×15,000, **(D)**.

949
950
Figure S2. Analysis of AL-224 proteins in water fraction 3 of cardiac tissue fibril extract using 2D SDS-PAGE, 2D western blotting and LC-MS/MS.
A, Coomassie-stained gel and **B**, the corresponding 2D western blot probed with a primary anti-human λ light chains antibody. Two most prominent spots are circled and marked #15 and #16. These spots were excised and analyzed by LC-MS/MS, along with the entire water fraction 3. **C**, Amino acid sequence of AL-224L shows the position (in green) and sequence (underlined bold) of the peptides identified in spots #15, #16 and in the entire water fraction 3 (indicated).

Patient-derived cardiac amyloid AL224L cryo-EM dataset

Figure S3. Cryo-EM data processing workflow.
General processing pipeline to generate an EM map of an AL-224L single-filament amyloid polymorph.

Figure S4. **Global resolution and map-model fit evaluation of amyloid structure.**
A, Fourier shell correlation (FSC) plot at a gold standard 0.143 cutoff of cardiac amyloid AL-224L.
B-E, AL-224L model superimposed over the EM map to show the map-model fit at EM map at thresholds as
indicated.

A

Left-handed map and model

B

Right-handed map and model

966
Figure S5. **Backbone corrugations reveal the AL-224L amyloid to be right-handed.**
A, Slice-view showing a superimposition of the left-handed amyloid model over the highly sharpened left-
handed amyloid map. Green check icon indicates that the right-handed map and model have correct
handedness. **B,** Slice-view showing a superimposition of the right-handed amyloid model over the highly
sharpened right-handed amyloid map. Green check icon indicates that the right-handed map and model have
correct handedness.

Figure S6. Amyloid and short chain fatty acid (SCFA) content in fractions 1-5 extracted from autopsied cardiac tissue AL-224.
SCFA were measured by ELISA in technical and biological duplicates; mean values $\pm$ SEM are shown. Amyloid was assessed by measuring fluorescence emission of a diagnostic dye thioflavin T (ThT) that shows increased fluorescence upon binding to amyloid. Methods provide experimental details.

979 **SUPPLEMENTAL DATA**

A

B

Supplemental Figure 1. **False Discovery Rate (FDR) Threshold Confidence Map.**
A, Slice-views of the unsharpened AL-224L amyloid input EM map to FDR thresholding. White boxes indicate
the 3D area that was sampled to produce a noise estimate to generate a confidence map. **B,** Traditional
unsharpened AL-224L amyloid EM map (blue) within an unsharpened AL-224L amyloid EM confidence map
(purple) generated with FDR thresholding and viewed at a threshold of 0.99999999, equivalent to a
0.000001% false discovery rate for visible voxels.