Chain-length regulation by WzzE is necessary for, but genetically separable from, cyclic enterobacterial common antigen synthesis

Joseph F. Carr[a], Jennifer S. Rudolf[b]*, Daniel J. Warzecha[c]*, Yohannes H. Rezenom[d], & Angela M. Mitchell[e]#

[a] Department of Biology, Texas A&M University, College Station, Texas, USA (jfcarr@tamu.edu)

[b] Department of Biology, Texas A&M University, College Station, Texas, USA (jsrudolf@tamu.edu)

[c] Department of Biology, Texas A&M University, College Station, Texas, USA (dwarzecha03@tamu.edu)

[d] Chemistry Mass Spectrometry Facility, Texas A&M University, College Station, Texas, USA

[e] Department of Biology, Texas A&M University, College Station, Texas, USA (amitchell@bio.tamu.edu)

* JSR and DJW contributed equally to this work. Author order was determined alphabetically.

Address correspondence to Angela M. Mitchell, amitchell@bio.tamu.edu

Running title: Biosynthesis of cyclic enterobacterial common antigen

Keywords: gram negative envelope, enterobacterial common antigen, O antigen, Wzz, glycan biosynthesis, chain length regulation

Funding Statement: This work was supported by the National Institute of Allergy and Infectious Disease under award number R01-AI155915 (to A.M.M.).

Conflict Statement: The authors declare that they have no conflicts of interest.

Abstract

Enterobacterial common antigen (ECA) is a conserved glycan that supports outer membrane

impermeability and intrinsic antibiotic resistance in Enterobacterales. ECA exists in outer

membrane diacylglycerol-phosphate- and lipopolysaccharide-linked forms, and a cyclic

periplasmic form (ECA_{CYC}). Intriguingly, ECA_{CYC} both affects the outer-membrane permeability

barrier and functions in regulation of diacylglycerol-phosphate-linked ECA levels. While the

length of linear ECA polymers generated by WzyE is regulated by the co-polymerase WzzE,

WzzE is also required for ECA_{CYC} biogenesis and no ECA_{CYC} is synthesized in its absence. To

uncover WzzE functions necessary for ECA_{CYC} biosynthesis, we generated plasmid-borne and

chromosomal *wzzE* mutants in *Escherichia coli* K-12 and quantified their effects on linear chain-

length regulation and on ECA_{CYC} synthesis. Mutations affecting linear ECA chain-length

regulation in either transmembrane helix 2 or the periplasmic domain abolished ECA_{CYC}

synthesis. Notably, two variants at residue F104 exhibited identical linear ECA chain-length

regulation which was similar to wild type, but resulted in sharply different ECA_{CYC} production:

chromosomal $WzzE_{F104Y}$ produced near wild-type levels of ECA_{CYC}, whereas $WzzE_{F104H}$

produced approximately two-fold less ECA_{CYC}. This difference in ECA_{CYC} production vs. linear

chain-length regulation demonstrates ECA_{CYC} synthesis is genetically separable from this

regulation. Thus, WzzE-mediated chain-length regulation is necessary for ECA_{CYC} biogenesis,

but not sufficient.

Introduction

Enterobacterales is an order of gram-negative bacteria that includes numerous genera

with human pathogens (e.g., *Escherichia, Salmonella, Klebsiella, Yersinia*) (Doern, 2024,

Adeolu et al., 2016). These organisms cause a variety of diseases including respiratory, soft

tissue, and urinary tract infections, as well as endocarditis (Ramirez and Giron, 2024). To

survive in their enteric environment, members of this family take advantage of their outer

membrane (OM) which acts as a robust permeability barrier composed of an asymmetric bilayer

with phospholipids in the inner leaflet and predominantly lipopolysaccharides (LPS) in the outer

leaflet (Maher and Hassan, 2023, Silhavy et al., 2010). LPS molecules form an extensive

hydrophilic network of lateral interactions that excludes many large and hydrophobic

compounds—such as many antibiotics, bile salts, and detergents—from entering the cell

(Farfour et al., 2022, Aminov, 2010).

Beyond LPS, the outer membrane is decorated with other components critical for

fitness, including enterobacterial common antigen (ECA) (Bennett and Mitchell, 2025, Rai and

Mitchell, 2020). Identified in 1962 by Calvin M. Kunin et al. as a "common antigen" associated

with urinary tract infections, ECA is an invariant carbohydrate restricted to *Enterobacterales* and

absent only in endosymbionts with significantly reduced genomes (Rai and Mitchell, 2020,

Jenkins et al., 2017, Kunin, 1963, Kunin et al., 1962). ECA has been shown to play a role in the

pathogenesis in several members within the order of *Enterobacterales* (Aguiniga et al., 2016,

Kalynych et al., 2014, Gilbreath et al., 2012, Armbruster et al., 2017, Holmes et al., 2023,

Guérin et al., 2020). For example, *Salmonella enterica* serovar Typhimurium LT2 strains lacking

ECA display reduced virulence, heightened sensitivity to bile salts, and establish a low-level

persistent infection in mouse models (Kalynych et al., 2014, Gilbreath et al., 2012).

ECA is defined by recurring trisaccharide repeat units (RUs) of GlcNAc (*N*-acetyl-D-

glucosamine), ManNAcA (*N*-acetyl-D-mannosaminuronic acid), and Fuc4NAc (4-acetamido-4,6-

dideoxy-D-galactose) (Mannel and Mayer, 1978). There are three forms of ECA, (i) ECA$_{LPS}$,

which replaces O-antigen (Oag) on the LPS core; (ii) ECA_{PG}, a cell-surface form linked to

diacylglycerol-phosphate; and (iii) ECA_{CYC}, a soluble, cyclic form found in the periplasm

(Goździewicz et al., 2015, Kunin, 1963, Mannel and Mayer, 1978, Kajimura et al., 2005). ECA_{PG}

and ECA_{LPS} vary in the number of RUs with a range of 1 to 14 and has a modal value of 5 to 7

at 37°C (Mitchell et al., 2018, Barr et al., 1999), whereas ECA_{CYC} has a defined number of

exactly 4 RUs in *Escherichia coli* K-12 (Kajimura et al., 2005). Structural analysis studies

including crystallography, nuclear magnetic resonance (NMR), and molecular dynamics indicate

that ECA_{CYC} adopts either a rhomboidal or square conformations (Färnbäck et al., 2003, Erbel

et al., 2004). We have previously found that ECA_{CYC} contributes to OM integrity in *E.coli* K-12

and mediates resistance to detergents and bile salts (Mitchell et al., 2018).

ECA biosynthesis genes reside mainly in the *wec* operon and employs a pathway

analogous to the Wzy/Wzz-dependent pathway of Oag synthesis (Eade et al., 2021, Rick et al.,

1985, Barr and Rick, 1987, Barr et al., 1989, Bertani and Ruiz, 2018, Barr et al., 1999, Erbel et

al., 2003). After a single undecaprenyl-phosphate-linked ECA RU is synthesized, WzxE flips the

intermediate to the outer leaflet of the inner membrane (Rai and Mitchell, 2020). WzyE and

WzzE form a complex where the translocated undecaprenyl-phosphate-linked ECA subunits are

polymerized by WzyE and the number of RU in the resulting chain is regulated by WzzE (Leo et

al., 2021b, Maczuga et al., 2022b, Nath and Morona, 2015, Weckener et al., 2023). WzzE is an

octameric inner-membrane protein with a large periplasmic domain (**Figure S1**) (Tocilj et al.,

2008, Kalynych et al., 2012b, Collins et al., 2017, Kalynych et al., 2015). A cryo-EM structure

from *Pectobacterium atrosepticum* suggest that WzyE resides within the central lumen formed

by WzzE's transmembrane helices (Weckener et al., 2023).

WzzE is a member of the class 1 polysaccharide co-polymerase (PCP1) family of chain

length regulators, which includes WzzB, FepE, Wzz among others (Wiseman et al., 2021,

Morona et al., 2009, Wiseman et al., 2023). Despite low sequence identity, PCP1 proteins share

a conserved architecture consisting of two transmembrane helices and a bell-shaped

periplasmic domain. (Tocilj et al., 2008, Kalynych et al., 2012b, Collins et al., 2017, Kalynych et

al., 2015). In *Shigella flexneri*, WzzB C-terminal transmembrane helix two (TMH2) is known to

interact with WzyB, the polymerase, and is necessary for Oag chain-length regulation (Ascari et

al., 2022, Leo et al., 2021a). WzzB and WzyB co-purify, and the TMH2 of WzzE can maintain

this interaction and contribute to Oag chain length regulation (Leo et al., 2021b). It has also

been shown that mutations disrupting Oag chain-length regulation are found throughout Wzz

and are not restricted to TMH2 (Whitfield et al., 2020).

WzzE plays a more complex role in ECA synthesis than other PCP1s do in Oag

production. Loss of WzzE generates linear ECA with increased heterogeneity—producing ECA

with both shorter and longer RUs—rather than the pronounced shortening typical of PCP1

deletions in Oag pathways (Leo et al., 2021b, Ascari et al., 2022, Huszczynski et al., 2019,

Kalynych et al., 2011). In contrast to linear ECA, strains lacking WzzE fail to produce ECA_{CYC},

indicating that WzzE is required for ECA_{CYC} synthesis despite a lack of known catalytic activity

(Rai and Mitchell, 2020, Mitchell et al., 2018, Kajimura et al., 2005). In addition, WzzE plays a

role in regulating levels of ECA_{PG} in *E. coli* K-12. ElyC regulates production of ECA_{PG} in such a

way that loss of ElyC causes large increases in ECA_{PG} production (Rai et al., 2021). This

regulation is dependent on WzzE and likely on ECA_{CYC}, and changes in ECA_{PG} levels with loss

of ElyC are much less in the absence of WzzE (Mitchell et al., 2018, Rai et al., 2021). Finally,

ECA takes on aberrant activity in the absence of the anterograde phospholipid transporter,

YhdP, causing increased sensitivity to vancomycin and the detergent SDS, and this phenotype

is reversed when *wzzE* is absent (Mitchell et al., 2018).

Several non-exclusive hypotheses for how PCPs regulate chain length have been

proposed (Islam and Lam, 2014, Leo et al., 2021b, Wiseman et al., 2021, Wiseman et al., 2023,

Kintz and Goldberg, 2011, Papadopoulos and Morona, 2010, Morona et al., 1995, Weckener et

al., 2023). (i) The inner or outer surface of the PCP may act as a molecular ruler for chain length

regulation (Kintz and Goldberg, 2011). Specifically, the PCP may bind the extending

carbohydrate, facilitating catalysis until the polymer length breaks the interaction (Kalynych et

al., 2012a). (ii) The PCP could serve as a scaffold on which multiple polymerase molecules

assemble (Tocilj et al., 2008). (iii) The elongating carbohydrate could extend in the lumen of the

PCP until it reaches the top of the periplasmic domain and is exposed (Islam and Lam, 2014).

(iv) The PCP could interact with the polymerase to sequester short glycan chains from

downstream ligases (i.e., WaaL), increasing the processivity of polymerization (Hong et al.,

2023). Finally, (v) the loops at the top of the periplasmic bell may move in an alternating up and

down arrangement which allows for the polysaccharide chain to extend using a "ratchet"-type

mechanism (Wiseman et al., 2023). However, the mechanism(s) used by WzzE for ECA_{CYC}

synthesis is unknown. Here, we used site-directed mutagenesis and biochemical analysis to

test whether WzzE requires only polymerase-interaction/chain-length regulation to produce

ECA_{CYC}, or if an additional function of WzzE contributes to ECA_{CYC} biogenesis. Our findings

demonstrate that polymerase interaction and chain-length regulation are necessary for ECA_{CYC}

biosynthesis. Moreover, we found that ECA_{CYC} synthesis genetically separated from linear ECA

regulation, suggesting an additional function of WzzE in ECA_{CYC} synthesis.

Results

Plasmid-based *wzzE* allows analysis of *wzzE* mutant function. In order to define the

functions of WzzE required for ECA_{CYC} synthesis in *E. coli* K-12, we examined the effects of

wzzE mutations analogous to known residues affecting PCP1 Oag regulation. We first

generated a set of *wzzE* constructs constitutively expressed by the *tet* promoter on the low copy

number pZS21 plasmid (Lutz and Bujard, 1997), which does not contain a Tet repressor (**Figure**

1A). These included wild-type *wzzE* (p*wzzE*) and C-terminal fusions bearing a glycine-serine

linker and triple FLAG tag (p*wzzE*-FLAG), a translation fusion to chloramphenicol

acetyltransferase (p*wzzE-cat*), and a p*wzzE-cat* triple FLAG tag fusion (p*wzzE-cat*-FLAG).

Based on the topology of WzzE, we anticipated that the C-terminal fusions would be exposed in

the cytoplasm, thereby allowing Cat to function. Thus, these constructs allowed us to potentially

identify defects in WzzE's folding or stability as measured by chloramphenicol resistance. This

approach was informed by a translational coupling methodology previously used a tool in *E. coli*

(Mendez-Perez et al., 2012, Sieber, 2003), in which an unstable or degraded protein construct

was shown to not allow chloramphenicol resistance. Additionally, we hypothesized that the

formation of the functional cat trimer (Andreeva et al., 2000) is more likely if WzzE is properly

multimerized, allowing Cat monomers to come into contact (with up to two Cat trimers

potentially formed per octamer), giving insight into potential disruptions in octamerization. This

assay is particularly useful because the addition of the bulky Cat (26KDa) protein likely imposes

stress on WzzE folding. However, the chloramphenicol resistance results remain suggestive

due to the potential limitations of this assay, such as the multimerization of Cat affecting

octamer formation (i.e., a loss of chloramphenicol resistance is more indicative of a loss of

stability and/or multimerization than chloramphenicol resistance is indicative of multimerization).

We next assessed whether these constructs complemented Δ*wzzE.* In the absence of

yhdP, ECA causes sensitivity to vancomycin in a *wzzE* dependent manner (Mitchell et al.,

2018). Therefore, we expressed the *wzzE* constructs in a Δ*wzzE* Δ*yhdP* strain and tested for

vancomycin sensitivity. Our results demonstrated each construct complemented Δ*wzzE* by

restoring vancomycin sensitivity (**Figure 1B**)*.* Additionally, p*wzzE-cat* conferred chloramphenicol

resistance to the strain indicating that the Cat fusion was functional. To determine if the p*wzzE-*

FLAG construct could be detected, we immunoblotted for FLAG using GroEL as a loading

control (**Figure 1C**) and determined that the fusion protein could be clearly detected.

To determine whether linear ECA regulation was restored in the *wzzE* plasmid

constructs, we first performed immunoblots for linear ECA (**Figure 1D**) using a polyclonal ECA

antibody (Leo et al., 2021b). Our results revealed a range of ECA bands consistent with

previous publications (Mitchell et al., 2018, Rai et al., 2021, Morris and Mitchell, 2023), with

increasing molecular weight indicating molecules with more RUs. The distribution of ECA chain

lengths was wider in a Δ*wzzE* mutant (Lane 3) than in wild-type cells (Lane 2) with an increase

in longer and shorter ECA bands, indicating loss of chain-length regulation. We also observed

more linear ECA with Δ*wzzE* as we would expect from previous findings (Rai et al., 2021,

Mitchell et al., 2018). For simplicity, we will refer to these changes as "loss of chain-length

regulation". Expression of p*wzzE* (Lane 5) but not the empty vector (Lane 4) restored chain-

length regulation, albeit with a lower modal chain length than the wild-type strain and much less

linear ECA overall. These changes may suggest overactivity of WzzE due to multi-copy

expression. The same pattern was observed for the p*wzzE-cat* construct (Lane 7); however, the

p*wzzE*-FLAG and p*wzzE-cat*-FLAG constructs resulted in chain-length regulation closer to that

of wild type.

Finally, to test whether the plasmid-based *wzzE* constructs rescue ECA_{CYC} synthesis, we

expressed the p*wzzE* and empty vector plasmids in a Δ*wzzE* Δ*wecH* strain. WecH has been

shown to non-stoichiometrically acetylate ECA (Kajimura et al., 2006) and deleting *wecH* allows

ECA_{CYC} to be quantitated as a single mass species (Kajimura et al., 2006, Mitchell et al., 2018).

Unlike the two linear forms of ECA, ECA_{CYC} is uncharged and cannot be detected by

immunoblotting. Therefore, we grew cultures with either a normal (^{14}N) or heavy (^{15}N) nitrogen

source, combined labeled and unlabeled samples, purified ECA_{CYC}, and assayed ECA_{CYC} levels

by liquid chromatography-mass spectrometry (LC-MS). ECA_{CYC} has a defined size of 4 RU in *E.*

coli K-12 (Kajimura et al., 2005) and this molecule has an monoisotopic mass of 2428.9 Da in

the absence of acetylation. Labeling with heavy nitrogen causes a 12 Da increase in this weight

due to the presence of 12 nitrogen atoms in ECA_{CYC} (one in each sugar monomer) (Mitchell et

al., 2018) (**Figure S2AB**). For this type of assay, each peak consists of multiple minor peaks

due to the stochiometric incorporation of naturally occurring isotopes (e.g., ^{14}C) (see **Figure**

1EF, **S2C**). When labeled and unlabeled samples carrying p*wzzE* were combined, we detected

peaks at both the unlabeled and labeled m/z (**Figure 1E**). However, when the labeled p*wzzE*

sample was combined with an unlabeled sample carrying the empty vector, only the ^{15}N labeled

ECA$_{CYC}$ peak was detected (**Figure 1F**). ECA$_{CYC}$ production from the p*wzzE* and p*wzzE-cat*

constructs was very similar (**Figure S2C**). These data confirm plasmid-based expression of

wzzE can facilitate the synthesis of ECA$_{CYC}$, making this an appropriate system to investigate

the role of WzzE in ECA$_{CYC}$ biosynthesis.

Conserved glycines in TMH2 are necessary for WzzE function. After validating our WzzE

functional assays, we next examined the contribution of TMH2 on WzzE function. TMH2 has

been shown to mediate interactions between WzzB and Wzy in O-antigen synthesis (Leo et al.,

2021b), and prior mutagenesis studies of WzzB identified a conserved GXXXG motif in the

TMH2 region of WzzB from *Shigella flexneri Y* that was important for function (Daniels and

Morona, 1999), likely due to effects on helix-helix packing (Teese and Langosch, 2015, Prakash

et al., 2011, Mueller et al., 2014). A corresponding double mutant in *Shigella flexneri Y*

(G305A/G311A WzzB; G306A/G312A in *E. coli*) disrupts Oag chain length regulation

(Papadopoulos et al., 2016); therefore, we hypothesized that the analogous mutations to

GXXXG motif located in WzzE would play a significant role in mediating the helix-helix packing

of the transmembrane region and its interaction with WzyE.

To test this, we generated TMH2 mutations (**Figure S1B**) in the p*wzzE* constructs

described in **Figure 1A**. We first sought to determine whether the levels of WzzE protein

produced by p*wzzE*-FLAG or p*wzzE-cat*-FLAG were altered by the mutations (**Figure 2AB,**

Figure S3A). Our results showed a small but significant decrease in protein abundance for our

WzzE$^{GG333/4LL}$ in the WzzE-FLAG fusion (**Figure 2AB**), whereas no decrease is apparent in the

WzzE-Cat-FLAG fusion (**Figure S3A**). As the levels of the WzzE$^{GG333/4LL}$ variant are 82% of wild

type on average and as high as 98% in individual replicates, the biological relevance of this

decrease is unclear. To determine whether the mutations might have caused alterations to

protein stability or multimerization, we utilized our p*wzzE-cat* construct to assay

chloramphenicol resistance conferred by the fusion protein. None of our mutations showed any

chloramphenicol sensitivity (**Figure 2C, Figure S3B**), suggesting that they are largely stable

and multimerized, despite the small decrease in abundance for WzzE$^{GG333/4LL}$. Next, we set out

to determine if the mutations affected WzzE's ability to regulate ECA chain length by performing

ECA immunoblots. Three of the TMH2 mutations, WzzE^{A323G}, WzzE$^{GG333/4AA}$, and WzzE^{G339A}

(Lanes 4,5, and 7) retained wild-type ECA chain length regulation (**Figure 2DE**). In contrast,

WzzE$^{GG333/4LL}$ (Lane 6) markedly disrupted chain-length regulation, resembling or exceeding the

defect observed in our empty-vector control (Lane 3). These results identify the TMH2 GXXXG

region as a key determinant of WzzE-mediated ECA chain-length regulation

The TMH2 GXXXG motif affects ECA$_{CYC}$ synthesis. Given the strong defect of WzzE$^{GG333/4LL}$

in regulating ECA chain length, we assessed whether the TMH2 mutations impair ECA$_{CYC}$

biogenesis. We utilized LC-MS to detect and quantitate ECA$_{CYC}$ in purified samples as in **Figure**

1EF. We expressed the TMH2 mutants or controls in a $\Delta wzzE$ $\Delta wecH$ strain background and

labeled the wild type p$wzzE$ strain with ^{15}N, while the mutants were grown with ^{14}N. The

quantification demonstrated WzzE^{A323G}, WzzE$^{GG333/4AA}$, and WzzE^{G339A} did not significantly

change ECA$_{CYC}$ levels, whereas the WzzE$^{GG333/4LL}$ mutant produced significantly reduced levels

of ECA$_{CYC}$, similar to those of the empty vector control (**Figure 3A**). These data demonstrate

that the TMH2-dependent chain-length regulation is required for ECA$_{CYC}$ synthesis.

244   To visualize how these mutations might alter WzzE function, we modeled the

WzyE:WzzE(8) complex for wild-type WzzE and WzzE$^{GG333/4LL}$ using AlphaFold 3 (Abramson et

al., 2024) (**Figure 3B** & **S4**, **Table S1**). Alignment of monomers revealed a ~2 Å expansion

between TMH1 and TMH2 in WzzE$^{GG333/4LL}$ consistent with steric disruption of helix-helix

packing (**Figure 3B**). This structural shift provides a potential mechanistic basis for the loss of

both chain-length regulation and ECA$_{CYC}$ biosynthesis in our GXXXG mutant.

The periplasmic domain of WzzE functions in chain length regulation. After assessing the

importance of WzzE TMH2 for ECA$_{CYC}$ biosynthesis, we next sought to determine whether

chain-length regulating activities within the periplasmic domain of WzzE are also required for

ECA$_{CYC}$ biosynthesis. Twenty-one periplasmic residues are conserved among WzzE, FepE, and

WzzB in *E. coli*, suggesting that they are important for PCP1 function (Tocilj et al., 2008). We

also hypothesized that non-conserved residues could contribute to WzzE-mediated chain-length

regulation or ECA$_{CYC}$ synthesis. Therefore, we made mutations in a conserved cluster of

predominantly hydrophobic residues located in the core of the α/β domains—spanning strands

β1 and β4, helix α2 and the N terminus of helix α6 (see **Figure S1B**). Additionally, we targeted

non-conserved residues throughout WzzE in positions similar to those that conferred changes

to Oag chain length in other PCP1s (Tocilj et al., 2008).

We first characterized the periplasmic mutants (**Figure S1B**) by examining WzzE protein

levels using the p*wzzE*-FLAG construct. WzzE$^{\Delta F104}$ (an in-frame, single codon deletion) had

undetectable WzzE-FLAG protein levels, whereas all other mutants retained wild-type protein

accumulation (**Figure 4AB**). When we examined the levels of WzzE-Cat-FLAG for WzzE$^{\Delta F104}$,

we observed a faint band at the expected molecular weight of the fusion and a second, lower-

molecular-weight band, indicating cleavage of the C-terminal tag (**Figure S3C**). Given the

cleavage of the tags, no conclusion could be drawn for the stability of WzzE$^{\Delta F104}$. All other

protein levels were similar to those of wild type.

We next evaluated whether the *wzzE* mutations might affect protein stability and/or

multimerization by measuring chloramphenicol resistance conferred by the p*wzzE*-cat fusion.

WzzE^{K162A}, WzzE^{L259A}, and WzzE^{F104Y} caused partial or full chloramphenicol sensitivity in the

p*wzzE*-Cat fusion (**Figure S3D**) but showed full chloramphenicol resistance in the p*wzzE*-Cat-

FLAG fusion, which better complements linear chain length regulation (**Figure 4C**). These

differences suggest these variants may exhibit mild instability or reduced multimerization.

All other mutants conferred chloramphenicol resistance similar to wild type in both fusions

(**Figure 4C, S3D**). WzzE$^{\Delta F104}$ showed wild-type levels of chloramphenicol resistance, likely

reflecting cleavage of the C-terminal tag. We also tested an additional 18 periplasmic-domain

mutations that caused chloramphenicol sensitivity in the WzzE-Cat background, emphasizing

the importance of this domain for WzzE structural stability and multimerization (**Figure S5**).

After characterizing the effect of the periplasmic mutations on WzzE levels and stability,

we next examined how these periplasmic mutations affect ECA chain-length regulation (**Figure**

4DE). WzzE$^{\Delta F104}$ and WzzE^{K162A} (lanes 4 and 7) disrupted chain-length regulation when

expressed from p*wzzE* (**Figure 4D**). However, when expressed from p*wzzE*-cat-Flag,

WzzE$^{\Delta F104}$, exhibited partial impairment of chain-length regulation, while WzzE^{K162A} showed wild-

type chain-length regulation (**Figure 4E**). These differences parallel the chloramphenicol

resistance phenotypes observed for the corresponding fusions (**Figure 4C**, **S3D**). In contrast,

WzzE^{F104Y}, WzzE^{F104H}, WzzE^{K162S}, WzzE^{K162T}, WzzE^{L259A}, and WzzE^{L259I} (lanes 5, 6, 8-11)

demonstrated wild-type chain length regulation in both the p*wzzE* and p*wzzE-cat-FLAG*

constructs. Together, these mutants provide a useful range of phenotypes for dissecting the

requirements for ECA$_{CYC}$ synthesis.

Periplasmic residues of WzzE are important for ECA$_{CYC}$ biogenesis. We next assessed the

impact of our periplasmic mutants on ECA$_{CYC}$ biosynthesis by quantifying ECA$_{CYC}$ levels using

LC-MS. Both the WzzE$^{\Delta F104}$ and WzzE^{K162A} mutants exhibited impaired linear chain-length

regulation and produced little to no ECA$_{CYC}$ (**Figure 5A**). This loss of ECA$_{CYC}$ production could

results from either (i) insufficient protein accumulation and/or stability or (ii) disruption of chain-

length-regulating activities in these mutants. In contrast, WzzE^{F104Y}, WzzE^{K162S}, WzzE^{K162T},

WzzE^{L259A}, and WzzE^{L259I} produced ECA$_{CYC}$ at levels comparable to wild type. WzzE^{F104H}

showed highly variable ECA$_{CYC}$ levels despite maintaining wild type chain-length regulation and

protein levels, suggesting that this mutation may have a specific ECA$_{CYC}$ synthesis effect. These

data highlight the importance of WzzE's periplasmic domain on ECA$_{CYC}$ synthesis, likely through

its structural role, its chain-length-regulating activity, and possibly a function specific to ECA$_{CYC}$

synthesis. Given the differing effects of the F104 mutations on ECA$_{CYC}$ synthesis and linear ECA

chain-length regulation, we examined their potential structural consequences using AlphaFold 3

(Abramson et al., 2024) and aligned wild type and WzzE$^{\Delta F104}$ complexes (**Figure 5B** & **S4**,

Table S1). These models predict that F104 forms contacts with a neighboring loop, and

alterations in these interactions may underlie the functional differences observed for

substitutions at this residue.

ECA$_{CYC}$ biogenesis is genetically separable from chain-length regulation. Our plasmid-

based expression analyses of *wzzE* identified several point mutations with distinct effects on

ECA$_{CYC}$ production. Notably, substitution of the TMH2 residues (WzzE$^{GG333/4LL}$) abolished chain-

length regulation and ECA$_{CYC}$ synthesis, while maintaining protein stability and near-wild-type

protein accumulation. Additionally, several mutations to residue F104 also produced variable

effects on WzzE accumulation, chain-length regulation, and ECA$_{CYC}$ production. However,

plasmid-based expression of wild-type *wzzE* yielded lower levels of linear ECA and reduced

modal chain-length compared expression of *wzzE* from its native locus (**Figure 1D**), a

difference likely contributing to some of the phenotypes observed in the plasmid system. We

also noted substantial variability in ECA$_{CYC}$ levels among periplasmic mutants, which we

attributed to non-native gene expression. To circumvent these limitations, we used CRISPR-

Cas9 to introduce select point mutations into the native *wzzE* locus and assayed their effect on

linear ECA and ECA$_{CYC}$.

LC-MS analysis of purified ECA$_{CYC}$ revealed that chromosomal WzzE$^{\Delta F104}$ and

WzzE$^{GG333/4LL}$ mutants produced little to no ECA$_{CYC}$, similar to what is observed with $\Delta wzzE$

(**Figure 6A**). In contrast, WzzE^{F104H} produced measurable levels of ECA$_{CYC}$ that were about 2-

fold lower than the wild type, whereas WzzE^{F104Y} showed no significant change in ECA$_{CYC}$ levels

relative to the wild type. We then assayed linear ECA chain length regulation by immunoblot

(**Figure 6B**). Consistent with the plasmid-based system, WzzE$^{\Delta F104}$ and WzzE$^{GG333/4LL}$ failed to

regulate linear ECA. The magnitude of chain-length dysregulation in these mutants exceeded

that of Δ*wzzE*, potentially suggesting an aberrant interaction of non-functional WzzE and ECA

synthesis that differs from the absence of WzzE. WzzE^{F104H} and WzzE^{F104Y} each displayed

intact and nearly wild-type chain-length regulation (**Figure 6B**), with a possible slight increase in

modal chain length that was not present in all replicates. However, the linear ECA regulation of

WzzE^{F104H} and WzzE^{F104Y} was consistently identical between these two mutants.

In our ECA immunoblot protocol, we take steps to minimize ECA degradation (see Experimental

Procedures), as ECA can be delipidated by mild acid hydrolysis (BASU et al., 1987). One of

these steps is to avoid long incubations, such as those used to proteinase K (ProK) treat LPS

samples (May and Silhavy, 2018). However, protein in our samples cause ECA to spread over a

larger area of the gel and potentially could mask small changes in chain length regulation. To

improve detection of small changes in chain length while minimizing ECA degradation, we

pretreated our immunoblot samples with various concentrations of ProK and incubation times at

56 °C found in the literature (Leo et al., 2020, Jiang et al., 2020, Maczuga et al., 2022a, Liu et

al., 2020, Peters et al., 1985), as well as treatments optimized based on our initial results. The

resulting blots and comparative signal quality are shown in (**Figure S5**). We selected a one-

hour treatment with a high concentration of ProK to move forward with as this treatment resulted

in loss of protein signal while minimizing ECA degradation. After treatment with ProK in these

optimized conditions, both WzzE^{F104H} and WzzE^{F104Y} showed ECA chain length that was

extremely similar to wild type (**Figure 6C**). Overall, the identical regulation of linear ECA chain-

length by WzzE^{F104H} and WzzE^{F104Y} but differing effects on ECA$_{CYC}$ synthesis demonstrate that

levels of ECA$_{CYC}$ production are genetically separable from linear chain-length regulation.

We were surprised that the dysregulation of ECA chain length, and even more the levels

of ECA, in the WzzE^{ΔF104} and WzzE$^{GG333/4LL}$ mutants was still greater than the Δ*wzzE* control,

and so we wondered whether, in a Δ*wzzE* strain, the remaining PCP1s could partially regulate

ECA chain length through interaction with WzyE. In this model, nonfunctional WzzE mutants

might sequester WzyE from the other PCP1s, exacerbating dysregulation. Therefore, we

assayed whether we could cause further disruption of ECA chain-length regulation in a Δ*wzzE*

strain by deleting *wzzB* and/or *fepE*. Deletion of Δ*wzzB*, Δ*fepE*, and Δ*wzzB* Δ*fepE*, or both had

no detectable effect on chain regulation in a Δ*wzzE* strain (**Figure 6D**), though the triple mutant

showed a modest increase in linear ECA levels. Taken together, these findings demonstrate that

mutations in *wzzE* that disrupt linear ECA chain-length regulation can also abolish ECA$_{CYC}$

synthesis, and that normal ECA$_{CYC}$ production is genetically separable from regulation of linear

ECA chain-length.

Discussion

In this study, we constructed and characterized a set of *wzzE* site-directed mutants to

define which functions of WzzE are required for ECA$_{CYC}$ biogenesis. Although previous

mutagenesis studies have identified regions of *Shigella flexneri* WzzB important for activity

(Nath and Morona, 2015, Ascari et al., 2022), no prior work has examined how WzzE

contributes to the production of cyclic ECA. Here, we show that substitutions at F104 within the

periplasmic domain and to the GXXXG motif in TMH2 can compromise both ECA chain-length

regulation and ECA$_{CYC}$ synthesis. Moreover, our data demonstrate that regulation of linear ECA

chain length and the level of ECA$_{CYC}$ production are genetically separable, as we have identified

two mutations at F104 with divergent effects on ECA$_{CYC}$ levels, but produce identical regulation

of linear ECA chain length. These data show regulation of chain length by WzzE is necessary

but not sufficient for production of normal levels of ECA$_{CYC}$.

Despite low sequence conservation among PCP1s, their overall structures are highly

similar, consisting of two transmembrane domains and an α-helix/β-sheet domain within the

periplasmic space that extends ~100 Å above the membrane (Tocilj et al., 2008, Kalynych et al.,

2012b, Collins et al., 2017, Kalynych et al., 2015). PCP1s also harbor a conserved GXXXG

motif that has been shown to contribute to chain-length regulation. In WzzB, a G305A/G311A

double mutant in *Shigella flexneri* WzzB that encompassed and extended outside of this motif

resulted in short-chain O-antigen (Oag) (Daniels and Morona, 1999, Papadopoulos et al., 2016),

whereas single, double, and triple glycine to alanine mutations restricted to the GXXXG motif

did not affect chain-length (Collins et al., 2017, Daniels and Morona, 1999). Motivated by these

findings, we focused on residues within and around the GXXXG motif, characterized their effect

on chain-length regulation and ECA_{CYC} biogenesis, and identified two adjacent, conserved

glycine residues in the GXXXG motif of WzzE—analogous to G305/G306 in WzzB—as critical

for ECA chain-length regulation and ECA_{CYC} biosynthesis. Interestingly Ala substitutions at

these positions had no detectable effect on chain-length regulation or ECA_{CYC} production,

whereas introduction of adjacent Leu residues led to pronounced dysregulation of ECA chain-

length and significantly reduced levels of ECA_{CYC}.

GXXXG motifs often mediate dimerization and multimerization of transmembrane

domains (Parrish et al., 2015, Faingold et al., 2012), suggesting that the motif may facilitate

positioning of TMH1 and TMH2 of WzzE. In support of this idea, our structural modeling

suggests increased distance between TMH1 and TMH2 in the $WzzE^{GG333/4LL}$ mutant (**Figure 3D**,

S6A). Alignments of WzzE monomers predicted to closely interact with WzyE, shows that WzyE

rotates in the plane perpendicular to the membrane in $WzyE-WzzE^{GG333/4LL}$ model compared to

the wild type (**Figure S6BC**). The alignment also shows the distance in the structure between

the TMH of opposite WzzE monomers was 2.0 to 2.7 Å smaller in the $WzzE^{GG333/4LL}$ model than

in the wild-type model. In compensation, WzyE takes on a more condensed structure (1.4 Å

narrower) in the $WzzE^{GG333/4LL}$ model. As this type of rotation and structure change could impair

WzyE function, this prediction provides a possible hypothesis that $WzzE^{GG3334/LL}$ has impaired

interaction with WzyE. Alternatively, disruption of TMH packing may interfere with an intrinsic

and lesser defined activity of WzzE.

In contrast to mutations in TMH2, many substitutions introduced into the periplasmic

domain rendered WzzE unstable, as evidenced by loss of chloramphenicol resistance conferred

by the WzzE-Cat fusion. This observation underscores the importance of the periplasmic

domain for WzzEs structural integrity and/or multimerization. Prior work on other PCP1s

suggests that multiple surface-exposed residues within the periplasmic domain affect chain-

length regulation, producing either shorter or longer Oag chains (Kalynych et al., 2012b,

Papadopoulos and Morona, 2010, Daniels and Morona, 1999, Hong and Payne, 1997, Tran and

Morona, 2013). More recently, increasing evidence suggests that the inner surface of the WzzE

octamer may bind to the elongating carbohydrate to facilitate growth of the chain (Wiseman et

al., 2023, Hong et al., 2023, Weckener et al., 2023).

WzzE's internal cavity possesses two highly concentrated, negatively charged bands,

separated by a flexible neutrally charged loop that may bind the growing carbohydrate chain

(Wiseman et al., 2023). A loop at the apex of the periplasmic domain has been suggested to

"ratchet" up and down to facilitate carbohydrate movement over this binding surface (Wiseman

et al., 2023). F104 is buried near the base of the periplasmic "bell" and may affect this binding

surface indirectly. In contrast, K162 is positioned adjacent to the lower negatively charged band,

and L259 residues within the proposed ratchet loop. Mutations to L259 did not affect chain-

length regulation; however, loss of charge at K162 did prevent chain-length regulation and

ECA_{CYC} synthesis, perhaps supporting this model of ECA chain-length elongation. As to how the

$wzzE_{K162A}$ and $wzzE_{\Delta F104}$ mutations lead to loss of ECA_{CYC} production, we speculate that the

chain-length regulating activity of WzzE—related to the carbohydrate binding and ratcheting

mechanism—is necessary for ECA_{CYC} synthesis.

In contrast, the $wzzE_{F104H}$ mutation reduces ECA_{CYC} levels while maintaining

comparatively wild-type chain-length regulation. Moreover, its chain-length regulation is

indistinguishable from $wzzE_{F104Y}$, which produces wild-type ECA_{CYC} levels. These results

demonstrate that chain-length regulation alone is insufficient for normal ECA_{CYC} synthesis and

suggest that WzzE plays an additional role specific to ECA_{CYC} synthesis beyond that of chain-

length regulation. One possible mechanism would be that WzzE bends ECA molecules of the

correct size into an orientation that can be spontaneously cyclized. Another possibility is that

WzzE allows the binding of another protein (i.e., a cyclase) that can cyclize ECA chains of the

appropriate length. Finally, it is possible that WzzE has specific features that promote the

formation of ECA chains with 4 RUs that can later be cyclized to form ECA_{CYC}. Further, genetic

and biochemical experiments will be necessary to distinguish between these possibilities and to

solidify an additional role for WzzE in ECA_{CYC} biogenesis.

Unexpectedly, we consistently observed that WzzE $^{\Delta F104}$ and WzzE$^{GG333/4LL}$ mutants had

greater linear ECA levels than what was observed in our *wzzE* deletion. We confirmed that this

phenotype does not arise from compensation by other PCP1s in *E.coli*, although we did notice a

modest increase in linear ECA in strains lacking all other PCP1 genes. Therefore, why WzzE

$^{\Delta F104}$ and WzzE$^{GG333/4LL}$ have elevated linear ECA levels remains unclear. One explanation is that

inactive WzzE octamers partially sequester growing ECA chains, increasing variability in chain

length. Regardless, we believe that the phenotypic distinction between Δ*wzzE* strain and the

mutants strengthens the conclusion that these mutants produce inactive proteins which

accumulate but are incapable of regulating chain-length or supporting ECA_{CYC} biosynthesis.

Overall, our analysis interrogates only a small fraction of WzzE residues, and additional

mutations that selectively impair ECA_{CYC} biosynthesis likely exist. Identification of such mutants

will clarify the molecular mechanism of ECA_{CYC} biosynthesis. Nonetheless, our findings reveal

fundamental requirements for ECA_{CYC} and chain-length regulation. We propose coordinated

activity between WzyE and WzzE, together with periplasmic charge elements within WzzE that

facilitate chain elongation, is required for ECA_{CYC} biogenesis. Importantly, we demonstrate that

chain-length regulation can be genetically uncoupled from ECA_{CYC} synthesis, indicating that

WzzE possesses a second function beyond chain-length regulation that is important for ECA_{CYC}

biosynthesis. These insights help to define key features of WzzE that enable synthesis of a

cyclic carbohydrate.

Experimental Procedures

Strains and growth conditions. The strains used in this study are listed in **Table S2**. Cultures

were inoculated and grown at 37 °C in LB Lennox unless otherwise noted, and, when

necessary, were supplemented with 20 mg/L chloramphenicol, 25 mg L^{-1} kanamycin, 10 mg L^{-1}

tetracycline, or a combination thereof. Insertion-deletion alleles containing kanamycin resistance

cassettes from the Keio collection (Baba et al., 2006) were moved to isogenic backgrounds

using P1vir transduction (Silhavy et al., 1984). Clean deletions were constructed using the Flp

recombinase-FRT system as described (Datsenko and Wanner, 2000). Chromosomal

mutagenesis in *wzzE* was performed using a CRISPR-Cas9 system harboring a

pCRISPR::protospacer plasmid in W3110 containing a heat shock-inducible λcI857 lysogen, as

previously described (Pyne et al., 2015, Jiang et al., 2013). The pCas9 (Addgene #42876) and

pCRISPR (Addgene #42975) plasmids were a kind gift from Professor Luciano Marraffini

(Rockefeller University). Oligonucleotides used to generate the protospacer and repair

templates used for mutagenesis are listed in **Table S3**. Prior to chromosomal mutagenesis,

metE::Tn*10* was linked to *wzzE* (Nichols et al., 1998), allowing the mutations to be transferred

into MG1655 via P1vir transduction. Strains carrying *metE*::Tn*10* were supplemented with 1µm

cobalamin (vitamin B$_{12}$) when grown in minimal media (Banerjee et al., 1989).

Plasmids, cloning, and site directed mutagenesis. All primers used in this study are listed in

Table S4. To subclone the p*wzzE* expression plasmid, chromosomal *wzzE* with an ochre stop

codon was PCR-amplified using primers pZS21-wzzE_fwd and wzzE-pZS21_rev. The low copy

number plasmid pZS21 (Lutz and Bujard, 1997) was amplified using primers pZS21_fwd and

pZS21_rev. The plasmid was assembled using HiFi Assembly Master Mix (New England

Biolabs) according to the manufacturer's instructions. To construct p*wzzE-cat*, *wzzE* lacking a

stop codon was amplified using primers pZS21-wzzE_fwd and wzzE-cat_rev. The

chloramphenicol acetyltransferase gene (*cat*) was amplified from pBAD33 (Guzman et al., 1995)

using primers wzzE-cat_fwd and cat-pZS21_rev. These fragments were assembled into pZS21,

amplified as described above, using HiFi Assembly. To construct p*wzzE*-FLAG, p*wzzE* was

amplified using primers p*wzzE*_x3F_Fwd and p*wzzE*_x3F_Rvr, which contained sequence

homology to a glycine-serine-linked FLAG tag that incorporating a stop codon. The glycine-

serine-linked FLAG tag was supplied as a gene-block fragment (**Table S3**) and assembled into

the p*wzzE* plasmid via HiFi Assembly. To generate the p*wzzE-cat*-FLAG plasmid, *cat* was

amplified using primers WzzE:Cmr Fp and WzzE:Cmr Rp and the p*wzzE*-FLAG plasmid

backbone was amplified using Cmr: GS-x3F_FP and wzzE-cat_rev. The fragments were

assembled using HiFi Assembly.

Site-direct mutations were generated using Q5 PCR mutagenesis (New England

Biolabs). Briefly, non-overlapping primers containing the mutation of interest were used to

amplify the indicated pZS21-sanA plasmids (pZS21-*wzzE*, pZS21-*wzzE-cat*, and pZS21-*wzzE*-

FLAG). PCR products were digested with DpnI (New England Biolabs), phosphorylated using

T4 polynucleotide kinase (New England Biolabs) and self-ligated with T4 DNA ligase (New

England Biolabs). Ligations were transformed into Turbo cloning cells (New England Biolabs),

and the mutations were confirmed by sequencing (Plasmidsaurs). Primers used all site-directed

mutants are listed in **Table S4**. Modeling of the predicted mutant structures was performed

using AlphaFold 3 (Abramson et al., 2024) with eight monomers of WzzE and one of WzyE.

Relevant confidence scores are given in **Table S1** and **Figure S2**.

Antibiotic Sensitivity. Efficiency of plating (EOP) assays were performed by plating 10-fold serial

dilutions of overnight cultures onto LB plates containing the indicated antibiotics using a 48-pin

tool. Plates were incubated overnight at 30 °C before imaging.

Immunoblot Analysis. Immunoblots for WzzE-FLAG and ECA were conducted as previously

described with minor modification (Mitchell et al., 2018, Morris and Mitchell, 2023). For analysis

of WzzE-FLAG protein levels, samples were normalized to equal OD_{600} from overnight cultures,

lysed by boiling for 5 minutes in sample loading buffer, and loaded on 12% SDS-PAGE gels.

Levels of WzzE-FLAG were quantitated by densitometry using ImageJ as has been previously

described (Mitchell et al., 2018) on immunoblots with the lowest exposure at which WzzE bands

could be detected. Samples for ECA analysis were prepared and immunoblotted as previously

described with minor modification (Mitchell et al., 2018, Morris and Mitchell, 2023). To avoid

ECA degradation, ultrapure water, commercial lysis reagents, commercial running buffer, and

precast gels were used for ECA immunoblots. In addition, all western equipment was washed

before and after ECA immunoblots with Extran 300 Detergent (Millipore Sigma) and rinsed 10

times with ultrapure water. To determine whether proteins affect the apparent chain length of

ECA, we pre-treated samples with ProK. Samples were subjected to different pre-treatment

conditions, varying both in the concentration of ProK and time of treatment (**Figure S5A**), based

on protocols used in previous (Peters et al., 1985, Liu et al., 2020, Leo et al., 2020, Maczuga et

al., 2022a) papers. All treatments were performed at 56 °C. Following ProK treatment, samples

were boiled again for 10min, and loaded on 15% SDS-PAGE gels. The outcomes of each

treatment condition and their respective effects on the resolution and clarity of the immunoblot, as

well as effects on ECA levels are shown in (**Figure S5B**). Antibodies used for blotting were as

follows: M2 αFLAG (Millipore Sigma, 1:50,000 dilution), αGroEL (Millipore Sigma, 1: 60,000

dilution), αECA (a gift from Professor Renato Morona, University of Adelaide, 1:30,000), rabbit

α-mouse (Prometheus, 1:100,000 dilution), goat α-rabbit (Prometheus, 1:100,000 dilution), or

αBamA (a gift from Professor Thomas Silhavy, Princeton University, 1:50,000 dilution).

Quantification of Cyclic ECA. Quantification of ECA_{CYC} was performed as previously described,

with minor modifications (Mitchell et al., 2018). Briefly, cells were grown in M63 glucose

minimum media lacking nitrogen and supplemented with either 0.2% $(^{14}NH_4)_2SO_4$ or

$(^{15}NH_4)_2SO_4$. ECA_{CYC} was purified as previously described. Sample analysis was performed

using a Thermo Scientific Q Exactive Focus and liquid chromatography unit (UltiMate 3000 RS).

Purified samples were dissolved in 0.1% formic acid and separated by injecting 20 μL of sample

into the Synergi Hydro-RP (4.6 x 250 mm; 4.0 μm) column (Phenomenax). The mobile phase

consisted of 0.1% Formic acid (eluent A) and methanol (eluent B). The flow rate was set at

800 μL/min with the following gradient: 0–5.0 min 5% B, 5.0-12.0 min 5-60% B, 12-14 min 60-

90% B, 14-17 min 90% B, and returned to 5% B in 0.1 min and held for 5 min at 5% B. The Q

Exactive Focus ESI source was operated in full MS (100-1000 m/z). The mass resolution was

tuned to 70000 FWHM at m/z 200. The spray voltage was set to 3.3 kV in negative mode, and

the sheath gas and auxiliary gas flow rates were set to 60 and 20 arbitrary units, respectively.

The transfer capillary and auxiliary gas heater temperatures were held at 380 and 300 °C,

respectively. The S-Lens RF level was set at 70 V. Exactive Series 2.11 /Xcalibur 4.2.47

software was used for data acquisition and processing. For relative quantification, the ratio of

the areas of the ^{14}N to ^{15}N ECA_{CYC} peaks was calculated.

Acknowledgements

Thank you to the members of the Mitchell Lab for helpful and productive discussions and for

critical reading of our manuscript. We also thank Drs. Jolene Ramsey, Deborah Siegele,

Jennifer Herman, and Ryland Young (Texas A&M University), as well as members of the

Ramsey lab, for their insightful comments. The ECA antibody was a kind gift from Professor

Renato Morana (University of Adelaide). The pCas9 and pCRISPR plasmids were a kind gift

from Professor Luciano Marraffini (Rockefeller University) through the Addgene Repository. This

work was supported by the National Institute of Allergy and Infectious Disease under award

number R01-AI155915 (to A.M.M.).

References

ABRAMSON, J., ADLER, J., DUNGER, J., EVANS, R., GREEN, T., PRITZEL, A.,

RONNEBERGER, O., WILLMORE, L., BALLARD, A. J., BAMBRICK, J., BODENSTEIN,

S. W., EVANS, D. A., HUNG, C.-C., O'NEILL, M., REIMAN, D., TUNYASUVUNAKOOL,

K., WU, Z., ŽEMGULYTė, A., ARVANITI, E., ... & JUMPER, J. M. 2024. Accurate

structure prediction of biomolecular interactions with AlphaFold 3. *Nature,* 630**,** 493-500.

ADEOLU, M., ALNAJAR, S., NAUSHAD, S. & S. GUPTA, R. 2016. Genome-based phylogeny

and taxonomy of the '*Enterobacteriales*': proposal for *Enterobacterales* ord. nov. divided

into the families *Enterobacteriaceae*, *Erwiniaceae* fam. nov., *Pectobacteriaceae* fam.

nov., *Yersiniaceae* fam. nov., *Hafniaceae* fam. nov., *Morganellaceae* fam. nov., and

Budviciaceae fam. nov. *International Journal of Systematic and Evolutionary*

Microbiology, 66**,** 5575-5599.

AGUINIGA, L. M., YAGGIE, R. E., SCHAEFFER, A. J. & KLUMPP, D. J. 2016.

Lipopolysaccharide Domains Modulate Urovirulence. *Infection and Immunity,* 84**,** 3131-

3140.

AMINOV, R. I. 2010. A brief history of the antibiotic era: lessons learned and challenges for the

future. *Front Microbiol,* 1**,** 134.

ANDREEVA, A. E., BORISSOVA, B. E., MIRONOVA, R., GLYKOS, N. M., KOTSIFAKI, D.,

IVANOV, I., KRYSTEVA, M. & KOKKINIDIS, M. 2000. Crystallization of type I

chloramphenicol acetyltransferase: an approach based on the concept of ionic strength

reducers. *Acta Crystallogr D Biol Crystallogr,* 56**,** 101-3.

ARMBRUSTER, C. E., FORSYTH-DEORNELLAS, V., JOHNSON, A. O., SMITH, S. N., ZHAO,

L., WU, W. & MOBLEY, H. L. T. 2017. Genome-wide transposon mutagenesis of *Proteus*

mirabilis: Essential genes, fitness factors for catheter-associated urinary tract infection,

and the impact of polymicrobial infection on fitness requirements. *PLOS Pathogens,* 13**,**

e1006434.

ASCARI, A., TRAN, E. N. H., EIJKELKAMP, B. A. & MORONA, R. 2022. Identification of the

Shigella flexneri Wzy Domain Modulating Wzz(pHS-2) Interaction and Detection of the

Wzy/Wzz/Oag Complex. *J Bacteriol,* 204**,** e00224-22.

584 BABA, T., ARA, T., HASEGAWA, M., TAKAI, Y., OKUMURA, Y., BABA, M., DATSENKO, K. A.,

585   TOMITA, M., WANNER, B. L. & MORI, H. 2006. Construction of *Escherichia coli* K-12 in-

586   frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol,* 2**,** 2006.0008.

587 BANERJEE, R. V., JOHNSTON, N. L., SOBESKI, J. K., DATTA, P. & MATTHEWS, R. G. 1989.

588   Cloning and sequence analysis of the *Escherichia coli* metH gene encoding cobalamin-

589   dependent methionine synthase and isolation of a tryptic fragment containing the

590   cobalamin-binding domain. *J Biol Chem,* 264**,** 13888-95.

591 BARR, K., KLENA, J. & RICK, P. 1999. The Modality of Enterobacterial Common Antigen

592   Polysaccharide Chain Lengths Is Regulated by o349 of the wec Gene Cluster of

593   *Escherichia coli K-12. Journal of bacteriology,* 181**,** 6564-8.

594 BARR, K., NUNES-EDWARDS, P. & RICK, P. D. 1989. In vitro synthesis of a lipid-linked

595   trisaccharide involved in synthesis of enterobacterial common antigen. *J Bacteriol,* 171**,**

596   1326-32.

597 BARR, K. & RICK, P. D. 1987. Biosynthesis of enterobacterial common antigen in *Escherichia*

598   *coli.* In vitro synthesis of lipid-linked intermediates. *J Biol Chem,* 262**,** 7142-50.

599 BASU, S., KUHN, H.-M., NESZMELYI, A., HIMMELSPACH, K. & MAYER, H. 1987. Chemical

600   characterization of enterobacterial common antigen isolated from *Plesiomonas*

601   *shigelloides* ATCC 14029. *European Journal of Biochemistry,* 162**,** 75-81.

602 BENNETT, H. C. & MITCHELL, A. M. 2025. Recent advances in understanding of

603   enterobacterial common antigen synthesis and regulation. *Open Biology,* 15**,** 250055.

604 BERTANI, B. & RUIZ, N. 2018. Function and Biogenesis of Lipopolysaccharides. *EcoSal Plus,*

605   8**,** ESP-0001-2018.

606 COLLINS, R. F., KARGAS, V., CLARKE, B. R., SIEBERT, C. A., CLARE, D. K., BOND, P. J.,

607   WHITFIELD, C. & FORD, R. C. 2017. Full-length, Oligomeric Structure of Wzz

608   Determined by Cryoelectron Microscopy Reveals Insights into Membrane-Bound States.

609   *Structure,* 25**,** 806-815.e3.

DANIELS, C. & MORONA, R. 1999. Analysis of *Shigella flexneri* wzz (Rol) function by mutagenesis and cross-linking: wzz is able to oligomerize. *Mol Microbiol,* 34**,** 181-94.

DATSENKO, K. A. & WANNER, B. L. 2000. One-step inactivation of chromosomal genes in *Escherichia coli K-12* using PCR products. *Proceedings of the National Academy of Sciences,* 97**,** 6640-6645.

DOERN, C. D. 2024. Chapter 2 - Classification of medically important bacteria. *In:* TANG, Y.-W., HINDIYEH, M. Y., LIU, D., SAILS, A., SPEARMAN, P. & ZHANG, J.-R. (eds.) *Molecular Medical Microbiology (Third Edition).* Academic Press.

EADE, C. R., WALLEN, T. W., GATES, C. E., OLIVERIO, C. L., SCARBROUGH, B. A., REID, A. J., JORGENSON, M. A., YOUNG, K. D. & TROUTMAN, J. M. 2021. Making the enterobacterial common antigen glycan and measuring its substrate sequestration. *ACS Chem Biol,* 16**,** 691-700.

ERBEL, P. J., BARR, K., GAO, N., GERWIG, G. J., RICK, P. D. & GARDNER, K. H. 2003. Identification and biosynthesis of cyclic enterobacterial common antigen in *Escherichia coli*. *J Bacteriol,* 185**,** 1995-2004.

ERBEL, P. J., SEIDEL, R., MACINTOSH, S. E., GENTILE, L. N., AMOR, J. C., KAHN, R. A., PRESTEGARD, J. H., MCINTOSH, L. P. & GARDNER, K. H. 2004. Cyclic enterobacterial common antigen: potential contaminant of bacterially expressed protein preparations. *J Biomol NMR,* 29**,** 199-204.

FAINGOLD, O., COHEN, T. & SHAI, Y. 2012. A GxxxG-like Motif within HIV-1 Fusion Peptide Is Critical to Its Immunosuppressant Activity, Structure, and Interaction with the Transmembrane Domain of the T-cell Receptor*. *Journal of Biological Chemistry,* 287**,** 33503-33511.

FARFOUR, E., DORTET, L., GUILLARD, T., CHATELAIN, N., POISSON, A., MIZRAHI, A., FOURNIER, D., BONNIN, R. A., DEGAND, N., MORAND, P., JANVIER, F., FIHMAN, V., CORVEC, S., BROUTIN, L., LE BRUN, C., YIN, N., HéRY-ARNAUD, G., GRILLON, A.,

BILLE, E., ... & ON BEHALF OF THE GMC STUDY, G. 2022. Antimicrobial Resistance in

Enterobacterales Recovered from Urinary Tract Infections in France. *Pathogens,* 11**,**

356.

FäRNBäCK, M., ERIKSSON, L., SENCHENKOVA, S. Y., ZYCH, K., KNIREL, Y. A.,

SIDORCZYK, Z. & WIDMALM, G. 2003. Crystal Structure of a Cyclic Enterobacterial

Common Antigen. *Angewandte Chemie International Edition,* 42**,** 2543-2546.

GILBREATH, J. J., COLVOCORESSES DODDS, J., RICK, P. D., SOLOSKI, M. J., MERRELL,

D. S. & METCALF, E. S. 2012. Enterobacterial common antigen mutants of *Salmonella*

enterica serovar Typhimurium establish a persistent infection and provide protection

against subsequent lethal challenge. *Infect Immun,* 80**,** 441-50.

GOźDZIEWICZ, T. K., ŁUKASIEWICZ, J. & ŁUGOWSKI, C. 2015. [The structure and

significance of enterobacterial common antigen (ECA)]. *Postepy Hig Med Dosw (Online),*

69**,** 1003-12.

GUéRIN, F., LALLEMENT, C., GOUDERGUES, B., ISNARD, C., SANGUINETTI, M., CACACI,

M., TORELLI, R., CATTOIR, V. & GIARD, J.-C. 2020. Landscape of in vivo Fitness-

Associated Genes of *Enterobacter cloacae* Complex. *Frontiers in Microbiology,* 11**,** 1609.

GUZMAN, L. M., BELIN, D., CARSON, M. J. & BECKWITH, J. 1995. Tight regulation,

modulation, and high-level expression by vectors containing the arabinose P_{BAD}

promoter. *J Bacteriol,* 177**,** 4121-30.

HOLMES, C. L., WILCOX, A. E., FORSYTH, V., SMITH, S. N., MORICZ, B. S.,

UNVERDORBEN, L. V., MASON, S., WU, W., ZHAO, L., MOBLEY, H. L. T. &

BACHMAN, M. A. 2023. *Klebsiella pneumoniae* causes bacteremia using factors that

mediate tissue-specific fitness and resistance to oxidative stress. *PLOS Pathogens,* 19**,**

e1011233.

HONG, M. & PAYNE, S. M. 1997. Effect of mutations in *Shigella flexneri* chromosomal and

plasmid-encoded lipopolysaccharide genes on invasion and serum resistance. *Mol*

Microbiol, 24**,** 779-91.

HONG, Y., HU, D., VERDEROSA ANTHONY, D., QIN, J., TOTSIKA, M. & REEVES PETER, R.

2023. Repeat-Unit Elongations To Produce Bacterial Complex Long Polysaccharide

Chains, an O-Antigen Perspective. *EcoSal Plus,* 11**,** eesp-0020-2022.

HUSZCZYNSKI, S. M., COUMOUNDOUROS, C., PHAM, P., LAM, J. S. & KHURSIGARA, C. M.

2019. Unique regions of the polysaccharide copolymerase Wzz_2 from *Pseudomonas*

aeruginosa are essential for O-specific antigen chain length control. *Journal of*

Bacteriology, 201**,** e00165-19.

ISLAM, S. T. & LAM, J. S. 2014. Synthesis of bacterial polysaccharides via the Wzx/Wzy-

dependent pathway. *Can J Microbiol,* 60**,** 697-716.

JENKINS, C., RENTENAAR, R. J., LANDRAUD, L. & BRISSE, S. 2017. 180 -

Enterobacteriaceae. In: COHEN, J., POWDERLY, W. G. & OPAL, S. M. (eds.) *Infectious*

Diseases (Fourth Edition). Elsevier.

JIANG, W., BIKARD, D., COX, D., ZHANG, F. & MARRAFFINI, L. A. 2013. RNA-guided editing

of bacterial genomes using CRISPR-Cas systems. *Nature biotechnology,* 31**,** 233-239.

JIANG, X. E., TAN, W. B., SHRIVASTAVA, R., SEOW, D. C. S., CHEN, S. L., GUAN, X. L. &

CHNG, S.-S. 2020. Mutations in enterobacterial common antigen biosynthesis restore

outer membrane barrier function in Escherichia coli tol-pal mutants. *Molecular*

Microbiology, 114**,** 991-1005.

KAJIMURA, J., RAHMAN, A., HSU, J., EVANS, M. R., GARDNER, K. H. & RICK, P. D. 2006. O

acetylation of the enterobacterial common antigen polysaccharide is catalyzed by the

product of the *yiaH* Gene of *Escherichia coli* K-12. *Journal of Bacteriology,* 188**,** 7542-

7550.

KAJIMURA, J., RAHMAN, A. & RICK, P. D. 2005. Assembly of cyclic enterobacterial common

antigen in *Escherichia coli* K-12. *J Bacteriol,* 187**,** 6917-6927.

KALYNYCH, S., CHERNEY, M., BOSTINA, M., ROUILLER, I. & CYGLER, M. 2015. Quaternary

structure of WzzB and WzzE polysaccharide copolymerases. *Protein Science,* 24**,** 58-

69.

KALYNYCH, S., MORONA, R. & CYGLER, M. 2014. Progress in understanding the assembly

process of bacterial O-antigen. *FEMS Microbiol Rev,* 38**,** 1048-65.

KALYNYCH, S., RUAN, X., VALVANO, M. A. & CYGLER, M. 2011. Structure-guided

investigation of lipopolysaccharide O-antigen chain length regulators reveals regions

critical for modal length control. *Journal of Bacteriology,* 193**,** 3710-3721.

KALYNYCH, S., VALVANO, M. A. & CYGLER, M. 2012a. Polysaccharide co-polymerases: the

enigmatic conductors of the O-antigen assembly orchestra. *Protein Eng Des Sel,* 25**,**

797-802.

KALYNYCH, S., YAO, D., MAGEE, J. & CYGLER, M. 2012b. Structural characterization of

closely related O-antigen lipopolysaccharide (LPS) chain length regulators. *J Biol Chem,*

287**,** 15696-705.

KINTZ, E. N. & GOLDBERG, J. B. 2011. Site-directed mutagenesis reveals key residue for O

antigen chain length regulation and protein stability in *Pseudomonas aeruginosa* Wzz2.

J Biol Chem, 286**,** 44277-44284.

KUNIN, C. M. 1963. Separtion, chacterization, and Biological Significance of a Common Antigen

in *Enterobacteriaceae*. *J Exp Med,* 118**,** 565-86.

KUNIN, C. M., BEARD, M. V. & HALMAGYI, N. E. 1962. Evidence for a Common Hapten

Associated with Endotoxin Fractions of *E. coli* and other *Enterobacteriaceae*. *Proc Soc*

Exp Biol Med, 111**,** 160-166.

LEO, V., TEH, M. Y., TRAN, E. N. H. & MORONA, R. 2021a. Identification of a Region in

Shigella flexneri WzyB Disrupting the Interaction with Wzz(pHS2). *J Bacteriol,* 203**,**

e0041321.

LEO, V., TRAN, E. & MORONA, R. 2020. Polysaccharide co-polymerase WzzB/WzzE chimeras

reveal transmembrane 2 region of WzzB is important for interaction with WzyB. *J*

Bacteriol, 203.

LEO, V., TRAN, E. & MORONA, R. 2021b. Polysaccharide co-polymerase WzzB/WzzE

chimeras reveal transmembrane 2 region of WzzB is important for interaction with WzyB.

J Bacteriol, 203**,** 00598-20.

LIU, Q., SHEN, X., BIAN, X. & KONG, Q. 2020. Effect of deletion of gene cluster involved in

synthesis of Enterobacterial common antigen on virulence and immunogenicity of live

attenuated Salmonella vaccine when delivering heterologous *Streptococcus*

pneumoniae antigen PspA. *BMC Microbiology,* 20**,** 150.

LUTZ, R. & BUJARD, H. 1997. Independent and tight regulation of transcriptional units in

Escherichia coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic*

Acids Res, 25**,** 1203-10.

MACZUGA, N., TRAN, E. N. H. & MORONA, R. 2022a. Subcellular localization of the

enterobacterial common antigen GT-E-like glycosyltransferase, WecG. *Mol Microbiol,*

118**,** 403-416.

MACZUGA, N., TRAN, E. N. H., QIN, J. & MORONA, R. 2022b. Interdependence of *Shigella*

flexneri O Antigen and Enterobacterial Common Antigen Biosynthetic Pathways. *J*

Bacteriol, 204**,** e0054621.

MAHER, C. & HASSAN, K. A. 2023. The Gram-negative permeability barrier: tipping the

balance of the in and the out. *mBio,* 14**,** e01205-23.

MANNEL, D. & MAYER, H. 1978. Isolation and Chemical Characterization of the Enterobacterial

Common Antigen. *European Journal of Biochemistry,* 86**,** 361-370.

MAY, K. L. & SILHAVY, T. J. 2018. The *Escherichia coli* phospholipase PldA regulates outer

membrane homeostasis via lipid signaling. *mBio,* 9**,** e00379-18.

MENDEZ-PEREZ, D., GUNASEKARAN, S., ORLER, V. J. & PFLEGER, B. F. 2012. A

translation-coupling DNA cassette for monitoring protein translation in *Escherichia coli*.

Metabolic Engineering, 14**,** 298-305.

MITCHELL, A. M., SRIKUMAR, T. & SILHAVY, T. J. 2018. Cyclic Enterobacterial Common

Antigen Maintains the Outer Membrane Permeability Barrier of *Escherichia coli* in a

Manner Controlled by YhdP. *mBio,* 9**,** 01321-18.

MORONA, R., PURINS, L., TOCILJ, A., MATTE, A. & CYGLER, M. 2009. Sequence–structure

relationships in polysaccharide co-polymerase (PCP) proteins. *Trends in biochemical*

sciences, 34**,** 78-84.

MORONA, R., VAN DEN BOSCH, L. & MANNING, P. A. 1995. Molecular, genetic, and

topological characterization of O-antigen chain length regulation in *Shigella flexneri*. *J*

Bacteriol, 177**,** 1059-68.

MORRIS, K. N. & MITCHELL, A. M. 2023. Phosphatidylglycerol Is the Lipid Donor for Synthesis

of Phospholipid-Linked Enterobacterial Common Antigen. *Journal of Bacteriology,* 205**,**

e00403-22.

MUELLER, B. K., SUBRAMANIAM, S. & SENES, A. 2014. A frequent, GxxxG-mediated,

transmembrane association motif is optimized for the formation of interhelical Cα–H

hydrogen bonds. *Proceedings of the National Academy of Sciences,* 111**,** E888-E895.

NATH, P. & MORONA, R. 2015. Detection of Wzy/Wzz interaction in *Shigella flexneri*.

Microbiology (Reading), 161**,** 1797-1805.

NICHOLS, B. P., SHAFIQ, O. & MEINERS, V. 1998. Sequence analysis of Tn10 insertion sites

in a collection of Escherichia coli strains used for genetic mapping and strain

construction. *J Bacteriol,* 180**,** 6408-11.

PAPADOPOULOS, M. & MORONA, R. 2010. Mutagenesis and chemical cross-linking suggest
that Wzz dimer stability and oligomerization affect lipopolysaccharide O-antigen modal
chain length control. *J Bacteriol,* 192**,** 3385-93.

PAPADOPOULOS, M., TRAN, E. N. H., MURRAY, G. L. & MORONA, R. 2016. Conserved
transmembrane glycine residues in the *Shigella flexneri* polysaccharide co-polymerase
protein WzzB influence protein-protein interactions. *Microbiology (Reading),* 162**,** 921-
929.

PARRISH, H. L., GLASSMAN, C. R., KEENEN, M. M., DESHPANDE, N. R., BRONNIMANN, M.
P. & KUHNS, M. S. 2015. A Transmembrane Domain GGxxG Motif in CD4 Contributes to
Its Lck-Independent Function but Does Not Mediate CD4 Dimerization. *PLOS ONE,* 10**,**
e0132333.

PETERS, H., JüRS, M., JANN, B., JANN, K., TIMMIS, K. N. & BITTER-SUERMANN, D. 1985.
Monoclonal antibodies to enterobacterial common antigen and to Escherichia coli
lipopolysaccharide outer core: demonstration of an antigenic determinant shared by
enterobacterial common antigen and E. coli K5 capsular polysaccharide. *Infect Immun,*
50**,** 459-66.

PRAKASH, A., JANOSI, L. & DOXASTAKIS, M. 2011. GxxxG motifs, phenylalanine, and
cholesterol guide the self-association of transmembrane domains of ErbB2 receptors.
Biophys J, 101**,** 1949-58.

PYNE, M. E., MOO-YOUNG, M., CHUNG, D. A. & CHOU, C. P. 2015. Coupling the
CRISPR/Cas9 System with Lambda Red Recombineering Enables Simplified
Chromosomal Gene Replacement in *Escherichia coli*. *Appl Environ Microbiol,* 81**,** 5103-
14.

RAI, A. K., CARR, J. F., BAUTISTA, D. E., WANG, W. & MITCHELL, A. M. 2021. ElyC and cyclic
enterobacterial common antigen regulate synthesis of phosphoglyceride-linked
enterobacterial common antigen. *mBio,* 12**,** e02846-21.

RAI, A. K. & MITCHELL, A. M. 2020. Enterobacterial Common Antigen: Synthesis and Function

of an Enigmatic Molecule. *mBio,* 11**,** mBio.01914-20.

RAMIREZ, D. & GIRON, M. 2024. Enterobacter Infections. *StatPearls.* Treasure Island (FL):

StatPearls Publishing Copyright © 2024, StatPearls Publishing LLC.

RICK, P. D., MAYER, H., NEUMEYER, B. A., WOLSKI, S. & BITTER-SUERMANN, D. 1985.

Biosynthesis of enterobacterial common antigen. *J Bacteriol,* 162**,** 494-503.

SIEBER, V. 2003. Selection for Soluble Proteins via Fusion with Chloramphenicol

Acetyltransferase. *In:* ARNOLD, F. H. & GEORGIOU, G. (eds.) *Directed Enzyme*

Evolution: Screening and Selection Methods. Totowa, NJ: Humana Press.

SILHAVY, T. J., BERMAN, M. L. & ENQUIST, L. W. 1984. Experiments with gene fusions. *(No*

Title).

SILHAVY, T. J., KAHNE, D. & WALKER, S. 2010. The bacterial cell envelope. *Cold Spring Harb*

Perspect Biol, 2**,** a000414.

TEESE, M. G. & LANGOSCH, D. 2015. Role of GxxxG Motifs in Transmembrane Domain

Interactions. *Biochemistry,* 54**,** 5125-35.

TOCILJ, A., MUNGER, C., PROTEAU, A., MORONA, R., PURINS, L., AJAMIAN, E., WAGNER,

J., PAPADOPOULOS, M., VAN DEN BOSCH, L., RUBINSTEIN, J. L., FéTHIèRE, J.,

MATTE, A. & CYGLER, M. 2008. Bacterial polysaccharide co-polymerases share a

common framework for control of polymer length. *Nat Struct Mol Biol,* 15**,** 130-8.

TRAN, E. N. H. & MORONA, R. 2013. Residues located inside the *Escherichia coli* FepE

protein oligomer are essential for lipopolysaccharide O-antigen modal chain length

regulation. *Microbiology (Reading),* 159**,** 701-714.

WECKENER, M., WOODWARD, L. S., CLARKE, B. R., LIU, H., WARD, P. N., LE BAS, A.,

BHELLA, D., WHITFIELD, C. & NAISMITH, J. H. 2023. The lipid linked oligosaccharide

polymerase Wzy and its regulating co-polymerase, Wzz, from enterobacterial common

antigen biosynthesis form a complex. *Open Biology,* 13**,** 220373.

WHITFIELD, C., WILLIAMS, D. M. & KELLY, S. D. 2020. Lipopolysaccharide O-antigens-

bacterial glycans made to measure. *J Biol Chem,* 295**,** 10593-10609.

WISEMAN, B., NITHARWAL, R. G., WIDMALM, G. & HöGBOM, M. 2021. Structure of a full-

length bacterial polysaccharide co-polymerase. *Nat Commun,* 12**,** 369.

WISEMAN, B., WIDMALM, G. & HöGBOM, M. 2023. Alternating L4 loop architecture of the

bacterial polysaccharide co-polymerase WzzE. *Commun Biol,* 6**,** 802.

 Figures and Legends

Figure 1: Plasmid-based expression of *wzzE* partially complements a chromosomal *wzzE*

deletion. (A) Diagrams of constructs for expression of wild-type WzzE, WzzE with a C-terminal

glycine-serine linker and triple FLAG tag, WzzE translationally fused with chloramphenicol

acetyltransferase (Cat), and WzzE fused with Cat and triple FLAG are shown. The indicated

constructs were expressed from the low copy number pZS21 plasmid (Lutz and Bujard, 1997).

(B) The vancomycin sensitivity of a Δ*yhdP* strain can be suppressed by loss of WzzE (Mitchell

et al., 2018). When the Δ*wzzE* Δ*yhdP* strain carried the pZS21 empty plasmid, it remained

resistance to vancomycin as assayed by EOP but became sensitive when *wzzE, wzzE*-FLAG,

or *wzzE-cat* was expressed, indicating all three constructs complemented Δ*wzzE*. In addition,

translational fusion of *cat* and *wzzE* conferred chloramphenicol resistance, suggesting this

construct could readout WzzE stability and multimerization based on chloramphenicol

resistance. **(C)** WzzE-FLAG was detectable by immunoblotting. GroEL serves as a loading

control. **(D)** ECA chain-length regulation was assayed through immunoblotting using an ECA

antibody. Plasmid-base expression of WzzE restored chain-length regulation of linear ECA,

although lower levels of ECA and shorter chain-length were observed than in the wild-type strain

for the p*wzzE* and p*wzzE* constructs, while the p*wzzE*-FLAG and p*wzzE-cat*-FLAG constructs

complemented better. The Δ*wecA* strain produces no ECA and serves as negative control. X:

nonspecific band. **(E-F)** The strain carrying a plasmid with wild-type *wzzE* was grown with ^{15}N to

label ECA and then combined with an unlabeled strain carrying either wild-type *wzzE* (E) or the

empty vector (F). ECA$_{CYC}$ was purified and then detected by LC-MS. MS traces are shown for

the indicated samples. The mass, charge, and identity of the ECA$_{CYC}$-associated peaks are

shown. Expression of *wzzE* from p*wzzE* restored ECA$_{CYC}$ synthesis in the Δ*wzzE* strain. The

strains contained Δ*wecH* to prevent non-stochiometric acetylation of ECA. All images are

representative of at least three independent experiments.

Figure 2: Alteration of WzzE TMH2 can prevent ECA chain length regulation without affecting WzzE stability. (A) The effect of the indicated *wzzE* site-directed mutations on WzzE-FLAG protein levels was assessed using immunoblotting. GroEL serves as a loading control. **(B)** Densitometry was used for quantification of the immunoblots in (A). Levels of WzzE-FLAG compared to GroEL are shown normalized to the wild-type *wzzE* complemented strain. All variants show similar protein levels to wild-type WzzE, except for a small, but significant decrease in the WzzE$^{GG333/4LL}$ variant. Data are shown as the average of three biological replicates ± SEM and individual datapoints. * $p<0.05$ by Mann-Whitney test. **(C)** Strains carrying the indicated plasmids were assayed for their chloramphenicol resistance by EOP as a proxy for WzzE stability and multimerization ability. All *wzzE* mutants retained full chloramphenicol resistance, suggesting WzzE stability and multimerization. **(D)** Chain-length regulation of the linear ECA forms by the indicated WzzE variants was assayed by immunoblotting. Short and long exposures are shown. "X" indicates non-specific bands. WzzE$^{GG333/4LL}$ showed loss of

861   chain-length regulation while the other variants retained wild-type function. The Δ*wecA* strain

862   serves as negative control. **(E)** Chain-length regulation of the WzzE-Cat-FLAG variants was

863   assayed by immunoblot. Similar phenotypes were observed to those found in the WzzE

864   variants. "X" indicates non-specific band. All images are representative of at least three

865   independent experiments.

Figure 3. Conserved glycines in WzzE TMH 2 are required for ECA$_{CYC}$ synthesis. (A) The

strain carrying a plasmid with wild-type *wzzE* was grown with ^{15}N to label ECA and then

combined with an unlabeled strain carrying the indicated *wzzE* mutant or control. ECA$_{CYC}$ was

purified and the relative amount of unlabeled to labeled ECA$_{CYC}$ was determined by LC-MS.

WzzE$^{GG333/4LL}$ showed a loss of the ability to produce ECA$_{CYC}$. All plasmids were expressed in a

Δ*wzzE* Δ*wecH* strain background to prevent non-stochiometric acetylation of ECA. Data are

shown as the average of three biological replicates ± SEM and as individual datapoints. *

$p < 0.05$ by Mann-Whitney test. **(B)** The predicted structure of a WzzE octamer with WzyE was

calculated using AlphaFold 3 (Abramson et al., 2024). The structures of wild-type WzzE and

WzzE$^{GG333/4LL}$ monomers were aligned with the mutated residues shown in red. The insets show

the predicted distance between the two WzzE TMH in the wild type WzzE and WzzE$^{GG333/4LL}$.

Figure 4: The periplasmic region of WzzE contributes to WzzE stability and chain length regulation. (A) Immunoblotting was used to determine protein levels for the indicated WzzE-FLAG variants. GroEL serves as a loading control. **(B)** WzzE-FLAG levels from (A) were quantified by densitometry. WzzE-FLAG levels compared to GroEL are shown normalized to the wild-type *wzzE* complemented strain. All variants are similar to wild type WzzE except for WzzE$^{\Delta F104}$ which shows significantly reduced protein levels. Data are shown as the average of three biological replicates ± SEM and individual datapoints. * $p < 0.05$ by Mann-Whitney test. **(C)** The indicated strains were assayed for WzzE-Cat-FLAG stability as measured by chloramphenicol resistance. All *wzzE* point mutations showed chloramphenicol resistance indicating stability and multimerization, with the exception of WzzE$^{\Delta F104}$ where the tag is cleaved (**Figure S3C**). **(D)** Linear ECA chain-length regulation by WzzE variants was assayed by immunoblotting with ECA antibody. Long and short exposures are shown. The periplasmic point mutants have variable effects on chain length regulation as well as ECA$_{CYC}$ biogenesis. The WzzE$^{\Delta F104}$ and WzzE^{K162A} variants showed loss of ECA chain-length regulation while the other

895   variants had wild-type chain-length regulation. The Δ*wecA* strain serves as negative control. **(E)**

896   The effect of WzzE-Cat-FLAG variants on ECA chain-length regulation was assayed by

897   immunoblot. The variants show similar phenotypes as in (D) except for WzzE^{K162A}, which shows

898   normal chain-length regulation and WzzE$^{\Delta F104}$ which shows partial chain-length dysregulation.

899   All images are representative of at least three independent experiments.

Figure 5: The periplasmic domain of WzzE contributes to ECA$_{CYC}$ synthesis. (A) The strain

carrying wild-type *wzzE* was grown with ^{15}N to label ECA and then combined with an unlabeled

strain carrying the indicated *wzzE* mutant or control strain. After ECA$_{CYC}$ purification, the relative

amount of unlabeled to labeled ECA$_{CYC}$ was determined by LC-MS. WzzE$^{\Delta F104}$ and WzzE^{K162A}

had little to no ECA$_{CYC}$ production while WzzE^{F104H} showed a variable phenotype. All strains

included Δ*wecH* to prevent non-stochiometric acetylation of ECA. Data are shown as the

average of three biological replicates ± SEM and as individual datapoints. * $p < 0.05$ by Mann-

Whitney test. **(B)** Structures predicted as in **Figure 3** for wild-type WzzE and WzzE$^{\Delta F104}$ were

aligned. The mutated residue is shown in red on a WzzE monomer (B). The inset shows the

interaction between alpha helix 2 and an adjoining loop in wild-type WzzE.

Figure 6: Levels of ECA$_{CYC}$ production and chain-length regulation are genetically

separable. CRISPR-Cas9 was used to make the indicated mutations to *wzzE* in its native locus.

All strains contain *metE*::Tn*10*. **(A)** The strain wild type for *wzzE* was grown with ^{15}N to label

ECA and then combined with an unlabeled strain with the indicated *wzzE* mutant or control

strain. After ECA$_{CYC}$ purification, the ratio of unlabeled to labeled ECA$_{CYC}$ was determined by

LC-MS. All variants other than WzzE^{F104Y} produced significantly reduced ECA$_{CYC}$ levels. Data

are shown as the average of three biological replicates ± SEM and as individual datapoints. *

$p < 0.05$ vs. wild type by Mann-Whitney test; ‡ $p < 0.05$ between the indicated strains by Mann-

Whitney test. **(B)** Immunoblotting for ECA was conducted on the indicated chromosomal

mutants. A short and long exposure are shown. The WzzE^{ΔF104} and WzzE$^{GG333/4LL}$ variants

resulted in linear ECA dysregulation, while the other mutations more closely resembled wild-

type linear ECA regulation. The Δ*wecA* strain serves as negative control. **(C)**. ECA

Immunoblotting was performed on the indicated chromosomal mutants after ProK treatment to

more clearly assess chain length. WzzE mutants ΔF104 and GG333/4LL resulted in

dysregulation of linear ECA, whereas F104Y and F104H strongly resembled wild-type linear

ECA regulation. **(D)** Immunoblotting was used to assay ECA chain-length regulation in the

indicated strains. Deletion of *wzzB* and/or *fepE* in combination with Δ*wzzE* did not cause the

increased levels of linear ECA observed with the loss-of-function point mutants; however, the

Δ*wzzE* Δ*wzzB* Δ*fepE* strain did have a small increase in linear ECA levels compared to Δ*wzzE*

alone. All images are representative of at least three independent experiments.