1

Chain-length regulation by WzzE is necessary for, but genetically separable from, cyclic enterobacterial common antigen synthesis

Joseph F. Carr[a], Jennifer S. Rudolf[a]*, Daniel J. Warzecha[a]*, Yohannes H. Rezenom[b], &

Angela M. Mitchell[a]#

[a] Department of Biology, Texas A&M University, College Station, Texas, USA

[b] Chemistry Mass Spectrometry Facility, Texas A&M University, College Station, Texas, USA

* JSR and DJW contributed equally to this work. Author order was determined alphabetically.

Address correspondence to Angela M. Mitchell, amitchell@bio.tamu.edu

Running title: Biosynthesis of cyclic enterobacterial common antigen

Abstract

Bacteria of the order *Enterobacterales* (e.g., *Escherichia, Klebsiella, Yersinia*) produce an

invariant glycan, enterobacterial common antigen (ECA), that plays a role in maintaining outer

membrane (OM) impermeability. ECA is found as the headgroup of a phospholipid and attached

to LPS in the OM, and as a periplasmic cyclic form (ECA_{CYC}). WzyE polymerizes ECA repeat

units to form the ECA chain with final chain length regulated by WzzE, a class 1 polysaccharide

co-polymerase (PCP1). PCP1 function studies have shown interaction of the PCP1

transmembrane helices with the polymerase and a secondary activity found in the periplasmic

domain of the PCP1 are required for chain-length regulation. However, WzzE is also necessary

for ECA_{CYC} biosynthesis, and it remains unclear why loss of WzzE prevents ECA_{CYC} production

but not linear ECA production. Here, we constructed plasmid-based and chromosomal *wzzE*

mutants in *E. coli* K-12 with alterations that affect chain-length regulation of other PCP1 and

assessed their effects on ECA_{CYC} biogenesis. Our data show loss of chain-length regulation and

ECA_{CYC} synthesis from mutations altering transmembrane helix 2 and mutations altering the

periplasmic domain of WzzE. We also identified two WzzE variants to the same residue with

identical, near wild type linear ECA chain-length regulation but differing effects on ECA_{CYC}

production. Specifically, $WzzE^{F104Y}$ produces wild-type levels of ECA_{CYC}, while $WzzE^{F104H}$

produces twofold less ECA_{CYC} than wild type, demonstrating ECA_{CYC} synthesis is genetically

separable from chain length regulation. Overall, our results show that chain-length regulation by

WzzE is necessary but not sufficient for normal production of ECA_{CYC}.

Importance

The gram-negative cell envelope acts as a permeability barrier excluding many

antibiotics. Therefore, understanding envelope synthesis pathways has the potential to identify

targets for new antimicrobials. Here, we investigated synthesis requirements for an invariant

carbohydrate found throughout *Enterobacterales,* cyclic enterobacterial common antigen

(ECA$_{CYC}$), which is important for maintaining the envelope permeability barrier and relies on a

polymerase (WzyE) and co-polymerase (WzzE) for synthesis. We found ECA$_{CYC}$ synthesis

depends on WzzE regions necessary for interaction with WzyE and for chain-length regulation.

However, we can genetically separate phenotypes of ECA$_{CYC}$ production and chain-length

regulation. These data shed light on mechanisms contributing to the envelope permeability

barrier and on why the WzyE-WzzE system synthesizes a cyclic carbohydrate while other Wzy-

Wzz systems do not.

Introduction

Enterobacterales is an order of gram-negative bacteria containing numerous genera

with human pathogens (e.g., Escherichia, Salmonella, Klebsiella, Yersinia) (1, 2). These

bacteria can result in diseases ranging from respiratory infections, soft tissue infections, urinary

tract infections, and endocarditis (3). To survive in their enteric environment, members of this

family have a cell envelope and, specifically, an outer membrane which acts as a robust

permeability barrier. The outer membrane is an asymmetric bilayer that contains phospholipids

in its inner leaflet and predominantly lipopolysaccharides (LPS) in the outer leaflet (4, 5). The

LPS creates a strong hydrophilic network of lateral interactions that excludes large compounds

and hydrophobic compounds, such as many antibiotics, bile salts, and detergents, from entering

the cells (6, 7).

In addition to LPS, the outer membrane is decorated with lower-abundant components,

including enterobacterial common antigen (ECA) (8). Initially discovered in 1962 by Calvin M.

Kunin et al. as a "common antigen" associated with urinary tract infections, ECA is an invariant

carbohydrate moiety restricted to the order of Enterobacterales and found in all

Enterobacterales except for endosymbionts with significantly reduced genomes (8-11).

Interestingly, ECA_{CYC} plays a role in maintaining the envelope permeability barrier in E. coli K-12

and is important for the cell's resistance to detergents including bile salts (12). In addition, ECA

has been shown to play a role in the pathogenesis of several Enterobacterales (13-18). For

example, in Salmonella enterica serovar Typhimurium, strains lacking ECA exhibit reduced

virulence, increased sensitivity to bile salts (14, 15), and establish a low-level persistent

infection in mouse models (15).

ECA is defined by recurring trisaccharide repeat units (RUs) of GlcNAc (N-

acetylglucosamine), ManNAcA (N-acetyl-D-mannosaminuronic acid), and Fuc4NAc (4-

acetamido-4,6-dideoxy-D-galactose) (19). There are three forms of ECA, (i) ECA_{LPS}, which is

ECA covalently linked to the LPS core in place of O-antigen; (ii) ECA_{PG}, a cell surface form

linked to diacylglycerol-phosphate; and (iii) ECA_{CYC}, a soluble, cyclic form found within the

periplasmic space between the inner and outer membranes (10, 19-21). In contrast to the other

forms of ECA which have variable number of RU, ECA_{CYC} has a defined size of 4 RUs in

Escherichia coli K-12 (21). ECA_{PG} and ECA_{LPS} have 1 to 14 RUs and a modal value of 6 to 7 at

37 °C (12, 22). Structural analysis by crystallography, nuclear magnetic resonance (NMR), and

molecular dynamics suggest ECA_{CYC} can exist in one of two different three-dimensional

conformations, a rhomboidal shape and square shape (23, 24).

The ECA synthesis genes are located in the *wec* operon. ECA is synthesized through a

Wzz/Wzy-dependent pathway analogous to many O-antigen biosynthesis pathways (22, 25-30).

After a single trisaccharide ECA RU is assembled and linked to an undecaprenyl-phosphate

carrier (8), it is flipped to the outer leaflet of the inner membrane by the flippase, WzxE (26).

Then, a complex forms in which translocated undecaprenyl-phosphate-linked ECA subunits are

polymerized by WzyE with the number of RU in the final chain regulated by WzzE (31-34).

WzzE is an inner membrane protein with a large periplasmic domain that assembles into a

homo-octamer (**Figure S1**) (35-38). It belongs to the class 1 polysaccharide co-polymerase

(PCP1) family which includes WzzB, FepE, Wzz, and plays a crucial role in controlling

carbohydrate chain length during polymerization (39-41). Although PCP1 proteins share low

sequence similarity (35), they exhibit a conserved architecture, characterized by a large bell-

shaped periplasmic domain and two transmembrane helices (35-38). The C-terminal

transmembrane helix (TMH2) of Wzz interacts with Wzy, the polymerase, and this reaction is

necessary for O-antigen chain-length regulation (42, 43). However, mutations to Wzz that can

disrupt O-antigen chain-length regulation are distributed throughout Wzz's structure and are not

confined to TMH2 (44). In *Shigella flexneri*, WzyB and WzzB co-purify together and the second

transmembrane region (TMH2) of WzzB and WzzE plays a role in this interaction and in chain

length regulation (31). A cryo-EM study in *Pectobacterium atrosepticum* suggested WzyE lies

within the central lumen between the transmembrane helices of WzzE (34).

The role of WzzE is more complex in ECA synthesis than the role of other PCP1 in O-

antigen synthesis. In the absence of WzzE, linear ECA is made with increased variability in

number of RU, with both longer and shorter chain lengths being observed (22), whereas PCP1

deletion in the context of O-antigen production often leads to a large decrease in chain length

with some longer species observed (31, 45-47). In contrast to linear ECA, strains lacking WzzE

no longer produce ECA_{CYC}, demonstrating WzzE is necessary for ECA_{CYC} synthesis despite a

lack of demonstrated catalytic activity for PCP1 proteins (8, 12). Finally, in *E. coli* K-12, ElyC

regulates production of ECA_{PG} in such a way that loss of ElyC causes large increases in ECA_{PG}

production (48). This regulation is dependent on the presence of WzzE and likely on levels of

ECA_{CYC}, and changes in ECA_{PG} levels with loss of ElyC are much less in the absence of WzzE.

Because of this and freeing up of ECA precursors from the synthesis of ECA_{CYC} levels of linear

ECA are higher in strains with deletion of *wzzE* than they are in wild type cells (48, 49).

Several non-exclusive hypotheses for how PCPs regulate chain length have been

proposed (31, 39, 40, 50-54). (i) The inner or outer surface of the PCP may act as a molecular

ruler for chain length regulation (51). Specifically, the PCP may bind the extending carbohydrate

facilitating catalysis until the polymer length breaks the interaction (55). (ii) The PCP could serve

as a scaffold on which multiple polymerase molecules assemble (35). (iii) The elongating

carbohydrate could extend in the lumen of the PCP until it reaches the top of the periplasmic

domain and is exposed (50). (iv) The PCP could interact with the polymerase to sequester short

glycan chains from downstream ligases (i.e., WaaL), increasing the processivity of the

polymerization (56). Finally, (v) the loops at the top of the periplasmic bell may move in an

alternating up and down arrangement which allows for the polysaccharide chain to extend using

a "ratchet"-type mechanism (57).

However, the mechanism(s) necessitating WzzE for ECA_{CYC} synthesis are completely

unknown. Therefore, we set out to determine what functions of WzzE are required to produce a

cyclic carbohydrate in *E. coli* K-12. We used a site-directed mutagenesis approach and

biochemical analysis to determine whether the regions important for WzzE-WzyE interaction,

ECA chain-length regulation, and any additional mechanisms are necessary for ECA$_{CYC}$

biogenesis. Our findings demonstrate regions important for interaction between WzzE and

WzyE and for chain length regulation, are required for ECA$_{CYC}$ biosynthesis. However, our

results also demonstrate WzzE chain-length regulation can be genetically separated from

synthesis of ECA$_{CYC}$.

Results

Plasmid-based *wzzE* allows analysis of *wzzE* mutant function. To identify the functions of

WzzE required for ECA$_{CYC}$ synthesis in *E. coli* K-12, we examined the effects of *wzzE* mutations

analogous to those known to impact specific functions in other PCP1. We then assessed their

influence on the synthesis of both linear ECA and ECA$_{CYC}$. We first created *wzzE* constructs

constitutively expressed by the *tet* promoter on the low copy number pZS21 plasmid (58), which

does not contain a Tet repressor (**Figure 1A**). These constructs included wild-type *wzzE*

(p*wzzE*), *wzzE* fused with a 3′ glycine-serine linker and triple FLAG tag (p*wzzE*-FLAG) for

detection via immunoblot, *wzzE* translationally fused with chloramphenicol acetyltransferase

(p*wzzE-cat*), and *wzzE* fused to *cat* and triple FLAG (p*wzzE-cat*-FLAG). Based on the topology

of WzzE, we anticipated that the C-terminal fusions would be exposed in the cytoplasm, thereby

allowing Cat to function. Thus, this construct allowed us to identify defects in WzzE folding,

stability, and octamerization, as measured by their effect on Cat through chloramphenicol

resistance. This approach was informed by a translational coupling methodology previously

used a tool in *E. coli* (59, 60). This assay is particularly useful because the addition of the bulky

Cat (26KDa) protein likely imposes stress on WzzE folding. Additionally, the formation of the

functional cat trimer (61) is more likely if WzzE is properly multimerized.

After producing the constructs, we tested the function of fusions with each tag by

determining whether they complemented a *wzzE* deletion. In the absence of *yhdP*, ECA causes

severe sensitivity to vancomycin in a *wzzE* dependent manner (12). Therefore, we expressed

the *wzzE* constructs in a Δ*wzzE* Δ*yhdP* strain and tested for vancomycin sensitivity. Our results

demonstrated each construct complemented Δ*wzzE* by restoring vancomycin sensitivity in the

Δ*wzzE* Δ*yhdP* strain (**Figure 1B**)*.* Additionally, p*wzzE-cat* conferred chloramphenicol resistance

to the strain. To determine if the p*wzzE*-FLAG construct could be detected, we immunoblotted

for FLAG and for GroEL as a loading control (**Figure 1C**) and determined that the fusion protein

could be clearly detected.

To determine whether the function of WzzE as a chain-length regulator is complemented

by the p*wzzE* construct, we performed immunoblots for linear ECA (**Figure 1D**). Our results

revealed a range of ECA bands, with increasing molecular weight indicating molecules with

more RUs. The distribution of ECA chain lengths was wider in a Δ*wzzE* mutant (Lane 3) than in

wild-type cells (Lane 2) with an increase in longer and shorter ECA bands, indicating loss of

chain-length regulation. We also observed more linear ECA with Δ*wzzE* as we would expect

from previous findings (48, 62). For simplicity, we will refer to these changes as "loss of chain-

length regulation". However, expression of p*wzzE* (Lane 5) but not the empty vector (Lane 4)

restored chain-length regulation, albeit with a lower modal chain length than the wild-type strain

and much less linear ECA overall. These changes may suggest overactivity of WzzE due to

multi-copy expression. The same pattern was observed for the p*wzzE-cat* construct (Lane 7);

however, the p*wzzE*-FLAG and p*wzzE-cat*-FLAG constructs resulted in chain-length regulation

closer to that of wild type.

Finally, to confirm that ECA$_{CYC}$ synthesis is restored by the plasmid-expressed *wzzE*, we

expressed the p*wzzE* and empty vector plasmids in a Δ*wzzE* Δ*wecH* strain. WecH has been

shown to non-stoichiometrically acetylate ECA (63) and deleting *wecH* allows ECA$_{CYC}$ to be

quantitated as a single mass species (12, 63). Unlike the two linear forms of ECA, ECA$_{CYC}$ is

uncharged and cannot be detected by immunoblotting. Therefore, we then grew cultures with

either a normal (^{14}N) or heavy (^{15}N) nitrogen source, combined labeled and unlabeled samples,

purified ECA$_{CYC}$, and assayed ECA$_{CYC}$ levels by liquid chromatography-mass spectrometry (LC-

MS). ECA$_{CYC}$ has an monisotopic mass of 2428.9 Da and the labeling with heavy nitrogen

causes a 12 Da increase in this weight due to the presence of 12 nitrogen atoms in ECA$_{CYC}$ (12)

(**Figure S2AB**). For this type of assay, each peak consists of multiple minor peaks due to the

stochiometric incorporation of naturally occurring isotopes (e.g., ^{14}C) (see **Figure 1EF**, **S2C**).

When labeled and unlabeled samples carrying p*wzzE* were combined, we detected peaks at

both the unlabeled and labeled m/z (**Figure 1E**). However, when the labeled p*wzzE* sample was

combined with an unlabeled sample carrying the empty vector, only the ^{15}N labeled ECA$_{CYC}$

peak was detected (**Figure 1F**). ECA$_{CYC}$ production from the p*wzzE* and p*wzzE-cat* constructs

was very similar (**Figure S2C**). These data confirm plasmid-based expression of *wzzE* can

facilitate the synthesis of ECA$_{CYC}$, making this an appropriate system to investigate the role of

WzzE in ECA$_{CYC}$ biosynthesis.

Conserved glycines in TMH2 are necessary for WzzE function. After validating assays for

WzzE function, we set off to determine the role of TMH2 in WzzE function. TMH2 has been

shown to be important for interaction between WzzB and Wzy in O-antigen synthesis (31).

Previous mutagenesis studies of WzzB suggested a conserved GXXXG motif in the TM2 region

of WzzB from *Shigella flexneri Y* was important for function (64), likely through effects on helix-

helix packing (65-67). A WzzB double mutant G305A/G311A WzzB in *Shigella flexneri Y*

(G306A/G312A in *E. coli*) targeting this motif conferred short-chain O antigen (68). Based on

these studies, we hypothesized that the GXXXG motif located in TMH2 of WzzE plays a

significant role in mediating the helix-helix packing of the transmembrane region and interaction

with WzyE.

To test this hypothesis, we created TMH2 mutations (**Figure S1B**) in the p*wzzE*

constructs described in **Figure 1A**. We first sought to determine whether the levels of WzzE

protein produced by p*wzzE*-FLAG or p*wzzE-cat*-FLAG were altered by the mutations (**Figure**

2AB, Figure S3A). Our results showed a small but significant decrease in protein abundance

for our WzzE$^{GG333/4LL}$ in the WzzE-FLAG fusion (**Figure 2AB**), whereas no decrease is apparent

in the WzzE-Cat-FLAG fusion (**Figure S3A**). As the levels of the WzzE$^{GG333/4LL}$ variant are 82%

of wild type on average and as high as 98% in individual replicates, the decreased level of this

protein may not be biologically relevant to its function. To determine whether the mutations

might have caused alterations to protein stability or multimerization, we utilized our p*wzzE-cat*

construct to assay chloramphenicol resistance conferred by the fusion protein. None of our

mutations showed any chloramphenicol sensitivity (**Figure 2C, Figure S3B**), suggesting that

they are largely stable and multimerized, despite the small decrease in abundance for

WzzE$^{GG333/4LL}$. Next, we set out to determine if the mutations affected WzzE's ability to regulate

ECA chain length by performing ECA immunoblots. Three of the changes to TMH2, WzzE^{A323G},

WzzE$^{GG333/4AA}$, and WzzE^{G339A} (Lanes 4,5, and 7) did not affect ECA chain length regulation

(**Figure 2D**). However, WzzE$^{GG333/4LL}$ (Lane 6) showed a similar or greater disruption of chain

length regulation to that observed in the empty vector control (Lane 3).

The TMH2 GXXXG motif affects ECA$_{CYC}$ synthesis. Given the effect of WzzE$^{GG333/4LL}$ on ECA

chain length regulation, we next decided to examine the effect of the TMH2 mutations on

ECA$_{CYC}$ biogenesis. Therefore, we utilized LC-MS to detect and quantitate ECA$_{CYC}$ in purified

samples as in **Figure 1EF**. We expressed the TMH2 mutants or controls in a Δ*wzzE* Δ*wecH*

strain background and labeled the wild-type *wzzE* strain with ^{15}N, while the mutants were grown

with ^{14}N. Our quantification demonstrated WzzE^{A323G}, WzzE$^{GG333/4AA}$, and WzzE^{G339A} did not

significantly change ECA$_{CYC}$ levels, whereas the WzzE$^{GG333/4LL}$ mutant produced significantly

reduced levels of ECA$_{CYC}$, similar to those of the empty vector control (**Figure 3A**). Together

with our chain length regulation data, these results demonstrate the chain-length regulating

activities of WzzE found in TMH2 are required for ECA$_{CYC}$ synthesis.

To visualize the potential effect of the mutants on WzzE and its complex with WzyE, we

predicted the structure of the WzyE:WzzE(8) complex for wild-type WzzE and WzzE[GG333/4LL]

using AlphaFold 3 (69) (**Figure 3B** & **S4**, **Table S1**) and aligned wild-type WzzE and

WzzE[GG333/4LL] monomers. When we closely examined the transmembrane region of the

structure, the model suggested a 2 Å increase in distance between TMH1 and TMH2 in

WzzE[GG333/4L] likely due to steric interference (**Figure 3B**). This suggests a possible hypothesis

that the arrangement of the TMHs is responsible for the phenotype of this mutant.

The periplasmic domain of WzzE functions in chain length regulation. After assessing the

importance of WzzE TMH2 for ECA$_{CYC}$ biosynthesis, we wanted to determine whether the

chain-length regulating activities found in the periplasmic domain of WzzE were necessary for

ECA$_{CYC}$ biosynthesis. There are 21 periplasmic residues conserved between WzzE, FepE, and

WzzB in *E. coli*, which are likely to be important for the functions of the PCP1 proteins (35).

Furthermore, we hypothesized that other non-conserved residues might also contribute to

WzzE's role in chain-length regulation or ECA$_{CYC}$ synthesis functions. Therefore, we made

mutations in a conserved cluster of predominantly hydrophobic residues that lie within the core

of the α/β domains in strands β1 and β4, as well as helix α2 and the N terminus of helix α6 (see

Figure S1B). We also targeted non-conserved residues throughout WzzE in positions similar to

those that conferred changes to O-antigen chain length in other PCP1s (35).

We started characterizing the periplasmic mutants (**Figure S1B**) by investigating WzzE

protein levels from the p*wzzE*-FLAG construct. One of the mutants (WzzE[ΔF104]) had

undetectable WzzE-FLAG protein levels, while the other mutants retained wild-type levels of

accumulation (**Figure 4AB**). When we tested the levels of WzzE-Cat-FLAG produced from this

fusion, we detected a faint band at the molecular weight of the fusion and a second lower

molecular weight band for WzzE[ΔF104], suggesting that the C-terminal tags were cleaved from the

protein (**Figure S3C**). Given the cleavage of the tags, no conclusion can be drawn for the

stability of WzzE$^{\Delta F104}$. All other protein levels were similar to wild type. Then, we tested whether

the *wzzE* constructs showed changes to protein stability and/or multimerization through

chloramphenicol resistance using the p*wzzE-cat* construct. WzzE^{K162A}, WzzE^{L259A}, and

WzzE^{F104Y} resulted in partial or full chloramphenicol sensitivity in the WzzE-Cat fusion (**Figure**

S3D) but showed full chloramphenicol resistance in the WzzE-Cat-FLAG fusion that better

complements for linear chain length regulation (**Figure 4C**). Thus, it is possible these mutants

could be slightly unstable or have a decreased ability to multimerize. The other mutants

conferred resistance to chloramphenicol similar to wild type in both fusions (**Figure 4C, S3D**).

WzzE$^{\Delta F104}$ showed wild-type levels of chloramphenicol resistance, likely due to cleavage of the

C-terminal tags. We tested an additional 18 mutations in the periplasmic region that caused

chloramphenicol sensitivity in the WzzE-Cat background, emphasizing the importance of the

periplasmic domain on the structural stability of WzzE (**Figure S5**).

274   After characterizing the effect of the periplasmic mutations on WzzE levels and stability,

we investigated their effect on ECA chain-length regulation (**Figure 4D**). The WzzE$^{\Delta F104}$ and

WzzE^{K162A} mutants (lanes 4 and 7) prevented regulation of ECA chain length. In contrast,

WzzE^{F104Y}, WzzE^{F104H}, WzzE^{K162S}, and WzzE^{K162T} (lanes 5, 6, 8 and 9) demonstrated wild-type

ECA chain length regulation. The WzzE^{L259A} and WzzE^{L259I} mutants did not affect ECA chain-

length regulation despite the lack of chloramphenicol resistance observed for WzzE^{L259A}. It may

be that this protein is partially destabilized but able to function in the absence of the Cat fusion.

Overall, these mutants provide an interesting range of phenotypes useful for the investigation of

ECA$_{CYC}$ synthesis.

Periplasmic residues of WzzE are important for ECA$_{CYC}$ biogenesis. We then assessed the

impact of the periplasmic mutants on ECA$_{CYC}$ biosynthesis by quantifying ECA$_{CYC}$ levels using

LC-MS. Both the WzzE$^{\Delta F104}$ and WzzE^{K162A} mutants exhibited impaired linear chain-length

regulation and produced little to no ECA$_{CYC}$ (**Figure 5A**). This lack of ECA$_{CYC}$ production could

be due to either (i) insufficient protein accumulation and/or stability, or (ii) the disruption of chain-

length regulating activities in these mutants. In contrast, WzzE^{F104Y}, WzzE^{K162S}, WzzE^{K162T},

WzzE^{L259A}, and WzzE^{L259I} had similar levels of ECA$_{CYC}$ biosynthesis to wild type. WzzE^{F104H}

showed a highly variable level of ECA$_{CYC}$ despite having wild type chain-length regulation and

protein levels, suggesting that this mutation might have some sort of variable effect on ECA$_{CYC}$

synthesis. These data highlight the importance of the periplasmic domain of WzzE on ECA$_{CYC}$

synthesis, likely through its structural role, chain-length regulating activity, and possibly a

function specific to ECA$_{CYC}$ synthesis.

295   Given the varying effects of the F104 mutations on ECA$_{CYC}$ synthesis and linear ECA

chain length regulation, we decided to visualize the possible effect of these mutations on

WzzE's structure using AlphaFold 3 (69) and aligned the wild type and mutant complexes

(**Figure 5B** & **S4**, **Table S1**). When examining these structures, we observed that F104 is

predicted to make contacts with a neighboring loop. It is possible that alterations in these

contacts are responsible for the changes in function for variants at this residue.

ECA$_{CYC}$ biogenesis is genetically separable from chain-length regulation. Our results,

using plasmid-based expression of WzzE, identified several point mutations with differing effects

on ECA$_{CYC}$ function. Most interestingly, we identified a change to TMH2 (WzzE$^{GG333/4LL}$) that

prevented chain-length regulation and ECA$_{CYC}$ synthesis, despite maintaining protein stability

and near wild type protein accumulation. Additionally, several mutations to residue F104

exhibited varying effects on WzzE accumulation, chain-length regulation, and ECA$_{CYC}$

production. However, plasmid-based expression of wild-type *wzzE* resulted in lower levels of

linear ECA and lower chain length than expression of *wzzE* from its native locus (**Figure 1D**),

which may have influenced the phenotypes of the mutants assayed. In addition, we observed a

high level of variability in ECA$_{CYC}$ levels produced in several of the periplasmic mutants, which

we reasoned could be because of the non-native expression of WzzE. Therefore, we used

CRISPR-Cas9 to produce selected point mutations to *wzzE* at its native locus and assayed their

effect on linear ECA and ECA$_{CYC}$.

We first performed LC-MS on purified ECA$_{CYC}$ to determine the effect of the

chromosomal mutants on ECA$_{CYC}$ biosynthesis. The WzzE$^{\Delta F104}$ and WzzE$^{GG333/4LL}$ chromosomal

mutants produced very little to no ECA$_{CYC}$, similar to what is observed with Δ*wzzE* (**Figure 6A**).

In contrast, WzzE^{F104H} produced measurable levels of ECA$_{CYC}$ that were about 2-fold lower than

the wild type. Finally, WzzE^{F104Y} did not cause a significant change in ECA$_{CYC}$ levels when

compared to the wild type. We then assayed linear ECA chain length regulation by immunoblot

(**Figure 6B**). Consistent with the plasmid-based system, WzzE$^{\Delta F104}$ and WzzE$^{GG333/4LL}$ were not

able to regulate linear ECA chain length and dysregulated chain length was observed.

Interestingly, the levels of linear ECA and chain-length dysregulation were even greater than

that of Δ*wzzE*, suggesting that these mutants produced a non-functional protein that interacted

with ECA synthesis differently than loss of the WzzE protein. WzzE^{F104H} and WzzE^{F104Y} each

had identical, intact ECA chain-length regulation, which was close to that of wild-type WzzE but

with some increase in chain-length. The identical regulation of linear ECA chain-length by these

mutants but differing effects of ECA$_{CYC}$ synthesis demonstrate that levels of ECA$_{CYC}$ production

are genetically separable from linear chain-length regulation.

We were surprised that the dysregulation of ECA chain length, and even more the levels

of ECA, in the WzzE$^{\Delta F104}$ and WzzE$^{GG333/4LL}$ mutants was greater than the Δ*wzzE* control and we

wondered whether in a Δ*wzzE* strain the remaining PCP1s could partially regulate ECA chain

length through interaction with WzyE. In this scenario, the nonfunctional WzzE mutants could

sequester WzyE from the other PCP1s. Therefore, we assayed whether we could cause further

disruption of ECA chain-length regulation in a Δ*wzzE* strain by deleting *wzzB* and/or *fepE*.

However, our results demonstrated Δ*wzzB*, Δ*fepE*, and Δ*wzzB* Δ*fepE* in a Δ*wzzE* strain has no

effect on ECA chain-length regulation (**Figure 6C**). Nevertheless, there was a small increase in

the amount of linear ECA observed in the Δ*wzzE* Δ*wzzB* Δ*fepE* strain. Taken together our

results demonstrate that mutations to *wzzE* disrupting linear ECA chain length regulation can

prevent synthesis of ECA_{CYC}, and that production of normal levels of ECA_{CYC} is genetically

separable from regulation of ECA chain length.

Discussion

In this study, we constructed and characterized a set of *wzzE* site-directed mutants to

identify what functions of WzzE are necessary to produce ECA_{CYC}. Previous mutagenesis

studies have suggested regions of *Shigella flexneri* WzzB that are important for proper function

(33, 42). However, despite the multitude of hypotheses for PCP1s function, there have been no

reports on WzzE's role in producing a cyclic carbohydrate, ECA_{CYC}. Here, we demonstrate that

changes to amino acids F104 and K162 in the periplasmic region and to the GXXXG motif in

TMH2 can impair ECA chain-length regulation and ECA_{CYC} production, suggesting WzzE's

ability to regulate ECA chain length is necessary for ECA_{CYC} synthesis. Moreover, our data

demonstrate that regulation of linear ECA chain length and the level of ECA_{CYC} production are

genetically separable, as we have identified two mutations to the same WzzE residue (F104)

which have differing effects on ECA_{CYC} levels but produce identical regulation of linear ECA

chain length. This shows that regulation of chain length by WzzE is necessary but not sufficient

for production of normal levels of ECA_{CYC}.

Interestingly, despite the low sequence identity amongst PCP1s, they have highly similar

three-dimensional structures that are marked by two transmembrane domains and an α-helix/β-

sheet domain within the periplasmic space that extends ~100 Å (35-38). Additionally, PCP1s

have a conserved GXXXG motif that has been identified to be important for chain length

regulation, as a G305A/G311A double mutant in *Shigella flexneri* WzzB that encompassed and

extended outside of this motif resulted in short-chain O antigen production (64, 68). Notably, this

same study showed that single, double, and triple Gly to Ala mutations in this motif did not affect

O-antigen chain length apart from the G305A/G311A mutation (37, 64). Given this, we focused

on residues within and around the GXXXG motif, characterized their effect on chain-length

regulation and ECA_{CYC} biogenesis, and identified two adjacent, conserved glycine residues in

the GXXXG motif of WzzE—analogous to G305/G306 in WzzB—as critical for ECA_{CYC}

biosynthesis. We observed no changes to either chain length regulation or ECA_{CYC} production

when we introduced an alanine to both sites. However, we did observe dysregulation of ECA

chain length and prevention of ECA_{CYC} synthesis by introducing two adjacent leucines.

In other proteins, GXXXG motifs have been shown to mediate dimerization and

multimerization of transmembrane domains (70, 71). Thus, this motif may be required for proper

interaction between TMH1 and TMH2 and the long side chain on the leucine could push TMH1

and TMH2 further apart, reducing their interaction. In support of this idea, our structural

modeling suggests increased distance between TMH1 and TMH2 in the $WzzE^{GG333/4LL}$ mutant

(**Figure 3D**, **S6A**). When we aligned the monomers of WzzE that most closely interacted with

WzyE, we found that the angle of WzyE was rotated in the plane perpendicular to the

membrane in $WzyE-WzzE^{GG333/4LL}$ model compared to the wild type (**Figure S6BC**). The

distance in the structure between the TMH of opposite WzzE monomers was 2.0 to 2.7 Å

smaller in the $WzzE^{GG333/4LL}$ model than in the wild-type model. In compensation, WzyE takes on

a more condensed structure (1.4 Å narrower) in the $WzzE^{GG333/4LL}$ model. As this type of rotation

and structure change could impair WzyE function, this prediction provides a possible hypothesis

that $WzzE^{GG3334/LL}$ has impaired interaction with WzyE. Alternatively, the shift of the TMH may

affect the less clearly defined activity of WzzE that regulates chain length.

In contrast to the TMH2 mutants, many of the mutations that we made in the periplasmic

domain of WzzE cause the protein to become unstable as assayed by chloramphenicol

resistance conferred by the WzzE-Cat fusion, suggesting the periplasmic domain is critical for

WzzE's structural integrity and/or multimerization. Additionally, the periplasmic domain has been

proposed to be the "chain-length regulating" portion of WzzE. Studies with other PCP1s have

shown that multiple, surface-exposed regions on both the interior and exterior space of the

periplasmic domain affect chain-length regulation, resulting in either shorter or longer O-antigen

chains (36, 52, 64, 72, 73). More recently, increasing evidence suggests that the inner surface

of the WzzE octamer may bind to the elongating carbohydrate to facilitate growth of the chain

(34, 40, 56).

The internal cavity of WzzE possesses two highly concentrated negatively charged

bands of residues separated by a band of neutrally-charged-flexible-loops in between, which

may act to bind the growing carbohydrate chain (57). A loop at the top of the periplasmic domain

has been suggested to "rachet" up and down to facilitate carbohydrate movement over this

binding surface (57). While F104 is found buried inside the bottom of the "bell" of the WzzE

periplasmic domain and may have indirect effects on carbohydrate binding, K162 is located

near the lower negatively charged band and the membrane, and L259 is located in the loop that

has been proposed to drive the ratchet mechanism. Our mutations to L259 did not affect chain-

length regulation; however, loss of charge at K162 did prevent chain-length regulation and

ECA_{CYC} synthesis, supporting this model of ECA chain-length elongation. As to how the

$wzzE_{K162A}$ and $wzzE_{F104\Delta}$ mutations lead to loss of ECA_{CYC} production, we speculate that the

chain-length regulating activity of WzzE—related to the carbohydrate binding and ratcheting

mechanism—is necessary for ECA_{CYC} synthesis.

In contrast, the $wzzE_{F104H}$ mutation reduces ECA_{CYC} synthesis while maintaining chain-

length regulation. There is a small difference in linear ECA chain length between this mutant

and the wild type; however, the chain length and regulation thereof is identical to that of the

$wzzE_{F104Y}$ mutant which produces wild-type levels of $ECA_{CYC.}$ These data demonstrate that

chain-length regulation by WzzE is not sufficient for normal levels of ECA_{CYC} production and are

an indication that WzzE may play a role in ECA_{CYC} synthesis beyond that of chain-length

regulation. One possible mechanism would be that WzzE bends ECA molecules of the correct

size into an orientation that can be spontaneously cyclized. Another possibility is that WzzE

allows the binding of another protein (i.e., a cyclase) that can cyclize ECA chains of the

appropriate length. Finally, it is possible that WzzE has specific features that promote the

formation of ECA chains with 4 RUs that can later be cyclized to form ECA_{CYC}. Further, genetic

and biochemical experiments will be necessary to distinguish between these possibilities and to

solidify an additional role for WzzE in ECA_{CYC} biogenesis.

Lastly, we would like to draw attention to the observation that the $WzzE^{F104\Delta}$ and

$WzzE^{GG333/4LL}$ mutants had greater linear ECA levels than is observed when deleting *wzzE*

entirely. We further showed that this is not due to the influence of other PCP1s found in *E. coli*,

though there is a slight increase in linear ECA levels in a triple mutant of all the PCP1 genes.

Thus, why $WzzE^{F104\Delta}$ and $WzzE^{GG333/4LL}$ have increased linear ECA levels remains a mystery. It

is possible that the inactive WzzE octamers sequester some growing ECA chains resulting in

the increased variability in chain length. Although this can be investigated in the future, the

difference between the Δ*wzzE* strain and these mutants provides evidence that the mutants

produced a protein that could not regulate chain length and could not synthesize ECA_{CYC},

increasing the strength of our conclusion that chain-length regulation is important for ECA_{CYC}

synthesis.

Through this work, we have examined only a very small number of the residues in

WzzE, so it is quite likely there are mutations that cause effects solely for ECA_{CYC} production.

Future work to identify these mutations will provide the need evidence to determine the

mechanism of ECA_{CYC} synthesis. Nevertheless, our work provides insight into requirements for

carbohydrate cyclization and chain-length regulation. We propose coordinated action between

WzyE and WzzE, as well as chain-length regulation, is necessary for ECA_{CYC} biosynthesis and

suggest that charged residues in the periplasmic base help to facilitate chain elongation and

ECA_{CYC} biosynthesis. We also demonstrated that chain-length regulation can be genetically

separated from synthesis of normal levels of ECA_{CYC}, suggesting that WzzE may play a role in

ECA_{CYC} biosynthesis beyond that of chain-length regulation. Thus, our work provides insight into

the characteristics of WzzE that make it necessary for synthesis of a cyclic carbohydrate.

Materials and Method

Strains and growth conditions. The strains used in this study are listed in **Table S2**. Cultures

were inoculated and grown at 37 °C in LB Lennox unless otherwise noted, and as necessary,

cultures were supplemented with 20 mg/L chloramphenicol, 25 mg/L kanamycin, 10 mg/L

tetracycline, or a combination thereof. Insertion-deletion alleles containing kanamycin resistance

cassettes from the Keio collection (74) were moved to isogenic backgrounds using P1vir

transduction (75). Clean deletions were constructed utilizing the Flp recombinase-FRT system

as has described (76). Chromosomal mutagenesis in WzzE was constructed using a CRISPR

Cas-9 system harboring a pCRISPR::protospacer plasmid in W3110 containing a heat shock

inducible λcI857 lysogen as has been described (77, 78). The pCas9 (Addgene #42876) and

pCRISPR (Addgene #42975) plasmids were a kind gift from Professor Luciano Marraffini

(Rockefeller University). Oligos used to form the protospacer and repair templates used to make

the desired mutations are listed in **Table S3**. Prior to chromosomal mutagenesis, *metE*::Tn*10*

was linked to *wzzE* (79) allowing the mutations to be moved to MG1655 through P1vir

transduction. Strains carrying *metE*::Tn*10* were supplemented with 1μm cobalamin (vitamin

B12) when grown in minimal media (80).

Plasmids, cloning, and site directed mutagenesis. All primers used in this study are listed in

Table S4. To sub-clone the p*wzzE* expression plasmid, chromosomal *wzzE* with an ochre stop

codon was PCR amplified using primers pZS21-wzzE_fwd and wzzE-pZS21_rev. The low copy

number pZS21 plasmid (58) was amplified using primers pZS21_fwd and pZS21_rev. The

plasmid was assembled using HiFi Assembly Master Mix (New England Biolabs) as per the

manufacturer's instructions. To construct p*wzzE-cat*, *wzzE* without a stop codon was amplified

using primers pZS21-wzzE_fwd and wzzE-cat_rev. The chloramphenicol acetyltransferase

gene, *cat*, was amplified from pBAD33 (81) using primers wzzE-cat_fwd and cat-pZS21_rev.

These fragments were assembled into pZS21 amplified as before using HiFi Assembly. To

construct p*wzzE*-FLAG, p*wzzE* was amplified using primers p*wzzE*_x3F_Fwd and

p*wzzE*_x3F_Rvr, which contained sequence homology to a glycine serine linked FLAG tag that

incorporated a stop codon. The glycine serine linked FLAG tag was a gene block fragment

(**Table S3**) assembled into the p*wzzE* plasmid through HiFi Assembly. To clone the p*wzzE-cat-*

FLAG plasmid, *cat* was amplified using the WzzE:Cmr Fp and WzzE:Cmr Rp and the p*wzzE-*

FLAG plasmid was amplified using the Cmr: GS-x3F FP and wzzE-cat_rev. The fragments were

assembled using HiFi Assembly.

Site-direct mutations were constructed using Q5 PCR mutagenesis (New England

Biolabs). Briefly, non-overlapping primers containing the mutation of interest were used to

amplify the indicated pZS21-sanA plasmids (pZS21-*wzzE*, pZS21-*wzzE-cat*, and pZS21-*wzzE-*

FLAG). The PCR products were then digested with DpnI (New England Biolabs),

phosphorylated with T4 polynucleotide kinase (New England Biolabs) and self-ligated using T4

DNA ligase (New England Biolabs). Ligations were transformed into Turbo cloning cells (New

England Biolabs) and the mutations were confirmed by sequencing (Plasmidsaurs). The primers

used for each site-directed mutant are described in **Table S4**. Modeling of the predicted mutant

structure was performed using AlphaFold 3 (69) with eight monomers of WzzE and one of

WzyE. Relevant confidence scores are given in **Table S1** and **Figure S2**.

Antibiotic Sensitivity. Efficiency of plating (EOP) assays were performed by plating 10-fold

dilutions of overnight cultures on LB plates containing the indicated antibiotics using a 48-pin

tool. Plates were incubated overnight at 30 °C before imaging.

Immunoblot Analysis. Immunoblots for WzzE-FLAG and ECA were conducted as previously

described with minor modification (12, 82). For analysis of WzzE-FLAG protein levels, samples

with equal OD_{600} from overnight cultures of the indicated strains were lysed by boiling for 5

minutes in sample loading buffer and were loaded on 12% SDS-PAGE gels. Samples for ECA

analysis were prepared and immunoblotted as previously described. The antibodies used for

blotting were as follows: M2 αFLAG (Millipore Sigma, 1:50,000 dilution), αGroEL (Millipore

Sigma, 1: 60,000 dilution), αECA (a gift from Professor Renato Morona (University of Adelaide),

1:30,000), rabbit α-mouse (Prometheus, 1:100,000 dilution), goat α-rabbit (Prometheus,

1:100,000 dilution). Levels of WzzE-FLAG were quantitated by densitometry using ImageJ as

has been previously described (12) on immunoblots with the lowest exposure at which WzzE

bands could be detected.

Quantification of Cyclic ECA. Quantification of ECA_{CYC} was performed as previously

described with minor modifications (12). Briefly, cells were grown in M63 glucose minimum

media without nitrogen supplemented with 0.2% $(^{14}NH_4)_2SO_4$ or $(^{15}NH_4)_2SO_4$ and ECA was

purified as previously described. Sample analysis was performed using a Thermo Scientific Q

Exactive Focus and liquid chromatography unit (UltiMate 3000 RS). Purified samples were

dissolved in 0.1% formic acid and separated by injecting 20 μL of sample into the Synergi

Hydro-RP (4.6 x 250 mm; 4.0 μm) column (Phenomenax). The mobile phase consisted of 0.1%

Formic acid (eluent A) and methanol (eluent B). The flow rate was set at 800 μL/min with the

following gradient: 0–5.0 min 5% B, 5.0-12.0 min 5-60% B, 12-14 min 60-90% B, 14-17 min 90%

B, and returned back to 5% B in 0.1 min and held for 5 min at 5% B. The Q Exactive Focus ESI

source was operated in full MS (100-1000 m/z). The mass resolution was tuned to 70000

FWHM at m/z 200. The spray voltage was set to 3.3 kV in negative mode, and the sheath gas

and auxiliary gas flow rates were set to 60 and 20 arbitrary units, respectively. The transfer

capillary and auxiliary gas heater temperatures were held at 380 and 300 °C, respectively. The

S-Lens RF level was set at 70 V. Exactive Series 2.11 /Xcalibur 4.2.47 software was used for

data acquisition and processing. For relative quantification, the ratio of the areas of the ^{14}N to

^{15}N ECA_{CYC} peaks was calculated.

Acknowledgements

Thank you to the members of the Mitchell Lab for helpful and productive discussions and for

critical reading of our manuscript. We also thank Drs. Jolene Ramsey, Deborah Siegele,

Jennifer Herman, and Ryland Young (Texas A&M University), as well as members of the

Ramsey lab, for their insightful comments. The ECA antibody was a kind gift from Professor

Renato Morana (University of Adelaide). The pCas9 and pCRISPR plasmids were a kind gift

from Professor Luciano Marraffini (Rockefeller University) through the Addgene Repository. This

work was supported by the National Institute of Allergy and Infectious Disease under award

number R01-AI155915 (to A.M.M.).

References

1. Doern CD. 2024. Chapter 2 - Classification of medically important bacteria, p 9-21. *In*

Tang Y-W, Hindiyeh MY, Liu D, Sails A, Spearman P, Zhang J-R (ed), Molecular Medical

Microbiology (Third Edition) doi:https://doi.org/10.1016/B978-0-12-818619-0.00029-0.

Academic Press.

2. Adeolu M, Alnajar S, Naushad S, S. Gupta R. 2016. Genome-based phylogeny and

taxonomy of the '*Enterobacteriales*': proposal for *Enterobacterales* ord. nov. divided into

the families *Enterobacteriaceae*, *Erwiniaceae* fam. nov., *Pectobacteriaceae* fam. nov.,

Yersiniaceae fam. nov., *Hafniaceae* fam. nov., *Morganellaceae* fam. nov., and

Budviciaceae fam. nov. International Journal of Systematic and Evolutionary

Microbiology 66:5575-5599.

3. Ramirez D, Giron M. 2024. Enterobacter Infections, StatPearls. StatPearls Publishing

Copyright © 2024, StatPearls Publishing LLC., Treasure Island (FL).

4. Maher C, Hassan KA. 2023. The Gram-negative permeability barrier: tipping the balance

of the in and the out. mBio 14:e01205-23.

5.   Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. Cold Spring Harb

Perspect Biol 2:a000414.

6.   Farfour E, Dortet L, Guillard T, Chatelain N, Poisson A, Mizrahi A, Fournier D, Bonnin

RA, Degand N, Morand P, Janvier F, Fihman V, Corvec S, Broutin L, Le Brun C, Yin N,

Héry-Arnaud G, Grillon A, Bille E, ..., On Behalf Of The Gmc Study G. 2022.

Antimicrobial Resistance in *Enterobacterales* Recovered from Urinary Tract Infections in

France. Pathogens 11:356.

7.   Aminov RI. 2010. A brief history of the antibiotic era: lessons learned and challenges for

the future. Front Microbiol 1:134.

8.   Rai AK, Mitchell AM. 2020. Enterobacterial Common Antigen: Synthesis and Function of

an Enigmatic Molecule. mBio 11:mBio.01914-20.

9.   Jenkins C, Rentenaar RJ, Landraud L, Brisse S. 2017. 180 - *Enterobacteriaceae*, p

1565-1578.e2. *In* Cohen J, Powderly WG, Opal SM (ed), Infectious Diseases (Fourth

Edition) doi:https://doi.org/10.1016/B978-0-7020-6285-8.00180-5. Elsevier.

10.   Kunin CM. 1963. Separtion, chacterization, and Biological Significance of a Common

Antigen in *Enterobacteriaceae*. J Exp Med 118:565-86.

11.   Kunin CM, Beard MV, Halmagyi NE. 1962. Evidence for a Common Hapten Associated

with Endotoxin Fractions of *E. coli* and other *Enterobacteriaceae*. Proc Soc Exp Biol Med

111:160-166.

12.   Mitchell AM, Srikumar T, Silhavy TJ. 2018. Cyclic Enterobacterial Common Antigen

Maintains the Outer Membrane Permeability Barrier of *Escherichia coli* in a Manner

Controlled by YhdP. mBio 9:01321-18.

13.   Aguiniga LM, Yaggie RE, Schaeffer AJ, Klumpp DJ. 2016. Lipopolysaccharide Domains

Modulate Urovirulence. Infection and Immunity 84:3131-3140.

14.   Kalynych S, Morona R, Cygler M. 2014. Progress in understanding the assembly

process of bacterial O-antigen. FEMS Microbiol Rev 38:1048-65.

15.   Gilbreath JJ, Colvocoresses Dodds J, Rick PD, Soloski MJ, Merrell DS, Metcalf ES.

2012. Enterobacterial common antigen mutants of *Salmonella enterica* serovar

Typhimurium establish a persistent infection and provide protection against subsequent

lethal challenge. Infect Immun 80:441-50.

16.   Armbruster CE, Forsyth-DeOrnellas V, Johnson AO, Smith SN, Zhao L, Wu W, Mobley

HLT. 2017. Genome-wide transposon mutagenesis of *Proteus mirabilis*: Essential genes,

fitness factors for catheter-associated urinary tract infection, and the impact of

polymicrobial infection on fitness requirements. PLOS Pathogens 13:e1006434.

17.   Holmes CL, Wilcox AE, Forsyth V, Smith SN, Moricz BS, Unverdorben LV, Mason S, Wu

W, Zhao L, Mobley HLT, Bachman MA. 2023. *Klebsiella pneumoniae* causes bacteremia

using factors that mediate tissue-specific fitness and resistance to oxidative stress.

PLOS Pathogens 19:e1011233.

18.   Guérin F, Lallement C, Goudergues B, Isnard C, Sanguinetti M, Cacaci M, Torelli R,

Cattoir V, Giard J-C. 2020. Landscape of in vivo Fitness-Associated Genes of

Enterobacter cloacae Complex. Frontiers in Microbiology 11:1609.

19.   Mannel D, Mayer H. 1978. Isolation and Chemical Characterization of the

Enterobacterial Common Antigen. European Journal of Biochemistry 86:361-370.

20.   Goździewicz TK, Łukasiewicz J, Ługowski C. 2015. [The structure and significance of

enterobacterial common antigen (ECA)]. Postepy Hig Med Dosw (Online) 69:1003-12.

21.   Kajimura J, Rahman A, Rick PD. 2005. Assembly of cyclic enterobacterial common

antigen in *Escherichia coli* K-12. J Bacteriol 187:6917-6927.

22.   Barr K, Klena J, Rick P. 1999. The Modality of Enterobacterial Common Antigen

Polysaccharide Chain Lengths Is Regulated by o349 of the wec Gene Cluster of

Escherichia coli K-12. Journal of bacteriology 181:6564-8.

23.   Färnbäck M, Eriksson L, Senchenkova Sy, Zych K, Knirel YA, Sidorczyk Z, Widmalm G.

2003. Crystal Structure of a Cyclic Enterobacterial Common Antigen. Angewandte

Chemie International Edition 42:2543-2546.

24.   Erbel PJ, Seidel R, Macintosh SE, Gentile LN, Amor JC, Kahn RA, Prestegard JH,

McIntosh LP, Gardner KH. 2004. Cyclic enterobacterial common antigen: potential

contaminant of bacterially expressed protein preparations. J Biomol NMR 29:199-204.

25.   Eade CR, Wallen TW, Gates CE, Oliverio CL, Scarbrough BA, Reid AJ, Jorgenson MA,

Young KD, Troutman JM. 2021. Making the enterobacterial common antigen glycan and

measuring its substrate sequestration. ACS Chem Biol 16:691-700.

26.   Rick PD, Mayer H, Neumeyer BA, Wolski S, Bitter-Suermann D. 1985. Biosynthesis of

enterobacterial common antigen. J Bacteriol 162:494-503.

27.   Barr K, Rick PD. 1987. Biosynthesis of enterobacterial common antigen in *Escherichia*

coli. In vitro synthesis of lipid-linked intermediates. J Biol Chem 262:7142-50.

28.   Barr K, Nunes-Edwards P, Rick PD. 1989. In vitro synthesis of a lipid-linked trisaccharide

involved in synthesis of enterobacterial common antigen. J Bacteriol 171:1326-32.

29.   Bertani B, Ruiz N. 2018. Function and Biogenesis of Lipopolysaccharides. EcoSal Plus

8:ESP-0001-2018.

30.   Erbel PJ, Barr K, Gao N, Gerwig GJ, Rick PD, Gardner KH. 2003. Identification and

biosynthesis of cyclic enterobacterial common antigen in Escherichia coli. J Bacteriol

185:1995-2004.

31.   Leo V, Tran E, Morona R. 2020. Polysaccharide co-polymerase WzzB/WzzE chimeras

reveal transmembrane 2 region of WzzB is important for interaction with WzyB. J

Bacteriol 203:00598-20.

32.   Maczuga N, Tran ENH, Qin J, Morona R. 2022. Interdependence of *Shigella flexneri* O

Antigen and Enterobacterial Common Antigen Biosynthetic Pathways. J Bacteriol

204:e0054621.

33.   Nath P, Morona R. 2015. Detection of Wzy/Wzz interaction in *Shigella flexneri*.

Microbiology (Reading) 161:1797-1805.

34.   Weckener M, Woodward LS, Clarke BR, Liu H, Ward PN, Le Bas A, Bhella D, Whitfield

C, Naismith JH. 2023. The lipid linked oligosaccharide polymerase Wzy and its

regulating co-polymerase, Wzz, from enterobacterial common antigen biosynthesis form

a complex. Open Biology 13:220373.

35.   Tocilj A, Munger C, Proteau A, Morona R, Purins L, Ajamian E, Wagner J, Papadopoulos

M, Van Den Bosch L, Rubinstein JL, Féthière J, Matte A, Cygler M. 2008. Bacterial

polysaccharide co-polymerases share a common framework for control of polymer

length. Nat Struct Mol Biol 15:130-8.

36.   Kalynych S, Yao D, Magee J, Cygler M. 2012. Structural characterization of closely

related O-antigen lipopolysaccharide (LPS) chain length regulators. J Biol Chem

287:15696-705.

37.   Collins RF, Kargas V, Clarke BR, Siebert CA, Clare DK, Bond PJ, Whitfield C, Ford RC.

2017. Full-length, Oligomeric Structure of Wzz Determined by Cryoelectron Microscopy

Reveals Insights into Membrane-Bound States. Structure 25:806-815.e3.

38.   Kalynych S, Cherney M, Bostina M, Rouiller I, Cygler M. 2015. Quaternary structure of

WzzB and WzzE polysaccharide copolymerases. Protein Science 24:58-69.

39.   Wiseman B, Nitharwal RG, Widmalm G, Högbom M. 2021. Structure of a full-length

bacterial polysaccharide co-polymerase. Nat Commun 12:369.

40.   Wiseman B, Widmalm G, Högbom M. 2023. Alternating L4 loop architecture of the

bacterial polysaccharide co-polymerase WzzE. Commun Biol 6:802.

41.   Morona R, Purins L, Tocilj A, Matte A, Cygler M. 2009. Sequence–structure relationships

in polysaccharide co-polymerase (PCP) proteins. Trends in biochemical sciences 34:78-

84.

42. Ascari A, Tran ENH, Eijkelkamp BA, Morona R. 2022. Identification of the *Shigella*

flexneri Wzy Domain Modulating Wzz(pHS-2) Interaction and Detection of the

Wzy/Wzz/Oag Complex. J Bacteriol 204:e00224-22.

43. Leo V, Teh MY, Tran ENH, Morona R. 2021. Identification of a Region in *Shigella flexneri*

WzyB Disrupting the Interaction with Wzz(pHS2). J Bacteriol 203:e0041321.

44. Whitfield C, Williams DM, Kelly SD. 2020. Lipopolysaccharide O-antigens-bacterial

glycans made to measure. J Biol Chem 295:10593-10609.

45. Ascari A, Tran ENH, Eijkelkamp BA, Morona R. 2022. Identification of the *Shigella*

flexneri Wzy Domain Modulating Wzz$_{pHS-2}$ Interaction and Detection of the Wzy/Wzz/Oag

Complex. Journal of Bacteriology 0:e00224-22.

46. Huszczynski SM, Coumoundouros C, Pham P, Lam JS, Khursigara CM. 2019. Unique

regions of the polysaccharide copolymerase Wzz$_2$ from *Pseudomonas aeruginosa* are

essential for O-specific antigen chain length control. Journal of Bacteriology 201:e00165-

19.

47. Kalynych S, Ruan X, Valvano MA, Cygler M. 2011. Structure-guided investigation of

lipopolysaccharide O-antigen chain length regulators reveals regions critical for modal

length control. Journal of Bacteriology 193:3710-3721.

48. Rai AK, Carr JF, Bautista DE, Wang W, Mitchell AM. 2021. ElyC and cyclic

enterobacterial common antigen regulate synthesis of phosphoglyceride-linked

enterobacterial common antigen. mBio 12:e02846-21.

49. Mitchell AM, Srikumar T, Silhavy TJ. 2018. Cyclic enterobacterial common antigen

maintains the outer membrane permeability barrier of *Escherichia coli* in a manner

controlled by Yhdp. mBio 9:e01321-18.

50. Islam ST, Lam JS. 2014. Synthesis of bacterial polysaccharides via the Wzx/Wzy-

dependent pathway. Can J Microbiol 60:697-716.

51.   Kintz EN, Goldberg JB. 2011. Site-directed mutagenesis reveals key residue for O

antigen chain length regulation and protein stability in *Pseudomonas aeruginosa* Wzz2.

J Biol Chem 286:44277-44284.

52.   Papadopoulos M, Morona R. 2010. Mutagenesis and chemical cross-linking suggest that

Wzz dimer stability and oligomerization affect lipopolysaccharide O-antigen modal chain

length control. J Bacteriol 192:3385-93.

53.   Morona R, van den Bosch L, Manning PA. 1995. Molecular, genetic, and topological

characterization of O-antigen chain length regulation in *Shigella flexneri*. J Bacteriol

177:1059-68.

54.   Weckener M, Woodward LS, Clarke BR, Liu H, Ward PN, Le Bas A, Bhella D, Whitfield

C, Naismith JH. 2023. The lipid linked oligosaccharide polymerase Wzy and its

regulating co-polymerase, Wzz, from enterobacterial common antigen biosynthesis form

a complex. Open Biol 13:220373.

55.   Kalynych S, Valvano MA, Cygler M. 2012. Polysaccharide co-polymerases: the

enigmatic conductors of the O-antigen assembly orchestra. Protein Eng Des Sel 25:797-

802.

56.   Hong Y, Hu D, Verderosa Anthony D, Qin J, Totsika M, Reeves Peter R. 2023. Repeat-

Unit Elongations To Produce Bacterial Complex Long Polysaccharide Chains, an O-

Antigen Perspective. EcoSal Plus 11:eesp-0020-2022.

57.   Wiseman B, Widmalm G, Högbom M. 2023. Alternating L4 loop architecture of the

bacterial polysaccharide co-polymerase WzzE. Communications Biology 6:802.

58.   Lutz R, Bujard H. 1997. Independent and tight regulation of transcriptional units in

Escherichia coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. Nucleic

Acids Res 25:1203-10.

59.   Mendez-Perez D, Gunasekaran S, Orler VJ, Pfleger BF. 2012. A translation-coupling

693   DNA cassette for monitoring protein translation in *Escherichia coli*. Metabolic

694   Engineering 14:298-305.

695   60.   Sieber V. 2003. Selection for Soluble Proteins via Fusion with Chloramphenicol

696   Acetyltransferase, p 45-55. *In* Arnold FH, Georgiou G (ed), Directed Enzyme Evolution:

697   Screening and Selection Methods doi:10.1385/1-59259-396-8:45. Humana Press,

698   Totowa, NJ.

699   61.   Andreeva AE, Borissova BE, Mironova R, Glykos NM, Kotsifaki D, Ivanov I, Krysteva M,

700   Kokkinidis M. 2000. Crystallization of type I chloramphenicol acetyltransferase: an

701   approach based on the concept of ionic strength reducers. Acta Crystallogr D Biol

702   Crystallogr 56:101-3.

703   62.   Mitchell AM, Silhavy TJ. 2019. Envelope stress responses: balancing damage repair and

704   toxicity. Nature Reviews Microbiology 17:417-428.

705   63.   Kajimura J, Rahman A, Hsu J, Evans MR, Gardner KH, Rick PD. 2006. O acetylation of

706   the enterobacterial common antigen polysaccharide is catalyzed by the product of the

707   *yiaH* Gene of *Escherichia coli* K-12. Journal of Bacteriology 188:7542-7550.

708   64.   Daniels C, Morona R. 1999. Analysis of *Shigella flexneri* wzz (Rol) function by

709   mutagenesis and cross-linking: wzz is able to oligomerize. Mol Microbiol 34:181-94.

710   65.   Teese MG, Langosch D. 2015. Role of GxxxG Motifs in Transmembrane Domain

711   Interactions. Biochemistry 54:5125-35.

712   66.   Prakash A, Janosi L, Doxastakis M. 2011. GxxxG motifs, phenylalanine, and cholesterol

713   guide the self-association of transmembrane domains of ErbB2 receptors. Biophys J

714   101:1949-58.

715   67.   Mueller BK, Subramaniam S, Senes A. 2014. A frequent, GxxxG-mediated,

716   transmembrane association motif is optimized for the formation of interhelical Cα–H

717   hydrogen bonds. Proceedings of the National Academy of Sciences 111:E888-E895.

68.   Papadopoulos M, Tran ENH, Murray GL, Morona R. 2016. Conserved transmembrane

glycine residues in the *Shigella flexneri* polysaccharide co-polymerase protein WzzB

influence protein-protein interactions. Microbiology (Reading) 162:921-929.

69.   Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L,

Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C, O'Neill M, Reiman D,

Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O, Bridgland A,

Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB,

Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S,

Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A, Bapst

V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of

biomolecular interactions with AlphaFold 3. Nature 630:493-500.

70.   Parrish HL, Glassman CR, Keenen MM, Deshpande NR, Bronnimann MP, Kuhns MS.

2015. A Transmembrane Domain GGxxG Motif in CD4 Contributes to Its Lck-

Independent Function but Does Not Mediate CD4 Dimerization. PLOS ONE

10:e0132333.

71.   Faingold O, Cohen T, Shai Y. 2012. A GxxxG-like Motif within HIV-1 Fusion Peptide Is

Critical to Its Immunosuppressant Activity, Structure, and Interaction with the

Transmembrane Domain of the T-cell Receptor*. Journal of Biological Chemistry

287:33503-33511.

72.   Hong M, Payne SM. 1997. Effect of mutations in *Shigella flexneri* chromosomal and

plasmid-encoded lipopolysaccharide genes on invasion and serum resistance. Mol

Microbiol 24:779-91.

73.   Tran ENH, Morona R. 2013. Residues located inside the *Escherichia coli* FepE protein

oligomer are essential for lipopolysaccharide O-antigen modal chain length regulation.

Microbiology (Reading) 159:701-714.

74.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M,

Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene

knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

75.   Silhavy TJ, Berman ML, Enquist LW. 1984. Experiments with gene fusions. (No Title).

76.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in

Escherichia coli K-12 using PCR products. Proceedings of the National Academy of

Sciences 97:6640-6645.

77.   Pyne ME, Moo-Young M, Chung DA, Chou CP. 2015. Coupling the CRISPR/Cas9

System with Lambda Red Recombineering Enables Simplified Chromosomal Gene

Replacement in *Escherichia coli*. Appl Environ Microbiol 81:5103-14.

78.   Jiang W, Bikard D, Cox D, Zhang F, Marraffini LA. 2013. RNA-guided editing of bacterial

genomes using CRISPR-Cas systems. Nature biotechnology 31:233-239.

79.   Nichols BP, Shafiq O, Meiners V. 1998. Sequence analysis of Tn10 insertion sites in a

collection of *Escherichia coli* strains used for genetic mapping and strain construction. J

Bacteriol 180:6408-11.

80.   Banerjee RV, Johnston NL, Sobeski JK, Datta P, Matthews RG. 1989. Cloning and

sequence analysis of the *Escherichia coli* metH gene encoding cobalamin-dependent

methionine synthase and isolation of a tryptic fragment containing the cobalamin-binding

domain. J Biol Chem 264:13888-95.

81.   Guzman LM, Belin D, Carson MJ, Beckwith J. 1995. Tight regulation, modulation, and

high-level expression by vectors containing the arabinose P_{BAD} promoter. J Bacteriol

177:4121-30.

82.   Morris KN, Mitchell AM. 2023. Phosphatidylglycerol Is the Lipid Donor for Synthesis of

Phospholipid-Linked Enterobacterial Common Antigen. Journal of Bacteriology

205:e00403-22.

Figures and Legends

Figure 1: Plasmid-based expression of *wzzE* partially complements a chromosomal *wzzE*

deletion. (A) Diagrams of constructs for expression of wild-type WzzE, WzzE with a C-terminal

glycine-serine linker and triple FLAG tag, WzzE translationally fused with chloramphenicol

acetyltransferase (Cat), and WzzE fused with Cat and triple FLAG are shown. The indicated

constructs were expressed from the low copy number pZS21 plasmid (58). **(B)** The vancomycin

sensitivity of a Δ*yhdP* strain can be suppressed by loss of WzzE (12). When the Δ*wzzE* Δ*yhdP*

strain carried the pZS21 empty plasmid, it remained resistance to vancomycin as assayed by

EOP but became sensitive when *wzzE, wzzE*-FLAG, or *wzzE-cat* was expressed, indicating all

three constructs complemented Δ*wzzE*. In addition, translational fusion of *cat* and *wzzE*

conferred chloramphenicol resistance, suggesting this construct could readout WzzE stability

and multimerization based on chloramphenicol resistance. **(C)** WzzE-FLAG was detectable by

immunoblotting. GroEL serves as a loading control. **(D)** ECA chain-length regulation was

assayed through immunoblotting using an ECA antibody. Plasmid-base expression of WzzE

restored chain-length regulation of linear ECA, although lower levels of ECA and shorter chain-

length were observed than in the wild-type strain for the p*wzzE* and p*wzzE* constructs, while the

p*wzzE*-FLAG and p*wzzE-cat*-FLAG constructs complemented better. The Δ*wecA* strain

produces no ECA and serves as negative control. X: nonspecific band. **(E-F)** The strain carrying

a plasmid with wild-type *wzzE* was grown with ^{15}N to label ECA and then combined with an

unlabeled strain carrying either wild-type *wzzE* (E) or the empty vector (F). ECA$_{CYC}$ was purified

and then detected by LC-MS. MS traces are shown for the indicated samples. The mass,

charge, and identity of the ECA$_{CYC}$-associated peaks are shown. Expression of *wzzE* from

p*wzzE* restored ECA$_{CYC}$ synthesis in the Δ*wzzE* strain. The strains contained Δ*wecH* to prevent

non-stochiometric acetylation of ECA. All images are representative of at least three

independent experiments.

Figure 2: Alteration of WzzE TMH2 can prevent ECA chain length regulation without

affecting WzzE stability. (A) The effect of the indicated *wzzE* site-directed mutations on WzzE-

FLAG protein levels was assessed using immunoblotting. GroEL serves as a loading control.

(B) Densitometry was used for quantification of the immunoblots in (A). Levels of WzzE-FLAG

compared to GroEL are shown normalized to the wild-type *wzzE* complemented strain. All

variants show similar protein levels to wild-type WzzE. Data are shown as the average of three

biological replicates ± SEM and individual datapoints. * $p < 0.05$ by Mann-Whitney T-test. **(C)**

Strains carrying the indicated plasmids were assayed for their chloramphenicol resistance by

EOP as a proxy for WzzE stability and multimerization ability. All *wzzE* mutants retained full

chloramphenicol resistance, suggesting WzzE stability and multimerization. **(D)** Chain-length

regulation of the linear ECA forms by the indicated WzzE variants was assayed by

809   immunoblotting. Short and long exposures are shown. "X" indicates non-specific bands.

810   WzzE$^{GG333/4LL}$ showed loss of chain-length regulation while the other variants retained wild-type

811   function. The Δ*wecA* strain serves as negative control. All images are representative of at least

812   three independent experiments.

Figure 3. Conserved glycines in WzzE TMH 2 are required for ECA$_{CYC}$ synthesis. (A) The
strain carrying a plasmid with wild-type *wzzE* was grown with ^{15}N to label ECA and then
combined with an unlabeled strain carrying the indicated *wzzE* mutant or control. ECA$_{CYC}$ was
purified and the relative amount of unlabeled to labeled ECA$_{CYC}$ was determined by LC-MS.
WzzE$^{GG333/4LL}$ showed a loss of the ability to produce ECA$_{CYC}$. All plasmids were expressed in a
ΔwzzE ΔwecH strain background to prevent non-stochiometric acetylation of ECA. Data are
shown as the average of three biological replicates ± SEM and as individual datapoints. * p <
0.05 by Mann-Whitney T-test. **(B)** The predicted structure of a WzzE octamer with WzyE was
calculated using AlphaFold 3 (69). The structures of wild-type WzzE and WzzE$^{GG333/4LL}$
monomers were aligned with the mutated residues shown in red. The insets show the predicted
distance between the two WzzE TMH in the wild type WzzE and WzzE$^{GG333/4LL}$.

Figure 4: The periplasmic region of WzzE contributes to WzzE stability and chain length

regulation. (A) Immunoblotting was used to determine protein levels for the indicated WzzE-

FLAG variants. GroEL serves as a loading control. **(B)** WzzE-FLAG levels from (A) were

quantified by densitometry. WzzE-FLAG levels compared to GroEL are shown normalized to the

wild-type *wzzE* complemented strain. All variants are similar to wild type WzzE except for

WzzE$^{\Delta F104}$ which shows significantly reduced protein levels. Data are shown as the average of

three biological replicates ± SEM and individual datapoints. * p <0.05 by Mann-Whitney T-test.

(C) The indicated strains were assayed for WzzE-Cat-FLAG stability as measured by

chloramphenicol resistance. All *wzzE* point mutations showed chloramphenicol resistance

indicating stability and multimerization, with the exception of WzzE$^{\Delta F104}$ where the tags are

cleaved (**Figure S3C**). **(D)** Linear ECA chain-length regulation by WzzE variants was assayed

by immunoblotting with ECA antibody. Long and short exposures are shown. The periplasmic

point mutants have variable effects on chain length regulation as well as ECA$_{CYC}$ biogenesis.

841   The WzzE$^{\Delta F104}$ and WzzE^{K162A} variants showed loss of ECA chain-length regulation while the

842   other variants had wild-type chain-length regulation. The Δ*wecA* strain serves as negative

843   control. All images are representative of at least three independent experiments.

Figure 5: The periplasmic domain of WzzE contributes to ECA_{CYC} synthesis. (A) The strain

carrying wild-type *wzzE* was grown with ^{15}N to label ECA and then combined with an unlabeled

strain carrying the indicated *wzzE* mutant or control strain. After ECA_{CYC} purification, the relative

amount of unlabeled to labeled ECA_{CYC} was determined by LC-MS. $WzzE^{\Delta F104}$ and $WzzE^{K162A}$

had little to no ECA_{CYC} production while $WzzE^{F104H}$ showed an variable phenotype. All strains

included $\Delta wecH$ to prevent non-stochiometric acetylation of ECA. Data are shown as the

average of three biological replicates ± SEM and as individual datapoints. * $p < 0.05$ by Mann-

Whitney T-test. **(B)** Structures predicted as in **Figure 3** for wild-type WzzE and $WzzE^{F104\Delta}$ were

aligned. The mutated residue is shown in red on a WzzE monomer (B). The inset shows the

interaction between alpha helix 2 and an adjoining loop in wild-type WzzE.

Figure 6: Levels of ECA$_{CYC}$ production and chain-length regulation are genetically
859

separable. CRISPR-Cas9 was used to make the indicated mutations to *wzzE* in its native locus.
860

All strains contain *metE*::Tn*10*. **(A)** The strain wild type for *wzzE* was grown with ^{15}N to label
861

ECA and then combined with an unlabeled strain with the indicated *wzzE* mutant or control
862

strain. After ECA$_{CYC}$ purification, the ratio of unlabeled to labeled ECA$_{CYC}$ was determined by
863

LC-MS. All variants other than WzzE^{F104Y} produced significantly reduced ECA$_{CYC}$ levels. **(B)**
864

Immunoblotting for ECA was conducted on the indicated chromosomal mutants. A short and
865

long exposure are shown. The WzzE$^{\Delta F104}$ and WzzE$^{GG333/4LL}$ variants resulted in ECA chain
866

length dysregulation, while the other mutations more closely resembled wild-type WzzE chain

length regulation. The Δ*wecA* strain serves as negative control. **(C)** Immunoblotting was used to

assay ECA chain-length regulation in the indicated strains. Deletion of *wzzB* and/or *fepE* in

combination with Δ*wzzE* did not cause the increased levels of linear ECA observed with the

loss-of-function point mutants; however, the Δ*wzzE* Δ*wzzB* Δ*fepE* strain did have a small

increase in linear ECA levels compared to Δ*wzzE* alone.