The basolateral amygdala complex and perirhinal cortex represent focal and peripheral

states of information processing in rats.

Francesca S. Wong, A. Simon Killcross, R. Fred Westbrook & Nathan M. Holmes[*]

School of Psychology, University of New South Wales, Sydney, Australia

* For correspondence: n.holmes@unsw.edu.au

Competing interests: The authors declare that no competing interests exist.

Funding: This work was supported by an Australian Government Research Training

Fellowship to FSW, an Australian Research Council (ARC) Discovery Project Grant to RFW

(DP2201036501) and an ARC Future Fellowship to NMH (FT190100697).

Abstract. This study used variations of a sensory preconditioning protocol in male and

female rats to test a theory that the basolateral amygdala complex (BLA) and perirhinal

cortex (PRh) represent focal and peripheral states of attention, respectively. It specifically

tested predictions derived from the theory regarding when learning about a stimulus that

signals danger will be disrupted by BLA or PRh infusions of the *N-methyl-D-aspartate*

receptor (NMDAR) antagonist, DAP5. Consistent with the theory, the effects of these

infusions depended on the novelty/familiarity of the conditioned stimulus as well as the

manner in which it was paired with foot shock. When a stimulus was novel, its conditioning

required activation of NMDAR in the BLA and not the PRh (Experiments 2A and 2B)

regardless of whether the stimulus-shock pairings were contiguous or separated in time.

When a pre-exposed and, thereby, familiar stimulus was presented contiguously with shock,

its conditioning again required activation of NMDAR in the BLA and not the PRh

(Experiments 1A, 1B, 3A and 3B). However, when a pre-exposed stimulus was indirectly

paired with shock - because it was associatively activated at the time of shock or separated

from the shock by another stimulus - its conditioning required activation of NMDAR in the

PRh and not the BLA (Experiments 1A, 1B, 3A and 3B). These findings are discussed in

relation to theories of information processing that distinguish between focal and peripheral

states of attention/memory, and past studies that have examined the substrates of learning and

memory in the PRh and BLA.

Introduction

Sensory preconditioning is a laboratory protocol that can be used to examine how the

mammalian brain processes sensory and emotional information. It typically involves three

stages. In stage 1, rats are exposed to pairings of two relatively harmless stimuli, S2 and S1

(e.g., a sound and light). In stage 2, rats are exposed to pairings of one of these stimuli, the S1

(e.g., the light), with foot shock. Finally, in stage 3, rats display fear or defensive responses

(e.g., freezing) when tested with the S1 that had been paired with shock and the S2 that had

never been paired with shock. Importantly, the fear of S2 is not due to generalization from S1

or any innately aversive properties of the S2, as it is absent among controls exposed to

explicitly unpaired presentations of the relevant stimuli in either stage 1 or stage 2 (Holmes et

al., 2013; Holmes & Westbrook, 2017; Kikas et al., 2019; Michalscheck et al., 2021; Parkes

& Westbrook, 2010; Rizley & Rescorla, 1972; Wong et al., 2019, 2025). Instead, where it is

observed, fear of the S2 reflects integration of the sensory and emotional information

encoded in stages 1 and 2. That is, the S2-S1 (sensory) association formed in stage 1 is

integrated with the S1-shock (emotional) association formed in stage 2 to drive fear responses

during presentations of the S2 alone in stage 3.

Over the past 10 years, our laboratory has examined the neural substrates of the

different types of information that rats encode in this sensory preconditioning protocol. Our

work has focused on two regions of the medial temporal lobe - the perirhinal cortex (PRh)

and basolateral amygdala complex (BLA) - and revealed that the involvement of these two

regions can be quite distinct. For example, the S2-S1 association that forms in stage 1 is

encoded through activation of *N-methyl-D-aspartate* receptors (NMDARs) in the PRh, not

the BLA; whereas the S1-shock association that forms in stage 2 is encoded through

activation of NMDARs in the BLA, not the PRh (Campeau et al., 1992; Fanselow & Kim,

1994; Miserendino et al., 1990; Holmes et al., 2013, 2018; Rodrigues et al., 2001; see also

Bang & Brown, 2008; Phillips & LeDoux, 1995; Romanski & LeDoux, 1992a, 1992b;

Wilensky et al., 2006). In a recent pair of studies, we extended these findings in two ways.

First, we showed that S1 does not just form an association with shock in stage 2; it also

mediates an association between S2 and the shock. Thus, S2 enters testing in stage 3 already

conditioned, able to elicit fear responses (Wong et al., 2019). Second, we showed that this

mediated S2-shock association requires NMDAR-activation in the PRh, as well as

communication between the PRh and BLA (Wong et al., 2025). These findings raise two

critical questions: 1) *why* is the PRh engaged for mediated conditioning of S2 but not for

direct conditioning of S1; and 2) more generally, *what* determines whether the BLA and/or

PRh is engaged for conditioning of the S1 and/or S2?

One explanation for the different substrates of conditioning to S2 and S1 in our

sensory preconditioning protocol is that the two stimuli access different states of

attention/memory which are differentially supported by the BLA and PRh. That is, during

stage 2 of our protocol, the BLA may function like a focal state of attention in processing

information about the presented S1 and foot shock. By contrast, because the PRh encodes the

S2-S1 pre-exposures in stage 1, this region may function like a peripheral state of attention in

processing information about the S2 that is retrieved from long-term memory. Such a

characterization explains why the direct S1-shock association requires activation of

NMDARs in the BLA but not the PRh (S1 and shock co-occur in the focus of attention); and

the mediated S2-shock association requires activation of NMDARs in the PRh as well as

communication between the PRh and BLA (S2 conditions in the periphery of attention as the

shock occurs in its focus). However, it also implies that the substrates of conditioning to S2

will depend on how the S2-S1 and S1-shock pairings are presented. For example, different

types of arrangements may influence the substrates of conditioning to S2 by influencing its

novelty and/or its predictive value at the time of the shock, on the supposition that familiar

stimuli are processed in the periphery of attention and, thereby, the PRh (Bogacz & Brown,

2003; Brown & Banks, 2015; Brown & Bashir, 2002; Martin et al., 2013; McClelland et al.,

2014; Morillas et al., 2017; Murray & Wise, 2012; Robinson et al., 2010; Suzuki & Naya,

2014; Voss et al., 2009; Yang et al., 2023) whereas novel stimuli are processed in the focus

of attention and, thereby, the amygdala (Holmes et al., 2018; Qureshi et al., 2023;

Roozendaal et al., 2006; Rutishauser et al., 2006; Schomaker & Meeter, 2015; Wright et al.,

2003). This can be achieved by presenting the S2-S1 and S1-shock pairings together as S2-

S1-shock sequences. Under such circumstances, conditioning to the S2 should cease to occur

in the periphery of attention as it is novel and no longer requires retrieval of information from

long-term memory. Instead, this conditioning should occur in the focus of attention and,

thereby require activation of NMDARs in the BLA and not the PRh (e.g., Williams-Spooner

et al., 2022).

Different types of arrangements may also influence the substrates of conditioning to

S2 as a function of its "distance" or temporal separation from the shock. The supposition here

is that shock/danger and its immediate antecedents are processed in the focus of attention

(and, thereby, the amygdala), regardless of whether the antecedents are novel or familiar;

whereas stimuli that are temporally separated from the shock *can be* processed in the

periphery of attention (and, thereby, the PRh) and *are more likely to be* processed there if

they are familiar. This can be assessed by pre-exposing rats to S2-S1 pairings prior to

conditioning with S2-S1-shock sequences. Under these circumstances, conditioning to the

familiar S1 that is contiguous with shock should occur in the focus of attention and, thereby,

require activation of NMDARs in the BLA and not the PRh; whereas conditioning to the familiar S2 that is separated from the shock (by S1) should occur in the periphery of attention and, thereby, require activation of NMDARs in the PRh and not the BLA.

The present series of experiments used these variations of a sensory preconditioning protocol to test the implications of our theory that the BLA and PRh represent focal and peripheral states of attention, respectively. The first implication is that, in our standard sensory preconditioning protocol (S2-S1 pairings in stage 1, S1-shock pairings in stage 2), the substrates of conditioning to S2 and S1 will be doubly dissociable at the level of NMDARs in the PRh and BLA. That is, during the session of S1-shock pairings in stage 2, mediated conditioning of the absent S2 should occur in the periphery of attention and hence, require NMDAR-activation in the PRh but *not* the BLA, whereas direct conditioning of the presented S1 should occur in the focus of attention and hence, require NMDAR-activation in the BLA but not the PRh. This was supported by the initial experiments which found that a stage 2 infusion of the NMDAR antagonist, DAP5 into the PRh disrupts mediated conditioning of S2 but spares direct conditioning of S1 (Experiment 1A); whereas a stage 2 infusion of DAP5 into the BLA disrupts direct conditioning of S1 but spares mediated conditioning of S2 (Experiment 1B). Subsequent experiments then used variations of this protocol to examine whether the engagement of NMDAR in the PRh or BLA for Pavlovian fear conditioning is influenced by the novelty/predictive value of the stimuli at the time of the shock (second implication of theory) as well as their distance or separation from the shock (third implication of theory; Table 1). Following the arguments above, we predicted that:

  1) When S2 or S1 is novel and paired with shock, its conditioning will be disrupted by a DAP5 infusion into the BLA but unaffected by a DAP5 infusion into the PRh (Experiments 2A and 2B);

2) When S2 or S1 is familiar and directly paired (i.e., co-occurs) with shock, its

   conditioning will again be disrupted by a DAP5 infusion into the BLA but not the PRh

   (Experiments 1A, 1B, 3A and 3B);

3) However, when S2 or S1 is familiar and indirectly paired with shock - because it is

   associatively activated at the time of shock or separated from the shock by another

   stimulus - its conditioning will be unaffected by a DAP5 infusion into the BLA but

   disrupted by a DAP5 infusion into the PRh (Experiments 1A, 1B, 3A and 3B).

Results

Experiments 1A and 1B – Encoding of the mediated S2-shock association requires NMDAR-activation in the PRh but not the BLA; Encoding of the direct S1-shock association requires NMDAR-activation in the BLA but not the PRh

We propose that the BLA and PRh represent focal and peripheral states of attention/memory, respectively. The first implication of this proposal is that, during stage 2 of our standard sensory preconditioning protocol, the substrates of conditioning to the absent S2 and presented S1 will be doubly dissociable at the level of NMDARs in the PRh and BLA: mediated conditioning of the absent S2 should occur in the periphery of attention and hence, require NMDAR-activation in the PRh but *not* the BLA, whereas direct conditioning of the presented S1 should occur in the focus of attention and hence, require NMDAR-activation in the BLA but not the PRh. Experiments 1A and 1B tested this implication. The design of these experiments is shown in Figure 1A. The two experiments differed only in that we assessed the involvement of the PRh in Experiment 1A and the BLA in Experiment 1B. In each experiment, two groups of rats were exposed to a session of S2-S1 pairings in stage 1 and, 24 h later, a session of S1-shock pairings in stage 2. Immediately prior to the conditioning

session in stage 2, rats in each experiment received an infusion of DAP5 or vehicle only into

either the PRh (Experiment 1A) or BLA (Experiment 1B). Finally, all rats were tested with

presentations of the S2 alone and S1 alone in stage 3. Based on our past findings and the

reasoning given above, we expected that mediated conditioning of the S2 would require

activation of NMDARs in the PRh (Wong et al., 2025) but not the BLA, whereas the direct

conditioning of the S1 would require activation of NMDARs in the BLA but not the PRh

(e.g., Rodrigues et al., 2001; Williams-Spooner et al., 2022). Hence, we anticipated that the

stage 2 infusion of DAP5 into the PRh would disrupt the test level of freezing to S2 without

affecting the test level of freezing to S1; and that the stage 2 infusion of DAP5 into the BLA

would disrupt the test level of freezing to S1 without affecting the test level of freezing to S2.

Experiment 1A – NMDAR-activation in the PRh is required for acquisition of the mediated

S2-shock association but not the direct S1-shock association

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,19)}$ = 2.876; p = .106).

Conditioning of the S1 was successful. Freezing to S1 increased across the four S1-shock

pairings in stage 2 ($F_{(1,19)}$ = 74.708; p < .001; n_p^2 = .797; 95% CI: [1.658, 2.717]). The rate of

this increase did not differ between the two groups ($F_{(1,19)}$ = .008; p = .930), and there was no

significant between-group differences in overall freezing to the S1 ($F_{(1,19)}$ = 1.803; p = .195).

Test. Figure 1B shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) and S1 alone (right panel) during testing in Experiment 1A, averaged across the eight

presentations of each stimulus. Both groups showed equivalent levels of freezing to the

directly conditioned S1 ($F_{(1,19)}$ = 1.423; p = .248), indicating that the PRh infusion of DAP5

had no effect on this conditioning. However, rats that received a PRh infusion of DAP5

(Group DAP5) in stage 2 exhibited significantly less freezing to the preconditioned S2 than

rats that received a PRh infusion of vehicle only (Group VEH; $F_{(1,19)} = 7.136$; $p = .015$; $n_p^2 =$

.273; 95% CI: [0.176, 1.447]).

Experiment 1B – NMDAR-activation in the BLA is required for acquisition of the direct

S1-shock association but not the mediated S2-shock association

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,26)} = 1.590$; $p = .219$).

Conditioning of the S1 was successful. Freezing to S1 increased across the four S1-shock

pairings in stage 2 ($F_{(1,26)} = 104.494$; $p < .001$; $n_p^2 = .801$; 95% CI: [1.633, 2.456]). The rate

of this increase did not differ between groups ($F_{(1,26)} = .259$; $p = .615$), however there was a

significant between-group difference in overall freezing to the S1 ($F_{(1,26)} = 6.068$; $p = .021$;

$n_p^2 = .189$; 95% CI: [0.094, 1.046]), indicating that the BLA infusion of DAP5 disrupted

conditioning to the S1.

Test. Figure 1D shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) and S1 alone (right panel) during testing in Experiment 1B, averaged across the eight

presentations of each stimulus. Both groups showed equivalent levels of freezing to the

preconditioned S2 ($F_{(1,26)} = .013$; $p = .910$). However, rats that received a BLA infusion of

DAP5 (Group DAP5) in stage 2 exhibited significantly less freezing to the directly

conditioned S1 than rats that received a BLA infusion of vehicle only (Group VEH; $F_{(1,26)} =$

4.775; $p = .038$; $n_p^2 = .155$; 95% CI: [0.040, 1.308]).

Experiment 1A replicated previous findings that a stage 2 infusion of DAP5 into the

PRh disrupts the acquisition of sensory preconditioned freezing to S2 without affecting the

acquisition of freezing to S1 (Wong et al., 2025). By contrast, Experiment 1B showed that a

BLA infusion of DAP5 disrupts the acquisition of freezing to S1 (replication) without

affecting the acquisition of sensory preconditioned freezing to S2 (new result). Taken

together, these findings indicate that the involvement of NMDARs in direct and mediated

conditioning is doubly dissociable at the levels of the BLA and PRh. The direct conditioning

of a stimulus that co-terminates with shock requires activation of NMDARs in the BLA but

does not require activation of NMDARs in the PRh. Conversely, the mediated conditioning

of a stimulus representation that is retrieved from long-term memory (and, hence, indirectly

paired with shock) requires activation of NMDARs in the PRh but occurs independently of

NMDAR-activation in the BLA. These findings are exactly consistent with the predictions

derived from our proposal that the BLA and PRh represent focal and peripheral states of

attention/memory, respectively. The BLA represents a focal state and, thereby, encodes the

directly conditioned S1-shock association, whereas the PRh represents a peripheral state and,

thereby, encodes the indirect or mediated S2-shock association.

Experiments 2A and 2B – The roles of the PRh and BLA in Pavlovian fear conditioning

when the S2-S1 and S1-shock pairings are combined into an S2-S1-shock sequence.

A second implication of our proposal is that the engagement of NMDAR in the PRh

for conditioning to S2 will be influenced by its novelty/familiarity at the time of the shock.

We addressed this implication by combining the S2-S1 and S1-shock pairings into S2-S1-

shock sequences that were administered in a single training session (Experiments 2A and

2B). The consequence of doing so is that, in contrast to the previous experiments in which the

S2 and S1 were both familiar at the time of their conditioning with foot shock, here the S2

and S1 were both novel at the time of their conditioning with foot shock. Immediately prior

to the session containing the S2-S1-shock sequences, rats received an infusion of DAP5 or

vehicle only into either the PRh (Experiment 2A) or BLA (Experiment 2B). Finally, all rats

were tested with presentations of the S2 alone and S1 alone in stage 3. We hypothesized that,

because S2 was novel and presented in sequence with S1 and shock in stage 2, it would be

processed in a focal state of attention rather than the periphery. Hence, whereas conditioning

to a representation of the pre-exposed S2 in the previous experiment required NMDAR-

activation in the PRh but not the BLA, we predicted that conditioning to the novel S2 in this

experiment would require NMDAR-activation in the BLA but not the PRh. Specifically, we

expected that the stage 2 infusion of DAP5 into the PRh would have no effect on the test

level of freezing to either S2 or S1; and that the stage 2 infusion of DAP5 into the BLA

would disrupt the test level of freezing to both S2 and S1.

Experiment 2A – When the S2-S1 and S1-shock pairings are combined into S2-S1-shock

sequences, the PRh is no longer engaged for conditioning of the S2

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,15)} = 3.026$; $p = .102$).

Conditioning was successful. Freezing to the S2 and S1 increased linearly across the four S2-

S1-shock sequences in stage 2 (smaller $F_{(1,15)} = 47.193$; $p < .001$; $n_p^2 = .759$; 95% CI: [1.453,

2.760]). The linear x Group interaction was significant for freezing to the S2 ($F_{(1,15)} = 5.117$;

$p < .039$; $n_p^2 = .254$; 95% CI: [0.056, 1.868]), but not for freezing to the S1 ($F_{(1,15)} = .009$; $p =$

.926). This suggests that the PRh infusion of DAP5 slowed acquisition of freezing to the S2

without affecting freezing to the S1. However, there were no significant between-group

differences in overall freezing to the S2 ($F_{(1,15)} = 4.125$; $p = .06$) or S1 ($F_{(1,15)} = 2.726$; $p =$

.120).

Test. Figure 2B shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) and S1 alone (right panel) during drug-free testing in Experiment 2A, averaged across

the eight presentations of each stimulus. Both groups showed equivalent levels of freezing to

the S2 ($F_{(1,15)} = 1.292$; $p = .274$) and S1 ($F_{(1,15)} = .838$; $p = .374$).

Experiment 2B – When the S2-S1 and S1-shock pairings are combined into S2-S1-shock

sequences, the BLA is engaged for conditioning of the S2

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,15)}$ = .576; p = .460).

Conditioning was again successful. Freezing to the S2 and S1 (smaller $F_{(1,15)}$ = 62.211; $p <$

.001; n_p^2 = .806; 95% CI: [1.093, 1.902]) increased linearly across the four S2-S1-shock

sequences in stage 2. The linear x Group interaction was significant for freezing to the S2

($F_{(1,15)}$ = 9.477; p = .008; n_p^2 = .387; 95% CI: [0.337, 1.852]) and S1 ($F_{(1,15)}$ = 12.771; p =

.003; n_p^2 = .460; 95% CI: [0.548, 2.167]), suggesting that the BLA infusion of DAP5 slowed

acquisition of freezing to both stimuli. However, there were no significant between-group

differences in overall freezing to the S2 ($F_{(1,15)}$ = 2.999; p = .104) or S1 ($F_{(1,15)}$ = 3.747; p =

.072).

Test. Figure 2D shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) and S1 alone (right panel) during drug-free testing in Experiment 2B, averaged across

the eight presentations of each stimulus. Rats that received a BLA infusion of DAP5 (Group

DAP5) in stage 2 exhibited significantly less freezing to both the S2 ($F_{(1,15)}$ = 8.660; p = .010;

n_p^2 = .366; 95% CI: [0.349, 2.182]) and S1 ($F_{(1,15)}$ = 12.089; p = .003; n_p^2 = .446; 95% CI:

[0.507, 2.114]) compared to the vehicle-infused controls.

Experiments 2A and 2B showed that combining the S2-S1 and S1-shock pairings into

S2-S1-shock sequences altered the roles of the PRh and BLA in conditioning to the S2. In the

previous experiments, the S2-S1 (stage 1) and S1-shock (stage 2) pairings were separated by

24 hours and acquisition of freezing to S2 required NMDAR-activation in the PRh but not the

BLA. Here, the acquisition of freezing to S2 (and S1) across S2-S1-shock sequences required

NMDAR-activation in the BLA but not the PRh. These findings are consistent with our

proposal that the BLA and PRh function like different states of attention/memory. The BLA

represents a focal state and, thus, processes information about novel stimuli that are presented

in a sequence that co-terminates with shock: hence it was engaged for conditioning to both

the S2 and S1 in the current experiment. By contrast, the PRh represents a peripheral state

and maintains the trace or representation of a pre-exposed stimulus: hence, it was not

engaged for conditioning to either the S2 or S1 in the current experiment.

Experiments 3A and 3B – The roles of the PRh and BLA in Pavlovian fear conditioning

when rats are pre-exposed to S2-S1 pairings prior to conditioning with S2-S1-shock

sequences.

A third implication of our proposal is that pre-exposure can alter the way that stimuli

are processed in the focus/periphery of attention and, thereby, the substrates of their

conditioning in the PRh and BLA. The additional implication is based on decades of work

showing that pre-exposure to a stimulus can reduce its rate of conditioning (Holmes & Harris,

2009, 2010; Kennedy et al., 2021; Leung et al., 2013a, 2013b; Lubow, 1973; Lubow &

Moore, 1959) while increasing the context dependence of any responding that develops

(Holmes & Westbrook, 2014; Rosenberg et al., 2011; Westbrook et al., 2000). We addressed

this implication by pre-exposing rats to S2-S1 pairings in stage 1 followed by conditioning

with S2-S1-shock sequences in stage 2. We specifically examined whether pre-exposing rats

to S2-S1 pairings reduces processing of the S2 in the focus of attention when it is presented

as part of S2-S1-shock sequences in stage 2: i.e., whether it reduces the ability of S2 to access

the focus of attention (unlikely given the S1 results in Experiments 1A and 1B) and/or

accelerates the rate at which S2 transitions to the periphery when it is replaced by

presentation of the S1. If so, the substrates of conditioning to S2 in the current experiments

should differ from the substrates of conditioning to S2 in Experiments 2A and 2B. That is,

whereas conditioning of the novel S2 in Experiments 2A and 2B required NMDAR-

activation in the BLA and not the PRh, conditioning of the pre-exposed S2 in the current

experiments should require NMDAR-activation in the PRh and not the BLA.

The design of Experiments 3A and 3B is shown in Figure 3. Briefly, rats were

exposed to a session of S2-S1 pairings in stage 1 and, 24 h later, a session of S2-S1-shock

sequences in stage 2. Immediately prior to the latter session, rats received an infusion of

DAP5 or vehicle only into either the PRh (Experiment 3A) or BLA (Experiment 3B). Finally,

all rats were tested with presentations of the S2 alone and S1 alone in stage 3. We

hypothesized that pre-exposure to S2-S1 pairings prior to conditioning with S2-S1-shock

sequences would *not* affect the substrates of directly conditioned fear to S1: that is, just as the

direct conditioning of a pre-exposed S1 in Experiments 1A and 1B required NMDAR-

activation in the BLA but not the PRh, here again we expected that direct conditioning of the

pre-exposed S1 would require NMDAR-activation in the BLA but not the PRh. Such a result

would suggest that, with the level of pre-exposure used here, the S1 accesses the focus of

attention: hence its conditioning requires activation of NMDAR in the BLA and not the PRh.

By contrast, we hypothesized that pre-exposure to S2-S1 pairings prior to conditioning with

S2-S1-shock sequences would alter the substrates of conditioning to S2: that is, whereas

conditioning to the novel S2 in the previous experiments required NMDAR-activation in the

BLA but not the PRh, here we expected that conditioning of the pre-exposed S2 would

require NMDAR-activation in the PRh but not the BLA. Such a result would suggest that a

pre-exposed S2 more rapidly transitions from the focus of attention to its periphery when it is

replaced by S1: hence its conditioning ceases to require activation of NMDAR in the BLA

and now requires activation of NMDAR in the PRh.

Experiment 3A – When rats are pre-exposed to S2-S1 pairings prior to conditioning with

S2-S1-shock sequences, the PRh is engaged for acquisition of freezing to S2 but not S1

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,22)} = 3.143$; $p = .090$).

Conditioning was successful. Freezing to the S2 and S1 increased across the four S2-S1-

shock sequences in stage 2 (smaller $F_{(1,22)}$ = 110.127; p < .001; n_p^2 = .833; 95% CI: [1.877,

2.802]). The rate of this increase did not differ between groups (largest $F_{(1,22)}$ = 1.397; p =

.250), and there were no significant between-group differences in overall freezing to the S2

and S1 (largest $F_{(1,22)}$ = .620; p = .439).

Test. Figure 3B shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) and S1 alone (right panel) during testing in Experiment 3A, averaged across the eight

presentations of each stimulus. Both groups showed equivalent levels of freezing to the

directly conditioned S1 ($F_{(1,22)}$ = .226; p = .639). However, rats that received a PRh infusion

of DAP5 (Group DAP5) in stage 2 exhibited significantly less freezing to the S2 than rats that

received a PRh infusion of vehicle only (Group VEH; $F_{(1,22)}$ = 4.644; p = .042; n_p^2 = .174;

95% CI: [0.025, 1.330]).

Experiment 3B – When rats are pre-exposed to S2-S1 pairings prior to conditioning with

S2-S1-shock sequences, the BLA is engaged for acquisition of freezing to S1 but not S2

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,18)}$ = 2.045; p = .170).

Conditioning was again successful. Freezing to the S2 and S1 increased across the four S2-

S1-shock sequences in stage 2 (smaller $F_{(1,18)}$ = 36.409; p < .001; n_p^2 = .669; 95% CI: [1.035,

2.141]). The rate of this increase did not differ between groups (largest $F_{(1,18)}$ = 1.474; p =

.240), and there were no significant between-group differences in overall freezing to the S2

($F_{(1,18)}$ = 1.056; p = .318) and S1 ($F_{(1,18)}$ = .071; p = .793).

Test. Figure 3D shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) and S1 alone (right panel) during testing in Experiment 3B, averaged across the eight

presentations of each stimulus. Both groups showed equivalent levels of freezing to the S2

($F_{(1,18)}$ = 2.078; p = .167). However, rats that received a BLA infusion of DAP5 (Group

DAP5) in stage 2 exhibited significantly less freezing to the directly conditioned S1 than rats

that received a BLA infusion of vehicle (Group VEH; $F_{(1,18)} = 5.899$; $p = .026$; $n_p^2 = .247$;

95% CI: [0.114, 1.571]).

Experiments 3A and 3B showed that pre-exposure can alter the way that stimuli are

processed in the focus/periphery of attention and, thereby, the substrates of their conditioning

in the PRh and BLA. In the preceding experiments (2A and 2B) where rats were conditioned

with S2-S1-shock sequences in the absence of any stimulus pre-exposure, the acquisition of

freezing to S1 and S2 required NMDAR-activation in the BLA but not the PRh. By contrast,

in the present experiments where rats were conditioned with S2-S1-shock sequences 24 h

after having been pre-exposed to S2-S1 pairings, the substrates of conditioning to S1 and S2

were distinct: acquisition of freezing to S1 required NMDAR-activation in the BLA but not

the PRh, whereas acquisition of freezing to S2 required NMDAR-activation in the PRh but

not the BLA. These findings are consistent with our proposal that the BLA and PRh function

like different states of attention. The BLA functions like a focal state and, thus, processes

information about stimuli that are directly paired with shock: hence it was engaged for

conditioning to the pre-exposed S1 which co-terminated with foot shock but was not engaged

for conditioning to the pre-exposed S2 which was separated from the shock by S1. By

contrast, the PRh represents a peripheral state and, as such, maintains a trace/representation

of a pre-exposed stimulus when it is replaced by another stimulus: hence, it was not engaged

for conditioning to the pre-exposed S1 which co-terminated with foot shock but was engaged

for conditioning to the pre-exposed S2 which was separated from the shock by S1.

Experiments 4A and 4B – The roles of the PRh and BLA in Pavlovian fear conditioning

when rats are pre-exposed to S2-S1 pairings prior to conditioning with S2-[trace]-shock

sequences.

In Experiments 3A and 3B, conditioning of the pre-exposed S1 required NMDAR-

activation in the BLA and not the PRh; whereas conditioning of the pre-exposed S2 required

NMDAR-activation in the PRh and not the BLA. We attributed these findings to the fact that

the pre-exposed S2 was separated from the shock by S1 during conditioning of the S2-S1-

shock sequences in stage 2: hence, at the time of the shock, S2 was no longer processed in the

focal state of attention supported by the BLA; instead, it was processed in the peripheral state

of attention supported by the PRh.

Experiments 4A and 4B employed a modification of the protocol used in Experiments

3A and 3B to examine whether a pre-exposed S1 influences the processing of a pre-exposed

S2 across conditioning with S2-S1-shock sequences. The design of these experiments is

shown in Figure 4A. Briefly, in each experiment, two groups of rats were exposed to a

session of S2-S1 pairings in stage 1 and, 24 hours later, a session of S2-[trace]-shock pairings

in stage 2, where the duration of the trace interval was equivalent to that of S1 in the

preceding experiments. Immediately prior to the trace conditioning session in stage 2, one

group in each experiment received an infusion of DAP5 or vehicle only into either the PRh

(Experiment 4A) or BLA (Experiment 4B). Finally, all rats were tested with presentations of

the S2 alone in stage 3. If the substrates of conditioning to S2 are determined only by the

amount of time between presentations of this stimulus and foot shock in stage 2, the results

obtained in Experiments 4A and 4B should be the same as those obtained in Experiments 3A

and 3B: acquisition of freezing to S2 will require activation of NMDARs in the PRh and not

the BLA. If, however, the presence of S1 in the preceding experiments (Experiments 3A and

3B) accelerated the rate at which processing of S2 transitioned from the focus of attention to

its periphery, the results obtained in Experiments 4A and 4B will differ from those obtained

in Experiments 3A and 3B. That is, in contrast to the preceding experiments where

acquisition of freezing to S2 required NMDAR-activation in the PRh and not the BLA, here

acquisition of freezing to S2 should require NMDAR-activation in the BLA but not the PRh.

Experiment 4A – When rats are exposed to S2-S1 pairings in stage 1 and trace

conditioning of S2 in stage 2, the PRh is NOT required for encoding of the S2-shock

association

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,27)}$ = 1.095; p = .305).

Conditioning was successful. Freezing to the S2 increased across the four S2-[trace]-shock

pairings in stage 2 ($F_{(1,27)}$ = 120.496; p < .001; n_p^2 = .817; 95% CI: [1.966, 2.871]). The rate

of this increase did not differ between the two groups ($F_{(1,27)}$ = .332; p = .569), and there was

no significant difference between them in overall freezing to the S2 ($F_{(1,27)}$ = .323; p = .575).

Test. Figure 4B shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) during testing in Experiment 4A, averaged across the eight presentations of this

stimulus. Both groups showed equivalent levels of freezing to the trace conditioned S2 ($F_{(1,27)}$

= 1.505; p = .230).

Experiment 4B – When rats are exposed to S2-S1 pairings in stage 1 and trace

conditioning of S2 in stage 2, the BLA is required for encoding of the S2-shock association

Conditioning. The baseline levels of freezing during conditioning and test sessions

were low (< 10%) and did not differ between groups (largest $F_{(1,24)}$ = 3.823; p = .062).

Conditioning was successful. Freezing to the S2 increased across the four S2-[trace]-shock

pairings in stage 2 ($F_{(1,24)}$ = 47.799; p < .001; n_p^2 = .666; 95% CI: [1.069, 1.979]). The rate of

this increase did not differ between the two groups ($F_{(1,24)}$ = 3.421; p = .077), however there

was a significant between-group difference in overall freezing to the S2 ($F_{(1,24)}$ = 7.040; p =

.014; n_p^2 = .227; 95% CI: [0.168, 1.348]).

Test. Figure 4D shows the mean (± SEM) levels of freezing to the S2 alone (left

panel) during testing in Experiment 4B, averaged across the eight presentations of this

stimulus. Rats that received a BLA infusion of DAP5 (Group DAP5) in stage 2 exhibited

significantly less freezing to the trace conditioned S2 than rats that received a BLA infusion

of vehicle only ($F_{(1,24)} = 7.390$; $p = .012$; $n_p^2 = .235$; 95% CI: [0.213, 1.554]).

Together with the results of the preceding experiments (3A and 3B), the present

findings show that when a pre-exposed S2 is paired with shock, the substrates of its

conditioning are determined by more than just the time between its offset and the shock: they

are additionally determined by what happens in the interval between the two events. When

the interval is blank, as it was in the present experiments (4A and 4B), the acquisition of

freezing to S2 requires NMDAR-activation in the BLA, not the PRh. When the interval is

occupied by a presentation of the pre-exposed S1, as it was in the preceding experiments (3A

and 3B), the acquisition of freezing to S2 requires NMDAR-activation in the PRh, not the

BLA. These findings can be understood in terms of our proposal that the PRh and BLA

function like different states of attention: the presence of S1 in the preceding experiments

(Experiments 3A and 3B) accelerated the rate at which processing of S2 transitioned to a

peripheral state of attention, hence its conditioning in those experiments required NMDAR-

activation in the PRh and not the BLA. This and other implications of our proposal will be

considered further in the General Discussion.

435   **General Discussion**

436   This series of experiments used variations of a sensory preconditioning protocol in

437   rats to test the implications of our proposal that the BLA and PRh represent different states of

438   attention: the BLA functions like a focal state in processing stimuli that are physically

present, regardless of their associative history; whereas the PRh functions like a peripheral

state in supporting the traces or representations of pre-exposed stimuli that are not physically

present. The first implication was that, in our standard sensory preconditioning protocol, the

substrates of conditioning to S2 and S1 would be doubly dissociable at the level of NMDARs

in the PRh and BLA: during the session of S1-shock pairings in stage 2, mediated

conditioning of the absent S2 would occur in the periphery of attention and hence, require

NMDAR-activation in the PRh but *not* the BLA; whereas direct conditioning of the presented

S1 would occur in the focus of attention and hence, require NMDAR-activation in the BLA

but not the PRh. In these experiments, rats were exposed to a session of S2-S1 pairings in

stage 1 and, 24 hours later, to a session of S1-shock pairings in stage 2. We have previously

shown that, across the session of S1-shock pairings, the S1 is directly associated with foot

shock and mediates an association between the *absent-but-expected* S2 and foot shock. Here,

we replicated our previous findings that formation of the direct S1-shock association requires

activation of NMDARs in the BLA but not the PRh (Williams-Spooner et al., 2022; Wong et

al., 2025); and extended these findings to show that formation of the mediated S2-shock

association requires activation of NMDARs in the PRh but not the BLA. Specifically, we

found that a BLA infusion of the NMDAR antagonist, DAP5, immediately prior to the

training session in stage 2 disrupted freezing to the S1 at test without affecting the level of

freezing to S2; and, conversely, a PRh infusion of DAP5 immediately prior to stage 2

disrupted freezing to the S2 at test without affecting the level of freezing to S1. These

findings were taken to mean that the BLA regulates conditioning to a stimulus that is directly

paired with danger and, hence, processed in the focus of attention; whereas the PRh regulates

conditioning to a stimulus that is indirectly paired with danger and, hence, processed in a

peripheral state of attention/memory. Consequently, blocking NMDAR in the BLA impaired

direct conditioning of the S1 but spared mediated conditioning of the S2; and blocking

NMDAR in the PRh impaired mediated conditioning of the S2 but spared direct conditioning

of the S1.

The second implication of our proposal was that combining the S2-S1 and S1-shock

pairings into S2-S1-shock sequences should selectively alter the substrates of conditioning to

S2. Under these circumstances, the novel S2 would be more effectively maintained in the

focus of attention along with the S1: hence, conditioning to both stimuli would occur in the

BLA and not the PRh. This prediction was supported in Experiments 2A and 2B. Here, rats

were exposed to S2-S1-shock sequences and the acquisition of freezing to both S1 and S2

was disrupted by a DAP5 infusion into the BLA but unaffected by a DAP5 infusion into the

PRh. That is, combining the S2-S1 pairings and S1-shock pairings into S2-S1-shock

sequences had no impact on the substrates of conditioning to S1: as in Experiments 1A and

1B where the S2-S1 and S1-shock pairings were separated by 24 hours, acquisition of

freezing to S1 required NMDAR-activation in the BLA but not the PRh. It did, however, alter

the substrates of conditioning to S2: in contrast to Experiments 1A and 1B where acquisition

of freezing to S2 required NMDAR-activation in the PRh but not the BLA, here acquisition

of freezing to S2 required NMDAR-activation in the BLA but not the PRh.

The remaining implications of our proposal concerned the effects of pre-exposing rats

to S2-S1 pairings 24 hours prior to conditioning with S2-S1-shock sequences. We predicted

that, relative to the results of Experiments 2A and 2B, pre-exposing rats to S2-S1 pairings

would again alter the substrates of conditioning to S2 without affecting the substrates of

conditioning to S1. Specifically, we predicted that at the end of each S2-S1-shock sequence

in conditioning (i.e., the time of the shock), the pre-exposed (and thereby familiar) S2 would

be processed in the periphery of attention while the S1 is processed in its focus: hence,

conditioning to S2 would occur in the PRh and not the BLA, whereas conditioning to S1

would occur in the BLA and not the PRh. These predictions were supported in Experiments

3A and 3B. Rats were pre-exposed to S2-S1 pairings and, 24 hours later, exposed to S2-S1-

shock sequences under the influence of a DAP5 infusion into the PRh or BLA. The test

results revealed that the acquisition of freezing to S2 was disrupted by a DAP5 infusion into

the PRh but unaffected by a DAP5 infusion into the BLA; whereas the acquisition of freezing

to S1 was disrupted by a DAP5 infusion into the BLA but unaffected by a DAP5 infusion

into the PRh. That is, pre-exposing rats to S2-S1 pairings prior to conditioning with S2-S1-

shock sequences again had no effect on the substrates of conditioning to S1: as in

Experiments 2A and 2B where rats received no pre-exposure, acquisition of freezing to S1

required NMDAR-activation in the BLA but not the PRh. It did, however, alter the substrates

of conditioning to S2: in contrast to Experiments 2A and 2B where acquisition of freezing to

the novel S2 required NMDAR-activation in the BLA but not the PRh, here acquisition of

freezing to the pre-exposed S2 required NMDAR-activation in the PRh but not the BLA.

The results of Experiments 2A, 2B, 3A and 3B provide support for our idea that the

BLA and PRh function like focal and peripheral states of attention (respectively) by showing

that the locus of conditioning in one region or the other is determined by novelty/familiarity

of the stimulus (or whether it has been pre-exposed or not) and the nature of stimulus-shock

pairings (whether they are direct or indirect). The results of Experiments 4A and 4B extended

these findings by showing that the substrates of conditioning to a pre-exposed stimulus are

also influenced by what happens in the period between presentations of the stimulus and

shock. Here, rats were pre-exposed to S2-S1 pairings and, 24 hours later, exposed to S2-

[trace]-shock pairings under the influence of a DAP5 infusion into the PRh or BLA. Thus,

these experiments were like the preceding ones (3A and 3B) in that S2 was separated from

the shock by 10 s during its conditioning; but critically differed from the preceding

experiments in the fact that the 10 s interval between the S2 and shock was *not* filled by a

presentation of the S1. The consequences of leaving this interval blank were evident at test,

where the pattern of results produced by infusions of DAP5 into the PRh and BLA was

exactly opposite to that obtained in Experiments 3A and 3B. That is, whereas Experiments

3A and 3B showed that freezing to S2 was disrupted by a DAP5 infusion into the PRh but

unaffected by a DAP5 infusion into the BLA, here the test level of freezing to S2 was

disrupted by a DAP5 infusion into the BLA but unaffected by a DAP5 infusion into the PRh.

These results were taken to mean that the substrates of conditioning to a pre-exposed S2 are

influenced by more than just the interval between presentations of this stimulus and foot

shock: it is also influenced by events that occur in that interval. If the interval is blank, the

pre-exposed S2 may remain in a focal state of attention: hence its conditioning requires

NMDAR-activation in the BLA and not the PRh (4A and 4B). If the interval is occupied by

S1, the pre-exposed S2 more rapidly transitions to a peripheral state of attention: hence its

conditioning requires NMDAR-activation in the PRh and not the BLA (3A and 3B).

       Taken together, the results of the present study are generally consistent with theories

of information processing that distinguish between focal and peripheral states of attention

(e.g., Atkinson & Shiffrin, 1968; Posner & Snyder, 1975; Shiffrin & Schneider, 1977).

Importantly, these theories were developed to explain the different characteristics of

information processing; they were not developed to explain changes in the substrates of

learning and memory with variations in the circumstances of training. They do, however,

provide an excellent framework for thinking about the present findings. For example,

according to one type of theory (Wagner, 1981), stimuli are represented as a series of

elements that transition between states of activity and inactivity; and these transitions are

governed by formal rules. Briefly, when a stimulus is presented, its elements are activated to

the focal state of attention (A1) from which they decay to a peripheral state (A2) before

returning to a state of inactivity. Importantly, the capacity of A1 is limited and much less than

that of A2; stimuli processed in A1 can prime their associates to A2; and the rate at which

stimulus elements decay from A1 to A2 is influenced by their familiarity (a result of context-induced priming) as well as other stimuli that are present (i.e., the presence of other stimuli increases the processing demand on A1 and, thereby, the rate of information decay to A2). Given these features and our proposal that the BLA and PRh represent the focus and periphery of attention, a version of this theory can accommodate the full pattern of results in this study (Holland, 1983). When rats are exposed to S2-S1-shock sequences in the absence of any pre-exposure (Experiments 2A and 2B), the elements of both S2 and S1 overlap with those of the foot shock in the focal state of attention. Hence, conditioning to each stimulus requires activation of NMDAR in the BLA and not the PRh. When rats are conditioned with S1-shock pairings in stage 2 after pre-exposure to S2-S1 pairings in stage 1 (Experiments 1A and 1B), S1 elements are activated to the focus of attention and prime S2 elements to the peripheral state. Hence, conditioning to S1 requires NMDAR-activation in the BLA and not the PRh, whereas conditioning to S2 requires NMDAR-activation in the PRh and not the BLA. When rats are conditioned with S2-S1-shock sequences in stage 2 after pre-exposure to S2-S1 pairings in stage 1 (Experiments 3A and 3B), the S2 elements are initially processed in the focus of attention but are rapidly shunted to the periphery during presentations of the S1. Hence, conditioning to S1 requires NMDAR-activation in the BLA and not the PRh, whereas conditioning to S2 requires NMDAR-activation in the PRh and not the BLA. Finally, when rats are conditioned with S2-[trace]-shock pairings after pre-exposure to S2-S1 pairings in stage 1 (Experiments 4A and 4B), the S2 elements are initially processed in the focus of attention and naturally decay to the periphery during the trace interval. However, as the trace interval is relatively short, most S2 elements remain in the focal state at the time of foot shock: hence, conditioning to S2 requires NMDAR-activation in the BLA and not the PRh. With a longer trace interval, the theory predicts that there will be a greater accumulation of S2 elements in the peripheral state: hence, conditioning to S2 would cease to require

NMDAR-activation in the BLA and now require NMDAR-activation in the PRh. This will be

tested in future research.

An additional point to consider in relation to Experiments 3A, 3B, 4A and 4B is the

level of surprise that rats experienced following presentations of the familiar S2 in stage 2.

Specifically, in Experiments 3A and 3B, S2 was followed by the expected S1 (low surprise)

and its conditioning required activation of NMDA receptors in the PRh and not the BLA. By

contrast, in Experiments 4A and 4B, S2 was followed by omission of the expected S1 (high

surprise) and its conditioning required activation of NMDA receptors in the BLA and not the

PRh. This raises the possibility that surprise, or prediction error, also influences the way that

S2 is processed in focal and peripheral states of attention. When prediction error is low, S2 is

processed in the peripheral state of attention: hence, learning under these circumstances

requires NMDA receptor activation in the PRh and not the BLA. By contrast, when

prediction error is high, S2 is preserved in the focal state of attention: hence, learning under

these circumstances requires NMDA receptor activation in the BLA and not the PRh. The

impact of prediction error on the processing of S2 could be assessed using two types of

designs. In the first design, rats are pre-exposed to S2-S1 pairings in stage 1 and this is

followed by S2-S3-shock pairings in stage 2. The important feature of this design is that, in

stage 2, the S2 is followed by surprise in omission of S1 and presentation of S3. Thus, if a

large prediction error maintains processing of the familiar S2 in the BLA, we might expect

that its conditioning in this design would require NMDA receptor activation in the BLA (in

contrast to the results of Experiment 3B) and no longer require NMDA receptor activation in

the PRh (in contrast to the results of Experiment 3A). In the second design, rats are pre-

exposed to S2 alone in stage 1 and this is followed by S2-[trace]-shock pairings in stage 2.

The important feature of this design is that, in stage 2, the S2 is not followed by the surprising

omission of any stimulus. Thus, if a small prediction error shifts processing of the familiar S2

to the PRh, we might expect that its conditioning in this design would no longer require

NMDA receptor activation in the BLA (in contrast to the results of Experiment 4B) but,

instead, require NMDA receptor activation in the PRh (in contrast to the results of

Experiment 4A). Future studies will use both designs to determine whether prediction error

influences the processing of S2 in the focus versus periphery of attention and, thereby,

whether learning about this stimulus requires NMDA receptor activation in the BLA or PRh.

Finally, the present findings add to a body of work which shows that the neural

substrates of Pavlovian conditioned fear critically depend on the type of conditioning (Keidar

et al., 2023; Lay et al., 2018; Leake et al., 2024; Leidl et al., 2018; Williams-Spooner et al.,

2019, 2022); and extend our previous work in sensory preconditioning by identifying how the

mediated S2-shock association is encoded at the level of the PRh and BLA. Specifically, our

previous work showed that formation of this association requires neuronal activity in the PRh

(Wong et al., 2019), as well as communication between this region and the BLA (Wong et

al., 2025). Here, we have extended these findings by showing that NMDAR-activation is a

critical component of the processing requirement in the PRh; but is not a component of the

processing requirement in the BLA. Future work will identify the nature of the processing

requirement in the BLA and exactly how the distinct signaling requirements in the PRh and

BLA combine to generate the mediated S2-shock association.

In summary, the present study has shown that conditioning to a novel stimulus

requires activation of NMDAR in the BLA but does not require activation of NMDAR in the

PRh. By contrast, the substrates of conditioning to a pre-exposed stimulus are influenced by

several factors, including its contiguity with shock and events interpolated between its

presentations and shock. The overall pattern of results is consistent with our proposal that the

BLA and PRh function like focal and peripheral states of attention/memory, respectively; and

generally consistent with information processing theories which specify how events come to

614 be represented in the focal and peripheral states (Holland, 1983; Wagner, 1981). Future

615 research will test the additional implications of these theories and continue to examine

616 commonalities and differences in the processes through which mediated and direct

617 associations are encoded and stored in the medial temporal lobe.

Materials and methods

Subjects

Subjects were experimentally naïve male and female Long-Evans rats, obtained from

a colony maintained by the Biological Resources Centre at the University of New South

Wales. The rats were housed by sex in plastic cages (40 cm wide x 22 cm high x 67 cm long),

with four rats per cage and food and water continuously available. The cages were located in

an air-conditioned and humidity-controlled colony room maintained on a 12 h light-dark

cycle (lights on at 7 a.m. and off at 7 p.m.) at a temperature of approximately 21°C. All

procedures were approved by the Animal Care and Ethics Committee at the University of

New South Wales and conducted in accordance with the Australian code for the care and use

of animals for scientific purposes (published by the National Health and Medical Research

Council of Australia). Please note that these subject details (source, housing conditions, and

animal maintenance) are exactly the same as those reported in past studies by our laboratory

(e.g., Kennedy et al., 2024; Wong et al., 2025).

Surgery

Prior to behavioral training and testing in each experiment, rats were surgically

implanted with cannulas targeting either the perirhinal cortex (PRh; Experiments 1A, 2A, 3A

and 4A) or basolateral amygdala complex (BLA; Experiments 1B, 2B, 3B and 4B). Rats were

anesthetized using isoflurane, which was delivered in a steady stream of oxygen. The rat was

then mounted onto a stereotaxic apparatus (David Kopf Instruments) and incisions made over

the skull. Two holes were drilled through the skull and 26-gauge guide cannulas (Plastic

Ones) were implanted into the brain, one in each hemisphere. For Experiments 1A, 2A, 3A

and 4A, the tips of the cannulas targeted the PRh at coordinates 4.30 mm posterior to

Bregma, 5.00 mm lateral to the midline, 8.4 mm ventral to Bregma, and angled at

approximately 9° (Paxinos & Watson, 2007). For Experiments 1B, 2B, 3B and 4B, the tips of

the cannulas targeted the BLA at coordinates 2.40 mm posterior to Bregma, 4.90 mm lateral

to the midline, 8.40 mm ventral to Bregma (Paxinos & Watson, 2007). Guide cannulas were

secured in place with four jeweller's screws and dental cement. A dummy cannula was kept

in each guide cannula at all times except during drug infusions. Immediately after surgery,

rats received a subcutaneous (SC) injection of a prophylactic dose (0.1 ml/kg) of Duplocillin

(Merck & Co, NJ, USA). Rats were allowed seven days to recover from surgery, during

which they were monitored and weighed daily. Please note that these surgery details are

exactly the same as those reported in our earlier studies (e.g., Wong et al., 2019; Wong et al.,

2025).

Drug infusions

In each experiment, the NMDA receptor antagonist, DAP5, or vehicle was infused

bilaterally into the PRh or BLA. For these infusions, infusion cannulas were connected to 25

µl Hamilton syringes via polyethylene tubing. These syringes were fixed to an infusion pump

(Harvard Apparatus). The infusion procedure began by removing the dummy caps from the

guide cannulas on each rat and inserting 33-gauge infusion cannulas in its place. The pump

was programmed to infuse a total of 0.5 µl of DAP5 at a rate of 0.25 µl/min, which resulted

in a total infusion time of two min. The infusion cannulas remained in place for an additional

two min after the infusion was complete to allow for diffusion of the drug into the PRh or

BLA tissue and, thereby, avoid reuptake when the infusion cannulas were withdrawn. This

resulted in a total infusion time of four min. After the additional two min, the infusion

cannulas were removed and replaced with dummy cannulas. The day prior to infusions, the

dummy cannulas were removed and the infusion pump was activated to familiarize the rats

with the procedure and thereby minimize any non-specific effect of the procedure on the day

of infusions. Please note that these drug and infusion details are exactly the same as those

reported in the earlier study by Wong et al. (2025).

Drugs

The NMDA receptor antagonist, DAP5 (Sigma, Australia) was prepared in the

manner described by Williams-Spooner et al. (2022). Briefly, it was dissolved in ACSF to a

concentration of 10 μg/μl and injected into the PRh or BLA as described above.

Histology

After behavioral training and testing, rats were euthanized with a lethal dose of

sodium pentobarbital. The brains were extracted and sectioned into 40 μm coronal slices.

Every second slice was mounted onto a glass microscope slide and stained with cresyl violet.

The placement of the cannula tip was determined under a microscope using the boundaries

defined by Paxinos and Watson (2006). Please note that the details for histology are exactly

the same as those reported in the earlier studies by Wong et al. (2019, 2025).

Rats with misplaced cannulas were excluded from statistical analysis. Figure 5 shows

placement of the most ventral portion of these cannulas in the BLA for all rats that were

included in the statistical analyses of data from Experiments 1B, 2B, 3B and 4B. Figure 6

shows placement of the most ventral portion of these cannulas in the PRh for all rats that

were included in the statistical analyses of data from Experiments 1A, 2A, 3A and 4A. The

numbers of rats excluded from each experiment based on misplaced cannulas were 3 rats in

Experiment 1A, 12 rats in Experiment 1B, 9 rats in Experiment 2A, 11 rats in Experiment 2B,

6 rats in Experiment 3A, 6 rats in Experiment 3B, 3 rats in Experiment 4A, and 5 rats in

Experiment 4B. A further nine rats were excluded from all experiments due to issues with

their head caps and/or health issues prior to the infusion day. The final *n*s for each group are

shown in the figure legend for each experiment.

Behavioral apparatus

All experiments were conducted in four identical chambers (30 cm wide x 26 cm long

x 30 cm high). The side walls and ceiling were made of aluminum, and the front and back

walls of clear plastic. The floor consisted of stainless steel rods, each two mm in diameter and

spaced 13 mm apart (center-to-center). A waste tray containing bedding material was located

under the floor. At the end of each session, the chambers were cleaned with water and any

soiled bedding was removed from the waste trays and replaced with fresh bedding. Each

chamber was enclosed in a sound- and light-attenuating wooden cabinet. The cabinet walls

were painted black. A speaker and LED lights within a fluorescent tube were mounted onto

the back wall of each cabinet. The speaker was used to deliver a 1000 Hz square-wave tone

stimulus, presented at 75 dB when measured at the center of the chamber (digital sound

meter: Dick Smith Electronics, Australia). The LED lights were used to deliver a flashing

light stimulus, presented at 3.5 Hz. A custom-built, constant current generator was used to

deliver a 0.5 s duration, 0.8 mA intensity shock to the floor of each chamber. Each chamber

was illuminated with an infrared light source, and a camera mounted on the back wall of each

cabinet was used to record the behavior of each rat. The cameras were connected to a monitor

and a high-definition digital image recorder located in an adjacent room. This room also

contained the computer that controlled stimulus and shock presentations through an interface

and appropriate software (MATLAB, MathWorks, USA). Please note that the behavioral

apparatus used for this study was exactly the same as that used in the earlier studies by Wong

et al. (2019, 2025) and a previous study by Kennedy et al. (2024).

Behavioral procedure

 Experiments 1A and 1B

Context exposure. On Days 1 and 2, rats received two 20 min exposures to the chambers, one

in the morning and the other approximately three hours later in the afternoon.

Stage 1: Sensory preconditioning. On Day 3, rats were exposed to eight paired presentations

of the stimuli designated S2 and S1. For half the rats, the S2 was a sound and the S1 was a

light. For the remaining rats, the S2 was a light and the S1 was a sound. Each presentation of

the S2 was 30 s in duration, each presentation of the S1 was 10 s in duration, and the stimuli

were paired in such a way that the offset of S2 co-occurred with the onset of S1. The first S2

presentation occurred five min after rats were placed into the chambers, and the interval

between each S2-S1 pairing was five min (measured from the offset of the S1 to onset of the

next S2). After the last S1 presentation, rats remained in the chambers for an additional one

min. They were then returned to their home cages.

Stage 2: First-order conditioning of the S1. On Day 4, rats were exposed to four paired

presentations of the S1 and foot shock. The first S1 presentation occurred five min after rats

were placed in chambers, each 10 s S1 co-terminated with the 0.5 s foot shock, and the

interval between the S1-shock pairings was five min. Rats remained in the chambers for an

additional one min after the final S1-shock pairing and were then returned to their home

cages.

Context extinction. On Day 5, rats received two 20 min exposures to the chambers, one in the

morning and the other three hours later in the afternoon. These exposures were intended to

extinguish any freezing elicited by the context alone so that the level of freezing to the S2 and

S1 could be assessed unconfounded by context elicited freezing. On Day 6, rats received a

further 10 min extinction exposure to the context.

Testing. On Day 6, approximately two hours after the context extinction session, rats were

tested for their levels of freezing to the preconditioned S2. On Day 7, rats were tested for

their levels of freezing to the conditioned S1. For each test, the first stimulus was presented

two min after rats were placed in the chambers. Each test consisted of eight stimulus alone

presentations, with a three min interval between each presentation. Each S2 presentation was

30 s in duration and each S1 presentation was 10 s. Rats remained in the chamber for an

additional min after the final stimulus presentation.

Experiments 2A and 2B

Context exposure. On Days 1 and 2, rats received two 20 min exposures to the chambers in

the manner described for the previous experiments.

Serial-order conditioning of S2 and S1. On Day 3, rats were exposed to four S2-S1-shock

sequences. For half the rats, the S2 was a sound and the S1 was a light. For the remaining

rats, the S2 was a light and the S1 was a sound. Each presentation of the S2 lasted for 30 s,

each presentation of the S1 lasted for 10 s, and each presentation of foot shock lasted for 0.5

s. The stimuli were presented in such a way that offset of the 30 s S2 co-occurred with onset

of the 10 s S1, which then co-terminated with the 0.5 s foot shock. The first S2 presentation

occurred five min after rats were placed into the chambers, the interval between each S2-S1-

shock sequence was five min (measured from the offset of the S1 to onset of the next S2),

and rats remained in the chambers for an additional one min after exposure to the final S2-S1-

shock sequence. They were then returned to their home cages.

Context extinction. On Day 4, rats received two 20 min exposures to the chambers, one in the

morning and the other three hours later in the afternoon. These exposures were intended to

extinguish any freezing elicited by the context alone so that the level of freezing to the S2 and

S1 could be assessed unconfounded by context elicited freezing. On Day 5, rats received a further 10 min extinction exposure to the context.

Testing. On Day 5, approximately two hours after the context extinction session, rats were tested for their levels of freezing to the S2. On Day 6, rats were tested for their levels of freezing to the S1. The details of these test sessions were identical to those described for the previous experiments.

Experiments 3A and 3B

Context exposure. On Days 1 and 2, rats received two 20 min exposures to the chambers in the manner described for Experiments 1A and 1B.

Stage 1: Sensory preconditioning. On Day 3, rats received a single session of sensory preconditioning in which they were exposed to eight paired presentations of the stimuli designated S2 and S1. For half the rats, the S2 was a sound and the S1 was a light. For the remaining rats, the S2 was a light and the S1 was a sound. The details for this session were identical to those described for the sensory preconditioning session in Experiments 1A and 1B.

Stage 2: Serial-order conditioning of the S2 and S1. On Day 4, rats received a single session in which they were exposed to four S2-S1-shock sequences. The details for this session were identical to those described for the serial-order conditioning session in Experiments 2A and 2B.

Context extinction. On Day 5, rats received two 20 min sessions of context extinction; and this was followed by an additional 10 min context extinction session on Day 6. The details and purpose of these sessions was exactly as described for each of the previous experiments.

Testing. On Day 6, approximately two hours after the context extinction session, rats were tested for their levels of freezing to the S2. On Day 7, rats were tested for their levels of

782 freezing to the S1. The details of these test sessions were exactly as described for each of the

783 previous experiments.

Experiments 4A and 4B

Context exposure. On Days 1 and 2, rats received two 20 min exposures to the chambers in

the manner described for Experiments 1A and 1B.

Stage 1: Sensory preconditioning. On Day 3, rats received a single session of sensory

preconditioning in which they were exposed to eight paired presentations of the stimuli

designated S2 and S1. For half the rats, the S2 was a sound and the S1 was a light. For the

remaining rats, the S2 was a light and the S1 was a sound. The details for this session were

identical to those described for the sensory preconditioning session in Experiments 1A and

1B.

Stage 2: Trace conditioning of the S2. On Day 4, rats received a single session in which they

were exposed to four S2-[10 s]-shock pairings. The stimuli were presented in such a way that

offset of the 30 s S2 was followed by a 10 s trace interval, which co-terminated with a 0.5 s

foot shock. The first S2 presentation occurred five min after rats were placed into the

chambers, the interval between each S2-[10 s]-shock pairing was five min, and rats remained

in the chambers for an additional one min after exposure to the final S2-[10 s]-shock pairing.

They were then returned to their home cages.

Context extinction. On Day 5, rats received two 20 min sessions of context extinction; and

this was followed by an additional 10 min context extinction session on Day 6. The details

and purpose of these sessions was exactly as described for each of the previous experiments.

Testing. On Day 6, approximately two hours after the context extinction session, rats were

tested for their levels of freezing to the S2. On Day 7, rats were tested for their levels of

freezing to the S1. The details of these test sessions were exactly as described for each of the

previous experiments.

Scoring and statistics

Conditioning and test sessions were recorded. Freezing, defined as the absence of all

movements except those required for breathing (Fanselow, 1980), was used as a measure of

conditioned fear. Freezing data were collected using a time-sampling procedure in which

each rat was scored as either 'freezing' or 'not freezing' every two seconds by an observer

that was unaware of the rat's group allocation. A second naïve observer also scored all of the

test data. The correlation between the two observers' scores was high (Pearson > 0.9). The

data were analyzed using a set of planned orthogonal contrasts (Hays, 1963), with the Type 1

error rate controlled at $\alpha = 0.05$. Standardized 95% confidence intervals (CIs) are reported for

all significant results, and Cohen's d is reported as a measure of effect size (where 0.2, 0.5,

and 0.8 is a small, medium and large effect size, respectively). The required number of rats

per group was determined during the design stage of the study. It was based on our prior

studies of sensory preconditioning which indicated that eight subjects per group (or per

comparison in the contrast testing procedure) provides sufficient statistical power to detect

effect sizes > 0.5 with the recommended probability of 0.8-0.9. Please note that the details for

scoring and data analysis are exactly the same as those reported in the earlier study by Wong

et al (2025).

825   **References**

Atkinson RC, Shiffrin RM. (1968). Human memory: a proposed system and its control

processes. *Psychology of Learning and Motivation*, *2*, 89-195.

Bauer EP, Schafe GE, LeDoux JE. (2002). NMDA receptors and L-type voltage-gated

calcium channels contribute to long-term potentiation and different components of fear

memory formation in the lateral amygdala. *Journal of Neuroscience*, *22*, 5239-49.

Bogacz R, Brown MW. (2003). Comparison of computational models of familiarity

discrimination in the perirhinal cortex. *Hippocampus*, *13*(4), 494-524. doi:

10.1002/hipo.10093.

Brown MW, Banks PJ. (2015). In search of a recognition memory engram. *Neuroscience and*

Biobehavioral Reviews, *50*, 12-28. doi: 10.1016/j.neubiorev.2014.09.016.

Brown MW, Bashir ZI. (2002). Evidence concerning how neurons of the perirhinal cortex

may effect familiarity discrimination. *Philos Trans R Soc Lond B Biol Sci*, 357(1424),

1083-95. doi: 10.1098/rstb.2002.1097.

Campeau S, Miserendino MJ, Davis M. (1995). Intra-amygdala infusion of the N-methyl-D-

aspartate receptor antagonist AP5 blocks acquisition but not expression of fear-

potentiated startle to an auditory conditioned stimulus. *Behavioral Neuroscience*, *106*,

569-74. doi: 10.1037/0735-7044.106.3.569.

Fanselow MS, Kim JJ. (1994). Acquisition of contextual Pavlovian fear conditioning is

blocked by application of an NMDA receptor antagonist D,L-2-amino-5-

phosphonovaleric acid to the basolateral amygdala. *Behavioral Neuroscience*, *108*, 210-

12. doi: 10.1037/0735-7044.108.1.210.

Gewirtz JC, Davis M. (1997). Second-order fear conditioning prevented by blocking NMDA

receptors in amygdala. *Nature*, *388*, 471-74.

Holland PC. (1983). Representation-mediated overshadowing and potentiation of conditioned

aversions. *Journal of Experimental Psychology: Animal Behavior Processes*, *9*, 1-13.

Holmes NM, Harris JA. (2009). The loss of latent inhibition across compound conditioning.

Journal of Experimental Psychology: Animal Behavior Processes, *35*(3), 328-39. doi:

10.1037/a0014537.

Holmes NM, Parkes SL, Killcross AS, Westbrook RF. (2013). The basolateral amygdala is

critical for learning about neutral stimuli in the presence of danger, and the perirhinal

cortex is critical in the absence of danger. *Journal of Neuroscience*, 33, 13112-25. doi:

10.1523/JNEUROSCI.1998-13.2013.

Holmes NM, Raipuria M, Qureshi OA, Killcross S, Westbrook F. (2018). Danger changes the

way the mammalian brain stores information about innocuous events: a study of

sensory preconditioning in rats. *eNeuro*, *5*(1), ENEURO.0381-17.2017. doi:

10.1523/ENEURO.0381-17.2017.

Holmes NM, Westbrook RF. (2017). A dangerous context changes the way that rats learn

about and discriminate between innocuous events in sensory preconditioning. *Learning*

and Memory, *24*, 440-48. doi: 10.1101/lm.044297.116.

Holmes NM, Westbrook RF. (2013). ABA renewal is greater when extinction occurs in the

same context as cue pre-exposure. *Journal of Experimental Psychology: Animal*

Learning and Cognition, *40*(3), 369-79. doi: 10.1037/xan0000024.

Keidar T, Williams-Spooner MJ, Wong FS, Westbrook RF, Holmes NM. (2023). NMDA

receptors in the basolateral amygdala complex are engaged for Pavlovian fear

conditioning when an animal's predictions about danger are in error. *Journal of*

Neuroscience, *43*(39), 6679-96. doi: 10.1523/JNEUROSCI.0067-23.2023.

Kennedy NGW, Holmes NM, Westbrook RF. (2021). Latent inhibition is facilitated when a

target stimulus is preexposed in compound with a nontarget stimulus, but only when the

two stimuli coterminate: a test of the Hall-Rodriguez theory. *Journal of Experimental*

Psychology: Animal Learning and Cognition, *47*(2), 91-103. doi: 10.1037/xan0000289.

Lay BPP, Westbrook RF, Glanzman DL, Holmes NM. (2018). Commonalities and

differences in the substrates underlying consolidation of first- and second-order

conditioned fear. *Journal of Neuroscience*, *38*(8), 1926-41. doi:

10.1523/JNEUROSCI.2966-17.2018.

Leake J, Leidl DM, Lay BPP, Fam JP, Giles MC, Qureshi OA, Westbrook RF, Holmes NM.

(2024). What is learned determines how Pavlovian conditioned fear is consolidated in

the brain. *Journal of Neuroscience*, *44*(2), e0513232023. doi:

10.1523/JNEUROSCI.0513-23.2023.

Leidl DM, Lay BPP, Chakouch C, Westbrook RF, Holmes NM. (2018). Protein synthesis in

the basolateral amygdala complex is required for consolidation of a first-order fear

memory, but not for consolidation of a higher-order fear memory. *Neurobiology of*

Learning and Memory, *153*(B), 153-65. doi: 10.1016/j.nlm.2018.04.001.

Leung HT, Killcross AS, Westbrook RF. (2013a). A further assessment of the Hall-

Rodriguez theory of latent inhibition. *Journal of Experimental Psychology: Animal*

Behavior Processes, *39*(2), 117-25. doi: 10.1037/a0031724.

Leung HT, Killcross AS, Westbrook RF (2013b). Error correction in latent inhibition and its

disruption by opioid receptor blockade with naloxone. *Neuropsychopharmacology*,

38(12), 2439-45. doi: 10.1038/npp.2013.144.

Lubow RE. (1973). Latent inhibition. *Psychological Bulletin*, *79*(6), 398-407. doi:

10.1037/h0034425.

Lubow RE, Moore AU. (1959). Latent inhibition: the effect of nonreinforced pre-exposure to

the conditional stimulus. *Journal of Comparative and Physiological Psychology*, *52*,

415-9. doi: 10.1037/h0046700.

Martin CB, McLean DA, O'Neil EB, Köhler S. (2013). Distinct familiarity-based response

patterns for faces and buildings in perirhinal and parahippocampal cortex. *Journal of*

Neuroscience, *33*(26), 10915-23. doi: 10.1523/JNEUROSCI.0126-13.2013.

McLelland VC, Chan D, Ferber S, Barense MD. (2014). Stimulus familiarity modulates

functional connectivity of the perirhinal cortex and anterior hippocampus during visual

discrimination of faces and objects. *Frontiers in Human Neuroscience*, *8*:117. doi:

10.3389/fnhum.2014.00117.

Miserendino MJ, Sananes CB, Melia KR, Davis M. (1990). Blocking of acquisition but not

expression of conditioned fear-potentiated startle by NMDA antagonists in the

amygdala. *Nature*, 345, 716-18. doi: 10.1038/345716a0.

Morillas E, Gómez-Chacón B, Gallo M. (2017). Flavor and object recognition memory

impairment induced by excitotoxic lesions of the perirhinal cortex. *Neurobiology of*

Learning and Memory, *144*, 230-34. doi: 10.1016/j.nlm.2017.08.002.

Murray EA, Wise SP. (2012). Why is there a special issue on perirhinal cortex in a journal

called hippocampus? The perirhinal cortex in historical perspective. *Hippocampus*,

22(10), 1941-51. doi: 10.1002/hipo.22055.

Parkes SL, Westbrook RF. (2010). The basolateral amygdala is critical for the acquisition and

extinction of associations between a neutral stimulus and a learned danger signal but

not between two neutral stimuli. *Journal of Neuroscience*, *30*, 12608-18. doi:

10.1523/JNEUROSCI.2949-10.2010.

Phillips RG, LeDoux JE. (1995). Lesions of the fornix but not the entorhinal or perirhinal

cortex interfere with contextual fear conditioning. *Journal of Neuroscience*, *15*, 5308-

15. doi: 10.1523/JNEUROSCI.15-07-05308.1995.

Posner MI, Snyder CRR. (1975). Attention and Cognitive Control. In *Information Processing*

and Cognition: The Loyola Symposium, (pp.55-85), 1975, Hillsdale, NJ: Lawrence

Erlbaum Associates.

Qureshi OA, Leake J, Delaney AJ, Killcross S, Westbrook RF, Holmes NM. (2023). Danger

changes the way the brain consolidates neutral information; and does so by interacting

with processes involved in the encoding of that information. *Journal of Neuroscience*,

43(16), 2934-49. doi: 10.1523/JNEUROSCI.1796-22.2023.

Robinson J, Whitt EJ, Horsley RR, Jones PM. (2010). Familiarity-based stimulus

generalization of conditioned suppression in rats is dependent on the perirhinal cortex.

Behahvioral Neuroscience, *124*(5), 587-99. doi: 10.1037/a0020900.

Rodrigues SM, Schafe GE, LeDoux JE. (2001). Intra-amygdala blockade of the NR2B

subunit of the NMDA receptor disrupts the acquisition but not the expression of fear

conditioning. *Journal of Neuroscience*, *21*, 6889-96.

Romanski LM, LeDoux JE. (1992a). Bilateral destruction of neocortical and perirhinal

projection targets of the acoustic thalamus does not disrupt auditory fear conditioning.

Neuroscience Letters, *142*, 228-32. doi: 10.1016/0304-3940(92)90379-L.

Romanski LM, LeDoux JE. (1992b). Equipotentiality of thalamo-amygdala and thalamo-

cortico-amygdala circuits in auditory fear conditioning. *Journal of Neuroscience*, *12*,

4501-09. doi: 10.1523/JNEUROSCI.12-11-04501.1992.

Roozendaal B, Okuda S, Van der Zee EA, McGaugh JL. (2006). Glucocorticoid enhancement

of memory requires arousal-induced noradrenergic activation in the basolateral

amygdala. *Proceedings of the National Academy of Sciences USA*, *103*(17), 6741-46.

doi: 10.1073/pnas.0601874103.

Rosenberg H, Holmes NM, Harris JA, Westbrook RF. Pre-exposure enhances recovery of

conditioned responding after extinction. *Learning and Behavior*, *39*(3), 212-23. doi:

10.3758/s13420-011-0019-5.

Rutishauser U, Mamelak AN, Schuman EM. (2006). Single-trial learning of novel stimuli by

individual neurons of the human hippocampus-amygdala complex. *Neuron*, *49*(6), 805-

13. doi: 10.1016/j.neuron.2006.02.015.

Schomaker J, Meeter M. (2015). Short- and long-lasting consequences of novelty, deviance

and surprise on brain and cognition. *Neuroscience and Biobehavioral Reviews*, *55*, 268-

79. doi: 10.1016/j.neubiorev.2015.05.002.

Shiffrin RM, Schneider W. (1977). Controlled and automatic human information processing:

II. Perceptual learning, automatic attending and a general theory. *Psychological

Review, 84*(2), 127–190. doi: 10.1037/0033-295X.84.2.127.

Suzuki WA, Naya Y. (2014). The perirhinal cortex. *Annual Review of Neuroscience*, 37, 39-

53. doi: 10.1146/annurev-neuro-071013-014207.

Voss JL, Hauner KK, Paller KA. (2009). Establishing a relationship between activity

reduction in human perirhinal cortex and priming. *Hippocampus*, *19*(9), 773-8. doi:

10.1002/hipo.20608.

Wagner AR. (1981). SOP: A model of automatic memory processing in animal behavior. In

In: Spear, N.E., Miller, R.R. (Eds.), Information Processing in Animals: Memory

Mechanisms. (pp. 5-47).

Westbrook RF, Jones ML, Bailey GK, Harris JA. (2000). Contextual control over

conditioned responding in a latent inhibition paradigm. *Journal of Experimental*

Psychology: Animal Behavior Processes, *26*(2), 157-73. doi: 10.1037//0097-

7403.26.2.157.

Wilensky AE, Schafe GE, Kristensen MP, LeDoux JE. (2006). Rethinking the fear circuit:

the central nucleus of the amygdala is required for the acquisition, consolidation, and

expression of Pavlovian fear conditioning. *Journal of Neuroscience*, *26*, 12387-96. doi:

10.1523/JNEUROSCI.4316-06.2006.

Williams-Spooner MJ, Delaney AJ, Westbrook RF, Holmes NM. (2022). Prediction error

determines whether NMDA receptors in the basolateral amygdala complex are involved

in Pavlovian fear conditioning. *Journal of Neuroscience*, *42*(21), 4360-4379. doi:

10.1523/JNEUROSCI.2156-21.2022.

Wong FS, Westbrook RF, Holmes NM. (2019). 'Online' integration of sensory and fear

memories in the rat medial temporal lobe. *eLife*, 8:e47085. doi: 10.7554/eLife.47085.

Wong FS, Thomas AB, Killcross S, Laurent V, Westbrook RF, Holmes NM. (2025).

Integration of sensory and fear memories in the rat medial temporal lobe. *eLife*,

13:RP101965. doi: 10.7554/eLife.101965.

Wright CI, Martis B, Schwartz CE, Shin LM, Fischer H H, McMullin K, Rauch SL. (2003).

Novelty responses and differential effects of order in the amygdala, substantia

innominata, and inferior temporal cortex. *Neuroimage*, *18*(3), 660-9. doi:

10.1016/s1053-8119(02)00037-x.

Yang H, McRae K, Köhler S. (2023). Perirhinal cortex automatically tracks multiple types of

familiarity regardless of task-relevance. *Neuropsychologia*, *187*, 108600. doi:

10.1016/j.neuropsychologia.2023.108600. Epub 2023 May 29.

Tables and Figures

992   **Table 1.** Designs and predictions for each experiment in the series.

EXPERIMENT	TRAINING		PREDICTIONS	
	STAGE 1	STAGE 2	FOR S2	FOR S1
1A	S2-S1	● S1-shock	Disrupted	intact
1B	S2-S1	● S1-shock	Intact	disrupted
2A	...	● S2-S1-shock	Intact	intact
2B	...	● S2-S1-shock	Disrupted	disrupted
3A	S2-S1	● S2-S1-shock	Disrupted	intact
3B	S2-S1	● S2-S1-shock	Intact	disrupted
4A	S2-S1	● S2-[trace]-shock	Intact	
4B	S2-S1	● S2-[trace]-shock	Disrupted	

993   *Notes.* ● *= PRh infusion of DAP5;* ● *= BLA infusion of DAP5. In Experiments 1A, 1B, 3A and 3B, S2*
994   *is familiar and indirectly paired with shock, whereas S1 is familiar and directly paired with shock. In*
995   *Experiments 2A and 2B, S2 is novel and indirectly paired with shock, whereas S1 is novel and directly*
996   *paired with shock. In Experiments 4A and 4B, S2 is familiar and more-directly paired with shock*
997   *(compared to Experiments 3A and 3B).*

Figure 1. The neural substrates of the mediated S2-shock and direct S1-shock associations that form in stage 2 of sensory preconditioning are doubly dissociable at the level of NMDAR-activation in the PRh and BLA. (A, C) Schematics of the protocols used to assess the involvement of NMDAR in acquisition of freezing to S2 and S1 in the PRh (Experiment 1A) and BLA (Experiment 1B). The protocols differed only in whether rats received an infusion of the NMDAR antagonist, DAP5, into the PRh (in A) or BLA (in C). (B) Test results showing that a stage 2 infusion of DAP5 into the PRh disrupts freezing to S2 without affecting freezing to S1. (D) Test results showing that a stage 2 infusion of DAP5 into the BLA disrupts freezing to S1 without affecting freezing to S2. The bars in each histogram show the mean level of freezing in each group across repeated presentations of S2 or S1 under conditions of extinction. The error bars show the standard error of the mean, and the overlaid data points represent the mean level of freezing for individual rats. The final group

1013   sizes were $n = 10$ for Group VEH and $n = 11$ for Group DAP5 in Experiment 1A; and $n = 15$

1014   for Group VEH and $n = 13$ for Group DAP5 in Experiment 1B.

1015

1016

Figure 2. Combining the S2-S1 and S1-shock pairings into S2-S1-shock sequences alters the substrates of conditioning to the S2 in the PRh and BLA. **(A, C)** Schematics of the protocols used to assess when NMDAR are involved in acquisition of freezing to S2 and S1 in the PRh (Experiment 2A) and BLA (Experiment 2B). The protocols differed only in whether rats received an infusion of the NMDAR antagonist, DAP5, into the PRh (in A) or BLA (in C). **(B)** Test results showing that a stage 2 infusion of DAP5 into the PRh has no effect on levels of freezing to either S2 or S1. **(D)** Test results showing that a stage 2 infusion of DAP5 into the BLA disrupts levels of freezing to both the S2 and S1. The final group sizes were $n = 8$ for Group VEH and $n = 9$ for Group DAP5 in Experiment 2A; and $n = 8$ for Group VEH and $n = 9$ for Group DAP5 in Experiment 2B.

1029

Figure 3. Pre-exposing rats to S2-S1 pairings prior to a session of S2-S1-shock sequences re-

engages NMDAR in the PRh for conditioning of the S2. **(A, C)** Schematics of the protocols

used to assess when NMDAR are involved in acquisition of freezing to S2 and S1 in the PRh

(Experiment 3A) and BLA (Experiment 3B). The protocols differed only in whether rats

received an infusion of the NMDAR antagonist, DAP5, into the PRh (in A) or BLA (in C).

(B) Test results showing that a stage 2 infusion of DAP5 into the PRh disrupts freezing to S2

without affecting freezing to S1. **(D)** Test results showing that a stage 2 infusion of DAP5

into the BLA disrupts freezing to S1 without affecting freezing to S2. The final group sizes

were $n = 12$ for Group VEH and $n = 12$ for Group DAP5 in Experiment 3A; and $n = 10$ for

Group VEH and $n = 10$ for Group DAP5 in Experiment 3B.

1040

Figure 4. Pre-exposing rats to S2-S1 pairings prior to a session of S2-[trace]-shock pairings re-engages NMDAR in the BLA for conditioning of the S2. **(A, C)** Schematics of the protocols used to assess when NMDAR are involved in acquisition of freezing to S2 and S1 in the PRh (Experiment 4A) and BLA (Experiment 4B). **(B, D)** Test results showing that a stage 2 infusion of DAP5 has no effect on the level of freezing to S2 when injected into the PRh (in B) but disrupts the level of freezing to S2 when injected into the BLA (in D). The final group sizes were $n = 14$ for Group VEH and $n = 15$ for Group DAP5 in Experiment 3A; and $n = 12$ for Group VEH and $n = 14$ for Group DAP5 in Experiment 3B.

-1.72

-1.80

-1.92

-2.04

-2.16

-2.28

-2.40

-2.52

-2.64

-2.76

1051

1052 **Figure 5.** Cannula placements in the PRh for rats in Experiments 1B, 2B, 3B and 4B. The

1053 most ventral portion of the cannulas are marked on coronal sections based on the atlas of

1054 Paxinos and Watson (2006).

1055

1056

-3.48

-3.60

-3.72

-3.84

-3.96

-4.08

-4.20

-4.36

-4.44

-4.56

-4.68

1057

1058 **Figure 6.** Cannula placements in the PRh for rats in Experiments 1A, 2A, 3A and 4A. The

1059 most ventral portion of the cannulas are marked on coronal sections based on the atlas of

1060 Paxinos and Watson (2006).

A.

B.

Test with Preconditioned Sound

Test with Conditioned Light

C.

D.

Test with Preconditioned Sound

Test with Conditioned Light

A. Stage 1 — Stage 2 — Test

PRh infusion: DAP5 or VEH

B.

Test with Preconditioned Sound

Test with Conditioned Light

C. Stage 1 — Stage 2 — Test

BLA infusion: DAP5 or VEH

D.

Test with Preconditioned Sound

Test with Conditioned Light

A.

B.

C.

D.

A.

| Stage 1 | Stage 2 | Test |

24 h → 10 s 48 h →

PRh infusion: DAP5 or VEH

B.   Test with Preconditioned Sound

C.

| Stage 1 | Stage 2 | Test |

24 h → 10 s 48 h →

BLA infusion: DAP5 or VEH

D.   Test with Preconditioned Sound

-1.72

-1.80

-1.92

-2.04

-2.16

-2.28

-2.40

-2.52

-2.64

-2.76

-3.48

-3.60

-3.72

-3.84

-3.96

-4.08

-4.20

-4.36

-4.44

-4.56

-4.68