Bilateral equalization of synaptic output in olfactory glomeruli of Xenopus tadpoles

Marta Casas[1,2], Beatrice Terni[1,2] and Artur Llobet[1,2*]

[1] Laboratory of Neurobiology, Department of Pathology and Experimental Therapy, Institute of Neurosciences, University of Barcelona, 08907 L'Hospitalet de Llobregat, Barcelona, Spain

[2] Bellvitge Biomedical Research Institute (IDIBELL), 08907 L'Hospitalet de Llobregat, Barcelona, Spain

* Corresponding Author:
Artur Llobet
Laboratory of Neurobiology
Department of Pathology and Experimental Therapy
Faculty of Medicine and Health Sciences – University of Barcelona
08907 L'Hospitalet de Llobregat
Barcelona, Spain
E-mail: allobet@ub.edu
Phone: +34-934024279
ORCID: 0000-0001-5797-6782

Odorants stimulate olfactory sensory neurons (OSNs) to create a bilateral sensory map defined by a set of glomeruli present in the left and right olfactory bulbs. Using *Xenopus tropicalis* tadpoles we challenged the notion that glomerular activation is exclusively determined ipsilaterally. Glomerular responses evoked by unilateral stimulation were potentiated following transection of the contralateral olfactory nerve. The gain of function was observed as early as 2 hours after injury and faded away with a time constant of 4 days. Potentiation was mediated by the presence of larger and faster calcium transients driving glutamate release from OSN axon terminals. The cause was the reduction of the tonic presynaptic inhibition exerted by dopamine D_2 receptors. Inflammatory mediators generated by injury were not involved. These findings reveal the presence of a bilateral modulation of glomerular output driven by dopamine that compensates for imbalances in the number of operative OSNs present in the two olfactory epithelia. Considering that the constant turnover of OSNs is an evolutionary conserved feature of the olfactory system and determines the innervation of glomeruli, the compensatory mechanism here described may represent a general property of the vertebrate olfactory system to establish an odor map.

1 **INTRODUCTION**
2
Olfactory glomeruli are spherical regions of neuropil that contain the first synapse for

processing olfactory information and form the glomerular layer of the olfactory bulb. Despite

variations in number or size among vertebrate species, glomeruli show an evolutionarily

conserved synaptic connectivity (Shepherd, Chen, and Greer 2004). Glomerular input comes

from axon terminals of olfactory sensory neurons (OSNs) expressing the same odorant

receptor that transfer information to the dendrites of mitral cells. Synaptic transmission

between OSNs and mitral cells is regulated by inhibitory and excitatory contacts established

with juxtaglomerular neurons. This connectivity allows that upon exposure to odorants only

a defined set of glomeruli is activated, creating an odor map (Mombaerts et al. 1996).

Glomeruli exhibit an anatomical and functional symmetrical distribution in the left and right

olfactory bulbs, therefore, the sensory map created in the olfactory bulb is bilateral and is

generated by homologous glomeruli reflecting the contribution of the two nasal cavities

(Lodovichi 2021; Belluscio and Katz 2001).

OSNs originate in the olfactory epithelium and are subject to a continuous turnover

throughout life (Holl 2018). An efficient neuronal replacement is accomplished by the rapid

connection of newborn OSNs to the olfactory bulb. For example, in *X. tropicalis* tadpoles

OSNs can establish functional synapses in a time window of 4 days (Terni et al. 2017) and, in

mice, it takes a week for newborn OSNs to get inserted in the olfactory bulb through a *plug-*

and-play mechanism (Browne, Crespo, and Grubb 2022). The balance existing between the

elimination of OSNs and their insertion in olfactory bulb circuitry likely determines a range

of input neurons innervating a glomerulus, rather than a constant, precise figure. In this

context, how the formation of an odor map accounts for possible variations in the number of

input neurons is unknown. Considering that OSNs exclusively form ipsilateral synapses

(Shepherd, Chen, and Greer 2004) and release glutamate with near-maximal release

probability (Murphy et al. 2004), the possible crosstalk balancing the individual contribution

of bilaterally distributed glomeruli remains unknown, because their high output gain is

assumed to be defined unilaterally.

Here, we take advantage of the experimental capacities offered by *Xenopus tropicalis*

tadpoles to assess whether the output of a genetically labelled glomerulus (Terni et al. 2017;

Terni and Llobet 2021), is solely determined by ipsilateral stimulation. In vivo recordings

carried out after disrupting the contribution of the contralateral pathway, challenged this

notion. Our results show that the tonic inhibition of glomerular responses exerted by

dopamine D_2 receptors is bilaterally shaped to compensate for differences in the number of

OSNs innervating the contralateral olfactory bulb. Considering the evolutionary conserved

developmental, morphological, and functional features of the *Xenopus* tadpole olfactory

system (Menini 2010; Manzini, Schild, and Di Natale 2022), the homeostatic mechanism

here described might represent a general rule to the formation of an integrated odor map in

vertebrates.

MATERIALS AND METHODS

Animals

Ethical procedures were approved by the regional government (Generalitat de

Catalunya, experimental procedure #10753). *Xenopus tropicalis* were housed and raised

according to the standard protocols of the animal facilities of the University of Barcelona.

Tadpoles were obtained by natural mating and kept in tanks at 25 ℃. *Xenopus* larvae are not

sexually dimorphic in aquatic stages and sex differences were not considered in the current

study. *Xenopus* water conductivity was adjusted to 700 $\mu s \cdot cm^{-1}$, pH=7.5. Two to three weeks

old tadpoles, found at stage 47-52 of the Nieuwkoop–Faber criteria (Nieuwkoop and Faber

1956),were used for the experiments. The animals of a defined experimental group came

from >4 different natural matings. Experimental group size typically ranged from 8 to 15

tadpoles as it allowed to evaluate statistical differences using parametric tests.

The *X. tropicalis* transgenic line *Dre.mxn1:GFP* (RRID:NXR_1111) was used for the

visualization of a discrete population of OSNs (Terni et al. 2017; Terni and Llobet 2021) to

carry out electrophysiological recordings in a genetically defined glomerulus. Labelling of

OSNs was consistent in all animals of this transgenic line. Calcium imaging was performed

in the *X. tropicalis* transgenic line *ElasGFP:Tubb2b-GCaMP6s* (RRID:NXR_1123). In vivo

detection of reactive oxygen species (ROS) was carried out in tadpoles of the *X.laevis*

transgenic line *Hsa.UBC-Gal4;UAS:HyPer-YFP* (RRID: NXR_0127).

For surgical procedures, *Xenopus* tadpoles were anesthetized in 0.02% MS-222 and

transferred to a wet nitrocellulose filter paper placed under a stereomicroscope. Iridectomy

scissors were used for the unilateral transection of an olfactory nerve or the bilateral

transection of optic nerves. Tadpoles were returned to water tanks after surgery.

Electrophysiology

X. tropicalis tadpoles were anesthetized in 0.02% MS-222 and the portion of skin

covering the olfactory bulb was removed. Animals were transferred to a well fabricated in a

dish coated with silicone elastomer. A coverslip restricted tadpole movements and left

olfactory placodes and bulbs accessible (Terni et al. 2018). The dish was placed on the stage

of an upright microscope (Zeiss, Axioexaminer A1, Oberkochen, Germany) and was

continuously perfused with *Xenopus* Ringer containing (in mM):100 NaCl, 2 KCl, 1 $CaCl_2$, 2

$MgCl_2$, 10 glucose, 10 HEPES, 240 mOsm/kg, pH=7.8, supplemented with 1 µM d-

tubocurarine to prevent muscle contractions. All salts and drugs were from Sigma-Aldrich

(Sant Louis, MO). CGP-36742 was from Novartis Pharmaceuticals (Basel, Switzerland). To

carry out extracellular recordings a borosilicate pipette filled with *Xenopus* ringer was

targeted to the olfactory glomerulus showing GFP fluorescence. Signals were acquired using

a Geneclamp 500A amplifier (Molecular Devices, San Jose, CA) and digitized at 10 kHz

using a National Instruments NI-USB-6341 DAC board (National Instruments, Austin, TX)

controlled by mafPC software (courtesy of M. A. Xu-Friedman, University at Buffalo, NY).

The recorded changes of the local field potential (LFP) were low pass filtered below 100 Hz

and analyzed with Igor Pro 9.0 (Wavemetrics, OR).

Methionine was chosen as odor stimulus based on its broad capacity to stimulate

OSNs in *Xenopus laevis* tadpoles (Manzini and Schild 2004). A puff of 200 µM methionine

solution, obtained by diluting a 10 mM stock solution prepared in *Xenopus* Ringer (pH=7.8),

was applied on the ipsilateral olfactory epithelium to the recorded glomerulus. The amino

acid was delivered through a 0.25 mm diameter fused silica capillary (World Precision

Instruments, Hertfordshire, UK) positioned on the top of a nasal cavity. The timing of

application was controlled via a TTL pulse. The characteristic odor-evoked glomerular

response was obtained by averaging LFP changes triggered by sequential stimulations

applied at 2 min intervals. The local application of antagonists was carried out by transiently

applying 20 psi pressure to the pipette holder inlet. Approximately 1 µL of the pipette

solution was delivered to the glomerulus.

Imaging

Quantification of OSNs present in the olfactory epithelium of *Dre.mxn1:GFP*

tadpoles was performed in an inverted LSM880 confocal microscope (Zeiss) using animals

anesthetized in 0.02% MS-222. The number of neurons was estimated from maximal

intensity projections in confocal Z-stacks. To selectively eliminate GFP positive OSNs the

2Phatal method (Hill et al. 2017) was adapted to confocal microscopy. Tadpoles were

immersed for 15 minutes in *Xenopus* water containing 5 µg/mL Hoechst 33342 and

anesthetized in 0.02% MS-222. Animals were placed dorsally on a 25 mm glass #1.5

coverslip acting as the bottom of an imaging chamber and transferred to the microscope

stage. The right olfactory epithelium, located contralaterally to the recording site, was

imaged with a 10X Plan Apo objective, NA=0.45 (Zeiss). The digital zoom was set to 3.5.

Photobleaching was carried out using ZenBlue software (Zeiss) in 4 to 6 regions of interest

(ROIs) containing two or more GFP positive OSNs. Animals were next returned to water

tanks.

The *X. tropicalis* transgenic line *ElasGFP:Tubb2b-GCaMP6s*, where the neuronal β-

tubulin promoter drives the expression of the calcium sensor GCaMP6s, was used to

visualize the activation of olfactory glomeruli by odorants. As for in vivo electrophysiology,

tadpoles were anesthetized in 0.02% MS-222, placed in a recording dish and transferred to

the stage of an upright microscope (Zeiss, Axioexaminer A1). Tadpoles were continuously

perfused with *Xenopus* Ringer containing 1 μM d-tubocurarine. Imaging was carried out at

76 Hz using a Maico MEMS confocal unit (Hamamatsu Photonics, Hamamatsu City, Japan).

A 40X, 0.75 NA water immersion W-N-Achroplan objective (Zeiss) was used. Odorant

stimulation was performed as described for electrophysiology experiments. The timing of

methionine application was synchronized with image acquistion (HCImage software,

Hamamatsu Photonics; RRID:SCR_015041) using a Master 8 stimulator (AMPI, Jerusalem,

Israel). In experiments where imaging was coupled to recordings of the LFP, the electrode

was targeted to the lateral glomerular cluster. All signals were synchronized using a Master 8

stimulator.

Raw fluorescence image sequences were used to construct ΔF/F movies according to

the relationship $(F-F_0)/F_0$, where F_0 corresponded to the basal fluorescence levels obtained

during 1 s before stimulation. ΔF/F sequences were next subsampled by averaging groups of

10 frames to identify putative glomeruli, which were defined as 10-20 μm diameter round

structures consistently responding to sequential stimulations applied at 2 min intervals. The

selected ROIs were transferred to the original raw fluorescence movie to quantify increases in

basal intracellular calcium levels. A ROI showing $\Delta F/F$ increases evoked by methionine

application to the ipsilateral olfactory epithelium that were ≥ 3 SDs above baseline levels was

considered as a single glomerulus. Calcium imaging of the dorsolateral pallium was carried

out as in the olfactory bulb but using ROIs measuring 30 µm diameter. The temporal

response was quantified using Igor Pro 9 software (Wavemetrics, OR).

The production of ROS in the *X. laevis* HyPer-YFP was evaluated after transection of

a single olfactory nerve. Tadpoles were imaged in an inverted confocal LSM 900 microscope

(Zeiss) using a Plan-Apochromat 20X, 0.8NA objective. Hyper-YFP was excited at 405 nm

and 488 nm. ROS levels were estimated from the relationship obtained between HyPer-YFP

excited at 488 nm and 405 nm (Love et al. 2013).

Uncaging of Rubi-glutamate

Rubi-glutamate was added to the pipette solution at a final concentration of 300 µM.

Procedures were carried out considering the high sensibility to light of the caged compound

(Fino et al. 2009). To prevent spontaneous activation of Rubi-glutamate, glass pipettes were

coated with beeswax. The recording electrode was positioned above the lateral region of the

glomerular layer using dim transmitted light. Next, the GFP labelled glomerulus was targeted

using blue light attenuated with a ND filter (0.5 OD). Approximately 1 µL of the pipette

solution was injected in the glomerulus in the absence of light. Upon certifying that the

recording of LFP signal was stable for 2 min, a TTL signal opened a shutter (Lambda SC,

Sutter Instrument, Novato, CA) for 500 ms to deliver blue light (470 ± 20 nm) through the

epifluorescence port. A diaphragm restricted light application to the region targeted by the

electrode.

Histological procedures

Tadpoles were fixed for immunohistochemistry in 4% PFA during 4-7 days at 4ºC

and immersed in 30% sucrose. Animals were next embedded in O.C.T. freezing medium

(Tissue-Tek®, Sakura Finetek, Zoeterwoude, the Netherlands), snap-frozen in isopentane in a

Bright Clini-RF rapid freezer and stored at -80ºC until use. Sagittal sections (15 µm thick)

were obtained using a cryostat (Leica, Reichert-Jung, Heidelberg, Germany) and mounted on

superfrost plus slides (VWR International GmbH, Barcelona). For immunofluorescent

staining the sections were blocked for 2 hr at room temperature with PBS solution containing

0.2% Triton X-100 and 10% NGS. They were next incubated in a moist chamber overnight

at 4ºC in PBS with 0.2% Triton X-100 and 2% NGS containing anti-tyrosine hydroxylase

(mouse monoclonal, Immunostar cat. no. 22941, 1:250; RRID:AB_57226) and anti-GFP

(rabbit polyclonal, A6455, Invitrogen, 1:300, RRID: AB-221570). After three washes with

PBS, sections were incubated with appropriate secondary antibodies followed by DAPI

staining (1:10000) and mounted with fluoromount (Sigma-Aldrich, Sant Louis, MO). Some

sections were obtained from tadpoles whose nasal cavities were injected with CM-DiI

(Thermofisher Scientific, Walthman, MA, cat. no. C7001) 24h before fixation.

Statistical analysis

For statistical analysis, two tailed paired and unpaired t-test were used to evaluate

differences between two experimental groups. Comparisons among three or more groups

were performed using one way ANOVA followed by multiple comparison Tukey's HSD test.

Average values are expressed as mean±s.e.m. Statistical analysis was carried out using Igor

Pro software (RRID:SCR_000325).

RESULTS

Characterization of odor-evoked responses in a genetically defined glomerulus

The *Xenopus tropicalis* line *Dre.mxn1:GFP* allows the identification of an olfactory

glomerulus located laterally and innervated by a discrete population of OSNs (Terni et al.

2017; Terni and Llobet 2021). We targeted an electrode to the left GFP labelled glomerulus

and recorded changes in the LFP evoked by ipsilateral stimulation with an amino acid acting

as a waterborne odorant (Figs. 1A and B). An olfactory response was triggered by 100 ms

application of 200 µM methionine to the olfactory epithelium, thus confirming the ability of

Xenopus tadpoles to detect amino acids (Manzini et al. 2007). Stimulation evoked a negative

deflection of the LFP that resembled responses obtained in the glomerular layer of rats upon

sniffing odors (Chaigneau et al. 2007). A main difference between negativities recorded in *X.*

tropicalis and those found in rodents is that the latter are respiration-locked, while in tadpoles

there was a single deflection that recovered with a half time ranging from 1 to 4 s.

An estimate of the spatial extent of the LFP signal was obtained by evaluating

responses after changing the position of the recording electrode. The characteristic profile of

LFP changes disappeared when the pipette was directed to the mitral cell layer (Fig. 1C), thus

illustrating they were confined to the glomerular layer. The consistent success rate (85%,

n=269) obtained by targeting the GFP labelled glomerulus with the recording electrode

decreased to 50% (n=36) when random locations linearly spaced by 50 µm were tested within

the glomerular layer. The example shown in Fig. 1D illustrates how the characteristic

response to methionine was obtained only in one out of four positions tested. These

observations could be explained by the capacity of the electrode to detect activated glomeruli

(Manzini et al. 2007). The high success rate of the recordings guided by fluorescence support

that the GFP labeled glomerulus responded to methionine. Considering many olfactory

glomeruli in *Xenopus* tadpoles are broadly tuned (Manzini et al. 2007), it is conceivable that

the recorded glomerulus was also activated by other waterborne odorants, which were not

tested in the current study.

The LFP signal likely sampled the entire volume of 15765 ± 2119 µm^3 (n=33) defined

by fluorescence but, as the lateral cluster of glomeruli is markedly involved in the detection

of amino acids (Weiss, Manzini, and Hassenklöver 2021), the influence of adjacent glomeruli

could not be ruled out from the start. Considering the labelled glomerulus had a diameter of

~30 µm (Fig. 1E) and there was a 50 µm axial resolution to obtain independent readouts (Fig.

1D), only a portion of surrounding glomeruli could theoretically participate in the observed

LFP signal. The glomerular layer of the amphibian olfactory bulb is loosely organized

compared to mammals, as glomerular units show different sizes and lack ensheathing

astrocytes (Nezlin and Schild 2000; Gaudin and Gascuel 2005). The absence of a

homogeneous distribution of glomeruli, added to the dorsorostral location of GFP labelled

axon terminals (Fig. 1E), restricted the number of adjacent glomeruli that could theoretically

contribute to the recorded LFP signal to those found ventrally or caudally. Even if 50% of

such units were activated by methionine, the decay of the LFP following a relationship

inversely related to distance to the recording site, as reported for the mammalian olfactory

bulb (Karnup et al. 2006), would minimize their contribution. Altogether indicates that

methionine application to the olfactory epithelium reliably activated the lateral glomerulus

labeled by GFP in the *Dre.mxn1:GFP* line.

Glomerular responses were assayed by repetitive stimulation at two-minute intervals

and all negativities showed a comparable profile (Fig. 1B). The peak of the negative

deflection was reached ~1 s after olfactory stimulation. The characteristic LFP response of

any given tadpole was obtained by averaging 5 to 12 consecutive stimulations. Local

application of 100 μM CNQX decreased LFP negativities by approximately 50% (Fig. 2A).

The reduction was comparable when 100 μM D-AP5 was used instead of CNQX (Fig. 2B).

The concomitant application of both drugs produced an additive effect that reduced responses

by 70% (Fig. 2C p=0.0004, paired t-test), thus evidencing their synaptic origin, as well as the

involvement of AMPA and NMDA receptors. These results are comparable to those obtained

in rats (Chaigneau et al. 2007; Lecoq, Tiret, and Charpak 2009), which validates our

experimental configuration to record LFP responses in a region defined by a genetically

labelled glomerulus.

The onset phase of negativities was particularly sensitive to CNQX and AP5,

suggesting that it was reflecting the involvement of receptors activated by glutamate release

from OSN axon terminals. This possibility was tested by inducing a jump in glutamate

concentration inside the glomerulus. We targeted the electrode to the lateral GFP

glomerulus, injected *Xenopus* ringer solution containing 300 μM Rubi-glutamate and

uncaged it using a flash of blue light (Fino et al. 2009). A transient negativity was induced by

a 500 ms pulse of light, which was reproduced by repeated stimulations (Fig. 2D). The

average amplitude of LFP_{peaks} was of 30±3 μV (n=21, 3 tadpoles). This observation

supported that the initial phase of LFP negativities was caused by glutamate secreted by OSN

axon terminals that converged in a glomerulus.

The contribution of inhibitory neurotransmission mediated by $GABA_A$ receptors was

negligible because no variations in LFP changes evoked by methionine were observed after

local application of 1 mM picrotoxin (Fig. 2E). To certify that ipsilateral stimulation of

OSNs was the trigger of recorded responses, we investigated the effect of methionine

application to the contralateral epithelium. Negativities disappeared (Fig. 2F), thus

demonstrating that glomerular activation was exclusively unilateral. In 37% of the tadpoles

studied (n=174) the negativity was preceded by a transient positivity (Fig. 1B).   When

present, this LFP deflection was not modified by glutamate and GABA$_A$ receptor antagonists

(Figs. 2A and E), thus indicating it was unrelated to synaptic mechanisms.   As the positivity

was caused by the application of olfactory stimuli and always occurred before the synaptic

response, it is possible to consider its origin in cellular structures conveying peripheral

information to the olfactory bulb such as the layer of nerve fibers.

Glomerular responses depend on the number of input neurons.

The amplitude of LFP negativities decreases during the reinnervation of the

glomerular layer after olfactory nerve transection (Terni et al. 2017), which suggests that LFP

changes are related to the number of OSNs axons projecting to glomeruli.   If odor-evoked

responses were indeed related to glomerular input, the smallest negativities should take place

in the earliest developmental stages.   This possibility was confirmed by observing that GFP

positive neurons found in the olfactory epithelium increased with normal development and

their number followed an empirical linear relationship related to olfactory nerve width for

tadpoles found between NF stages 47 and 54 (Fig. 3A, r^2=0.96).   The peak of the LFP

response also varied related to olfactory nerve width but followed an exponential fit.

Negativities reached a steady-state value (Fig. 3A), thus suggesting the achievement of a

consolidated glomerular activation in this developmental time interval.   The presence of

constant glomerular responses for NF stages >50 coincides with the consolidation of the

basic structure of the olfactory bulb, which remains constant through larval life (Byrd and

Burd 1991).   LFP changes were empirically described by the function $LFP_{peak}=LFP_{steady\text{-}}$

$_{state}+Ae^{-r \cdot ONW}$ that associated the maximum change in the LFP (in μV) to olfactory nerve

width (ONW, expressed in μm). The fit revealed an increase rate (r) of 0.13 μm^{-1} starting at a

diameter of 21 μm, which corresponds to the thinnest olfactory nerves present during

development around NF stage 40. Considering the linear correlation found between the

number of GFP positive OSNs and the width of olfactory nerves, it was possible to rewrite

the exponential fit as a function of the number of OSNs. The estimated peak negativity in the

LFP could thus be obtained according to $LFP_{peak}=108-1866 \cdot e^{-0.11* \cdot OSNs}$ for a given number of

OSNs in the olfactory epithelium. On average, an olfactory epithelium contained 30 ± 2 GFP

positive OSNs, ranging from 24 to 39 labelled neurons (n=44) and, this relationship provided

a quantitative association of the amplitude of odor-evoked responses to the insertion of OSNs

in a glomerulus during normal development.

Glomerular responses are potentiated by the injury of the contralateral olfactory

pathway.

Transection of both olfactory nerves causes a silencing of the capacity of *X. tropicalis*

tadpoles to perceive odors (Terni et al. 2017). Since animals having a single nerve still

display normal olfactory guided behavior, we investigated how information is transmitted in

the absence of the mirror pathway. Olfactory nerve transection caused a potentiation of

evoked LFP negativities in the recorded contralateral olfactory glomerulus (Fig. 3B) that was

already observed two hours after injury (Fig. 3C). All tadpoles investigated showed odor-

evoked responses above the expected LFP_{peak} level for the range of developmental stages

analyzed. By reaching a maximum potentiation between 24 and 48 hours after transection,

LFP_{peak} responses returned to control values following an exponential recovery phase that

took place with a time constant of 100 hours (Fig. 3D). The overall duration of potentiation

coincided with the approximately two weeks required to reform the glomerular layer upon

olfactory nerve transection (Terni et al. 2017), suggesting that the increase of glomerular

output occurred while the circuitry of the contralateral olfactory bulb was being remodeled

after injury.

LFP$_{peak}$ responses had a strong synaptic contribution because they were dependent on

the number of OSNs innervating the glomerulus (Fig. 3A), were sensitive to the application

of glutamate receptor blockers (Figs. 2A-C) and could be triggered by glutamate uncaging

(Fig. 2D). These results complement those obtained in rats showing a correlation of the

amplitude of LFP negativities to calcium influx in OSN terminals (Lecoq, Tiret, and Charpak

2009) and EPSPs recorded in mitral cells (Chaigneau et al. 2007). We thus hypothesized that

the observed potentiation arises from glutamatergic synapses mediating glomerular

activation.

Tissue repair mechanisms are not involved in the potentiation of odor-evoked responses

Since inflammatory mediators can affect synaptic functions (Wu et al. 2015), we

evaluated the involvement of molecules released by injury using two different strategies.

First, to observe if sensory nerve damage were enhancing LFP$_{peak}$ responses, we sectioned

both optic nerves (Fig. 4A). Even though optic and olfactory nerves show substantial

functional and anatomical differences, the nature of the injury, as well as the distance to the

recorded site was comparable between optic and olfactory nerve transection. On these bases

we assumed that the effect generated by diffusible inflammatory mediators, or the capacity to

mobilize cells mediating inflammation, should be comparable in both experimental

conditions. LFP negativities recorded 24h to 48h after bilateral optic nerve transection

showed an amplitude of 86±6 μV (n=17). This value was expected by normal development

but was significantly lower (p=1.92·10^{-5}, unpaired t-test), than the amplitude of 186±16 μV

(n=14) found in tadpoles subjected to transection of the contralateral olfactory nerve (Fig.

4B).

Secondly, we investigated the release of reactive oxygen species (ROS), which are

key to trigger tissue repair mechanisms in tadpoles (Love et al. 2013). The *Xenopus laevis*

line HyPer-YFP displays a ubiquitous expression of the H_2O_2 sensor HyPerYFP that allows

the in vivo detection of ROS created by wounds (Love et al. 2013; Niethammer et al. 2009).

Two hours after unilateral olfactory nerve transection there was an increase in H_2O_2 levels

that was restricted to the cells present at the injury site (Fig. 4C). ROS levels remained

unaltered in the olfactory epithelium and in both olfactory bulbs. Moreover, the amplitude of

odor-evoked changes in the LFP were unaffected by the presence 200 μM apocynin or 2 μM

diphenyleneiodonium, two blockers of ROS production (Fig. 4D, p=0.36, ANOVA followed

by Tukey's test) that block tail regeneration at these concentrations (Love et al. 2013). These

findings together suggested that the enhancement of glomerular responses was unlikely

caused by inflammatory mediators or the activation of fundamental tissue repair mechanisms

present in tadpoles and, revealed the presence of an intrinsic compensatory mechanism

operating in the olfactory system.

Potentiation of glomerular responses is of presynaptic origin

The activation of individual glomeruli was imaged in the transgenic *X. tropicalis* line

tubb2b:GCaMP6s to gain insight on a putative synaptic mechanism acting on the bilateral

balance of odor-evoked responses. Ipsilateral application of 200 μM methionine to the

olfactory epithelium activated a defined set of glomeruli, which were identified as round, 10-

20 μm diameter structures. Same glomeruli reacted to three or more consecutive methionine

stimulations (Fig. 5A). Responses detected as calcium transients showed comparable

amplitudes and temporal profiles (Fig. 5B), suggesting that the simultaneous activation of

sparse glomeruli contributed to the elaboration of an olfactory map. Two lines of evidence

support that calcium increases originated in presynaptic terminals of OSNs. First, because

the *tubb2b* promoter primarily targets OSNs over olfactory bulb interneurons. A detailed

characterization carried out in *X. laevis* tadpoles shows that *tubb2b* efficiently drives

expression of a genetically encoded fluorescent reporter in OSNs but fails to label

periglomerular neurons positive for calretinin or tyrosine hydroxylase and only targets 24%

of mitral cells (Daume et al. 2022). The convergence of multiple fluorescent OSN axon

terminals on a single glomerulus and a minimal presence of labelled juxtaglomerular and

mitral cells, makes *X. tropicalis tubb2b:GCaMP6s* tadpoles well-suited to detect the

presynaptic component of glomerular responses. Second, because the simultaneous imaging

of glomerular activation and recording of LFP negativities revealed a temporal correlation of

the two signals (Fig. 5C). This observation matches the findings of a previous study carried

out in rats, where a comparable relationship was described between responses detected in

OSN axon terminals labelled with a high affinity calcium dye and LFP signals (Lecoq, Tiret,

and Charpak 2009). Although tadpoles of the *tubb2b-GCaMP6s* line consistently showed

glomeruli activated by methionine, the coupling of imaging to LFP signals had a low success

rate. Even though the electrode was targeted to the lateral glomerular cluster, which is the

region expected to concentrate responses to amino acids (Weiss, Manzini, and Hassenklöver

2021), LFP negativities were detected in only 33% of animals tested (n=17/52). This figure

is below the 50% found after the performance of recordings in random locations of the

glomerular layer (Fig. 1D) but it is higher than the ~25% success rate found in *X. laevis*

tadpoles (Manzini et al. 2007). This observation reinforces the selectivity of LFP recordings

to detect the output of a single glomerulus, which in this set of experiments was coupled to

the fluorescent response found in closest apposition to the tip of the pipette. The readouts of

glomerular input and output were thus reported by transient increases in GCaMP6s

fluorescence and LFP signals, respectively.

A hallmark of contralateral olfactory nerve transection was the development of larger

LFP negativities with a faster onset. The time constant to reach the LFP_{peak} shortened

significantly (p=0.017, unpaired t-test) from 1.05±0.1 s (n=46 tadpoles) to 0.71±0.1 s (n=39

tadpoles), whilst the recovery phase was unaffected (Fig. 5D, p=0.79, unpaired t-test).

Changes in glomerular output likely had a presynaptic origin, because calcium transients

were larger and showed faster onset kinetics after contralateral nerve transection (Figs. 5E

and F). The average amplitude increased more than two-fold (p=0.02, unpaired t-test) and

the time constant describing the rise time shortened significantly (p=0.0003, unpaired t-test)

in tadpoles with a transected olfactory nerve compared to control animals (Fig. 5F). The

slow kinetics of GCaMP6s prevented a precise temporal association between calcium buildup

and the onset of LFP changes (Fig. 5C) and a certain saturation of the high affinity calcium

indicator in axon terminals was also expected. But, despite these limitations to quantitative

analysis, the changes reported by GCaMP6s showed an overall enhancement of cytosolic

calcium levels mediating neurotransmitter release. We next sought to investigate which

regulatory mechanisms acting on OSN axon terminals were under bilateral control to alter

glomerular output.

Presynaptic inhibition driven by dopamine D_2 receptors is affected by contralateral

glomerular input

Presynaptic inhibition regulates neurotransmission in olfactory glomeruli (McGann

2013). In mice, discrete populations of juxtaglomerular neurons that release GABA or

dopamine activate $GABA_B$ or D_2 receptors present in presynaptic terminals of OSNs to lower

glutamate release (Wachowiak et al. 2005; Wachowiak and Cohen 1999; Ennis et al. 2001).

Evidence points out that this is a tonically active mechanism (Pírez and Wachowiak 2008),

thus meaning the normal processing of odor information relies on a certain level of inhibition

of glomerular output. Since several previous works identified neurons immunoreactive for

tyrosine hydroxylase or GABA innervating the glomerular layer of *X. laevis* tadpoles

(González et al. 1994; Daume et al. 2022; Nezlin and Schild 2000), dopamine and/or GABA

could also be mediating glomerular inhibition in *X. tropicalis* tadpoles.

404   We identified TH positive neurons (TH +) at the border of the glomerular and the

mitral cell layers (Fig. 6A). The morphological characteristics of these neurons were

reminiscent of type-1 TH neurons described in the olfactory bulb of adult frogs, associated

with a dopaminergic phenotype (Boyd and Delaney 2002). The processes of TH+ neurons

ramified within the glomerular layer suggesting, the establishment of relationships to several

glomeruli (Fig. 6B). The lateral glomerulus formed by GFP positive axon terminals was

contacted by TH+ puncta, thus indicating its function was modulated by dopamine signaling.

412   Odor-evoked responses were enhanced by raclopride, a D_2 antagonist, in tadpoles

with intact olfactory pathways (Fig. 7A). The average negativity of 73 ± 5 µV (n=8)

experimented a gradual increase in the presence of 300 nM raclopride that reached 128 ± 16

µV, 20 min after its application (Fig. 7B, p=0.012, paired t-test). The potentiation of odor-

evoked responses caused by the D_2 antagonist was not observed in tadpoles with the

contralateral olfactory nerve transected. Baseline negativities of 133 ± 24 µV (n=11) were

minimally increased by raclopride, changing to 156 ± 33 µV (n=11, p=0.16, paired t-test), 20

min after drug application.

420   The time course of the increase in LFP_{peaks} observed in control tadpoles was well

described by a Hill equation, reporting a time required to reach the 50% of the maximal

increase of approximately 20 min (Fig. 7C). The time window describing the effect of

raclopride could be attributed to disinhibition, since the primary target of presynaptic D_2

receptor activation is the decrease of cAMP levels and a consequent lowering of

neurotransmitter release (Kaneko and Takahashi 2004). The potentiation mediated by

raclopride was comparable to the increase in LFP_{peaks} caused by olfactory nerve transection

(Fig. 7B, p=0.46, unpaired t-test). These results indicated that the tonic inhibition of

glomerular output was comparably reduced either by directly inhibiting D_2 receptors, or, by

surgically transecting the contralateral olfactory nerve. Further evidence for the involvement

of D_2 receptors was obtained by simultaneously imaging GCaMP6s fluorescence and LFP

signals (Fig. 7D). In the illustrated example raclopride shortened the time constant (τ) to

reach the calcium peak from 1.3 s to 0.5 s and potentiated LFP changes by 50%, thus

reproducing the effect of contralateral olfactory nerve transection on glomerular input and

output, respectively (Figs. 3C and 5F). On average, raclopride reduced the onset time

constant of calcium transients from 1.27 s to 0.8 s (n=4 tadpoles, p=0.02, paired t-test),

evidencing that the potentiation of glomerular responses was mediated by a faster buildup of

presynaptic calcium levels that took place after antagonizing the inhibitory action of D_2

receptors.

To explore a concomitant inhibition mediated by presynaptic $GABA_B$ receptors

(McGann 2013) we investigated the effect of the selective $GABA_B$ antagonist CGP-36742.

LFP_{peaks} remained unchanged (Figs. 7E and F, p=0.77, paired t-test). A significant inhibition

by GABA was thus ruled out, revealing dopamine was the main neurotransmitter inhibiting

glomerular output in the glomerular layer of *Xenopus* tadpoles.

Effect on glomerular responses of the partial elimination of mirror olfactory sensory

neurons.

To shed light on the involvement of chemotopy in the contralaterally driven

potentiation of glomerular output we modified the two-photon chemical apoptotic targeted

ablation (2Phatal) technique (Hill et al. 2017) to eliminate groups of GFP positive OSNs

using a conventional confocal microscope (Fig. 8A). Tadpoles were placed during 15

minutes in *Xenopus* water containing 5 µg/mL Hoechst 33342, which is a membrane

permeable dye with affinity for nucleic acids, to label all cells found within the olfactory

epithelium. Damage was exerted by photobleaching groups of cells found in regions of

interest (ROIs) that contained two or more GFP positive OSNs. Efficient photobleaching of

the nuclear label caused cell death, which was certified by the observation of condensed

nuclei after 24 hours (Fig. 8A). The use of the confocal microscope precluded the

achievement of single cell ablation but supported the robust elimination of cells within ROIs.

Olfactory epithelial cells found outside the selected areas remained unaffected.

Photobleaching was typically carried out in 4 to 6 ROIs, thus resulting in the elimination of

10 to 15 OSNs. Considering an epithelium contained on average ~30 GFP positive OSNs,

the protocol led to an estimated 30 to 50% reduction of OSNs innervating the right GFP

labeled glomerulus. Since the viability of tadpoles subjected to a more extensive reduction of

fluorescent OSNs was compromised, all animals investigated still had a significant number of

functional GFP labelled OSNs after photoablation.

A characteristic of the recorded odor-evoked glomerular responses was their

reproducibility. For a given tadpole, LFP negativities occurred after successive stimulations

with a comparable profile (Figs. 1B, 2A, 4A), and showed an average variance in their

amplitude of 375 ± 92 μV^2 (n=17). Variance of the responses obtained increased more than

three-fold in tadpoles subjected to transection of the olfactory nerve (see for example Fig.

3B), reaching an average value of 1488 ± 280 μV^2 (n=14). An enhancement of variance was

thus associated to the potentiation observed in response to olfactory nerve transection.

Incubation of tadpoles with Hoechst 33342 neither modified the amplitude nor the variance

of LFP_{peaks} (Fig. 8B). LFP negativities in tadpoles presenting a decimated population of

contralateral GFP positive OSNs showed an amplitude of 78 ± 7 μV (n=14) and a variance of

771 ± 351 μV^2, both being comparable to controls (Fig. 8C). These values were also similar to

tadpoles where regions lacking GFP positive OSNs were photobleached.

These results suggest that potentiation driven by damage of contralateral OSNs was

unrelated to chemotopy, which could be consistent with the innervation of several glomeruli

by a single TH+ neuron (Boyd and Delaney 2002). However, a reduction between 1/3 and

1/2 of mirror contralateral input might be insufficient to drive a significant increase in

LFP_{peaks} amplitude or variance, because only the complete ablation of GFP labelled neurons

could replicate the effect of transection. Consequently, it was not possible to rule out that the

gain of function caused by contralateral injury was related to a graded contribution of each

OSN to the total input of topographically related glomeruli.

Pallial neurons are involved in the potentiation of glomerular responses induced by

contralateral injury.

To shed light on the circuit mediating the contralaterally driven potentiation of

glomerular responses we considered three theoretical scenarios. First, a direct bilateral

innervation of glomerular layers by OSNs. Second, an interhemispheric association of

glomeruli mediated by olfactory bulb projection neurons. And third, the involvement of the

lateral pallium, which is considered to play a key role in olfactory processing in amphibia

(Moreno et al. 2008; Roth et al. 2007).

The axons of OSNs labelled in the *Dre.mxn1:GFP X. tropicalis* formed the lateral

glomerulus and showed some projections to medial glomerular structures (Fig. 1E), but we

did not find evidence for crossings at the level of the anterior commissure or midline. The

possibility that GFP labelled neurons displayed the properties of those innervating the γ-

glomerulus (Kludt et al. 2015) was thus ruled out. In premetamorphic stages most OSNs

project to ipsilateral glomeruli (Weiss et al. 2021) and, considering the lack of support for

interhemispheric connections among olfactory bulb neurons as reported in zebrafish (Kermen

et al. 2020), we thus evaluated the third possibility by investigating the capacity of pallial

neurons to respond to odor stimulation.

Regions of interest measuring 30 μm diameter were selected in the dorsolateral

pallium of *tubb2b:GCaMP6*s tadpoles. All of them displayed synchronous, rhythmic calcium

transients that occurred at frequencies ranging from 0.1 to 0.5 Hz. Stimulation of the

ipsilateral olfactory epithelium with 200 μM methionine evoked a response in all selected

regions that was consistently observed during consecutive stimulations (Fig. 9A). These

observations confirmed the involvement of the dorsolateral pallium in the processing of

olfactory information.

If pallial neurons were participating in the potentiation of contralateral glomerular

output, their damage should induce a similar gain of function to that evoked by olfactory

nerve transection (Fig. 3A). Using iridectomy scissors we made an injury at the level of the

dorsal pallium in the medial to the lateral direction and recorded odor-evoked responses 1 to

2 days afterwards (Fig. 9B). A ~70% increase in odor-evoked responses was observed. The

average amplitude of LFPpeaks was of 135±11 μV (n=25), significantly higher than

responses found in control tadpoles (Fig. 9C, p=6·10^{-5}, unpaired Student's test). These results

showed that pallial neurons participated in the control of glomerular output and suggested

that they could modulate the activity of contralateral dopaminergic neurons mediating

presynaptic inhibition of OSN axon terminals.

DISCUSSION

Odor-evoked responses were recorded as changes of the LFP in an olfactory

glomerulus labelled in the *Dre.mxn1:GFP X. tropicalis* line (Fig. 1) and showed the

following characteristics: *i)* were mediated by glutamatergic synapses (Figs. 2A-C), *ii)* were

triggered by ipsilateral stimulation of OSNs (Fig. 2F), and, *iii)* showed an amplitude related

to the number of input OSNs (Fig. 3A). These characteristics are comparable to responses

recorded in the glomerular layer of living rats where negativities are supported by the

activation of glutamate receptors and locked to the respiration frequency (Lecoq, Tiret, and

Charpak 2009). Glomerular responses were potentiated by the complete silencing of OSNs

projecting to the contralateral olfactory bulb, thus showing the processing of information in

the olfactory glomeruli of *Xenopus* tadpoles is not exclusively unilateral and is shaped

contralaterally.   The observed potentiation was not related to inflammatory mediators

associated to injury, because it was caused by a release of the inhibition made by D_2

dopamine receptors present in OSN axon terminals.

The similarities between the recordings obtained (Fig. 1B) and those described in rats

(Chaigneau et al. 2007), indicate the presence of evolutionary conserved morphological and

functional features between both species and therefore, the applicability to vertebrates of the

findings here described. Our results achieved using glutamate blockers and uncaging of

Rubi-glutamate show that the onset of LFP changes detected in glomeruli is determined by

glutamate release from OSNs. The ionic bases supporting the recovery phase of negativities

are uncertain but, considering responses obtained by glutamate uncaging were locked to the

period defined by the light pulse, the participation of sustained neurotransmitter release is

conceivable. This view is also supported by imaging of OMP-synaptopHluorin mice, where

continuous synaptic vesicle exocytosis occurs over several seconds in OSN axon terminals

(Petzold et al. 2008).

The mammalian and amphibian olfactory bulbs display comparable anatomical and

cellular organization (Manzini, Schild, and Di Natale 2022) but, it is yet unknown whether

the three classes of juxtaglomerular neurons present in rodents: periglomerular, short-axon

and external tufted, also exist in *Xenopus*. In mice, short-axon cells are the main source of

dopamine to olfactory glomeruli (Kiyokage et al. 2010), however, there is a lack of evidence

for an equivalent neuronal type in the olfactory bulb of *Xenopus*. Here, we show that the

processes of TH + neurons extensively innervate the glomerular layer, thus indicating a major

role for dopaminergic signaling, in agreement with morphological observations (González et

al. 1994). Our results indicate that a major role of dopamine is mediating the presynaptic

inhibition of OSN axon terminals through D2 receptors. In rodents, presynaptic inhibition of

OSNs is mediated by GABA acting through $GABA_B$ receptors and dopamine (McGann 2013)

and both neurotransmitters are typically co-released by specific types of juxtaglomerular

neurons (Kosaka, Pignatelli, and Kosaka 2020). Since we did not find evidence for the

participation of $GABA_B$ receptors in the inhibition glomerular responses, the data obtained

support a segregation of dopaminergic and gabaergic pathways, in agreement with

morphological evidence (Boyd and Delaney 2002).

To understand how tonic dopamine release in the glomerular layer of *Xenopus*

tadpoles is affected by the number of contralateral OSNs, it is necessary to have a

comprehensive understanding of the different types of interneurons present in the olfactory

bulb of *X. tropicalis* tadpoles, which is currently lacking. The dopaminergic neurons present

in the border of the glomerular and mitral cell layers of *X. tropicalis* tadpoles send their

projections to glomeruli, which fit with the characteristics of type-1 TH+ interneurons

described in adult frogs (Boyd and Delaney 2002). The dendrites of type-1 TH+ interneurons

innervate glomeruli and when type-1 TH+ neurons contain an axon, it recurves and projects

to the glomerular layer. In mice, there are five different types of morphologically distinct

juxtaglomerular dopaminergic neurons (Kosaka, Pignatelli, and Kosaka 2020), and, those that

are anaxonic release dopamine from their dendritic tree, while those that are axon bearing

release it from their axon (Dorrego-Rivas et al. 2025) Considering that an axon was not

observed in all type-1 TH+ interneurons (Boyd and Delaney 2002), it is conceivable that

axon-bearing and anaxonic dopaminergic neurons could also exist in *Xenopus* tadpoles and

both innervate glomeruli. If the majority of type-1 TH+ were spontaneously active, as

reported in mice for dopaminergic juxtaglomerular neurons (Pignatelli et al. 2005), dopamine

would thus play a key modulatory role of glomerular neurotransmission in *X. tropicalis*

tadpoles.

Our results support that dopamine release is affected by the number of operative

contralateral OSNs so that, in the extreme situation where all mirror olfactory pathways are

silenced, dopamine secretion decreases and most of the inhibition is released (Fig. 9D). The

consequence is an immediate increase of glomerular responses in the non-damaged pathway,

which results in a compensated output signal. The lateral pallium as it is a main center for

olfactory processing in frogs (Moreno et al. 2008) and we provide evidence for its

participation in the bilateral homeostasis of glomerular output. The injury of contralateral

pallial neurons evoked a potentiation of glomerular responses that reproduced the effect of

olfactory nerve transection, which supports a regulation of dopamine release by the pallium.

Considering the capacity of pallial neurons to establish ipsilateral and contralateral

relationships (Roth et al. 2007; Westhoff and Roth 2002), our results are consistent with the

presence of a pallial connectivity controlling the firing of TH+ neurons that innervate the

glomerular layer of the olfactory bulb.

The in vivo evidence presented shows that presynaptic D_2 receptors tonically exert an

inhibitory action on calcium buildup in OSN axon terminals. This finding is well aligned

with previous observations obtained in mouse brain slices (Ennis et al. 2001) and describes a

relevant biological scenario for the role of dopamine inhibition in the glomerular layer of the

olfactory bulb. The likely target of dopamine are N-type voltage gated calcium channels

because they are present in presynaptic terminals in olfactory glomeruli (Weiss et al. 2014)

and they can be inhibited through multiple mechanisms activated by D_2 receptors (Kisilevsky

and Zamponi 2008).

Altogether, our results illustrate a homeostatic mechanism used to compensate a

reduced contribution of contralateral OSNs via an enhancement of glomerular output. It is

possible that the observed compensation upon transection of an olfactory nerve takes

advantage of dopaminergic signaling pathways used for encoding of olfactory information.

The normal activity of presynaptic D_2 receptors present in OSN axon terminals could be

contralaterally regulated through a homeostatic loop controlled by pallial neurons to correct

for imbalances in glomerular input. Considering the evolutionary conserved features of the

*Xenopu*s olfactory bulb, the mechanism here described could be broadly applicable to

vertebrates.

FIGURE LEGENDS

Figure 1. Recording of odor-evoked responses in a genetically defined glomerulus. A)

Schematic diagram illustrating the experimental preparation. A puff of 200 µM methionine

was applied for 100 ms to the left olfactory epithelium of *Dre.mxn1:GFP Xenopus tropicalis*

tadpoles. Changes in the Local Field Potential (LFP) were measured ipsilaterally with an

electrode targeted to the lateral glomerulus formed by axon terminals of olfactory sensory

neurons (OSNs) expressing GFP. The inset shows a confocal projection illustrating labelled

OSNs (s: soma, a:axon) and the bilateral formation of glomeruli (g). The *Dre.mxn1* promoter

also drives GFP expression in the endothelial cells of some blood vessels (e). **B)**

Representative glomerular odor-evoked response (black) obtained by averaging individual

responses (gray) of the LFP following stimulation (arrow). The asterisk shows a positivity

that was evident in 37% of the recordings and preceded the characteristic negativity

associated with glomerular activation. **C)** Negative deflections of LFP (individual responses,

gray; average trace, black) were observed when the recording electrode was placed in the

GFP labelled glomerulus but disappeared in the mitral cell layer (ML, red traces; GL,

glomerular layer). **D)** Experiment showing LFP recordings performed in four different

locations of the glomerular layer spaced by 50 µm. The characteristic odor-evoked response

was observed in only one of the positions tested. The representative glomerular odor-evoked

response (black) was obtained by averaging individual responses (gray) following stimulation

with methionine (arrow). **E)** Hyperstack projections of the left olfactory bulb of three

different *Dre.mxn1:GFP X. tropicalis* tadpoles. The lateral glomerular cluster (L) is always

evident, and some medial projections (M) are apparent in two of the illustrated examples.

The color scale indicates dorsoventral disposition.

Figure 2. Odor-evoked responses are mediated by glutamatergic neurotransmission. A)

The representative glomerular odor-evoked response (black) was obtained by averaging

individual responses (gray) of the Local Field Potential (LFP) following ipsilateral

stimulation of the olfactory epithelium using 200 μM methionine (arrow). In this example,

the pipette solution contained 100 μM CNQX and upon local injection of 1 μL, there was a

reduction in the amplitude of LFP negativities. **B)** Mean LFP changes obtained under control

conditions (n=21) were reduced after the application of 100 μM CNQX (n=8), 100 μM AP5

(n=5), or both 100 μM CNQX and 100 μM AP5 (n=8). **C)** Box plot illustrating how the

initially recorded peak negativities were affected by the application of 100 μM CNQX, or,

100 μM AP5 together with 100 μM CNQX. Boxes represent the median (horizontal line),

25th to 75th quartiles, and ranges (whiskers) of the indicated experimental groups. Statistical

differences were evaluated using paired t-test. **D)** Rubi-glutamate was injected into the GFP

labelled glomerulus and locally uncaged with a 500 ms pulse of blue light (circle). The

example shows the change in LFP induced by two flashes (squares) delivered at an interval of

9 seconds. **E)** Application of picrotoxin, a $GABA_A$ antagonist, did not modify odor-evoked

changes. Recordings show average responses (n=9). Statistical differences were evaluated

using paired t-test**. F)** Odor-evoked LFP changes were exclusively triggered by ipsilateral

stimulation. Individual responses are indicated in gray, with the representative average

response in black. Contralateral stimuli (yellow) did not modify the LFP, as shown in the

average representative response (brown).

Figure 3. **Potentiation of odor-evoked responses by transection of the contralateral**

olfactory nerve. **A)** The number of olfactory sensory neurons (OSNs) and the amplitude of

odor-evoked negative deflections of the Local Field Potential (LFP) were related to olfactory

nerve width according to linear and exponential functions, respectively. Individual data points

are represented by circles (n=48). Each bin indicates the mean ± s.e.m. of n=6 tadpoles. The

dotted line indicates the steady-state LFP amplitude reached during development. **B)**

Representative odor-evoked responses obtained in a control tadpole (black) and in a different

animal (red), 24 h after contralateral nerve transection. Gray traces indicate individual

responses to the application of 200 μM methionine solution (arrow). **C)** Odor-evoked LFP

changes exhibit amplitudes above the expected values (dotted line as in A) after contralateral

olfactory nerve transection at the indicated time points. The dots represent the mean ± s.e.m.

obtained 2 to 7 hours (n=10), 1 to 2 days (n=14), and 10 to 11 days (n=11) post-injury. There

was a 75% increase in animals recorded 1 to 2 days after transection of the contralateral

olfactory nerve (red arrow) compared to control tadpoles (dotted line). **D)** Dots (mean ±

s.e.m.) connected by a line illustrate odor-evoked glomerular responses at the indicated times

after injury. The superimposed violin plot displays individual data. Most LFP_{peak} values are

above the level expected for the developmental period studied (dotted line as in A).

Figure 4. The potentiation of odor-evoked responses is not mediated by injury derived

cues. **A)** Odor-evoked Local Field Potential (LFP) changes were recorded by an electrode

targeted to the GFP-positive glomerulus of *Dre.mxn1:GFP* tadpoles one to two days after

bilateral transection of optic nerves. **B)** Peak LFP negativities recorded in tadpoles with

sectioned optic nerves (n=18) did not exhibit the characteristic potentiation observed after

transection of the contralateral olfactory nerve (n=14), as they remained within the range of

values observed during normal development (solid line, as in Fig. 3A). Bins indicate

mean±s.e.m., circles show individual values. **C)** Imaging of reactive oxygen species (ROS)

two hours after transecting one olfactory nerve (arrow). The ratio between the fluorescence

emitted by HyPer-YFP when excited at 488 nm and 405 nm is indicated in pseudocolor.

Notice that ROS were increased at the injury site but remained at basal levels in both

olfactory bulbs as indicated by the box plot. Each circle shows values collected in a single

tadpole. **D)** Block of ROS production by incubating tadpoles with 200 µM apocynin (n=10)

or 2 µM diphenyleneiodonium (DPI, n=5) did not modify the amplitude of odor-evoked LFP

responses recorded 24 h after contralateral olfactory nerve transection (n=10). Boxes

represent the median (horizontal line), 25th to 75th quartiles, and ranges (whiskers) of the

indicated experimental groups.

Figure 5. The presynaptic component of glomerular activation is affected by damage to

contralateral olfactory sensory neurons. A) Application of a puff of 200 μM methionine to

the olfactory epithelium activates a set of glomeruli in the ipsilateral olfactory bulb (arrows)

of *tubb2b:GCaMP6*s tadpoles. Images show the relative changes in GCaMP6s fluorescence

(ΔF/F) obtained after two sequential stimulations carried out in a single tadpole. **B)** Time

course of the responses detected in the glomeruli indicated in A). **C)** An example showing

the simultaneous recording of Local Field Potential (LFP) and changes in GCaMP6s

fluorescence in the region targeted by the electrode. Colored traces and gray traces show the

change in GCaMP6s fluorescence (ΔF/F) and LFP respectively observed after three

sequential applications of 200 μM methionine. Black traces show the average ΔF/F and LFP

responses. **D)** Kinetics of the change in LFP observed in tadpoles with the contralateral

olfactory nerve transected between 2 h and 48h prior to recording. The differences are

illustrated by representative recordings obtained in two different tadpoles. **E)** Intracellular

calcium increases detected in glomeruli of control tadpoles with intact olfactory pathways (35

glomeruli, 10 tadpoles, black), and, in tadpoles subjected to the transection of the

contralateral olfactory nerve (10 glomeruli, 3 tadpoles, red). Each trace indicates the

response of a glomerulus to a single stimulus. Solid lines and error bars indicate mean ±

s.e.m. **F)** Calcium transients detected in tadpoles with an olfactory nerve transected showed a

larger amplitude and a rising phase with a shorter time constant (τ). Boxes in D) and F)

represent the median (horizontal line), 25th to 75th quartiles, and ranges (whiskers) of the

indicated experimental groups. Statistical differences in D) and F) were evaluated using

paired and unpaired t-tests, respectively. Circles in D) indicate tadpoles and in F) refer to

glomeruli.

Figure 6. Tyrosine hydroxylase positive neurons project to the glomerular layer of the

olfactory bulb. A) Cell bodies of neurons expressing tyrosine hydroxylase (TH+, magenta)

were sparsely distributed at the border of the glomerular (GL) and mitral cell (ML) layers of

the olfactory bulb and sent their neuronal processes (arrows) to innervate the glomerular layer

(asterisks). **B)** Projections of TH+ neurons (arrows) contacted axon terminals of olfactory

sensory neurons labelled with DiI (cyan, asterisks). **C)** The lateral glomerular cluster (L)

labeled in Dre.mxn1:GFP *Xenopus tropicalis* tadpoles was contacted by projections

(asterisks) of processes emerging from TH+ neurons (arrows).

Figure 7. Contralateral input modulates presynaptic inhibition mediated by dopamine

D_2 receptors and is involved in the potentiation of glomerular responses. **A)** Recordings

obtained in a control tadpole showing how the amplitude of Local Field Potential (LFP)

responses (gray traces) obtained during a baseline period of 8 minutes increased in a time-

dependent manner after local application of 300 nM raclopride, a D_2 receptor antagonist. **B)**

Box plot showing the effect of 300 nM raclopride (blue) on the amplitude of LFP responses

recorded in tadpoles with full capacity to process odors (control, n=8, black) and tadpoles

subjected to the transection of the contralateral olfactory nerve (n=11, red). Boxes represent

the median (horizontal line), 25th to 75th quartiles, and ranges (whiskers) of the indicated

experimental groups. The effect of raclopride was evaluated using the paired t-test and the

comparison between control and transected groups was performed using the unpaired t-test.

C) Relative change in LFP responses induced by 300 nM raclopride in control tadpoles and

in tadpoles subjected to the transection of the contralateral olfactory nerve. Dots represent

mean ± s.e.m. The solid black line illustrates the fit to a Hill equation, defining a T_{50} at 20

minutes. **D)** Simultaneous recording of LFP and changes in GCaMP6s fluorescence in a

tubb2b:GCaMP6s tadpole. Gray traces and light blue traces show individual responses to

sequential stimulations before and after application of 300 nM raclopride, respectively**.**

Average responses are shown in black and dark blue. **E)** Application of CGP-36742, a

GABA$_B$ receptor antagonist, did not modify LFP responses. The box plot compares the

amplitude of LFP changes recorded before (gray) and 20 min after local application of 300

μM CGP-36742 (green). Statistical differences were evaluated using paired t-test. **F)** Time

course of relative LFP changes induced by 300 μM CGP-36742. Dots represent mean ± s.e.m

(n=10). The inset shows recordings obtained in a tadpole in baseline conditions (gray) and

after injection of CGP-36742 (green).

Figure 8. Effect on odor-evoked responses of selective photoablation of olfactory

sensory neurons innervating the homologous contralateral glomerulus. A) Odor-evoked

Local Field Potential (LFP) changes were recorded one day after the selective elimination of

olfactory sensory neurons (OSNs) located in the right nasal cavity. After the identification of

GFP-positive OSNs in the epithelium labeled with the nuclear marker Hoechst 33342,

regions containing ≥2 fluorescent neurons were identified and photobleached. Cell targeting

was confirmed by the suppression of the nuclear label. Only cells found within the

photobleached areas exhibited fragmented nuclei (arrows) 24 hours after photobleaching. **B)**

Examples showing odor-evoked responses recorded in a tadpole incubated with Hoechst

33342 (black, average), and in a different tadpole 24 hours after the photobleaching of

selected regions in the contralateral olfactory epithelium (violet, average). **C)** Photoablation

of GFP positive neurons did not modify the amplitude or variance of contralateral odor-

evoked responses recorded in the glomerulus innervated by cognate neurons. Boxes

represent the median (horizontal line), 25th to 75th quartiles, and ranges (whiskers) of the

indicated experimental groups. Statistical differences were evaluated using ANOVA followed

by Tukey's test. Circles indicate values obtained from single tadpoles.

Figure 9. Pallial neurons are involved in the potentiation of glomerular responses driven

by contralateral injury. **A)** Rhythmic calcium transients were detected in six different

regions of interest (ROIs) located in the dorsolateral pallium of a *X. tropicalis*

*tubb2b:GCaMP6*s tadpole. Three consecutive stimulations carried out by applying of 200

µM methionine to the ipsilateral olfactory epithelium (dotted line) evoked a synchronous

response in the ROIs investigated. **B,C)** Odor-evoked responses were recorded 24-48 h after

making a tangential injury in the contralateral dorsolateral pallium. The amplitude of LFP

changes significantly increased in injured tadpoles. Boxes represent the median (horizontal

line), 25th to 75th quartiles, and ranges (whiskers) of the indicated experimental groups.

Statistical differences were evaluated using unpaired t-test. **D)** Model proposed for the

bilateral modulation of glomerular output in the olfactory bulb of *Xenopus* tadpoles. A

population of juxtaglomerular neurons releases dopamine to inhibit glomerular output by

activating presynaptic D2 receptors present in OSNs (dotted box). The constant presence of

dopamine within glomeruli is favored by the activity of the contralateral olfactory bulb.

When the contribution of the contralateral pathway is suppressed, dopamine release

diminishes, and glomerular responses become potentiated. The contralateral modulation of

the tonic activity of dopaminergic juxtaglomerular neurons corrects for input differences and

equalizes the synaptic output of olfactory glomeruli to achieve a bilaterally balanced transfer

of information. The activity of dopaminergic interneurons is likely controlled by pallial

neurons through a yet undetermined connectivity, taking advantage of their participation in

the processing of olfactory information.

ACKNOWLEDGEMENTS

This work was sponsored by the Ministry of Science, Innovation and Universities

(MICIU/AEI), grant PID2021-124536NB-I00 (A.L), co-funded by the European Regional

Development Fund (ERDF), "a way of making Europe". The work was also supported by

two Whitman Fellowships awarded to A.L in 2023 and 2024 (Marine Biological Laboratory,

University of Chicago). The authors thank the institutional support from the María de Maeztu

Unit of Excellence, Institute of Neurosciences, University of Barcelona, CEX2021-001159-M

(Ministry of Science, Innovation and Universities) and the CERCA Program of Generalitat de

Catalunya. A.L. is a Serra Húnter fellow. The authors thank Francisco Ciruela for suggesting

the use of raclopride.

REFERENCES

Belluscio, L., and L. C. Katz. 2001. "Symmetry, stereotypy, and topography of odorant representations in mouse olfactory bulbs." *J Neurosci* 21 (6):2113-22. doi: 10.1523/JNEUROSCI.21-06-02113.2001.

Boyd, J. D., and K. R. Delaney. 2002. "Tyrosine hydroxylase-immunoreactive interneurons in the olfactory bulb of the frogs Rana pipiens and Xenopus laevis." *J Comp Neurol* 454 (1):42-57. doi: 10.1002/cne.10428.

Browne, L. P., A. Crespo, and M. S. Grubb. 2022. "Rapid presynaptic maturation in naturally regenerating axons of the adult mouse olfactory nerve." *Cell Rep* 41 (10):111750. doi: 10.1016/j.celrep.2022.111750.

Byrd, C. A., and G. D. Burd. 1991. "Development of the olfactory bulb in the clawed frog, Xenopus laevis: a morphological and quantitative analysis." *J Comp Neurol* 314 (1):79-90. doi: 10.1002/cne.903140108.

Chaigneau, E., P. Tiret, J. Lecoq, M. Ducros, T. Knöpfel, and S. Charpak. 2007. "The relationship between blood flow and neuronal activity in the rodent olfactory bulb." *J Neurosci* 27 (24):6452-60. doi: 10.1523/JNEUROSCI.3141-06.2007.

Daume, D., T. Offner, T. Hassenklöver, and I. Manzini. 2022. "Patterns of." *Front Neuroanat* 16:914281. doi: 10.3389/fnana.2022.914281.

Dorrego-Rivas, A., D. J. Byrne, Y. Liu, M. Cheah, C. Arslan, M. Lipovsek, M. C. Ford, and M. S. Grubb. 2025. "Strikingly different neurotransmitter release strategies in dopaminergic subclasses." *Elife* 14. doi: 10.7554/eLife.105271.

Ennis, M., F. M. Zhou, K. J. Ciombor, V. Aroniadou-Anderjaska, A. Hayar, E. Borrelli, L. A. Zimmer, F. Margolis, and M. T. Shipley. 2001. "Dopamine D2 receptor-mediated presynaptic inhibition of olfactory nerve terminals." *J Neurophysiol* 86 (6):2986-97. doi: 10.1152/jn.2001.86.6.2986.

Fino, E., R. Araya, D. S. Peterka, M. Salierno, R. Etchenique, and R. Yuste. 2009. "RuBi-Glutamate: Two-Photon and Visible-Light Photoactivation of Neurons and Dendritic spines." *Front Neural Circuits* 3:2. doi: 10.3389/neuro.04.002.2009.

Gaudin, A., and J. Gascuel. 2005. "3D atlas describing the ontogenic evolution of the primary olfactory projections in the olfactory bulb of Xenopus laevis." *J Comp Neurol* 489 (4):403-24. doi: 10.1002/cne.20655.

González, A., O. Marín, R. Tuinhof, and W. J. Smeets. 1994. "Ontogeny of catecholamine systems in the central nervous system of anuran amphibians: an immunohistochemical study with antibodies against tyrosine hydroxylase and dopamine." *J Comp Neurol* 346 (1):63-79. doi: 10.1002/cne.903460105.

Hill, R. A., E. C. Damisah, F. Chen, A. C. Kwan, and J. Grutzendler. 2017. "Targeted two-photon chemical apoptotic ablation of defined cell types in vivo." *Nat Commun* 8:15837. doi: 10.1038/ncomms15837.

Holl, A. M. 2018. "Survival of mature mouse olfactory sensory neurons labeled genetically perinatally." *Mol Cell Neurosci* 88:258-269. doi: 10.1016/j.mcn.2018.02.005.

Kaneko, M., and T. Takahashi. 2004. "Presynaptic mechanism underlying cAMP-dependent synaptic potentiation." *J Neurosci* 24 (22):5202-8. doi: 10.1523/JNEUROSCI.0999-04.2004.

Karnup, S. V., A. Hayar, M. T. Shipley, and M. G. Kurnikova. 2006. "Spontaneous field potentials in the glomeruli of the olfactory bulb: the leading role of juxtaglomerular cells." *Neuroscience* 142 (1):203-21. doi: 10.1016/j.neuroscience.2006.05.068.

Kermen, F., P. Lal, N. G. Faturos, and E. Yaksi. 2020. "Interhemispheric connections between olfactory bulbs improve odor detection." *PLoS Biol* 18 (4):e3000701. doi: 10.1371/journal.pbio.3000701.

Kisilevsky, A. E., and G. W. Zamponi. 2008. "D2 dopamine receptors interact directly with N-type calcium channels and regulate channel surface expression levels." *Channels (Austin)* 2 (4):269-77. doi: 10.4161/chan.2.4.6402.

Kiyokage, E., Y. Z. Pan, Z. Shao, K. Kobayashi, G. Szabo, Y. Yanagawa, K. Obata, H. Okano, K. Toida, A. C. Puche, and M. T. Shipley. 2010. "Molecular identity of periglomerular and short axon cells." *J Neurosci* 30 (3):1185-96. doi: 10.1523/JNEUROSCI.3497-09.2010.

Kludt, E., C. Okom, A. Brinkmann, and D. Schild. 2015. "Integrating temperature with odor processing in the olfactory bulb." *J Neurosci* 35 (20):7892-902. doi: 10.1523/JNEUROSCI.0571-15.2015.

Kosaka, T., A. Pignatelli, and K. Kosaka. 2020. "Heterogeneity of tyrosine hydroxylase expressing neurons in the main olfactory bulb of the mouse." *Neurosci Res* 157:15-33. doi: 10.1016/j.neures.2019.10.004.

Lecoq, J., P. Tiret, and S. Charpak. 2009. "Peripheral adaptation codes for high odor concentration in glomeruli." *J Neurosci* 29 (10):3067-72. doi: 10.1523/JNEUROSCI.6187-08.2009.

Lodovichi, C. 2021. "Topographic organization in the olfactory bulb." *Cell Tissue Res* 383 (1):457-472. doi: 10.1007/s00441-020-03348-w.

Love, N. R., Y. Chen, S. Ishibashi, P. Kritsiligkou, R. Lea, Y. Koh, J. L. Gallop, K. Dorey, and E. Amaya. 2013. "Amputation-induced reactive oxygen species are required for successful Xenopus tadpole tail regeneration." *Nat Cell Biol* 15 (2):222-8. doi: 10.1038/ncb2659.

Manzini, I., C. Brase, T. W. Chen, and D. Schild. 2007. "Response profiles to amino acid odorants of olfactory glomeruli in larval Xenopus laevis." *J Physiol* 581 (Pt 2):567-79. doi: jphysiol.2007.130518 [pii]
10.1113/jphysiol.2007.130518.

Manzini, I., and D. Schild. 2004. "Classes and narrowing selectivity of olfactory receptor neurons of Xenopus laevis tadpoles." J *Gen Physiol* 123 (2):99-107. doi: jgp.200308970 [pii]
10.1085/jgp.200308970.

Manzini, I., D. Schild, and C. Di Natale. 2022. "Principles of odor coding in vertebrates and artificial chemosensory systems." Ph*ysiol Rev 1*02 (1):61-154. doi: 10.1152/physrev.00036.2020.

McGann, J. P. 2013. "Presynaptic inhibition of olfactory sensory neurons: new mechanisms and potential functions." Ch*em Senses 3*8 (6):459-74. doi: 10.1093/chemse/bjt018.

Menini, A. 2010. "The Neurobiology of Olfaction." In.

Mombaerts, P., F. Wang, C. Dulac, S. K. Chao, A. Nemes, M. Mendelsohn, J. Edmondson, and R. Axel. 1996. "Visualizing an olfactory sensory map." Ce*ll 8*7 (4):675-86. doi: 10.1016/s0092-8674(00)81387-2.

Moreno, N., R. Morona, J. M. López, L. Dominguez, M. Muñoz, and A. González. 2008. "Anuran olfactory bulb organization: embryology, neurochemistry and hodology." Br*ain Res Bull 7*5 (2-4):241-5. doi: 10.1016/j.brainresbull.2007.10.027.

Murphy, G. J., L. L. Glickfeld, Z. Balsen, and J. S. Isaacson. 2004. "Sensory neuron signaling to the brain: properties of transmitter release from olfactory nerve terminals." J *Neurosci 2*4 (12):3023-30. doi: 10.1523/JNEUROSCI.5745-03.2004.

Nezlin, L. P., and D. Schild. 2000. "Structure of the olfactory bulb in tadpoles of Xenopus laevis." Ce*ll Tissue Res 3*02 (1):21-9.

Niethammer, P., C. Grabher, A. T. Look, and T. J. Mitchison. 2009. "A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish." N*ature 459* (7249):996-9. doi: 10.1038/nature08119.

Nieuwkoop, P.D, and J. Faber. 1956. N*ormal table of Xenopus laevis (Daudin). A systematical and chronological survey of the development from the fertilized egg till the end of metamorphosis.:* Amsterdam: North-Holland Publishing Company. Guilders.

Petzold, GC, DF Albeanu, TF Sato, and VN Murthy. 2008. "Coupling of neural activity to blood flow in olfactory glomeruli is mediated by astrocytic pathways." N*euron 58* (6):897-910. doi: S0896-6273(08)00403-0 [pii]

10.1016/j.neuron.2008.04.029.

Pignatelli, A., K. Kobayashi, H. Okano, and O. Belluzzi. 2005. "Functional properties of dopaminergic neurones in the mouse olfactory bulb." J P*hysiol 564* (Pt 2):501-14. doi: 10.1113/jphysiol.2005.084632.

Pírez, N., and M. Wachowiak. 2008. "In vivo modulation of sensory input to the olfactory bulb by tonic and activity-dependent presynaptic inhibition of receptor neurons." J N*eurosci 28* (25):6360-71. doi: 10.1523/JNEUROSCI.0793-08.2008.

Roth, G., F. Laberge, S. Mühlenbrock-Lenter, and W. Grunwald. 2007. "Organization of the pallium in the fire-bellied toad Bombina orientalis. I: Morphology and axonal projection pattern of neurons revealed by intracellular biocytin labeling." J C*omp Neurol 50*1 (3):443-64. doi: 10.1002/cne.21255.

Shepherd, G.M., W.R. Chen, and C.A. Greer. 2004. The *Synaptic Organization of the Brain: O*xford University Press.

Terni, B., and A. Llobet. 2021. "Axon terminals control endolysosome diffusion to support synaptic remodelling." Lif*e Sci Alliance 4* (8). doi: 10.26508/lsa.202101105.

Terni, B., P. Pacciolla, H. Masanas, P. Gorostiza, and A. Llobet. 2017. "Tight temporal coupling between synaptic rewiring of olfactory glomeruli and the emergence of odor-guided behavior in Xenopus tadpoles." J C*omp Neurol 525* (17):3769-3783. doi: 10.1002/cne.24303.

Terni, B., P. Pacciolla, M. Perelló, and A. Llobet. 2018. "Functional Evaluation of Olfactory Pathways in Living Xenopus Tadpoles." J V*is Exp (1*42). doi: 10.3791/58028.

Wachowiak, M., and L. B. Cohen. 1999. "Presynaptic inhibition of primary olfactory afferents mediated by different mechanisms in lobster and turtle." J N*eurosci 19* (20):8808-17. doi: 10.1523/JNEUROSCI.19-20-08808.1999.

Wachowiak, M., J. P. McGann, P. M. Heyward, Z. Shao, A. C. Puche, and M. T. Shipley. 2005. "Inhibition [corrected] of olfactory receptor neuron input to olfactory bulb glomeruli mediated by suppression of presynaptic calcium influx." J N*europhysiol 94* (4):2700-12. doi: 10.1152/jn.00286.2005.

Weiss, J., M. Pyrski, P. Weissgerber, and F. Zufall. 2014. "Altered synaptic transmission at olfactory and vomeronasal nerve terminals in mice lacking N-type calcium channel Cav2.2." Eur *J Neurosci 40* (10):3422-35. doi: 10.1111/ejn.12713.

Weiss, L., I. Manzini, and T. Hassenklöver. 2021. "Olfaction across the water-air interface in anuran amphibians." Cel*l Tissue Res 38*3 (1):301-325. doi: 10.1007/s00441-020-03377-5.

Weiss, L., P. Segoviano Arias, T. Offner, S. J. Hawkins, T. Hassenklöver, and I. Manzini. 2021. "Distinct interhemispheric connectivity at the level of the olfactory bulb emerges during Xenopus laevis metamorphosis." Cel*l Tissue Res 386* (3):491-511. doi: 10.1007/s00441-021-03527-3.

Westhoff, G., and G. Roth. 2002. "Morphology and projection pattern of medial and dorsal
pallial neurons in the frog Discoglossus pictus and the salamander Plethodon jordani."
J C*omp Neurol 44*5 (2):97-121. doi: 10.1002/cne.10136.
Wu, Y., L. Dissing-Olesen, B. A. MacVicar, and B. Stevens. 2015. "Microglia: Dynamic
Mediators of Synapse Development and Plasticity." Tre*nds Immunol 36* (10):605-
613. doi: 10.1016/j.it.2015.08.008.
955

Figure 1

Figure 2

Figure 3

Figure 4

A

B
O.N. Transection
Optic Nerves Transection

LFP Peak (μV)

O.N. width (μm)

50 μV

1 s

C
OB
ON

max
min

100 μm

F_{HyPer} OB left / OB right

Control   O.N. Trans.

D
LFP Peak (μV)

n.s.

n.s.

O.N. Trans.   + Apocynin   + DPI

Figure 5

Figure 6

Figure 7

A Met

50 μV

1 s

Baseline
(0-8 min)

Raclopride
(4 min 14 min 24 min 34 min 42 min)

B
n.s. p=0.46
n.s. p=0.2
*p<0.05

LFP Peak (μV)

Control   Contr. trans.

C
LFP Peak$_{Rac.}$ / LFP Peak$_{basal}$

Control
O.N. Transection

Raclopride

Time (min)

D Met   Baseline
Raclopride

ΔF/F GCaMP6s

LFP Peak (μV)

Time (s)

E
n.s., p=0.78

LFP Peak (μV)

Control   CGP-36742

F
Baseline
CGP-36742 (4 min-42 min)

LFP Peak$_{CGP}$ / LFP Peak$_{basal}$

50 μV

1 s

CGP-36742

Time (min)

Figure 8

Figure 9