Bilateral equalization of synaptic output in olfactory glomeruli of Xenopus tadpoles

Marta Casas[1,2], Beatrice Terni[1,2] and Artur Llobet[1,2*]

[1] Laboratory of Neurobiology, Department of Pathology and Experimental Therapy, Institute of Neurosciences, University of Barcelona, 08907 L'Hospitalet de Llobregat, Barcelona, Spain

[2] Bellvitge Biomedical Research Institute (IDIBELL), 08907 L'Hospitalet de Llobregat, Barcelona, Spain

* Corresponding Author:
Artur Llobet
Laboratory of Neurobiology
Department of Pathology and Experimental Therapy
Faculty of Medicine and Health Sciences – University of Barcelona
08907 L'Hospitalet de Llobregat
Barcelona, Spain
E-mail: allobet@ub.edu
Phone: +34-934024279
ORCID: 0000-0001-5797-6782

Odorants stimulate olfactory sensory neurons (OSNs) to create a bilateral sensory map defined by a set of glomeruli present in the left and right olfactory bulbs. Using *Xenopus tropicalis* tadpoles we challenged the notion that glomerular activation is exclusively determined ipsilaterally. Glomerular responses evoked by unilateral stimulation were potentiated following transection of the contralateral olfactory nerve. The gain of function was observed as early as 2 hours after injury and faded away with a time constant of 4 days. Potentiation was mediated by the presence of larger and faster calcium transients driving glutamate release from OSN axon terminals. The cause was the reduction of the tonic presynaptic inhibition exerted by dopamine D_2 receptors. Inflammatory mediators generated by injury were not involved. These findings reveal the presence of a bilateral modulation of glomerular output driven by dopamine that compensates for imbalances in the number of operative OSNs present in the two olfactory epithelia. Considering that the constant turnover of OSNs is an evolutionary conserved feature of the olfactory system and determines the innervation of glomeruli, the compensatory mechanism here described may represent a general property of the vertebrate olfactory system to establish an odor map.

1 **INTRODUCTION**
2

Olfactory glomeruli are spherical regions of neuropil that contain the first synapse for

processing olfactory information and form the glomerular layer of the olfactory bulb. Despite

variations in number or size among vertebrate species, glomeruli show an evolutionarily

conserved synaptic connectivity (Shepherd, Chen, and Greer 2004). Glomerular input comes

from axon terminals of olfactory sensory neurons (OSNs) expressing the same odorant receptor

that transfer information to the dendrites of mitral cells. Synaptic transmission between OSNs

and mitral cells is regulated by inhibitory and excitatory contacts established with

juxtaglomerular neurons. This connectivity allows that upon exposure to odorants only a

defined set of glomeruli is activated, creating an odor map (Mombaerts et al. 1996). Glomeruli

exhibit an anatomical and functional symmetrical distribution in the left and right olfactory

bulbs, therefore, the sensory map created in the olfactory bulb is bilateral and is generated by

homologous glomeruli reflecting the contribution of the two nasal cavities (Lodovichi 2021;

Belluscio and Katz 2001).

OSNs originate in the olfactory epithelium and are subject to a continuous turnover

throughout life (Holl 2018). An efficient neuronal replacement is accomplished by the rapid

connection of newborn OSNs to the olfactory bulb. For example, in *X. tropicalis* tadpoles

OSNs can establish functional synapses in a time window of 4 days (Terni et al. 2017) and, in

mice, it takes a week for newborn OSNs to get inserted in the olfactory bulb through a *plug-*

and-play mechanism (Browne, Crespo, and Grubb 2022). The balance existing between the

elimination of OSNs and their insertion in olfactory bulb circuitry likely determines a range of

input neurons innervating a glomerulus, rather than a constant, precise figure. In this context,

how the formation of an odor map accounts for possible variations in the number of input

neurons is unknown. Considering that OSNs exclusively form ipsilateral synapses (Shepherd,

Chen, and Greer 2004) and release glutamate with near-maximal release probability (Murphy

et al. 2004), the possible crosstalk balancing the individual contribution of bilaterally

distributed glomeruli remains unknown, because their high output gain is assumed to be

defined unilaterally.

Here, we take advantage of the experimental capacities offered by *Xenopus tropicalis*

tadpoles to assess whether the output of a genetically labelled glomerulus (Terni et al. 2017;

Terni and Llobet 2021), is solely determined by ipsilateral stimulation. In vivo recordings

carried out after disrupting the contribution of the contralateral pathway, challenged this notion.

Our results show that the tonic inhibition of glomerular responses exerted by dopamine D_2

receptors is bilaterally shaped to compensate for differences in the number of OSNs innervating

the contralateral olfactory bulb. Considering the evolutionary conserved developmental,

morphological, and functional features of the *Xenopus* tadpole olfactory system (Menini 2010;

Manzini, Schild, and Di Natale 2022), the homeostatic mechanism here described might

represent a general rule to the formation of an integrated odor map in vertebrates.

MATERIALS AND METHODS

Animals

Ethical procedures were approved by the regional government (Generalitat de

Catalunya, experimental procedure #10753). *Xenopus tropicalis* were housed and raised

according to the standard protocols of the animal facilities of the University of Barcelona.

Tadpoles were obtained by natural mating and kept in tanks at 25 ºC. *Xenopus* water

conductivity was adjusted to 700 µs·cm⁻¹, pH=7.5. Tadpoles at stage 47-52 of the Nieuwkoop–

Faber criteria (Nieuwkoop and Faber 1956) were used for the experiments. The *X. tropicalis*

transgenic line *Dre.mxn1:GFP* (RRID:NXR_1111) was used for the visualization of a discrete

population of OSNs (Terni et al. 2017; Terni and Llobet 2021) and for carrying out

electrophysiological recordings in a genetically defined glomerulus. Labelling of OSNs was

consistent in all animals of this transgenic line. Calcium imaging was performed in the *X.*

tropicalis transgenic line *ElasGFP:Tubb2b-GCaMP6s* (RRID:NXR_1123). In vivo detection

of reactive oxygen species (ROS) was carried out in tadpoles of the *X.laevis* transgenic line

Hsa.UBC-Gal4;UAS:HyPer-YFP (RRID: NXR_0127).

To apply surgical procedures, *Xenopus* tadpoles were anesthetized in 0.02% MS-222

and transferred to a wet nitrocellulose filter paper placed under a stereomicroscope. Iridectomy

scissors were used for the unilateral transection of an olfactory nerve or the bilateral transection

of optic nerves. Tadpoles were returned to water tanks after surgery.

Electrophysiology

X. tropicalis tadpoles were anesthetized in 0.02% MS-222 and the portion of skin

covering the olfactory bulb was removed. Animals were transferred to a well fabricated in a

dish coated with silicone elastomer. A coverslip restricted tadpole movements and left

olfactory placodes and bulbs accessible (Terni et al. 2018). The dish was placed on the stage

of an upright microscope (Zeiss, Axioexaminer A1, Oberkochen, Germany) and was

continuously perfused with *Xenopus* Ringer containing (in mM):100 NaCl, 2 KCl, 1 $CaCl_2$, 2

$MgCl_2$, 10 glucose, 10 HEPES, 240 mOsm/kg, pH=7.8, supplemented with 1 μM d-

tubocurarine to prevent muscle contractions. All salts and drugs were from Sigma-Aldrich

(Sant Louis, MO). CGP-36742 was from Novartis Pharmaceuticals (Basel, Switzerland). To

carry out extracellular recordings a borosilicate pipette filled with *Xenopus* ringer was targeted

to the olfactory glomerulus showing GFP fluorescence. Signals were acquired using a

Geneclamp 500A amplifier (Molecular Devices, San Jose, CA) and digitized at 10 kHz using

a National Instruments NI-USB-6341 DAC board (National Instruments, Austin, TX)

controlled by mafPC software (courtesy of M. A. Xu-Friedman, University at Buffalo, NY).

The recorded changes of the LFP were low pass filtered below 100 Hz and analyzed with Igor

Pro 9.0 (Wavemetrics, OR).

Methionine was chosen as odor stimulus based on its broad capacity to stimulate OSNs

in *Xenopus laevis* tadpoles (Manzini and Schild 2004). A puff of 200 μM methionine solution,

obtained by diluting a 10 mM stock solution prepared in *Xenopus* Ringer (pH=7.8), was applied

on the ipsilateral olfactory epithelium to the recorded glomerulus. The amino acid was

delivered through a 0.25 mm diameter fused silica capillary (World Precision Instruments,

Hertfordshire, UK) positioned on the top of a nasal cavity. The timing of application was

controlled via a TTL pulse. The characteristic odor evoked glomerular response was obtained

by averaging LFP changes triggered by sequential stimulations applied at 2 min intervals. The

local application of antagonists was carried out by transiently applying 20 psi pressure to the

pipette holder inlet. Approximately 1 μL of the pipette solution was delivered to the

glomerulus.

Imaging

Quantification of OSNs present in the olfactory epithelium of *Dre.mxn1:GFP* tadpoles

was performed in an inverted LSM880 confocal microscope (Zeiss) using animals anesthetized

in 0.02% MS-222. The number of neurons was estimated from maximal intensity projections

in confocal Z-stacks. To selectively eliminate GFP positive OSNs the 2Phatal method (Hill et

al. 2017) was adapted to confocal microscopy. Tadpoles were immersed for 15 minutes in

Xenopus water containing 5 μg/mL Hoechst 33342 and anesthetized in 0.02% MS-222.

Animals were placed dorsally on a 25 mm glass #1.5 coverslip acting as the bottom of an

imaging chamber and transferred to the microscope stage. The right olfactory epithelium,

located contralaterally to the recording site, was imaged with a 10X Plan Apo objective,

NA=0.45 (Zeiss). The digital zoom was set to 3.5. Photobleaching was carried out using

ZenBlue software (Zeiss) in 4 to 6 regions of interest (ROIs) containing two or more GFP

positive OSNs. Animals were next returned to water tanks.

The *X. tropicalis* transgenic line *ElasGFP:Tubb2b-GCaMP6s*, where the neuronal β-

tubulin promoter drives the expression of the calcium sensor GCaMP6s, was used to visualize

the activation of olfactory glomeruli. As for in vivo electrophysiology, tadpoles were

anesthetized in 0.02% MS-222, placed in a recording dish and transferred to the stage of an

upright microscope (Zeiss, Axioexaminer A1). Tadpoles were continuously perfused with

Xenopus Ringer containing 1 μM d-tubocurarine. Imaging was carried out at 76 Hz using a

Maico MEMS confocal unit (Hamamatsu Photonics, Hamamatsu City, Japan). A 40X, 0.75

NA water immersion W-N-Achroplan objective (Zeiss) was used. Odorant stimulation was

performed as described for electrophysiology experiments. The timing of methionine

application was synchronized with image acquistion (HCImage software, Hamamatsu

Photonics) using a Master 8 stimulator (AMPI, Jerusalem, Israel). In experiments where

imaging was coupled to recordings of the LFP, the electrode was targeted to the lateral

glomerular cluster. All signals were synchronized using a Master 8 stimulator.

Raw fluorescence image sequences were used to construct ΔF/F movies according to

the relationship $(F-F_0)/F_0$, where F_0 corresponded to the basal fluorescence levels obtained

during 1 s before stimulation. ΔF/F sequences were next subsampled by averaging groups of

10 frames to identify putative glomeruli, which were defined as 10-20 μm diameter round

structures consistently responding to sequential stimulations applied at 2 min intervals. The

selected ROIs were transferred to the original raw fluorescence movie to quantify increases in

basal intracellular calcium levels. A ROI showing ΔF/F increases evoked by methionine

application to the ipsilateral olfactory epithelium that were ≥3 SDs above baseline levels was

considered as a single glomerulus. The temporal response was quantified using Igor Pro 9

software (Wavemetrics, OR).

The production of ROS in the *X. laevis* HyPer-YFP was evaluated after transection of

a single olfactory nerve. Tadpoles were imaged in an inverted confocal LSM 900 microscope

(Zeiss) using a Plan-Apochromat 20X, 0.8NA objective. Hyper-YFP was excited at 405 nm

and 488 nm. ROS levels were estimated from the relationship obtained between HyPer-YFP

excited at 488 nm and 405 nm (Love et al. 2013).

Uncaging of Rubi-glutamate

Rubi-glutamate was added to the pipette solution at a final concentration of 300 μM.

Procedures were carried out considering the high sensibility to light of the caged compound

(Fino et al. 2009). To prevent spontaneous activation of Rubi-glutamate, glass pipettes were

coated with beeswax. The electrode was positioned above the lateral region of the glomerular

layer using dim transmitted light. Next, the GFP labelled glomerulus was targeted using blue

light attenuated with a ND filter (0.5 OD). Approximately 1 μL of the pipette solution was

injected in the glomerulus. Upon certifying that the recording of LFP signal was stable for 2

min, a TTL signal opened a shutter (Lambda SC, Sutter Instrument, Novato, CA) for 500 ms

to deliver blue light (470±20 nm) through the epifluorescence port. A diaphragm restricted

light application to the region targeted by the electrode.

Statistical analysis

For statistical analysis, paired and unpaired t-test were used to evaluate differences

between two experimental groups. Comparisons among three or more groups were performed

using one way ANOVA followed by multiple comparison Tukey's HSD test. Average values

are expressed as mean±s.e.m. Statistical analysis was carried out using Igor Pro software.

RESULTS

Characterization of odor evoked responses in a genetically defined glomerulus

The *Xenopus tropicalis* line *Dre.mxn1:GFP* allows the identification of an olfactory

glomerulus located laterally and innervated by a discrete population of OSNs (Terni et al. 2017;

Terni and Llobet 2021). We targeted an electrode to the left GFP labelled glomerulus and

recorded changes in the Local Field Potential (LFP) evoked by ipsilateral stimulation with an

amino acid acting as a waterborne odorant (Figs. 1A and B). An olfactory response was

triggered by 100 ms application of 200 μM methionine to the olfactory epithelium, thus

confirming the ability of *Xenopus* tadpoles to detect amino acids (Manzini et al. 2007).

Stimulation evoked a negative deflection of the LFP that resembled responses obtained in the

glomerular layer of rats upon sniffing odors (Chaigneau et al. 2007). A main difference

between negativities recorded in *X. tropicalis* and those found in rodents is that the latter are

respiration-locked, while in tadpoles there was a single deflection that recovered with a half

time ranging from 1 to 4 s.

An estimate of the spatial extent of the LFP signal was obtained by evaluating responses

after changing the position of the recording electrode. The characteristic profile of LFP

changes disappeared when the pipette was directed to the mitral cell layer (Fig. 1C), thus

illustrating they were confined to the glomerular layer. The consistent success rate (85%,

n=269) obtained by targeting the GFP labelled glomerulus with the recording electrode

decreased to 50% (n=36) when random locations linearly spaced by 50 μm were tested within

the glomerular layer. The example shown in Fig. 1D illustrates how the characteristic response

to methionine was obtained only in one out of four positions tested. These observations could

be explained by the capacity of the electrode to detect only activated glomeruli (Manzini et al.

2007). The high success rate of the recordings guided by fluorescence support that the GFP

labeled glomerulus responded to methionine. Considering many olfactory glomeruli in

Xenopus tadpoles are broadly tuned (Manzini et al. 2007), it is conceivable that the recorded

glomerulus was also activated by other waterborne odorants, which were not tested in the

current study.

The LFP signal likely sampled the entire volume of 15765 ± 2119 μm^3 (n=33) defined

by fluorescence but, as the lateral cluster of glomeruli is markedly involved in the detection of

amino acids (Weiss, Manzini, and Hassenklöver 2021), the influence of adjacent glomeruli

cannot be ruled out from the start. Considering the labelled glomerulus had a diameter of ~30

μm (Fig. 1E) and there was a 50 μm axial resolution to obtain independent readouts (Fig. 1D),

only a portion of surrounding glomeruli could theoretically participate in the observed LFP

signal. The glomerular layer of the amphibian olfactory bulb is loosely organized compared to

mammals, as glomerular units show different sizes and lack ensheathing astrocytes (Nezlin and

Schild 2000; Gaudin and Gascuel 2005). The absence of a homogeneous distribution of

glomeruli, added to the dorsorostral location of GFP labelled axon terminals (Fig. 1E),

restricted the number of adjacent glomeruli that could theoretically contribute to the recorded

LFP signal to those found ventrally or caudally. Even if 50% of such units were activated by

methionine, the decay of the LFP following a relationship inversely related to distance to the

recording site, as reported for the mammalian olfactory bulb (Karnup et al. 2006), would

minimize their contribution. Altogether indicates that methionine application to the olfactory

epithelium reliably activated the lateral glomerulus labeled by GFP in the *Dre.mxn1:GFP* line.

Glomerular responses were assayed by repetitive stimulation at two-minute intervals

and all negativities showed a comparable profile (Fig. 1B). The peak of the negative deflection

was reached ~1 s after olfactory stimulation. The characteristic LFP response of any given

tadpole was obtained by averaging 5 to 8 consecutive stimulations. Local application of 100

μM CNQX decreased LFP negativities by approximately 50% (Fig. 2A). The reduction was

comparable when 100 μM D-AP5 was used instead of CNQX (Fig. 2B). The concomitant

application of both drugs produced an additive effect that reduced responses by 70% (Fig. 2C

p=0.0004, paired t-test), thus evidencing their synaptic origin, as well as the involvement of

AMPA and NMDA receptors. These results are comparable to those obtained in rats

(Chaigneau et al. 2007; Lecoq, Tiret, and Charpak 2009), which validates our experimental

configuration to record LFP responses in a region defined by a genetically labelled glomerulus.

The onset phase of negativities was particularly sensitive to CNQX and AP5,

suggesting that it was reflecting the involvement of receptors activated by glutamate release

from OSN axon terminals. A way to test this possibility was to induce a jump in glutamate

concentration inside the glomerulus and assay whether a change in the LFP signal occurred.

We targeted the electrode to the lateral GFP glomerulus, injected *Xenopus* ringer solution

containing 300 μM Rubi-glutamate and uncaged it using a flash of blue light (Fino et al. 2009).

A transient negativity was induced by a 500 ms pulse of light, which was reproduced by

repeated stimulations (Fig. 2D). The average amplitude of LFP_{peaks} was of 30±3 μV (n=21, 3

tadpoles). This observation supported that the initial phase of LFP negativities was caused by

the marked increase in the extracellular concentration of glutamate secreted by OSN axon

terminals converging in a glomerulus.

The contribution of inhibitory neurotransmission mediated by $GABA_A$ receptors was

negligible because no variations in LFP changes evoked by methionine were observed after

local application of 1 mM picrotoxin (Fig. 2E). To certify that ipsilateral stimulation of OSNs

was the trigger of recorded responses, we investigated the effect of methionine application to

the contralateral epithelium. Negativities disappeared (Fig. 2F), thus demonstrating that

glomerular activation was exclusively unilateral. In 37% of the tadpoles studied (n=174) the

negativity was preceded by a transient positivity (Fig. 1B). When present, this LFP deflection

was not modified by glutamate and $GABA_A$ receptor antagonists (Figs. 2A and E), thus

indicating it was unrelated to synaptic mechanisms. As the positivity was caused by the

application of olfactory stimuli and always occurred before the synaptic response, it is possible

to consider its origin in cellular structures conveying peripheral information to the olfactory

bulb such as the layer of nerve fibers.

Odor evoked responses depend on the number of input neurons.

The amplitude of LFP negativities decreases during the reinnervation of the glomerular

layer after olfactory nerve transection (Terni et al. 2017), which suggests that LFP changes are

related to the number of OSNs axons projecting to glomeruli. If odor evoked responses were

indeed related to glomerular input, the smallest negativities should take place in the earliest

developmental stages. This possibility was confirmed by observing that GFP positive neurons

found in the olfactory epithelium increased with development and their number followed an

empirical linear relationship related to olfactory nerve width for tadpoles found between NF

stages 47 and 54 (Fig. 3A, r^2=0.96). The peak of the LFP response was also related to olfactory

nerve width but followed an exponential fit. Negativities reached a steady-state value (Fig.

3A), thus suggesting the achievement of a consolidated glomerular activation in this

developmental time interval. The presence of constant glomerular responses for NF stages >50

coincides with the consolidation of the basic structure of the olfactory bulb, which remains

constant through larval life (Byrd and Burd 1991). LFP changes were empirically described

by the function $LFP_{peak}=LFP_{steady-state}+Ae^{-r \cdot ONW}$ that associated the maximum change in the LFP

(in μV) to olfactory nerve width (ONW, expressed in μm). The fit revealed an increase rate (r)

of 0.13 μm^{-1} starting at a diameter of 21 μm, which corresponds to the thinnest olfactory nerves

present during development around NF stage 40. Considering the linear correlation found

between the number of GFP positive OSNs and the width of olfactory nerves, it was possible

to rewrite the exponential fit as a function of the number of OSNs. The estimated peak

negativity in the LFP could thus be obtained according to $LFP_{peak}=108-1866 \cdot e^{-0.11* \cdot OSNs}$ for a

given number of OSNs in the olfactory epithelium. On average, an olfactory epithelium

contained 30 ± 2 GFP positive OSNs, ranging from 24 to 39 labelled neurons (n=44) and, this

relationship provided a quantitative association of the amplitude of odor evoked responses to

the insertion of OSNs in a glomerulus during normal development.

Odor evoked responses are potentiated by the injury of the contralateral olfactory

pathway.

Transection of both olfactory nerves causes a silencing of the capacity of *X. tropicalis*

tadpoles to perceive odors (Terni et al. 2017). Since animals having a single nerve still display

normal olfactory guided behavior, we investigated how information is transmitted in the

absence of the mirror pathway. Olfactory nerve transection caused a potentiation of evoked

LFP negativities in the recorded contralateral olfactory glomerulus (Fig. 3B) that was already

observed two hours after injury (Fig. C). All tadpoles investigated showed odor evoked

responses above the expected LFP_{peak} level for the range of developmental stages analyzed.

By reaching a maximum potentiation between 24 and 48 hours after transection, LFP_{peak}

responses returned to control values following an exponentially recovery phase that took place

with a time constant of 100 hours (Fig. 3D). The overall duration of potentiation coincided

with the approximately two weeks required to reform the glomerular layer upon olfactory nerve

transection (Terni et al. 2017), suggesting that the increase of glomerular output occurred while

the circuitry of the contralateral olfactory bulb was being remodeled after injury.

LFP_{peak} responses had a strong synaptic contribution because they were dependent on

the number of OSNs innervating the glomerulus (Fig. 3A), were sensitive to the application of

glutamate receptor blockers (Figs. 2A-C) and could be triggered by glutamate uncaging (Fig.

2D). These results complement those obtained in rats showing a correlation of the amplitude

of LFP negativities to calcium influx in OSN terminals (Lecoq, Tiret, and Charpak 2009) and

EPSPs recorded in mitral cells (Chaigneau et al. 2007). We thus hypothesized that the observed

potentiation affected to glutamatergic synapses mediating glomerular activation.

Tissue repair mechanisms are not involved in the potentiation of odor evoked responses

Since inflammatory mediators affect to synaptic functions (Wu et al. 2015), we

evaluated the involvement of molecules released by injury using two different strategies. First,

to observe if sensory nerve damage were enhancing LFP_{peak} responses, we sectioned both optic

nerves (Fig. 4A). Even though optic and olfactory nerves show substantial functional and

anatomical differences, the nature of the injury, as well as the distance to the recorded site was

comparable between optic and olfactory nerve transection. On these bases we assumed that

the effect generated by diffusible inflammatory mediators, or the capacity to mobilize cells

mediating inflammation, should be comparable in both experimental conditions. LFP

negativities recorded 24h to 48h after bilateral optic nerve transection showed an amplitude of

86 ± 6 μV (n=17). This value was expected by normal development but was significantly lower

($p=1.92\cdot10^{-5}$, unpaired t-test), than the amplitude of 186 ± 16 μV (n=14) found in tadpoles

subjected to transection of the contralateral olfactory nerve (Fig. 4B).

Secondly, we investigated the release of reactive oxygen species (ROS), which are key

to trigger tissue repair mechanisms in tadpoles (Love et al. 2013). The *Xenopus laevis* line

HyPer-YFP displays a ubiquitous expression of the H_2O_2 sensor HyPerYFP that allows the in

vivo detection of ROS created by wounds (Love et al. 2013; Niethammer et al. 2009). Two

hours after unilateral olfactory nerve transection there was an increase in H_2O_2 levels that was

restricted to the cells present at the injury site (Fig. 4C). ROS levels remained unaltered in the

olfactory epithelium and in both olfactory bulbs. Moreover, the amplitude of odor evoked

changes in the LFP were unaffected by the presence 200 μM apocynin or 2 μM

diphenyleneiodonium, two blockers of ROS production (Fig. 4D, p=0.36, ANOVA followed

by Tukey's test) that block tail regeneration at these concentrations (Love et al. 2013). All

these findings together suggested that the enhancement of glomerular responses was unlikely

caused by inflammatory mediators or the activation of fundamental tissue repair mechanisms

operating in tadpoles and, revealed the presence of an intrinsic compensatory mechanism

operating in the olfactory system.

Potentiation of odor evoked responses is of presynaptic origin

The activation of individual glomeruli was imaged in the transgenic *X. tropicalis* line

tubb2b:GCaMP6s to gain insight on a putative synaptic mechanism acting on the bilateral

balance of odor evoked responses. Ipsilateral application of 200 μM methionine to the

olfactory epithelium activated a defined set of glomeruli, which were identified as round, 10-

20 μm diameter structures. Same glomeruli reacted to three or more consecutive methionine

stimulations (Fig. 5A). Responses detected as calcium transients showed comparable

amplitudes and temporal profiles (Fig. 5B), suggesting that the simultaneous activation of

sparse glomeruli contributed to the elaboration of an olfactory map. Two lines of evidence

support that calcium increases originated in presynaptic terminals of OSNs. First, because the

tubb2b promoter primarily targets OSNs over olfactory bulb interneurons. A detailed

characterization carried out in *X. laevis* tadpoles shows that *tubb2b* efficiently drives expression

of a genetically encoded fluorescent reporter in OSNs but fails to label periglomerular neurons

positive for calretinin or tyrosine hydroxylase and only targets 24% of mitral cells (Daume et

al. 2022). The convergence of multiple fluorescent OSN axon terminals on a single glomerulus

and a minimal presence of labelled juxtaglomerular and mitral cells, makes *X. tropicalis*

tubb2b:GCaMP6s tadpoles well-suited to detect the presynaptic component of glomerular

responses. Second, because the simultaneous imaging of glomerular activation and recording

of LFP negativities revealed a temporal correlation of the two signals (Fig. 5C). This

observation matches the findings of a previous study carried out in rats, where a comparable

relationship was described between responses detected in OSN axon terminals labelled with a

high affinity calcium dye and LFP signals (Lecoq, Tiret, and Charpak 2009). Although tadpoles

of the *tubb2b-GCaMP6s* line consistently showed glomeruli activated by methionine, the

coupling of imaging to LFP signals had a low success rate. Even though the electrode was

targeted to the lateral glomerular cluster, which is the region expected to concentrate responses

to amino acids (Weiss, Manzini, and Hassenklöver 2021), LFP negativities were detected in

only 33% of animals tested (n=17/52). This figure is below the 50% found after the

performance of recordings in random locations of the glomerular layer (Fig. 1D) but it is higher

than the ~25% success rate found in *X. laevis* tadpoles (Manzini et al. 2007). This observation

reinforces the selectivity of LFP recordings to detect the output of a single glomerulus, which

in this set of experiments was coupled to the fluorescent response found in closest apposition

to the tip of the pipette. Altogether indicates that the readouts of glomerular input and output

were reported by transient increases in GCaMP6s fluorescence and LFP signals, respectively.

A hallmark of contralateral olfactory nerve transection was the development of faster

LFP negativities. The time constant to reach the LFP_{peak} shortened significantly (p=0.017,

unpaired t-test) from 1.05±0.1 s (n=46 tadpoles) to 0.71±0.1 s (n=39 tadpoles), whilst the

recovery phase was unaffected (Fig. 5D, p=0.79, unpaired t-test). The faster glomerular output

likely had a presynaptic origin, because calcium transients were larger and showed faster onset

kinetics after contralateral nerve transection (Figs. 5E and F). The average amplitude increased

more than two-fold (p=0.02, unpaired t-test) and the time constant describing the rise time

shortened significantly (p=0.0003, unpaired t-test) in tadpoles with a transected olfactory nerve

compared to control animals (Fig. 5F). The slow kinetics of GCaMP6s prevented a precise

temporal association between calcium buildup and the onset of LFP changes (Fig. 5C) and a

certain saturation of the high affinity calcium indicator in axon terminals was also expected.

But, despite these limitations to quantitative analysis, the changes reported by GCaMP6s

showed an overall enhancement of cytosolic calcium levels mediating neurotransmitter release.

We next sought to investigate which regulatory mechanisms acting on OSN axon terminals

were controlled bilaterally to alter glomerular output.

Dopamine D_2 receptors modulated by contralateral glomerular input mediate

presynaptic inhibition

Presynaptic inhibition regulates neurotransmission in olfactory glomeruli (McGann

2013). Discrete populations of juxtaglomerular neurons that release GABA or dopamine

activate $GABA_B$ or D_2 receptors present in presynaptic terminals of OSNs to lower glutamate

release (Wachowiak et al. 2005; Wachowiak and Cohen 1999). Evidence points out that this is

a tonically active mechanism (Pírez and Wachowiak 2008), thus meaning the normal

processing of odor information relies on a certain level of inhibition of glomerular output. To

evaluate the possible involvement of dopamine or GABA in the potentiation of LFP_{peaks}

observed after contralateral olfactory nerve transection, we locally applied antagonists of D_2

and $GABA_B$ receptors to the recorded glomerulus. Odor evoked responses were enhanced by

raclopride, a D_2 antagonist, in tadpoles with their olfactory pathways intact (Fig. 6A). The

average negativity of 73 ± 5 µV (n=8) experimented a gradual increase in the presence of 300

nM raclopride that reached 128 ± 16 µV, 20 min after its application (Fig. 6B, p=0.012, paired

t-test). The potentiation of odor evoked responses caused by the D_2 antagonist was not

observed in tadpoles with the contralateral olfactory nerve transected. Baseline negativities of

133 ± 24 µV (n=11) were minimally increased by raclopride, changing to 156 ± 33 µV (n=11,

p=0.16, paired t-test), 20 min after drug application.

The time course of the increase in LFP_{peaks} observed in control tadpoles was well

described by a Hill equation, reporting a time required to reach the 50% of the maximal increase

of approximately 20 min (Fig. 6C). The time window describing the effect of raclopride could

be attributed to disinhibition, since the primary target of presynaptic D_2 receptor activation is

the decrease of cAMP levels and a consequent lowering of neurotransmitter release (Kaneko

and Takahashi 2004). The potentiation mediated by raclopride was comparable to the increase

in LFP_{peaks} caused by olfactory nerve transection (Fig. 6B, p=0.46, unpaired t-test). These

results indicated that the tonic inhibition of glomerular output was comparably reduced either

by directly inhibiting D_2 receptors, or, by surgically transecting the contralateral olfactory

nerve. Further evidence for the involvement of D_2 receptors was obtained by simultaneously

imaging GCaMP6s fluorescence and LFP signals (Fig. 6D). In the illustrated example

raclopride shortened the time constant (τ) to reach the calcium peak from 1.3 s to 0.5 s and

potentiated LFP changes by 50%, thus reproducing the effect of contralateral olfactory nerve

transection on glomerular input and output, respectively (Figs. 3C and 5F). On average

raclopride reduced τ from 1.27 s to 0.8 s (n=4 tadpoles, p=0.02, paired t-test), evidencing that

the potentiation of glomerular responses was mediated by a faster buildup of presynaptic

calcium levels that took place after antagonizing the inhibitory action of D_2 receptors.

Finally, we explored whether presynaptic $GABA_B$ receptors activated by

juxtaglomerular neurons (McGann 2013) could mediate a synergic effect to D_2 receptors. This

possibility was investigated by applying the selective $GABA_B$ antagonist CGP-36742. LFP_{peaks}

remained unchanged (Figs. 6E and F, p=0.77, paired t-test). A significant tonic inhibition by

GABA was thus ruled out, revealing dopamine was the main neurotransmitter inhibiting

glomerular output in the glomerular layer of *Xenopus* tadpoles.

Effect on glomerular responses of the partial elimination of mirror olfactory sensory

neurons.

To shed light on the synaptic connectivity involved in the contralaterally driven

potentiation of glomerular output we modified the 2Phatal technique (Hill et al. 2017) to

eliminate groups of GFP positive OSNs using a conventional confocal microscope (Fig. 7A).

Tadpoles were placed during 15 minutes in *Xenopus* water containing 5 µg/mL Hoechst 33342,

which is a membrane permeable dye with affinity for nucleic acids, to label all cells found

within the olfactory epithelium. Damage was exerted by photobleaching groups of cells found

in regions of interest (ROIs) that contained two or more GFP positive OSNs. Efficient

photobleaching of the nuclear label caused cell death, which was certified by the observation

of condensed nuclei after 24 hours (Fig. 7A). The use of the confocal microscope precluded

the achievement of single cell ablation but supported the robust elimination of cells within

ROIs. Olfactory epithelial cells found outside the selected areas remained unaffected.

Photobleaching was typically carried out in 4 to 6 ROIs, thus resulting in the elimination of 10

to 15 OSNs. Considering an epithelium contained on average ~30 GFP positive OSNs, the

protocol led to an estimated 30 to 50% reduction of OSNs innervating the right GFP labeled

glomerulus. Since the viability of tadpoles subjected to a more extensive reduction of

fluorescent OSNs was compromised, all animals investigated still had a significant number of

functional GFP labelled OSNs after photoablation.

A characteristic of the recorded odor evoked glomerular responses was their

reproducibility. For a given tadpole, LFP negativities occurred after successive stimulations

with a comparable profile (Figs. 1B, 2A, 4A), and showed an average variance in their

amplitude of 386 ± 96 μV^2 (n=17). Variance of the responses obtained increased more than

three-fold in tadpoles subjected to transection of the olfactory nerve (see for example Fig. 3B),

reaching an average value of 1488 ± 280 μV^2 (n=14). An enhancement of variance was thus

associated to the potentiation observed in response to olfactory nerve transection. Incubation

of tadpoles with Hoechst 33342 neither modified the amplitude nor the variance of LFP_{peaks}

(Fig. 7B). LFP negativities in tadpoles presenting a decimated population of contralateral GFP

positive OSNs showed an increase in LFP_{peaks} from 80 ± 5 μV (n=17) to 89 ± 7 μV (n=10, p=0.32,

unpaired t-test) and in variance to 1024 ± 475 μV^2 (n=10, p=0.21, unpaired t-test).

The non-significant changes in LFP_{peaks} amplitude and variance induced by selective

photoablation could be attributed to the limited suppression of GFP labeled OSNs and it was

not possible to establish through this experimental approach the effect of the complete

suppression of cognate neurons. LFP_{peaks} variance and amplitude were linearly related (r=0.65)

considering control and unilaterally transected tadpoles together (Fig. 7C) and in this context,

the reduction between 1/3 and 1/2 of the mirror contralateral glomerular input induced a right-

shift compared to control values. Considering that the photoablation procedure left most of

OSNs present in the olfactory epithelium intact, the observed displacement suggests that

topographical relationships established bilaterally could be relevant to establish a

contralaterally driven potentiation. Moreover, our results support that bilateral modulation of

glomerular output occurs within a graded scale that is inversely related to the contralateral

innervation by OSNs.

DISCUSSION

Odor evoked responses were recorded as changes of the LFP in an olfactory glomerulus

labelled in the *Dre.mxn1:GFP X. tropicalis* line (Fig. 1) and showed the following

characteristics: *i)* were mediated by glutamatergic synapses (Figs. 2A-C), *ii)* were triggered by

ipsilateral stimulation of OSNs (Fig. 2F), and, *iii)* showed an amplitude related to the number

of input OSNs (Fig. 3A). These characteristics are comparable to responses recorded in the

glomerular layer of living rats where negativities are supported by the activation of glutamate

receptors and locked to the respiration frequency (Lecoq, Tiret, and Charpak 2009).

Glomerular responses were potentiated by the complete silencing of OSNs projecting to the

contralateral olfactory bulb, thus showing the processing of information in the olfactory

glomeruli of *Xenopus* tadpoles is not exclusively unilateral and is shaped contralaterally. The

observed potentiation was not related to inflammatory mediators associated to injury, because

it was caused by a release of the inhibition made by D_2 dopamine receptors present in OSN

axon terminals.

The similarities between the recordings obtained (Fig. 1B) and those described in rats

(Chaigneau et al. 2007), indicate the presence of evolutionary conserved morphological and

functional features between both species and therefore, the applicability to vertebrates of the

findings here described. Our results achieved using glutamate blockers and uncaging of Rubi-

glutamate show that the onset of LFP changes detected in glomeruli is determined by glutamate

release from OSNs. The ionic bases supporting the recovery phase of negativities are uncertain

but, considering responses obtained by glutamate uncaging were locked to the period defined

by the light pulse, the participation of sustained neurotransmitter release is conceivable. This

view is also supported by imaging of OMP-synaptopHluorin mice, where continuous synaptic

vesicle exocytosis occurs over several seconds in OSN axon terminals (Petzold et al. 2008).

The mammalian and amphibian olfactory bulbs display comparable anatomical and

cellular organization (Manzini, Schild, and Di Natale 2022) but, it is yet unknown whether the

three classes of juxtaglomerular neurons present in rodents: periglomerular, short-axon and

external tufted, also exist in *Xenopus*. In mice, short-axon cells are the main source of

dopamine to olfactory glomeruli (Kiyokage et al. 2010), however, there is a lack of evidence

for such a discrete population in the olfactory bulb of *Xenopus* tadpoles. Dopaminergic neurons

are distributed around olfactory glomeruli after NF stage 49 (González et al. 1994) and in adult

Xenopus, type-I tyrosine hydroxylase positive cells extensively innervate the glomerular layer

(Boyd and Delaney 2002). These morphological observations indicate that dopaminergic

neurons exist in the glomerular layer of amphibia, but their specific functional properties are

unknown. As we did not find evidence for the $GABA_B$ mediated inhibition described in rats

(Wachowiak et al. 2005), our results support that dopamine is the main inhibitory

neurotransmitter in the glomerular layer of *Xenopus* tadpoles.

To understand how tonic dopamine release in the glomerular layer of *Xenopus* tadpoles

is affected by the number of contralateral OSNs, it is necessary to have a comprehensive

understanding of olfactory bulb connectivity in *X. tropicalis* tadpoles which is currently

lacking. Interneurons of the amphibian olfactory bulb might not display the degree of

specialization found in rats or mice, however, there is evidence suggesting that type-1 tyrosine

hydroxylase positive interneurons are morphologically similar to external tufted cells present

in rodents (Boyd and Delaney 2002). External tufted cells show rhythmic activity (Hayar et al.

2004), are responsible for constructing a map of homologous glomeruli through intrabulbar

connections (Lodovichi, Belluscio, and Katz 2003) and act as gatekeepers of glomerular output

(Banerjee et al. 2015). Therefore, if the dopaminergic neurons present in the glomerular layer

of *Xenopus* tadpoles play analogous roles to external tufted cells, our results call for

considering this group of interneurons as candidates to exert the contralateral regulation of

glomerular output.

Here, we show that the formation of an odor map in *Xenopus* tadpoles relies on a

balanced contribution of the glomerular responses occurring in both olfactory bulbs. Tonic

activation of presynaptic D_2 receptors, likely mediated by dopaminergic juxtaglomerular

neurons, drives to a constant inhibition of glomerular output. Dopamine release is affected by

the number of operative contralateral OSNs so that, in the extreme situation where all mirror

olfactory pathways are silenced, dopamine secretion decreases and most of the inhibition is

released (Fig. 7D). The consequence is an immediate increase of glomerular responses in the

non-damaged pathway, which results in a compensated output signal. A possible direct

contralateral targeting by OSNs, as described for the γ-glomerulus present in *X. laevis* tadpoles

(Kludt et al. 2015), was ruled because the population labelled in the *Dre.mxn1:GFP* showed

exclusively a unilateral distribution. Interhemispheric connections targeting dopaminergic

juxtaglomerular neurons could be established by axons of mitral cells as reported in zebrafish

(Kermen et al. 2020). However, the participation of neurons of the lateral pallium cannot be

disregarded as it is a main center for olfactory processing (Moreno et al. 2008). Either one of

these projecting neurons could be targeting contralateral dopaminergic neurons found in the

glomerular layer of *X. tropicalis* tadpoles (Fig. 7D). Although the existence of an

intraglomerular circuit controlling dopamine release cannot be disregarded (Banerjee et al.

2015), and, considering the differences existing between olfactory bulb interneurons present in

amphibia and mammals (Imamura, Ito, and LaFever 2020), we modeled the simplest scenario

based in only one cell type of juxtaglomerular neuron.

The in vivo evidence presented shows that presynaptic D_2 receptors tonically exert an

inhibitory action on calcium buildup in OSN axon terminals. This finding is well aligned with

previous observations obtained in mouse brain slices (Ennis et al. 2001) and describes a

relevant biological scenario for the role of dopamine inhibition in the glomerular layer of the

olfactory bulb. Both the suppression of contralateral OSN inputs and the local injection of the

D_2 antagonist raclopride enhanced glomerular output, a finding that can be explained by the

tight coupling existing between calcium influx and neurotransmitter release in OSN axon

terminals (Wachowiak et al. 2005). The likely target of dopamine are N-type voltage gated

calcium channels because they are present in presynaptic terminals in olfactory glomeruli

(Weiss et al. 2014) and they can be inhibited through multiple mechanisms activated by D_2

receptors (Kisilevsky and Zamponi 2008).

Taken altogether our results illustrate a homeostatic mechanism used to compensate a

reduced contribution of contralateral OSNs characterized by an enhancement of glomerular

output. It is possible that the observed compensation maximizes the role of a pathway relevant

to the normal encoding of olfactory information upon transection of an olfactory nerve. Under

physiological conditions the contralateral modulation of presynaptic D_2 receptors could be used

to correct imbalances in the firing of mitral cells collecting information from bilaterally related

glomeruli. Although we can only speculate about such functional role, our lines of evidence

show that the mechanism here described is efficiently used for achieving a bilateral

compensation of glomerular output in a vertebrate species.

FIGURE LEGENDS

Figure 1. Recording of odor evoked responses in a genetically defined glomerulus. A)

Schematic diagram illustrating the experimental preparation. A puff of 200 µM methionine was

applied during 100 ms to the left olfactory epithelium of *Dre.mxn1:GFP Xenopus tropicalis*

tadpoles. Changes in the Local Field Potential (LFP) were measured ipsilaterally with an

electrode targeted to the lateral glomerulus formed by axon terminals of olfactory sensory

neurons (OSNs) expressing GFP. The inset shows a confocal projection illustrating labelled

OSNs (s: soma, a:axon) and the bilateral formation of glomeruli (g). The *Dre.mxn1* promoter

also drives GFP expression in the endothelial cells of some blood vessels (e). **B)** Representative

glomerular odor evoked response (black) obtained by averaging individual responses (gray) of

the LFP following stimulation (arrow). The asterisk shows a positivity that appeared in 37% of

the recordings and preceded the characteristic negativity associated to glomerular activation.

C) Negative deflections of LFP (individual responses, gray; average trace, black) were

observed when the recording electrode was placed in the GFP labelled glomerulus but

disappeared in the mitral cell layer (MC, red traces). **D)** Experiment showing LFP recordings

performed in four different locations of the glomerular layer spaced by 50 µm. The

characteristic odor evoked response was observed in only one of the positions tested. The

representative glomerular odor evoked response (black) was obtained by averaging individual

responses (gray) following stimulation with methionine (arrow). E) Hyperstack projections of

the left olfactory bulb of three different *Dre.mxn1:GFP Xenopus tropicalis* tadpoles. The

lateral glomerular cluster (L) is always obvious, and some medial projections (M) are evident

in two of the illustrated examples. Color scale indicates dorsoventral disposition.

Figure 2. Odor evoked responses are mediated by glutamatergic neurotransmission. A)

The representative glomerular odor evoked response (black) was obtained by averaging

individual responses (gray) of the Local Field Potential **(**LFP) following stimulation (arrow).

In this example, the pipette solution contained 100 µM CNQX and upon local injection of 1

µL, there was a reduction in the amplitude of LFP negativities. **B)** Mean LFP changes obtained

under control conditions (n=21) were reduced after the application of 100 µM CNQX (n=8),

100 µM AP5 (n=5), or both 100 µM CNQX and 100 µM AP5 (n=8). **B)** Box plot illustrating

how the initially recorded peak negativities were affected by the application of 100 µM CNQX,

or, 100 µM AP5 together with 100 µM CNQX. Boxes represent the median (horizontal line),

25th to 75th quartiles, and ranges (whiskers) of the indicated experimental groups. Statistical

differences were evaluated using paired t-test. **C**) Rubi-glutamate was injected in the GFP

labelled glomerulus and locally uncaged with a 500 ms pulse of blue light (circle). The

example shows the change in LFP induced by two flashes (squares) delivered at an interval of

9 seconds. **E)** Application of picrotoxin, a GABA$_A$ antagonist, did not modify odor evoked

changes. Recordings show average responses (n=9). Statistical differences were evaluated

using paired t-test**. F)** Odor evoked LFP changes were exclusively triggered by ipsilateral

stimulation. Individual responses are indicated in gray, with the representative average

response in black. Contralateral stimuli (yellow) did not modify the LFP, as shown in the

average representative response (brown).

Figure 3.   Potentiation of odor evoked responses by transection of the contralateral

olfactory nerve. A) The number of olfactory sensory neurons (OSNs) and the amplitude of

odor evoked negative deflections of the Local Field Potential (LFP) were related to olfactory

nerve width according to linear and exponential functions, respectively. Individual data points

are represented by circles (n=48). Each bin indicates the mean ± standard error of n=6 tadpoles.

The dotted line indicates the steady-state LFP amplitude reached during development. **B)**

Representative odor evoked response (red; individual responses, gray) obtained 24 h after

contralateral nerve transection. A solution of 200 µM methionine was used as stimulus (arrow).

C) Odor evoked LFP changes exhibit amplitudes above the expected values (dotted line as in

A) after contralateral olfactory nerve transection at the indicated time points. The dots represent

the mean ± s.e.m obtained 2 to 7 hours (n=10), 1 to 2 days (n=14), and 10 to 11 days (n=11)

post-injury. There was a 75% increase in animals recorded 1 to 2 days after transection of the

contralateral olfactory nerve (red arrow) compared to control tadpoles (dotted line). **D)** Dots

(mean ± s.e.m.) connected by a line illustrate odor evoked glomerular responses at the indicated

times after injury. The superimposed violin plot displays individual data. Most LFP_{peak} values

are above the level expected for the developmental period studied (dotted line as in A).

Figure 4. The potentiation of odor evoked responses is not mediated by injury derived

cues. A) Odor evoked Local Field Potential (LFP) changes were recorded by an electrode

targeted to the GFP-positive glomerulus of *Dre.mxn1:GFP* tadpoles one to two days after

bilateral transection of optic nerves. **B)** Peak LFP negativities recorded in tadpoles with

sectioned optic nerves did not exhibit the characteristic potentiation observed after transection

of the contralateral olfactory nerve, as they remained within the range of values observed

during normal development (solid line, as in Fig. 3A). Bins indicate mean±s.e.m., circles show

individual values. **C)** Imaging of reactive oxygen species (ROS) two hours after transectioning

one olfactory nerve (arrow). The ratio between the fluorescence emitted by HyPer-YFP when

excited at 488 nm and 405 nm is indicated in pseudocolor. Notice that ROS were increased at

the injury site but remained at basal levels in both olfactory bulbs as indicated by the box plot.

Each circle shows values collected in a single tadpole. **D)** Block of ROS production by

incubating tadpoles with 200 μM apocynin (n=10) or 2 μM diphenyleneiodonium (DPI, n=5)

did not modify the amplitude of odor evoked LFP responses recorded 24 h after contralateral

olfactory nerve transection (n=10). Boxes represent the median (horizontal line), 25th to 75th

quartiles, and ranges (whiskers) of the indicated experimental groups.

Figure 5. The presynaptic component of glomerular activation is affected by damage to

contralateral olfactory sensory neurons. A) Application of a puff of 200 μM methionine to

the olfactory epithelium activates a set of glomeruli in the ipsilateral olfactory bulb (arrows) of

*tubb2b:GCaMP6*s tadpoles. Images show the relative changes in GCaMP6s fluorescence

(ΔF/F) obtained after two sequential stimulations carried out in a single tadpole. **B)** Time

course of the responses detected in the glomeruli indicated in A). **C)** Example showing the

simultaneous recording of Local Field Potential (LFP) and changes in GCaMP6s fluorescence

in the region targeted by the electrode. Colored traces and gray traces show the change in

GCaMP6s fluorescence (ΔF/F) and LFP respectively observed after three sequential

applications of 200 μM methionine. Black traces show the average ΔF/F and LFP responses.

D) Kinetics of the change in LFP observed in tadpoles with the contralateral olfactory nerve

transected between 2 h and 48h prior to recording. **E)** Intracellular calcium increases detected

in glomeruli of control tadpoles with intact olfactory pathways (35 glomeruli, 10 tadpoles,

black), and, in tadpoles subjected to the transection of the contralateral olfactory nerve (10

glomeruli, 3 tadpoles, red). Each trace indicates the response of a glomerulus to a single

stimulus. Solid lines and error bars indicate mean ± s.e.m. **F)** Calcium transients detected in

tadpoles with an olfactory nerve transected showed a larger amplitude and a rising phase with

a shorter time constant (τ). Boxes in D) and F) represent the median (horizontal line), 25th to

75th quartiles, and ranges (whiskers) of the indicated experimental groups. Statistical

differences in D) and F) were evaluated using paired and unpaired t-tests, respectively. Circles

in D) indicate values obtained for a single tadpole and in F) refer to a single glomerulus.

Figure 6. Contralateral input modulates presynaptic inhibition mediated by dopamine D₂

receptors and is involved in the potentiation of glomerular responses. **A)** Recordings

obtained in a control tadpole showing how the amplitude of Local Field Potential (LFP)

responses (gray traces) obtained during a baseline period of 8 minutes increase in a time-

dependent manner after local application of 300 nM raclopride, a D_2 receptor antagonist. **B)**

Box plot showing the effect of 300 nM raclopride (blue) on the amplitude of LFP responses

recorded in tadpoles with full capacity to process odors (control, n=8, black) and tadpoles

subjected to the transection of the contralateral olfactory nerve (n=11, red). Boxes represent

the median (horizontal line), 25th to 75th quartiles, and ranges (whiskers) of the indicated

experimental groups. The effect of raclopride weas evaluated using the paired t-test and the

comparison between control and transected groups was performed using the unpaired t-test.

C) Relative change in LFP responses induced by 300 nM raclopride in control tadpoles and in

tadpoles subjected to the transection of the contralateral olfactory nerve. Dots represent mean

± s.e.m. The solid black line illustrates the fit to a Hill equation, defining a T_{50} at 20 minutes.

D) Simultaneous recording of LFP and changes in GCaMP6s fluorescence in a

tubb2b:GCaMP6s tadpole. Gray traces and light blue traces show individual responses to

sequential stimulations before and after application of 300 nM raclopride, respectively**.**

Average responses are shown in black and dark blue. **E)** Application of CGP-36742, a GABA_B

receptor antagonist, did not modify LFP responses. The box plot compares the amplitude of

LFP changes recorded before (gray) and 20 min after local application of 300 μM CGP-36742

(green). Statistical differences were evaluated using paired t-test. **E)** Time course of relative

LFP changes induced by 300 μM CGP-36742. Dots represent mean ± s.e.m (n=10).

Figure 7. Changes in odor evoked responses caused by selective photoablation of

olfactory sensory neurons innervating the homologous contralateral glomerulus. A) Odor

evoked Local Field Potential (LFP) changes were recorded one day after the selective

elimination of olfactory sensory neurons (OSNs) located in the right nasal cavity. After the

identification of GFP-positive OSNs in the epithelium labeled with the nuclear marker Hoechst

33342, regions containing ≥2 fluorescent neurons were identified and photobleached. Cell

targeting was confirmed by the suppression of the nuclear label. Only cells found within the

photobleached areas exhibited fragmented nuclei (arrows) 24 hours after photobleaching. **B)**

Examples showing odor evoked responses recorded in a tadpole incubated with Hoechst 33342

(black, average), and in a different tadpole 24 hours after the photobleaching of selected regions

in the contralateral olfactory epithelium (violet, average). **C)** Plot of variance against the

amplitude of LFP_{peaks} in three experimental groups of tadpoles: control (n=17), photoablated

(n=10) and subjected to unilateral olfactory nerve transection (n=14). Circles show individual

values and dots indicate mean±s.e.m. The line shows a fit through values of control and

unilaterally transected tadpoles (r=0.65). **D)** Model proposed for the bilateral modulation of

glomerular output in the olfactory bulb of *Xenopus* tadpoles. Juxtaglomerular neurons display

tonic dopamine release that inhibits glomerular output by activating presynaptic D2 receptors

present in OSNs (dotted box). The constant presence of dopamine within glomeruli is favored

by the activity of the contralateral olfactory bulb. When the contribution of the contralateral

pathway is suppressed, dopamine release diminishes, and glomerular responses become

potentiated. The contralateral modulation of the tonic activity of dopaminergic

juxtaglomerular neurons corrects for input differences and equalizes the synaptic output of

olfactory glomeruli to achieve a bilaterally balanced transfer of information. Axons sent by

mitral cells could mediate the pathway described but, the involvement of neurons found in

higher brain regions and participating in the processing of olfactory information, such as those

of the lateral pallium, could also be considered.

ACKNOWLEDGEMENTS

This work was sponsored by the Ministry of Science, Innovation and Universities (MICIU/AEI), grant PID2021-124536NB-I00 (A.L), co-funded by the European Regional Development Fund (ERDF), "a way of making Europe". The work was also supported by two Whitman Fellowships awarded to A.L in 2023 and 2024 (Marine Biological Laboratory, University of Chicago). The authors thank the institutional support from the María de Maeztu Unit of Excellence, Institute of Neurosciences, University of Barcelona, CEX2021-001159-M (Ministry of Science, Innovation and Universities) and the CERCA Program of Generalitat de Catalunya. A.L. is a Serra Húnter fellow.

REFERENCES

Banerjee, A., F. Marbach, F. Anselmi, M. S. Koh, M. B. Davis, P. Garcia da Silva, K. Delevich, H. K. Oyibo, P. Gupta, B. Li, and D. F. Albeanu. 2015. "An Interglomerular Circuit Gates Glomerular Output and Implements Gain Control in the Mouse Olfactory Bulb." *Neuron* 87 (1):193-207. doi: 10.1016/j.neuron.2015.06.019.

Belluscio, L., and L. C. Katz. 2001. "Symmetry, stereotypy, and topography of odorant representations in mouse olfactory bulbs." *J Neurosci* 21 (6):2113-22. doi: 10.1523/JNEUROSCI.21-06-02113.2001.

Boyd, J. D., and K. R. Delaney. 2002. "Tyrosine hydroxylase-immunoreactive interneurons in the olfactory bulb of the frogs Rana pipiens and Xenopus laevis." *J Comp Neurol* 454 (1):42-57. doi: 10.1002/cne.10428.

Browne, L. P., A. Crespo, and M. S. Grubb. 2022. "Rapid presynaptic maturation in naturally regenerating axons of the adult mouse olfactory nerve." *Cell Rep* 41 (10):111750. doi: 10.1016/j.celrep.2022.111750.

Byrd, C. A., and G. D. Burd. 1991. "Development of the olfactory bulb in the clawed frog, Xenopus laevis: a morphological and quantitative analysis." *J Comp Neurol* 314 (1):79-90. doi: 10.1002/cne.903140108.

Chaigneau, E., P. Tiret, J. Lecoq, M. Ducros, T. Knöpfel, and S. Charpak. 2007. "The relationship between blood flow and neuronal activity in the rodent olfactory bulb." *J Neurosci* 27 (24):6452-60. doi: 10.1523/JNEUROSCI.3141-06.2007.

Daume, D., T. Offner, T. Hassenklöver, and I. Manzini. 2022. "Patterns of." *Front Neuroanat* 16:914281. doi: 10.3389/fnana.2022.914281.

Ennis, M., F. M. Zhou, K. J. Ciombor, V. Aroniadou-Anderjaska, A. Hayar, E. Borrelli, L. A. Zimmer, F. Margolis, and M. T. Shipley. 2001. "Dopamine D2 receptor-mediated presynaptic inhibition of olfactory nerve terminals." *J Neurophysiol* 86 (6):2986-97. doi: 10.1152/jn.2001.86.6.2986.

Fino, E., R. Araya, D. S. Peterka, M. Salierno, R. Etchenique, and R. Yuste. 2009. "RuBi-Glutamate: Two-Photon and Visible-Light Photoactivation of Neurons and Dendritic spines." *Front Neural Circuits* 3:2. doi: 10.3389/neuro.04.002.2009.

Gaudin, A., and J. Gascuel. 2005. "3D atlas describing the ontogenic evolution of the primary olfactory projections in the olfactory bulb of Xenopus laevis." *J Comp Neurol* 489 (4):403-24. doi: 10.1002/cne.20655.

González, A., O. Marín, R. Tuinhof, and W. J. Smeets. 1994. "Ontogeny of catecholamine systems in the central nervous system of anuran amphibians: an immunohistochemical study with antibodies against tyrosine hydroxylase and dopamine." *J Comp Neurol* 346 (1):63-79. doi: 10.1002/cne.903460105.

Hayar, A., S. Karnup, M. Ennis, and M. T. Shipley. 2004. "External tufted cells: a major excitatory element that coordinates glomerular activity." *J Neurosci* 24 (30):6676-85. doi: 10.1523/JNEUROSCI.1367-04.2004.

Hill, R. A., E. C. Damisah, F. Chen, A. C. Kwan, and J. Grutzendler. 2017. "Targeted two-photon chemical apoptotic ablation of defined cell types in vivo." *Nat Commun* 8:15837. doi: 10.1038/ncomms15837.

Holl, A. M. 2018. "Survival of mature mouse olfactory sensory neurons labeled genetically perinatally." *Mol Cell Neurosci* 88:258-269. doi: 10.1016/j.mcn.2018.02.005.

Imamura, F., A. Ito, and B. J. LaFever. 2020. "Subpopulations of Projection Neurons in the Olfactory Bulb." *Front Neural Circuits* 14:561822. doi: 10.3389/fncir.2020.561822.

Kaneko, M., and T. Takahashi. 2004. "Presynaptic mechanism underlying cAMP-dependent synaptic potentiation." *J Neurosci* 24 (22):5202-8. doi: 10.1523/JNEUROSCI.0999-04.2004.

Karnup, S. V., A. Hayar, M. T. Shipley, and M. G. Kurnikova. 2006. "Spontaneous field potentials in the glomeruli of the olfactory bulb: the leading role of juxtaglomerular cells." *Neuroscience* 142 (1):203-21. doi: 10.1016/j.neuroscience.2006.05.068.

Kermen, F., P. Lal, N. G. Faturos, and E. Yaksi. 2020. "Interhemispheric connections between olfactory bulbs improve odor detection." *PLoS Biol* 18 (4):e3000701. doi: 10.1371/journal.pbio.3000701.

Kisilevsky, A. E., and G. W. Zamponi. 2008. "D2 dopamine receptors interact directly with N-type calcium channels and regulate channel surface expression levels." *Channels (Austin)* 2 (4):269-77. doi: 10.4161/chan.2.4.6402.

Kiyokage, E., Y. Z. Pan, Z. Shao, K. Kobayashi, G. Szabo, Y. Yanagawa, K. Obata, H. Okano, K. Toida, A. C. Puche, and M. T. Shipley. 2010. "Molecular identity of periglomerular and short axon cells." *J Neurosci* 30 (3):1185-96. doi: 10.1523/JNEUROSCI.3497-09.2010.

Kludt, E., C. Okom, A. Brinkmann, and D. Schild. 2015. "Integrating temperature with odor processing in the olfactory bulb." *J Neurosci* 35 (20):7892-902. doi: 10.1523/JNEUROSCI.0571-15.2015.

Lecoq, J., P. Tiret, and S. Charpak. 2009. "Peripheral adaptation codes for high odor concentration in glomeruli." *J Neurosci* 29 (10):3067-72. doi: 10.1523/JNEUROSCI.6187-08.2009.

Lodovichi, C. 2021. "Topographic organization in the olfactory bulb." *Cell Tissue Res* 383 (1):457-472. doi: 10.1007/s00441-020-03348-w.

Lodovichi, C., L. Belluscio, and L. C. Katz. 2003. "Functional topography of connections linking mirror-symmetric maps in the mouse olfactory bulb." *Neuron* 38 (2):265-76. doi: 10.1016/s0896-6273(03)00194-6.

Love, N. R., Y. Chen, S. Ishibashi, P. Kritsiligkou, R. Lea, Y. Koh, J. L. Gallop, K. Dorey, and E. Amaya. 2013. "Amputation-induced reactive oxygen species are required for successful Xenopus tadpole tail regeneration." *Nat Cell Biol* 15 (2):222-8. doi: 10.1038/ncb2659.

Manzini, I., C. Brase, T. W. Chen, and D. Schild. 2007. "Response profiles to amino acid odorants of olfactory glomeruli in larval Xenopus laevis." *J Physiol* 581 (Pt 2):567-79. doi: jphysiol.2007.130518 [pii]
10.1113/jphysiol.2007.130518.

Manzini, I., and D. Schild. 2004. "Classes and narrowing selectivity of olfactory receptor neurons of Xenopus laevis tadpoles." *J Gen Physiol* 123 (2):99-107. doi: jgp.200308970 [pii]
10.1085/jgp.200308970.

Manzini, I., D. Schild, and C. Di Natale. 2022. "Principles of odor coding in vertebrates and artificial chemosensory systems." *Physiol Rev* 102 (1):61-154. doi: 10.1152/physrev.00036.2020.

McGann, J. P. 2013. "Presynaptic inhibition of olfactory sensory neurons: new mechanisms and potential functions." *Chem Senses* 38 (6):459-74. doi: 10.1093/chemse/bjt018.

Menini, A. 2010. "The Neurobiology of Olfaction." In.

Mombaerts, P., F. Wang, C. Dulac, S. K. Chao, A. Nemes, M. Mendelsohn, J. Edmondson, and R. Axel. 1996. "Visualizing an olfactory sensory map." *Cell* 87 (4):675-86. doi: 10.1016/s0092-8674(00)81387-2.

Moreno, N., R. Morona, J. M. López, L. Dominguez, M. Muñoz, and A. González. 2008.
"Anuran olfactory bulb organization: embryology, neurochemistry and hodology."
B*rain Res Bull 7*5 (2-4):241-5. doi: 10.1016/j.brainresbull.2007.10.027.

Murphy, G. J., L. L. Glickfeld, Z. Balsen, and J. S. Isaacson. 2004. "Sensory neuron signaling
to the brain: properties of transmitter release from olfactory nerve terminals." J
*Neurosci 2*4 (12):3023-30. doi: 10.1523/JNEUROSCI.5745-03.2004.

Nezlin, L. P., and D. Schild. 2000. "Structure of the olfactory bulb in tadpoles of Xenopus
laevis." Ce*ll Tissue Res 3*02 (1):21-9.

Niethammer, P., C. Grabher, A. T. Look, and T. J. Mitchison. 2009. "A tissue-scale gradient of
hydrogen peroxide mediates rapid wound detection in zebrafish." Na*ture 4*59
(7249):996-9. doi: 10.1038/nature08119.

Nieuwkoop, P.D, and J. Faber. 1956. No*rmal table of Xenopus laevis (Daudin). A systematical*
and chronological survey of the development from the fertilized egg till the end of
metamorphosis.: Amsterdam: North-Holland Publishing Company. Guilders.

Petzold, GC, DF Albeanu, TF Sato, and VN Murthy. 2008. "Coupling of neural activity to
blood flow in olfactory glomeruli is mediated by astrocytic pathways." Ne*uron 5*8
(6):897-910. doi: S0896-6273(08)00403-0 [pii]
10.1016/j.neuron.2008.04.029.

Pírez, N., and M. Wachowiak. 2008. "In vivo modulation of sensory input to the olfactory bulb
by tonic and activity-dependent presynaptic inhibition of receptor neurons." J N*eurosci*
28 (25):6360-71. doi: 10.1523/JNEUROSCI.0793-08.2008.

Shepherd, G.M., W.R. Chen, and C.A. Greer. 2004. The *Synaptic Organization of the Brain:*
*Ox*ford University Press.

Terni, B., and A. Llobet. 2021. "Axon terminals control endolysosome diffusion to support
synaptic remodelling." Lif*e Sci Alliance 4* (8). doi: 10.26508/lsa.202101105.

Terni, B., P. Pacciolla, H. Masanas, P. Gorostiza, and A. Llobet. 2017. "Tight temporal coupling
between synaptic rewiring of olfactory glomeruli and the emergence of odor-guided
behavior in Xenopus tadpoles." J C*omp Neurol 52*5 (17):3769-3783. doi:
10.1002/cne.24303.

Terni, B., P. Pacciolla, M. Perelló, and A. Llobet. 2018. "Functional Evaluation of Olfactory
Pathways in Living Xenopus Tadpoles." J V*is Exp (1*42). doi: 10.3791/58028.

Wachowiak, M., and L. B. Cohen. 1999. "Presynaptic inhibition of primary olfactory afferents
mediated by different mechanisms in lobster and turtle." J N*eurosci 19* (20):8808-17.
doi: 10.1523/JNEUROSCI.19-20-08808.1999.

Wachowiak, M., J. P. McGann, P. M. Heyward, Z. Shao, A. C. Puche, and M. T. Shipley. 2005.
"Inhibition [corrected] of olfactory receptor neuron input to olfactory bulb glomeruli
mediated by suppression of presynaptic calcium influx." J N*europhysiol 94* (4):2700-
12. doi: 10.1152/jn.00286.2005.

Weiss, J., M. Pyrski, P. Weissgerber, and F. Zufall. 2014. "Altered synaptic transmission at
olfactory and vomeronasal nerve terminals in mice lacking N-type calcium channel
Cav2.2." Eur *J Neurosci 40* (10):3422-35. doi: 10.1111/ejn.12713.

Weiss, L., I. Manzini, and T. Hassenklöver. 2021. "Olfaction across the water-air interface in
anuran amphibians." Cel*l Tissue Res 38*3 (1):301-325. doi: 10.1007/s00441-020-
03377-5.

Wu, Y., L. Dissing-Olesen, B. A. MacVicar, and B. Stevens. 2015. "Microglia: Dynamic
Mediators of Synapse Development and Plasticity." Tre*nds Immunol 36* (10):605-613.
doi: 10.1016/j.it.2015.08.008.

Figure 1

Figure 2

Figure 3

Figure 4

A

50 μV

1 s

B

O.N. Transection
Optic Nerves Transection

LFP Peak (μV)

0   40   80
O.N. width (μm)

C

OB
ON

max
min

100 μm

2.0

F_{HyPer} OB left / OB right
1.5
1.0
0.5
0.0

Control   Transected

D

LFP Peak (μV)
300
200
100
0

n.s.   n.s.

Transected   + Apocynin   + DPI

Figure 5

Figure 6

Figure 7

A

GFP Hoechst 33342

0 h

Pre-bleach

10 μm

24 h

100 μm

Post-bleach

B

Hoechst 33342

24h after photobleaching

50 μV

1 s

C

Control
O.N. Transection
Photobleaching

D

Bilateral processing
of olfactory information

Unilateral processing
of olfactory information

Mitral
cells
Juxt.
neurons
OSNs