Development of a method for large-scale single-molecule analysis of tau proteoforms

Joly J [1+], Budamagunta V [1+], Zhang Z [1+], Nortman B [1+], Jouzi M [1+], Bhatnagar R [1], Egertson JD [1], Flaster ME [1], Grothe R [1], Guha S [1], Kaneshige K [1], McVey K [1], Nelson N [1], Perera RT [1], Tan SJ [1], Arnott D [2], Lipka J [2], Rougé L [2], Wendorff TJ [2], Kirkpatrick D [2], Rohou A [2], Butler DC [3], Lotz S [3], Forton A [4], Sartori ER [4], Brereton PS [1], Chen K [1], Darcy MA [1], Golnabi HR [1], Hartley R [1], Indermuhl PF [1], Inman CE [1], Jin KM [1], Katsyuk S [1], Kota R [1], Lowry B [1], Menger JC [1], Miller DA [1], Newman MR [1], Ogunyemi A [1], Robinson JK [1], Steiner N [1], Sun J [1], Wang L [1], Wang Z [1], Wilcox SK [1], Nautilus Biotechnology*, Kapp GT [1], Patel S [1], Temple S [3], Bertucci T [3], Blanchard J [4], Huhmer A [1], Sankar S [1], Juneau K [1#], Mallick P [1#]

Affiliations:
[1] Nautilus Biotechnology, San Carlos, CA 94070
[2] Genentech, South San Francisco, CA 94080
[3] Neural Stem Cell Institute, 150 New Scotland Ave. Albany NY 12208
[4] The Ronald Loeb Center for Alzheimer's Disease at Mt. Sinai, New York, NY 10029;
* A list of authors and their affiliations appears at the end of the paper
+ These authors contributed equally
# To whom correspondence should be addressed

ABSTRACT

Proteins exist as diverse proteoforms resulting from a combination of genetic variation, alternative splicing, and post-translational modifications. Current methods struggle to capture this complexity at the single-molecule level. Here we introduce Iterative Mapping of Proteoforms (IMaP), a method that enables the massively-parallel interrogation of millions to billions of single-protein molecules through iterative probing with fluorescently labeled antibodies. Using 12 site-specific antibodies, the method is capable of measuring 2^{12} (4,096) potential proteoform groups. We used IMaP to measure proteoform group profiles of the tau protein, a key player in neurodegenerative diseases, using two pan anti-tau antibodies (Tau-13, Tau-216), three isoform-specific antibodies (Anti-0N, Anti-2N, Anti-4R), and seven phosphosite-specific antibodies (Anti-pT181, Anti-pS202+pT205, Anti-pT205, Anti-pS214, Anti-pT217, Anti-pT231, and Anti-pS396). The method demonstrates high sensitivity (detecting proteoforms at 0.1% abundance), high reproducibility (median CV <5.5%), and broad dynamic range (>3 orders of magnitude), outperforming conventional techniques in resolving closely related proteoform groups. We demonstrated that the method can be used on relevant biological samples by examining various neuronal models (iNeuron cells, organoids, miBrains, and mouse brains) and human samples. This examination revealed 130 distinct tau proteoform groups with as many as six phosphorylation events. The non-random distribution of these phosphorylation events suggests ordered and site-specific modification processes rather than random, stochastic accumulation. Certain combinations of phosphorylation events were more abundant than others; for example, pT217 preferentially co-occurred with pT181. In validating the applicability of the assay to human disease samples, we noted a specific pattern of multiple phosphorylation events in an advanced Alzheimer's disease patient that suggests a sequential pathway of pathological tau modification. Iterative Mapping of Proteoforms provides insights into proteoform complexity at the single-molecule level, with significant implications for understanding protein regulation in neurodegenerative diseases and beyond.

INTRODUCTION

Cellular regulation is driven by a complex interplay between protein synthesis, transport, modification, and degradation. The most prevalent proteomics methods to measure these processes, shotgun mass spectrometry (MS) [1] and affinity-based profiling [2] [3], focus on estimating a sample's protein composition by measuring the abundance of peptide fragments or the binding of specific affinity reagents.

While these methods can also quantify peptides or individual sites that contain sequence variants or post-translational modifications (PTMs), they are unable to capture one of the most critical aspects of protein-based regulation—combinations of protein alterations. Specifically, a combination of genetic polymorphisms, RNA splice variants, proteolytic processing, and a multiplicity of PTMs may collectively generate millions of different proteoforms [4]. Protein alterations can influence a protein's structure [5], localization [6], interactions [7], stability [8], and enzymatic function [7,9]. However, outside of a few specific examples [10], such as histones [11], very little is known about how proteoforms or a diverse set of proteoforms drive cellular function. Fundamental questions remain, including: 'which of the near-infinite number of possible proteoforms exist?,' 'is there an order and timing by which a given protein molecule acquires multiple modifications?,' and 'how do molecularly heterogeneous sets of proteoforms work in concert to drive cell behavior?'

Despite the many unknowns, recent studies have established connections between proteoform alterations and complex biological processes [12]. Proteoforms have also been implicated in cancers and neurodegenerative disorders [13]. As a result, scientists increasingly emphasize the need to study proteins at the proteoform level rather than relying solely on measurements of protein abundance [12].

The current gap in understanding about the functional impacts of proteoforms arises because their measurement is extremely challenging. Proteoforms are typically present at low abundance, making them difficult to detect directly within complex biological matrices and requiring prefractionation or enrichment[14]. This challenge is compounded by the limitations of conventional separation techniques, which often fail to distinguish structurally similar proteoforms [15]. Even when separated, mass spectrometry—despite its power—struggles to resolve isotopomers, in which PTMs occur at different sites but result in indistinguishable masses or when structural isomers have identical fragmentation patterns [16]. These analytical hurdles are further exacerbated by the requirement for large amounts of highly purified material, which is rarely accessible for low abundance proteoforms in biological samples. Identification is also constrained by

the lack of a comprehensive reference database of human proteoforms, making confident assignment difficult even when high-quality data are available [12]. Together, these challenges hinder systematic proteoform discovery and quantification, slowing efforts to decipher their biological significance.

To date, the most prevalent approach to measure proteoforms is based on top-down mass spectrometry (TDMS), which analyzes intact proteins without enzymatic digestion. TDMS can identify sequence variations and discover novel PTMs and associated proteoforms [13]. TDMS is highly effective for measuring the relative abundance of proteoforms, proteoform families (sets of proteoforms originating from a single gene), or proteoform groups (sets of related proteoforms that share a defined set of observed alterations). TDMS typically achieves this by analyzing ensembles of millions of molecules, but this approach results in reduced sensitivity because of signal dilution across multiple charge states for each isotopologue and lack of chromatographic resolution [17]. Recently, single-ion MS methods [18] that enable the detection of proteoform families have been introduced[19]. However, they are not typically able to easily resolve proteoforms that are not separable by mass or ion mobility. Despite its significant value in proteomics research, the difficulty of analyzing larger proteins creates a discovery bias towards smaller proteins and the complexity of TDMS workflows significantly constrains its practicality for widespread application.

An emerging class of single-molecule methods are anticipated to advance proteoform measurement. Single molecule counting methods have theoretical benefits for being able to measure rare proteoforms extremely sensitively and possibly with simpler experimental workflows than currently achieved with TDMS. In addition, it may be possible to examine modifications existing at multiple sites across a protein [20]. A number of single-molecule peptide sequencing methods cannot measure proteoforms as they rely on protein digestion, similar to shotgun mass spectrometry [21,22]. However, two classes of methods have emerged for single-molecule analysis of proteoforms and proteoform groups: protein fingerprinting and nanopore-based sequencing. Protein fingerprinting methods have been shown to use single-molecule fluorescence resonance energy transfer (FRET) [23] to distinguish isoforms of alpha-Synuclein. Though this analysis enabled isoform-specific identification and quantification in defined mixtures, it has not yet been applied for biological inquiry. In addition, it is unclear how it might examine a multiplicity of PTMs within the same protein molecule. Nanopore-based protein sequencing technologies [24] aim to identify and characterize full-length proteoforms. Recent nanopore technology improvements may allow for sequencing of closely-spaced phosphorylation in individual reads of single molecules and improved identification by rereading individual protein molecules, providing proof-of concept for a

platform for highly accurate protein barcode sequencing and full-length proteoform identification [25] . However, detection of complex and highly charged PTMs remains technically challenging and underexplored. While early nanopore protein sequencing prototypes exist, single-molecule library preparation methods from biological samples that are compatible with nanopore sequencing have not yet been demonstrated. It also remains unclear whether current nanopore systems provide the scale and throughput needed to measure the millions to billions of molecules required to capture the full dynamic range of biological systems.

Here we introduce and apply a new single-molecule method, which we call Iterative Mapping of Proteoforms (IMaP), to overcome the key limitations of methods for quantifying proteoform groups. Iterative Mapping of Proteoforms is designed to reproducibly quantify pre-defined sets of proteoform groups across a wide range of abundances via massively parallel measurement of millions to billions of individual protein molecules.

We developed an assay, based upon the single-molecule Iterative Mapping of Proteoforms method, to measure tau proteoform groups encoded by the human *MAPT* gene. For simplicity, we refer to this "Iterative Mapping of Proteoforms of tau" assay below as the "tau proteoform assay." Tau was chosen as a first exemplar given its critical importance in neurodegenerative diseases and its diverse landscape of splice variants and PTMs. Specifically, tau exists as six splicing isoforms in the adult human brain [26,27] and undergoes diverse PTMs including proteolysis [28], phosphorylation [29], acetylation [30], and ubiquitination [29], resulting in millions of potential tau proteoforms. Recently, the measurement of phosphorylated tau has been shown to have greater diagnostic power than measurements of total tau[31,32]. While much is known about tau, and its various splice-forms and PTMs, very little is known about its proteoform landscape, and how this varies in diverse biological systems.

We first evaluated the tau proteoform assay through a series of control samples of known composition and enriched biological lysates known to contain tau proteoforms associated with epitope-specific probes within in our assay. Next, we demonstrated that the assay can be applied to biological samples by measuring the complex landscape of tau proteoform groups in neuronal tissue model systems of tauopathies, including fronto-temporal dementia (FTD) and Alzheimer's disease (AD). We resolved isoform-specific variation in maturing tau across model systems. We additionally identified which combinations of phosphorylation events are more frequently observed. To demonstrate that the assay can be applied to human-derived brain tissue samples, we analyzed a small number of samples from patients that were either cognitively normal or had AD-

associated impairment. As part of our analysis, we examined if phosphorylation patterns are random or instead suggestive of a specific order of occurrence, finding evidence for both specificity and ordering. These measurements also potentially shed light on routes by which specific proteoforms are generated and provide a foundation for future research investigating which tau proteoforms contribute the most to neural dysfunction and neurodegeneration, and how.

RESULTS

Single-molecule Iterative Mapping of Proteoforms

The method we introduce here, Iterative Mapping of Proteoforms (IMaP) (**Figure 1**), consists of four steps that are outlined in **Figure 1**. In the first step (**Figure 1A**), protein is extracted from tissue lysate, denatured, and functionalized with an amine-reactive methyltetrazine-PEG4-STP (4-Sulfo-2,3,5,6-tetrafluorophenyl) ester (mTz) click reagent. If needed, a sample can be enriched using protein-specific antibodies. Next, functionalized protein is conjugated to specialized nanoparticles that were designed and built using DNA-origami approaches[33]. At most one protein molecule can be conjugated to each nanoparticle. At no point is the protein digested. Next (**Figure 1B**), protein-nanoparticle conjugates are deposited into a lane of a custom microfluidics flow cell with a functionalized glass surface containing millions to billions of optically-resolvable landing pads, as previously described[34]. As the DNA-origami particles are fluorescently labeled, we can determine which coordinates of the array contain a protein-nanoparticle conjugate and which are empty by imaging in the 488-wavelength. In the final step, single molecules are iteratively probed in a massively parallel manner using a series of site-specific fluorescently labeled [35] antibodies (**Figure 1C**). Each iteration consists of a bind, rinse, image, and remove cycle. Specifically, a fluorescently labeled antibody probe is introduced into the flow cell and allowed to incubate. Next, unbound antibody is rinsed from the flow cell and the flow cell is optically imaged [34] to determine where the antibody has bound. Following imaging, bound antibodies are removed from the flow cell, preparing the flow cell for a subsequent iteration. As fluorescent images are acquired, positions of each single-molecule binding event are recorded. Collectively, probing of the single-molecule array produces a series of binary bind/no-bind outcome patterns (single molecule binding traces), which are used to determine the potential identity of the proteoform at each coordinate in the flow cell. The cumulative relative binding across a typical experiment is shown across 36 cycles in **Supp. Fig 1**. Note that the binding signal is stable across the 36 cycles and that there is no significant build-up of background signal. By counting the observed binding patterns and correcting for

antibody false positive and false negative binding, we estimate the quantity of each proteoform within each sample (**Figure 1D**). Proteoform abundances are then reported in parts per million. We note that Iterative Mapping is a fully general method that may have diverse applications beyond the proteoform analysis described herein.

Iterative Mapping of Proteoforms resolves tau proteoforms

We applied the Iterative Mapping of Proteoforms method to build a tau proteoform assay incorporating 12 antibodies that included two pan anti-tau antibodies (Tau-13, Tau-216), three isoform-specific antibodies (Anti-0N, Anti-2N, Anti-4R), and seven phosphosite-specific antibodies (Anti-pT181, Anti-pS202+pT205, Anti-pT205, Anti-pS214, Anti-pT217, Anti-pT231, and Anti-pS396) spanning key isoforms and phosphorylations in tau (**Figure 2A, Supp. Table S3**). 1N isoforms were inferred by lack of binding of the 0N and 2N antibodies. 3R isoforms were inferred by lack of binding of the 4R antibody. Phosphorylation at pS202 was inferred by binding of the Anti-pS202+pT205 antibody and lack of binding of the Anti-pT205 antibody. All other forms and modifications were measured directly from observed binding. An assay with 12 antibodies potentially measures up to 2^{12} (4,096) proteoform groups. Because 0N and 2N cannot both bind concurrently, and because we restrict our focus here to full-length tau, a tau proteoform assay restricted to molecules that have not been truncated enables us to measure 768 ($6*2^7$) potential proteoform groups (**Figure 2A, Supp. Table S3**) of full-length tau. Given the targeted nature of the assay, there may exist additional protein alterations beyond those measured. As such, the assay measures proteoform groups rather than full proteoforms. The antibodies used herein were extensively validated for on-target, off-target, and non-specific binding as described in **Methods**. We included four control samples within each experiment comprising phosphorylated peptides, non-phospho peptides, and full-length tau proteoforms to measure the on and off target binding of each probe within that experiment (**Methods**). Representative images and a set of derived binary single-molecule bound/not-bound traces from the assay are shown in **Figure 2B.** Estimates of proteoform group are extracted from these traces for each molecule. For example, 'Proteoform 2' in **Figure 2B** shows binding with antibodies targeting 0N, pT181, and pS396, thereby identifying the molecule at that position as part of the proteoform group defined by the 0N-3R isoform with a double phosphorylation at sites pT181 and pS396. As noted above, quantification is derived from the assay by counting up the identifications and normalizing for false-positive and false-negative binding as described in **Methods**.

We developed a set of recombinant tau proteoform standards to enable characterization of assay performance. We specifically used recombinantly expressed and purified 0N4R

and 2N4R isoforms of tau, and derivatives thereof (2N-PKA and 0N-ERK2) that had been phosphorylated by either protein kinase A (PKA) or Mitogen-activated protein kinase 1 (ERK2) (**Supp. Methods**). PKA preferentially modifies 2N4R at pS214 with approximately 99% completeness. ERK2 is less specific, dominantly modifying 0N4R at pT181, pS202/T205, and pS396 with approximately 90% phosphorylation efficiency (**Supp. Table S2**). Consequently, samples made with these standards contain a mixture of three dominant proteoforms and a series of proteoforms that arise from incomplete modification (**Figure 2C-left**). To mimic cell-derived tau, these standards can be mixed into a background cell-line that does not contain tau. Studies herein used Expi293F cells for spike-in experiments. The individual estimates of the composition of each standard are shown in the right panel in **Figure 2C**. Each standard has been assigned a color of either navy blue, orange, or light blue. These standards were mixed in a variety of ratios and assayed, demonstrating that the various proteoform groups were readily resolved (**Figure 2C-right**).

To examine how this proteoform reference standard mixture can be measured by other common analytical approaches, we performed western blots (WB). As WB relies on electrophoretic separation and antibody visualization, we expected it to be less precise in resolving complex tau proteoforms. Analysis of an aliquot of a 1:2:3 mixture of three tau proteoforms—0N4R, 2N4R, and 2N4R-PKA—using silver staining successfully resolved the 0N and 2N isoforms (**Supp. Fig. 2A, lanes 2-3**). Detection with the pan-tau and isoform-specific antibodies confirmed the presence of the 0N4R and 2N4R tau isoforms (**Supp. Fig. 2B**) but was unable to differentiate the 2N4R from the 2N4R-PKA proteoform in the lane with a pure standard or in the mixtures. Similarly, the epitope-specific probe anti-pS214 specifically stained the 2N4R-PKA standard (but not the highly phosphorylated control standard 0N4R-ERK2 lacking pS214) as well as the protein standard mixtures (**Supp. Fig. 2C**). However, the specific proteoforms in the mixture and their relative phosphorylation site occupancy remained unresolved. These results affirm that WB provides limited resolution of tau isoforms, while the tau proteoform assay enables detailed proteoform characterization, including the detection of phosphorylated proteoforms in mixtures, establishing it as a higher resolution method for analyzing tau proteoforms.

Accuracy, reproducibility, and dynamic range of the tau proteoform assay

We characterized the quantitative accuracy and reproducibility of the tau proteoform assay using mixtures of three tau protein standards (0N4R, 2N4R, and 2N4R-PKA) at molar ratios of 1:2:3 and 1:1:1. The mean absolute percent error for the 4R tau proteoforms was 9.97% (**Figure 3A**) across 21 replicates. We further examined

reproducibility by analyzing the protein standard mixtures in a range of reproducibility challenges using a nested tree design. Overall, the median CV amongst technical replicates of prepared nanoparticle protein conjugates within the same experimental run was 1.5%. In addition to examining reproducibility within runs of the prepared control mixture within the same experiment, we examined the reproducibility across five batches of flow cells (CV=2.6%) and across three instruments (CV=3.6%). We additionally examined reproducibility of full end-to-end experiments that included library preparation performed 3X by the same operator (CV=5.5%) or as performed by three different operators (CV=5.5%). Though 90% of the intra-run variation ranges from 0-2.3%, percent, 90% of the interoperator variation ranges from 0 to 15.6%. Overall, the assay demonstrated strong reproducibility (**Figure 3B**) across a reproducibility challenge consisting of 297 sets of measurements.

Tau protein is relatively low in abundance in neuronal tissue, constituting approximately 0.09% to 0.7% of total protein [36]. To address this, we developed a bead-based enrichment method using a combination of three distinct anti-tau antibodies (**Figure 1A, Suppl. Table S3**), which is compatible with the single-molecule library preparation workflow. Comparing proteoform abundances between the unenriched mixture and the enriched samples revealed a strong correlation (R^2 = 0.95, stddev 0.2%-1.03%), indicating consistent relative recovery across proteoforms with minimal bias against the specific tau species tested (**Suppl. Fig. 3**).

We established the dynamic range and linearity of tau proteoform quantification using spike-in experiments. In this context, dynamic range refers to the quantification of decreasing percentages of a specific proteoform within a background of another proteoform. A wide dynamic range is critical in proteoform assays as prior studies have shown tau isoforms can vary significantly in abundance [29]. We serially diluted tau standard 2N4R-PKA (representing the tau proteoform 2N4R-pS214) into 0N3R, and performed seven detection cycles with five (anti-Tau13, anti-2N, anti-pS214, anti-4R, anti-Tau216) detection antibodies. Results indicate a linear dynamic range of 3.1, where the limit of quantitation was defined as the lowest point on the titration curve at which the signal to noise ratio is greater than 10 and the coefficient of variation is under 30%. (**Figure 3C**).

In summary, the tau proteoform assay, based upon the Iterative Mapping of Proteoforms method, quantifies tau proteoforms relevant to tauopathy, enabling reproducible detection of biologically meaningful variation using commercially available antibodies. Its analytical performance is linear across approximately three orders of magnitude for

all tau isoforms, with a limit of detection of approximately 1000 ppm. This surpasses WB and offers insights beyond the reach of peptide-based approaches.

Proteoform landscape of tau in models of neurodegenerative disease

Tau plays a central role in maintaining neuronal structure by stabilizing microtubules, particularly in axons. It has been extensively studied due to its pathological aggregation and hyperphosphorylation in FTD and AD, where it forms neurofibrillary tangles—a hallmark of neurodegeneration. Research studies to better understand tau's physiological function and the molecular mechanisms driving its pathological transformation use a wide range of model systems, including iPSC-derived neurons, brain organoids, transgenic mice, and postmortem human brain tissue. Despite extensive study, the proteoform landscape of tau across model systems has yet to be established. To ensure that the assay had sample compatibility with the key sample types used in the field, we applied the tau proteoform assay to investigate what range of proteoforms were present, the characteristics of those proteoforms (e.g. were they dominantly singly modified species), and how those proteoforms differed across model systems. Our major goal in the below studies was to demonstrate that the assay could potentially be used to ask questions across a diversity of relevant model systems. As such, any biological findings introduced in the following sections should be treated as exploratory rather than definitive, particularly in light of the low number of biological replicates examined.

The model systems in the study included a mix of cell-derived models: 2D induced *NGN2*-induced neurons [37] (iNeurons) from four iPSC lines/donors and engineered 3D immuno-glial-neurovascular human multicellular integrated brain (miBrain) model with *APOE3* or *APOE4* genotypes to reflect AD disease risk from two lines, one donor at 8 and 16 weeks after hydrogel setup containing 9 and 17 week aged neurons [38]. We additionally examined patient iPSC-derived cerebral cortical organoids comprised of approximately 16 subtypes of cortical neurons, astrocytes and progenitor cells, from 8 lines derived from 4 donors at 3 and 6 months carrying *MAPT* mutations (IVS10+16, V337M) and their isogenic controls (WT/WT) as a model for frontotemporal dementia (FTD) [39,40] (**Suppl. Table S4**), one hemisphere of brain tissue from a healthy hTau mouse (Mouse brain extract) and a frontal cortex tissue sample from a cognitively normal human (Human brain extract) (see **Methods** for more details about how samples were generated).

We compared the landscape of tau proteoform groups across the model systems, directly assessing their relative abundance. Samples from each model system were analyzed in at least three technical replicates. We note that this study does not attempt

to estimate the biological variation of these model systems, but instead solely attempts to perform the first ever characterization of the proteoform landscape. A heat map of the normalized relative abundance of the tau proteoforms (mean across technical replicates) that were abundant at greater than a 0.1% level across the model systems is shown (**Figure 4A**). Models and proteoforms have been clustered by abundance (z-score normalized per-proteoform group). Each row corresponds to a different proteoform and each column represents a model system. The human brain and mouse brain samples had a lower number of phosphorylations. The stem-cell based models (iNeurons, Organoids and MiBrains) all showed higher extents of multiple phosphorylation. Notably, the cellular models were dominated by the 0N isoform, which is potentially most reflective of the highly phosphorylated tau species described in fetal tissue[41]. The mutant organoids at 3 months, in particular, showed the largest number of phosphorylated species.

Given the quantitative nature of the tau proteoform assay, we compared proteoform counts, uniqueness, and phosphosite occupancy across the diverse set of model systems and tissue samples analyzed. We define a proteoform group as observed if it appears in triplicate in at least one model system at an abundance >0.1% of all full-length tau species (non-proteolytically cleaved). We observed a broad range in the number of detected tau proteoforms, with most samples derived from stem-cell based models yielding between 50–65 distinct proteoform groups and only 23 tau proteoform groups observed in the healthy human brain sample (**Figure 4B**). Across all sample types, we observed 130 unique full-length (non-proteolytically cleaved) tau proteoform groups out of the 768 targeted by our assay (**Supp. Table 5**). We quantified the number of proteoforms shared across the models and found that out of all tau proteoforms detected, 34 proteoforms were found in only one model system, with fewer proteoforms shared amongst two or more model systems (**Suppl. Fig. 4A**). Overall, there was significant variation across the diverse models, with iNeurons appearing to be a large outlier relative to the organoid models (**Suppl. Fig. 5A**). Seven of the eleven proteoforms common to all model systems were phosphorylated, while all proteoforms found in only one model system carried phosphorylation. When comparing the proportion of phosphorylated to unphosphorylated tau molecules across all model systems, we observed the highest modification rates in WT/IVS10-16 cortical organoids (82%), followed by miBrain APOE4 genotype (77%), WT/V337M mutant cortical organoids (76%), miBrain *APOE3* genotype (72%) and lower modification rates for the hTau mouse (37%) and human brain extract (13%) (**Supp. Fig. 4B**).

Next, we investigated the phosphosite occupancy across all model systems and found that the number of phosphorylations per tau molecule was much lower in animal model

systems than in cellular model systems, where up to six co-occurring modifications per tau molecule were detected (**Figure 4C**). In contrast, tau proteoforms from mouse and human brain tissues displayed up to three concurrent phosphorylation events at significantly lower levels. Phosphorylation site occupancy analysis revealed pS396 as the most occupied epitope in cellular systems (60%), while pT181 showed the highest occupancy in animal models (25%). Notably, pT205 and pS202+pT205 were each occupied at ~40% in cellular systems, exceeding their occupancy in the animal model (**Figure 4D**, **Supp. Fig. S4C**).

Age-dependent changes in the tau isoform landscape

Tau splicing is regulated developmentally and only the shortest isoform (0N3R) is present in the fetal brain and maturing cellular models. This isoform may contribute to cytoskeletal plasticity during early neural development by enabling dynamic cytoskeletal adjustments needed in immature neurons [42,27]. Adult tau isoforms containing 4R, on the other hand, are more efficient at promoting microtubule assembly[43]. The inclusion of three isoform-specific probes targeting 0N, 2N, and 4R tau allows our assay to resolve isoform-dependent biological differences. To assure that we can detect and quantify known isoform changes in tissue, we applied the tau proteoform assay to iPSC-derived organoid models containing the *MAPT* mutation [40] (IVS10+16), which alters *MAPT* splicing and results in increased levels of exon 10-containing mRNA (4R). This mutation skews the normally balanced 3R/4R tau ratio of 1:1 in the adult human brain towards 4R but does not affect tau's primary sequence [44]. Nevertheless, this imbalance leads to FTD-related neurodegeneration[44]. Comparing wild-type organoids (WT/WT) at 3 and 6 months of age, we detected an age-dependent increase (3.0-fold) in the mature 4R tau isoform with a similar, age-dependent increase (2.1-fold) for the mutant organoids (**Figure 4E**). Compared to wild-type organoids (WT/WT) at 3 months of age, organoids with the heterozygous *MAPT* mutation (WT/IVS10+16) at the same age had 9.04-fold more 4R tau isoform, increasing to 19.3-fold for the 6-month-old mutant organoids.

Similarly, we determined the age-dependent tau isoform ontogenesis when we measured tau isoforms isolated from 2- and 4-month-old miBrains. The engineered 3D immuno-glial-neurovascular human multicellular integrated brain (miBrain) model combines the enhanced neuronal maturation of organoids with the co-assembly of all six major central nervous system (CNS) cell classes into an engineered human brain mimic supporting the study of genetic risks factors like apolipoprotein E4 (APOE4) in sporadic AD. Here, the ability to detect full length tau protein allowed us to capture all six isoforms and determine the age-dependent ontogenesis of tau with relative

abundances shifting from primarily 0N isoforms to the longer 1N and 2N forms (**Figure 4F**). The ability to directly probe the 4R isoform revealed the presence of the most mature 2N4R tau isoform at 7223 ppm, above the noise level of 1762 ppm. During aging of the miBrains, we observed a marked shift in tau isoform abundance from predominantly 0N3R to more mature isoforms, including 1N3R and 2N3R/4R. This transition from shorter to longer tau isoforms had been previously shown in humans [43]. Comparing the relative isoform abundance of miBrains to human brain, the increased abundance of 2N4R in miBrains at 4 months of maturation more closely aligns with the adult human brain [29].

In summary, we recapitulated age-dependent changes in tau isoforms in a *MAPT* mutant model system of tauopathy. Furthermore, our assay captured isoform-specific changes in miBrain models. These findings show that our assay captures known biological phenotypes while offering greater specificity and broader tau proteoform coverage than traditional methods.

Co-occurrence of multiple phosphorylation across model systems

Multisite phosphorylation may alter tau's conformation and interactions with other proteins. Early stages of abnormal tau processing are believed to be characterized by sequential appearance of pT181 and pT231 followed by pS202 and pT205 then pS214 and pT217 [45]. However, it has not been possible to directly observe if these modifications co-occur on individual molecules or instead are simply spread throughout many molecules. In general, co-occurrence of phosphorylation at one site is influenced by its own relative abundance and its preferred association with another site. Additionally, hyperphosphorylation of tau has often been mentioned as a hallmark of disease progression. However, it is unclear if hyperphosphorylation implies a greater percentage of molecules with any arbitrary phosphorylation, a greater percentage of molecules with a specific phosphorylation, or instead a greater percentage of tau molecules with multiple specific phosphorylations.

As part of demonstrating the potential applications of the tau proteoform assay, we investigated if it were possible to examine if tau has a specific pattern of multiple phosphorylation or if, instead, we simply see a relatively random multiplicity. With 7 anti-phosphoantibodies, there are 21 possible patterns of double phosphorylation and 35 patterns of triple or quadruple phosphorylation independent of isoform. While we observed 18 out of 21 patterns of double phosphorylation across the total set of samples, we only observed 16 of the 35 possible patterns of triple phosphorylation (**Supp. Table S6**). To assess whether multiple phosphorylations follow a random or

preferential path, we calculated how abundant phosphoforms might be expected from a random phosphorylation path. We explored two random models, one in which modification was uniform random, and a site-dependent model that maintained the observed total extent of modification at each site. In **Figure 4G** we highlight this difference in observed phosphorylation patterns (orange) vs those expected at random by either a uniform random model (light blue) or a site independent phosphorylation model (dark blue). The assessment is made irrespective of the isoform in which the modification occurs and defined as phosphoform. Some phosphoforms with double, triple, and quadrupole phosphorylation include sites T205, T217, and pT231 which exhibited greater than expected abundance across model systems, whereas many others showed abundances that agreed with an independent phosphorylation model (**Suppl. Fig. S4D**). For example, for the mutant organoid (IVS10+16) FTD model, we observed 3-fold higher abundance of triple (pT205-pT231-pS396) and quadruple (pT181-pT205-pT231-pTS396) phosphoforms than expected in an independent phosphorylation model (**Figure 4G**).

We next assessed the data for evidence of interactions between sites – where the phosphorylation status of one site influences the phosphorylation of another on the same molecule. We first looked at co-occurrence of phosphosite pairs. Some phosphosites co-occur with other phosphosites at higher rates than others. For example, pS396 occurs with each other phosphosite 66-71% of the time (**Figure 4H**). This is partially due to the extensive pS396 phosphorylation. We also assessed whether some phosphosites were more likely to be observed with another phosphosite rather than without it. Some phosphosite pairs appeared to have stronger interactions than others **(Figure 4H, Suppl. Fig. S4E**). For example, the presence of pT181 increases the probability of observing pT217 8-fold (**Suppl. Fig. S4E**). Other phosphosites also notably increase the probability of observing pT217 including pS202+pT205 (5.2-fold), pT205 (4.3-fold), and pS214 (3.3-fold). Generally, pT217 appears to be the site most tightly coupled to the phosphorylation status of other sites on the same molecule. Conversely, pS396 appears to be the least tightly coupled to the status of the other sites on the same molecule, perhaps because it is physically further away, and outside of the proline-rich region of tau. This is despite pS396 being the highest abundance phosphosite observed in the data. Phosphosite dependency trends appear largely conserved across model systems with some evidence of higher codependency in brain extracts (data not shown). The dependence of pT217 on modification of other sites, without those sites having as substantial a dependence on pT217, suggests that pT217 may be a later modification, or suggest the activity of a phosphatase.

Collectively, the results highlight a complex and evolving landscape of tau phosphoforms, where the co-occurrence of distinct phosphorylation events suggests the existence of preferential and coordinated modification pathways and the conservation of co-occurrence across model systems implies conservation of said pathways. Notably, some of these co-occurring phosphorylations span the entire length of the protein—from the N- to the C-terminus—underscoring the necessity of a methodology capable of resolving combinatorial PTM patterns across the full-length, intact tau molecule.

Tau proteoform assay compatibility with normal and cognitively impaired brain samples

To validate that the IMaP method was compatible with human clinical samples we analyzed a small cohort of 7 human samples that included aged asymptomatic patients and patients with Alzheimer's Disease and Related Dementias (ADRD) (**Figures 5A, Supp. Fig. 5, Supp. Table S7**), Previous studies characterizing tau PTMs in individuals with AD have revealed considerable heterogeneity [29]. We sought to determine whether our tau proteoform assay could measure human clinical tissue samples and if tau proteoforms could potentially distinguish between cognitively impaired cases and cognitively normal controls. We measured tau proteoforms from brain samples of a small cohort of five cognitively impaired patients with ADRD and two cognitively normal subjects. Clustering of impaired and normal samples revealed three distinct groups, with the normal samples clustering together. Nine proteoforms differentiated the normal samples from the ADRD samples. These nine proteoforms were significantly lower in abundance in ADRD samples (**Figure 5B**). These proteoforms contained phosphorylation at pT181, pT231, or both sites and were found on 0N3R, 0N4R, 1N3R, and 1N4R isoforms. While this is a relatively small sample size, these results suggest that tau proteoforms from brain tissue may differ between ADRD and normal.

In addition, one patient with ADRD exhibited higher phosphorylation than the remaining patients. This patient also exhibited the most severe pathology, with an ABC score of 44 (A3, B3, C3), reflecting the highest levels of Aβ/amyloid plaques, neurofibrillary tangles, and neuritic plaques (each scored 0–3, with 3 indicating severe pathology)[46]. This patient harbored tau proteoforms with quadruply phosphorylated 1N3R tau at pT181-pT217-pT231-pS396. The doubly and triply phosphorylated forms of 1N3R that are a subset of the quadruply phosphorylated form suggest a regulated, rather than random, process driving these phosphorylation events (**Figure 5C**). Three of four triply phosphorylated 1N3R proteoforms showed higher than expected abundance, whereas four of six doubly phosphorylated 1N3R proteoforms showed lower than expected abundance if phosphorylation events were randomly distributed (**Supp. Fig. 6**). These data suggest a

preference for specific hyperphosphorylated tau proteoforms in the patient with the most severe ADRD.

DISCUSSION

We have introduced a new method, Iterative Mapping of Proteoforms, for the large-scale analysis of intact single-molecule proteoforms and applied it to develop an assay for quantification of defined sets of proteoform groups of tau. As shown in **Figure 3** we observed a strong accuracy (R^2 = 0.985) across defined mixtures and median CVs below 5.5% across a wide range of reproducibility challenges. Currently, the greatest source of variation is in library preparation across operators. It is possible that further improvements in reproducibility could come from improvements in the robustness of the library preparation method and in improvements in training to further reduce operator-to-operator variation. Despite being a first demonstration of a new method, this CV is notably lower than those reported for existing, mature proteomics platforms [47,48]. The assay resolved distinct proteoforms across greater than three orders of magnitude in dynamic range, providing a precise readout of relative stoichiometry of tau phosphorylation, and outperformed conventional methods like western blotting in distinguishing closely related proteoforms. In addition, for the first time, we reveal the molecular heterogeneity of tau proteoforms in model systems of neurodegenerative diseases iteratively mapping millions of single molecules to quantify the prevalence of distinct proteoforms.

Application of the tau proteoform assay allowed us to begin investigating several longstanding questions in the field of tau biology and proteoform analysis. First, we observe that tau molecules frequently harbor combinations of more than two phosphorylated sites, with certain proteoforms displaying up to six concurrent modifications. We additionally demonstrated that disease models do not just contain a greater percentage of phosphorylated tau molecules (82% for organoids vs 36% for healthy mouse brain and 13% for healthy human brain), but also that the number of phosphorylations-per-protein-molecule is larger (median 3/molecule in organoid vs 1.5/molecule for healthy human brain) reflecting the developmental or adult biology of the samples or their *in vivo* and *in vitro* origin, respectively.

Importantly, phosphorylations are not randomly distributed. We observed a non-uniform distribution of multi-phosphorylated tau species, with certain combinations of phosphorylation occurring disproportionately, suggesting an ordered and regulated process rather than stochastic modification. This is exemplified by the preferential co-occurrence of pT217 with pT181 with other co-occurrence of pT217 less frequently or

not observed, implying temporal or hierarchical control in phospho-tau biogenesis. It is not possible to know whether this arises because of the action of kinases or phosphatases.

We further established that the extent of proteoform detection is limited not by assay sensitivity, but by biological diversity. Of the 768 possible tau proteoform groups detectable by our antibody panel, only a subset—ranging from 23 to 63, depending on the model—were actually observed. This suggests that tau modification is constrained by biological regulation, and not all theoretically possible proteoforms are realized in vivo. Previous work on histone revealed a similar extent of modification quantifying between 70-200 proteoforms for histone H4 and H3.2, respectively[49]. Nonetheless, our approach dramatically expands the number of resolvable proteoforms compared to traditional methods.

Finally, our method and assay fill key gaps in the field. It enables (1) direct observation of site-specific co-occurring PTMs on individual molecules, (2) quantification of isoform-specific changes over time in human-relevant systems like organoids and miBrains, and (3) comparative assessment of model systems against human disease samples. Moreover, in at least one ADRD patient, we infer a plausible sequence of phosphorylation events leading to a distinct, highly-modified tau species, reinforcing the notion of ordered PTM accumulation in disease progression.

While Iterative Mapping of Proteoforms provides high-resolution quantification of proteoform groups, as demonstrated here with tau, several limitations should be considered. First, this study focused exclusively on full-length, RIPA-soluble tau, in contrast to most prior work that emphasizes sarkosyl-soluble and -insoluble fractions. This may particularly impact the interpretation of samples with a highly variable extent of truncation as our estimates of relative fold-change could be impacted by biological changes in total % of intact proteoforms. This can be addressed either through relating our estimates of proteoform relative abundance to all the measurements on the flow cell, all the measurements of tau of any length, or through using additional cycles for reagents that map truncations of tau explicitly.

Second, the targeted nature of this application of the method inherently limits its scope to known epitopes. Because Iterative Mapping of Proteoforms relies on predefined antibodies to detect specific isoforms and PTMs, it cannot identify novel sites or unanticipated modifications. This contrasts with discovery-based techniques like TDMS, which can uncover unexpected PTMs but lack single-molecule resolution and accurate stoichiometric co-occurrence information.

Third, this application of the method depends on antibodies with high specificity and compatibility with single-molecule detection. Cross-reactivity and inadequate epitope resolution—especially for adjacent or closely spaced modifications—can compromise accuracy. To mitigate this, we applied a dual-screening process for antibody validation and incorporated both null-lane controls and correction algorithms during data analysis, which effectively reduced non-specific signals to below 1%. Nonetheless, discriminating between closely spaced PTMs (e.g. neighboring phosphorylation sites) remains a fundamental challenge for antibody-based approaches. One possibility may also be to use highly non-specific affinity reagents, such as pan pTyr reagents in the assay to get a general gestalt of phosphorylation as a complement to the site-specific reagents used herein.

Finally, the current library preparation protocol requires lysine-based bioconjugation, which introduces random attachment points and molecular orientations on the array. While our results suggest that random orientation under denaturing conditions does not impede detection, future efforts to implement site-specific (e.g., N-terminal) labeling could further standardize molecule presentation and reduce potential sources of variability. Collectively, these limitations define the current boundaries of the tau proteoform assay and present opportunities for further refinement.

Iterative Mapping of Proteoforms establishes a new benchmark for high-resolution, single-molecule proteoform analysis, and several future directions hold promise for expanding its impact across neurodegeneration research and beyond. While protein sequencing technologies and TDMS offer broad discovery capabilities for detecting proteoform modifications, IMaP provides a scalable, targeted approach with unprecedented resolution of individual proteoforms beyond the reach of TDMS. The method is not limited to 12 antibodies and can be further expanded with additional probing cycles to include a wider array of PTMs. In stability studies (data not shown) we have demonstrated binding and removal of antibodies for over 150 cycles without loss of specific antibody binding or significant loss or damage of deposited proteins. Tau-specific modifications including acetylation, ubiquitination, and proteolytic truncations implicated in tau pathology are of high interest. Incorporating these dimensions will improve the granularity of tau proteoform characterization, particularly in disease progression and therapeutic response contexts.

With an observed median CV less than 5.5%, the reproducibility of the current assay now supports scaling to larger, biological studies and clinically annotated cohorts. Applying the tau proteoform assay to a broader range of brain samples—from

cognitively healthy individuals to those across the full spectrum of AD and other tauopathies—may provide deep insight into the spatial and regional distribution of tau proteoforms. Leveraging pathologically characterized brain bank tissues could further link specific proteoform signatures to tau burden across multiple brain regions, generating maps of proteoform pathology that could inform disease staging and therapeutic targeting.

A critical next frontier is the adaptation of the assay for detection of tau proteoforms in biofluids such as cerebrospinal fluid (CSF) and blood, where concentrations are expected to be orders of magnitude lower than in tissue. Achieving single-molecule resolution in these matrices would enable the discovery of clinically relevant biomarkers and facilitate longitudinal studies to monitor the emergence and clearance of specific tau species over time. Such capabilities could support early diagnosis, stratification of disease subtypes, and assessment of drug efficacy in both preclinical and clinical settings.

Finally, the versatility of the method allows it to be readily adapted for other key proteins implicated in neurodegenerative disease—including α-synuclein, amyloid-β, and TDP-43—as well as proteoforms central to other human diseases such as cancer and autoimmune disorders. By applying the same principles, single-molecule Iterative Mapping of Proteoforms could help unravel the structural and functional heterogeneity of diverse protein populations, offering a transformative tool for proteoform-centric biology and precision medicine.

Figures & Figure Legends.

Figure 1: Single-molecule Iterative Mapping of Proteoforms workflow

Figure 1A: Proteins are functionalized on lysine residues with methyltetrazine-PEG4-STP (4-Sulfo-2,3,5,6-tetrafluorophenyl) ester, conjugated by a cycloaddition reaction to a DNA origami nanoparticle containing a single trans-cyclooctene (TCO) moiety.

Figure 1B: Tau protein nanoparticle-conjugates are deposited as single proteins on a glass surface chip. Between 4 and 12 sample lanes are configured in a flow cell for an experiment.

Figure 1C: A series of fluorescently labeled antibodies are used to iteratively probe each molecule via a series of incubation, rinse, imaging, and washing steps.

Figure 1D: Proteoforms are determined from the pattern of antibody binding and quantified by counting. A normalizing computation (not shown) is used to account for false-positive and false-negative binding of antibodies.

Figure 2: Application of iterative mapping to measure proteoforms of tau

Figure 2A: Tau isoforms generated through tau gene expression, alternative RNA splicing, and protein translation. N1 and N2 tau isoforms (orange, dark navy blue) are generated by alternate splicing of exon 2 and 3 and alternate splicing of exon 10 produces the 3R and 4R isoforms (light blue). Approximate epitope positions for isoform and proteoform specific detection antibodies to probe the molecular heterogeneity used for Iterative Mapping of proteoforms and approximate epitope binding sites for antibodies used for enrichment.

Figure 2B: Primary imaging data capturing a representative area of 130 μm x 130 μm of flow cell surface showing 16,000 landing pads. Each illuminated pixel is a tau single nanoparticle-conjugate deposited in the hyper-dense array. Over a series of

fluorescently labeled antibody nanoparticle-conjugate binding and imaging cycles a series of single molecule binding traces are recorded for each tau proteoform. Green halo indicates positive binding event.

Figure 2C: Tau proteoforms standards (0N3R, 2N4R-PKA, 0N4R-ERK) and expected epitope combination (A-J). The right panel shows the relative abundance of individual proteoforms in parts per million. The left panel shows the epitope profile of mixtures of tau proteoforms resolved by interactive mapping of proteoforms. The mixtures of protein standards are: 33.3% 0N-ERK2 : 66.6% 0N3R and 33.3% 0N-ERK2 : 66.6% 2N4R-PKA. NOTE: 2N4R-PKA represents tau proteoform 2N4R treated with Protein Kinase A, which is phosphorylated to >99.9% at Serine 214 (pS214), 0N-ERK2 denotes 0N4R treated with ERK2, giving rise to multiple phosphorylations. Error bars represent standard deviation and each sample has three technical replicates.

Figure 3: Accuracy, reproducibility, and dynamic range of the tau proteoform assay

Figure 3A. Quantification accuracy and relative abundance were assessed using two mixtures of tau proteoform standards, each containing 0N4R, 2N4R, and 2N4R-PKA at defined molar ratios of 1:2:3 and 1:1:1, respectively. The results shown represent 21 technical replicates comparing the measured ratios between the two mixtures, with error bars indicating the standard deviation. The mean absolute percent error across all measurements was 9.97%.

Figure 3B. Here we examined the reproducibility of quantitation. Overall, the median CV amongst technical replicates of prepared nanoparticle protein conjugates within the same experimental run was 1.5%. In addition to examining reproducibility within runs of the control mixture within the same experiment, we examined the reproducibility across batches of five flow cell batches (CV=2.6%) and across three instruments (CV=3.6%). We additionally examined reproducibility of full end-to-end experiments including library preparation performed by the same operator 3X (CV=5.5%) and as performed by three different operators (CV=5.5%). In total, estimates of reproducibility are drawn from 297 measurements.

Figure 3C: Dynamic range was assessed using the 2N4R-PKA proteoform standard representing the proteoform 2N4R-pS214 with serial dilution into an orthogonal background of the 0N3R proteoform employing five (anti-Tau13, anti-2N, anti-pS214,

anti-4R, anti-Tau216) detection antibodies (dark blue circle) and four (anti-Tau13, anti-2N, anti-pS214, anti-Tau-216) detection antibodies (light blue). The limit of quantitation (LOQ) was defined as the lowest point on the titration curve at which the signal to noise ratio is greater than 10 and the coefficient of variation is under 30%. Error bars represent the mean of nine technical replicates across three experiments.

Figure 4: Examination of the landscape of tau proteoforms in models of neurodegenerative disease

Figure 4A: Model systems (columns) and tau proteoforms (rows) are hierarchically clustered based on proteoform abundance. The blue-to-orange scale indicates relative abundance by z-score normalized per-proteoform of the mean of all replicates across each model system. The accompanying black/beige heatmap to the right denotes the presence (black) or absence (beige) of specific epitopes within each proteoform.

Figure 4B: Number of proteoforms detected above 1000 ppm relative abundance for each of the tissues investigated. Data represent the RIPA buffer-soluble fraction of neuronal tissue as indicated. Number of phosphorylations are in Suppl. Table S5. For iNeurons n=3 technical replicates, for WT/WT-3Mon and 6Mon n = 12 (2 donors x 2 conditions x 3 technical replicates), for WT/V337M 3Mon and 6Mon n = 6 (1 donor x 2 conditions x 3 technical replicates), for WT/IVS10+16 3Mon and 6Mon n = 6 (1 donor x 2 conditions x 3 technical replicates), for MiBrains *APOE3* and *APOE4* n=8 (2 x 4 technical replicates), and for Human Brain Extracts and Mouse Brain Extracts n=9 technical replicates each. Error bars are standard deviation.

Figure 4C: Number of tau protein molecules phosphorylated in human brain, mouse brain, miBrains, cortical organoids, and iNeurons. Data represent the average of at least two biological with 3 technical replicates with exception of the human brain extract (n=1). For iNeurons n=3 technical replicates, for Organoids n=24, for MiBrains n=8, and for Human Brain Extracts and Mouse Brain Extracts n=9. Error bars are standard deviation.

Figure 4D: Bar graph showing the percent occupancy for each tau epitope targeted in the assay. The same number of replicates are used as in 4B. Error bars are standard deviation.

Figure 4E: Relative abundance (in ppm) of tau isoforms in maturing organoids at 3 and 6 months comparing IVS10+16 mutant (WT/IVS10+16) against isogenic controls (WT/WT). Representative result (donor GIH178C1) of two iPSC donor cell lines analyzed with an average of 8 organoids processed in two independent analyses. Cortical organoids details in Supplemental Table S4. Error bars are standard deviation.

Figure 4F: Bar graph shows the relative abundance in ppm of the 3R and 4R tau protein in maturing miBrains of the APOE4 genotype. Tau epitopes on the x-axis reflect the isoform specific probes used to assess full length tau protein isoforms (3R Tau putative). Noise level for the 3R tau was determined at 1474 ppm and at 485 ppm for the 2- and 4-month-old miBrains, respectively. For the 4R tau, corresponding noise levels for 2- and 4-month-old miBrains were at 1762 ppm and 702 ppm, respectively. Data shows four technical replicates of lysate from four miBrains at (57 days) and the average of 6 miBrains (112 days) processed independently.

Figure 4G: Abundances are calculated independent of tau isoform and defined as phosphoforms. Orange dots represent observed phosphoform abundance, while dark blue and light blue dots indicate expected abundance based on models of independent

and random phosphorylation, respectively. Statistical significance between observed and expected abundance levels from the independent phosphorylation model is shown (*p<0.05, **p<0.01, ***p<0.001). Error bars are standard deviation. N=12.

Figure 4H: Co-occurrence of tau phosphorylation. Here we quantify across all model systems how often a given site is phosphorylated (y-axis) assuming the presence of a phosphorylation on another site (x-axis). The pattern of co-occurrence is distinctively non-uniform suggesting a preference for multiple phosphorylation.

Figure 5: Application of tau proteoform assay to human samples

Figure 5A: Tau proteoforms contrast normal from cognitively impaired brain samples. A heat map of the normalized relative abundance of the proteoforms of tau (mean across technical replicates) from samples of a small cohort with two cognitively normal patients (patient 127 & 204) and five cognitively impaired ADRD patients (18, 213, 122, 12, 203). Each patient is represented by three technical replicate measurements. Proteoforms

plotted were at greater than 10% in abundance. Presence and severity of AD-related lesions for each patient is indicate in the lower panel following the ABC score. ABC = Thal, Braak, CERAD score for amyloid plaque deposition, tau protein tangles, and cortical neuritic plaques, respectively.

Figure 5B: Relative abundance of multiple phosphorylations in human brain with two cognitively normal patients (patient 127 & 204) and five cognitively impaired ADRD patients. Tau proteoforms with multiple phosphorylations comparing cognitively normal to ADRD impaired. Star indicates p-values adjusted for multiple hypotheses using a Bonferroni correction.

Figure 5C: Hyperphosphorylation of 1N3R in patient 18 at pT181, pT217, pT231, and pS396 potentially follows a preferential path from double to quadruple phosphorylation. Node fill color is the abundance of the proteoform as estimated by the mean of technical replicates. Edge weights correspond to estimated relative contribution of a given node to its parent nodes.

MATERIALS & METHODS

Screening and validation process for suitable antibodies

To assess the binding characteristics and specificity of the probe affinity reagents used for Iterative Mapping of Proteoforms we characterized the affinity reagents using a two-step process. We first assess the specificity of the antibodies to individual epitopes using control peptides (**Supplemental Table S1**). In a second step we confirm the specificity of the reagents (**Supplemental Table S3)** on the single molecule proteoform detection system to verify that those detection antibodies accurately recognized the specific epitopes of tau including phosphorylation and multiple phosphorylation sites.

As an example, to demonstrate that we accurately detect a mixture of three tau epitopes, we create phosphorylated peptides that represent tau epitopes of pT181, pS215, pS396 and immobilize them in a flow cell at a ratio of 1:3:1 to detect them with the corresponding antibodies. As negative control, corresponding non-phospho peptides or full length 2N4R proteins were used. Results from these specificity assessments using the negative control demonstrated that single molecule on-platform epitope recognition is qualitatively accurate and shows minimal off-target detection (cross-reactivity) with false positive detection below 1%.

Enrichment antibodies were selected for high affinity towards pan Tau proteins while still suitable for elution with cognate epitope peptides. They were able to deplete >90% Tau after overnight incubation with a protein mixture containing about 1 ng/ul Tau in ~100 ul volume. The three antibodies used in this study recognize N-, middle, and C-terminus of Tau.

Preparation of detection antibody conjugates

Modification of Anti-Tau Antibodies

Antibodies were acquired as carrier-free formulations in PBS pH 7.2-7.4 (**Supplementary Table S3**), or else buffer exchanged into PBS pH 7.4 using Amicon Ultra 100k MWCO Regenerated Cellulose Spin Filters. Modification is performed by first adjusting the pH of the antibody solution using 0.5M Sodium Phosphate (pH 8.0, 0.25 ml per of Antibody). A freshly prepared solution of sulfo-NHSLC-LC-Biotin is added to the Antibody solution in 20-fold molar excess and incubated (25C, shaking, protected from light) for 2 hours. Quenching of the modification reaction is performed for 0.5 hours (25C, shaking, dark) with a final concentration of L-Arginine (in the quench reaction) of

10mM. Modified antibodies are purified using desalting column (Zeba, Thermo Scientific). The extent of biotinylation is measured using MALDI Mass spectrometry

Conjugation of Anti-Tau Antibodies to fluorescent labels
Fluorescent labels containing up to four streptavidin groups are incubated with 2 molar equivalents of biotinylated anti-Tau antibody with equivalents calculated relative to the measured number of Streptavidin attached per nanoparticle. The conjugation proceeds for 2 hours at 25C, shaking in the dark. Ab-conjugated nanoparticles are purified from excess unconjugated antibody via HPLC- Size Exclusion Chromatography (SEC) using an Agilent 1260 Infinity II HPLC system equipped with DAD and fraction collector. The SEC column used is a Shodex OHpak SB-2006M (P/N: F6516017). Method for purification consists of isocratic flow at 2.5mL/min of 1x HPLC O.B. (200mM NaCl, 11mM $MgCl_2$, 5mM Tris-HCl, 1mM EDTA) for 1.5 column volumes. Fractions corresponding to the nanoparticle peak are pooled and concentrated using Amicon Ultra 100k MWCO Regenerated Cellulose Spin Filters. Purified, concentrated sample is stored in DNA Low-Bind tubes at 4C.

Tau proteoform standards and mixtures
To assess the accuracy and performance of our platform to correctly identify and quantify specific tau proteoforms, we created four highly characterized tau protein standards and mixtures thereof. Tau proteoform standards, included a) unphosphorylated tau 0N4R isoform, b) the unphosphorylated 2N4R isoform, c) phosphorylated tau 0N4R isoform treated with ERK2 (multiple phosphorylation) and d) the tau 2N4R isoform treated with PKA (single phosphorylation at serine 214). We established the exact composition of four distinct tau proteoforms by LCMS assessing the site and percentage of phosphorylation of tau proteoform standards as described in the (Supplemental Tables and Figures, Table S2).

Additionally, we purchased tau standards 0N4R (BioLegend 843001), 2N4R (R&D Systems (SP-495). The tau standards were used throughout the study as proteoform controls and used to prepare tau proteoform mixtures as indicated in the experimental details. As binding controls for the assay, we created DNA-origami nanostructure conjugates of the following tau proteoforms standards 2N4R, 0N4R, 0N3R, triply and quadruply phosphorylated peptides (see **Suppl. Table S1**), a short peptide as a null target, and standard mixtures of 0N4R:2N4R:2N4R-PKA tau proteoforms at different ratios as indicated in the experimental details. Before mTz modification, tau proteoform standards commercially purchased recombinant Tau proteins were first precipitated with four volumes acetone and resuspended in PBS, supplemented with 1% SDS.

Cells and Tissues

Postnatal-like iPSC-derived neurons (iNeurons) were generated using the *NGN2* protocol and cultured as previously as described [50]. Cortical organoids from patient-derived iPSC using a guided differentiation protocol with dual SMAD and WNT inhibition in a 96-slitwell format, as previously described [51]. 10 organoids at 3 months and 6 months maturity, harboring the original disease genes or reverted to the wild type by CRISPR techniques (**Supplemental Table S4**), were harvested as 3 vials (3-4 organoids each) for protein extraction and library preparation in parallel. [38,5138,5238,5138,5238,5338,52]. 4-6 miBrains harboring apolipoprotein E3 or E4 genes (*APOE3/E4*), were harvested individually at days 57 and 112 after seeding and processed for protein extraction and proteoform analysis. hTau mouse brains at 3 months age used in Figures 4 and 5 were obtained from Quest Pharmaceutical Services (Newark, DE), from transgenic mice expressing a human tau derived from a human PAC, H1 haplotype, and with the murine Tau gene knocked out by a targeted disruption of exon1. The healthy human frontal cortex brain sample denoted Human brain extract as used in Figures 4 and 5 was obtained from Analytical Biological Services Inc. (New Castle, DE) from a postmortem Caucasian male donor died at the age of 66 with lung cancer without neurological disorder. A cohort of seven post-mortem human brain samples from the cortical region was obtained from the Mount Sinai Neuropathology Brain Bank and Research CoRE. These samples are used Figure 5. Donor information and disease stages are listed in the table **Supplementary Table S6**.

A detailed description of all the biological samples and their lysis/protein extraction procedures can be found in **Supplementary Information**.

Single-molecule Iterative Mapping of Proteoforms Methodology

The method we introduce here, Iterative Mapping of Proteoforms (IMaP) consists of four steps that are detailed below and diagrammed in **Figure 1**. The first step is library preparation (detailed in **Single molecule library preparation**) which aims to take protein molecules from a sample and conjugate each molecule to a DNA-origami nanostructure.

In one instantiation of this workflow recombinant protein is used directly functionalized with an amine-reactive mTz click reagent (as described in *Functionalization of recombinant Tau proteins).* In another instantiation of this workflow protein is extracted from tissue lysate prior to functionalization (as described in *Functionalization of cell lysates).* If needed, a sample can be enriched using protein-specific antibodies (detailed in Optional *Tau Enrichment).* Next, protein is conjugated to DNA-origami-based nanoparticles. When used alongside enrichment, functionalization can either occur using

protein directly (detailed in *DNA nanostructure conjugation for un-enriched samples*) or on bead (detailed in *DNA nanostructure conjugation for enriched samples*). At no point is the protein digested. Next (**Figure 1B**), protein-nanoparticle conjugates are deposited into a lane of a custom microfluidics flow cell [34]. Flow cell fabrication is described in Supplemental Methods in the section (*Flow cell preparation and assembly*). Single molecules are iteratively probed (as described in the section *Sample analysis*) in a massively parallel manner using a series of site-specific fluorescently labeled [35] antibodies (**Figure 1C**). Antibody labeling is described above in the section (**Preparation of detection antibody conjugates**).

Each iteration consists of a bind, rinse, image, and remove cycle. Specifically, a fluorescently labeled antibody probe is introduced into the flow cell and allowed to incubate. Next, unbound antibody is rinsed from the flow cell and the flow cell is optically imaged [34] to determine where the antibody has bound. Following imaging, bound antibodies are removed from the flow cell, preparing the flow cell for a subsequent iteration. As fluorescent images are acquired, positions of each single-molecule binding event are recorded. Collectively, probing of the single-molecule array produces a series of binary bind/no-bind outcome patterns (single molecule binding traces), which are used to determine the potential identity of the proteoform at each coordinate in the flow cell. Analysis of the images and sample quantification are described in the sections *Processing of Images* and *Proteoform Analysis* respectively.

Single molecule library preparation
Functionalization of recombinant Tau proteins
Protein samples are resuspended in PBS, supplemented with 1% SDS and then treated with 100 uM methyltetrazine-PEG4-STP (4-Sulfo-2,3,5,6-tetrafluorophenyl) ester (mTz) (Vector labs, CCT-1399) (with 1% DMSO) at 25C for 3 hours on a thermomixer shaking at 850 rpm protected from light. The reaction was quenched with Dithiobutylamine (Sigma Aldrich 774405) at 5 mM for 1 hr at 25C, followed by iodoacetamide (Pierce A39271) at 10 mM for 15 min at 25C. The protein was precipitated by 9 volumes of absolute ethanol, washed once with ethanol at room temperature, air dried, then solubilized in (200mM NaCl, 15mM $MgCl_2$, 5mM Tris-HCl, 1mM EDTA, 1% SDS), and quantified using the Pierce 660nm Protein Assay Reagent following manufacturer's manual (Thermo Scientific).

DNA nanostructure conjugation for un-enriched samples
For recombinant proteins, or eluted proteins mTz modified proteins were incubated with DNA nanoparticles (at 50~100 nM) bearing a single TCO (*trans*-cyclooctene) functional group[53] at an approximate molar ratio of 5:1 in a buffer (200mM NaCl, 15mM $MgCl_2$,

5mM Tris-HCl, 1mM EDTA) overnight (approximately 16 hours) at 25C protected from light. Spin filtration with a 100kDa MWCO cut-off Amicon Ultracel (Millipore Sigma UFC910008) was used to remove excess unconjugated proteins and exchange to a new buffer (5mM Tris-HCl pH 7.5, 200mM NaCl, 15mM $MgCl_2$, 1mM EDTA, 0.05% CHAPS). A typical reaction of about 20 ul was first diluted to 500 ul, concentrated to about 20 ul, and repeated three times in total. The purified Tau-conjugated nanoparticles were quantified with a Nanodrop instrument using the dsDNA settings.

Functionalization of cell lysates.
Neuronal tissue/cell lysates and Expi293 lysates with recombinant Tau spiked-in were first precipitated with 4 volumes of pre-chilled (-20C) acetone. The pellets were briefly washed with cold acetone, air dried, then resuspended in PBS supplemented with 1% SDS. Lysates were quantified with Pierce 660 Assay, adjusted to 2 mg/ml, then treated with 200 uM methyltetrazine-PEG4-STP (4-Sulfo-2,3,5,6-tetrafluorophenyl) ester (mTz) with 2% DMSO at 25C for 3 hours on a thermomixer shaking at 850 rpm protected from light. The reaction was quenched with Dithiobutylamine (Sigma Aldrich 774405) at 5 mM for 1 hr at 25C, followed by iodoacetamine (Pierce A39271) at 10 mM for 15 min at 25C. The protein was precipitated by 9 volumes of absolute ethanol, washed once with ethanol at room temperature, air dried, then solubilized in PBS supplemented with 1% SDS.

Optional Tau Enrichment
When used with biologic samples, enrichment of tau is performed using a magnetic bead-based pulldown. Anti-Tau monoclonal antibodies targeting N-, middle, and C-terminus of Tau, optionally together with mouse IgG (Invitrogen 31903) as the negative control, are immobilized individually using Dynabeads™ Antibody Coupling Kit (Invitrogen 14311D) at 10 μg antibody per mg beads according to the manufacturer's instructions. Anti-Tau beads were mixed in equimolar ratios prior to use. Modified lysates were diluted by 5 volumes of IP Buffer (50mM Tris-HCl pH 8.0, 150mM NaCl, 1% Triton X-100) supplemented with 1x Protease/Phosphatase Inhibitor cocktail (Cell Signaling Technology #5872), to bring final SDS to 0.16%, then mixed with antibody containing beads at an IgG:Tau ratio of around 15~50:1 at room temperature overnight (approx. 16hrs), on a tube rotisserie. The beads are then extensively washed with PBS and IP Buffer 2~3 times each,

DNA nanostructure conjugation for enriched samples
While still on the beads, enriched tau is conjugated with 2.5 molar excess of DNA nanoparticles (relative to the estimated Tau input) bearing a single TCO (*trans-*cyclooctene) functional group in a buffer (5mM Tris-HCl pH7.5, 200mM NaCl, 15mM

MgCl$_2$, 1mM EDTA) overnight at room temperature on a tube rotisserie. Beads were washed three times with buffer (5mM Tris-HCl pH 7.5, 200mM NaCl, 15mM MgCl$_2$, 1mM EDTA, 0.05% CHAPs), and eluted with the same buffer supplemented with synthetic peptides encompassing the epitopes of each of the Tau antibodies used at room temperature for 90 minutes on a tube rotisserie. Tau-conjugated nanoparticles were quantitated based on 488 florescence if the nanoparticles were labeled, or by Quant-iT PicoGreen dsDNA Reagent (Fisher Scientific P7581) following manufacture suggested procedure, using the unconjugated nanoparticles as standards, aliquoted and stored at -80C before use.

Sample analysis

Experimental Set-up and Imaging
Flow cells, running buffer (50mM HEPES pH7.4 + 120mM NaCl+5mM KCl + 10mM MgCl2+ 0.1% Tween-20 + 0.5% Procilin 150+ 10mM Sodium Sulfite +10mM Sodium L-Ascorbate), remove buffer (5.2 M Guanidine HCl, 20 mM MgCl2, 20 mM Sodium Acetate, pH 4.30 +/- 0.10), DNA nanoparticle conjugated Tau molecules suspended in a buffer comprising 5 mM Tris, pH 8.0, 120 mM NaCl, 12.5 mM MgCl2, 1 mM EDTA, 0.05% CHAPS and affinity reagents diluted in 10 mM HEPES, pH 7.4 120 mM NaCl, 12.5 mM MgCl2, 5 mM KCl, 0.1% Tween-20, 1% Pluronic F-127, 1 mg/mL ssDNA, 1% BSA, 0.1% ProClin-150 were all loaded onto the instrument prior to assay initiation. For the duration of the assay, the running and remove buffers were held at room temperature (~21° C) while labeled probes were maintained at 15° C and the flow cell surface was maintained at 23° C. The assay was performed on the instrument in an automated fashion involving sequential steps. In brief, the flow cell surface was primed with running buffer, followed by injection of the DNA nanoparticle-conjugated Tau proteins. Deposition onto the flow cell surface was mediated by the annealing of complementary oligonucleotides between the nanoparticle and pre-immobilized DBCO-oligonucleotides on the flow cell surface. After a 30-minute incubation, the unbound material was washed away from the flow cell using running buffer. Subsequently, sequential probing was carried out using fluorescently labeled antibodies targeting distinct epitopes. Each probe was incubated for 17 minutes, followed by a wash step with running buffer to remove unbound probes. A 17-minute incubation with remove buffer was then used to dissociate the bound probes, resetting the surface for the next cycle. This process was repeated for 36 total cycles, covering 12 unique probes.

Images of the flow cell surface were acquired at key checkpoints of each cycle using a custom-built high-resolution imaging system. This system included an Olympus UPLXAPO20X objective lens, Thorlabs TTL180-A 180mm focal length tube lens and

dual Teledyne Dalsa Linea HS4K TDI line scan cameras. Excitation was provided by 488nm and 638nm lasers. With an assortment of dichroic and bandpass filters, multicolor imaging was enabled. Autofocus was maintained using a "Dover DOF-5" objective focusing stage integrated with a WDI ATF4 autofocus module. A high precision Dover SmartStage linear stage was used for positioning the flow cells under the objective lens and enabling continuous scanning of the flow cell surface throughout multiple cycles. A custom designed flow cell holder was mounted on the Dover SmartStage to maintain flow cell position and temperature stability of the flow cells under the microscope during imaging. Images of the flow cell surface were acquired after protein conjugated DNA nanoparticle deposition, following each probe wash step and after every remove buffer incubation step. These image stacks were then processed to perform alignment, spot detection and downstream data extraction for quantitation and analysis.

Data analysis

Processing of Images
The input images were processed as previously described [34]. Briefly, patterned array and single pad boundaries were determined using subarray fluorescent patterns excited by the 488 nm laser. The mapping between 488 nm and 647 nm laser acquired images uses subarray locations in the 488 nm scan and fiducial beads in the 647nm scans. A calibration flow cell is used to allow mapping of locations directly from 488 nm images to 647 nm images. A reference base scan is taken for both the 488 nm and 647 nm images. The 647 nm-reference scan is used to align all subsequent 647 nm images. Each image was normalized against the background. Background intensity estimates were calculated by tiling the image and computing the 5[th] percentile for each tile. Then, the tiles are used to solve for a 2-dimensional paraboloid. The image is then normalized against this smooth paraboloid. The primary feature score was computed for each candidate object by summing the logarithm of 5 pixels forming a 3x3 t-shape centered on each landing pad in the background normalized image.

Proteoform Analysis
The Iterative Mapping of Proteoforms algorithm analyzes a series of antibody binding measurements acquired on a sample with unknown proteoforms and determines the abundance of each epitope in the sample. The algorithm uses an expectation-maximization approach to estimate the probability of each proteoform in the sample. The inputs to this algorithm are the series of antibody binding measurements and the rates at which the antibodies bind to their target epitopes. The model employs a Poisson binomial distribution, where the likelihood that the proteoform contains a given epitope

is determined by the antibody binding rates. The output of the model is partial counts corresponding to the probability that each candidate proteoform produced the observed series of binding measurements. The number of candidate proteoforms is 2^m, where m represents the number of antibodies used in the experiment. To account for sample-to-sample and run-to-run variation, the algorithm also estimates the binding rates for each antibody within each sample. We use control proteoforms to determine the affinity of labeled antibody binding in the presence and absence of the target epitope, and set a prior on antibody binding rates based on these control proteoforms. A uniform prior is set on proteoform abundance estimates. To account for potential protein degradation during sample preparation, we discard any landing pads that do not contain at least one positive binding measurement for an antibody that targets the N-terminal region of Tau and at least one positive binding measurement for an antibody that targets the C-terminal region of Tau. This ensures that we analyze landing pads that contain full length Tau. A two sided t-test was used to assess the significance of differences in abundance between proteoforms across samples. In the case of the cognitively impaired and non-cognitively impaired cohorts, p-values were adjusted to correct for multiple hypotheses using a Bonferroni correction.

Expected Multiply Phosphorylated Tau Molecules
Two models for expected phosphorylation abundance were constructed. A "Random Phosphorylation" model was built by assuming that all epitopes were equally likely to be phosphorylated. An "Independent Phosphorylation" model was built by assuming that phosphorylation events occur independent from phosphorylation elsewhere on the molecule. This process was modeled as a joint Bernoulli model, such that the probability of observing each combination of post translational modifications is equal to the product of the probabilities of the presence and absence of each individual phosphorylation event. In the case of Supplementary Figure S4D, isoform information was ignored. In Supp. Fig. S5, isoform information was included such that the expected phosphorylation values were based on the singly phosphorylated occupancies of the 1N3R isoform.

Co-Occurrence of Tau Phosphorylation
To assess whether post translational modifications were co-occurring at a higher than expected rate, we calculated the likelihood ratio comparing two conditional probabilities. We compared the rate at which post translational modification Y occurs given post translational modification X to the rate at which post translational modification Y occurs when post translational modification X is not present. Notably, this analysis ignored isoform information.

REFERENCES

1. Zhang, Y., Fonslow, B. R., Shan, B., Baek, M.-C. & Yates, J. R. Protein Analysis by Shotgun/Bottom-up Proteomics. *Chem. Rev.* **113**, 2343–2394 (2013).

2. Assarsson, E. *et al.* Homogenous 96-Plex PEA Immunoassay Exhibiting High Sensitivity, Specificity, and Excellent Scalability. *PLoS ONE* **9**, e95192 (2014).

3. Gold, L. *et al.* Aptamer-Based Multiplexed Proteomic Technology for Biomarker Discovery. *PLoS ONE* **5**, e15004 (2010).

4. Aebersold, R. *et al.* How many human proteoforms are there? *Nat Chem Biol* **14**, 206–214 (2018).

5. Lee, J. M., Hammarén, H. M., Savitski, M. M. & Baek, S. H. Control of protein stability by post-translational modifications. *Nat Commun* **14**, 201 (2023).

6. Breckels, L. M. *et al.* Advances in spatial proteomics: Mapping proteome architecture from protein complexes to subcellular localizations. *Cell Chemical Biology* **31**, 1665–1687 (2024).

7. Yang, Y.-H., Wen, R., Yang, N., Zhang, T.-N. & Liu, C.-F. Roles of protein post-translational modifications in glucose and lipid metabolism: mechanisms and perspectives. *Mol Med* **29**, 93 (2023).

8. Li, W., Li, H.-L., Wang, J.-Z., Liu, R. & Wang, X. Abnormal protein post-translational modifications induces aggregation and abnormal deposition of protein, mediating neurodegenerative diseases. *Cell Biosci* **14**, 22 (2024).

9. Kokkinidis, M., Glykos, N. M. & Fadouloglou, V. E. Catalytic activity regulation through post-translational modification: the expanding universe of protein diversity. in *Advances in Protein Chemistry and Structural Biology* vol. 122 97–125 (Elsevier, 2020).

10. Zhang, J. *et al.* Top-Down Quantitative Proteomics Identified Phosphorylation of Cardiac Troponin I as a Candidate Biomarker for Chronic Heart Failure. *J. Proteome Res.* **10**, 4054–4065 (2011).

11. Jenuwein, T. & Allis, C. D. Translating the Histone Code. *Science* **293**, 1074–1080 (2001).

12. Smith, L. M. *et al.* The Human Proteoform Project: Defining the human proteome. *Sci. Adv.* **7**, eabk0734 (2021).

13. Gregorich, Z. R. & Ge, Y. Top-down proteomics in health and disease: Challenges and opportunities. *Proteomics* **14**, 1195–1210 (2014).

14. Doucette, A. A., Tran, J. C., Wall, M. J. & Fitzsimmons, S. Intact proteome fractionation strategies compatible with mass spectrometry. *Expert Rev Proteomics* **8**, 787–800 (2011).

15. Capriotti, A. L., Cavaliere, C., Foglia, P., Samperi, R. & Laganà, A. Intact protein separation by chromatographic and/or electrophoretic techniques for top-down proteomics. *Journal of Chromatography A* **1218**, 8760–8776 (2011).

16. Claes, B. S. R., Takeo, E., Fukusaki, E., Shimma, S. & Heeren, R. M. A. Imaging Isomers on a Biological Surface: A Review. *Mass Spectrometry* **8**, A0078–A0078 (2019).

17. Fulcher, J. M. *et al.* Discovery of Proteoforms Associated With Alzheimer's Disease Through Quantitative Top-Down Proteomics. *Molecular & Cellular Proteomics* **24**, 100983 (2025).

18.   Deslignière, E. *et al.* Ultralong transients enhance sensitivity and resolution in Orbitrap-based single-ion mass spectrometry. *Nat Methods* **21**, 619–622 (2024).

19.   Kafader, J. O. *et al.* Multiplexed mass spectrometry of individual ions improves measurement of proteoforms and their complexes. *Nat Methods* **17**, 391–394 (2020).

20.   Prabakaran, S., Lippens, G., Steen, H. & Gunawardena, J. Post-translational modification: nature's escape from genetic imprisonment and the basis for dynamic information encoding. *WIREs Mechanisms of Disease* **4**, 565–583 (2012).

21.   Reed, B. D. *et al.* Real-time dynamic single-molecule protein sequencing on an integrated semiconductor device. *Science* **378**, 186–192 (2022).

22.   Floyd, B. M. & Marcotte, E. M. Protein Sequencing, One Molecule at a Time. *Annu. Rev. Biophys.* **51**, 181–200 (2022).

23.   Filius, M. *et al.* Full-length single-molecule protein fingerprinting. *Nat. Nanotechnol.* **19**, 652–659 (2024).

24.   Motone, K. *et al.* Multi-pass, single-molecule nanopore reading of long protein strands. *Nature* **633**, 662–669 (2024).

25.   Nova, I. C. *et al.* Detection of phosphorylation post-translational modifications along single peptides with nanopores. *Nat Biotechnol* **42**, 710–714 (2024).

26.   Buchholz, S. & Zempel, H. The six brain-specific TAU isoforms and their role in Alzheimer's disease and related neurodegenerative dementia syndromes. *Alzheimer's & Dementia* **20**, 3606–3628 (2024).

27.   Goedert, M., Spillantini, M. G., Jakes, R., Rutherford, D. & Crowther, R. A. Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron* **3**, 519–526 (1989).

28.   Boyarko, B. & Hook, V. Human Tau Isoforms and Proteolysis for Production of Toxic Tau Fragments in Neurodegeneration. *Front. Neurosci.* **15**, 702788 (2021).

29.   Wesseling, H. *et al.* Tau PTM Profiles Identify Patient Heterogeneity and Stages of Alzheimer's Disease. *Cell* **183**, 1699-1713.e13 (2020).

30.   Chakraborty, P. *et al.* Acetylation discriminates disease-specific tau deposition. *Nat Commun* **14**, 5919 (2023).

31.   Blennow, K., Hampel, H., Weiner, M. & Zetterberg, H. Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol* **6**, 131–144 (2010).

32.   Palmqvist, S. *et al.* Discriminative Accuracy of Plasma Phospho-tau217 for Alzheimer Disease vs Other Neurodegenerative Disorders. *JAMA* **324**, 772 (2020).

33.   Yang, Y. *et al.* Programmable Site-Specific Functionalization of DNA Origami with Polynucleotide Brushes. *Angew Chem Int Ed* **60**, 23241–23247 (2021).

34.   Aksel, T. *et al.* High-density and scalable protein arrays for single-molecule proteomic studies. Preprint at https://doi.org/10.1101/2022.05.02.490328 (2022).

35.   Stawicki, C. M. *et al.* Modular fluorescent nanoparticle DNA probes for detection of peptides and proteins. *Sci Rep* **11**, 19921 (2021).

36.   Khatoon, S., Grundke-Iqbal, I. & Iqbal, K. Brain Levels of Microtubule-Associated Protein τ Are Elevated in Alzheimer's Disease: A Radioimmuno-Slot-Blot Assay for Nanograms of the Protein. *Journal of Neurochemistry* **59**, 750–753 (1992).

37.   Shan, X. *et al.* Fully defined *NGN2* neuron protocol reveals diverse signatures of neuronal maturation. *Cell Reports Methods* **4**, 100858 (2024).

38.   Stanton, A. E. *et al.* Engineered 3D Immuno-Glial-Neurovascular Human miBrain Model. Preprint at https://doi.org/10.1101/2023.08.15.553453 (2023).

39.   Bowles, K. R. *et al.* ELAVL4, splicing, and glutamatergic dysfunction precede neuron loss in *MAPT* mutation cerebral organoids. *Cell* **184**, 4547-4563.e17 (2021).

40.   Karch, C. M. *et al.* A Comprehensive Resource for Induced Pluripotent Stem Cells from Patients with Primary Tauopathies. *Stem Cell Reports* **13**, 939–955 (2019).

41.   Brion, J. P., Octave, J. N. & Couck, A. M. Distribution of the phosphorylated microtubule-associated protein tau in developing cortical neurons. *Neuroscience* **63**, 895–909 (1994).

42.   Yang, Y.-H., Wen, R., Yang, N., Zhang, T.-N. & Liu, C.-F. Roles of protein post-translational modifications in glucose and lipid metabolism: mechanisms and perspectives. *Mol Med* **29**, 93 (2023).

43.   Yu, Y. *et al.* Developmental regulation of tau phosphorylation, tau kinases, and tau phosphatases. *Journal of Neurochemistry* **108**, 1480–1494 (2009).

44.   Liu, F. & Gong, C.-X. Tau exon 10 alternative splicing and tauopathies. *Mol Neurodegeneration* **3**, 8 (2008).

45.   Luna-Muñoz, J., Chávez-Macías, L., García-Sierra, F. & Mena, R. Earliest Stages of Tau Conformational Changes are Related to the Appearance of a Sequence of Specific Phospho-Dependent Tau Epitopes in Alzheimer's Disease1. *JAD* **12**, 365–375 (2007).

46.   Hyman, B. T. *et al.* National Institute on Aging–Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease. *Alzheimer's & Dementia* **8**, 1–13 (2012).

47.   Bennett, KeirynL. *et al.* The 2012/2013 ABRF Proteomic Research Group Study: Assessing Longitudinal Intralaboratory Variability in Routine Peptide Liquid Chromatography Tandem Mass Spectrometry Analyses*. *Molecular & Cellular Proteomics* **14**, 3299–3309 (2015).

48.   Eldjarn, G. H. *et al.* Large-scale plasma proteomics comparisons through genetics and disease associations. *Nature* **622**, 348–358 (2023).

49.   Young, N. L. *et al.* High Throuput Characterization of Combinatorial Histone Codes. *Molecular & Cellular Proteomics* **8**, 2266–2284 (2009).

50.   Shan, X. *et al.* Fully defined *NGN2* neuron protocol reveals diverse signatures of neuronal maturation. *Cell Reports Methods* **4**, 100858 (2024).

51.   Bertucci, T. *et al.* Improved Protocol for Reproducible Human Cortical Organoids Reveals Early Alterations in Metabolism with *MAPT* Mutations. Preprint at https://doi.org/10.1101/2023.07.11.548571 (2023).

52.   Stanton, A. E. *et al.* Engineered 3D Immuno-Glial-Neurovascular Human miBrain Model. Preprint at https://doi.org/10.1101/2023.08.15.553453 (2023).

53.   Devaraj, N. K., Upadhyay, R., Haun, J. B., Hilderbrand, S. A. & Weissleder, R. Fast and Sensitive Pretargeted Labeling of Cancer Cells through a Tetrazine/ *trans* -Cyclooctene Cycloaddition. *Angew Chem Int Ed* **48**, 7013–7016 (2009).

Consortia

The members of Nautilus Biotechnology are Jamie E. Anderson, Fletcher Bain, Filip Bartnicki, Issa Beekun, Giovanni Bellesia, Rajat Bhatnagar, Keith Bjornson, Phillip S. Brereton, Stephen Brown, Vivekananda Budamagunta, Kaitlyn Burke, Chieh-Li Chen, Kevin Chen, Priscilla L. Chen, Krista Cosert, Danya Cruver, Michael A Darcy, Wade Dugdale, Ahana Dutta, Jarrett D. Egertson, Michael E. Flaster, Tyler J. Ford, Keith Frankie, Taryn E. Gillies, Keith Gneshin, Hamid R. Golnabi, Sam Grillo, Rob Grothe, Cynthia Y. Guerrero, Ross Hartley, Brooks D. Hoehn, Jr., Huber Martin, Andreas FR Huhmer, Christina E. Inman, James Joly, Kevin M. Jin, Maryam Jouzi, Kara Juneau, Greg Kapp, Kota Kaneshige, Stanislav Katsyuk, Eun Kim, Rajesh Kota, Jamie Rose Kuhar, Jonathan B. Leano, Darrel Lee, Winnie Leung, Cara Li, Bryant Lowry, Parag Mallick, Mara Manskie, Carlos F Martinez, Kevin McVey, Jonathan C. Menger, Denise A. Miller, Eva Monsen, Anna Mowry, Grant Napier, Nicholas Nelson, Nicole Newland, Maureen R. Newman, Brittany Nortman, Adedeji Ogunyemi, Jayetha Panakkadan, Sujal Patel, Rukshan T. Perera, Nguyen Pham, Keshav Prabhu, Maximilian O Press, Hongji Qian, Torri E. Rinker, Adam J. Roberts, Julia K. Robinson, Aimee A. Sanford, Ali Najafi Sohi, Subra Sankar, Parag Shekher, Genee Victoria Skulich, Irene Tang Sparck, Isaac B. Sprague, Noah Steiner, Robert Stephenson, Juanfeng Sun, Kentaro Suzuki, Scott M. Tabakman, Steven J Tan, Nicholas Tardif, Sonal S. Tonapi, Jasmine Trinh, Maria Villancio-Wolter, Jacinto Villanueva, Zhiqiang Wang, Lisen Wang, Gwen Weld, Sheri K. Wilcox, Katherine Winters, Feng Yan, Yvonne Yeung, Zhengjian Zhang, All members are affiliated with Nautilus Biotechnology, San Carlos, CA 94070.

Author contributions

VB, SG, MJ, and JJ designed and executed the experiments on the Nautilus platform, RW, BN, and ZZ designed the Tau enrichment method, developed the sample preparation methods, developed the single molecule library preparation methods, and prepared the single molecule libraries for analysis. ST prepared peptide and proteoform standards. PB and JS prepared affinity reagents. JJ performed the data analysis. GK, KJ, PM conceptualized experiments and contributed to the manuscript. TB, SL, NeuraCell prepared and performed the organoid experiments. ST contributed to the manuscript. DA performed the mass spectrometry analysis. AFH, BN, CEI, KJ, MEF, PM, RG, SKW, VB, ZZ contributed to conceptualization. ANS, BDH, BN, GB, HQ, JDE, JKR, JT, JV, KJ, MEF, MJ, MRN, PI, PM, RG, RK, RTP, SG, SJT, SKW, SMT, VB, ZZ contributed to methodology. AJR, CC, DAM, DL, EM, HRG, KC, KM, KW, LW, MEF, MM, MV, NT, RB, RG, RH, SB, SK, WL contributed to software. AO, BN, JS, KJ, KM, KP, KW, MEF, MJ, MV, PSB, RG, RTP, SJT, SST, TER, VB, ZW, ZZ contributed to validation. FB, JDE, JS, KM, MJ, RG, ZZ contributed to formal analysis. AD, AO, BDH, BN, FB, FY, JBL, JS, JT, KB, LW, MAD, MEF, MJ, MV, NS, PI, RG, RTP, SG, SG, SMT, VB, ZZ contributed to investigation. AAS, AD, BL,

BN, CYG, DL, EK, FB, FY, GVS, JBL, JCM, JV, KB, KC, KF, KK, KM, KMJ, KS, MAD, MOP, NS, PSB, RS, TER, WD, YY, ZZ contributed to resources. AO, CL, FB, FY, KC, KM, MJ, MOP contributed to data curation. AFH, CEI, JJ, PM, VB, ZZ contributed to writing the original draft of the manuscript. AFH, DCB, BN, JDE, JJ, JRK, KJ, KS, ST, PM, TJF, VB, ZZ contributed to reviewing and editing the manuscript. JDE, JJ and PM contributed to visualization. FY, GW, HM, JRK, JV, KJ, MJ, PM, SKW, SMT, SP, ZZ contributed to supervision. DL, GW, JCM, JEA, MAD, PM, SKW, SMT, SP, YY, ZZ contributed to project administration. AM, PM, SP contributed to funding acquisition.

Competing interest declaration

The authors have declared the following conflict of interest: The authors listed in the Consortium Nautilus Biotechnology are current or past employees of Nautilus Biotechnology and have financial interest in Nautilus Biotechnology. Arnott D., Lipka J., Pandya NJ., Rougé L., Wendorff T.J., Kirkpatrick D., Rohou A., are current or past employees of Genentech Corporation and have financial interest in Genentech Corporation.

Acknowledgements

MAPT Human induced pluripotent stem cells lines (**Supplemental Table S4**) were provided through the generous support of the Tau Consortium and the Rainwater Charitable Foundation and are available at https://www.neuralsci.org/tau. We are indebted to the NSCI NeuraCell core facility for organoid production and technical support.

We additionally acknowledge funding from The Regenerative Research Foundation (NSCI investigators); The Rainwater Charitable Foundation and the Tau Consortium and Cure PSP (ST, TB, DCB); NIH: RF1NS123568 (ST, DCB), 1RF1NS142335 (ST, TB).

Additional information

Supplementary Information is available for this paper. Correspondence and requests for materials should be addressed to paragm@nautilus.bio or karaj@nautilus.bio