1
2
3
4
5 **Variation in social feeding behavior and interactions among *Caenorhabditis* nematodes**
6

7 Haskell D[1], Palo J[1], Eugene RFH[1], Large CRL[1], Hart MP[1]
8
9 Affiliations
10 [1]Department of Genetics, Perelman School of Medicine, University of Pennsylvania
11 * corresponding author Michael P. Hart – email: hartmic@pennmedicine.upenn.edu

Abstract

The ability to respond to complex stimuli and environmental cues is essential for organisms to survive and reproduce. Responding to a wide range of stimuli requires a neuronal network that can integrate cues and execute behavioral responses. Evolution of behaviors occurs ubiquitously in most established ecological niches, even among closely related species. To uncover the genetic and neuronal drivers of evolving behaviors, we have taken advantage of the large and [relatively] ancient divergence in the *Caenorhabditis* genus to ask how different *Caenorhabditis* nematodes respond to environmental stimuli and whether behavioral traits shared or distinct. Here, we assayed foraging behaviors of twelve members of the *Caenorhabditis* clade, including members of both the *elegans* and *japonica* supergroup, and the basal taxon *C. monodelphis*. For each species, we analyzed social feeding and bordering behaviors, which are well characterized in *C. elegans.* These behaviors are the functional readout of complex sensory integration of multiple sensory cues including pheromones, touch, O_2/CO_2 concentration, and attractive and noxious stimuli. We hypothesized that the evolutionary divergence between species would correlate to divergence in these behaviors. We observed wide variation in aggregate social feeding and bordering behaviors of hermaphrodite and female animals, but the variation did not correlate with evolutionary relatedness of the species. Combination of both sexes of individual species increased aggregation behavior of select species that had lower levels of aggregation in single sex assays. Addition a second species with *C. elegans* significantly reduced aggregate feeding behavior of *C. elegans*, but not the other species. Intraspecies and interspecies interactions can modify behavioral paradigms. Overall, we find that foraging and social feeding behaviors vary widely across *Caenorhabditis* species, likely due to species-specific responses and integration of context sensory cues. In general, the clade represents a compelling model to dissect evolution of behavior across diverse environments and a large timescale.

Introduction

Behavior is how organisms interact with their environment and respond to both internal

and external stimuli. Simple forms of life, especially those that represent unicellular phyla, rely

heavily on the detection of chemical and tactile stimuli through direct contact[1]. In contrast, broad

sensory specialization is heavily emphasized in Eukaryotes, which range not only in the

complexity of their specialization (including the evolution of differentiated and highly specialized

neuronal tissues) but also in the manner and breadth of their behavioral responses to environmental

stimuli [2,3]. The evolution of complex behavior can be loosely correlated with the evolution of

multicellularity and complex tissues[4]. Likewise, tissue differentiation also gave rise to neural

tissues, which in their simplest form, link networks of sensory cells together forming a basic neural

circuit. In more complex Eukaryotes, more sophisticated neural networks, utilizing multiple

sensory and neuronal cell types form the basis of complex organs such as neural ganglion, nerve

tracts, and the brain [5,6]

Changes in behavioral traits occurs regularly in most eukaryotic organisms, especially

between related species[7]. Logically, rapid shifts in behavior should be tightly coupled with the

congruent alteration of their physiological and genetic drivers as highlighted by several studies in

Drosophila and related species [8,9]. Small changes in gene regulatory networks, modest alterations

in neural circuits, or mutations influencing receptor/ligand interactions are all sufficient to alter

behavioral paradigms in an often-unpredictable manner. Without the ability to identify step-wise

drivers of shifting behavioral paradigms it can be challenging to understand how selective

pressures lead to modification of underlying genetic and molecular factors, which ultimately

dictate the neurological activity underlying complex responses to stimulus.

Compounding the study of behavioral evolution is the complex nature of behavior itself.

To truly differentiate between behavioral paradigms requires dissection at multiple levels of

organismal complexity, starting with changes to the genome and epigenome, and culminating in

intra-circuit alterations leading to modification in the complex network of integrated and often co-

dependent behaviors. While comprehensive dissection of genetic and molecular drivers of

altered/novel behaviors is rarely achieved, we can utilize existing genetic information in clades

such as *Caenorhabditis* to inform research to begin circuit level descriptions.

Amongst common lab models, the nematode *Caenorhabditis elegans* is a particularly

powerful tool for studying behavior. In addition to the many genetic tools, a fully mapped neural

connectome (chemical, electrical, and peptidergic) provide a powerful model to link stereotyped

behaviors with alterations of neuronal activity at both a single cell and circuit level. In addition,

the *Caenorhabditis* clade of nematodes represents a unique opportunity to study evolution of

behavior in a diverse and evolutionarily ancient lineage **(Figure 1)**. Nematodes of this clade are

found in a worldwide dispersal, with populations found in nearly every ecological niche probed

for their presence[10]. In the wild, members of the *Caenorhabditis* nematodes persist in a free-living

state generally colonizing bacteria-rich substrates, such as rotting fruit, allowing them to utilize

the local microbiota as food source. Considering their global dispersion, individual members of

the clade experience significant variation in local environment, including differences in

temperature, humidity, concentration of oxygen, and dominant microbiota[11]. Notably, these

species are rarely isolated from homogenous populations; *elegans* for example is frequently co-

isolated with *briggsae* and/or *remanei* from the same substate. Altogether, the *Caenorhabditis*

clade presents a compelling model through which to query the evolution of behavior across a large

evolutionary timescale and a wide range of environments.

Due to high levels of variation among local environments, we chose to first assay foraging

behaviors, which we reasoned would be ubiquitously present among the clade given likely shared

feeding strategies. One of the most well-characterized feeding behaviors is termed social feeding

and is defined by social interactions that occur between individuals feeding in the same local

environment[12]. Although not well defined, one can speculate that this behavior has costs and

benefits for individual animals. Animals may be attracted to high density food that is resource rich

for obvious nutrient seeking reasons, but animals populating a food/bacteria rich location can also

result in spreading of the food source (bacteria) to surrounding areas. Further, group feeding could

facilitate short-distance signaling between animals (perhaps through pheromones) and/or increase

opportunities for mating, but it seems clear that this behavior is environment and context

dependent[13,14]. Early surveys of wild *elegans* isolates showed a surprisingly high level of

variability in social feeding behaviors. Contrary to the widely used N2 Bristol lab strain, which

displays solitary feeding, many wild isolates from a range of environments show higher levels of

social feeding[12]. de Bono & Bargmann attributed the solitary feeding behavior of N2 Bristol strain

to a gain of function missense mutation in the *npr-1* gene (ortholog of neuropeptide Y receptor).

This mutation results in an amino acid change in the third intracellular loop of the NPR-1 G-protein

coupled receptor , whereas loss of function mutants in the *npr-1* gene result in social feeding in

elegans[12]. For the purpose of this study we utilize the *npr-1(ad609)* loss of function mutant to

induce social feeding in the N2 lab control strain, which represents the most reliable and robust

positive control for social feeding behaviors[15].

Subsequent work further elucidated the role of NPR-1 and other neuronal drivers of solitary

and social feeding, including that *elegans* social feeding behavior is driven primarily through the

integration of environmental cues including pheromones, O_2/CO_2 concentrations, and perception

of touch and noxious stimuli [13,16–19]. Sensory neurons including ADL, ASH, ASK, ADE and AWB

form a circuit to detect the local sensory environment which is then integrated primarily through

the interneuron RMG to alter locomotion paradigms[16,17]. Genetic dissection of this sensory circuit

has uncovered distinct roles for the neuropeptide receptor NPR-1[12,15] and gap junction connections

between involved sensory neurons and the RMG interneuron[16,20]. More recent work has implicated

glutamate signaling from ADL and ASH sensory neurons, where conserved synaptic adhesion

molecules (NRX-1 and NLG-1) contribute to social feeding, and *nrx-1* impacts architecture and

function of ASH chemical synaptic connections[15]. Therefore, social feeding behavior relies on co-

dependent signaling of neuropeptides, and electrical and chemical synapses. Together, these

mechanisms serve to fine-tune the balance between solitary and social feeding behaviors[15]. Given

the complexity of this behavioral circuit, we set out to test whether social feeding behavior is

conserved throughout the *Caenorhabditis* clade of nematodes.

In this study, we compared social feeding behaviors of twelve *Caenorhabditis* species to

multiple strains of *elegans* (**Figure 1**). This clade of nematodes is not only relatively ancient, but

also has significant divergence over evolutionary time, with an estimated genomic divergence of

> 20 million years between *elegans* and the four most-closely related species[21]. Two large

subgroups, *elegans* and *japonica*, have emerged based on genomic data providing multiple

lineages to interrogate. *monodelphis*, representing the basal taxon of the clade, is estimated to have

diverged over 100 million years ago, an evolutionary timescale far beyond that of the split between

mice and humans [22]. Lastly, the true strength of this model lies the opportunity to study "wild"

populations of nematodes (representing both hermaphroditic and gonochoristic species) in a

controlled lab setting applicable to medium-throughput behavioral assays. When considering the

evolution of behavior within the *Caenorhabditis* clade we set out to compare differences amongst

the species, providing a base on from which future work investigating differences in sensory cues,

neuronal function, and circuitry can be launched.

180
Figure 1. The *Caenorhabditis* clade of nematodes is an evolutionary and diverse lineage. (A)
Phylogenetic tree of *Caenorhabditis* clade showing the two large subgroups (*elegans* and
japonica) and the basal taxon *monodelphis* (adapted from *Caenorhabditis* Genome Project).
Species included in behavioral assays are highlighted and color-coded based on reproductive
strategy (red = hermaphroditic or orange = gonochoristic).

Using the twelve members of the *Caenorhabditis* clade we have found significant variation

in social feeding behaviors of hermaphrodites and females amongst the different species, with all

but two showing reduced levels of aggregate feeding compared to the *elegans* social feeding

control strain, and half of the species display significantly more social feeding that the *elegans*

solitary control strain. Furthermore, by testing more ecological relevant populations in our assays

we have begun to tease out the impact of intraspecies and interspecies interactions. Mixing males
with females of individual species altered aggregate feeding behavior in two species when
compared to homogeneous female populations. Lastly, combining a second species with social
feeding *elegans* was sufficient to significantly reduce aggregation feeding of *elegans*. This work
represents the first characterization of foraging and social feeding behaviors across the
Caenorhabditis clade of nematodes and lays the groundwork for the dissection of ecological,
cellular and circuit level mechanisms of conserved behaviors.

Table 1. Reference table for *Caenorhabditis* species information.

Species	Subgroup	Strain	*RS	*GH	Genome Accession	*MC	Behavioral Characterization
elegans							
N2	elegans		♀	lab strain	PRJNA13758	yes	yes
CB4856							
briggsae	elegans	AF16	♀	lab strain	PRJNA10731	yes	yes
imperialis	elegans	EG5942	MF	30x inbred	GCA_963572205.1	n/a	no
nigoni	elegans	JU1422	MF	25x inbred	PRJNA384657	n/a	no
remanei	elegans	PB4641	MF	20x inbred	PRJNA577507	yes	some
tropicalis	elegans	JU1373	♀	n/a	PRJNA53597	n/a	no
japonica	japonica	DF5081	MF	20x inbred	GCA_963572235.1	yes	no
afra	japonica	JU1286	MF	20x inbred	GCA_963570955.1	n/a	no
sulstoni	japonica	JU2788	MF	25x inbred	PRJEB12601	n/a	no
virilis	drosophilae	JJU1968	MF	25x inbred	GCA_964036255.1	n/a	no
drosophilae	drosophilae	DF5112	MF	20x inbred	PRJEB67483	n/a	no
uteleia	guadalopensis	JU2585	MF	25x inbred	GCA_963573275.1	n/a	no
monodelphis	n/a	JU1667	MF	20x inbred	GCA_964197825.1	yes	no

*RS - Reproductive Strategy; ♀ represents hermaphroditic species, MF represents gonochoristic species

* GS- Genetic Heterogeneity; number of times inbred according to CGC strain information

*MC - Morphological Characterization

<u>Methods</u>

Strains Maintenance

Animals were maintained under normal growth conditions (~20° C) on normal NGM media[23].

Plates were seeded with *E. coli* OP50 to provide food. Strains utilized in this study are listed in

Table 1. *npr-1*(*ad609*) and *nrx-1*(*wy778*) mutations were confirmed in *elegans* strains and were

used as positive controls. All other species were obtained from the CGC and maintained by

chunking.

Social feeding behavior assay

Standard 6-well culture plates were filled with 6 mLs of standard NGM media and then seeded

with 75 µL of *E. coli* OP50 and allowed to dry overnight. For the standard assay, 50 animals were

picked onto clean plates and then moved onto the seeded wells, transferring as little bacteria as

possible to prevent buildup. For standard social feeding and bordering assays (**Figure 2B&C**) the

50 animals were either female or hermaphrodites depending on the species. For the mixed sex

assays, we combined 25 males with 25 females/hermaphrodites (depending on species).

Regardless of the composition of the assay, worms were picked during and placed on the assay

during their fourth larval stage. Female or hermaphrodite animals were staged based on the

presence of half-moon shape characteristic of a L4 developing vulva. Males were staged based on

relative size, germline features, and morphological development of the tail. After placing the

animals on the 6-well plate, a small amount of Tween20 was used to thinly coat the lid to prevent

condensation forming during imaging. The plates are imaged using the WormWatcher automated

imaging platform (Tau Scientific) for a total of 20 hours with a cluster of 10 images (over 1 minute

timecourse) being taken every hour. To quantify social feeding behavior each well was manually

scored from blinded images, where a worm was considered aggregating if was contacting two or

more other worms[12,15]. Bordering was assayed by calculating the percentage of solitary animals

that we in direct contact with the border region of the bacterial lawn (with the border defined as

approximately half a body length in width). Data shown is from the 15-hour mark, which placed

the animals of all species in the middle of Day 1 of adulthood.

Mixed species assays

In order to assay the effect of mixed species, we first generated a fluorescently tagged

elegans strain to differentiate them from the other species while imaging behavior. *npr-1(ad609)*

elegans were injected using standard technique with plasmid *myo-2p*::*mScarlet* in order to label

pharyngeal muscle (25 ng/µL). Strong expression of the transgene allows visualization of a single

worm when in an aggregate. To perform the assay a standard 5cm NGM plates was seeded with

25 µl of *E. coli* OP50 and allowed to dry. To mix the populations, 15 *elegans* expressing *myo-*

2p::*mScarlet* were placed on the bacteria lawn along with 15 of *monodelphis* or *nigoni,*. The plates

were placed under a fluorescent dissecting scope and allowed to sit for 45 minutes to mitigate any

effects of plate movement. The plate was then imaged using white light (for a reference image)

and green light (~546 nm, to capture the *mScarlet* fluorescence). Consistent with previous assays,

animals touching two or more individuals were scored as aggregating.

Orthology analysis

We used the genome sequence and gene annotations from Wormbase [24] and the *Caenorhabditis*

Genome Project to find orthologous gene sets (orthogroups) in non-*elegans Caenorhabditis*

species. First, we extracted the longest isoform from every gene using AGAT[25]. Then, using the

protein sequence, we constructed orthogroups using OrthoFinder[26,27] using the sequence aligner,

Diamond, in ultra-sensitive mode[28].

Statistics and reproducibility:

The number for each experiment was based on previous studies and effect size, with each

experiment performed with at least 3 independent replicates and each trial performed with matched

controls. All data were analyzed and plotted in GraphPad Prism 10 and statistical significance was

determined using one-way ANOVA with Tukey's post-hoc test. For comparisons of two data sets,

a two-tailed unpaired t-test was used to compare significance. Error bars on figures represent

standard error of the mean (SEM) and p-values are shown in each figure to indicate significance

($P < 0.05$).

<u>Results</u>

Social feeding behaviors differ across the *Caenorhabditis* clade

We assayed aggregation and bordering behavior using a quantitative imaging setup[15].

Assays were performed as previously described, with 50 female animals being used instead of 50

hermaphrodites in species that are gonochoristic. In our assay we included four *elegans* control

strains; N2 'Bristol' that are solitary, *npr-1(ad609)* that are social feeding, *npr-1(ad609);nrx-*

1(wy778) that display intermediate social feeding, and CB4856 'Hawaiian' *elegans* that display

intermediate social feeding. We observed wide variation in mean aggregation behavior amongst

members of the genus, with some species like *imperialis* and *virilis* showing relatively high levels

of aggregation, although not to the level of the *elegans* social control (*elegans npr-*

1(ad609))(**Figure 2A-B**). Interestingly, *monodelphis* essentially phenocopies the *elegans* solitary

controls (N2) with little to no aggregation (**Figure 2A-B**). Outside these three examples, we find

that the majority of the species have an intermediate level of aggregation, consistent with a

modulated aggregation response. Further, we observe that within some individual species, a large

level of variation between replicates/experiments, namely in *japonica, uteleia, and remanei*

(**Figure 2A-B**). Somewhat surprisingly, there was no clear pattern between evolutionary

relatedness and the observed level of aggregation. In addition to aggregating, when social feeding

behavior occurs, *elegans* aggregate in groups near or in the border of the bacterial lawn where the

bacteria is the thickest[12] (**Figure 2A**). When we examined bordering behavior we noted two

distinct trends. First, a general downward trend in bordering tendency the more distantly related

the species are to *elegans*, and second, that the overall trend of widespread variation is also

observed when considering this bordering phenotype (**Figure 2A&C**).

Figure 2. Aggregation behavior is diverse across *Caenorhabditis* clade. (A) Representative images of select *Caenorhabditis* species aggregation feeding and bordering behaviors. (B) Quantification of number of aggregating animals in 13 members of the *Caenorhabditis* clade. Significant variation is present among members of the clade, and also within individual species. The 4 *elegans* groups assayed represent the negative control (N2 'Bristol'), positive control *npr-1(ad609)*, modulated positive control *npr-1(ad609);nrx-1(wy778)*, and wild isolate CB4856 'Hawaiian', respectively. Bar graphs represent the mean values across biological replicates (each replicate indicated by a dot). Error bars show stand error of mean. One-way ANOVA with Tukey's post-hoc test was used for statistical comparisons, with p-values indicated in standard format (

291 p=0.05(*), p=0.01(**), p=0.001(***), p= <0.001 (****)). Hermaphroditic species are labeled in
292 red, while gonochoristic species are labeled in black. **(C)** Quantification of % solitary animals on
293 border in 13 members of *Caenorhabditis* clade. Significant variation is present among members
294 of the clade, and also within individual species. Graphical representations and statistical analysis
295 are consistent with previous panel.
296

To visualize variation among species in both aggregation and bordering phenotypes we
generated heat-maps showing the adjusted p-values generated by one-way ANOVA of all
combinatorial comparisons (**Figure 3A & B**, respectively). Plotting the phenotypes in a XY plot
reveals a relatively broad dispersal, without a clear clustering among closely related species
(**Figure 3C**). Overall, the plot emphasizes that while that *elegans* N2 and *npr-1*, and *monodelphis*
are relative extremes compared to the rest of the species, such extreme can occur both in
domesticated and "wild" populations. Consistent with their independent evolution of
hermaphrodism[31], there does not seem to be any significant clustering of the hermaphroditic
species. Overall, in characterizing two social feeding behaviors we identified unanticipated levels
of variation both among and within species, suggesting out assays may be subject to one of more
additional variables that are contributing to variation. Given the wide range of environments from
which these species were isolated, we speculated that the lab conditions may represent less than
optimal conditions for several species and therefore introduce an additional element of variation.

Figure 3. Statistical analysis of phenotypes. (A) Heat-map of ANOVA-generated p-values of all
pairwise comparisons in aggregation assay. **(B)** Heat-map of ANOVA-generated p-values of all
pairwise comparisons in bordering assay. **(C)** XY data plot showing clustering off species when
comparing aggregation and bordering phenotypes. As anticipated, *elegans* N2 and *monodelphis*,
and *elegans npr-1* represent the low and high extremes of the cluster, respectively. Consistent with
the figures above, hermaphroditic species are labeled in red.
317

Ecological drivers modulate social feeding behaviors

Given the differences in environment and reproductive strategies amongst members of the
genus we reasoned that the makeup of the individual populations may influence social feeding
behaviors. To test this hypothesis, we repeated the standard aggregation assays using mixed sex
populations (25 males + 25 females, or 25 hermaphrodites in hermaphroditic species with presence
of males) (**Figure 4A & B**). For this assay we focused on a subset of the species displaying high
level of variation in the female/hermaphrodite only assays, in part to test if a more "ecologically
relevant" populations might reduce some of the observed variation. Overall, for most species the
mixed sex populations did not alter aggregation behavior compared the single sex populations
(**Figure 4A & B**). However, *monodelphis* and *tropicalis,* showed significantly higher aggregation
in the mixed sex population assay compared to the single sex hermaphrodite/female population
assay (**Figure 4A**). These results again highlight the impact of local environment and context on
variation in behaviors, whereas these assays reflect a modulated response to real-time sensory-
environment landscape.

To further explore other ecologically relevant population dynamics (such as interspecies
cohabitation and interactions, for example escape from predation[32]), we tested whether combining
additional species with *elegans* would influence their social feeding behavior. We first tested
whether a reduced population size (30 animals) or the addition of a *myo-2p* fluorescent transgene
would impact the aggregation behaviors of each species. Aggregation in both *nigoni* and
monodelphis were significantly reduced compared to *elegans* (15 *myo-2p+* and 15 *myo-2p-*
)(**Figure 4C**) and significantly different from one another, which is consistent with the 50 animal
aggregation assay results (**Figure 2**). Furthermore, the inclusion of the *myo-2p* transgene did not
impact *elegans npr-1*(*ad609*) aggregation behavior. Next, we mixed 15 *myo-2p+ elegans npr-
1(ad609)* social feeding controls with 15 *myo-2p- elegans, monodelphis,* or *nigoni* on a seeded
NGM plate and allowed the mixed populations to acclimate for 45 minutes. After the acclimation
period, the plates were imaged and assayed for aggregation behavior. We aimed to understand if
the mixing of two distinct populations would alter the aggregate feeding dynamics of either or both
species. In the mixed species population assays, we noted that addition of *nigoni* to the *myo-2p+*
elegans population significantly reduced aggregation behavior of elegans, while with the addition
of *monodelphis* the reduction was not significant but trended strongly downward (**Figure 4D&E**).
Interestingly, we did not observe an impact of *elegans* on behavior of the other species. We

reasoned that disruption of *elegans* aggregation behavior could be the result of a number of
changes to environmental cues present on the plates that result from mixing populations.

Figure 4. Ecological drivers can modulate social feeding behaviors. (A) Representative images and **(B)** quantification of number of aggregating animals of *Caenorhabditis* species in mixed sex populations. *tropicalis* and *monodelphis* both showed significant differences in aggregation when males were added to the population. Bar graphs represent the mean values across biological replicates (each replicate indicated by a dot). Error bars show stand error of mean. Standard T-test was used for individual comparisons; p-values are indicated in standard format (p=0.05(*), p=0.01(**), p=0.001(***), p= <0.001 (****)). Hermaphroditic species are labeled in red, while gonochoristic species are labeled in black. **(C)** Quantification of 30 animal single-species controls. **D)** Quantification of mixed species assays, where 15 *elegans myo-2p+* animals were mixed with 15 *elegans myo-2p-, monodelphis,* or *nigoni*. Combining *elegans* and *nigoni* significantly lowers aggregation of *myo-2* expressing *elegans* compared to their *elegans* alone control (left), while addition of monodelphis caused an insignificant but trending reduction (left). *monodelphis* and *nigoni* have significantly lower aggregation than their *elegans myo-2p–* control (right). **(E)** Representative images of mixed species assays. *myo-2p* expressing *elegans* are indicated by arrows (top left window).

Caenorhabditis clade has high level of conservation in NPR-1 receptor protein

Given the variation in the social feeding behaviors in the *Caenorhabditis* genus, we wondered about mechanisms that might result in species differences. Morphological differences likely drive some of the changes, however, there has been very little characterization in most of the species we assayed. There are several papers examining the development of the vulva [33–35], tail morphology[36–38] and body size[39], however there is still a large deficit in the literature. This lack of species characterization is especially apparent when considering the nematode nervous systems, with the only other *Caenorhabditis* species with any significant neuronal characterization being *briggsae*[40,41]. To bypass the lack of circuit-based knowledge within the clade, we leveraged several known genetic drivers of social feeding behavior in *elegans*, notably the neuropeptide receptor NPR-1. To compare *npr-1* across species, we first queried both existing and draft genome assemblies for orthologues using OrthoFinder[27]. Given the size and relative domain conservation of the NPR family of genes in *elegans* we were somewhat surprised to identify a single orthologue of NPR-1 in all species except for *nigoni*, which had multiple related paralogues and was thus excluded from our analysis. To characterize sequence conservation among species we used CLUSTAL V5.0 analysis to perform a multiple sequence alignment to produce an NPR-1 alignment (**Figure 5A**). The alignment showed there is a high level of conservation among the species, even when considering the most evolutionary divergent species, like *monodelphis*. We queried three residues in NPR-1 that when mutated have been shown to induce social behaviors[12] and found them to be well conserved and wildtype. Interestingly, when compared to a consistent

protein length among species, *uteleia* also appears to have a large expansion of its *C-terminal* tail to include a novel Hox DNA-binding homeodomain. **(Figure 5B)**. However, given the draft nature of the *uteleia* genome assembly, this expansion could also be an artifact of an adjacent or overlapping gene feature.

Figure 5. Protein alignment of NPR-1 orthologues in amongst *Caenorhabditis* nematode species. **(A)** NPR-1 orthologues were queried from existing *Caenorhabditis* genomes or assemblies based on sequence similarity. A single orthologue was identified in each species with the exception of *nigoni* which was excluded due to the lack of a confident orthologue. Overall,

397 NPR-1 is relatively well conserved across the clade, with the exception of *uteleia* which showed a
398 large expansion of amino acid sequence to include a novel Hox homeodomain **(B)**.
399

<u>Discussion</u>

Here, we examined several social foraging behaviors of an evolutionarily diverse clade of
Caenorhabditis nematodes to better understand how such behaviors differ and may have evolved
between species. Across these divergent species of nematodes, we noted several surprising trends
within a set of social feeding behaviors that that underpin a shared general feeding strategy.
Notably, we confirmed of *elegans* (*npr-1*) is a relative extreme when compared to other species
but can be partially phenocopied by several wild isolates and other species. In contrast, most of
the species we assayed (including the wild *elegans* isolate CB4856 'Hawaiian' strain) showed
intermediate levels of aggregation, while basal taxon *monodelphis* showing little to no aggregation.
Furthermore, we uncovered a surprising trend when examining bordering behavior among species,
notably that the phenotype seems to generally decline the further away the species is in relatedness
to *elegans*. While this trend is not perfect, it does again highlight the idea that N2 'Bristol' *elegans*
may represent a behavioral outlier when compared with wild *elegans* isolates and most other
species.

We noted that moderate to high levels of variation were present in most species when
considering both aggregation and bordering phenotypes. The extensive variation present within
several of the species raises the possibility that an active mechanism is driving heterogeneity
beyond the expected variation present in most behaviors. This unexpected variation could be in
part due to nuanced genetic heterogeneity present between individuals, maintained in part through
selection pressure. *elegans* strains have been artificially selected and lab-evolved (in part because
they reproduce clonally), while in comparison, most of the species we assayed represent far more
heterogeneous populations. Several species have been backcrossed and inbred extensively from
their wild isolates, effectively rendering them lab strains **(Table 1)** after undergoing selection for
domestication. We can speculate that some of the species likely retain higher levels of genetic
heterogeneity, especially among genetic elements such as single nucleotide variations (SNPs).
While the vast majority of SNPs are unlikely to be the causative agent directly underlying a
phenotypic change, single nucleotide changes can drastically alter behavior, as highlighted by
single nucleotide variation in the neuropeptide receptor NPR-1. Many wild isolates of *elegans*
naturally aggregate and display social feeding, but in the lab N2 Bristol strain, a single adenine to

guanine substitution in NPR-1 is sufficient to produce a gain of function phenotype, eliminating

social behavior and inducing solitary feeding[12]. Lastly, it is not unreasonable to speculate that

SNPs present within individuals may have a modulatory effect within particular behavioral

circuits, especially if they are closely associated with critical neuronal genes or regulatory

elements. This concept is highlighted in recent work showing mutations in the *daf-19* transcription

factor are able to dramatically alter chemosensory and oxygen-sensing circuits leading to a

disruption social feeding behaviors in both *C. elegans* and *P. pacificius*[42].

Outside of genetic mechanisms, it is very likely that some of the variation seen in social

feeding behaviors is a result of the lab environment in which these assays are being performed. In

the lab we make every effort to maintain temperature at 20 °C with low to moderate humidity,

which is ideal for the growth and maintenance of lab strains like *elegans* and *briggsae*. However,

many of the other *Caenorhabditis* species were isolated from drastically different climates and

environments and likely underwent adaptation of their environmental-sensory circuits to these

divergent conditions in contrast to those in which we assayed them. While testing ecologically

relevant temperatures and humidity levels for each species is logistically beyond the scope of this

study, one might imagine that social feeding behaviors might be alter if those conditions were to

more accurately reflect what occurs in the wild.

Furthermore, the substate on which we feed the animals is resoundingly artificial, using a

monoculture of a lab strain of *E. coli* (OP50) on an agarose surface is hardly representative of the

natural ecology of wild feeding substates. When considering the natural environments in which

these nematodes normally reside, the microbiomes can also be dramatically heterogenous

depending on location. Most natural feeding substates contain a range of bacterial species, some

of which are viable food sources and some that may represent nematode pathogens. To reconcile

these differences, recent research has characterized food choice preferences in *elegans*, uncovering

their preference for *bacillus* species over the traditional lab *E. coli* strains[43]. Even within *E. coli*

species, *elegans* tend to prefer nutrient dense strains, likely to supplement for essential nutrients,

and potentially leading to changes in metabolism[43,44]. It is not surprising then that different food

sources can lead to dramatic variations in behavior. Best studied is the attraction/avoidance

response to food, which allows the animals to avoid potentially pathogenic bacteria while

optimizing nutrient uptake[45,46]. Lastly, *elegans* has been shown to modulate their feeding behaviors

based on the density of food, where both solitary and social strains disperse when food density if

strongly diminished[13], likely as part of a co-dependent circuit detecting both starvation cues and

increased oxygen[47].

The last major factor potentially influencing these behaviors is chemical signaling in the

form of environmental secretions and touch of other animals. Although the nature of these

chemical modulators is widely unknown, they represent an intriguing mechanism by which

individuals can modulate interactions with other animals and their environment. For example, it is

well known that several species of nematodes have complex and diverse secretomes, which they

use to modulate interactions with their environment. Several studies have shown that parasitic

nematodes secrete protein or chemical modulators to downregulate their host's immune response[48].

Similarly, the *elegans* genome encodes a large number of putative secreted proteins[49], some of

which were isolated via mass spectrometry, identifying a number of novel and nematode specific

protein factors of unknown function. In parallel, the chemical secretome of *elegans* revealed a

complex biochemical landscape, heavily enriched in putative pheromones, especially ascaosides[50].

Given the unknown function of many of these proteins and chemicals it is not unreasonable to

assume at least a subset may be influencing the behaviors of both individuals and populations of

individuals. Several ascaroside pheromones for example are well documented to alter population

behaviors, modulating hermaphrodite-male attractions and repulsion, mating, and dauer formation

in the absence of food[51]. Furthermore, pheromone secretion detected in the environment can also

induce the production of EV's in neuronal tissues of the affected animals, potentially altering

behavior[52]. Taken together we must consider that unknown signaling molecules are modulating

the interaction between individuals in our social feeding assays. Among the individuals of the same

species, these signals may provide a real-time readout of the local sensory environment helping

animals to respond and alter their behavior accordingly. Among individuals of different species,

the interactions are potentially even more complex. Differences in pheromone usage or chemical

secretome might produce novel interactions leading to interesting behavioral variations.

Overall, this work represents the first broad characterization of social feeding behaviors

across a large and evolutionarily ancient lineage of nematodes. Given the range of distinct

environmental conditions in which these species naturally reside, the variation we observed among

species across several behavioral paradigms in not altogether surprising. The current circuit

mechanism underlying the social feeding behaviors is still trying to fully reconcile how behavior

is tuned in response to various sensory cues like food identify/density[13,43,44], and O_2/CO_2 balance[53].

Complicating matters is recent evidence in *P. pacificus* showing that even within an established
circuit single cues such as elevation (and corresponding depletion of O_2) can override the
established behavioral paradigm and introduce unanticipated variation[14]. While the idea of
context-dependent variation is likely to complicate dissection of the genetic and molecular
mechanisms of behavioral variation, it also presents and exciting opportunity to study real-time
plasticity within neuronal circuits.

Supplementary Figure 1. Aggregation behavior is diverse across *Caenorhabditis* clade. (A) Representative images of additional species in the *Caenorhabditis* clade. **(B)** Multiple representative images of *nigoni* and *japonica* showing variation in aggregation and bordering behaviors.

Acknowledgements
The authors thank the labs of John I. Murray, Colin C. Conine, Chris Fang-Yen, David M. Raizen, and Meera V. Sundaram for their feedback on this project. We also thank Yun Ding, John I. Murray, and members of the Hart lab for comments on the manuscript. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). This work was supported in part by NIH 1R01MH096881 and NIH 1R35GM146782 (MPH), and the David and Gail Gasser fund (RFHE). CRLL was supported by NSF grant PRFB2305513.

Author Contributions
DH and MPH conceived and designed the study and experiments, and DH, JP, and RFHE conducted all experiments. DH processed, analyzed, and interpreted all data with help from CRLL. DH wrote the manuscript with assistance from MPH, and all authors reviewed, revised, and approved the manuscript.

Competing Interests
The authors declare no conflicts of interest.

References
1. Bagorda, A. & Parent, C. A. Eukaryotic Chemotaxis at a glance. *J. Cell Sci.* **121**, 2621–2624 (2008).
2. Archibald, J. M. Endosymbiosis and Eukaryotic Cell Evolution. *Curr. Biol. CB* **25**, R911-921 (2015).
3. Valencia-Montoya, W. A., Pierce, N. E. & Bellono, N. W. Evolution of Sensory Receptors. *Annu. Rev. Cell Dev. Biol.* **40**, 353–379 (2024).
4. Wan, K. Y. & Jékely, G. Origins of eukaryotic excitability. *Philos. Trans. R. Soc. B Biol. Sci.* **376**, 20190758.
5. Baluška, F. & Mancuso, S. Deep evolutionary origins of neurobiology: Turning the essence of 'neural' upside-down. *Commun. Integr. Biol.* **2**, 60–65 (2009).
6. Hartenstein, V. & Stollewerk, A. The Evolution of Early Neurogenesis. *Dev. Cell* **32**, 390–407 (2015).
7. Martins, E. P. *Phylogenies and the Comparative Method in Animal Behavior*. (Oxford University Press, 1996).
8. Ding, Y. *et al.* Neural Evolution of Context-Dependent Fly Song. *Curr. Biol.* **29**, 1089-1099.e7 (2019).
9. Hernández, D. G. *et al.* A framework for studying behavioral evolution by reconstructing ancestral repertoires. *eLife* **10**, e61806 (2021).
10. Cutter, A. D. Caenorhabditis evolution in the wild. *BioEssays* **37**, 983–995 (2015).
11. Schulenburg, H. & Félix, M.-A. The Natural Biotic Environment of Caenorhabditis elegans. *Genetics* **206**, 55–86 (2017).
12. de Bono, M. & Bargmann, C. I. Natural variation in a neuropeptide Y receptor homolog modifies social behavior and food response in C. elegans. *Cell* **94**, 679–689 (1998).
13. de Bono, M., Tobin, D. M., Davis, M. W., Avery, L. & Bargmann, C. I. Social feeding in Caenorhabditis elegans is induced by neurons that detect aversive stimuli. *Nature* **419**, 899–903 (2002).

14. Moreno, E., McGaughran, A., Rödelsperger, C., Zimmer, M. & Sommer, R. J. Oxygen-
induced social behaviours in Pristionchus pacificus have a distinct evolutionary history and
genetic regulation from Caenorhabditis elegans. *Proc. Biol. Sci.* **283**, 20152263 (2016).
15. Cowen, M. H. *et al.* Conserved autism-associated genes tune social feeding behavior in C.
elegans. *Nat. Commun.* **15**, 9301 (2024).
16. Macosko, E. Z. *et al.* A hub-and-spoke circuit drives pheromone attraction and social
behaviour in C. elegans. *Nature* **458**, 1171–1175 (2009).
17. Laurent, P. *et al.* Decoding a neural circuit controlling global animal state in C. elegans.
eLife **4**, e04241 (2015).
18. Cheung, B. H. H., Cohen, M., Rogers, C., Albayram, O. & de Bono, M. Experience-
dependent modulation of C. elegans behavior by ambient oxygen. *Curr. Biol. CB* **15**, 905–
917 (2005).
19. Bretscher, A. J., Busch, K. E. & de Bono, M. A carbon dioxide avoidance behavior is
integrated with responses to ambient oxygen and food in Caenorhabditis elegans. *Proc. Natl.*
Acad. Sci. U. S. A. **105**, 8044–8049 (2008).
20. Jang, H. *et al.* Dissection of neuronal gap junction circuits that regulate social behavior in
Caenorhabditis elegans. *Proc. Natl. Acad. Sci.* **114**, E1263–E1272 (2017).
21. Memar, N. *et al.* Twenty million years of evolution: The embryogenesis of four
Caenorhabditis species are indistinguishable despite extensive genome divergence. *Dev.*
Biol. **447**, 182–199 (2019).
22. Ponting, C. P. & Goodstadt, L. Separating derived from ancestral features of mouse and
human genomes. *Biochem. Soc. Trans.* **37**, 734–739 (2009).
23. Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).
24. Sternberg, P. W. *et al.* WormBase 2024: status and transitioning to Alliance infrastructure.
Genetics **227**, iyae050 (2024).
25. Dainat, J. *et al.* NBISweden/AGAT: AGAT-v1.4.1. Zenodo
https://doi.org/10.5281/zenodo.13799920 (2024).
26. Emms, D. M. & Kelly, S. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157
(2015).
27. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).
28. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using
DIAMOND. *Nat. Methods* **12**, 59–60 (2015).
29. Hart, M. P. & Hobert, O. Neurexin controls plasticity of a mature, sexually dimorphic
neuron. *Nature* **553**, 165–170 (2018).
30. Hart, M. P. Stress-Induced Neuron Remodeling Reveals Differential Interplay Between
Neurexin and Environmental Factors in *Caenorhabditis elegans*. *Genetics* **213**, 1415–1430
(2019).
31. Kiontke, K. *et al.* Caenorhabditis phylogeny predicts convergence of hermaphroditism and
extensive intron loss. *Proc. Natl. Acad. Sci.* **101**, 9003–9008 (2004).
32. Quach, K. T. & Chalasani, S. H. Intraguild predation between Pristionchus pacificus and
Caenorhabditis elegans: a complex interaction with the potential for aggressive behaviour. *J.*
Neurogenet. **34**, 404–419 (2020).
33. Félix, M.-A. *et al.* Evolution of Vulva Development in the Cephalobina (Nematoda). *Dev.*
Biol. **221**, 68–86 (2000).

34. Sharanya, D. *et al.* Genetic Control of Vulval Development in Caenorhabditis briggsae. *G3*
GenesGenomesGenetics **2**, 1625–1641 (2012).

35. Sommer, R. J. & Sternberg, P. W. Changes of induction and competence during the
evolution of vulva development in nematodes. *Science* **265**, 114–118 (1994).

36. Emmons, S. W. From cell fates to morphology: Developmental genetics of the
Caenorhabditis elegans male tail. *BioEssays* **14**, 309–316 (1992).

37. Fitch, D. H. A. Evolution of Male Tail Development in Rhabditid Nematodes Related to
Caenorhabditis Elegans. *Syst. Biol.* **46**, 145–179 (1997).

38. Slos, D., Sudhaus, W., Stevens, L., Bert, W. & Blaxter, M. Caenorhabditis monodelphis sp.
n.: defining the stem morphology and genomics of the genus Caenorhabditis. *BMC Zool.* **2**, 4
(2017).

39. Flemming, A. J., Shen, Z.-Z., Cunha, A., Emmons, S. W. & Leroi, A. M. Somatic
polyploidization and cellular proliferation drive body size evolution in nematodes. *Proc.*
Natl. Acad. Sci. **97**, 5285–5290 (2000).

40. Ortiz, C. O. *et al.* Searching for neuronal left/right asymmetry: genomewide analysis of
nematode receptor-type guanylyl cyclases. *Genetics* **173**, 131–149 (2006).

41. Toker, I. A. *et al.* Molecular patterns of evolutionary changes throughout the whole nervous
system of multiple nematode species. 2024.11.23.624988 Preprint at
https://doi.org/10.1101/2024.11.23.624988 (2024).

42. Moreno, E. *et al.* DAF-19/RFX controls ciliogenesis and influences oxygen-induced social
behaviors in Pristionchus pacificus. *Evol. Dev.* **20**, 233–243 (2018).

43. Abada, E. A. *et al.* C. elegans behavior of preference choice on bacterial food. *Mol. Cells* **28**,
209–213 (2009).

44. Brooks, K. K., Liang, B. & Watts, J. L. The influence of bacterial diet on fat storage in C.
elegans. *PloS One* **4**, e7545 (2009).

45. Meisel, J. D. & Kim, D. H. Behavioral avoidance of pathogenic bacteria by *Caenorhabditis*
elegans. *Trends Immunol.* **35**, 465–470 (2014).

46. Lei, M. *et al.* Loss of cilia in chemosensory neurons inhibits pathogen avoidance in
Caenorhabditis elegans. *Microbes Infect.* **26**, 105370 (2024).

47. Rogers, C., Persson, A., Cheung, B. & de Bono, M. Behavioral motifs and neural pathways
coordinating O2 responses and aggregation in C. elegans. *Curr. Biol. CB* **16**, 649–659
(2006).

48. Tritten, L., Ballesteros, C., Beech, R., Geary, T. G. & Moreno, Y. Mining nematode protein
secretomes to explain lifestyle and host specificity. *PLoS Negl. Trop. Dis.* **15**, e0009828
(2021).

49. Suh, J. & Hutter, H. A survey of putative secreted and transmembrane proteins encoded in
the C. elegans genome. *BMC Genomics* **13**, 333 (2012).

50. Zhang, X., Noguez, J. H., Zhou, Y. & Butcher, R. A. Analysis of Ascarosides from
Caenorhabditis elegans Using Mass Spectrometry and NMR Spectroscopy. in *Pheromone*
Signaling: Methods and Protocols (ed. Touhara, K.) 71–92 (Humana Press, Totowa, NJ,
2013). doi:10.1007/978-1-62703-619-1_6.

51. Edison, A. S. Caenorhabditis elegans Pheromones Regulate Multiple Complex Behaviors.
Curr. Opin. Neurobiol. **19**, 378–388 (2009).

52. Szczepańska, A. *et al.* Pheromone-based communication influences the production of
somatic extracellular vesicles in C. elegans. *Nat. Commun.* **15**, 2715 (2024).

53. Gray, J. M. *et al.* Oxygen sensation and social feeding mediated by a C. elegans guanylate
cyclase homologue. *Nature* **430**, 317–322 (2004).
642