Thienopyrimidine amide analogs target MmpL3 in *Mycobacterium tuberculosis*

Vanessa Pietrowski Baldin[a], Christopher L. Harding[a],   Diana Quach[b], Joseph Sugie[b], Joe

Pogliano[b], Tanya Parish[a,c]

[a]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle,

United States.

[b]Linnaeus Bioscience, Inc., San Diego, USA.

[c]Department of Pediatrics, University of Washington School of Medicine, Seattle, USA.

Abstract

Objectives: The identification of novel agents with mechanisms of action distinct from those

currently utilized in tuberculosis treatment remains a significant challenge. The mycobacterial

protein MmpL3 has emerged as a promising drug target due to its essential role in the synthesis

of the cell wall of *Mycobacterium tuberculosis*. We previously identified novel thienopyrimidine

amides with good anti-tubercular activity.

Methods: We profiled a subset of thienopyrimidine amides determining activity against

intracellular bacteria and bactericidal activity against replicating bacteria. We ran assays to

determine mode of action by measuring cell wall stress, ATP production, and bacterial

cytological profiling. We determined activity against a strain of M. tuberculosis with mutations in

MmpL3. We isolated and sequenced resistant mutants.

Results: We tested five analogs against a strain of *M. tuberculosis* with mutations in MmpL3 and

determined that they lost potency. Analogs induced P_{iniBAC}, a reporter for cell wall stress, and led

to an ATP boost characteristic of cell wall inhibitors. Bacterial cytological profiling of a

representative compound revealed a morphological profile consistent with other MmpL3

inhibitors.

Conclusions: Together, our data support MmpL3 as the most probable drug target for the TPA

analogs and add to the growing list of scaffolds that can inhibit this vulnerable transporter.

Introduction

Despite the availability of drug treatment regimens, tuberculosis remains a major global health concern, with an estimated 8.2 million new cases reported in 2023, including 400,000 cases of drug resistant tuberculosis (1). Currently, no vaccine provides effective protection against pulmonary TB. Additionally, the bacteria can enter a latent phase, where they display no symptoms, or cause symptoms that mimic other pathologies (2). Together, these factors highlight the urgent need for intensified efforts in TB control and the development of novel therapies, particularly those targeting drug-resistant bacilli and aiming to shorten the duration of treatment.

Mycobacterium tuberculosis is the causative agent of TB in humans. *M. tuberculosis* has a thick cell wall, which is composed by a highly hydrophobic bilayer of mycolic acids linked to arabinogalactan in turn linked to peptidoglycan. This structure forms a unique barrier against drugs and the host immune defenses (3–5). Hence, cell wall biosynthesis represents an attractive target for the development of new inhibitors.

MmpL3 (Rv0206c) is an essential transporter which is highly conserved across the *Mycobacterium* genus (4, 6, 7) and is the only member of the MmpL family that is essential in *M. tuberculosis* (2, 4, 7). Its essentiality is attributed to its role in the translocation of trehalose monomycolate (TMM) to the periplasmic space, where it serves as a precursor for the synthesis of trehalose dimycolate (TDM), a critical component of the cell envelope (4, 8). Given the important role of MmpL3 in constructing the mycobacterial cell wall, it has emerged as a high-value, druggable target, and at the moment, one of the most studied for anti-TB drug development (6). There are numerous MmpL3 inhibitors with diverse chemical structures which have potent antibacterial activity (5, 9).

We previously identified and characterized novel thienopyrimidine amide (TPA) analogs that

inhibit *M. tuberculosis* growth and had low cytotoxicity. From this series, we identified two

subsets. One subset (TPA-L) had increased activity against a LepB (signal peptidase)

hypomorph, indicating that the signal peptidase or protein secretion is the target or mode of

action (10, 11). The second subset of molecules (TPA-M) had excellent potency but were

equipotent against wild-type and hypomorph strains of *M. tuberculosis* suggesting that the target

was not protein secretion. In this paper we explored the biological profile of the TPA-M series of

molecules and identified MmpL3 as the most likely intracellular target.

Material and Methods

Bacterial strains and culture conditions

M. tuberculosis H37Rv-LP ATCC 25618 (wild-type) (12)   and   *M. tuberculosis* LP-0497754-

RM301 (MmpL3 F255L, V646M, F644I) (13) were cultured in Middlebrook 7H9 broth medium

supplemented with 10% oleic acid, albumin, dextrose, and catalase (OADC) enrichment

(BBL/Beckton and Dickson) and 0.05% w/v Tween 80 (7H9-Tw-OADC). Nutrient-starved cells

were generated by resuspending cells in phosphate-buffered saline (PBS) with 0.05% w/v

Tyloxapol (PBS-Tyl) at OD 1.0 and incubating for 7 days at 37°C without agitation. Avirulent *M.*

tuberculosis mc^{2}6206 (Δ*pan*CD Δ*leu*CD) was grown in 7H9-Tw-OADC supplemented with 0.5%

wt/vol glycerol, 0.2% wt/vol Casamino Acids, 48 µg/mL pantothenate, and 50 µg/mL leucine at

30°C.

Determination of minimum inhibitory concentration (MIC)

TPA molecules (**Figure 1**) were synthesized as previously described (10). Compound stocks

were resuspended in DMSO and stored at -20°C. MICs were determined in 96- or 384-well

microplates as described (14). Briefly, *M. tuberculosis* was grown to mid-log phase; cultures

were dispensed into plates containing test compounds to a final OD_{590} of 0.02 and incubated at

37°C for 5 days (H37Rv) or 7 days (mc^{2}6206). The OD_{590} was measured and IC_{90} determined

as the concentration at which 90% of growth was inhibited as compared to controls (DMSO

only).

Determination of activity against intracellular bacilli

THP-1 cells were cultivated in RPMI-1640 medium supplemented with 10% FBS and incubated

at 37°C, 5% CO_2. Cells were treated with 80 nM PMA for 24 h prior to infection, harvested using

Accumax™ solution, and resuspended in fresh cRPMI at a final density of 9×10^5 cells/mL. Cells

were infected overnight at a multiplicity of 1:1 with *M. tuberculosis* expressing LuxABCDE and

exposed to compounds in 96-well plates for 72 h at 37°C and 5% CO_2. Bacterial viability was

assessed by luminescence; IC_{50} was determined as the concentration at which 50% of growth

was inhibited as compared to controls (DMSO only).

Determination of bactericidal activity

87 *M. tuberculosis* H37Rv-LP was cultured in 7H9-Tw-OADC or starved for 7 days in PBS-Tyl,

88 adjusted to a theoretical OD_{590} of 0.02 and exposed to compounds in 96-well plates. Cultures

89 were spotted onto 7H10 agar supplemented with v/v 10% OADC (7H10-OADC) on day 0, 7 and

90 14. The MBC was defined as the lowest concentration at which no visible growth was observed.

91 Niclosamide was included as a control.

Induction of cell wall stress

M. tuberculosis strain carrying the P_{iniBAC}-lux reporter plasmid (15) was grown to mid-log in

GAST/Fe protein-free medium containing 0.3 g/L Bacto Casitone, 0.05 g/L ferric ammonium

citrate, 4 g/L dibasic anhydrous potassium phosphate, 2 g/L citric acid, 1 g/L L-alanine, 1.2 g/L

magnesium chloride, 0.6 g/L potassium sulfate, 2 g/L ammonium chloride, 1.8 mL of 10 M

sodium hydroxide, 10 mL glycerol, 5 mL 10% Tween 80, plus 15 µg/mL kanamycin. Cultures

were adjusted to a theoretical OD_{590} of 0.02 and used to inoculate 96-well plates containing

compounds. After 72 hours of incubation at 37°C, 1 vol of 10 mg/mL luciferin in of 1M HEPES

buffer pH 7.8, 1M DTT was added and RLU was read after 25 min of incubation at RT.

Ethambutol was included as a positive control.

Determination of ATP levels

103 ATP was measured using the BacTiter-Glo Assay kit (Promega) according to the manufacturer's

104 instructions. Log phase *M. tuberculosis* was exposed to compounds for 24 h, .50 µL BacTiter-

105 Glo™ was added, incubated at RT for 10 min and RLU read. Growth was measured after 120 h

106 by OD. Q203, was included as a positive control.

Isolation of resistant mutants

Log phase *M. tuberculosis* was plated on 7H10-OADC plates with 5X solid MIC and 10X solid

MIC. Plates were incubated at 37°C until isolated colonies appeared. Potential resistant mutants

were picked and streaked onto 5X MIC. MIC in liquid medium was measured to confirm

resistance. Genomic DNA from three isolates was extracted from cultured cells by heat

inactivation for 10 min at 100°C followed by 0.22 µm filtration. PCR amplification was performed

using 10 µL of extract DNA in a final volume of 50 µL containing 1 µL of Pfu polymerase

(Agilent), 5 µL of 10X Pfu amplification buffer (Agilent), 2.5 µL of primers MmpL3_seq_1_fwd

(5'-gattcgctacctgagcag-3') and MmpL3_seq_11_rev (0.5 µM) (5'-catttactgcagccgctg-3') and 4 µL

of 10 mM dNTP mix. PCR amplification was conducted, products were purified using the Qiagen

PCR purification kit, and sequencing was performed by Plasmidsaurus using Oxford Nanopore

Technology with custom analysis and annotation.

Bacterial Cytological Profiling (BCP)

M. tuberculosis mc^{2}6206 for bacterial cytological profiling was prepared as described (16). Briefly, cultures were adjusted to an OD_{600} of ~0.06-0.08 and incubated at 30°C for 18-20 hours before exposure to compounds at 1X and 5X MIC for 48h and 120 hours. Cells were fixed using a mixture of 100 µL of 16% paraformaldehyde, 3 µL of 8% glutaraldehyde, and 20 µL of 0.4 M phosphate buffer pH 7.5, washed twice with 200 µL of warm medium, concentrated to approximately 30 µL, and stained for 30 min. Full-field fluorescence microscopy images of the samples were preprocessed from their original proprietary microscope file format into a common image format (TIFF). The original image dimensions (3×2048×2048) were cropped by trimming 124 pixels from each edge to avoid optical artifacts, resulting in dimensions of 3×1800×1800. A 600-pixel square sliding window was passed over this image with a step size of 60 pixels to produce 400 sub-images, which were each passed to a trained convolutional neural network (CNN). Each full-field image, consisting of 400 sub-images, generated a vector that describes a point in latent space. Similarity scores were calculated by comparing the location of unknown compound-treated cells in this space to control compound-treated cells using a scaled version of the average minimum distance.

Results and Discussion

TPA analogs are active against intracellular mycobacteria

We previously demonstrated that the TPA series of compounds are active against aerobically-grown *M. tuberculosis* (10). A subset of analogs had potent anti-tubercular activity. Since these analogs had equipotent activity against wild-type and LepB hypomorph strains, we hypothesized that they did not target protein secretion. We wanted to determine the mode of action and target of these potent molecules.

142　We had already noted that compounds were active against replicating bacteria, so we expanded

143　our work to look at activity against intracellular bacteria and non-replicating bacteria. We

144　selected several active analogs (Figure 1). Molecules were tested for activity against

145　intracellular *M. tuberculosis* in THP-1 cells. All compounds were active against intracellular *M.*

146　*tuberculosis*, with activity similar to the MICs (Table 1).

148　TPA analogs are bactericidal against replicating bacilli

We assessed the bactericidal activity of the TPA analogs against *M. tuberculosis* wild type

following a 14-day exposure under replicating conditions as well as against non-replicating

bacteria generated by nutrient starvation. All analogs were bactericidal against replicating *M.*

tuberculosis with MBC/MIC ratio of <4. No bactericidal activity was seen against non-replicating

cells (MBC > 100 µM) (**Table 1**). These data are in contrast to what we previously noted with

inhibitors of LepB which showed bactericidal activity against non-replicating bacteria further

supporting a different target or mode of action (17).

156　TPA analogs induce cell wall stress in *M. tuberculosis*

We were interested in determining the mode of action of our molecules. As a first step, we

determined whether molecules induced cell wall stress. We used a reporter strain of M.

tuberculosis expressing luciferase under the control of P_{iniBAC}, since Induction of *iniBAC*

expression is a marker of cell envelope stress (18). We saw induction of PiniBAC for all

analogs, which predominantly occurred at concentrations close to the MIC. These data suggest

that TPA analogs cause cell wall stress (Figure 2 and Figure S1).

TPA analogs boost ATP in M. tuberculosis

A link between induction of *iniBAC* and a burst in ATP production has previously been proposed (19). To determine if our molecules also affect ATP levels, we measured intracellular ATP levels following treatment with TPA analogs. As anticipated, increasing compound concentrations led to a boost in ATP levels (Figure 3 and Figure S2). Again we saw the boost occurring at concentrations around the MIC. These data are consistent with the response seen with other MmpL3 inhibitors (3, 20).

MmpL3 mutation leads to resistance to TPA analogs

We hypothesized that our molecules could be targeting MmpL3, since this is a highly promiscuous target and cell wall stress and elevated intracellular ATP levels are features associated with MmpL3 inhibition in *M. tuberculosis* (3). We compared TPA activity against the wild-type *M. tuberculosis* and strain with mutations in MmpL3 which is resistant to a wide range of MmpL3 inhibitors (13, 20). The MmpL3 mutant strain demonstrated increased resistance to all analogs, with at least a 4-fold increase in IC_{90} (Table 1). Interestingly, this strain was highly resistant to three of the five analogs which lost all activity (IC_{90} >50-100 μM). Thus our data are consistent with MmpL3 being the drug target for this compound series.

Resistant strains have a non-synonymous mutation in *mmpL3*

In order to determine whether there might be additional targets, we isolated resistant mutants using TPN-0089300. Three strains were selected with an IC_{90} of >100 μM. Since MmpL3 mutations lead to resistance, we sequenced *mmpL3* in three resistant isolates (RM3, RM4, and RM6). All isolates had the same non-synonymous mutation in MmpL3 of F644L. This mutation has been associated with resistance to other MmpL3 inhibitors and is similar to one of the mutations in the MmpL3 mutant strain we used to test for resistance (F644I) (4, 8, 13, 21).

Cytological profiling

We determined the phenotypic effect of a representative TPA analog using bacterial cytological

profiling (16). We confirmed that TPN-0089300 was active against the avirulent *M. tuberculosis*

strain mc^{2}6206 (MIC of 1 μg/mL). Bacterial cells were exposed to TPN-0089300 for 48 and 120

hours at 2X and 5X MIC (2 μg/mL and 5 μg/mL respectively). Morphological profiling

demonstrated that bacterial morphology changed with cells becoming more rounded and losing

membrane integrity after 120h (Figure 4). Based on the similarity score, this profile was

identified as a strong match to other MmpL3 inhibitors (Figure 5) (16, 20).

Conclusions

We provide strong evidence that a subset of TPA analogs, which we now term TPA-M, target

MmpL3 in *M. tuberculosis*. TPA-M compounds were bactericidal for replicating bacilli, induced

cell walls stress and boosted ATP production. Mutations in MmpL3 conferred resistance and

bacterial cytological profiling identified MmpL3 as the most likely mode of action. We

demonstrated that the TPA-M analogs do not kill starved, non-replicating bacilli. This is

consistent with other MmpL3 inhibitors that have little effect against non-replicating *M.*

tuberculosis (3, 22). Cells treated with MmpL3 inhibitors develop dimples at the dividing pole,

where the protein localizes during active cell division and the lack of activity against non-

replicating cells has been attributed to the absence of MmpL3 localization at the pole in the non-

replicating state (7, 23). Taken together our data suggest that MmpL3 is the primary target of the

TPA-M series.

Acknowledgments

207 The authors kindly thank Grace Liu, Renee Allen, and Lauren Ames for technical assistance.

Funding Sources

This work was supported by the Department of Defense office of the Congressionally Directed Medical Research Programs under award number under award number PR180794 and by NIAID of the National Institutes of Health under award number R01AI182006. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest

The authors declare no competing interests.

References

1. World Health Organization. 2024. Global tuberculosis report 2024. https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2024

2. Diab A, Dickerson H, Al Musaimi O. 2025. Targeting the Heart of Mycobacterium: Advances in Anti-Tubercular Agents Disrupting Cell Wall Biosynthesis. Pharmaceuticals 18:70.

3. Ames L, Allen R, Boshoff HI, Cleghorn L, Engelhart C, Schnappinger D, Parish T. Common biological features of Mycobacterium tuberculosis MmpL3 inhibitors.

4. Kapp E, Calitz H, Streicher EM, Dippenaar A, Egieyeh S, Jordaan A, Warner DF, Joubert J, Malan SF, Sampson SL. 2023. Discovery and biological evaluation of an adamantyl-amide derivative with likely MmpL3 inhibitory activity. Tuberculosis 141:102350.

5. North EJ, Schwartz CP, Zgurskaya HI, Jackson M. 2023. Recent advances in mycobacterial membrane protein large 3 inhibitor drug design for mycobacterial infections. Expert Opin Drug Discov 18:707–724.

6. Adams O, Deme JC, Parker JL, Fowler PW, Lea SM, Newstead S. 2021. Cryo-EM structure

and resistance landscape of M. tuberculosis MmpL3: An emergent therapeutic target.

Structure 29:1182-1191.e4.

7. Williams JT, Abramovitch RB. 2023. Molecular Mechanisms of MmpL3 Function and

Inhibition. Microb Drug Resist 29:190–212.

8. Williams JT, Haiderer ER, Coulson GB, Conner KN, Ellsworth E, Chen C, Alvarez-Cabrera

N, Li W, Jackson M, Dick T, Abramovitch RB. Identification of New MmpL3 Inhibitors by

Untargeted and Targeted Mutant Screens Defines MmpL3 Domains with Differential

Resistance.

9. Chauhan M, Barot R, Yadav R, Joshi K, Mirza S, Chikhale R, Srivastava VK, Yadav MR,

Murumkar PR. 2024. The *Mycobacterium tuberculosis* Cell Wall: An Alluring Drug Target for

Developing Newer Anti- TB Drugs—A Perspective. Chem Biol Drug Des 104:e14612.

10. Lynde BE, Chemaly DM, Baldin VP, Greve E, Harding CL, Graner JM, Hardy M, Chowdhury

S, Parish T. 2025. Novel 3-Aminothieno[2,3- *b*]pyridine-2-carboxamides with Activity

against *Mycobacterium tuberculosis*. ACS Med Chem Lett 16:241–249.

11. Bonnett SA, Ollinger J, Chandrasekera S, Florio S, O'Malley T, Files M, Jee J-A, Ahn J,

Casey A, Ovechkina Y, Roberts D, Korkegian A, Parish T. 2016. A Target-Based Whole Cell

Screen Approach To Identify Potential Inhibitors of *Mycobacterium tuberculosis* Signal

Peptidase. ACS Infect Dis 2:893–902.

12. Ioerger TR, Feng Y, Ganesula K, Chen X, Dobos KM, Fortune S, Jacobs WR, Mizrahi V,

Parish T, Rubin E, Sassetti C, Sacchettini JC. 2010. Variation among genome sequences of

H37Rv strains of Mycobacterium tuberculosis from multiple laboratories. J Bacteriol

192:3645–3653.

13. McNeil MB, O'Malley T, Dennison D, Shelton CD, Sunde B, Parish T. 2020. Multiple

Mutations in Mycobacterium tuberculosis MmpL3 Increase Resistance to MmpL3 Inhibitors.

mSphere 5:e00985-20.

14. Ollinger J, Bailey MA, Moraski GC, Casey A, Florio S, Alling T, Miller MJ, Parish T. 2013. A

Dual Read-Out Assay to Evaluate the Potency of Compounds Active against

Mycobacterium tuberculosis. PLoS ONE 8:e60531.

15. Naran K, Moosa A, Barry CE, Boshoff HIM, Mizrahi V, Warner DF. 2016. Bioluminescent

Reporters for Rapid Mechanism of Action Assessment in Tuberculosis Drug Discovery.

Antimicrob Agents Chemother 60:6748–6757.

16. Quach D, Sharp M, Ahmed S, Ames L, Bhagwat A, Deshpande A, Parish T, Pogliano J,

Sugie J. 2025. Deep learning–driven bacterial cytological profiling to determine antimicrobial

mechanisms in *Mycobacterium tuberculosis*. Proc Natl Acad Sci 122:e2419813122.

17. Bonnett SA, Dennison D, Files M, Bajpai A, Parish T. 2018. A class of hydrazones are

active against non-replicating Mycobacterium tuberculosis. PloS One 13:e0198059.

18. Alland D, Steyn AJ, Weisbrod T, Aldrich K, Jacobs WR. 2000. Characterization of the

Mycobacterium tuberculosis iniBAC Promoter, a Promoter That Responds to Cell Wall

Biosynthesis Inhibition. J Bacteriol 182:1802–1811.

19. Shetty A, Dick T. 2018. Mycobacterial Cell Wall Synthesis Inhibitors Cause Lethal ATP

Burst. Front Microbiol 9:1898.

20. Allen R, Ames L, Baldin VP, Butts A, Henry KJ, Durst G, Quach D, Sugie J, Pogliano J,

Parish T. 2024. An arylsulfonamide that targets cell wall biosynthesis in *Mycobacterium*

tuberculosis. Antimicrob Agents Chemother 68:e01037-24.

21. Ioerger TR, O'Malley T, Liao R, Guinn KM, Hickey MJ, Mohaideen N, Murphy KC, Boshoff

HIM, Mizrahi V, Rubin EJ, Sassetti CM, Barry CE, Sherman DR, Parish T, Sacchettini JC.

2013. Identification of New Drug Targets and Resistance Mechanisms in Mycobacterium

tuberculosis. PLoS ONE 8:e75245.

22. Rao SPS, Lakshminarayana SB, Kondreddi RR, Herve M, Camacho LR, Bifani P, Kalapala

SK, Jiricek J, Ma NL, Tan BH, Ng SH, Nanjundappa M, Ravindran S, Seah PG, Thayalan P,

Lim SH, Lee BH, Goh A, Barnes WS, Chen Z, Gagaring K, Chatterjee AK, Pethe K, Kuhen

K, Walker J, Feng G, Babu S, Zhang L, Blasco F, Beer D, Weaver M, Dartois V, Glynne R,

Dick T, Smith PW, Diagana TT, Manjunatha UH. 2013. Indolcarboxamide Is a Preclinical

Candidate for Treating Multidrug-Resistant Tuberculosis. Sci Transl Med 5.

23. Lun S, Tasneen R, Chaira T, Stec J, Onajole OK, Yang TJ, Cooper CB, Mdluli K, Converse

PJ, Nuermberger EL, Raj VS, Kozikowski A, Bishai WR. 2019. Advancing the Therapeutic

Potential of Indoleamides for Tuberculosis. Antimicrob Agents Chemother 63:e00343-19.

	IC$_{90}$ (µM)	IC$_{90}$ (µM)			MBC (µM)	
Molecule ID	Intracellular	Wild-type	RM301	Fold-change	Replicating	Non-replicating
TPN-0102024	1.5 ± 0.21	1.5 ± 1.0	>100	>70	4.1 ± 1.8	>100
TPN-0099994	0.26 ± 0.03	0.7 ± 0.19	3.5 ± 2.4	6	2.3 ± 1.3	>100
TPN-0099934	2.7 ± 0.21	7.5 ± 3.3	>50	7	21 ± 7.2	>100
TPN-0099730	1.3 ± 0.47	1.2 ± 0.5	>100	>90	3.6 ± 2.4	>100
TPN-0089300	0.57 ± 0.33	1.2 ± 0.6	4.2 ± 0.6	4	3.1 ± 0	>100

291
292

Table 1. Activity of TPA analogs against *M. tuberculosis*.

Intracellular IC$_{90}$ = concentration that leads to 90% inhibition of bacterial replication in THP-1 macrophages. Data are the average and standard deviation of two independent replicates. IC$_{90}$ = concentration that leads to 90% inhibition of bacterial growth in aerobic culture. Data are the average and standard deviation of a minimum of 3 replicates. Fold-change = comparison of IC$_{90}$ for wild-type and RM301 (MmpL3 F255L, V646M, F644I) strains. MBC = minimum bactericidal concentration – concentration at which no viable bacteria were detected after 14 days. Data are the average and standard deviation of three replicates.

303
304
305
306 **Figure 1. Structure of analogs used in this study.** (1) TPN-0102024 (2) TPN-0099994 (3) TPN-

307 0099934 (4) TPN-0099730 (5) TPN-0089300

310 **Figure 2. Exposure to TPA analogs induces cell wall stress in *M. tuberculosis*.**

311 *M. tuberculosis* P$_{iniBAC}$-Lux was exposed to compounds for 72h and luminescence was read.
312 Data are representative of two independent experiments (see Fig S1).

Figure 3. TPA analogs boost ATP in *M. tuberculosis*.

M. tuberculosis was exposed to compounds for 24h and ATP was measured using BacTiter-Glo. Growth was measured by OD after 5 d. Data are representative of two independent experiments (see Fig S2). Q203 was used as a control.

Figure 4. Cytological profiling in response to TPN-0089300. *M. tuberculosis* mc²6206 was

treated for 48 and 120 h with TPN-0089300 or DMSO (control) and stained. Membranes are

shown in red, DNA in blue and green indicates loss of membrane integrity.

	ATP_PMF			CW1		CW2				DNA			FA_MA				MEMB		TRSC		TRSL			DMSO
	Bedaquiline	CCCP	Q203	Cycloserine	Meropenem	AU1235	Ethambutol	TBA-7371	SQ109	Gatifloxacin	Mitomycin_C	Novobiocin	Cerulenin	Ethionamide	Isoniazid	Pretomanid	Nisin	Thioridazine	Actinomycin_D	Rifabutin	Clarithromycin	Gentamicin	Sutezolid	DMSO
TPN-0089300 (2X MIC)	12	18	28	9	11	90	84	85	52	4	19	4	8	21	35	11	21	19	26	26	27	22	18	18
TPN-0089300 (5X MIC)	18	22	34	16	18	70	40	49	53	8	27	9	14	38	54	17	36	29	59	48	38	31	38	21

328 **Figure 5. Similarity scores for cytological profiling in response to TPN-0089300.**