Genetic variation in reproductive life-history traits is not correlated with estrogen-receptor positive breast
cancer risk

Euan A Young, E Yagmur Erten, Erik Postma, Virpi Lummaa, Bert van der Vegt, Geertruida H de Bock,
and Hannah L Dugdale

Author names, ORCID ID

Euan A. Young[1]*, 0000-0001-9370-9681

E Yagmur Erten[1], 0000-0001-9736-0405

Erik Postma[2], 0000-0003-0856-1294

Virpi Lummaa[3], 0000-0002-2128-7587

Bert van der Vegt[4], 0000-0002-2613-1506

Geertruida H de Bock[5], 0000-0003-3104-4471

Hannah L. Dugdale[1]*, 0000-0001-8769-0099

1 - Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, The
Netherlands

2 - Centre for Ecology and Conservation, University of Exeter, Penryn, United Kingdom

3 - Department of Biology, University of Turku, Turku, Finland

4 – Department of Pathology & Medical Biology, University of Groningen, University Medical Center
Groningen, Groningen, The Netherlands

5- Department of Epidemiology, University Medical Center Groningen, University of Groningen,
Groningen, The Netherlands

* Corresponding author emails: e.a.young@rug.nl, h.l.dugdale@rug.nl

ABSTRACT

Estrogen-receptor positive breast cancer (ER+ BC) is one of the most prevalent cancers, but the
evolutionary processes shaping genetic variation in ER+ BC risk are poorly understood. Both evolutionary
life-history theory and evidence studying individual ER+ BC risk variants suggest that increased genetic
ER+ BC risk is indicative of life-history trade-offs, and is associated with faster maturation, earlier
reproduction, and/or increased reproductive success, but how well this pattern is replicated across the full
polygenic architecture of these complex traits is unclear. Here, we estimate genome-wide genetic
correlations between ER+ BC risk and three reproductive life-history traits (age at menarche, age at first
birth, and the number of children born) for Lifelines biobank women using genomic restricted maximum
likelihood analyses. We complement these by performing linkage disequilibrium score regression on
European population and family-based genome-wide association study data. Regardless of the data or
method used, genetic correlations were low and not statistically different from zero. These results
therefore suggest that while some important loci underlying ER+ BC risk may demonstrate antagonistic
pleiotropy symptomatic of life-history trade-offs, the available genomic data suggests these findings are
not replicated, on average, across the genome. Consequently, our results suggest that genetic variation
does not overall shape the observed phenotypic relationships between these life-history traits and ER+ BC
risk. Future studies incorporating local-genetic correlations or gene-network data could further our
understanding how as to how genetic variation in these traits maps to phenotypic associations and aid our
understanding of the evolutionary processes shaping this important disease.

KEYWORDS: life-history trade-offs, genetic correlations, evolutionary medicine, quantitative genetics,
SNP heritability, antagonistic pleiotropy, GWAS data

INTRODUCTION

Breast cancer is the fourth most deadly cancer worldwide (GBD, 2021) and accounts for the greatest proportion of new cancer cases (Sung et al., 2021). Estrogen receptor-positive breast cancer (ER+ BC) is the most common subtype, accounting for ~80% of all breast cancer cases and is most common after age ~50 (Yasui & Potter, 1999). While lifestyle can influence an individual's risk of developing ER+ BC (Colditz et al., 2004), common germline genetic variation explains ~14% of the variation in a woman's risk (Zhang et al., 2020), and over 200 genome-wide significant risk variants have now been identified (Michailidou et al., 2013, 2017; Zhang et al., 2020). However, we have a poor understanding of why genetic variation promoting a life-threatening disease persists in contemporary human populations. Examining how genetic variation in ER+ BC risk is associated with human life-history traits would give insight into the evolutionary processes maintaining ER+ BC risk variants, the disease's overall genetic architecture, and help quantify the extent to which these risk variants will become more or less prevalent in future generations.

Evolutionary life-history theory and the antagonistic pleiotropy theory of aging hypothesize that genetic risk variants for diseases such as cancer may persist in populations due to their beneficial effects at younger ages, as a part of life-history trade-offs (Jacqueline et al., 2017; Promislow & Harvey, 1990; Roff, 1992; Stearns, 1977, 1992; Williams, 1957). For example, genetic variants promoting a faster life-history (earlier maturation and reproduction) could be selected for if they result in higher reproductive success, even if they increase the risk of cancer in later life (Figure 1, Dujon et al., 2022; Jacqueline et al., 2017). Indeed, although organisms have been under selective pressure to evolve cancer suppression mechanisms since the dawn of multicellularity over half a billion years ago (Merlo et al., 2006; Nunney, 2013), cancer remains common across the tree of life suggesting the presence of evolutionary constraints (Aktipis et al., 2013; Boddy et al., 2015), with a recent study also suggesting the presence of a reproduction-cancer suppression trade-off amongst bird species (Kapsetaki et al., 2024). However, whether such life-history trade-offs also shape genetic risk for ER+ BC is unclear.

In humans, the genetics of reproductive traits can play an important role in health and disease (reviewed in Benonisdottir et al., 2024), including genetic predispositions to earlier reproduction and higher numbers of children being associated with faster aging (Long & Zhang, 2023; Wang et al., 2013) and increased risk for specific diseases [(Byars et al., 2017; Trumble et al., 2023) reviewed in (Byars & Voskarides, 2020)]. Cancer appears to be no exception, with a host of tumor suppressor genes also being linked to the regulation of the timing of puberty and fertility (Boddy et al., 2015; Byars & Voskarides, 2020; Crespi & Summers, 2006; Dujon et al., 2022; Jacqueline et al., 2017). These include: *p53*, where genetic variants increasing cancer risk also appear to increase conception likelihoods (Kang et al., 2009); and more repeats

of the *CAG* repeat region of the androgen receptor being associated with both increased fertility and higher prostate cancer risk (Summers & Crespi, 2008). Breast cancer (and in particular ER+ BC) is a good candidate for antagonistically pleiotropic genes maintained by life-history trade-offs considering that ER+ breast cancer and these life-history traits share hormonal pathways underpinned by genes (Ghosh et al., 2020; Jacqueline et al., 2017; Jasienska et al., 2017). One genealogical study from Utah found that carriers of *BRCA1/2* mutations also had more children (Smith et al., 2012) suggesting that these trade-offs may have shaped the evolution of some of the most important breast cancer risk variants. More contemporary studies examining all genome-wide significant loci for age at menarche, found that genetic variants lowering age at menarche were also associated with higher overall breast cancer risk (Xiang et al., 2025) and, specifically, higher ER+ BC risk (Day et al., 2017; Zhao et al., 2024).

Yet, these associations between human life-history traits and ER+ BC risk expected from life-history theory (and evidenced in genetic studies) are not entirely consistent with the phenotypic associations from epidemiological studies. While earlier ages at menarche have been associated phenotypically with higher ER+ BC risk, earlier ages at first birth and higher numbers of children are actually associated with lower ER+ BC risk (Figure 1, Aktipis et al., 2015; Kelsey et al., 1993; Momenimovahed & Salehiniya, 2019). This may be due to these trends resulting in women spending less time in amenorrhea – which is induced by breastfeeding – and therefore experiencing more menstrual cycles and having a higher lifetime exposure to estrogen (Eaton et al., 1994; Short, 1976; Strassmann, 1999). Estrogen in turn has proliferative effects on mammary cells and thereby increases the risk of tumor development (Dall et al., 2018). ER+ BC risk being associated with fewer children also suggests that ER+ BC risk variants are – contrary to life-history theory and genetic evidence – under negative selection in contemporary populations. These contrasting findings are not mutually exclusive due to the importance of environmental factors on both ER+ BC risk and human life-history traits (such as lifestyle factors and socioeconomic and cultural backgrounds, Barban et al., 2016; Colditz et al., 2004) which may mask underlying genetic trade-offs. Moreover, previous genetic studies have largely focused on genome-wide significant variants which fail to capture the full scope of genetic variation in both ER+ BC risk and human life-history traits which are polygenic traits. Specifically, genome-wide significant variants explain less than 50% of the total variance in genomic risk for ER+ BC (Zhang et al., 2020), amount to only 25% of the total genetic variation in age at menarche (Day et al., 2017), with even fewer genome-wide significant variants for age at first birth and number of children born being detected despite significant heritability for both traits (Barban et al., 2016; Mills et al., 2021; Tropf et al., 2015).

To help understand these inconsistent results, we need to estimate genetic correlations using whole-genome data. Genetic correlations estimated from genomic data reflect the proportion of genetic variance

shared by traits and are a measure of the overall contribution of pleiotropy to the covariation of these traits (van Rheenen et al., 2019). We can therefore estimate genetic correlations between ER+ BC risk and human life-history traits to quantify to what extent life-history trade-offs shape ER+ BC risk on average across all loci, whether these trade-offs contribute genetically to the observed phenotypic trends, or whether observed phenotypic associations reflect underlying genetic factors (Figure 1). To the best of our knowledge, studies estimating genome-wide genetic correlations between ER+ BC risk and human life-history traits are limited to two studies estimating genetic correlations between ER+ BC risk and age at menarche (Jiang et al., 2019; Prince et al., 2024), one of which also estimated the genetic correlations between other life-history traits, including age at first birth and number of children (Prince et al., 2024). These studies revealed negligible genetic correlations between all life-history traits and ER+ BC risk, except for a small (r_g=-0.048) and statistically non-significant (p = 0.062) genetic correlation between number of children and ER+ BC risk. This suggests that these phenotypic trends do not translate into evolutionary change (Haldane, 1954; Lande & Arnold, 1983; Robertson, 1966). However, these studies did not fully account for confounding factors such as population stratification, which can bias genetic correlations (Berg et al., 2019; Border et al., 2022), particularly those involving human life-history traits (Howe et al., 2022; Tan et al., 2024). Thus, it is still poorly understood how the frequency of genetic variants underlying an important disease will change in future generations, and what adaptive processes shape the prevalence of these genetic variants.

To estimate genetic correlations between human reproductive life-history traits and ER+ BC risk we first perform a genomic restricted maximum likelihood (GREML) analysis (Lee et al., 2012; Yang et al., 2010) using the Dutch biobank Lifelines (Scholtens et al., 2015) linked to national pathology records (Casparie et al., 2007). We then repeat our analyses by estimating genetic correlations using linkage disequilibrium (LD) score regressions of the largest available GWAS data (Bulik-Sullivan, Finucane, et al., 2015). We first use family-based GWAS data, which exploits the random Mendelian segregation among siblings to remove environmental confounding from genetic correlations (Kong et al., 2018; Young et al., 2022; Young, 2024). We then repeat our LD score regression analysis using population GWAS data as family-based GWAS data can suffer from reduced power and can underestimate genetic effects in the presence of assortative mating (Kemper et al., 2021).

Figure 1: Predicted direction of genetic correlations between ER+ BC risk and reproductive life-history traits (age at menarche, age at first birth, and number of children) under different hypotheses. Circles below the x-axis indicate negative genetic correlations, i.e. genetic predispositions for lower/earlier values for the life-history trait are correlated with increased ER+ BC risk. Circles above the x-axis indicate positive genetic correlations, i.e. genetic predispositions for higher/later values for the life-history trait are correlated with increased ER+ BC risk. Circles on the x axis line show no correlation. Circle colors indicate the expected direction of genetic correlations under different hypotheses: *genetic life-history trade-offs* as predicted by antagonistic pleiotropy theory of ageing and life-history theory, genetic predispositions to ER+ BC are indicative of a reproduction-survival trade-off and/or faster life history; and red, *genetic correlations match established phenotypic correlations (e.g. due to estrogen exposure)*, genetic correlations match the existing evidence from phenotypic studies of how life-history traits shape ER+ BC risk phenotypically, potentially due to their influence on a woman's lifetime estrogen exposure.

RESULTS

GREML analyses of Lifelines data

We first performed GREML analyses on data from the Lifelines biobank ($n = 97,131$ women) using the

Global screening array genomic data ($n = 535,107$ SNPs). Univariate analyses revealed statistically

significant heritability (±SE) for age at menarche (43.1±5.8%, n = 8,071 women) and age at first birth

(24.2±7.4%, n = 6,260 women), suggesting that genetic variance plays a significant role in shaping these

traits (p < 0.05), but the heritability of the number of children was estimated with less precision and was

not statistically significant (12.8±9.8%, p = 0.337, n = 4,602 women, Table 1a). This was also true for

ER+ BC risk, where the heritability was estimated as 11.8% (±5.2%, p = 0.190, n = 182 cases and n =

8,700 controls, Table 1a). These heritability estimates were largely replicated using other genomic data

available on Lifelines (the Affymetrix, n = 441,001 SNPs, and CytoSNP genomic array data, n = 244,041

SNPs, Table S1), but the heritability estimate for ER+ BC risk became statistically significant when using

the Affymetrix array data (p = 0.04, Table S1), probably because these contained a slightly higher number

of ER+ BC cases (n = 198 cases, n = 8,972 controls). Moreover, the heritability estimate for age at

menarche was considerably lower (but remained statistically significant) when using the CytoSNP data

(17.0% ±5.2%, p < 0.001, n = 6,155 women, Table S1), most likely because of the lower genomic

coverage of the CytoSNP data (Table S2).

Table 1: Heritability estimates (and standard error, SE) of traits from **a)** GREML analyses using Lifelines

genotype data and **b)** LD score regression on GWAS data. P-values from LRT tests (for GREML) and z-

scores (for LD score regressions) and results are presented in bold when significant (< 0.05). Results from

Lifelines Global screening array data are shown. Results from analyses of other Lifelines genotype data

are shown in Table S1.

| Trait | a) GREML on Lifelines data | | b) LD score regression on family-based GWAS data | |
	Heritability±SE	P-value	Heritability±SE	P-value
ER+ BC risk	11.8±5.2%	0.190	**10.7±0.9%**	**< 0.001**
Age at menarche	**43.1±5.8%**	**< 0.001**	14.1±5.9%	**0.008**
Age at first reproduction	**24.2±7.4%**	**< 0.001**	1.6±1.9%	0.202
Number of children	12.8±9.8%	0.337	**8.6±1.7%**	**< 0.001**

Genetic correlations (±SE) between ER+ BC risk and life-history traits were all low and statistically non-

significant in the Lifelines data (p > 0.01$\dot{6}$). Figure 2a, Table S3). For the Global screening array data

these were 0.05±0.17 for age at menarche (n = 176 cases, n = 7,895 controls), 0.10±0.24 for age at first

birth (n = 155 cases, n = 6,105 controls), and 0.10±0.35 for number of children (n = 146 cases, n = 4,456

controls, p > 0.016, Fig 2a, Table S2). Precision of these estimates was also low, but a power analysis

estimated that we had 80% power to detect genetic correlations between ER+BC risk and the life-history

trait stronger than ±0.29 for age at menarche, ±0.40 for age at first birth, and ±0.52 for the number of

children (Figure S1). These null findings were replicated when using both the Affymetrix and CytoSNP

data (Table S3), with slightly reduced power in the CytoSNP data (Figure S1), which was expected given

its lower sample size.

Figure 2: Genetic correlations between ER+ BC risk and life-history traits from: (**a**) a genomic relatedness matrix (GREML) analyses using Lifelines biobank data and LD score regression analyses using (**b**) family-based GWAS data and (**c**) population GWAS data. Circles above the dotted line indicate positive genetic correlations and circles below indicate negative genetic correlations. Error bars show 95% confidence intervals approximated from SEs. P-values testing a non-zero genetic correlation are shown above each trait.

To help us understand whether our results from the GREML analysis of Lifelines biobank data were

limited by power or confounded by gene-environment correlations, we repeated our analysis using LD

score regression using family-based GWAS data. For this analysis, we used European GWAS data

from the largest available study on ER+ BC (sample sizes: $n = 11,792,542$ SNPs, $n = 69,501$ cases,

and $n = 95,042$ controls, Michailidou et al., 2017) and the most recent family-based GWAS data study

including life-history traits ($n = 5,671,230$-$6,551,047$ SNPs, Tan et al., 2024). Albeit estimated with

more precision, our heritability estimates (±SE) using direct genetic effects from family-based GWAS

analyses were largely consistent with our results from the Lifelines biobank for ER+ BC risk

($10.7 \pm 0.9\%$, $n = 1,129,448$ SNPs) and the number of children ($8.6\% \pm 1.7\%$, $n = 670,251$ SNPs),

further suggesting that genetic variation plays a small but non-zero role in shaping among individual

variation in these traits (Table 1). However, contrary to results from the Lifelines biobank, the

heritability of age at menarche was lower ($14.1 \pm 5.9\%$, versus $43.1 \pm 5.8\%$ using the Global screening

array data, $n = 580,219$ SNPs), although it remained statistically significant, whereas age at first birth

was not shaped by genetic variation when using GWAS data ($1.6\% \pm 1.9$, $n = 708,027$ SNPs, Table 1).

Overall, this suggests that the estimates of the heritability of age at first birth using GREML analyses

may have been inflated through factors such as unaccounted population stratification or the presence

of assortative mating. A supplementary analysis of ER+ BC risk GWAS data also suggested little

evidence of bias due to environmental confounding, including factors such as population stratification

(Supplementary note 1, Table S5).

Using family-based GWAS summary statistics repeated the findings of no genetic correlations

between life-history traits and ER+ BC risk, though our estimates were smaller and more precise

(Figure 2b, Table S4, $p > 0.01\dot{6}$). Specifically, we estimated the genetic correlations (±SE) between

ER+ BC risk and our life-history traits as: age at menarche, 0.079 (± 0.15, $p = 0.602$, $n = 551,452$

SNPs); age at first birth, -0.052 (± 0.178, $p = 0.764$, $n = 668,389$ SNPs); and the number of children

ever born, -0.010 (± 0.037, $p = 0.795$, $n = 635,154$ SNPs). Taking the 95% confidence intervals, these

estimates allowed us to rule out genetic correlations between life-history traits and ER+ BC risk >0.17

and <-0.39 for age at menarche, >0.37 and <-0.22 for age at first birth above, and >0.06 or <-0.08 for

the number of children.

Finally, we estimated genetic correlations using the population GWAS summary statistics for life-

history traits uncorrected for population stratification from Tan et al. (2024) as family-based GWAS

data can suffer from lower power and may underestimate genetic effects in the presence of assortative

mating (Kemper et al., 2021). Using population GWAS data, we again repeated the findings of no

statistically significant genetic correlations ($p > 0.01\dot{6}$), but estimates were generally estimated with

more precision and, in some cases, varied quantitatively (Figure 2c, Table S6). Specifically, the genetic

correlation (±SE) between the life-history traits and ER+ BC risk was estimated as: for age at
menarche, -0.015 (±0.036, p = 0.67, n = 886,799 SNPs), for age at first birth, 0.054 (±0.054, p = 0.32,
n = 922,056 SNPs), and for number of children, -0.0063 (±0.066, p = 0.92, n = 952,783 SNPs, Table
S6).

DISCUSSION

We estimated genetic correlations between three important human reproductive life-history traits (age
at menarche, age at first birth, and number of children) and ER+ BC risk to help understand the extent
to which life-history trade-offs shape genetic predispositions to ER+ BC. Our study builds on research
showing evidence for antagonistic pleiotropy between ER+ BC risk and life-history traits at important
loci underlying these traits (Day et al., 2017; Smith et al., 2012; Xiang et al., 2025; Zhao et al., 2024)
to include the full contribution of pleiotropy to the covariance between these traits. Across all three
life-history traits and all analyses we find no significant genetic correlations with ER+ BC risk. These
results suggest that overall the importance of reproductive life-history trade-offs in shaping genetic
ER+ BC risk is low and that genetic variation does not contribute overall to the observed phenotypic
trends from epidemiological studies (Aktipis et al., 2015; Kelsey et al., 1993; Momenimovahed &
Salehiniya, 2019). These results are in line with previous non-significant estimates from two studies
(Jiang et al., 2019; Prince et al., 2024), with our analyses suggesting that these patterns hold for
contemporary European ancestry populations regardless of the data or approach used, even when
accounting for environmental confounding using family-based GWAS data.

Our study adds to previous research where studies focusing only on genome-wide significant loci
show evidence for genetic life-history trade-offs shaping (ER+) BC risk (Day et al., 2017; Smith et al.,
2012; Xiang et al., 2025; Zhao et al., 2024), whereas studies focusing on whole-genome genetic
correlations find no evidence (Jiang et al., 2019; Prince et al., 2024). These results are not mutually
exclusive, and our results do not refute the importance of life-history trade-offs being important in
shaping individual ER+ BC risk variants. Especially in the case of high-prevalence loci of strong
effect such as BRCA1/2 mutations (Smith et al., 2012) which are believed to be under strong negative
selection in the absence of such pleiotropy (Pavard & Metcalf, 2007). However, because genetic
correlations measure the average contribution of pleiotropy to the covariance between two traits (van
Rheenen et al., 2019), together these results suggest that loci showing strong antagonistic pleiotropy
are not repeatedly found across all loci underlying these complex, highly polygenic traits. Thus, rather
than supporting the antagonistic pleiotropy theory of ageing, our results suggest that, in general,
genetic ER+ BC risk variants may have accumulated due to weak selection at older ages, providing
indirect support for the mutation accumulation theory of ageing (Charlesworth, 1980, 2001; Haldane,
1941; Medawar, 1952). Antagonistic pleiotropy between diseases acting in early- versus late-life can

also maintain genetic variation in disease risk, also in line with the antagonistic pleiotropy disease of
ageing (Rodríguez et al., 2017). Thus, while our results suggest a minor role for reproductive life-
history trade-offs in maintaining genetic variation in ER+ BC risk, studies estimating genetic
correlations between ER+ BC risk and earlier onset diseases could help resolve the relative importance
of the antagonistic pleiotropy or mutation accumulation theories of ageing for shaping genetic
variation in this disease.

It is unclear whether these low and non-significant genetic correlations are due to: (1) little overall
overlap between the genetic architecture of ER+ BC and these human life-history traits; and/or (2)
variation in the direction of the associations across individual loci important for these sets of traits
(van Rheenen et al., 2019). The first explanation is viable considering the polygenicity of these traits
and was previously suggested as an explanation by Jiang et al. (2019). However, the current methods
and data available cannot rule out the role of important breast cancer risk genes showing either
agonistic or antagonistic, pleiotropy with life-history traits. With increasing power to detect genome-
wide significant variants, future studies may be able to help assess the relative numbers of
antagonistically versus agonistically pleiotropic variants and the degree of shared genetic architecture
between ER+ BC risk and life-history traits.

Our results also provide no evidence for strong, consistent directional selection across ER+ BC risk
loci, suggesting that these variants will not significantly increase or decrease in prevalence in future
generations due to selection. This is despite demonstrating that contemporary European populations
retain the ability to evolve through natural selection (Corbett et al., 2018; Fisher, 1930; Haldane,
1954). In fact, our heritability estimate of ~8.6% for the number of children is higher than the average
estimates of heritability of fitness among wild bird and mammal species (3%, Bonnet et al., 2022).
Thus, while previous phenotypic trends would suggest that ER+ BC risk variants are under negative
selection (Aktipis et al., 2015), our results suggest that these phenotypic trends do not translate to the
genetic level (Haldane, 1954; Lande & Arnold, 1983; Robertson, 1966). Again, this does not rule out
selection on individual loci, and this picture may change with additional, higher-power, GWAS
studies. However, for traits such as the number of children, where the number of loci of large effect are
relatively few (Barban et al., 2016), studies examining local genetic correlations, for example using
ρHESS (Shi et al., 2017), would be best placed to detect finer-scale variation in the direction and
strength of selection on genomic ER+ BC risk and assess the importance of selection and
reproduction-cancer suppression trade-offs in shaping genetic ER+ BC risk. Additionally, recent
research suggests that genetic variation in breast cancer is enriched in gene-expression hubs involved
in estrogen metabolism and signaling (Stone et al., 2025). Given that these hubs are also likely to be
important in regulating these life-history traits (Jacqueline et al., 2017; Jasienska et al., 2017),
incorporating gene-networks and gene-expression data could further our understanding of how

variation at the genetic level could map to phenotypic trends, with studies also showing that
transcriptomic correlations are generally higher than genetic correlations (Gerring et al., 2022).

No (or at most very weak) genetic correlations between life-history traits and ER+ BC risk also
suggest that the observed phenotypic trends from epidemiological studies are largely due to non-
genetic factors (Aktipis et al., 2015; Kelsey et al., 1993; Momenimovahed & Salehiniya, 2019). This
provides indirect support for the hypothesis that this is due to an evolutionary mismatch due to the
rapid non-genetic changes in these traits over recent generations (Coe & Steadman, 1995; Eaton et al.,
1994; Short, 1976; Strassmann, 1999). Changes in diet and nutrition have caused earlier ages at
menarche (Duan et al., 2020; Nguyen et al., 2020; Rees, 1993) and recent changes in cultural
preferences towards smaller families have driven later ages at first birth and fewer children (Aitken,
2022; Kirk, 1996). These changes may have led to higher lifetime exposure than a woman's cancer
suppression mechanisms have evolved to handle (Coe & Steadman, 1995; Eaton et al., 1994; Short,
1976; Strassmann, 1999). In line with this hypothesis, differences in birth rates among countries
predict breast cancer prevalence (You et al., 2018), with our findings of low heritability of these traits
and low or no genetic correlations further suggesting that these phenotypic associations are
environmental in origin. However, the extent to which changes in these reproductive life-history traits
have also driven the increased prevalence in ER+ BC risk is unclear, which longitudinal studies would
be best placed to answer.

Our study suffers from a few key limitations. First, although we found little evidence in support of
life-history trade-offs for ER+ BC risk, exploring genetic correlations between estrogen receptor-
negative breast cancer (ER- BC) risk and life-history traits may find different results. While ER- BC
risk is not as strongly linked to reproductive life-history traits as ER+ BC risk at the phenotypic level
(Aktipis et al., 2015), ER- BC risk does show higher heritability than ER+ BC (Bauer et al., 2007), and
there is some evidence that ER- BC risk is genetically correlated with earlier ages of reproduction
(Prince et al., 2024), indicative of a fast life-history, as previously hypothesized (Hidaka & Boddy,
2016). Second, we could not account for age at menopause in our analysis due to insufficient sample
sizes. Age at menopause is an important life-history trait and late ages at menopause are a risk factor
for ER+ breast cancer (Kelsey & Bernstein, 1996) with previous studies showing some evidence for
genetic variants underlying age at menopause also influencing cancer risk (Benonisdottir et al., 2024).
Moreover, our study focused on reproductive life-history trade-offs, but body size – due to its impact
on an organism's number of cells – is another important life-history trait affecting overall cancer
susceptibility, including potentially ER+ BC risk (Aktipis et al., 2013; Nunney, 2018, 2024). Height
has been identified as a potential risk factor for breast cancer (Green et al., 2011), although current
evidence of this being due to genetic factors is lacking (Jiang et al., 2019), but warrants further study.
Finally, despite using both individual-level biobank data and the largest available GWAS data for life-
history traits and ER+ BC risk, some of our genetic correlations were still somewhat imprecise. These

power limitations were mostly constrained to the GREML analyses of Lifelines biobank data, where
some traits in some analyses did not show significant heritability (e.g., ER+ BC risk and number of
children when using the Global screening array and CytoSNP array data). However, even when some
of these traits showed higher and statistically significant heritability (e.g., using the Affymetrix data),
the finding of non-significant genetic correlations was replicated, although we could only rule out
genetic correlations larger than ±0.29-0.52 for the Global screening array data. Estimating genetic
correlations with GREML is computationally demanding but further studies could aim to conduct
similar analyses using larger biobanks. Nevertheless, our analyses of GWAS data were less power-
limited and more precise. Our study therefore provides a robust assessment of the strength of genetic
correlations between ER+ BC risk and these human life-history traits, given the current GWAS data
and methods available.

These findings provide an important step towards a more complete understanding of the genetic and
evolutionary processes shaping ER+ BC risk, which is a deadly form of cancer with increasing
prevalence. Our results suggest that genetic predispositions towards ER+ BC risk are not overall
shaped by life-history trade-offs in European ancestry populations but whether this is also true across
other geographic regions is unknown. Studies of populations with higher birth rates, increasing the
chance of detecting reproduction-survival trade-offs, have previously been used to identify
antagonistically pleiotropic genes underlying disease and fertility (e.g., Trumble et al., 2023), could be
particularly fruitful. Moreover, although our results suggest that the prevalence of ER+ BC risk
variants in future generations is unlikely to change due to natural selection, we highlight the
importance of examining the gene-by-environment interactions in the context of ER+ BC risk, which
has so far received limited attention (but see Middha et al. 2023). This is an important avenue of
research, with the broad range of non-genetic ER+ BC risk factors and the number of existing variants
giving a high potential of gene-by-environment interactions. Finally, we hope that our multifaceted
approach can provide a template for future studies using the latest developments in genetic analyses to
test evolutionary hypotheses.

METHODS

Lifelines biobank

Lifelines is a multi-disciplinary prospective population-based cohort study examining, in a unique
three-generation design, the health and health-related behaviors of 167,729 persons living in the North
of the Netherlands. It employs a broad range of investigative procedures in assessing the biomedical,
socio-demographic, behavioral, physical and psychological factors which contribute to the health and
disease of the general population, with a special focus on multi-morbidity and complex genetics. We

focused on data from 97,131 women born 1919-1979 and recruited 2007-2013 (https://data-
catalogue.lifelines.nl/, accessed 08-05-2024, *Lifelines*, 2024).

Life-history traits

Age at menarche describes the age in which a female first started menstruating (Rees, 1993), can vary
among individuals due to both environmental (e.g., nutritional status) and genetic effects, and is linked
to future disease risk, including breast cancer (Parent et al., 2003; Perry et al., 2015). We quantified
age at menarche using the lifelines baseline question: "How old were you when you had your first
menstruation?" (N=85,210 women). For quality control, we removed 153 values >3 standard
deviations from the mean. This gave a mean age at menarche of 13.1 (SD = 1.7, range = 8-18, N =
85,057, Figure S2A), representative of the Netherlands (Parent et al., 2003).

Age at first birth and number of children born are likewise influenced by both genetic factors and
environmental factors which are then further constrained by the social environment, cultural
background, and any remaining individual preferences (Barban et al., 2016; Mills et al., 2021). We
quantified age at first birth and the number of children ever born using questions for a participant's
birth date and the years of birth of her children focusing on full-term pregnancies. For these variables,
we used data from baseline questionnaires, supplemented with follow-up questions during wave two
of the Lifelines data collection (2014-2018). For age at first birth, we screened the questionnaire data
removing any years of birth of children before an individual was born and after an individual had
answered the questionnaire (n = 16). We used the birth dates of participants to calculate ages at first
birth (mean = 26.9, 95% CIs = 18.9-36.3, Figure S2B) which occurred 1942-2017 (N = 64,779). The
number of children ever born was then calculated only for women who were asked for the years of
birth of their children after age 45 to avoid biasing the number of childless women and the total
number of children born. This gave a mean number of children born of 2.0 (range 0-8, N = 51,672,
Figure S2C). All data were handled in R 4.2.2 (2022-10-31, R Core Team, 2020) using *dplyr* 1.1.4
(Wickham et al., 2019).

ER+ BC diagnoses

ER+ BC diagnoses were determined by linking Lifelines participants to the Dutch Nationwide
Pathology Databank (Palga) which contains the national records of histo- and cytopathology, with
complete coverage of the Netherlands since 1991 (www.palga.nl, Casparie et al., 2007). The Palga
databank details age and dates of diagnoses, and pathology reports describing the subtype, nature, and
grade of the tumor (Casparie et al., 2007). Linkage of Lifelines participants to the Palga databank was
successfully done for all, except 244 Lifelines participants, producing 10,114 unique pathology reports
on breast tumors across 4,310 individuals.

We subtyped these records according to estrogen receptor status (ER-status) using the report

conclusions written by pathologists contained in the Palga excerpts (see Supplementary note 2 for

further details). According to national Dutch guidelines, ER-status is considered positive when >=

10% of tumor cells show are ER+ (IKNL, 2012). We focused on women whose first cancer diagnosis

was breast cancer to remove possible metastatic tumors of different origin (n = 628). We also excluded

15 male diagnoses. This left 3,667 Lifelines participants with their first-cancer diagnosis being breast

cancer, diagnosed from 1981-2023, and 2,177 ER+ diagnoses (Figure S3), 415 ER- diagnoses, with

1,090 being unable to be assigned an ER-status. The ER+ BC diagnoses had a median age of diagnosis

of 51 (range = 24-93),

Genotype data

Genotype data on European-Ancestry Lifelines participants were available on three different

platforms: the Illumina CytoSNP-12v2 array (hence CytoSNP array data), the Infinium Global

Screening Array MultiEthnic Disease Version 1.0 (hence Global screening array data), and the

Affymetrix FinnGen Thermo Fisher Axiom Custom array (hence Affymetrix array data). Full details

about the quality control of each genomic dataset and imputation can be found online at:

https://wiki.lifelines.nl/. In brief, single nucleotide polymorphisms (SNPs) were filtered according to

minimum allele frequency, Hardy-Weinberg equilibrium tests, Plink call rates, and ancestry outliers.

Imputation was also performed using Minimac 2012.10.3 (Howie et al., 2012) with *The Genome of*

The Netherlands release 5 (The Genome of the Netherlands Consortium, 2014) and the 1000 Genomes

phase1 v3 reference panels (The 1000 Genomes Project Consortium, 2010).

We aimed to combine the different Lifelines genotypes in one GREML analysis and therefore

harmonized these three datasets using Genotype Harmonizer 1.4.25 (available at

https://github.com/molgenis/systemsgenetics/wiki/Genotype-Harmonizer). This software uses linkage

disequilibrium patterns to address strand issues, aligning genotype data to one reference dataset, for

which we used the Global screening array. The harmonized data contained 1,032,001 variants across

80,045 individuals.

GREML analyses of Lifelines data

We performed GREML analyses (using GCTA/1.93.2beta-GCCcore-7.3.0, Yang et al., 2011) which

leverages the genomic relatedness between distantly related individuals in a population to examine

how genetic variation associated with one trait covaries with another (Yang et al., 2010). We first

performed a univariate GREML on ER+ BC risk, age at menarche, age at first birth, and number

children of ever born to estimate the heritability (h^2) of each trait, which is the proportion of total

phenotypic variance in a trait (σ_P^2) due to genetic factors (σ_G^2):

$$h^2 = \frac{\sigma_G^2}{\sigma_P^2}$$

435   With GREML, the heritability of a given phenotype (**y**) is estimated using a linear mixed model
436   conditioned upon a vector of fixed effects (β) (such as age and sex) and where the variation explained
437   by the genomic relatedness matrix (g) is divided by the residual variance (ε) to estimate the
438   heritability in the equation:

$$y = X\beta + g + \varepsilon$$

439   with parameters estimated using the restricted maximum likelihood approach (see Yang et al., 2010,
440   for comprehensive methods).

441   We then estimated the genetic correlation ($r(G)$) between ER+ BC risk and each life-history trait in
442   three bivariate models (Lee et al., 2012). $r(G)$ is estimated as the genetic covariance between traits
443    $Cov(G_{t1,t2})$ scaled by the genetic variance in trait:

$$r\left(G_{t1,t2}\right) = \frac{Cov\left(G_{t1,t2}\right)}{\sqrt{V_{Gt1}} * \sqrt{V_{Gt2}}}$$

We created the genomic relatedness matrices (GRMs) for all individuals for which we had genomic
data using the *--make-grm* argument in the *gcta64* function. We created GRMs for each chromosome
first, then merged them into one GRM using *--mgrm*. Finally, the merged GRM was pruned of
combinations of closely related individuals while maximizing the sample size using *--grm-cutoff* at a
relatedness threshold of 0.05 (*n*-women = 25,068) to remove bias in genetic variances due to gene-
environment correlations.

The age of the individual when they were either first diagnosed with breast cancer using the Palga
databank, or (for individuals not diagnosed with ER+ BC) when they were last known to be alive and
not have ER+ BC, was included as a linear and squared covariate to control for potential age effects
while maximizing available sample sizes. The undiagnosed age could be the age of the individual at
the date the Palga records were up to date until (February 21st, 2023) or the date of death for an
individual if this occurred prior (n = 2,293). Dates of death were included in Lifelines as recorded by
the Dutch government and were known even in cases of emigration. These dates were decimalized
using *lubridate* 1.9.3 in R (Grolemund & Wickham, 2011) and ages were calculated using the birth
year and month available in Lifelines. We also included the participant's birthdate as a linear and
squared covariate, to account for any temporal trends in variables, and the top 20 principal components
of the pruned GRM used in each model to account for possible population stratification (Yang et al.,
2014). Principal components were generated using the *--pca* argument in the *gcta64* function.

However, our bivariate models struggled to converge and univariate models provided apparently
biased heritability estimates inconsistent with the literature (e.g., 0% for age at menarche, Table S1d).
We posited that this was due to poor coverage of loci between the different genomic samples. For
example, although we had over 1 million SNPs in total, the number of SNPs measured in individuals
belonging to, for example, both the Affymetrix and CytoSNP array samples was ~2% of the total
number of SNPs available (Table S2). This could result in biased genomic relatedness estimates and
increase the chance of missing causal loci, downwardly biasing heritability estimates in the univariate
models (Yang et al., 2010). We therefore created GRMs based on each genomic dataset separately
where the number of loci measured between individuals was larger and more consistent (Table S2).
We present the analysis using the Global screening array data in the main text – because it was the
largest in both number of SNPs and individuals – and use the analyses of the other cohorts as
replicates. Our sample sizes of informative individuals for our analyses using the Global screening
array were: age at menarche, 8,071; age at first birth, 6,260; number of children ever born, 4,602; and
182 ER+ BC cases and 8,700 controls. We also performed a power analysis for our genetic
correlations estimated from GREML analyses for each of our traits and genomic datasets based upon
heritability estimates from univariate analyses and case and control sample sizes (Figure S1, following
Visscher et al., 2014)

Likelihood ratio tests (LRTs) were used to test if heritabilities (in univariate models) and genetic
correlations (in bivariate models) were statistically different from 0. For genetic correlations, we
adjusted for multiple testing across our three life-history traits using Bonferroni correction to a cutoff
of 0.05, giving an adjusted significance threshold of 0.01$\dot{6}$. To aid convergence, we also used the
Expectation–maximization algorithm, increased the number of iterations to 10,000, and standardized
life-history trait values to a mean of zero and standard deviation of one.

LD score regression of GWAS data

We then replicated our findings from the genomic relatedness analyses using LD score regression to
estimate heritabilities and genetic correlations between our traits from previously published GWAS
summary statistics (Bulik-Sullivan, Loh, et al., 2015). LD score regression combines genome-wide
association study (GWAS) data (i.e., statistical associations between all genetic variants and a trait
from a particular study) across different analyses to examine how strongly correlated the effects of
genetic variants are across different traits, controlling for their genetic linkage. For ER+ BC risk, we
used   GWAS   summary   statistics   from   Michailidou   et   al.,   (2017,   available   at:
https://www.ebi.ac.uk/gwas/studies/GCST004988) which contained associations of 11,792,542 SNPs
with ER+ BC risk based upon 69,501 cases and 95,042 controls (see Michailidou et al., 2017, for full
methods). We cleaned these data using the *munge_sumstats* function from the software using *ldsc*
1.0.1 (Bulik-Sullivan, Loh, et al., 2015) supported by *Miniconda3*. Specifically, we removed SNPs not

found in the HapMap3 to control for imputation quality in the absence of imputation quality scores (n

= 10,643,464), with MAF < 0.01 (n = 10,735), missing values (n = 8,108), that were strand ambiguous

(n = 379), and duplicated identifiers (n = 408). This left 1,129,448 SNPs with a mean χ^2 value of

1.537, 2,217 of which held genome-wide significance.

For our life-history traits, we used SNP summary statistics from Tan et al. (2024): a recent family-

based GWAS study which aimed to separate direct genetic effects of a given SNP on a trait from

indirect genetic effects and factors such as assortative mating and population stratification (Kong et

al., 2018; A. I. Young et al., 2022). This can be achieved by conditioning genetic associations found

across individuals upon the parental genotypes in the GWAS model regression; see Tan et al., (2024)

for a full description of methods.

We downloaded the Tan et al., (2024) summary statistics from the SSGAC data portal

(https://thessgac.com/), which contained the summary statistics of 5,671,230 SNPs for age at

menarche, 6,551,047 for age at first birth, and 6,412,325 for number of children ever born. We again

used the *munge_sumstats* function to clean the data prior to analyses and, again, selected only cites

used in HapMap3, and – as advised by Tan et al., (2024) – filtered for SNPs with an effective sample

size greater than the median for each trait to ensure summary statistics were included for only loci

whose genetic effects were well estimated. The number of SNPs with known associations for each trait

after cleaning were 580,219 for age at menarche, 708,027 for age at first birth, and 670,251 for

number of children ever born. The mean and number of genome-wide significant SNPs for these traits

were 1.127 and 26, 1.078 and 0, and 1.051 and 0, respectively.

Heritabilities and genetic correlations were estimated using the software *ldsc* (1.0.1, Bulik-Sullivan,

Finucane, et al., 2015; Bulik-Sullivan, Loh, et al., 2015). *ldsc* estimates the heritability of a trait by

regressing the χ^2-statistics from SNPs on linkage disequilibrium scores using the equation:

$$E\left[\chi_j^2\right] \approx \frac{N_j h_g^2}{M} l_j + 1$$

where $E\left[\chi_j^2\right]$ is the expected χ^2-statistics for the association between the trait and SNP j, N_j is the

number of individuals measured for the trait for SNP j, h_g^2 is the squared correlation between the

phenotype and the best linear unbiased predictor estimated from the SNPs, M is the total number of

SNPs, and l_j denotes the LD score of SNP j (Bulik-Sullivan, Loh, et al., 2015). In all models, we used

European ancestry LD scores from the 1000 Genome project (The 1000 Genomes Project Consortium,

2010) available from online at: https://zenodo.org/records/7768714. We removed loci in the major

histocompatibility complex because of its unusual genetic architecture.

Genome-wide genetic correlations were estimated using the following equation where the expected

value of $z_{1j} z_2 l_j$ for a SNP j is

$$E[z_{1j}z_2 l_j] = \frac{\sqrt{N_1 N_2}\varrho_g}{M} l_j + \frac{\varrho N_s}{\sqrt{N_1 N_2}}$$

529   where N_i is the sample size for the study i, ϱ_g is the genetic covariance, l_j is the LD score, N_s is the

530   number of total individuals in both studies, ϱ is the phenotypic correlation among the overlapping

531   samples (Bulik-Sullivan, Finucane, et al., 2015). We can then normalize the genetic covariance using

532   the heritabilities from trait 1, h_1^2, and trait 2, h_2^2, to estimate the genetic correlation, r_g, with the

533   equation:

$$r_g = \frac{\varrho_g}{\sqrt{h_1^2 h_2^2}}$$

We used the same LD scores and GWAS data as above. The sample size for SNPs with available LD

scores, and GWAS summary statistics for both traits for our three bivariate models were: age at

menarche = 551,453, age at first birth = 668,389, and number of children ever born = 635,154.

Significance of the genetic correlations was determined by z-score derived p-values provided by *ldsc,*

adjusted to a significance threshold of $0.01\dot{6}$. We also report the heritabilities estimated for each trait in

the genetic correlation models and the mean z-scores of each SNPs multiplied association with both

traits. As a supplementary analysis we also estimated our genetic correlations using the genetic effects

from Tan *et al.* (2024) that did not correct for environmental confounding (i.e., population genetic

effects), but this did not produce qualitatively different results (Table S6).

544   DATA AVAILABILITY

545   Data may be obtained from Lifelines and are not publicly available. More information about how to

546   request Lifelines data and the conditions of use can be found on their website (https://www.lifelines-

547   biobank.com/researchers/working-with-us/step-1-prepare-and-submit-your-application). Code for

548   replicating data wrangling and analysis is available for reviewers here:

549   https://dataverse.nl/privateurl.xhtml?token=93828d52-1a89-4e46-8244-7e21eaf987cf.

551   FUNDING

The Lifelines initiative has been made possible by subsidy from the Dutch Ministry of Health, Welfare

and Sport, the Dutch Ministry of Economic Affairs, the University Medical Center Groningen

(UMCG), Groningen University and the Provinces in the North of the Netherlands (Drenthe,

Friesland, Groningen). E.A.Y.'s PhD was funded by the University of Groningen, through a Rosalind

Franklin Fellowship awarded to H.L.D. V.L. was funded by the Strategic Research Council of the

Academy of Finland (grant nos. 345185 and 345183) and an European Research Grant (ERC-2022-
ADG 101098266).

ACKNOWLEDGEMENTS

We thank the Lifelines data management team and especially Sylvia Gerritsma for their support during
this project and Mariana Escobar Rodríguez, Grigory Sivolenko, Thijs Janzen, Raphaël Scherrer,
Judith Vonk, Harold Snieder, and Ilja Nolte for their advice on the data handling and analysis. Finally,
we would also like to thank the University of Exeter's quantitative genetics club and Luca G. Hahn for
conceptual feedback and discussions regarding the results. We thank the Center for Information
Technology of the University of Groningen for their support and for providing access to the Hábrók
high performance computing cluster.

ETHICAL APPROVAL INFORMATION

The Lifelines Cohort Study is conducted following the principles of the Declaration of Helsinki and
with approval from the Medical Ethics Committee of the University Medical Center Groningen,
Groningen (number 2007/152). Written informed consent was given by each Lifelines participant and
the research presented here were approved by Lifelines board (number OV17_00378), adhering to the
Lifelines User Code of Conduct. All data was pseudo-anonymized before being received and was
handled in accordance with the General Data Protection and Regulation act. This research was
approved by the scientific commission of Palga (number 2023-126). Further, data were accessed only
within the secure Lifelines workspace or cluster environment on the University Medical Center
Groningen cluster *Gearshift* (https://docs.gcc.rug.nl/gearshift/cluster/).

REFERENCES

Aitken, R. J. (2022). The changing tide of human fertility. *Human Reproduction*, *37*(4), 629–638.
https://doi.org/10.1093/humrep/deac011

Aktipis, C. A., Boddy, A. M., Gatenby, R. A., Brown, J. S., & Maley, C. C. (2013). Life history trade-
offs in cancer evolution. *Nature Reviews Cancer*, *13*(12), 883–892.
https://doi.org/10.1038/nrc3606

Aktipis, C. A., Ellis, B. J., Nishimura, K. K., & Hiatt, R. A. (2015). Modern reproductive patterns

associated with estrogen receptor positive but not negative breast cancer susceptibility.

Evolution, Medicine and Public Health, *2015*(1), 52–74. https://doi.org/10.1093/emph/eou028

Barban, N., Jansen, R., De Vlaming, R., Vaez, A., Mandemakers, J. J., Tropf, F. C., Shen, X., Wilson,

J. F., Chasman, D. I., Nolte, I. M., Tragante, V., Van Der Laan, S. W., Perry, J. R. B., Kong, A.,

Ahluwalia, T. S., Albrecht, E., Yerges-Armstrong, L., Atzmon, G., Auro, K., … Mills, M. C.

(2016). Genome-wide analysis identifies 12 loci influencing human reproductive behavior.

Nature Genetics, *48*(12), 1462–1472. https://doi.org/10.1038/ng.3698

Bauer, K. R., Brown, M., Cress, R. D., Parise, C. A., & Caggiano, V. (2007). Descriptive analysis of

estrogen receptor (ER)-negative, progesterone receptor (PR)-negative, and HER2-negative

invasive breast cancer, the so-called triple-negative phenotype: A population-based study from

the California cancer Registry. *Cancer*, *109*(9), 1721–1728. https://doi.org/10.1002/cncr.22618

Benonisdottir, S., Straub, V. J., Kong, A., & Mills, M. C. (2024). Genetics of female and male

reproductive traits and their relationship with health, longevity and consequences for

offspring. *Nature Aging*, *4*(12), 1745–1759. https://doi.org/10.1038/s43587-024-00733-w

Berg, J. J., Harpak, A., Sinnott-Armstrong, N., Joergensen, A. M., Mostafavi, H., Field, Y., Boyle, E.

A., Zhang, X., Racimo, F., Pritchard, J. K., & Coop, G. (2019). Reduced signal for polygenic

adaptation of height in UK Biobank. *eLife*, *8*, e39725. https://doi.org/10.7554/eLife.39725

Boddy, A. M., Kokko, H., Breden, F., Wilkinson, G. S., & Aktipis, C. A. (2015). Cancer susceptibility

and reproductive trade-offs: A model of the evolution of cancer defences. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *370*(1673), 20140220.

https://doi.org/10.1098/rstb.2014.0220

Bonnet, T., Morrissey, M. B., Villemereuil, P. D., Alberts, S. C., Arcese, P., Bailey, L. D., Boutin, S.,

Brekke, P., Brent, L. J. N., Camenisch, G., Charmantier, A., Clutton-Brock, T. H., Cockburn,

A., Coltman, D. W., Courtiol, A., Davidian, E., Evans, S. R., Ewen, J. G., Festa-bianchet, M.,

… Kruuk, L. E. B. (2022). Genetic variance in fitness indicates rapid on-going adaptive

evolution in wild animals. *Science*, *376*(6596), 1012–1016.

Border, R., Athanasiadis, G., Buil, A., Schork, A. J., Cai, N., Young, A. I., Werge, T., Flint, J., Kendler, K. S., Sankararaman, S., Dahl, A. W., & Zaitlen, N. A. (2022). Cross-trait assortative mating is widespread and inflates genetic correlation estimates. *Science*, *378*(6621), 754761.

Bulik-Sullivan, B. K., Finucane, H. K., Anttila, V., Gusev, A., Day, F. R., Loh, P.-R., ReproGen Consortium, Psychiatric Genomics Consortium, Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3, Duncan, L., Perry, J. R. B., Patterson, N., Robinson, E. B., Daly, M. J., Price, A. L., & Neale, B. M. (2015). An atlas of genetic correlations across human diseases and traits. *Nature Genetics*, *47*(11), 1236–1241. https://doi.org/10.1038/ng.3406

Bulik-Sullivan, B. K., Loh, P.-R., Finucane, H. K., Ripke, S., Yang, J., Consortium, S. W. G. of the P. G., Patterson, N., Daly, M. J., Price, A. L., & Neale, B. M. (2015). LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nature Genetics*, *47*(3), 291–295. https://doi.org/10.1038/ng.3211

Byars, S. G., Huang, Q. Q., Gray, L.-A., Bakshi, A., Ripatti, S., Abraham, G., Stearns, S. C., & Inouye, M. (2017). Genetic loci associated with coronary artery disease harbor evidence of selection and antagonistic pleiotropy. *PLOS Genetics*, *13*(6), e1006328. https://doi.org/10.1371/journal.pgen.1006328

Byars, S. G., & Voskarides, K. (2020). Antagonistic Pleiotropy in Human Disease. *Journal of Molecular Evolution*, *88*(1), 12–25. https://doi.org/10.1007/s00239-019-09923-2

Casparie, M., Tiebosch, A. T. M. G., Burger, G., Blauwgeers, H., Van De Pol, A., Van Krieken, J. H. J. M., & Meijer, G. A. (2007). Pathology Databanking and Biobanking in The Netherlands, a Central Role for PALGA, the Nationwide Histopathology and Cytopathology Data Network and Archive. *Analytical Cellular Pathology*, *29*(1), 19–24. https://doi.org/10.1155/2007/971816

Charlesworth, B. (1980). *Evolution in age-structured populations*. Cambridge University Press Cambridge.

639   Charlesworth, B. (2001). Patterns of age-specific means and genetic variances of mortality rates

640   predicted by the mutation-accumulation theory of ageing. *Journal of Theoretical Biology*,

641   *210*(1), 47–65.

642   Coe, K., & Steadman, L. B. (1995). The human breast and the ancestral reproductive cycle: A

643   preliminary inquiry into breast cancer etiology. *Human Nature*, *6*(3), 197–220.

644   https://doi.org/10.1007/BF02734139

645   Colditz, G. A., Rosner, B. A., Chen, W. Y., Holmes, M. D., & Hankinson, S. E. (2004). Risk Factors

646   for Breast Cancer According to Estrogen and Progesterone Receptor Status. *JNCI Journal of*

647   *the National Cancer Institute*, *96*(3), 218–228. https://doi.org/10.1093/jnci/djh025

648   Corbett, S., Courtiol, A., Lummaa, V., Moorad, J., & Stearns, S. (2018). The transition to modernity

649   and chronic disease: Mismatch and natural selection. *Nature Reviews Genetics*, *19*(7), 419–

650   430. https://doi.org/10.1038/s41576-018-0012-3

651   Crespi, B. J., & Summers, K. (2006). Positive selection in the evolution of cancer. *Biological Reviews*

652   *of the Cambridge Philosophical Society*, *81*(3), 407–424.

653   https://doi.org/10.1017/S1464793106007056

654   Dall, G. V., Hawthorne, S., Seyed-Razavi, Y., Vieusseux, J., Wu, W., Gustafsson, J.-A., Byrne, D.,

655   Murphy, L., Risbridger, G. P., & Britt, K. L. (2018). Estrogen receptor subtypes dictate the

656   proliferative nature of the mammary gland. *Journal of Endocrinology*, *237*(3), 323–336.

657   https://doi.org/10.1530/JOE-17-0582

658   Day, F. R., Thompson, D. J., Helgason, H., Chasman, D. I., Finucane, H., Sulem, P., Ruth, K. S.,

659   Whalen, S., Sarkar, A. K., Albrecht, E., Altmaier, E., Amini, M., Barbieri, C. M., Boutin, T.,

660   Campbell, A., Demerath, E., Giri, A., He, C., Hottenga, J. J., … Smith, G. D. (2017). Genomic

661   analyses identify hundreds of variants associated with age at menarche and support a role for

662   puberty timing in cancer risk. *Nature Genetics*, *49*(6), 834–841.

663   https://doi.org/10.1038/ng.3841

664   Duan, R., Chen, Y., Qiao, T., Duan, R., Chen, M., Zhao, L., Gong, Y., & Cheng, G. (2020). Modern

665   dietary pattern is prospectively associated with earlier age at menarche: Data from the CHNS

666   1997–2015. *Nutrition Journal*, *19*(1), 95. https://doi.org/10.1186/s12937-020-00622-z

Dujon, A. M., Boutry, J., Tissot, S., Lemaître, J.-F., Boddy, A. M., Gérard, A.-L., Alvergne, A., Arnal,

A., Vincze, O., Nicolas, D., Giraudeau, M., Telonis-Scott, M., Schultz, A., Pujol, P., Biro, P.

A., Beckmann, C., Hamede, R., Roche, B., Ujvari, B., & Thomas, F. (2022). Cancer

Susceptibility as a Cost of Reproduction and Contributor to Life History Evolution. *Frontiers*

in Ecology and Evolution, *10*(May), 1–18. https://doi.org/10.3389/fevo.2022.861103

Eaton, S. B., Pike, M. C., Short, R. V., Lee, N. C., Trussell, J., Hatcher, R. A., Wood, J. W., Worthman,

C. M., Konner, M. J., Hill, K. R., Bailey, R., & Hurtado, A. M. (1994). Women's Reproductive

Cancers in Evolutionary Context. *The Quarterly Review of Biology*, *69*(3), 353–367.

https://doi.org/10.1086/418650

Fisher, R. A. (1930). *The genetical theory of natural selection*. Oxford University Press.

GBD. (2021). *Global Burden of Disease Study 2021* [Data set]. Global Burden of Disease

Collaborative Network, Institute for Health Metrics and Evaluation (IHME).

https://vizhub.healthdata.org/gbd-results/

Gerring, Z. F., Thorp, J. G., Gamazon, E. R., & Derks, E. M. (2022). An analysis of genetically

regulated gene expression and the role of co-expression networks across 16 psychiatric and

substance use phenotypes. *European Journal of Human Genetics*, *30*(5), 560–566.

https://doi.org/10.1038/s41431-022-01037-6

Ghosh, M. G., Thompson, D. A., & Weigel, R. J. (2020). PDZK1 and GREB1 Are Estrogen-regulated

Genes Expressed in Hormone-responsive Breast Cancer. *Cancer Research*, *60*, 6367–6375.

Green, J., Cairns, B. J., Casabonne, D., Wright, F. L., Reeves, G., & Beral, V. (2011). Height and

cancer incidence in the Million Women Study: Prospective cohort, and meta-analysis of

prospective studies of height and total cancer risk. *The Lancet Oncology*, *12*(8), 785–794.

https://doi.org/10.1016/S1470-2045(11)70154-1

Grolemund, G., & Wickham, H. (2011). Dates and Times Made Easy with {lubridate}. *Journal of*

Statistical Software, *40*(3), 1–25. https://www.jstatsoft.org/v40/i03/

Haldane, J. (1941). *New paths in genetics.* Allen & Unwin.

Haldane, J. (1954). *The measurement of natural selection*. *1*, 480–487.

Hidaka, B. H., & Boddy, A. M. (2016). Is estrogen receptor negative breast cancer risk associated with

a fast life history strategy? *Evolution, Medicine and Public Health*, *2016*(1), 17–20.

https://doi.org/10.1093/EMPH/EOV034

Howe, L. J., Nivard, M. G., Morris, T. T., Hansen, A. F., Rasheed, H., Cho, Y., Chittoor, G., Ahlskog,

R., Lind, P. A., Palviainen, T., Van Der Zee, M. D., Cheesman, R., Mangino, M., Wang, Y., Li,

S., Klaric, L., Ratliff, S. M., Bielak, L. F., Nygaard, M., … Davies, N. M. (2022). Within-

sibship genome-wide association analyses decrease bias in estimates of direct genetic effects.

Nature Genetics, *54*(5), 581–592. https://doi.org/10.1038/s41588-022-01062-7

Howie, B., Fuchsberger, C., Stephens, M., Marchini, J., & Abecasis, G. R. (2012). Fast and accurate

genotype imputation in genome-wide association studies through pre-phasing. *Nature*

Genetics, *44*(8), 955–959. https://doi.org/10.1038/ng.2354

IKNL. (2012). *Breast Cancer Guideline*. The Netherlands Comprehensive Cancer Organization.

https://www.nabon.nl/wp-content/uploads/2022/10/Dutch-Breast-Cancer-Guideline-2012.pdf

Jacqueline, C., Biro, P. A., Beckmann, C., Moller, A. P., Renaud, F., Sorci, G., Tasiemski, A., Ujvari,

B., & Thomas, F. (2017). Cancer: A disease at the crossroads of trade-offs. *Evolutionary*

Applications, *10*(3), 215–225. https://doi.org/10.1111/eva.12444

Jasienska, G., Bribiescas, R. G., Furberg, A. S., Helle, S., & Núñez-de la Mora, A. (2017). Human

reproduction and health: An evolutionary perspective. *The Lancet*, *390*(10093), 510–520.

https://doi.org/10.1016/S0140-6736(17)30573-1

Jiang, X., Finucane, H. K., Schumacher, F. R., Schmit, S. L., Tyrer, J. P., Han, Y., Michailidou, K.,

Lesseur, C., Kuchenbaecker, K. B., Dennis, J., Conti, D. V., Casey, G., Gaudet, M. M.,

Huyghe, J. R., Albanes, D., Aldrich, M. C., Andrew, A. S., Andrulis, I. L., Anton-Culver, H.,

… Lindström, S. (2019). Shared heritability and functional enrichment across six solid

cancers. *Nature Communications*, *10*(1), 431. https://doi.org/10.1038/s41467-018-08054-4

Kang, H. J., Feng, Z., Sun, Y., Atwal, G., Murphy, M. E., Rebbeck, T. R., Rosenwaks, Z., Levine, A. J.,

& Hu, W. (2009). Single-nucleotide polymorphisms in the p53 pathway regulate fertility in

humans. *Proceedings of the National Academy of Sciences of the United States of America*,

106(24), 9761–9766. https://doi.org/10.1073/pnas.0904280106

Kapsetaki, S. E., Compton, Z. T., Dolan, J., Harris, V. K., Mellon, W., Rupp, S. M., Duke, E. G.,

Harrison, T. M., Aksoy, S., Giraudeau, M., Vincze, O., McGraw, K. J., Aktipis, A., Tollis, M.,

Boddy, A. M., & Maley, C. C. (2024). Life history traits and cancer prevalence in birds.

Evolution, Medicine, and Public Health, *12*(1), 105–116.

https://doi.org/10.1093/emph/eoae011

Kelsey, J. L., & Bernstein, L. (1996). Epidemiology and prevention of breast cancer. *Annual Review of*

Public Health, *17*(1), 47–67.

Kelsey, J. L., Gammon, M. D., & John, E. M. (1993). Reproductive factors and breast cancer.

Epidemiologic Reviews, *15*(1), 36–47. https://doi.org/10.1093/oxfordjournals.epirev.a036115

Kemper, K. E., Yengo, L., Zheng, Z., Abdellaoui, A., Keller, M. C., Goddard, M. E., Wray, N. R.,

Yang, J., & Visscher, P. M. (2021). Phenotypic covariance across the entire spectrum of

relatedness for 86 billion pairs of individuals. *Nature Communications*, *12*(1), 1050.

https://doi.org/10.1038/s41467-021-21283-4

Kirk, D. (1996). Demographic transition theory. *Population Studies*, *50*(3), 361–387.

Kong, A., Thorleifsson, G., Frigge, M. L., Vilhjalmsson, B. J., Young, A. I., Thorgeirsson, T. E.,

Benonisdottir, S., Oddsson, A., Halldorsson, B. V., Masson, G., & others. (2018). The nature

of nurture: Effects of parental genotypes. *Science*, *359*(6374), 424–428.

Lande, R., & Arnold, S. J. (1983). The measurement of selection on correlated characters. *Evolution*,

37(6), 1201226. https://www.jstor.org/stable/2408842

Lee, S. H., Yang, J., Goddard, M. E., Visscher, P. M., & Wray, N. R. (2012). Estimation of pleiotropy

between complex diseases using single-nucleotide polymorphism-derived genomic

relationships and restricted maximum likelihood. *Bioinformatics*, *28*(19), 2540–2542.

https://doi.org/10.1093/bioinformatics/bts474

Lifelines. (2024). [Data set]. https://data-catalogue.lifelines.nl/

Long, E., & Zhang, J. (2023). Evidence for the role of selection for reproductively advantageous

alleles in human aging. *Science Adanvces*, *9*(eadh4990), 1–11.

Medawar, P. (1952). An unsolved problem in biology. Delineates of Theory of Mutation

Accumulation. *Lewis, HK London, UK. Reprinted in: Medawar, PB (1957). The Uniqueness of*

the Individual. Methuen. London, UK, 44–70.

Merlo, L. M. F., Pepper, J. W., Reid, B. J., & Maley, C. C. (2006). Cancer as an evolutionary and

ecological process. *Nature Reviews Cancer*, *6*(12), 924–935. https://doi.org/10.1038/nrc2013

Michailidou, K., Lindström, S., Dennis, J., Beesley, J., Hui, S., Kar, S., Lemaçon, A., Soucy, P., Glubb,

D., Rostamianfar, A., Bolla, M. K., Wang, Q., Tyrer, J., Dicks, E., Lee, A., Wang, Z., Allen, J.,

Keeman, R., Eilber, U., … Easton, D. F. (2017). Association analysis identifies 65 new breast

cancer risk loci. *Nature*, *551*(7678), 92–94. https://doi.org/10.1038/nature24284

Michailidou, K., The Breast and Ovarian Cancer Susceptibility Collaboration, Hereditary Breast and

Ovarian Cancer Research Group Netherlands (HEBON), kConFab Investigators, Australian

Ovarian Cancer Study Group, The GENICA (Gene Environment Interaction and Breast

Cancer in Germany) Network, Hall, P., Gonzalez-Neira, A., Ghoussaini, M., Dennis, J., Milne,

R. L., Schmidt, M. K., Chang-Claude, J., Bojesen, S. E., Bolla, M. K., Wang, Q., Dicks, E.,

Lee, A., Turnbull, C., … Easton, D. F. (2013). Large-scale genotyping identifies 41 new loci

associated with breast cancer risk. *Nature Genetics*, *45*(4), 353–361.

https://doi.org/10.1038/ng.2563

Mills, M. C., Tropf, F. C., Brazel, D. M., van Zuydam, N., Vaez, A., Agbessi, M., Ahsan, H., Alves, I.,

Andiappan, A. K., Arindrarto, W., Awadalla, P., Battle, A., Beutner, F., Jan Bonder, M.,

Boomsma, D. I., Christiansen, M. W., Claringbould, A., Deelen, P., Esko, T., … Day, F. R.

(2021). Identification of 371 genetic variants for age at first sex and birth linked to

externalising behaviour. *Nature Human Behaviour*. https://doi.org/10.1038/s41562-021-

01135-3

Momenimovahed, Z., & Salehiniya, H. (2019). Epidemiological characteristics of and risk factors for

breast cancer in the world. *Breast Cancer: Targets and Therapy*, *Volume 11*, 151–164.

https://doi.org/10.2147/BCTT.S176070

Nguyen, N., Fan, H.-Y., Tsai, M.-C., Tung, T.-H., Huynh, Q., Huang, S.-Y., & Chen, Y. (2020).

Nutrient Intake through Childhood and Early Menarche Onset in Girls: Systematic Review

and Meta-Analysis. *Nutrients*, *12*(9), 2544. https://doi.org/10.3390/nu12092544

Nunney, L. (2013). The real war on cancer: The evolutionary dynamics of cancer suppression.

Evolutionary Applications, *6*(1), 11–19. https://doi.org/10.1111/eva.12018

Nunney, L. (2018). Size matters: Height, cell number and a person's risk of cancer. *Proceedings of the

Royal Society B: Biological Sciences*, *285*(1889), 20181743.

https://doi.org/10.1098/rspb.2018.1743

Nunney, L. (2024). The effect of body size and inbreeding on cancer mortality in breeds of the

domestic dog: A test of the multi-stage model of carcinogenesis. *Royal Society Open Science*,

11(1), 231356. https://doi.org/10.1098/rsos.231356

Parent, A.-S., Teilmann, G., Juul, A., Skakkebaek, N. E., Toppari, J., & Bourguignon, J.-P. (2003). The

Timing of Normal Puberty and the Age Limits of Sexual Precocity: Variations around the

World, Secular Trends, and Changes after Migration. *Endocrine Reviews*, *24*(5), 668–693.

https://doi.org/10.1210/er.2002-0019

Pavard, S., & Metcalf, C. J. E. (2007). Negative Selection on BRCA1 Susceptibility Alleles Sheds

Light on the Population Genetics of Late-Onset Diseases and Aging Theory. *PLoS ONE*,

2(11), e1206. https://doi.org/10.1371/journal.pone.0001206

Perry, J. R. B., Murray, A., Day, F. R., & Ong, K. K. (2015). Molecular insights into the aetiology of

female reproductive ageing. *Nature Reviews Endocrinology*, *11*(12), 725–734.

https://doi.org/10.1038/nrendo.2015.167

Prince, C., Howe, L. D., Sanderson, E., Sharp, G. C., Fraser, A., Lloyd-Lewis, B., & Richmond, R. C.

(2024). Evaluating the causal impact of reproductive factors on breast cancer risk: A

multivariable mendelian randomization approach. *medRxiv*.

https://doi.org/10.1101/2024.02.02.24301815

Promislow, D. E. L., & Harvey, P. H. (1990). Living fast and dying young: A comparative analysis of

life□history variation among mammals. *Journal of Zoology*, *220*(3), 417–437.

https://doi.org/10.1111/j.1469-7998.1990.tb04316.x

R Core Team. (2020). *R: A language environment for statistical computing* [Computer software]. R

Foundation for Statistical Computing. http://www.r-project.org/.

Rees, M. (1993). Menarche when and why? *The Lancet*, *342*(8884), 1375–1376.

https://doi.org/10.1016/0140-6736(93)92746-G

Robertson, A. (1966). A mathematical model of the culling process in dairy cattle. *Animal Science*,

8(1), 95–108.

Rodríguez, J. A., Marigorta, U. M., Hughes, D. A., Spataro, N., Bosch, E., & Navarro, A. (2017).

Antagonistic pleiotropy and mutation accumulation influence human senescence and disease.

Nature Ecology and Evolution, *1*(3), 1–5. https://doi.org/10.1038/s41559-016-0055

Roff, D. A. (1992). *The evolution of life histories: Theory and analysis*. Chapman & Hall; WorldCat.

Scholtens, S., Smidt, N., Swertz, M. A., Bakker, S. J., Dotinga, A., Vonk, J. M., Van Dijk, F., Van Zon,

S. K., Wijmenga, C., Wolffenbuttel, B. H., & Stolk, R. P. (2015). Cohort Profile: LifeLines, a

three-generation cohort study and biobank. *International Journal of Epidemiology*, *44*(4),

1172–1180. https://doi.org/10.1093/ije/dyu229

Shi, H., Mancuso, N., Spendlove, S., & Pasaniuc, B. (2017). Local Genetic Correlation Gives Insights

into the Shared Genetic Architecture of Complex Traits. *The American Journal of Human*

Genetics, *101*(5), 737–751. https://doi.org/10.1016/j.ajhg.2017.09.022

Short, R. V. (1976). The evolution of human reproduction. *Proceedings of the Royal Society of*

London. Series B. Biological Sciences, *195*(1118, A Discussion on Contraceptives of the

Future), 3–24. https://doi.org/10.1098/rspb.1976.0095

Smith, K. R., Hanson, H. A., Mineau, G. P., & Buys, S. S. (2012). Effects of BRCA1 and BRCA2

mutations on female fertility. *Proceedings of the Royal Society B: Biological Sciences*,

279(1732), 1389–1395. https://doi.org/10.1098/rspb.2011.1697

Stearns, S. C. (1977). The Evolution of Life History Traits: A Critique of the Theory and a Review of

the Data. *Annual Review of Ecology and Systematics*, *8*(1), 145–171.

https://doi.org/10.1146/annurev.es.08.110177.001045

Stearns, S. C. (1992). *The Evolution of Life Histories*. Oxford University Press; WorldCat.

Stone, K. L., Platig, J., Quackenbush, J., & Fagny, M. (2025). The Importance of Regulatory Network

Structure for Complex Trait Heritability and Evolution. *Molecular Biology and Evolution*,

42(8), msaf174. https://doi.org/10.1093/molbev/msaf174

Strassmann, B. I. (1999). Menstrual Cycling and Breast Cancer: An Evolutionary Perspective. *Journal*

of Women's Health, *8*(2), 193–202. https://doi.org/10.1089/jwh.1999.8.193

Summers, K., & Crespi, B. (2008). The androgen receptor and prostate cancer: A role for sexual

selection and sexual conflict? *Medical Hypotheses*, *70*(2), 435–443.

https://doi.org/10.1016/j.mehy.2007.04.044

Sung, H., Ferlay, J., Siegel, R. L., Laversanne, M., Soerjomataram, I., Jemal, A., & Bray, F. (2021).

Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide

for 36 Cancers in 185 Countries. *CA: A Cancer Journal for Clinicians*, *71*(3), 209–249.

https://doi.org/10.3322/caac.21660

Tan, T., Jayashankar, H., Guan, J., Nehzati, S. M., Mir, M., Bennett, M., Agerbo, E., Ahlskog, R., Pinto

De Andrade Anapaz, V., Asvold, B. O., Benonisdottir, S., Bhatta, L., Boomsma, D. I.,

Brumpton, B., Campbell, A., Chabris, C. F., Cheesman, R., Chen, Z., China Kadoorie Biobank

Collaborative Group, … Young, A. S. (2024). *Family-GWAS reveals effects of environment*

and mating on genetic associations. https://doi.org/10.1101/2024.10.01.24314703

The 1000 Genomes Project Consortium. (2010). A map of human genome variation from population-

scale sequencing. *Nature*, *467*(7319), 1061–1073. https://doi.org/10.1038/nature09534

The Genome of the Netherlands Consortium. (2014). Whole-genome sequence variation, population

structure and demographic history of the Dutch population. *Nature Genetics*, *46*(8), 818–825.

https://doi.org/10.1038/ng.3021

Tropf, F. C., Stulp, G., Barban, N., Visscher, P. M., Yang, J., Snieder, H., & Mills, M. C. (2015).

Human fertility, molecular genetics, and natural selection in modern societies. *PLoS ONE*,

10(6), 1–14. https://doi.org/10.1371/journal.pone.0126821

Trumble, B. C., Charifson, M., Kraft, T., Garcia, A. R., Cummings, D. K., Hooper, P., Lea, A. J.,

Rodriguez, D. E., Koebele, S. V., Buetow, K., Beheim, B., Minocher, R., Gutierrez, M.,

Thomas, G. S., Gatz, M., Stieglitz, J., Finch, C. E., Kaplan, H., & Gurven, M. (2023).

Apolipoprotein-ε4 is associated with higher fecundity in a natural fertility population. *SCIENCE ADVANCES*.

van Rheenen, W., Peyrot, W. J., Schork, A. J., Lee, S. H., & Wray, N. R. (2019). Genetic correlations of polygenic disease traits: From theory to practice. *Nature Reviews Genetics*, *20*(10), 567–581. https://doi.org/10.1038/s41576-019-0137-z

Visscher, P. M., Hemani, G., Vinkhuyzen, A. A. E., Chen, G.-B., Lee, S. H., Wray, N. R., Goddard, M. E., & Yang, J. (2014). Statistical Power to Detect Genetic (Co)Variance of Complex Traits Using SNP Data in Unrelated Samples. *PLoS Genetics*, *10*(4), e1004269. https://doi.org/10.1371/journal.pgen.1004269

Wang, X., Byars, S., & Stearns, S. (2013). Genetic links between post-reproductive lifespan and family size in Framingham. *Evolution Medicine & Public Health*, *2013*(1), 241–253. https://doi.org/10.1093/emph/eot013

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Williams, G. C. (1957). Pleiotropy, Natural Selection, and the Evolution of Senescence. *Evolution*, *11*(4), 398–411. https://www.jstor.org/stable/2406060

Xiang, Y., Tanwar, V., Singh, P., La Follette, L., Narayan, V. P., & Kapahi, P. (2025). Early menarche and childbirth accelerate aging-related outcomes and age-related diseases: Evidence for antagonistic pleiotropy in humans. *eLife*, *13*, RP102447. https://doi.org/10.7554/eLife.102447

Yang, J., Benyamin, B., McEvoy, B. P., Gordon, S., Henders, A. K., Nyholt, D. R., Madden, P. A., Heath, A. C., Martin, N. G., Montgomery, G. W., Goddard, M. E., & Visscher, P. M. (2010). Common SNPs explain a large proportion of the heritability for human height. *Nature Genetics*, *42*(7), 565–569. https://doi.org/10.1038/ng.608

Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: A Tool for Genome-wide Complex Trait Analysis. *The American Journal of Human Genetics*, *88*(1), 76–82. https://doi.org/10.1016/j.ajhg.2010.11.011

885   Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M., & Price, A. L. (2014). Advantages and

886   pitfalls in the application of mixed-model association methods. *Nature Genetics*, *46*(2), 100–

887   106. https://doi.org/10.1038/ng.2876

888   Yasui, Y., & Potter, J. D. (1999). The shape of age–incidence curves of female breast cancer by

889   hormone-receptor status. *Cancer Causes & Control*, *10*, 431–437.

890   You, W., Symonds, I., Rühli, F. J., & Henneberg, M. (2018). Decreasing Birth Rate Determining

891   Worldwide Incidence and Regional Variation of Female Breast Cancer. *Advances in Breast*

892   *Cancer Research*, *07*(01), 1–14. https://doi.org/10.4236/abcr.2018.71001

893   Young, A. I., Nehzati, S. M., Benonisdottir, S., Okbay, A., Jayashankar, H., Lee, C., Cesarini, D.,

894   Benjamin, D. J., Turley, P., & Kong, A. (2022). Mendelian imputation of parental genotypes

895   improves estimates of direct genetic effects. *Nature Genetics*, *54*(6), 897–905.

896   https://doi.org/10.1038/s41588-022-01085-0

897   Young, A. S. (2024). Genome-wide association studies have problems due to confounding: Are family-

898   based designs the answer? *PLOS Biology*, *22*(4), e3002568.

899   https://doi.org/10.1371/journal.pbio.3002568

900   Zhang, H., Ahearn, T. U., Lecarpentier, J., Barnes, D., Beesley, J., Qi, G., Jiang, X., O'Mara, T. A.,

901   Zhao, N., Bolla, M. K., Dunning, A. M., Dennis, J., Wang, Q., Ful, Z. A., Aittomäki, K.,

902   Andrulis, I. L., Anton-Culver, H., Arndt, V., Aronson, K. J., … García-Closas, M. (2020).

903   Genome-wide association study identifies 32 novel breast cancer susceptibility loci from

904   overall and subtype-specific analyses. *Nature Genetics*, *52*(6), 572–581.

905   https://doi.org/10.1038/ng.1049

906   Zhao, Z., Zhang, J., & Tian, X. (2024). Relationship between age at menarche and breast cancer in

907   individuals, as well as in first-degree kin and estrogen receptor status: A Mendelian

908   randomization study. *Frontiers in Oncology*, *14*, 1408132.

909   https://doi.org/10.3389/fonc.2024.1408132