The gut microbiome shapes pharmacology and treatment outcomes for a key anti-inflammatory therapy

Vanya Sofia Villa Soto[*,1], Alexandra L. Degraeve[*,1], Chloe M. Heath[*,1], Diego A. Orellana[1], Erin R. Reilly[2], Mohana Mukherjee[1], Jacob G. Brockert[3], Darren S. Dumlao[3], Rebecca B. Blank[5], Noah Perlmutter[1], Steven Yu[1], Judith Ashouri[1], Jose U. Scher[5], Andrew D. Patterson[2], Peter J. Turnbaugh[3,4], Renuka R. Nayak[1,6,#]

[1]Department of Medicine, Division of Rheumatology, University of California, San Francisco, San Francisco, CA, USA
[2]Department of Microbiology, Pennsylvania State University, University Park, PA, USA
[3]Department of Microbiology & Immunology, University of California, San Francisco, San Francisco, CA, USA
[4]Chan-Zuckerberg Biohub-San Francisco, San Francisco, CA, USA
[5]Department of Medicine, Division of Rheumatology, New York University, New York, NY, USA
[6]Lead contact

*these authors contributed equally

#Corresponding Author:
Renuka R. Nayak: renuka.nayak@ucsf.edu

Highlights

- The gut microbiome impacts methotrexate (MTX) pharmacology in mice
- The human gut microbiome contributes to interindividual variation in MTX pharmacology
- Human gut microbes produce novel MTX metabolites, pMABG and 6-MPDA
- Microbial metabolism of MTX is linked to treatment outcomes

Abstract

The human gut microbiome encodes a formidable metabolic repertoire that harvests nutrients from the diet, but these same pathways may also metabolize medications. Indeed, large screens have revealed extensive microbial metabolism of drugs *in vitro*, but the pharmacologic and clinical repercussions of microbiota-mediated metabolism *in vivo* remain to be discerned. As a proof-of-concept, we investigate how human gut microbes contribute to *in vivo* pharmacology and efficacy of a key anti-inflammatory drug, methotrexate (MTX). Specifically, we demonstrate that the gut microbiome shapes drug pharmacology *in vivo* in mice, both by directly metabolizing the drug and by inducing host pathways that promote drug metabolism. Moreover, interindividual variation in the human gut microbiome contributes to variation in pharmacokinetic (PK) profiles. When we quantified metabolites produced by microbes, we unexpectedly identified novel MTX metabolites, one of which, p-methylaminobenzoyl-L-glutamic acid (pMABG), was a major byproduct of microbial metabolism both *in vitro* and *in vivo*. Further, we find that a large proportion of patient-associated microbes are capable of metabolizing MTX. Finally, we show that microbial metabolism of MTX is linked to PK profiles and disease outcomes in a mouse model of inflammatory arthritis. Taken together, these findings provide evidence that the human gut microbiome causally contributes to drug pharmacology *in vivo* for a key anti-inflammatory drug through known and novel mechanisms. Our studies provide a framework for elucidating the clinical relevance of drug microbial metabolism in the context of treatment response. These results are a first step towards understanding and manipulating the human gut microbiome in the treatment of autoimmunity and the advancement of precision medicine for millions of patients taking MTX for immune or inflammatory conditions.

Introduction

The human gut microbiome encodes for millions of genes, many of which provide critical benefits for the host, but these microbial pathways may lead to unintended consequences in the setting of drug therapy. For example, microbes facilitate nutrient extraction[1], but they also metabolize therapeutic drugs[2]. Indeed, this extensive metabolic capacity of microbes has been demonstrated in recent large-scale *in vitro* drug screens: >400 therapeutic drugs were metabolized by individual microbial isolates as well as by complex bacterial communities from healthy individuals *ex vivo*[3,4]. However, the *in vivo* repercussions (i.e., therapeutic efficacy or toxicity) of this metabolism on the host remain largely unexplored. Thus, translational studies are needed to elucidate how the microbiome contributes to treatment outcomes in the host. They are essential for developing microbiome-based interventions to transform treatment[5,6].

Toward this goal, we chose to investigate an anchor therapy in the treatment of autoimmune and inflammatory disease: methotrexate (MTX). From a microbiome perspective, MTX is an ideal drug to study because it is an antimetabolite: it structurally mimics an essential vitamin, folate, that is required by nearly all forms of life[7,8]. Microbes encode enzymes to synthesize[99] and break down folate (unlike mice or humans)[10,11]. Importantly, these same folate metabolism enzymes act on MTX[12]. Thus, compelling *in vitro* evidence supports the hypothesis that the microbiome shapes MTX pharmacokinetics (PK) and impacts treatment outcomes.

From a clinical perspective, MTX is widely used to treat millions of patients suffering from autoimmunity, cancer, and other autoimmune diseases (e.g., dermatomyositis, lupus, psoriasis and psoriatic arthritis)[13]. Hence, it is designated by the World Health Organization (WHO) as an essential medicine. Currently, oral MTX is first-line treatment for rheumatoid arthritis (RA)[14] and confers multiple unique therapeutic benefits to patients[15]. For these reasons, treatment guidelines recommend maximally up-titrating oral MTX or, if that fails, using subcutaneous MTX, which is more bioavailable[14]. Thus, MTX is a central therapeutic in the treatment of autoimmunity, and maximizing its efficacy is important for multiple reasons.

Patients, however, show extensive interindividual variation in MTX response, and the underlying molecular mechanisms remain incompletely understood despite decades of investigation[16-18]. Host and environmental factors have been explored with the goal of identifying modifiable factors to improve response[16]. While genetic factors and smoking have been found to contribute to MTX responsiveness, other factors likely remain unidentified[16].

Our group focuses on the impact of the microbiome on the treatment of autoimmunity. We recently showed that among RA patients, pre-treatment microbiomes differ between MTX-responders (MTX-R) versus MTX-non-responders (MTX-NR)[19]. In *ex vivo* studies, pre-treatment RA microbiota variably depleted MTX, and depletion was negatively associated with future MTX response[19]. We identified multiple human gut bacterial strains that directly deplete MTX *in vitro*[20]. Together, these findings show that human gut microbes directly metabolize MTX and that this metabolism is associated with clinical response in patients. However, demonstration that microbial metabolism causally impacts MTX PK and shapes treatment outcomes remains to be determined. Such knowledge is required to translate fundamental basic science findings and pave the way for rationally manipulating the gut microbiome to maximize efficacy of this central therapeutic.

The role of the gut microbiome in MTX pharmacology has been previously explored in animal models[21], but whether human gut microbiota metabolize MTX has been debated[22,23] or

metabolism was thought to be minimal[24]. These early studies were conducted when MTX was
given intravenously, but currently it is administered orally, theoretically allowing for more
interactions between MTX and the gut microbiome. Furthermore, early studies were done before
the development of highly sensitive analytical chemistry platforms, which could distinguish
between MTX and its metabolites. Hence, for historical reasons, previous studies may have
underestimated MTX-microbiota interactions and the ability of the microbiota to shape drug
pharmacology and drug response.
107
Given these clinical associations and supportive *ex vivo* and *in vitro* findings, here we test
the causal contributions of gut microbial metabolism to MTX response *in vivo*. Specifically, we
test the extent to which the human gut microbiome contributes to MTX PK, identify the spectrum
of human microbes that metabolize MTX and the metabolites they produce, and examine the
impacts of microbial metabolism on treatment outcomes. Distinct from prior studies[21,25] largely
examining fecal and urinary drug levels and murine microbiota, we quantify levels of MTX and its
metabolites in the circulation of gnotobiotic mice colonized with human gut microbes. We find that
the microbiome causally contributes to MTX PK and that this metabolism is linked to treatment
outcomes in mice. These findings raise the possibility that microbial metabolism can be modulated
to impact MTX outcomes in the host. These results are a first step towards manipulating the
human gut microbiome in order to advance precision medicine for millions of patients globally
taking MTX for autoimmunity.
120

Results

122
123 **The gut microbiome impacts MTX pharmacology *in vivo***
124
We first sought to determine whether the gut microbiome contributes to MTX PK *in vivo*.
To test this, we compared plasma MTX levels in germ-free (GF, N=6) vs. specific-pathogen-free
(SPF, N=8) C57BL/6J mice given MTX 50 mg/kg oral dose (**Supplementary Figure 1A**). We also
quantified levels of 3 previously reported MTX metabolites: deoxyaminopteroic acid (DAMPA), 7-
hydroxy-methotrexate (7-OH-MTX), and polyglutamated MTX ($MTXPG_n$) (**Figure 1A**)[15]. Of these
previously reported MTX metabolites, DAMPA is the only one that is exclusively produced by
microbes[26]. 7-OH-MTX is produced by the host liver enzyme aldehyde oxidase (AO)[27,28]; to the
best of our knowledge, this is not produced by microbes. Polyglutamated MTX metabolites,
ranging from 2 to 7 additional glutamates, are primarily produced once MTX is absorbed into host
cells and can accumulate intracellularly over several days to months[29]. We collected tail vein
blood at 0 hours (prior to administration) and multiple time points after dosing and profiled plasma
concentrations. From these PK curves, Area Under the Curve (AUC) and peak concentrations
were calculated for each mouse.
138
Peak plasma concentrations of MTX were significantly higher in GF mice relative to SPF
mice (**Figure 1B-C**), suggesting that the microbiome reduces MTX absorption from the gut. Peak
plasma DAMPA concentrations were significantly higher in SPF mice and notably absent in GF
mice (**Figure 1C**), consistent with previous reports indicating that DAMPA is exclusively
microbially produced[30]. This finding also shows that while DAMPA is produced in the gut, it can
enter host circulation. The mean AUC of 7-OH-MTX was higher in SPF mice compared to GF
mice (**Figure 1D**). To discern whether this might be due to increased AO enzymatic activity in
SPF mice, we compared the product-substrate ratio between 7-OH-MTX and MTX. The ratio of
7-OH-MTX to MTX ($7\text{-OH-MTX}_{AUC}/\text{MTX}_{AUC}$) was significantly higher in SPF mice (**Figure 1E**),
suggesting that the microbiome increases MTX liver metabolism in the host. Finally, no significant
differences were observed for polyglutamated MTX (**Supplementary Figure 1B**), which was

either absent or present at very low levels, likely because multiple weeks are required for
accumulation of this metabolite[29].
152
We next quantified fecal levels of MTX and its metabolites. MTX levels were significantly
higher in SPF mice compared to GF mice (**Figure 1F-H**). Consistent with it being a microbiota-
produced metabolite, DAMPA was only seen in the feces of SPF mice (**Figure 1F-H**) and was
present at high levels, suggesting incomplete absorption of DAMPA. Similar to plasma levels,
fecal 7-OH-MTX concentrations were higher in SPF in mice than in GF mice (**Figure 1F-H**). These
findings in the stool support the hypothesis that the gut microbiota impacts drug metabolism to
shape MTX pharmacology.
160
Taken together, these results suggest that the mouse gut microbiome contributes to MTX
PK *in vivo* through microbiota- and host-mediated mechanisms. We show for the first time that
DAMPA, which is exclusively produced by gut microbes, enters into host circulation. We also
show for the first time that levels of 7-OH-MTX, which is made by the liver and excreted into the
gut, is impacted by the microbiota. Thus, the microbiome contributes to MTX pharmacology
directly, through microbial metabolism, and indirectly, through mechanisms of absorption,
metabolism, and/or enterohepatic circulation.
168
169　## Parenterally administered MTX undergoes microbial metabolism
170
Next, we considered whether the route of administration might mitigate microbiota-
mediated metabolism of MTX. Because MTX can be administered orally or subcutaneously in
patients with autoimmune disease, we sought to determine whether gut microbial metabolism of
MTX could be bypassed by administering MTX through non-oral routes. We treated SPF
C57BL/6J mice with MTX 50 mg/kg either via oral gavage (PO) or by intraperitoneal injection (IP)
(N=3 / route of administration). Peak plasma MTX levels and AUC were significantly higher in
mice given IP MTX compared to PO (**Supplementary Figure 2A-C**), a finding that is consistent
with prior reports that parenteral MTX has higher bioavailability[31]. Similarly, peak 7-OH-MTX
levels and AUC were elevated in mice given IP dosing (**Supplementary Figure 2A-C**); the ratio
of 7-OH-MTX to MTX did not differ by route (**Supplementary Figure 2D**). Plasma concentrations
of DAMPA were detected in both groups and did not significantly differ (**Supplementary Figure
2A-C**). These findings suggest injection-dosing of MTX does not bypass gut microbial
metabolism, likely due to enterohepatic circulation of the parent compound[32].
184
185　## Interindividual variation in human gut microbiota contributes to MTX PK *in vivo*
186
Having established a role for the murine microbiome in MTX PK, we next sought to
determine the impact of the human gut microbiome on MTX PK and to ascertain the clinical
relevance of our findings. Specifically, we asked whether natural interindividual variation in human
gut microbial communities of patients contributes to MTX PK *in vivo*. We focused on RA patients
because MTX is first-line therapy for this common autoimmune disease affecting 1% of the
worldwide population[14][14]. We colonized GF C57BL/6J mice with fecal communities from two
different treatment-naïve RA patients (Patient 170 vs. Patient 317, N=11 mice per donor
colonization group) and treated with oral MTX 50 mg/kg. Mice colonized with Patient 170
microbiota had significantly higher peak plasma concentrations of MTX than mice colonized with
microbiota from Patient 317 (**Figure 2A-B**). We were unable to detect significant differences
between 7-OH-MTX levels or 7-OH-MTX/MTX ratios between colonization states, suggesting
similar levels of first-pass liver metabolism (**Figure 2B-E**). While there was a trend toward lower
plasma levels of DAMPA in mice colonized with Patient 170 microbiota (consistent with the higher
MTX levels in this colonization group), we were unable to demonstrate that this difference was

statistically significant (**Figure 2B-D**). These findings show that interindividual variation in RA patient microbiota contributes to differences in MTX PK *in vivo*.

Identification of novel MTX metabolites produced directly by human gut microbes

We were perplexed that despite finding differences in plasma MTX concentrations in mice colonized with different microbiota, we did not find concomitant statistically significant differences in either DAMPA or 7-OH-MTX (**Figure 2**). This led us to ask whether MTX may be metabolized into novel compounds that were not captured/quantified in our studies. In prior studies, we found that type strains of *Clostridium asparagiforme*, *Clostridium innocuum*, and *Clostridium symbiosum* depleted MTX from the culture medium as assessed by high-performance liquid chromatography (HPLC)[20]. But when we profiled these cultures via liquid chromatography-mass spectrometry (LC-MS), which is able to disambiguate between MTX metabolites, we were surprised to find that while MTX was depleted by all three strains, only *C. symbiosum* produced DAMPA, and in relatively small amounts (**Figure 3A**). This led to two non-mutually exclusive hypotheses: (1) DAMPA is further metabolized by *C. asparagiforme* and *C. innocuum*, which is why we do not detect it, or (2) these two species metabolize MTX into metabolites other than DAMPA.

To determine the metabolic fate of MTX, we performed stable isotope labeling experiments followed by global mass spectrometry profiling. Specifically, we incubated unlabeled and d_3-labeled MTX with 3 human gut bacterial isolates, 2 of which metabolize the drug (*C. asparagiforme* and *C. symbiosum*) and 1 which does not (*Bacteroides thetaiotaomicron*)[20]. After 72 hours of incubation, stable isotope profiling identified 2 novel MTX metabolites. While DAMPA was not detected in this isotope labeling study perhaps because it was not produced in sufficient amounts, we found evidence of a novel metabolite carrying the heavy isotope (**Figure 3B**): p-methylaminobenzoyl-L-glutamic acid (pMABG) with a mass-to-charge (*m/z*) value of 281 (unlabeled) or 284 (d_3-labeled). Based on chemical logic, we surmised that a second unlabeled, novel compound should be produced upon generation of pMABG: 6-methylpteridine-2,4-diamine (6-MPDA) with a *m/z* value of 177. We obtained authentic standards of these compounds and confirmed that the fragmentation pattern of our novel metabolites matched those of the authentic standards (**Figure 3C**). We next asked which of these novel metabolites are produced by other species previously shown to metabolize MTX[20], and found that many of these species produced predominantly pMABG rather than DAMPA (**Figure 3D**), suggesting that pMABG is a major microbial metabolite of MTX.

Novel MTX metabolites are produced by complex human microbial communities and detected *in vivo* in mice

We next sought to determine whether these novel metabolites are produced by complex microbial communities from RA patients. We incubated fecal slurries from 17 treatment-naïve patients with vehicle control versus MTX 100 ⌠g/mL for 24 hours and profiled the supernatant by using a newly developed and validated targeted LC-MS method. We found that RA microbiota produced significant amounts of DAMPA and pMABG, with pMABG being the major metabolite (**Figure 4A**).

To determine whether pMABG is produced by the host or exclusively produced by microbiota, we returned to the fecal samples harvested from our initial gnotobiotic mouse experiment and found high levels of pMABG in the stool of SPF mice but not GF mice (**Figure 4B**). These findings suggest that pMABG is exclusively microbially produced.

Taken together, these results show that human gut microbes metabolize MTX into at least
3 compounds: DAMPA, pMABG, and 6-MPDA. DAMPA has not been previously shown to be
directly produced by human gut bacteria (as we have shown here with *C. symbiosum in vitro* and
in vivo in mice, **Figure 2**), and the other 2 have never been reported to be MTX metabolites
before. Thus, these studies have uncovered novel metabolites of a widely used antirheumatic
drug that has been used for over 4 decades, demonstrating that it is only with recent advances in
anaerobic microbiology and analytical chemistry that we are now able to identify novel MTX
metabolites.
259
260   ## Identification of multiple novel MTX metabolizing bacteria from RA patients
261
We next sought to determine whether bacterial species harvested from RA patient
microbiota could similarly metabolize MTX into known and novel metabolites. To build a culture
collection of microbial strains from RA patients, we harvested bacterial isolates from 12 treatment-
naïve RA patients. Patients provided stool samples prior to initiating treatment. We used rich
media to maximize recovery of multiple species and strains as well as a spore-forming enrichment
method[33] to increase the odds of harvesting *Firmicutes*, which our prior studies suggested are
enriched for metabolizing species[20]. Colonies with varying morphologies were identified and
picked for further testing (**Supplementary Figure 3**).
270
We harvested a total of 360 bacterial isolates from RA patient microbial communities
(median of 24 isolates per patient; range: 24 – 48). Of these, 306 (306/360, 85%) grew in liquid
media and could be further studied; 54 (54/360, 15%) grew on the agar media but not in liquid
media (**Supplementary Figure 4A**). Of the 306 bacterial isolates that grew in liquid media, 221
(221/306, 72**%**) grew in MTX 50 mg/mL and could be assessed for MTX metabolism capacity.
The remaining 85 (85/306, 28%) likely did not tolerate the growth inhibitory effects of MTX 50
ug/mL.
278
279   Next, we sought to identify these strains. Of the 360 total isolated strains, we were able to
280   identify bacterial species by full-length 16S sequencing for 241 isolates (241/360, 67%). These
281   represented 36 unique species (**Supplementary Figure 4B**). From each patient, we identified 2-
282   15 unique species (median = 7).
283
Next, we asked which of our harvested isolates metabolize MTX *in vitro*. Of the 221
isolates that grew in MTX, we identified 63 (63/221 = 29% of the isolates that grew in MTX or
63/360 = 18% of total isolates harvested) that could metabolize MTX, defined as strains showing
at least a 60% depletion of MTX in the supernatant (**Figure 5A**). These represented at least 12
unique species as determined by full-length 16S sequencing (**Figure 5A**). Other than *Clostridium*
innocuum[20]*,* none have previously been reported to metabolize MTX. Five unique species
seemed to exhibit strain-level variation in MTX metabolism, in which multiple isolates of the same
species variably showed 60% depletion. These included *Clostridium cibarium*, *Coprobacillus*
cateniformis, *Romboutsia hominis*, *Enterococcus faecalis*, and *Intestinibacter bartletti*. These
findings suggest that multiple patient-associated gut bacterial species can metabolize MTX.
294
295   ## A novel metabolite is a major product of microbial MTX metabolism
296
Next, we sought to identify the metabolites produced by each of these 12 unique species.
For 10 of the 12 strains, we validated significant MTX depletion compared to sterile and negative
controls as quantified by HPLC (see Methods)(**Supplementary Table 1, Figure 5B**); 1 showed
impaired growth (*E. faecalis*, 70% compared to vehicle control) and so we may have been limited
in our detection of MTX metabolism, and 1 (*Enterococcus lactis*) failed to validate. For all 10

validated strains, LC-MS revealed that pMABG was the dominant metabolite produced (**Figure**
5C). While *E. faecalis* did not show significant depletion of MTX by HPLC (**Figure 5B**), we did
detect modest pMABG production (**Figure 5C**), suggesting that it is a MTX metabolizer. For all
11 of the metabolizing strains, we did not detect 7-OH-MTX or DAMPA production compared to
sterile controls (**Supplementary Figure 4C**). This may be either because these are not produced
or because they are further metabolized and shuttled into other anabolic pathways. Given that 6-
MPDA is expected to be a byproduct of pMABG production, we were expecting to see elevated
levels of 6-MPDA in metabolizers. However, we were not able to detect significant elevations of
6-MPDA compared to controls (**Supplementary Figure 4C**), suggesting either that our method
for extracting and quantifying this small molecule needs refinement or that a compound other than
6-MPDA is produced upon pMABG generation (e.g., 6-MPDA bonded to a formyl or methyl
group). While *Hungatella hominis* showed significant MTX depletion by HPLC compared to sterile
controls, the extent of metabolism was modest (12% MTX depletion), and we were not able to
detect any of the known or novel MTX metabolites by LC-MS. This might suggest that *H. hominis*
makes metabolites that remain unknown or that the known metabolites are further metabolized
into compounds that are not captured by our targeted method.
318
Taken together, we have identified at least 11 MTX metabolizing microbes harvested from
RA patients (11/36, representing 31% of identifiable isolates) and found that the major metabolite
produced is pMABG. These findings reveal that a significant percentage of strains isolated from
RA patients have the capacity to metabolize MTX, with some strains exhibiting strain-level
variation. These findings suggest that variation in gut microbiome composition and/or strain level
variation could contribute to variation in MTX response via microbial drug metabolism.
325
326 ## MTX response in an arthritis model is associated with the microbiome
327
Concurrent with our *in vivo* pharmacologic and *in vitro* metabolism studies above, we
sought to develop an animal model system to test the impact of microbial metabolism on MTX
efficacy *in vivo*. For this, we turned to the SKG mouse model of inflammatory arthritis, which
recapitulates multiple features of human RA disease, including production of rheumatoid factor
and CCP antibodies, development of chronic inflammatory arthritis of the small joints, elevated
inflammatory markers, and female predominance[34-37]. Multiple previous studies have shown that
gut microbes are important for the pathogenesis of autoimmunity in this model[34,35,38], but none
have explored its significance in therapeutic drug response.
336
Because prior studies showed that MTX alleviates SKG arthritis when given via the
intraperitoneal route[39], we first tested whether oral MTX similarly alleviated arthritis. In order to
increase our sample size for this pilot experiment, we pooled SKG mice that were bred at two
different facilities: Parnassus ("Parn.") and Mount Zion ("Zion"). The mice from Zion were
transferred to Parnassus, were cohoused with Parnassus-reared mice, and were permitted to
acclimate to the Parnassus colony for 2 weeks ("Experiment 1"). Arthritis was induced in mice
with a single IP injection of zymosan, and then mice were treated with oral MTX 10 mg/kg (N=8)
or vehicle control (VEH, N=8) every other weekday for 3 weeks. The Clinical Arthritis Score (CAS)
was significantly lower in mice treated with MTX compared to VEH mice (**Figure 6A-B**). These
findings confirmed that oral MTX, similar to IP MTX[39], alleviates arthritis in SKG mice.
347
Because we observed variation in treatment response in this pilot experiment, we stratified
the data by facility-of-origin (**Figure 6C**). Remarkably, we saw that while arthritis development
was similar for both groups of mice in the VEH arm (**Figure 6C, left panel**), the response to MTX
was significantly better in Zion-reared mice compared to Parnassus-reared mice (**Figure 6C,**
right panel); evaluation of the AUC of the arthritis curves also supported this finding (**Figure 6D**).

This difference was reproduced in a second set of mice that were not co-housed ("Experiment
2"). Mice reared at Zion showed significantly lower arthritis scores on Days 7 through 14 (**Figure
6E, left panel**), which is also reflected in the arthritis AUC (**Figure 6E, right panel**). Together,
these results suggest that facility-of-origin of the mice impacted treatment response to MTX. Of
note, this murine phenotype recapitulates what is seen in RA: patients possess similar arthritis
severity but can vary significantly in their response to MTX. These findings demonstrate that the
SKG arthritis model responds to oral MTX and that it can be used to dissect mechanisms
underlying variable treatment response.
361
We next tested for differences between the gut microbiomes of mice reared in these two
facilities. We studied two sets of mice to understand the impact of co-housing: "representative"
mice that did not undergo co-housing ("Representative") as well as mice that were co-housed for
2 weeks prior to arthritis induction and treatment (Experiment 1). We performed 16S rRNA gene
amplicon sequencing (16S-seq) on stool samples from Zion-reared SKG mice (N=5), Parnassus-
reared SKG mice (N=4), and mice from our pilot co-housing experiment prior to the start of any
treatments (Experiment 1, Zion-reared N=7; Parnassus-reared N=9). Alpha diversity was
significantly higher in Zion-reared mice (**Supplementary Figure 5A, left panel**) compared to
Parnassus-reared mice. This difference was effectively eliminated by co-housing
(**Supplementary Figure 5A, right panel**), with Parnassus-co-housed mice acquiring multiple
amplicon sequence variants (ASVs, representative of individual species, **Supplementary Table
2**). Community composition (beta diversity) differed by facility-of-origin in the "Representative"
mice (ANOSIM p=0.013, PERMANOVA p=0.017; **Supplementary Figure 5B**, open circles). In
the co-housed mice, this difference persisted, albeit in a reduced form (ANOSIM p=0.01,
PERMANOVA p=0.054; **Supplementary Figure 5B**, closed circles). We next identified
differentially abundant taxa. There were no significant phylum-level differences by facility-of-origin
in either group. While 123 ASVs were differentially abundant in the "Representative" mice (with
109 higher in Zion and 14 higher in Parnassus), after 2 weeks of co-housing, only 8 ASVs were
differentially abundant by facility-of-origin (with 6 higher in Zion and 2 higher in Parnassus,
Supplementary Figure 5C). These findings confirm that the microbiome of Zion- and Parnassus-
reared mice differed and that that co-housing facilitated transfer of microbes from Zion mice to
Parnassus mice. Given that we observed that treatment response to MTX was better in
Parnassus-cohoused-with-Zion mice compared to Parnassus-reared mice (**Figure 6F-G**), this
suggests that transfer of microbes from Zion to Parnassus improved treatment response in this
mouse model.
387

MTX response is linked to MTX metabolism *in vivo*

389
Next, we tested whether the mechanism for differential MTX response was linked to
microbial MTX metabolism and modulation of MTX PK *in vivo*. We re-derived the SKG model in
our gnotobiotic facility. This enables us to selectively examine the contribution of gut microbes
and control for other factors. We colonized GF SKG mice with cecal microbiota from either
Parnassus- vs. Zion-reared mice (N=5-6 per colonization group). We found that peak plasma
concentrations of MTX were higher in mice colonized with Parnassus microbiota (**Figure 7A-B**).
The ratio of 7-OH-MTX to MTX (7-OH-MTX$_{AUC}$/MTX$_{AUC}$) was numerically higher in Parnassus-
colonized mice (**Figure 7D**). Microbial metabolism was also dependent on colonization state:
DAMPA levels and AUC were higher in mice colonized with Parnassus microbiota (**Figure 7A-
C**). pMABG levels were numerically higher in Parnassus-colonized mice, but we were unable to
detect a significant difference in the plasma (**Figure 7A-C**).
In the stool, MTX levels and AUC were significantly higher in Parnassus-colonized mice
(**Figure 7E-G**) suggesting greater intestinal elimination. AUC of 7-OH-MTX was significantly
higher in Parnassus-colonized mice (**Figure 7E-G**). Similarly, fecal DAMPA and pMABG levels

were significantly higher in Parnassus-colonized mice (**Figure 7E**) as was the product-substrate
ratio of fecal pMABG to MTX, suggesting increased microbial metabolism.
Taken together, this suggests that while they may permit increased MTX absorption
relative to Zion-microbiota, Parnassus-microbiota led to increased microbiota- and host-
dependent MTX metabolism as detected in the blood and stool. These findings suggest that
microbiota-dependent metabolism of MTX contributes to MTX PK and is linked to clinical
outcomes in inflammatory arthritis.
411

412 ## Discussion
413

Large-scale *in vitro* screens have identified multiple therapeutic drugs that are metabolized
by human gut microbiota[3,4], but the pharmacologic and clinical repercussions of these microbial
transformations remain largely unknown. Here, we advance our understanding of how the gut
microbiome interacts with a key anti-inflammatory drug, MTX, a WHO essential drug used by
multiple wide-ranging clinical disciplines. MTX has been extensively studied for over 50 years. By
taking a microbiologic lens, we uncover novel pharmacology that fundamentally transforms our
understanding of how the gut microbiome interacts with MTX to shape therapeutic outcomes. We
show that the gut microbiome contributes to MTX PK *in vivo* through both direct and indirect
mechanisms, that natural variation in human microbiota leads to differential MTX PK profiles, that
multiple human-associated microbes metabolize MTX into known (DAMPA) and novel (pMABG
and 6-MPDA) metabolites, and that microbial metabolism *in vivo* in mice is linked to treatment
outcomes. While it has long been known that there is extensive interindividual variation in the
bioavailability of MTX[31], the mechanisms underpinning this variation remain incompletely
elucidated. Here, we provide evidence from multiple orthogonal systems (*in vitro*, *ex vivo*, and *in*
vivo mouse studies) to suggest that the gut microbiome plays a pivotal role. Given that the
microbiome is modifiable through diet, probiotics, transplant, and other strategies, these findings
provide support for targeting the microbiome to maximize the efficacy of this important anti-
inflammatory drug.
432

Our findings place the microbiome on the map as a determinant of MTX response. While
one of the very first MTX metabolites to be discovered was microbiota-produced DAMPA, the
extent of microbial metabolism was reported to be very minimal[24]; our findings run counter to this.
This may be because these early studies were performed in cancer patients receiving intravenous
MTX, and DAMPA was quantified in urine, not the gut, where it is produced[22]. Given that MTX is
administered orally for autoimmunity, these early findings in cancer patients may not generalize.
Our results suggest that MTX undergoes significant metabolism by the gut microbiota, into both
DAMPA and pMABG, and that the highest levels of these are found in the gut, where they are
produced. Importantly, we show that DAMPA and pMABG are found in the blood as well. In some
of our studies (mice colonized with their native microbiota), we find that levels of DAMPA
approximate concentrations achieved by host-produced metabolites, namely 7-OH-MTX,
suggesting that DAMPA may be as important as its host-produced counterpart and should not be
considered a minor metabolite. Further, the microbiota regulates liver metabolism of MTX into 7-
OH-MTX, thereby indirectly impacting MTX pharmacology. Indeed, prior studies show that AO,
the enzyme responsible for 7-OH-MTX production, is upregulated in colonized mice relative to GF
mice[40] and that interindividual variation in AO in humans impacts 7-OH-MTX production[28]. Given
the extensive variation in gut microbiotas across patients, we suspect that the gut microbiome
plays a critical role in MTX bioavailability and drug metabolism. Developing microbiota-directed
interventions will allow patients to derive maximal benefit from this central drug. Future studies
are needed to understand natural variation in the production of these metabolites and their
association with response in patients, the physiologic impacts of microbiota-produced metabolites

on the host and its commensal microbiome, as well as how best to specifically modulate the
human microbiome to improve drug response.
456
Our findings showcase how taking a microbiome perspective leads to unexpected
pharmacological insights. For instance, we were surprised to discover novel MTX metabolites
given that the pharmacology of MTX had been investigated for decades. Stable isotope labeling
combined with *in vitro* microbiologic assessment allowed us to identify novel MTX metabolites;
GF mouse studies revealed that one of these novel metabolites, pMABG, is exclusively
microbially produced. For the smaller metabolite, 6-MPDA, though we were able to detect this in
the *in vitro* stable isotope study, we had difficulty quantifying it in our mouse experiments, in either
the blood or the feces. This may be due to its small molecular weight, suggesting that we need to
further optimize methods to quantify these small molecules. Another possibility is that it is
produced and immediately shuttled into host or microbial biosynthetic pathways. As for pMABG,
no living organisms have been reported to produce this compound. Interestingly, the folate
analog, p-aminobenzoylglutamic acid (pABG), is produced and even used by organisms as a
nutrient source[41]. Thus, pathways that produce folate-derived pABG may also act on MTX to
produce pMABG. The identity of these enzymes remain unknown; indeed, there are still many
unknowns with respect to folate metabolism[42]. Similar to the way in which MTX opened up a
"Golden Era" in the discovery of folate metabolism[43], microbial metabolites of MTX may uncover
novel pathways in microbes that are of fundamental importance. While the identity of these
microbial enzymes remains to be determined, our findings reveal that human gut microbes
produce novel MTX metabolites *in vitro* and *in vivo*, and that these compounds, which are
produced in the gut, can enter into circulation.
477
Finally, our work showcases the translational implications of basic science studies focused
on *in vitro* microbial drug metabolism[3,4] and suggests that microbiome modification can be
plausibly leveraged to improve drug efficacy *in vivo*. Taken together with our previous findings
that microbiome function and *ex vivo* metabolism is associated with clinical response to MTX[19],
the findings reported here in preclinical models provide strong evidence for the role of the
microbiome in shaping treatment outcomes for patients taking MTX.
484

Limitations

486
We found that SPF mice reared in two different campus facilities experienced different
responses to MTX, that response was transferrable by co-housing, and that GF colonized with
these different microbiotas showed different PK profiles as well as metabolite production. A
natural extension would be to evaluate MTX response in an arthritis experiment with GF colonized
mice with Zion- vs. Parnassus-microbiota. However, when we performed power calculations in
preparation for this experiment, we learned that it would be technically challenging and costly
because arthritis scores are lower in gnotobiotic mice for multiple reasons, leading to "floor
effects"[44,45]. For instance, it is well-known that gnotobiotic SKG mice experience significantly
reduced arthritis in the gnotobiotic setting[35,38]. Moreover, in the setting of MTX treatment, both
colonization arms (i.e., Zion vs. Parnassus) experience further reductions in arthritis scores;
detecting a difference between colonization states will require large numbers of gnotobiotic mice
(upwards of 30-40 per colonization group) and expensive isolators. While these logistical and
financial challenges may be overcome in future work requiring extensive optimization, the
preponderance of evidence presented in this and previous reports[19,20] serve to support the main
conclusions presented here. Another limitation is that while we were able to detect pMABG in the
blood and stool of mice and in *ex vivo* cultures of RA patients incubated with MTX, the extent to
which this metabolite is found in human circulation is unknown.

Conclusion

In conclusion, we have shown that the human gut microbiome metabolizes a key anti-inflammatory drug, MTX, and shapes the pharmacology of this drug *in vivo*. The microbiome directly metabolizes MTX and also affects host metabolic pathways that shape drug pharmacology. We identified novel metabolites of MTX, one of which likely is a major metabolite found in high abundance in the stool. This metabolism is associated with treatment response in a mouse model of inflammatory disease. Taken together with prior human studies[19,20], these findings provide a strong foundation for future microbiome-directed interventions to improve MTX outcomes. Further, we provide a framework to investigate additional drugs that undergo microbial metabolism and evaluate impacts *in vivo*. These findings support the notion that microbial enzymes that are beneficial to the host for nutrient harvest (e.g., folate metabolism) may also lead to unintended metabolism of therapeutic drugs and suggest that the metabolic capacity of the microbiome should be considered when personalizing therapies for patients.

Material and methods

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mouse experiments

Animal strains and care

GF and SPF C57BL/6J and BALB/c SKG mice (males and/or females, ages 6-13 weeks) were born at UCSF. SPF mice were kept in SPF conditions in the Animal Barrier Facility at UCSF, while GF mice were maintained within the UCSF Gnotobiotics Core Facility (gnotobiotics.ucsf.edu) and housed in gnotobiotic isolators for the duration of the experiment (Class Biologically Clean). The mice were housed at temperatures ranging from 19 °C to 26 °C and humidity ranging from 30 to 70% with a 12 h/12 h light/dark cycle. Mice had *ad libitum* access to water and standard chow diet (Lab Diet, cat # 5058, USA) in the Parnassus SPF facility and (Lab Diet, cat # 5053, USA) in the Mount Zion (MTZ) SPF facility, while standard autoclaved chow diet (Lab Diet, cat # 5021, USA) was used in the gnotobiotic facility.

Similarly, GF and SPF SKG mice were born at UCSF. GF SKG mice were rederived by Cesarean delivery into the University of California, San Francisco (UCSF) Gnotobiotic Core Facility and maintained in GF isolators for breeding. Stool pellets from isolators in the Gnotobiotic Core Facility were screened monthly by PCR for 16S bacterial and 18S (Internal transcribed spacer) fungal ribosomal sequences. Mice were randomly housed based on sex.

Several independent experiments were carried out and, for each experiment, mice were randomly assigned to groups. Key details for each mouse experiment can be found in **Supplementary Table 3**. All mouse experiments were approved by and conformed to the standards of the University of California San Francisco Institutional Animal Care and Use Committee.

Mice colonization

For humanization of GF mice with a human microbiome, mice were colonized with stool from treatment-naïve human donors with RA that were collected as part of a prior study[19]. Human stool was diluted 1:10 g/mL in prereduced sterile PBS + 0.05% cysteine and homogenized in an anaerobic chamber (Coy Laboratory Products) using pre-equilibrated reagents and supplies.

570 Insoluble material was separated from supernatant by centrifugation at 50 *g* for 1 min. Aliquots of
571 supernatant (100 µL per mouse) were gavaged into mice. Mice were colonized for at least
572 2 weeks before initiation of treatment with MTX.
573
For colonization of mice with murine microbiota, cecal contents from SPF BALB/c SKG mice
(either Zion or Parnassus) were harvested after euthanasia. The gastrointestinal tract was excised
and regions proximal and distal to the cecum were tied off with dental floss. This was brought into
an anaerobic chamber where the cecum was opened and the contents were placed into a conical.
Cecal contents were diluted 1:10 g/mL in prereduced sterile PBS + 0.05% cysteine and
homogenized in an anaerobic chamber using pre-equilibrated reagents and supplies. Insoluble
material was separated from supernatant by centrifugation at 50 *g* for 1 min. Aliquots of
supernatant (100 µL per mouse) were gavaged into GF BALB/c SKG mice. Mice were
colonized for at least 2 weeks before initiation of treatment with MTX.
583
584 ## Human subjects
585
Human subject samples were used for colonizing GF mice with human microbiota and also for *ex*
vivo profiling. These patients were enrolled either as part of a prior study[19] and as part of an
ongoing study NYU-UCSF collaboration. Consecutive patients from the New York University
Langone Medical Center's rheumatology clinics and offices were screened for the presence of
RA based on ACR criteria[46]. After informed consent was signed, each patient's medical history
(according to chart review and interview/questionnaire), diet, and medications were determined.
A screening musculoskeletal examination and laboratory assessments were also performed or
reviewed. All RA patients who met the study criteria were offered enrollment. The criteria for
inclusion in the study required that patients meet the American College of
Rheumatology/European League Against Rheumatism 2010 classification criteria for RA[46],
including seropositivity for rheumatoid factor and/or anti–citrullinated protein antibodies, and that
all subjects be age 18 years or older. New-onset RA was defined as disease duration of a
minimum of 6 weeks and up to 6 months since diagnosis, and absence of any treatment with
disease-modifying anti-rheumatic drugs, biologic therapy or steroids (ever). The exclusion criteria
applied to all groups were as follows: recent (<3 months prior) use of any antibiotic therapy,
current extreme diet (e.g., parenteral nutrition or macrobiotic diet), known inflammatory bowel
disease, known history of malignancy, current consumption of probiotics, any gastrointestinal tract
surgery leaving permanent residua (e.g., gastrectomy, bariatric surgery, colectomy), or significant
liver, renal, or peptic ulcer disease. This study was approved by the Institutional Review Board of
New York University School of Medicine protocols. All new onset rheumatoid arthritis patients
were recruited using established protocols from a previously described study[19]. Pre-treatment
stool samples were collected at baseline. Clinical and demographic data was de-identified and
recorded in RedCap by the designated study personnel.
609
610 ## METHOD DETAILS
611
612 ## Methotrexate administration and pharmacokinetic studies
613
For each experiment evaluating the PK profile of MTX, a single dose of 50 mg/kg (Pharmaceutical-
grade, Fresenius Kabi, NDC 63323-122-50, cat # 102250) was administered by oral gavage (PO),
except for the route of administration experiment where half of the mice were treated by
intraperitoneal injection (IP). MTX was dissolved at 50 mg/mL in a vehicle (VEH) containing 66%
of PBS 1X and 33% of NaOH 0.1 M. MTX solution was filtered through a 0.22 µm filter. Blood
samples were collected from the tail of awake freely moving mice at the following time points: 0
(before treatment), 15, 30, 60, 120, 240, and 440 min after treatment; for blood collection, we

used microhematocrit capillary tubes (Fisher #22-362-566). For some experiments, a 24-h time
point was also included. After centrifugation at 3,500 *g* for 15 min at 4 °C, plasma was collected.

In parallel to blood sampling, stool samples were collected whenever possible. Plasma and stool

samples were stored at −80 °C for targeted metabolomics (*see below*). From the concentration–
time curve obtained from each individual mouse, *max* and *auc* functions from the 'MESS'
package (v0.5.12) in R were used to calculate individual peak concentration and AUC values.
Of note, a value of 0 nmol/mg of dry feces was inferred for a metabolite when no stool sample
could be obtained from a mouse for the initial 0 min time point and all other mice showed a
value of 0.
630
631   **Targeted metabolomics for PK studies**
632
Plasma
Plasma quantification of MTX and its metabolites was performed by targeted metabolomic
analysis using liquid chromatography-tandem mass spectrometry (LC–MS). Samples were
analyzed using a SCIEX ExionLC autosampler (Shimadzu SIL-30AC series) in sequence with a
hybrid triple quadrupole – linear ion trap mass spectrometer (either SCIEX 6500 QTRAP or a
SCIEX 7500 QTRAP, depending on machine availability) (Shimadzu, USA). 5 µL of each sample
was diluted with 45 µL HPLC-grade H_2O and extracted with 200 µL of HPLC-grade methanol
(containing either aminopterin or 2-amino-3-bromo-5-methylbenzoic acid as internal standard).
Samples were incubated for at least 30 min at -20 °C before being centrifuged at 13,000 rpm for
10 min. 100 µL of supernatant was dried using a SpeedVac concentrator (Savant, cat # DNA 120)
and then reconstituted in 100 µL of HPLC-grade H_2O. To minimize fluctuations in pressure and
clogging, a filtering step was implemented using a 0.2 µm PVDF hydrophilic membrane filter plate
(Corning, cat# 3505) at 3,200 rpm for 5 min.
646
Feces
Feces quantification of MTX and its metabolites was performed similarly to plasma samples (using
the SCIEX 7500 QTRAP). Fecal samples were lyophilized; wet and dry weight was recorded
before and after lyophilization. Dried feces were extracted with 500 uL of extraction solvent
containing 80% methanol in H_2O and 0.1% formic acid (with 2-amino-3-bromo-5-methylbenzoic
acid as internal standard). Samples were homogenized using the Precellys 24-bead homogenizer
pre-cooled with dry ice at 6,400 rpm for 3 cycles of 20 s. Samples were centrifuged at 4 °C for 15
min at 16,000 rpm. 400 uL of supernatant was transferred into new tubes, and speed vacuumed
until completely dry. Dried samples were re-suspended in 200 uL of HPLC-grade H_2O and a 1:10
dilution was performed. To minimize fluctuations in pressure and clogging, a filtering step was
implemented using a 0.2 µm PVDF hydrophilic membrane filter plate (Corning, cat# 3505) at
3,200 rpm for 5 min.
659
MRM method development
Multiple reaction monitoring (MRM) was established for each compound using concentrated pure
standards directly on a SCIEX 7500 or 6500 Triple Quad Mass Spectrometer using a syringe
pump model. Intense peaks were used to identify the molecular fragmentation (Q3 mass), and
collision energy (CE) ramping was used to determine the optimal detection of the standards on
the instrument, with reported optimization values referenced in **Supplementary Table 4**.
666
Two different methods were used to quantify MTX and its metabolites. Method A was used initially
to quantify MTX, 7-OH-MTX and DAMPA. Method B was developed in order to quantify these as
well as the novel MTX metabolites: pMABG and 6-MPDA. 5 µL of reconstituted samples were

injected into the system. Analytes were chromatographically separated using either a Synergi 4
µm Fusion-RP 80 Å, 2 x 50 mm HPLC column (Phenomenex, cat# 00B-4424-B0) for method A,
or a Kinetex F5 2.6 µm, 2.1 × 150 mm LC column (Phenomenex, cat # 00F-4723-AN) for method
B.

In method A, a mobile phase scheme consisting of [A] H_2O + 0.1% formic acid and [B] methanol
+ 0.1% formic acid was used. LC flow was set to a constant rate of 600 µL/min, and the timed
linear gradient program consisted of: time = 0 min, 5% B, time = 1 min, 5% B, time = 5 min, 40%
B, time = 5.25 min, 100% B, time = 5.75 min, 100% B, time = 6 min, 5% B, and time = 8 min, 5%
B. Data was collected in positive polarity mode using MRM with the following source parameters:
curtain gas (CUR) = 35 psi; nebulizer gas (GS1) = 50 psi; heater gas (GS2) = 70 psi; collision gas
= 9; source temperature (TEM) = 450 °C; ion spray voltage (ISV) = 1,500 V. Other method
parameters used include a cycle time of 0.525 s, dwell time of 100 ms/MRM pair, settling time =
0 ms, and pause time of 5.007 ms.

In method B, a mobile phase scheme consisting of [A] H_2O + 0.1% formic acid and [B] acetonitrile
+ 0.1% formic acid was used. LC flow was set to a constant rate of 200 µL/min, and the timed
linear gradient program consisted of: time = 0 min, 0.2% B, time = 1 min, 0.2% B, time = 2 min,
10% B, time = 4 min, 25% B, time = 6 min, 98% B, time = 8 min, 98% B, time = 8.1 min, 0.2% B,
and time = 10 min, 0.2% B. Data was collected in positive and negative polarity modes using
MRM with the following source parameters: CUR = 35 psi; GS1 = 50 psi; GS2 = 70 psi; collision
gas = 9; Temperature = 450 °C; ISV = 2,200 V (positive) and -2,200 V (negative). Other method
parameters used include a cycle time of 0.435 s, dwell time of 10 ms/MRM pair, settling time =
15 ms, and pause time of 5.007 ms.

Raw data was processed using a built-in SCIEX OS software package (version 2.1.6.59781) for
peak picking, alignment, and quantitation. Peak areas of the standard curves were converted to
concentrations at ng/mL using linear regression, then expressed as nM (plasma samples) or
nmol/mg of dry feces (stool samples).

Stable isotope profiling
Labeled methotrexate was purchase (d3-MTX, Santa Cruz # sc-218707) along with unlabeled
MTX. Stock concentrations were dissolved in DMSO at a concentration of 90 mg/mL. Labeled
MTX, unlabeled MTX, or DMSO was further diluted to achieve a final concentration of 100 ug/mL
into the experimental plate. For each of *Bacteroides thetaiotaomicron* DSM 2079, *Clostridium*
asparagiforme DSM 15981, and *Clostridium symbiosum* DSM 934, a single colony of each isolate
was subcultured in Bacto Brain Heart Infusion media (BD Biosciences, 37 g/L) supplemented
with L-cysteine-HCl (0.05%, w/v), menadione (1 µg/mL), and hemin (5 µg/mL) (referred to
hereafter as BHI+) for 48 h in an anaerobic chamber at 37 °C with an atmosphere composed of
2-3% H2, 20% CO2, and the balance N2. This subculture was diluted down to an OD600 of ~0.1,
which was then further diluted 100-fold, and then used to inoculate the experimental plate
containing the drug. Sterile controls were also tested for each drug. After 72 h, 100 uL aliquots
were centrifuged at 1,900 rcf for 5 min at room temperature, and 40 uL of the supernatant was
submitted for global LC-MS profiling. Each drug-bacterial species pair and sterile control was
tested in quadruplicate.

Global profiling of isotope labeled *in vitro* monocultures
Samples were thawed and combined with 80 µL of LC-MS-grade methanol: acetonitrile (1:1 v/v
and 1µM chlorpropamide) with approximately 10 µL of 0.1 mm zirconia beads. Samples were then
thoroughly homogenized in BeadBlaster 24 Microtube Homogenizer 2 x 30 s at 4,260 rpm.

Samples were incubated at -20 °C for 1 h. Then samples were vortexed and centrifuged at 12,000
g at 4 °C for 15 min. Supernatant was transferred to autosampler vials for LC-MS analysis.
722
Samples (5uL) were separated by reverse phase HPLC using a Prominence 20 UFLCXR system
(Shimadzu, Columbia, MD) with a Waters (Milford, MA) BEH C18 column (100 mm x 2.1 mm 1.7
um particle size) maintained at 55 °C and using a 20 min aqueous acetonitrile gradient, at a flow
rate of 250 uL/min. Solvent A was HPLC-grade H_2O with 0.1% formic acid and Solvent B was
HPLC-grade acetonitrile with 0.1% formic acid. The initial conditions were 97% A and 3% B,
increasing to 45% B at 10 min, 75% B at 12 min where it was held at 75% B until 17.5 min before
returning to the initial conditions. The eluate was delivered into a 5600 (QTOF) TripleTOF using
a Duospray™ ion source (all Sciex, Framingham, MA).The capillary voltage was set at 5.5 kV in
positive ion mode with a declustering potential of 80 V. The mass spectrometer was operated with
a 100 ms TOF scan from 50 to 1000 *m/z*, and 16 MS/MS product ion scans (100 ms) per duty
cycle using a collision energy of 50 V with a 20 V spread.
734
RA patient fecal sample *ex vivo* profiling
All work was carried out in an anaerobic chamber . For each patient, stool was aliquoted into a
pre-equilibrated cryovial, diluted in pre-reduced PBS at 10 mL/g of stool, and vortexed to
homogenize the sample. The sample was spun at ~20 *g* for 1 min on a mini-centrifuge to facilitate
settling of sediments, and the sediment-free supernatant ("fecal slurry") was then aliquoted into a
new pre-equilibrated cryovial for evaluation of *ex vivo* growth. Growth was evaluated by
inoculating liquid BHI+ with 1:50 dilution of this fecal slurry. Samples were treated with MTX
100 μg/mL or the vehicle control and incubated with shaking for 24 h at 37 °C, and then stored at
-80 °C for future profiling. For LC-MS profiling, samples were thawed, centrifuged at 4 °C at 13,000
g for 5 min. 200 μL of supernatant was mixed with 800 μL of LC-MS-grade methanol: acetonitrile
(1:1 v/v) and incubated at -20 °C for 1 h. Samples were vortexed, and then centrifuged at 15,000
g for 15 min at 4 °C. Supernatant was transferred to autosampler vials for targeted metabolomics.
747
Ex vivo **fecal sample targeted metabolomics**
The analysis was performed using a Nexera 20 HPLC system (Shimadzu) coupled to a TripleTOF
5600 mass spectrometer (Sciex) equipped with a DuoSpray ion source, operating in electrospray
ionization (ESI) mode with positive ion detection. Chromatographic separation was achieved on
a Waters XSelect HSS T3 column (2.1 × 100 mm, 2.5 μm particle size) maintained at 40.0 °C.
The mobile phase was delivered at a flow rate of 0.200 mL/min. Solvent A consisted of LC/MS-
grade water with 0.1% formic acid, and Solvent B was LC/MS-grade acetonitrile containing 0.1%
formic acid. The gradient elution program was as follows: 0.00-5.00 min, 5.0% B; 9.00 min, 90.0%
B; 14.00 min, 100.0% B; 14.50–20.00 min, 5.0% B. The injection volume was 2.0 μL. Mass
spectrometric data (MS1 and MS2) were acquired using a declustering potential (DP) of 70 V.
MS1 data were collected over a mass range of 100–1000 Da. The ion source parameters were
set as follows: ion spray voltage (IS), 5500 V; curtain gas (CUR), 30 psi; nebulizer gas (GS1),
60 psi; heater gas (GS2), 50 psi; and source temperature (TEM), 500 °C. High-resolution multiple
reaction monitoring (MRM) was employed with a total cycle time of 500 ms and unit resolution in
Q1. The following precursor ions and their respective collision energies (CE) were used for MS2
analysis:

● pMABG (m/z 281.13, CE 55 eV, 8.50 min)
● MTX (m/z 455.20, CE 50 eV, RT 8.35 min)
● DAMPA (m/z 326.15, CE 34 eV, RT 8.60 min)

Data were analyzed using MultiQuant software (Sciex, v3.0.3) with the MQ4 quantitation algorithm. Quantification was performed against a standard curve of authentic standards at nine concentrations ranging from 1 nM to 1 μM.

The following fragment ions were used for quantitation:

- MTX: m/z 308.13
- DAMPA: m/z 176.08
- pMABG: m/z 134.06

Masses were integrated with a 10 mDa tolerance.

Isolation of microbes from patient stool samples

Bacterial isolates were harvested from MTX-naïve RA patient fecal slurries prepared as previously described[19].For isolate harvesting, fecal slurries were diluted 1:20 in BHI+ media then serially diluted 1:10 in BHI+ media 7 times. Serial dilutions were streaked on BHI+ agar plates. To enrich for *Firmicutes*, another set of serial dilutions were exposed to aerobic conditions and 70% (v/v) ethanol for 4 h prior to streaking on BHI+ agar plates supplemented with taurocholate as previously described[34]. Isolates were grown at 37 °C for 48 h in an anaerobic chamber. Unique colony morphologies were identified and picked for growth in liquid BHI+ media at 37° C for 48 h before reconstitution in 2:1 of 80% (v/v) glycerol and placement in -80 °C for long-term storage.

Bacterial DNA extraction and full-length 16S rRNA sequencing

Full-length Sanger sequencing of the 16S rRNA gene was performed by either extracting DNA from turbid, 48-h cultures; verification was done by colony PCR. *Liquid cultures*: For each isolate, bacterial liquid cultures (2 mL) incubated for 48 h in an anaerobic chamber were pelleted by centrifugation for 5 min at 2,000 rcf and the supernatant was discarded. The bacterial pellet was resuspended in 200 uL Instagene Matrix (Cat # 732-6030) and incubated at 56 °C and 100 °C for 30 min and 8 min respectively. Samples were vortexed before and after each incubation. *Colony PCR*: After streaking glycerol stocks and incubating for 48 h at 37 °C under anaerobic conditions, an individual colony was picked and placed in a PCR strip-tube containing PCR master mix. The PCR master mix consisted of 25 uL Amplitaq gold (Life Tech/Thermo Fisher 438881), 1 uL forward primer (8F, 5`-AGAGTTTGATCCTGGCTCAG-3'), 1 uL reverse primer (1542R, 5′-AAGGAGGTGATCCAGCCGCA-3'), and 18 uL molecular grade H_2O per reaction. PCR amplification was as follows: 10 min at 95 °C, 29x (30 s at 95 °C, 30 s at 50 °C, 30 s at 72 °C), and ending at 4 °C.

PCR purification was achieved by using a PureLink PCR Purification Kit (Invitrogen K3100-01) or Axygen AxyPrep Mag kit (#MAG-PCR-CL-5) according to the manufacturer's instructions. Purified DNA was quantified using Nanodrop (Thermo Scientific NanoDrop 2000 Spectrophotometer). In PCR strip-tubes, 10 uL of the mix template and 5 uL of each primer (either F or R) was mixed and submitted to GENEWIZ for Sanger Sequencing. Forward and reverse reads were combined into a contig that was analyzed using BLAST (BLAST: Basic Local Alignment Search Tool (nih.gov)) where the highest "percent identity" was chosen for species identification.

MTX metabolism screen

RA-associated strains were grown for 48 h in an anaerobic chamber at 37 °C and were subcultured at a 1:50 dilution (2 uL of bacterial culture in 98 uL culture media) in a 96-well U bottom plate. Bacteria were grown in BHI+ with MTX 50 ug/mL or vehicle control (DMSO). After 72 h of incubation at 37 °C, the plates were placed in the -80 °C freezer until extraction.

HPLC-based quantification of MTX levels of *in vitro* studies

816 For bacterial samples, plates were spun down at 2,400 rpm 10 min to pellet cells. 30 uL of the
817 bacterial supernatant was extracted with 150 uL of HPLC-grade methanol. The samples were
818 incubated at -20 °C for 30 min. After incubation, the plates were centrifuged at 3,000 rpm for 30
819 min. After centrifugation, 60 uL of supernatant from each well was transferred into 200 uL plastic
820 inserts in HPLC glass vials. HPLC-grade methanol was used as blank.
821
The samples were profiled by reversed phase HPLC using an Agilent 1260 Infinity II system with
a Kinetex 2.6 um C18 column and a 8 min gradient, at a flow rate of 0.6 mL/min. For the mobile
phase, solvent A was 0.1% formic acid and solvent B was 100% HPLC-grade methanol. The initial
condition was 90% B, decreasing to 70% B by 1 min, 0% B by 7 min, and 90% B by 7.5 min where
it was held until 8 min. Authentic standards of MTX were used for peak identification and a
standard curve was generated for quantification (100, 50, 25, 12.5, 6.25 ug/mL). When MTX
depletion is reported (usually as a percentage), we normalize each sample to sterile controls on
the same plate. Occasionally, we noted MTX percentages > 100%, which we hypothesized could
be due to evaporation effects.
831
Validation of MTX metabolism for bacterial strains *in vitro*
Glycerol stocks from harvested RA-associated strains were streaked on BHI+ agar plated in an
anaerobic chamber. A single colony was subcultured in BHI+ liquid media for 48 h at 37 °C. This
subculture was diluted down to an OD600 of 0.08 - 0.1, which was then further diluted 100-fold,
and then exposed to MTX at a final concentration of either 50 ug/ml in quadruplicate along with
vehicle controls (DMSO). Plates were incubated at 37 °C for 72 h. An Eon Microplate
Spectrophotometer (BioTek Instruments, Inc) was used to measure OD600 at 0 and 72 h to
confirm bacterial growth. Data was exported using the Gen5 (Version 3.12) software. After 72 h,
aliquots were prepared for extraction and profiling on HPLC (*see above,* "HPLC-based
quantification of MTX levels of *in vitro* studies") and LC-MS platforms.
842
LC-MS analysis of *in vitro* isolates
A single sample of each bacterial species incubated with MTX (50 - 100 ug/ml) or sterile control
was selected for profiling on LC-MS. From each sample, 60 uL of supernatant was transferred
and stored at -20 °C. Based on MTX depletion and detection of novel peaks on HPLC, a sample
from each bacterial species was selected: 60 uL was dried and reconstituted in 1 mL HPLC-grade
H_2O. The samples were filtered using a 0.2 µm PVDF hydrophilic membrane filter plate (Corning,
cat# 3505) at 3,200 rpm for 5 min. 100 uL of the samples were transferred into glass inserts and
vials for LC-MS profiling. LC-MS targeted metabolomics (*see above,* "Targeted metabolomics for
PK studies" method B) was used for quantifying MTX and its metabolites with a standard curve
of authentic standards at concentrations ranging from 0.2 to 200 ng/mL .
853
SKG arthritis experiments
To evaluate the therapeutic effect of MTX on an arthritis mouse model, BALB/c SKG mice were
given once 100 uL of zymosan emulsion (10 mg/mL) through IP injection to induce arthritis.
Zymosan (Sigma, #Z4250-1G) was emulsified in sterile 1X PBS. Mice were either treated with
MTX (10 mg/kg of body weight, 3 times a week) or the vehicle control (VEH). MTX was dissolved
at 2 mg/mL in a vehicle (VEH) containing 66% of PBS 1X and 33% of NaOH 0.1 M. MTX solution
was filtered using a 0.22 µm filter and stored briefly in -20 °C until use.
861
Three times a week, starting the day of zymosan injection, mice were gavaged with MTX or VEH,
and clinical arthritis scores (CAS) were evaluated. Experimenters evaluating CAS were blinded
to the treatment. The wrist, ankle, and digits (4 digits on each upper limb and 5 digits on each
lower limb) were scored. Joint swelling was monitored by inspection and scored as follows: 0, no
joint swelling; 0.1,swelling of one finger joint; 0.5, mild swelling of wrist or ankle; 1.0, severe

swelling of wrist or ankle. Scores for all fingers of forepaws and hindpaws, wrists and ankles were
summed for each mouse. For example, an ankle that maintained its natural, angular shape was
given a 0, an ankle with a round presentation was given a 0.5, and an ankle that presented a
horseshoe shape due to severe swelling was given a 1. Similarly, if the wrist maintained its
natural, slim appearance, it was given a 0, while some swelling was given a 0.5, and severe
swelling that caused the wrist to bulge was given a 1. Each digit of fore paws and hind paws were
scored as either 0 or 0.1 for no inflammation or inflammation, respectively. Each day, the score
of each mouse's digits, ankles, and wrists were added up and recorded as the CAS. The
maximum CAS that a mouse could receive was 5.8 (1.5 for each lower limb plus 1.4 for each
upper limb). CAS were collected over the course of the experiment and analyzed over time and
stratified by microbiome or treatment. Additionally, individual AUCof CAS curves were calculated
using the *auc* function from the MESS package (v0.5.12) in R.
879
16S-seq of SKG mouse gut microbiota
Broadly, aliquots of mouse fecal samples were homogenized using a bead-beating (Mini-
Beadbeater-24, BioSpec) method followed by DNA extraction and purification as previously
described[47]. DNA was extracted using ZymoBIOMICS 96 well MagBead ZymoBIOMICS 96
MagBead DNA Kit (Cat#D4302) as per the manufacturer's protocol. GoLay-barcoded 515F/806R
primers[48] were used to carry out 16S rRNA gene PCR according to the methods of the Earth
Microbiome Project (earthmicrobiome.org). The following were combined: 2 µL of DNA, 25 µL of
AmpliTaq Gold 360 Master Mix (Life Technologies), 5 µL of primers (2 µM each GoLay-barcoded
515/806R), and 18 µL H_2O. Amplification was as follows: 10 min at 95 °C, 25x (30 s at 95 °C, 30
s at 50 °C, 30 s at 72 °C), and 7 min at 72 °C. For the dose-response gnotobiotic experiment,
amplicons were quantified with PicoGreen (Quant-It dsDNA; Life Technologies) and pooled at
equimolar concentrations. Libraries were quantified (NEBNext Library Quantification Kit; New
England Biolabs) and sequenced with a 600 cycle MiSeq Reagent Kit (251x151; Illumina) with
~10% PhiX. For the remaining sequencing experiments, samples underwent primary PCR for
amplification and secondary PCR to add flow cell adaptors and indices as previously described[49].
DNA was normalized to equal concentrations based on PicoGreen quantification and pooled.
Pooled libraries were purified and concentrated with MinElute PCR Purification kit (Qiagen
#28004), run on 1% gel, size-selected and purified using MinElute Gel Extraction kits (Qiagen,
#28604). Pooled libraries were run at the Chan Zuckerberg Biohub using Illumina MiSeq platform.

QUANTIFICATION AND STATISTICAL ANALYSIS

General

Power calculations were performed for PK and arthritis experiments whenever pilot data were available. All statistical analyses were performed using the R environment. "N.S." indicates "not significant." In general, Student's t-tests were used to infer significance ($p \leq 0.05$) unless otherwise indicated in the figure legend. Whenever provided, boxplots depict the median, hinges represent the first and third quartiles, and whiskers show the range. Whenever provided, error bars represent the standard error of the mean (SEM).

16S rRNA amplicon analysis of mouse samples

Reads that were not already demultiplexed were demultiplexed using QIIME[50] v1.9.1 (split_libraries_fastq.py). QIIME2[51] (v 2020.2) was used to trim reads, denoise the data, and create a feature table using the following commands: *qiime cutadapt trim-paired*, *qiime dada2 denoise-paired*, *qiime feature-table filter-samples*. Within QIIME2, taxonomy was assigned using the DADA2[52] implementation of the RDP Naïve Bayesian classifier[53] using the DADA2-formatted SILVA v128 training set; taxonomic assignments to the species level were chosen by exact match of the ASV to the reference database, and where multiple matches were present, all are reported. For figures and in the text, genus and species assignments are reported when available, and when neither was available, we included the family taxonomic assignment; we also report a unique ASV identifier. This ASV identifier can be used to look up the full taxonomic assignment from kingdom to species and the associated sequence variant. The QIIME2R package was used to import QIIME2 artefacts into R (https://github.com/jbisanz/qiime2R). Diversity metrics were generated using Vegan (v2.5-6) and Phyloseq (v1.30.0)[54], with principal coordinate analysis (PCoA) or principal components analysis (PCA) carried out with Ape (v5.3) or Vegan, respectively. Analyses were carried out on either: (1) centered log2-ratio (clr) normalized taxonomic abundances calculated as $A_{clr}=[\log_2(A_1/g_a), \log_2(A_2/g_a),\ldots \log_2(A_n/g_a),]$, where A is a vector of non-zero read counts and g_a is the geometric mean of all values of A, or (2) relative abundance calculated as proportion of reads. ANOSIM and PERMANOVA were used to detect changes in community composition using counts from rarefied data and Bray-Curtis distances. DESeq2 (v1.26.0)[55] was used to determine differentially abundant taxa on raw count data. For each sample, fastq files are available in NCBI's Sequence Read Archive (SRA), accession number [*to be determined upon publication*].

Author Contributions

Conceptualization: RRN. Funding acquisition: JUS, PJT, RRN. Investigation: VSVS, ALD, CMH, DAO, ERR, MM, NP, SY, JA, RRN. Methodology: ALD, DAO, MM, ERR, ADP, JGB, DSD. Project administration: RRN. Resources: RBB, JUS. Supervision: JA, JUS, ADP, PJT, RRN. Validation: VSVS, ALD, CMH, RRN. Visualization: VSVS, ALD, CMH, DAO, RRN. Writing – original draft: RRN. Writing – review & editing: all authors.

Acknowledgements

This work was supported by the following: A.D. was supported by a Postdoctoral Fellowship from the Belgian American Educational Foundation (B.A.E.F.); D.A.O. was supported by: 3K08AR073930-05S1 (NIAMS/OD) and ImmunoDiverse SRA Fellowship. R.B.B. was supported by NIH/NCATS UL1TR001445. E.R.R was supported by PSU/NIDDK (T32DK120509). This was work was supported by multiple grants: 2R01HL122593 (P.J.T), 1R01AR074500 (J.U.S and P.J.T), UCSF Breakthrough Program for Rheumatoid Arthritis-related Research (BPRAR; partially funded by the Sandler Foundation; P.J.T), Arthritis Foundation Center of Excellence (R.R.N); K08AR073930 (R.R.N); R03AR082036 (R.R.N.), I01CX002557 (R.R.N.), R35GM151349

(R.R.N.), Russell/Engleman Rheumatology Research Center (R.R.N.), and the Arthritis National
Research Foundation (R.R.N.), Benioff Center for Microbiome Medicine (R.R.N. and P.J.T).

We acknowledge support from the following core facilities: UCSF QMAC, UCSF Gnotobiotics
Core, PSU Metabolomics Core Facility at The Huck Institutes of the Life SciencesPenn State
University (Sergei Koshkin).

We would like to thank William Seaman and members of the Nayak Lab for their helpful comments
and suggestions on the manuscript.

Conflict of Interest Statement
The authors have no conflict of interest relevant to this work.

Supplementary Table 1. RA-associated human gut strains that metabolize MTX.

#	Species	HPLC quantification (Student's t-test as compared to DMSO)	Metabolite produced	Growth in MTX (relative to DMSO)	Confirmed?
1	*Paeniclostridium sordellii*	*$p < 0.0001$	pMABG	81%	Confirmed
2	*Terrisporobacter petrolearius*	*$p = 0.0003$	pMABG	86%	Confirmed
3	*Intestinibacter bartlettii*	*$p < 0.0001$	pMABG	138%	Confirmed
4	*Romboutsia hominis*	*$p < 0.0001$	pMABG	189%	Confirmed
5	*Clostridium perfringens*	*$p < 0.0001$	pMABG	71%	Confirmed
6	*Clostridium tertium*	*$p < 0.0001$	pMABG	44%	Confirmed
7a	*Clostridium cibarium RRN216**	N.S.	pMABG (strain-specific)	3%	Confirmed
7b	*Clostridium cibarium RRN216**	N.S.	None detected	14%	N/A
8	*Hungatella hominis*	*$p = 0.019$	None detected	26%	Confirmed
9	*Coprobacillus cateniformis*	$p = 0.052$	pMABG	46%	Confirmed
10	*Clostridium innocuum*	*$p < 0.0001$	pMABG	54%	Confirmed
11	*Enterococcus faecalis*	N.S.	Small mount pMABG	70%	Limited by growth
12	*Enterococcus lactis*	N.S.	None detected	107%	Not confirmed

963
964   *** RRN number indicates the unique** identifier **of the patient.**

965   **Supplementary Table 2. Differentially abundant amplicon sequence variants (ASVs) in**
966   **Zion vs. Parnassus mice (not co-housed).**
967
968   *See Excel sheet.*

969 **Supplementary Table 3. Details of mouse experiments.**

#	Phenotype	Description	#mice/ group	Strain	Microbiome	Dose	Route	Sex	Age at time of treatment	Colonization time prior to treatment	Duration of drug treatment
1	PK	GF vs. SPF	6	C57BL/6J	GF	50 mg/kg	PO	F	6-7 weeks old	not relevant	1 single dose of MTX
			8		SPF						
2	PK	IP vs. PO	3	C57BL/6J	SPF	50 mg/kg	PO	M	8 weeks old	not relevant	1 single dose of MTX
			3				IP				
3	PK	Patient 170 vs. Patient 317	11	C57BL/6J	Humanized with Patient 170	50 mg/kg	PO	M	9-13 weeks old	2 weeks	1 single dose of MTX
			11		Humanized with Patient 317						
4	Arthritis (#1)	vehicle (VEH) vs. MTX, co-housed Parn. & Zion	8	BALB/c SKG	SPF from PSB or MTZ	(VEH)	PO	F	10-13 weeks old	2 weeks of co-housing	MTX 3 times a week
			8		SPF from PSB or MTZ	10 mg/kg					
5	Arthritis (#2)	Parn. origin vs Zion origin (not co-housed)	6	BALB/c SKG	SPF from PSB	10 mg/kg	PO	F	11-12 weeks old	not relevant (no co-housing)	MTX 3 times a week
			3		SPF from MTZ						
6	PK	Parn. vs. Zion	5	BALB/c SKG	GF colonized with Parn. microbiota	50 mg/kg	PO	F	13-24 weeks old	2 weeks	1 single dose of MTX
			6		GF colonized with Zion microbiota						

970
971 F: female; GF: germ-free, IP: intraperitoneal; M: male; MTX: methotrexate; Parn.: Parnassus;
972 PO: per os (= oral); PK: pharmacokinetic; SPF: specific-pathogen-free; Zion: Mount Zion.

973 **Supplementary Table 4. Details of LC-MS targeted metabolomics for *in vitro* and *in vivo***
974 **experiments.**

Compound	Q1 mass	Q3 mass	EP	CE	CXP	Mode	LC-MS/MS method
methotrexate (MTX)	455.2	308.2	10	20	14	Positive	A
				29	15	Positive	B
methotrexate diglutamate (MTXPG2)	584.2	308.2	10	20	14	Positive	A
				36	15	Positive	B
deoxyaminopteroic acid (DAMPA)	326.2	175.1	10	20	14	Positive	A
				26	15	Positive	B
7-hydroxy-methotrexate (7-OH-MTX)	471.2	324.1	10	15	14	Positive	A
				25	15	Positive	B
6-methylpteridine-2,4-diamine (6-MPDA)	177.0	160.0	10	18	14	Positive	A
	177.0	97.8	10	21	15	Positive	B
aminopterin (APT, *IS1*)	441.2	294.0	10	20	14	Positive	A
				30	15	Positive	B
2-amino-3-bromo-5-methylbenzoic acid (2A3Br5MBA, *IS2*)	231.0	104.0	10	25	15	Positive	B
p-methylaminobenzoylglutamic acid (pMABG)	278.9	150.2	-10	-25	-15	Negative	B

975
976 CE: collision energy; CXP: collision cell exit potential; EP: entrance potential; IS: internal standard.

977
Figure 1. The gut microbiome impacts MTX PK *in vivo*. (**A**) Structures of methotrexate (MTX)
and its known metabolites, 7-hydroxy-MTX (7-OH-MTX), deoxyaminopteroic acid (DAMPA), and
polyglutamated MTX (MTXPG$_n$). (**B**) Plasma concentrations of MTX and its known metabolites
were quantified by LC-MS in germ-free (GF, N=6) or colonized (SPF, N=8) C57BL/6J female mice

after a one-time oral administration of MTX 50 mg/kg. Multiple plasma samples were taken from
each mouse over an 8 h time course. (**C**) Peak plasma concentration, and (**D**) Area Under the
Curve (AUC) were compared between GF and SPF mice. (**E**) Ratio of AUC of 7-OH-MTX to MTX
in the plasma over 8 h. (**F**) Stool concentrations of MTX and its metabolites were quantified at
multiple time points. (**G**) Peak stool concentrations, (**H**) AUC and (**I**) ratio of AUC of DAMPA to
MTX are shown. Results from Student's t-test (**B-I**) are reported. Error bars are SEM.

Supplementary Figure 1, related to Figure 1. Experimental design and polyglutamated MTX. (**A**) Experimental design of a pharmacokinetic (PK) study with germ-free (GF, N=6) and colonized (SPF, N=8) mice. (**B**) MTX-PG$_2$ levels in plasma.

993
Supplementary Figure 2. Parenteral administration of MTX does not preclude microbial
metabolism of MTX into DAMPA. (**A**) Plasma concentration of MTX and its known metabolites
were quantified by LC-MS in SPF C57BL/6J male mice after a one-time administration of MTX 50
mg/kg, given either by intraperitoneal injection (IP, N=3) or orally (PO, N=3). Multiple plasma
samples were taken from each mouse over an 8 h time course. (**B**) Peak concentration and (**C**)
AUC were compared between IP vs. PO treated mice. (**D**) Ratio of AUC of 7-OH-MTX to MTX in
the plasma over 8 h. Results from Student's t-test are reported. Error bars are SEM.

1001
Figure 2. Interindividual variation of the human gut microbiome contributes to MTX PK ***in***
vivo. (**A**) Experimental design of PK study in GF C57BL/6J male mice colonized with donor
microbiota from two RA patients (Patient 170 and Patient 317; N=11 per colonization group) given
a one-time oral administration of MTX 50 mg/kg. (**B**) Plasma concentration of MTX and its known
metabolites were quantified by LC-MS. Multiple plasma samples were taken from each mouse
over a 24 h time course (first 8 h depicted for clarity). (**C**) Peak concentration and (**D**) AUC were
compared between mice colonized with gut microbiomes from Patient 170 or Patient 317. (**E**)
Ratio of AUC of 7-OH-MTX to MTX in the plasma over 24 h. Results from Student's t-test are
reported. Error bars are SEM.

1011
Figure 3. **Human gut microbes produce novel MTX metabolites**. (**A**) MTX and DAMPA peak
intensities (peak analyte areas) in the supernatant of three type strains incubated with MTX over
63 h. (**B**) Stable isotope profiling of three type strains incubated with MTX or d_3-labeled MTX
("MTX-L") for 72 h (N=4 replicates per species per compound). Feature peak intensities that
correspond to labeled and unlabeled metabolites are depicted in the bar plot. Intensity is
normalized to the mean (average of all samples) and relative intensity is reported. The mass-to-
charge (*m/z*) value of each peak is listed on top, and the treatments are denoted on the right-
hand side. (**C**) Chemical structures of the 2 novel microbiota-derived metabolites of MTX identified
by stable isotope labeling experiments (upper panel). The purple shading indicates the location
of the deuterium label. Novel metabolites were confirmed using authentic standards: mirror plots
comparing the sample (red) with the authentic standard (blue) are shown (lower panel). (**D**)
Quantification of MTX and its microbial metabolites from the supernatant of multiple human gut
bacterial isolates incubated with MTX (100 ug/mL) for 72 h *in vitro*. Dashed grey lines represent
the amount of compound in sterile control wells. St, sterile; *Bt*, *Bacteroides thetaiotaomicron*; *Bp*,
Blautia producta; *Ll*, *Lactonifactor longoviformis*; *Ee*, *Eubacterium eligens*; *Cs*, *Clostridium
symbiosum*; *Csy*, *Clostridium symbiosum*; *Csp*, *Clostridium sporogenes*; *Ci*, *Clostridium
innocuum*; *Ca*, *Clostridium asparagiforme*.

1029

1030
Figure 4. **Novel metabolites are produced by human gut microbiota *ex vivo* and by murine**
microbiota *in vivo*. (**A**) Microbial communities from treatment-naïve RA patients produce novel
MTX metabolites *ex vivo* to variable extents. Fecal slurries were prepared in pre-reduced saline
and incubated with 100 mg/mL of MTX or vehicle control for 24 h in an anaerobic chamber.
"Sterile" refers to sterile media with MTX. (**B**) As in Figure 1, fecal samples from GF (N=6) or
colonized (SPF, N=8) C57BL/6J female mice were profiled on LCMS to quantify the novel
metabolites (6-MPDA unable to be quantified). Results from Student's t-test are reported. Error
bars are SEM.

1039
Figure 5. **Multiple human gut bacterial strains isolated from treatment-naïve RA patients**
can metabolize MTX *in vitro*. (**A**) Results of our MTX metabolism screen showing the phylum

and species identification (left), the number of strains per species tested (middle), and the
proportion of MTX remaining in the supernatant relative to sterile controls (right). The RA patient
donor of each strain is also shown (indicated by different color circles). Strains that depleted MTX
by at least 60% (red dashed line) were scored as "metabolizers" and are highlighted in orange.
(**B**) Validation of MTX depletion by each human isolate. Each isolate was incubated with MTX
100 ug/mL for 72 h (N=4 replicates per species) and MTX was quantified by HPLC. RA-associated
species that significantly deplete MTX are highlighted in orange. (**C**) LC-MS quantification of
pMABG in the same bacterial supernatant as in (**B**). For (**B**) and (**C**), the dashed line indicates
abundance of the compound in sterile controls. The phylum is indicated by the color of the bar.
Results of Student's t-test are reported.

1052
Supplementary Figure 3, related to Figure 5. **Strain harvesting workflow for identifying RA-**
associated microbial strains that metabolize MTX. (**Top panel, Steps 1 through 4**) Fecal
slurries from RA patients are streaked out at multiple dilutes on rich media (BHI+), and media
supplemented with taurocholate, a bile acid which induces spores to germinate. This latter step
was devised to enrich for *Firmicutes*, which we found are enriched for MTX metabolizing species.
(**Middle panel**) We picked 24 colonies with different morphologies per patient and created
glycerol stocks. These glycerol stocks underwent 16S full-length Sanger sequencing in order to
identify bacterial species (Step 5 and 6A). (**Bottom panel**) In Step 5, we also inoculated harvested
isolates in either MTX 50 ug/mL or vehicle control (DMSO). We quantified growth to ensure that
microbes grew at this concentration , incubating samples for 72 h. Cell pellets were spun down,
and supernatant was extracted in methanol and profiled by HPLC to quantify the amount of MTX
remaining compared to sterile controls (Step 6B). Figure generated with BioRender.

1065
Supplementary Figure 4, related to Figure 5. Many RA-associated bacterial species
metabolize MTX into known and novel metabolites *in vitro*. (A) Flow chart of RA-associated
microbes that were harvested and screened for MTX metabolism. (**B**) Unique species identified
by full-length 16S Sanger sequencing. The contribution of each treatment-naïve RA patient donor
is indicated by the color of the bar. (**C**) LC-MS quantification of MTX and its metabolites in the
supernatant of RA-associated isolates incubated with MTX for 72 h (N=1 bacterial sample). The
phylum is indicated by the color of the bar. Type strains of *Bacteroides* species (green) were used
as negative controls.

1074
Figure 6. MTX response in a model of inflammatory arthritis is associated with the
microbiome and is transferable. (**A**) Clinical arthritis scores (CAS) of female SKG mice treated
with vehicle (VEH, N=8) or oral MTX 10 mg/kg (N=8) three times weekly. (**B**) AUC of CAS (from
day 0 to day 21). (**C**) CAS over time and (**D**) AUC of CAS stratified by treatment (VEH vs. MTX)
and facility-of-origin (Zion vs. Parnassus). (**E**) CAS of Zion (N=3) and Parnassus (N=6) female
SKG mice treated with MTX three times weekly in the absence of co-housing. (**F**) Comparison of
CAS in two experiments of mice treated with MTX over a 14-day time frame. (**G**) Comparison of
CAS AUC in two experiments. Results of Student's t-test are reported. Error bars are SEM.

Figure 7. MTX response in a model of inflammatory arthritis is linked to microbial metabolism of MTX. (**A**) Plasma concentration of MTX and its metabolites were quantified by LC-MS in female GF SKG mice colonized with either Zion or Parnassus (Parn.) microbiota (N=6 per colonization state). After a one-time oral administration of MTX 50 mg/kg, multiple plasma samples were taken from each mouse over a 24 h time course. (**B**) Peak concentration and (**C**) AUC were compared between Zion and Parnassus colonized mice. (**D**) Ratio of AUC of 7-OH-MTX to MTX in the plasma over 24 h. (**E**) Stool concentrations of MTX and its metabolites were quantified at multiple time points, and (**F**) peak stool concentrations, (**G**) AUC and (**H**) product-substrate ratios of DAMPA (top panel) and pMABG (bottom panel) relative to MTX are shown. Results from Student's t-test (regular black) or Wilcoxon test (italic blue, data are non-normal by Shapiro-Wilk test) are reported. Error bars are SEM.

1095
Supplementary Figure 5, related to Figure 6. The microbiome of SKG mice reared at two
different facilities is associated with MTX response. (**A**) Alpha diversity (number of observed
ASVs) by facility-of-origin in mice that are not co-housed ("Representative", left panel) and in mice
that are co-housed for 2 weeks ("Co-housed", right panel). (**B**) Principal components analysis of
Euclidean distances of clr-transformed abundances (beta diversity) of Co-housed mice (solid
circles) and Representative mice from each facility that were not co-housed (open circles). (**C**)
Relative abundance of ASVs that were differentially abundant by facility-of-origin (DESeq,
$p_{adj}<0.05$) in mice that were co-housed. Results of PERMANOVA reported in B. All results in **C**
are significant by DESeq.

Reference list

1. Backhed, F., Ding, H., Wang, T., Hooper, L.V., Koh, G.Y., Nagy, A., Semenkovich, C.F., and Gordon, J.I. (2004). The gut microbiota as an environmental factor that regulates fat storage. Proc Natl Acad Sci U S A *101*, 15718-15723. 10.1073/pnas.0407076101.

2. Verdegaal, A.A., and Goodman, A.L. (2024). Integrating the gut microbiome and pharmacology. Sci Transl Med *16*, eadg8357. 10.1126/scitranslmed.adg8357.

3. Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R., and Goodman, A.L. (2019). Mapping human microbiome drug metabolism by gut bacteria and their genes. Nature *570*, 462-467. 10.1038/s41586-019-1291-3.

4. Javdan, B., Lopez, J.G., Chankhamjon, P., Lee, Y.J., Hull, R., Wu, Q., Wang, X., Chatterjee, S., and Donia, M.S. (2020). Personalized mapping of drug metabolism by the human gut microbiome. Cell *181*, 1661-1679 e1622. 10.1016/j.cell.2020.05.001.

5. Turjeman, S., Rozera, T., Elinav, E., Ianiro, G., and Koren, O. (2025). From big data and experimental models to clinical trials: Iterative strategies in microbiome research. Cell *188*, 1178-1197. 10.1016/j.cell.2025.01.038.

6. Porcari, S., Ng, S.C., Zitvogel, L., Sokol, H., Weersma, R.K., Elinav, E., Gasbarrini, A., Cammarota, G., Tilg, H., and Ianiro, G. (2025). The microbiome for clinicians. Cell *188*, 2836-2844. 10.1016/j.cell.2025.04.016.

7. de Crecy-Lagard, V., El Yacoubi, B., de la Garza, R.D., Noiriel, A., and Hanson, A.D. (2007). Comparative genomics of bacterial and plant folate synthesis and salvage: predictions and validations. BMC Genomics *8*, 245. 10.1186/1471-2164-8-245.

8. Gorelova, V., Bastien, O., De Clerck, O., Lespinats, S., Rebeille, F., and Van Der Straeten, D. (2019). Evolution of folate biosynthesis and metabolism across algae and land plant lineages. Sci Rep *9*, 5731. 10.1038/s41598-019-42146-5.

9. Engevik, M.A., Morra, C.N., Roth, D., Engevik, K., Spinler, J.K., Devaraj, S., Crawford, S.E., Estes, M.K., Kalkum, M., and Versalovic, J. (2019). Microbial Metabolic Capacity for Intestinal Folate Production and Modulation of Host Folate Receptors. Front Microbiol *10*, 2305. 10.3389/fmicb.2019.02305.

10. Goldman, P., and Levy, C.C. (1967). Carboxypeptidase G: purification and properties. Proc Natl Acad Sci U S A *58*, 1299-1306. 10.1073/pnas.58.4.1299.

11. Valerino, D.M., Johns, D.G., Zaharko, D.S., and Oliverio, V.T. (1972). Studies of the metabolism of methotrexate by intestinal flora. I. Identification and study of biological properties of the metabolite 4-amino-4-deoxy-N 10 -methylpteroic acid. Biochem Pharmacol *21*, 821-831.

12. Goldman, P., and Levy, C.C. (1968). The enzymatic hydrolysis of folate analogues. Biochem Pharmacol *17*, 2265-2270. 10.1016/0006-2952(68)90033-6.

13. Cronstein, B.N., and Aune, T.M. (2020). Methotrexate and its mechanisms of action in inflammatory arthritis. Nat Rev Rheumatol *16*, 145-154. 10.1038/s41584-020-0373-9.

14. Fraenkel, L., Bathon, J.M., England, B.R., St Clair, E.W., Arayssi, T., Carandang, K., Deane, K.D., Genovese, M., Huston, K.K., Kerr, G., et al. (2021). 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. Arthritis Care Res (Hoboken) *73*, 924-939. 10.1002/acr.24596.

15. Nayak, R.R., and Orellana, D.A. (2024). The impact of the human gut microbiome on the treatment of autoimmune disease. Immunol Rev *325*, 107-130. 10.1111/imr.13358.

16. Romao, V.C., Canhao, H., and Fonseca, J.E. (2013). Old drugs, old problems: where do we stand in prediction of rheumatoid arthritis responsiveness to methotrexate and other synthetic DMARDs? BMC Med *11*, 17. 10.1186/1741-7015-11-17.

17. Szostak, B., Machaj, F., Rosik, J., and Pawlik, A. (2020). Using pharmacogenetics to
predict methotrexate response in rheumatoid arthritis patients. Expert Opin Drug Metab
Toxicol *16*, 617-626. 10.1080/17425255.2020.1777279.
18. Eektimmerman, F., Swen, J.J., Madhar, M.B., Allaart, C.F., and Guchelaar, H.J. (2020).
Predictive genetic biomarkers for the efficacy of methotrexate in rheumatoid arthritis: a
systematic review. Pharmacogenomics J *20*, 159-168. 10.1038/s41397-019-0098-9.
19. Artacho, A., Isaac, S., Nayak, R., Flor-Duro, A., Alexander, M., Koo, I., Manasson, J.,
Smith, P.B., Rosenthal, P., Homsi, Y., et al. (2021). The pretreatment gut microbiome is
associated with lack of response to methotrexate in new-onset rheumatoid arthritis.
Arthritis Rheumatol *73*, 931-942. 10.1002/art.41622.
20. Bustion, A.E., Nayak, R.R., Agrawal, A., Turnbaugh, P.J., and Pollard, K.S. (2023).
SIMMER employs similarity algorithms to accurately identify human gut microbiome
species and enzymes capable of known chemical transformations. Elife *12*.
10.7554/eLife.82401.
21. Zaharko, D.S., Bruckner, H., and Oliverio, V.T. (1969). Antibiotics alter methotrexate
metabolism and excretion. Science *166*, 887-888.
22. Donehower, R.C., Hande, K.R., Drake, J.C., and Chabner, B.A. (1979). Presence of 2,4-
diamino-N10-methylpteroic acid after high-dose methotrexate. Clin Pharmacol Ther *26*,
63-72.
23. Stewart, M.J., Watson, I.D., Farid, Y.Y., and Skellern, G.G. (1986). An investigation into
the source of the deglutamated metabolites of methotrexate in patients treated with high
dose infusions. Ann Clin Biochem *23 (Pt 2)*, 210-215. 10.1177/000456328602300213.
24. Grim, J., Chladek, J., and Martinkova, J. (2003). Pharmacokinetics and
pharmacodynamics of methotrexate in non-neoplastic diseases. Clin Pharmacokinet *42*,
139-151. 10.2165/00003088-200342020-00003.
25. Letertre, M.P.M., Munjoma, N., Wolfer, K., Pechlivanis, A., McDonald, J.A.K., Hardwick,
R.N., Cherrington, N.J., Coen, M., Nicholson, J.K., Hoyles, L., et al. (2020). A two-way
interaction between methotrexate and the gut microbiota of male Sprague-Dawley rats. J
Proteome Res. 10.1021/acs.jproteome.0c00230.
26. Widemann, B.C., Sung, E., Anderson, L., Salzer, W.L., Balis, F.M., Monitjo, K.S.,
McCully, C., Hawkins, M., and Adamson, P.C. (2000). Pharmacokinetics and
metabolism of the methotrexate metabolite 2, 4-diamino-N(10)-methylpteroic acid. J
Pharmacol Exp Ther *294*, 894-901.
27. Jordan, C.G., Rashidi, M.R., Laljee, H., Clarke, S.E., Brown, J.E., and Beedham, C.
(1999). Aldehyde oxidase-catalysed oxidation of methotrexate in the liver of guinea-pig,
rabbit and man. J Pharm Pharmacol *51*, 411-418. 10.1211/0022357991772619.
28. Kitamura, S., Sugihara, K., Nakatani, K., Ohta, S., Ohhara, T., Ninomiya, S., Green,
C.E., and Tyson, C.A. (1999). Variation of hepatic methotrexate 7-hydroxylase activity in
animals and humans. IUBMB Life *48*, 607-611. 10.1080/713803569.
29. Dervieux, T., Furst, D., Lein, D.O., Capps, R., Smith, K., Walsh, M., and Kremer, J.
(2004). Polyglutamation of methotrexate with common polymorphisms in reduced folate
carrier, aminoimidazole carboxamide ribonucleotide transformylase, and thymidylate
synthase are associated with methotrexate effects in rheumatoid arthritis. Arthritis
Rheum *50*, 2766-2774. 10.1002/art.20460.
30. Bleyer, W.A. (1978). The clinical pharmacology of methotrexate: new applications of an
old drug. Cancer *41*, 36-51.

31.   Hoekstra, M., Haagsma, C., Neef, C., Proost, J., Knuif, A., and van de Laar, M. (2004).
1201   Bioavailability of higher dose methotrexate comparing oral and subcutaneous
1202   administration in patients with rheumatoid arthritis. J Rheumatol *31*, 645-648.
1203   32.   Steinberg, S.E., Campbell, C.L., Bleyer, W.A., and Hillman, R.S. (1982). Enterohepatic
1204   circulation of methotrexate in rats in vivo. Cancer Res *42*, 1279-1282.
1205   33.   Browne, H.P., Forster, S.C., Anonye, B.O., Kumar, N., Neville, B.A., Stares, M.D.,
1206   Goulding, D., and Lawley, T.D. (2016). Culturing of 'unculturable' human microbiota
1207   reveals novel taxa and extensive sporulation. Nature *533*, 543-546.
1208   10.1038/nature17645.
1209   34.   Sakaguchi, N., Takahashi, T., Hata, H., Nomura, T., Tagami, T., Yamazaki, S.,
1210   Sakihama, T., Matsutani, T., Negishi, I., Nakatsuru, S., and Sakaguchi, S. (2003).
1211   Altered thymic T-cell selection due to a mutation of the ZAP-70 gene causes
1212   autoimmune arthritis in mice. Nature *426*, 454-460. 10.1038/nature02119.
1213   35.   Rehaume, L.M., Mondot, S., Aguirre de Carcer, D., Velasco, J., Benham, H., Hasnain,
1214   S.Z., Bowman, J., Ruutu, M., Hansbro, P.M., McGuckin, M.A., et al. (2014). ZAP-70
1215   genotype disrupts the relationship between microbiota and host, leading to
1216   spondyloarthritis and ileitis in SKG mice. Arthritis Rheumatol *66*, 2780-2792.
1217   10.1002/art.38773.
1218   36.   Sakaguchi, S., Benham, H., Cope, A.P., and Thomas, R. (2012). T-cell receptor
1219   signaling and the pathogenesis of autoimmune arthritis: insights from mouse and man.
1220   Immunol Cell Biol *90*, 277-287. 10.1038/icb.2012.4.
1221   37.   Tanaka, A., Maeda, S., Nomura, T., Llamas-Covarrubias, M.A., Tanaka, S., Jin, L., Lim,
1222   E.L., Morikawa, H., Kitagawa, Y., Akizuki, S., et al. (2023). Construction of a T cell
1223   receptor signaling range for spontaneous development of autoimmune disease. J Exp
1224   Med *220*. 10.1084/jem.20220386.
1225   38.   Maeda, Y., Kurakawa, T., Umemoto, E., Motooka, D., Ito, Y., Gotoh, K., Hirota, K.,
1226   Matsushita, M., Furuta, Y., Narazaki, M., et al. (2016). Dysbiosis contributes to arthritis
1227   development via activation of autoreactive T Cells in the intestine. Arthritis Rheumatol
1228   *68*, 2646-2661. 10.1002/art.39783.
1229   39.   Nagate, T., Kawai, J., and Nakayama, J. (2009). Therapeutic and preventive effects of
1230   methotrexate on zymosan-induced arthritis in SKG mice. J Vet Med Sci *71*, 713-717.
1231   40.   Larsson, E., Tremaroli, V., Lee, Y.S., Koren, O., Nookaew, I., Fricker, A., Nielsen, J.,
1232   Ley, R.E., and Backhed, F. (2012). Analysis of gut microbial regulation of host gene
1233   expression along the length of the gut and regulation of gut microbial ecology through
1234   MyD88. Gut *61*, 1124-1131. 10.1136/gutjnl-2011-301104.
1235   41.   Maynard, C., Cummins, I., Green, J., and Weinkove, D. (2018). A bacterial route for folic
1236   acid supplementation. BMC Biol *16*, 67. 10.1186/s12915-018-0534-3.
1237   42.   Zheng, Y., and Cantley, L.C. (2019). Toward a better understanding of folate metabolism
1238   in health and disease. J Exp Med *216*, 253-266. 10.1084/jem.20181965.
1239   43.   Huennekens, F.M. (1996). In search of dihydrofolate reductase. Protein Sci *5*, 1201-
1240   1208. 10.1002/pro.5560050626.
1241   44.   McHorney, C.A., and Tarlov, A.R. (1995). Individual-patient monitoring in clinical
1242   practice: are available health status surveys adequate? Qual Life Res *4*, 293-307.
1243   10.1007/BF01593882.
1244   45.   Stucki, G., Liang, M.H., Stucki, S., Katz, J.N., and Lew, R.A. (1999). Application of
1245   statistical graphics to facilitate selection of health status measures for clinical practice
1246   and evaluative research. Clin Rheumatol *18*, 101-105. 10.1007/s100670050065.
1247   46.   Aletaha, D., Neogi, T., Silman, A.J., Funovits, J., Felson, D.T., Bingham, C.O., 3rd,
1248   Birnbaum, N.S., Burmester, G.R., Bykerk, V.P., Cohen, M.D., et al. (2010). 2010

Rheumatoid arthritis classification criteria: an American College of
Rheumatology/European League Against Rheumatism collaborative initiative. Arthritis
Rheum *62*, 2569-2581. 10.1002/art.27584.
47.   Nayak, R.R., Alexander, M., Deshpande, I., Stapleton-Gray, K., Rimal, B., Patterson,
A.D., Ubeda, C., Scher, J.U., and Turnbaugh, P.J. (2021). Methotrexate impacts
conserved pathways in diverse human gut bacteria leading to decreased host immune
activation. Cell Host Microbe *29*, 362-377 e311. 10.1016/j.chom.2020.12.008.
48.   Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N.,
Owens, S.M., Betley, J., Fraser, L., Bauer, M., et al. (2012). Ultra-high-throughput
microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J *6*,
1621-1624. 10.1038/ismej.2012.8.
49.   Gohl, D.M., Vangay, P., Garbe, J., MacLean, A., Hauge, A., Becker, A., Gould, T.J.,
Clayton, J.B., Johnson, T.J., Hunter, R., et al. (2016). Systematic improvement of
amplicon marker gene methods for increased accuracy in microbiome studies. Nat
Biotechnol *34*, 942-949. 10.1038/nbt.3601.
50.   Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello,
E.K., Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I., et al. (2010). QIIME allows
analysis of high-throughput community sequencing data. Nat Methods *7*, 335-336.
10.1038/nmeth.f.303.
51.   Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A.,
Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., et al. (2019). Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nat
Biotechnol *37*, 852-857. 10.1038/s41587-019-0209-9.
52.   Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., and Holmes,
S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. Nat
Methods *13*, 581-583. 10.1038/nmeth.3869.
53.   Wang, Q., Garrity, G.M., Tiedje, J.M., and Cole, J.R. (2007). Naive Bayesian classifier
for rapid assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ
Microbiol *73*, 5261-5267. 10.1128/AEM.00062-07.
54.   McMurdie, P.J., and Holmes, S. (2013). phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. PLoS One *8*, e61217.
10.1371/journal.pone.0061217.
55.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550. 10.1186/s13059-014-
0550-8.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supp. Figure 1

A

B

Supp Figure 2

Supp Figure 3

Supp Figure 4

Supp Figure 5