Structural Impact of Alzheimer's Disease Mutations on Transmembrane TREM2-DAP12 Interactions: An Atomistic Perspective

Zhiwen Zhong[1,2], Martin B. Ulmschneider[2], and Christian D. Lorenz[3,*]

[1]Department of Physics, King's College London, London WC2R 2LS, UK

[2]Department of Chemistry, King's College London, London, SE1 1DB, UK

[3]Department of Engineering, King's College London, London WC2R 2LS, UK

[*]To whom correspondence should be addressed: chris.lorenz@kcl.ac.uk

ABSTRACT

Triggering receptor expressed on myeloid cells 2 (TREM2) is an immunomodulatory receptor that plays a critical role in microglial activation through its association with the adaptor protein DNAX-activation protein 12 (DAP12). Genetic studies have identified rare TREM2 variants as key risk factors for Alzheimer's disease (AD), with mutations affecting both the extracellular and transmembrane domains. While the TREM2-DAP12 complex is essential for microglial activation and is implicated in Alzheimer's disease pathology, the structural and mechanistic effects of transmembrane domain mutations, particularly within different TREM2 isoforms, remain unclear. In this study, we employ multi-scale molecular dynamics simulations to investigate the impact of six TREM2 mutations (four in isoform 1, two in isoform 2) and one wild-type control (isoform 2) on complex stability. By integrating coarse-grain and all-atom simulations with unsupervised machine learning techniques, we reveal distinct conformational states and residue-level interactions that govern complex formation. Our findings demonstrate how mutations such as W191X in TREM2 isoform 2 disrupt critical interactions, leading to destabilisation of the complex and potential impairments in neuroimmune signalling. These results provide new insights into the structural mechanisms linking TREM2 mutations to AD and offer a foundation for developing targeted therapeutic strategies.

KEYWORDS

Alzheimer's disease, TREM2, MD simulation, AlphaFold, Machine learning

INTRODUCTION

Alzheimer's disease (AD) is a progressive neurodegenerative condition characterised mainly by β-amyloid (Aβ) plaque deposition in the brain (Jonsson et al. 2013b; Huang and Mucke 2012). Microglia, the brain's resident immune cells, are frequently found surrounding Aβ plaques in both AD patients (D'Andrea et al. 2004) and transgenic mouse models (Dickson 1999), where they contribute to the clearance of Aβ and modulation of neuroinflammation. Recent genome-wide association studies (GWAS) have identified rare variants in TREM2 (Triggering Receptor Expressed on Myeloid cells 2) as significant genetic risk factors for late-onset AD, with the R47H mutation in the extracellular domain conferring a particularly strong effect (Guerreiro et al. 2013b; Jonsson et al. 2013a). TREM2 functions as an immunomodulatory receptor on microglial surfaces and initiates intracellular signalling via its transmembrane adaptor DAP12 (DNAX-activating protein of 12 kDa) (Colonna 2023; Ulland and Colonna 2018).

TREM2 interacts with a diverse range of ligands such as phospholipids, sulfatides, bacterial lipopolysaccharide (LPS) and DNA (Wang et al. 2015; Cannon et al. 2012; Daws et al. 2003; Kawabori et al. 2015), and the nature of these interactions can differentially modulate its signalling strength and outcome (Kober and Brett 2017). Ligand engagement at the TREM2 extracellular domain initiates the signalling cascade through association with adaptor proteins DAP10/12 (Ulland et al. 2017). Subsequently, the TREM2 ectodomain is shed by ADAM10/17 metalloproteases, followed by intramembrane cleavage of the C-terminal fragment (CTF) by γ-secretase-a process whose structural mechanism has recently been elucidated by cryo-EM (Zhou et al. 2019; Thornton et al. 2017; Feuerbach et al. 2017). This proteolytic sequence effectively terminates TREM2 signalling, enabling microglia to revert to a homeostatic state. Following ligand binding, TREM2 couples with DAP12 via their transmembrane domains (Steiner et al. 2020a; Hamerman et al. 2006;

Peng et al. 2010), triggering phosphorylation of DAP12's immunoreceptor tyrosine-based activation motifs (ITAMs) and recruitment of kinases including SYK, PLCγ, and PI3K (Kleinberger et al. 2014; Guerreiro et al. 2013a; Paloneva et al. 2002).

The TREM2-DAP12 interaction is initiated through electrostatic pairing between lysine 186 (K186) of TREM2 and conserved aspartate residues (D50) in DAP12, forming a stable trimeric transmembrane complex (Call et al. 2010a). Additional stabilisation is provided by W194 in TREM2, which forms a hydrogen bond with T54 in DAP12 (Zhong et al. 2024). While most functional studies of TREM2 focus on extracellular domain mutations, emerging evidence implicates transmembrane domain variants in AD risk. Notably, the W191X (X represents truncation) nonsense mutation in TREM2 isoform 2 has been associated with increased AD susceptibility in African American populations (Jin et al. 2015). TREM2 exists in at least three isoforms by alternative splicing (Moutinho et al. 2023; Bharadwaj et al. 2024; Giraldo et al. 2013), with isoform 1 (230 residues) being the canonical and most abundant form (Celarain et al. 2016). Isoform 2 (219 residues), was found to be expressed at lower levels than isoform 1 in the hippocampus of AD patients (Ma et al. 2016), while isoform 3 (222 residues) remains even less studied (Jin et al. 2014).

Structural biology approaches like cryo-EM and X-ray crystallography offer valuable but static views of protein complexes (Callaway 2024; Shoemaker and Ando 2018; Beck et al. 2024). Machine-learning-based prediction models, such as AlphaFold2 and AlphaFold3 (Jumper et al. 2021b; Abramson et al. 2024), provide high-quality structures but often lack the conformational dynamics critical for understanding membrane-embedded interactions (Malhotra et al. 2025). Atomistic molecular dynamics (MD) simulations offer complementary insights by capturing time-resolved structural fluctuations within biologically realistic environments (Karplus and McCammon 2002; Hospital et al. 2015). In this study, we build upon our previous work resolving the TREM2-DAP12 complex in a lipid bilayer (Zhong et al. 2024), investigating the structural consequences of AD-associated and designed point mutations in the TREM2 transmembrane domain (TMD). We

investigate seven different forms of TREM2, including four (K186A, K186X, W194A & W194X) mutants of isoform 1 as well as the wild type and two mutants (W191A & W191X) of isoform 2. All simulations are conducted with the TREM2-DAP12 complex embedded within a model lipid membrane consisting of an 80:20 mixture of POPC and cholesterol. By combining multi-scale MD simulations with unsupervised machine learning analysis, we provide an atomistic perspective on how these mutations modulate the TREM2-DAP12 complex stability and their potential implications in Alzheimer's disease pathogenesis.

RESULTS

Unsupervised Learning Reveals Distinct Conformational Clusters in TREM2-DAP12 Complexes

Previous studies have shown that TREM2 interacts with the signalling adaptor DAP12, primarily through a key salt bridge between K186 in TREM2 and D50 in DAP12 (Ulland and Colonna 2018; Steiner et al. 2020b). The cartoons shown in Fig. 1A represent the membrane-embedded α-helices structures of TREM2 and DAP12, which we reported in previous work (Zhong et al. 2024). A focused sequence alignment of the transmembrane regions is presented in Fig. 1B (with the full alignment shown in Supplemental Figure S1). We observed that residue W191 in the TREM2-219 isoform corresponds positionally to K186 in TREM2-230. Since K186 is known to be critical for TREM2-DAP12 interaction, and W191 has been potentially linked to Alzheimer's disease, this suggests that W191 may play a key role in mediating the interaction between TREM2-219 and DAP12.

Fig. 1C illustrates the ligand binding to the extracellular domain of TREM2 and the subsequent intracellular signalling cascade initiated by DAP12. Upon proteolytic cleavage and formation of the C-terminal fragment (CTF), TREM2 remains associated with DAP12, enabling ITAMs phosphorylation and SYK recruitment (Fig. 1D). To investigate how specific mutations affect the structural dynamics of the TREM2-DAP12 complex, we introduced several point mutations into TREM2 and simulated each variant in complex with DAP12. The details of system preparation

and mutation design are described in the Methods section. Coarse-grain molecular dynamics (CG-MD) simulations were performed for 300 μs on TREM2-230 variants and 200 μs on TREM2-219 variants.

Using the trajectories from the CG-MD simulations, we implemented an unsupervised machine learning pipeline to characterise the conformational landscapes of the TREM2-DAP12 complexes. High-dimensional contact map data from the trajectories were first embedded into a two-dimensional space using the Uniform Manifold Approximation and Projection (UMAP) algorithm. Then the Hierarchical Density-Based Spatial Clustering of Applications with Noise (HDBSCAN) algorithm was applied to identify distinct conformational states in the UMAP embeddings. The resulting clusters represent dominant structural states sampled by each mutant. These clusters are visualised in Figure 2A and 2D for the TREM2-230 and TREM2-219 constructs, respectively. The relative populations of these conformational states are depicted as doughnut plots in Fig. 2B and Fig. 2E. The temporal evolution of the cluster assignments across the various simulation trajectories, shown in Fig. 2C and Fig. 2F, reveals differences in the stability and dynamics of the sampled conformations of the various mutants.

These results indicate different conformations on different timescales. The detailed contact maps and representative conformational structures are shown in Supplemental Figures S2-S8.

Characterization of TREM2-DAP12 Interaction Interfaces and Mutant Effects

To further understand the interaction interfaces of the different TREM2-DAP12 complexes, we identified representative structures from the dominant conformational clusters observed in the CG-MD simulations (Table 1) and then converted them to all-atom models using the CHARMM-GUI converter (Qi et al. 2014; Jo et al. 2008). We simulated these resulting systems for 300 ns. From these trajectories, we constructed contact maps using a 3 Å cutoff distance (Zhong et al. 2024), between C-α atoms to define residue-residue interactions (Supplemental Figures S9-S15). Residue pairs that maintained contact for over 60% of the trajectory were labelled as interacting residues, while those with contact persistence above 90% were defined as important residues (Fig. 3, blue

and red dots, respectively). A comprehensive summary of the interacting residue pairs that persist for 60%, 70%, 80%, and 90% of the trajectories is provided in Supplemental Table S1.

In isoform TREM2-230 (Fig. 3A), the K186A mutant exhibited fewer persistent contacts than are found in the wild-type, underscoring the role of K186 in stabilising the TREM2-DAP12 complex. The key contact sites between the K186X mutant and DAP12 are found to be in the C-terminal regions of both proteins. Notably, the W194A-230 mutant displayed the highest number of contacts among the isoform 1 variants, with interaction hotspots spanning from the middle to the C-terminal region of TREM2 and across the whole length of DAP12. In contrast, the W194X-230 mutant showed markedly reduced contact frequency, suggesting impaired complex formation.

In isoform TREM2-219 (Fig. 3B), WT-219 complexes retained substantial interactions ranging from the middle to the C-terminal regions of both TREM2 and DAP12. The W191A mutation preserved these contacts with a prominent clustering toward the C-terminal regions. Strikingly, the W191X-219 mutant exhibited no detectable contacts within the defined threshold, indicating a complete loss of interaction. This observation is consistent with the hypothesis that W191 is critical for maintaining the TREM2-DAP12 interface and is in line with genetic studies linking W191 variants to Alzheimer's disease susceptibility.

The rectangles overlaid in Fig. 3 denote the approximate boundaries of the transmembrane α-helices in each complex, providing structural context for the observed contact distributions.

Mechanistic Insights into Residue-Residue Interactions and Mutant-Induced Disruptions in TREM2-DAP12 Complex Stability

To gain mechanistic insight into the specific residue-residue interactions stabilising TREM2-DAP12 complexes, we analysed interaction types across all mutant systems (Fig. 4A). Within the K186A-230 mutant, we found that π-π stacking interaction between W194 of TREM2 and Y62 of DAP12 was observed in the K186A-230 mutant (Fig. 4B) to play an important in the formation of the complex, occurring 37.60% of the total simulation time-32.28% in a parallel (sandwich-like) orientation and 5.32% in a T-shaped configuration (Fig. 4C). The details of the π-π stacking

measurement can be found in Supplemental Figure S16. In addition, a hydrophobic interaction between W194 and V61 further supports trimer stability (Fig. 5A).

In the W194A-230 mutant, a stable hydrogen bond forms between K186 in TREM2 and D50 in both DAP12 monomers (Fig. 4D). This hydrogen bond persists for 92.34% of the simulation, with a hydrogen bond to one DAP12 monomer present 39.10% of the time and hydrogen bonds to both DAP12 monomers occurring simultaneously in 53.24% of the frames (Fig. 4E). The details of the number of hydrogen bonds as a function of time can be found in Supplemental Figure S17. The further interaction network that stabilises the W194A-230 complex includes multiple hydrophobic contacts and a salt bridge that contribute to tight trimer packing (Fig. 5C, top panel). To facilitate comparison between the wild-type and mutant forms of isoform 1, the previously reported key interacting residues in WT-230 (Zhong et al. 2024) are also highlighted in Fig. 4F.

In contrast, the two truncated TREM2-230 complexes are primarily stabilised by hydrophobic interactions. In the case of the W194X-230 complex, the hydrophobic contact between L178 (TREM2) and T54 (DAP12) is found to play an important role in the stabilisation of the complex. While in the K186X-230 complex, we found that the complex is a relatively stable conformation driven by hydrophobic interactions involving I185 of TREM2 with Y62 and R66 of DAP12 (Fig. 5A).

For the WT-219 system, complex stability is maintained through a hydrophobic interaction between L198 (TREM2) and A58 (DAP12), along with a salt bridge and hydrogen bond between E202 (TREM2) and R66 (DAP12), with a 70% occupancy reported in Table S1 and shown in Fig. 5C (middle panel). The W191A-219 mutant retains the L198-A58 hydrophobic interaction and features an additional hydrogen bond between E202 (TREM2) and Y62 (DAP12) (Fig. 5C, bottom panel).

In stark contrast, the W191X-219 mutant completely disrupts all residue-level contacts, decreasing complex stability (Fig. 4A). Structural analysis reveals that TREM2 reorients horizontally along the membrane interface, while DAP12 retains a vertical orientation (Fig. 5B). Analysis of the z-axis positioning over time (Fig. 5D and Fig. 5E) further confirms this dissociation: in WT-219,

both proteins remain colocalised in the upper membrane leaflet, whereas in W191X-219, TREM2 shifts toward the lower leaflet while DAP12 remains within the center of the membrane.

The K186X-230 and W191X-219 mutants exhibited significantly greater center-of-mass distances between TREM2 and DAP12 in the $x - y$ plane, measuring 13.02 Å and 11.20 Å, respectively (Supplementary Figure S18). In contrast, smaller distances were observed in the K186A-230 (7.86 Å), W194A-230 (8.08 Å), W194X-230 (9.83 Å), WT-219 (6.06 Å), and W191A-230 (6.07 Å) complexes. These findings suggest that the increased spatial separation in K186X-230 and W191X-230 reflects destabilised or weakened interactions between TREM2 and DAP12.

Impact of Point Mutations on Structural Flexibility and Stability of the TREM2-DAP12 Complex

To investigate how point mutations affect structural flexibility, we analysed the root mean square fluctuation (RMSF) profiles of each TREM2-DAP12 complex (Fig. 6). Fig. 6 A shows RMSF values for the 230-residue constructs. Compared to the WT-230 complex, the K186A-230 mutant displays increased flexibility in the N-terminal region of TREM2. This likely results from the loss of K186-mediated anchoring interactions. In the K186X-230 complex, higher fluctuations are observed in one of the DAP12 chains, suggesting that truncation of TREM2 disrupts the symmetric stability of the trimeric interface. The W194A-230 variant also exhibits elevated dynamics in the N-terminal domain of TREM2, possibly due to the loss of the aromatic W194 residue, which contributes to π-stacking and hydrophobic stabilisation. W194X-230 shows localised fluctuation increases but remains relatively constrained compared to K186X.

In the 219-residue systems (Fig. 6B), the WT-219 complex exhibits extensive flexibility in the cytoplasmic region of TREM2 and the DAP12 interface, maybe due to the lack of interaction of the N-terminus of TREM2 with either DAP12 or lipids (Fig. 5B). W191A-219 displays a slightly more rigid profile, possibly due to compensatory stabilising hydrogen bonds. W191X-219 shows moderate flexibility across both proteins. These RMSF profiles underscore the dynamic nature of the TREM2-DAP12 interaction and reveal how specific mutations perturb stability through altered

₂₁₄ flexibility in both transmembrane and cytoplasmic domains.

₂₁₅

DISCUSSION

₂₁₇   The W191X mutation introduces a premature stop codon within the transmembrane domain of

₂₁₈ the TREM2-219 isoform, leading to truncation of the receptor's C-terminal tail. Genetic studies

₂₁₉ have previously associated this nonsense variant with an increased risk of Alzheimer's disease,

₂₂₀ particularly in African American populations (Giraldo et al. 2013; Jin et al. 2015). However, the

₂₂₁ biophysical consequences of W191X on TREM2-DAP12 complex formation and function have

₂₂₂ remained elusive.

₂₂₃   Our simulations provide structural and mechanistic insights into how this mutation disrupts

₂₂₄ interprotein signalling. In contrast to the WT-219 and W191A-219 systems, which maintain

₂₂₅ robust interactions through hydrophobic, hydrogen bond, and salt bridge networks, the W191X-

₂₂₆ 219 complex shows a complete loss of residue-level contacts at the TREM2-DAP12 interface

₂₂₇ (Fig. 3B & Fig. 4A). This disruption is further supported by the W191X variant, which exhibits a

₂₂₈ striking decoupling of the two helices, both laterally (11.2 Å in the x-y plane) and vertically (along

₂₂₉ the z-axis) orientation (Fig. 5D, Fig. 5E & Supplemental Figure S18). Notably, the TREM2 helix in

₂₃₀ W191X-219 reorients almost horizontally along the bilayer interface-a configuration incompatible

₂₃₁ with canonical receptor-adaptor signalling (Colonna 2023; Ulland and Colonna 2018).

₂₃₂   At the molecular level, the truncation eliminates distal cytoplasmic residues, including E202-a

₂₃₃ key site for salt bridge formation with R66 of DAP12 in WT-219 and W191A-219 systems (Fig. 5C).

₂₃₄ The loss of this interaction likely destabilises the helix alignment necessary for signal transduction

₂₃₅ via ITAMs on DAP12. These findings collectively suggest that W191X disrupts not only static

₂₃₆ contacts but also the dynamic coherence required for functional signalling.

₂₃₇   Importantly, our results also provide comparative insights into mutation-induced destabilisation

₂₃₈ across isoforms. The K186X-230 mutant, though distinct in sequence and location, mirrors the

₂₃₉ disruption seen in W191X-219, with increased centre-of-mass separation and reduced interhelical

₂₄₀ stability (Fig. 5A, Supplemental Figure S18). This parallel underscores the critical role of C-

terminal residues in preserving the spatial geometry of the receptor-adaptor complex.

Beyond the immediate biophysical implications, these findings have broader relevance for understanding TREM2's role in neuroimmune signalling. Misalignment or dissociation of TREM2 from DAP12 could impair SYK recruitment and ITAMs phosphorylation, mechanisms essential for microglial activation and phagocytosis (Wang et al. 2022; Zhang et al. 2025). Given TREM2's critical role in regulating microglial responses to amyloid plaques, including early-stage plaque seeding and clustering (Parhizkar et al. 2019; Ulrich et al. 2017), structural disruptions such as W191X may mechanistically link altered microglial function to increased Alzheimer's disease risk.

In summary, this study establishes a molecular framework for understanding how C-terminal truncation mutations, such as W191X, destabilise the TREM2-DAP12 complex. It underscores the importance of structural integrity in maintaining signalling-competent receptor assemblies and provides new insights for targeting TREM2 variants in the context of neurodegenerative diseases.

METHODS

Approach to Build Systems

The TREM2 mutants are made with ColabFold v1.5.2 (Mirdita et al. 2022; Jumper et al. 2021a) using the mutated sequence (structures confidence can be found in Supplemental Figure S19-S25), and the DAP12 dimer is using PDB 2L34 (Call et al. 2010b). The complexes are made by merging the TREM2 mutants and DAP12 together as a reference using HADDOCK (Dominguez et al. 2003; Van Zundert et al. 2016), AlphaFold2 (Jumper et al. 2021a), AlphaFold3 (Abramson et al. 2024), and the Wild-type complex structure, we simulated a stable conformation. The sequences of each system can be found in Supplemental Table S2. Each system was then equilibrated using all-atom molecular dynamics simulations for 100 ns. The systems utilised in this paper were generated through the use of the CHARMM-GUI Membrane Builder (Jo et al. 2008) and the MARTINI Maker (Siewert et al. 2007) for all-atom (AA) and coarse-grain (CG) simulations, respectively. The membrane consists of POPC and cholesterol at a ratio of 80 : 20. The solvent employed in every simulation was water, and to maintain a neutral charge and a 0.15 M salt concentration, Na^+ and

Cl⁻ ions were added to the system. The initial configurations for each system are shown in Supplemental Figure S26, while the complete simulation workflow is outlined in Supplemental Figure S27.

Coarse-grain Molecular Dynamics Simulations

Coarse-grain simulations of 300 μs and 200 μs were performed for isoform 1 and isoform 2, respectively, to obtain equilibrated systems embedded in a POPC/cholesterol lipid bilayer. Simulations were carried out using GROMACS version 2019.2 (Van Der Spoel et al. 2005). The MARTINI22P force field (Qi et al. 2015) was applied for all systems. Each membrane system underwent energy minimisation followed by equilibration at 310.15 K. The MARTINI22P model maps three to five non-hydrogen atoms onto coarse-grain beads, with bead types defined by the atomic composition and chemical characteristics. Beads are categorised into four main classes-charged, polar, non-polar, and apolar-with five subtypes per class, distinguished by polarity and hydrogen-bonding potential. An overview of the CG simulation setups for various TREM2-DAP12 complexes is provided in Table 2.

All-atom Molecular Dynamics Simulations

Following 300 μs or 200 μs of coarse-grain simulations, each system was converted to an all-atom representation using the CHARMM-GUI Martini to All-atom Converter (Qi et al. 2014; Jo et al. 2008), and subsequently simulated for 300 ns. GROMACS version 2020.3 (Van Der Spoel et al. 2005) was used to perform these AA simulations. The same POPC/cholesterol bilayer composition was retained, with lipids described by the CHARMM36 force field (Guvench et al. 2011; Klauda et al. 2010) and water represented by the CHARMM-modified TIP3P model (MacKerell Jr et al. 1998). Each system was first energy-minimised and then equilibrated at 310.15 K and 1 bar, following CHARMM-GUI protocols (MacKerell Jr et al. 1998; Best et al. 2012). The equilibration began with steepest descent minimisation, followed by a 125 ps NVT ensemble run using the Nosè-Hoover thermostat to maintain temperature (310.15 K) with a 1 fs time step. This was followed by a 300 ns production simulation in the NPT ensemble, employing the Nosè-Hoover

thermostat and the Parrinello-Rahman barostat to sustain the temperature at 310.15 K and pressure

at 1 atm (NOSÉ 2002; Parrinello and Rahman 1981). Bond constraints involving hydrogens were

applied using the LINCS algorithm (Hess et al. 1997). Periodic boundary conditions were used in

all three dimensions. Both equilibration and production phases used a 1 fs time step. Electrostatic

and Lennard-Jones (LJ) interactions were truncated at 12 Å, with a switching function applied to

taper LJ interactions to zero starting from 10 Å. Table 3 summarises the AA simulation setups for

the TREM2-DAP12 complexes. Three independent replicas were conducted.

Simulation Analysis Techniques

All simulations were analysed using in-house Python (3.10) scripts built primarily on the

MDAnalysis package (Gowers et al. 2016; Michaud-Agrawal et al. 2011). Visualisations were

generated using Matplotlib (Hunter 2007), and trajectory inspection was carried out with VMD

1.9.3 (Humphrey et al. 1996).

Hydrogen Bond Analysis

Hydrogen bond probabilities were calculated using the Hydrogen Bond Analysis module in

MDAnalysis (Smith et al. 2019), via custom Python scripts. A donor-acceptor distance cutoff of

3 Å and a donor-hydrogen-acceptor angle cutoff of 150 ° were applied.

π-π Stacking Analysis

π-π stacking interactions were evaluated using an in-house Python script based on PySoftK

(López-Ríos de Castro et al. 2025). Ring centroids were computed as the mean position of ring

atoms, and ring plane orientation was defined using three non-collinear atoms. Interactions were

classified using a centroid distance cutoff of 6.5 Å (Piovesan et al. 2016) and an interplanar angle

criterion: $\leq 30°$ for parallel-displaced, and $60°$-$120°$ for T-shaped stacking.

Conformational Clustering of TREM2-DAP12 Complexes

To identify stable conformational states of the TREM2-DAP12 complexes from coarse-grain simulations, we first extracted inter-residue distance features from backbone beads (BB) spaced every five residues along the transmembrane helices of TREM2 and both DAP12 chains. At each simulation frame, we computed pairwise distance matrices between the selected BB atoms of TREM2 and those of each DAP12 chain independently using MDAnalysis. These matrices were flattened into feature vectors and concatenated across frames to form a high-dimensional dataset representing the dynamic spatial organisation of the transmembrane helices.

We then applied UMAP for dimensionality reduction, followed by clustering with HDBSCAN. These hyperparameters were selected iteratively to yield physically meaningful conformational clusters. The resulting low-dimensional embeddings captured dominant structural states sampled by each system and were used to initialise subsequent all-atom simulations. Previously, we have used this general approach to characterising the conformations taken by antimicrobial peptides (Serian et al. 2024), apo-c3 (al Badri et al. 2022), interacting lipids (Smith et al. 2020) and polymers (Ziolek et al. 2021; Ziolek et al. 2022; López-Ríos de Castro et al. 2023; López-Ríos de Castro et al. 2024). For isoform 1, UMAP was performed with `n_neighbors` set to 50, and subsequent clustering used HDBSCAN with `min_cluster_size` = 50 and `cluster_selection_epsilon` = 2.0. For isoform 2, the same `min_cluster_size` was used, with `n_neighbors` = 10 and `cluster_selection_epsilon` = 1.5.

Contact Map Analysis

Residue-level contact maps between TREM2 and DAP12 were calculated using a custom Python script with MDAnalysis. Contacts were defined as any pair of heavy atoms from TREM2 and DAP12 residues within 3 Å. At each frame, a binary contact matrix was generated and accumulated over the trajectory. The final contact map represents the frequency of residue-residue contacts over time, enabling identification of stable interaction interfaces.

Root Mean Square Fluctuation (RMSF) Analysis

RMSF was calculated using a custom Python script with MDAnalysis and NumPy. Alpha carbon atoms were selected, and the trajectory was iterated to compute the mean atomic positions and displacement over time. The RMSF for each residue was computed as:

$$\text{RMSF}_i = \sqrt{\frac{1}{T} \sum_{t=1}^{T} \|\vec{r}_i(t) - \langle \vec{r}_i \rangle\|^2}$$

where $\vec{r}_i(t)$ is the position of atom i at frame t, and $\langle \vec{r}_i \rangle$ is its time-averaged position. This analysis was performed across the full trajectory for each system to assess per-residue flexibility.

ACKNOWLEDGEMENT

We thank Dr. Robert M. Ziolek for his valuable feedback on our manuscript. We are grateful to the UK Materials and Molecular Modelling Hub for computational resources, which is partially funded by EPSRC (EP/T022213/1, EP/W032260/1 and EP/P020194/1). We also appreciate the King's Computational Research, Engineering and Technology Environment (CREATE) at King's College London for providing us access to computational resources (e Research 2025). Z.Z acknowledges receiving PhD studentships from King's College London and China Scholarship Council.

CONTRIBUTION

Z.Z, M.B.U, and C.D.L designed research; Z.Z performed research; Z.Z and C.D.L analysed data; Z.Z and C.D.L wrote the paper.

COMPETING INTERESTS

The authors declare no conflict of interest.

DATA AVAILABILITY

Example trajectories and system setup for analysis are available in a Zenodo repository, accessible at https://doi.org/10.5281/zenodo.15746936.

SUPPLEMENTARY INFORMATION

Supplementary Figures and Tables are available as a PDF.

REFERENCES

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A. J., Bambrick, J., et al. (2024). "Accurate structure prediction of biomolecular interactions with alphafold 3." *Nature*, 630(8016), 493–500.

al Badri, M. A., Smith, P., al Jamal, K. T., and Lorenz, C. D. (2022). "Nanomaterial functionalization modulates hard protein corona formation: Atomistic simulations applied to graphitic materials." *Adv. Mater. Interfaces*, 9(1), 2101236.

Beck, M., Covino, R., Hänelt, I., and Müller-McNicoll, M. (2024). "Understanding the cell: Future views of structural biology." *Cell*, 187(3), 545–562.

Best, R. B., Zhu, X., Shim, J., Lopes, P. E., Mittal, J., Feig, M., and MacKerell Jr, A. D. (2012). "Optimization of the additive charmm all-atom protein force field targeting improved sampling of the backbone ϕ, ψ and side-chain $\chi 1$ and $\chi 2$ dihedral angles." *Journal of chemical theory and computation*, 8(9), 3257–3273.

Bharadwaj, S., Groza, Y., Mierzwicka, J. M., and Malỳ, P. (2024). "Current understanding on trem-2 molecular biology and physiopathological functions." *International Immunopharmacology*, 134, 112042.

Call, M. E., Wucherpfennig, K. W., and Chou, J. J. (2010a). "The structural basis for intramembrane assembly of an activating immunoreceptor complex." *Nature immunology*, 11(11), 1023–1029.

Call, M. E., Wucherpfennig, K. W., and Chou, J. J. (2010b). "The structural basis for intramembrane assembly of an activating immunoreceptor complex." *Nature Immunology*, 11(11), 1023–1029.

Callaway, E. (2024). "These'movies' of proteins in action are revealing the hidden biology of cells.." *Nature*, 627(8004), 480–482.

Cannon, J. P., O'Driscoll, M., and Litman, G. W. (2012). "Specific lipid recognition is a general feature of cd300 and trem molecules." *Immunogenetics*, 64, 39–47.

Celarain, N., Sánchez-Ruiz de Gordoa, J., Zelaya, M. V., Roldán, M., Larumbe, R., Pulido, L., Echavarri, C., and Mendioroz, M. (2016). "Trem2 upregulation correlates with 5-hydroxymethycytosine enrichment in alzheimer's disease hippocampus." *Clinical epigenetics*, 8,

1–10.

Colonna, M. (2023). "The biology of trem receptors." *Nature Reviews Immunology*, 23(9), 580–594.

Daws, M. R., Sullam, P. M., Niemi, E. C., Chen, T. T., Tchao, N. K., and Seaman, W. E. (2003). "Pattern recognition by trem-2: binding of anionic ligands." *The Journal of Immunology*, 171(2), 594–599.

Dickson, D. W. (1999). "Microglia in alzheimer's disease and transgenic models: how close the fit?." *The American journal of pathology*, 154(6), 1627–1631.

Dominguez, C., Boelens, R., and Bonvin, A. M. (2003). "Haddock: a protein- protein docking approach based on biochemical or biophysical information." *Journal of the American Chemical Society*, 125(7), 1731–1737.

D'Andrea, M. R., Cole, G. M., and Ard, M. D. (2004). "The microglial phagocytic role with specific plaque types in the alzheimer disease brain." *Neurobiology of aging*, 25(5), 675–683.

e Research (2025). "King's computational research, engineering and technology environment (create), <https://doi.org/10.18742/rnvf-m076>.

Feuerbach, D., Schindler, P., Barske, C., Joller, S., Beng-Louka, E., Worringer, K. A., Kommineni, S., Kaykas, A., Ho, D. J., Ye, C., et al. (2017). "Adam17 is the main sheddase for the generation of human triggering receptor expressed in myeloid cells (htrem2) ectodomain and cleaves trem2 after histidine 157." *Neuroscience letters*, 660, 109–114.

Giraldo, M., Lopera, F., Siniard, A. L., Corneveaux, J. J., Schrauwen, I., Carvajal, J., Muñoz, C., Ramirez-Restrepo, M., Gaiteri, C., Myers, A. J., et al. (2013). "Variants in triggering receptor expressed on myeloid cells 2 are associated with both behavioral variant frontotemporal lobar degeneration and alzheimer's disease." *Neurobiology of aging*, 34(8), 2077–e11.

Gowers, R. J., Linke, M., Barnoud, J., Reddy, T. J., Melo, M. N., Seyler, S. L., Domanski, J., Dotson, D. L., Buchoux, S., Kenney, I. M., et al. (2016). "Mdanalysis: a python package for the rapid analysis of molecular dynamics simulations." *Proceedings of the 15th python in science conference*, Vol. 98, SciPy Austin, TX, 105.

Guerreiro, R., Bilgic, B., Guven, G., Brás, J., Rohrer, J., Lohmann, E., Hanagasi, H., Gurvit, H.,

and Emre, M. (2013a). "A novel compound heterozygous mutation in trem2 found in a turkish frontotemporal dementia-like family." *Neurobiology of aging*, 34(12), 2890–e1.

Guerreiro, R., Wojtas, A., Bras, J., Carrasquillo, M., Rogaeva, E., Majounie, E., Cruchaga, C., Sassi, C., Kauwe, J. S., Younkin, S., et al. (2013b). "Trem2 variants in alzheimer's disease." *New England Journal of Medicine*, 368(2), 117–127.

Guvench, O., Mallajosyula, S. S., Raman, E. P., Hatcher, E., Vanommeslaeghe, K., Foster, T. J., Jamison, F. W., and MacKerell Jr, A. D. (2011). "Charmm additive all-atom force field for carbohydrate derivatives and its utility in polysaccharide and carbohydrate–protein modeling." *Journal of chemical theory and computation*, 7(10), 3162–3180.

Hamerman, J. A., Jarjoura, J. R., Humphrey, M. B., Nakamura, M. C., Seaman, W. E., and Lanier, L. L. (2006). "Cutting edge: inhibition of tlr and fcr responses in macrophages by triggering receptor expressed on myeloid cells (trem)-2 and dap12." *The Journal of Immunology*, 177(4), 2051–2055.

Hess, B., Bekker, H., Berendsen, H. J., and Fraaije, J. G. (1997). "Lincs: A linear constraint solver for molecular simulations." *Journal of computational chemistry*, 18(12), 1463–1472.

Hospital, A., Goñi, J. R., Orozco, M., and Gelpí, J. L. (2015). "Molecular dynamics simulations: advances and applications." *Advances and Applications in Bioinformatics and Chemistry*, 37–47.

Huang, Y. and Mucke, L. (2012). "Alzheimer Mechanisms and Therapeutic Strategies." *Cell*, 148(6), 1204–1222.

Humphrey, W., Dalke, A., and Schulten, K. (1996). "Vmd: visual molecular dynamics." *Journal of molecular graphics*, 14(1), 33–38.

Hunter, J. D. (2007). "Matplotlib: A 2d graphics environment." *Computing in Science & Engineering*, 9(3), 90–95.

Jin, S. C., Benitez, B. A., Karch, C. M., Cooper, B., Skorupa, T., Carrell, D., Norton, J. B., Hsu, S., Harari, O., Cai, Y., et al. (2014). "Coding variants in trem2 increase risk for alzheimer's disease." *Human molecular genetics*, 23(21), 5838–5846.

Jin, S. C., Carrasquillo, M. M., Benitez, B. A., Skorupa, T., Carrell, D., Patel, D., Lincoln, S.,

Krishnan, S., Kachadoorian, M., Reitz, C., et al. (2015). "Trem2 is associated with increased risk for alzheimer's disease in african americans." *Molecular neurodegeneration*, 10, 1–7.

Jo, S., Kim, T., Iyer, V. G., and Im, W. (2008). "Charmm-gui: a web-based graphical user interface for charmm." *Journal of computational chemistry*, 29(11), 1859–1865.

Jonsson, T., Stefansson, H., Steinberg, S., Jonsdottir, I., Jonsson, P. V., Snaedal, J., Bjornsson, S., Huttenlocher, J., Levey, A. I., Lah, J. J., et al. (2013a). "Variant of trem2 associated with the risk of alzheimer's disease." *New England Journal of Medicine*, 368(2), 107–116.

Jonsson, T., Stefansson, H., Steinberg, S., Jonsdottir, I., Jonsson, P. V., Snaedal, J., Bjornsson, S., Huttenlocher, J., Levey, A. I., Lah, J. J., Rujescu, D., Hampel, H., Giegling, I., Andreassen, O. A., Engedal, K., Ulstein, I., Djurovic, S., Ibrahim-Verbaas, C., Hofman, A., Ikram, M. A., Duijn, C. M. v., Thorsteinsdottir, U., Kong, A., and Stefansson, K. (2013b). "Variant of TREM2 associated with the risk of alzheimer's disease." *New England Journal of Medicine*, 368(2), 107–116 Publisher: Massachusetts Medical Society _eprint: https://www.nejm.org/doi/pdf/10.1056/NEJMoa1211103.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli, P., and Hassabis, D. (2021a). "Highly accurate protein structure prediction with AlphaFold." *Nature*, 596(7873), 583–589.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021b). "Highly accurate protein structure prediction with alphafold." *nature*, 596(7873), 583–589.

Karplus, M. and McCammon, J. A. (2002). "Molecular dynamics simulations of biomolecules." *Nature structural biology*, 9(9), 646–652.

Kawabori, M., Kacimi, R., Kauppinen, T., Calosing, C., Kim, J. Y., Hsieh, C. L., Nakamura, M. C., and Yenari, M. A. (2015). "Triggering receptor expressed on myeloid cells 2 (trem2) deficiency

attenuates phagocytic activities of microglia and exacerbates ischemic damage in experimental stroke." *Journal of Neuroscience*, 35(8), 3384–3396.

Klauda, J. B., Venable, R. M., Freites, J. A., O'Connor, J. W., Tobias, D. J., Mondragon-Ramirez, C., Vorobyov, I., MacKerell Jr, A. D., and Pastor, R. W. (2010). "Update of the charmm all-atom additive force field for lipids: validation on six lipid types." *The journal of physical chemistry B*, 114(23), 7830–7843.

Kleinberger, G., Yamanishi, Y., Suárez-Calvet, M., Czirr, E., Lohmann, E., Cuyvers, E., Struyfs, H., Pettkus, N., Wenninger-Weinzierl, A., Mazaheri, F., et al. (2014). "Trem2 mutations implicated in neurodegeneration impair cell surface transport and phagocytosis." *Science translational medicine*, 6(243), 243ra86–243ra86.

Kober, D. L. and Brett, T. J. (2017). "Trem2-ligand interactions in health and disease." *Journal of molecular biology*, 429(11), 1607–1629.

López-Ríos de Castro, R., Santana-Bonilla, A., Ziolek, R. M., and Lorenz, C. D. (2025). "Automated analysis of soft matter interfaces, interactions, and self-assembly with pysoftk." *J. Chem. Inf. Model.*, 65(4), 1679–1684.

López-Ríos de Castro, R., Ziolek, R. M., , Ulmschneider, M. B., and Lorenz, C. D. (2024). "Therapeutic peptides are preferentially solubilized in specific microenvironments within peg-plga polymer nanoparticles." *Nano Lett.*, 24(6), 2011–2017.

López-Ríos de Castro, R., Ziolek, R. M., and Lorenz, C. D. (2023). "Topology-controlled self-assembly of amphiphilic block copolymers." *Nanoscale*, 15(37), 15230–15237.

Ma, L., Allen, M., Sakae, N., Ertekin-Taner, N., Graff-Radford, N. R., Dickson, D. W., Younkin, S. G., and Sevlever, D. (2016). "Expression and processing analyses of wild type and p. r47h trem2 variant in alzheimer's disease brains." *Molecular neurodegeneration*, 11, 1–9.

MacKerell Jr, A. D., Bashford, D., Bellott, M., Dunbrack Jr, R. L., Evanseck, J. D., Field, M. J., Fischer, S., Gao, J., Guo, H., Ha, S., et al. (1998). "All-atom empirical potential for molecular modeling and dynamics studies of proteins." *The journal of physical chemistry B*, 102(18), 3586–3616.

Malhotra, Y., John, J., Yadav, D., Sharma, D., Rawal, K., Mishra, V., Chaturvedi, N., et al. (2025). "Advancements in protein structure prediction: A comparative overview of alphafold and its derivatives." *Computers in Biology and Medicine*, 188, 109842.

Michaud-Agrawal, N., Denning, E. J., Woolf, T. B., and Beckstein, O. (2011). "Mdanalysis: a toolkit for the analysis of molecular dynamics simulations." *Journal of computational chemistry*, 32(10), 2319–2327.

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022). "ColabFold: making protein folding accessible to all." *Nature Methods*, 19(6), 679–682.

Moutinho, M., Coronel, I., Tsai, A. P., Di Prisco, G. V., Pennington, T., Atwood, B. K., Puntambekar, S. S., Smith, D. C., Martinez, P., Han, S., et al. (2023). "Trem2 splice isoforms generate soluble trem2 species that disrupt long-term potentiation." *Genome medicine*, 15(1), 11.

NOSÉ, S. U. I. (2002). "A molecular dynamics method for simulations in the canonical ensemble." *Molecular physics*, 100(1), 191–198.

Paloneva, J., Manninen, T., Christman, G., Hovanes, K., Mandelin, J., Adolfsson, R., Bianchin, M., Bird, T., Miranda, R., Salmaggi, A., et al. (2002). "Mutations in two genes encoding different subunits of a receptor signaling complex result in an identical disease phenotype." *The American Journal of Human Genetics*, 71(3), 656–662.

Parhizkar, S., Arzberger, T., Brendel, M., Kleinberger, G., Deussing, M., Focke, C., Nuscher, B., Xiong, M., Ghasemigharagoz, A., Katzmarski, N., et al. (2019). "Loss of trem2 function increases amyloid seeding but reduces plaque-associated apoe." *Nature neuroscience*, 22(2), 191–204.

Parrinello, M. and Rahman, A. (1981). "Polymorphic transitions in single crystals: A new molecular dynamics method." *Journal of Applied physics*, 52(12), 7182–7190.

Peng, Q., Malhotra, S., Torchia, J. A., Kerr, W. G., Coggeshall, K. M., and Humphrey, M. B. (2010). "Trem2-and dap12-dependent activation of pi3k requires dap10 and is inhibited by ship1." *Science signaling*, 3(122), ra38–ra38.

Piovesan, D., Minervini, G., and Tosatto, S. C. (2016). "The ring 2.0 web server for high quality

residue interaction networks." *Nucleic acids research*, 44(W1), W367–W374.

Qi, Y., Cheng, X., Han, W., Jo, S., Schulten, K., and Im, W. (2014). "Charmm-gui pace cg builder

for solution, micelle, and bilayer coarse-grained simulations." *Journal of chemical information*

and modeling, 54(3), 1003–1009.

Qi, Y., Ingólfsson, H. I., Cheng, X., Lee, J., Marrink, S. J., and Im, W. (2015). "Charmm-gui

martini maker for coarse-grained simulations with the martini force field." *Journal of chemical*

theory and computation, 11(9), 4486–4494.

Serian, M., Mason, A. J., and Lorenz, C. D. (2024). "Emergent conformational and aggregation

properties of synergistic antimicrobial peptide combinations." *Nanoscale*, 16(44), 20657–20669.

Shoemaker, S. C. and Ando, N. (2018). "X-rays in the cryo-electron microscopy era: structural

biology's dynamic future." *Biochemistry*, 57(3), 277–285.

Siewert, J. M. et al. (2007). "The martini force field: coarse grained model for biomolecular

simulations." *The Journal of Physical Chemistry B*, 111(27), 7812–7824.

Smith, P., Quinn, P. J., and Lorenz, C. D. (2020). "Two coexisting membrane structures are de-

fined by lateral and transbilayer interactions between sphingomyelin and cholesterol." *Langmuir*,

36(33), 9786–9799.

Smith, P., Ziolek, R. M., Gazzarrini, E., Owen, D. M., and Lorenz, C. D. (2019). "On the interaction

of hyaluronic acid with synovial fluid lipid membranes." *Physical Chemistry Chemical Physics*,

21(19), 9845–9857.

Steiner, A., Schlepckow, K., Brunner, B., Steiner, H., Haass, C., and Hagn, F. (2020a). "γ-secretase

cleavage of the alzheimer risk factor trem 2 is determined by its intrinsic structural dynamics."

The EMBO journal, 39(20), e104247.

Steiner, A., Schlepckow, K., Brunner, B., Steiner, H., Haass, C., and Hagn, F. (2020b). "γ-secretase

cleavage of the alzheimer risk factor trem 2 is determined by its intrinsic structural dynamics."

The EMBO journal, 39(20), e104247.

Thornton, P., Sevalle, J., Deery, M. J., Fraser, G., Zhou, Y., Ståhl, S., Franssen, E. H., Dodd,

R. B., Qamar, S., Gomez Perez-Nievas, B., et al. (2017). "Trem 2 shedding by cleavage at the

h157-s158 bond is accelerated for the alzheimer's disease-associated h157y variant." *EMBO molecular medicine*, 9(10), 1366–1378.

Ulland, T. K. and Colonna, M. (2018). "Trem2—a key player in microglial biology and alzheimer disease." *Nature reviews neurology*, 14(11), 667–675.

Ulland, T. K., Song, W. M., Huang, S. C.-C., Ulrich, J. D., Sergushichev, A., Beatty, W. L., Loboda, A. A., Zhou, Y., Cairns, N. J., Kambal, A., et al. (2017). "Trem2 maintains microglial metabolic fitness in alzheimer's disease." *Cell*, 170(4), 649–663.

Ulrich, J. D., Ulland, T. K., Colonna, M., and Holtzman, D. M. (2017). "Elucidating the role of trem2 in alzheimer's disease." *Neuron*, 94(2), 237–248.

Van Der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A. E., and Berendsen, H. J. (2005). "Gromacs: fast, flexible, and free." *Journal of computational chemistry*, 26(16), 1701–1718.

Van Zundert, G., Rodrigues, J., Trellet, M., Schmitz, C., Kastritis, P., Karaca, E., Melquiond, A., van Dijk, M., De Vries, S., and Bonvin, A. (2016). "The haddock2. 2 web server: user-friendly integrative modeling of biomolecular complexes." *Journal of molecular biology*, 428(4), 720–725.

Wang, S., Sudan, R., Peng, V., Zhou, Y., Du, S., Yuede, C. M., Lei, T., Hou, J., Cai, Z., Cella, M., et al. (2022). "Trem2 drives microglia response to amyloid-β via syk-dependent and-independent pathways." *Cell*, 185(22), 4153–4169.

Wang, Y., Cella, M., Mallinson, K., Ulrich, J. D., Young, K. L., Robinette, M. L., Gilfillan, S., Krishnan, G. M., Sudhakar, S., Zinselmeyer, B. H., et al. (2015). "Trem2 lipid sensing sustains the microglial response in an alzheimer's disease model." *Cell*, 160(6), 1061–1071.

Zhang, L., Xiang, X., Li, Y., Bu, G., and Chen, X.-F. (2025). "Trem2 and strem2 in alzheimer's disease: from mechanisms to therapies." *Molecular Neurodegeneration*, 20(1), 43.

Zhong, Z., Ulmschneider, M. B., and Lorenz, C. D. (2024). "Unraveling the molecular dance: Insights into trem2/dap12 complex formation in alzheimer's disease through molecular dynamics simulations." *ACS omega*, 9(26), 28715–28725.

Zhou, R., Yang, G., Guo, X., Zhou, Q., Lei, J., and Shi, Y. (2019). "Recognition of the amyloid

precursor protein by human γ-secretase." *Science*, 363(6428), eaaw0930.

Ziolek, R. M., Santana-Bonilla, A., López-Ríos de Castro, R., K uhn, R., Green, M., and Lorenz, C. D. (2022). "Conformational heterogeneity and interchain percolation revealed in an amorphous conjugated polymer." *ACS Nano*, 16(9), 14432–14442.

Ziolek, R. M., Smith, P., Pink, D. L., Dreiss, C. A., and Lorenz, C. D. (2021). "Unsupervised learning unravels the structure of four-arm and linear block copolymer micelles." *Macromolecules*, 54(8), 3755–3768.

FIGURES AND TABLES

Fig. 1. TREM2-DAP12 complex: sequence features and signalling mechanisms. (A) Schematic of the transmembrane domains of TREM2 (isoform 1, 230 aa) and the DAP12 dimer, highlighting key residues involved in membrane embedding and interaction. (B) Sequence alignment of TREM2 isoform 1 (230 aa) and isoform 2 (219 aa), emphasising differences in the transmembrane regions. (C) Ligand engagement induces conformational changes in TREM2, followed by proteolytic cleavage of the ectodomain by ADAM10/17. (D) TREM2-DAP12 association via salt bridges between transmembrane residues triggers downstream phosphorylation of DAP12 ITAM motifs, leading to recruitment of SYK and other kinases.

Fig. 2. Unsupervised clustering reveals distinct conformational states of TREM2-DAP12 mutants. (A) UMAP (Uniform Manifold Approximation and Projection) plots showing the clustering of conformational states for four TREM2-230 mutants: K186A, W194X, W194A, and K186X. Each point represents a structural snapshot colored by its assigned cluster. (B) Corresponding doughnut charts depicting the relative abundance (percentage) of each conformational cluster identified in panel A. (C) Cluster assignments over time for each mutant in panel A, illustrating the temporal evolution and stability of conformational states across simulation trajectories. (D) UMAP plots of three TREM2-219 constructs: wild-type (WT), W191A, and W191X, showing the distribution of their structural ensembles. (E) Doughnut charts representing the relative population of clusters for the constructs shown in panel D, indicating how mutations alter conformational preferences. (F) Temporal evolution of cluster assignments for the TREM2-219 constructs shown in panel D, highlighting the stability and transitions between conformational states over time.

Fig. 3. Residue-level interaction analysis of TREM2-DAP12 transmembrane mutants. (A) Pairwise residue interaction maps for TREM2 mutants (K186A, K186X, W191A, W191X) in the 230 construct, plotted with TREM2 residues on the x-axis and DAP12 residues on the y-axis. Blue dots indicate interacting residues, and red dots highlight residues identified as important for the interaction. (B) Residue interaction maps for selected TREM2 constructs in the 219 background: wild-type (WT), W191A, and W191X. These maps reveal the altered interaction profiles resulting from specific point mutations and truncations, underscoring the differential contribution of individual residues to the TREM2-DAP12 interface.

F

TREM2	L194	K186	K186	W194	W194	W194	W194	W198
DAP12	I46	D50	T54	T54	I57	A58	V61	Y62
WT-230	🔵	🟩⭐	🟩	🟩	🔵	🔵	🔵	🔵

Fig. 4. Molecular interaction profiles of TREM2-DAP12 transmembrane mutants. (A) Table summarising key residue-residue interactions between TREM2 and DAP12 for various mutants and constructs. Interaction types are colour-coded: hydrophobic (blue circle), π-π stacking (red triangle), hydrogen bonds (green square), and salt bridges (yellow star). (B) Representative structural models illustrating π-π stacking interactions between W194 (TREM2) and Y62 (DAP12) observed in K186A-230. (C) Bar graph quantifying the frequency of different π-π interaction geometries (parallel, T-shaped, or none) across simulation frames. (D) Structural snapshots showing the formation of a salt bridge between K186 (TREM2) and D50 (DAP12) in W194A-230. (E) Histogram showing the distribution of hydrogen bond numbers formed between K186 and D50 across simulations. These analyses demonstrate that specific point mutations significantly alter interaction patterns at the TREM2-DAP12 interface. (F) The Table shows the important residue pairs of WT-230.

Fig. 5. Structural insights into the TREM2-DAP12 transmembrane interactions and stability across mutants. (A) Representative snapshots of the TREM2-DAP12 transmembrane complexes embedded in a lipid bilayer for the TREM2-230 mutants: K186A, K186X, W194A, and W194X. Interacting residues at the interface are shown in surface representation. (B) Structural models of TREM2-219 constructs (WT, W191A, W191X) in complex with DAP12, highlighting the critical interfacial residues. (C) Zoomed-in views of key electrostatic and hydrogen-bond interactions formed between TREM2 and DAP12 residues in W194A-230, WT-219, and W191A-219. Positively and negatively charged regions are indicated. (D, E) Distance measurements over time between the centre of mass of selected TREM2 and DAP12 transmembrane residues from molecular dynamics simulations. (D) shows the distance with the WT-219; (E) shows constructs with the W191X-219 construct. These structural and dynamic analyses reveal that specific point mutations and truncations significantly alter the interfacial interactions and complex stability.

Fig. 6. Root Mean Square Fluctuation (RMSF) profiles of TREM2-DAP12 complexes highlight dynamic changes due to mutations. (A) RMSF plots of residue-wise flexibility for DAP12 and TREM2 in the 230-residue constructs. WT-230 and four mutants (K186A, K186X, W194A, W194X) are compared. Shaded regions denote the DAP12 (gray) and TREM2 (blue) chains. Mutations lead to distinct increases in flexibility, particularly in the transmembrane and cytoplasmic regions. (B) RMSF profiles for the 219-residue constructs: WT-219, W191A-219, and W191X-219.

TABLE 1. Cluster assignments and times for each variant

Variant	Replica 1	Replica 2	Replica 3
K186A-230	Cluster 4 290 μs	Cluster 1 295 μs	Cluster 3 298 μs
K186X-230	Cluster 2 297 μs	Cluster 4 299 μs	Cluster 4 300 μs
W194A-230	Cluster 4 297 μs	Cluster 4 299 μs	Cluster 4 300 μs
W194X-230	Cluster 2 297 μs	Cluster 2 299 μs	Cluster 2 300 μs
WT-219	Cluster 1 198 μs	Cluster 2 199 μs	Cluster 2 200 μs
W191A-219	Cluster 2 198 μs	Cluster 1 199 μs	Cluster 1 200 μs
W191X-219	Cluster 2 197 μs	Cluster 2 198 μs	Cluster 2 200 μs

TABLE 2. Details of each of the coarse-grain MD simulation systems.

Variant	n_{beads}	n_{lipids}	n_{waters}	n_{ions}	Final box dimensions (nm)
K186A-230	24024	310	6713	152	$100.6 \times 100.6 \times 127.4$
K186X-230	19022	310	5075	115	$100.2 \times 100.2 \times 107.0$
W194A-230	23904	310	6673	153	$100.4 \times 100.4 \times 127.2$
W194X-230	19618	310	5267	118	$100.3 \times 100.3 \times 108.7$
WT-219	24875	310	6089	158	$100.5 \times 100.5 \times 130.6$
W191A-219	18802	310	4980	116	$100.5 \times 100.5 \times 106.1$
W191X-219	24558	320	6883	154	$100.7 \times 100.7 \times 128.7$

TABLE 3. Details of each of the all-atom MD simulation systems.

Variant	n_{atoms}	n_{lipids}	n_{waters}	n_{ions}	Final box dimensions (nm)
K186A-230	120584	310	26969	152	$97.9 \times 97.9 \times 130.1$
K186X-230	100554	310	20405	115	$97.7 \times 97.7 \times 106.9$
W194A-230	120055	310	26793	153	$97.4 \times 97.4 \times 130.7$
W194X-230	103026	310	21188	118	$96.1 \times 96.1 \times 113.5$
WT-219	123893	310	28060	158	$98.3 \times 98.3 \times 132.0$
W191A-219	99717	310	20020	116	$99.2 \times 99.2 \times 103.7$
W191X-219	122945	315	27642	154	$98.5 \times 98.5 \times 130.3$