Chromatin accessibility differences between the hybrids of *nasuta-albomicans* complex of *Drosophila*

Radhika Padma, Ajinkya Bharatraj Patil, Nagarjun Vijay*

Computational Evolutionary Genomics Lab, Department of Biological Sciences, IISER Bhopal, Bhauri, Madhya Pradesh, India

*Correspondence: nagarjun@iiserb.ac.in

Running Title: Accessibility differences in hybrids of *Drosophila* species

Abstract

Understanding how chromosomal rearrangements (CRs) interact with epigenetic changes to drive

speciation is a fundamental question in evolutionary biology. CRs are key contributors to

chromosome evolution and can play a pivotal role in reproductive isolation. The *Drosophila nasuta-*

albomicans species complex presents an ideal model to explore this, as *D. albomicans* possesses neo-

sex chromosomes formed by Robertsonian fusion of chr3L, chr3R, and a sex chromosome, unlike its

sister species *D. nasuta*. In this study, we investigate CRs influence on accessible chromatin (AC) and

its relationship with genetic differentiation. We used ATAC-seq to decipher ACs of testis in two

hybrids of *D. albomicans* and *D. nasuta*, and genome-wide fixation index (F_{ST}) scans of *D.*

albomicans and *D. nasuta* to identify regions of genetic differentiation. Our analyses revealed that

chromosome 4 (Muller F) harbors the largest number of differentially accessible regions (~97 Kb),

which coincide with peaks in F_{ST}. Moreover, changes in ACs were associated with differential

transcription factor (TF) binding across the genome. These results suggest that CRs can drive

epigenomic divergence in hybrids, particularly on Muller F, and chromatin-level changes may play a

key role in reproductive isolation. Our study provides an example of how chromosomal and

epigenetic architecture interact in the early stages of speciation.

Keywords: Chromatin accessibility, Drosophila, Epigenetics, Genomic islands, Pioneer TF,

Speciation

Introduction

Genome regulation, a curator of the genome, uses multiple means to initiate and sustain tissue-

specific functionalities in a multicellular organism, thus playing a pivotal role in evolutionary

diversification (Wray 2003). For example, beak morphological differences seen in Darwin's Finches

(Abzhanov et al. 2004); pelvic reduction in sticklebacks (Chan et al. 2010); wing patterns in dipteran

insects (Prud'homme et al. 2007); regulatory divergence within and between species of

Drosophila (Wittkopp et al. 2008). Incidentally, epigenetic components govern genome regulation:

DNA methylation, histone tail modifications, chromatin accessibility, and DNA architecture

(Holtzman and Gersbach 2018). We are primarily interested in chromatin accessibility, a dynamic

property that determines and sustains cellular identity (Klemm et al. 2019). For instance, accessible

chromatin (AC) allows transcriptional and regulatory machinery to interact with the DNA physically,

and they, in turn, are regulated by the nucleosomes (Allis and Jenuwein 2016).

Nucleosome distribution across the genome is an integral part of genome regulation, determining

cells' developmental trajectory and tissue-specific identity. Thus, nucleosome density and depletion

across the genome represent inactive heterochromatin (Sun et al. 2001) and active euchromatin (Lee

et al. 2004). Not all the heterochromatin regions found in the genome are perennial; based on the

nature of inactivation (Richards and Elgin 2002), they can be (a) constitutive, where the inactivation

is permanent and consistent across cell types, and (b) facultative, when the inactivation is transient and a hallmark of cell-type and locus-specific, for example, X-chromosome inactivation (Allshire and Madhani 2018). The constitutive heterochromatin guards the genome against deleterious effects of rearrangement, recombination, and transposons (Janssen et al. 2018). They can breakdown due to chromosomal rearrangement (CR), leading to position effect variegation (PEV); a phenomenon first identified in Drosophila, later in others (Muller 1930; McClintock 1953), which modifies the gene expression of organisms (Elgin and Reuter 2013; Harewood and Fraser 2014). Even though heterochromatin is inaccessible to the transcription factors (TFs), a class of TFs known as pioneer TFs bind to them and change the accessibility (Balsalobre and Drouin 2022; Duan et al. 2021).

CRs as an influential evolutionary player, reshuffle and disrupt gene order/synteny (Bhutkar et al. 2008), resulting in changing the gene expression (Harewood and Fraser 2014; Stewart and Rogers 2019), regulation (Goettel and Messing 2010), phenotype (Muller 1930), adaptation (Guerrero and Kirkpatrick 2014), and speciation (Rieseberg 2001; Ayala and Coluzzi 2005; Kirkpatrick and Barton 2006). Numerous events of CR in insects hint at their possible role in the diversification of insects (de Vos et al. 2020; Lukhtanov et al. 2011; Mathers et al. 2021; Bhutkar et al. 2008; Ranganath and Hfigele 1982). The role of CR in promoting genetic variation and differentiation is complex (Navarro and Barton 2003; Ayala and Coluzzi 2005). Therefore, understanding the evolution of interactions between CRs, chromatin organization, and genome regulation is an important question.

Affordable DNA sequencing technologies allowed large-scale investigation of various aspects of biology, for example, how genetic changes accumulate between species (Holsinger and Weir 2009; Corbett-Detig and Hartl 2012; Poelstra et al. 2014; Kou et al. 2020; Hollox et al. 2022). These studies identified high levels of genetic differentiation in short genomic regions commonly called speciation islands (Turner et al. 2005). These speciation islands are interpreted either as barriers to gene flow (Feder et al. 2012; Poelstra et al. 2014) or due to varying degrees of linked selection (Comeron et al. 2012; Burri et al. 2015; Wolf and Ellegren 2017). In contrast to these widely studied genetic differentiation islands, no study has demonstrated an extreme accumulation of between-species epigenetic changes in genomic islands, to the best of our knowledge. Hence, it is unknown whether such epigenetic islands of speciation exist.

We are interested in effects of CR event seen in *Drosophila albomicans* in hybrids of *D. albomicans* and *D. nasuta* (**Fig. 1A**).Where, *D. albomicans* has a centric fusion of three Muller Elements (CF-3M) leads to forming a neo-sex-chromosome (Ranganath and Hfigele 1982). On the other hand, *D. nasuta,* a closely related species of *D. albomicans,* does not display this CF-3M CR, making them an ideal system for understanding the changes driven by a CR in hybrids. Both species belong to the *D. immigrans* species group in the subgenus *Drosophila.* The *D. immigrans* species group has experienced a burst of speciation events during a short time, resulting in species with little to no

morphological differences and reproductive isolation (Wilson et al. 1969; Kitagawa et al. 1982). In
recent years, *D. albomicans* have garnered ample attention leading to the exploration of the genome
(Zhou et al. 2012), expression profile across tissue types (Gibilisco et al. 2016; Nozawa et al. 2021),
male achiasmy (Satomura and Tamura 2016; Wei and Bachtrog 2019), inversions (Mai and Bachtrog
2021), and transposon density (Wei et al. 2021). However, the effect of centric fusion on regulome
and its significance on speciation is understudied. Similarly, the association of speciation islands with
regulation is not known yet. With this study, we hope to understand the impact of CRs on the genome
and how it changes the chromatin landscape and regulation in hybrids.

To explore the accessibility variation between the hybrids of *D. albomicans* and *D. nasuta*, we have
used the ATAC-seq of adult testes with biological replicates. The genome-wide ATAC-seq landscape
revealed the presence of accessibility variation across chromosomes and implicated Muller F as one
of the prime candidates for an epigenetic speciation island. Further population genetic study of public
datasets found this epigenetic island overlaps with islands of genetic differentiation. Additionally, the
prediction of transcription factor binding sites (TFBS) suggests the accessibility difference between
the two hybrids. Our findings determine that genetic differentiation, accessibility differences, and
TFBS variation are interlinked.

Materials and Methods

ATAC-seq data generation

The hybrids identifying with *D. albomicans* and *D. nasuta* (see **Supplementary Text** for details)
were collected from the Drosophila Stock Center, University of Mysore, India, and maintained on a
standard culture medium at 25°C. The testis of two species was harvested from 5-10 days old males
with PBS (pH 7.4) and stored in Recovery™ Cell Culture Freezing Medium (cat no 12648010,
Gibco). The study consists of four samples, replicates from each species, and each sample consists of
25-30 individual testis. We used Active motif's ATAC-seq kit (cat no. 53150) to create ATAC-seq
libraries of PE-150 with 450±50 bp insert size sequenced on the Illumina NovaSeq 6000. The
sequenced data is available at ENA under the project PRJEB46771 (**Supplemental Table S1**).

Data analysis

Our analysis involves three steps (a) **pre-core analysis**, (b) **core analysis**, and (c) **advanced analysis**
(**Supplemental Fig. 1**), and the analysis pipeline is available on GitHub
(https://github.com/RadPa/ATAC-seq.git).

Pre-core analysis

The raw reads passed through customary FastQC v0.11.9 (Andrews and others 2010), which reported
the presence of adapters. Thus, Cutadapt v2.8 (Martin 2011) trimmed these adapters, and MultiQC

v1.11 (Ewels et al. 2016) aggregated the fastqc reports. The adapter pruned reads aligned to the *D.*

albomicans (RefSeq: GCF_009650485.1_drosAlbom15112) genome using Bowtie v2-2.4.2

(Langmead et al. 2009) with the following parameters -X 750 -very-sensitive. After read alignment,

Samtools v1.13 (Li et al. 2009) removed mitochondrial reads and MarkDuplicates of GATK v4.2.2.0

(Van der Auwera and O'Connor 2020) marked the duplicates. After the initial quality checks, 80% of

reads mapped to the genome of *D. albomicans*, followed by various filtering (**Supplemental Table**

S1). We assessed the library quality and fragment length distribution of aligned reads using the

ATACseqQC (Ou et al. 2018), an R package   (**Supplemental Fig. 2 and 3**). Additionally,

ATACseqQC shifted the aligned reads, i.e., + strand reads offset by +4 bp and – strand reads offset by

−5 bp.

Blacklist

ENCODE (Amemiya et al. 2019) suggests the removal of unstructured high signal regions of the

genome called blacklist for functional genomic analysis before peak calling. We used public DNA-

seq datasets for *D. albomicans* and *D. nasuta* (**Supplemental Table S2**) to establish a blacklist for the

species. The datasets were mapped to *D. albomicans* genome (RefSeq:

GCF_009650485.1_drosAlbom15112) using bwa-mem v0.7.17 (Li and Durbin 2009), Samtools v1.13

used for post-processing of aligned reads, and MarkDuplicates of GATK v4.2.2.0 (Van der Auwera

and O'Connor 2020) removed the optical duplicates. The umap v1.1.1 tool (Karimzadeh et al. 2018)

(https://github.com/hoffmangroup/umap) identifies uniquely mappable regions for the genome, and

Blacklist (https://github.com/Boyle-Lab/Blacklist) utilizes umap regions to build blacklist regions. For

our analysis, we used merged blacklist regions of *D. albomicans* and *D. nasuta*, using bedtools

v2.30.0 (Quinlan and Hall 2010) to represent both species. We estimated 4.9% of the genome as

blacklist and showed the importance of using more datasets to detect such regions (**Supplemental**

Table S3).

Core Analysis

Peak calling

We have used HMMRATAC v1.2.10 (Tarbell and Liu 2019) to identify the accessible chromatin

regions of testis in our datasets. The program uses the reads and genome as requirements; we have

used the shifted reads from ATACseqQC, *D. albomicans* chromosome size, and blacklist, as

described above. Most of the parameters were unchanged except for minimum mapping quality

scores, changed to 20; fold change ranges, -u and -l (**Supplemental Table S4**). We selected the

sensitive models producing the highest number of peaks per dataset for the downstream analysis

(**Supplemental Fig. 4**). The peak files were filtered for nucleosome-free regions, as HMMRATAC

identifies both nucleosome-enriched and nucleosome-free regions in the datasets; based on a
minimum of 10 read support.

Differential Accessibility between two hybrids

To find the presence of differential accessibility, we concatenated and sorted peak files from four
datasets by Unix utility and merged using bedtools. Then bedtools multicov command generated the
count matrix for all the ACs across datasets; the count matrix filtered for regions less than 10 reads
support across four datasets. DESeq2 (Love et al. 2014) and edgeR (Robinson et al. 2010) used the
count-matrix to identify differential accessible regions between two hybrids. The DACs from the
analysis intersected with gene annotations that were enhanced with AGAT v0.8.0 (Dainat et al. 2021)
(https://github.com/NBISweden/AGAT) and sorted by gff3sort v1.0.0 (Zhu et al. 2017)
(https://github.com/billzt/gff3sort).

Advanced analysis

Population genetic analysis

We utilized publicly available whole-genome sequencing (WGS) datasets of various populations of *D.*
albomicans and *D. nasuta* (**Supplemental Table S2**) to understand population genetic statistics. We
mapped the reads to the *D. albomicans* genome (RefSeq: GCF_009650485.1_drosAlbom15112).
Population-specific read groups were added to the bam alignments using AddOrReplaceReadGroups,
and further sequencing duplicates were removed using MarkDuplicates of GATK v4.2.0 (Van der
Auwera and O'Connor 2020). We estimated the folded site frequency spectrum (SFS) for each
population using genotype likelihoods generated by ANGSD v 0.935-53-gf475f10 (Korneliussen et al.
2014). These SFS outputs of each population were used to calculate nucleotide diversity statistics
using the saf2theta command of realSFS. Nucleotide diversity stats were converted into 50Kb non-
overlapping window format using thetastat module of ANGSD. Two-dimensional SFS was obtained
using the realSFS module of ANGSD using overlapping sites for pairwise comparisons. These 2D-
SFS for each pairwise comparison between populations were used to generate per base and 50 kb non-
sliding window F_{ST} metrics using the F_{ST} stats2 module of ANGSD. Generally, sex chromosomes
harbor a higher differentiation than autosomes (Ellegren, H. et al. 2012). We also observed a similar
pattern of higher differentiation within species pairs than between species pairs. We have excluded
chrX from the analyses due to the heterogeneity in population sampling of male and female
individuals.

Transcription factor binding site analysis
To find the footprint of ATAC-seq across samples, we have used ATAC-seq specific digital genomic
footprinting method, TOBIAS v0.13.2 (Bentsen et al. 2020), which discovers differential binding of

transcription factor (TF) between samples. The unshifted filtered reads of replicates merged using Samtools merge, as TOBIAS does not support biological replicates. TOBIAS uses aligned reads, accessible regions identified by HMMRATAC, annotation by UROPA (Kondili et al. 2017), genome, and TF motifs from the JASPAR2022 (Castro-Mondragon et al. 2022) to find the differentially bound TF between two species. The TF network was created using TOBIAS CreateNetwork and visualized in Cytoscape 3.9.1 (Shannon et al. 2003). The network represents only 118 TFs, because of limited annotation in *D. albomicans.*

Results

Differentially accessible Muller F

We found 42,534 Accessible Chromatin (AC) regions across four samples, spanning 77.8% of the genome assembly with varying accessibility (**Fig. 1B**). The autosomes had a higher degree of accessibility (mean: 85.4%) than chrX (64.1%). All the autosomes had a comparable level of accessibility (see **Fig. 1C and Supplemental Table S5**). DESeq2 (149 regions) and edgeR (98 regions) identified the Differentially Accessible Chromatin (DAC) regions between hybrids of *D. albomicans* and *D. nasuta* (**Supplemental Table S6 and S7**). Both tools identified 87 common regions as DAC, which comprise 58.4% and 88.7% of all DACs determined by DESeq2 and edgeR, respectively. The remaining 41.6% and 11.2% of DACs are unique to DESeq2 and edgeR (**Supplemental Table S8**). Most of the unique DACs determined by DESeq2 are present in chr2L (25.8%) and chrX (17.7%), whereas edgeR finds unique regions in chr2L (54.5%). We observed uneven distribution of DACs across the genome, with most regions located on the chr4 (Muller F) (**Supplemental Table S5**), even though it constitutes only 2.4% of the whole genome. Muller F hosts DACs that span 97Kb of its 3.9Mb (**Fig. 2** and **Supplemental Fig. 5**), which entails 48.4% of all DAC regions identified by DESeq2 and 56.41% of edgeR (**Supplemental Table S5**). Details comparing the tools and variation of ACs and DACs across chromosomes are in **Supplemental Table S5**.

Greater accessibility in hybrid identifying with *D. nasuta*

DESeq2 found 149 DAC regions, constituting more than 200Kb, of which 7 (~32Kb) and 142 (~169Kb) are highly accessible in hybrids identifying with *D. albomicans* and *D. nasuta,* respectively (**Fig. 1B and Supplemental Table S6**). Similarly, edgeR identified 98 DAC regions spanning ~130Kb (**Supplemental Table S7)**, where 93 (~124 Kb) and 5 (~22 Kb) regions are highly accessible in hybrids identifying with *D. nasuta* and *D. albomicans,* respectively (**Supplemental Fig. 6**)**.** We have selected DESeq2 results for the subsequent analysis. Out of 149 DACs, 97 DACs (~ 85Kb) occur in genomic regions which lack annotated genes. As a result, 52 DACs host the genes, albeit 48 genes, as the gene length varies, and in some cases, a single DAC hosts more than one gene (**Fig. 2**). The size of the DACs varies, and Muller F hosts a few of the largest DACs; the largest contiguous

DAC is 41kb and contains three annotated genes: *Tdg*, *Pur-α*, and *CG32006*; where *Tdg* and *Pur-α*

are involved in DNA mismatch repairing.

There are 24 DAC regions with more than a three-fold change, where 11 of them are associated with

unplaced scaffolds, 10 of them with Muller F, and chr2L and chr3 have one each. But none of the 21

DACs in Muller F and unplaced scaffolds contain annotated genes; though chr2L does have one

annotated gene, it is uncharacterized (see **Fig. 2B**). A single DAC region of chr3 contains *Sugb* (Sugar

baby) and has higher accessibility in hybrid identifying with *D. nasuta* (see **Fig. 2C**). Genes such as

Ten-a (Tenascin accessory) and *beat-Ib* with roles in axon guidance also have higher accessibility in

D. nasuta (see **Fig. 2A** and **2B**). The gene *qless* involved in the synthesis of the isoprenoid side chain

of Coenzyme Q is also more accessible in hybrid identifying with *D. nasuta* (see **Fig. 2E**). Only one

DAC region with > 1 fold-change has higher accessibility in hybrid identifying with *D. albomicans*

and contains a long non-coding RNA (lncRNA) and the centrosomal protein-encoding gene *CG9170*

(see **Fig. 2A**). The DACs also harbored nine long non-coding RNA (lncRNA); Muller F has four.

These observations showed that the hybrid identifying with *D. nasuta* had a higher magnitude and a

larger accessible genomic fraction, with 95.3% of all DAC found by DESeq2.

DACs overlap the high F_{ST} region in Muller F

The overall autosomal genetic differentiation between populations of the same species (Mean F_{ST}

ranges: 0.1324-0.3971) was much less than between-species differentiation (Mean F_{ST} ranges: 0.37-

0.50) (**Fig. 3A, Supplemental Fig. 6, and Supplemental Fig. 7**). The within-species comparisons in

D. albomicans have relatively higher levels of autosomal differentiation (Mean F_{ST} ranges: 0.15-0.42)

than *D. nasuta* (Mean F_{ST} ranges: 0.18-0.22) (**Supplemental Fig. 7 and Supplemental Table S9**).

Muller F showed the highest differentiation among the autosomes in between-species comparisons

(Mean F_{ST} ranges: 0.5346-0.6162). However, Muller F retains the genome-wide pattern of lower

differentiation between *D. nasuta* populations (Mean F_{ST} ranges: 0.1980-0.2542) compared to *D*

.albomicans populations (Mean F_{ST} ranges: 0.1235-0.4041) (**Figure S7 and Supplemental Table**

S9). Differentiation along Muller F is heterogeneous, with the initial 1.9 Mb having an $F_{ST} < 0.3$ and

the subsequent region having an $F_{ST} > 0.75$.

We identified several overlaps between the differential chromatin accessibility landscape and the high

between-species genomic differentiation regions mostly occurring on Muller F (see **Supplementary**

Figure S6). The highly differentiated distal end of Muller F has reduced genetic diversity (θw) in

both *D. albomicans* and *D. nasuta* (see **Fig. 3B**). In contrast to Muller F, the differentiation peak on

chr2R has a lower genetic diversity (θw) in hybrid identifying with *D. albomicans* compared to

hybrid identifying with *D. nasuta* (see **Fig. 3C**). Numerous DAC regions, including the longest

overlap with the island of elevated genomic differentiation at the end of Muller F. To ensure that these

results are biologically meaningful, we excluded the repeat regions, and the results remained

unchanged. Moreover, we found that the distal end of Muller F has a lower repeat density and higher
exon density than the initial part (**Supplementary Figure S8**).

Greater affinity towards pioneer TFs in hybrid identifying with *D. nasuta*

TOBIAS identified TFBS regions for 150 TF motif profiles across peak regions between two species.
We found that TFs occupy 4.8% of the accessible genome across two species, where TFs occupy
3.22% in hybrid identifying with *D. albomicans*, 3.20% in hybrid identifying with *D. nasuta*, and
1.6% region in either one of the hybrids. TOBIAS calculated the TF activity score across two species
and showed variation in the number of TF-occupied regions (**Table 2** and **Supplemental Table S10,**
S22). The chrX had the fewest TF-bound sites than the Muller F (**Table 3** and **Supplemental Table**
S11), explaining their lowest accessibility (**Figure 1B** and **Table 1**).

We found the highest TF activity scores for 46 TF (**Figure 4A)** and differential binding scores across
hybrids (**Figure 4B**). Furthermore, hybrid identifying with *D. albomicans* showed higher binding
affinity towards 28 TFs; hb, CG11294, CG32105, abd-A, al, H2.0, HGTX, lab, Lim1, otp, repo, unpg,
zen2, mirr, br, C15, lms, Hmx, NK7.1, hbn, slou, CG11085, CG15696, CG32532, CG34031, unc-4,
HHEX, and ubx. Whereas hybrid identifying with *D. nasuta* preferred eight TFs; Trl, Clamp, btd,
odd, opa, Cf2, twi, and zld: a pioneer TF zld (Duan et al. 2021) chromatin remodelers; Trl and
Clamp. The TF-TF network of 118 TFs (**Figure 4C and 4D**) shows that the higher affinity towards a
specific TF changes the number of interacting partners. For example, the Trl network in hybrid
identifying with *D. albomicans* and *D. nasuta* show variation in unique partners and the number of
partners.

Discussion

For the first time, we describe the former sex-chromosome – Muller F (Vicoso and Bachtrog 2013),
driving the differential accessibility between hybrids of the *nasuta-albomicans* complex of
Drosophila: *D. albomicans* and *D. nasuta.* There are few studies on the reversal of sex-chromosomes
to autosomes, an unprecedented event in most organisms (Landeen and Presgraves 2013). Our study
provides evidence for the existence of species that shows the transition of Muller F from sex-
chromosome-like to autosome-like behavior in spermatogenesis. We found that hybrid identifying
with *D. nasuta* has higher accessibility, though hybrid identifying with *D. albomicans* has lesser
accessibility at the Muller F, consistent with another study (Vicoso and Bachtrog 2013). Our results
showed a consistent pattern, where Muller F of hybrid identifying with *D. nasuta* is distinct from
hybrid identifying with *D. albomicans* and other *Drosophila* species. For instance, TFBS prediction
between hybrids implies hybrid identifying with *D. nasuta* binds differentially to the pioneer TFs,
thus supporting the accessibility difference. Additionally, Muller F has higher genetic differentiation
than other autosomes studied, hinting at the presence of genomic islands of differentiation.

The *nasuta-albomicans* complex of *Drosophila*

Even though *D. albomicans* and *D. nasuta* are morphologically indistinguishable, few features set them apart. For instance, CRs (Ranganath and Hfigele 1982), followed by heterochromatin (Ranganath and Hfigele 1982; Wei et al. 2022), and later repeats and transposon density (Wei et al. 2022). A plethora of reasons recommends the consideration of testis in a variety of studies across species (Clark et al. 2007; Zhao et al. 2014; Xia et al. 2020; Witt et al. 2019). Furthermore, they are among the most studied adult tissue systems across species, especially in molecular and evolutionary biology. And are known to be highly transcriptionally active compared to any tissue of an organism (Witt et al. 2019). Our study shows greater accessibility across the genome, as expected from a highly dynamic tissue of an organism, more than 80% in all except one – least accessible chrX. Notably, the chrX undergoes inactivation during spermatogenesis, as illustrated by many studies (Hense et al. 2007; Richler et al. 1992; Turner 2007; Royo et al. 2010). Even though DACs are present across the genome, they nexus on a specific chromosome, Muller F (**Fig. 1B and C**). Even though we set out to study the CRs' role in changing the chromatin accessibility landscape, we were intrigued by Muller F because of their compelling presence in DACs.

Muller F is one of the peculiar chromosomes found in *Drosophila.* They are primarily composed of heterochromatin, hosting less number of genes due to their petite size and lengthier genes; they enlist a vast number of repeats and a higher density of transposons (Funikov et al. 2020; Leung et al. 2015; Wei et al. 2022). For example, distal regions of Muller F of *D. albomicans,* 1.9Mb, showed a higher gene density, 113 genes with an average length of 10.8Kb. Additionally, Muller F's idiosyncratic lineage of evolution (Vicoso and Bachtrog 2013) piques interest in exploring why they host nearly 50% of all identified DACs (**Fig. 1B and C**) and the functional role of the genes it contains. Muller F, in our study, hosts one of the largest DAC with a three-fold difference between hybrids identifying with *D. albomicans* and *D. nasuta*, entailing 41kb of 3.9Mb of its length, and has three annotated genes: *Tdg*, *Pur-α*, and *CG32006*. Recent studies have observed that the *D. nasuta* accommodate high repeat contents compared to *D. albomicans* (Wei et al. 2022), and it's known that the repeats elicit the DNA mismatches (Li et al. 2002; Iyer et al. 2015), repaired by the enzymes like *Tdg* and *Pur-α* (Hardeland et al. 2003; White et al. 2009). This information led us to speculate that the repeats load and the repairing genes could explain the Muller F three-fold differential accessibility.

This observation (**Fig. 1B and C**) led us to suspect that the Muller F could harbor species' genomic islands of differentiation and potentially be a reproductive barrier (Feder et al. 2012). To discern our speculation, we scanned for genome-wide divergence in public datasets of the species (**Supplemental Table S2**). The F_{ST} across the chromosomes showed few regions across the genome, chr2L, chr3, Muller F, chr2R, and chrX; generally, sex chromosomes exhibit high divergence (Wolf and Ellegren

2017). All these regions have higher F$_{ST}$; though sex chromosomes show high divergence, only the

distal end of the Muller F showed fixed divergence (**Fig. 3A, B, and C**), and they also juxtaposed

with the DACs in our study. We have detected differentiation across various genome regions, hinting

that these species might be experiencing low or no gene flow, a characteristic of allopatric species

(Feder et al. 2012).

After assessing the DAC and fixed divergence across two hybrids, we want to identify if these

elements affect the TF binding. Indeed, we found differences, specifically in hybrid identifying with

D. nasuta, involving two pioneer TFs: ZLD and CLAMP. Pioneer TFs are the only TFs that can bind

to nucleosomal DNA, and modulate for chromatin accessibility, thus associated with open chromatin

and active transcription (McPherson et al. 1993; Zaret 2020). ZLD and CLAMP are the drivers of

Drosophila zygotic genome activation (ZGA), and ZLD is necessary for hundreds of regions' open

chromatin (Schulz et al. 2015). Thus, pioneer TFs contribute towards higher accessibility observed in

hybrid identifying with *D. nasuta*. Additionally, hybrid identifying with *D. nasuta* also has a

higher binding affinity for TRL or GAF (Trithorax-like), which belongs to Trithorax (TrxG) group

proteins. TrxG and Polycomb group (PcG) can regulate their target at multiple levels, modifying the

local chromatin to higher chromatin order and global genome structure (Schuettengruber et al. 2017).

TRL is involved in modifying the accessibility of promoters, allowing other TFs to bind and start

transcription (Farkas et al. 1994). We believe that the accessibility changes seen in hybrid

identifying with *D. nasuta* are a culmination of the higher activity of these TFs.

Limitations: During the analysis of the ATAC-seq data using different reference genomes, it was

noticed that differential mapping bias towards the parental species affects specific genomic regions.

Since the hybrids used in this study have differing levels of hybrid ancestry (see **Supplementary**

Text), disentangling the effect of mapping bias from biological differences in accessibility remains

challenging. Future studies with high-quality reference genomes and sophisticated tools to overcome

the challenges of mapping bias in hybrids will be crucial for further progress. Although the mapping

bias observed could be responsible for some of the biological patterns observed, we can rule out the

existence of a distinctive biological signal. Therefore, we urge caution in the interpretation of these

results.

Conclusion

We set out to understand the interplay between epigenetics, chromosomal rearrangement, and

speciation using ATAC-seq and DNA-seq of members of the *nasuta-albomicans* complex of

Drosophila and hybrids. Our results showed chromatin accessibility differentiation, one of the many

components of epigenetics, between the two hybrids. Though we found few DACs in the

Robertsonian fused chromosome, Muller F, one of the smallest assembled chromosomes, drew our

attention. Due to the density of DACs identified and because it's one of the notable karyotypic differences found in *D. albomicans* and *D. nasuta*. *D. albomicans* hosts a lengthier and heterochromatinized version of Muller F than one in *D. nasuta*. The result reemphasizes the observation with the juxtaposed region of significantly higher accessibility and genetic differentiation in hybrid identifying with *D. nasuta*. The higher accessibility in hybrid identifying with *D. nasuta* is probably due to their strong affinity towards binding pioneer TFs and chromatin modifiers, like CLAMP, TRL, and ZLD. Our main objective was to understand how CRs and ACs affect the speciation in general and precisely how they shape the genome of hybrids of closely related species, *D. albomicans* and *D. nasuta*. Our findings conclude that the first difference that could arise in heterochromatin. We urge future studies to carefully observe the interaction of karyotype, diversity of accessibility and genetic variation, and TF motifs to understand the speciation.

Funding

This article was funded by the WOS-A Project - Life Sciences, Department of Science and Technology, India (Grant no. SR/WOS-A/LS-225/2018(G)).

Authors contributions.

R.P. designed the study, obtained funding and performed all the experiments. R.P. wrote the manuscript with inputs from A.B.P and N.V. R.P. analyzed all the ATAC-seq data. A.B.P performed the population genetic analysis and provided crucial inputs on the integration of diverse analysis. All authors reviewed the manuscript.

Competing interests.

The authors declare no competing financial interests.

Data accessibility.

The associated data is available in an easy-to-view format on the GitHub repository: https://github.com/RadPa/Chromatin-accessibility-of-hybrids-of-D.-albomicans-and-D.-nasuta.git.

References:

Abzhanov A, Protas M, Grant BR, Grant PR, Tabin CJ. 2004. Bmp4 and Morphological Variation of Beaks in Darwin's Finches. *Science (80-)* **305**: 1462–1465. www.science.org/cgi/content/full/305/5689/1457/ (Accessed March 10, 2022).

Allis CD, Jenuwein T. 2016. The molecular hallmarks of epigenetic control. *Nat Rev Genet 2016 178* **17**: 487–500. https://www.nature.com/articles/nrg.2016.59 (Accessed May 3, 2022).

Allshire RC, Madhani HD. 2018. Ten principles of heterochromatin formation and function. *Nat Rev*
Mol Cell Biol **19**: 229–244. https://www.nature.com/articles/nrm.2017.119 (Accessed March 12,
2022).

Amemiya HM, Kundaje A, Boyle AP. 2019. The ENCODE Blacklist: Identification of Problematic
Regions of the Genome. *Sci Rep* **9**: 9354. https://www.nature.com/articles/s41598-019-45839-z
(Accessed October 27, 2021).

Andrews S, others. 2010. FastQC: a quality control tool for high throughput sequence data. 2010.
Https://WwwBioinformaticsBabrahamAcUk/Projects/Fastqc/
http://www.bioinformatics.babraham.ac.uk/projects/.
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (Accessed November 20, 2021).

Ayala FJ, Coluzzi M. 2005. Chromosome speciation: Humans, Drosophila, and mosquitoes. *Proc Natl*
Acad Sci **102**: 6535–6542. http://www.pnas.org/cgi/doi/10.1073/pnas.0501847102 (Accessed
August 24, 2019).

Balsalobre A, Drouin J. 2022. Pioneer factors as master regulators of the epigenome and cell fate. *Nat*
Rev Mol Cell Biol 2022 1–16. https://www.nature.com/articles/s41580-022-00464-z (Accessed
May 22, 2022).

Bentsen M, Goymann P, Schultheis H, Klee K, Petrova A, Wiegandt R, Fust A, Preussner J, Kuenne
C, Braun T, et al. 2020. ATAC-seq footprinting unravels kinetics of transcription factor binding
during zygotic genome activation. *Nat Commun* **11**: 4267.
https://www.nature.com/articles/s41467-020-18035-1 (Accessed August 18, 2021).

Bhutkar A, Schaeffer SW, Russo SM, Xu M, Smith TF, Gelbart WM. 2008. Chromosomal
Rearrangement Inferred From Comparisons of 12 Drosophila Genomes. *Genetics* **179**: 1657–
1680. https://academic.oup.com/genetics/article/179/3/1657/6063947 (Accessed October 25,
2019).

Burri R, Nater A, Kawakami T, Mugal CF, Olason PI, Smeds L, Suh A, Dutoit L, Bureš S,
Garamszegi LZ, et al. 2015. Linked selection and recombination rate variation drive the
evolution of the genomic landscape of differentiation across the speciation continuum of
Ficedula flycatchers. *Genome Res* **25**: 1656–1665.
https://genome.cshlp.org/content/25/11/1656.full (Accessed April 28, 2022).

Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, Lemma RB, Turchi L, Blanc-Mathieu R,
Lucas J, Boddie P, Khan A, As Manosalva P ´Erez N, et al. 2022. JASPAR 2022: the 9th
release of the open-access database of transcription factor binding profiles. *Nucleic Acids Res*
50: D165–D173. https://academic.oup.com/nar/article/50/D1/D165/6446529 (Accessed

February 17, 2022).

Chan YF, Marks ME, Jones FC, Villarreal G, Shapiro MD, Brady SD, Southwick AM, Absher DM,
Grimwood J, Schmutz J, et al. 2010. Adaptive Evolution of Pelvic Reduction in Sticklebacks by
Recurrent Deletion of a Pitx1 Enhancer. *Science (80-)* **327**: 302–305.
https://www.science.org/doi/10.1126/science.1182213 (Accessed March 10, 2022).

Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis M,
Gelbart W, Iyer VN, et al. 2007. Evolution of genes and genomes on the Drosophila phylogeny.
Nature **450**: 203–218. https://www.nature.com/articles/nature06341 (Accessed February 17,
2022).

Comeron JM, Ratnappan R, Bailin S. 2012. The Many Landscapes of Recombination in Drosophila
melanogaster ed. D.A. Petrov. *PLoS Genet* **8**: e1002905.
https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1002905 (Accessed April
28, 2022).

Corbett-Detig RB, Hartl DL. 2012. Population Genomics of Inversion Polymorphisms in Drosophila
melanogaster. *PLOS Genet* **8**: e1003056.
https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1003056 (Accessed March
12, 2022).

Dainat J, Hereñú D, pascal-git. 2021. NBISweden/AGAT: AGAT-v0.8.0.
https://zenodo.org/record/5336786 (Accessed October 27, 2021).

de Vos JM, Augustijnen H, Bätscher L, Lucek K. 2020. Speciation through chromosomal fusion and
fission in Lepidoptera. *Philos Trans R Soc B Biol Sci* **375**: 20190539.
https://royalsocietypublishing.org/doi/abs/10.1098/rstb.2019.0539 (Accessed September 28,
2021).

Duan JE, Rieder LE, Colonnetta MM, Huang A, McKenney M, Watters S, Deshpande G, Jordan WT,
Fawzi NL, Larschan EN. 2021. Clamp and zelda function together to promote drosophila
zygotic genome activation. *Elife* **10**.

Elgin SCR, Reuter G. 2013. Position-Effect Variegation, Heterochromatin Formation, and Gene
Silencing in Drosophila. *Cold Spring Harb Perspect Biol* **5**: a017780–a017780.
http://cshperspectives.cshlp.org/lookup/doi/10.1101/cshperspect.a017780 (Accessed October 25,
2019).

Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis results for multiple
tools and samples in a single report. *Bioinformatics* **32**: 3047–3048.
https://academic.oup.com/bioinformatics/article/32/19/3047/2196507 (Accessed October 27,

2021).

Farkas G, Gausz J, Galloni M, Reuter G, Gyurkovics H, Karch F. 1994. The Trithorax-like gene
encodes the Drosophila GAGA factor. *Nature* **371**: 806–808.
https://pubmed.ncbi.nlm.nih.gov/7935842/ (Accessed February 23, 2022).

Feder JL, Egan SP, Nosil P. 2012. *The genomics of speciation-with-gene-flow*. Elsevier Current
Trends https://www.sciencedirect.com/science/article/pii/S0168952512000492 (Accessed
October 1, 2019).

Funikov SY, Rezvykh AP, Kulikova DA, Zelentsova ES, Protsenko LA, Chuvakova LN, Tyukmaeva
VI, Arkhipova IR, Evgen'ev MB. 2020. Adaptation of gene loci to heterochromatin in the course
of Drosophila evolution is associated with insulator proteins. *Sci Rep* **10**: 11893.
https://www.nature.com/articles/s41598-020-68879-2 (Accessed September 1, 2021).

Gibilisco L, Zhou Q, Mahajan S, Bachtrog D. 2016. Alternative Splicing within and between
Drosophila Species, Sexes, Tissues, and Developmental Stages ed. G.S. Barsh. *PLOS Genet* **12**:
e1006464. https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1006464
(Accessed November 17, 2021).

Goettel W, Messing J. 2010. Divergence of gene regulation through chromosomal rearrangements.
BMC Genomics **11**: 1–19. https://bmcgenomics.biomedcentral.com/articles/10.1186/1471-2164-
11-678 (Accessed November 20, 2021).

Guerrero RF, Kirkpatrick M. 2014. LOCAL ADAPTATION AND THE EVOLUTION OF
CHROMOSOME FUSIONS. *Evolution (N Y)* **68**: 2747–2756.
https://onlinelibrary.wiley.com/doi/full/10.1111/evo.12481 (Accessed November 8, 2021).

Hardeland U, Bentele M, Jiricny J, Schär P. 2003. The versatile thymine DNA-glycosylase: A
comparative characterization of the human, Drosophila and fission yeast orthologs. *Nucleic
Acids Res* **31**: 2261–2271. https://academic.oup.com/nar/article/31/9/2261/1080396 (Accessed
May 27, 2022).

Harewood L, Fraser P. 2014. The impact of chromosomal rearrangements on regulation of gene
expression. *Hum Mol Genet* **23**: R76–R82.
https://academic.oup.com/hmg/article/23/R1/R76/2900848 (Accessed September 28, 2021).

Hense W, Baines JF, Parsch J. 2007. X Chromosome Inactivation during Drosophila Spermatogenesis
ed. M.A.F. Noor. *PLoS Biol* **5**: e273.
https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.0050273 (Accessed
February 7, 2022).

Hollox EJ, Zuccherato LW, Tucci S. 2022. *Genome structural variation in human evolution*. Elsevier

http://www.cell.com/article/S0168952521001852/fulltext (Accessed March 12, 2022).

Holsinger KE, Weir BS. 2009. Genetics in geographically structured populations: defining, estimating

and interpreting FST. *Nat Rev Genet* **10**: 639–650. https://www.nature.com/articles/nrg2611

(Accessed March 12, 2022).

Holtzman L, Gersbach CA. 2018. Editing the epigenome: Reshaping the genomic landscape. *Annu*

Rev Genomics Hum Genet **19**: 43–71. https://www.annualreviews.org/doi/abs/10.1146/annurev-

genom-083117-021632 (Accessed May 3, 2022).

Iyer RR, Pluciennik A, Napierala M, Wells RD. 2015. DNA Triplet Repeat Expansion and Mismatch

Repair. *Annu Rev Biochem* **84**: 199–226.

https://www.annualreviews.org/doi/abs/10.1146/annurev-biochem-060614-034010 (Accessed

May 31, 2022).

Janssen A, Colmenares SU, Karpen GH. 2018. Heterochromatin: Guardian of the Genome. *Annu Rev*

Cell Dev Biol **34**: 265–288. https://doi.org/10.1146/annurev-cellbio-100617- (Accessed March

11, 2022).

Karimzadeh M, Ernst C, Kundaje A, Hoffman MM. 2018. Umap and Bismap: quantifying genome

and methylome mappability. *Nucleic Acids Res* **46**: e120–e120.

https://academic.oup.com/nar/article/46/20/e120/5086676 (Accessed October 27, 2021).

Kirkpatrick M, Barton N. 2006. Chromosome Inversions, Local Adaptation and Speciation. *Genetics*

173: 419–434. https://academic.oup.com/genetics/article/173/1/419/6061572 (Accessed March

12, 2022).

Kitagawa O, Wakahama K-I, Fuyama Y, Shimada Y, Takanashi E, Hatsumi M, Uwabo M, Mita Y.

1982. Genetic studies of the Drosophila nasuta subgroup, with notes on distribution and

morphology. *Japanese J Genet* **57**: 113–141.

http://www.jstage.jst.go.jp/article/ggs1921/57/2/57_2_113/_article (Accessed November 16,

2021).

Klemm SL, Shipony Z, Greenleaf WJ. 2019. *Chromatin accessibility and the regulatory epigenome*.

Nature Publishing Group www.nature.com/nrg (Accessed October 17, 2019).

Kondili M, Fust A, Preussner J, Kuenne C, Braun T, Looso M. 2017. UROPA: A tool for Universal

RObust Peak Annotation. *Sci Rep* **7**: 1–12. https://www.nature.com/articles/s41598-017-02464-y

(Accessed February 17, 2022).

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: Analysis of Next Generation Sequencing

Data. *BMC Bioinformatics* **15**.
https://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-014-0356-4 (Accessed
October 27, 2021).

Kou Y, Liao Y, Toivainen T, Lv Y, Tian X, Emerson JJ, Gaut BS, Zhou Y. 2020. Evolutionary
Genomics of Structural Variation in Asian Rice (Oryza sativa) Domestication. *Mol Biol Evol* **37**:
3507–3524. https://academic.oup.com/mbe/article/37/12/3507/5873527 (Accessed March 12,
2022).

Landeen EL, Presgraves DC. 2013. Evolution: From autosomes to sex chromosomes - And back.
Curr Biol **23**: R848–R850. https://linkinghub.elsevier.com/retrieve/pii/S0960982213010270
(Accessed June 2, 2022).

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. *Genome Biol* **10**: R25.
https://genomebiology.biomedcentral.com/articles/10.1186/gb-2009-10-3-r25 (Accessed
October 27, 2021).

Lee CK, Shibata Y, Rao B, Strahl BD, Lieb JD. 2004. Evidence for nucleosome depletion at active
regulatory regions genome-wide. *Nat Genet* **36**: 900–905.
https://www.nature.com/articles/ng1400 (Accessed March 11, 2022).

Leung W, Shaffer CD, Reed LK, Smith ST, Barshop W, Dirkes W, Dothager M, Lee P, Wong J,
Xiong D, et al. 2015. Drosophila Muller F Elements Maintain a Distinct Set of Genomic
Properties Over 40 Million Years of Evolution. *G3 Genes|Genomes|Genetics* **5**: 719–740.
http://www.g3journal.org/lookup/suppl/doi:10.1534/g3.114.015966/-/DC1 (Accessed November
25, 2020).

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**: 1754–1760.
https://academic.oup.com/bioinformatics/article/25/14/1754/225615 (Accessed November 20,
2021).

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009.
The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.
https://academic.oup.com/bioinformatics/article/25/16/2078/204688 (Accessed October 27,
2021).

Li YC, Korol AB, Fahima T, Beiles A, Nevo E. 2002. Microsatellites: Genomic distribution, putative
functions and mutational mechanisms: A review. *Mol Ecol* **11**: 2453–2465.
https://onlinelibrary.wiley.com/doi/full/10.1046/j.1365-294X.2002.01643.x (Accessed May 31,

2022).

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**: 550.
https://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-0550-8 (Accessed
October 27, 2021).

Lukhtanov VA, Dincă V, Talavera G, Vila R. 2011. Unprecedented within-species chromosome
number cline in the Wood White butterfly Leptidea sinapis and its significance for karyotype
evolution and speciation. *BMC Evol Biol* **11**: 109.
https://bmcecolevol.biomedcentral.com/articles/10.1186/1471-2148-11-109 (Accessed
November 16, 2021).

Mai D, Bachtrog D. 2021. Molecular characterization of inversion breakpoints in the Drosophila
nasuta species group. *bioRxiv* 2021.06.01.446624.
https://www.biorxiv.org/content/10.1101/2021.06.01.446624v1 (Accessed March 13, 2022).

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**: 10. https://github.com/marcelm/cutadapt (Accessed November 20, 2021).

Mathers TC, Wouters RHM, Mugford ST, Swarbreck D, van Oosterhout C, Hogenhout SA. 2021.
Chromosome-Scale Genome Assemblies of Aphids Reveal Extensively Rearranged Autosomes
and Long-Term Conservation of the X Chromosome ed. A. Ouangraoua. *Mol Biol Evol* **38**: 856–
875. https://academic.oup.com/mbe/article/38/3/856/5910553 (Accessed November 16, 2021).

McClintock B. 1953. Induction of Instability at Selected Loci in Maize. *Genetics* **38**: 579–599.
https://pubmed.ncbi.nlm.nih.gov/17247459/ (Accessed November 20, 2021).

McPherson CE, Shim EY, Friedman DS, Zaret KS. 1993. An active tissue-specific enhancer and
bound transcription factors existing in a precisely positioned nucleosomal array. *Cell* **75**: 387–
398.

Muller HJ. 1930. Types of visible variations induced by X-rays inDrosophila. *J Genet* **22**: 299–334.
https://link.springer.com/article/10.1007/BF02984195 (Accessed November 20, 2021).

Navarro A, Barton NH. 2003. Chromosomal Speciation and Molecular Divergence--Accelerated
Evolution in Rearranged Chromosomes. *Science (80-)* **300**: 321–324.
https://www.science.org/doi/abs/10.1126/science.1080600 (Accessed April 28, 2022).

Nozawa M, Minakuchi Y, Satomura K, Kondo S, Toyoda A, Tamura K. 2021. Shared evolutionary
trajectories of three independent neo-sex chromosomes in Drosophila. *Genome Res* **31**: 2069–
2079. https://genome.cshlp.org/content/31/11/2069.full (Accessed November 17, 2021).

Ou J, Liu H, Yu J, Kelliher MA, Castilla LH, Lawson ND, Zhu LJ. 2018. ATACseqQC: a
Bioconductor package for post-alignment quality assessment of ATAC-seq data. *BMC*
Genomics **19**: 169. https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-018-4559-
3 (Accessed June 28, 2019).

Poelstra JW, Vijay N, Bossu CM, Lantz H, Ryll B, Müller I, Baglione V, Unneberg P, Wikelski M,
Grabherr MG, et al. 2014. The genomic landscape underlying phenotypic integrity in the face of
gene flow in crows. *Science (80-)* **344**: 1410–1414.

Prud'homme B, Gompel N, Carroll SB. 2007. Emerging principles of regulatory evolution. *Proc Natl*
Acad Sci **104**: 8605–8612. www.nasonline.org/adaptation_and_complex_design. (Accessed
March 10, 2022).

Quinlan AR, Hall IM. 2010. BEDTools: A flexible suite of utilities for comparing genomic features.
Bioinformatics **26**: 841–842. https://academic.oup.com/bioinformatics/article/26/6/841/244688
(Accessed May 30, 2019).

Ranganath HA, Hfigele K. 1982. The Chromosomes of Two Drosophila Races: D. nasuta nasuta and
D. n. albomicana I. Distribution and Differentiation of Heterochromatin *. *Chromosom* **85**: 83–
92.

Richards EJ, Elgin SCR. 2002. Epigenetic Codes for Heterochromatin Formation and Silencing:
Rounding up the Usual Suspects. *Cell* **108**: 489–500.

Richler C, Soreq H, Wahrman J. 1992. X inactivation in mammalian testis is correlated with inactive
X–specific transcription. *Nat Genet* **2**: 192–195. https://www.nature.com/articles/ng1192-192
(Accessed February 17, 2022).

Rieseberg LH. 2001. Chromosomal rearrangements and speciation. *Trends Ecol Evol* **16**: 351–358.
http://www.cell.com/article/S0169534701021875/fulltext (Accessed September 27, 2021).

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.
https://academic.oup.com/bioinformatics/article/26/1/139/182458 (Accessed February 17, 2022).

Royo H, Polikiewicz G, Mahadevaiah SK, Prosser H, Mitchell M, Bradley A, de Rooij DG, Burgoyne
PS, Turner JMA. 2010. Evidence that Meiotic Sex Chromosome Inactivation Is Essential for
Male Fertility. *Curr Biol* **20**: 2117–2123.
https://linkinghub.elsevier.com/retrieve/pii/S0960982210014351 (Accessed February 17, 2022).

Satomura K, Tamura K. 2016. Ancient Male Recombination Shaped Genetic Diversity of Neo-Y
Chromosome in Drosophila albomicans. *Mol Biol Evol* **33**: 367–374.

https://academic.oup.com/mbe/article/33/2/367/2579414 (Accessed November 17, 2021).

Schuettengruber B, Bourbon H-M, Di Croce L, Cavalli G. 2017. Genome Regulation by Polycomb
and Trithorax: 70 Years and Counting. *Cell* **171**: 34–57.
https://linkinghub.elsevier.com/retrieve/pii/S0092867417308905 (Accessed February 23, 2022).

Schulz KN, Bondra ER, Moshe A, Villalta JE, Lieb JD, Kaplan T, McKay DJ, Harrison MM. 2015.
Zelda is differentially required for chromatin accessibility, transcription factor binding, and gene
expression in the early Drosophila embryo. *Genome Res* **25**: 1715–1726.
https://genome.cshlp.org/content/25/11/1715.full (Accessed March 22, 2022).

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T.
2003. Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction
Networks. *Genome Res* **13**: 2498–2504. https://genome.cshlp.org/content/13/11/2498.full
(Accessed February 17, 2022).

Stewart NB, Rogers RL. 2019. Chromosomal rearrangements as a source of new gene formation in
Drosophila yakuba ed. H.S. Malik. *PLOS Genet* **15**: e1008314.
https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1008314 (Accessed
September 1, 2021).

Sun F-L, Cuaycong MH, Elgin SCR. 2001. Long-Range Nucleosome Ordering Is Associated with
Gene Silencing in Drosophila melanogaster Pericentric Heterochromatin. *Mol Cell Biol* **21**:
2867–2879. /pmc/articles/PMC86916/ (Accessed March 11, 2022).

Tarbell ED, Liu T. 2019. HMMRATAC: a Hidden Markov ModeleR for ATAC-seq. *Nucleic Acids*
Res **47**: e91–e91. https://academic.oup.com/nar/article/47/16/e91/5519166 (Accessed October
27, 2021).

Turner JMA. 2007. Meiotic sex chromosome inactivation. *Development* **134**: 1823–1831.
https://journals.biologists.com/dev/article/134/10/1823/52756/Meiotic-sex-chromosome-
inactivation (Accessed February 17, 2022).

Turner TL, Hahn MW, Nuzhdin S V. 2005. Genomic Islands of Speciation in Anopheles gambiae ed.
N. Barton. *PLoS Biol* **3**: e285.
https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.0030285 (Accessed April
28, 2022).

Van der Auwera GA, O'Connor BD. 2020. Genomics in the Cloud.
https://www.oreilly.com/library/view/genomics-in-the/9781491975183/ (Accessed October 27,
2021).

Vicoso B, Bachtrog D. 2013. Reversal of an ancient sex chromosome to an autosome in Drosophila.

Nature **499**: 332–335. https://www.nature.com/articles/nature12235 (Accessed March 19, 2021).

Wei KH-C, Mai D, Chatla K, Bachtrog D. 2021. Dynamics and impacts of transposable element

proliferation during the Drosophila nasuta species group radiation. *bioRxiv* 2021.08.12.456169.

https://www.biorxiv.org/content/10.1101/2021.08.12.456169v1 (Accessed February 18, 2022).

Wei KH-C, Mai D, Chatla K, Bachtrog D. 2022. Dynamics and Impacts of Transposable Element

Proliferation in the Drosophila nasuta Species Group Radiation ed. R. Rogers. *Mol Biol Evol* **39**.

https://academic.oup.com/mbe/article/39/5/msac080/6575839 (Accessed May 30, 2022).

Wei KHC, Bachtrog D. 2019. Ancestral male recombination in Drosophila albomicans produced

geographically restricted neo-Y chromosome haplotypes varying in age and onset of decay ed.

K.A. Dyer. *PLOS Genet* **15**: e1008502. https://doi.org/10.1371/journal.pgen.1008502 (Accessed

November 26, 2020).

White MK, Johnson EM, Khalili K. 2009. Multiple roles for Purα in cellular and viral regulation. *Cell*

Cycle **8**: 414–420. https://www.tandfonline.com/doi/abs/10.4161/cc.8.3.7585 (Accessed May 30,

2022).

Wilson FD, Wheeler MR, Harget M, Kambysellis M. 1969. Cytogenetic relations in the Drosophila

nasuta subgroup of the immigrans group of species. *Univ Texas Publ* **6918**: 207–253.

https://ci.nii.ac.jp/naid/10015117093 (Accessed November 16, 2021).

Witt E, Benjamin S, Svetec N, Zhao L. 2019. Testis single-cell RNA-seq reveals the dynamics of de

novo gene transcription and germline mutational bias in Drosophila. *Elife* **8**.

https://elifesciences.org/articles/47138 (Accessed August 2, 2021).

Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression differences

within and between Drosophila species. *Nat Genet* **40**: 346–350.

https://www.nature.com/articles/ng.77 (Accessed March 10, 2022).

Wolf JBW, Ellegren H. 2017. Making sense of genomic islands of differentiation in light of

speciation. *Nat Rev Genet* **18**: 87–100. http://www.nature.com/articles/nrg.2016.133 (Accessed

May 23, 2019).

Wray GA. 2003. The Evolution of Transcriptional Regulation in Eukaryotes. *Mol Biol Evol* **20**: 1377–

1419. https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msg140 (Accessed

December 19, 2019).

Xia B, Yan Y, Baron M, Wagner F, Barkley D, Chiodin M, Kim SY, Keefe DL, Alukal JP, Boeke JD,

et al. 2020. Widespread Transcriptional Scanning in the Testis Modulates Gene Evolution Rates.

Cell **180**: 248-262.e21. https://linkinghub.elsevier.com/retrieve/pii/S0092867419313777

(Accessed February 17, 2022).

Zaret KS. 2020. Pioneer Transcription Factors Initiating Gene Network Changes. *Annu Rev Genet* **54**.

/pmc/articles/PMC7900943/ (Accessed February 15, 2022).

Zhao L, Saelao P, Jones CD, Begun DJ. 2014. Origin and spread of de novo genes in Drosophila

melanogaster populations. *Science (80-)* **343**: 769–772. https://www.science.org (Accessed

February 17, 2022).

Zhou Q, Zhu H, Huang Q, Zhao L, Zhang G, Roy SW, Vicoso B, Xuan Z, Ruan J, Zhang Y, et al.

2012. Deciphering neo-sex and B chromosome evolution by the draft genome of Drosophila

albomicans. *BMC Genomics* **13**: 109. https://link.springer.com/articles/10.1186/1471-2164-13-

109 (Accessed November 17, 2021).

Zhu T, Liang C, Meng Z, Guo S, Zhang R. 2017. GFF3sort: a novel tool to sort GFF3 files for tabix

indexing. *BMC Bioinformatics* **18**: 482.

https://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-017-1930-3 (Accessed

October 27, 2021).

Figure Legends:

Figure 1: Karyotype and testes' chromatin landscape of *D. albomicans* and *D. nasuta*.

A Karyotype of male and female *D. albomicans* and *D. nasuta*. The karyotype of *D. nasuta* shows

ancestral chromosome complement, and *D.albomicans* shows rearranged chromosomes. Where

rearrangement of chromosomes leads to the formation of neo-sex-chromosomes by the fusion of

Muller elements A, C, and D. The chromosome arms colored with corresponding Muller elements in

both the species. **B** Accessible chromatin (ACs) of testes in hybrids identifying with *D. albomicans*

and *D. nasuta,* the outermost and second outermost tracks represent chromosomes, and non-accessible

chromatin density, respectively. The innermost track represents differential accessibility (DACs), blue

and red bars represent high and low accessibility regions in hybrid identifying with *D. albomicans*.

The second innermost track represents the transposable element density. The chromosome colors

depicted Muller elements, except for chromosome 3; due to the fusion of Muller elements (C and D)

and unplaced scaffolds, the scaffold of the largest size (i.e., U2) represented in a different color from

others. **C** Chromosome sizes with ACs and DACs by DESeq2 across *D. albomicans* genome.

Figure 2: **Volcano plots of differentially accessible chromatins (DACs)**

A to **F** DACs of chromosome X, 2L, 3, 4, and 2R, respectively, the x-axis denotes the log2-fold-

change, while the y-axis denotes -log10 of the p-value. **C** shows the y-axis break between 38 and 76,

and **E** shows the y-axis break between 8 and 16. The chromosomes are represented in Muller colors,

except chromosome 3 illustrated in turquoise, unplaced scaffolds in tan, and scaffold 2 in misty rose. The points represent the ACs, and the labels are the genes residing in the DA; if a DAC entails more than one gene, then the genes are separated by a comma. The point size represents the length of DACs with a padj < 0.1, and PSI-BLAST found the gene's symbol against *D. melanogaster*.

Figure 3: Genomic landscape of differentiation in *D. albomicans* and *D. nasuta* populations.

A Pairwise genetic differentiation (FST) in 50-kb non-overlapping windows across the genome between *D. albomicans* population pair (CHN vs TAI) in top panel, between *D. nasuta* population pair (IND vs KEN) in second panel and between *D. albomicans* (IND) and *D. nasuta* (IND) population pair in third panel. Colours represent respective Muller elements; horizontal dotted red line demarcates FST estimate of 0.9. Chromosome 4 in third panel is highlighted in grey to show high differentiation region in the later half of chromosome. **B** Zoomed in view of the chromosome 4 for the same population pair of *D. albomicans* and *D. nasuta* to show their levels of genetic diversity depicted here as Watterson's theta (θw) with blue and black lines respectively. The red dots denotes significant differential accessibility regions, whereas size of the dots is proportional to the length of the region. Most of the differential accessible regions overlap with high differentiation and low genomic diversity regions in later half of the chromosome. **C** Zoomed in view of the chromosome 2R for the same population pair of *D. albomicans* and *D. nasuta* to show their levels of genetic diversity depicted here as Watterson's theta (θw) with blue and black lines respectively. The red and blue dots denotes significant differential accessibility regions, whereas size of the dots is proportional to the length of the region. Unlike chromosome 4, there is a drastic difference in genetic diversity levels between two species.

Figure 4: Transcription factor (TF) motifs across two species

A Heatmap of TF activity score across two hybrids, the scores were transformed into Z-score, where each row represents a TF in a species. Blue indicates low activity, and red indicates the higher TFs **B** TF motif activity score in testes across two species, by TOBIAS. Each dot in the volcano plot represents the activity scores of the TF motifs. The points were colored and labeled when the score meets the -log10(p-value) above 95% quantile and binding scores smaller/larger than the 5% and 95% quantiles **C** and **D** TF-TF network Trl in hybrids identifying with *D. albomicans* and *D. nasuta,* respectively. The color of node/TFs represents the binding scores of individual TFs, and the gradient of blue and red represents the higher binding score in hybrids identifying with *D. albomicans* and *D. nasuta,* respectively. The border colors around TFs show that they are unique to the individual species, and the edges/connection width represents the weights calculated by TOBIAS.