Combining multiple genetic estimates of N_e

Robin S. Waples

School of Aquatic and Fishery Sciences, University of Washington, Seattle

Running head: Combining N_e estimates

Corresponding author:
Robin Waples (robinw3@uw.edu (206) 789-4648)

September 2025

Abstract
Researchers often use multiple genetic methods to estimate contemporary effective
population size (N_e), but few formally combine estimates despite potential benefits for increasing
precision. Maximizing benefits requires an optimal, inverse-variance weighting scheme.
Methods should be estimating the same parameter, which can be appropriate either for estimates
using the same method applied to different time periods, or estimates using different methods
applied to the same time period. Previous approaches focused on $\mathrm{var}(\widehat{N}_e)$ for weighting, but that
is problematical because $\widehat{N}_e$ is highly skewed and can be infinitely large. A new approach is
described using weights inversely proportional to $\mathrm{var}(1/\widehat{N}_e)$, which is the drift signal that N_e-
estimation methods respond to. The distribution of $1/\widehat{N}_e$ is close to normal even when $\widehat{N}_e$
assumes extreme values. Benefits are maximized under three general conditions: estimators
have approximately equal variances; they are uncorrelated or have weak positive correlations;
individual estimates have low precision (i.e., if data are limited and/or true N_e is large).
Analytical and numerical results demonstrate that: (1) existing theory allows robust estimates of
$\mathrm{var}(1/\widehat{N}_e)$ for the temporal and LD methods, which provide independent information about N_e –
both of which facilitate optimally combining those methods; (2) estimates for the LD and sibship
methods are essentially uncorrelated when data are limited but can be strongly positively
correlated in genomics-scale datasets. General theory predicting $\mathrm{var}(1/\widehat{N}_e)$ for the sibship
method is lacking, but values for specific scenarios have been published. New software
(*COMBONE*) is introduced to calculate combined estimates.
21
22

1 | INTRODUCTION

Effective population size (N_e) is one of the most important parameters in evolutionary biology, and interest in estimating N_e has grown rapidly in recent years (Palstra and Fraser 2012; Wang et al. 2016; Clarke et al. 2024; Waples 2025). This upward trend is likely to continue into the foreseeable future, given that (1) parties to the Convention on Biological Diversity recently committed to tracking genetic diversity for all species, and (2) effective size was identified as a key metric for monitoring within-population genetic diversity (CBD 2022; Mastretta-Yanes et al. 2024). A wide range of genetic methods for estimating N_e have been proposed, and collectively the time frames considered by these methods extend from the present to thousands of generations in the past (reviewed by Nadachowska-Brzyska et al. 2022). For applied conservation and management, the most relevant estimates are those for contemporary N_e, which roughly corresponds to the time frame encompassed by the samples. The three most widely-used methods to estimate contemporary N_e are the temporal method (which measures changes in allele frequency between two or more samples spaced in time) and two methods that require only a single sample: the sibship method of Wang (2009), and a bias-corrected version of the method based on linkage disequilibrium (LD; Waples and Do 2008). Two other single-sample methods—the heterozygote-excess method of Pudovkin et al. (1996) and the molecular coancestry method of Nomura (2008)—have not performed well in performance evaluations (Do et al. 2014; Wang 2016) and are not widely used.

Precision is often a limiting factor for genetic methods to estimate contemporary N_e, especially when true N_e is large. Furthermore, many authors report N_e estimates from more than one method. This suggests that formally combining information from multiple methods could lead to more robust overall estimates. To illustrate potential benefits, consider two fundamental properties of a variance. First, the variance of the product of a random variable (X) and a constant (a) is

$$var(aX) = a^2 var(X). \tag{1}$$

Second, the variance of the sum of two random variables is

$$var(X + Y) = var(X) + var(Y) + 2cov(X,Y), \tag{2}$$

where $cov(X,Y)$ is the covariance of X and Y. Combining Equations 1 and 2, we have

$$var(aX + bY) = a^2 var(X) + b^2 var(Y) + 2ab * cov(X,Y). \tag{3}$$

Now assume that both X and Y are unbiased estimators of the same parameter θ, and let a and b be standardized such that $a+b=1$. Then $Z = aX + bY$ is a weighted average of X and Y, with a and b being relative weights. Consider a simple example where $var(X) = var(Y)$, so X and Y are weighted equally ($a=b=0.5$). Substituting into Equation 3:

$$var(Z) = 0.5^2 var(X) + 0.5^2 var(Y) + 2(0.5)(0.5)cov(X,Y)$$

$$= 0.25var(X) + 0.25var(X) + (0.5)cov(X,Y)$$

$$= 0.5var(X) + (0.5)cov(X,Y).$$

69 If X and Y are independent (hence $cov(X,Y) = 0$), then $var(Z) = 0.5var(X)$; that is, the
70 variance of the weighted mean would be half that of either individual estimator. More generally,
71 Cochrane (1937; see also Shahar 2017) showed that if X and Y are independent and normally
72 distributed, and if the weights are chosen to be inversely proportional to variances, such that
73
74 $$a = W_1 = \frac{1/var(X)}{1/var(X) + 1/var(Y)} \text{ and } b = W_2 = \frac{1/var(Y)}{1/var(X) + 1/var(Y)},$$

75
then $Z = W_1X + W_2Y$ is an unbiased estimator of θ and $var(Z) = var(X)var(Y)/[var(X) +$
$var(Y)]$, which is less than or equal to the smaller of $var(X)$ and $var(Y)$.
Waples and Do (2010) showed how to combine estimates from the LD and temporal
methods, but combined estimates are rarely presented, even by authors who report results from
more than one method. One difficulty is that Waples and Do (2010) calculated weights based on
published formulas for var($\hat{N}_e$) for the temporal method (by Pollak 1983) and for the LD method
(by Hill 1981). However, the distribution of $\hat{N}_e$ can be highly skewed and the estimates can take
infinitely large or even negative values, which makes calculating var($\hat{N}_e$) problematical except in
some scenarios where the distribution of $\hat{N}_e$ is quasi-normal.
The goal here is to develop a more robust way for combining multiple estimates of
effective size, with the weights being inversely proportional to var($1/\hat{N}_e$) rather than var($\hat{N}_e$).
This simple change has several advantages compared to the approach outlined by Waples and Do
(2010). First, $1/N_e$ is the drift signal that all genetic methods to estimate contemporary effective
size are sensitive to (Wang 2001, 2009). Second, $1/\hat{N}_e$ is approximately normally distributed
even when $\hat{N}_e$ decidedly is not (Figure 1), which facilitates setting confidence intervals (CIs).
Third, var($1/\hat{N}_e$) has a robust analytical solution for both the temporal and LD methods, as a
function of parameters that are known (sample sizes of loci and individuals; elapsed time
between samples) or can be estimated (true N_e). Here I consider how to combine multiple
estimates of effective size, both within and between methods.
95
96
97 **2 | METHODS**
98
2.1 | Assumptions
Methods to estimate contemporary N_e generally make the following assumptions, which
also are adopted here: generations are discrete; mutation is unimportant over the time frames
considered; samples are drawn from a single, isolated population closed to immigration and
emigration; and the genetic markers assayed are selectively neutral, such that observed variation
reflects the action of random genetic drift. We revisit these assumptions in Discussion.
105
2.2 | Expected Values and Variances for Effective Size Estimators
Consider first the standard temporal method, which compares allele frequencies within a
single population at two times separated by t generations. A widely-used index of change in
allele frequency (P) is temporal F (hereafter just F). For a given genetic marker, parametric F is
the squared difference in population allele frequency between times 0 and t, standardized to
minimize effects of initial allele frequency:
112

$$F = \frac{(P_0 - P_t)^2}{\bar{P}(1-\bar{P})},$$ (4)

where $\bar{P} = (P_0 + P_t)/2$ is the mean frequency over time. In general, P_0 and P_t are unknown and
must be estimated, from two samples of S_0 and S_t individuals. Several authors (Nei and Tajima
1981; Pollak 1983; Jorde and Ryman 2007) have proposed estimators of F; differences among
these estimators are relatively small and they are collectively referred to here as $\hat{F}$. The core
premise of the temporal method is that the expected value of $\hat{F}$ ($E(\hat{F})$) is a simple function of N_e
and t. Two sampling plans must be considered because they affect the magnitude of allele-
frequency change between the samples (Nei and Tajima 1989; Waples 1989):

Plan I: $E(\hat{F}) \approx \frac{t}{2N_e} + \frac{1}{S} - \frac{1}{N}$; (5a)

Plan II: $E(\hat{F}) \approx \frac{t}{2N_e} + \frac{1}{S}$, (5b)

where N is census size. The contribution to $E(\hat{F})$ from sampling error is approximately $1/(2S_0)$
at time 0 and $1/(2S_t)$ at time t. For simplicity, we assume that $S_0=S_t=S$, so the total magnitude of
sampling error is $1/S$. If $S_0 \neq S_t$, the harmonic mean of the two sample sizes should be used in
place of S.
In Plan I, individuals for genetic analysis are sampled after reproduction or non-lethally
before reproduction; in Plan II, individuals are sampled before reproduction and not allowed to
reproduce. The key difference is that under Plan I, some individuals in the initial sample can
contribute genes to future generations, which creates positive correlations that have to be
accounted for by the $1/N$ term in Equation 5a. Here, N is the total number of individuals subject
to (presumably random) sampling at time 0; N at the time of the second sample is not relevant.
Temporal F is closely related to Wright's (1965) F_{ST} and related measures of genetic
differentiation among populations. Considerable interest has focused on variability of F_{ST} among
loci, as markers with unusually high F_{ST} might reflect effects of natural selection. Lewontin and
Krakauer (1973) proposed that the random drift variance of F_{ST} among loci is

$$var(F_{ST}) = 2[E(F_{ST})]^2/n,$$ (6)

where n is the number of independent alleles used to compute each F_{ST}, and that a significantly
higher variance than given by Equation 6 could be taken as evidence for the action of selection.
Other authors (Robertson 1975; Ewens and Feldman 1976) soon pointed out that various factors
besides natural selection can inflate $var(F_{ST})$, and accounting for these factors greatly
complicates identification of true outlier loci (Beaumont and Nichols 1996; Lotterhos and
Whitlock 2014). The situation is different with temporal F, which measures allele-frequency
differences within rather than among populations and is not affected by these complicating
factors (Gaines and Whittam 1980).
The analogue to Equation 6 that is applicable to the temporal method is

$$var(\hat{F}) = 2[E(\hat{F})]^2/n_F,$$ (7)

where n_F is the number of independent alleles used to compute $\hat{F}$. Most genetic datasets now use
diallelic SNPs, which have one independent allele per locus, so for our purposes $n_F = L$ = number
of loci used to compute $\hat{F}$. Loci on the same chromosome do not assort independently, and as
the number of loci per chromosome becomes large, it becomes more and more likely (and
eventually inevitable) that any new locus will be close to, and provide largely redundant
information for, one or more existing loci. This lack of independence, or pseudoreplication,
reduces the total information content of large datasets. This lack of independence can be
quantified by n'_F, which is the effective degrees of freedom associated with $\hat{F}$. The method
described by Waples et al. (2022) was used to estimate n'_F as a function of L, S, N_e, and genome
size.
In equations 5a and 5b, $E(\hat{F})$ has two components: a drift signal ($t/(2N_e)$ for Plan II and
$t/(2N_e) - 1/N$ for Plan I) and noise from sampling error ($1/S$ for both plans). The standard
temporal estimator of effective size is obtained by rearranging these equations as follows
(Waples 1989):

$\quad\quad$ Plan I: $\hat{N}_{eTemp} = \dfrac{t}{2(\hat{F}-\frac{1}{S}+\frac{1}{N})} = \dfrac{t}{2\hat{F}\prime}$, with $\hat{F}\prime = \hat{F} - \dfrac{1}{S} + \dfrac{1}{N}$; $\hfill$ (8a)

$\quad\quad$ Plan II: $\hat{N}_{eTemp} = \dfrac{t}{2(\hat{F}-\frac{1}{S})} = \dfrac{t}{2\hat{F}\prime}$, with $\hat{F}\prime = \hat{F} - \dfrac{1}{S}$, $\hfill$ (8b)

where $\hat{F}\prime$ is the raw $\hat{F}$ adjusted for sampling error and (for Plan I) the $1/N$ term that accounts for
temporal correlations.
Given the above, it is straightforward to derive the theoretical expectation for
$var(1/\hat{N}_{eTemp})$:

$\quad\quad$ $1/\hat{N}_{eTemp} = \dfrac{2\hat{F}\prime}{t}$; $\hfill$ (9)

$\quad\quad$ $var\left(\dfrac{1}{\hat{N}_{eTemp}}\right) = var\left(\dfrac{2}{t}\hat{F}\prime\right) = \left(\dfrac{2}{t}\right)^2 var(\hat{F}\prime) = \dfrac{4}{t^2} var(\hat{F}\prime)$. $\hfill$ (10)

Now adding or subtracting constants does not affect a variance, so $var(\hat{F}\prime) = var(\hat{F})$, and using
the result from Equation 7 for $var(\hat{F})$,

$\quad\quad$ $var\left(\dfrac{1}{\hat{N}_{eTemp}}\right) = \dfrac{4}{t^2} var(\hat{F}\prime) = \dfrac{4}{t^2} var(\hat{F}) = \dfrac{4}{t^2}\dfrac{2[E(\hat{F})]^2}{n'_F} = \dfrac{8[E(\hat{F})]^2}{t^2 n'_F}$. $\hfill$ (11)

Equation 11 applies to both sampling plans, with the understanding that $E(\hat{F})$ differs between
the plans as specified in Equations 5a,b. This means that, all else being equal, $var(1/\hat{N}_{eTemp})$
will be smaller for Plan I sampling because $E(\hat{F})$ is smaller.
The moment-based LD method, first described by Hill (1981) and reviewed by Waples
(2024), has many parallels to the temporal method. The drift signal is captured by r^2, which is
the squared correlation of alleles at different loci. If the recombination fraction (c) between a
pair of loci on the same chromosome is known, the LD method can provide insights into changes
of effective size in the past (Tenesa et al. 2007; Hollenbeck et al. 2016; Santiago et al. 2020).

For estimating contemporary N_e, focus is on pairs of loci on different chromosomes, which assort independently ($c=0.5$), in which case the estimate applies primarily to the parents of the individuals sampled (Waples 2005). Parametric LD in theory applies to $E(r^2)$ in an infinite number of progeny (Weir and Hill 1980). Finite sampling leads to the LD estimator $\hat{r}^2$, and the Burrows method (Weir 1979; Weir and Hill 1980) is the most widely-used approach for calculating $\hat{r}^2$. The simplest way to calculate the Burrows coefficient is to use the Pearson product–moment correlation of the genotypic vectors for two loci (Gao et al., 2008).

For unlinked loci, the expectation of $\hat{r}^2$ is approximately

$$E(\hat{r}^2) \approx \frac{1}{3N_e} + \frac{1}{S-1}, \tag{12}$$

with the first term accounting for genetic drift and the second term for sampling error. Equation 12 is applicable to unphased genotypic data, which is the most common type of data generated by NGS methods for non-model species. Hill (1981) and Weir and Hill (1980) indicated that the contribution to $\hat{r}^2$ from sampling a finite number of individuals was approximately $1/S$, but this underestimates effects of sampling error and leads to downward bias in $\widehat{N}_{eLD}$ (England et al. 2006; Waples 2006). The term $1/(S-1)$ closely approximates the bias adjustment used in LDNe and implemented in NeEstimator (Do et al. 2014) (unpublished data). Rearranging Equation 12 leads to the LD estimator of contemporary N_e

$$\widehat{N}_{eLD} = \frac{1}{3\left(\hat{r}^2 - \frac{1}{S-1}\right)} = \frac{1}{3\hat{r}^{2\prime}}, \text{ with } \hat{r}^{2\prime} = \hat{r}^2 - \frac{1}{S-1}. \tag{13}$$

Inverting Equation 13 leads to

$$1/\widehat{N}_{eLD} = 3\hat{r}^{2\prime}. \tag{14}$$

The analogue to Equation 7 that is applicable to the contemporary LD method is

$$var(\hat{r}^{2\prime}) = var(\hat{r}^2) \approx 2[E(\hat{r}^2)]^2/n_{LD}, \tag{15}$$

where n_{LD} is the number of pairs of loci used to compute $\hat{r}^2$. As with the temporal method, physical linkage creates lack of independence for the LD method, but a larger issue is lack of independence caused by overlapping pairs of the same loci. Results from Waples et al. (2022) were used to estimate n'_{LD}, which is the effective number of pairs of loci. Making this substitution, and taking the variance of both sides of Equation 14,

$$var\left(\frac{1}{\widehat{N}_{eLD}}\right) = var(3\hat{r}^2) = 9var(\hat{r}^2) = 18[E(\hat{r}^2)]^2/n'_{LD}. \tag{16}$$

2.3 | Combining Estimates

Combining two or more estimates can be appropriate when they are estimating the same parameter. Considerations are different for combining estimates within and between methods. In either case, an optimal weighting scheme (in terms of minimizing variance of the combined estimate) accounts for the covariance structure of the estimators (Keller and Olkin 2004). In practice, the covariances will rarely be known. Fortunately, in some important scenarios the

241 estimators can be treated as independent, in which case optimal weights are proportional to
242 reciprocals of variances. In the simulated data, performance of the inverse-variance weighting
243 approach was evaluated under scenarios where the estimators were and were not independent.
244
2.3.1 | Combining estimates within a single method
If multiple samples of individuals provide information about N_e in a single generation for
the same population, the simplest approach is to combine the individuals into a single sample.
This will reduce the variance of $1/\widehat{N}_e$ by reducing $E(\widehat{F})$ or $E(\widehat{r}^2)$ in Equations 5a,b or 12,
respectively.
If multiple samples of individuals provide information about N_e in different generations
for the same population, using perhaps different sample sizes of individuals and loci, then
combining the estimates to increase precision can be appropriate under two scenarios:
(1) N_e can be considered to be constant over time;
(2) N_e might vary, but it is of interest to compute a central tendency for N_e over the period
encompassed by the samples.
The general approach is the same for these two scenarios (and for both the temporal and LD
methods) and is described below.
Assume one has $j>1$ separate estimates of $1/\widehat{N}_e$ ($1/\widehat{N}_{e(i)}$, $i=1,j$), derived from Equations
like 9 or 14. Associated with each estimate is a theoretical variance, given by Equations 11 or
16. Denote these variances as V_1, V_2 … V_j. The combined estimate ($1/\widehat{N}_{eCombo}$) is a weighted
mean of the individual estimates, with weights being proportional to reciprocals of variances:
262

263
$$\frac{1}{\widehat{N}_{eCombo}} = \frac{W_1}{\widehat{N}_{e(1)}} + \frac{W_2}{\widehat{N}_{e(2)}} + \cdots \frac{W_j}{\widehat{N}_{e(j)}} = \sum_{i=1}^{j} \frac{W_i}{\widehat{N}_{e(i)}}. \tag{17}$$

264
265 In Equation 17, the weights are standardized such that $W_i = \frac{1/V_i}{\sum(\frac{1}{V_i})}$ and $\sum(W_i) = 1$.

266   By extension of Equation (2), the variance of the combined estimate is the sum of the
267 variances plus 2 times all the pairwise covariances:
268

269
$$var(\frac{1}{\widehat{N}_{eCombo}}) = \sum_{i=1}^{j} var(\frac{W_i}{\widehat{N}_{e(i)}}) + 2\sum_{i\neq j} W_i W_j cov(\frac{1}{\widehat{N}_{e(i)}}, \frac{1}{\widehat{N}_{e(j)}})$$

270
271
$$= \sum_{i=1}^{j} (W_i)^2 var(\frac{1}{\widehat{N}_{e(i)}}) + 2\sum_{i\neq j} W_i W_j cov(\frac{1}{\widehat{N}_{e(i)}}, \frac{1}{\widehat{N}_{e(j)}}). \tag{18}$$

272
273 If the different estimates are independent, then the covariance terms drop out and the variance of
274 the combined estimate simplifies to
275

276
$$var(\frac{1}{\widehat{N}_{eCombo}}) = \sum_{i=1}^{j} (W_i)^2 var(\frac{1}{\widehat{N}_{e(i)}}). \tag{19}$$

277
278 The term $\Sigma(W_i)^2$ is minimized when all weights are equal.
279   For the temporal method the raw weights are of the form
280

$$W_{Temp} \propto \frac{1}{var\left(\frac{1}{\widehat{N}_{eTemp}}\right)} = \frac{1}{\frac{8[E(\hat{F})]^2}{t^2 n\prime_F}} = \frac{t^2 n_F}{8[E(\hat{F})]^2}$$

$$= \frac{t^2 n\prime_F}{8\left[\frac{t}{2N_e}+\frac{1}{S}-\frac{1}{N}\right]^2} (Plan\ I) \qquad\qquad (20a)$$

$$or\ \frac{t^2 n\prime_F}{8\left[\frac{t}{2N_e}+\frac{1}{S}\right]^2} (Plan\ II)\ . \qquad\qquad (20b)$$

After dividing by the sum of all the raw weights, the standardized weights sum to 1 and can be
used in Equation 19. For the LD method, the raw weights are of the form

$$W_{LD} \propto \frac{1}{var\left(\frac{1}{\widehat{N}_{eLD}}\right)} = \frac{1}{\frac{18[E(\hat{r}^2)]^2}{n\prime_{LD}}} = \frac{n\prime_{LD}}{18[E(\hat{r}^2)]^2} = \frac{n\prime_{LD}}{18\left[\frac{1}{3N_e}+\frac{1}{S-1}\right]^2}. \qquad (21)$$

Inspection of Equations 20a,b and 21 indicates that optimal weights for different estimates are
functions of true N_e, which is unknown. This suggests an iterative process to calculate the
weights, as described below.

It is of interest to ask how $var\left(\frac{1}{\widehat{N}_e}\right)$ for a combo estimate compares to the variance for a
single estimate. For the LD method, if there are j estimates that all apply to the same constant or
harmonic mean N_e, then (assuming constant S and $n\prime_{LD}$) all estimates have the same $E(\hat{r}^2)$ and
the same theoretical variance and hence the same weight ($W = 1/j$). It follows that

$$\frac{1}{\widehat{N}_{eLDCombo}} = \frac{1}{j}\frac{1}{\widehat{N}_{eLD(1)}} + \frac{1}{j}\frac{1}{\widehat{N}_{eLD(2)}} + \cdots \frac{1}{j}\frac{1}{\widehat{N}_{eLD(j)}}$$

$$var\left(\frac{1}{\widehat{N}_{eLDCombo}}\right) = var\left(\frac{1}{j}\frac{1}{\widehat{N}_{eLD(1)}} + \frac{1}{j}\frac{1}{\widehat{N}_{eLD(2)}} + \cdots \frac{1}{j}\frac{1}{\widehat{N}_{eLD(j)}}\right)$$

$$= \frac{1}{j^2}\sum var\left(\frac{1}{\widehat{N}_{eLD(i)}}\right) + 2\sum_{i\neq j} cov\left(\frac{1}{\widehat{N}_{eLD(i)}},\frac{1}{\widehat{N}_{eLD(j)}}\right)\ . \qquad (22)$$

Genetic drift is a Markov process, so it is reasonable to assume independence of LD estimates
across time. Ignoring the covariance terms, and substituting for $var\left(\frac{1}{\widehat{N}_{eLD(i)}}\right)$ from Equation 16,

$$var\left(\frac{1}{\widehat{N}_{eLDCombo}}\right) = \frac{1}{j^2}\sum \frac{18[E(\hat{r}^2)]^2}{n\prime_{LD}} = j\frac{1}{j^2}\frac{18[E(\hat{r}^2)]^2}{n\prime_{LD}} = \frac{1}{j}\frac{18[E(\hat{r}^2)]^2}{n\prime_{LD}}\ . \qquad (23)$$

Thus, assuming independence, combining information from j equally-weighted LD estimates
reduces the overall variance of $\frac{1}{\widehat{N}_{eLD(i)}}$ by the factor $1/j$.

For the temporal method, consider the general case where samples are taken not only in
generations 0 and t, but also in intermediate generation $t/2$. From Equation 11, the variance of
the estimate using the two samples maximally-separated in time is

317
$$var\left(\frac{1}{\widehat{N}_{eTemp(0,t)}}\right) = \frac{8[E(\hat{F}_{0,t})]^2}{t^2 n'_F} = \frac{8\left[\frac{t}{2N_e}+\frac{1}{S}\right]^2}{t^2 n'_F},$$

319 assuming Plan II sampling. The intermediate sample allows one to consider another estimate
320 that combines information from the two shorter estimates:

322
$$\frac{1}{\widehat{N}_{eTempNested}} = \frac{W_1}{\widehat{N}_{eTemp(0,\frac{t}{2})}} + \frac{W_2}{\widehat{N}_{eTemp(\frac{t}{2},t)}};$$

324
$$var\left(\frac{1}{\widehat{N}_{eTempNested}}\right) = var\left(\frac{W_1}{\widehat{N}_{eTemp(0,\frac{t}{2})}} + \frac{W_2}{\widehat{N}_{eTemp(\frac{t}{2},t)}}\right) + 2W_1 W_2 cov\left(\frac{1}{\widehat{N}_{eTemp(0,\frac{t}{2})}}, \frac{1}{\widehat{N}_{eTemp(\frac{t}{2},t)}}\right).$$

326 Assuming fixed sample sizes of individuals and loci, $W_1 = W_2 = 0.5$, and $E\left(\hat{F}_{0,t/2}\right) =$
327 $E\left(\hat{F}_{t/2,t}\right) = \frac{t/2}{2N_e} + \frac{1}{S} = \frac{t}{4N_e} + \frac{1}{S}$. Therefore, letting C stand for the covariance term,

329
$$var\left(\frac{1}{\widehat{N}_{eTempNested}}\right) = 0.5^2 \frac{8[E(\hat{F}_{0,t/2})]^2}{(t/2)^2 n'_F} + 0.5^2 \frac{8[E(\hat{F}_{t/2,t})]^2}{(t/2)^2 n'_F} + C$$

331
$$= \frac{2[\frac{t}{4N_e}+\frac{1}{S}]^2}{(t/2)^2 n'_F} + \frac{2[\frac{t}{4N_e}+\frac{1}{S}]^2}{(t/2)^2 n'_F} + C = \frac{4[\frac{t}{4N_e}+\frac{1}{S}]^2}{t^2 n'_F/4} + C = \frac{16[\frac{t}{4N_e}+\frac{1}{S}]^2}{t^2 n'_F} + C, \tag{24}$$

which can be compared with the variance for an estimate based on 2 samples taken *t* generation
apart (Equation 11).

2.3.2 | Combining estimates across methods
Combining estimates from different methods might be appropriate under either of the
scenarios described above for combining estimates within methods. However, probably the
greatest benefits from using different methods arise when they estimate N_e in the same time
periods, in which case no assumptions are necessary regarding whether N_e varies over time and,
if so, to what degree. Care must be taken to ensure that the different methods are estimating N_e
in the same generation(s), following rules outlined in Waples (2005). A combined estimate using
information from both the temporal and LD methods is of the form

345
$$\frac{1}{\widehat{N}_{eCombo}} = \frac{W_{Temp}}{\widehat{N}_{eTemp}} + \frac{W_{LD}}{\widehat{N}_{eLD}}, \tag{25}$$

347 where $W_{Temp} = \dfrac{\frac{1}{var(1/\widehat{N}_{eTemp})}}{\frac{1}{var(1/\widehat{N}_{eTemp})}+\frac{1}{var(1/\widehat{N}_{eLD})}}$ and $W_{LD} = \dfrac{\frac{1}{var(1/\widehat{N}_{eLD})}}{\frac{1}{var(1/\widehat{N}_{eTemp})}+\frac{1}{var(1/\widehat{N}_{eLD})}}$.

349 Comparing the variances gives insight into the relative precision of the two methods:

351
$$R = \frac{var(1/\widehat{N}_{eTemp})}{var(1/\widehat{N}_{eLD})} = \frac{\frac{8[E(\hat{F})]^2}{t^2 n'_F}}{\frac{18[E(\hat{r}^2)]^2}{n'_{LD}}} = \frac{4}{9t^2} \frac{n'_{LD}}{n'_F} \frac{[E(\hat{F})]^2}{[E(\hat{r}^2)]^2} . \tag{26}$$

353 Assuming Plan II sampling for the temporal method, this becomes

$$R = \frac{var(1/\hat{N}_{eTemp})}{var(1/\hat{N}_{eLD})} = \frac{4}{9t^2} \frac{n\prime_{LD}}{n\prime_F} \frac{[\frac{t}{2N_e}+\frac{1}{S}]^2}{[\frac{1}{3N_e}+\frac{1}{S-1}]^2} = \frac{4}{9t^2} \frac{n\prime_{LD}}{n\prime_F} \frac{\frac{t^2}{4N_e^2}+\frac{1}{S^2}+\frac{t}{SN_e}}{\frac{1}{9N_e^2}+\frac{1}{(S-1)^2}+\frac{2}{3N_e(S-1)}}. \tag{27}$$

When $R>1$, $var(1/\hat{N}_e)$ is larger for the temporal method (so the LD method is more precise),
and the opposite is true for $R<1$.

More generally, assume that X and Y both are unbiased estimators of the same parameter
θ, and further assume that the variance of X (V_1) is equal to or less than the variance of Y. It
follows that $var(Y) = KV_1$, where $K \geq 1$. Then the standardized weight for X is $W_1 = (1/V_1)/(1/V_1$
$+ 1/(KV_1)) = (1/V_1)/((K+1)/KV_1) = K/(K+1)$ and $W_2 = 1 - W_1 = 1/(K+1)$. Then, if X and Y are
uncorrelated,

$$var(W_1 X + W_2 Y) = W_1^2 V_1 + (1 - W_1)^2 KV_1 = V_1(\frac{K}{K+1})^2 + KV_1(\frac{1}{K+1})^2$$

$$= \frac{K^2 V_1 + KV_1}{(K+1)^2} = \frac{KV_1(K+1)}{(K+1)^2} = V_1 \frac{K}{(K+1)}. \tag{28}$$

In the special case where the two variances are equal, $K=1$ and $K/(K+1) = 0.5$, so the variance of
the combined estimate is reduced by 50% compared to either individual variance, as in the
example in the Introduction. If $V_1 < V_2$, the reduction is <50% compared to the smaller variance
(Figure S1), and the variance reduction is further restricted if the different estimators are
positively correlated.

2.4 | Computer Simulations
Computer simulations were used to empirically evaluate theoretical results. R code to
conduct the simulations will be posted on Zenodo on acceptance. In all simulations, Wright-
Fisher populations with constant $N=N_e$ were modeled. Populations were initiated with a flat
distribution of allele frequencies in the range 0.1-0.5 at a variable number of diallelic (SNP) loci.
Within each of 100-1000 replicates, within-method simulations for the temporal method modeled
10 generations of genetic drift. At each generation from $t=0$ to $t=10$, 10 replicate samples of S
individuals were taken for genetic analysis, according to either Plan I or Plan II protocols. All of
these replicate samples were based on the same population pedigree, so they avoided adding
spurious demographic variance as described by Waples and Faulkner (2009). Within each
replicate, the full matrix of pairwise comparisons of samples was used to generate the full matrix
of $\hat{F}$ values (for comparisons of generations 0 and 1, 0 and 2, 1 and 2 … 0 and 10). For each
type of comparison, means and variances of $\hat{F}$ (and the resulting $1/\hat{N}_e$ values) were computed
across the 10 replicate samples, and these means and variances were then averaged across
replicates. Various types of within-method combined estimates were computed as described in
the text.
The procedure for within-method LD simulations was similar, except that a 7-generation
burnin period was added at the start of each replicate to allow an approximately stable amount of
LD to accumulate. The 10 generations of samples were not compared with each other, as in the
temporal method; instead, data for each generation were analyzed separately. For each

generation, means and variances were computed across the 10 replicate samples, as described
above.
Combined estimates across methods are only reasonable when the two methods are
estimating the same parameter. Accordingly, these simulations focused on a scenario where two
temporal samples of progeny from generations t and t +1, and a single LD sample of progeny
from generation t +1, are all used to estimate N_e in generation t +1. After initialization and a 7-
generation burnin, an initial temporal sample was taken, followed by a single generation of
genetic drift and a second temporal sample, which was also used for the LD estimate. As
described above, the samples were replicated and means and variances computed across sample
replicates and then averaged across replicate simulations.
Correlations between estimates were evaluated three ways. For the temporal and LD
methods, correlations were calculated between simulated data series for both within-method and
between-method evaluations. Second, simulation code made available in Supporting
Information for Waples (2021) was used to quantify correlations between LD and sibship
estimates. These simulations assumed that all sibship reconstructions were done without error,
so results for the sibship method are optimistic except perhaps for large datasets using ≥1000
SNPs. Finally, the R function MASS::mvrnorm was used to simulate random normal data with
specified means, variances, and correlations.
In addition to empirical variances, which measure precision, two more performance
metrics were used with the simulated data. The root-mean-squared error (RMSE; square root of
the mean squared deviation from the true value) combines information about both precision and
bias. Mathematically, RMSE can be calculated as $\mathrm{var}(1/\widehat{N}_e) + [\mathrm{bias}(1/\widehat{N}_e)]^2$. Confidence interval
coverage records the percentage of replicates for which the 95% confidence interval around
$\widehat{N}_{eCombo}$ contained the true value of N_e. Confidence intervals assumed that $1/\widehat{N}_e$ was normally
distributed, so the 95% CIs were $1/\widehat{N}_e$ +/- 1.96*sd($1/\widehat{N}_e$).

3 | RESULTS

422   I begin by summarizing some of the main analytical results from Methods, followed by
423 presentation of simulation results.

3.1 | Var($1/\widehat{N}_e$)

For the temporal method, $\mathrm{var}(1/\widehat{N}_{eTemp})$ is proportional to squared $E(\widehat{F})$ (Equation 11),
which varies according to the sampling plan as shown in Equations 5a,b. All else being equal,
$E(\widehat{F})$ is smaller for Plan I sampling by the magnitude $1/N$, so Plan I sampling leads to higher
precision. $\mathrm{var}(1/\widehat{N}_{eTemp})$ is reduced by the number of generations between samples and by the
effective number of alleles used in the calculation of $\widehat{F}$ (Equation 11). In simulations, Equation
11 accurately predicted $\mathrm{var}(1/\widehat{N}_{eTemp})$ for samples spaced 1-10 generations apart (Figure S2, top
panel).
For the LD method, $\mathrm{var}(1/\widehat{N}_{eTemp})$ is proportional to squared $E(\hat{r}^2)$ and it is inversely
proportional to the number of effectively independent pairs of loci used in the calculation of $\hat{r}^2$
(Equation 16). In the simulations, Equation 16 tended to slightly underestimate (by a few
percent) $\mathrm{var}(1/\widehat{N}_{eLD})$ for single samples of progeny (Figure S2, bottom panel).

3.2 | Combining Estimates Within Methods

439   Before combining multiple estimates of $1/\hat{N}_e$, it is necessary to calculate appropriate
440   weights, which should be standardized so they sum to 1. We use weights that are inversely
441   proportional to var($1/\hat{N}_e$), so temporal weights (Equations 20a,b) are inversely proportional to
442   Equation 11 and LD weights (Equation 21) are inversely proportional to Equation 16.
443   Standardized weights are of the form
444

$W_i^* = \dfrac{W_i}{\Sigma W_i}$.

Per Equation 2, variances of combined estimates are the sum of the individual weighted
variances, plus covariance terms for each pair of estimates being averaged. Single-sample
estimates are simpler to analyze in this regard because they are arguably independent across
generations, in which case the covariance terms vary randomly around 0 and can be ignored. If
so, the combined LD estimate that leverages information from separate LD estimates in j
generations has expected variance that is $(1/j)$ times the variance of an estimates based on a
single-generation of data (Equation 23). Simulation results (Figure S3, top panel) were in
excellent agreement with this prediction, indicating that it is reasonable to ignore the covariance
terms in var($1/\hat{N}_{eLDCombo}$). RMSE of the combined LD estimates also declined systematically as
information from more samples was utilized (Figure S3, bottom panel).
Combining multiple temporal estimates is trickier because the estimates typically are
positively correlated. Two general scenarios can be considered. In the first (nested) scenario, a
single pair of samples from generations 0 and t is supplemented by an additional sample at
generation $t/2$. var($1/\hat{N}_{eTemp}$) for the two extreme samples is given by Equation 11. The other
two pairs of samples each span half the total time ($t/2$ generations) and have the same weights
and the same $E(\hat{F})$, and the combined variance is given by Equation 24. Even ignoring the
covariance term, var($1/\hat{N}_{eTempNested}$) is generally larger than the variance from the two samples
spanning the full t generations (Equation 11), unless S is relatively large compared to N_e and/or t
is very large (Table S1). In this nested scenario, however, the covariance term will be positive
because the two estimates based on the shorter time frame share the same sample at generation
$t/2$. This covariance will become relatively less important as t increases. In the simulations,
var($1/\hat{N}_{eTempNested}$) was markedly higher than the single estimate for $t=2$ and only modestly
higher for $t=10$ (Figure S4, top).   Thus, splitting a long temporal comparison into two shorter
ones does not improve precision.
In the second scenario, all 3 estimates (two spanning $t/2$ generations and one spanning all
t generations) are combined into a single, comprehensive estimate, using weights calculated as in
Equations 20a,b. The estimates spanning short and long time periods are highly positively
correlated, as they not only share samples but also duplicate coverage of overlapping generations
of genetic drift. These positive correlations increase the variance of the comprehensive estimates
and limit the benefits of adopting this scenario. Nevertheless, simulations showed that precision
did increase somewhat by using all of the data (compared to the single estimate,
var($1/\hat{N}_{eTempComprehensive}$) was reduced by 13% for $t=2$, 23% for $t=5$, and 24% for $t=10$; Figure
S4, bottom).

3.3 | Combining Estimates Across Methods
3.3.1 | Temporal and LD methods
483

484   The general formula for combining estimates across temporal and LD methods is
485 (Equation 22)
486

487    $\frac{1}{\widehat{N}_{eCombo}} = \frac{W_{Temp}}{\widehat{N}_{eTemp}} + \frac{W_{LD}}{\widehat{N}_{eLD}}$,

488
489 where the raw weights are calculated as in Equations 20 and 21 and standardized weights are of
490 the form
491

492    $W_i^* = \frac{W_i}{W_{Temp}+W_{LD}}$.

493
494 It is useful to express the theoretical variances of the two methods as a ratio (R), which provides
495 insights into relative precision (Equation 26):
496

497    $R = \frac{var(1/\widehat{N}_{eTemp})}{var(1/\widehat{N}_{eLD})}$    .

498
499 $R>1$ indicates higher variance (less precision) for the temporal method, and $R<1$ indicates the
500 reverse. If temporal estimates span a single generation ($t=1$), the LD method has greater
501 precision unless at least 10^4 SNPs are used (Figure 2, top). Datasets in the range of 100-1000
502 SNPs are in the "sweet spot" for the LD method, where the benefits of being able to use
503 information from many pairwise combinations of loci are maximized. As more and more SNPs
504 are used, lack of independence limits further increases in precision more for the LD method than
505 it does for the temporal method. All else being equal, relative precision of the LD method is
506 higher when true N_e is larger. When temporal estimates span multiple generations ($t>1$), relative
507 precision of the temporal method increases proportionally, resulting more frequently in $R<1$
508 (Figure 2, bottom). In this scenario, however, a single LD estimate applies to only one of the
509 (potentially many) generations over which N_e is estimated by the temporal method, so that must
510 be kept in mind in interpreting what $\widehat{N}_{eCombo}$ represents.
511   The scenario with $t=1$ is of particular interest because in that case both the temporal and
512 LD methods estimate effective size in a single generation, which facilitates estimation of the key
513 ratio N_e/N. Fortunately, the two estimators are effectively uncorrelated, even for relatively large
514 values of L and S (Table 1). This means that the covariance term can be ignored in estimating
515 $var(1/\widehat{N}_{eCombo})$. For the scenarios considered in Table 1, the LD method was always more
516 precise (smaller $var(1/\widehat{N}_e)$) than the temporal method, but properly combining the two estimates
517 still provided benefits (reduced $var(1/\widehat{N}_e)$ and $RMSE(1/\widehat{N}_e)$ for $\widehat{N}_{eCombo}$; Tables 1C and 1D).
518   The benefit of combining estimates is maximized when they are uncorrelated and have
519 equal variances (Equation 28). This is illustrated in Figure 3, which shows how $var(1/\widehat{N}_{eCombo})$
520 varies in simulated data as a function of the weight assigned to the LD method (W_{LD}). In the top
521 panel (using 5000 SNPs), the optimal weights are $W_{LD} = 0.63$ and $W_{Temp} = 0.37$ and the optimal
522 combined estimate has a variance that is 37% lower than that of the LD estimates alone (and
523 63% lower than the variance of the temporal method alone). Conversely, if the variances are
524 very different, the benefits of combining estimates can be marginal at best (maximum
525 improvement <10% for the scenario in bottom panel of Fig 4, where 500 SNPs are used, which is
526 in the "sweet spot" for relative precision of the LD method).

Figure 3 also illustrates the consequences of improperly weighting the individual
estimates. In the top scenario, $\text{var}(1/\widehat{N}_{eCombo}) \leq \text{var}(1/\widehat{N}_{eLD})$ as long as W_{LD} is anything larger
than about 0.25.   But in the bottom scenario, with very skewed weights, $\text{var}(1/\widehat{N}_{eCombo})$ can be
greater than $\text{var}(1/\widehat{N}_{eLD})$ unless the weight assigned to the LD estimate is 0.8 or larger.
If information from two methods is being combined and both methods use information
from multiple samples, estimates within methods should be combined first, and an overall
estimate of $1/N_e$ can be computed as a weighted mean of the two combined within-method
estimates. Note that because of positive correlations among nested estimates, estimating the
variances associated with comprehensive temporal estimates can be challenging.
Effects of positive correlations are illustrated with simulated data in Figure 4. Using a
target N_e of 1000, two series of random normal variates were simulated, both having expected
value = $1/N_e$ = 1/1000. The variance of series B was twice that of series A, so A was weighted
more heavily in computing the combined estimate $1/\widehat{N}_{eCombo} = W_A*A + (1- W_A)*B$. When the
two series were simulated independently (white vertical bars in top panel in Figure 4), the
distribution of $\widehat{N}_{eCombo}$ was considerably tighter than for series A (black bars) or B (not shown),
with fewer estimates >2x the true value. But when the two series were simulated jointly with
expected positive correlation of 0.5 (gray bars), computing a combined estimate led to little or no
increase in precision compared to that provided by Series A alone.

3.3.2 | The sibship method
The other widely-used single-sample estimator is the sibship method developed by Wang
(2009). Analyses directly comparable to those described above for the temporal and LD methods
are difficult because no analytical formula for $var\left(\dfrac{1}{\widehat{N}_e}\right)$ analogous to Equations 11 or 16 exists
for the sibship method. Instead, uncertainty in N_e estimation has been handled in two ways. For
individual point estimates, confidence intervals can be generated as part of the maximum-
likelihood estimation process (Wang and Santure 2009). Second, for selected simulation
scenarios, Wang (2009, 2016) has published empirical variances of $\dfrac{1}{\widehat{N}_e}$ or RMSE values. Wang's
comparisons have shown that a) precision of the sibship and LD methods is much higher than for
either the coancestry or heterozygote excess methods, and b) precision generally was a bit
higher for the sibship than the LD method, but the difference narrowed as more loci were used.
Most of these comparisons involved relatively small numbers of loci (<100).
Unlike the temporal and LD methods, the sibship method has a hard upper limit to
performance, because once enough loci are used to correctly identify all siblings within a sample,
adding more loci cannot further increase accuracy or precision. Waples (2021) showed that,
because of this limitation, in genomics-scale datasets ($\geq$1000 SNPs) precision of the LD method
will generally exceed that of the sibship method, even if all siblings are correctly identified.
Waples (2021) also examined the correlations between $1/\widehat{N}_e$ for the sibship and LD methods and
found that correlation coefficients were generally low for small datasets but could be very high
for genomics-scale datasets. Typical results are shown in Table 2: correlations were generally
<0.1 for 100 or fewer loci but could be as high as 0.9 when more than a thousand SNPs were
used. For the same number of loci, correlations were generally lower for smaller sample sizes of
individuals. For example, with true N_e=200 and using at least 1000 SNPs, Pearson's r was 0.79
or higher for S=50 or 100 but $\leq$0.21 for S=25.

3.4 | Confidence Intervals

Two major factors can degrade performance of confidence intervals: 1) incorrectly
calculating the variance of the estimator; 2) using a biased estimator. Superior performance
occurs when all estimators are unbiased and variances are correctly estimated. Both of these
conditions were met for the combined estimates shown in Figure 4 when Series A and Series B
were simulated independently (white vertical bars). 95.0% of the nominally-95% CIs included
the true N_e, 2.6% of the CIs were too low, and 2.4% were too high (first row of data in Table S2).
Results were similar when Series A and Series B were simulated with expected correlation 0.5
(gray vertical bars) and the expected variance of the combined estimate was computed using
Equation 3 to account for the resulting covariance term (second row of data in Table S2). But
when correlated data were simulated but the resulting covariance was ignored in computing the
estimated variance, the CIs around $1/\widehat{N}_{eCombo}$ were too narrow and included the true value less
than 90% of the time (third row of data in Table S2).
The effect of bias on performance of confidence intervals can be seen for within-method
analyses using the LD method. For parameter values used in the simulations, $\widehat{N}_{eLD}$ has an
upward bias of about 5% (Figure S2), which has been documented before (Waples and Do 2010)
and is the result of a slight over-correction for downward bias in the original version (Hill 1981)
of the LD method. When only a single estimate is considered, based on 50 individuals scored for
1000 SNPs, this modest bias has little effect on CI performance (93.8% coverage; first line in
Table S3). But when multiple samples are used and the width of the CIs shrinks, coverage
declines. When information from 10 LD samples is combined, coverage has dropped to 80.7%,
and almost all CIs that don't include the true N_e are too high (last line of Table S3). This latter
result occurs because as the estimates become more precise and the width of CIs shrinks, they
converge on a value larger than the true N_e. An estimator with a downward bias would have the
opposite effect: CIs that did not contain the true N_e would tend to be too low.
Table S3 shows that in the presence of bias, CI performance degrades as more samples
are combined to increase precision. A similar phenomenon occurs when one increases the
number of loci. For the analyses reported in Table 1, an LD estimate based on a single sample
was combined with a temporal estimate based on two samples spanning a single generation.
With only 100 loci, coverage of the CIs for the combined estimates was actually >95% (Table
1D), despite the bias contributed by the LD method. But when 5000 loci were used, coverage
dropped below the expected 95%, and substantially so for the smaller sample sizes.
603
3.5 | Details Regarding Estimation
Combining multiple estimates to arrive at a single estimate of effective size involves the
following general steps:
(1) Generate 2 or more individual estimates of $1/N_e$;
(2) Estimate the variance associated with each estimate;
(3) Assign weights based on reciprocals of the variances from Step 2;
(4) Compute the weighted mean of the two estimators to obtain $1/\widehat{N}_{eCombo}$;
(5) Generate CIs for $1/\widehat{N}_{eCombo}$;
(6) Invert the values in Step 5 to generate the point estimate $\widehat{N}_{eCombo}$ and its CIs.
In Step 2, variances for the LD and temporal methods can be estimated using Equations 11 and
16, respectively. Alternatively, if a series of estimates is available for a single method, the
empirical variance of $1/\widehat{N}_e$ could be used. Step 3 uses Equations 20-21 to compute weights, and
Step 4 uses the generalized Equation 17 to calculate $\widehat{N}_{eCombo}$. Step 5 takes advantage of the fact
that $1/\widehat{N}_e$ will be approximately normally distributed as long as the number of datapoints used in

its calculation is at least 50. In contemporary datasets, 50 loci or independent alleles is generally
easy to achieve for the temporal method and even easier for the LD method, which uses pairs of
loci.

Optimal weights for $1/\hat{N}_e$ for the temporal and LD methods depend on the respective
variances, which are functions of true N_e (cf Equations 11 and 16). As true N_e is unknown, an
iterative process is used:

    1. Compute an initial $1/\hat{N}_{eCombo}$ using equal weights;
624
    2. Compute initial weights assuming true $N_e=\hat{N}_{eCombo}$;
625
    3. Repeat Steps 1 and 2 until convergence, using the updated weights in Step 1 and the
626
       updated $\hat{N}_{eCombo}$ in Step 2.
627
628 Evaluations indicate that 3 iterations is sufficient for convergence of most scenarios, but some
629 can require 5 or more.

3.5 | Software
631

632   COMBONE, coded in R, is introduced to carry out analyses described here. The R code,
633 together with a sample input file and a ReadMe text file, is available in Supporting Information.

3.6 | A Worked Example
635

A worked example uses the following input data for the temporal and LD methods (which
are the first row of data from the sample input file): $L = 9100$ SNPs; $S = 40$ individuals; $\hat{N}_{eTemp}$
$= 89$ and $\hat{N}_{eLD} = 114$. The initial equally-weighted estimate of N_e is $\hat{N}_{eCombo0}= 1/(0.5/89 +$
$0.5/114) = 99.96$. Properly weighting the two estimates requires knowing the effective numbers
of independent datapoints (effective degrees of freedom) associated with each. Embedded in
COMBONE is code from Waples et al. (2022) that estimates n'_{LD} and n'_F as a function of N_e, S,
genome size, and L. For illustration, we assume the focal species has 16 chromosomes, in which
case the initial effective degrees of freedom estimates are $n'_F = 3118$ for the temporal method
and $n'_{LD} = 7940$ for the LD method. Based on Equation 11, the initial estimate for
$\text{var}(1/\hat{N}_{eTemp})$ is $V1_{Temp} = (8*(1/(2*N_e) + 1/S))/n'_F$. Substituting the above values, $V1_{Temp} =$
$(8*(1/(2*99.96) + 1/40))/3118 = 7.698 \times 10^{-5}$ and inverting that gives raw $W1_{Temp} = 12991$.
Similarly, using Equation 16, the initial estimate for $\text{var}(1/\hat{N}_{eLD})$ is $V1_{LD} =(18*(1/(3*N_e) + 1/(S-$
$1)))/n'_{LD} = (18*(1/(3*99.96) + 1/(39)))/7940 = 6.569 \times 10^{-5}$, and inverting that gives the initial raw
weight of 15223. After standardization so that the weights sum to 1, the standardized weights are
$W1*_{Temp} = 0.460$ and $W1*_{LD} = 1-0.460 = 0.540$. With the first estimated weights, we update the
combined estimate of N_e as $\hat{N}_{eCombo1} = 1/(0.460/89 + 0.540/114) = 100.96$. In the second
iteration, the estimated variances and effective degrees of freedom are updated with the new
$\hat{N}_{eCombo1}$, leading to very slightly different weights ($W2*_{Temp} = 0.461$ and $W2*_{LD} = 0.539$.) The
new weights change only the third decimal place of $\hat{N}_{eCombo2}$, and the third iteration produces
little more in the way of change. So the best combined estimate is $\hat{N}_{eCombo} = 101$, with the two
methods providing roughly equal amounts of information.
The final step is to calculate CIs for the combined estimate. After the third iteration, the
final estimates of $\text{var}(1/\hat{N}_e)$ were 2.302×10^{-6} for the temporal method and 1.899×10^{-6} for the LD
method. From Equation 19, assuming independence (as is appropriate for temporal/LD
combinations), $\text{var}(1/\hat{N}_{eCombo}) = W_{Temp}^2 \text{var}\left(\frac{1}{\hat{N}_{eTemp}}\right) + W_{LD}^2 \text{var}\left(\frac{1}{\hat{N}_{eLD}}\right) = (0.461)^2 *$
$2.302 \times 10^{-6} + (0.539)^2 * 1.899 \times 10^{-6} = 1.041 \times 10^{-6}$, so the standard deviation of $1/\hat{N}_{eCombo}$

is $\sqrt{1.041 \times 10^{-6}} = 0.001$. 95% CIs for $1/\widehat{N}_{eCombo}$ are thus $1/101$ +/- 1.96*0.001 = 0.0099 +/-
0.00196 = [0.00794, 0.01186]. Inverting these gives the upper 95% CI for $\widehat{N}_{eCombo}$ as 126 and
the lower CI as 84. With this much data, and combining information from two independent
methods, it is possible to make fairly precise statements about effective population size.
666
4 | Discussion

Using genetic methods to estimate effective populations size has become something of a
cottage industry—and a fairly large one. It is relatively easy to use the same or overlapping
datasets to estimate effective size using more than one method, and many authors do so, but it
remains relatively rare to report combined estimates that leverage information from more than
one method. This is unfortunate, as combining estimates can substantially improve precision,
which is often the limiting factor in N_e estimation.
The approach outlined here improves on previous approaches by focusing on the variance
of $1/\widehat{N}_e$ rather than the variance of $\widehat{N}_e$, which is problematical because the distribution of $\widehat{N}_e$ is
highly non-normal. $\mathrm{Var}(1/\widehat{N}_e)$ is closely related to the variances of temporal $\widehat{F}$ and $\hat{r}^2$, and the
distributions of these metrics are well understood and effectively normal if the number of
independent datapoints used in their calculation is larger than about 50. As a consequence, it is
straightforward to calculate the variance of a combined estimate $1/\widehat{N}_{eCombo}$ and use that to set
realistic confidence intervals for $\widehat{N}_e$ using standard statistical theory. Benefits of doing this
include fewer very high or infinite estimates and tighter confidence intervals with higher lower
bounds, all of which improve the value of effective size estimates for applied conservation and
management.
A precondition for combining estimates is that the different methods must be estimating
the same parameter. This can potentially be appropriate either in combining estimates for
different time periods within methods (see Section 2.3.1 for details) or for combining estimates
for different methods for the same time period(s) (see Section 2.3.2 for details). The latter
scenario is particularly well-suited to increasing precision of the estimated N_e/N ratio.
Benefits of combining data from multiple estimators are maximized under three general
conditions:
1. The variances (and hence weights) of individual estimators are equal. In this scenario,
$\widehat{N}_{eCombo}$ is the unweighted harmonic mean of the individual $\widehat{N}_e$ values.
2. The estimators are uncorrelated. Strictly speaking, precision is maximally increased when
the estimators are negatively correlated, but that is not likely to occur in real datasets.
Therefore, in practical terms, the ideal situation occurs when positive correlations are low
or non-existent.
3. Individual estimates have low precision, which occurs when data are limited and/or true
N_e is large.
699
Condition 1 (equal variances) is ideal because the reduction in variance of the combined
estimate diminishes as the weights become more uneven (Equation 28; Figure S1). When
weights are very unequal, most of the useful information is provided by just one of the methods
and combining them provides relatively little benefit (Figure 3, bottom panel). The temporal and
LD methods make it easy to properly weight combined estimates because $\mathrm{var}(1/\widehat{N}_e)$ is
straightforward to estimate. At present there is not a simple, generic way to calculate appropriate
weights for Wang's sibship method. For some specific parameter combinations, Wang (2009,
2016) and Waples (2021) have provided estimates of $\mathrm{var}(1/\widehat{N}_e)$ or related quantities. In other

scenarios, giving equal weights to sibship and LD estimates might be reasonable as a first
approximation.
Condition 2 (uncorrelated estimators) appears to be a valid assumption for the following
scenarios: (a) combining temporal and LD estimators for the same time periods; (b) combining
LD estimates for different time periods; (c) combining LD and sibship estimates when N_e is not
too small and relatively few genetic markers are used. Although I am not aware of a direct
evaluation, it seems likely that estimates from the temporal and sibship methods are also
essentially uncorrelated. When the number of genetic markers is large ($\geq$1000 SNPs), both the
LD and sibship methods become very good at detecting essentially the same signal of genetic
drift, with the consequence that positive correlations reduce the benefits of combining estimates.
Positive correlations also limit benefits from combining temporal estimates based on overlapping
samples and/or time periods, but this is a complex topic that merits further study. One common
scenario has already been examined in detail. If a time-series of consecutive samples is taken as
part of a systematic genetic monitoring program, precision of temporal estimates of N_e in
individual generations can be improved by using information from comparisons than span one
additional generation on either side of the focal generation (using the program *MAXTEMP*
described by Waples et al. 2025).
Condition 3 represents good news for researchers, as it means that the greatest
improvements to precision can be achieved when they are needed the most.
All of the above analyses have focused on precision. Most population genetics theory
begins with simplifying assumptions to make analyses tractable (models for estimating effective
size are no exception), and violations of these core assumptions can lead to bias. Two of the
most consequential assumptions for N_e-estimation methods are that (1) generations are discrete,
and (2) the focal population is isolated. Both of these assumptions are commonly violated in real
populations. Table 3 of Waples et al. (2025) provides a compilation of numerous published
papers that have evaluated sensitivities to these and other potential sources of bias, and
researchers interested in combining estimates from multiple methods should also consider their
respective biases.
Performance of confidence intervals (summarized in Tables 1, S2, and S3) illustrates
some potential tradeoffs between precision and bias. If any estimator has even modest bias, the
bias will have more pronounced effects on CI coverage as precision increases and CI width
shrinks. Thus, it is possible for combined estimates of N_e to have increased precision but lower
CI coverage. Researchers should be aware of this possibility and consider an experimental
design that maximizes the chances of achieving their objectives.
742

References

Beaumont, M.A. and Nichols, R.A., 1996. Evaluating loci for use in the genetic analysis of population structure. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *263*(1377), pp.1619-1626.

Clarke, S.H., Lawrence, E.R., Matte, J.M., Gallagher, B.K., Salisbury, S.J., Michaelides, S.N., Koumrouyan, R., Ruzzante, D.E., Grant, J.W. and Fraser, D.J., 2024. Global assessment of effective population sizes: Consistent taxonomic differences in meeting the 50/500 rule. *Molecular Ecology*, *33*(11), p.e17353.

Cochran, W. G. (1937). Problems arising in the analysis of a series of similar experiments, Supplement to the Journal of the Royal Statistical Society 4 102- 118.

Convention on Biological Diversity (CBD). 2022. "*Decision Adopted by the Conference of the Parties to the Convention on Biological Diversity CBD/COP/DEC/15/4 Kunming-Montreal Global Biodiversity Framework*. CBD/COP/DEC/15/4. https:// www. cbd. int/ doc/ decis ions/ cop-15/cop-15-dec-04-en.pdf.

Do, C., R.S. Waples, D. Peel, G.M. Macbeth, B.J. Tillet, and J.R. Ovenden. 2014. NeEstimator V2: re-implementation of software for the estimation of contemporary effective population size (Ne) from genetic data. Molecular Ecology Resources 14:209-214 (DOI: 10.1111/1755-0998.12157).

England, P. R., Cornuet, J.-M., Berthier, P., Tallmon, D. A., & Luikart, G. (2006). Estimating effective population size from linkage disequilibrium: Severe bias using small samples. *Conservation Genetics*, *7*, 303–308.

Ewens, W.J. and Feldman, M.W., 1976. The theoretical assessment of selective neutrality. *Population genetics and ecology*, pp.303-337. S. Karlin and E. Nevo, eds. Academic Press, NY.

Gaines, M.S. and Whittam, T.S., 1980. Genetic changes in fluctuating vole populations: selective vs. nonselective forces. Genetics, 96(3), pp.767-778.

Gao, X., Starmer, J., & Martin, E. R. (2008). A multiple testing correction method for genetic association studies using correlated single nucleotide polymorphisms. *Genetic Epidemiology: The Official Publication of the International Genetic Epidemiology Society*, *32*(4), 361–369.

Hill, W.G., 1981. Estimation of effective population size from data on linkage disequilibrium. Genetics Research, 38(3), pp.209-216.

Hollenbeck, C., Portnoy, D., & Gold, J. (2016). A method for detecting recent changes in contemporary effective population size from linkage disequilibrium at linked and unlinked loci. *Heredity*, *117*, 207–216.

Jorde, P.E. and Ryman, N., 2007. Unbiased estimator for genetic drift and effective population size. *Genetics*, *177*(2), pp.927-935.

Keller, T. and Olkin, I., 2004. Combining correlated unbiased estimators of the mean of a normal distribution. *Lecture Notes-Monograph Series*, pp.218-227.

Lewontin, R.C. and Krakauer, J., 1973. Distribution of gene frequency as a test of the theory of the selective neutrality of polymorphisms. *Genetics*, *74*(1), pp.175-195.

Lotterhos, K.E. and Whitlock, M.C., 2014. Evaluation of demographic history and neutral parameterization on the performance of FST outlier tests. Molecular ecology, 23(9), pp.2178-2192.

Mastretta-Yanes, A., J. M. Da Silva, C. E. Grueber, et al. 2024. "Multinational Evaluation of Genetic Diversity Indicators for the Kunming-Montreal Global Biodiversity Framework." *Ecology Letters* 27, no. 7: e14461.

Nadachowska-Brzyska, K., M. Konczal, and W. Babik. 2022. "Navigating the Temporal Continuum of Effective Population Size." *Methods in Ecology and Evolution* 13: 22‑41.

Nei, M. and Tajima, F., 1981. Genetic drift and estimation of effective population size. *Genetics*, *98*:625-640.

Nomura, T., 2008. Estimation of effective number of breeders from molecular coancestry of single cohort sample. *Evolutionary applications*, *1*(3), pp.462-474.

Palstra, F. P., and D. J. Fraser. 2012. "Effective/Census Population Size Ratio Estimation: A Compendium and Appraisal." *Ecology and Evolution* 2: 2357‑2365.

Pollak E (1983) A new method for estimating the effective population size from allele frequency changes. Genetics, 104,531–548.

Pudovkin, A.I., Zaykin, D.V. and Hedgecock, D., 1996. On the potential for estimating the effective number of breeders from heterozygote-excess in progeny. *Genetics*, *144*(1), pp.383-387.

Robertson, A., 1975. Gene frequency distributions as a test of selective neutrality. *Genetics*, *81*(4), pp.775-785.

Santiago, E., Novo, I., Pardiñas, A. F., Saura, M., Wang, J., & Caballero, A. (2020). Recent demographic history inferred by high-resolution analysis of linkage disequilibrium. *Molecular Biology and Evolution*, *37*(12), 3642–3653.

Shahar, D.J., 2017. Minimizing the variance of a weighted average. *Open Journal of Statistics*, *7*(2), pp.216-224.

Tenesa, A., Navarro, P., Hayes, B. J., Duffy, D. L., Clarke, G. M., Goddard, M. E., & Visscher, P. M. (2007). Recent human effective population size estimated from linkage disequilibrium. *Genome Research*, *17*, 520–526.

Wang, J., 2001. A pseudo-likelihood method for estimating effective population size from temporally spaced samples. *Genetics Research*, *78*(3), pp.243-257.

Wang J (2009) A new method for estimating effective population size from a single sample of multilocus genotypes. Molecular Ecology, 18, 2148–2164.

Wang, J., 2016. A comparison of single-sample estimators of effective population sizes from genetic marker data. *Molecular Ecology*, *25*(19), pp.4692-4711.

Wang, J. and Santure, A.W., 2009. Parentage and sibship inference from multilocus genotype data under polygamy. *Genetics*, *181*(4), pp.1579-1594.

Wang, J., E. Santiago, and A. Caballero. 2016. "Prediction and Estimation of Effective Population Size." *Heredity* 117: 193–206.

Waples, R.S. 1989. A generalized approach for estimating effective population size from temporal changes in allele frequency. Genetics 121:379-391.

Waples, R.S. 2005. Genetic estimates of contemporary effective population size: To what time periods do the estimates apply? Molecular Ecology 14:3335-3352.

Waples, R.S. 2006. A bias correction for estimates of effective population size based on linkage disequilibrium at unlinked gene loci. Conservation Genetics 7:167-184.

Waples, RS. 2021. Relative precision of the sibship and LD methods for estimating effective population size with genomics-scale datasets. Journal of Heredity 112:535:539.

Waples, R.S. 2024. Practical application of the linkage disequilibrium method for estimating contemporary effective population size: a review. Molecular Ecology Resources 24:e13879

Waples, R.S. 2025. The Idiot's Guide to effective population size. Molecular Ecology
https://doi.org/10.1111/mec.17670

Waples, R.S., and C. Do. 2008. *LDNE*: A program for estimating effective population size from data on linkage disequilibrium. Mol. Ecol. Resources 8:753-756.

Waples, R.S., and C. Do. 2010. Linkage disequilibrium estimates of contemporary N_e using highly variable genetic markers: A largely untapped resource for applied conservation and evolution. Evolutionary Applications 3:244-262.

Waples, R.S., and J.R. Faulkner. 2009. Modeling evolutionary processes in small populations: Not as ideal as you think. Molecular Ecology 18:1834-1847.

Waples, R.S., Waples, R.K., and Ward, E.J., 2022. Pseudoreplication in genomics-scale datasets. Molecular Ecology Resources 22:503-518.

Waples, R.S., M.M. Masuda, M.E.F. LaCava, A.J. Finger. 2025. *MaxTemp*: A method to maximize precision of the temporal method for estimating N_e in genetic monitoring programs. Molecular Ecology Resources 25:e14057.

Weir, B.S., 1979. Inferences about linkage disequilibrium. *Biometrics*, pp.235-254.

Weir, B. S., & Hill, W. G. (1980). Effect of mating structure on variation in linkage disequilibrium. *Genetics*, *95*(2), 477–488.

Wright, S., 1965. The interpretation of population structure by F-statistics with special regard to systems of mating. *Evolution*, pp.395-420.

Data Accessibility and Benefit-Sharing
Data accessibility statement

No data were used in this paper except those that a) were generated by simulations, or b) have been previously published. Code to conduct simulations described here will be posted on Zenodo on acceptance.

Benefit-sharing statement

This is a theoretical paper, so this statement is not applicable.

Table 1. Performance metrics for combined estimates based on the LD and temporal estimators. Results are for simulations using sample sizes of individuals (S) ranging from 25 to 100 and numbers of unlinked, diallelic (SNP) loci ranging from 100 to 5000. True N_e was 200. A: Correlations between $1/\widehat{N}_e$ for the two methods. B: Confidence interval coverage—percentage of replicates for which the 95% confidence interval around $\widehat{N}_{eCombo}$ contained the true value. C: ratio of $\mathrm{var}(1/\widehat{N}_{eCombo})$ to $\mathrm{var}(1/\widehat{N}_e)$ for LD method. D: ratio of the root-mean-squared error (RMSE) of $1/\widehat{N}_{eCombo}$ to RMSE$(1/\widehat{N}_e)$ for LD method. For the scenarios modeled, $\mathrm{var}(1/\widehat{N}_e)$ and RMSE$(1/\widehat{N}_e)$ were both smaller for the LD method than for the temporal method.

A

S	Number of loci		
	100	1000	5000
25	-0.005	-0.007	0.004
50	0.000	-0.006	-0.002
100	-0.010	0.004	0.002

C

S	Number of loci		
	100	1000	5000
25	0.94	0.87	0.62
50	0.94	0.89	0.65
100	0.95	0.96	0.87

B

S	Number of loci		
	100	1000	5000
25	94.8	94.8	73.1
50	96.3	94.0	74.5
100	99.1	99.2	95.1

D

S	Number of loci		
	100	1000	5000
25	0.97	0.92	0.76
50	0.97	0.94	0.79
100	0.97	0.98	0.92

Table 2. Correlations between $1/\widehat{N}_e$ for the sibship and LD methods in simulated data. Simulations used the software that is available in Supporting Information for Waples (2021), which assumes that all sibling assignments are made without error. True N_e was 200 (top) and 2000 (bottom). S is the number of individuals sampled.

N_e=200	Number of loci			
S	10	100	1000	2500
25	0.024	0.085	0.171	0.211
50	0.034	0.089	0.844	0.892
100	0.029	0.306	0.791	0.839

N_e=2000	Number of loci			
S	10	100	1000	2500
25	0.011	0.117	0.335	0.183
50	-0.011	0.060	0.302	0.285
100	0.001	0.032	0.730	0.904

Figure 1. Distribution of $\hat{N}_e$ (top) and $1/\hat{N}_e$ (bottom) in simulated data using the temporal method. The data are from Figure 6 of Waples (2025) for a scenario in which true N_e was 500 and temporal samples of 50 individuals each were taken one generation apart and scored for 100 diallelic loci. In the top panel, negative $\hat{N}_e$ values (farthest left bar) are interpreted as infinity (inf; no evidence for genetic drift). Note the log scale on the X axis of the top panel.

Figure 2. Relative precision of the temporal and LD methods for estimating effective size. The Y axis shows theoretical ratio (R) of var($1/\widehat{N}_e$) for the temporal method to var($1/\widehat{N}_e$) for the LD method, as a function of true N_e, the number of SNPs used to estimate effective size, and elapsed time between temporal samples. Temporal estimates used 2 samples of 50 individuals; LD estimates used one sample of 50 progeny. Top: Effects of N_e, with all temporal samples taken 1 generation apart. Bottom: results for temporal samples taken $t = 1$, 3, or 10 generations apart, with N_e fixed at 200. R was computed using Equation 23, assuming Plan II sampling for the temporal method. The horizontal dotted line indicates equal variances for the two methods. These results account for pseudoreplication (lack of independence) caused by physical linkage and (for the LD method) overlapping pairs of the same loci, as quantified by Waples et al. (2022). These results assume 16 chromosomes of 50Mb (mean chromosome numbers for plants, invertebrates, and vertebrates are 13, 11, and 25, respectively; Li et al. 2011). Note the log scale on both axes.

Figure 3. Variance of a combined estimate (var($1/\widehat{N}_{eCombo}$); Y axis), as a function of the relative weight for the LD method (W_{LD}; X axis). Combined estimates were computed as $1/\widehat{N}_{eCombo} = W_{LD}/\widehat{N}_{eLD} + W_{Temp}/\widehat{N}_{eTemp}$, where $W_{Temp} = 1- W_{LD}$. Filled circles are results from simulations; vertical lines show the theoretical optimum weighting scheme based on Equations 11, 16, and 28. Horizontal dotted lines show the range of values of W_{LD} that produce var($1/\widehat{N}_{eCombo}$) $\leq$ var($1/\widehat{N}_{eLD}$). Simulation parameters were $N_e = 200$, $S = 50$, t=1 for the temporal method, and using L=5000 (top) or 500 (bottom) unlinked, diallelic loci.

Figure 4. Top: Distribution of $\hat{N}_e$ in random simulated data. Two series of 2000 paired, random normal datapoints were simulated. Both series (A and B) had an expected value of $1/N_e = 1/1000$, and the variance of B was twice that of A. A third, combo series ($C=1/\hat{N}_{eCombo}$) was generated as the weighted average of A and B, with weights of 0.67 and 0.33, respectively, based on reciprocals of variances. The inverse of the simulated data are the $\hat{N}_e$ values shown in the graph, with true N_e (1000) indicated by a dotted vertical line. The last group of bars on the right include all $\hat{N}_e > 2000$. Black bars show the distribution of Series A. White bars show the distribution of $\hat{N}_{eCombo}$ when A and B are modeled as uncorrelated (realized correlation coefficient = -0.02), and gray bars show the distribution of $\hat{N}_{eCombo}$ when A and B are modeled with expected correlation 0.5 (realized correlation coefficient = 0.51). Bottom: RMSE of $1/\hat{N}_e$ for Series A and for $1/\hat{N}_{eCombo}$ in the uncorrelated and correlated scenarios.