Post-translational modifications of microtubules are crucial for malaria parasite transmission

Kodzo Atchou[1,2*], Magali Roques[1], Ruth Rehmann[1], Reto Caldelari[1], Melanie Schmid[1,2], Simone
Grossi[1], Bianca Manuela Berger[1,2], Torsten Ochsenreiter[1], Friedrich Frischknecht[3], Volker
Heussler[1*]

[1] Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland

[2] Graduate School for Cellular and Biomedical Sciences, University of Bern, 3012 Bern, Switzerland

[3] Department of Parasitology, University of Heidelberg Medical School, Im Neuenheimer Feld 324, 69120,
Heidelberg, Germany

* Corresponding authors

Abstract

Microtubules, composed of α- and β-tubulin polymers, are essential components of the eukaryotic
cytoskeleton. They maintain cellular shape and structural integrity and play critical roles in cell division and
in intracellular vesicular transport. In *Plasmodium,* the parasite that causes malaria, nuclear replication
during the liver stage is among the fastest known in eukaryotic cells and relies heavily on microtubules for
DNA segregation and cytoskeletal organization. Despite their importance, the role of microtubules in liver
stage development remains largely unexplored. Here, we investigated microtubule dynamics during liver
stage development using a combination of cell and molecular biology techniques, expansion microscopy,
and live-cell imaging. By employing antibodies specific for α-tubulin post-translational modifications (PTMs),
we found that the *Plasmodium* sporozoites subpellicular microtubules (SSPM) persist during liver infection,
giving rise to liver stage parasite microtubule bundles (LSPMB). These LSPMB form multimeric tubulin
structures within hepatocytes and are redistributed to the hemi-spindle poles of parasite nuclei during
schizogony. Deletion of the C-terminal region encompassing all known *Plasmodium* α-tubulin PTM sites
prevented sporozoite migration from the mosquito midgut to the salivary glands, effectively blocking
parasite transmission. Using *Plasmodium* microtubule-specific depolymerisation drugs, we found that while
LSPMB are stable in sporozoites, they exhibit dynamic behavior during hepatocyte infection. Given the
regulatory role of PTMs in microtubule dynamics, we generated parasite mutants by substituting and
deleting key α-tubulin C-terminal residues involved in PTMs. Substitution of the polyglutamylation site with
alanine and deletion of the C-terminal tyrosination/detyrosination motifs impaired parasite growth during
liver infection. Together, our findings reveal extensive microtubule remodeling during liver stage
development and establish α-tubulin C-terminal modifications as critical regulators of both intracellular
development and parasite transmission of *Plasmodium* parasites.

Introduction

Plasmodium parasites are responsible for malaria and cause yearly half a million deaths in sub-Saharan
countries. Malaria continues to be a major worldwide health concern despite recent successful attempts to
lower its prevalence (1–4). The infection starts when an infected female *Anopheles* mosquito, during a
blood meal, injects salivary gland sporozoites into the skin of the vertebrate host. Some of the motile
sporozoites escape the host immune defence and invade blood vessels. They are carried by the blood flow
to the liver, where they invade and infect hepatocytes (5). Approximately 50% of the parasites are often
killed by host cell responses (6–8). Those that managed to establish a productive invasion form a
parasitophorous vacuole (PV) in which the parasite undergoes massive growth and asexual replication
known as schizogony. One single parasite gives rise to thousands of daughter parasites called merozoites
in 3 to 6 days depending on the *Plasmodium* species (9–11). Fully developed merozoites exit the liver cells

in merosome vesicles through adjacent blood vessels (11–14). The merosomes then rupture and release
merozoites into the blood stream. The parasites immediately invade erythrocytes, where they undergo
again asexual replication to form the blood stage merozoites. Some merozoites differentiate into sexual
stages, termed gametocytes, which are ingested by female mosquitos during a blood meal. In the mosquito
midgut, the gametocytes develop into microgametes and macrogametes, which fertilize and form a zygote.
The later undergoes differentiation into motile ookinetes, which penetrate the gut wall, forming oocysts (15–
18). The oocyst produces the midgut sporozoites from the sporoblasts. The sporozoites migrate to the
salivary gland from day 11 post-infection in *P. berghei* (Frischknecht et al., 2004; Frischknecht &
Matuschewski, 2017; jerome P. Vanderberg, 1974; J. P. Vanderberg, 1975). During the liver stage
development, the parasite relies on the fast and massive asexual replication which is essential for the
transmission and pathogenesis in erythrocyte stage, representing key targets for malaria control strategies.
Investigating the parasite microtubule dynamics during the rapid replication could provide important insight
on the parasite biology and to develop strategies for halting the massive proliferation during the clinically
asymptomatic liver stage.
Microtubules are cytoskeletal filaments of about 25nm outer diameter, made up of usually 13 chains of
protofilaments composed of globular α-tubulin and β-tubulin heterodimers. They are crucial components of
the cytoskeleton and offer structural support to cells and also function as tracks for the movement of
organelles, vesicles, and other cellular components (23,24). Microtubules are involved in important cellular
activities such as cell division and the production of cilia and flagella to allow cell motility. Microtubules
exhibit inherent dynamics, alternating between polymerisation (rapid growth) and spontaneous
depolymerisation (shrinkage) in a process termed dynamic instability (23,25,26). The dynamics of
microtubules and their interactions with other cellular structures are regulated by specific microtubule-
associated proteins (MAPs), which selectively bind to either the plus or minus ends of microtubules (27–
30).
Tubulin is subject to seven identified post-translational modifications (PTMs), among which are acetylation
of the lysine residue at position 40 in the luminally exposed part of α-tubulin, glycylation, which entails the
incorporation of polyglycine side chains of variable length, phosphorylation, palmitoylation, tyrosination/
detyrosination and polyglutamylation. These modifications are extensively preserved and enabling cells to
modulate microtubule characteristics, dynamics, and interactions with microtubule-associated proteins
(MAPs) and motor proteins (31,32). Tyrosinated microtubules are often associated with microtubule
stability. These stable microtubules represent a specialised subset that exhibits enhanced resistance to
depolymerization, even in the presence of destabilizing agents. They play a crucial role in maintaining
cellular architecture, facilitating intracellular transport, and supporting specialized cellular functions. In
contrast to dynamic microtubules, which continuously undergo cycles of polymerization and
depolymerization due to dynamic instability, stable microtubules persist for extended periods. They are also
frequently polyglutamylated, a modification that regulates their interactions with motor proteins and other
cellular components. These microtubules are predominantly localized in structures requiring sustained
structural integrity, such as cilia, flagella, neuronal axons, and motile cell types (33–36). Glutamylation, or
polyglutamylation, is a post-translational modification characterized by the addition of multiple glutamate
side chains to glutamate residues on both α- and β-tubulin. This modification occurs primarily at the C-
terminal tails and is catalyzed by a family of enzymes known as polyglutamylases. Polyglutamylation plays
a critical role in regulating the interactions of microtubules with MAPs and motor proteins, thereby
influencing microtubule dynamics and cellular functions (37–40). The properties and functions of tubulin
are largely encoded within their protein sequences, contributing to molecular diversity that influences key
microtubule characteristics. This variation has been conceptualized as the "tubulin code", a regulatory
framework through which cells control microtubule diversity to achieve specific physiological functions (24).
Nevertheless, the precise mechanisms by which cells establish and interpret this code remain largely
unknown. Although tubulin is highly conserved across evolution, it exhibits structural and functional
variations between species, cell types, and even within single cells.

In eukaryotic cells, tubulins are key components of the microtubule cytoskeleton, and their genetic
complexity reflects the organism's level of cellular specialization. In yeast, two distinct genes encode α-
tubulin, while a single gene encodes β-tubulin. In contrast, the human genome contains nine genes
encoding α-tubulin and an equal number encoding β-tubulin, reflecting a greater level of complexity and
functional specialization in higher eukaryotes (24,41,42).
The *Plasmodium* genome encodes two α-tubulins (α1-tubulin and α2-tubulin) and one β-tubulin (43–47).
The α1-tubulin gene encodes a protein of 450 amino acids, while α2-tubulin encodes 453 amino acids (48).
The two isoforms share 94% amino acid identity. However, α2-tubulin lacks a terminal tyrosine residue,
which is present in α1-tubulin. This difference suggests potential variation in post-translational modifications
and functional roles (44,49). In *P. berghei,* the α1-tubulin gene is expressed across most stages of
the *Plasmodium* life cycle, whereas α2-tubulin expression appears restricted to male gametocytes,
gametes, and newly formed zygotes, indicating distinct functional roles for the two isoforms (32,50).
Deletion of α1-tubulin blocks sporozoite formation in oocysts and lowering α1-tubulin expression leads to
less infective sporozoites (50).
There is a significant degree of amino acid conservation between human and *Plasmodium* tubulins, with
approximately 83.7% identity for α-tubulin and 88.5% for β-tubulin. However, *Plasmodium* tubulin shows
greater similarity to plant tubulin than to mammalian tubulin. This raises the potential for developing
compounds that selectively target parasite microtubules while sparing the host cell cytoskeleton, offering a
promising avenue for therapeutic intervention (43,45–47,51).
Recent studies on *Plasmodium falciparum* blood stage schizonts have revealed that hemi-spindles lack
polyglutamylation, whereas subpellicular microtubules do exhibit this post-translational modification (52–
54). Similarly, SPMTs appear to be stabilized by microtubule inner proteins, while spindle microtubules do
not show these luminal proteins (55). These differences may reflect functional specialisation of microtubule
subpopulations in different cellular structures and developmental stages.
The liver stage of *Plasmodium* exhibits one of the fastest rates of replication known among eukaryotic cells,
a process likely dependent on microtubule dynamics. However, to date, no studies have investigated
microtubule dynamics during the liver stage. This study was therefore designed to explore the dynamics of
parasite tubulin during the asexual liver stage, using recently developed expansion microscopy protocols
(54,56–58).
122
Results

Expansion microscopy reveals the presence of the sporozoites subpellicular microtubules during
the liver stage of infection

To investigate microtubule organization during the *Plasmodium* parasite liver stage, HeLa cells were
infected with sporozoites and fixed them at different time points using cold methanol. We then performed
IFA to stain host and parasite microtubules using an α-tubulin antibody (Figure 1A). The IFA revealed a
marked accumulation of microtubules in proximity of the PVM (stained with anti-UIS4) from early infection
(24 hours post-infection (hpi)) to the schizogony (30 to 54 hpi) (Figure 1A inlets). To examine these
structures in greater detail, we applied the recently developed pre-staining expansion microscopy (PS-ExM)
(56). By expanding the cells five-fold, PS-ExM enabled visualization of a large multimeric tubulin structure
within the parasite, detectable from 6 hours post-infection (hpi) to 48 hpi (Figure 1B). Based on observations
of sporozoites prior to liver infection, we hypothesize that this structure originates from the sporozoite
subpellicular microtubules (SSPM) (Figure 1C top panel). We refer to this structure as the liver stage
parasite microtubule bundles (LSPMB). Notably, the LSPMB progressively shrank over time and was no
longer detectable by the end of schizogony (54 hpi) (Figure 1A, B). Using the 3D and 4D imaging software
IMARIS, we quantified the LSPMB volume at various stages of liver infection. The results confirmed a
consistent decrease in LSPMB size between 6 and 56 hpi (Figure 1D).

To verify that the LSPMB is not an artifact of the IFA and PS-ExM, we performed live cell imaging of infected
HeLa cells. Cells were incubated with the tubulin tracker SPY650 for one hour prior to imaging. The LSPMB
was clearly visible in sporozoites before invasion (Movie 1, ~10 hpi) and persisted until late liver stage
development, gradually shrinking during schizogony (Figure 1E; Movies 2–4, ~36 hpi to 48 hpi). These live
imaging results corroborate our findings from IFA and expansion microscopy, validating the presence and
dynamic nature of the LSPMB during liver stage development.

The liver stage parasite subpellicular microtubules are dynamic

The SSPM has previously been described as highly stable structures, enabling sporozoite formation and
invasion (50,55,59). These microtubules, similar to the corresponding microtubules in *T. gondii*, are
resistant to conventional depolymerizing agents (59). To investigate whether the LSPMB retain this stability
during liver stage development, we treated infected HeLa cells with the microtubule-depolymerizing agents
Amiprophos-methyl (AMP) and oryzalin herbicide. These compounds are known to selectively target
dynamic *Plasmodium* microtubules without affecting mammalian cells (43,45,47,51,60) (Figure 2 and
Supplementary Figure 1A and B). We first validated drug specificity and cytotoxicity by testing various
concentrations on HeLa cells over one week. Neither compound affected host cell viability (Supplementary
Figure 1A and B) or disrupted host microtubules (Supplementary Figure 1Dand E). However, oryzalin
showed a stronger inhibitory effect on parasite growth compared to AMP (Supplementary Figure C) and
was therefore selected for further experiments (Figure 2A). Sporozoites were dissected into medium
containing 10 μM or 20 μM oryzalin and then used to infect HeLa cells (Figure 2A and B), which were
incubated with the same drug concentrations. At 48 hpi, the cells were fixed and analyzed by IFA and PS-
ExM. To test reversibility, oryzalin was washed out at 24 hpi prior to the onset of schizogony (Figure 2B).
At 10 μM, oryzalin significantly impaired nuclear replication, spindle formation, and LSPMB volume when
present the entire cultivation time (Figure 2C, D and E). Partial recovery of hemi-spindles and nuclear
replication was observed, suggesting that the effect of 10 μM oryzalin is reversible at this stage (Figure 2C,
D, and E). However, at 20 μM, neither the hemi-spindles nor the LSPMB recovered after drug removal,
indicating irreversible microtubule disruption at higher concentrations (Figure 2A and B). To investigate
whether the microtubule depolymerization by oryzalin affects the parasite growth and survival during the
liver stage, we measured parasite size within infected HeLa cells at 6, 24 and 48 hpi. Treatment with 10 μM
oryzalin led to a significant reduction in parasite size and survival (Figure 2 F, G, H, and I). We further
examined merozoite formation by staining for MSP1 at 56 hpi. Parasites treated with 10 μM oryzalin failed
to express MSP1, indicating a block in merozoite development and suggesting that intact microtubules are
essential for DNA replication and merozoite formation (Figure 2J). Interestingly, at lower concentrations of
0.5 μM, oryzalin treatment led to an unexpected elongation of the LSPMB and hemi-spindles up to four
times longer than controls suggesting that microtubule turnover is tightly regulated (Supplementary Figure
1F, G and H). In conclusion, oryzalin-mediated depolymerization confirms that both the LSPMB and the
hemi-spindles are dynamic structures during the liver stage. This dynamic behaviour was further supported
by live-cell imaging using the SPY650 tubulin tracker (Movie 5, 6, and 7).
Given the observed microtubule remodeling in response to oryzalin, we next sought to determine whether
known microtubule-associated proteins, specifically those regulating plus-end dynamics, are expressed
during the liver stage and might contribute to the observed plasticity. Only a limited number of motor proteins
and microtubule associated proteins, such as centrin and the kinesin protein family, have been identified
during the *Plasmodium* liver stage (11,61–63). The recently characterized EB1 protein, known to regulate
microtubules plus-end dynamics and length, has so far been reported for being expressed during the sexual
blood stage (Mauer et al., 2023; Yang et al., 2023; Zeeshan et al., 2023). However, these studies did not
assess EB1 expression during the asexual mosquito and liver stages. Given our observations of dynamic
microtubule behavior and excessive polymerization following oryzalin treatment, we explored whether EB1
might be expressed during the liver stage. To this end, we generated a parasite line expressing endogenous

EB1 fused to eGFP via double homologous recombination (Supplementary Figure 2C and D). As expected,
EB1-eGFP was detected in sexual blood stages (Supplementary Figure 2E and F) consistent with previous
reports (64,65). Interestingly, we also observed EB1 expression in both midgut and salivary gland
sporozoites (Supplementary Figure 2G). Importantly, EB1 was also detected during liver stage
development, where it localized to the LSPMB and hemi-spindles (Supplementary Figure 2H–I).
These findings suggest that EB1 may play a broader role in *Plasmodium* microtubule regulation than
previously recognized, potentially contributing to the dynamic remodeling of microtubules during the liver
stage.
195
The liver stage parasite subpellicular microtubule carries tubulin post-translational modifications

Microtubule functions are often regulated by different post-translational modifications (PTMs), which
influence their stability, interactions, and dynamics. To gain insight into the dynamic nature of LSPMB during
the liver stage development, we examined α-tubulin PTMs using immunofluorescence (IFA) and expansion
microscopy (Figure 3 and Movie 5, 6 and 7). To investigate whether the LSPMB is present in cell types
other than HeLa cells, we infected primary hepatocytes with *Plasmodium* sporozoites and fixed the cells at
different timepoints (6, 24, 36, 48, 54 and 56 hpi). We also analyzed sporozoites isolated from mosquito
salivary glands. Using antibodies specific for polyglutamylated (anti-IN105) (66) (Figure 3A) and tyrosinated
(anti-Ty) α-tubulin (Figure 3B), we observed that sporozoite microtubules, including the LSPMB, are both
polyglutamylated and tyrosinated. Quantitative colocalization analysis confirmed a strong overlap between
LSPMB and polyglutamylated tubulin in both sporozoites and liver stage parasites (Supplementary Figure
2A). Interestingly, following hepatocyte invasion, the LSPMB lost tyrosination (i.e., became detyrosinated)
while retaining polyglutamylation. This detyrosination persisted throughout liver stage development,
including early and late schizogony (24–56 hpi). During this period, polyglutamylation was observed on the
LSPMB and at the poles of the hemi-spindles, but not along the polymerizing hemi-spindles themselves,
which eventually disappeared along with the LSPMB (Figure 3A and C). Given previous findings implicating
polyglutamylation in hemi-spindle pole formation in erythrocytic merozoites (31), we investigated whether
there is a structural relationship between the LSPMB and the hemi-spindle poles. Using PS-ExM and the
tubulin tracker SPY650, we visualized direct connections between LSPMB filaments and hemi-spindle
poles. Live cell imaging confirmed this connection between LSPMB and hemi-spindle poles, mainly during
the shrinkage of the LSPMB (Fig 3 F; Movie 8, 9 and 10, Supplementary Movie 1). These observations
suggest that the LSPMB may be redistributed to form or support the hemi-spindle poles during schizogony.
We also stained for γ-tubulin and observed its localization at the hemi-spindle poles in both the parasite
and host cell (Supplementary Figure 2B), supporting the idea that *Plasmodium* may utilize γ-tubulin for
nuclear division and microtubule assembly, as reported in other systems (67–70).
To further define the spatial localization of the LSPMB, we used *P. berghei* parasites expressing the plasma
membrane protein PMP1 fused to superfolder GFP (sfGFP) previously generated (71). ImageJ RGB
profiling revealed that the LSPMB lies between the parasite plasma membrane (PM) and the
parasitophorous vacuole membrane (PVM), suggesting that the sporozoite subpellicular microtubules
undergo subcellular relocalization during liver stage development (Figure 3D and E). At later stages,
particularly the cytomere stage, we observed reappearance of tyrosinated tubulin near the parasite nuclei
(Figure 3B). During merozoite formation, this tyrosinated tubulin appeared as punctate structures within the
forming merozoites (Figure 3B).
Tyrosination has previously been associated with stable microtubules involved in motility, such as those in
sporozoites, flagella, and spermatozoa (33,34,50,55). The absence of tyrosination during early intracellular
liver stage development when the parasite is non-motile may facilitate its transformation from a banana-
shaped sporozoite to a rounded liver stage form (Figure 1 and 3). LSPMB detyrosination may contribute to
microtubule dynamic instability, providing both energy and tubulin monomers for rapid growth and spindle

formation during schizogony. Finally, the reappearance of tyrosinated microtubules during merozoite
formation may serve to stabilize the microtubules required for merozoite motility and erythrocyte invasion.

The *Plasmodium* α-tubulin C-terminal modification are important for sporozoite formation

To investigate the functional role of parasite α-tubulin PTMs during the liver stage development, we
generated *P. berghei* mutant parasite lines with targeted mutations in the α-tubulin C-terminal region, where
most PTMs are predicted to occur (Figure 4A and Supplementary Figure 3A and B). Specifically, we created
three mutant lines: PsΔ: a point mutation replacing the polyglutamylation site glutamic acid (E445) with
alanine (A), TyΔ: deletion of the tyrosination/detyrosination motif (YEADY) from position 449, CtΔ: deletion
of the entire C-terminal tail (EGEDEGYEADY) to prevent all possible C-terminal tail PTMs (Supplementary
Figure 3A to C).
Using AlphaFold predictions, we confirmed that the mutations were unlikely to affect tubulin folding
(Supplementary Figure 3C). The mutants were generated via double homologous recombination and
verified by integration PCR and sequencing of the mutated α-tubulin gene (Figure 4A and B; Supplementary
Figure 3B). The development of mutant parasites was investigated during the mosquito, and liver stages.
Significant defects emerged during mosquito stage development. The PsΔ and TyΔ mutants were able to
form oocysts (Figure 4 C, Supplementary Fig 4A) and gliding sporozoites, albeit in reduced numbers (Figure
4D, E and G; Supplementary Fig 4B; and Supplementary Movie 2, 3, 4 and 5). Quantification of salivary
gland sporozoites at 18 days post-blood feeding showed that PsΔ and TyΔ mutants produced
approximately half as many sporozoites as the control line (NoΔ) (Figure 4D). We next examined sporozoite
motility, which typically consists of continuous circular gliding (16,72–76). PsΔ and TyΔ salivary gland
sporozoites were assessed and compared to controls. While PsΔ have a similar gliding velocity compared
to controls, TyΔ sporozoites displayed increased gliding velocity (2.5 μm/s vs. 1.4 μm/s) (Figure 4G),
suggesting a regulatory role of the tyrosination motif in fine-tuning motility. To assess whether these motility
phenotypes influenced infectivity, we performed in vivo infection assays with PsΔ and TyΔ. All mutant lines
sporozoites were able to infect mice similarly to control parasites (Supplementary Figure 4C). However, the
CtΔ mutant failed to produce salivary gland sporozoites entirely (Figure 4D). Since CtΔ mutant parasites
failed to reach the salivary glands, we analyzed hemolymph sporozoites for this line compared to the
controls (Figure 4F). These showed severely impaired motility, lacking the typical gliding pattern and
exhibiting significantly reduced speed (Figure 4F and H). Given that PsΔ and TyΔ mutants produced motile
salivary gland sporozoites, we infected HeLa cells with PsΔ and TyΔ mutant sporozoites and CtΔ mutant
hemolymph sporozoites to assess their liver stage development and identify any downstream phenotypic
effects.

The *Plasmodium* α-tubulin C-terminal mutants undergo liver stage development

Using polyglutamylation-specific antibodies, we found that PsΔ mutant parasites still exhibited
polyglutamylated liver stage parasite microtubule bundles (LSPMBs) in both sporozoites and liver stage
forms, although the signal intensity was significantly reduced compared to controls (Figure 5A, B and C and
G). As tyrosination was previously observed only in sporozoites and late liver stages, we performed PS-
ExM on sporozoites and stained them with α-tubulin tyrosination-specific antibodies. TyΔ salivary gland
sporozoites and CtΔ hemolymph sporozoites both showed detectable tyrosination signals, although the
signal in CtΔ parasites was markedly weaker than in controls (Figure 5D, E and F). Interestingly, as PTM-
specific antibodies continued to recognize signals in the PsΔ and TyΔ mutants, it is possible that the
parasite compensates by relocating the PTM sites to adjacent residues within the C-terminal sequence.
Alternatively, the antibody signal may stem from β-tubulin modifications rather than α-tubulin.
To assess parasite survival and growth during liver stage development, we performed quantitative
microscopy (Figure 5H to O). Both PsΔ and TyΔ mutant parasites successfully infected HeLa cells and
progressed through liver stage development (Figure 5B, D, H, I, J and K). This was further confirmed by

staining the parasitophorous vacuole membrane (PVM) using anti-UIS4 antibodies (Figure 5A, B and C).
Parasite size was measured at 6 hpi and 48 hpi to assess developmental progression. No significant
differences were observed at 6 hpi (Figure 5I). However, by 48 hpi, mutant parasites were significantly
smaller than controls, indicating impaired growth (Figure 5J). Survival rates of PsΔ and TyΔ mutants were
comparable to controls (Figure 5K), with infection rates of 15% for PsΔ (matching the control) and 27%
for TyΔ, exceeding the control. As expected, the CtΔ hemolymph sporozoites failed to infect and to develop
liver stage forms (Figure 5C, L, M, N and O). The sporozoites failed to establish productive infections with
a typical PVM at all examined time points (Figure 5C, M, N and O).
To confirm the phenotypes, we compared our mutants with previously generated *P. berghei* α1-tubulin
mutants described by Spreng et al. (2019), including parasites with intron deletions (intronΔ), C-terminal
tyrosination deletions 540ADY (C-termΔ), and α2-tubulin exon swaps. These mutants showed similar
phenotypes, including a complete absence of salivary gland sporozoites in α2-tubulin exon swap parasites,
mirroring the CtΔ mutant phenotype (Figure 4D and H; Supplementary Figure 3D and E). These results
underscore the importance of α-tubulin C-terminal PTMs in sporozoite formation and migration from the
midgut to the salivary glands.
The failure of CtΔ parasites to generate viable sporozoites highlights the essential role of α-tubulin PTMs
in sporozoite development and migration. Co-mutation of both α- and β-tubulin PTM sites may prove lethal
to parasite development in the mosquito stage. Future studies should focus on identifying and
characterizing the enzymes responsible for tubulin PTMs in *Plasmodium*, and genetic knockout strategies
should be pursued to further elucidate the function of these modifications.

Discussion

Our study provides the first in-depth characterization of microtubule architecture and dynamics during
the *Plasmodium* liver stage. Using tubulin trackers and α-tubulin antibodies, we identified two major
microtubule populations: a large, multimeric subpellicular network that we termed the liver stage parasite
microtubule bundles (LSPMB), and the polymerizing hemi-spindles involved in nuclear division. The
LSPMB appears to originate from the SSPM and persists until late schizogony. These observations align
with earlier electron microscopy studies by Meis et al. (77) who reported the presence of subpellicular
microtubules and inner membranes in liver stage parasites up to the point of mitotic spindle formation.
The behavior and function of microtubules are closely regulated by PTMs. We found that α1-tubulin
expressed during the liver stage carries residues associated with polyglutamylation and tyrosination. Using
PTM-specific antibodies, we confirmed that *Plasmodium* sporozoites display both modifications on their
subpellicular microtubules. Tyrosination, a well-known marker of stable microtubules, has been described
in highly motile systems such as cilia, flagella, and spermatozoa (33,34,50,55). This is consistent with the
need for high motility in sporozoites, which must traverse multiple tissue barriers during transmission
(16,72,73).
Polyglutamylation, in contrast, introduces negative charges on the tubulin surface that enhance interactions
with MAPs, and is commonly associated with dynamic microtubule behavior (78–80). This modification has
previously been linked to merozoite formation and intracellular trafficking in *Plasmodium* blood stages (31).
The co-occurrence of tyrosination and polyglutamylation in sporozoites likely ensures a microtubule
network that is both stable and functionally interactive during mosquito stage migration.
Upon hepatocyte invasion, we observed a loss of tyrosination while polyglutamylation was maintained,
suggesting a functional switch from stable to dynamic microtubules (Figure 6). This transition may facilitate
morphological transformation from a motile, slender sporozoite into a rounded liver stage form and could
also support the parasite's rapid replication. The liver stage is one of the fastest-replicating phases in
the *Plasmodium* life cycle, with replication rates among the highest observed in eukaryotic cells (81,82).
During this phase, the parasite synthesizes proteins de novo and hijacks host cell lipids and proteins for
membrane biogenesis (83–86). We propose that the parasite brings its SSPM into hepatocytes to be

repurposed as the LSPMB (Figure 6). The observed microtubule disassembly over time may support tubulin
turnover and safes energy for intracellular development.
In addition to their structural roles, LSPMB filaments were occasionally found in contact with hemi-spindle
poles, suggesting that LSPMB contribute to spindle formation during schizogony. This connection, although
infrequent in fixed cells, was frequently observed during live imaging suggesting a transient connection. We
also detected expression of EB1, a microtubule plus-end binding protein previously believed to be restricted
to the sexual blood stage (64,65,87), during mosquito and liver stages. Its localization to LSPMB and hemi-
spindles suggests a broader role for EB1 in regulating microtubule dynamics throughout the parasite life
cycle.
To directly test the function of tubulin PTMs, we generated *P. berghei* mutants with targeted C-terminal
modifications of α1-tubulin. These included a glutamate-to-alanine substitution at a polyglutamylation site,
deletion of the tyrosination motif, and complete deletion of the C-terminal tail. Despite the mutations, PTM-
specific antibody staining remained detectable in the first two mutants, suggesting compensatory
mechanisms such as modification of adjacent residues or of β-tubulin. The tyrosination antibody used (anti-
Ty, T9028, Sigma-Aldrich) is a purified monoclonal antibody targeting C-terminal tyrosine residues on
tubulins, whereas the monoclonal GT335 antibody (AG-20B-0020-C100, AdipoGen Life Sciences)
recognizes the epitope formed at the branching point of polyglutamylation side chains (GEGE*EE, * marking
the modification site) and detects both short and long glutamate side chains. The polyclonal antibody (AG-
25B-0030-C050, AdipoGen Life Sciences) was raised against at least three glutamate residues and is
thereby specific to long polyglutamylation side chains, though both antibodies may cross-react with β-
tubulin or other proteins containing genetically encoded C-terminal modifications (66,88). The reduced
fluorescence signal observed (Figure 5) in mutant parasite microtubules suggests either antibody cross-
reactivity with β-tubulin or compensatory PTM relocation to adjacent α-tubulin residues. In the CtΔ mutant
(lacking the entire C-terminal tail), PTM relocation is abolished, and residual signal likely stems from β-
tubulin which was insufficient to sustain sporozoite motility and transmission. These possibilities warrant
further investigation. Previous studies have also demonstrated partial redundancy between α1- and α2-
tubulin modified (49,50,89). However, the complete C-terminal deletion mutant failed to produce salivary
gland sporozoites and could not establish liver stage infection. These results are consistent with findings
from Spreng et al. (2019), who showed that α1-tubulin mutants lacking the C-terminal region produced
defective sporozoites with altered microtubule numbers and impaired motility.
We also observed that the LSPMB relocates during liver stage development. Using expansion microscopy
on PMP1::GFP-expressing parasites (71), we found that the LSPMB localizes between the parasite plasma
membrane and the PVM. This differs from earlier reports of SSPM positioning (77,90–92) but both notions
are not mutually exclusive. It is possible that the LSPMB first localizes within the parasite but then re-
localizes at later stages or even re-localizes dynamically. Its localization between the PM and the PVM
positions it well for interaction with the host cell indicating an interaction with host cell structures such as
ER tubules or host microtubules (71,84,93,94). However, whether this re-localization plays a role in nutrient
acquisition or host-pathogen signaling remains to be explored.
Finally, during late schizogony and merozoite formation, we observed the reappearance of tyrosinated
tubulin. This may reflect the need for stable microtubules in newly formed merozoites, which must be
competent for host cell invasion.
In conclusion, our study reveals a previously unrecognized complexity in the microtubule biology
of *Plasmodium* liver stages. The identification of the LSPMB as a dynamic structure regulated by tubulin
PTMs and repurposed from the sporozoite cytoskeleton opens new avenues for understanding how the
parasite controls its intracellular architecture. These PTMs appear critical for parasite shape, replication,
and stage transition, and may represent valuable targets for therapeutic intervention. Future work should
aim to identify the enzymes responsible for these modifications and explore their potential as drug targets.
The behavior and function of microtubules are closely tied to their PTMs. We found that α1-tubulin
expressed during the liver stage carries residues predicted to undergo polyglutamylation and tyrosination.

Using PTM-specific antibodies, we confirmed that sporozoite microtubules are both tyrosinated and
polyglutamylated. Tyrosination is typically associated with stable microtubules and has been described in
highly motile structures such as cilia and spermatozoa. This is consistent with the sporozoite's need for
high motility gliding at approximately 1.5 µm/s as it migrates from the mosquito midgut to the salivary glands
and eventually traverses skin and vasculature to reach hepatocytes.

Methods

Ethics statement

All experimental procedures were performed in full compliance with the Swiss Animal Protection Act
(TSchG) and were approved by the Canton of Bern's Animal Experimentation Commission (Authorization,
BE118/22). Personnel involved in animal studies held the Federation of European Laboratory Animal
Science Associations (FELASA) accredited certifications. Female C57BL/6 and BALB/c mice of 6 to 10
weeks old and 20-30 g at infection were sourced from Janvier Labs (France) or bred in the animal facility
at the University of Bern's Institute of Cell Biology. Animals were maintained under standard housing
conditions with daily health monitoring by trained staff.

P. berghei lines used in this study and the generation of transgenic parasites

The *P. berghei* ANKA parasite line was utilized for both murine and cellular infection models. To generate
the α1-tubulin C-terminal mutant and eb1::eGFP parasite lines, the C-terminal coding and 3' untranslated
regions (UTRs) were cloned into the double homologous recombination vector pDHR, which contains the
3'UTR PbDHFR, the ef1α promoter region, the selectable markers hdhfr::yfcu, and additional 3'UTR
PbDHFR sequences. The pDHR vector backbone originated from the plasmid pG362 (kindly provided by
the Waters lab) (95).
First, the pG362 plasmid was digested with SacII and PstI to remove the AID, selectable marker, and GFP
sequences. The amplified product (linker-eGFP-3'UTR PbDHFR) from the pOB277 vector was then cloned
using T4 DNA ligase (NEB, M0202L). For subsequent cloning, restriction sites were added to the forward
(KpnI and ApaI) and reverse (EcoRI and SmaI) primers. The plasmid was then digested with EcoRI and
SmaI to clone the amplified product: ef1α promoter region-hdhfr::yfcu marker (a fusion gene of the positive
selection marker human dihydrofolate reductase (hdhfr) and the negative selection marker yeast cytosine
deaminase-uridyl phosphoribosyl transferase (yfcu) 3'UTR PbDHFR from the genomic DNA of the *P.*
berghei GIMO parasites (RMgm-687) parental line (96).
The complete α1-tubulin gene sequence (PBANKA_0417700), including 5' and 3' UTRs, was retrieved from
PlasmoDB (https://plasmodb.org). The linker-eGFP region of the pDHR vector was removed using ApaI
and NotI restriction enzymes. The 5'UTR of the α1-tubulin gene was then subcloned into the digested pDHR
using Gibson assembly. The resulting vector was digested with SmaI, and the PbHsp70 promoter and
mCherry regions from RMgm-687, along with the 3'UTR of the α1-tubulin gene, were subcloned into the
vector at the ApaI and SmaI restriction sites, respectively, using Gibson assembly.
Site-directed mutagenesis (QuickChange II XL Kit) was used to introduce the C-terminal modifications, with
primer sequences listed in Tables 2 and 3. The final plasmid was linearized with ScaI before being
transfected into the PbWT line (*P. berghei* ANKA line cl15cy1) (97). Similarly, the end-binding protein eB1
(PBANKA_0405600) with an eGFP tag was engineered using identical cloning strategies. The 5' and 3'
UTRs were retrieved from PlasmoDB (https://plasmodb.org) and subcloned into the pDHR vector at the
ApaI and SmaI restriction sites, respectively, using Gibson assembly. The final vector was linearized with
NdeI before being transfected into the Pb1868 line (constitutively expressing Hsp70-mCherry-ef1α-NLuc)
(98). Primer sequences are listed in Table 1.
Vector transfections were performed in schizont cultures as previously described (16,99). Briefly, *P. berghei*
schizonts were cultured and purified (16,99,100). Electroporation of schizonts from the respective parasite

lines was performed using the Amaxa® Human T Cell Nucleofector® system with the U33 program,
followed by intravenous injection into naïve mice. The construct was integrated by double-crossover
homologous recombination at the locus of interest. Pyrimethamine drug selection was administered via
drinking water, and parasitemia was monitored daily from day 7 post-infection until mice tested positive
(99).

Parasitaemia measurement and cloning of transgenic parasite lines

To measure the parasitemia ratio of infected mice, approximately 2 µL of blood from the tail vein of each
infected mouse was added to 500 µL of PBS and analyzed by flow cytometry using the NovoCyte (Agilent,
2023) as described (101). The data were analyzed using NovoExpress Flow Cytometry Analysis Software
(version 1.6.2, Build 230717.50551; Agilent Technologies).
To clone the different engineered transgenic parasite lines, mice were infected with the transgenic
parasites. Three days post-infection, parasitemia was assessed and when parasitemia reached at least
1%, blood was collected from the infected mouse via heart puncture following euthanasia as described
(16). The collected blood was passaged into a new naïve mouse. When parasitemia in the newly infected
mouse reached 1-2%, 2 µL of tail vein blood was collected and mixed with RPMI-1640. Maintaining low
parasitemia was crucial to prevent both double erythrocyte infection by *P. berghei* and gametocyte
formation. Single *P. berghei-* infected erythrocytes were then sorted into individual wells of 96-well plates
(Cellvis) containing 200 µL of RPMI-1640 supplemented with 10% FCS using a BD FACSDiscover S8 with
CellView. Supplementing the medium with 10% FCS allowed the sorted infected erythrocytes to recover
from the sorting process. Five mice were then intravenously infected with a single sorted infected
erythrocyte each. The mice underwent drug selection via drinking water, and parasitemia was monitored
daily from day 7 post-infection until parasites were detected. Clonality of the parasites in positive mice was
verified by integration PCR using primers listed in Tables 1, 2, and 3.

Cell culture

Human epithelial HeLa cells (European Cell Culture Collection) were maintained in Minimum Essential
Medium (MEM; 1-31F01-I BioConcept) supplemented with Earle's salts and 10% heat-inactivated fetal
bovine serum (FBS; Sigma-Aldrich), hereafter referred to as complete MEM (cMEM). Primary hepatocytes
were cultured in William's Medium E (WME) complete medium. All cells were incubated at 37°C under 5%
CO_2. For infection assays, 40,000 HeLa cells were seeded onto coverslips in 24-well plates and allowed to
adhere overnight. Cells were then infected with *P. berghei* sporozoites and maintained in cMEM containing
2.5 µg/mL amphotericin B (PAA Laboratories, P11-001) to prevent fungal contamination. Medium was
exchanged every 24 hours.

Primary hepatocyte isolation

The mice primary hepatocytes were isolated according to the method described (8). Briefly, the cells were
isolated from 12 to16 week old Balb/c or C57BL/6 mice using a two-step perfusion technique (8,83,102).
Mice were euthanized via CO_2 inhalation (1–2 minutes), and after securing the animal in a supine position,
a midline abdominal incision was made to expose the portal vein or inferior vena cava (IVC). A peristaltic
pump was pre-rinsed with 70% ethanol and HEPES buffer, then set to 50 for initial cannulation of the portal
vein (or IVC) using a Teflon-coated needle. Upon successful insertion, the flow was increased to maximum
(999), and the opposing vein was severed to allow perfusion with HEPES buffer for 5 minutes, during which
periodic clamping (10 seconds per minute) of the cut vein was performed. The tubing was subsequently
changed to a collagenase solution, and perfusion was held for 7 minutes with repeated clamping. The liver
was excised, transferred to a collagenase-containing well, and mechanically dissociated using filter tips and
gentle shaking with forceps. Cell suspensions were sequentially collected, pooled, and washed with

William's Medium E (WME) by centrifugation at 50g for 2 minutes. The pellet was resuspended in WME,
optionally passed through a 70 μm nylon strainer, and washed twice more. The final pellet was resuspended
in WME complete medium (WME supplemented with L-glutamine, penicillin-streptomycin, amphotericin B,
insulin-transferrin, and aprotinin), and cells were counted and seeded at densities appropriate for T25
flasks, GB dishes, or 24-well plates. After initial adhesion (1 hour), culture medium was supplemented or
exchanged after 2–4 hours.

Immunofluorescence assay (IFA) of fixed primary hepatocytes and HeLa cells

Immunofluorescence was carried out as previously described (56,86,103). Cells were seeded onto
coverslips in culture 24 well-plates and allowed to adhere overnight before infection. Infection was
maintained for 24 hours post-infection (hpi), after which cells were fixed using cold methanol at -20°C for
15 minutes and subsequently washed twice with Phosphate buffered saline (PBS). Blocking was performed
by incubating cells in 10% FCS in PBS for 20 minutes at room temperature to reduce non-specific binding.
Primary antibodies staining was carried out by incubating cells in 10% of FCS diluted in PBS for 1 hour at
room temperature, followed by three washes with PBS. Secondary antibodies incubation was performed in
10% of FCS as well for 1 hour at room temperature in the dark. The antibodies used are listed in the
enumerated session bellow. After three additional washes with PBS, nuclei were counterstained with DAPI.
Coverslips were then mounted onto slides with DAKO mounting medium and fluorescence imaging was
performed using either Nikon W1 or LIPSI and Crest V3 confocal microscope equipped with filters for DAPI
(4,6-Diamidin-2-phenylindol), AF488, AF594, and Cy5. Single-stain controls were carried out for every
staining to ensure that bleed-through was negligible.

Sporozoite isolation from mosquito salivary glands and haemolymph

Anopheles mosquitoes were fed on *Plasmodium*-infected mice exhibiting a gametocytaemia of 0.2 to 1%
and the sporozoite isolations were performed from day 18 post-infection as previously described (16,61).
For salivary gland sporozoites, mosquitoes were anesthetized with chloroform, sterilized in 70% ethanol,
and rinsed twice in PBS. The mosquitoes were then dissected, and salivary glands were extracted under a
binocular microscope in Iscove's Modified Dulbecco's Medium (IMDM) pooled in 50 μL of ice-cold IMDM
and mechanically homogenized using a pestle (Sigma-Aldrich, Z359947) powered with cordless motor
(Sigma-Aldrich, Z359971) as described (16,61). For hemolymph sporozoite extraction, glass capillaries
were heat-pulled to fine tips. The abdominal terminal segment of the mosquito was excised, and
hemolymph was flushed from the thorax by PBS perfusion via capillary insertion (75,104) . Hemolymphs
extracted were centrifuged (11,000 × g, 1 min), and pellets were resuspended in PBS or IMDM.
Sporozoite quantification was performed using a Neubauer hemacytometer. Midgut, hemolymph, or
salivary gland suspensions were mixed 1:1 with RPMI-1640 supplemented 10% FCS, and sporozoites
within four corner grids were counted. Total yields (sporozoites/mL and per mosquito) were calculated
based on dilution factors and dissection numbers.

Sporozoites IFA and gliding motility assay

To perform the sporozoite immunofluorescence assay (IFA), 12 mm cover glasses (0111530, Paul
Marienfeld, Germany) were coated with 0.1 mg/mL poly-D-lysine (A38904-01, Gibco, USA) (56,105) and
incubated overnight at room temperature. The following day, sporozoites were isolated from mosquito
midguts, hemolymph, or salivary glands as described above. The extracted sporozoites were suspended
in RPMI-1640 medium to a final volume of 200 μL in an Eppendorf tube and centrifuged at 1000 rpm for 3
min. The pellet was fixed by resuspension in 200 μL of cold methanol and incubated for 15 min at −20°C.
After fixation, sporozoites were washed twice with PBS (1000 rpm centrifugation) and transferred onto poly-
D-lysine-coated cover glasses in a 24-well plate (400024, Bioswisstec, Switzerland). The plate was

incubated for 30 min at room temperature to facilitate sporozoite adhesion before proceeding with either
the IFA or expansion microscopy protocol.
For the gliding motility assay, 5 µL of the sporozoite suspension in RPMI-1640 (supplemented with 10%
FCS) was added to 96-well glass-bottom plates (P96-1.5H-N, Cellvis, USA) (56,72,73). The plate was
centrifuged (1000 rpm, 1 min) and immediately live imaged using a Leica DM5500 widefield fluorescence
microscope.

Expansion microscopy

Expansion microscopy (PS-ExM) was performed as previously established (56–58). Infected cells (HeLa
cells, primary hepatocytes and sporozoites) were seeded on 13 mm coverslips in 24-well plates and fixed
with either 4% paraformaldehyde (PFA; F8775, Sigma) or cold methanol. The cells were anchored by
incubation in PBS containing 0.7% formaldehyde and 1% acrylamide at 37°C for 2 hours. Gelation was
then performed by adding 5 µL each of 10% tetramethylethylenediamine (TEMED; 17919, Thermo Fisher)
and 10% ammonium persulfate (APS) to 90 µL of monomer solution containing 19% sodium acrylate
(408220, Sigma), 10% acrylamide (A4058, Sigma), and 0.1% N, N'-methylenebisacrylamide (M1533,
Sigma). A 35 µL of monomer solution was dispensed onto parafilm in a humidity chamber, and coverslips
containing the anchored cells were inverted onto droplets. Following 5 minutes incubation on ice to improve
gel penetration, gels were polymerized at 37°C for 30 minutes. Gel-embedded samples were transferred
to denaturation buffer (200 mM SDS, 200 mM NaCl, 50 mM Tris-HCl, pH 9) in 6-well plates and agitated
(200 rpm) at room temperature for 15 minutes, followed by incubation in fresh buffer at 95°C for 1 hour with
gentle shaking. Gel expansion was performed in autoclaved MilliQ water (50 mL) for 1 hour with optional
buffer exchange. For immunostaining, gels were incubated with primary antibodies in 2% BSA for 2 hours
at 37°C, washed with PBS-Tween 0.1%, then labelled with secondary antibodies plus 1 µg/mL DAPI for
nuclear staining in 2% BSA for 2 hours at 37°C. Final gel expansion was conducted in MilliQ water
containing 0.2% propyl gallate (02370, Sigma) to prevent photobleaching. Expanded samples were
mounted on poly-D-lysine-coated (A38904, Gibco USA) glass-bottom dishes with cellular surfaces facing
downward and imaged with confocal microscopy.

Antibodies used for IFA and expansion microscopy

The primary antibodies used were: anti-UIS4 chicken (1:1000 IFA/1:500 expansion; Proteogenix), anti-UIS4
rabbit (1:1000 IFA/1:500 expansion; P. Sinnis, Baltimore), polyglutamylation IN105 rabbit (1:1000 IFA/1:500
expansion; AG-25B-0030-C050, AdipoGen Life Sciences), anti-GFP rabbit (1:1000 IFA/1:500 expansion;
SP3005P, Origene), anti-α-tubulin guinea pig (1:500 IFA/1:125 expansion; AA345, ABCD), anti-β-tubulin
guinea pig (1:500 IFA/1:125 expansion; AA344, ABCD), anti-tyrosinated tubulin mouse (1:1000 IFA/1:500
expansion; T9028, Sigma Aldrich), anti-polyglutamylation GT335 mouse (AG-20B-0020-C100, AdipoGen
Life Sciences), anti-γ-tubulin mouse (1:1000 IFA/1:500 expansion; T6557, Sigma Aldrich), and anti-GFP
mouse (1:1000 IFA/1:500 expansion; 814460001, Roche).
Secondary antibodies included anti-chicken IgY Alexa Fluor 594 (1:1000 IFA/1:500 expansion; A11042,
Invitrogen Molecular Probes), anti-rabbit IgG Alexa Fluor 594 (1:1000 IFA/1:500 expansion; A21207, Life
Technologies) and 488 (1:1000 IFA/1:500 expansion; A11034, Sigma Aldrich), anti-rabbit IgG ATTO 647N
(1:1000 expansion only; 40839, Sigma Aldrich), anti-guinea pig IgG Alexa Fluor 647 (1:1000 IFA/1:500
expansion; SAB4600033, Sigma Aldrich) and 594 (1:1000 IFA/1:500 expansion; A11076, Invitrogen
Molecular Probes), and anti-mouse IgG ATTO 647N (1:1000 expansion only; 50185, Sigma Aldrich) and
Alexa Fluor 488 (1:1000 IFA/1:500 expansion; A11001, Invitrogen Molecular Probes).

In vivo assay

To assess the infectivity of parasite mutant sporozoites to hepatocytes, *Anopheles stephensi* mosquitoes
infected with different parasite mutants were allowed to feed on anesthetized naïve mice as previously
described (Atchou et al., 2025). Three mice per parasite line and five mosquitoes per mouse were used.
The blood stage infection progression was monitored daily via flow cytometry, starting from the day of
infection until mice tested positive.

Oocyst size measurement

To quantify oocyst size, *Anopheles stephensi* mosquitoes were allowed to feed on anesthetized mice
infected with the parasite (Rogerio et al. 2006; Hegge et al., 2009). At 7- and 11-days post-feeding, ten
mosquitoes per time point were dissected, and their midguts were extracted and placed in 1× PBS. The
midguts were then transferred to a 21-well microscopy slide, covered with a coverslip, and imaged using a
Nikon Crest V3 confocal microscope (10× objective). The oocyst area was measured using Fiji software
(ImageJ). Each experiment was performed in triplicate for every parasite line.

Liver stage parasite size measurement

To quantify the parasite size during liver stage development, HeLa cells (40,000 cells/well, triplicate) were
seeded in 96-well plates and infected 24 hours later with 20,000 sporozoites per well. After a 2-hour
incubation at 37°C, infected cells from each well were washed and redistributed into eight wells across two
96 well plates (4 wells/plate) to optimize cell density for imaging (Pino et al., 2017; Atchou et al., 2025;
Sturm et al., 2006). Parasite development was analyzed at 6, 24, and 48 hpi using an InCell Analyzer 2000
with 10x objective. The image processing parameters such as kernel size, sensitivity, and size thresholds
were optimized in InCell Developer software to distinguish individual parasites from background signals.
Parasite growth was assessed based on size measurements, while survival rates were calculated as
percentages relative to 6 hpi baseline counts.

Live cell imaging of *Plasmodium* microtubule dynamism

HeLa cells were seeded in a 96-well plate and infected the following day with *P. berghei* salivary gland
sporozoites. After a 2-hour incubation at 37°C, cells were washed with MEM medium to remove uninfected
parasites and residual mosquito salivary gland materials. The infected cells were then reseeded in a dark
24-well glass-bottom plate (Cellvis) and maintained in fresh cMEM at 37°C with 5% CO_2 (16,106). Prior to
live imaging, cells were incubated for 1 hour in cMEM containing the SPY650-tubulin probe (1:1000 dilution;
SC503, Spirochrome, Switzerland). Images were acquired using a Nikon W1 LIPSI spinning disk
microscope with a 100X objective. Marked infected HeLa cells were imaged at defined time points to
generate time-lapse movies capturing tubulin dynamics and parasite development.

Image processing and data analysis

All the live cell imaging, expansion microscopy, and IFA images in this study were acquired with Nikon W1
LIPSI and Crest V3, both in spinning disk mode. The microscope images were processed and analyzed
using FIJI software, Huygens deconvolution Professional version (Huygens Remote Manager v3.8,
Scientific Volume Imaging, The Netherlands), or, when specified, the 3D and 4D IMARIS software (Imaris
10.2.0; June 18, 2024). The 3D/4D image analysis software was used to segment the parasite LSPMB,
hemi-spindle, and poles into either isospots or isosurfaces, as previously described (16,56). The counts,
volumes, and coefficients of the isospots or isosurfaces were retrieved in .csv files, which were then
imported into GraphPad Prism for statistical analysis. All data were analyzed using GraphPad Prism version
10.4.1 for macOS (GraphPad Software, San Diego, California, USA). Statistical comparisons were
performed using one-way ANOVA with Dunn's multiple comparisons test (*$p \leq 0.05$, $p \leq 0.01$, *$p \leq 0.001$).
All experiments in the study were carried out in triplicate.

Acknowledgments

The authors gratefully acknowledge the Microscopy Imaging Center (MIC) at the University of Bern, Switzerland, for their technical support and access to microscopy equipment. K.A. acknowledges the Swiss Confederation for the award of a Government Excellence Scholarship. *This work was funded by the Swiss National Science Foundation (SNSF) (grant number 310030_212795) and an Multidisciplinary Center for Infectious Diseases (MCID), University of Bern, Switzerland, grant to Volker Heussler (MA-09).*

References

1. Wambani J, Okoth P. Impact of Malaria Diagnostic Technologies on the Disease Burden in the Sub-Saharan Africa. 2022; Available from: https://onlinelibrary.wiley.com/doi/10.1155/2022/7324281

2. Poespoprodjo JR, Douglas NM, Ansong D, Kho S, Anstey NM. Malaria. Lancet [Internet]. 2023 Dec 16 [cited 2025 Mar 10];402(10419):2328–45. Available from: https://pubmed.ncbi.nlm.nih.gov/37924827/

3. Varo R, Chaccour C, Bassat Q. Update on malaria. Med Clin (Barc) [Internet]. 2020 Nov 13 [cited 2025 Mar 10];155(9):395–402. Available from: https://pubmed.ncbi.nlm.nih.gov/32620355/

4. Venkatesan P. WHO world malaria report 2024. Lancet Microbe [Internet]. 2025 Apr 1 [cited 2025 Jun 11];6(4):101073. Available from: https://www.thelancet.com/action/showFullText?pii=S2666524725000011

5. Tavares J, Formaglio P, Thiberge S, Mordelet E, Van Rooijen N, Medvinsky A, et al. Role of host cell traversal by the malaria sporozoite during liver infection. Journal of Experimental Medicine [Internet]. 2013 May 6 [cited 2025 May 18];210(5):905–15. Available from: www.jem.org/cgi/doi/10.1084/jem.20121130

6. Bindschedler A, Wacker R, Egli J, Eickel N, Schmuckli-Maurer J, Franke-Fayard BM, et al. Plasmodium berghei sporozoites in nonreplicative vacuole are eliminated by a PI3P-mediated autophagy-independent pathway. Cell Microbiol [Internet]. 2021 Jan 1 [cited 2025 Mar 10];23(1). Available from: https://pubmed.ncbi.nlm.nih.gov/32979009/

7. Marques-da-Silva C, Schmidt-Silva C, Kurup SP. Hepatocytes and the art of killing Plasmodium softly. Trends Parasitol [Internet]. 2024 Jun 1 [cited 2025 May 18];40(6):466–76. Available from: https://www.cell.com/action/showFullText?pii=S1471492224000862

8. Prado M, Eickel N, De Niz M, Heitmann A, Agop-Nersesian C, Wacker R, et al. Long-term live imaging reveals cytosolic immune responses of host hepatocytes against Plasmodium infection and parasite escape mechanisms. Autophagy [Internet]. 2015 [cited 2023 Jan 19];11(9):1561–79. Available from: https://pubmed.ncbi.nlm.nih.gov/26208778/

9. Zanghi G, Vaughan AM. Plasmodium vivax pre-erythrocytic stages and the latent hypnozoite. Parasitol Int. 2021 Dec 1;85:102447.

10. Prudêncio M, Rodriguez A, Mota MM. The silent path to thousands of merozoites: the Plasmodium liver stage. Nature Reviews Microbiology 2006 4:11 [Internet]. 2006 Nov [cited 2025 May 18];4(11):849–56. Available from: https://www.nature.com/articles/nrmicro1529

635   11.   Roques M, Bindschedler A, Beyeler R, Heussler VT. Same, same but different: Exploring
636   Plasmodium cell division during liver stage development. PLoS Pathog [Internet]. 2023 Mar 1
637   [cited 2025 Mar 12];19(3):e1011210. Available from:
638   https://pmc.ncbi.nlm.nih.gov/articles/PMC10062574/

639   12.   Shears MJ, Sekhar Nirujogi R, Swearingen KE, Renuse S, Mishra S, Jaipal Reddy P, et al.
640   Proteomic Analysis of Plasmodium Merosomes: The Link between Liver and Blood Stages in
641   Malaria. J Proteome Res [Internet]. 2019 Sep 6 [cited 2025 May 18];18(9):3404–18.
642   Available from: https://pubs.acs.org/doi/full/10.1021/acs.jproteome.9b00324

643   13.   De Niz M, Heussler VT. Rodent malaria models: insights into human disease and parasite
644   biology. Curr Opin Microbiol [Internet]. 2018 Dec 1 [cited 2025 May 18];46:93–101. Available
645   from: https://pubmed.ncbi.nlm.nih.gov/30317152/

646   14.   Sturm A, Amino R, Van De Sand C, Regen T, Retzlaff S, Rennenberg A, et al. Manipulation of
647   host hepatocytes by the malaria parasite for delivery into liver sinusoids. Science (1979)
648   [Internet]. 2006 Sep 1 [cited 2024 May 16];313(5791):1287–90. Available from:
649   https://www.science.org/doi/10.1126/science.1129720

650   15.   Heussler V, Spielmann T, Frischknecht F, Gilberger T. Plasmodium. Molecular Parasitology:
651   Protozoan Parasites and their Molecules [Internet]. 2016 Jan 1 [cited 2025 Mar 10];241–84.
652   Available from: https://link.springer.com/chapter/10.1007/978-3-7091-1416-2_9

653   16.   Atchou K, Caldelari R, Roques M, Schmuckli-Maurer J, Beyeler R, Heussler V. Expanding
654   the fluorescent toolkit: Blue fluorescent protein-expressing Plasmodium berghei for enhanced
655   multiplex microscopy. Portugaliza HP, editor. PLoS One [Internet]. 2025 Mar 3 [cited 2025
656   Mar 10];20(3):e0308055. Available from: https://pubmed.ncbi.nlm.nih.gov/40029851/

657   17.   Stanway RR, Bushell E, Chiappino-Pepe A, Roques M, Sanderson T, Franke-Fayard B, et al.
658   Genome-Scale Identification of Essential Metabolic Processes for Targeting the Plasmodium
659   Liver Stage. Cell [Internet]. 2019 Nov 14 [cited 2025 Mar 10];179(5):1112-1128.e26. Available
660   from: https://pubmed.ncbi.nlm.nih.gov/31730853/

661   18.   Meibalan E, Marti M. Biology of Malaria Transmission. Cold Spring Harb Perspect Med
662   [Internet]. 2017 Mar 1 [cited 2025 Mar 10];7(3). Available from:
663   https://pubmed.ncbi.nlm.nih.gov/27836912/

664   19.   Frischknecht F, Baldacci P, Martin B, Zimmer C, Thiberge S, Olivo-Marin JC, et al. Imaging
665   movement of malaria parasites during transmission by Anopheles mosquitoes. Cell Microbiol
666   [Internet]. 2004 Jul 1 [cited 2025 May 18];6(7):687–94. Available from:
667   https://onlinelibrary.wiley.com/doi/full/10.1111/j.1462-5822.2004.00395.x

668   20.   Vanderberg JP. Development of infectivity by the Plasmodium berghei sporozoite. Journal of
669   Parasitology. 1975;61(1):43–50.

670   21.   Vanderberg jerome P. Studies on the Motility of Plasmodium Sporozoites*. J Protozool
671   [Internet]. 1974 Oct 1 [cited 2025 May 18];21(4):527–37. Available from:
672   https://onlinelibrary.wiley.com/doi/full/10.1111/j.1550-7408.1974.tb03693.x

673   22.   Frischknecht F, Matuschewski K. Plasmodium Sporozoite Biology. Cold Spring Harb
674   Perspect Med [Internet]. 2017 May 1 [cited 2025 May 18];7(5):a025478. Available from:
675   https://pmc.ncbi.nlm.nih.gov/articles/PMC5411682/

676   23.   Mitchison T, Kirschner M. Dynamic instability of microtubule growth. Nature 1984 312:5991
677   [Internet]. 1984 [cited 2025 Mar 11];312(5991):237–42. Available from:
678   https://www.nature.com/articles/312237a0

679   24.   Janke C, Magiera MM. The tubulin code and its role in controlling microtubule properties and
680   functions. Nat Rev Mol Cell Biol. 2020 Jun 1;21(6):307–26.

681   25.   Nogales E. Structural insights into microtubule function. Annu Rev Biophys Biomol Struct
682   [Internet]. 2001 Jun 1 [cited 2025 Mar 11];30(Volume 30, 2001):397–420. Available from:
683   https://www.annualreviews.org/content/journals/10.1146/annurev.biophys.30.1.397

684   26.   Roll-Mecak A. The Tubulin Code in Microtubule Dynamics and Information Encoding. Dev
685   Cell. 2020 Jul 6;54(1):7–20.

686   27.   Bodakuntla S, Jijumon AS, Villablanca C, Gonzalez-Billault C, Janke C. Microtubule-
687   Associated Proteins: Structuring the Cytoskeleton. Trends Cell Biol. 2019 Oct 1;29(10):804–
688   19.

689   28.   Lee G. Non-motor microtubule-associated proteins. Curr Opin Cell Biol. 1993;5(1):88–91.

690   29.   Peng N, Nakamura F. Microtubule-associated proteins and enzymes modifying tubulin.
691   Cytoskeleton [Internet]. 2023 Mar 1 [cited 2025 Mar 11];80(3–4):60–76. Available from:
692   https://onlinelibrary.wiley.com/doi/full/10.1002/cm.21748

693   30.   Peng N, Nakamura F. Microtubule-associated proteins and enzymes modifying tubulin.
694   Cytoskeleton (Hoboken) [Internet]. 2023 Mar 1 [cited 2025 Mar 11];80(3–4):60–76. Available
695   from: https://pubmed.ncbi.nlm.nih.gov/36798013/

696   31.   Fennell BJ, Al-shatr ZA, Bell A. Isotype expression, post-translational modification and stage-
697   dependent production of tubulins in erythrocytic Plasmodium falciparum. Int J Parasitol. 2008
698   Apr 1;38(5):527–39.

699   32.   Rawlings DJ, Fujioka H, Fried M, Keister DB, Aikawa M, Kaslow DC. α-Tubulin II is a male-
700   specific protein in Plasmodium falciparum. Mol Biochem Parasitol. 1992 Dec 1;56(2):239–50.

701   33.   Arregui C, Busciglio J, Caceres A, Barra HS. Tyrosinated and detyrosinated microtubules in
702   axonal processes of cerebellar macroneurons grown in culture. J Neurosci Res [Internet].
703   1991 [cited 2025 Mar 11];28(2):171–81. Available from:
704   https://pubmed.ncbi.nlm.nih.gov/1674546/

705   34.   Webster DR, Gundersen GG, Bulinski JC, Borisy GG. Differential turnover of tyrosinated and
706   detyrosinated microtubules. Proc Natl Acad Sci U S A [Internet]. 1987 [cited 2025 Mar
707   11];84(24):9040–4. Available from: https://pubmed.ncbi.nlm.nih.gov/3321065/

708   35.   Sanyal C, Pietsch N, Ramirez Rios S, Peris L, Carrier L, Moutin MJ. The detyrosination/re-
709   tyrosination cycle of tubulin and its role and dysfunction in neurons and cardiomyocytes.
710   Semin Cell Dev Biol [Internet]. 2023 Mar 15 [cited 2025 Mar 11];137:46–62. Available from:
711   https://pubmed.ncbi.nlm.nih.gov/34924330/

712   36.   Girão H, Macário-Monteiro J, Figueiredo AC, Silva E Sousa R, Doria E, Demidov V, et al. α-
713   tubulin detyrosination fine-tunes kinetochore-microtubule attachments. Nat Commun
714   [Internet]. 2024 Dec 1 [cited 2025 Mar 11];15(1):9720. Available from:
715   https://pubmed.ncbi.nlm.nih.gov/39521805/

37. Gagnon C, White D, Cosson J, Huitorel P, Eddé B, Desbruyères E, et al. The polyglutamylated lateral chain of alpha-tubulin plays a key role in flagellar motility. J Cell Sci [Internet]. 1996 Jun 1 [cited 2025 Mar 11];109(6):1545–53. Available from: https://dx.doi.org/10.1242/jcs.109.6.1545

38. Edde B, Rossier J, Le Caer JP, Prome JC, Desbruyeres E, Gros F, et al. Polyglutamylated alpha-tubulin can enter the tyrosination/detyrosination cycle. Biochemistry [Internet]. 1992 Feb 1 [cited 2025 Mar 11];31(2):403–10. Available from: https://pubmed.ncbi.nlm.nih.gov/1370628/

39. Ruse CI, Chin HG, Pradhan S. Polyglutamylation: biology and analysis. Amino Acids [Internet]. 2022 Apr 1 [cited 2025 Mar 11];54(4):529–42. Available from: https://pubmed.ncbi.nlm.nih.gov/35357568/

40. Zheng P, Obara CJ, Szczesna E, Nixon-Abell J, Mahalingan KK, Roll-Mecak A, et al. ER proteins decipher the tubulin code to regulate organelle distribution. Nature [Internet]. 2022 Jan 6 [cited 2025 Mar 11];601(7891):132–8. Available from: https://pubmed.ncbi.nlm.nih.gov/34912111/

41. Neff NF, Thomas JH, Grisafi P, Botstein D. Isolation of the beta-tubulin gene from yeast and demonstration of its essential function in vivo. Cell [Internet]. 1983 [cited 2025 Mar 11];33(1):211–9. Available from: https://pubmed.ncbi.nlm.nih.gov/6380751/

42. Baum P, Thorner J, Honig L. Identification of tubulin from the yeast Saccharomyces cerevisiae. Proc Natl Acad Sci U S A. 1978;75(10):4962–6.

43. Hirst WG, Fachet D, Kuropka B, Weise C, Saliba KJ, Reber S. Purification of functional Plasmodium falciparum tubulin allows for the identification of parasite-specific microtubule inhibitors. Current Biology. 2022 Feb 28;32(4):919-926.e6.

44. Holloway SP, Gerousis M, Delves CJ, Sims PFG, Scaife JG, Hyde JE. The tubulin genes of the human malaria parasite Plasmodium falciparum, their chromosomal location and sequence analysis of the α-tubulin II gene. Mol Biochem Parasitol. 1990 Dec 1;43(2):257–70.

45. Fennell BJ, Naughton JA, Dempsey E, Bell A. Cellular and molecular actions of dinitroaniline and phosphorothioamidate herbicides on Plasmodium falciparum: tubulin as a specific antimalarial target. Mol Biochem Parasitol [Internet]. 2006 Feb [cited 2025 Mar 11];145(2):226–38. Available from: https://pubmed.ncbi.nlm.nih.gov/16406111/

46. Dow GS, Armson A, Boddy MR, Itenge T, McCarthy D, Parkin JE, et al. Plasmodium: assessment of the antimalarial potential of trifluralin and related compounds using a rat model of malaria, Rattus norvegicus. Exp Parasitol [Internet]. 2002 Mar 1 [cited 2025 Mar 11];100(3):155–60. Available from: https://pubmed.ncbi.nlm.nih.gov/12173400/

47. Dempsey E, Prudêncio M, Fennell BJ, Gomes-Santos CS, Barlow JW, Bell A. Antimitotic herbicides bind to an unidentified site on malarial parasite tubulin and block development of liver-stage Plasmodium parasites. Mol Biochem Parasitol [Internet]. 2013 Apr [cited 2025 Mar 11];188(2):116–27. Available from: https://pubmed.ncbi.nlm.nih.gov/23523992/

48. Kooij TWA, Franke-Fayard B, Renz J, Kroeze H, Van Dooren MW, Ramesar J, et al. Plasmodium berghei alpha-tubulin II: a role in both male gamete formation and asexual blood stages. Mol Biochem Parasitol [Internet]. 2005 Nov [cited 2025 Mar 11];144(1):16–26. Available from: https://pubmed.ncbi.nlm.nih.gov/16115694/

49.　Holloway SP, Sims PFG, Delves CJ, Scaife JG, Hyde JE. Isolation of alpha-tubulin genes
from the human malaria parasite, Plasmodium falciparum: sequence analysis of alpha-
tubulin. Mol Microbiol [Internet]. 1989 [cited 2025 Mar 11];3(11):1501–10. Available from:
https://pubmed.ncbi.nlm.nih.gov/2693901/

50.　Spreng B, Fleckenstein H, Kübler P, Di Biagio C, Benz M, Patra P, et al. Microtubule number
and length determine cellular shape and function in Plasmodium . EMBO J [Internet]. 2019
Aug 24 [cited 2025 Mar 11];38(15). Available from:
https://www.embopress.org/doi/10.15252/embj.2018100984

51.　Zhang G, Niu G, Hooker–Romera D, Shabani S, Ramelow J, Wang X, et al. Targeting
plasmodium α-tubulin-1 to block malaria transmission to mosquitoes. Front Cell Infect
Microbiol [Internet]. 2023 [cited 2025 Mar 11];13:1132647. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC10064449/

52.　Valenstein ML, Roll-Mecak A. Graded Control of Microtubule Severing by Tubulin
Glutamylation. Cell [Internet]. 2016 Feb 25 [cited 2025 Mar 11];164(5):911–21. Available
from: https://www.cell.com/action/showFullText?pii=S0092867416000593

53.　Bonnet C, Boucher D, Lazereg S, Pedrotti B, Islam K, Denoulet P, et al. Differential Binding
Regulation of Microtubule-associated Proteins MAP1A, MAP1B, and MAP2 by Tubulin
Polyglutamylation. Journal of Biological Chemistry [Internet]. 2001 Apr 20 [cited 2025 Mar
11];276(16):12839–48. Available from:
https://www.jbc.org/action/showFullText?pii=S0021925819343935

54.　Bertiaux E, Balestra AC, Bournonville L, Louvel V, Maco B, Soldati-Favre D, et al. Expansion
microscopy provides new insights into the cytoskeleton of malaria parasites including the
conservation of a conoid. PLoS Biol [Internet]. 2021 Mar 11 [cited 2025 Mar 11];19(3).
Available from: https://pubmed.ncbi.nlm.nih.gov/33705377/

55.　Hentzschel F, Binder AM, Dorner LP, Herzel L, Nuglisch F, Sema M, et al. Microtubule inner
proteins of Plasmodium are essential for transmission of malaria parasites. Proceedings of
the National Academy of Sciences [Internet]. 2025 Feb 11 [cited 2025 Mar
12];122(6):e2421737122. Available from:
https://www.pnas.org/doi/abs/10.1073/pnas.2421737122

56.　Atchou K, Berger BM, Heussler V, Ochsenreiter T. Pre-gelation staining expansion
microscopy for visualisation of the Plasmodium liver stage. J Cell Sci [Internet]. 2023 [cited
2025 Mar 11];136(22):jcs261377. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC10729816/

57.　Liffner B, Absalon S. Expansion microscopy of apicomplexan parasites. Mol Microbiol
[Internet]. 2024 Apr 1 [cited 2025 Mar 11];121(4):619–35. Available from:
https://pubmed.ncbi.nlm.nih.gov/37571814/

58.　Liffner B, Diaz AKC, Blauwkamp J, Anaguano D, Frolich S, Muralidharan V, et al. Atlas of
Plasmodium falciparum intraerythrocytic development using expansion microscopy. Elife
[Internet]. 2023 Dec 18 [cited 2025 Mar 11];12. Available from:
https://pubmed.ncbi.nlm.nih.gov/38108809/

59.   Ferreira JL, Pražák V, Vasishtan D, Siggel M, Hentzschel F, Binder AM, et al. Variable
799   microtubule architecture in the malaria parasite. Nat Commun [Internet]. 2023 Dec 1 [cited
800   2025 Mar 12];14(1):1216. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC9984467/

801   60.   Aguayo-Ortiz R, Dominguez L. Unveiling the Possible Oryzalin-Binding Site in the α-Tubulin
802   of Toxoplasma gondii. ACS Omega [Internet]. 2022 Jun 7 [cited 2025 Mar 12];7(22):18434–
803   42. Available from: https://pubs.acs.org/doi/full/10.1021/acsomega.2c00729

804   61.   Beyeler R, Jordan M, Dorner L, He B, Cyrklaff M, Roques M, et al. Putative prefoldin
805   complex subunit 5 of Plasmodium berghei is crucial for microtubule formation and parasite
806   development in the mosquito. Mol Microbiol [Internet]. 2024 Mar 1 [cited 2025 Mar
807   12];121(3):481–96. Available from: https://pubmed.ncbi.nlm.nih.gov/38009402/

808   62.   Roques M, Stanway RR, Rea EI, Markus R, Brady D, Holder AA, et al. Plasmodium centrin
809   PbCEN-4 localizes to the putative MTOC and is dispensable for malaria parasite
810   proliferation. Biol Open [Internet]. 2019 Jan 1 [cited 2025 Mar 12];8(1):bio036822. Available
811   from: https://pmc.ncbi.nlm.nih.gov/articles/PMC6361220/

812   63.   Zeeshan M, Shilliday F, Liu T, Abel S, Mourier T, Ferguson DJP, et al. Plasmodium kinesin-8X
813   associates with mitotic spindles and is essential for oocyst development during parasite
814   proliferation and transmission. PLoS Pathog [Internet]. 2019 [cited 2025 Mar
815   12];15(10):e1008048. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC6786531/

816   64.   Yang S, Cai M, Huang J, Zhang S, Mo X, Jiang K, et al. EB1 decoration of microtubule lattice
817   facilitates spindle-kinetochore lateral attachment in Plasmodium male gametogenesis.
818   Nature Communications 2023 14:1 [Internet]. 2023 May 19 [cited 2025 Mar 12];14(1):1–20.
819   Available from: https://www.nature.com/articles/s41467-023-38516-3

820   65.   Mauer S, Camargo N, Abatiyow BA, Gargaro OR, Kappe SHI, Kumar S. Plasmodium
821   microtubule-binding protein EB1 is critical for partitioning of nuclei in male gametogenesis.
822   mBio [Internet]. 2023 Aug 31 [cited 2025 Mar 12];14(4):e00822-23. Available from:
823   https://pmc.ncbi.nlm.nih.gov/articles/PMC10470552/

824   66.   Magiera MM, Janke C. Investigating Tubulin Posttranslational Modifications with Specific
825   Antibodies. Methods Cell Biol. 2013 Jan 1;115:247–67.

826   67.   Oakley BR. gamma-Tubulin. Curr Top Dev Biol [Internet]. 2000 [cited 2025 Mar 11];49:27–54.
827   Available from: https://pubmed.ncbi.nlm.nih.gov/11005013/

828   68.   Maessen S, Wesseling JG, Smits MA, Konings RNH, Schoenmakers JGG. The γ-tubulin
829   gene of the malaria parasite Plasmodium falciparum. Mol Biochem Parasitol. 1993 Jul
830   1;60(1):27–35.

831   69.   Haase R, Puthenpurackal A, Maco B, Guérin A, Soldati-Favre D. γ-tubulin complex controls
832   the nucleation of tubulin-based structures in Apicomplexa. Mol Biol Cell [Internet]. 2024 Sep
833   1 [cited 2025 Jun 21];35(9):ar121. Available from:
834   https://pmc.ncbi.nlm.nih.gov/articles/PMC11449391/

835   70.   Barik S, Andrews J. Host–Parasite Interactions in Toxoplasma gondii-Infected Cells: Roles of
836   Mitochondria, Microtubules, and the Parasitophorous Vacuole. Int J Mol Sci [Internet]. 2024
837   Dec 1 [cited 2025 Jun 21];25(24):13459. Available from:
838   https://pmc.ncbi.nlm.nih.gov/articles/PMC11677533/

839   71.   Burda PC, Caldelari R, Heussler VT. Manipulation of the Host Cell Membrane during
840   Plasmodium Liver Stage Egress. mBio [Internet]. 2017 Mar 1 [cited 2025 Mar 13];8(2).
841   Available from: https://pubmed.ncbi.nlm.nih.gov/28400525/

842   72.   Hopp CS, Chiou K, Ragheb DRT, Salman AM, Khan SM, Liu AJ, et al. Longitudinal analysis
843   of plasmodium sporozoite motility in the dermis reveals component of blood vessel
844   recognition. Elife. 2015 Aug 13;4(AUGUST2015).

845   73.   Beyer K, Kracht S, Kehrer J, Singer M, Klug D, Frischknecht F. Limited Plasmodium
846   sporozoite gliding motility in the absence of TRAP family adhesins. Malar J [Internet]. 2021
847   Dec 1 [cited 2023 Apr 12];20(1):1–12. Available from:
848   https://malariajournal.biomedcentral.com/articles/10.1186/s12936-021-03960-3

849   74.   Klug D, Frischknecht F. Motility precedes egress of malaria parasites from oocysts. Elife.
850   2017 Jan 24;6.

851   75.   Braumann F, Klug D, Kehrer J, Song G, Feng J, Springer TA, et al. Conformational change of
852   Plasmodium TRAP is essential for sporozoite migration and transmission . EMBO Rep
853   [Internet]. 2023 Jul 5 [cited 2025 May 13];24(7). Available from:
854   https://www.embopress.org/doi/10.15252/embr.202357064

855   76.   Douglas RG, Reinig M, Neale M, Frischknecht F. Screening for potential prophylactics
856   targeting sporozoite motility through the skin. Malar J [Internet]. 2018 Aug 31 [cited 2025 May
857   13];17(1). Available from: https://pubmed.ncbi.nlm.nih.gov/30170589/

858   77.   Meis JFGM, Verhave JP, Jap PHK, Meuwissen JHET. Transformation of sporozoites of
859   Plasmodium berghei into exoerythrocytic forms in the liver of its mammalian host. Cell Tissue
860   Res [Internet]. 1985 Aug [cited 2025 Mar 13];241(2):353–60. Available from:
861   https://link.springer.com/article/10.1007/BF00217180

862   78.   Short B. Polyglutamylation makes the cut. J Cell Biol [Internet]. 2010 Jun 14 [cited 2025 Mar
863   12];189(6):920. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC2886350/

864   79.   Genova M, Grycova L, Puttrich V, Magiera MM, Lansky Z, Janke C, et al. Tubulin
865   polyglutamylation differentially regulates microtubule-interacting proteins. EMBO J [Internet].
866   2023 Mar [cited 2025 Mar 12];42(5):e112101. Available from:
867   https://pmc.ncbi.nlm.nih.gov/articles/PMC9975938/

868   80.   Pathak NH, Drummond IA. Polyglutamylation and the fleer gene. Methods Cell Biol.
869   2009;94:317–32.

870   81.   Jayabalasingham B, Bano N, Coppens I. Metamorphosis of the malaria parasite in the liver is
871   associated with organelle clearance. Cell Res [Internet]. 2010 Sep [cited 2025 Mar
872   12];20(9):1043. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC4137911/

873   82.   Graewe S, Stanway RR, Rennenberg A, Heussler VT. Chronicle of a death foretold:
874   Plasmodium liver stage parasites decide on the fate of the host cell. FEMS Microbiol Rev
875   [Internet]. 2012 Jan 1 [cited 2025 Mar 12];36(1):111–30. Available from:
876   https://dx.doi.org/10.1111/j.1574-6976.2011.00297.x

877   83.   de Niz M, Caldelari R, Kaiser G, Zuber B, Heo W Do, Heussler VT, et al. Hijacking of the host
878   cell Golgi by Plasmodium berghei liver stage parasites. J Cell Sci [Internet]. 2021 May 1

[cited 2025 Mar 12];134(10):jcs252213. Available from:
https://pmc.ncbi.nlm.nih.gov/articles/PMC8186485/

84.   O'neal AJ, Butler LR, Rolandelli A, Gilk SD, Pedra JHF. Lipid hijacking: A unifying theme in
vector-borne diseases. Elife. 2020 Oct 1;9:1–31.

85.   Fraser M, Curtis B, Phillips P, Yates PA, Lam KS, Netzel O, et al. Harnessing cholesterol
uptake of malaria parasites for therapeutic applications. EMBO Mol Med [Internet]. 2024 Jul
15 [cited 2025 Mar 12];16(7):1515–32. Available from:
https://www.embopress.org/doi/10.1038/s44321-024-00087-1

86.   Schmuckli-Maurer J, Bindschedler AF, Wacker R, Würgler OM, Rehmann R, Lehmberg T, et
al. Plasmodium berghei liver stage parasites exploit host GABARAP proteins for TFEB
activation. Communications Biology 2024 7:1 [Internet]. 2024 Nov 21 [cited 2025 Mar
12];7(1):1–13. Available from: https://www.nature.com/articles/s42003-024-07242-x

87.   Zeeshan M, Rea E, Abel S, Vukušić K, Markus R, Brady D, et al. Plasmodium ARK2 and
EB1 drive unconventional spindle dynamics, during chromosome segregation in sexual
transmission stages. Nature Communications 2023 14:1 [Internet]. 2023 Sep 13 [cited 2025
Mar 12];14(1):1–19. Available from: https://www.nature.com/articles/s41467-023-41395-3

88.   Van Dijk J, Miro J, Strub JM, Lacroix B, Van Dorsselaer A, Edde B, et al. Polyglutamylation Is
a Post-translational Modification with a Broad Range of Substrates. Journal of Biological
Chemistry. 2008 Feb 15;283(7):3915–22.

89.   Morrissette N, Abbaali I, Ramakrishnan C, Hehl AB. The Tubulin Superfamily in
Apicomplexan Parasites. Microorganisms [Internet]. 2023 Mar 1 [cited 2025 Mar 11];11(3).
Available from: https://pubmed.ncbi.nlm.nih.gov/36985278/

90.   Kappe SHI, Kaiser K, Matuschewski K. The Plasmodium sporozoite journey: a rite of
passage. Trends Parasitol. 2003 Mar 1;19(3):135–43.

91.   Ferreira JL, Heincke D, Wichers JS, Liffner B, Wilson DW, Gilberger TW. The Dynamic Roles
of the Inner Membrane Complex in the Multiple Stages of the Malaria Parasite. Front Cell
Infect Microbiol. 2021 Jan 8;10:611801.

92.   Harding CR, Meissner M. The inner membrane complex through development of Toxoplasma
gondii and Plasmodium. Cell Microbiol [Internet]. 2014 [cited 2025 Mar 13];16(5):632.
Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC4286798/

93.   Burda PC, Roelli MA, Schaffner M, Khan SM, Janse CJ, Heussler VT. A Plasmodium
Phospholipase Is Involved in Disruption of the Liver Stage Parasitophorous Vacuole
Membrane. PLoS Pathog [Internet]. 2015 Mar 1 [cited 2025 Mar 13];11(3):e1004760.
Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC4364735/

94.   Kaiser G, De Niz M, Zuber B, Burda PC, Kornmann B, Heussler VT, et al. High resolution
microscopy reveals an unusual architecture of the Plasmodium berghei endoplasmic
reticulum. Mol Microbiol [Internet]. 2016 Dec 1 [cited 2025 Mar 13];102(5):775–91. Available
from: https://pubmed.ncbi.nlm.nih.gov/27566438/

95.   Philip N, Waters AP. Conditional degradation of plasmodium calcineurin reveals functions in
parasite colonization of both host and vector. Cell Host Microbe [Internet]. 2015 Jul 8 [cited

2025 Jun 21];18(1):122–31. Available from:
https://www.cell.com/action/showFullText?pii=S1931312815002504

96.   Lin J wen, Annoura T, Sajid M, Chevalley-Maurel S, Ramesar J, Klop O, et al. A novel "Gene Insertion/Marker Out" (GIMO) method for transgene expression and gene complementation in rodent malaria parasites. PLoS One. 2011 Dec 27;6(12).

97.   Janse CJ, Ramesar J, Waters AP. High-efficiency transfection and drug selection of genetically transformed blood stages of the rodent malaria parasite Plasmodium berghei. Nature Protocols 2006 1:1 [Internet]. 2006 Jun 29 [cited 2025 Jun 21];1(1):346–56. Available from: https://www.nature.com/articles/nprot.2006.53

98.   De Niz M, Stanway RR, Wacker R, Keller D, Heussler VT. An ultrasensitive NanoLuc-based luminescence system for monitoring Plasmodium berghei throughout its life cycle. Malaria Journal 2016 15:1 [Internet]. 2016 Apr 21 [cited 2025 Jun 20];15(1):1–24. Available from: https://malariajournal.biomedcentral.com/articles/10.1186/s12936-016-1291-9

99.   Salman AM, Mogollon CM, Lin JW, Van Pul FJA, Janse CJ, Khan SM. Generation of transgenic rodent malaria parasites expressing human malaria parasite proteins. Methods in Molecular Biology. 2015;1325:257–86.

100.   Govindarajalu G, Rizvi Z, Kumar D, Sijwali PS. Lyse-Reseal Erythrocytes for Transfection of Plasmodium falciparum. Sci Rep [Internet]. 2019 Dec 1 [cited 2024 Jan 23];9(1). Available from: /pmc/articles/PMC6934678/

101.   Pino P, Caldelari R, Mukherjee B, Vahokoski J, Klages N, Maco B, et al. A multistage antimalarial targets the plasmepsins IX and X essential for invasion and egress. Science (1979) [Internet]. 2017 Oct 27 [cited 2025 Jun 20];358(6362):522–8. Available from: https://www.science.org/doi/10.1126/science.aaf8675

102.   Kaur I, Vasudevan A, Rawal P, Tripathi DM, Ramakrishna S, Kaur S, et al. Primary Hepatocyte Isolation and Cultures: Technical Aspects, Challenges and Advancements. Bioengineering [Internet]. 2023 Feb 1 [cited 2025 Apr 25];10(2):131. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC9952008/

103.   Bindschedler A, Schmuckli-Maurer J, Buchser S, Fischer TD, Wacker R, Davalan T, et al. LC3B labeling of the parasitophorous vacuole membrane of Plasmodium berghei liver stage parasites depends on the V-ATPase and ATG16L1. Mol Microbiol [Internet]. 2024 Jun 1 [cited 2024 Nov 13];121(6):1095–111. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/mmi.15259

104.   Blight J, Sala KA, Atcheson E, Kramer H, Turabi A El, Real E, et al. Dissection-independent production of Plasmodium sporozoites from whole mosquitoes. Life Sci Alliance [Internet]. 2021 Jul 1 [cited 2025 Apr 25];4(7):e202101094. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC8321652/

105.   Liffner B, Silva TLA e, Vega-Rodriguez J, Absalon S. Mosquito Tissue Ultrastructure-Expansion Microscopy (MoTissU-ExM) enables ultrastructural and anatomical analysis of malaria parasites and their mosquito. BMC Methods 2024 1:1 [Internet]. 2024 Oct 7 [cited 2025 Apr 25];1(1):1–15. Available from: https://bmcmethods.biomedcentral.com/articles/10.1186/s44330-024-00013-4

960   106.   Rankin KE, Graewe S, Heussler VT, Stanway RR. Imaging liver-stage malaria parasites. Cell
961   Microbiol [Internet]. 2010 May [cited 2023 Jan 19];12(5):569–79. Available from:
962   https://pubmed.ncbi.nlm.nih.gov/20180802/

964   **Figures**

Figure 1: Immunofluorescence assay and expansion microscopy (PS-ExM) reveals liver stage parasite subpellicular microtubule. **(A)**, and **(B)**, show maximum projection of z-stacks confocal images of infected HeLa cells and **(C)** sporozoites. Host and parasite microtubules were stained with anti-α-tubulin (magenta), and nuclei with DAPI (blue). The parasitophorous membrane vacuole (PVM) was stained with anti-UIS4 (green). **(A)** HeLa cells were infected with *Plasmodium* salivary glands sporozoites and methanol-fixed at 24, 30, 48, and 56 hpi. IFA was performed. **(B)** Fixed infected HeLa cells were expanded using the PS-ExM protocol. Inlets show LSPMB structures at 6, 30, and 48 hpi. Scale bar: 10 μm. **(C)** To determine whether the LSPMB originates from the sporozoite stage, PS-ExM was performed on salivary gland sporozoites, and parasite liver stage at 2 and 6 hpi. **(D)** LSPMB volume was quantified from PS-ExM z-stacks using 3D and 4D analysis in IMARIS software. Normalised mean volumes were: 1294.12 μm³ in sporozoites, 628.34 μm³ at 2 hpi, 414.84 μm³ at 24 hpi, 231.4 μm³ at 48 hpi, to 5.6 μm³ at 54 hpi. **(E)** Live cell confocal imaging of infected HeLa cells incubated with tubulin tracker SPY650. Images show microtubules at 10, 30, 36, and 48 hpi. Host and parasite microtubules are shown in magenta; the parasite expresses cytosolic GFP (green). *Scale bar: 5 μm.*

Figure 2: Microtubules are essential for *Plasmodium* liver stage development. **(A** and **B)** show the effect of oryzalin on parasite microtubules during liver stage development. Sporozoites isolated from mosquito salivary glands were incubated in medium containing either 10 or 20 µM of oryzalin. HeLa cells were infected and fixed at 48 hpi with cold methanol prior to PS-ExM. **(A)** confocal z-stack projections of control cells (DMSO) and cells treated with 10 and 20 µM of Oryzalin. **(B)** confocal z-stack projections of infected cells where oryzalin was washed out at 24 hpi. Microtubule was stained with anti-α-tubulin (magenta), the PVM with anti UIS4 (green), and nuclei with DAPI (blue). Scale bar: 10 µm. ''+'' indicates cells treated with Oryzalin; '' - '' indicates cells from which the drug has been removed at 24 hpi. **(C)** Normalized total parasite DNA volume at 48 hpi. Mean DNA volume was 1952.02 µm³ in the DMSO control, 827.26 µm³ in 10 µM oryzalin-treated cells, and 1411.73 µm³ after drug removal. **(D)** Normalized total hemi-spindle volume. The mean volume was 539.3 µm ³ in the control, 0 µm³ with oryzalin treatment, and 89.08 µm³ after drug removal. **(E)** Normalized LSPMB volume at 48 hpi. Mean volume was 5.92 µm³ in the control, 2.3 µm³ with oryzalin, and 1.65 µm³ after drug removal. **(F, G and H)** Parasite size measurement at 6, 24, and 48 hpi, respectively. At 24 and 48 hpi, mean parasite size was 145.9 µm² in the control, 71.75 µm² with oryzalin, and 115.3 µm² after drug removal. **(I)** Parasite survival rate from 6 to 48 hpi. A slight increase in parasite death was observed in oryzalin-treated cells compared to the DMSO control. The red line on the graphs represents the mean and standard deviation. **(C**, **D**, **E**, **F**, and **G)**, and **(H)**: each dot represents an individual quantified parasite. **(J)** Confocal z-stack projection of IFA images at 56 hpi in cells treated with 10 µM oryzalin and DMSO control. Cells were fixed with cold methanol and stained for MSP1 (magenta), α-tubulin (grey) and nuclei (DAPI, blue). Scale bar: 5 µm. Statistical comparisons were performed using one-way ANOVA with Dunnett's multiple comparisons test (*$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$).

Figure 3: LSPMB is polyglutamylated

(A and **B)** show maximum projection of confocal z-stacks form sporozoites and infected primary hepatocytes at different time points during liver stage development. Primary hepatocytes were isolated from mice and infected with sporozoites. Cells were fixed and stained with antibodies against **(A)** polyglutamylated tubulin (anti-IN105), **(B)** tyrosinated tubulin (anti-Ty), and α-tubulin (anti-α-tubulin). Nuclei were stained with DAPI (blue), and the PVM with anti-UIS4 (grey). ROIs in **(A)** show the LSPMB at 6 and 36 hpi and polyglutamylated hemi-spindle poles at 48 hpi. ROIs in (B) display LSPMB and α-tubulin evolution across liver stages. **(C)** Normalized LSPMB volume and the hemi-spindle pole counts from 6 hpi to 56 hpi, quantified using IMARIS 3D/4D analysis. LSPMB volume decreased from 1038.9 μm^3 at 6 hpi to 0 μm^3 at 56 hpi, while hemi-spindle poles count increased from 0 at 6 hpi to a median of 928 at 56 hpi, showing a perfect Spearman anti-correlation coefficient of -1. n = 15 per time point. **(D)** Single-plane PS-ExM confocal image of infected HeLa cells at 36 hpi with parasites expressing PMP1::GFP. Cells were PFA-fixed and stained with anti-GFP (magenta), anti-IN105 (green), and anti-UIS4 (PVM). The yellow arrow indicates the direction of Cell RGB Profiler analysis from outside the PVM to inside the parasite. **(E)** Pixel intensity profiles (arbitrary units) for PVM (grey), LSPMB (green), and parasite plasma membrane (magenta) in expanded (PS-ExM) cells (E) and non-expanded (NE) cells. Overlap was observed in NE cells, while PS-ExM resolved LSPMB localization between the PVM and the parasite plasma membrane. n=10, in triplicate. Scare bar: 10 µm. **(F)** Single-plane confocal image of infected primary hepatocytes fixed with cold methanol. Cells were stained with anti-IN105 (green), α-tubulin (magenta), and DAPI (blue). The ROI shows a LSPMB tubule in direct contact with a parasite hemi-spindle pole. MZP: Maximum Z projection. Scale bars: 10 µm.

Figure 4: *Plasmodium* α1-tubulin C-terminal post-translational modification is important for the mosquito stage development. **(A)** Schematic of the double homologous recombination (DHR) construct targeting the *Plasmodium* α1-tubulin gene (PBANKA_0417700) for C-terminal mutagenesis. The cassette (blue box) includes the 3'UTR *P. berghei* dihydrofolate reductase (PbDHFR), elongation factor-1 alpha (ef1α) promoter, human dihydrofolate reductase (hdhfr), yeast FCU (yfcu), 3'UTR PbDHFR, Hsp70 promoter, mCherry and the 3' UTR Hsp70, with the 3'UTR region of the PBANKA_0417700 gene. This cassette was flanked by a 5′ homologous region (5′HR) containing the α1-tubulin coding region including the stop codon (with introduced mutations in the C-terminal region, cyan box) and a part of the 3'UTR of the gene (3'HR). The construct was transfected into the PbWT (*P. berghei* ANKA line cl15cy1). Black arrows indicate PCR primer binding sites and expected product sizes. Primer sequences are listed in Table 2 and 3. **(B)** Integration PCR results for the mutant parasite lines, showing expected band sizes on a 1% agarose gel: 5'HR = 4.4 kb, 3'HR = 2.7 kb, whole locus (WL) = 8.8 kb. Ladder: 1 kb DNA marker. Genotypes: WT, (wild-type); PsΔ (glutamine-to-alanine substitution); TyΔ (deletion the C-terminal tyrosination/detyrosination residues); CtΔ (complete C-terminal deletion), and NoΔ (construct control with no mutation). To ensure that all samples contained the same amount of DNA, a control gene representing a portion of tubulin (Ctrl tub about 470 bp) was used for integration PCR. **(C)** Representative confocal images of live oocsts at 11 days post infection (dpi). Nuclei were stained with Hoechst 33342 (hot cyan) and cytosolic mCherry (red) was expressed under the HSp70 promoter. BF, brightfield. **(D)** Normalized salivary sporozoite counts at 18 dpi. The control (NoΔ) averaged 14,033 sporozoites count per mosquito (C./mosq.), compared to 7,300 (PsΔ), 8,066 (TyΔ), and 200 (CtΔ). Experiment performed in triplicate; n = 20 per replicate. **(E** and **F)** Wide-field microscope motility assay images of salivary gland sporozoites showing the typical circular movement (maximum z projection in Fiji) of the control, PsΔ, and TyΔ salivary gland (SG) sporozoites **(E)** and the control and CtΔ hemolymph (H) sporozoites **(F)**. For CtΔ, hemolymph sporozoites were used due to the absence of salivary gland sporozoites; these lacked typical motility. **(G))** Quantification of sporozoite gliding velocity for PsΔ, and TyΔ salivary gland sporozoites. Mean velocities: NoΔ = 1.3 µm/s, PsΔ = 1.38 µm/s, TyΔ = 1.9 µm/s, CtΔ = 0.5 µm/s. Statistical analysis: one-way ANOVA with Dunnett's multiple comparisons test (*$p \le 0.05$, **$p \le 0.01$, ***$p \le 0.001$). **(H)** Quantification of sporozoite gliding velocity of CtΔ haemolymph sporozoites. The average gliding speed for control hemolymph sporozoites was 1.4 µm/s, whereas CtΔ sporozoites exhibited a slower speed of 0.5 µm/s. Statistical comparison tests were performed using two tailed unpaired t tests (*$p \le 0.05$, **$p \le 0.01$, ***$p \le 0.001$).

Figure 5: Liver stage development of *Plasmodium* α1-tubulin mutant parasites.

(A, B and **C)** Maximum projections of PS-ExM-acquired z-stack confocal images of infected HeLa cells at 6 and 36 hpi, showing the control (NoΔ), PsΔ, and CtΔ mutant parasites. **(D and E)** Maximum projections of PS-ExM-acquired z-stack confocal images of control and TyΔ salivary gland sporozoites, as well as CtΔ hemolymph sporozoites (E). In **(A, B and C)**, cells were stained with antibodies against polyglutamylated tubulin (anti-IN105), while **(D and E)** antibodies against tyrosinated tubulin (anti-Ty) were used. In all panels, α-tubulin was labeled with anti-α-tubulin, nuclei with DAPI (blue), and the parasitophorous vacuole membrane (PVM) with anti-UIS4 (grey). **(F** and **G)** represent the fluorescence intensity (arbitrary units, a.u.) of anti-Ty in sporozoites (F) and anti-IN105 at 36 hpi in the liver stage (G). In **(F)**, the mean fluorescence intensity of CtΔ (22 a.u.) was significantly reduced compared to TyΔ (33.4 a.u.) and NoΔ (49.6 a.u.), while in **(G)**, PsΔ (79.9 a.u.) showed a significant reduction compared to NoΔ (121 a.u.). **(H** and **L)** showed infection rates of mutant parasites in HeLa cells at 6 hpi, determined by pre-counted sporozoites and quantitative microscopy. PsΔ (23.7%) and TyΔ (27.3%) exhibited typical infection rates compared to control (NoΔ) (16%) and WT (14%) (H), whereas CtΔ hemolymph sporozoites had a drastically reduced infection rate (2%) relative to controls hemolymph sporozoites (7.4%) (L). **(K** and **O)** Parasite survival from 6 to 48 hpi, normalized to 100% at 6 hpi. At 48 hpi, survival rates were 63% (NoΔ), 60% (WT), 50% (PsΔ), and 47% (TyΔ) (K). In contrast, CtΔ hemolymph sporozoites showed a significantly lower survival rate (2%) compared to controls (52%) (O), indicating impaired viability. (**I** and **J, M** and **N)** Parasite size measurements at 6 and 48 hpi for control, PsΔ, and TyΔ salivary gland sporozoites (I and J), as well as control and CtΔ hemolymph mutants (M and N). At 48 hpi, PsΔ (208.16 μm²) and TyΔ (203.7 μm²) were significantly smaller than NoΔ (237.9 μm²) and WT (235.7 μm²) (J). The CtΔ hemolymph mutants failed to establish liver stage infection, with an average size (88.1 μm²) markedly smaller than controls (187.4 μm²) (N). Each dot represents an individual parasite. PsΔ: Glutamine-to-alanine substitution at the polyglutamylation site; TyΔ: Deletion of C-terminal tyrosination/detyrosination residues; CtΔ: Complete deletion of PBANKA_0417700 α-tubulin C-terminal PTM residues and NoΔ: Control construct with no mutation. WT: wild type parasite. The experiments were conducted using, **(F, H, I, J, K)**: One-way ANOVA with Dunnett's multiple comparisons test (*p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001). **(G, L, M, N, O)**: Two-tailed unpaired t-tests (*p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001). All experiments were performed in triplicate.

Figure 6: Model of the tubulin behavior during Plasmodium berghei salivary gland sporozoite and exo-erythrocytic stage.

The graphical abstract summarizes the key finding of this study.

It illustrates the development of *Plasmodium* parasites within primary hepatocytes, from invasion to the cytomere stage. α-tubulin was stained with anti-α-tubulin (magenta), nuclei with DAPI (blue), and the parasitophorous vacuole membrane (PVM) with anti-UIS4 (grey). The model highlights the evolution of sporozoite subpellicular microtubules (SSPM) into liver stage subpellicular microtubules (LSPMB), along with their associated tubulin post-translational modifications (PTMs). We propose that polyglutamylated LSPMB originates from the SSPM and is recycled during liver stage development to support hemi-spindle pole formation, membrane biogenesis, and potentially other functions. Upon hepatocyte invasion, the SSPM loses tyrosination, a modification that initially confers stability and supports sporozoite motility. This detyrosination likely facilitates parasite shape transformation, increases microtubule dynamics, and enables interactions with ER structures and MAPs. Tyrosination reappears in late stages, possibly stabilizing merozoite microtubules for erythrocyte invasion. Microtubule-stabilizing drugs did not affect the SSPM but inhibited liver stage microtubule polymerization, confirming their dynamic nature. Inhibition of microtubule dynamics also prevented merozoite formation. Finally, C-terminal tubulin PTM residues were shown to be essential for salivary gland sporozoite development and successful liver stage infection.

1112
S1 Figure Effect of *Plasmodium*-specific microtubule inhibitors on HeLa cells
HeLa cells were seeded in 6-well plates and treated the following day with various concentrations of
microtubule-targeting drugs. Media were replaced daily with fresh drug-supplemented medium. After
five days, surviving cells were detached and counted. **(A)** Effect of amiprophos-methyl (AMP) on HeLa
cell viability. A significant reduction in cell count was observed only at 50 µM compared to the DMSO
control. **(B)** effect of the oryzalin on HeLa cells viability. No significant differences in cell count were
observed at any tested concentrations tested relative to DMSO. **(C)** represent the parasite size at 48
hpi in infected HeLa cells treated with oryzalin and AMP at 50 µM. oryzalin showed a significant
inhibitory effect on parasite growth compared to AMP and DMSO control. Statistical comparison tests
were performed using a one-way ANOVA test (*$p \le 0.05$, **$p \le 0.01$, ***$p \le 0.001$). **(D** and **E)** Maximum
projection confocal IFA z-stacks of HeLa cells treated with amiprophos-methyl (D) and oryzalin (E)
tested at different concentrations, respectively. Cells were fixed with cold and stained with anti-α-
tubulin (magenta) and DAPI (blue). **(F)** Confocal IFA z-projection of HeLa cells treated with low-dose
oryzalin (0.5 µM), showing unusually elongated microtubule structures. **(G)** PS-ExM confocal z-stack
of oryzalin-treated cells showing the very long LSPMB and hemi-spindles. **(H)** Quantification of
LSPMB and hemi-spindle lengths. Oryzalin-treated microtubules were on average four times longer
than those in DMSO-treated controls. Each dot represents a measurement from a single cell. LMT,
Long microtubule; NMT, normal microtubule. Statistical comparison tests were performed using two
tailed unpaired t tests (*$p \le 0.05$, **$p \le 0.01$, ***$p \le 0.001$).

S2 Figure *Plasmodium* end-binding protein 1 (eB1) is expressed in mosquito and liver stage

(A) Pearson correlation coefficients between polyglutamylated LSPMB and α-tubulin LSPMB at 6,
36, and 48 hpi during liver stage development. On average, 92% of polyglutamylated tubulin
colocalized with the LSPMB. **(B)** Maximum projection of PS-ExM confocal z-stacks of infected HeLa
cells fixed at 30 and 48 hpi. Cells were stained with anti-γ-tubulin (green), anti-α-tubulin (magenta),
anti-UIS4 (grey, PVM), and DAPI (blue, nuclei). White arrows indicate host centrosomes; yellow
arrows highlight the LSPMB. Parasite hemi-spindle poles were also detected with γ-tubulin. **(C)**
Schematic of the double homologous recombination construct used to tag *Plasmodium* eB1
(PBANKA_0405600) with eGFP at the C-terminus. The expression cassette (blue box) includes
3′UTR *Pb*DHFR, EF1α promoter, hDHFR, yeast fcu, and the 3′UTR of PBANKA_0405600, flanked by
the 5′HR (eB1 coding region) and 3′UTR. The construct was transfected into the PbmCherry 1868
mother line. **(D)** Integration PCR confirming correct insertion of the eGFP-tagged eB1 cassette. PCR
product sizes: 5′HR = 1.5 kb, 3′HR = 1.3 kb, whole locus (WL) = 6 kb. Primer sequences listed in
Table 1. Lane labels: Tr (transgenic), WT (wild type), ctrl (control). Ladder: 1 kb DNA marker. **(E)** Flow
cytometry profile of erythrocytes infected with eB1-GFP-expressing parasites. Gate P1 includes all
analyzed cells; P11 identifies mCherry-positive (PE-Texas Red-H) infected erythrocytes; R10
identifies GFP+/mCherry+ gametocytes. **(F)** Summary table showing parasitemia percentages across
gates defined in (E). **(G)** Representative confocal live imaging of *P. berghei* eB1-GFP oocysts at 7
and 12 days post-infection (dpi), showing eB1-GFP (green), parasite mCherry (red), and Hoechst-
stained nuclei (blue). Scale bar: 5 μm. **(H)** Confocal IFA image of *P. berghei* eB1-GFP sporozoites
stained with anti-GFP (green), anti-α-tubulin (magenta), and anti-circumsporozoite protein (CSP,
grey). **(I)** Maximum projection of PS-ExM confocal z-stacks of HeLa cells infected with *P. berghei* eB1-
GFP parasites at 36 and 48 hpi. Cells were stained with anti-GFP (green), anti-α-tubulin (magenta),
and anti-UIS4 (grey). Scale bar: 10 μm.

A

— Residue 40 (K40) where Acetylation happens
— C-terminal (Y, tyrosinated and E, Detyrosinated)
— Residue where polyglutamylation happens

>**XP_034420234.1 alpha tubulin 1 [_Plasmodium berghei ANKA_] 453aa**
MREVISIHVGQAGIQVGNACWELFCLEHGIQPDGQMPPD**Q**AGRANDDAFNTFFSETGAGKHVPRCVFVDL
EPTVVDEVRTGTYRQLFHPEQLISGKEDAANNFARGHYTIGKEVIDVCLDRIRKLADNCTGLQGFLMFSA
VGGGTGSGFGCLMLERLSVDYGKKSKLNFCCWPSPQVSTAVVEPYNSVLSTHSLLEHTDVAIMLDNEAIY
DICKKNLDIERPTYTNLNRLIAQVISSLTASLRFDGALNVDVTEFQTNLVPYPRIHFMLSSYAPVVSAEK
AYHEQLSVSEITNSAFEPANMMAKCDPRHGKYMACCLMYRGDVVPKDVNAAVATIKTKRTIQFVDWCPTG
FKCGINYQPPTVVPGGDLAKVMRAVCMISNSTAIAEVFSRMDQKFDLMYAKRAFVHWYVGEGMEEGEFSE
AREDLAALEKDYEEVGIETNDGEG**E**DEGYEAD**Y**

>**XP_034420468.1 alpha tubulin 2 [_Plasmodium berghei ANKA_] 450aa**
MREVISIHVGQAGIQIGNACWELFCLEHGIQPDGQMPSD**Q**VVAGGDDAFNTFFSETGAGKHVPRCVFVDL
EPTVVDEVRTGTYRQLFHPEQLISGKEDAANNFARGHYTIGKEIVDICLDRVRKLADNCTGLQGFLMFNA
VGGGTGSGLGCLLLERLAIDYGKKSKLNFCSWPSPQVSTAVVEPYNSVLSTHSLLEHTDVAIMLDNEAIY
DICKKNLDIERPTYTNLNRLIAQVISSLTASLRFDGALNVDVTEFQTNLVPYPRIHFMLSSYAPIISAEK
AYHEQLSVSEITNSAFEPASMMAKCDPRHGKYMACCLMYRGDVVPKDVNAAVATIKTKRSIQFVDWCPTG
FKCGINYQPPTVVPGGDLAKVMRAVCMISNSTAIAEVFSRMDQKFDLMYAKRAFVHWYVGEGMEEGEFSE
AREDLAALEKDYEEVGIETNDGEG**E**DEGY**E**

B

2059

_Pb_ PBANKA_0417700.	GAAACAAATGATGGCGAAGGAGAAGATGAAGGATATGAGGCAGACTATTAA
_Pb_ PBANKA_0417700 No"	GAAACAAATGATGGCGAAGGAGAAGATGAAGGATATGAGGCAGACTATTAA
_Pb_ PBANKA_0417700 Ps"	GAAACAAATGATGGCGAAGGAG**CG**GATGAGAAGATGAGGCAGACTATTAA
_Pb_ PBANKA_0417700 Ty"	GAAACAAATGATGGCGAAGGAGAAGATGAAGGA---------------TAA
_Pb_ PBANKA_0417700 Ct"	GAAACAAATGATGGC-------------------------------TAA

C

D

E

S3 Figure *Plasmodium* α-tubulin proteins and mutant parasite characterization. **(A)** Comparison of *Plasmodium* α1- and α2-tubulin protein sequences highlighting predicted post-translational modification (PTM) sites. Polyglutamylation site E445 is conserved in both isoforms (green). α1-tubulin contains a predicted tyrosination site at Y453 (magenta), while α2-tubulin has a predicted detyrosination site at E450 (magenta). Neither isoform has the conserved acetylation site K40 (yellow). C-terminal PTM regions are underlined in red. **(B)** Sequence alignment of PBANKA_0417700 gene showing mutations introduced in the tubulin mutant parasites. Red highlights indicate substitutions or deletions. Dashes (—) indicate deleted nucleotides. Ps″: glutamine (E) substituted with alanine (A) at the polyglutamylation site; Ty″: deletion of the C-terminal tyrosination/detyrosination motif; Ct″: deletion of the entire C-terminal PTM region; No″: unmodified construct control. **(C)** Structural comparison of α1-tubulin mutant proteins using AlphaFold 3 predictions, aligned with PyMOL (Version 2.6.0). Green: wild type (No″); cyan: Ps″ mutant; magenta: Ty″ mutant; and yellow: Ct″ mutant. All predicted structures retained overall folding. **(D)** Maximum projection of confocal z-stacks of HeLa cells infected with mutant parasites (adapted from Freedy et al.). Microtubules were stained with anti-α-tubulin (magenta) and polyglutamylated tubulin with anti-IN105 (green). Nuclei were stained with DAPI (blue); the parasitophorous vacuole membrane (PVM) was stained with anti-UIS4 (grey). IntronΔ: line with deleted introns in the *Plasmodium* α1-tubulin gene; C-termΔ: line with deletion of C-terminal residues 540ADY. **(E)** In vivo infection kinetics in mice infected with mutant parasites. Parasitemia was measured daily from day 1 to day 5 post-infection. No significant differences were observed between mutant lines (intronΔ and C-termΔ) and wild type (WT). ExonΔ: mutant line in which α1-tubulin exons were replaced with α2-tubulin introns. This line failed to produce salivary gland sporozoites. Liver stage development in this line was delayed by 4 days due to hemolymph sporozoites used. Experiments were performed in triplicate.

S4 Figure Plasmodium α-tubulin mutant parasite oocyst formation and sporozoite infectivity.
(A) Comparison of oocyst area at day 7 and 11 post-infection. No significant difference was observed
between the WT (NoΔ) and the PsΔ or TyΔ mutants at day 7 post-infection. However, by day 11, the
CtΔ mutant exhibited a significantly reduced oocyst size compared to the WT. **(B)** Normalized midgut
sporozoites count per mosquito (C./mosq.) at day 14 post-feeding. No significant difference in
sporozoite numbers was observed between NoΔ and PsΔ, whereas CtΔ showed a drastic reduction.
In contrast, TyΔ displayed both a significantly larger oocyst size at day 11 and a higher sporozoite
count at day 14 compared to the WT. **(C)** Blood parasitemia levels in mice infected with mutant
parasites, measured from day 0 to day 7 post-infection. All mutants (PsΔ and TyΔ) established
infections and progressed similarly to the WT (NoΔ). PsΔ: glutamine-to-alanine substitution at
polyglutamylation site; TyΔ: deletion of C-terminal tyrosination/detyrosination residues; CtΔ: complete
deletion of PBANKA_0417700 α-tubulin C-terminal PTM residues; NoΔ: control construct with no
mutation. The experiments were conducted in triplicate. Statistical multiple comparisons were
performed using one-way ANOVA with Dunnett's multiple comparisons test (*$p \leq 0.05$, **$p \leq 0.01$, ***p
≤ 0.001).

Movies

Movie 1: time-lapse live imaging of *P. bergei* salivary gland sporozoites expressing GFP cytosolically.
Sporozoites were isolated and incubated with tubulin tracker SPY650. Microtubules are shown in
magenta; sporozoites are shown in green. Scale bar: 5μm.

Movies 2, 3, and **4**: Time-lapse live imaging of HeLa cells infected with *P. berghei* liver stage parasites
expressing cytosolic GFP. Infected cells were incubated with tubulin tracker SPY650 for one hour
prior to imaging. The movies show the development and dynamics of the LSPMB during liver stage
infection. Microtubules are shown in magenta; parasites are shown in green. **Movie 2**: imaging from
10 hpi to 25 hpi. **Movie 3:** imaging from 30 hpi to 35 hpi, and **Movie 4:** imaging from 37 hpi to 49 hpi.
Scale bar: 5 μm.

Movies 5, **6**, and **7** Time-lapse live imaging of HeLa cells infected with *P. berghei* liver stage parasites
expressing cytosolic GFP. Infected cells were incubated with tubulin tracker SPY650 for one hour
prior to imaging. The movies highlight the high dynamicity of the LSPMB during liver stage
development, starting from early schizogony (~30 hpi). Microtubules are shown in magenta; parasites
are shown in green. Scale bar: 5 μm.

Movies 8, **9**, and **10** Time-lapse live imaging of HeLa cells infected with *P. berghei* liver stage
parasites expressing cytosolic GFP. Infected cells were incubated with tubulin tracker SPY650 for one
hour prior to imaging. The movies reveal the interaction between the LSPMB and the parasite's hemi-
spindles and spindle poles during liver stage development. Microtubules are shown in magenta;
parasites are shown in green. **Movie 8**: LSPMB (white arrow) localizes at the periphery of the parasite
and connects to hemi-spindles. **Movie 9:** LSPMB (white arrows) connects to the parasite's hemi-
spindle poles. **Movie 10:** shrinkage of the LSPMB during the liver stage development (ROI marked
by a white arrow). Scale bar: 5 μm.

Supplementary Movie 1: 3D projection of PS-ExM images showing LSPMB connected to the hemi-
spindle pole in *P. berghei*-infected primary hepatocytes. Fixed and expanded primary hepatocytes
were stained with anti-IN105 (polyglutamylated tubulin, green), anti-α-tubulin (magenta), and anti-
UIS4 (PVM, grey). The movie shows the LSPMB in contact with the parasite's hemi-spindle pole.
Image analysis was performed using IMARIS 3D and 4D visualization software.

Supplementary Movie 2: *Plasmodium berghei* salivary gland (SG) sporozoite motility of *P. berghei*
NoΔ (control), PsΔ, and TyΔ mutants, as well as hemolymph sporozoites of *P. berghei NoΔ* and CtΔ
mutants. *Scale bar: 5 μm*

Tables

Table 1: *Plasmodium berghei* eB1egfp parasites integration PCR primer

Integration	Mutants	Primer pair	Primer Sequence	Expected size
5'HR		upstream of the construct	GGTTGTAAGTCGAAAAGAGCT GATAGAA	1.5kb
		within the ef1alpha	TCGATGCCCTTCAGCTCG	
3'HR		within yfcu	GAGGTGTTAAGCCAGAGAG	1.3kb
		downstream the 3'UTR	CGGTTCAATAAATGAGGGATTA GC	
Whole locus		Forward	GGTTGTAAGTCGAAAAGAGCT GATAGAA	6kb
		Reverse	CGGTTCAATAAATGAGGGATTA GC	
Control		Forward	TGGAGCAGGAAATAACTGGG	470bp
		Reverse	ACCTGACATAGCGGCTGAAA	

1230

1231   **Table 2:** *Plasmodium berghei* α1-tubulin mutant parasites integration PCR primer

Integration	Mutants	Primer pair	Primer Sequence	Expected size
5'HR	PsΔ TyΔ CtΔ NoΔ	upstream of the construct	CGTATGCTCTACGCACTTCGC	4.4kb
		within the ef1alpha	GGTGCTTTGAGGGGTGAGC	
3'HR	PsΔ TyΔ CtΔ NoΔ	within mCherry	CCTACAAGGCCAAGAAGC	2.7kb
		Within the 3'UTR	CCATGTTCCCATGTGTATCTAC	
Whole locus	PsΔ TyΔ CtΔ NoΔ	Forward	CGTATGCTCTACGCACTTCGC	8.8kb
		Reverse	CCATGTTCCCATGTGTATCTAC	
Control	PsΔ TyΔ CtΔ NoΔ	Forward	TGGAGCAGGAAATAACTGGG	470bp
		Reverse	ACCTGACATAGCGGCTGAAA	

1232

1233

1234

1235

1236

1237

1238

1239

1240   **Table 3:** *Plasmodium berghei* α1-tubulin mutagenesis QuickChangeII primers sequences

Method/Kits	Mutations	Primer pair	Primer Sequence
Mutagenesis/ QuickChangeII	PsΔ	Primer	GTCTGCCTCATATCCTTCATCCGCTCCTTCGCCATCATTT GTTTC
		antisense	GAAACAAATGATGGCGAAGGAGCGGATGAAGGATATGAG GCAGAC
	TyΔ	Primer	GTAAGAAAAACGGCGGCCTTATCCTTCATCTTCTCCTTCG
		antisense	CGAAGGAGAAGATGAAGGATAAGGCCGCCGTTTTTCTTA C
	CtΔ	Primer	GTAAGAAAAACGGCGGCCTTAGCCATCATTTGTTTCAATT CC
		antisense	GGAATTGAAACAAATGATGGCTAAGGCCGCCGTTTTTCTT AC

1241

1242