Evolutionary repurposing of a DNA segregation machinery into a cytoskeletal system

controlling cyanobacterial cell shape

Benjamin L. Springstein[1,#], Manjunath G. Javoor[1,†], Daniela Megrian[2,†], Roman Hajdu[1],

Dustin M. Hanke[3], Florian K. M. Schur[1], Martin Loose[1,#]

[1] Institute of Science and Technology Austria (ISTA), 3400 Klosterneuburg, Austria

[2] Institut Pasteur de Montevideo, Bioinformatics Unit, 11200 Montevideo, Uruguay

[3] Institute of General Microbiology, Kiel University, 24118 Kiel, Germany

[#] Address correspondence to Benjamin L. Springstein (benjamin.springstein@ista.ac.at) and

Martin Loose (martin.loose@ista.ac.at).

[†] MJ and DM contributed equally

Running Title: Evolutionary repurposing of the ParMR system for cell shape control

Key words: ParM, ParR, DNA segregation, cell shape, cytoskeleton, bacteria, MinC,

Nostocales cyanobacteria

Abstract:

Bacteria, despite their diversity, use conserved cytoskeletal systems for their intracellular organization. In unicellular bacteria, the ParMRC DNA partitioning apparatus is well known for forming actin-like filaments that push low copy number plasmids to opposite cell poles. In multicellular cyanobacteria such as *Anabaena* sp. PCC 7120, the presence of a chromosomally encoded ParMR system suggests it may play an analogous role in chromosome segregation. However, we show here that it instead constitutes a novel cytoskeletal system – termed CorMR – whose loss does not impair DNA segregation but leads to severe morphological defects, revealing a role in maintaining cell shape. Using live-cell imaging and *in vitro* reconstitution experiments, we demonstrate that CorM forms dynamically instable filaments. These filaments are recruited to the membrane by CorR, which has acquired a conserved N-terminal amphipathic helix specific to multicellular cyanobacteria. Cryo-EM analysis reveals that CorM forms antiparallel double-stranded filaments, in contrast to the polar, parallel filament pair seen in ParM. Furthermore, CorMR filaments are excluded from the cell poles and division plane by MinC, via a long N-terminal extension also specific to multicellular cyanobacteria. Comparative genomics suggests that cyanobacterial multicellularity co-evolved with the chromosomal relocation of ParMR genes, acquisition of the amphipathic helix in CorR, and functional extension of MinC, supporting an evolutionary link among these features. Our findings uncover a striking repurposing of ancestral biochemical systems, expanding the known roles of ParMR and Min systems beyond plasmid segregation and division site selection to include regulation of cell shape.

Importance:

The ParMRC system is a well-characterized machinery for the segregation of low-copy-number plasmids to the cell poles and, until now, was primarily described to be encoded on plasmids. We found that during the emergence of multicellularity in cyanobacteria, a chromosome-encoded ParMR system has undergone extensive evolutionary and functional repurposing, resulting in a previously undescribed bacterial cytoskeletal system that controls cell shape. We also found that this new cytoskeleton, called CorMR, is under the control of the MinCDE system, which has evolved to not only antagonize Z-ring formation, but also to regulate CorM polymerization. Such interplay between different cytoskeletal systems highlights an intricate regulatory mechanism linking cell division and morphology. As this repurposing of existing cytoskeletal system coincided with the emergence of multicellularity in cyanobacteria, it may have played a key role for the evolution of cyanobacteria with distinct cell shapes. Overall, our discovery highlights the plasticity of bacterial polymeric systems and gives insight into the evolution of cellular and multicellular complexity.

Introduction:

Bacteria employ different polymeric protein systems to control cell shape, growth and division[1]. Protein filaments homologous to the eukaryotic cytoskeleton segregate plasmid DNA (ParM)[2], control cell division (FtsZ and FtsA)[3–5], or mediate cell elongation (MreB)[6], as also reviewed in [1]. In *E. coli* and many other rod-shaped species, a spatiotemporal oscillation of the Min system generates an inhibitory gradient that restricts FtsZ polymerization to the midcell[7].

While these discoveries provided profound insights into how bacterial cells are organized, prokaryotes exhibit immense evolutionary diversity, and we are only beginning to grasp the full spectrum of their organizational strategies. This is particularly true for Cyanobacteria, one of the most morphologically complex phyla described to date[8]. Especially multicellular cyanobacteria exhibit a remarkable diversity in cell and cell filament (termed trichome) morphologies, some of which are capable of cellular differentiation[9]. Among them, multicellular cyanobacteria of the order *Nostocales*, including the model organism *Anabaena*

sp. PCC 7120 (hereafter *Anabaena*), form long trichomes and can differentiate heterocysts -

specialized cells capable of nitrogen fixation[10]. Like all cyanobacteria, *Anabaena* has a hybrid

Gram phenotype, containing an outer membrane and an unusually thick cell wall with several

layers of peptidoglycan (PG)[11]. Multicellularity in cyanobacteria arises through incomplete cell

division, resulting in a shared outer membrane, a continuous PG layer and metabolically

connected cytoplasms[10]. Septal junctions - functional analogs to eukaryotic gap junctions -

promote cell-cell communication through nanopores in the septal PG[10,12,13]. The emergence of

such complex, multicellular species likely requires the evolution of distinct protein systems to

organize the intracellular space, to maintain cell morphology and to control cell division.

The cellular physiology of multicellular cyanobacteria makes them a powerful model for

identifying potential novel solutions for these challenges. For example, cyanobacteria lack

FtsA[14,15], the predominant FtsZ membrane anchor in Gram negative bacteria such as *E. coli*[16],

and instead recruit FtsZ to the membrane by the cyanobacteria-specific protein ZipN[17]. Despite

the presence of thylakoid membranes, the Min system oscillates in unicellular cyanobacteria,

which possess an additional, cyanobacteria-specific protein – Cdv3 – that recruits MinC to

midcell. This suggests that both Cdv3 and MinE contribute to the spatial regulation of MinC's

inhibitory effect on Z-ring formation[18]. In contrast, much less is known about the role of MreB

in *Anabaena*. Unlike in other rod-shaped bacteria, MreB is not essential for viability but is

crucial for preserving the barrel-like shape of *Anabaena* cells[19–21]. It also contributes to septal

PG synthesis and is important for maintaining cell–cell communication between vegetative

cells and heterocysts[20,22].

Another characteristic of cyanobacteria is their remarkable diversity in chromosome

copy numbers, ranging from one to over 200 copies per cell[23], suggesting different strategies

to distribute the genetic information to their daughters. For instance, the highly polyploid

unicellular *Synechocystis sp.* PCC 6803 most likely relies on a random distribution of its up to

~200 chromosomes during cell division[23,24]. In contrast, the maintenance of chromosomes and

low copy number plasmids probably depends on active DNA segregation systems[25]. For

instance, the unicellular *Synechococcus elongatus* PCC 7942 aligns its 3-6 chromosomes

along the long cell axis prior to cell division, through a mechanism that remains largely unknown[26,27], although the involvement of a ParA-like protein has been speculated[28]. In *Anabaena*, which contains ~8 chromosome copies, a dedicated segregation mechanism seems likely, but none has been described to date[21].

In the context of a recent comparative analysis[19], we discovered a chromosomal ParMR system in multicellular cyanobacteria such as *Anabaena.* As ParMR systems typically ensure maintenance of low copy plasmids during bacterial cell division[29], we hypothesized that the chromosomal systems in multicellular cyanobacteria may have evolved to control segregation of chromosomes. Typically, the ParMR system consists of the actin-like ATPase ParM, the DNA-binding adaptor protein ParR, and the centromere-like DNA element *parC*[29,30]. The system gets initiated upon binding of ParR (generally encoded in an operon with *parM*) to short, centromere-like *parC* elements on the plasmid, thereby forming ParRC complexes. *E. coli* ParM forms parallel (i.e., polar), helical double-stranded filaments (a -filament being composed of two protofilaments), which undergo cycles of polymerization and catastrophic depolymerization[30–33]. Through this dynamic instability, ParM searches and captures ParRC complexes, which then stabilizes one filament end. Antiparallel alignment of two ParRC-bound, parallel ParM double-filaments gives rise to a cytoskeletal structure analogous to the mitotic spindle in eukaryotic cells, pushing newly replicated plasmids to opposite cell poles by incorporating new ParM subunits at the ParRC-bound filament ends[29,30,32]. Whether the chromosomal ParMR systems of multicellular cyanobacteria segregate chromosomes has not been shown so far.

Here, we set out to understand the function of the chromosome-encoded ParMR system in *Anabaena*. To our surprise, we found that this system – and those of other multicellular cyanobacteria - have undergone extensive evolutionary diversification such that they now control cell shape. This functional repurposing of an existing bacterial cytoskeleton – from DNA segregation to cell shape control - reveals an unexpected evolutionary innovation in bacterial intracellular organization coinciding with the emergence of multicellularity.

Results:

A chromosomal ParMR system not involved in DNA segregation

While looking for proteins involved in division, growth and cell-cell communication in multicellular cyanobacteria[19], we discovered a chromosome-encoded ParMR system in *Anabaena* (which we first called chrParMR; Fig. 1A). This finding was surprising, as ParMR systems have until recently[34] only been described on plasmids, with a single complete chromosomal ParMR system reported in *K. pneumoniae*[35] and few chromosomal ParR homologs in select *Enterobacteriaceae* species[36] - though the function of these chromosomal systems remains unknown. *Anabaena* has about 8 chromosome copies and how they are maintained during cell division remains unknown[21,23]. We therefore speculated that the chrParMR system may function as a chromosome segregation machinery.

Compared to other plasmid-encoded ParMR such as the *E. coli* ParMR (EcParMR) from the R1 plasmid[37], the chrParMR of *Anabaena* does not have *parC*-like DNA repeats[38–40]. In addition, the proteins are longer and share only limited sequence identity of 20-30% with the other ParMR proteins, contrasting the relatively high sequence identity of the plasmid-encoded EcParM with the chromosomal *K. pneumoniae* ParM (KpParM) ($\sim$54 %) (Fig. 1B). Structure predictions indicated that chrParM contains an extensive N-terminal intrinsically disordered region (IDR), which is absent in EcParM (Fig S1A) and KpParMR. Both chrParR and EcParR are ribbon-helix-helix (RHH) domain containing proteins (Fig 1C, S1B), which are typically dimers and bind DNA[41]. chrParR is predicted to lack helices H3 and H4, which are characteristic for canonical ParR proteins, while helix H5 appears to be divided into two segments - a long H5a and a short H5b segment - connected by an IDR that is not present in EcParR (Fig 1C, S1B). Notably, in *E. coli*, helix H5 mediates ParMR interactions[42].

We first conducted a phylogenetic analysis to identify other chromosomal ParMR systems within cyanobacteria and other bacterial species. As the sequence conservation among ParR homologs (Fig. 1B) is low we performed this analysis looking for ParM homologs by sequence similarity and identifying ParR homologs based on their genomic context and 3D structure prediction. We limited our search to fully sequenced species with closed

chromosomes and plasmids to unambiguously distinguish plasmid-encoded from

chromosomal ParMR systems. The phylogeny reconstructed from the identified ParM protein

sequences (Fig. 1D) shows that all cyanobacterial sequences are monophyletic, suggesting a

common origin. However, we observed three distinct clades of ParMR systems in

Cyanobacteria (Fig. 1D; Supplementary File 1), including two evolutionarily distant clades

whose proteins are mostly plasmid-encoded (yellow and salmon). Of these only the ParMR

proteins from the yellow clade structurally resembled previously described plasmid segregation

systems, based on their 3D structure prediction and automatic annotation[36,43]. One well-

separated clade containing only chromosomal ParMR systems (green) is restricted to the

orders *Nostocales* (including *Anabaena*) and *Oscillatoriales*. Within our representative

sampling of 148 fully assembled cyanobacterial genomes, we found a chromosomal ParMR

system in every *Nostocales* and *Oscillatoriales* genus analyzed, suggesting that its

evolutionary emergence likely traces back to an ancestor of these two multicellular

cyanobacterial orders. Notably, extended N-terminal IDRs were restricted to the green clade

of chromosomal ParMs and are largely absent in the plasmid-encoded clades (Fig. S1C). ParM

sequences identified in other bacterial phyla do not show monophyly within phyla, suggesting

horizontal gene transfer between species. Surprisingly, we found chromosomal ParMR

systems in a broad spectrum of bacterial species, challenging the notion of ParMR as a sole

plasmid segregation system. Chromosomal ParM sequences appear scattered throughout the

phylogenetic tree and are widely distributed. Indeed, when we searched for proteins identical

to *E. coli* ParM from the R1 plasmid (WP_000959884.1) in all *E. coli* assemblies available in

the NCBI RefSeq Genome database, we found 365 occurrences in plasmids and 12 in

chromosomes, indicating that such exchanges occur even among different strains of the same

species. When present in the chromosome, we identified a transposase gene

(WP_001067858.1) in the same genomic context as *parM*, supporting the incorporation of the

gene from a mobile element.

Together, these findings suggest that during bacterial evolution, the ParMR system has

likely been exchanged between bacterial chromosomes and plasmids multiple times during

evolution undergoing both vertical and horizontal transfers between species. In contrast, our

results indicate that the evolutionary history of the ParMR systems in Cyanobacteria is more

constrained, with two ancestral duplications and an inferred event of chromosomal integration

that has been vertically inherited from an ancestor of *Nostocales* and *Oscillatoriales*.

To test the hypothesis that chrParMR is involved in chromosome segregation in

Anabaena, we generated a Δ*parMR* mutant and assessed general DNA distribution and

segregation using DAPI staining. Contrary to our initial hypothesis, we observed no obvious

defects in cellular DNA distribution or any anuclear cells (Fig. 1E, I) as are commonly observed

in strains deficient in other types of chromosome segregation systems[44–46]. Instead, the mutant

showed a striking change in cell morphology, losing the characteristic barrel-like shape of the

wild type (WT) and forming enlarged and rounded cells instead (Fig. 1E). Δ*parMR* cells

displayed a near 1:1 aspect ratio, with width (3.88 ± 0.56 µm) approximately equal to length

(4.53 ± 0.47 µm), contrasting WT cells (length: 4.8 ± 0.96 µm, width: 2.89 ± 0.35 µm) (Fig. 1F-

H). This morphological change was accompanied by a slight reduction in fitness (Fig. S1D),

likely due to defects in septum maturation or completion as identified by scanning electron

microscopy (SEM) (Fig. S1E). Overall, the Δ*parMR* phenotype resembles a previously

described *Anabaena* Δ*mreB* strain[22], suggesting a similar function in cell shape control.

Complementation of the mutant only partially rescued the aberrant cell shape phenotype (Fig.

S1F), likely due to high levels of protein expression from the *parMR* promoter. We could not

recapitulate the native *parMR* expression levels through heterologous expression since the

native *parMR* promoter is also employed by *E. coli* and strains carrying plasmids with P_{parMR}-

parM-parR could not be generated, likely due to the toxic effects of ParR overexpression (Fig.

S1G). Based on these observations, we conclude that the *Anabaena* chrParMR system

represents a unique variant of the ParMR family that plays a role in cell shape control, rather

than in DNA segregation.

Anabaena's chrParM forms dynamically instable cortical filaments _in vivo_

To localize chrParM within _Anabaena_, we used a CRISPR-Cpf1 system and generated a strain in which the 5' end of chr_parM_ is translationally fused to mNeonGreen (mNG), resulting in sole expression of the tagged protein[47]. We note that the mNG-chrParM fusion protein is not fully functional, as evidenced by a mild alteration in cell morphology (Fig. 2A, S2A). C-terminal fusions could not be generated due to the presence of the chr_parR_ ribosomal binding site within the 3' end of chr_parM_. Unlike EcParM, which forms cytoplasmic filament bundles[2], fluorescence imaging of live cells showed that chrParM assembled into filament arrays along the inner cell membrane, with no association to the intracellular thylakoid membranes or the cellular DNA (Fig. 2A-C). These filaments aligned perpendicular to the cell's long axis, with a mean angle of 88.3 ± 12.15° (Fig. 2D). Time lapse microscopy of dividing cells revealed a cell cycle-dependent distribution of chrParM filaments. During cell elongation chrParM is more and more excluded from the cell poles and the future division sites. Right after division chrParM is completely absent from the new cell poles (Fig. 2E; Supplementary Movie 1), suggesting active mechanisms limiting its cellular distribution. Using total internal reflection fluorescence (TIRF) microscopy, we then imaged the polymerization dynamics of chrParM in living cell. Consistent with previously described plasmid-encoded ParMs[48,49], chrParM exhibits dynamic instability _in vivo_ (Fig. 2F,G; Supplementary Movie 2), with a filament growth rate of 18.96 ± 3.609 nm/s for mNG-chrParM (Fig. 2H), approximately half that of EcParM[48]. Filaments also underwent rapid shrinkage at a rate of at least 52.06 ± 23.54 nm/s, about one-fifth of the shrinkage rate observed in _E. coli_[48]. Expressing an mNG fusion of a chrParM variant lacking the N-terminal IDR (i.e., the first 40 amino acids; $^{\Delta1\text{-}40}$chrParM) resulted in more severe morphological defects than the strain expressing mNG-tagged full-length chrParM (Fig. S2A), indicating that the N-terminal IDR is essential for maintaining proper cell shape in _Anabaena_. While mNG-$^{\Delta1\text{-}}$ ^{40}chrParM still predominantly localizes perpendicular to the long cell axis, we observed a generally less ordered mesh-like filament arrangement with occasional localizations to the division site (Fig. S2B). Interestingly, although dynamic instability was retained (Fig. S2C), the N-terminal deletion significantly increased the filament growth rate to 36.14 ± 8.035 nm/s (Fig.

S2D). Thus, chrParM maintained the polymerizing properties of EcParM, including the dynamic

instability, but exhibits a vastly different cellular localization pattern with dynamic properties

dependent on the N-terminal IDR absent in plasmid-encoded ParMs.

To assess the role of chrParR for chrParM localization, we ectopically expressed mNG-

chrParM from a replicative plasmid in ΔparMR. Here, in the absence of chrParR, we observed

mNG-chrParM filaments appearing in the cytoplasm of the cells without a clear localization

pattern (Fig. 2I), suggesting that chrParR is essential for the organization of chrParM into a

cortical array. Genomic replacement of chrParR with an mNG-tagged variant (chrParR-mNG)

as the sole gene copy confirmed its localization to the cell envelope (Fig. S2E). The

accumulation of chrParR-mNG signal at the cell poles is likely an artifact resulting from the

close proximity of two neighboring cell membranes at these sites, where signal from adjacent

cells merges due to the resolution limits of fluorescence microscopy. However, the swollen and

rounded phenotype of this strain indicates that the fusion protein is not fully functional,

suggesting that an unmodified C-terminus is essential for chrParM binding (Fig. S1E, F) similar

to what was previously reported for EcParR[36,42].

In summary, *Anabaena*'s chrParM and chrParR proteins form an evolutionarily distinct

ParMR system, whose deletion results in a pronounced cell shape defect. Using fluorescent

versions of the protein, we found that this actin-related protein assembles into a cortical array

of membrane-associated filaments. Together, these findings strongly suggest that chrParMR

is not involved in DNA segregation. According to their intracellular localization and cellular

function, we therefore propose to rename chrParM and chrParR as CorM and CorR,

respectively, to reflect their cortical cytoskeletal organization.

In vitro characterization of the CorMR system

To biochemically characterize CorM, we purified the protein via heterologous

expression in *E. coli*. CorM retained ATPase activity but in contrast to EcParM, showed

negligible GTPase activity (Fig. S3A), and was unable to polymerize in the presence of GTP

and only weakly polymerized in the presence of AMP-PNP, a non-hydrolyzable ATP analog

(Fig. S3B-D). ATP hydrolysis by CorM increased with protein concentration (Fig. S3E), while

high-speed pelleting assays did not indicate a clear critical concentration for polymerization

under the tested conditions (Fig. S3F). Interestingly, deletion of the N-terminal IDR ($^{\Delta 1-40}$CorM)

led to elevated ATPase activity (Fig. S3G), indicating that the IDR modulates polymerization

kinetics.

We next confirmed interaction between CorM and CorR (Fig. S3H) in high-speed co-

pelleting assays (Fig. S3I, J), as well as through bacterial two-hybrid (B2H) screens (Fig. S3K).

Using CorR mutants, we confirmed that this interaction is mediated by CorR's helix H5b (Fig.

S3K), consistent with the *E. coli* ParMR system,

In plasmid-based ParMR systems, DNA segregation depends on ParR binding to *parC*

repeats (see for example [33,36,50]). To determine whether CorR also binds to DNA, we performed

electrophoresis mobility shift assays (EMSA) using the promoter region of *corMR* and *E. coli*

parMR as bait. CorR failed to bind DNA even at high concentrations (Fig. 3A), whereas EcParR

produced a clear mobility shift (Fig. 3A).

Sequence analysis using AmphipaseeK[51] revealed an N-terminal amphipathic helix

(AH) in CorR's first 10 amino acids, located just upstream of the (RHH)$_2$ domain (Fig. 3B). We

confirmed membrane binding of CorR in vesicle co-sedimentation assays, where the protein

was only found in the pellet in the presence of multilaminar vesicles (MLVs) (Fig. 3C, S4A).

Furthermore, CorM co-sedimented with MLVs mostly only in the presence of CorR, confirming

that CorR recruits CorM to lipid membranes. Although deletion of CorM's N-terminal IDR did

not impair its recruitment to the pellet by CorR, $^{\Delta 1-40}$CorM exhibited increased pelleting

compared to the full-length CorM, suggesting that the N-terminal IDR contributes to filament

length control or bundling (Fig. S4B, C). In contrast, deletion of CorR's N-terminal AH ($^{\Delta 1-}$

^{10}CorR) abolished CorR's ability to bind MLVs and to recruit CorM (Fig. 3C, S4A). Meanwhile,

a CorR variant lacking the C-terminal α-helix (amino acid residues 150-162; CorR$^{\Delta 150-162}$)

retained lipid-binding capacity but failed to recruit CorM (Fig. 3C, S4A), consistent with our

B2H results (Fig. S1E). Together with our observation of cytoplasmic CorM filament bundles

in cells missing CorR, this suggests that CorR acts as a membrane tether for CorM filaments.

Next, we were wondering if other bacterial ParRs have acquired an N-terminal AH

suggesting membrane- instead of DNA-binding. However, by analyzing the protein sequence

of all cyanobacterial ParRs, we found that only CorR homologs from *Nostocales* and

Oscillatoriales cyanobacteria (Fig 3D) contain an AH, with strong sequence conservation in

Nostocales but not in *Oscillatoriales* species (Fig. S4D, S5). This suggests that the acquisition

of *corR* in the chromosome and the emergence of this membrane binding motif have a common

origin.

Together, these findings provide the first evidence for a ParR homolog that has lost its

DNA-binding capacity and instead evolved the ability to bind membranes, mediated by a newly

acquired amphipathic helix at the N-terminus of the $(RHH)_2$ domain.

CorM assembles into antiparallel, double-stranded helical filaments

Given the unexpected behavior of the CorMR system, we aimed to gain structural insights into

the molecular architecture of CorM filaments and their interaction with CorR. To this end, we

performed single particle analysis cryo-electron microscopy (cryo-EM) of filaments consisting

of solely CorM, $^{\Delta 1-40}$CorM or CorM assembled in the presence of CorR (CorMR) (Fig. 4A, S6.1-

6.4). We included $^{\Delta 1-40}$CorM to investigate if the unusually long N-terminal part of CorM might

play a structural role in filament assembly. For all three samples, we obtained electron

microscopy density maps at resolutions that allowed unambiguous assignment of individual

CorM subunits (Fig. 4B, S6.1-6.4), revealing in each case that CorM filaments consist of two

antiparallel, unstaggered protofilaments forming a left-handed helix. Our structures also show

that both deletion of the N-terminal amino acids as well as the presence of CorR in the *in vitro*

assembly do not visibly alter filament architecture (Fig. S6.1-6.4). However, we observed that

CorM alone forms shorter filaments in our *in vitro* preparation (Fig. S6.3) and that these

filaments had a slightly differing rise and twist (46.5 Å/-18.5°) as compared to the other two

structures (CorMR: 46.427 Å/-15.75°; $^{\Delta 1-40}$CorM 47.294 Å/-15.344°) (Fig. S6.1-6.2). Since the

CorMR sample reached the highest resolution (3.9 Å as compared to 4.4 Å for $^{\Delta 1-40}$CorM and

7.1 Å for CorM alone), we used this electron microscopy density map to build a molecular

model of the CorM filament (Fig. 4C, D). Upon closer inspection of the maps fitted with the molecular model, we found that the 40 N-terminal amino acids do not adopt a structured conformation in either of the CorM or CorMR samples. Similarly, within the CorMR filament structure all visible density was occupied by CorM alone, which suggests that CorR only binds to the filament ends, as has been shown for other ParR systems[30]. Given the non-staggered nature of the filament assembly, the lateral association of the two CorM protofilaments (designated n and n* in Figure 4D) is only established by interactions of always only two neighboring subunits. Analysis of our model using PDBePISA[52] shows only weak hydrogen bond interactions across a small lateral interface (interface area 452.2 Å^2) (Figure 4E). Hence, it seems CorM forms weaker lateral interactions compared to EcParM, where multiple lateral interactions between the protofilaments are determined by salt bridges[32]. This potentially has implications in filament stability or dynamic behavior of CorM. Further cryo-EM processing to obtain a higher resolution structure and a structure of filament ends will be required to further characterize the lateral interactions as well as how CorR interacts with CorM.

In contrast, the longitudinal interactions between CorM subunits along the filament are extensive (interface area 1466.7 Å^2) and involve multiple salt bridges and hydrogen bonds (Fig. 4F), similar to EcParM filaments[32]. Interestingly, EcParM filaments have been described to form parallel, polar and staggered helices, which is drastically different from the antiparallel, non-polar and non-staggered assembly of CorM (Figure 4G). Instead, the filament architecture of CorM more closely resembles the antiparallel, non-polar filament doublets formed by MreB (Figure 4G), although MreB does not adopt a helical conformation and does not show dynamic instability.

In summary, our cryo-EM analysis shows a novel filament architecture with in the ParM family formed by our CorM filaments, underscoring the uniqueness of the CorMR system relative to plasmid-encoded, DNA-segregating ParMR systems.

Reconstitution of CorM dynamics *in vitro* on supported lipid bilayers

We next asked whether CorM dynamics could be reconstituted on membranes *in vitro*. To this end, we used supported lipid bilayers (SLBs), as previously described for the cytoskeletal system formed by FtsZ and FtsA[4] (Fig. 5A). Indeed, we successfully recapitulated the *in vivo* behavior of CorM and CorR *in vitro* (Fig. 5B), demonstrating that these two proteins are sufficient to drive filament formation and dynamics.

When Alexa488-labeled CorM was added to SLBs in the presence of CorR, it initially assembled into individual filaments that tended to swivel on the membrane surface. Over time, these filaments formed stable and increasingly dense networks, eventually giving rise to a highly complex filamentous mesh (Fig. 5B). We saw the same filament assembly phenotype over a range of CorM concentrations (from 1-2.5 µM). Fluorescence recovery after photobleaching (FRAP) experiments confirmed the dynamic nature of CorM filaments *in vitro* (Fig. S7A; Supplementary Movies 4), and filament turnover rates increased with rising concentrations of CorR (half recovery times ($T_{1/2}$) for different CorR concentrations: 0.125 µM: 10.64 ± 0.87 s; 0.25 µM: 6.97 ± 0.24 s; 0.75 µM: 4.94 ± 0.14 s) (Fig. S7B).

The $^{\Delta 1-40}$CorM mutant retained dynamic behavior, although FRAP recovery was significantly slower than for the WT protein ($T_{1/2}$ WT: 10.64 ± 0.87 s; $T_{1/2}$ $^{\Delta 1-40}$CorM: 24.65 ± 1.22 s) (Fig. S7A, C, D; Supplementary Movie 5). Notably, filament architecture differed substantially: while initial filament formation resembled that of WT CorM, $^{\Delta 1-40}$CorM ultimately formed a much more ordered and aligned network (Fig. S7E), again suggesting that the IDR modulates filament architecture over time.

Consistent with observations in live cells, CorM filaments displayed dynamic instability *in vitro*, undergoing repeated cycles of growth and sudden shrinkage events (Fig. 5C-E; Supplementary Movie 6). Growth was observed at both filament ends (Fig. 5F–I; Supplementary Movie 7), with a mean growth velocity of 76.63 ± 19.23 nm/s (Fig 5I), comparable to EcParM[49]. Dual-color imaging, where individual monomers were labeled with a different fluorophore than the rest of the filament, revealed that while filaments grew at both ends, individual monomers within the filament remained static (Fig. 5F-H). Intriguingly, when

we quantified bidirectional growth, we saw that one filament end grew slightly faster than the

other (faster end: 35.85 ± 5.53 nm/s; slower end: 23.53 ± 5.36 nm/s). On average resulting in

a net unipolar growth rate of 29.69 ± 8.28 nm/s (Fig. 5I). As the two ends are structurally

identical, we assume that the observed kinetic difference is likely due to only one filament end

being engaged by CorR at a given time. End-binding of CorR would also be consistent with

previous reports on the ParMR system[42] and would also align with the observed swiveling

motion of CorM filaments on membrane surfaces. Due to its small size and the essential roles

of its N- and C-termini, we were unable to generate a fluorescently labeled version of CorR to

verify this assumption. Shrinkage occurred at a much higher rate, with mean disassembly

velocities of -1313.42 ± 671.54 nm/s (Fig. 5I), approximately eight times faster than previously

reported for EcParM[49].

Control of CorMR filaments by the Min system

Plasmid-encoded ParMR systems typically function independently of host regulation and no

other regulators or interactors have been described to date. In contrast, the distinct filament

pattern of CorM *in vivo* (Fig. 2A) and the pronounced phenotype observed upon CorMR

deletion (Fig 1D) suggest active spatial regulation of CorM polymerization and possible

interactions with proteins involved in cell wall remodelling.

To identify putative binding partners for CorM, we performed an *in silico* screen using

AlphaFold-Multimer[53,54] of CorM against the entire *Anabaena* proteome (Fig. 6A;

Supplementary File 2). This approach uses structural predictions of pairwise protein-protein

interactions and ranks them according to their interface-predicted template modeling score

(ipTM score). Consistent with our experimental data, the top-ranking predictions included

CorM-CorR (rank 8) and CorM-CorM (rank 12) interactions, both with ipTM scores above 0.65.

Notably, MinC - a well-characterized inhibitor of FtsZ polymerization and an element of the Min

system[55] - ranked second in the list, with an ipTM score of 0.77 (Fig. 6A, Fig. S8A). This

unexpected prediction raised the possibility that, in multicellular cyanobacteria, MinC may

regulate not only the Z-ring formation but also the CorMR system.

To experimentally validate these interactions, we first used B2H assays, which confirmed that CorM interacts with MinC and revealed that CorR also interacts with MinC (Fig. S8B). However, neither CorM nor CorR interacted with MinD or MinE (Fig. S8D). Additionally, CorR was found to interact with Cdv3 (Fig. S8C), a cyanobacteria-specific component of the Min system[14,18]. MinC interacted with Cdv3 and – as expected from prior findings in *E. coli,* with MinD and MinE (Fig. S8E).

Interestingly, we found that MinC proteins from multicellular cyanobacteria - particularly of *Nostocales* - are substantially longer on average than their counterparts in unicellular cyanobacteria or other bacteria (Fig. 6B, S9; Supplementary File 3). Multiple sequence alignments showed that this length increase is always due to MinC's N-terminal extension, which is unique to *Nostocales* cyanobacteria (Fig. S9).

Structural predictions suggest that this region is intrinsically disordered and includes a short α-helix (Fig. 6C, S8A). We superimposed the predicted structures of CorM-CorR and CorM-MinC (compare Fig. S3H and S8A) and found that of MinC's N-terminal α-helix binds to the same pocket on CorM as the C-terminal α-helix of CorR (Fig 6C). This suggests that CorR and MinC compete for CorM binding. Notably, this binding pocket of CorM is well conserved among chromosomal CorM homologs while highly variable in plasmid-encoded ParMs (Fig. S10A), highlighting its likely relevance in regulating CorM's function.

In vivo, we found that MinC oscillates from pole to pole (Fig. 6D, E; Supplementary Movie 8), as previously described for unicellular bacteria such as *E. coli*[56,57] and the rod-shaped cyanobacterium *S. elongatus*[18]. This provides the first evidence that robust Min system oscillation can also occur in multicellular bacteria. As described in *S. elongatus*, MinC forms local maxima at the cell poles and at the sites of active constriction (Fig. S10B). This localization pattern is mutually exclusive with CorM, which is excluded from these sites, suggesting spatial antagonism. Unfortunately, due to the strong chlorophyll A autofluorescence of cyanobacterial thylakoid membranes, dual-color imaging using green and red fluorophores is currently not feasible, limiting our ability to directly co-visualize MinC and CorM.

To test for the functional relevance of the CorM and MinC interaction, we performed *in*

vitro pelleting assays with CorM under polymerizing conditions while titrating MinC. We found

that MinC potently depolymerized CorM in a concentration dependent manner (Fig. 6F, S10C).

This effect was abolished when we mutated conserved residues in the N-terminal α-helix of

MinC (S50A+L55D, Fig. S10D) or deleted the entire helix (residues 46-67) (Fig. 6G, S10E).

No effect was observed upon addition of a synthetic peptide mimicking MinC's N-terminal α-

helix, suggesting that additional elements of MinC are needed to promote CorM disassembly

(Fig. 6G). Depolymerization of CorM filaments was also observed *in vitro* on SLBs upon

addition of WT MinC but not with the mutant variants or buffer control (Fig. 6H, I;

Supplementary Movies 9-12). Moreover, MinC addition reduced CorM ATPase activity (Fig.

S10F), suggesting a sequestering mechanism of filament depolymerization as described for

inhibitory proteins of actin[58–60], tubulin[61] or FtsZ[62] polymerization and providing a mechanistic

link between MinC and filament disassembly. Finally, using the B2H system, we confirmed that

the N-terminal helix is essential for MinC's ability to interact with CorM (Fig. S10G).

Taken together, these findings reveal a novel regulatory role for the Min system in

multicellular cyanobacteria. Beyond its established role in regulating Z-ring positioning, MinC

modulates the CorMR system by directly competing with CorR for CorM binding, thereby

spatially restricting CorM polymerization and contributing to cell shape control.

Discussion:

This study uncovers the functional repurposing of a protein system - from plasmid segregation

to cell shape control. In multicellular cyanobacteria, the canonical plasmid segregating ParMR

system has undergone substantial evolutionary adaptation to form the cortical CorMR

cytoskeleton on the cell membrane (Fig. 7A). This adaptation seems to be specific to

Nostocales and *Oscillatoriales* cyanobacteria, most of which retain plasmid-encoded ParMR

systems in addition to the chromosomal CorMR system.

The evolutionary history of ParMR appears to be complex, shaped by multiple

horizontal transfers, gene duplications, and transfer between plasmids and chromosomes.

These results challenge the notion that ParMR systems solely act as plasmid segregation

systems, and detailed, lineage-specific analyses will be necessary to determine whether

ParMR systems have acquired distinct functions independently in other groups. In

Cyanobacteria, however, the evolutionary trajectory seems more straightforward, involving two

ancestral duplications and some lineage-specific chromosomal acquisition, with a single

chromosomal incorporation that has been transmitted vertically. Besides CorMR, two broadly

distributed plasmid-associated ParMR clades are observed: one is known to mediate plasmid

copy number regulation and partitioning, while the function of the other remains unknown.

Observations from both bacterial and cyanobacterial phylogenetic relationships raise the

question of whether the ParMR system is more functionally plastic than previously recognized

and might have been repurposed multiple times during evolution.

The transition from plasmid segregation to cell shape control is accompanied by several

structural and functional innovations. Most prominently, the associated adaptor protein, CorR,

has acquired an N-terminal amphipathic helix that enables membrane association while

classical ParR proteins bind DNA. Our cryo-EM structure also reveals that CorM filaments form

antiparallel filament bundles, in contrast to ParM which forms polar double-stranded

filaments[30]. We also find that the canonical cell division regulator MinC has acquired a novel

regulatory role. In *Nostocales*, MinC has an extended N-terminal intrinsically disordered region

containing short α-helical motif, which binds a conserved pocket in CorM. Here, MinC

competes with CorR binding on the one hand and also with CorM polymerization, potentially

through a sequestration mechanism similar to what was reported for actin[58–60], tubulin[61] or

FtsZ[62]. In contrast to MinC's effect on FtsZ, which does not affect FtsZ's GTPase activity, we

observed a reduction in CorM's ATPase activity in the presence of MinC, suggesting that

additional mechanisms beyond mere sequestration may be involved. As a result, MinC is able

to efficiently depolymerize and remove CorM filaments from supported membranes *in vitro* as

well as from the cell poles *in vivo*. This dual functionality of a Min gradient - regulating both

FtsZ and CorM - is, to our knowledge, unprecedented and suggests an expanded role for the

Min system in the cytoskeletal organization of multicellular cyanobacteria (Fig. 7A, B).

The cell shape phenotype of Δ*corMR* and Δ*mreB*[22] observed in *Anabaena* suggests a

role for CorMR as a spatial coordinator of PG biogenesis, while the precise molecular

interactions between CorMR and the PG synthesis machinery are currently under investigation

and will be addressed in subsequent studies. This function is also reminiscent of plant root

cortical microtubules (prcMTs), which guide cellulose synthase complexes to shape plant

cells[63–65]. Both systems operate through transversely oriented polymers and likely act in

concert with cell wall synthases to control cell shape. Although CorM is an actin homolog and

prcMTs are built from tubulin, the mechanistic parallels suggest an important role of dynamic

instability to orient a cortical cytoskeleton perpendicular to the cell axis, a potential case of

convergent evolution of their functions.

Interestingly, while Δ*corMR* and Δ*mreB*, Δ*mreC* or Δ*mreD* mutants exhibit similar

morphological phenotypes - a swollen or rounded and enlarged cell[20,22] - neither system is

essential for viability in *Anabaena*, even under nitrogen starvation conditions that typically

fragment cell filaments. This observation contrasts with the essential role of MreBCD in rod-

shaped unicellular cyanobacteria and in most other bacteria[26,66–68], highlighting the robustness

and redundancy of morphogenetic systems in multicellular cyanobacteria.

CorM and MreB form antiparallel, double-stranded filaments, where individual

protofilaments have opposite orientation[69]. In contrast to MreB filaments, whose motion relies

on PG synthesis[6,70], we find that CorMR filaments exhibit dynamic instability both *in vivo* and

in vitro, similar to its ancestor ParM. Unlike canonical polymeric systems composed of actin,

tubulin, ParM, or FtsZ, which are intrinsically polar and stabilized through nucleotide caps at

their structurally distinct plus ends[32,49,71,72], CorM filaments are antiparallel, and thus both

filament ends are structurally and kinetically identical. Accordingly, the classic nucleotide cap

model for dynamic instabilities of microtubules[73] cannot be applied here and additional

mechanistic studies are necessary to reveal how this is achieved in a non-polar, antiparallel

filament.

Despite the symmetry of CorM filaments, we observed slightly different growth

velocities at the two ends *in vitro* (Fig. 5I) and apparent uni-directional growth *in vivo* (Fig. 2F,

G). We believe that these asymmetric growth kinetics arises from the interaction with CorR.

Based on the previous observations of a clash between the C-terminal peptide of ParR with a

ParM monomer within the ParM filament[42] and the stabilization of the ATP cap at the filament

end during growth[49], we believe that CorR competes with the addition of another CorM

monomer during polymerization. Such a competition between CorR and CorM at one of the

two ends could not only explain asymmetric growth kinetics, but also the characteristic

swiveling of membrane-bound CorM filaments and accelerated CorM turnover at higher CorR

concentrations (Fig. S7B).

The CorMR system exemplifies evolutionary innovation, having been repurposed from a

plasmid segregation module into a membrane cytoskeleton that controls cell shape (Fig. 7C).

In addition, our findings establish the chromosomal CorMR as a novel membrane-associated

actin system in bacteria that assembles into non-polar, dynamically instable filaments. Our

findings highlight the plasticity of bacterial cytoskeletal systems and demonstrate how pre-

existing modules can be co-opted and functionally diversified to serve novel cellular roles,

offering insights into the evolution of cellular and multicellular complexity.

Supplemental material:

Supplementary Movie 1: Live cell time lapse microscopy of mNG-CorM

Anabaena cells expressing mNG-CorM were grown on 1.5% agar pads and illuminated with

an in-house plant growth light. Images were acquired at 15-minute intervals. Scale bar: 5 µm.

Supplementary Movie 2: Dynamic instability of mNG-CorM *in vivo*

Polymerization and sudden shrinkage of mNG-CorM filaments in *Anabaena* cells recorded by

TIRF microscopy. Images were acquired at 10-second intervals. Scale bar: 1 µm.

Supplementary Movie 3: Dynamic instability of mNG-$^{\Delta 1-40}$CorM *in vivo*

Polymerization and sudden shrinkage of mNG-$^{\Delta 1-40}$CorM filaments in *Anabaena* cells recorded

by TIRF microscopy. Images were acquired at 5-second intervals. Scale bar: 1 µm.

Supplementary Movie 4: FRAP of CorM *in vitro*

FRAP experiment of Alexa488-CorM filaments *in vitro* on SLBs. Images were acquired at 1-

second intervals. Scale bar: 10 µm.

Supplementary Movie 5: FRAP of $^{\Delta1\text{-}40}$CorM *in vitro*

FRAP experiment of Alexa488-$^{\Delta1\text{-}40}$CorM filaments *in vitro* on SLBs. Images were acquired at

1-second intervals. Scale bar: 10 µm.

Supplementary Movie 6: CorM polymerization dynamics *in vitro*

TIRF microscopy of Alexa488-CorM filaments on supported lipid bilayers. Individual filaments

can only be tracked during the early stages of polymerization. Images were acquired at 0.15-

second intervals. Scale bar: 2 µm.

Supplementary Movie 7: CorM dual color filament tracking *in vitro*

TIRF microscopy of Alexa488-CorM filaments with single Cy5-CorM monomers on supported

lipid bilayers. Images were acquired at 1-second intervals. Scale bar: 1 µm.

Supplementary Movie 8: Live cell time lapse microscopy of mNG-MinC

Anabaena cells expressing mNG-MinC were grown on 1.5% agar pads and images were

acquired at 1-minute intervals.

Supplementary Movie 9: CorM + MinC *in vitro*

TIRF microscopy of pre-assembled Alexa488-CorM filaments. MinC was added at equimolar

concentration after 30 seconds. Images were acquired at 1-second intervals. Scale bar: 5 µm.

Supplementary Movie 10: CorM + MinC$^{S50A+L55D}$ *in vitro*

TIRF microscopy of pre-assembled Alexa488-CorM filaments. MinC$^{S50A+L55D}$ was added at

equimolar concentration after 30 seconds. Images were acquired at 1-second intervals. Scale

bar: 5 µm.

Supplementary Movie 11: CorM + $^{\Delta46\text{-}67}$MinC *in vitro*

TIRF microscopy of pre-assembled Alexa488-CorM filaments. $^{\Delta46\text{-}67}$MinC was added at

equimolar concentration after 30 seconds. Images were acquired at 1-second intervals. Scale

bar: 5 µm.

Supplementary Movie 12: CorM + buffer ctrl. *in vitro*

TIRF microscopy of pre-assembled Alexa488-CorM filaments. Buffer was added at a volume

equivalent to that of MinC after 30 seconds. Images were acquired at 1-second intervals. Scale

bar: 5 µm.

<u>Supplementary File 1</u>: Taxonomic sampling and ParM and ParR NCBI protein identifiers for

the Cyanobacteria and Bacteria databases.

<u>Supplementary File 2</u>: CorM AlphaFold pulldown against *Anabaena* proteome

AlphaFold results testing CorM against the whole *Anabaena* proteome. To ease readability,

only one out of five predicted dimer models provided by AlphaFold is given.

<u>Supplementary File 3</u>: Taxonomic sampling and MinC NCBI protein identifiers for the

Cyanobacteria and Bacteria databases.

<u>Supplementary Tables</u>: Strains, plasmids and primers used in this study.

Figure legends:

Figure 1: A ParMR system functioning in cell shape control

(**A**) Schematic representation of the chromosomal *parMR* operon from *Anabaena*. The

promoter region is predicted to be about 400 bp long. Numbers below indicate genomic locus

position. (**B**) Sequence identity and similarity (in brackets) in percent of chrParMR from

Anabaena with chromosome-encoded KpParMR from *K. pneumoniae* and plasmid-encoded

ParMR from *E. coli* R1 plasmid. (**C**) Schematic representation of the chromosome and plasmid-

encoded ParM and ParR proteins from *Anabaena* and *E. coli*, respectively. chrParM contains

a N-terminal 40 amino acid (aa) long intrinsically disordered region (IDR) in addition to globular

domain observed in both, chrParM and EcParM. chrParR is longer than EcParR and lacks

helix H3 and H4. Instead, helix H5 appears to be split into one large helix (H5a) and one small

helix (H5b). Numbers indicate aa positions. (**D**) Maximum-likelihood phylogeny of bacterial

ParM. All cyanobacterial sequences (outlined with a dashed line) form a monophyletic clade.

Chromosome-encoded sequences are marked with a green background for Cyanobacteria,

and with a black dot for other bacterial phyla. The scale bar represents the average number of

substitutions per site. For the detailed tree, see Source Data. ParM proteins from the yellow

clade structurally resemble previously characterized plasmid segregation systems, whereas

the functions of ParM proteins from the green and pink clades remain unknown. (**E**)

Micrographs of *Anabaena* WT and Δ*parMR* cells showing an overlay of Direct Illumination

Acquisition (DIA) with chlorophyll a (chlA) autofluorescence images and DAPI staining from

the same cells. (**F-H**) Scatter plots showing the mean cell property values ± standard deviation

(SD) for (**F**) cell width (WT: 4.8 ± 0.96 μm; Δ*parMR*: 4.53 ± 0.47 μm), (**G**) cell length (WT: 2.89

± 0.35 μm; Δ*parMR*: 3.88 ± 0.56 μm) and (**H**) cell aspect ratio (WT: 0.62 ± 0.13; Δ*parMR*: 0.86

± 0.09) of *Anabaena* WT and Δ*parMR* cells. WT: n=222 cells; Δ*parMR*: n=259 cells. (**I**) Mean

DAPI fluorescence intensity from individual cells plotted against the normalized cell length (0

= one cell pole, 1 = opposite cell pole). Values for WT are shown in cyan and values for Δ*parMR*

are shown in grey. Shaded areas correspond to the SD around the mean. The intensity profiles

show a peak near the center of the cell, indicating midcell enrichment of DAPI signal, with

reduced signal toward the poles. WT: n=46 cells; Δ*parMR*: n=49 cells.

Figure 2: CorM forms cortical filaments transverse to the long cell axis

(**A**) Maximum intensity projection micrographs of *Anabaena* cells expressing mNG-ParM from

its native locus as the sole copy within the cell. Left: mNG-ParM fluorescence signal, middle:

overlap of mNG-ParM and chlorophyll A (chlA) autofluorescence (magenta), right: side view of

the left image. (**B**) Cross section view of the same cells as in (**A**) showing the middle plane of

the cells, highlighting that mNG-ParM and chlA signals do not overlap. Beige bar indicates

region used to measure the intensity plot profile from mNG-ParM and chlA signal as shown in

(C). (**D**) Angles of mNG-ParM filaments (n=386 filaments) in respect to the long cell axis.

Cartoon indicates process of angle measurement. (**E**) Dynamic localization of mNG-ParM

during the cell division cycle (see Supplementary Movie 1). The kymograph (center) shows the

mean fluorescence intensity of mNG-ParM over time from 56 dividing cells. Top and bottom

cell sketches indicate the chrParM patterns before and after cell division, respectively. Purple

horizontal bars indicate specific time points (~2.5 h, ~7.5 h, ~12.5 h), corresponding to intensity

profiles shown on the right. These profiles display the average distribution of mNG-ParM

intensities at three key division stages: during growth, before constriction, and after division.

Each line profile is the average signal from 3 micrographs taken within a 45-minute time

window (images were taken every 15 min). Intensity profiles correspond the normalized mean

fluorescence intensity signal from mNG-ParM, while shaded purple regions indicate the cell

boundaries. The SD of the cell boundaries is shaded in light purple (n=56). (**F**) Time series

from Supplementary Movie 2 displaying TIRF micrographs of individual mNG-ParM filaments

over a period of 180 seconds. White triangle marks the initial position of a mNG-ParM filament,

and the red triangle indicates its furthest observed position during the time series. (**G**)

Kymograph of individual mNG-ParM filaments from TIRF microscopy. White arrows indicate

growing filaments whereas magenta arrows indicate filament disassembly/shrinkage. (**H**)

Scatter plot showing the growth velocities (mean 18.96 ± 3.61 nm/s) of mNG-ParM filaments

as recorded by TIRF microscopy. Error bars represent the mean ± SD of 80 independent

filaments measured. (**I**) Micrograph showing merged DIA and mNG-ParM fluorescence images

of Δ*parMR* carrying a low copy number plasmid heterologously expressing mNG-ParM from a

synthetic *ntcA* promoter integrated into the neutral *thrS2* site on the *Anabaena* chromosome[74]

resulting in medium expression levels. Unlike in the WT, mNG-ParM loses its transverse

localization in the absence of ParR.

Figure 3: CorR binds to lipid membranes and recruits CorM

(**A**) EMSA assay with increasing concentrations of CorR or *E. coli* ParR using the promoter

region of *corMR* (400 bp) or *parC* (380 bp), respectively, as DNA bait. Concentrations in µM

are given above the individual lanes. Blot shows a representative example of six individual

experiments. (**B**) Left shows an AlphaFold3 prediction of a CorR dimer with C-terminal CorM

interaction site and N-terminal amphipathic helix (AH) indicated. The AH within the first 10 aa

was identified using Amphipaseek[51]. Notably, AlphaFold3 does not predict the N-terminal AH

as a helical structure but rather as an unstructured region. This may be because AHs often

exist as nascent helices that only adopt their proper fold upon contact with the membrane.

Right shows the helical wheel projection of the first 11 aa of CorR and *E. coli* ParR. Arrows

indicate the hydrophobic moment of the predicted helix indicating the amphipathicity of the

respective proteins. The high hydrophobic moment of CorR indicates the potential to bind to

lipid membranes at the interface between the lipid bilayer and the cytosolic phase. (**C**)

Representative Coomassie-stained SDS-PAGE gel of a pelleting assay with indicated protein

combinations with or without MLVs. 30 % of the total pellet [P] and 12.5 % of the total

supernatant [S] samples were separated on the gel. The respective quantifications are shown

in Fig. S3B. CorR is enriched in the pellet fraction in the presence of MLVs. CorM alone is not

efficiently pelleted at 21,000 x *g* but gets enriched in the pellet when co-incubated with CorR,

indicating that CorR recruits CorM to the MLV-containing pellet. CorR lacking the N-terminal

AH ($^{\Delta1-10}$CorR) fails to bind MLVs and is thus not enriched in the MLV-containing pellet fraction.

In contrast, CorR$^{\Delta150-162}$ retains the N-terminal AH and can be recruited to the pellet via MLV

interactions but cannot recruit CorM to the pellet due to the absence of the C-terminal peptide

required for CorM binding. Dotted lines separate individual gels stiched together for better

visualization. (**D**) Maximum-likelihood phylogeny based on the concatenation of CorM/ParM

and CorR/ParR sequences from Cyanobacteria. The presence of a predicted AH in CorR/ParR

cyanobacterial homologs is indicated with a red dot. *Anabaena* CorR is indicated with a red

circle. The scale bar represents the average number of substitutions per site. For the detailed

tree, see Source Data.

Figure 4: Cryo-EM structure of *in vitro* reconstituted CorM filaments

(**A**) Cryo-EM micrograph showing CorMR filaments formed in vitro at a protein concentration

of 5 µM for each CorM and CorR. (**B**) Cryo-EM density map of *in vitro* reconstituted wild type

CorM filament in the presence of CorR (CorMR) at 3.9 Å. The details of the cryo-EM processing

pipeline for all the three obtained structures are provided in Fig. S6.1-S6.3. (**C-D**) Refined

model of the CorM filament built by fitting the Alphafold-predicted CorM monomer into

CorM+CorR cryo-EM density map. ADP molecules are shown. (**E**) The two antiparallel

protofilaments are annotated as n and n*. Rectangles outline the enlarged lateral and

longitudinal interaction interfaces in **F**. Both of the interface interactions were determined using

PDBePISA (Proteins, Interfaces, Structure and Assemblies)[52]. (**G**) Comparison of surface

representations of CorM (this study), with previously published *E. coli* ParM (PDB entry 5AEY)

and *C. crescentus* MreB (PDB entry 4CZJ) models, respectively. Individual protofilaments are

colored in dark or light green to illustrate their relative arrangement. Similarities and differences

in polarity, subunit arrangement and helical parameters are given below the models.

Figure 5: *In vitro* reconstitution of CorMR polymerization on supported membranes

(**A**) Illustration of the experimental assay to reconstitute CorM polymerization with CorR on

supported lipid bilayers (SLBs). CorR binds to the SLB through its AH and recruits CorM

filaments to the membrane. CorM filaments grow from both sites through bidirectional addition

of monomers to an existing filament. (**B**) TIRF micrographs of purified and Alexa488-labeled

CorM (0.25 µM) together with unlabeled CorM (0.75 µM) recruited to the SLB through

interaction with CorR (0.25 µM). Initially, CorM forms individual filaments (left image) that then

eventually form a denser network of filaments (middle image), which ultimately merge into a

dense, seemingly highly interconnected meshwork of thin filaments (right image). (**C**) Time

series from Supplementary Movie 6 displaying TIRF micrographs of individual Alexa488-

labeled CorM filaments showing dynamic instability. Blue triangles mark the initial position of

an individual CorM filament that continues to grow throughout the whole time series. Orange

triangles mark the initial position of an individual CorM filament that grows initially and then

rapidly collapses before regrowing. The purple triangles indicate the furthest observed position

of another filament which ultimately undergoes rapid and complete depolymerization during

the time series. (**D**) Kymographs of the respective filaments marked with blue, orange and

purple triangles in (C) showing growth, catastrophe and rescue events. The filament indicated

with blue triangles only grows during the 120 sec time series shown in (C) and eventually also

undergoes depolymerization. The filament indicated with purple triangles grows bidirectionally

and then completely collapses. (**E**) CorM filament undergoing an initial polymerization phase

followed by rapid depolymerization. Growth is represented by the increase in distance between the two poles of the filament over time. The filament polymerizes steadily at a rate of 96.95 nm/s (green line) followed by a rapid depolymerization phase with a rate of -958.07 nm/s (orange line). (**F**) Time series from Supplementary Movie 7 displaying TIRF micrographs of an individual Alexa488-labeled CorM filament supplemented with Cy5-CorM over a period of 32 seconds. Alexa488-CorM: 0.5 µM; Cy5-CorM: 4.48 pM; CorR: 0.25 µM. The Cy5-CorM monomer within the Alexa488-CorM filament remains static while Alexa488-CorM monomers are added to both filament poles. (**G**) Kymograph of CorM filament depicted in (F), confirming the immobile nature of the Cy5-CorM monomer and the bipolar extension of the CorM filament. (**H**) Start (triangle) and end (cross) positions of the two ends of the filament ends (light and dark green) shown in (F) as well as the position of the Cy5-CorM monomer (orange) during the time lapse. (**I**) Growth and shrinkage properties of CorM filaments from *in vitro* time-lapse TIRF microscopy. Scatter plots show the mean ± SD filament velocities of Alexa488-CorM during bipolar growth (76.63 ± 19.23 nm/s; n=13), unipolar growth (29.69 ± 8.28 nm/s; n=10), and shrinkage (-1313.42 ± 671.54 nm/s; n=5), as determined by single-filament tracking from TIRF microscopy. Bipolar growth and shrinkage were measured from single and dual color time-lapse imaging series using Alexa488-CorM or Alexa488-CorM with low levels of Cy5-CorM, while unipolar growth was extracted from dual color time-lapse imaging series only.

Figure 6: Control of CorMR filaments by the Min system

(**A**) *in silico* pulldown to identify interactions of CorM (bait) against the whole *Anabaena* proteome (5326 open reading frames included in the analysis). Accuracy of the predicted interaction is given as interface predicted template modelling (ipTM) score as described by AlphaFold2[75]. ipTM scores for CorM, CorR and MinC are highlighted in the plot. (**B**) Scatter plots of MinC sequence lengths (in aa) from all bacteria, cyanobacteria and *Nostocales* cyanobacteria. (**C**) (top) AlphaFold3 prediction of an *Anabaena* MinC dimer. (bottom) A zoom-in into a well-conserved pocket of CorM predicted to bind to the C-terminal α-helix of CorR (blue) and the N-terminal α-helix of MinC (orange), suggesting a competition of CorR and MinC

for binding to CorM. Conservation is based on a multiple sequence alignment from all identified

cyanobacterial chromosomal CorMs (green clade). Coloring by sequence conservation is

based on the entropy-based measure AL2CO from ChimeraX[76] using the default coloring

scheme (cyan: poor conservation; white: intermediate conservation; red: highly conserved;

grey: no conservation). (**D**) *in vivo* time lapse microscopy of *Anabaena* cells expressing mNG-

MinC from a synthetic *ntcA* promoter integrated into the neutral *thrS2* site on the *Anabaena*

chromosome. Orange arrows indicate the positions of mNG-MinC fluorescence maxima in the

uppermost cell over time, highlighting its pole to pole oscillatory behavior throughout the time-

lapse analysis. Grey bar spanning the cell filament indicates region used to generate the

kymograph show in (E). (**F**) Representative Coomassie-stained SDS-PAGE gel from a high-

speed ultracentrifugation-based pelleting assay of 5 µM CorM with or without indicated MinC

concentrations in the presence of 2.5 mM ATP (i.e., under polymerizing conditions). 40 % of

the total pellet [P] and 11.1 % of the total supernatant [S] samples were separated on the gel.

CorM runs at around 45 kDa while MinC runs at ~55 kDa. The respective quantifications are

shown in Fig. S8G. (**G**) Representative Coomassie-stained SDS-PAGE gel from a high-speed

ultracentrifugation-based pelleting assay of 5 µM CorM with or without 5 µM MinC WT,

MinC$^{S50A+L55D}$, $^{\Delta46-67}$MinC or a synthesized peptide mimicking MinC's N-terminal α-helix in the

presence of 2.5 mM ATP (i.e., under polymerizing conditions). 40 % of the total pellet [P] and

11.1 % of the total supernatant [S] samples were separated on the gel. The respective

quantifications are shown in Fig. S10A. (**H**) Snapshots from TIRF microscopy time lapse series

extracted from Supplementary Movies 8-11 of pre-formed CorMR filaments (0.75 µM

Alexa488-CorM, 0.25 µM CorM and 0.25 µM CorR) on SLBs supplemented with 1 µM MinC

WT, MinC$^{S50A+L55D}$, $^{\Delta46-67}$MinC or buffer. Only MinC WT can depolymerize CorMR filaments on

the SLBs. We note that CorMR filaments re-assemble several minutes after addition of MinC

WT, indicating that CorM depolymerization by MinC is a transient effect. (**I**) Normalized

fluorescence intensity over time from the time lapse series shown in (H), with the initial value

set arbitrarily to 1. A decrease in fluorescence signal is observed only upon addition of MinC

WT, while both MinC mutants and the buffer control samples cause an increase in fluorescence

signal, likely due to mixing after addition, which replenishes the local concentration of CorM

monomers near the membrane.

Figure 7: A quadruple evolutionary event regulated cell shape in *Anabaena*

(**A**) Cyanobacterial phylogenetic species tree with annotations indicating the presence of a

chromosomal CorMR system, an amphipathic helix (AH) in CorR homologs, an N-terminal

intrinsically disordered region (IDR) in CorM homologs, and a long MinC protein (>300 aa).

Monophyletic families were collapsed into single branches. Presence of the respective feature

within a family is assumed if at least 50 % of the species within the clade display the feature.

The complete tree is available in the Supplementary Figure 5 and Source Data, and detailed

information is provided in Supplementary File 1 and 3. (**B**) Proposed model for the regulation

of the chromosomal CorMR system in *Anabaena*. CorM assembles into cortical-like filaments

oriented transverse to the cell's long axis. These filaments are dynamically unstable and are

recruited to the cytoplasmic membrane by CorR, which has acquired membrane-binding

affinity through its N-terminal AH. CorM polymers are excluded from the cell poles and division

sites due to the regulatory actions of MinC, which localizes to these regions and actively

depolymerizes CorM filaments. (**C**) A model for the evolutionary diversification and cytoskeletal

innovation of an existing plasmid segregation system (ParMR), which was functionally

repurposed and evolved into CorMR, thereby contributing to the emergence of complex

multicellular (cyano)bacterial species.

Supplementary figure legends:

Figure S1: Properties of the chromosomal ParMR

(**A**) AlphaFold3-predicted structure of chromosomal ParM (chrParM) from *Anabaena* (pTM

score: 0.83) and the resolved X-ray structure of plasmid-encoded ParM from *E. coli* (PDB ID:

4A62). (**B**) AlphaFold3-predicted structure of a chromosomal ParR (chrParR) dimer from

Anabaena (pTM score: 0.34; ipTM score: 0.34) and the resolved X-ray structure of the plasmid-

encoded ParR dimer from *E. coli* (PDB ID: 2JD3). For A and B, predicted aligned error (PAE)

maps and pIDDT score-colored structure predictions for AlphaFold3 predictions are also shown. (**C**) Mean amino acid residues ± SD within the N-terminal IDR from ParM homologs from the green (42.05 ± 11.03), yellow (9.66 ± 11.32) and pink (0.71 ± 2.149) clades of cyanobacterial ParM's showing that only the green clade has any substantial N-terminal IDR. (**D**) Representative growth curve analysis of *Anabaena* WT and Δ*parMR* revealing that the mutant lacking ParMR has a slight growth defect. OD_{750} values were recorded every day or every other day for a total of 11 days. Afterwards, cultures became too sticky to properly measure the OD_{750} value. Error bars represent the mean ± SD of three technical replicates. The experiment was performed twice with similar results. (**E**) Scanning electron microscopy images of (left) *Anabaena* WT and Δ*parMR* revealing cell division defects in 30 % of actively dividing Δ*parMR* cells (6 of 20 cells). These cells displayed irregular or incomplete septa, being partially constricted but not fully separated, a phenotype not observed in the WT (16 cells). Left Δ*parMR* image shows defects in proper constrictions, which are not observed in another subset of Δ*parMR* cells as indicated on the right Δ*parMR* image. (**F**) Complementation attempts of the Δ*parMR* phenotype. DIA micrograph of Δ*parMR* carrying a replicative plasmid expressing *parM-parR* from a medium level, unregulated synthetic promoter (P_{ntcA}) derived from the promoter of the global regulator of nitrogen homeostasis NtcA from *Anabaena*[77]. For comparison Δ*parMR* is shown. The swollen phenotype of Δ*parMR* is moderated in the complementation strain but expression of *parM-parR* from P_{ntcA} is not sufficient to provide the same cellular levels of ParMR as the native promoter can. However, attempts to clone such construct failed because *E. coli* can employ the P_{parMR} promoter, and we found that excess levels of ParMR are toxic for *E. coli*. (**G**) Micrographs of *E. coli* C41 (DE3) co-expressing mNG-ParM from P_{T7} and ParR-mScarlet-I from P_{BAD} grown over night in TB medium supplemented with 0.02 % L-Ara and 0.25 mM IPTG at 37 °C. Cells containing ParR-mScarlet-I signal are strikingly malformed, forming huge balloon-like structures. These cells lack any discernible mNG-ParM signal, indicating that they cannot tolerate both proteins together for extended periods. Micrograph with enhanced contrast reveals red fluorescent cell debris, likely originating from burst cells expressing ParR-mScarlet-I. mNG-ParM localizes as filamentous

bundles in cells with WT-like morphology that lacked ParR-mScarlet-I signal, suggesting that

specifically ParR is responsible for the aberrant cell phenotype.

Fig. S2: ParM's N-terminal IDR is not involved in cellular ParM localization or dynamics

(**A**) Scatter plots showing the mean cell property values ± SD for cell width (WT: 4.8 ± 0.96

µm; Δ*parMR*: 4.53 ± 0.47 µm; mNG-ParM: 3.73 ± 0.65 µm; mNG-$^{\Delta 1\text{-}40}$ParM: 3.42 ± 0.49 µm),

cell length (WT: 2.89 ± 0.35 µm; Δ*parMR*: 3.88 ± 0.56 µm; mNG-ParM: 4.91 ± 0.56 µm; mNG-

$^{\Delta 1\text{-}40}$ParM: 4.06 ± 0.29 µm) and cell aspect ratio (WT: 0.62 ± 0.13; Δ*parMR*: 0.86 ± 0.09; mNG-

ParM: 0.76 ± 0.09; mNG-$^{\Delta 1\text{-}40}$ParM: 0.84 ± 0.09) of *Anabaena* WT, Δ*parMR*, mNG-ParM or

mNG-$^{\Delta 1\text{-}40}$ParM expressing cells. WT: n=222 cells; Δ*parMR*: n=259 cells; mNG-ParM: n=67

cells; mNG-$^{\Delta 1\text{-}40}$ParM: n=144 cells. Values for WT and Δ*parMR* are the same as in Fig. 1F-H.

(**B**) mNG-$^{\Delta 1\text{-}40}$ParM and merged mNG-$^{\Delta 1\text{-}40}$ParM and chlA autofluorescence micrographs of

Anabaena cells expressing mNG-$^{\Delta 1\text{-}40}$ParM as the sole copy obtained with a Nikon SoRa

spinning disk confocal microscope. (**C**) Time series from Supplementary Movie 3 displaying

TIRF micrographs of individual mNG-$^{\Delta 1\text{-}40}$ParM filaments over a period of 230 seconds. White

triangles mark the initial position of a single mNG-$^{\Delta 1\text{-}40}$ParM filament and red triangles mark the

furthest observed position of this filament during the time series. Blue triangles mark the initial

position of another single mNG-$^{\Delta 1\text{-}40}$ParM filament, yellow triangles mark the furthest observed

position of this filament during the time series and purple triangles indicate the position this

filament retracted to upon a dynamically instable shrinkage event. (**D**) Scatter plot showing the

mean ± SD of the growth velocities of mNG-ParM (18.96 ± 3.61 nm/s; n=80) and mNG-$^{\Delta 1\text{-}}$

^{40}ParM (35.75 ± 8.19 nm/s; n=58) filaments, revealing the N-terminal IDR is responsible to limit

the growth kinetics *in vivo*. (**E**) ParR-mNG, DIA and merged ParR-mNG and chlA

autofluorescence micrographs of *Anabaena* expressing ParR-mNG as the sole copy. This

strain's cellular morphology is markedly altered, resembling the Δ*parMR* strain. This suggests

that the ParR-mNG fusion is not fully functional, likely because ParR's C-terminal peptide is

essential for interaction with ParM and is sterically hindered by the C-terminal mNG fusion.

Note that membrane localization of proteins in *Anabaena* looks mostly diffuse at the periphery

as is the case for ParR-mNG. For comparison, *Anabaena* cells expressing soluble mNG are

shown on the right side. Micrographs were obtained by a standard Nikon TI2 epi-fluorescence

microscope.

Fig. S3: Biochemical properties of CorM reveal functional divergence from EcParM and

regulation by the N-terminal IDR

(**A**) NTPase activity assay using the Malachite Green Phosphate Assay Kit (Sigma Aldrich)

shows that CorM readily hydrolyzes ATP but, unlike the EcParM, is not able to meaningfully

hydrolyze GTP. NTPase activity is quantified by the degree of phosphate release during the

conversion of NTP to NDP during NTPase activity. Error bars represent the mean ± SD of two

independent replicates (n=2). (**B**) Representative Coomassie-stained SDS-PAGE gel from a

high-speed ultracentrifugation-based pelleting assay comparing CorM with EcParM (both 5

μM) in the presence of buffer (ctrl.), ATP, ADP, GTP and a non-hydrolyzable ATP analog

(AMP-PNP). Nucleotides were provided at 2.5 mM each. 40 % of the total pellet [P] and 11.1

% of the total supernatant [S] samples were separated on the gel (**C, D**) Quantification of the

pelleting efficiency as displayed in (B), confirming that CorM only polymerizes in the presence

of ATP but not with GTP or AMP-PNP. Error bars represent the mean ± SD of three

independent replicates (n=3). (**E**) ATPase activity assay showing that CorM ATPase activity is

a concentration dependent process that seemingly begins to plateau at higher concentrations.

Error bars represent the mean ± SD of three independent replicates (n=3). (**F**) The degree of

CorM polymerization is not a concentration-dependent process and is not altered upon high-

levels of ParM (n=1). (**G**) ATPase activity assay comparing CorM with a mutant lacking the first

40 aa ($^{\Delta 1-40}$CorM) reveals that the N-terminal IDR limits CorM's ATPase activity. Error bars

represent the mean ± SD of two independent replicates with 3 technical replicates each. (**H**)

AlphaFold3-predicted interaction between CorM and a CorR dimer, together with the

corresponding PAE map, showing interaction of CorR's C-terminal α-helix H5b with CorM, as

well as a pIDDT score-colored CorM-CorR complex (pTM score: 0.59; ipTM score: 0.51). (**I**)

Representative Coomassie-stained SDS-PAGE gel from a high-speed ultracentrifugation-

based pelleting assay comparing ParM and ParR, alone or in combination (5 µM each),

incubated with 2.5 mM ATP or without nucleotide (ctrl.) (n=2). 34 % of the total pellet [P] and

9.7 % of the total supernatant [S] samples were separated on the gel (**J**) Quantification of the

pelleting efficiency from (C), showing pellet and supernatant values for ParM (left graph) and

ParR (right graph) individually. CorR is recruited to the pellet through polymerized CorM (ATP-

containing sample). Data represent the mean ± SD from two independent experiments (n=2)

(**K**) B2H assays testing for the interaction of chrParM with chrParR from Anabaena fused to

either the T25 or T18 subunit, including a ParR variant lacking the predicted C-terminal α-helix,

which is known to facilitate the interaction of plasmid-encoded ParR with ParM in *E. coli*.

chrParR lacking amino acids 150-162 is no longer able to interact with ParM. Negative control:

ParM-T18 with empty pKNT25 plasmid. Positive control: Zip/Zip control. Error bars represent

the mean ± SD of three independent replicates (n=3).

Fig. S4: CorR binds to lipid membranes

(**A**) Quantification analysis of the CorM with CorR pelleting assay from Fig. 3C showing the

values for CorM and CorR separately. P indicates values for pellet samples and S indicates

values for supernatant samples. Error bars represent the mean ± SD from at least three

independent replicates. (**B**) Representative Coomassie-stained SDS-PAGE gel from a

pelleting assay with indicated protein combinations with or without MLVs (proteins are provided

as 5 µM each). 30 % of the total pellet [P] and 12.5 % of the total supernatant [S] samples

were separated on the gel. (**C**) Quantifications from pelleting assay shown in (B), showing

values for CorM and CorR separately. Error bars represent the mean ± SD from three

independent replicates (n=3). (**D**) Amino acid sequence logo (using WebLogo[78]) from the

multiple sequence alignment shown in Fig. S5, showing only the first 20 aa of CorR's N-

terminus. To enhance readability, CorR sequences from *Oscillatoriales* and *Nostocales* are

shown independently, which also highlights that the N-terminal CorR region is strongly

conserved among *Nostocales* species and less so among *Oscillatoriales* species.

Figure S5: Conservation of CorR's amphipathic helix

Cyanobacterial phylogenetic species tree with multiple sequence alignment of CorR homologs containing a N-terminal AH. Since AHs were only found among chromosomal CorMR homologs in *Nostocales* and *Oscillatoriales*, only sequences from these species are shown. The N-terminal region comprising the AH (roughly 10 aa) is highly conserved among *Nostocales* species and less among *Oscillatoriales* species.

Fig. S6: Cryo-EM image processing strategy applied to obtain structure of CorM filament (CorMR filaments (6.1), $^{\Delta 1\text{-}40}$CorM (6.2), and CorM (6.3)) and the final cryo-EM maps (6.4)

Schematics of the image processing strategy used for determining all the three obtained CorM filament structures using cryoSPARC v4.4. Representative micrographs, 2D class averages, 3D class averages, and final electron microscopy density maps are shown. For helical refinement jobs, the initial estimates of helical rise and helical twist were obtained using online helical indexing tool (HI3D- Helical Indexing using cylindrical projection of a 3D map[79]). These values are marked with * under each helical refinement job. Two side views of local resolution filtered EM maps and Fourier Shell Correlation (FSC) curves are displayed. The surface representation of the final cryo-EM maps of CorM, $^{\Delta 1\text{-}40}$CorM, and CorMR and their estimated resolutions (using FSC 0.143 criteria) are shown in 6.4.

Fig. S7: The N-terminal IDR prevents filament bundling and increases protein dynamics

(**A**) Representative FRAP snapshots from TIRF microscopy time series (Supplementary Movies 4-5) of pre-formed CorMR or $^{\Delta 1\text{-}40}$CorMR filaments (0.75 µM unlabeled CorM WT or $^{\Delta 1\text{-}40}$CorM, 0.25 µM Alexa488-CorM or Alexa488-$^{\Delta 1\text{-}40}$CorM and 0.125 µM unlabeled CorR) on SLBs subjected to photobleaching. (**B**) Dependency of the half-recovery time on CorR concentration. Plot showing the half-recovery time ($T_{1/2}$) in seconds as a function of CorR concentration (0.125 µM, 0.25 µM, and 0.75 µM). Increasing concentrations of CorR lead to a dose-dependent decrease in $T_{1/2}$, suggesting higher CorM dynamics with higher CorR levels. Error bars represent the mean ± SD from individual measurements: 0.125 µM: 10.64 ± 0.87

nm/s (n=9); 0.25 µM: 6.97 ± 0.24 nm/s (n=6); 0.75 µM: 4.94 ± 0.14 nm/s (n=5). Half-recovery times comparing CorM WT and $^{\Delta1-40}$CorM in the presence of 0.125 µM CorR, showing that the N-terminal IDR in CorM modulates dynamics and reaction kinetics. Error bars represent the mean ± SD from individual measurements: CorM values are the same as in (B); $^{\Delta1-40}$CorM: 24.65 ± 1.22 nm/s (n=7). (**D**) Representative recovery curves from FRAP experiments comparing CorM WT and $^{\Delta1-40}$CorM in the presence of 0.125 µM CorR. (**E**) Representative snapshots of $^{\Delta1-40}$CorM (protein concentrations same as in A) filaments in the course of an imaging session. Like CorM WT, $^{\Delta1-40}$CorM initially forms individual filaments. These filaments then form thick bundles but eventually assemble into a well-defined network of filaments, which appears much more ordered compared to CorM WT (see Fig. 5B for comparison).

Fig. S8: Properties of the *Anabaena* Min system and it's modulating function on CorM
(A) (top) AlphaFold3-predicted interaction between CorM and MinC, together with the corresponding PAE map, showing interaction of MinC's N-terminal α-helix with CorM, as well as a plDDT score-colored CorM-MinC complex (pTM score: 0.51; ipTM score: 0.76). (bottom) AlphaFold3-predicted MinC dimer (pTM score: 0.46; ipTM score: 0.47), including the corresponding PAE map and a plDDT score-colored dimer. (**B-D**) B2H assays testing for the interaction of CorMR with (**B**) MinC, (**C**) Cdv3 or (**D**) MinD and MinE fused to either the T25 or T18 subunit. Both, CorM and CorR interact with MinC but only CorR interacts with Cdv3. Neither CorM nor CorR interacts with MinD or MinE. Negative control: (B, D) CorM-T25 with empty pUT18C plasmid or (C) CorM-T18 with empty pKNT25 plasmid. Positive control: Zip/Zip control Error bars represent the mean ± SD of three independent replicates (n=3). (**E**) B2H assays testing for the interaction of MinC with MinD, MinE and Cdv3 fused to either the T25 or T18 subunit. Results show that MinC interacts with all three proteins. Negative control: MinC-T25 with empty pUT18C plasmid. Positive control: Zip/Zip control. Error bars represent the mean ± SD of three independent replicates (n=3).

Fig. S9: In *Nostocales* cyanobacteria MinC exhibits an extended N-terminal domain

MinC phylogenetic tree, including cyanobacteria and other bacterial species. Blue-shaded cubes next to the tree indicates MinC sequence lengths. Note that specifically *Nostocales* cyanobacteria contain long MinC sequences (>300 aa), which is represented by an N-terminal IDR with an α-helix in between. On the right, a multiple sequence alignment of all listed MinCs is shown with two different conservation patterns. Orange shades indicate conservation exclusively among the N-terminal IDR of *Nostocales* MinCs whereas grey shades indicate conservation exclusively among the globular domain of MinC. Black dots indicate Ultrafast Bootstrap supports (UFB) >= 85. The scale bar represents the average number of substitutions per site.

Fig. S10: MinC's N-terminal α-helix mediates CorM depolymerization

(**A**) AlphaFold3-predicted structures of (top) plasmid-encoded ParM (#BAB78165.1; All7081) from the large low copy number Alpha plasmid from *Anabaena* or (bottom) plasmid-encoded EcParM (WP_000959884.1) from the R1 plasmid. Conservation is based on multiple sequence alignments from all identified cyanobacterial plasmid-encoded ParMs or all identified other bacterial plasmid-encoded ParMs. Coloring by sequence conservation is based on the entropy-based measure AL2CO from ChimeraX[76] using the default coloring scheme (cyan: poor conservation; white: intermediate conservation; red: highly conserved). (**B**) mNG-MinC fluorescence, chlA autofluorescence and merged micrographs of *Anabaena* expressing mNG-MinC from P_{ntcA} integrated into the neutral chromosomal *thrS2* sites. Fluorescence intensity profiles show high levels of MinC at the cell poles and midcell in diving cells and only at the cell poles in vegetative cells. (**C**) Quantifications from pelleting assay shown in Fig. 6F showing values for CorM and MinC separately. Different MinC protein concentrations are indicated below. Error bars represent the mean ± SD of two independent replicates (n=2). (**D**) Sequence logo showing the conservation of aa residues within the N-terminal α-helix of *Nostocales* cyanobacteria. A multiple sequence alignment of all tested MinC sequences from cyanobacteria and other bacteria is shown in Fig. S9. (**E**) Quantifications from pelleting assay

987   shown in Fig. 6G showing values for CorM. Both MinC helix mutants, $MinC^{S50A+L55D}$ and $^{\Delta46-67}MinC$ lost the ability to depolymerize CorM filaments. A synthesized peptide mimicking MinC's N-terminal α-helix had no influence on CorM polymerization, indicating that other aspects of the interaction of CorM with MinC are essential for active depolymerization. Error bars represent the mean ± SD of three independent replicates (n=3). (**F**) CorM ATPase activity assay with or without indicated MinC concentrations resolved over time, showing that MinC acts on CorM by decreasing its enzymatic activity. Data represents the mean values for each data point from three independent replicates (n=3). (**G**) B2H assays testing for the interaction of CorM with MinC helix mutants ($MinC^{S50A+L55D}$ and $^{\Delta46-67}MinC$) fused to either the T25 or T18 subunit. Results show that CorM interaction with MinC is lost in both MinC helix mutants. Negative control: CorM-T25 with empty pUT18C plasmid. Positive control: Zip/Zip control. Error bars represent the mean ± SD of three independent replicates (n=3).

Acknowledgements:

We thank all members of the Martin Loose lab at ISTA for helpful discussions and Marko Kojic for critical reading of the manuscript. This research was supported by the Scientific Service Units (SSU) of ISTA through resources provided by the Imaging & Optics Facility (IOF), the Scientific Computing (SciComp) and the Electron Microscopy Facility (EMF), as well as the Lab Support Facility (LSF). This project has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No.101034413 awarded to BLS as well as an ERC grant (ActinID, 101076260) from the European Union awarded to FKMS. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

We are grateful for Antonia Herrero (Sevilla University) for sharing her extensive BACTH plasmid library and other plasmids as well as cyanobacterial strains. Likewise, we would like

to thank Tal Dagan and Fabian Nies (both Kiel University) for sharing cyanobacterial strains

and plasmids and for valuable discussions.

We would further like to express our gratitude to Nicolas Sapay and Alexis Michon for providing

the Amphipaseek code, which enabled us to perform our large-scale amphipathic helix screen

of cyanobacterial CorR proteins. Finally, we also want to thank Jesse Hansen for advice in

cryo-EM data processing.

1020

1021   **Author contributions:**

1025

1026   **Competing interests:**

1027   The authors declare no competing interests.

1028

Data availability:

The electron microscopy density maps of the different CorM filament structures have been

deposited in the Electron Microscopy Data Bank under accession codes: **EMD-54060** (CorM

in the presence of CorR), **EMD-54061** (CorM) and **EMD-54062** ($^{\Delta 1-40}$CorM). The model of the

CorM filament refined into EMD-54060 has been deposited in the Protein Data Bank under

accession code: **PDB 9RMI** (Extended PDB ID pdb_00009RMI).

Source Data for phylogenetic analysis will be available in a Mendeley repository under

accession code 10.17632/d6hmm7bjnt.1 upon publication, but is temporarily accessible from:

https://data.mendeley.com/preview/d6hmm7bjnt?a=9155e04d-88b1-4719-8ec6-

0a300aa479f6

Supplementary Movies will be available from an ISTA repository but are temporarily available

from the authors upon request. All other data are either directly enclosed to this manuscript or

available from the authors upon reasonable request.

Material and methods:

Bacterial strains and growth conditions

Escherichia coli strain DH5α was used for routine cloning procedures and chemically competent *E. coli* were transformed with plasmid DNA by the standard heat shock procedure. BL21 (DE3) (NEB, USA) or C41 (DE3) (Merck, USA) were used for heterologous recombinant protein expression. Strain BTH101 was used for bacterial adenylate cyclase two-hybrid system (BACTH) analysis. All *E. coli* strains listed in Supplementary Table 1 were grown in standard LB or TB medium containing the appropriate antibiotics at standard concentrations. *Anabaena* was a gift from Tal Dagan (Kiel University, Germany) and cells were routinely grown photoautotrophically in BG11 medium at constant illumination with an intensity of 1200 Lux (using a Osram T8 L 15W/840 LUMILUX Cool White G13 light bulb) or at RT with ambient light exposure for extended culturing. When appropriate, liquid growth medium or agar plates (containing 1.5 % agarose) were supplemented with 50 µg/ml neomycin or 5% sucrose.

Plasmid construction

All plasmids generated in this study (see Supplementary Table 1) were constructed using an in-house Gibson assembly mix or the Q5® Site-Directed Mutagenesis Kit (NEB, USA). Primers employed for plasmid construction are listed in Supplementary Table 2. In each case, plasmid sequence integrity was verified by Sanger sequencing by Microsynth Austria GmbH (Austria). In addition, we are grateful for Antonia Herrero, Fabian Nies and Tal Dagan for a wealth of plasmids, particularly for the BACTH plasmid library from Antonia Herrero.

pBAD33.1-Km: The nptII cassette was amplified from pRL25C using primers (oBS218+oBS219) and integrated into PCR-amplified pBAD33.1 (using primers oBS216+oBS217). pBS1: *corM* was amplified from *Anabaena* genomic DNA using primers oBS8+oBS9 and then integrated into HindIII+BamHI-digested pKNT25 by Gibson assembly. pBS2: *corM* was amplified from *Anabaena* genomic DNA using primers oBS10+oBS11 and then integrated into BamHI+EcoRI-digested pKT25 by Gibson assembly. pBS3: *corM* was amplified from *Anabaena* genomic DNA using primers oBS8+oBS9 and then integrated into

HindIII+BamHI-digested pUT18 by Gibson assembly. pBS4: *corM* was amplified from

Anabaena genomic DNA using primers oBS12+oBS13 and then integrated into BamHI+EcoRI-

digested pUT18C by Gibson assembly. pBS9: corR was amplified from *Anabaena* genomic

DNA using primers oBS18+oBS19 and then integrated into HindIII+BamHI-digested pKNT25

by Gibson assembly. pBS10: *corR* was amplified from *Anabaena* genomic DNA using primers

oBS20+oBS21 and then integrated into BamHI+EcoRI-digested pKT25 by Gibson assembly.

pBS11: *corR* was amplified from *Anabaena* genomic DNA using primers oBS18+oBS19 and

then integrated into HindIII+BamHI-digested pUT18 by Gibson assembly. pBS12: *corR* was

amplified from *Anabaena* genomic DNA using primers oBS22+oBS23 and then integrated into

BamHI+EcoRI-digested pUT18C by Gibson assembly. pBS41: *corR* was amplified from

Anabaena genomic DNA using primers oBS69+oBS70 and integrated into PCR-amplified

pTB146 (using primers oBS59+oBS60) by Gibson assembly. pBS47: *parR* from the *E. coli*

NR1 plasmid was amplified from a gBlock (Twist Biosciences, USA) using primers

oBS81+oBS82 and ligated into PCR-amplified pTB146 (using primers oBS59+oBS60) by

Gibson assembly. pBS71: *corM* was amplified from *Anabaena* genomic DNA using primers

oBS62+oBS126 and ligated into PCR-amplified pTB146 (using primers oBS59+oBS60) by

Gibson assembly. pBS75: *parM* from the *E. coli* NR1 plasmid was amplified from a gBlock

(Twist Biosciences, USA) using primers oBS80+oBS130 and ligated into PCR-amplified

pTB146 (using primers oBS59+oBS60) by Gibson assembly. Plasmid pBS80 was constructed

using Gibson assembly to ligate three PCR-amplified fragments. The vector was amplified from

plasmid pTB146 using primers oBS59+oBS150, mNeonGreen (mNG) was amplified from a

gBlock (Twist Biosciences, USA) using primers oBS151+oBS119 and *corM* was amplified from

Anabaena genomic DNA using primers oBS120+oBS62. pBS103 was generated by annealing

of primers oBS199+oBS200 and ligating the annealed primers into AarI-digested pCpf1b using

T4 ligase (NEB, USA). pBS105 was generated by annealing of primers oBS203+oBS204 and

ligating the annealed primers into AarI-digested pCpf1b using T4 ligase (NEB, USA). pBS108

was constructed by Gibson assembly, ligating two PCR-amplified fragments - amplified from

Anabaena genomic DNA using primers oBS205+oBS206 and oBS207+oBS208 - into BamHI-

digested pBS105. Plasmid pBS135 was constructed using Gibson assembly to ligate three

PCR-amplified fragments. The vector was amplified from plasmid pTB146 using primers

oBS59+oBS150, *parM* from the *E. coli* NR1 plasmid was amplified from a gBlock (Twist

Biosciences, USA) using primers oBS284+oBS285 and mNG was amplified from a gBlock

(Twist Biosciences, USA) using primers oBS154+oBS155. pBS136 was generated by

annealing of primers oBS286+oBS287 and ligating the annealed primers into AarI-digested

pCpf1b using T4 ligase (NEB, USA). pBS138 was constructed by Gibson assembly, ligating

two PCR-amplified fragments - amplified from *Anabaena* genomic DNA using primers

oBS290+oBS291 and oBS292+oBS293 - into BamHI-digested pBS136. pBS154 was

generated by annealing of primers oBS401+oBS402 and ligating the annealed primers into

AarI-digested pCpf1b using T4 ligase (NEB, USA). pBS159 was generated by Gibson

assembly-based ligation of two PCR products into PCR-amplified (using primers

oBS143+oBS144); one amplifying *corR* from *Anabaena* genomic DNA using primers

oBS145+oBS147 and the 2nd amplifying *mScarlet-I* from a gBlock (Twist Biosciences) using

primers oBS148+oBS149. pBS166 was generated by Gibson assembly-based ligation of three

PCR products into BamHI-digested pBS138; one amplifying a synthetically optimized *ntcA*

promoter (P$_{ntcA}$)[80] using primers oBS294+oBS297 and pCSL135[81] as a template, the 2nd one

amplifying *corR* using primers oBS338+oBS345 and *Anabaena* genomic DNA as a template

and the 3rd amplifying SCFP3 using primers oBS346+oBS347 and a gBlock (Twist

Biosciences, USA) as a template. pBS184 was generated by Gibson assembly-based ligation

of three PCR products into BamHI-digested pBS138, one amplifying P$_{ND}$ using primers

oBS294+oBS297 and pCSL135 as a template, the 2nd one amplifying *corR* using primers

oBS338+oBS376 and *Anabaena* genomic DNA as a template and the 3rd amplifying mNG

using primers oBS377+oBS296 and a gBlock (Twist Biosciences, USA) as a template. pBS190

was generated by annealing of primers oBS385+oBS386 and ligating the annealed primers

into AarI-digested pCpf1b using T4 ligase (NEB, USA). pBS191 was generated by Gibson

assembly-based ligation of three PCR products into BamHI-digested pBS190. The first product

amplified 1000 bp upstream of *corM* using primers oBS205+oBS117 and *Anabaena* genomic

DNA as a template. The 2nd product was amplified sequentially by first amplifying mNG using

primers oBS118+oBS119 and a gBock as a template and then using this product as a new

template for the following PCR using primers oBS118+oBS378. The 3rd PCR amplified parts

of the *corM* open reading frame (ORF) using primers oBS388+oBS389 and *Anabaena*

genomic DNA as a template. pBS196 was generated by Gibson assembly-based ligation of

three PCR products into BamHI-digested pBS154, one amplifying a ~1000 bp fragment that

encompassed *corR* and parts of *corM* using primers oBS403+oBS404 and *Anabaena* genomic

DNA as a template, the 2nd one amplifying mNG using primers oBS405+oBS406 and pBS184

as a template and the 3rd amplifying 1000 bp downstream of *corR* using primers

oBS407+oBS208 and *Anabaena* genomic DNA as a template. pBS201: *minD* was amplified

from *Anabaena* genomic DNA using primers oBS415+oBS416 and then ligated into

HindIII+BamHI-digested pKNT25. pBS202: *minD* was amplified from *Anabaena* genomic DNA

using primers oBS417+oBS418 and then ligated into BamHI+EcoRI-digested pKT25. pBS203:

minD was amplified from *Anabaena* genomic DNA using primers oBS415+oBS416 and then

ligated into HindIII+BamHI-digested pUT18. pBS204: *minD* was amplified from *Anabaena*

genomic DNA using primers oBS419+oBS420 and then ligated into BamHI+EcoRI-digested

pUT18C. pBS205: *minE* was amplified from *Anabaena* genomic DNA using primers

oBS421+oBS422 and then ligated into HindIII+BamHI-digested pKNT25. pBS206: *minE* was

amplified from *Anabaena* genomic DNA using primers oBS423+oBS424 and then ligated into

BamHI+EcoRI-digested pKT25. pBS208: *minD* was amplified from *Anabaena* genomic DNA

using primers oBS425+oBS426 and then ligated into BamHI+EcoRI-digested pUT18C.

pBS213: *cdv3* was amplified from *Anabaena* genomic DNA using primers oBS433+oBS434

and then ligated into HindIII+BamHI-digested pKNT25. pBS214: *cdv3* was amplified from

Anabaena genomic DNA using primers oBS435+oBS436 and then ligated into BamHI+EcoRI-

digested pKT25. pBS231 was generated by site-directed mutagenesis using primers

oBS60+oBS464 and pBS41 as a template. pBS233 generated by site-directed mutagenesis

using primers oBS60+oBS475 and pBS71 as a template. pBS316: *minC* was amplified from

Anabaena genomic DNA using primers oBS648+oBS649 and ligated into PCR-amplified

pTB146 (using primers oBS59+oBS60) by Gibson assembly. pBS326 was generated by

Gibson assembly-based ligation of three PCR products into BamHI-digested pBS138, one

amplifying P_{ND} using primers oBS294+oBS229 and pCSL135 as a template, the 2nd one

amplifying mNG using primers oBS118+oBS119 and a gBlock as a template and the 3rd

amplifying *minC* using primers oBS508+oBS669 and *Anabaena* genomic DNA as a template.

pBS338: $^{\Delta150-162}$*corR* was amplified from *Anabaena* genomic DNA using primers

oBS20+oBS700 and then ligated into BamHI+EcoRI-digested pKT25. pBS339: $^{\Delta150-162}$*corR*

was amplified from *Anabaena* genomic DNA using primers oBS22+oBS701 and then ligated

into BamHI+EcoRI-digested pUT18C. pBS346 was generated by site-directed mutagenesis

using primers oBS59+oBS711 and pBS41 as a template. pBS353 was generated by site-

directed mutagenesis using primers oBS720+oBS721 and pBS316 as a template. pBS354

was generated by site-directed mutagenesis using primers oBS722+oBS723 and pBS316 as

a template. pBS386 was generated by site-directed mutagenesis using primers

oBS720+oBS721 and pCSAV269 (a gift from Antonia Herrero, Sevilla University) as a

template. pBS387 was generated by site-directed mutagenesis using primers

oBS722+oBS723 and pCSAV269 as a template. pBS396 was generated by Gibson assembly-

based ligation of two PCR products into BamHI-digested pBS103. The first product amplified

1000 bp upstream of *corM* and *mNG* using primers oBS205+oBS119 and pBS191 as a

template. The 2nd product amplified 882 bp downstream of codon 40 of *corM* using primers

oBS651+oBS389 and pBS191.

1174

Strain construction

Transformation into *Anabaena* WT or mutant strains was achieved using a triparental mating,

essentially as described before in [47], but with the following modification: instead of *E. coli*

HB101, we used *E. coli* DH5α. Over-night grown *E. coli* and *Anabaena* were mixed and plated

onto nitrocellulose filters (0.45 µm pore size) placed on BG11 plates supplemented with 5%

LB pre-mixed powder. Cells were incubated for 6 - 8 hours at 30 °C under low light and then

transferred to BG11 plates containing 50 µg/ml neomycin (BG11+Nm) and exposed to full light.

Colonies routinely appeared after 10 - 12 days. These were picked and re-streaked on

BG11+Nm plates to isolate single colonies and eliminate residual *E. coli* for several rounds.

For heterologous expression of fluorescently labeled proteins, newly emerging colonies were

screened by fluorescence microscopy and propagated on BG11+Nm plates instead.

Strain BS326 was generated by Cpf1-based heterologous insertion of the P_{ntcA}-*mNG-minC*

construct into the *thrS2* open reading frame (encoding threonine-tRNA ligase; NCBI Ref. Seq.:

WP_010998854.1), a chromosomal neutral site[20,74]. Double homologous recombination using

~1000 bp flanking regions cloned into plasmid pBS326 replaced the *thrS2* segment encoding

residues Q188–F228 with P_{ntcA}-*mNG-minC*. Complete segregation was verified by colony PCR

using construct-specific primers.

For colony PCR, colonies were either picked with a pipette tip, briefly spotted onto a fresh

BG11+Nm plate, and then resuspended directly in the PCR reaction mixture, or resuspended

in 10 µl sterile PBS, of which 1 µl was used for each 10 µl PCR reaction.

For the Δ*corMR* (strain BS108), plasmid pBS108 was conjugated into *Anabaena* WT. Arising

colonies were re-streaked twice onto fresh BG11+Nm plates. Deletion of *corM-corR* was

confirmed by colony PCR using site- and gene-specific primers, followed by Sanger

sequencing of the encompassing regions.

Strain BS191, expressing mNG-CorM as the sole copy of CorM, was generated by conjugating

plasmid pBS191 into *Anabaena* WT. Resulting colonies were picked, re-streaked on

BG11+Nm plates, screened for mNG-CorM fluorescence by microscopy, and verified by

Sanger sequencing of a PCR-amplified fragment encompassing the P_{corMR}-*mNG-corM-corR*

region from a single colony. Strain BS396 was obtained in the same manner as BS191, using

plasmid pBS396 instead of pBS191. Sanger sequencing confirmed the presence of P_{corMR}-

mNG-$^{\Delta1–40}$corM-corR.

In all cases, curing of pCpf1b was performed by sucrose counter-selection. For this, strains

were grown in BG11 medium, sonicated in a VWR Ultrasonic Cleaner USC-TH for 10 min at

maximum intensity, and then plated in 1:10 serial dilutions on BG11 plates supplemented with

5% sucrose. Arising colonies were screened by streaking on BG11 and BG11+Nm plates;

those that failed to grow on BG11+Nm (indicating loss of pCpf1) were selected for further
experiments.

1212

BACTH analysis

For protein-protein interaction analysis, the BACTH system (Euromedex, France) was
employed. Genes were cloned into the expression vectors pKNT25, pKT25, pUT18, and
pUT18C using GIBSON assembly, resulting in N- or C-terminal translational fusions to the T25
or T18 subunit of the *E. coli* adenylate cyclase. *E. coli* BTH101 cells were co-transformed with
about 5 ng of the respective plasmid combinations and then plated on LB agar containing 200
µg/ml X-gal, 0.5 mM IPTG, 100 µg/ml ampicillin (Amp), and 50 µg/ml Kanamycin (Km). As a
positive control, plasmids pKT25-zip and pUT18C-zip and as a negative control empty plasmid
pUT18 and a pKNT25 plasmid containing a test gene were also co-transformed into BTH101
cells. The plates were incubated at RT for 2-3 days. To then measure beta-galactosidase as a
surrogate measure to quantify protein-protein interaction strengths, a modified protocol from
Karimova et al.[82] was employed. For each combination, three independent colonies were
picked and grown in 300 µl LB supplemented with 0.5 mM IPTG, Amp, and Km in a 2 ml 96-
well deep well plate for 2-3 days at RT and 1000 rpm in an orbital microplate shaker.
Afterwards, 50 µl of cells were diluted with 50 µl 2x M63 medium and OD_{595} values were
recorded in a 96-well plate. Cells were then lysed by adding 10 µl of PopCulture® Reagent
(Merck, USA) supplemented with 10µl/ml Lysonase™ Bioprocessing Reagent (Merck, USA).
Lysis was conducted for at least 30 min at RT and rigorous shaking with 650 rpm in the 96-
well plate on an orbital microplate shaker (We find that even hour-long lysis did not decrease
experimental readout). Subsequently, 20 µl of lysate was added to 105 µl of PM2 assay
medium supplemented with 100 mM β-mercaptoethanol in a new 96-well plate and incubated
at RT with mild shaking for 20 min before the reaction was stopped by adding 50 µl of 1 M
Na_2CO_3. Finally, OD_{405} values were recorded, and beta-galactosidase activity was calculated
as described in the BACTH System Kit (Euromedex).

1237

Microscopy

Experiments were performed using a Nikon Eclipse Ti2 microscope equipped with iLas2

(GATACA) 360° / Ring / Azimuthal TIRF module, a TI2-N-ND-P Perfect Focus System and

either a CFI Apochromat TIRF 60x oil immersion objective (NA 1.49, WD 0.13 mm) or a CFI

Plan Apo Lambda D 100x oil immersion objective (NA 1.45, WD 0.13 mm).

Fluorophores and fluorescent proteins were excited using the Omicron LightHUB Ultra laser

system for TIRF imaging and Fluorescence Recovery After Photobleaching (FRAP), equipped

with 405nm, 488nm, 561nm and 640nm lasers. Emitted fluorescence signals were filtered

using a Nikon 525/50 ET bandpass and a 635 nm long pass filter for dual-color imaging. For

the dual-color experiments, a Cairn TwinCam camera splitter equipped with a TwinCam Cube

dichroic mirror for green/red,far-red was used in addition.

For epifluorescence microscopy, LED-FITC-A Filter Cube EX 474/27, EM 525/45 and LED-

Cy5-A Filter Cube EX 628/40, BA 692/40 filter cubes were used, and bright field images were

obtained with a CoolLED pE300 White MB light source. The microscope was operated using

the NIS elements AR (v.5.42.04) software and images were recorded using one or two Andor

iXon Life 888 Back Illuminated EMCCD cameras.

High-resolution spinning disk confocal microscopy was conducted using a Nikon Ti2 CSU W1-

01 SoRa+NIR microscope, equipped with a Nipkow disk-based Super-resolution via optical

Re-assignment (SoRa) module and a Nikon CFI Plan Apochromat 100× silicone immersion

objective (NA 1.35, WD 0.3 mm). Excitation was provided by an Omicron LightHUB UltraN

laser system with 405 nm, 488 nm, and 561 nm lasers, guided through a confocal quad-band

dichroic mirror (405/488/561/640 nm). Emission signals were collected through a quad-band

emission filter (440/521/607/700 nm) and imaged using a Teledyne Photometrics BSI scientific

CMOS camera.

1262

DAPI staining

For DAPI staining, cells grown on BG11 agar plates were re-suspended in PBS supplemented with 10 µg/ml DAPI (final), incubated for 1-2 min and then spotted and immobilized on BG11 2% agar pads prior to imaging.

Time lapse microscopy of mNG-CorM in Anabaena cells

For time-resolved tracking of mNG-CorM filaments in *Anabaena* cells of strain BS191, cells were grown on BG11 plates, re-suspended in sterile BG11 and spotted on BG11 pads containing 1.5 % low melting agarose (TopVision Low Melting Point Agarose, Thermo Fisher Scientific) supplemented with 10 mM $NaHCO_3$ in a Grene Frame (65 µl; Thermo Fisher Scientific) to avoid evaporation. Cells were visualized in a Nikon Eclipse Ti2 microscope equipped with iLas2 system and mNG-CorM filaments were recorded in epifluorescence mode (50 ms exposure, 10 % blue LED light intensity). Cells were grown at RT and illuminated using a custom-made plant light equipped with three LEDs emitting blue (450 nm), red (655 nm), and far-red (725 nm) light. The light intensities at the sample position were 285.20 mW/m² (blue), 41.25 mW/m² (red), and 38.65 mW/m² (far-red). Images were acquired every 15 minutes with the perfect focus system enabled to maintain the correct focal plane.

Protein purification

Purification of CorM

CorM with three N-terminal Gly (GGG-CorM) to make it amenable for sortagging was purified as a N-terminal His_6-SUMO fusion protein expressed from pBS71 in *E. coli* C41 (DE3) cells. For this, cells were grown at 37 °C, 220 rpm in Terrific Broth (TB) supplemented with Amp and expression was induced at an OD_{600} 0.5-0.7 with 0.5 mM isopropyl-β-thiogalactopyranoside (IPTG) for 4 h at 37 °C. Cells were harvested by centrifugation (8,000 x *g,* 20 min at 4 °C) and cell pellets were scooped out and stored in 50 ml reaction tubes at -70 °C. Cell pellets were resuspended in HEPES lysis buffer (HLB; 50 mM HEPES, 150 mM KCl, 5 mM $MgCl_2$, 1 mM DTT, 25 mM imidazole and 10 % glycerol) supplemented with 1x EDTA-free protease inhibitor

cocktail (Roche, Switzerland). Cells were lysed by sonication using Q700 Sonicator equipped

with a 1/2 in horn tip (amplitude 40, 1 sec on + 1 sec off with 10 min on time) and subsequently

incubated with 10 µl DNase I (2500 U/ml; Thermo Fisher Scientific, USA) per every 1 L of initial

bacterial culture volume for 15 min. The lysate was then centrifuged for 30 min at 60,000 x g

at 4 °C and the supernatant transferred to HisPur™ Ni-NTA Resin (1 ml per 1 L of initial

bacterial culture volume; Thermo Fisher Scientific, USA) pre-equilibrated with HLB in

incubated with mild agitation for 1-2 h at 4 °C. The mixture was loaded onto a gravity flow

column, the flow through was discarded and the resin was washed with 3x 20 ml HLB. Proteins

were eluted with HEB (HLB with 250 mM imidazole) in 1 ml steps. Eluates were briefly tested

for protein content using a quick and dirty Bradford (Bio-Rad Protein Assay Dye Reagent;

BioRad, USA) assay by adding 2 µl of each elution fraction to 200 µl Bradford assay solution.

Protein fractions that resulted in a discernible blue color shift were pooled and dialyzed over

night at 4 °C against HLB in Pur-A-Lyzer™ Maxi 6000 dialysis tubes (Sigma Aldrich, USA)

supplemented with His-tagged SUMO protease (Ulp1) (ratio of 1:100). The next day, reverse

affinity was performed to remove His-SUMO and His-Ulp1. For this, the solution was subjected

to Ni-NTA (half the volume as used for the initial purification) pre-equilibrated in HLB, incubated

for 1 h at 4 °C and mild rotation. The resin was applied to a gravity flow column and the flow

through was collected followed by 3x washing with 1x column volume (CV) of HLB. The flow

through and the wash fractions were pooled, concentrated using a Vivaspin 20 10KMWCO

centrifugal concentrator (Sartorius, Germany) and subjected to size-exclusion chromatography

(SEC) on an Äkta pure system using a HiLoad 16/600 Superdex 75 PG column (Cytiva, USA)

equilibrated with HEPES storage buffer (HSB, HLB without imidazole). Protein-containing

fractions were pooled, concentrated using another Vivaspin 20 10KMWCO centrifugal

concentrator and then finally flash-frozen in liquid nitrogen and then stored at -70 °C.

1315

Purification of $^{\Delta 1-40}$CorM

$^{\Delta 1-40}$CorM with three N-terminal Gly (GGG-$^{\Delta 1-40}$CorM) to make it amenable for sortagging was

purified as a N-terminal His$_6$-SUMO fusion protein expressed from pBS233 in *E. coli* C41 (DE3)

cells for 3 h at 37 °C with 0.5 mM IPTG. Cell pellets were re-suspended in HLB and lysed as

described for CorM. Cell-free lysates were then subjected to Ni-NTA purification using the Äkta

pure system equipped with a HisTrap^{TM} HP 5 ml column and proteins were eluted in HEB.

Pooled fractions were then subjected to Ulp1 cleavage and dialyzed overnight at 4 °C against

HLB. After subsequent reverse affinity, the flow through and two wash fractions were pooled,

concentrated using a Vivaspin 20 10KMWCO centrifugal concentrator and subjected to

another round of reverse affinity due to remaining contamination with His6-SUMO. The flow

through and two wash fractions were then pooled, and protein purity was confirmed by SDS

polyacrylamide gel electrophoresis (SDS-PAGE) followed by flash-freezing in liquid nitrogen

and storage at -70 °C.

1329

Purification of EcParM from *E. coli* R1 plasmid

ParM from *E. coli* NR1 plasmid was expressed from pBS75 with three N-terminal Gly (GGG-

ParM) in C41 (DE3) for 3 h at 37 °C with 0.5 mM IPTG. Lysis and subsequent Ni-NTA-based

purification were done as described for CorM. After the reverse affinity, ParM was concentrated

using a Vivaspin 20 10KMWCO centrifugal concentrator and subjected to buffer exchange to

HSB using a PD-10 desalting column. ParM was again concentrated using a Vivaspin 20

10KMWCO centrifugal concentrator and flash-frozen in liquid nitrogen and then stored at -70

°C.

1338

Purification of CorR

CorR was expressed as a N-terminal His$_6$-SUMO fusion protein from pBS41 in BL21 (DE3) for

3 h at 37 °C with 0.5 mM IPTG or in C41 (DE3) for 18 h at 18 °C with 0.5 mM IPTG. Lysis and

subsequent Ni-NTA-based purification were done as described for GGG-CorM. After the

reverse affinity, flow through and wash fractions were pooled, concentrated using a Vivaspin

20 10KMWCO centrifugal concentrator and subjected to anion exchange chromatography

(AIEX) on Äkta pure system using HiTrap 1ml Q HP anion exchange chromatography column

(Cytiva, USA) equilibrated with no salt buffer (50 mM HEPES pH 7.4, 5 mM MgCl$_2$, 10 %

glycerol). For this, CorR was diluted 1/5 in no salt buffer and subjected to a linear gradient of

no salt and high salt buffer (50 mM HEPES pH 7.4, 1 M KCl, 5 mM $MgCl_2$, 10 % glycerol).

CorR elutes as numerous, not well-defined peaks, likely representing different oligomeric

states of CorR. CorR-containing fractions were pooled, concentrated using a Vivaspin 20

5KMWCO centrifugal concentrator, flash-frozen in liquid nitrogen and stored at -70 °C.

1352

1353   **Purification of $^{\Delta 1-10}$CorR**

1354 $^{\Delta 1-10}$CorR was expressed as a N-terminal His_6-SUMO fusion protein from pBS231 in BL21

1355   (DE3) for 3 h at 37 °C with 0.5 mM IPTG. Lysis and subsequent purification steps were done

1356   as described for CorR.

1357

1358   **Purification of $^{\Delta 150-162}$CorR**

1359 $^{\Delta 150-162}$CorR was expressed as a N-terminal His_6-SUMO fusion protein from pBS346 in BL21

1360   (DE3) for 16 h at 18 °C with 0.5 mM IPTG. Lysis and subsequent purification steps were done

1361   as described for CorR.

1362

Purification of ParR

ParR from *E. coli* NR1 plasmid was expressed as a N-terminal His_6-SUMO fusion protein from

pBS47 in BL21 (DE3) for 3 h at 37 °C with 0.5 mM IPTG. After overnight cleavage with Ulp1,

ParR was subjected to SEC using HiLoad 16/600 Superdex 75 PG equilibrated with HSB.

Protein containing fractions were pooled, supplemented with 10 mM imidazole and subjected

to reverse affinity. The flow through and wash fractions were pooled, concentrated using a

Vivaspin 20 5KMWCO centrifugal concentrator, flash frozen in liquid nitrogen and stored at -

70 °C.

1371

Purification of MinC

MinC with three N-terminal Gly (GGG-MinC) to make it amenable for sortagging was purified

as an N-terminal His_6-SUMO fusion protein expressed from pBS316 in *E. coli* BL21 (DE3) or

C41 (DE3) cells for 3 h at 37 °C with 0.5 mM IPTG. Lysis and subsequent Ni-NTA-based

purification were done as described for CorM. After the reverse affinity, MinC was concentrated

using a Vivaspin 20 10KMWCO centrifugal concentrator and was then subjected to SEC using

HiLoad 16/600 Superdex 75 PG column equilibrated with HSB. One major peak was detected

and pooled and proteins subjected to AIEX as described for CorR with a gradient from no salt

to high salt buffer. A single peak was identified, pooled, flash-frozen in liquid nitrogen and

stored at -70 °C.

1382

Purification of MinC$^{S50A+L55D}$

MinC$^{S50A+L55D}$ with three N-terminal Gly (GGG-MinC$^{S50A+L55D}$) to make it amenable for

sortagging was purified as a N-terminal His$_6$-SUMO fusion protein expressed from pBS353 in

E. coli BL21 (DE3) cells for 4 h at 37 °C with 0.5 mM IPTG. Lysis and subsequent Ni-NTA-

based purification were done as described for CorM. After the reverse affinity, MinC$^{S50A+L55D}$

was concentrated using a Vivaspin 20 10KMWCO centrifugal concentrator and was then

subjected to AIEX as described for CorR. Due to remaining contamination with His6-SUMO,

another reverse affinity step was included, the flow through and wash steps were collected,

pooled and then dialyzed overnight at 4 °C against HSB in Pur-A-Lyzer™ Maxi 6000 dialysis

tubes. Afterwards, MinC$^{S50A+L55D}$ was concentrated using a Vivaspin 20 10KMWCO centrifugal

concentrator, flash-frozen in liquid nitrogen and stored at -70 °C.

1394

1395   **Purification of $^{\Delta46-67}$MinC**

1396 $^{\Delta46-67}$MinC with three N-terminal Gly (GGG-$^{\Delta46-67}$MinC) to make it amenable for sortagging was

1397   purified as a N-terminal His$_6$-SUMO fusion protein expressed from pBS354 in *E. coli* BL21

1398   (DE3) cells for 4 h at 37 °C with 0.5 mM IPTG. $^{\Delta46-67}$MinC was purified as described for

1399   MinC$^{S50A+L55D}$.

1400

Peptide labelling

For labelling of peptide for subsequent sortagging reactions, we obtained dried CLPETGG peptides from Biomatik (USA). 1.5 mg of the peptide was dissolved in 50 mM HEPES pH 7.4 to a final concentration of 100 mM. Alexa Fluor™ 488 (Alexa488; Thermo Fisher Scientific) C5 Maleimide and Sulfo-Cyanine5 maleimide (Lumiprobe, Germany) were dissolved in DMSO to obtain a 150 mM solution. For extended storage, dissolved dyes were stored at -70 °C. A roughly 2x molar excess of TCEP was added to the reconstituted peptide (i.e., 10 µl of peptide and 5 µl of 0.4 M TCEP) to obtain a 66 mM peptide solution. Afterwards, a roughly 3x molar excess of dye was added to the peptide+TCEP mix (i.e., 20 µl of dye was added leading to a peptide concentration of roughly 28 mM and a dye concentration of roughly 86 mM). The reaction was then incubated over night at RT protected from light. The next day, 2 µl of 2-mercaptoethanol was added to quench unreacted maleimide dye. The 25 mM peptide solution was then flash-frozen in liquid nitrogen and stored at -70 °C.

Protein labelling - Sortagging

For Sortase-based N-terminal fluorescence labelling, we routinely prepare 1 or 2 ml reactions of 50 µM GGG-CorM, 0.5 mM labelled peptides (Alexa488-LPETGG or Cy5-LPETGG) and 10 µM Sortase 7M in HSB, which are then incubated overnight at 4 °C in the dark with gentle rotation. The next day, 3 µl of 2-mercaptoethanol was added to the reaction to quench all unreacted dye and incubated for 15 min at 4 °C. The reactions were supplemented with 25 mM imidazole (final) and 300 µl pre-equilibrated Ni-NTA slurry was added. After incubation for 60 min at 4 °C with mild rotation, the solution was applied to a gravity flow column and the flow through was collected. The Ni-NTA was washed 2x with 1x column volume HDB and pooled with the flow through. To remove unbound labelled peptide and to exchange the buffer to HSB, Cy5-CorM was then subjected to a PD-10 desalting column (Cytiva, USA) and eluted protein was concentrated using a Vivaspin 2 10KMWCO centrifugal concentrator (Sartorius, Germany). For Alexa488-CorM, unbound labelled peptide and imidazole traces were removed by size-exclusion chromatography on a Superdex 75 Increase 10/300 GL (Cytiva, USA) using

HSB. Protein-containing peaks were then again concentrated using a Vivaspin 2 10KMWCO

centrifugal concentrator and in both cases protein concentration was measured using

NanoDrop One and labelling efficiency was calculated based on the ratio of the concentration

of fluorescence dye (equals labelled protein) to the protein concentration. In both cases,

labelling efficiency was about 30 %. Finally, labelled proteins were flash-frozen in liquid

nitrogen and stored at -70 °C.

1435

Piranha treatment of coverslips for TIRF microscopy

TIRF-suitable glass coverslips (18X18 mm NR 1.5 high precision cover glasses; Marienfield,

Germany) in a custom-made Teflon holder were subjected to piranha solution (mixture of 50

ml 30 % H_2O_2 and 150 ml H_2SO_4) for 2 h at RT until the solution cooled down to RT. Cover

slips were then washed 10x with 500 ml MiliQ H_2O, sonicated in a sonicator bath (VWR

Ultrasonic Cleaner USC - TH) for 20 min and washed again 4x with 500 ml MiliQ H_2O and

finally stored at RT in 500 ml MiliQ H_2O for no longer than 2 weeks. For TIRF microscopy,

coverslips were dried by compressed air and reaction chambers made of 0.5 ml reaction tubes

(cut in half to create a chamber without the conical end) were glued onto the coverslips with a

UV-reactive glue (Norland Optical Adhesive 63) for 2 min under UV light exposure.

1446

Preparation of multilaminar vesicles and small unilaminar vesicles

For pelleting assays and TIRF microscopy experiments we used DOPC (1,2-dioleoyl-sn-

glycero-3-phosphocholine) and DOPG (1,2-dioleoyl-sn-glycero-3-phospho-(1′-rac-glycerol)) *E.*

coli polar lipids (Avanti, USA) dissolved in chloroform (stock concentration 2.5 mg) at a ratio of

67:33 mol %. Lipids were mixed in a glass vial and then dried in a stream of nitrogen gas.

Residual chloroform was removed from the lipids by vacuum desiccation for at least 3 h or

overnight. Afterwards, lipids were resuspended by vortexing in HEPES reaction buffer (HRB;

50 mM HEPES [pH 7.4], 150 mM KCl, 5 mM $MgCl_2$) to obtain a 5 mM large multilaminar

vesicles (MLVs) solution. To prepare small MLVs, this solution was then incubated at 37 °C

for 30 min and then subjected to 7x freeze-thaw cycles using a dry ice-isopropanol bath and

hot tap water. Small MLVs were then subjected to centrifugation at 21,000 x g, 4 °C, 30 min

and the supernatant was transferred to a new reaction tube, which was then overlayed with

argon gas and stored at 4 °C until further use. For preparation of small unilaminar vesicles

(SUVs), small MLVs were subjected to tip-sonication (Q700 Sonicator equipped with a 1.6 mm

tip) on ice with the following settings: amplitude 1; 1 s on time, 4 s off time; total 5 min on time.

Afterwards, SUVs were centrifuged for 20 min at 21,000 x g, 4°C and the supernatant was

transferred to a new tube and overlayed with argon gas.

1464

Supported lipid bilayer preparation

For the preparation of supported lipid bilayers (SLBs), 100 µl reactions were prepared in

reaction chambers containing 89 µl HRB, 0.5 mM SUVs and vesicle rupture was induced by

adding 5 mM $CaCl_2$ to the SUVs. For fusion of the ruptured vesicles with the piranha-treated

glass coverslips, chambers were incubated for 20 min at 37 °C followed by extensive washing

with 6x 200 µl HRB to remove unfused vesicles. Chambers were again washed 3x with 200 µl

HRB right before each individual experiment. SLBs were prepared fresh for each experiment

day.

1473

Total Internal Reflections Fluorescence microscopy

For Total Internal Reflections Fluorescence (TIRF) microscopy, 0.75 µM unlabeled CorM or $^{\Delta 1\text{-}}$

^{40}CorM, 0.25 µM Alexa488-CorM or Alexa488-$^{\Delta 1\text{-}40}$CorM, 0.25 µM CorR and an oxygen

scavenger mix composed of 1 µM glucose oxidase from *Aspergillus niger* (Merck, USA), 332

nM catalase from bovine liver (Merck, USA), 11 mM D-glucose, 1 mM DTT and 1 mM Trolox

was used. The equal amount of buffer in the reaction chambers containing the SLBs was

removed and only then all components were added to the chambers. CorM polymerization on

SLBs was induced by the addition of 2.5 mM ATP and CorM polymers were observed by TIRF

imaging. To test for the effect of different CorR concentrations, the same kind of reactions were

prepared but with varying indicated CorR concentrations.

For dual color imaging to study the localization of individual CorM monomers within a CorM

protein filament, a mixture of either 1 µM Alexa488-CorM + 265 pM Cy5-CorM + 0.25 µM CorR

or 0.25 µM Alexa488-CorM + 265 pM Cy5-CorM + 0.75 µM CorM + 0.25 µM CorR was added

to the reaction chambers together with the oxygen scavenger mix. Time-lapse movies were

recorded between 2-5 min after addition of 2.5 mM ATP to the reaction.

To test for the effect of MinC, MinC$^{S50A+L55D}$ and $^{\Delta46-67}$MinC addition on CorM filaments *in vitro*,

CorM filaments were polymerized on SLBs in the presence of CorR and 2.5 mM ATP for about

5 min upon which equal molar amounts of MinC or MinC mutants (i.e., 1 µM) was added to the

chambers during time lapse imaging. Movies were recorded for 3 min each.

1493

1494   ***Fluorescence recovery after photobleaching experiments***

1495   For FRAP analysis, 3 - 6 pre-bleach frames were acquired, followed by photobleaching of a

1496   defined rectangular region of interest (ROI) using 80 % 488 nm laser power and a pixel dwell

1497   time of 100 µs/pixel. Fluorescence recovery within the ROI was subsequently recorded at a

1498   rate of 1 frame per second.

1499

Pelleting assay

Two types of pelleting assays were performed in this study. One did not involve MLVs and

relied on ultracentrifugation-based pelleting of protein polymers while the other employed the

intrinsic binding capacity of proteins to MLVs. Pelleting assays with MLVs were either

performed in a suspension with or without 0.93 mM small MLVs and proteins at indicated

concentration were incubated for 30 min at 30 °C in HRB supplemented with 2.5 mM ATP.

Unless otherwise stated, proteins were supplied at 5 µM. Afterwards, samples were

centrifuged for 20 min at 30 °C, 21,000 x *g*. For experiments without MLVs, proteins were

incubated in HRB with or without 2.5 mM of indicated nucleotide, incubated for 30 min at 30

°C and centrifuged for 45 min, 436,000 x *g* at 30 °C. In both cases, the supernatant (S) of each

reaction was collected and supplemented with an appropriate amount of 6x Laemmli SDS

sample buffer (Thermo Fisher Scientific, USA). The pellet (P) was resuspended in 50 µl 1x

Laemmli SDS sample buffer. Between 15-25 µl of pellet and supernatant samples were applied to SDS-PAGE using SurePAGE™ Bis-Tris 4-20 % gels (GenScript, USA). Upon protein separation, gels were stained with Quick Coomassie® Stain (Serva, Germany) and protein bands were recorded in a ChemiDoc MP or ChemiDoc Go system (BioRad, USA).

ATPase assay

For ATPase assays, we employed the Malachite Green Phosphate Assay Kit (Sigma Aldrich, USA) according to the manufacturer's instructions. Proteins at indicated concentrations were incubated in ATPase reaction buffer (25 mM HEPES [pH 7.4], 150 mM KCl, 1 mM $MgCl_2$) and if appropriate, supplemented with 1 mM ATP or GTP. Samples were incubated at 30 °C measurements were performed either as endpoint measurements after 2 h or in a time course manner (15, 30, 60, 120 min after ATP addition) of 1/8 dilutions and OD_{620} values were recorded in a Spectramax M2e (Molecular Devices, USA) plate reader. As controls, samples containing no nucleotide or nucleotide only (without protein) were included.

Electrophoretic mobility shift (EMSA) assay

For EMSA assays, test DNA sequences were amplified by PCR using Q5 polymerase (NEB, USA) with primer pairs (oBS249+oBS253 for P_{corMR} and oBS627+oBS729 for P_{parMR}) that each contained one unlabeled (oBS253 and oBS729) and one 5' Alexa488-labelled primer (oBS249 and oBS627). PCR products were purified by PCR clean-up using Monarch® PCR & DNA Cleanup Kit (NEB, USA) and stored at -20 °C until further use. 150 ng DNA per sample was then incubated for 30 min at RT with indicated concentrations of CorR or ParR in EMSA reaction buffer (ERB; 25 mM HEPES [pH 7.4], 150 mM KCl, 1 mM $MgCl_2$, 1 mM DTT, 0.1 mg/ml BSA, 0.1 mg/ml UltraPure™ Salmon Sperm DNA Solution (Thermo Fisher Scientific, USA), 5% glycerol). Samples were then loaded onto 5% Mini-PROTEAN® TBE Gel (BioRad, USA) without any loading dye and run in 1x TBE buffer (130 mM Tris, 45 mM boric acid, 2.5 mM EDTA) at 100 V for 1 h. Finally, Alexa488 signal was recorded in a ChemiDoc MP system (BioRad, USA).

in vivo CorM filament angle measurement

To measure the angle of CorM filaments in *Anabaena* cells, we used two approaches. For the first, we initially straightened *Anabaena* cells expressing mNG-CorM from obtained images using an in house ImageJ script ("straighten_2D"), and filament angles were then measured using CytoSpectre[83]. Alternatively, angles were measured manually in ImageJ using the angle tool by following an imaginary line from one cell pole to the other and then following each individual mNG-CorM filament from the point of contact with this imaginary pole-to-pole line. Both methods resulted in comparable results.

Sample preparation for cryo-EM

For CorM and CorMR samples, 5 µM of CorM alone or 5 µM of both CorM and CorR proteins were incubated in a buffer containing 25 mM HEPES (pH 7.4), 150 mM KCl, 1 mM $MgCl_2$, and 2 mM ATP for 2 hours at 30 °C. The samples were diluted 1/10 in the same buffer and 3 µL of sample were applied onto glow-discharged 200-mesh holey carbon girds (Quantifoil Micro Tools, Au R2/2).

In the case of $^{\Delta 1\text{-}40}$CorM, 5 µM of the protein was pre-mixed in the same buffer but without ATP. ATP was only subsequently added to this mixture and 3.5 µl of the sample were applied onto glow-discharged 200-mesh holey carbon girds (Quantifoil Micro Tools, Cu R1.2/1.3) after an incubation of 1 min at RT. These two modifications were necessary as $^{\Delta 1\text{-}40}$CorM otherwise readily formed extensive, interconnected protein networks within a few minutes after ATP addition that were unsuitable for cryo-EM analysis. All the samples were vitrified in liquid ethane (-185 °C) after 3-3.5s back-side blotting using a Leica GP2 plunger (Leica Microsystems) maintained at 80% humidity and 25 °C (chamber conditions). The grids were stored in liquid nitrogen conditions until imaging.

Cryo-electron microscopy

Single-particle datasets were acquired under cryogenic conditions on a Glacios TEM microscope (Thermo Fisher Scientific) operating at 200 kV equipped with Falcon 4i direct

electron detector. The automated data collection was set up using EPU version 3.7.0 (Thermo Fisher Scientific). The nominal magnification was set to 150,000x, resulting in a pixel size of 0.931 Å for the CorMR dataset and 0.9551 Å for the CorM and $^{\Delta 1-40}$CorM datasets. The defocus range was set to -1.0 to -3 μm. For all three samples, single-particle datasets were acquired in 4,096x4,096px movies in EER mode, with a cumulative dose ranging from 40-80 e$^-$/Å^2.

Single-particle image processing

All the processing steps were performed in CryoSPARC v4.4. For all the datasets, 4,096x4,096px movies in EER mode were motion corrected using patch motion correction, and patch CTF estimation was used to estimate CTF using default patch size in cryoSPARC. Manual curation was done to remove empty micrographs and micrographs containing crystalline ice. The subsequent steps used to process each dataset are summarized below and the details are provided in the schematics presented in Fig. S6.1-6.3.

For the CorMR dataset, initial particle picking was performed using filament tracer with a diameter of 90 Å on 2,305 manually curated micrographs. Particles were initially extracted with a box size of 256 px, followed by 2D classification was performed. Three classes were used as template to perform filament tracing with a diameter of 100 Å. Particles were extracted with a bigger box size (512px) and subjected to 2D classification. Using selected 2D classes (22 classed out of 100), ab-initio reconstruction and homogeneous refinement were performed without symmetry (C1 symmetry). The initial map obtained from homogeneous refinement was used to determine the helical twist and helical rise estimates using the online helical indexing tool (HI3D- Helical Indexing using cylindrical projection of a 3D map)[79]. Helical refinement was performed using an estimated helical twist of -18° and a helical rise 46.5 Å. Local refinement and local CTF refinements were performed after helical refinement and the map clearly displayed C2 symmetry. Two rounds of homogeneous refinement were performed after removing duplicate particles to obtain a final map at 3.92 Å with 124,109 particles, measured at the 0.143 FSC criterion.

For the $^{\Delta 1-40}$CorM dataset, particle picking was performed using filament tracer with a diameter

of 100 Å on 1,200 manually curated micrographs. Particles were extracted with a box size of

512px and subject to 2D classification. Using selected 2D classes (10 classed out of 75), ab-

initio reconstruction and homogeneous refinement were performed with C1 symmetry. The

initial map obtained from homogeneous refinement was used to determine the helical twist and

helical rise estimates using online helical indexing tool (HI3D- Helical Indexing using cylindrical

projection of a 3D map)[79]. Helical refinement was performed using the estimated helical twist

of -15.73° and helical rise 47.3 Å. The map from helical refinement had 287,700 particles, 3D

classification was performed to get 3 different classes. One class with good looking map

containing 44.2 % of the particles (127,289 particles) was selected and used for final helical

refinement to obtain a 4.42 Å map, measured at the 0.143 FSC criterion.

For the CorM dataset, initial particle picking was performed using filament tracer with a

diameter of 100 Å with 5 Å separation distance on 2,305 manually curated micrographs.

Particles were extracted with box size of 512 px and 2D classification was performed. 22

Classes out of 100 classes were used as template to perform filament tracing with a diameter

of 100 Å and 1 Å separation distance. Particles were extracted with a box size of 360px and

subjected to rigorous 2D classification (4 rounds). Using selected 2D classes, ab-initio

reconstruction and homogeneous refinement were performed with C1 symmetry. At this stage,

the particles were re-extracted with a bigger box size (512px). One more round of 2D

classification was performed to remove bad particles. Using selected 2D classes, ab-initio

reconstruction and homogeneous refinement were performed with C1 symmetry. The initial

map obtained from homogeneous refinement was used to determine the helical twist and

helical rise estimates using online helical indexing tool (HI3D- Helical Indexing using cylindrical

projection of a 3D map)[79]. Helical refinement was performed using estimated helical twist of -

18.5 ° and helical rise 46.5 Å. The map from helical refinement had 197,123 particles, 3D

classification was performed to get 3 different classes. Two classes with good looking map

containing 78.3 % of the particles (154,726 particles) were combined and used for final helical

refinement job to obtain a 7.14 Å map, measured at the 0.143 FSC criterion.

Model building and refinement

All model and EM-density visualizations were done in UCSF ChimeraX version 1.9 (2024-12-11)[76]. Model building and refinement was performed using the ISOLDE plugin (v.19)[84] in ChimeraX.

To build a model of the CorM filament, we first predicted CorM using the AlphaFold3 (AF3)[75] webserver, either in its APO-form or bound with ADP or ATP. These predicted models were then rigid body-fitted into the EM density of the filament structure obtained from the CorMR sample. This revealed that the APO-model showed the best fit. However, as there was clearly a density for the nucleotide visible, we manually placed an ADP molecule into the EM-density. As we only observed densities for residues Serine 43 to Asparagine 378 we removed the remaining residues.

To generate a filament-like model, we then rigid body-fitted 6 copies of the generated CorM(43-378)-ADP monomer into the EM-density of the filament. 5 non-crystallographic symmetry copies of the monomers were restrained to a centrally positioned (working) monomer (i.e. a monomer that was surrounded by 5 copies, hence having all its contact sites occupied). Fit and geometry of the working monomer were iteratively checked and locally refined in ISOLDE. Finally, the NCS copies were replaced with the working copy monomer to obtain the final CorM-filament model.

The accuracy of the model was checked using MolProbity[85]. The interface interactions within CorM filament structures were determined using PDBePISA (Proteins, Interfaces, Structure and Assemblies)[52].

Scanning electron microscopy (SEM)

For SEM analysis, cells were grown on BG11 plates, scraped off, and re-suspended in 1 ml MilliQ H_2O. Cells were pelleted by centrifugation (5000 × g, 4 min, RT), washed once by centrifugation with 1 ml Milli-Q H_2O, and then re-suspended in 1 ml of 0.1 M Tris base/HCl buffer containing 2.5% glutaraldehyde. Cells were fixed o/n at 4 °C. The next day, cells were washed 2x with 1 ml MilliQ H_2O by centrifugation (5000 × g, 4 min, RT). After washing, cells

were spotted for 30 seconds onto 12 mm round glass coverslips (Electron Microscopy

Sciences, USA) previously coated overnight with 0.01% poly-L-lysine (Sigma-Aldrich, USA).

Subsequently, cells were subjected to an ethanol dehydration series by sequential incubation

on the coverslips in 6-well plates with 40 %, 60 %, 80 %, and 100 % EtOH, each for 15 minutes.

Samples were then critical point dried using EM CPD300 (Leica Microsystems) and sputter-

coated with gold to a thickness of 5 nm using EM ACE600 coating device (Leica Microsystems)

prior to imaging using a FE-SEM Merlin VP Compact (Zeiss, Germany) at 5 kV with a

secondary electron detector.

1659

Protein sequence analyses

We assembled a database of complete cyanobacterial genomes, aiming to represent the

overall diversity of the phylum included in the NCBI Genome database[86] by including up to five

species per genus when available, resulting in a final dataset of 128 species. To identify

homologs of ParM, we used the *Nostoc* sp. PCC 7120 protein WP_044522477.1 as a query in

a jackhmmer search (HMMER v3.3.2)[87] against our cyanobacterial database. All hits were

aligned using MAFFT v7 (L-INS-i algorithm)[88], and the resulting alignment was manually

curated to remove outliers. From this alignment, we generated a hidden Markov model (HMM),

which we used for a second round of searches with hmmsearch[87]. The same approach was

applied to identify homologs of MinC (WP_190449770.1) and ParR (WP_010999216.1), both

from *Nostoc* sp. PCC 7120, but in the case of ParR, additional analyses were necessary due

to its low sequence conservation and variable length across homologs. To address these

limitations, ParR hits were filtered based on genomic context. In all reported cases, *parR* was

located adjacent to a predicted *parM* gene, and both genes were encoded in the same

orientation. Based on this observation, we only retained ParR hits found immediately next to a

ParM, and all others were discarded. We also examined the distribution of ParM across

Bacteria. Preliminary analyses showed that ParM is unevenly distributed and can be present

in some strains but absent in others within the same species. For this reason, we used a

different strategy than the one applied to Cyanobacteria. We searched all complete bacterial

chromosomes and plasmids deposited in RefSeq[89], excluding cyanobacterial entries, using

hmmsearch with the ParM HMM and an E-value cutoff of 1e−6. This search retrieved 3,693

chromosomal and 2,298 plasmid hits, the majority of which belonged to the phyla

Pseudomonadota and Bacillota, which together accounted for over 95% of all identified

sequences. To reduce redundancy, we randomly selected one hit per genus, aligned all

sequences using MAFFT v7 (L-INS-i algorithm)[88], and manually removed highly divergent or

poorly aligned entries. The resulting curated dataset consisted of 150 chromosomal and 78

plasmid sequences. Given the poor sequence conservation of ParR described above, we did

not rely on direct sequence similarity to search for its homologs across Bacteria. Instead, we

extracted the two neighboring genes (upstream and downstream) of each ParM hit, provided

they were in the same orientation, and predicted their three-dimensional structures using

AlphaFold2[54]. ParR candidates were then manually annotated based on structural similarity to

the ParR reference sequence. The presence of MinC homologs in other bacterial phyla was

assessed as previously described, but using a database containing taxa representative of most

bacterial phyla[90]. To predict amphipathic helices in all the selected ParR sequences, we used

Amphipaseek[51]. Helical wheels were generated using NetWheels[91] and amphipathicity was

calculated using PMIpred[92]. All protein alignments for ParM, ParR and MinC were performed

using MAFFT[88], and columns containing more than 20% gaps were removed using TrimAl[93].

Alignments were visualized in JalView[94]. Sequence logos were generated using WebLogo 3[78].

To analyze and display sequence conservation on protein structures, we used ChimeraX[95] in

combination with the AL2CO algorithm[96].

1700

Phylogenetic analyses

To reconstruct the phylogenies of ParM and MinC, we used all homologs identified in

Cyanobacteria and in the Bacteria datasets. Phylogenetic inference was performed with IQ-

TREE v2[97], using ModelFinder[98] to determine the best-fitting substitution model, and ultrafast

bootstrap approximation with 10,000 replicates[99] and a minimum bootstrap correlation

coefficient of 0.9999 to assess branch support. To improve resolution within the cyanobacterial

clade, we reconstructed a second phylogeny based on concatenated alignments of ParM and

ParR sequences, processed using the same strategy as described above. We also built a

species tree for Cyanobacteria using two conserved, assumed to be vertically inherited

markers: the DNA-directed RNA polymerase subunits β (BAB77960.1) and β′ (BAB77961.1).

Each sequence was aligned and trimmed separately using the same method explained above,

and the two alignments were concatenated for phylogenetic analysis. The resulting tree was

rooted using four species from *Candidatus* Melainabacteria, the sister clade of Cyanobacteria.

All phylogenetic trees were visualized and annotated using the iTOL web server v6[100].

1715

In vivo CorM filament tracking

mNG-CorM or mNG-$^{\Delta 1-40}$CorM filaments were tracked from time lapse movies with images

recorded usually every 5 or 10 seconds. Using Fiji[101], we used the line tool (width 3) and

followed the track of individual filaments of which we generated kymographs using the Reslice

plugin. Growth or shrinkage velocities were then calculated by distance traveled over time

(nm/s).

1722

1723 ***In vitro CorM filament tracking***

1724 Filament ends were manually tracked using the Tracking plugin in Fiji[101]. Distance between

1725 positions were then calculated using a custom Python script.

1726

1727 ***Intrinsically disordered region (IDR) prediction***

1728 Unstructured domains of cyanobacterial CorM or ParM homologs was performed using

1729 ALBATROSS[102] on a Google Colab notebook (IDRome constructor v. 3.0).

1730

AlphaFold-Multimer predictions

For prediction of protein-protein interactions, we tested CorM against the whole *Anabaena*

proteome (RefSeq ID: GCF_000009705.1) as extracted from NCBI. This 'candidates' list was

tested to predict interactions against the 'bait' CorM using AlphaFold-Multimer (v.2.3)[53,75]. This

pipeline produced five protein complexes per protein pair. Postprocessing of the results

includes a filtering step, which removes protein complexes with an ipTM standard deviation

larger than the 1.5xIQR outlier threshold, i.e. 0.095, indicate highly diverse structure

predictions. Unless stated otherwise, all depicted complex predictions and their corresponding

PAE plots are the top-scoring relaxed model generated by Alphafold-Multimer (v3) 64 and

visualized in ChimeraX.

1741

1742 ***Prediction of protein 3D structures***

1743 All shown Protein 3D structures were predicted using AF3[75] and further analyzed using UCSF

1744 ChimeraX (v. 1.9)[76]. For AF3 predictions, the respective top ranked model is shown in cartoon

1745 form and colored by pLDDT score confidence together with the corresponding Predicted

1746 Aligned Error (PAE) plot and the ipTM/pTM scores.

1747

Image and data analysis

Microscopy data analysis was performed using FIJI software (ImageJ2 for Mac; v.

2.14.0/1.54f)[101]. Time series micrographs were routinely processed using the ImageJ gaussian

blur (0.5) and walking average plugins, averaging four consecutive frames. Contrast was

optimized using the Enhance Contrast command (0.05-0.35% saturated pixels; normalize,

process all slices) and, where appropriate, background was subtracted using the Subtract

Background command (50 pixels Rolling ball radius) from ImageJ.

Quantification of protein signals from SDS gels was done using Fiji or Image Lab (BioRad,

USA) software. Protein signal intensities of the loaded pellet and supernatant samples were

recorded and extrapolated to the overall sample volume. Values are given as percentage of

proteins in the pellet or supernatant in respect to the whole protein amount of each reaction.

1759

In vivo timelapse cell division tracking

To analyze the temporal localization of mNG-CorM, time-lapse movies of individual cells were

manually stabilized using a custom Fiji macro to compensate for cell growth, division, rotation,

and lateral drift. Frame intensities were normalized to ensure accurate assessment of signal

localization, and each cell was cropped into a 100×100 pixel region. Initial stabilization was

performed by aligning movie frames to a line region of interest (ROI) positioned along the cell's

lateral axis, guided by mNG-CorM and thylakoid membrane fluorescence signals, to correct

for rotation. After stabilization, a kymograph was generated for each cell by drawing a line

along the longitudinal axis, spanning from pole to pole, with a line width of 1.25 µm to average

the mNG-CorM signal and compensate for gaps in fluorescence. These kymographs were

temporarily aligned based on the onset of cell division, defined by the sudden separation of

the mNG-CorM signal into two diverging trajectories. The aligned kymographs were then

stacked and averaged to produce the composite final image.

1773

Statistical analysis

1774

1775 For all plots throughout this work, error bars indicate the standard deviation (SD).

1776 For TIRF microscopy of *in vitro* experiments, we define a biological replicate as a reaction

1777 performed in a separate reaction chamber. In some cases, replicates were performed on the

1778 same cover glasses, which can hold up to three reaction chambers. A normal distribution was

1779 assumed for all replicates.

1780

References:

1. Wagstaff, J., and Löwe, J. (2018). Prokaryotic cytoskeletons: protein filaments

organizing small cells. Nat. Rev. Microbiol. *16*, 187–201.

https://doi.org/10.1038/nrmicro.2017.153.

2. Møller-Jensen, J., Borch, J., Dam, M., Jensen, R.B., Roepstorff, P., and Gerdes, K.

(2003). Bacterial Mitosis: ParM of Plasmid R1 Moves Plasmid DNA by an Actin-like

Insertional Polymerization Mechanism. Mol. Cell *12*, 1477–1487.

https://doi.org/10.1016/S1097-2765(03)00451-9.

3. Bi, E., and Lutkenhaus, J. (1991). FtsZ ring structure associated with division in

Escherichia coli. Nature *354*, 161–164. https://doi.org/10.1038/354161a0.

4. Loose, M., and Mitchison, T.J. (2014). The bacterial cell division proteins ftsA and ftsZ

self-organize into dynamic cytoskeletal patterns. Nat. Cell Biol. *16*, 38–46.

https://doi.org/10.1038/ncb2885.

5. Shiomi, D., and Margolin, W. (2007). Dimerization or oligomerization of the actin-like

FtsA protein enhances the integrity of the cytokinetic Z ring. Mol. Microbiol. *66*, 1396–

1415. https://doi.org/https://doi.org/10.1111/j.1365-2958.2007.05998.x.

6. Garner, E.C., Bernard, R., Wang, W., Zhuang, X., Rudner, D.Z., and Mitchison, T.

(2011). Coupled, circumferential motions of the cell wall synthesis machinery and

MreB filaments in B. subtilis. Science (80-.). *333*, 222–225.

https://doi.org/10.1126/science.1203285.

7. Ramm, B., Heermann, T., and Schwille, P. (2019). The E. coli MinCDE system in the

regulation of protein patterns and gradients. Cell. Mol. Life Sci. *76*, 4245–4273.

https://doi.org/10.1007/s00018-019-03218-x.

8. Flores, E., and Herrero, A. (2014). The cyanobacteria: morphological diversity in a

photoautotrophic lifestyle.

9. Rippka, R., Stanier, R.Y., Deruelles, J., Herdman, M., and Waterbury, J.B. (1979).

Generic Assignments, Strain Histories and Properties of Pure Cultures of

Cyanobacteria. Microbiology *111*, 1–61. https://doi.org/10.1099/00221287-111-1-1.

10. Herrero, A., Stavans, J., and Flores, E. (2016). The multicellular nature of filamentous

heterocyst-forming cyanobacteria. FEMS Microbiol. Rev. *40*, 831–854.

https://doi.org/10.1093/femsre/fuw029.

11. Hoiczyk, E., and Baumeister, W. (1995). Envelope structure of four gliding filamentous

cyanobacteria. J. Bacteriol. *177*, 2387–2395. https://doi.org/10.1128/jb.177.9.2387-

2395.1995.

12. Kieninger, A.K., and Maldener, I. (2021). Cell–cell communication through septal

junctions in filamentous cyanobacteria. Curr. Opin. Microbiol. *61*, 35–41.

https://doi.org/10.1016/j.mib.2021.02.002.

13. Weiss, G.L., Kieninger, A.-K., Maldener, I., Forchhammer, K., and Pilhofer, M. (2019).

Structure and Function of a Bacterial Gap Junction Analog. Cell *178*, 374-384.e15.

https://doi.org/10.1016/J.CELL.2019.05.055.

14. Miyagishima, S.Y., Wolk, P.P., and Osteryoung, K.W. (2005). Identification of

cyanobacterial cell division genes by comparative and mutational analyses. Mol.

Microbiol. *56*, 126–143. https://doi.org/10.1111/j.1365-2958.2005.04548.x.

15. Springstein, B.L., Weissenbach, J., Koch, R., Stücker, F., and Stucken, K. (2020). The

role of the cytoskeletal proteins MreB and FtsZ in multicellular cyanobacteria. FEBS

Open Bio *10*, 2510–2531. https://doi.org/https://doi.org/10.1002/2211-5463.13016.

16. Naha, A., Haeusser, D.P., and Margolin, W. (2023). Anchors: A way for FtsZ filaments

to stay membrane bound. Mol. Microbiol. *120*, 525–538.

https://doi.org/10.1111/mmi.15067.

17. Camargo, S., Picossi, S., Corrales-Guerrero, L., Valladares, A., Arévalo, S., and

Herrero, A. (2019). ZipN is an essential FtsZ membrane tether and contributes to the

septal localization of SepJ in the filamentous cyanobacterium Anabaena. Sci. Rep. *9*,

1–15. https://doi.org/10.1038/s41598-019-39336-6.

18. MacCready, J.S., Schossau, J., Osteryoung, K.W., and Ducat, D.C. (2017). Robust

Min-system oscillation in the presence of internal photosynthetic membranes in

cyanobacteria. Mol. Microbiol. *103*, 483–503. https://doi.org/10.1111/mmi.13571.

19. Springstein, B.L., Nürnberg, D.J., Weiss, G.L., Pilhofer, M., and Stucken, K. (2020).

Structural determinants and their role in cyanobacterial morphogenesis. Life *10*, 355.

https://doi.org/https://doi.org/10.3390/life10120355.

20. Velázquez-Suárez, C., Valladares, A., Luque, I., and Herrero, A. (2022). The Role of

Mre Factors and Cell Division in Peptidoglycan Growth in the Multicellular

Cyanobacterium Anabaena. MBio *13*. https://doi.org/10.1128/mbio.01165-22.

21. Hu, B., Yang, G., Zhao, W., Zhang, Y., and Zhao, J. (2007). MreB is important for cell

shape but not for chromosome segregation of the filamentous cyanobacterium

Anabaena sp. PCC 7120. Mol. Microbiol. *63*, 1640–1652.

https://doi.org/10.1111/j.1365-2958.2007.05618.x.

22. Velázquez-Suárez, C., Luque, I., and Herrero, A. (2020). The Inorganic Nutrient

Regime and the mre Genes Regulate Cell and Filament Size and Morphology in the

Phototrophic Multicellular Bacterium Anabaena. mSphere *5*, e00747-20.

https://doi.org/10.1128/mSphere.00747-20.

23. Griese, M., Lange, C., and Soppa, J. (2011). Ploidy in cyanobacteria. FEMS Microbiol.

Lett. *323*, 124–131. https://doi.org/10.1111/j.1574-6968.2011.02368.x.

24. Schneider, D., Fuhrmann, E., Scholz, I., Hess, W.R., and Graumann, P.L. (2007).

Fluorescence staining of live cyanobacterial cells suggest non-stringent chromosome

segregation and absence of a connection between cytoplasmic and thylakoid

membranes. BMC Cell Biol. *8*, 39. https://doi.org/10.1186/1471-2121-8-39.

25. Sengupta, M., and Austin, S. (2011). Prevalence and significance of plasmid

maintenance functions in the virulence plasmids of pathogenic bacteria. Infect. Immun.

79, 2502–2509. https://doi.org/10.1128/IAI.00127-11.

26. Jain, I.H., Vijayan, V., and O'Shea, E.K. (2012). Spatial ordering of chromosomes

enhances the fidelity of chromosome partitioning in cyanobacteria. Proc. Natl. Acad.

Sci. U. S. A. *109*, 13638–13643. https://doi.org/10.1073/pnas.1211144109.

27. Ohbayashi, R., Nakamachi, A., Hatakeyama, T.S., Watanabe, S., Kanesaki, Y.,

Chibazakura, T., Yoshikawa, H., and Miyagishima, S.Y. (2019). Coordination of

polyploid chromosome replication with cell size and growth in a cyanobacterium. MBio

10, 1–15. https://doi.org/10.1128/MBIO.00510-19.

28. Watanabe, S., Noda, A., Ohbayashi, R., Uchioke, K., Kurihara, A., Nakatake, S.,

Morioka, S., Kanesaki, Y., Chibazakura, T., and Yoshikawa, H. (2018). ParA-like

protein influences the distribution of multi-copy chromosomes in cyanobacterium

Synechococcus elongatus PCC 7942. Microbiology *164*, 45–56.

https://doi.org/10.1099/mic.0.000577.

29. Salje, J., Gayathri, P., and Löwe, J. (2010). The ParMRC system: Molecular

mechanisms of plasmid segregation by actin-like filaments. Nat. Rev. Microbiol. *8*,

683–692. https://doi.org/10.1038/nrmicro2425.

30. Gayathri, P., Fujii, T., Møller-Jensen, J., Van Den Ent, F., Namba, K., and Löwe, J.

(2012). A bipolar spindle of antiparallel ParM filaments drives bacterial plasmid

segregation. Science (80-.). *338*, 1334–1337.

https://doi.org/10.1126/science.1229091.

31. Van den Ent, F., Møller-Jensen, J., Amos, L.A., Gerdes, K., and Löwe, J. (2002). F-

actin-like filaments formed by plasmid segregation protein ParM. EMBO J. *21*, 6935–

6943. https://doi.org/10.1093/emboj/cdf672.

32. Bharat, T.A.M., Murshudov, G.N., Sachse, C., and Löwe, J. (2015). Structures of

actin-like ParM filaments show architecture of plasmid-segregating spindles. Nature

523, 106–110. https://doi.org/10.1038/nature14356.

33. Møller-Jensen, J., Jensen, R.B., Löwe, J., and Gerdes, K. (2002). Prokaryotic DNA

segregation by an actin-like filament. EMBO J. *21*, 3119–3127.

https://doi.org/10.1093/emboj/cdf320.

34. Ali, S., Koh, A., Popp, D., Tanaka, K., Kitaoku, Y., Miyazaki, N., Iwasaki, K., Mitsuoka,

K., Robinson, R.C., and Narita, A. (2025). Bacterial genome encoded ParMs. J. Biol.

Chem., 110351.

35. Gerdes, K., Moller-Jensen, J., and Jensen, R.B. (2002). Plasmid and chromosome

partitioning: Surprises from phylogeny. Mol. Microbiol. *37*, 455–466.

https://doi.org/10.1046/j.1365-2958.2000.01975.x.

36. Møller-Jensen, J., Ringgaard, S., Mercogliano, C.P., Gerdes, K., and Löwe, J. (2007).

Structural analysis of the ParR/parC plasmid partition complex. EMBO J. *26*, 4413–

4422. https://doi.org/10.1038/sj.emboj.7601864.

37. Breüner, A., Jensen, R.B., Dam, M., Pedersen, S., and Gerdes, K. (1996). The

centromere-like parC locus of plasmid R1. Mol. Microbiol. *20*, 581–592.

https://doi.org/10.1046/j.1365-2958.1996.5351063.x.

38. Dam, M., and Gerdes, K. (1994). Partitioning of plasmid R1 Ten direct repeats flanking

the parA promoter constitute a centromere-like partition site parC, that expresses

incompatibility. J. Mol. Biol. *236*, 1289–1298. https://doi.org/10.1016/0022-

2836(94)90058-2.

39.   Jiang, S., Narita, A., Popp, D., Ghoshdastider, U., Lee, L.J., Srinivasan, R.,

Balasubramanian, M.K., Oda, T., Koh, F., Larsson, M., et al. (2016). Novel actin

filaments from Bacillus thuringiensis form nanotubules for plasmid DNA segregation.

Proc. Natl. Acad. Sci. U. S. A. *113*, E1200–E1205.

https://doi.org/10.1073/pnas.1600129113.

40.   Brzoska, A.J., Jensen, S.O., Barton, D.A., Davies, D.S., Overall, R.L., Skurray, R.A.,

and Firth, N. (2016). Dynamic filament formation by a divergent bacterial actin-like

ParM protein. PLoS One *11*, 1–14. https://doi.org/10.1371/journal.pone.0156944.

41.   Schreiter, E.R., and Drennan, C.L. (2007). Ribbon-helix-helix transcription factors:

Variations on a theme. Nat. Rev. Microbiol. *5*, 710–720.

https://doi.org/10.1038/nrmicro1717.

42.   Salje, J., and Löwe, J. (2008). Bacterial actin: Architecture of the ParMRC plasmid

DNA partitioning complex. EMBO J. *27*, 2230–2238.

https://doi.org/10.1038/emboj.2008.152.

43.   Orlova, A., Garner, E.C., Galkin, V.E., Heuser, J., Mullins, R.D., and Egelman, E.H.

(2007). The structure of bacterial ParM filaments. Nat. Struct. Mol. Biol. *14*, 921–926.

https://doi.org/10.1038/nsmb1300.

44.   Lee, P.S., and Grossman, A.D. (2006). The chromosome partitioning proteins Soj

(ParA) and Spo0J (ParB) contribute to accurate chromosome partitioning, separation

of replicated sister origins, and regulation of replication initiation in Bacillus subtilis.

Mol. Microbiol. *60*, 853–869. https://doi.org/10.1111/j.1365-2958.2006.05140.x.

45.   Mohl, D.A., and Gober, J.W. (1997). Cell Cycle–Dependent Polar Localization of

Chromosome Partitioning Proteins in Caulobacter crescentus. Cell *88*, 675–684.

https://doi.org/10.1016/s0092-8674(00)81910-8.

46.   Niki, H., Ogura, T., Ichinose, C., Mori, H., Ezaki, B., and Jaffe, A. (1989). Chromosome

partitioning in Escherichia coli: novel mutants producing anucleate cells. J. Bacteriol.

171, 1496–1505.

47.   Ungerer, J., and Pakrasi, H.B. (2016). Cpf1 Is A Versatile Tool for CRISPR Genome

Editing Across Diverse Species of Cyanobacteria. Sci. Rep. *6*, 1–9.

https://doi.org/10.1038/srep39681.

48.   Campbell, C.S., and Mullins, R.D. (2007). In vivo visualization of type II plasmid

segregation: Bacterial actin filaments pushing plasmids. J. Cell Biol. *179*, 1059–1066.

https://doi.org/10.1083/jcb.200708206.

49.   Garner, E.C., Campbell, C.S., and Dyce Mullins, R. (2004). Dynamic Instability in a

DNA-Segregating Prokaryotic Actin Homolog. Science (80-.). *306*, 1021–1025.

50.   Garner, E.C., Campbell, C.S., Weibel, D.B., and Mullins, R.D. (2007). Reconstitution

of DNA Segregation Driven by Assembly of a Prokaryotic Actin Homolog. Science (80-

.). *315*, 1270–1274.

51.   Sapay, N., Guermeur, Y., and Deléage, G. (2006). Prediction of amphipathic in-plane

membrane anchors in monotopic proteins using a SVM classifier. BMC Bioinformatics

7, 1–11. https://doi.org/10.1186/1471-2105-7-255.

52.   Krissinel, E., and Henrick, K. (2007). Inference of Macromolecular Assemblies from

Crystalline State. J. Mol. Biol. *372*, 774–797.

https://doi.org/https://doi.org/10.1016/j.jmb.2007.05.022.

53.   Evans, R., O'Neill, M., Pritzel, A., Antropova, N., Senior, A., Green, T., Žídek, A.,

Bates, R., Blackwell, S., Yim, J., et al. (2022). Protein complex prediction with

AlphaFold-Multimer. bioRxiv, 2021.10.04.463034.

https://doi.org/10.1101/2021.10.04.463034.

54.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly

accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.

https://doi.org/10.1038/s41586-021-03819-2.

55.   de Boer, P.A.J., Crossley, R.E., and Rothfield, L.I. (1989). A division inhibitor and a

topological specificity factor coded for by the minicell locus determine proper

placement of the division septum in E. coli. Cell *56*, 641–649.

1959 https://doi.org/10.1016/0092-8674(89)90586-2.

1960 56. Hu, Z., and Lutkenhaus, J. (1999). Topological regulation of cell division in Escherichia

1961 coli involves rapid pole to pole oscillation of the division inhibitor MinC under the

1962 control of MinD and MinE. Mol. Microbiol. *34*, 82–90.

1963 https://doi.org/https://doi.org/10.1046/j.1365-2958.1999.01575.x.

1964 57. Raskin, D.M., and De Boer, P.A.J. (1999). Rapid pole-to-pole oscillation of a protein

1965 required for directing division to the middle of Escherichia coli. Proc. Natl. Acad. Sci.

1966 *96*, 4971–4976.

1967 58. Pollard, T.D. (2016). Actin and actin-binding proteins. Cold Spring Harb. Perspect.

1968 Biol. *8*, 1–17. https://doi.org/10.1101/cshperspect.a018226.

1969 59. Xue, B., Leyrat, C., Grimes, J.M., and Robinson, R.C. (2014). Structural basis of

1970 thymosin-β4/profilin exchange leading to actin filament polymerization. Proc. Natl.

1971 Acad. Sci. U. S. A. *111*, E4596–E4605. https://doi.org/10.1073/pnas.1412271111.

1972 60. Palmgren, S., Vartiainen, M., and Lappalainen, P. (2002). Twinfilin, a molecular

1973 mailman for actin monomers. J. Cell Sci. *115*, 881–886.

1974 https://doi.org/10.1242/jcs.115.5.881.

1975 61. Curmi, P.A., Andersen, S.S.L., Lachkar, S., Gavet, O., Karsenti, E., Knossow, M., and

1976 Sobel, A. (1997). The stathmin/tubulin interaction in vitro. J. Biol. Chem. *272*, 25029–

1977 25036. https://doi.org/10.1074/jbc.272.40.25029.

1978 62. Hu, Z., Mukherjee, A., Pichoff, S., and Lutkenhaus, J. (1999). The MinC component of

1979 the division site selection system in Escherichia coli interacts with FtsZ to prevent

1980 polymerization. Proc. Natl. Acad. Sci. U. S. A. *96*, 14819–14824.

1981 https://doi.org/10.1073/pnas.96.26.14819.

1982 63. Paredez, A.R., Somerville, C.R., and Ehrhardt, D.W. (2006). Visualization of Cellulose

1983 Synthase Demonstrates Functional Association with Microtubules. Science (80-.).

1984 *312*, 1491–1495. https://doi.org/10.1126/science.1126551.

1985 64. Sugimoto, K., Williamson, R.E., and Wasteneys, G.O. (2000). New techniques enable

1986 comparative analysis of microtubule orientation, wall texture, and growth rate in intact

roots of Arabidopsis. Plant Physiol. *124*, 1493–1506.

https://doi.org/10.1104/pp.124.4.1493.

65.   Paredez, A.R., Persson, S., Ehrhardt, D.W., and Somerville, C.R. (2008). Genetic

evidence that cellulose synthase activity influences microtubule cortical array

organization. Plant Physiol. *147*, 1723–1734. https://doi.org/10.1104/pp.108.120196.

66.   Savage, D.F., Afonso, B., Chen, A.H., and Silver, P.A. (2010). Spatially Ordered

Dynamics of the Bacterial Carbon Fixation Machinery. Science (80-.). *327*, 1258–

1261. https://doi.org/10.1126/science.1186090.

67.   Wachi, M., Doi, M., Tamaki, S., Park, W., Nakajima-Iijima, S., and Matsuhashi, M.

(1987). Mutant isolation and molecular cloning of mre genes, which determine cell

shape, sensitivity to mecillinam, and amount of penicillin-binding proteins in

Escherichia coli. J. Bacteriol. *169*, 4935–4940.

68.   Shi, H., Bratton, B.P., Gitai, Z., and Huang, K.C. (2018). How to Build a Bacterial Cell:

MreB as the Foreman of E. coli Construction. Cell *172*, 1294–1305.

https://doi.org/10.1016/j.cell.2018.02.050.

69.   van den Ent, F., Izoré, T., Bharat, T.A.M., Johnson, C.M., and Löwe, J. (2014).

Bacterial actin MreB forms antiparallel double filaments. Elife *2014*, 1–22.

https://doi.org/10.7554/eLife.02634.

70.   van Teeffelen, S., Wang, S., Furchtgott, L., Huang, K.C., Wingreen, N.S., Shaevitz,

J.W., and Gitai, Z. (2011). The bacterial actin MreB rotates, and rotation depends on

cell-wall assembly. Proc. Natl. Acad. Sci. *108*, 15822 LP – 15827.

https://doi.org/10.1073/pnas.1108999108.

71.   Mukherjee, A., and Lutkenhaus, J. (1998). Dynamic assembly of FtsZ regulated by

GTP hydrolysis. EMBO J. *17*, 462–469. https://doi.org/10.1093/emboj/17.2.462.

72.   Carlier, M.F., Pantaloni, D., and Korn, E.D. (1984). Evidence for an ATP cap at the

ends of actin filaments and its regulation of the F-actin steady state. J. Biol. Chem.

259, 9983–9986. https://doi.org/10.1016/s0021-9258(18)90914-2.

73.   Mitchison, T., and Kirschner, M. (1984). Dynamic instability of microtubule growth.

Nature *312*, 237–242.

74.   Rubio, M.Á., Napolitano, M., Ochoa De Alda, J.A.G., Santamaría-Gómez, J.,

Patterson, C.J., Foster, A.W., Bru-Martínez, R., Robinson, N.J., and Luque, I. (2015).

Trans-oligomerization of duplicated aminoacyl-tRNA synthetases maintains genetic

code fidelity under stress. Nucleic Acids Res. *43*, 9905–9917.

https://doi.org/10.1093/nar/gkv1020.

75.   Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,

Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure prediction of

biomolecular interactions with AlphaFold 3. Nature *630*, 493–500.

https://doi.org/10.1038/s41586-024-07487-w.

76.   Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H.,

and Ferrin, T.E. (2023). UCSF ChimeraX: Tools for structure building and analysis.

Protein Sci. *32*, e4792. https://doi.org/https://doi.org/10.1002/pro.4792.

77.   Vega-Palas, M.A., Flores, E., and Herrero, A. (1992). NtcA, a global nitrogen regulator

from the cyanobacterium Synechococcus that belongs to the Crp family of bacterial

regulators. Mol. Microbiol. *6*, 1853–1859. https://doi.org/https://doi.org/10.1111/j.1365-

2958.1992.tb01357.x.

78.   Crooks, G.E., Hon, G., Chandonia, J.-M., and Brenner, S.E. (2004). WebLogo: a

sequence logo generator. Genome Res. *14*, 1188–1190.

79.   Sun, C., Gonzalez, B., and Jiang, W. (2022). Helical indexing in real space. Sci. Rep.

12, 8162.

80.   Corrales-Guerrero, L., Camargo, S., Valladares, A., Picossi, S., Luque, I., Ochoa De

Alda, J.A.G., and Herrero, A. (2018). FtsZ of filamentous, heterocyst-forming

cyanobacteria has a conserved N-terminal peptide required for normal FtsZ

polymerization and cell division. Front. Microbiol. *9*, 1–20.

https://doi.org/10.3389/fmicb.2018.02260.

81.   Valladares, A., Picossi, S., Corrales-Guerrero, L., and Herrero, A. (2023). The role of

SepF in cell division and diazotrophic growth in the multicellular cyanobacterium

Anabaena sp. strain PCC 7120. Microbiol. Res. *277*, 127489.

https://doi.org/10.1016/j.micres.2023.127489.

82.   Karimova, G., Davi, M., and Ladant, D. (2012). The β-lactam resistance protein Blr, a

small membrane polypeptide, is a component of the Escherichia coli cell division

machinery. J. Bacteriol. *194*, 5576–5588. https://doi.org/10.1128/JB.00774-12.

83.   Kartasalo, K., Pölönen, R.P., Ojala, M., Rasku, J., Lekkala, J., Aalto-Setälä, K., and

Kallio, P. (2015). CytoSpectre: A tool for spectral analysis of oriented structures on

cellular and subcellular levels. BMC Bioinformatics *16*, 1–23.

https://doi.org/10.1186/s12859-015-0782-y.

84.   Croll, T.I. (2018). ISOLDE: a physically realistic environment for model building into

low-resolution electron-density maps. Biol. Crystallogr. *74*, 519–530.

85.   Williams, C.J., Headd, J.J., Moriarty, N.W., Prisant, M.G., Videau, L.L., Deis, L.N.,

Verma, V., Keedy, D.A., Hintze, B.J., and Chen, V.B. (2018). MolProbity: more and

better reference data for improved all-atom structure validation. Protein Sci. *27*, 293–

315.

86.   Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Farrell,

C.M., Feldgarden, M., Fine, A.M., Funk, K., et al. (2023). Database resources of the

National Center for Biotechnology Information in 2023. Nucleic Acids Res. *51*, D29–

D38. https://doi.org/10.1093/nar/gkac1032.

87.   Eddy, S.R. (2011). Accelerated Profile HMM Searches. PLOS Comput. Biol. *7*,

e1002195.

88.   Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment software

version 7: Improvements in performance and usability. Mol. Biol. Evol. *30*, 772–780.

https://doi.org/10.1093/molbev/mst010.

89.   Goldfarb, T., Kodali, V.K., Pujar, S., Brover, V., Robbertse, B., Farrell, C.M., Oh, D.-H.,

Astashyn, A., Ermolaeva, O., Haddad, D., et al. (2025). NCBI RefSeq: reference

sequence standards through 25 years of curation and annotation. Nucleic Acids Res.

53, D243–D257. https://doi.org/10.1093/nar/gkae1038.

90. Megrian, D., Taib, N., Jaffe, A.L., Banfield, J.F., and Gribaldo, S. (2022). Ancient origin

and constrained evolution of the division and cell wall gene cluster in Bacteria. Nat.

Microbiol. *7*, 2114–2127. https://doi.org/10.1038/s41564-022-01257-y.

91. AR, M., MS, C., and W, F. (2024). NetWheels: A Web Application to Create High

Quality Peptide Helical Wheel and Net Projections. J. Bioinforma. Syst. Biol. *7*, 98–

100. https://doi.org/10.26502/jbsb.5107082.

92. Van Hilten, N., Verwei, N., Methorst, J., Nase, C., Bernatavicius, A., and Risselada,

H.J. (2024). PMIpred: a physics-informed web server for quantitative protein-

membrane interaction prediction. Bioinformatics *40*, 1–10.

https://doi.org/10.1093/bioinformatics/btae069.

93. Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009). trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics *25*,

1972–1973. https://doi.org/10.1093/bioinformatics/btp348.

94. Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., and Barton, G.J. (2009).

Jalview Version 2—a multiple sequence alignment editor and analysis workbench.

Bioinformatics *25*, 1189–1191. https://doi.org/10.1093/bioinformatics/btp033.

95. Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I.,

Morris, J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for

researchers, educators, and developers. Protein Sci. *30*, 70–82.

https://doi.org/https://doi.org/10.1002/pro.3943.

96. Pei, J., and Grishin, N. V (2001). AL2CO: calculation of positional conservation in a

protein sequence alignment. Bioinformatics *17*, 700–712.

https://doi.org/10.1093/bioinformatics/17.8.700.

97. Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von

Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods

for Phylogenetic Inference in the Genomic Era. Mol. Biol. Evol. *37*, 1530–1534.

https://doi.org/10.1093/molbev/msaa015.

98. Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., and Jermiin, L.S.

(2017). ModelFinder: fast model selection for accurate phylogenetic estimates. Nat.

Methods *14*, 587–589. https://doi.org/10.1038/nmeth.4285.

99.   Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., and Vinh, L.S. (2018).

UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol. Biol. Evol. *35*, 518–

522. https://doi.org/10.1093/molbev/msx281.

100. Letunic, I., and Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. Nucleic Acids Res. *49*, W293–W296.

https://doi.org/10.1093/nar/gkab301.

101. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source

platform for biological-image analysis, https://doi.org/10.1038/nmeth.2019

https://doi.org/10.1038/nmeth.2019.

102. Lotthammer, J.M., Ginell, G.M., Griffith, D., Emenecker, R.J., and Holehouse, A.S.

(2024). Direct prediction of intrinsically disordered protein conformational properties

from sequence. Nat. Methods *21*, 465–476. https://doi.org/10.1038/s41592-023-

02159-5.

2115

Figure 1

Figure 2

A

P$_{parMR}$-mNG-ParM

P$_{parMR}$-mNG-ParM + chlA

side

5 µm

5 µm

B

cross section

2.5 µm

C

— mNG-ParM
— chlA
— DAPI

normalized intensity

Cell width (µm)

D

Angle to long axis (degrees)

E

Cell length ± Std. Dev.

x

t

~2.5 h

~7.5 h

~12.5 h

1 µm

t = ~2.5 h

During growth

t = ~7.5 h

Before constriction

t = ~12.5 h

After division

Normalized intensity

Distance (µm)

F

0 s   10 s   20 s

30 s   40 s   50 s

60 s   70 s   80s

90 s   100 s   120 s

130 s   140 s   150 s

160 s   170 s   180 s

P$_{parMR}$-mNG-ParM

1 µm

G

x

t

14 nm/s

21nm/s

1 min

0.5 µm

H

Growth velocity (nm/s)

ParM

I

P$_{ntcA}$-mNG-ParM

ΔparMR

5 µm

A

CorR [µM]

0 1 2.5 5 10 20

400 bp upstream *corM*

EcParR [µM]

0 1 2.5 5 10 20

parC (380 bp)

B

CorM binding

C C

N N

amphipathic helix

■ Basic ■ Acid
■ Polar/Uncharged ■ Nonpolar

Anabaena

V M
W   S
      1
S   V
K   F
Q K K

0.535

E. coli

I M
K   R
      1
K   A
T   M
D F R

0.157

C

CorM | CorR | Δ1-10CorR | CorM+CorR | CorM+Δ1-10CorR | Δ150-162CorR | CorM+Δ150-162CorR

MLVs ctrl. MLVs ctrl. MLVs ctrl. MLVs ctrl. MLVs ctrl. MLVs ctrl. MLVs ctrl.

kDa P S P S P S P S P S P S P S P S P S P S P S P S P S P S

55 —

CorM

35 —

25 —

CorR

15 —

D

● Predicted N-terminal AH
○ *Anabaena* CorR

Tree scale: 1

Figure 4

Feature	CorM	ParM	MreB
Polarity	Antiparallel	Parallel	Antiparallel
Staggered	No	Yes	No
Twist (°)	-15 to -18.5	-166 to -167	n/a
Rise (Å)	46-47.3	24-25	51

Figure 5

Figure 6

A

ParM
N-term IDR ParM (1-40)

Anabaena

Very high (pLDDT > 90)
Confident (90 > pLDDT > 70)
Low (70 > pLDDT > 50)
Very low (pLDDT < 50)

Scored residue

Aligned residue

Expected position error (Ångströms)

E. coli (PDB: 4A62)

B

ParR

H5b
H5a
(RHH)₂

Anabaena

Very high (pLDDT > 90)
Confident (90 > pLDDT > 70)
Low (70 > pLDDT > 50)
Very low (pLDDT < 50)

Scored residue

Aligned residue

Expected position error (Ångströms)

H5
H4
H3
(RHH)₂

E. coli (PDB: 2JD3)

C

Number of IDR residues

Green   Yellow   Pink
Clade

D

Optical density (OD)

WT
ΔparMR

Time (days)

E

WT   ΔparMR

2 µm   2 µm   2 µm

F

ΔparMR + P_ntcA-parM-parR   ΔparMR

5 µm

G

E. coli C41 (DE3)

P_T7-*mNG-parM*   P_BAD-*parR-mScarlet-I*
                        enhanced contrast   DIA

SUPPLEMENTARY 6.1

Cryo-EM image processing strategy applied to obtain CorM filament structure in the presence of CorR.

Cryo-EM image processing strategy applied to obtain N40_CorM filament structure.

SUPPLEMENTARY 6.3

Cryo-EM image processing strategy applied to obtain CorM filament structure.

A

7.1 Å
CorM

B

4.4 Å
$^{\Delta 1\text{-}40}$CorR

C

3.9 Å
CorMR

Tree scale: 1

Cyanobacteria   Nostocales

Protein length (# aa):   135   311   487

N-ter sequence identity (%):   80>   >60   >40

Globular domain sequence identity (%):   >80   >60   >40