CactEcoDB: Trait, spatial, environmental, phylogenetic and diversification data for the cactus family

Jamie B. Thompson[1*], Catherine Martinez[1,2], Jorge Avaria-Llautureo[1], Santiago Ramírez-Barahona[3], Gerardo Manzanarez-Villasana[3], Alastair Culham[1], Andrew Gdaniec[1], George Ryan[1], Chris Venditti[1], Georgia Keeling[4], Nicholas K. Priest[4*]

[1] School of Biological Sciences, University of Reading, Reading, UK
[2] School of Archaeology, Geography and Environmental Science, University of Reading, Reading, UK
[3] Departamento de Botánica, Instituto de Biología, Universidad Nacional Autónoma de México (UNAM), Circuito Exterior s/n, Ciudad de México, Mexico
[4] The Milner Centre for Evolution, Department of Life Sciences, University of Bath, Bath, UK
* Corresponding authors

Abstract

Integrated datasets linking traits, spatial distributions, environmental variables and phylogenies are essential for comparative research, but remain limited for many plant taxa, including those which are most threatened. Cactaceae are a morphologically and ecologically diverse succulent family that are iconic components of ecosystems across the Americas, and face high extinction risk. To support future comparative research, we present CactEcoDB (The Cactus Ecological Database), an Open Access dataset of curated spatial, ecological, trait, phylogenetic, and diversification data for over 1,000 cactus species. CactEcoDB includes species-level trait data, geographic occurrence records, environmental variables, range size estimates, speciation rates, and the largest time-calibrated phylogeny of cacti to date. By integrating these diverse data in a single platform, CactEcoDB enables analyses of trait evolution, trait–environment interactions, biogeography, climatic niche evolution, and macroevolutionary dynamics in one of the most celebrated and threatened plant families.

Background & Summary

Recent research highlights the importance of comparative methods for our understanding of the origins and ecological drivers of global biodiversity in large families of plants [1–12]. This type of comparative research relies on comprehensive datasets that adequately capture the complexity of trait evolution, environmental variation, and lineage diversification [13–35]. There is a growing appreciation of how this complexity, especially ecological interactions, coevolutionary dynamics, and environmental heterogeneity, shapes macroevolutionary and macroecological trajectories [9,10,12,36–41]. As global change accelerates and data availability expands unevenly across lineages, so does the need for integrative, high-quality datasets for threatened and unique groups

41 [25,26,42–50]. However, assembling such datasets remains difficult and time-consuming,
42 often requiring expert curation and synthesis across disparate data types and sources.
43
The cactus family (Cactaceae) is among the most morphologically and ecologically
distinctive plant lineages [51–57]. Comprising approximately 1,850 species, cacti are found
in an extraordinary range of habitats, from hyper-arid deserts and tropical dry forests to
high-elevation grasslands and volcanic islands [58]. Centers of species richness are found
in Mexico, the Andes, and Eastern Brazil [51–57], but their distribution extends from
southern Canada to Patagonia, with one species probably occurring naturally outside of
the Americas (*Rhipsalis baccifera*) [59]. Cacti are well known for their highly specialised
morphological and physiological adaptations to water-limited environments, including
stem succulence, heavily modified leaves, thick cuticles, and crassulacean acid
metabolism (CAM) photosynthesis [10,12,51,54]. These traits make them a classic example
of adaptive evolution in plants, and in particular, a model system for understanding
evolutionary responses to extreme environmental conditions.
56
Comparative research in the Cactaceae has bearing on diverse fields including plant
physiology [60–62], biogeography [51,63–66], ecology [67–69], evolutionary dynamics [10,12,51,54],
and conservation biology [53,66,70–73]. The growing clarity of phylogenetic relationships
within the family [52] has facilitated tests of the evolution of key traits such as growth
forms [60], pollinators [51,56], and chromosome count [74]. The power to detect drivers of
diversification is greatly aided by the remarkably wide distribution and range of growth
forms found in diverse environments across gradients including aridity, temperature,
and elevation [51,52,55,74].
65
Beyond their ecomorphological adaptations, cacti hold exceptional cultural and
economic significance [75–77]. Archaeological evidence shows that species such as
Opuntia ficus-indica were consumed more than 9,000 years ago in Mesoamerica [78,79],
where they continue to serve as important sources of food, forage, and dye [80–82]. Today,
cacti are cultivated globally for ornamental purposes, and their remarkably varied forms,
ranging from tiny globose species to massive columnars [60], make them among the most
collected and traded groups in horticulture [83] along with orchids. They are unusual in
horticulture in that the bulk of trade is of natural species, not highly bred and modified
cultivars. The deep cultural history, combined with their horticultural desirability, has led
to intense pressure on wild populations through unsustainable collection and illegal
trade [84,85]. As a result, Cactaceae is one of the most threatened plant families. A 2015
assessment found that 31% of cactus species are threatened with extinction, with
habitat loss, climate change, and overexploitation cited as primary drivers [70]. More
recent projections suggest that 60-90% may experience future range contractions under
near-future climate change scenarios [71].

The integration of phylogenetic, trait, and spatial data at the species level in cacti has
remained fragmented, limiting the ability of researchers to address broad ecological and
evolutionary questions, or to support conservation assessments grounded in
comprehensive ecological data [72,86]. To fill this gap, we present CactEcoDB, a curated
dataset that integrates spatial, ecological, trait, and phylogenetic information for over
1,000 species of Cactaceae. This is a greatly updated and expanded version of
previous datasets [10]. CactEcoDB includes biotic trait data (e.g., growth form, plant
height and pollinator), expert-curated geographic ranges, numerous important
environmental variables, and an updated phylogeny, the largest time-calibrated
molecular phylogeny for the group. As compared with the largest previous dataset [10],
CactEcoDB expands and improves the available data: i) it increases in species
sampling for plant height by 24.93%, spatial variables by ~6%, and adds eight new
variables; ii) it expands the number of classified growth forms from two to nine; iii) it
replaces highly incomplete GBIF data with expert ranges defined by the IUCN [53,66,70–
73,87], which were manually verified against independent data, especially the Plants of the
World Online (POWO) [88]; and iv) it provides a general-purpose resource for ecological,
evolutionary, biogeographic, and conservation-focussed research. Furthermore, we
have substantially expanded and updated several of the biotic traits with higher
resolution and species coverage than previously available [10]. By bringing together large
scale data covering all aspects of cactus diversity in an Open Access resource,
CactEcoDB enables new opportunities for research on trait-environment relationships,
climatic niche evolution, and conservation, in one of the most unique and threatened
plant families. Given the gradual increase of trait, spatial and phylogenetic data
generally, we welcome the submission of additional data, which will improve future
research into cacti.

Methods
The methods presented here include a brief description of methods used to generate
previous data [10], and our new efforts designed to improve and expand sampling of
spatial distributions, environmental data, and traits (Figure 2). Although (10) is a large
foundation of this work, we added ten new variables, recollected or added data to 39
traits (76.40% of CactEcoDB), and undertook thorough checking steps.

Phylogenetic reconstruction
We provide two phylogenetic hypotheses. One is the original phylogeny estimated by
Thompson et al. [10] (henceforth referred to as V1), and one is re-estimated using a
phylogenomic constraint that has been published since [89]. For both trees, the
supermatrix approach was used, to maximise taxonomic sampling using publicly-
available data in Genbank. 18 published plastid and nuclear loci were compiled from

GenBank, by identifying and clustering orthologous sequences with the OneTwoTree
pipeline [90], and aligned them using MAFFT [91] with quality inspection undertaken in
SeaView [92]. Partial sequences were merged with full-length sequences for each species
where homologous sequences clustered into a fragmented and full cluster, retaining the
longest version. Outgroup sequences from Anacampserotaceae, Portulacaceae, and
Talinaceae were added using MAFFT's "--add" function [93]. After alignment, poorly
aligned regions were trimmed using trimAl with the "gappyout" setting [94], and
concatenated all loci into a supermatrix using AMAS [95]. A Maximum Likelihood (ML)
phylogeny was reconstructed with RAxML v8 [95,96], applying a GTR+G model partitioned
by locus and assessing support with 300 bootstrap replicates. Time calibration was
performed under Penalized Likelihood criteria with treePL, using secondary constraints
on crown and stem age for the family derived from highest posterior densities from a
relaxed-clock fossil-calibrated phylogeny of angiosperms [5] (see [10] for full details), in the
absence of an informative fossil record of Cactaceae [55]. Cross-validation was used to
optimize the smoothing parameter across 100 priming runs, and the final dating was
based on the smoothing value with the lowest score.

To make V2, we replicated these steps but constrained relationships in the Maximum
Likelihood tree search according to a recently published phylogenomic backbone [89]. We
assessed relationships in this phylogenomic tree in the context of node support, and
identified 15 nodes to constrain. These were tribes Blossfeldioideae, Cacteae, Cereeae,
Copiapoeae, Cylindropuntieae, Fraileeae, Lymanbensonieae, Notocacteae, Opuntieae,
Phyllocacteae, Pterocacteae, Rhipsalideae, and subfamilies Maihuenioideae,
Pereskioideae and Leuenbergerioideae. This approach greatly improved relationships
compared to the old phylogeny, given the low sequence variation found in cacti which
commonly leads to weakly-supported nodes [10,52,55]. The number of nodes supported by
70% bootstrap support in V2 (BS) increased from V1 by 11.8% (32.3 to 36.1%), and the
number of nodes supported by 90% increased by 72.9% (11.9 to 20.6%). Eleven
species displayed pathologically long terminal branches when the constraint was
applied and were pruned.

Diversification rate estimation
To estimate speciation rates from the phylogenies we used three methods: 1). Bayesian
Analysis of Macroevolutionary Mixtures (BAMM) [97] (Figure 1), 2). Missing State
Speciation and Extinction (MiSSE) [98], and 3). the DR statistic [35]. Each of these have
different assumptions and approaches, and associated benefits and pitfalls. BAMM is a
Bayesian approach which discretises rates in the phylogeny into lineage-specific shift
regimes, and provides a posterior sample to account for uncertainty. It also allows
lineage-specific sampling fractions, which are of great importance given the
phylogenetic imbalance of molecular sequencing across cactus groups [10,52]. Although

BAMM has received criticism, notably regarding sensitivity to priors and the accuracy of
deep rate shifts [99,100], these have been defended statistically [101,102] and it remains one of
the most accurate methods [103]. MiSSE also discretises rates, but in a Maximum
Likelihood framework providing only one estimate per model, and accounts for
incomplete sampling with a global sampling fraction. While this does not account for
within-group variation in sequencing efforts, lineage-specific sampling fractions have
been criticised [104]. Finally, the DR statistic allows rapid estimation as it only makes use
of variation in terminal branch lengths and root-to-tip node count. However, it cannot
account for incomplete and imbalanced sampling, and is known to be noisy compared
to other metrics [105]. We recommend using BAMM estimates, as BAMM accounts for
incomplete sampling in a manner that does not assume that each lineage has been
sampled to an identical fraction, as well as providing a posterior sample of estimates
which accounts for uncertainty.
174
For phylogeny V1, we used the BAMM results from Thompson et al. [10]. We replicated
this process with V2. Prior to analysis with BAMM, outgroups were pruned. Incomplete
sampling was accounted for in BAMM by providing genus-level sampling fractions. Four
Markov Chain Monte Carlo (MCMC) chains were run for 50 million generations with
BAMM, sampling every 5,000 generations and discarding the first 10% as burn-in.
Priors were set using the BAMMtools package in R [106], and a conservative prior
allowing for a single rate shift was chosen to minimise overfitting. Convergence was
assessed using the coda R package, ensuring effective sample sizes (ESS) of >200 for
all key parameters. Following this, posterior distributions were summarized with
BAMMtools, the best shift configuration was estimated for visualisation, and mean
speciation rates at the tips were extracted for each species. As discussed in [10]), we
provide the entire event data file from BAMM to enable analysis of lineage
diversification dynamics, with the caveat that exact divergence timings in cacti are
highly uncertain due to the absence of informative fossils [55].
189
For the MiSSE analyses we used a global sampling fraction of 0.575 (phylogeny V1)
and 0.572 (phylogeny V2), respectively, and estimated 30 possible model structures
using the R package HiSSE [107], which were generated with the function
generateMiSSEGreedyCombinations from combinations of between one and ten
turnover parameters, and one and three extinction parameters. Using the function
MiSSEGreedy, we estimated MiSSE parameters on each of the 30 models, then
estimated ancestral diversification rates using the command MarginReconMisse.
Finally, we estimated tip rates using the function GetModelAveRates on the marginal
reconstruction results. We replicated this on a version of each phylogeny that has the
shortest branches pruned, using a threshold of 0.01 Mya, because formal -SSE models
can be biassed by tiny branch lengths [107,108]. In cacti, these could be real (recent

divergences) or an artefact, and it is difficult to unambiguously classify them. In V1, 41 species were pruned and in V2, 70 species were pruned. We present results from both the unpruned and pruned phylogenies. For the DR statistic analysis, we used the R function DR_statistic (github.com/Cactusolo/rosid_NCOMMS-19-37964-T).

Figure 1: The phylogeny (V1), showing estimated speciation rate variation in one method (BAMM). Phylogenetic branches in phylogeny V1 [10] are coloured according to speciation rates estimated with BAMM [97], and vary 32-fold. Arc segments of median speciation rate for thirteen morphologically varied cactus genera are indicated. This figure and figure legend is reproduced with permission from [10]. Cactus images are used under Creative Commons with modifications allowed. From left to right: images 1, 3, 8, 11, 12, and 13 used photos taken by Amante Darmanin, Forest & Kim Starr, John Tann, Renee Grayson, and Wendy Cutler, which are licensed under a Creative Commons Attribution 2.0 License (https://creativecommons.org/licenses/by/2.0/). Image 2 used a photo marked as being in the Public Domain (https://creativecommons.org/publicdomain/mark/1.0/). Images 4 and 10 used photos taken by Leonora Enking and Lyubo Gadzhev, which are licensed under a Creative Commons Attribution-ShareAlike 2.0 License (https://creativecommons.org/licenses/by-sa/2.0/). Images 5, 7 and 9 used photos marked as being in the Public Domain using the CC0 1.0 Universal Public Domain Dedication (https://creativecommons.org/publicdomain/zero/1.0/). Image 6 used a photo taken by Christer Johansson, which is licensed under a Creative Commons Attribution 3.0 Unported License (https://creativecommons.org/licenses/by/3.0/deed.en).

Trait data compilation

We compiled species-level trait data from a combination of authoritative textbooks [109],
primary taxonomic literature, specimen descriptions in online floras, and prior datasets
[10,51,60]. These data vary in their coverage across cactus species (Figure 2). Key traits
that we updated, expanded, and standardised in our new dataset (i.e. beyond [10]),
include plant height, growth form, pollination syndrome, and chromosome count. We
standardised traits previously recorded with different units, e.g. for height (to cm).
Where minimum and maximum plant heights were reported, we recorded maximum
values to reduce biases from juvenile, diseased, or partial specimens, as is standard
practice in comparative research into plant height [110]. We expanded and updated all
records for plant maximum size to improve species sampling by 187 species, from 750
to 937.

240
Figure 2: The distribution of species coverage by variable. *Speciation rate*
estimates, growth form and plant size (maximum height or length) have highest
coverage (1,063-982, 1,061 and 937 species, respectively), and the spatial
environmental variables sample 904 species. We only calculated range size for those
species in IUCN, given the occurrence scarcity in GBIF for the remaining species.
Pollinator and chromosome count have lower coverage with 327 and 378 species,
respectively. We compare the species coverage for variables in CactEcoDB against
those in the largest previous dataset [10].

249
250 We provide data on the binary growth form scoring system developed by Hernández-
251 Hernández et al. [51] and extended in sampling by [10]. This system included two main

252   categories grouping six subcategories: compact including smaller forms such as
253   globose and barrel) and arborescent (containing larger forms such as columnar and
254   shrubby).
255
However, we have devised a novel and more refined system using nine growth form
categories for CactEcoDB: arborescent, barrel, columnar, cushion forming, epiphytic,
geophytic, globose caespitose, globose solitary, and shrubby. For context, a previous
growth form system used in (Hernández-Hernández et al., 2011) scored species as
arborescent, shrubby, columnar, globose solitary, globose caespitose and barrel.
Briefly, arborescent cacti are those with a main trunk that branches above the base,
typically with a tree-like structure; shrubby cacti lack a dominant trunk and instead
exhibit basitonic branching from the base; columnar species are unbranched with
elongated stems, including tall or prostrate cylindrical forms; globose solitary and
globose caespitose refer to spherical-stemmed species under 0.5 m tall, growing singly
or in clumps, respectively; barrel cacti have spherical stems exceeding 0.5 m in height;
cushion-forming species grow in dense mats without a clearly assignable alternative
form; geophytic species grow at or below ground level but are not cushion-forming; and
epiphytic species are those growing on other plant hosts. We hypothesise that this
system of nine growth forms adequately explains growth form variation in cacti, but note
there is uncertainty and other systems exist [60,71]. We scored these species based on
botanical descriptions primarily drawn from [109], but also from other sources where
unavailable or inadequately described in [109] (Figure 3).
274

275
276
Figure 3: Examples of each growth form in CactEcoDB. *(a)* Rhipsalis baccifera*, an*
epiphytic cactus. (b) Echinocactus grusonii*, a barrel cactus. (c)* Cephalocereus
columna-trajani*, a columnar cactus. (d)* Mammillaria tetrancistra*, a globose solitary*
species. Note that we have not shown a globose caespitose species, which appears as
clusters of globose solitary species. (e) Maihuenia poeppigii*, a cushion-forming species.*
(f) Opuntia microdasys*, a shrubby species. (g)* Ariocarpus fissuratus*, a geophyte. (h)*
Pachycereus pringlei*, an arborescent species. Cactus images are used under Creative*
Commons licenses with modifications allowed. Image (a) used a photo taken by Maria
Vorontsova, which is licensed under the Creative Commons CC0 1.0 Universal Public
Domain Dedication (https://creativecommons.org/publicdomain/zero/1.0/). Images (b)
and (g) used photos taken by Dr. Hans-Günter Wagner, which are licensed under a
Creative Commons Attribution-ShareAlike 2.0 License
(https://creativecommons.org/licenses/by-sa/2.0/). Images (c) and (h) used photos taken
by Amante Darmanin, which are licensed under a Creative Commons Attribution 2.0
License (https://creativecommons.org/licenses/by/2.0/). Image (d) used a photo taken
by Jesse Pluim (BLM), which is marked as being in the Public Domain using the Public

We compiled pollination data and classified cactus species into five functional pollinator groups: bats, bees, birds (which are primarily Trochilidae), moths (which are primarily Sphingidae), and other insects (Figure 4). This level of resolution reflects the standard used in pollination research for cacti and other plant groups [51,111–114], and greatly improves on the binarisation approach (ancestral versus derived) established by [10,51]. In botanical data, observations and inferences of pollinators are typically reported at taxonomic levels higher than species, as we have provided here, and lower taxonomic levels are inconsistent and rarer. The pollination syndromes themselves are shaped by shared floral adaptations that map reliably to higher-level clades (e.g., orders or functional groups) rather than to individual pollinator species [115–117]. Species-level identification is uncommon in the botanical literature due to both data limitations and the diffuse, coevolutionary nature of many plant–pollinator interactions [118,119]. To assemble these classifications, we combined information from existing compilations at genus, tribe and whole-family levels (e.g. [120–122]) with targeted literature searches using query combinations such as "genus + pollinator" and "genus + pollination." All pollination assignments in [10], which used data primarily provided by [51], were re-assessed and 56 species were removed at the higher taxonomic level because their source could not be verified beyond [51]. By systematically validating records against traceable literature and retaining only those with adequate support, our dataset provides a foundation for comparative research as well as future data collection. Our pollination data covers 17.68% of cacti, a larger proportion than a recent compilation for Orchidaceae (~10%) [51,114].

Finally, we captured data on chromosome counts from the Chromosome Count Database (https://ccdb.tau.ac.il/) [123], which compiled data from other sources and involved a cleaning step. We retained all recorded counts instead of relying on the reported medians (Figure 5). As well as allowing within-species variation to be captured, this allowed the correction of some zero median counts reported for species with chromosome counts in the database. However, we also replicate the median chromosome count for each species as it is useful for macroevolutionary research [124–126]. We performed statistical and manual checks of collected trait data which flagged multiple records as spurious, which we either removed, confirmed or edited based on deeper literature searches (see Technical Validation).

Figure 4: The distribution of plant height, speciation rate, growth form, chromosome count, and pollinator. *(a) Cactus size (height or length) is bimodal, which is shaped by underlying growth form variation [10]. (b) Tip speciation rate is mostly relatively slow, with fewer relatively faster species. (c) The most common growth form is globose solitary, followed by globose caespitose and shrubby. The least common growth forms are cushion forming, geophytic and barrel. 222 species are polymorphic,*

which mostly consists of species which are both globose solitary and globose
caespitose (154 species), or both arborescent and shrubby (50 species). (d) Most
species have the base number of 11 chromosomes, and most chromosome count
variation is generated by polyploidy [74], reflected in the distribution (e.g. 22, 33, 44, 55).
(e) Most species are pollinated by derived pollinators (bats, birds, moths) rather than the
ancestral pollinators (bees) [51].

Distribution data

Previous distribution range datasets used data downloaded and cleaned from the
Global Biodiversity Information Facility (GBIF, https://www.gbif.org/). However, we have
greatly improved species and distributional sampling with a new sampling strategy
(Figure 5). For CactEcoDB, we have primarily used expert range polygons available in
the IUCN Red List of Threatened Species (https://www.iucnredlist.org/) [70,127], with some
occurrences newly-acquired and curated from GBIF to maximise species sampling,
following [128]. This approach increases species sampling by 54 species (~6%) and
provides more accurate estimates of geographic range sizes, which allows us to fully
characterise the environments of the entire ranges of species. Contrary to previous
datasets, our new data reduce geographic bias, especially for species with poor GBIF
records.

We downloaded IUCN ranges and manually checked them against the independent
data sources available in POWO (https://powo.science.kew.org/) which records native
and introduced ranges at a broad level [88]. This identified one IUCN range that had
sampled likely-introduced South American populations (for *Cylindropuntia tunicata*). We
corrected this accordingly and then we sampled coordinates randomly from within each
range with the number of coordinates sampled in relationship to the range size,
following an established protocol [128]. Specifically, we used a binning function to assign
sample sizes based on range area (in km²): species with ranges <10 km² were
sampled 20 times; 10–100 km², 50 times; 100–1,000 km², 150 times; 1,000–
10,000 km², 300 times; 10,000–100,000 km², 500 times; 100,000–1,000,000 km²,
700 times; 1,000,000–10,000,000 km², 900 times; and >10,000,000 km², 1,200 times.
This sampling approach ensured adequate environmental representation for both
narrow and wide-ranging species, while avoiding excessive data volume or downstream
computational demands, particularly for the most widespread taxa.

We curated the most up-to-date GBIF occurrences, for those species not covered by
the IUCN expert ranges. We used a rigorous and multi-step cleaning pipeline developed
by [129,130]. We queried all records under "Cactaceae", considering only "Preserved
Specimens" and removed species present in the IUCN expert range data. The code
checked taxonomy against POWO and flagged suspect records for removal based on

multiple criteria: i) records with missing or badly formatted coordinates; ii) records
associated with biodiversity institutions; iii) records with equal or zero latitude and
longitude; iv) records within a 10km radius of a country's capital; v) records within 5km
of a country's political centroid; vi) records in the sea [131]. As a final step, occurrence
records that were outside the range described by POWO for each species were
removed [88].
388

389
Figure 5: The spatial distribution of cactus richness and tip speciation rates. *(a)*
Species richness is plotted per grid cell (~50*50 miles), and shows Mexican, Eastern
Brazilian and Andean richness hotspots. (b) Median tip speciation rate (BAMM_V1 from
[10]) is plotted per grid cell, showing a relative decoupling between areas of highest
species richness and high speciation rate. This figure was presented in the
supplementary materials of [10], but with lower resolution and different underlying spatial
data.
397

Environmental data and verifying against GBIF data

We extracted spatial variables hypothesised to be important for cacti, succulents, and
angiosperms generally (summarised in [10]), from three different sources. From CHELSA
[132] (https://www.chelsa-climate.org/), we extracted the 19 bioclimatic variables, aridity
index, potential evapotranspiration and net primary productivity. From EarthEnv [133]
(https://www.earthenv.org/), we extracted elevation and six measures of topographic
complexity including slope, roughness, standard deviation of slope, and profile
curvature. From SoilGrids [134] (https://isric.org/), we extracted sand content, water
availability, and texture. We calculated median values per species for continuous
variables, since they are less likely to be influenced by outliers, and also provide the

entire distribution for each species across their range, to capture intraspecific variation.
The latter is especially valuable when analysing trait–environment relationships,
because it accounts for environmental heterogeneity within species ranges, instead of
relying on a point estimate. Incorporating intraspecific variance can lead to more
realistic ecological and evolutionary inferences [135–138]. As with the trait data, we
performed statistical validation analyses. We did this to identify the congruence between
our data and that of the largest previously-assembled dataset of cactus distributions [10]
(details in Technical Validation).
416
Data Records
All data files described below are available in the GitHub repository
(https://github.com/jamie-thompson/CactEcoDB) and Figshare
(https://doi.org/10.6084/m9.figshare.30940019) [139]. The dataset comprises cleaned
species-level trait, spatial, environmental, phylogenetic, and diversification rate data for
up to 1,063 species of Cactaceae (Figures 1, 2 and 5). Each file is formatted for ease of
integration into ecological and evolutionary analyses.
424
1. CactEcoDB_v1.csv
Format: Spreadsheet (.csv)
Description:
This file contains species-level data for Cactaceae, integrating trait, phylogenetic,
spatial, and environmental information. Specifically, it includes:
● Mean tip-specific speciation rates estimated using BAMM, MiSSE and DR
(lambda)
● Morphological and ecological trait data, such as maximum plant height (cm),
growth form (binary and refined), pollination syndrome, and geographic range
size (km^2)
● Summary statistics (typically medians) of environmental variables based on
georeferenced occurrence data
437
Fields:
● Diversification
○ BAMM_lamba_V1: Mean species-specific speciation rate estimated using
BAMM [97] for phylogeny V1
○ BAMM_lamba_V2: Mean species-specific speciation rate estimated using
BAMM for phylogeny V1
○ MiSSE_lamba_V1_full: Speciation rate estimated using MiSSE [98] for
phylogeny V1
○ MiSSE_lamba_V1_pruned: Speciation rate estimated using MiSSE for
phylogeny V1

- - - MiSSE_lamba_V2_full: Speciation rate estimated using MiSSE for phylogeny V2
    - MiSSE_lamba_V2_pruned: Speciation rate estimated using MiSSE for phylogeny V2
    - DR_lambda_V1: Speciation rate estimated using the DR statistic [35] for phylogeny V1
    - DR_lambda_V2: Speciation rate estimated using the DR statistic for phylogeny V2

- Morphological and ecological traits
    - growth_form_binary: Coarse growth form classification (arborescent vs. compact group), following [10,51]
    - growth_form_detailed: Detailed growth form description based on literature (arborescent, barrel, columnar, cushion forming, epiphytic, geophytic, globose caespitose, globose solitary, shrubby)
    - max_size_height_or_length: Maximum reported plant height or stem length (cm), building on [10,51]
    - pollinator_binary: Pollination syndrome grouped into ancestral (bee) vs. derived (bat, bird, moth, other insects), according to [10,51]
    - pollinator_detailed: Detailed pollination syndromes (bat, bird, moth, other insects)
    - chromosome_count_median: Median chromosome count (n)
    - chromosome_count_distribution: Summary information or distribution notes on chromosome counts (n) sourced from the Chromosome Counts Database [123]
    - range_size: Geographic range size estimates (km^2)

- Environmental variables (CHELSA bioclimatic variables [132], SoilGrids [134], and Global Multi-resolution Terrain Elevation Data (GMTED) [133,140])
    - Bio1–bio19: Standard bioclimatic variables, including temperature (bio1-bio11) and precipitation metrics, extracted from CHELSA [132]. Bio1 to bio11 are temperature-related variables, and bio12-bio19 are precipitation-related. For example:
        - bio1: Annual mean temperature (°C)
        - bio2: Mean diurnal temperature range (°C)
        - bio3: Isothermality (compares diurnal and annual temperature) (100 × (bio02 ÷ bio07))
        - bio4: Temperature seasonality, the standard deviation of mean monthly temperatures (100/°C)
        - bio5: Maximum temperature of warmest month (°C)

- bio6: Minimum temperature of warmest month (°C)
- bio7: Temperature annual range (°C)
- bio8: Mean temperature of wettest quarter (°C)
- bio9: Mean temperature of driest quarter (°C)
- bio10: Mean temperature of warmest quarter (°C)
- bio11: Mean temperature of coldest quarter (°C)
- bio12: Annual precipitation (kg m^{-2} $year^{-1}$)
- bio13: Precipitation of wettest month (kg m^{-2} month^{-1})
- bio14: Precipitation of driest month (kg m^{-2} month^{-1})
- bio15: Precipitation seasonality (unitless): Coefficient of variation (100 × SD ÷ mean) of monthly precipitation
- bio16: Precipitation of wettest quarter (kg m^{-2} month^{-1})
- bio17: Precipitation of driest quarter (kg m^{-2} month^{-1})
- bio18: Precipitation of warmest quarter (kg m^{-2} month^{-1})
- bio19: Precipitation of coldest quarter (kg m^{-2} month^{-1})
- (See CHELSA documentation for full descriptions, especially with regards to scales and offsets used in some variables [132])
- Aridity and productivity
  - aridity_index: Aridity index (mean annual precipitation/potential evapotranspiration)
  - pet: Potential evapotranspiration (kg m^{-2} $year^{-1}$)
  - npp: Net primary productivity (g C m^{-2} yr^{-1})
- Elevation and topography extracted from GMTED [133,140]
  - elevation: Median elevation across occurrence records (m)
  - slope: Median slope of the terrain (°)
  - sd_slope: Median standard deviation (sd) of slope
  - sd_elev: Median standard deviation of elevation
  - roughness: Median terrain roughness (largest inter-cell difference of a focal cell and its 8 surrounding cells, m)
  - prof_curve: Median standard deviation of profile curvature (sd of radians per meter)
  - tang_curve: Median standard deviation of tangential curvature (sd of radians per meter)
- Soil Variables extracted from SoilGrids [134]
  - soil_sand: Median proportion of sand particles (> 0.05/0.063 mm) in the fine earth fraction
  - soil_water: Median derived available soil water capacity (volumetric fraction)
  - texture: Soil texture class, classified into twelve categories (e.g. sand, silt, clay, loam, silty clay)

Notes:

• Speciation rates estimated with BAMM represent mean species-specific
estimates from the entire posterior sample. The full BAMM posterior event data
objects are available separately in CactEcoDB, and can be analyzed (e.g. rates
through time and deep rate shifts) using the R package BAMMtools [106].
• Plant height values are not statistical species means from population samples,
but are derived from maximum reported sizes in taxonomic or ecological
literature, as is common across plant research generally [10,110,141–144].
• Pollinator data was rarely specific to individual species of pollinator. Therefore,
we classified them into major taxonomic groups described in [10,51] as is standard
practice in pollinator research [51,111–114].

2. CactEcoDB_coords_envs.csv
Format: csv
Description: Curated distribution coordinates with associated environmental values for
every coordinate.
Fields: species, decimal latitude, decimal longitude, environmental variables (e.g. bio1-
19, aridity index, soil texture).
Notes: Most of these coordinates are randomly sampled from expert IUCN ranges,
some of them are curated GBIF occurrences. We have written permission from the
IUCN Red List to reproduce randomly-sampled coordinates from expert ranges.

3. CactEcoDB_phylogeny_V1.tre
Format: Newick (.tre)
Description: Time-calibrated phylogeny for 1,063 Cactaceae species, inferred using a
supermatrix approach and Penalized Likelihood dating.
Notes: Outgroups are pruned and tips are matched to species names in the trait and
spatial datasets to facilitate research.

4. CactEcoDB_phylogeny_V2.tre
Format: Newick (.tre)
Description: Time-calibrated phylogeny for 1,052 Cactaceae species, inferred using a
supermatrix approach and Penalized Likelihood dating.
Notes: This differs from the V1 Phylogeny, in that is uses a phylogenomic backbone to
improve relationships [89].

5. CactEcoDB_concatenated_alignments.fasta
Format: Fasta (.fasta)
Description: concatenated alignment of 18 loci, first used in Thompson et al. [10].

Notes: Locus regions are defined in CactEcoDB_phylo_partition.txt.
569
570   **6. CactEcoDB_nucleotide_accessions.csv**
571   Format: csv
572   Description: A species-by-locus matrix of GenBank accessions used to make both
573   phylogenies. Originally assembled by Thompson et al. [10]
574
575   **7. CactEcoDB_phylo_partition.txt**
576   Format: RAxML-format partition file (.txt)
577   Description: Defines the start and end of loci in
578   CactEcoDB_phylo_concatenated_alignment.fasta, used to apply a different nucleotide
579   substitution model to each locus.
580
8. CactEcoDB_phylogeny_V1_bootstrap.tre
Format: Newick (multiple phylogenies)
Description: The sample of 300 bootstrap replicates estimating alongside the maximum
likelihood search for phylogeny V1 estimated by Thompson et al. [10]
Notes: This can be used to assess topological support and replicate analyses while
accounting for phylogenetic uncertainty
587
9. CactEcoDB_phylogeny_V2_bootstrap.tre
Format: Newick (multiple phylogenies)
Description: The sample of 300 bootstrap replicates estimating alongside the maximum
likelihood search for phylogeny V2
Notes: This can be used to assess topological support and replicate analyses while
accounting for phylogenetic uncertainty
594
10. CactEcoDB_BAMM_eventdata_V1.txt
Format: txt
Description: The posterior sample of rate shift configurations (eventdata) estimated by
BAMM.
Fields: specific to BAMM [97], use the R package BAMMtools [106] to access and analyse
this.
Note: We do not provide this for MiSSE and DR, for which we provide tip-speciation
rates in the main CactEcoDB file. BAMM is a flexible system for rates through time and
rate shifts, therefore we provide the intermediary event data file. Tip-speciation rates
calculated from BAMM event data files are already present in the main CactEcoDB file.
605
11. CactEcoDB_BAMM_eventdata_V2.txt
Format: txt

Description: The posterior sample of rate shift configurations (eventdata) estimated by
BAMM.
Fields: specific to BAMM [97], use the R package BAMMtools [106] to access and analyse
this.
612
613 **12. CactEcoDB_variable_descriptions.txt**
614 Format: txt
615 Description: Definitions, units, and data sources for all variables in the dataset.
616 Notes: Intended to support reproducibility and proper variable interpretation.
617
618 **13. CactEcoDB_full_sources_shorthand.xlsx**
619 Format: xlsx
620 Description: References (in shorthand, see CactEcoDB_full_references_lookup.xlsx file)
621 for all variables in CactEcoDB.
622 Notes: Intended to support transparency, traceability and reliability.
623
624 **14. CactEcoDB_full_references_lookup.xlsx**
625 Format: xlsx
626 Description: A lookup table matching the shorthand reference style (e.g. IUCN,
627 CHELSA, BAMM) with full references.
628 Notes: Intended to support transparency, traceability and reliability.
629
15. CactEcoDB_code.zip
Format: compressed folder (ZIP) of code scripts
Description: Contains scripts (primarily R) for data assembly, random sampling of
spatial coordinates, taxonomic and spatial distribution cleaning, estimation of
diversification rates, post-processing (e.g. for BAMM), outlier detection, and
visualisation.
636
637 **Technical validation**
638 To ensure the robustness and reliability of CactEcoDB, we implemented multiple
639 validation steps integrating automated and manual approaches, which we applied to the
640 phylogenies, diversification estimates, trait data, species distributions, and
641 environmental summaries.
642
Phylogenies and diversification rate estimates
It is well known that phylogenetic relationships in cacti are difficult to resolve due to their
low sequence divergence [10,51,52,55,60,89]. There are different approaches available, such
as constructing phylogenies for individual groups (e.g. genera and tribes [64,65,145]), deeply
sampling entire genomes to maximise backbone resolution [89,146], and the supermatrix

approach, which maximises species sampling at the cost of the support provided by
genome-wide data [21,147-150]. Here, we have provided preliminary phylogenies using the
supermatrix approach, in order to maximise species sampling for comparative research,
but note that caution should be taken, as with all supermatrix phylogenies [151-155]. The
phylogenies were constructed using 18 multiple sequence alignments of commonly
sequenced nucleotide loci publicly available in Genbank. All alignments were manually
and statistically inspected, and corrected for quality and taxonomic coverage (see [10]).
Maximum Likelihood phylogenetic reconstruction was performed with partitioned models
to account for variation in substitution rates between loci, and bootstrap support was
calculated. In V1, a relaxed constraint was used to improve the likelihood computation
and in V2, we made use of a recently-published phylogenomic tree of cacti [89], in which
we implemented a much stricter topological constraint. Using both a largely data-driven
reconstruction (V1) and a topology guided by a recent phylogenomic framework (V2)
provides complementary strengths. V1 maximises the signal present in the sequence
data, while V2 enforces well-supported relationships among deeper nodes, thereby
improving overall phylogenetic stability. For users of the data wishing to explore the
phylogenetic uncertainty, we provide 300 bootstrap replicates for each topology, as well
as the full alignment and partition file. Time-calibration used secondary divergence
estimates, as there is no useful fossil record for cacti [55]. Resulting time-calibrated
phylogenies are the widest-sampled for cacti, encompassing all major lineages,
although sampling remains sparse in some poorly-sequenced lineages. Despite
incomplete sampling and inherent limitations of time-calibration, the topologies and
divergence timings are broadly consistent with other recent reconstructions, which
increasingly converge on a crown age of ~27-37 Mya for the family [5,51,55,146]. Our
phylogenies, like all divergence-time estimates lacking direct fossil calibration, carry
uncertainty in absolute node ages; however, their relative branch lengths and overall
temporal structure are robust and well suited for comparative analyses such as tip-rate
estimation, diversification modelling, and phylogenetic correction [10].
676
We provided multiple estimates of diversification rate to account for methodological and
topological differences. We provided BAMM estimates, followed established statistical
protocols when setting conservative priors [106], accounted for imbalance sampling efforts
at a high resolution of genus-level (fractions available at [10]), and ensured convergence
was reached. We provided MiSSE estimates as an alternative [98], which allows users to
test whether results change according to method, especially differences in the
implementation of corrections for incomplete taxonomic sampling. We also provide the
DR statistic [35], which makes fewer assumptions regarding the evolutionary model
underlying phylogenetic branching, but is noisy and cannot account for incomplete
sampling. Diversification rates estimated with MiSSE used a global sampling fraction to
account for incomplete sampling, and were repeated with and without the shortest

branches. We tested the congruence of all eight estimates (four per phylogeny: BAMM,
DR statistic, MiSSE with all species and MiSSE without short terminal branches).
Pairwise Pearson correlations of log-transformed tip rates revealed strong and
consistent agreement across approaches. All pairwise comparisons were strongly
significant ($p < 0.000001$), and correlation coefficients ranged from 0.24 to 87 (mean r =
0.61, median r = 0.64). We advise that researchers use multiple approaches to
investigating diversification rates, with an understanding of the uncertainties and
methodological differences between methods.
696
Trait data
As with similar trait datasets [14,25,26,28,29,156], when collecting trait data, we relied on
authoritative textbooks and peer reviewed literature, using secondary sources (e.g.
online floras) infrequently, and only when it provided a sensible value (i.e. not when
photographs of a cactus is clearly miniature but the recorded plant height is
unrealistically large). We used standardised measurement protocols to minimise errors,
such as unit conversion errors that are present in some databases.
704
Once collected, trait data were checked for plausibility and consistency through a
combination of automated flagging, literature cross-checking and expert review.
Generally, minimum and maximum plant height values were most frequently recorded
and individual level variation is not available, as is common in other groups [17,42,141]. We
retained maximum values, as is best-practice in comparative research on plant height to
avoid issues associated with diseased, juvenile or incomplete specimens [110,141]. Height
values that were extreme (e.g. 0cm, 2500cm) or contradictory were flagged and verified
against alternative sources when available. Growth form data were collected and
discussed between authors in accordance with our defined groupings. We flagged
records with outlying plant height for their growth form, on the principle that growth form
shapes height (i.e. globose species are much smaller than arborescent species, and
shrubby species and somewhere between). Outlier detection was performed within
each growth form, and polymorphic species were expanded so that their height values
were evaluated against every growth form they express. We defined outliers following
the standard interquartile method and inspected 75 that were detected. Of these, 11
were removed, three were changed, and 61 were kept as-is. Following this, an author
with 30 years experience in growing and over 30 field work trips to diverse cactus
ecosystems (AG), checked all the data. Pollinator data required significant additional
validation. A portion in Thompson et al. [10] originated from Hernández-Hernández et al.
[51], where many pollination syndromes were provided without traceable primary sources.
To assess the reliability of these data, we searched independently for evidence for each
species using published ecological studies, species-level monographs, and an
alternative collection which recorded sources [120], the latter of which we incorporated into

CactEcoDB. Where independent sources were available, we updated or confirmed the
pollinator data accordingly. For the 56 cases where no primary or secondary
documentation could be found, we removed these from the dataset. The retained
categories follow the definitions widely used in previous macroevolutionary work on
Cactaceae [51,60].
733
As a final validation step for plausibility, one author (AG), who has over 30 years
experience in the cactus horticulture industry and has undertaken 30 field trips to visit
diverse cactus habitats in the Americas, checked all trait data one by one. This expert
review step identified additional implausible datapoints that were not feasible by
statistical testing and literature checkers. Notably 28 species were flagged by AG, many
of which because their heights were likely misrecorded by Anderson [109] based solely on
single stem segment measurements. A similar issue was identified in Anderson [109] for
one species (*Opuntia chaffeyi*), where the reported height is ambiguously recorded and
may correspond to the size of a rhizome instead of the entire plant. These were not
possible to verify independently and were removed.
744
As with all comparative datasets, especially for plants [13,42,47,157], there are gaps owing to
imbalanced data collection efforts. The poorest-covered traits are pollinator and
chromosome count, because these are difficult to collect in the field compared to growth
form and plant height descriptions. Although we note that for pollinators, CactEcoDB
provides observations for a large proportion of species when compared to other recent
datasets for plant families (17.68% of species compared to ~10% for Orchidaceae [114]).
Gaps in pollinator observation data are fairly group-specific; there are relatively more
species with data in Cactoideae than in Opuntioideae, for example, and more in
Cactoideae tribes Cereeae and Phyllocacteae, than Cacteae. In contrast, chromosome
count data collection efforts seem less biased by group, and both growth form and plant
height are fairly well covered as commonly-recorded variables.
756
Spatial distributions and environmental variables
Spatial distributions were made of a combination of high resolution expert-ranges
(iucnredlist.org) and a smaller number of curated occurrences where expert ranges
were unavailable, following established protocols [128]. Randomly sampled coordinates
from expert ranges were visually inspected per species, which identified one unlikely
range that is the result of an introduced population. Occurrences from GBIF were
thoroughly checked following established protocols [109,129,130], which performed a six step
procedure, flagging coordinates based on spatial and statistical issues [131], and then
filtering according to independent estimates of distributions, broadly corresponding to
country scale [88]. These two complementary approaches maximise coverage and
reliability. Expert-defined range polygons provide consistent distributions for most

species, while rigorously filtered occurrence data extend sampling to species lacking
expert maps. As with all large-scale distribution datasets, ranges should be viewed as
the best current synthesis and may change as taxonomy is revised, population shifts
are documented, introductions are identified, and ongoing IUCN assessments and
occurrence data updates redefine the distributions. As with the phylogenies, we will
update the distributions in the future as new data compilations emerge.
774
Environmental variables were extracted from high-resolution raster datasets (e.g.
CHELSA [132], SoilGrids [134] for each coordinate, and species-level summaries were
calculated as medians to minimise the influence of outliers following [10]. Critically, we
statistically compared values of extracted spatial data between previous methods that
employed GBIF-derived data [10] and our current method, and found very strong
agreement, demonstrating that our approach of sampling from expert IUCN ranges is
appropriate. We estimated 1. linear regressions and 2. Pearson correlation coefficients
between variables in CactEcoDB and in the largest previous spatial dataset for cacti
(Thompson et al., 2024). We found that comparisons between all 31 environmental
variables are significant at $p < 0.05$, with a median slope and correlation coefficient of
0.89 and 0.88, respectively. Furthermore, 25 comparisons recovered a slope and
correlation coefficient of > 0.75. Our new approach also retrieved valid data for soil
texture, which is discrete and has strong agreement (72.60%) with data retrieved from
GBIF occurrences (Thompson et al., 2024).
789
Usage Notes
CactEcoDB is the first large-scale integrative database for the cactus family, and brings
together multiple data types that are rarely available in combination for any plant family,
including species-level traits, expert-curated spatial distributions, environmental
variables, time-calibrated phylogenies and speciation rate estimates. Such a synthesis
has not been previously available for cacti, despite their interest to biologists due to their
exceptional morphological and ecological diversity, and status as one of the most
threatened groups worldwide [70]. By assembling these diverse data sources, CactEcoDB
enables comparative, macroecological, macroevolutionary and conservation analyses
that were previously impractical or impossible due to fragmented data.
800
As is typical of large-scale biodiversity datasets, CactEcoDB reflects uneven data
availability across species and data types [47,158]. Trait coverage is extensive across the
family, but completeness varies by trait and specific group. For plant height, values
were derived from sources that only reported minimum and maximum height. In these
cases, maximum values were retained to reduce bias from juvenile, incomplete, or
diseased specimens. However, these records often lack associated metadata (e.g.,
number of individuals measured, habitat context), and users should treat them as

approximations of mature phenotypes rather than statistically derived species means.
The same can be true for chromosome count, which can vary intraspecifically.
810
811 CactEcoDB is designed to be a continually evolving resource. As new trait
812 measurements, spatial distributions, molecular sequences, and relevant environmental
813 maps become available, we will incorporate these into future versions. We therefore
814 welcome contributions of additional data and corrections to further improve the
815 accuracy, completeness, and utility of CactEcoDB.
816
In summary, CactEcoDB offers a comprehensive resource for large-scale ecological
and evolutionary analyses, but as with any aggregated dataset, its contents should be
interpreted in light of the inherent variability in data completeness and precision. Where
greater taxonomic or spatial resolution is required, users may find it helpful to consult
original sources or species-level references. Similar to other trait databases [15,42,157,159–
162], we recommend that users monitor ongoing changes in Cactaceae taxonomic
nomenclature.
824

Code availability

825
826 Code used to generate, process and analyse data, and produce figures is available at
827 Figshare (https://doi.org/10.6084/m9.figshare.30940019) [139] and in the GitHub
828 repository (https://github.com/jamie-thompson/CactEcoDB).
829

Funding

830
831 This work was funded by a Roger and Sue Whorrod OBE PhD studentship awarded to
832 J.B.T and N.K.P, and a British Cactus and Succulent Society research grant to J.B.T
833 and C.M.
834

Acknowledgments

835
836 We acknowledge Craig Hilton-Taylor for granting permission to reproduce IUCN spatial
837 coordinates, Eric Hagen for technical advice running MiSSE, Juriaan de Vos for advice
838 on the phylogenomic backbone, and Itay Mayrose for granting permission to reproduce
839 chromosome counts from the Chromosome Count Database.
840

References:

1. Pérez-Escobar, O. A. *et al.* The origin and speciation of orchids. *New Phytol* **242**,

700–716 (2024).

2. Sauquet, H. & Magallón, S. Key questions and challenges in angiosperm

macroevolution. *New Phytol* **219**, 1170–1187 (2018).

3.   Panero, J. L. & Crozier, B. S. Macroevolutionary dynamics in the early

diversification of Asteraceae. *Mol Phylogenet Evol* **99**, 116–132 (2016).

4.   Magallón, S., Sánchez-Reyes, L. L. & Gómez-Acevedo, S. L. Thirty clues to the

exceptional diversification of flowering plants. *Ann Bot* **123**, 491–503 (2019).

5.   Ramírez-Barahona, S., Sauquet, H. & Magallón, S. The delayed and

geographically heterogeneous diversification of flowering plant families. *Nat Ecol*

Evol **4**, 1232–1238 (2020).

6.   Cabral, A. *et al.* Seed-dispersing vertebrates and the abiotic environment shape

functional diversity of the pantropical Annonaceae. *New Phytol* **246**, 2263–2279

(2025).

7.   Onstein, R. E. *et al.* Frugivory-related traits promote speciation of tropical palms.

Nat Ecol Evol **1**, 1903–1911 (2017).

8.   Diversification rate shifts in the Cape Floristic Region: The right traits in the right

place at the right time. *Perspectives in Plant Ecology, Evolution and Systematics*

16, 331–340 (2014).

9.   Givnish, T. J. *et al.* Orchid phylogenomics and multiple drivers of their extraordinary

diversification. *Proc Biol Sci* **282**, (2015).

10.   Thompson, J. B., Hernández-Hernández, T., Keeling, G., Vásquez-Cruz, M. &

Priest, N. K. Identifying the multiple drivers of cactus diversification. *Nat Commun*

15, 7282 (2024).

11.   Thompson, J. B., Davis, K. E., Dodd, H. O., Wills, M. A. & Priest, N. K. Speciation

across the Earth driven by global cooling in terrestrial orchids. *Proc Natl Acad Sci U*

S A **120**, e2102408120 (2023).

12. Thompson, J., Ramírez-Barahona, S., Priest, N. & Hernández-Hernández, T. Did

succulents diversify in response to aridity? Evolutionary analyses of major

succulent lineages around the world. *bioRxiv* 2023.05.23.541957 (2023)

doi:10.1101/2023.05.23.541957.

13. Kattge, J. *et al.* TRY plant trait database - enhanced coverage and open access.

Glob Chang Biol **26**, 119–188 (2020).

14. Tavşanoğlu, Ç. & Pausas, J. G. A functional trait database for Mediterranean Basin

plants. *Sci Data* **5**, 180135 (2018).

15. Falster, D. *et al.* AusTraits, a curated plant trait database for the Australian flora.

Sci Data **8**, 254 (2021).

16. Wang, H. *et al.* The China plant trait database version 2. *Sci Data* **9**, 769 (2022).

17. Weigelt, P., König, C. & Kreft, H. GIFT – A Global Inventory of Floras and Traits for

macroecology and biogeography. *Journal of Biogeography* **47**, 16–43 (2020).

18. Upham, N. S., Esselstyn, J. A. & Jetz, W. Inferring the mammal tree: Species-level

sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS*

Biol **17**, e3000494 (2019).

19. Payne, A. R. D., Mannion, P. D., Lloyd, G. T. & Davis, K. E. Decoupling speciation

and extinction reveals both abiotic and biotic drivers shaped 250 million years of

diversity in crocodile-line archosaurs. *Nat Ecol Evol* **8**, 121–132 (2024).

20. Smith, S. A. & Brown, J. W. Constructing a broadly inclusive seed plant phylogeny.

Am J Bot **105**, 302–314 (2018).

21. Portik, D. M., Streicher, J. W. & Wiens, J. J. Frog phylogeny: A time-calibrated,

891  species-level tree based on hundreds of loci and 5,242 species. *Mol Phylogenet*

892  *Evol* **188**, 107907 (2023).

893 22. Waller, J. T., Willink, B., Tschol, M. & Svensson, E. I. The odonate phenotypic

894  database, a new open data resource for comparative studies of an old insect order.

895  *Sci Data* **6**, 316 (2019).

896 23. Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C. & Costa, G. C.

897  AmphiBIO, a global database for amphibian ecological traits. *Scientific Data* **4**, 1–7

898  (2017).

899 24. Pottier, P. *et al.* A comprehensive database of amphibian heat tolerance. *Scientific*

900  *Data* **9**, 600 (2022).

901 25. Wang, J. *et al.* CheloniansTraits: a comprehensive trait database of global turtles

902  and tortoises. *Scientific Data* **12**, 1–7 (2025).

903 26. Wei, P. *et al.* CaecilianTraits, an individual level trait database of Caecilians

904  worldwide. *Scientific Data* **12**, 1–7 (2025).

905 27. Chaudhary, V. B. *et al.* TraitAM, a global spore trait database for arbuscular

906  mycorrhizal fungi. *Scientific Data* **12**, 1–10 (2025).

907 28. Gómez-Gras, D. *et al.* The Octocoral Trait Database: a global database of trait

908  information for octocoral species. *Scientific Data* **12**, 1–32 (2025).

909 29. Báez, S. *et al.* FunAndes – A functional trait database of Andean plants. *Scientific*

910  *Data* **9**, 1–9 (2022).

911 30. Herberstein, M. E. *et al.* AnimalTraits - a curated animal trait database for body

912  mass, metabolic rate and brain size. *Scientific Data* **9**, 1–11 (2022).

913 31. Kissling, W. D. *et al.* PalmTraits 1.0, a species-level functional trait database of

palms worldwide. *Scientific Data* **6**, 1–13 (2019).

32.   Jetz, W. & Pyron, R. A. The interplay of past diversification and evolutionary

isolation with present imperilment across the amphibian tree of life. *Nature Ecology*

& Evolution **2**, 850–858 (2018).

33.   Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W. & Pyron, R. A. Fully-sampled

phylogenies of squamates reveal evolutionary patterns in threat status. *Biol.*

Conserv. **204**, 23–31 (2016).

34.   Stein, R. W. *et al.* Global priorities for conserving the evolutionary history of sharks,

rays and chimaeras. *Nature Ecology & Evolution* **2**, 288–298 (2018).

35.   Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. & Mooers, A. O. The global

diversity of birds in space and time. *Nature* **491**, 444–448 (2012).

36.   Givnish, T. J. *et al.* Adaptive radiation, correlated and contingent evolution, and net

species diversification in Bromeliaceae. *Mol Phylogenet Evol* **71**, 55–78 (2014).

37.   Siqueira, A. C., Morais, R. A., Bellwood, D. R. & Cowman, P. F. Trophic

innovations fuel reef fish diversification. *Nat Commun* **11**, 2669 (2020).

38.   Bouchenak-Khelladi, Y., Onstein, R. E., Xing, Y., Schwery, O. & Peter Linder, H.

On the complexity of triggering evolutionary radiations. *New Phytologist* **207**, 313–

326 (2015).

39.   Padilla-Morales, B. *et al.* Fast species diversification among dragonflies

(Anisoptera: Odonata: Insecta) inhabiting lentic environments regardless of wing

pigmentation. *Ecological Entomology* **47**, 314–322 (2022).

40.   Hembry, D. H. & Weber, M. G. Ecological interactions and macroevolution: a new

field with old roots. *Annu. Rev. Ecol. Evol. Syst.* **51**, 215–243 (2020).

41. Swiston, S. K. & Landis, M. J. Testing Relationships between Multiple Regional

Features and Biogeographic Processes of Speciation, Extinction, and Dispersal.

Syst Biol **74**, 282–300 (2025).

42. Reginato, M. *et al.* MelastomaTRAITs 1.0: A database of functional traits in

Melastomataceae, a large pantropical angiosperm family. *Ecology* **105**, e4308

(2024).

43. Etard, A., Morrill, S. & Newbold, T. Global gaps in trait data for terrestrial

vertebrates. *Global Ecology and Biogeography* **29**, 2143–2158 (2020).

44. Grenié, M., Bruelheide, H., Dawson, W., Essl, F., van Kleunen, M., Kühn, I., Kreft,

H., Pyšek, P., Weigelt, P. & Winter, M. Gaps in global alien plant trait data and how

to fill them. *Glob. Ecol. Biogeogr.* **34**, e70131 (2025).

45. Iversen, C. M. & McCormack, M. L. Filling gaps in our understanding of

belowground plant traits across the world: an introduction to a Virtual Issue. *New*

Phytol. **231**, 2097–2103 (2021).

46. Cutts, V. *et al.* Scientific floras can be reliable sources for some trait data in a

system with poor coverage in global trait databases. *Journal of Vegetation Science*

32, e12996 (2021).

47. Díaz, S. *et al.* The global spectrum of plant form and function: enhanced species-

level trait dataset. *Scientific Data* **9**, 1–18 (2022).

48. Cordeiro, A. L. *et al.* TropiRoot 1.0: Database of tropical root characteristics across

environments. *Ecology* **106**, e70074 (2025).

49. Guerrero-Ramírez, N. R. *et al.* Global root traits (GRooT) database. *Global Ecology*

and Biogeography **30**, 25–37 (2021).

50. Cornwell, W. K., Pearse, W. D., Dalrymple, R. L. & Zanne, A. E. What we (don't)

know about global plant diversity. *Ecography* **42**, 1819–1831 (2019).

51. Hernández-Hernández, T., Brown, J. W., Schlumpberger, B. O., Eguiarte, L. E. &

Magallón, S. Beyond aridification: multiple explanations for the elevated

diversification of cacti in the New World Succulent Biome. *New Phytol* **202**, 1382–

1397 (2014).

52. Guerrero, P. C., Majure, L. C., Cornejo-Romero, A. & Hernández-Hernández, T.

Phylogenetic Relationships and Evolutionary Trends in the Cactus Family. *J Hered*

110, 4–21 (2019).

53. Hultine, K. R. *et al.* Global change impacts on cacti (Cactaceae): current threats,

challenges and conservation solutions. *Ann Bot* **132**, 671–683 (2023).

54. Gibson, A. C. & Nobel, P. S. *The Cactus Primer*. (Harvard University Press, 1986).

55. Arakaki, M. *et al.* Contemporaneous and recent radiations of the world's major

succulent plant lineages. *Proc Natl Acad Sci U S A* **108**, 8379–8384 (2011).

56. del Carmen Mandujano, M., Carrillo-Angeles, I., Martínez-Peralta, C. & Golubov, J.

Reproductive Biology of Cactaceae. *Desert Plants* 197–230 (2010).

57. Godínez-Alvarez, H., Valverde, T. & Ortega-Baes, P. Demographic trends in the

Cactaceae. *Bot. Rev.* **69**, 173–201 (2003).

58. Barthlott, W. & Hunt, D. R. Cactaceae. in *Flowering Plants · Dicotyledons* 161–197

(Springer Berlin Heidelberg, Berlin, Heidelberg, 1993).

59. Calvente, A., Zappi, D. C., Forest, F. & Lohmann, L. G. Molecular Phylogeny,

Evolution, and Biogeography of South American Epiphytic Cacti. *International*

Journal of Plant Sciences (2011) doi:10.1086/660881.

60. Hernández-Hernández, T. *et al.* Phylogenetic relationships and evolution of growth

form in Cactaceae (Caryophyllales, Eudicotyledoneae). *Am J Bot* **98**, 44–61 (2011).

61. Ventura-Aguilar, R. I., Bosquez-Molina, E., Bautista-Baños, S. & Rivera-Cabrera, F.

Cactus stem (Opuntia ficus-indica Mill): anatomy, physiology and chemical

composition with emphasis on its biofunctional properties. *J Sci Food Agric* **97**,

5065–5073 (2017).

62. Edwards, E. J. & Donoghue, M. J. Pereskia and the origin of the cactus life-form.

Am Nat **167**, 777–793 (2006).

63. Amaral, D. T. *et al.* Tracking the xeric biomes of South America: The

spatiotemporal diversification of Mandacaru cactus. *Journal of Biogeography* **48**,

3085–3103 (2021).

64. Majure, L. C. *et al.* Pleistocene aridification underlies the evolutionary history of the

Caribbean endemic, insular, giant Consolea (Opuntioideae). *Am J Bot* **108**, 200–

215 (2021).

65. Majure, L. C., Barrios, D., Díaz, E., Bacci, L. F. & Piñeyro, Y. E. Phylogenomics of

the Caribbean melocacti: Cryptic species and multiple invasions. *TAXON* **71**, 993–

1012 (2022).

66. Amaral, D. T., Bonatelli, I. A., Romeiro-Brito, M., Moraes, E. M. & Franco, F. F.

Spatial patterns of evolutionary diversity in Cactaceae show low ecological

representation within protected areas. *Biol. Conserv.* **273**, 109677 (2022).

67. Fleming, T. H. & Valiente-Banuet, A. *Columnar Cacti and Their Mutualists:*

Evolution, Ecology, and Conservation. (University of Arizona Press, 2002).

68. Novoa, A., Le Roux, J. J., Robertson, M. P., Wilson, J. R. U. & Richardson, D. M.

Introduced and invasive cactus species: a global review. *AoB Plants* **7**, (2014).

69. Aranda-Pineda, J. A., Valverde, P. L., Búrquez, A., Golubov, J. & Mandujano, M. C. Effect of brousse tigrée on the dynamics of nurse–protégé interactions of a cactus in the Chihuahuan Desert. *Plant Species Biology* **36**, 450–462 (2021).

70. Goettsch, B. *et al.* High proportion of cactus species threatened with extinction. *Nat Plants* **1**, 15142 (2015).

71. Pillet, M. *et al.* Elevated extinction risk of cacti under climate change. *Nat Plants* **8**, 366–372 (2022).

72. Goettsch, B., Durán, A. P. & Gaston, K. J. Global gap analysis of cactus species and priority sites for their conservation. *Conserv Biol* **33**, 369–376 (2019).

73. Hultine, K. R. *et al.* The Role of Botanical Gardens in the Conservation of Cactaceae. *BioScience* **66**, 1057–1065 (2016).

74. Castro, J. P. *et al.* Karyotype characterization and evolution of chromosome number in Cactaceae with special emphasis on subfamily Cactoideae. *Acta Bot. Bras.* **34**, 135–148 (2019).

75. Yetman, D. *The Great Cacti: Ethnobotany & Biogeography*. (University of Arizona Press, 2007).

76. Pérez-Negrón, E., Dávila, P. & Casas, A. Use of columnar cacti in the Tehuacán Valley, Mexico: perspectives for sustainable management of non-timber forest products. *J Ethnobiol Ethnomed* **10**, 79 (2014).

77. Ortiz-Martínez, L. E., Manzanero-Medina, G. I., Golubov, J., Vásquez-Dávila, M. A. & Mandujano, M. C. Nopal de Monte: Cacti Named and Used by a Mixtec Community in Mountainous Oaxaca. *Ethnobotany of the Mountain Regions of*

Mexico 899–908 (2023).

78. Patrick Griffith, M. The origins of an important cactus crop, *Opuntia ficus-indica*

(Cactaceae): new molecular evidence. *American Journal of Botany* **91**, 1915–1921

(2004).

79. Kiesling, R. Origen, domesticación y distribución de *Opuntia ficus-indica. J. Prof.*

Assoc. Cactus Dev. **3**, 50–59 (1998)..

80. Food and Agriculture Organization of the United Nations. *Cactus (Opuntia Spp.) as*

Forage. (Food & Agriculture Org., 2001).

81. Food and Agriculture Organization of the United Nations. *Crop Ecology, Cultivation*

and Uses of Cactus Pear. (Food & Agriculture Org., 2018).

82. Donkin, R. A. *Spanish Red: An Ethnogeographical Study of Cochineal and the*

Opuntia Cactus. (1977).

83. Margulies, J. D., Moorman, F. R., Goettsch, B., Axmacher, J. C. & Hinsley, A.

Prevalence and perspectives of illegal trade in cacti and succulent plants in the

collector community. *Conserv Biol* **37**, e14030 (2023).

84. Novoa, A., Le Roux, J. J., Richardson, D. M. & Wilson, J. R. U. Level of

environmental threat posed by horticultural trade in Cactaceae. *Conserv Biol* **31**,

1066–1075 (2017).

85. Sajeva, M. *et al. CITES and Cacti: A User's Guide.* (Royal Botanic Gardens Kew,

2012).

86. Franco, F. F. *et al.* Evolutionary Genetics of Cacti: Research Biases, Advances and

Prospects. *Genes (Basel)* **13**, (2022).

87. Yesson, C. *et al.* How Global Is the Global Biodiversity Information Facility? *PLOS*

ONE **2**, e1124 (2007).

88. Brown, M. J. M. *et al.* rWCVP: a companion R package for the World Checklist of

Vascular Plants. *New Phytol* **240**, 1355–1365 (2023).

89. de Vos, J. M. *et al.* Phylogenomics and classification of Cactaceae based on

hundreds of nuclear genes. *Plant Systematics and Evolution* **311**, 28 (2025).

90. Drori, M. *et al.* OneTwoTree: An online tool for phylogeny reconstruction. *Mol Ecol*

Resour **18**, 1492–1499 (2018).

91. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version

7: improvements in performance and usability. *Mol Biol Evol* **30**, 772–780 (2013).

92. Gouy, M., Guindon, S. & Gascuel, O. SeaView version 4: A multiplatform graphical

user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol*

27, 221–224 (2010).

93. Katoh, K. & Frith, M. C. Adding unaligned sequences into an existing alignment

using MAFFT and LAST. *Bioinformatics* **28**, 3144–3146 (2012).

94. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*

25, 1972–1973 (2009).

95. Borowiec, M. L. AMAS: a fast tool for alignment manipulation and computing of

summary statistics. *PeerJ* **4**, e1660 (2016).

96. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis

of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

97. Rabosky, D. L. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. *PLoS One* **9**, e89543 (2014).

98. Vasconcelos, T., O'Meara, B. C. & Beaulieu, J. M. A flexible method for estimating

tip diversification rates across a range of speciation and extinction scenarios.

Evolution **76**, 1420–1433 (2022).

99. Moore, B. R., Höhna, S., May, M. R., Rannala, B. & Huelsenbeck, J. P. Critically

evaluating the theory and performance of Bayesian analysis of macroevolutionary

mixtures. *Proc Natl Acad Sci U S A* **113**, 9569–9574 (2016).

100.Meyer, A. L. S. & Wiens, J. J. Estimating diversification rates for higher taxa:

BAMM can give problematic estimates of rates and rate shifts. *Evolution* **72**, 39–53

(2018).

101.Rabosky, D. L., Mitchell, J. S. & Chang, J. Is BAMM Flawed? Theoretical and

Practical Concerns in the Analysis of Multi-Rate Diversification Models. *Syst Biol*

66, 477–498 (2017).

102.Rabosky, D. L. BAMM at the court of false equivalency: A response to Meyer and

Wiens. *Evol* **72**, 2246–2256 (2018).

103.Cooney, C. R. & Thomas, G. H. Heterogeneous relationships between rates of

speciation and body size evolution across vertebrate clades. *Nat Ecol Evol* **5**, 101–

110 (2021).

104.Beaulieu, J. M. The problem with clade-specific sampling fractions. Available at

https://rdrr.io/cran/hisse/f/inst/doc/Clade-specificsampling.pdf

105.Title, P. O. & Rabosky, D. L. Tip rates, phylogenies and diversification: What are

we estimating, and how good are the estimates? *Methods in Ecology and Evolution*

10, 821–834 (2019).

106.Rabosky, D. L. *et al.* BAMMtools: an R package for the analysis of evolutionary

dynamics on phylogenetic trees. *Methods in Ecology and Evolution* **5**, 701–707

(2014).

107. Beaulieu, J. M. & O'Meara, B. C. Detecting Hidden Diversification Shifts in Models

of Trait-Dependent Speciation and Extinction. *Syst Biol* **65**, 583–601 (2016).

108. Thompson, J. B. et al. Frequent shifts in pollination strategy are decoupled from

diversification in the terrestrial orchids. *EcoEvoRxiv*

https://doi.org/10.32942/X21P97 (2025).

109. Anderson, E. F. *The Cactus Family*. (Timber Press (OR), 2001).

110. Moles, A. T. *et al.* Global patterns in plant height. *Journal of Ecology* **97**, 923–932

(2009).

111. Stephens, R. E., Gallagher, R. V., Dun, L., Cornwell, W. & Sauquet, H. Insect

pollination for most of angiosperm evolutionary history. *New Phytol* **240**, 880–891

(2023).

112. Specht, C. D., Yockteng, R., Almeida, A. M., Kirchoff, B. K. & Kress, W. J.

Homoplasy, Pollination, and Emerging Complexity During the Evolution of Floral

Development in the Tropical Gingers (Zingiberales). *The Botanical Review* **78**,

440–462 (2012).

113. Barreto, E. *et al.* Macroevolution of the plant–hummingbird pollination system.

Biological Reviews **99**, 1831–1847 (2024).

114. Ackerman, J. D., Phillips, R. D., Tremblay, R. L., Karremans, A., Reiter, N., Peter,

C. I., Bogarín, D., Pérez-Escobar, O. A. & Liu, H. Beyond the various contrivances

by which orchids are pollinated: global patterns in orchid pollination biology. *Bot. J.*

Linn. Soc. **202**, 295–324 (2023).

115. Fenster, C. B., Armbruster, W. S., Wilson, P., Dudash, M. R. & Thomson, J. D.

Pollination syndromes and floral specialization. *Annu. Rev. Ecol. Evol. Syst.* **35**,

375–403 (2004).

116. Kessler, M., Abrahamczyk, S. & Krömer, T. The role of hummingbirds in the

evolution and diversification of Bromeliaceae: unsupported claims and untested

hypotheses. *Bot. J. Linn. Soc.* **192**, 592–608 (2020).

117. Rosas-Guerrero, V., Aguilar, R., Martén-Rodríguez, S., Ashworth, L.,

Lopezaraiza-Mikel, M., Bastida, J. M. & Quesada, M. A quantitative review of

pollination syndromes: do floral traits predict effective pollinators? *Ecol. Lett.* **17**,

388–400 (2014).

118. Waser, N. M., Chittka, L., Price, M. V., Williams, N. M. & Ollerton, J. Generalization

in Pollination Systems, and Why it Matters. *Ecology* **77**, 1043–1060 (1996).

119. Ollerton, J. Pollinator diversity: distribution, ecological function, and conservation.

Annu. Rev. Ecol. Evol. Syst. **48**, 353–376 (2017).

120. Gorostiague, P., Ollerton, J. & Ortega-Baes, P. Latitudinal gradients in biotic

interactions: Are cacti pollination systems more specialized in the tropics? *Plant*

Biol (Stuttg) **25**, 187–197 (2023).

121. Albuquerque-Lima, S., Taylor, N. P., Zappi, D. C. & Machado, I. C. Floral

specialization and bat pollination in subtribe Cereinae (Cactaceae): a morphological

approach. *Diversity* **15**, 207 (2023).

122. Schlumpberger, B. O. & Renner, S. S. Molecular phylogenetics of Echinopsis

(Cactaceae): Polyphyly at all levels and convergent evolution of pollination modes

and growth forms. *American Journal of Botany* **99**, 1335–1349 (2012).

123. Rice, A. *et al.* The Chromosome Counts Database (CCDB) - a community resource

of plant chromosome numbers. *New Phytol* **206**, 19–26 (2015).

124. Xavier, A., Yadav, R. & Gowda, V. Evolutionary patterns of variations in

chromosome counts and genome sizes show positive correlations with taxonomic

diversity in tropical gingers. *American Journal of Botany* **111**, e16334 (2024).

125. Osterman, W. H. A. *et al.* Rethinking pathways to the dioecy–polyploidy

association: Genera with many dioecious species have fewer polyploids. *American*

Journal of Botany **111**, e16318 (2024).

126. Román-Palacios, C., Medina, C. A., Zhan, S. H. & Barker, M. S. Animal

chromosome counts reveal a similar range of chromosome numbers but with less

polyploidy in animals compared to flowering plants. *J. Evol. Biol.* **34**, 1333–1339

(2021).

127. The. *IUCN Red List of Threatened Species* https://www.iucnredlist.org/en.

128. Avaria-Llautureo, J., Püschel, T. A., Meade, A., Baker, J., Nicholson, S. L. &

Venditti, C. The radiation and geographic expansion of primates through diverse

climates. *Proc. Natl Acad. Sci. USA* **122**, e2423833122 (2025).

129. Carta, A., Peruzzi, L. & Ramírez-Barahona, S. A global phylogenetic regionalization

of vascular plants reveals a deep split between Gondwanan and Laurasian biotas.

New Phytol **233**, 1494–1504 (2022).

130. Ramírez-Barahona, S. *et al.* Upslope plant species shifts in Mesoamerican cloud

forests driven by climate and land use change. *Science* **387**, 1058–1063 (2025).

131. Zizka, A. *et al.* CoordinateCleaner: Standardized cleaning of occurrence records

from biological collection databases. *Methods in Ecology and Evolution* **10**, 744–

751 (2019).

132.Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface

areas. *Sci Data* **4**, 170122 (2017).

133.Amatulli, G. *et al.* A suite of global, cross-scale topographic variables for

environmental and biodiversity modeling. *Sci Data* **5**, 180040 (2018).

134.Poggio, L. *et al.* SoilGrids 2.0: producing soil information for the globe with

quantified spatial uncertainty. *SOIL* **7**, 217–240 (2021).

135.Fisher-Reid, M. C. & Wiens, J. J. Is geographic variation within species related to

macroevolutionary patterns between species? *J Evol Biol* **28**, 1502–1515 (2015).

136.Avaria-Llautureo, J., Rivadeneira, M. M., Venditti, C. & Luna-Jorquera, G. Seabird

range contraction and dispersal under climate change. *EcoEvoRxiv*

https://doi.org/10.32942/X2B33W (2025).

137.Mitchell, N., Carlson, J. E. & Holsinger, K. E. Correlated evolution between climate

and suites of traits along a fast-slow continuum in the radiation of. *Ecol Evol* **8**,

1853–1866 (2018).

138.Nakagawa, S. & De Villemereuil, P. A General Method for Simultaneously

Accounting for Phylogenetic and Species Sampling Uncertainty via Rubin's Rules

in Comparative Analysis. *Syst Biol* **68**, 632–641 (2019).

139.Thompson, J., Martinez, C., Avaria-Llautureo, J., Ramirez-Barahona, S.,

Manzanarez-Villasana, G., Culham, A. et al. CactEcoDB: Trait, spatial,

environmental, phylogenetic and diversification data for the cactus family. *Figshare*

https://doi.org/10.6084/m9.figshare.30940019.v2 (2025).

140.Danielson, J. J. & Gesch, D. B. *Global Multi-Resolution Terrain Elevation Data*

1190 *2010 (GMTED2010).* (2011).

141. Boucher, F. C., Quatela, A.-S., Ellis, A. G. & Anthony Verboom, G. Diversification rate vs. diversification density: Decoupled consequences of plant height for diversification of Alooideae in time and space. *PLOS ONE* **15**, e0233597 (2020).

142. Mao, L. *et al.* The geographic and climatic distribution of plant height diversity for 19,000 angiosperms in China. *Biodiversity and Conservation* **29**, 487–502 (2019).

143. Liu, H. *et al.* Hydraulic traits are coordinated with maximum plant height at the global scale. *Sci Adv* **5**, eaav1332 (2019).

144. Mao, L., Chen, S., Zhang, J. & Zhou, G. Altitudinal patterns of maximum plant height on the Tibetan Plateau. *J Plant Ecol* **11**, 85–91 (2016).

145. Breslin, P. B., Wojciechowski, M. F. & Majure, L. C. Remarkably rapid, recent diversification of Cochemiea and Mammillaria in the Baja California, Mexico region. *Am J Bot* **109**, 1472–1487 (2022).

146. Zuntini, A. R. *et al.* Phylogenomics and the rise of the angiosperms. *Nature* **629**, 843–850 (2024).

147. Henríquez-Piskulich, P., Hugall, A. F. & Stuart-Fox, D. A supermatrix phylogeny of the world's bees (Hymenoptera: Anthophila). *Mol Phylogenet Evol* **190**, 107963 (2024).

148. Hinchliff, C. E. & Roalson, E. H. Using supermatrices for phylogenetic inquiry: an example using the sedges. *Syst Biol* **62**, 205–219 (2013).

149. McMahon, M. M. & Sanderson, M. J. Phylogenetic supermatrix analysis of GenBank sequences from 2228 papilionoid legumes. *Syst Biol* **55**, 818–836 (2006).

150. de Queiroz, A. & Gatesy, J. The supermatrix approach to systematics. *Trends Ecol.*

1213   *Evol.* **22**, 34–41 (2007).

151. Philippe, H., de Vienne, D. & Ranwez, V. *Pitfall in Supermatrix Phylogenomics.*
     (2017).

152. Smith, S. A., Beaulieu, J. M. & Donoghue, M. J. Mega-phylogeny approach for
     comparative biology: an alternative to supertree and supermatrix approaches. *BMC
     Evol Biol* **9**, 37 (2009).

153. Simmons, M. P. & Goloboff, P. A. Dubious resolution and support from published
     sparse supermatrices: the importance of thorough tree searches. *Mol Phylogenet
     Evol* **78**, 334–348 (2014).

154. Sanderson, M. J., McMahon, M. M., Stamatakis, A., Zwickl, D. J. & Steel, M.
     Impacts of Terraces on Phylogenetic Inference. *Syst Biol* **64**, 709–726 (2015).

155. Simmons, M. P. A confounding effect of missing data on character conflict in
     maximum likelihood and Bayesian MCMC phylogenetic analyses. *Mol Phylogenet
     Evol* **80**, 267–280 (2014).

156. Weng, J. *et al.* Comprehensive Dataset for Polychaetes in the IPC: Species
     Distribution, DNA Barcodes, and Functional Traits. *Scientific Data* **12**, 956 (2025).

157. Denelle, P., Weigelt, P. & Kreft, H. GIFT—An R package to access the Global
     Inventory of Floras and Traits. *Methods Ecol. Evol.* **14**, 2738–2748 (2023).

158. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171
     (2015).

159. Galán-Acedo, C., Arroyo-Rodríguez, V., Andresen, E. & Arasa-Gisbert, R.
     Ecological traits of the world's primates. *Scientific Data* **6**, 55 (2019).

160. Hurlbert, A. H., Olsen, A. M., Sawyer, M. M. & Winner, P. M. The Avian Diet

Database as a source of quantitative information on bird diets. *Scientific Data* **8**,

260 (2021).

161. Demetriou, J. *et al.* The Cyprus Database of Alien Species (CyDAS). *Scientific*

Data **12**, 1881 (2025).

162. De Wint, F. C. *et al.* Introducing a global database of entomopathogenic fungi and

their host associations. *Scientific Data* **11**, 1418 (2024).

1242