Model substrate particles uncover dynamics of microbial communities associated with

particulate organic matter decomposition in soil

Eva Simon[1,2], Lauren V. Alteio[1#*], Alexander König[1**], Bruna Imai[1***], Julia Horak[1****], Julia

Wiesenbauer[1,2], Joana Séneca[1,3,5], Bela Hausmann[3,5], Marc Mußmann[1], Barbara Kitzler[4],

and Christina Kaiser[1]

[1]Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna,

Austria

[2]Doctoral School in Microbiology and Environmental Science, University of Vienna, Vienna,

Austria

[3]Joint Microbiome Facility of the Medical University of Vienna and the University of Vienna,

Vienna, Austria

[4]Austrian Research Centre for Forests, Vienna, Austria

[5]Division of Clinical Microbiology, Department of Laboratory Medicine, Medical University of

Vienna, Vienna, Austria

Running Title: Model particle associated microbial communities

#Address correspondence to Lauren V. Alteio, lauren.alteio@ffoqsi.at

*Present address: Lauren V Alteio, Austrian Competence Centre for Feed and Food Quality,

Safety and Innovation, FFoQSI GmbH, Tulln, Austria

**Present address: Alexander König, University of Natural Resources and Life Sciences

(BOKU), Vienna, Austria

***Present address: Bruna Imai, MPI for Marine Microbiology, Bremen, Germany

***Present address: Julia Horak, Austrian Research Centre for Forests

Eva Simon and Lauren V Alteio contributed equally to this work. The author order was

determined   on   the   basis   of   seniority.

Two-part abstract

Abstract

Soil organic matter is the largest terrestrial reservoir of organic carbon. Its particulate fraction, particulate organic matter (POM), serves as a resource and surface for microbial colonization. Degradation of complex biopolymers like cellulose and chitin requires extracellular enzymes produced by phylogenetically diverse microbes. Despite their importance for carbon cycling, the structure and spatio-temporal dynamics of particle-associated microbial communities in soil and how specific substrates influence them remain poorly understood. This study investigated whether microbial communities associated with POM change in composition and richness over time and whether chitin and cellulose select for distinct fungal and bacterial taxa. We incubated self-manufactured millimeter-sized model POM substrate particles containing chitin or cellulose in soil under laboratory and field conditions. Over 50 days, we assessed particle-associated bacterial and fungal communities at multiple time points in the lab and after 47 days in the field. Our results show that community structure and temporal dynamics of particle-associated microbial communities were substrate-specific. While microbial biomass increased on both particle types, chitin-associated microbial exhibited stronger temporal changes. Communities on chitin and cellulose particles were enriched in specific bacterial and fungal genera compared to communities in the surrounding soil. In conclusion, our study demonstrates that microbial communities associated with model chitin particles underwent notable temporal changes. More specifically, microbial richness decreased, and community composition shifted over the incubation period. Both chitin and cellulose particles favored certain fungal, and especially bacterial genera compared to surrounding soil.

Importance

Particulate organic matter constitutes the largest terrestrial organic carbon storage. Microorganisms colonize, degrade, transform, and become part of organic matter. We use self-manufactured model chitin or cellulose particles to examine the temporal dynamics of bacterial and fungal communities associated with particulate organic matter in soil. Our findings suggest model particles can be used to investigate particle-associated microbial communities in soil. Furthermore, we show that temporal dynamics are substrate-specific. Our findings advance our understanding of particle-associated microbial communities and the impact of specific substrates on microorganisms in soil.

Keywords

Soil microbial communities, particulate organic matter, model particles, chitin, cellulose, temporal dynamics

Introduction

Soil organic matter is the largest terrestrial carbon reservoir (Lehmann and Kleber, 2015; Angst et al., 2021). Conceptually, soil organic matter can be divided into mineral-associated organic matter and particulate organic matter (Lavallee et al., 2020). Particulate organic matter serves as a nutrient source to microorganisms and a surface for colonization (Young et al., 2008). Particles are considered hotspots for microbial activity in soil (Witzgall et al., 2021) as they represent spatially isolated, nutrient-rich microhabitats in the otherwise nutrient-poor soil environment.

Microbes decompose particulate organic matter, as they can only take up small molecules (typically less than 600 Da) across their cell walls (Arnosti et al., 2014; Lehmann and Kleber, 2015). To acquire resources, they produce and excrete extracellular enzymes into their surroundings (Allison et al., 2010; Nunan et al., 2020). Extracellular enzymes are the agents of organic matter decomposition, as they break down large molecules into subunits and ultimately monomers, which can be taken up by microbes (Sinsabaugh et al., 2008; Gómez et al., 2020). By decomposing soil organic matter via extracellular enzymes, heterotrophic microorganisms mediate the recycling of carbon and nutrients (Sinsabaugh et al., 2008; Bardgett and Van Der Putten, 2014; Berlemont, 2017; Wu et al., 2024). Extracellular enzymes are substrate specific (Allison et al., 2010) and metabolically costly to produce (Nunan et al., 2020, Guseva et al., 2024). The complete breakdown of biopolymers requires multiple enzymes that act in synergy (Sinsabaugh et al., 2009; Talamantes et al., 2016, Guseva et al., 2024). Genetic potential to produce particular extracellular enzymes differs between microbial taxa (Berlemont and Martiny, 2013, 2015; Zimmerman et al., 2013; Arnosti et al., 2014). Consequently, phylogenetically diverse microbial communities are necessary to carry out all steps of biopolymer degradation.

Cellulose and chitin are the two most abundant natural polymers in nature (Das et al., 2010; Berlemont and Martiny, 2015). Cellulose is the major component of plant material and the most abundant biopolymer in nature, and chitin is the second most abundant polysaccharide (Das et al., 2010; Wieczorek et al., 2014). Both need to be enzymatically degraded prior to microbial uptake. While the potential to degrade these polysaccharides is widespread across bacterial taxa, the structure of degrader communities has been found to be driven by environmental conditions and substrate availability in situ (Berlemont and Martiny, 2015; D. Li et al., 2023). For instance, the ability to degrade cellulose has been shown to be more widespread amongst fungi than their ability to degrade chitin (Berlemont, 2017).

Spatial proximity of microbial cells to enzymatically degraded substrates is advantageous to enhance the return of enzyme investment. Microbial attachment to particulate organic matter and formation of spatially structured communities should theoretically minimize losses of

enzymatic breakdown products to the surrounding environment through diffusion. The presence of both, bacteria and fungi, has been detected on particulate organic matter in soil. More specifically, it has been shown that bacteria occur in biofilms on the surface of particulate organic matter and that fungal hyphae pervade bacterial biofilms (Witzgall et al., 2021). However, overall, little is known about microbial communities associated with particulate organic matter in soil, despite the relevance of particulate organic matter-degrading communities for carbon and nutrient cycling. We lack knowledge about the assembly and temporal dynamics of microbial communities on particulate organic matter, and whether specific substrates select distinct microbial communities and particular microbial taxa in soil.

This lack of knowledge is at least partly due to the complex nature of soil. It is challenging to study microbial communities associated with particulate organic matter in soil. Soils are opaque and characterized by high substrate heterogeneity (Young et al., 2008). Organic matter exhibits diverse chemical composition and is spatially heterogeneously distributed throughout the soil matrix (Lehmann et al., 2008, 2020; Cotrufo et al., 2013; Lavallee et al., 2020; Schlüter et al., 2022). Moreover, the extraction of particulate organic matter from soils requires chemical agents that potentially alter associated microbial communities and distort research findings. Investigating microbial communities of native particulate organic matter that have been chemically extracted from soils might lead to erroneous results.

In this study, we developed and carried out an experimental approach for studying microbial communities associated with particulate organic matter in soil. More specifically, we employed self-manufactured model substrate particles which either contained chitin or cellulose. Model particles mimic natural particulate organic matter, with the advantage that they contain one specific, known polymer. Model substrate particles have been used to study microbial community assembly and dynamics on particles in aquatic systems (Cordero and Datta, 2016; Datta et al., 2016; Ebrahimi et al., 2019), but, to our knowledge, not in soils. We incubated model substrate particles in sieved forest soil in the lab and the field and examined microbial communities of model particles and the soil in which the particles had been incubated, at various time points over a 50-day-long incubation in the lab and after 47 days of incubation in the field. Using this approach, we examined (i) whether model substrate particles select for specific subsets of the overall soil community, (ii) whether particle-associated microbial communities undergo temporal changes, and (iii) whether chitin and cellulose select for specific bacterial, archaeal, and fungal communities and particular microbial genera. We amplicon sequenced bacterial, archaeal, and fungal communities of model substrate particles and the soil in which the particles had been incubated and assessed their taxa richness and community composition. In addition, we determined microbial biomass on particles and in soil via digital droplet PCR.

Material and Methods

Model particle production

We manufactured three types of model particles in the lab: (i) model particles containing chitin ("model chitin particles"), (ii) model particles containing cellulose ("model cellulose particles"), and (iii) model particles containing neither chitin nor cellulose (""no substrate" model particles"). Model particles were designed to mimic particulate organic matter. Model particles to which neither chitin nor cellulose was added served as controls for the chosen scaffold material of model particles, gellan gum. We used Gelrite® gellan gum as a gelling agent for the model particles because of its physical properties (e.g., rapid and stable gelling in the presence of ionic cross-linkers, high transparency for visualization) and its resistance to enzymatic degradation, which makes it a stable and relatively biologically inert scaffold for embedding the selected substrates (chitin and cellulose) (Das et al., 2015). Gellan gum is biologically rather inert compared to other frequently used gelling agents, alginate and agarose (Wang et al., 2023). For model particle production, we followed a protocol by Mußmann (Mußmann et al., in prep). In detail, we dissolved 1.5 g of Gelrite gellan gum (Carl Roth) in 100 mL of pure MilliQ water in a beaker that was placed on a hotplate magnetic stirrer to ensure dissolution of gellan gum. We added a magnetic stirring stick to the beaker to ensure constant mixing of the solution. To facilitate polymer cross-linking and stability, 0.117 g 8 µM $CaCl_2$ was added. Additionally, we added 5 mL of 10 % w/v magnetite solution to the mixture to give particles a magnetic property. The magnetic feature of model particles facilitated the separation of particles from liquids in washing steps further downstream. We used approximately 1.4 g of chitin from shrimp shells (Sigma-Aldrich) or cellulose powder (Sigma) per plate. Produced model particles were of cylindric shape and were approximately 3 mm heigh and 2 mm wide. Model particles were UV-sterilized and stored in 20% ethanol at 4°C until use. For details of particle production see Mußmann et al. (submitted).

Soil collection and processing

Soil was collected in a beech-dominated temperate forest near Klausen - Leopoldsdorf, Austria (48.11925 N, 16.04343 E, 580 m altitude) in spring (April 26, 2021). The litter layer was removed prior to soil sampling. Soil cores were taken from the upper five centimeters of mineral soil at five previously established 5 x 5 m field plots (Simon et al., 2024) using a 5 cm x 8 cm metal corer. For each plot, two soil cores were pooled and sieved using a 2-millimetre-mesh sieve and homogenized into a single representative sample for each plot, producing five composite soil samples.

Set-up of laboratory and field incubation

We incubated model particles in sieved forest soil in the laboratory and in the field. For the lab incubation, we handcrafted mesocosms (3 cm in diameter and 5 cm in height) using 50 mL plastic Greiner vials. Each vial was cut at 5 cm height (30 mL mark on the side of the vial) and the part of the vial with the screw cap attached served as the mesocosm, the cap being the bottom of the mesocosm. For the field incubation, we sewed mesh bags from screen printing fabric (100% polyester, 120 threads per cm) with a mesh width of 40 μm, which allows the ingrowth of fungal hyphae but excludes the ingrowth of roots.

We set up four incubation treatments: i) 2-mm-sieved soil plus model chitin particles, ii) 2-mm-sieved soil plus model cellulose particles, iii) 2-mm-sieved soil plus model particles containing neither chitin nor cellulose, and iv) 2-mm-sieved soil without model particles (Fig. 1). We rinsed model particles, which had been stored in 20% ethanol, three times in 15 mL pure MilliQ water to remove residual ethanol. We added 1 g of model particles to 15 g of sieved soil to individual plastic bags and shook them to homogenize and distribute model particles evenly throughout the soil. Moreover, 15 g sieved soil which was incubated without model particles was shaken to guarantee proper homogenization. Soil (and model particles) was filled in lab mesocosms and field mesh bags (on April 28, 2021). Soil in mesocosms was compacted to resemble field density using a 50 mL syringe plunger. Mesocosms were covered with parafilm during incubation to prevent desiccation and they were weighed weekly to check and potentially adjust moisture content if necessary. Full mesh bags were sealed with a heat sealer and measured 7 cm x 10 cm in size. In total we set up 80 mesocosms and 20 mesh bags (Fig. 1).

Mesocosms were incubated at 11°C, matching the soil temperature in 5 centimeters soil depth measured on the day of soil sampling (April 26, 2021. Filled mesh bags were kept in closed plastic bags at 11°C in a lab incubator between setting them up and burying them in the field. Mesh bags were buried in the field at a depth of 5 cm (April 30, 2021).

Destructive harvest

Lab mesocosms were destructively harvested 1, 13, 29 and 50 days after the start of the lab incubation (start of incubation on April 28, 2021, destructive harvest on April 29, May 11, May 27, June 17, 2021). Field mesh bags were destructively harvested after 47 days of incubation in the field (filling of mesh bags on April 28, 2021, buried on April 30, 2021, destructive harvest on June 14, 2021). Upon destructive harvest, we pushed soil cores out of mesocosms and separated the model particles from soil by hand using tweezers. We washed the model particles to get rid of soil sticking to the particles. We submerged model particles in MilliQ water in a petri dish, carefully swayed petri dishes by hand, and discarded

liquid while holding back the particles with a strong magnet. We repeated the washing steps three times to obtain the particles and the directly attached microbial communities for subsequent analysis. The magnetic feature of particles facilitated the separation of model particles from the "dirty" liquid. Aliquots of washed model particles were (i) weighed for DNA extraction and (ii) fixed with ethanol for downstream microscopic analysis.

DNA extraction and sequencing

DNA extraction, library preparation, and amplicon sequencing were conducted as in Simon et al 2024. In short, we extracted DNA from a minimum of 20 model particles (150 mg ± 0.04) and 250 mg soil (253 mg ± 0.005) with the Qiagen PowerSoil kit (Qiagen, Valencia, CA, USA). Except for adding a bead beating step (30 sec), we followed the manufacturer´s protocol. For amplicon sequencing, we followed a two-step PCR barcoding approach (Pjevac et al., 2021) to amplify the V4 region of the 16S rRNA genes targeting bacteria and archaea, and the fungal internal transcribed spacer region 2 (ITS2) of the nuclear rRNA operon. We used primers 515F (Parada et al., 2016) and 806R (Apprill et al., 2015) for 16S rRNA genes. For the ITS2 region, a nested approach (Nilsson et al., 2019) was chosen, generating a PCR product using primers ITSOF-T (Taylor and McCormick, 2008; Tedersoo et al., 2008) and ITS4 (White et al., 1990), which was then used as a template for a second PCR using primers gITS7 (Ihrmark et al., 2012) and ITS4. Amplicons were sequenced on a MiSeq platform (Illumina, V3 chemistry, 600 cycles) at the Joint Microbiome Facility (JMF) of the Medical University of Vienna and the University of Vienna under project number JMF-2106-12. Raw data processing was performed as described previously (Pjevac et al., 2021).

Digital droplet PCR (ddPCR)

We quantified the number of copies of the V4 hypervariable region of the 16S rRNA gene and of the fungal ITS region of model particles and in soils with digital droplet PCR (ddPCR). Gene copy numbers can be used as a proxy for microbial biomass (Alteio et al., 2021). Prior to performing ddPCR, we measured DNA concentration of all DNA extracts using PicoGreen (Quant-iT PicoGreen dsDNA Assay Kit; Thermo Fisher Scientific). If the concentration was under the detection limit, we re-measured extracts using a Qubit high-sensitivity dsDNA assay (Thermo Fisher Scientific). For 16S ddPCR, DNA extracts were diluted to a concentration of 0.05 ng/µL and for ITS ddPCR to a concentration of 0.25 ng/µL. 2 µL of diluted DNA template were used as input for each ddPCR assay. Gene copy numbers per µL DNA extract were assessed with the QX200 Droplet Digital PCR (Bio-Rad) and analyzed with the QuantaSoft software (Bio-Rad).

Particle fixation and subsequent fluorescence microscopy

A fraction of model particles (~ 12 model particles) which had been incubated were fixed with
96% ethanol in 2 mL Eppendorf tubes. Tubes were stored at 4 °C overnight. Supernatant
was removed using a magnetic pulldown and model particles were resuspended in 1.5 mL of
1X PBS. These washing steps were repeated three additional times. Ethanol-fixed samples
were suspended in 1.5 mL of cold 96% ethanol. Model particles were inverted to mix.
Following fixation, samples were stored at -20 °C. DNA associated with model particles was
stained with the stain SYBR Green I nucleic acid stain (Invitrogen, Thermo Fisher). Selected
chitin particles stained with SYBR Green were analyzed using a Leica confocal laser
scanning microscope, coupled with a Thunder imaging system (Leica).

Data analyses

Amplicon sequence variants (ASVs) were inferred using the DADA2 R package v1.20
(https://www.ncbi.nlm.nih.gov/pubmed/27214047) applying the recommended workflow
(https://f1000research.com/articles/5-1492). 16S rRNA FASTQ reads 1 and 2 were trimmed
at 220 nucleotides (nt) and 150 nt respectively with allowed expected errors of 2. ITS FASTQ
reads 1 and 2 were trimmed at 230nt with allowed expected errors of 2 and 4 respectively.

16S rRNA ASV sequences were subsequently classified using DADA2 against the SILVA
database SSU Ref NR 99 release 138.1 (https://doi.org/10.5281/zenodo.4587955,
https://www.ncbi.nlm.nih.gov/pubmed/23193283) using a confidence threshold of 0.5. ITS
ASV sequences were classified against the UNITE all eukaryotes general FASTA release
version 8.2 (Abarenkov et al, 2020) using a confidence threshold of 0.5.

All data analyses were performed in R version 4.4.0 (R Core Team, 2021). We analyzed
samples that had yielded at least 3000 sequencing reads (16S rRNA genes: n = 172, ITS2
sequences: n = 179, see Table S1 and S2 for numbers of samples per treatment and time
point).

To compare the number of bacterial, archaeal, and fungal amplicon sequence variants
(ASVs) detected in the model particle fractions and in soil, we randomly rarefied samples to
3015 (16S rRNA genes) and 5676 (ITS2 sequences) reads to establish equal sequencing
depth across samples. To determine whether model substrate particles favored microbial
genera compared to the soil source community, we performed differential abundance
analysis based on relative abundance data using the R package ANCOMBC (v. 2.6.0) (Lin
and Peddada, 2020). Using the analysis of compositions of microbiomes with bias correction
(ANCOMBC), we determined genera which were significantly enriched and depleted in
microbial communities associated with model particles compared to the soil, in which
particles had been incubated ("surrounding soil"). In this study, we focus on the genera that

were significantly enriched on model particles compared to the "surrounding soil". In addition,
we examined temporal dynamics of particle-associated microbial communities by visualizing
changes of microbial community composition over the incubation period with principal
coordinate analysis (PCoA) plots. PCoA plots visualize Aitchison distances, Euclidean
distances calculated from centered-log-ratio transformed counts matrices, which account for
the compositional nature of amplicon sequencing data (Gloor et al., 2017).

Results

Microbial biomass increased on model particles over time, but not in the surrounding soil

We assessed microbial colonization and biomass of model particles using both droplet digital
PCR (ddPCR) and microscopic evaluation. Bacterial and archaeal, and fungal biomass
(significantly) increased on model particles over time, as evidenced by increasing gene copy
numbers on model chitin and cellulose particles over the 50 days and 47 days incubation
period for the lab mesocosms and the litter bags incubated in the field, respectively (Fig. 2A).
The number of 16S rRNA gene copies increased most on model chitin particles, from
45.13 ± 24.68 million gene copies per gram particles at the first time point to 328.61 ± 163.32
million after 50 and 262.19 ± 370.49 million after 47 incubation days, respectively. Bacterial
and archaeal gene copy numbers also slightly increased on "no substrate" model particles,
however not significantly (Fig. 2A). In contrast, fungal ITS copy numbers differed significantly
between particles after one day of incubation and 29 or 50 days of incubation (Fig. 2B). At
most time points, gene copy numbers were similarly small or large for model cellulose
particles and "no substrate" model particles, but larger on model chitin particles (Fig. 2A, 2B).
Particles incubated in the lab and the field were similarly densely populated by microbes after
47 and 50 days. In contrast, microbial biomass in the soil surrounding the model particles, as
well as soil incubated without additional substrate did not reveal a clear pattern over time
(Fig. 2C, 2D). Microscopy visualization of chitin particles supports the finding that particles
were more densely populated after several days of incubation than after one day (Fig. 2E).

ASV richness of model chitin particles decreased over the incubation period

To determine whether changes in microbial community structure occurred on model particles
and in soil across the different treatments, we investigated amplicon sequence variant (ASV)
richness of rarefied data. Overall, the richness of initial particle-associated microbial
communities was higher compared to communities associated with particles later in the
incubation period (Fig. 3). The number of bacterial and archaeal taxa associated with model
chitin particles dropped from 1583 ± 135 (median ± standard deviation) on the first day after
the incubation start to 908 ± 169 rarefied ASVs at the end of the incubation period (day 50,

Fig. 3A). Also, fungal richness of model chitin particles was lower after multiple days of incubation (332 ± 56.7 at day 50) than after one day of incubation (368 ± 44.8). Bacterial and archaeal, and fungal ASV richness detected on cellulose particles was slightly increased after 13, 29 and 50 days of incubation (16S rRNA: 13d: 1541±157, 29d: 1521±83.5, 50d: 1309±62.9; ITS2: 13d: 476±76.1, 29d: 498±77.3, 50d: 418±77) compared to particles after one day of incubation (16S rRNA genes: 1289±263 ASVs, ITS2: 277±88.5). Bacterial and archaeal ASV richness of "no substrate" model particles slightly decreased from 1566±234 (1d) to 1109±82.7 ASVs (50d), whereas fungal richness clearly increased over the incubation time (260±34 to 365±54.2 ASVs). In contrast to the substrate particles, microbial richness remained relatively more constant across time points and treatments (Fig. 3).

Chitin particle-attached microbial communities underwent temporal changes

Additionally, we investigated whether microbial community structure on model particles and the surrounding soil changed over the incubation period (Fig. 4). Differences in bacterial, archaeal, and fungal community composition between the time points were most pronounced for communities associated with model chitin particles (Fig. 4A, 4D). Communities that were detected on the first day of incubation were most similar to the communities of the soil which was used to set up lab mesocosms. Later communities (after more than one day of incubation (13, 29, and 50 days)) were further apart and hence differed more strongly from initial communities than communities after one incubation day. We found that bacterial and archaeal, and fungal communities that were associated with chitin particles, which had been incubated for 47 or 50 days in the field and the lab, were very similar in their composition (Fig. 4A, 4D). In comparison to model chitin particles, microbial communities associated with model cellulose particles did not show pronounced changes over the course of the incubation period (Fig. 4B, 4E). Bacterial and archaeal communities associated with "no substrate" model particles, the control particles that contained neither chitin nor cellulose, exhibited a similar, yet less pronounced temporal pattern as chitin particles (Fig. 4C). Bacterial and archaeal communities noticeably differed in their composition between "no substrate" model particles that were harvested after one day of incubation from particles that were harvested after multiple days of incubation (Fig. 4C). In comparison, fungal communities associated with control model particles (NS) did not show such a clear separation of the initial soil-communities and the communities associated with particles that were harvested after one incubation day from the communities of particles that were incubated for multiple days (Fig. 4F). Over all treatments, communities in soil did not exhibit strong changes over the incubation period. Composition of communities in soil was rather similar across the different harvest timepoints (1d, 13d, 29d, 50d, 47d) (Fig. 4G-4L).

Certain microbial genera were enriched on chitin and cellulose (particles)

Next, we compared relative abundances of microbial genera between the model substrate particles and the surrounding soil, which was used as the "baseline", to address whether substrates led to the enrichment of microbial genera on substrate particles using the differential abundance analysis R package ANCOMBC. In this study, we focused on genera which were significantly enriched in their relative abundances on model substrate particles compared to the soil (Fig. 5, 6). We did not only detect genera that were not enriched on model particles, but also genera that were significantly depleted on model particles compared to the surrounding soil (Fig. S1-S7). We observed that certain bacterial and fungal genera were significantly more abundant on model chitin and cellulose particles than in the soil in which these particles had been incubated (Fig. 5, 6). At the same time, other genera were significantly depleted in their relative abundances on model chitin and cellulose particles compared to soil (Fig. S1-S7). In the initial phase of colonization of particles (after one incubation day), few bacterial genera, *Paenibacillus* and *Bacillus* on chitin particles, and KD3-10, *Paenibacillus*, *Clostridium sensu stricto* 1, and *Micropepsis* on cellulose, were relatively significantly more abundant on model particles compared to the surrounding soil (Fig. 5A, 5B). *Asticcacaulis* and *Mucilaginibacter* were the two genera which were enriched on both model chitin and cellulose particles at all later harvest timepoints (13d, 29d, 50d, 47d) with the exception that *Asticaccaulis* was not detected on cellulose particles after 13 incubation days (Fig. 5). Most genera were observed amongst enriched genera only on one model substrate particle type or at certain time points. On chitin particles, bacterial genera *Alkanindiges*, *Silvimonas, Paucibacter, Roseatales,* and *Sphaerotilus* were amongst the most enriched genera at the later time points of the lab incubation (13d, 29d, 50d) (Fig. 5A). We show that a larger number of bacterial genera was significantly increased in their relative abundance on model chitin particles compared to soil than in the case of cellulose particles (Fig. 5). While the number of bacterial genera that were enriched on model chitin particles was similar for lab and field incubations after 47 and 50 days of incubation (Fig. 5A), more bacterial genera were enriched on cellulose particles that had been incubated in the field than in the lab (Fig. 5B). Most bacterial genera, whose relative abundances were significantly increased on particles compared to soil, matched between chitin and cellulose particles of field incubations (Fig. 5A, 5B).

We also found model substrate particles to be enriched in certain fungal genera compared to the surrounding soil (Fig. 6). On both model chitin and cellulose particles, *Trichoderma* was significantly relatively more abundant than in soil after one incubation day (Fig. 6A, 5B). Fungal genera that were significantly enriched on chitin particles differed among the harvest timepoints (Fig. 6A). *Pseudogymnascus* was relatively more abundant on cellulose particles than in soil after 13 and 29 days of lab and 47 days of field incubation (Fig. 6B). Cellulose

378 particles were also significantly enriched in *Ilyonectria* compared to corresponding soil after
379 one, 13 and 50 days (Fig. 6B). Overall, significant differences in genera between model chitin
380 particles and the soil were less pronounced for fungi than for bacteria (Fig. 6A, 5A).
381 Considerably fewer fungal than bacterial genera were significantly different in their relative
382 abundances between model substrate particles and the soil (Fig. 6, 5).

Discussion

Our work indicates that microbes that were detected on model particles were physically
attached to these particles, as they would otherwise have been removed during the washing
steps. These findings go in line with a previous study that proposed that bacteria and
archaea are widely surface-attached in soil as the energetic costs of motility in soils are high
(Nunan et al., 2020). At the same time, our study suggests that many bacterial and archaeal
taxa were able to move towards particles and attach to them within one day of particle
incubation in soil. It is likely that colonization of particles did not occur solely through adjacent
soil surfaces but rather that bacteria actively or passively moved to substrate particles. The
fast arrival of a large proportion of the microbial source community suggests that motility is
widespread among bacteria, archaea, and fungi in the here investigated forest soil. The
increase in microbial biomass associated with model substrate particles indicates either
continuous arrival of microbes on the particle surface or growth of already attached
microbes. Microbial growth would then be fostered by the degradation of chitin and cellulose
via extracellular enzymes which make resources available for microbial growth. It is possible
that a certain fraction of model particle-associated microbes was lost due to the employed
washing steps, which we cannot account for.

Analysis of changes in microbial richness revealed a decrease in bacterial, archaeal, and
fungal richness on model chitin particles over time. In contrast, bacterial and archaeal
richness remained unchanged on cellulose particles and "no substrate" particles, which
contained no additional substrate, during the incubation period and fungal richness even
increased on both model particle types. Bacterial, archaeal and fungal community
composition analysis showed that microbial communities associated with chitin particles
underwent temporal changes in community composition, whereas cellulose particles did not
exhibit clear temporal patterns. These findings indicate that chitin more strongly selects for
substrate-specific microbial consortia than cellulose. Communities associated with chitin
particles seemed to become more chitin-specific over time, which points towards microbial
communities undergoing temporal succession on particulate organic matter containing chitin
as has been shown by others (Enke et al., 2018). In contrast, cellulose seemingly did not
select for specific microbial communities as strongly as chitin. This might be a consequence
of the different chemical complexity of chitin, necessitating a specific suite of enzymes or

microbial interactions to be degraded. In contrast to model chitin particles, we did not detect
temporal patterns of microbial community composition for the soil, in which model chitin
particles had been incubated. Our results demonstrate that we might not be able to study the
effect of complex particulate substrates on microbial communities with composite soil
samples. Our results demonstrate that millimeter-sized model particles are a suitable study
system for investigating substrate-specific microbial communities in soil and their temporal
dynamics.

Fewer bacterial genera were enriched on model cellulose particles than on model chitin
particles in the lab incubation. This finding supports our claim that chitin exerts a stronger
selective pressure on microbial communities than cellulose does. The bacterial and fungal
genera that were enriched on model chitin and cellulose particles compared to the soil
metacommunity potentially were particularly well adapted to utilize or degrade chitin,
cellulose or their degradation by-products. Members of some bacterial and fungal genera
that were enriched on model substrate particles have been described to possess the genetic
potential to produce chitin or cellulose-degrading extracellular enzymes. After one incubation
day, chitin particles were enriched in the bacterial genus *Paenibacillus* and *Bacillus*, bacterial
genera which include known chitin degraders (Watanabe et al., 1990; Itoh et al., 2013).
Cellulose particles were also enriched in *Paenibacillus* as well as *Micropepsis* and
Clostridium sensu stricto after one incubation day, representing genera which contain known
degraders of cellulose or cellobiose, the repeating unit of cellulose (Harbison et al., 2017;
Kim et al., 2018; Yadav and Dubey, 2018; C.-J. Li et al., 2023). The bacterial genus
Paenibacillus was also significantly enriched on "no substrate" particles, which could indicate
that members of this genus are especially efficient in colonization of newly introduced
particulates to soil, irrespective of the specific substrate contained. The bacterial genus
Asticcacaulis (phylum Alphaproteobacteria) has been found to contain many sequences for
glycoside hydrolases, key enzymes targeting both cellulose and chitin, thereby supporting its
role as a generalist with regard to polymeric carbon degradation (Talamantes et al., 2016).
Members of *Mucilaginibacter* have been described to also possess the potential to produce
certain glycoside hydrolases and be capable of degrading chitin and cellulose (Wang et al.,
2020; Kim et al., 2022). Species of other significantly enriched genera on chitin and/or
cellulose particles, *Silvimonas* and *Collimonas,* have been found to possess chitinases (Yang
et al., 2005; Song et al., 2015, 2015; Ballhausen et al., 2016), and a member of *Roseateles*
was found to degrade cellulose (So et al., 2024). However, not all taxa detected on model
substrate particles were necessarily degrading the substrate themselves. Some taxa might
profit from the degradation products of enzymes produced by other microbes that are
present, or they might feed on metabolites of degraders. Additionally, it is important to note
that degradation potential for microorganisms is not necessarily conserved at the genus

level, and our ability to assign specific taxa to degradation of a specific substrate is limited
based on amplicon sequencing results. It would be advantageous if future studies would
investigate genes and extracellular enzyme activity of taxa which are associated with model
substrate particles to uncover taxa that degrade specific substrates.

Only a small number of fungal genera significantly increased in relative abundance on model
chitin and cellulose substrate particles compared to the surrounding soil. The ability to
degrade chitin or cellulose potentially is more widespread among fungal taxa than bacterial
taxa (Rytioja et al., 2014; Berlemont, 2017). Consequently, these specific substrates might
exhibit a lower selective pressure on fungi than on bacteria. Comparison of model particles,
which did not contain chitin nor cellulose, with the surrounding soil, shows that differences in
relative abundances of microbial genera between model substrate particles and the soil were
driven by the specific substrate and not the compounds used to produce model particles.

Field incubations showed different patterns of enrichment of microbial genera on model
substrate particles than lab incubations. Whereas more bacterial genera were enriched on
model chitin particles than on model cellulose particles in the lab, numbers were equal for
model substrate particles that had been incubated in the field. Moreover, more fungal genera
were enriched on model chitin and cellulose particles that were incubated in the field than in
the lab. For bacteria and fungi log fold changes were larger for field-incubated particles than
for lab-incubated particles. Differences in research findings can be expected between field
and lab incubations. Conditions in the lab are relatively more stable than in the field. In this
study, mesocosms containing model particles and soil were incubated at constant 11°C. In
contrast, the content of mesh bags that were buried in the field were exposed to temporal
fluctuations in soil temperature and soil moisture over the incubation period. Moreover, the
potential impact of plant exudates on microbial communities in the field is absent in the lab.
The here found differences between lab and field incubations underpin that caution must be
taken when translating findings of lab studies to nature.

Our study shows that model substrate particles are a suitable tool for studying substrate-
specific microbial communities on particulate organic matter in soil and their temporal
dynamics. To the best of our knowledge, model particles have been used to study particle-
associated microbial degrader communities in aquatic ecosystems (Cordero and Datta,
2016; Datta et al., 2016; Ebrahimi et al., 2019), however our study is the first to investigate
microbial communities associated with particulate organic matter in soil.

Our study underpins that self-manufactured model substrate particles provide a highly
flexible, standardized experimental system for the investigation of microbial degrader
communities (Mußmann et al., in prep, Cordero and Stocker, 2017; Enke et al., 2019). One
can choose the size and the shape of particles and add the substrate of interest to the

mixture, in particulate (Mußmann et al., in prep). Particles can be investigated under the
microscope due to the high transparency of the used gelling agent, Gelrite gellan gum. A
caveat of the here employed recipe for model particles may be the use of a carbon-
containing gelling agent as it could potentially be metabolized by microbes. It has been
shown that some microbes are able to degrade gellan gum (Das et al., 2015). However, all
polymers which could be used as an embedding matrix for substrates may be degraded by
some microorganisms and thus support microbial growth of some microorganisms. The low
increase in gene copies on gellan gum particles indicates that these particles were
unspecifically colonized by microbes. In contrast to the insignificant changes observed in this
study, Mußmann et al. (in prep) found significant but also substantially less growth on "no
substrate" particles compared to chitin and cellulose particles. Microbial biomass was overall
lower on "no substrate" model particles than on model cellulose particles and much lower
than on chitin particles at all time points. Moreover, our results suggest that the significant
enrichment of genera on model chitin and cellulose particles compared to the surrounding
soil was not the result of gellan gum, but a result of the presence of chitin or cellulose. Only
few genera including *Paenibacillus*, *Alkanindiges, Trichoderma,* and *Ilyonectria* were
enriched in their relative abundances on model chitin, cellulose, and "no substrate" particles.
Most genera that were enriched on model chitin and cellulose particles were not enriched on
"no substrate" particle controls. Especially after multiple days of incubation in the lab, we did
not find any fungal genera to be enriched on "no substrate" particles compared to the soil. In
addition, log fold changes of genera were much larger for chitin particles than for "no
substrate" particles. Nevertheless, we recommend including "no substrate" model particles
as controls for model particles which contain a specific substrate in all experiments. Also, it
might be advisable to shorten the here applied incubation duration in future experiments to
capture and study more fine-scale temporal dynamics of microbial communities associated
with particulate organic matter in soil.

Conclusion

This study demonstrates the value of lab-manufactured model substrate particles as a tool
for investigating substrate-specific microbial communities associated with particulate organic
matter and their temporal dynamics in soil. Our study demonstrates that complex substrates
like chitin or cellulose select for specific microbial communities of the soil microbiome.
Microbial communities associated with chitin undergo more strong temporal changes than
cellulose, demonstrating the specificity of microbial communities for substrates in soil. Future
studies could further advance our understanding which microbial taxa are able to degrade
complex biopolymers in soil by employing model substrate particles to isolate and
characterize substrate-specific microbial communities.

Acknowledgements

This research was funded by the European Research Council under the European Union's Horizon 2020 research and innovation program (grant agreement No. 819446). Lauren Alteio is funded by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 101060698. The authors would like to thank Benjamin Zwirzitz for his helpful advice and discussions throughout the course of this work. Marc Mußmann was supported by the Austrian Science Fund (FWF) project number P31010. We thank Sean Darcy, Ksenia Guseva, and Christian Ranits for their help in the field, and Petra Pjevac for advising and coordinating DNA extraction and amplicon sequencing. We additionally acknowledge the Life Science Computer Cluster (LiSC) of the University of Vienna for enabling the processing of the raw amplicon sequencing data.

Credit authorship contribution statement

Eva Simon: Formal analysis, Investigation, Visualization, Writing – original draft, Writing – review & editing. **Lauren Alteio**: Conceptualization, Formal analysis, Investigation, Methodology, Project administration, Writing – original draft, Writing – review & editing. **Alexander König:** Investigation, Writing – review & editing. **Bruna Imai**: Investigation, Writing – review & editing. **Julia Horak:** Investigation, Writing – review & editing. **Julia Wiesenbauer**: Investigation, Writing – review & editing. **Joana Séneca**: Data curation, Writing – review & editing. **Bela Hausmann**: Data curation, Writing – review & editing. **Marc Mußmann**: Methodology, Writing – review & editing. **Barbara Kitzler:** Resources, Writing – review & editing. **Christina Kaiser**: Conceptualization, Funding acquisition, Project administration, Supervision, Writing – review & editing.

Conflicts of Interest

The authors declare no conflict of interest.

Data Availability

Raw 16S rRNA genes and ITS2 amplicon sequencing data have been deposited in the NCBI Sequence Read Archive under the BioProject accession number PRJNA1208268. Bacterial, archaeal, and fungal gene copy numbers are available on the Zenodo repository (10.5281/zenodo.14614129).

Figure captions

Fig. 1 Schematic of the experimental set-up. (**A**) We set up four incubation treatments, incubating (i) self-manufactured model chitin (CH), (ii) cellulose (CE), (iii) "no substrate" (NS) particles and (iv) no model particles in 2-mm-sieved soil. We investigated five biological replicates by using soil sampled from five field plots. We set up four technical replicates per treatment and plot for the lab incubation, which we harvested at four time points, after one, 13, 29 or 50 days of incubation in the laboratory, and one technical replicate per treatment and plot for incubation in the field which we harvested after 47 days. (**B**) We mixed 1 g of model particles and 15 g 2-mm-sieved soil and filled it in lab mesocosms that were produced from 50 mL Greiner tubes or mesh bags for the field. Upon harvest of mesocosms and mesh bags, we manually separated model particles from the soil for subsequent analysis. (**C**) Self-manufactured model substrate particles. (**D**) Laboratory mesocosm filled with soil and model particles. (**E**) Mesh bag used for the field incubation.

Fig. 2 Bacterial, archaeal, and fungal biomass increased on model particles over the incubation period, but not in the surrounding soil. Boxplots show (**A, C**) 16S rRNA (16S) and (**B, D**) ITS gene copy numbers (per gram of particles or soil) (**A, B**) on model particles and (**C, D**) in the soil after zero, one, 13, 29, and 50 days of lab incubation, and 47 days of field incubation. Gene copy numbers can be interpreted as a proxy for microbial biomass associated with model particles or soil. (**A, B, C, D**) Letters above boxes denote significant differences across timepoints for each treatment and sample type, letters are based on Bonferroni-adjusted p-values of pairwise comparisons that were calculated by performing Dunn´s tests as post hoc test of Kruskal Wallis tests. (**E**) Fluorescence microscopy pictures show microbial DNA at model chitin particle surfaces after one, 13, 29, and 50 days of lab incubation. Yellow points represent DNA stained with SYBR Green I double stranded nucleic acid stain. Microscopy was performed using a Leica confocal laser scanning microscope coupled with a Thunder imaging system. Scale bars indicate 1 µm.

Fig. 3 Bacterial and archaeal, and fungal richness of model chitin particles significantly decreased over the incubation period. Boxplots (from left to right) show (**A**) bacterial and archaeal richness (number of ASVs), and (**B**) fungal richness that was detected in (i) the soil, which was used to set up incubations, (ii) on model particles, and (iii) in the surrounding soil after one, 13, 29, and 50 days of lab incubation and 47 days field incubation. Sequencing data was randomly rarefied at the minimum read count for bacteria and archaea (3015 reads), and for fungi (5676 reads) to guarantee equal sequencing depth of samples. Letters above boxes denote significant differences across timepoints for each treatment and sample type, letters are based on Bonferroni-adjusted p-values of pairwise comparisons that were calculated by performing Dunn´s tests as post hoc test of Kruskal Wallis tests. S –

sieved soil that was used to set up incubations, CH – chitin, CE – cellulose, NS – no

substrate (control to account for the effect of the scaffold material of model particles, gellan

gum), NP – soil without particles.

Fig. 4 Chitin particle-associated bacterial and archaeal communities underwent

temporal changes. PCoA plots show potential dissimilarity of (**A-C, G-I**) bacterial and

archaeal, and (**D-F, J-L**) fungal communities (i) associated with model chitin, cellulose or "no

substrate" particles or (ii) in the surrounding soil across harvest time points (day one to 50 or

47). Plots also include communities of the five sieved soils (at incubation start, 0d) that were

used to set up mesocosms and mesh bags. PCoA plots are based on Aitchison distances

(Euclidean distance of clr-transformed sequencing data). The colour of data points indicates

the harvest time point, the shape of points depicts the type of incubation (lab or field). CH –

chitin, CE – cellulose, NS – "no substrate".

Fig. 5 Model chitin and cellulose particles favored certain bacterial genera compared

to the soil, in which particles had been incubated. Plots show genera that were

significantly enriched in their relative abundance on (**A**) model chitin, (**B**) model cellulose

particles, and (**C**) "no substrate" model particles (controls) compared to the surrounding soil

at one day, 13, 29, and 50 days after the start of the lab incubation and 47 days after the

start of the field incubation. Bars represent the log fold change, a measure of effect size, in

the relative abundance of significantly enriched genera (FDR-adjusted P value < 0.05).

Differentially abundant genera were identified using the ANCOM-BC package (Lin and

Peddada, 2020). Error bars represent the standard error of the log fold change. We

controlled for false discovery rates with the Benjamini-Hochberg multiple test correction.

Aquaspirillum* refers to the *[Aquaspirillum] articum group*, *Rhizobiaceae spp.*** refers to the

Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium clade. Note that we did not detect

any genera that were significantly enriched on model "no substrate" particles after 47 days of

incubation in the field. See Fig. S1-S4 for positive and negative log fold changes of genera

on model particles compared to the surrounding soil.

Fig. 6 Model chitin and cellulose particles favored certain fungal genera compared to

the soil, in which particles had been incubated. Plots show genera which were

significantly enriched in their relative abundance on (**A**) model chitin, (**B**) model cellulose

particles, and (**C**) model "no substrate" particles (controls) compared to the surrounding soil

at one day, 13, 29, and 50 days after the start of the lab incubation and 47 days after the

start of the field incubation. Bars represent the log fold change, a measure of effect size, in

the relative abundance of significantly enriched genera (FDR-adjusted P value < 0.05).

Differentially abundant genera were identified using the ANCOM-BC package (Lin and

Peddada, 2020). Error bars represent the standard error of the log fold change. We

controlled for false discovery rates with the Benjamini-Hochberg multiple test correction. Note that we did not detect any genera which were significantly enriched on model "no substrate" particles after 13 and 50 days of incubation in the lab, and 47 days of incubation in the field. See Fig. S5-S7 for positive and negative log fold changes of genera on model particles compared to the surrounding soil.

References

Abarenkov, K., Zirk, A., Piirmann, T., Pöhönen, R., Ivanov, F., Nilsson, R. H., Kõljalg, U., 2020. UNITE general FASTA release for eukaryotes 2. Version 04.02.2020. UNITE Community, https://doi.org/10.15156/BIO/786371

Allison, S.D., Weintraub, M.N., Gartner, T.B., Waldrop, M.P., 2010. Evolutionary-Economic Principles as Regulators of Soil Enzyme Production and Ecosystem Function, in: Shukla, G., Varma, A. (Eds.), Soil Enzymology, Soil Biology. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 229–243. doi:10.1007/978-3-642-14225-3_12

Alteio, L.V., Séneca, J., Canarini, A., Angel, R., Jansa, J., Guseva, K., Kaiser, C., Richter, A., Schmidt, H., 2021. A critical perspective on interpreting amplicon sequencing data in soil ecological research. Soil Biology and Biochemistry 160, 108357. doi:10.1016/j.soilbio.2021.108357

Angst, G., Mueller, K.E., Nierop, K.G.J., Simpson, M.J., 2021. Plant- or microbial-derived? A review on the molecular composition of stabilized soil organic matter. Soil Biology and Biochemistry 156, 108189. doi:10.1016/j.soilbio.2021.108189

Apprill, A., McNally, S., Parsons, R., Weber, L., 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton 75, 129–137. doi:10.3354/ame01753

Arnosti, C., Bell, C., Moorhead, D.L., Sinsabaugh, R.L., Steen, A.D., Stromberger, M., Wallenstein, M., Weintraub, M.N., 2014. Extracellular enzymes in terrestrial, freshwater, and marine environments: Perspectives on system variability and common research needs. Biogeochemistry 117, 5–21. doi:10.1007/s10533-013-9906-5

Ballhausen, M.-B., Vandamme, P., Boer, W. de, 2016. Trait Differentiation within the Fungus-Feeding (Mycophagous) Bacterial Genus Collimonas. PLOS ONE 11, e0157552. doi:10.1371/journal.pone.0157552

Bardgett, R.D., Van Der Putten, W.H., 2014. Belowground biodiversity and ecosystem functioning. Nature 515, 505–511. doi:10.1038/nature13855

Berlemont, R., 2017. Distribution and diversity of enzymes for polysaccharide degradation in fungi. Scientific Reports 7, 222. doi:10.1038/s41598-017-00258-w

Berlemont, R., Martiny, A.C., 2015. Genomic Potential for Polysaccharide Deconstruction in
Bacteria. Applied and Environmental Microbiology 81, 1513–1519.
doi:10.1128/AEM.03718-14

Berlemont, R., Martiny, A.C., 2013. Phylogenetic Distribution of Potential Cellulases in
Bacteria. Applied and Environmental Microbiology 79, 1545–1554.
doi:10.1128/AEM.03305-12

Cordero, O.X., Datta, M.S., 2016. Microbial interactions and community assembly at
microscales. Current Opinion in Microbiology 31, 227–234.
doi:10.1016/j.mib.2016.03.015

Cordero, O.X., Stocker, R., 2017. A particularly useful system to study the ecology of
microbes. Environmental Microbiology Reports 9, 16–17. doi:10.1111/1758-
2229.12490

Cotrufo, M.F., Wallenstein, M.D., Boot, C.M., Denef, K., Paul, E., 2013. The Microbial
Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition
with soil organic matter stabilization: Do labile plant inputs form stable soil organic
matter? Global Change Biology 19, 988–995. doi:10.1111/gcb.12113

Das, N., Triparthi, N., Basu, S., Bose, C., Maitra, S., Khurana, S., 2015. Progress in the
development of gelling agents for improved culturability of microorganisms. Frontiers
in Microbiology 6. doi:10.3389/fmicb.2015.00698

Das, S.N., Sarma, P.V.S.R.N., Neeraja, C., Malati, N., Podile, A.R., 2010. Members of
Gammaproteobacteria and Bacilli represent the culturable diversity of chitinolytic
bacteria in chitin-enriched soils. World Journal of Microbiology and Biotechnology 26,
1875–1881. doi:10.1007/s11274-010-0369-8

Datta, M.S., Sliwerska, E., Gore, J., Polz, M.F., Cordero, O.X., 2016. Microbial interactions
lead to rapid micro-scale successions on model marine particles. Nature
Communications 7, 11965. doi:10.1038/ncomms11965

Ebrahimi, A., Schwartzman, J., Cordero, O.X., 2019. Cooperation and spatial self-
organization determine rate and efficiency of particulate organic matter degradation in
marine bacteria. Proceedings of the National Academy of Sciences of the United
States of America 116, 23309–23316. doi:10.1073/pnas.1908512116

Enke, T.N., Datta, M.S., Schwartzman, J., Barrere, J., Pascual-garcı, A., Cordero, O.X.,
Pascual-garcı, A., 2019. Modular Assembly of Polysaccharide- Degrading Marine
Microbial Communities Report Modular Assembly of Polysaccharide- Degrading
Marine Microbial Communities 1528–1535. doi:10.1016/j.cub.2019.03.047

Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V., Egozcue, J.J., 2017. Microbiome
datasets are compositional: And this is not optional. Frontiers in Microbiology 8, 2224.
doi:10.3389/fmicb.2017.02224

Gómez, E.J., Delgado, J.A., González, J.M., 2020. Persistence of microbial extracellular
enzymes in soils under different temperatures and water availabilities. Ecology and
Evolution 10, 10167–10176. doi:10.1002/ece3.6677

Cordero, O.X., Stocker, R., 2017. A particularly useful system to study the ecology of
microbes. Environmental Microbiology Reports 9, 16–17. doi:10.1111/1758-
2229.12490

Enke, T.N., Datta, M.S., Schwartzman, J., Barrere, J., Pascual-garcı, A., Cordero, O.X.,
Pascual-garcı, A., 2019. Modular Assembly of Polysaccharide- Degrading Marine
Microbial Communities Report Modular Assembly of Polysaccharide- Degrading
Marine Microbial Communities 1528–1535. doi:10.1016/j.cub.2019.03.047

Pjevac, P., Hausmann, B., Schwarz, J., Kohl, G., Herbold, C.W., Loy, A., Berry, D., 2021. An
Economical and Flexible Dual Barcoding, Two-Step PCR Approach for Highly
Multiplexed Amplicon Sequencing. Frontiers in Microbiology 12, 669776.
doi:10.3389/fmicb.2021.669776

Harbison, A.B., Price, L.E., Flythe, M.D., Bräuer, S.L., 2017. Micropepsis pineolensis gen.
nov., sp. nov., a mildly acidophilic alphaproteobacterium isolated from a poor fen, and
proposal of Micropepsaceae fam. nov. within Micropepsales ord. nov. International
Journal of Systematic and Evolutionary Microbiology 67, 839–844.
doi:10.1099/ijsem.0.001681

Ihrmark, K., Bödeker, I.T.M., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., Strid,
Y., Stenlid, J., Brandström-Durling, M., Clemmensen, K.E., Lindahl, B.D., 2012. New
primers to amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial
and natural communities. FEMS Microbiology Ecology 82, 666–677.
doi:10.1111/j.1574-6941.2012.01437.x

Itoh, T., Hibi, T., Fujii, Y., Sugimoto, I., Fujiwara, A., Suzuki, F., Iwasaki, Y., Kim, J.-K.,
Taketo, A., Kimoto, H., 2013. Cooperative Degradation of Chitin by Extracellular and
Cell Surface-Expressed Chitinases from Paenibacillus sp. Strain FPU-7. Applied and
Environmental Microbiology 79, 7482–7490. doi:10.1128/AEM.02483-13

Kim, E.S., Kim, B.-S., Kim, K.-Y., Woo, H.-M., Lee, S.-M., Um, Y., 2018. Aerobic and
anaerobic cellulose utilization by *Paenibacillus* sp. CAA11 and enhancement of its
cellulolytic ability by expressing a heterologous endoglucanase. Journal of
Biotechnology 268, 21–27. doi:10.1016/j.jbiotec.2018.01.007

Kim, J., Lee, B., Chhetri, G., Kim, I., So, Y., Jang, W., Seo, T., 2022. Identification of
Mucilaginibacter conchicola sp. nov., Mucilaginibacter achroorhodeus sp. nov. and
Mucilaginibacter pallidiroseus sp. nov. and emended description of the genus
Mucilaginibacter. International Journal of Systematic and Evolutionary Microbiology
72. doi:10.1099/ijsem.0.005431

Lavallee, J.M., Soong, J.L., Cotrufo, M.F., 2020. Conceptualizing soil organic matter into
particulate and mineral-associated forms to address global change in the 21st
century. Global Change Biology 26, 261–273. doi:10.1111/gcb.14859

Lehmann, J., Hansel, C.M., Kaiser, C., Kleber, M., Maher, K., Manzoni, S., Nunan, N.,
Reichstein, M., Schimel, J.P., Torn, M.S., Wieder, W.R., Kögel-Knabner, I., 2020.
Persistence of soil organic carbon caused by functional complexity. Nature
Geoscience 13, 529–534. doi:10.1038/s41561-020-0612-3

Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. Nature 528,
60–68. doi:10.1038/nature16069

Lehmann, J., Solomon, D., Kinyangi, J., Dathe, L., Wirick, S.U.E., Jacobsen, C., 2008.
Spatial complexity of soil organic matter forms at nanometre scales. Nature Geosc 1,
238–242. doi:10.1038/ngeo155

Li, C.-J., Zhang, Z., Zhan, P.-C., Lv, A.-P., Li, P.-P., Liu, L., Li, W.-J., Yang, L.-L., Zhi, X.-Y.,
2023. Comparative genomic analysis and proposal of *Clostridium yunnanense* sp.
nov., *Clostridium rhizosphaerae* sp. nov., and *Clostridium paridis* sp. nov., three novel
Clostridium sensu stricto endophytes with diverse capabilities of acetic acid and
ethanol production. Anaerobe 79, 102686. doi:10.1016/j.anaerobe.2022.102686

Li, D., Zhang, Z., Wang, J., Zhang, P., Liu, Y., Li, Y., 2023. Estimate of the degradation
potentials of cellulose, xylan, and chitin across global prokaryotic communities.
Environmental Microbiology 25, 397–409. doi:10.1111/1462-2920.16290

Lin, H., Peddada, S.D., 2020. Analysis of compositions of microbiomes with bias correction.
Nature Communications 11, 3514. doi:10.1038/s41467-020-17041-7

Mußmann, M. et al., in prep.. Substrate-coated model particles enable rapid *in situ*
enrichment of macromolecule-degrading microbes in anoxic marine sediments.

Nilsson, R.H., Anslan, S., Bahram, M., Wurzbacher, C., Baldrian, P., Tedersoo, L., 2019.
Mycobiome diversity: high-throughput sequencing and identification of fungi. Nature
Reviews Microbiology 17, 95–109. doi:10.1038/s41579-018-0116-y

Nunan, N., Schmidt, H., Raynaud, X., 2020. The ecology of heterogeneity: soil bacterial
communities and C dynamics. Philosphical Transactions B 375. doi:10.1007/s00248-
002-1044-x

Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series
and global field samples. Environmental Microbiology 18, 1403–1414.
doi:10.1111/1462-2920.13023

Pjevac, P., Hausmann, B., Schwarz, J., Kohl, G., Herbold, C.W., Loy, A., Berry, D., 2021. An
Economical and Flexible Dual Barcoding, Two-Step PCR Approach for Highly

Multiplexed Amplicon Sequencing. Frontiers in Microbiology 12, 669776.
doi:10.3389/fmicb.2021.669776

R Core Team, 2021. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria.

Rytioja, J., Hildén, K., Yuzon, J., Hatakka, A., de Vries, R.P., Mäkelä, M.R., 2014. Plant-
Polysaccharide-Degrading Enzymes from Basidiomycetes. Microbiology and
Molecular Biology Reviews 78, 614–649. doi:10.1128/MMBR.00035-14

Schlüter, S., Leuther, F., Albrecht, L., Hoeschen, C., Kilian, R., Surey, R., Mikutta, R., Kaiser,
K., Mueller, C.W., Vogel, H.J., 2022. Microscale carbon distribution around pores and
particulate organic matter varies with soil moisture regime. Nature Communications
13, 2098. doi:10.1038/s41467-022-29605-w

Simon, E., Guseva, K., Darcy, S., Alteio, L., Pjevac, P., Schmidt, H., Jenab, K., Ranits, C.,
Kaiser, C., 2024. Distinct microbial communities are linked to organic matter
properties in millimetre-sized soil aggregates. The ISME Journal 18, wrae156.
doi:10.1093/ismejo/wrae156

Sinsabaugh, R.L., Hill, B.H., Follstad Shah, J.J., 2009. Ecoenzymatic stoichiometry of
microbial organic nutrient acquisition in soil and sediment. Nature 462, 795–798.
doi:10.1038/nature08632

Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw, C.,
Contosta, A.R., Cusack, D., Frey, S., Gallo, M.E., Gartner, T.B., Hobbie, S.E.,
Holland, K., Keeler, B.L., Powers, J.S., Stursova, M., Takacs-Vesbach, C., Waldrop,
M.P., Wallenstein, M.D., Zak, D.R., Zeglin, L.H., 2008. Stoichiometry of soil enzyme
activity at global scale. Ecology Letters 11, 1252–1264. doi:10.1111/j.1461-
0248.2008.01245.x

So, Y., Chhetri, G., Kim, I., Park, S., Jung, Y., Seo, T., 2024. Roseateles caseinilyticus sp.
nov. and Roseateles cellulosilyticus sp. nov., isolated from rice paddy field soil.
Antonie Van Leeuwenhoek 117, 87. doi:10.1007/s10482-024-01988-4

Song, C., Schmidt, R., de Jager, V., Krzyzanowska, D., Jongedijk, E., Cankar, K.,
Beekwilder, J., van Veen, A., de Boer, W., van Veen, J.A., Garbeva, P., 2015.
Exploring the genomic traits of fungus-feeding bacterial genus Collimonas. BMC
Genomics 16, 1103. doi:10.1186/s12864-015-2289-3

Talamantes, D., Biabini, N., Dang, H., Abdoun, K., Berlemont, R., 2016. Natural diversity of
cellulases, xylanases, and chitinases in bacteria. Biotechnology for Biofuels 9, 133.
doi:10.1186/s13068-016-0538-6

Taylor, D.L., McCormick, M.K., 2008. Internal transcribed spacer primers and sequences for
improved characterization of basidiomycetous orchid mycorrhizas. New Phytologist
177, 1020–1033. doi:10.1111/j.1469-8137.2007.02320.x

Tedersoo, L., Jairus, T., Horton, B.M., Abarenkov, K., Suvi, T., Saar, I., Kõljalg, U., 2008.
Strong host preference of ectomycorrhizal fungi in a Tasmanian wet sclerophyll forest
as revealed by DNA barcoding and taxon-specific primers. New Phytologist 180,
479–490. doi:10.1111/j.1469-8137.2008.02561.x

Wang, X., Zohar-Perez, C., Zeng, Y., Zou, Y., Chen, Y., Wu, S., Wang, Y., Arazi, S.,
Nussinovitch, A., Achmon, Y., 2023. Assessment of the environmental impact of agar,
alginate, and gellan gum carbohydrate gum macro beads biodegradation in a
simulated agricultural field system. Environmental Technology & Innovation 30,
103034. doi:10.1016/j.eti.2023.103034

Wang, Z.Y., Wang, R.X., Zhou, J.S., Cheng, J.F., Li, Y.H., 2020. An assessment of the
genomics, comparative genomics and cellulose degradation potential of
Mucilaginibacter polytrichastri strain RG4-7. Bioresource Technology 297, 122389.
doi:10.1016/j.biortech.2019.122389

Watanabe, T., Oyanagi, W., Suzuki, K., Tanaka, H., 1990. Chitinase system of Bacillus
circulans WL-12 and importance of chitinase A1 in chitin degradation. Journal of
Bacteriology 172, 4017–4022. doi:10.1128/jb.172.7.4017-4022.1990

White, T.J., Bruns, T., Lee, S., Taylor, J., 1990. Amplification and direct sequencing of fungal
ribosomal RNA genes for phylogenetics, in: PCR Protocols. Elsevier, pp. 315–322.
doi:10.1016/B978-0-12-372180-8.50042-1

Wieczorek, A.S., Hetz, S.A., Kolb, S., 2014. Microbial responses to chitin and chitosan in
oxic and anoxic agricultural soil slurries. Biogeosciences 11, 3339–3352.
doi:10.5194/bg-11-3339-2014

Witzgall, K., Vidal, A., Schubert, D.I., Höschen, C., Schweizer, S.A., Buegger, F., Pouteau,
V., Chenu, C., Mueller, C.W., 2021. Particulate organic matter as a functional soil
component for persistent soil organic carbon. Nature Communications 12, 4115.
doi:10.1038/s41467-021-24192-8

Wu, H., Cui, H., Fu, C., Li, R., Qi, F., Liu, Z., Yang, G., Xiao, K., Qiao, M., 2024. Unveiling the
crucial role of soil microorganisms in carbon cycling: A review. Science of The Total
Environment 909, 168627. doi:10.1016/j.scitotenv.2023.168627

Yadav, S., Dubey, S.K., 2018. Cellulose degradation potential of *Paenibacillus lautus* strain
BHU3 and its whole genome sequence. Bioresource Technology 262, 124–131.
doi:10.1016/j.biortech.2018.04.067

Yang, H.-C., Im, W.-T., An, D.-S., Park, W.-S., Kim, I.S., Lee, S.-T., 2005. Silvimonas terrae
gen. nov., sp. nov., a novel chitin-degrading facultative anaerobe belonging to the
"Betaproteobacteria." International Journal of Systematic and Evolutionary
Microbiology 55, 2329–2332. doi:10.1099/ijs.0.63837-0

Young, I.M., Crawford, J.W., Nunan, N., Otten, W., Spiers, A., 2008. Chapter 4 Microbial
Distribution in Soils. Physics and Scaling, in: Advances in Agronomy. Elsevier Inc.,
pp. 81–121. doi:10.1016/S0065-2113(08)00604-4
Zimmerman, A., Martiny, A., Allison, S., 2013. Microdiversity of extracellular enzyme genes
among sequenced prokaryotic genomes. The ISME Journal 7, 1187–1199.
doi:10.1038/ismej.2012.176
848

A — Chitin

- **1d:** Trichoderma, Solicoccozyma
- **13d:** Umbelopsis, Mortierella
- **29d:** Clonostachys, Trichoderma, Umbelopsis, Mortierella, Dipodascus, Monocillium, Ostropales, Phragmocephala
- **50d:** Ophiostomatales
- **47d:** Metapochonia, Leptodontidium, Chaetosphaeriaceae, Pochonia, Pseudogymnoascus, Hyaloscypha, Cryptosporiopsis, Mortierella

B — Cellulose

- **1d:** Ilyonectria, Trichoderma
- **13d:** Pseudogymnoascus, Hypoxylon, Ilyonectria, Pezicula
- **29d:** Pseudogymnoascus, Mortierella, Infundichalara
- **50d:** Ilyonectria, Meripilaceae
- **47d:** Ceratobasidium, Pseudogymnoascus, Leotiomycetes, Nigrograna

C — No Substrate (particle control)

- **1d:** Ilyonectria, Apiotrichum, Trichoderma, Metapochonia, Solicoccozyma, Nectria
- **29d:** Byssocorticium, Botryobasidium, Arthrographis, Basidiodendron, Cyberlindnera, Diaporthales, Gymnascella, Nemania, Paramicrosporidium, Phaeosphaeriaceae, Preussia, Rhinocladiella