Mechanical loading reveals cell type-specific responses and role of PHGDH in endothelial cell growth

Sakari Mäntyselkä[1*], Erik Niemi[1*], Laura Ylä-Outinen[1], Kalle Kolari[1], Liina Uusitalo-Kylmälä[1], Alfredo Ortega-Alonso[1], Roosa-Maria Liimatainen[2], Vasco Fachada[1], Perttu Permi[2,3,4], Elina Kalenius[2], Juha J Hulmi[1#], Riikka Kivelä[1,5,6#]

[1]Faculty of Sport and Health Sciences, NeuroMuscular Research Center, University of Jyväskylä, Jyväskylä, Finland
[2]Department of Chemistry, University of Jyväskylä, Jyväskylä, Finland
[3]Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland
[4]Institute of Biotechnology, Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland
[5]Stem Cells and Metabolism Research Program, Research Programs Unit, Faculty of Medicine, University of Helsinki, Helsinki, Finland
[6]Wihuri Research Institute, Helsinki, Finland

* Equal contribution as first authors

# Equal contribution as senior authors

Correspondence to Riikka Kivelä (riikka.m.kivela@jyu.fi) & Juha J Hulmi (juha.hulmi@jyu.fi), NeuroMuscular Research Center, Faculty of Sport and Health Sciences, University of Jyväskylä, Rautpohjankatu 8, 40014 Finland

Keywords

Acetate/Cell-stretching/Exercise/Muscle/Serine

Abstract

Skeletal muscle and blood vessels are constantly subjected to mechanical forces, especially during exercise. We subjected human endothelial cells (ECs) and muscle cells (MCs) to cyclic mechanical stretch and investigated acute transcriptomic and metabolomic changes. Mechanical loading induced several responses typically seen after exercise in vivo. Both cell types released acetate in response to stretching, but surprisingly, many of the other changes were in opposite directions. For instance, electron transport chain genes were downregulated in ECs and upregulated in muscle cells. Additionally, ECs shifted their transcriptomic profile to a growth-oriented direction and ^{13}C-(U)-glucose tracing experiments revealed increased serine synthesis in ECs. Mechanistic experiments related to the serine synthesis pathway enzyme phosphoglycerate dehydrogenase (PHGDH) revealed its importance in EC anabolism. Our data suggest that mechanical loading induces cell type-specific responses in ECs and MCs, and several of these effects recapitulate exercise effects *in vivo*. In ECs, mechanical loading activates the serine synthesis pathway, and the serine-synthesizing enzyme PHGDH was highlighted as essential in regulating EC growth, a process indispensable in angiogenesis.

Introduction

Skeletal muscle is highly vascularized and myofibers and endothelial cells together
make up ~70% of the total nuclei count in skeletal muscle tissue (Englund *et al*,
2021; Pepe & Albrecht, 2023). Myofibers are contractile cells that both produce and
sense mechanical forces. Endothelial cells lining the blood vessels sense
mechanical forces produced by blood flow (shear stress) and contracting and
stretching muscle. In skeletal muscle, mechanical forces are produced and sensed
especially during exercise, which can be mimicked *in vitro* using contraction and
stretching models (Lautaoja *et al*, 2023). Previously, it has been demonstrated that
cyclic mechanical loading (i.e. cyclic stretching) induces a wide variety of molecular
and morphological responses in endothelial (reviewed in Jufri *et al*, 2015) and
muscle cells (reviewed in Ren *et al*, 2021). For example, mechanical loading has
been found to activate the mechanistic target of rapamycin complex 1 (mTORC1)
(Nakai *et al*, 2015) and to induce highly similar gene expression changes in
myofiber-like multinucleated cells (myotubes) as resistance exercise *in vivo* in
myofibers (Turner *et al*, 2021). Endothelial cells, in turn, have been shown to orient
in a 60-70° angle relative to the direction of stretch, secrete angiocrines, and activate
angiogenesis-related pathways (Große-Segerath *et al*, 2024; Jufri *et al*, 2015; Lorenz
et al, 2018), which also occur in response to exercise and increased blood flow *in
vivo* (Biernat *et al*, 2024; Sottiurai *et al*, 1989). However, there are only a few studies
(Große-Segerath *et al*, 2024; Hinkle *et al*, 2022; Ilaiwy *et al*, 2016) investigating the
remodeling of the global molecular landscape in response to mechanical loading,
and none in comparing responses in different cell types. It is recently acknowledged
that endothelial and muscle cell cross-talk is essential in regulating the development,
maintenance, and repair of muscle tissue (reviewed in Pepe & Albrecht, 2023).

Skeletal muscle responses to mechanical loading *in vivo* have mostly been
examined using mixed muscle samples, which lack cell-specificity as skeletal muscle
contains over 10 different cell types (Englund *et al*, 2021). *In vitro* models allow us to
study the metabolic responses in a cell type-specific manner. However, according to
our knowledge, there are no studies in which mechanical loading-induced responses
of muscle and endothelial cells have been compared. Exercise induces many
metabolic changes such as increased glycolysis and oxidative phosphorylation in the
whole muscle level (Smith *et al*, 2023). However, little is known about how exercise
or mechanical loading *per se* affects other metabolic pathways such as those
branching from glycolysis (e.g., the pentose phosphate and serine synthesis
pathways), and their role in exercise responses such as angiogenesis. Previously,
mechanical stretching of liver endothelial cells during vasodilation was demonstrated
to induce angiocrine signals that support liver growth and survival, and potentially
contribute to liver regeneration (Lorenz *et al*, 2018). Blood vessels in skeletal
muscles experience immediate vasodilation—and thus endothelial cell stretching—
when their physiological function is extensively needed during exercise.

Here we mechanically loaded two different human endothelial cell lines and primary
human skeletal muscle cells to mimic exercise on-a-dish and to study cell type-
specific responses. Transcriptomics and metabolomics analyses revealed unique

oxidative and glycolytic changes in these two cell types in response to mechanical
loading. Further mechanistic experiments identified the serine synthesis pathway
activation, and specifically phosphoglycerate dehydrogenase (PHGDH), as an
essential driver of endothelial cell anabolism.

Results

Mechanical loading induced transcriptomic changes in all cell types and altered exercise-related and mechanosensitive gene expression

We first investigated the transcriptomic changes in two different types of human
endothelial cell lines (HUVEC & HCMEC) as well as human skeletal muscle cells
(HSkM) (Fig. 1A). The mechanical loading did not induce detachment of the cells,
significant membrane damage, or cell death in either muscle or endothelial cells as
determined by total RNA yield and the extracellular lactate dehydrogenase (LDH)
activity assay (Supplementary Fig. 1A-B). Based on the differential gene expression
analysis, we found that all three cell types presented significant changes in their
transcriptome in response to the mechanical loading, as demonstrated by the
principal component analysis (PCA) (Fig. 1B). The transcriptomic response was the
largest in HUVECs, with the total number of differentially expressed genes being
approximately 3-fold larger than in HCMECs or HSkM cells (Fig. 1C).

In HUVECs, 717 genes were upregulated, and 779 genes were downregulated
($\log_2$FC ≥ 0.5 & adj. p ≤ 0.05). By indexing genes in order of magnitude and
consistency of change ($\log_2$FC and adj. p), the five most significantly upregulated
protein-coding genes were *STC1*, *ADAM19*, *INSR*, *AKAP12* and *GUCY1A1*. The five
most significantly downregulated protein-coding genes by the same indexing were
H2-4 clustered histones *H3C12*, *H4C9*, *H3C8*, *H3C13* and *H2AC16*. In HCMECs,
241 genes were upregulated, and 261 genes were downregulated. The five most
significantly upregulated protein-coding genes were *KITLG*, *ITGB8*, *FGF5*, *FBXO30*
and *SLC7A11* and, similarly to HUVECs, the five most significantly downregulated
ones were again H2-4 clustered histones *H4C9*, *H3C12*, *H4C6* and *H2BC7,* as well
as H1.3 linker histone *H1-3*. A total of 388 genes in HSkM cells were differentially
expressed, from which 262 genes were upregulated and 126 genes were
downregulated in response to mechanical loading. The five most significantly
upregulated protein-coding genes were *THBD, IL11, GPRC5A, SNAI1* and *RASD1*
and the five most significantly downregulated protein-coding genes were *CCN1,*
CCN2, *RGS4*, *DLC1* and *F3*. For the complete lists of differentially expressed genes
and their full names, please refer to Supplementary Table S1.

Moreover, when comparing overlaps in both up- and downregulated genes in all
three cell lines, we noticed that the two endothelial cell lines clearly showed
similarities in their gene expression, whereas the overlaps in gene expression
between endothelial cells and muscle cells were small (Fig. 1E). Only one gene was

upregulated (*CREB5*) in all three cell lines and similarly, only one gene was
downregulated (*FAM43A*) in all three cell lines in response to mechanical loading.
For the complete lists of overlapping differentially expressed genes, please refer to
Supplementary Table S2. These results demonstrated greater transcriptomic
responses to mechanical loading in endothelial cells than in muscle cells. Large
overlaps in gene expression could be seen between the endothelial cell lines,
whereas endothelial cells and muscle cells responded differently, highlighting the cell
type-specificity.

To evaluate the external validity of the model, we next checked the expression of
common exercise-induced and mechanosensitive genes. From HSkM cells, we
specifically focused on PGC1α encoding gene (*PPARGC1A*) and its upstream
regulator PPAR-delta (*PPARD*), myostatin (*MSTN*), *NR4A3, and VEGFA.* We
observed, similarly as shown in response to exercise in vivo (Pillon *et al*, 2020; Hood
et al, 2016; Fan & Evans, 2015; Wickramasinghe *et al*, 2022; Smith *et al*, 2024;
Hulmi *et al*, 2007; Kim *et al*, 2005), upregulation of *PPARD*, *NR4A3*, and *VEGFA,*
downregulation of *MSTN*, while no change in *PPARGC1A* (Supplementary Fig. 2A).
From endothelial cells, we examined the gene expression of common
mechanosensitive genes *KLF2/4*, *THDB*, *NRP2*, *YAP1*, *HIF1A*, NRF2 encoding gene
(*NFE2L2*) (Li *et al*, 2024; Davis *et al*, 2023; Peng *et al*, 2019) as well as *FLT1*, *KDR*
and *FLT4* which are genes encoding for VEGF receptors 1/2/3. We found that all
except *HIF1A* and *FLT4* were statistically significantly upregulated in HUVECs
(Supplementary Fig. 2B-C). In HCMECs none of the changes in these genes
reached statistical significance. In addition, we inspected the expression of selected
mechanotransduction genes that have been linked to angiogenesis in endothelial
cells (Nan *et al*, 2023). These genes included the VE-cadherin-encoding gene
(*CDH5*), and eight other genes (*CTNND1*, *CTNNA1, CTNNB*1, *SRC*, *PTPRB*,
PTPRJ, *PTPRM*, and *JUP*) related to VE-cadherin. We found that the gene
expression of α- and β-catenin coding genes *CTNNA1* and *CTNNB1* significantly
increased in HUVECs. We also observed that the expression of genes coding p120
(adj. p = 0.055), VE-PTP (adj. p = 0.07), and VE-Cadherin (adj. p = 0.08) tended to
increase (Supplementary Fig. 2D). Together, these findings suggest that our *in vitro*
mechanical loading model mimics exercise-related mechanical stress.

Fig. 1 Changes in gene expression of human endothelial cells (HUVECs, HCMECs) and muscle cells (HSkM) in response to mechanical loading. **A)** Schematic representation of

experimental design. The experiment was repeated for each cell line separately. Cells from
one stretching chamber were used as one sample for RNA-Seq. **B)** PCA plots of samples in
each cell line respectively. **C)** The total number of detected genes and the number of genes
that were either up- or downregulated ($\log_2$FC ≥ 0.5 and adj. *p* ≤ 0.05) for each cell line. **D)**
Volcano plots illustrating the amounts of differentially expressed and unchanged genes in all
three cell lines. Top 10 up- and downregulated genes by magnitude and consistency are
annotated **E)** Venn diagrams showing the number of shared up- and downregulated genes
in all three cell lines

Mechanical loading resulted in cell type-specific transcriptomic responses

Gene Set Enrichment Analysis (GSEA) results revealed (Supplementary Table S3)
that the two endothelial cell lines showed highly similar pathway enrichment after
mechanical loading. 20 Gene Ontology: Biological Processes (GO:BP) and four
Reactome pathways were similarly upregulated in HUVEC and HCMEC, and 27
GO:BP, 83 Reactome and three KEGG pathways were similarly downregulated.
Overlaps in pathway enrichment between endothelial cells and muscle cells towards
the same direction were seen in the upregulation of only two GO:BP pathways:
Ovulation and *Ovarian follicle development*. These pathways included TGF-β family
genes, which are known to regulate muscle growth. Interestingly, the GSEA results
of endothelial cells showed upregulation of cell growth-related pathways such as
Endothelial cell development, RNA-splicing and *Negative regulation of catabolic*
processes (Fig. 2A). Furthermore, both endothelial cell lines showed downregulation
of pathways related to *Chromatin remodeling* and *Cellular senescence* (Fig. 2B). For
muscle cells, upregulated pathways were mostly related to oxidative phosphorylation
and ribosome biogenesis, while pathways related to cell mechanical integrity, such
as *Actin organization* and *Cell matrix adhesion*, were downregulated (Fig. 2A-B).
Many of the GSEA findings were also reproduced by the gene set
overrepresentation analysis (ORA) (Supplementary Fig. 3).

Much to our surprise, when comparing gene set enrichment of the endothelial cell
lines to muscle cells, we found multiple pathways enriched in opposite directions in
response to mechanical loading (Supplementary Table S4). In HUVECs and HSkM
cells 12 GO:BP, three Reactome and two KEGG pathways were oppositely enriched
and the same was true for HCMECs and HSkM cells in 14 GO:BP, four Reactome
and three KEGG pathways. Most of these pathways were related to cellular
respiration, metabolism, electron transport chain (ETC), as well as chromatin
remodeling (Fig. 2C). Most prominently, pathways related to oxidative
phosphorylation were downregulated in endothelial cell lines, while upregulated in
muscle cells. Closer inspection revealed that gene expression in all five
mitochondrial ETC-complexes were affected in the same fashion (Fig. 2D).

Fig. 2 GSEA results of HUVECs, HCMECs and HSkM in response to mechanical loading. **A)** Selected positively enriched gene sets from GO:BP, Reactome and KEGG database terms. **B)** Selected negatively enriched gene sets from GO:BP, Reactome and KEGG database terms. **C)** Gene sets that were systematically downregulated in both endothelial cell lines and upregulated in muscle cells. **D)** Heatmap depicting the gene expression in the oxidative phosphorylation complex genes in response to mechanical loading. NES = Normalized enrichment score. GO:BP = Gene ontology:Biological process. Rank = Rank of gene set by NES in all statistically significant (adj. $p \leq 0.05$) up- or downregulated gene sets in their respective database. SASP = Senescence-associated secretory phenotype

To further compare our results to in vivo exercise, we applied the MetaMEx database (Pillon *et al*, 2020) to examine whether the *Endothelial Cell development* pathway, which was among the most upregulated GO:BP pathways in HUVECs, is commonly induced in response to acute aerobic exercise. Indeed, we found that many of these genes were induced also in human skeletal muscle tissue in response to acute aerobic exercise (Supplementary Fig. 4).

In summary, mechanical loading induced positive enrichment of pathways related to angiogenesis and ribosome biogenesis and changes in typical exercise-regulated genes, suggesting that mechanical loading alone recapitulates several common exercise effects seen *in vivo*. The pathway enrichment results together with the gene expression overlaps of individual genes, further reinforce the cell type-specific responses induced by mechanical loading. This was most prominently seen in the opposite expression of oxidative phosphorylation pathways and ETC genes, which were downregulated in endothelial cells and upregulated in muscle cells.

Mechanical loading resulted in greater metabolite changes in HUVECs compared to HSkM cells

Next, we compared metabolite changes in HUVECs and HSkM cells in response to mechanical loading. The HCMECs were excluded because these analyses required more sample material and resources than transcriptomics. We analyzed a targeted list of intracellular metabolites with UHPLC-MS/MS, and culture media metabolites with non-targeted ^{1}H NMR spectroscopy (Fig. 3A, Supplementary Table S5). In HSkM cells and HUVECs, 3/35 and 11/38 intracellular metabolites were altered in response to mechanical loading, respectively (Fig. 3B). The intracellular metabolite alterations were thus greater in HUVECs than in HSkM cells, which is in line with the transcriptome results. From the conditioned media (i.e., extracellular) metabolites, 3/29 were altered in HSkM cells, and 2/34 in HUVECs (Fig. 3B).

By comparing conditioned media to the media incubated without cells (i.e., cell-free media), we found that both HUVECs and HSkM cells released acetate into the media in response to mechanical loading (Fig. 3C-D). We also confirmed the acetate release into the media with UHPLC-MS (Supplementary Fig. 5). The acetate release results were even more prominent with UHPLC-MS than with ^{1}H NMR (Supplementary Fig. 5A-B). In addition, arginine and glutamine levels were greater in

the media of loaded HSkM cells and pyruvate levels in the media of loaded HUVECs
(Fig. 3C-D).

In muscle cells, intracellular levels of glucose 6-phosphate and phosphocreatine
were decreased while alanine levels were increased (Fig. 3E). In HUVECs,
intracellular levels of phosphoenolpyruvate, fructose 6-phosphate, and α-
ketoglutarate were decreased while phosphoserine, serine, asparagine, threonine,
proline, alanine, acetyl-CoA, and tryptophan levels increased in response to
mechanical loading (Fig. 3F). The two most increased metabolites were
phosphoserine and serine (Fig. 3F), while the serine synthesis pathway gene
expressions were not affected (Supplementary Fig. 2E). Proline and α-ketoglutarate
were the two most consistently (smallest *p*-values) changed metabolites. The
observed increase in proline and decrease in α-ketoglutarate may indicate that
proline was synthesized using α-ketoglutarate.

Next, we conducted a metabolite overrepresentation analysis using the significantly
altered intracellular metabolites. The overrepresentation analysis of HSkM cells did
not yield any significant results because of the low number of altered metabolites. In
HUVECs, the most enriched pathways were *Glycine and serine metabolism*,
Glutamate metabolism, and *Warburg effect* (Fig. 3G). These results, together with
the decreased gene expression of the electron transport chain complexes and
increased gene expression of growth-related pathways, suggest that mechanical
stretching may reprogram endothelial cell metabolism towards the Warburg effect
and biosynthesis of serine and other amino acids.

Fig. 3 Intracellular and culture media metabolite changes of human skeletal muscle cells (HSkM) and HUVECs in response to mechanical loading. **A)** Culture media metabolites were analyzed with ^{1}H NMR spectroscopy and intracellular metabolites were analyzed using the

targeted UHPLC-MS/MS method. Cells or media from three stretching chambers were
pooled for one sample. Also, media incubated without cells (i.e., Cell-free media) was
included as an additional reference. **B)** The number of increased, decreased, and
unchanged intracellular and culture media metabolites of HSkM cells and HUVECs. **C)**
Significantly changed media metabolites of HSkM cells. **D)** Significantly changed media
metabolites of HUVECs. **E)** Significantly changed intracellular metabolites in HSkM cells. **F)**
Significantly changed intracellular metabolites in HUVECs. **G)** Metabolite overrepresentation
analysis using the significantly altered intracellular metabolites of HUVECs. In panel G, the
number of metabolites indicates how many significantly altered metabolites from each
pathway were detected. Individual values (n = 7-8 per group) are plotted, and data are
presented as mean ± SEM. The dashed lines represent the average concentrations in cell-
free media samples. P-creatine = Phosphocreatine, G6P = Glucose 6-phosphate, PEP =
Phosphoenolpyruvate, F6P = Fructose 6-phosphate
291

Mechanical loading increased the synthesis of serine, alanine, and proline from glucose

As serine metabolism was the most highlighted pathway in mechanically loaded
HUVECs, we next conducted dynamic labeling experiments with ^{13}C-(U)-glucose
(Fig. 4A) to assess if the serine synthesis rate was affected. We found that there
were significantly more total (unlabeled + labeled), and newly synthesized serine and
phosphoserine at multiple time points after mechanical loading (Fig. 4B-E). This
indicates that the serine synthesis was, indeed, increased in response to mechanical
loading in HUVECs. There were no significant changes in the total serine level in the
media (Fig. 4F), whereas mechanically loaded HUVECs released significantly more
labeled serine into the culture media (Fig. 4G), supporting the interpretation that
serine synthesis was increased by mechanical loading.

We also assessed if the synthesis of alanine, proline, glycine, aspartate, asparagine,
glutamine, glutamate, and arginine were changed. We found that there were also
more labeled and total alanine and proline in loaded HUVECs and their media
(Supplementary Fig. 6A-G), indicating that alanine and proline synthesis were
increased together with serine synthesis in response to mechanical loading.

There was significantly less labeled glycine in loaded HUVECs at 3 h and 20 h time
points and total glycine at the 0 h time point (Supplementary Fig. 6H-I), but no
differences in labeled or total glycine release into media (Supplementary Fig. 6J-K)
compared to non-loaded controls. We did not find significant differences in labeled
aspartate and glutamate (Supplementary Fig. 7). In the case of asparagine, arginine,
and glutamine, ^{13}C-labeled molecules were not detected after isotope natural
abundance correction, suggesting that HUVECs did not use glucose to produce
these amino acids in these nutritional conditions (Supplementary Table S6).

Together, these results suggest that mechanical loading increases the synthesis of
serine, alanine, and proline in endothelial cells.

319
Fig. 4 Mechanical loading increased glucose-derived carbon incorporation into serine and
phosphoserine in HUVECs. **A)** A schematic diagram of the time points of the ^{13}C-(U)-glucose
experiment. **B)** Total intracellular serine level (unlabeled + labeled). **C)** Newly synthesized

(M+3) serine level. **D)** Total intracellular phosphoserine (P-serine) level (unlabeled +
labeled). **E)** Newly synthesized (M+3) phosphoserine level. **F)** Total media (extracellular)
serine level (unlabeled + labeled). **G)** Newly synthesized (M+3) media serine level. Sample
sizes were n = 3 per group per time point in intracellular analyses and n = 5 in media
analyses. Individual values are plotted, and data are presented as mean ± SEM. *p < 0.05,
**p < 0.01, ***p < 0.001*
329

Serine synthesis pathway enzyme phosphoglycerate dehydrogenase (PHGDH) is critical for endothelial cell anabolism

To assess the role of the serine synthesis pathway in HUVECs, we first inhibited the
rate-limiting serine synthesis pathway enzyme PHGDH (Figure 5A) and evaluated its
effects on proliferation. For this, we used two structurally distinct inhibitors, NCT-503
and CBR-5884. We discovered that both PHGDH inhibitors decreased the
proliferation of HUVECs (Fig. 5B) but did not induce cell death (Figure 5C). We also
tested whether the PHGDH inhibitors affect protein synthesis and glucose usage for
cellular biomass production. We found that both PHGDH inhibitors decreased protein
synthesis and glucose-derived carbon incorporation into proteins, RNA, and lipids
(Fig. 5D-F). Next, we used a lentiviral vector to overexpress PHGDH in HUVECs
(Fig. 5G). We found that the overexpression of PHGDH led to a ~10% increase in
protein synthesis ($p < 0.01$, Fig. 5H) and a trend for increased proliferation ($p =$
0.088, Fig. 5I). Finally, we tested whether mechanical loading affects protein
synthesis and proliferation of HUVECs. We found that mechanical loading did not
affect the total protein synthesis and proliferation of HUVECs (Supplementary Fig.
8). Overall, these data suggest that the PHGDH has an important role in endothelial
cell growth.

Figure 5. Effects of the serine synthesis pathway enzyme phosphoglycerate dehydrogenase (PHGDH) manipulation on cell proliferation, protein synthesis, and glucose usage for biomolecule synthesis in HUVECs. **A)** Schematic presentation of the serine synthesis

pathway. **B)** PHGDH inhibitors decreased the proliferation of HUVECs, as assessed by EdU
analysis. Each group was statistically compared to the DMSO vehicle control (n = 8 in the
DMSO control group and n = 4 in each inhibitor group). **C)** PHGDH inhibitors did not induce
cell death according to the lactate dehydrogenase (LDH) activity measured from conditioned
media as a marker of membrane damage and cell death (i.e., increased LDH activity would
indicate membrane damage and cell death) (n = 3). **D)** protein synthesis (^{14}C-valine
incorporation into proteins) in response to PHGDH inhibitors (n = 6). **E)** Glucose-derived
carbon incorporation into proteins, RNA, and lipids in response to NCT-503 (n = 9). **F)**
Glucose-derived carbon incorporation into proteins, RNA, and lipids in response to CBR-
5884 (n = 6). **G)** PHGDH level in response to overexpression treatment (n = 16). **H)** Protein
synthesis (^{14}C-phenylalanine incorporation into proteins) in response to PHGDH
overexpression treatment (n = 16, but in no treatment group n = 4). **I)** Cell proliferation in
response to PHGDH overexpression treatment (n = 24, but in no treatment group n = 6).
Individual values are plotted, and data are presented as mean ± SEM. *$p < 0.05$, **$p < 0.01$,
***$p < 0.001$, **** $p < 0.0001$

369
370

Discussion

In this study, we unraveled the effects of mechanical loading on skeletal muscle and
endothelial cells and identified both shared and cell type-specific responses. Notably,
both muscle and endothelial cells released acetate into the media and induced gene
expression changes typically seen after exercise. However, transcriptomic
responses exhibited several contrasting patterns, e.g. electron transport chain (ETC)
complex genes were upregulated in muscle cells but downregulated in endothelial
cells. Overall, endothelial cells demonstrated more pronounced responses to
mechanical loading, particularly in amino acid metabolism as well as transcriptomic
pathways related to angiogenesis and growth. Furthermore, using a ^{13}C-(U)-glucose
tracer, we observed that mechanical loading enhanced glucose-derived serine
synthesis in endothelial cells. Overexpression of the serine synthesis enzyme
PHGDH promoted protein synthesis of endothelial cells, whereas PHGDH inhibitors
decreased proliferation and anabolic activity, highlighting the possible role of serine
synthesis in endothelial cell growth and angiogenesis.

Both muscle and endothelial cells released acetate into the media in response to
mechanical loading, which was observed with two different analysis methods.
Previously, we found that electrically evoked myotube contractions also induced
acetate release from murine myotubes (Lautaoja *et al*, 2021). In humans, it has been
found that acetate is released into the bloodstream (van Hall *et al*, 2002) and urine
(Schranner *et al*, 2020) after exercise. Moreover, serum acetate concentrations have
been found to be greater in physically active individuals compared to inactive
individuals (Kujala *et al*, 2013). Therefore, acetate could be a possible exerkine (i.e.
a signaling molecule released in response to exercise) released by not only muscle
cells but also endothelial cells. One of the most relevant functions of acetate is acting
as an immunomodulator and promoter of T-cell effector function and regulating
differentiation of CD4$^+$ T-cell helper cells (Jing *et al*, 2023; Miller *et al*, 2023; Qiu *et
al*, 2019). This may partly explain the effects of exercise in enhancing immune
system function and increasing CD4$^+$ T-helper cell count in humans (Chastin *et al*,

2021). This is particularly interesting, as factors secreted by CD4[+] T-cells seem to be
important in angiogenesis, myogenesis, and muscle regeneration (Kwee *et al*, 2018;
Schiaffino *et al*, 2017). In addition, the potential source of acetate after mechanical
loading is interesting as acetate can be produced from (A) acetyl-CoA by certain
acyl-CoA thioesterases (Horibata *et al*, 2013), (B) from pyruvate by ketoacid
dehydrogenases and reactive oxygen species (Liu *et al*, 2018), and (C) by
deacetylation of proteins (Bose *et al*, 2019). Of these, at least reactive oxygen
species production is induced in muscle by exercise (Powers *et al*, 2024). Thus,
acetate's possible role as an exerkine and its links to the immune system or other
human physiological systems should be studied further.

Endothelial cells and muscle cells also demonstrated different, and even opposite
transcriptomic responses to mechanical loading, most prominently in oxidative
phosphorylation-related pathways and ETC complex genes. Oxidative
phosphorylation-related genes were upregulated in muscle cells, and skeletal muscle
is known to upregulate mitochondrial biogenesis and oxidative phosphorylation in
response to physiological stress, such as exercise (Smith *et al*, 2023). One of the
well-known mechanisms regulating these exercise adaptations is the AMPK-PGC1α
pathway (Hood *et al*, 2016). In our study, although PGC1α gene expression was
unchanged in muscle cells, we did observe an increase in *PPARD* expression after
mechanical loading. PPARD is upstream of PGC1α (Fan & Evans, 2015) and has
also been linked to enhanced oxidative metabolism in skeletal muscle
(Wickramasinghe *et al*, 2022). Furthermore, muscle cells also demonstrated an
upregulation of the *NR4A3* gene, which has recently been shown to be one of the
most exercise-responsive genes (Pillon *et al*, 2020), and essential in glucose
metabolism and anabolism, such as ribosome biogenesis in primary muscle cells
(Smith *et al*, 2024). In line with the upregulation of *NR4A3*, we also observed
increases in the *ribosome biogenesis* GO:BP pathway after mechanical loading.
Moreover, we saw a decrease in myostatin (*MSTN*) expression after mechanical
loading in muscle cells (Supplementary Table S1), which is a common response
after a single bout of resistance exercise (Hulmi *et al*, 2007; Kim *et al*, 2005). These
results demonstrate that mechanical loading of human skeletal muscle cells
recapitulates several of the common effects of exercise seen *in vivo*.

In endothelial cells, transcriptomic results showed that mechanical loading induced
upregulation of genes related to endothelial cell development, RNA splicing, and
negative regulation of catabolic processes, and moreover, downregulation of cellular
senescence-related genes. Many of the endothelial cell development genes were
also found to increase in response to acute aerobic exercise in human skeletal
muscle tissue, which is known to be highly vascularized (Englund *et al*, 2021; Pepe
& Albrecht, 2023; Pillon *et al*, 2020). Furthermore, our results showed upregulation of
mechanosensitive genes commonly associated with an atheroprotective endothelial
phenotype, which are typically induced by flow-mediated mechanical stress (Li *et al*,
2024; Davis *et al*, 2023; Peng *et al*, 2019). Our findings point out that a similar effect
was seen after cyclic stretching, which is comparable to skeletal muscle movement.
We also found that genes related to VEGF response, including VEGF receptors 1 &
2 (*FLT1 & KDR*), were upregulated in endothelial cells. Related to these, *VEGFA*

expression was also increased in mechanically loaded muscle cells, and it has been
hypothesized that skeletal muscle stem cells recruit ECs via VEGFA expression and
this interaction is important for muscle stem cell self-renewal (Verma *et al*, 2018). In
addition, similar cyclic stretch as used in this study, has previously been shown to
enhance endothelial cell proliferation (Li & Sumpio, 2005) and angiogenesis (Yung *et*
al, 2009). Mechanical loading also induced upregulation of insulin receptor signaling
pathways in endothelial cells, and more specifically, these pathways included genes
such as *FGF5, PDPK1,* and *KRAS* which were upregulated in both endothelial cell
lines. *FGF5* has previously been shown to promote angiogenesis (Seo *et al*, 2016),
and *PDPK1* and *KRAS* are linked to the PI3K/Akt signaling pathway (Manning &
Toker, 2017), which is known to regulate angiogenesis and endothelial cell growth
(Kobialka & Graupera, 2019). Since exercise-induced angiogenesis is a well-known
and documented physiological phenomenon in humans (Ross *et al*, 2023), our
results suggest that, similarly to muscle cells, mechanical loading of endothelial cells
also recapitulates the effects of exercise *in vivo*.

We found intriguing that the oxidative phosphorylation pathways and ETC complex
genes were coordinately downregulated in endothelial cells, which was completely
opposite to the response in muscle cells. It has previously been shown that
endothelial cells primarily rely on glycolysis for ATP production despite having
access to oxygen, and that vessel sprouting of endothelial cells *in vitro* is regulated
by glycolysis (De Bock *et al*, 2013). It is also known that proliferating cells similarly
rely on glycolysis to ensure adequate availability of metabolic intermediates for cell
growth, which is described as the Warburg effect (Lunt & Vander Heiden, 2011). In
the Warburg effect, glycolytic intermediates are used to produce e.g., nucleotides,
serine, alanine, glycine, and cysteine needed for growth. Thus, the transcriptomic
results collectively suggest a possible link between a metabolic shift towards
glycolytic ATP production and increased growth in mechanically loaded endothelial
cells.

In line with the transcriptomic results, the metabolite enrichment analysis showed
that the Warburg effect was the third most enriched pathway in endothelial cells.
Moreover, ^{13}C-glucose tracing analysis revealed that the abundance and synthesis
of growth-related amino acids: serine, phosphoserine, and alanine increased after
mechanical loading at multiple time points in endothelial cells. In the serine synthesis
pathway, PHGDH is an essential enzyme for serine production and anabolism in
many different cell types (Handzlik & Metallo, 2023; Mäntyselkä *et al*, 2024; Stegen
et al, 2022). Our further experiments revealed that overexpression of PHGDH in
endothelial cells increased protein synthesis, while commonly used non-competitive
PHGDH inhibitors decreased proliferation, protein synthesis, and glucose-derived
carbon incorporation into proteins, RNA, and lipids. This is similar to what we
recently demonstrated in muscle cells (Mäntyselkä *et al*, 2024). Loss of PHGDH has
previously been shown to cause lethal defects in angiogenesis in mice, and
knockdown of PHGDH impairs EC proliferation (Vandekeere *et al*, 2018). Moreover,
PHGDH knockdown impairs heme synthesis while PHGDH overexpression
increases endothelial spheroid sprouting in cultured endothelial cells (Vandekeere *et*
al, 2018). Interestingly, in this current study, we also observed upregulation of heme

signaling- and sprouting angiogenesis-related genes in response to mechanical
loading. These results indicate that mechanical stimuli and PHGDH overexpression
similarly affect endothelial cells to support their function, possibly via increased
serine production. The serine synthesis pathway has also been identified as a major
upregulated metabolic pathway in tumor endothelial cells, and the loss of PHGDH
reduced tumor vasculature (Zhang *et al*, 2023). Therefore, our results and the
existing evidence highlight that the serine synthesis pathway, especially PHGDH, is
crucial for endothelial cell function and anabolism. Further studies are needed to
investigate the importance of the serine synthesis pathway and PHGDH in
endothelial cells after exercise or exercise-mimicking stimuli.

The mechanism of how mechanical loading activates serine synthesis in endothelial
cells was not studied here. However, the mechanical loading did not increase the
serine synthesis pathway gene expression, indicating that the increased serine
synthesis is regulated by other mechanisms. Recently, it was found that epithelial
cadherin (E-cadherin) induces serine synthesis in cancer cells (Lee *et al*, 2024) and
others also found that shear stress (another form of mechanical loading) increases
glucose uptake in endothelial cells through the activation of vascular endothelial
cadherin (VE-cadherin) (Cronin *et al*, 2024). VE-cadherin and proteins linked to it are
important for mechanotransduction in endothelial cells (Kong *et al*, 2022; Nan *et al*,
2023). Here, we found increased expression of genes encoding proteins linked with
VE-cadherin. In further studies, the possible mechanism for the activation of serine
synthesis through VE-cadherin, or other upstream regulators, should be examined.

Strengths and limitations

We used two different human endothelial cell types and primary human skeletal
muscle cells. The loading induced several common exercise-induced and
mechanosensitive genes, suggesting an external validity of the model. Moreover,
several of the transcriptomic responses to mechanical loading were similar between
the two endothelial cell types (HUVEC and HCMEC), suggesting that many of the
changes can be applied to endothelial cells in general. We also replicated the main
metabolite results with different methods. For example, acetate release was
observed with both ^{1}H NMR spectroscopy and UHPLC-MS. In addition, the
increased serine, alanine, and proline content and synthesis were detected by
targeted metabolomics and isotopic tracing from multiple time points. We also
conducted mechanistic experiments to focus on the main findings regarding
increased serine and growth-directed gene expression revealed by the -*omics*
analyses.

The study also has limitations. We conducted the experiments with monolayers of
cells (i.e., two-dimensionally), and therefore, myotubes were not aligned
unidirectionally while the mechanical loading was unidirectional. In contrast, smaller
endothelial cells could change their alignment against the direction of stretch. This
difference in the ability to change alignment may in part explain why we detected
fewer molecular changes in myotubes compared to endothelial cells. Overall, the
data suggest that the mechanical loading model for two-dimensionally cultured cells
is suitable for endothelial cells, which grow in monolayers, also *in vivo*, while

myotubes may need to be loaded three-dimensionally to sufficiently mimic exercise
in future studies.

Conclusion

Mechanical loading induces common and cell type-specific transcriptomic and
metabolomic responses in endothelial and skeletal muscle cells. In general, several
of the effects recapitulated what is observed *in vivo* after exercise. Both endothelial
and muscle cells released acetate in response to mechanical loading, and therefore
acetate's potential role as an exerkine should be studied in the future. The
responses to mechanical loading were larger in endothelial cells, revealing increased
serine synthesis as one of the main metabolic changes. These results, together with
mechanistic experiments, highlight that mechanical stimulus, serine synthesis, and
PHGDH are interconnected and may be important for endothelial cell growth and
development. The potential importance of the mechanical loading and serine
synthesis pathway for improved vascularization should be studied *in vivo* in further
studies.

Materials and methods

Endothelial and muscle cells

We used human umbilical vein endothelial cells (HUVEC, Cat# C-12200,
PromoCell), human cardiac microvascular endothelial cells (HCMEC, Cat# C-12285,
PromoCell), and primary human skeletal muscle (HSkM) cells. The latter were
obtained as previously (Mäntyselkä *et al*, 2024) from two healthy 25-34-year-old
volunteers (1 male and 1 female). Vastus lateralis muscle biopsies were obtained
under local anesthesia with the Bergström biopsy needle. The human study
(Halonen *et al*, 2024) was conducted in accordance with the Declaration of Helsinki
and was approved as part of a larger study by the Ethics Committee of the University
of Jyväskylä (857/13.00.04.00/2021). All participants gave their written informed
consent for inclusion before they participated in the study.

Cell culture

The HUVEC and HCMEC cells were cultured in endothelial cell growth medium
(EGM) (Cat# C-22010, PromoCell) supplemented with EGM SupplementMix (Cat#
C-39215, PromoCell) and 1% (vol/vol) of 1000 U/ml penicillin-streptomycin (P/S,
Cat# 11548876, Gibco). HSkM cells were cultured in growth medium (GM)
consisting of high glucose (4.5 g/l) Dulbecco's Modified Eagle Medium (DMEM, Cat#
10313021, Gibco) supplemented with 20% (vol/vol) fetal bovine serum (FBS, Cat#
10270106, Gibco), 1% (vol/vol) of 1000 U/ml P/S and 1% (vol/vol) 200 mM L-
glutamine (Cat# 11500626, Gibco).

After reaching over 80% confluency in culture flasks, the cells were seeded (~20,000
cells/cm^2) on polydimethylsiloxane (PDMS) chambers (Cat# STB-CH-04, STREX).
PDMS chambers were pre-coated with FBS for 24 h and with 0.2% gelatin for an

additional 24 h. The PDMS chambers were placed in 150 mm diameter Petri dishes,
two chambers per dish. Cells were grown on PDMS chambers until reaching over
80% confluency. After that, HUVEC and HCMEC cells were mechanically loaded, as
described in the next section. In the case of HSkM cells, at 80% confluency the GM
was replaced with differentiation medium (DM) containing high glucose DMEM, 2%
(vol/vol) horse serum (HS, Cat# 26050088, Thermo Fisher Scientific), 1% P/S, 1% L-
glutamine and 1% (vol/vol) Insulin-Transferrin-Selenium-X supplement (ITS-X, Cat#
51500056, Gibco). To induce myoblast fusion into myotubes, the cells were cultured
in DM for 4-5 days and DM was changed every 48 h. Cells were cultured at 37 ℃ in
a 5% CO_2 atmosphere all the time.

Mechanical loading set-up

Mechanical loading of cultured cells was achieved with a commercial cell culture
pacing system (C-Pace EM 100, IonOptix) paired with a motor-powered mechanical
stimulation unit (C-Stretch 100, IonOptix). The C-Stretch mechanical stimulation unit
consists of a motor-driven tray that causes uniaxial elongation of the PDMS
chambers inserted in the device and exposes cells, that are attached to the bottom
of the chamber, to mechanical load.

As mentioned, cells were seeded onto 12 PDMS chambers for each stretch
experiment, with 6 chambers loaded into the C-Stretch unit in series, and 6
chambers left as static controls. The stretch unit was thoroughly cleaned with 70%
EtOH before use. Fresh media was changed for all chambers prior to starting the
experiment. The stretch unit was then placed in an incubator with conditions of 37 ℃
and 5% CO_2 atmosphere along with the 6 control chambers. A cyclic mechanical
stretch with a frequency of 1 Hz and 12% elongation was applied for 5 h. For each
one-second cycle, a trapezoidal waveform of 0.4 s stretch, 0.1 s hold, 0.4 s return,
and 0.1 s hold, was used to keep the velocity of the deformation of the silicone
chamber at a moderate speed.

Lactate dehydrogenase activity assay

602 Lactate dehydrogenase activity (LDH) was measured from conditioned media as an
603 indicator of possible membrane damage and cell death (Legrand *et al*, 1992). The
604 analysis was performed as previously (Mäntyselkä *et al*, 2024) using an LDH Activity
605 Assay Kit (Cat# 981906, Thermo Fisher Scientific).

RNA extraction and analyses

RNA isolation, library preparation and sequencing

RNA was extracted using a commercial column-based RNA isolation kit
(NucleoSpin® RNA, Cat# 740955.250, Macherey-Nagel) immediately after the 5-
hour mechanical loading. The PDMS chambers were washed with PBS, and cells
were lysed by pipetting back and forth vigorously with lysis solution (350 µl of
guanidium thiocyanate lysis buffer + 3.5 µl of β-mercaptoethanol). Extraction was
then continued according to the kit's instructions. An additional centrifugation step of
30 s at 11 000 g was included at the end to ensure complete removal of the wash
buffer. Bound RNA was then eluted into a volume of 60 µl of RNase-free water. The

collected elution was then aliquoted and samples were immediately transferred to -
80 °C. RNA quality control, library preparation and sample sequencing were
performed at Novogene (Cambridge, United Kingdom), utilizing an Illumina NovaSeq
6000 platform, producing 150bp paired-end reads. Filtering of raw sequence data
was also conducted by Novogene in the following way: (1) removing reads
containing adapters, (2) removing reads containing >10% of unsolvable bases and
(3) removing reads containing >50% of low quality bases (Qphred ≤ 5). Filtered RNA
sequencing data is available in the Gene Expression Omnibus (GEO) data repository
(accession number: GSE288064).

Read alignment, normalization and filtering

Reads were aligned to the human genome (build GRCh38.95) using the Bowtie2
software (Langmead & Salzberg, 2012), with alignment rates >80% for all the
samples. The aligned and mapped reads were later annotated and quantified using
the Rsubread package in R/Bioconductor software (Liao *et al*, 2014), with the human
GRCh38.110 genome build as annotation reference. The obtained gene counts were
transformed to counts-per-million (CPM) using the edgeR package in the
R/Bioconductor software (Robinson *et al*, 2010). Genes with a CPM < 1 in three
samples or more in each group (50%), were subsequently filtered out as described in
(Chen *et al*, 2016). The resulting expression data were finally normalized using the
trimmed mean of M-values (TMM) algorithm (Robinson & Oshlack, 2010).

Differential gene expression and pathway analyses

Differential gene expression (DGE) analyses were implemented through the
R/Bioconductor limma-voom pipeline (Ritchie *et al*, 2015). The voom-function of the
limma package was initially executed to format the data appropriately (Law *et al*,
2014), after which empirical Bayesian statistics were computed to rank genes based
on their evidence for differential expression. To account for type I errors induced by
multiple testing, the Benjamini-Hochberg method was used to adjust the tests' *p*-
values, retrieving an adjusted *p*-value (adj. *p*). Genes with a $\log_2$FC ≥ 0.5 and adj. *p*
≤ 0.05 were regarded as differentially expressed. For subsequent Gene Set
Enrichment Analysis (GSEA) genes were ranked in an unbiased way by multiplying
the $\log_2$FC by the -log10 of the adj. *p* to consider both the magnitude and
consistency of the change in gene expression. GSEA was performed with the
clusterProfiler R-package (Yu *et al*, 2012) with Gene Ontology: Biological Process
(GO:BP), Reactome, and the Kyoto Encyclopedia of Genes and Genomes (KEGG)
serving as reference databases. Ontologies and pathways were considered
significantly enriched if their corresponding tests yielded an adj. p-value <0.05. In
addition, gene set overrepresentation analysis (ORA) of the significantly up- or
downregulated genes ($\log_2$FC ≥ 0.5 & adj. *p* ≤ 0.05) was conducted using the
ShinyGO web-application v0.81 (https://bioinformatics.sdstate.edu/go/) (Ge *et al*,
2020) from GO:BP pathways. Pathway minimum size was set to 2 and maximum to
5000 and the FDR cutoff for significant pathway enrichment was set to 0.05. All
analyses implemented in R/Bioconductor included the core software version 4.3.1
with the R-studio interface version 2024.04.2.

Meta-analysis of skeletal muscle response to exercise (MetaMEx)

We examined gene expression of the *Endothelial cell development* pathway (GO:0001885, 72 genes, list checked on 10.06.2025) in response to acute aerobic exercise using the MetaMEx browser-based tool (https://www.metamex.eu/app/metamex) (Pillon *et al*, 2020). All the other settings were kept as a default, except that only the healthy population was included.

Metabolite extraction and analyses

^{1}H NMR spectroscopy metabolomics

Metabolites from the media samples were extracted as previously (Mäntyselkä *et al*, 2024; Lautaoja *et al*, 2021). Cell-free medium samples were also included as a reference to assess the uptake and release of metabolites. Dried samples were reconstituted with 200 µL of sodium phosphate buffer (150 mM, pH = 7.4) in 99.8% deuterium oxide (Acros Organic; Thermo Fisher Scientific) containing 0.5 mM 3-(trimethylsilyl)propanesulfonic-d_6 acid sodium salt (DSS-d_6; IS-2 Internal Standard, Chenomx Inc.) and transferred to round-bottom 3-mm OD NMR tubes (Cat# 341-PP-7, Wilmad). Data collection was carried out at 25 °C using the Bruker AVANCE III HD NMR spectrometer, operating at 800 MHz ^{1}H frequency equipped with a cryogenically cooled ^{1}H, ^{13}C, ^{15}N triple-resonance probe head (Bruker Corporation). The ^{1}H nuclear magnetic resonance (^{1}H NMR) spectra were collected as previously (Lautaoja *et al*, 2021). ^{1}H NMR spectra were processed, and metabolites were identified and quantified with the Chenomx 10.0 software (Chenomx).

UHPLC-MS/MS targeted metabolomics

Intracellular metabolites were collected and extracted as previously (Mäntyselkä *et al*, 2024) with minor modifications. In brief, 200 µL of ice-cold 80% methanol was added to each stretching chamber, and metabolites from three chambers were pooled for one sample. Dried samples were reconstituted with 40 µl of 50-50 acetonitrile-water mixture (vol/vol) before analysis. Metabolites were analyzed as previously (Mäntyselkä *et al*, 2024). Briefly, metabolites were separated with an Atlantis Premier BEH Z-HILIC UHPLC column (1.7 µm, 2.1 x 100 mm) which was used with an Agilent 1290 Infinity ultra-high-performance liquid chromatography (UHPLC) system. *Mobile phase A* was 15 mM ammonium bicarbonate (pH = 9.0) in water, and *mobile phase B* was 15 mM ammonium bicarbonate (pH = 9.0) in 90% acetonitrile-10% water. The flow rate was 0.7 mL/min, the column oven temperature was 40 °C, and the injection volume was 3 µl. The following gradient elution was used: the initial concentration of 90% *B* was held for 1 min and then was linearly decreased to 65% *B* over the next 5 min and was held for 1 min at 65% *B*. Then *B*% was increased to 90% over the next 1 min and was held 2 min at 90% *B* before next injection. The UHPLC system was coupled with an Agilent 6460 triple quadrupole mass spectrometer operated with dynamic multiple reaction monitoring mode (dMRM) as previously described (Mäntyselkä *et al*, 2024). A mixture of target metabolites (concentrations between 10 and 50 µM) was injected eight times at the

beginning of the analysis and after every 6-8 samples for quality control and
metabolite identification purposes.

Acetate was analyzed from media samples using the above-described
chromatographic method and the UHPLC-MS instrument which was operated on
negative polarization. For acetate, selected ion monitoring (e.g., pseudo-MRM
(Qasem *et al*, 2017)) of *m/z* 59 was done. The following parameters were used for
the selected ion monitoring experiment: fragmentor: 80; collision energy: 1; cell
accelerator voltage: 4. The other ion source parameters were the same as previously
reported (Mäntyselkä *et al*, 2024). Retention time and linearity of the signal were
tested with 10, 20, 40, 80, and 160 µM acetate standards that were prepared using
sodium acetate (Cat# 106268, Merck). The acetate standard was also used for a
spike-in experiment.

Batch correction, normalization, and statistical analyses of metabolomics data

713
714

Targeted UHPLC-MS/MS data and [1]H NMR data were measured within two batches
and therefore they were batch effect corrected using the EigenMS method on the
MetaboAnalyst 6.0 platform (https://www.metaboanalyst.ca/). In the case of UHPLC-
MS/MS, data collected with negative and positive modes were separately batch-
corrected before combining the data. Single missing values within different groups
were replaced with 1/5 of the minimum detected values. Metabolite
concentrations/abundances were $\log_2$ transformed for statistical testing with the
Student's t-test with significance set at $p < 0.05$ and fold change > I1.1I.

Metabolite overrepresentation analysis

724 MetaboAnalyst 6.0 platform was used for the metabolite overrepresentation analysis.
725 Names of the significantly altered metabolites were uploaded. Default settings were
726 used (SMPDB pathway and '*Only use metabolites sets containing at least 2 entries*'
727 option were used).

[13]C-glucose labeling experiments and analyses

Glucose-free and phenol red-free endothelial cell medium (Cat# H1168GPF, Cell
Biologics) medium including medium kit's supplements was supplemented with 5.5
mM U-$^{13}C_6$-D-glucose (Cat# CLM-1396-1, Cambridge Isotope laboratories, isotopic
purity 99%). This medium was changed to the cells just before the initiation of the
experiment. Intracellular metabolites were extracted with the same protocol as for
targeted UHPLC-MS/MS metabolomics, but this time two chambers were pooled
instead of three. Media metabolites were extracted with the same protocol as for [1]H
NMR metabolomics described above except now 100 µl of media from each
chamber was mixed with 200 µl of methanol. Intracellular and medium metabolites
were reconstituted into 45 µl and 400 µl of 50-50 acetonitrile-water solution,
respectively. Pooled quality control (QC) samples were prepared by transferring a
small aliquot of each sample into a vial (separate QC samples for intracellular and
medium metabolites). Selected amino acids were analyzed with the same method as
in targeted UHPLC-MS/MS metabolomics described above but the MRM transitions
were modified (Supplementary Table S7) allowing analysis of the isotopomers of the

selected amino acids. The data was processed and integrated with
MRMPROBS_v.3.66 software (Tsugawa *et al*, 2014). Phosphoserine and glycine
were processed and integrated with Agilent Qualitative Navigator B.08.00 software
because of their lower abundance. Isotope natural abundances were corrected with
IsoCor 2.2.2 software (Millard *et al*, 2019) as earlier reported (Yoon *et al*, 2023).

Chemical inhibitors

For chemical inhibitor experiments, two different PHGDH inhibitors NCT-503, (Cat#
HY-101966) and CBR-5884 (Cat# HY-100012) were obtained from
MedChemExpress. Inhibitor stock solutions were prepared in DMSO, and the DMSO
was also used as vehicle control. The inhibitors were used in concentrations of 5, 25,
and 50 μM as previously have been used for different cell types (Mäntyselkä *et al*,
2024; Zhang *et al*, 2018; Białopiotrowicz *et al*, 2020; Zhang *et al*, 2022). The exact
inhibitor concentration for each experiment is stated in the legends or caption of Fig.
5. Further details of inhibitor experiments are described in the next two sections.

Cell proliferation analysis with EdU labeling

Effects of chemical inhibitors (see the section above) on cell division capacity were
evaluated using a Click-iT EdU imaging kit (Cat# C10338, Invitrogen). HUVEC cells
were plated onto a 96-well plate in Endothelial cell GM and let to grow in
subconfluent culture for 24 h. Exposures were added to the wells, four parallel wells
for each condition, and eight parallel for the control condition (DMSO vehicle). After
20 h incubation of exposures, 5-ethynyl-20-deoxyuridine (EdU) was added for 4 h.
Cells were fixed with 4% PFA for 15 min, washed with 3% bovine serum albumin
(BSA) in PBS, and permeabilized with 0.5% Triton X-100 in PBS for 20 min RT, and
the Click reaction was done according to the manufacturer's instructions. Before
imaging, DAPI (0.5 μg/ml) was added to visualize all nuclei. Cells were imaged using
a Nikon A1R confocal microscope. The microscope was automated to find optimal
focus in each well, image every well, and take 2-9 views per well so that at least 300
nuclei were imaged. Detected nuclei (all nuclei + EdU stained nuclei) were analyzed
using an in-house built Fiji script (https://github.com/seiryoku-
zenyo/simple_ratio_DAPIvsPositive-nuclei/tree/main). Shortly, image data was
segmented using a deep learning algorithm, separately for DAPI-positive nuclei and
EdU-positive nuclei. Next, the ratio between DAPI and EdU nuclei was counted. The
data is presented as a fraction (%) of EdU-positive nuclei out of total (DAPI) nuclei.

Click-iT EdU imaging kit (Cat# C10339, Invitrogen) was also used to determine the
effect of PHGDH overexpression on HUVEC proliferation. Cells were cultured on 13
mm diameter coverslips (10 000 cell/cm^2) coated with 0.2% gelatin for 24 h after
which cells were incubated with EdU labeling solution for 5 h. Cells were fixed,
washed, and permeabilized as above. Click reaction buffer was prepared as
described by the manufacturer, added to the cells, and incubated for 30 min
protected from light. DAPI (0.5 μg/ml) was used to detect all the nuclei. Cells were
imaged with a confocal microscope (Zeiss LSM700) and the number of stained
nuclei was calculated with Fiji manually. The data is presented as a fraction (%) of
EdU-positive nuclei out of total (DAPI) nuclei.

To assess the effects of mechanical loading on HUVEC proliferation, cells were
mechanically loaded as described earlier, detached, and plated on 13 mm diameter
coverslips as above. The EdU analysis was performed as above.

Carbon-14 labeling

The following radioisotope tracers were used: ^{14}C-[U]-glucose (0.2 mCi/mL, Cat#
NEC042V250UC, PerkinElmer); ^{14}C-[U]-L-valine (271.0 µCi/mmol, Cat#
NEC291EU050UC, PerkinElmer); and ^{14}C-[1]-L-phenylalanine (0.1 mCi/mL, Cat#
NEC859S050UC, PerkinElmer). Only one tracer per experiment was used, and the
dilution of the tracers was 1 µL/mL (vol/vol) in the culture media. Two milliliters of
culture media were used per well of a 6-well plate, and one milliliter per stretching
chamber

HUVECs for PHGDH inhibitor experiments were seeded onto 0.2% gelatin-coated 6-
well plates at a density of 12,500 cells/cm². The cell population was expanded for 2
days, and the inhibition experiments were initiated by changing the media to
experimental media containing 25 µM of inhibitor and a radioisotope tracer. After
incubating cells for 48 h with inhibitor and tracer, proteins, RNA, and lipids were
extracted, and liquid scintillation counting was conducted as previously (Mäntyselkä
et al, 2024). When protein synthesis was measured during and after mechanical
loading, tracer was added to the medium just before the initiation of the 5 h
mechanical loading. After the loading, HUVECs were incubated for an additional 20
h in the medium, and then proteins were extracted.

PHGDH overexpression

The PHGDH overexpression construct pLJM5-WT PHGDH was a gift from David
Sabatini (Addgene plasmid #83901; http://n2t.net/addgene:83901; RRID:
Addgene_83901) (Pacold *et al*, 2016). Lentivirus particles were prepared and
distributed by Biomedicum Virus Core facility supported by HiLIFE and the Faculty of
Medicine, University of Helsinki, and Biocenter Finland. The control construct, clone
SHC002 (Scrambled shRNA in pLKO.1) was from Sigma Aldrich MISSION shRNA
library, distributed by Genome Biology Unit core facility supported by HiLIFE and the
Faculty of Medicine, University of Helsinki, and Biocenter Finland. The Biomedicum
Virus Core facility also determined the p24 capsid protein concentration with an
enzyme-linked immunosorbent assay (ELISA). p24 concentrations were used to
calculate the number of virus particles needed for transduction (1 pg of p24 = 100
transducing units).

HUVECs were cultured in T75 flasks. After reaching 50% confluence, 10 ml of
medium containing lentiviruses (MOI = 50) and polybrene (5 µg/ml), was added to
the cells. Polybrene (Cat# TR-1003, Sigma-Aldrich) was used to improve infection
efficiency. HUVECs were incubated with lentiviruses for 24 hours. After transduction,
the cells were resuspended in a fresh endothelial cell medium, cultured overnight,
and used for further experiments. HUVECs expressing scrambled shRNA and
HUVECs without virus treatment were used as controls for the PHGDH
overexpression construct.

Protein extraction and Western Blotting

HUVECs expressing either the PHGDH overexpression system or shControl gene were cultured in 6-well plates (5000 cells/cm^2) in endothelial cell medium supplemented with 1 µl/ml (vol/vol) ^{14}C-[1]-L-phenylalanine (43 mCi/mmol, Cat# NEC859S050UC, PerkinElmer). After 48 hours, cells were homogenized in lysis buffer containing 20 mM HEPES (pH 7.4), 1 mM EDTA, 5 mM EGTA, 10 mM MgCl$_2$, 100 mM β-glycerophosphate, 1 mM Na$_3$VO$_4$, 1 mM DTT, 1% TritonX-100 and supplemented with protease and phosphatase inhibitors (Cat# 78442, Thermo Fisher Scientific). Cell lysates were centrifuged for 5 min at 13 000 x g, 4 ℃, and the supernatant was collected. Separate aliquots of the supernatants were used for western blotting and analysis of ^{14}C-[1]-L-phenylalanine incorporation into proteins with liquid scintillation counting (see the Carbon-14 labeling -section). Western blot analysis of PHGDH was conducted by using low-fluorescence PVDF membranes and anti-PHGDH antibody (1:5000 dilution, Cat# HPA021241, Sigma-Aldrich) as previously (Mäntyselkä *et al*, 2024) with GAPDH (1:10 000 dilution, Cat# ab9485, Abcam) as a loading control.

Statistical testing in non-*omics* analyses

Statistical tests were conducted using GraphPad Prism10.3.1 (Dotmatics). The Student's t-test was used when two groups were compared whereas one-way ANOVA with Fisher's least significant difference test was used when a control group was compared to various treatment groups. Group difference was considered statistically significant with the $p \leq 0.05$.

Acknowledgments

We thank Associate Professor Thomas M. O'Connell for valuable advice regarding ^{13}C isotopic tracing and for providing the R script for ranking the significantly altered genes. We also thank all the laboratory technicians who helped us during this study. Figures 1A, 3A, the schematic part of 5A, and the graphical abstract were created with https://www.biorender.com/.

CRediT authorship contribution statement

Sakari Mäntyselkä: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Resources, Writing – Original Draft, Writing – Review & Editing, Visualization. **Erik Niem**i: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data Curation, Writing – Original Draft, Writing – Review & Editing, Visualization. **Laura Ylä-Outinen:** Conceptualization, Methodology, Investigation, Writing – Review & Editing**. Kalle Kolari:** Investigation, Writing – Review & Editing. **Liina Uusitalo-Kylmälä:** Formal analysis, Investigation, Writing – Review & Editing, Visualization. **Alfredo Ortega-Alonso:** Formal Analysis, Data Curation, Writing – Review & Editing. **Roosa-Maria Liimatainen:** Validation, Investigation, Review of the manuscript. **Vasco Fachada:** Software, Writing –

Review & Editing. **Perttu Permi:** Investigation, Writing – Review & Editing. **Elina**
Kalenius: Methodology, Writing – Review & Editing, Supervision. **Juha J Hulmi:**
Conceptualization, Resources, Writing – Original Draft, Writing – Review & Editing,
Supervision, Project administration. **Riikka Kivelä:** Conceptualization, Resources,
Writing – Original Draft, Writing – Review & Editing, Supervision, Project
administration, Funding acquisition.

Funding

876 This work was supported by the Research Council of Finland (R.K., Grant number:
877 348636), the Sigrid Jusélius Foundation (R.K.), the Finnish Foundation for
878 Cardiovascular Research (R.K., L. U-K.), the Emil Aaltonen Foundation (S.M., E.N.),
879 and the Finnish Cultural Foundation (K.K. and J.J.H).

Disclosure and competing interest statement

881 The authors declare that they have no known competing financial interests or
882 personal relationships that could have appeared to influence the work reported in
883 this paper.

Supplementary Data

A PDF file containing the Supplementary Figures 1-8.

Excel spreadsheets:

• Table S1: List of DEGs
• Table S2: Overlapping DEGs
• Table S3: GSEA results
• Table S4: Overlapping GSEA pathways
• Table S5: Metabolomics data
• Table S6: ^{13}C-(U)-glucose labeling data
• Table S7: MRM-transitions used in ^{13}C-(U)-glucose tracing

Data availability

896 Filtered non-aligned RNA sequencing data and non-normalized quantified reads are
897 available in the Gene Expression Omnibus (GEO) data repository (accession
898 number: GSE288064). Processed transcriptomics data are shared as
899 Supplementary Excel spreadsheets. Numerical metabolomics and isotopic tracing
900 data are shared as Supplementary Excel spreadsheets.

References

Białopiotrowicz E, Noyszewska-Kania M, Kachamakova-Trojanowska N, Łoboda A, Cybulska M, Grochowska A, Kopczyński M, Mikula M, Prochorec-Sobieszek M, Firczuk M, *et al* (2020) Serine Biosynthesis Pathway Supports MYC-miR-494-EZH2 Feed-Forward Circuit Necessary to Maintain Metabolic and Epigenetic Reprogramming of Burkitt Lymphoma Cells. *Cancers* 12: 580

Biernat K, Kuciel N, Mazurek J & Hap K (2024) Is It Possible to Train the Endothelium?-A Narrative Literature Review. *Life Basel Switz* 14: 616

Bose S, Ramesh V & Locasale JW (2019) Acetate Metabolism in Physiology, Cancer, and Beyond. *Trends Cell Biol* 29: 695–703

Chastin SFM, Abaraogu U, Bourgois JG, Dall PM, Darnborough J, Duncan E, Dumortier J, Pavón DJ, McParland J, Roberts NJ, *et al* (2021) Effects of Regular Physical Activity on the Immune System, Vaccination and Risk of Community-Acquired Infectious Disease in the General Population: Systematic Review and Meta-Analysis. *Sports Med Auckl NZ* 51: 1673–1686

Chen Y, Lun ATL & Smyth GK (2016) From reads to genes to pathways: differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Research* 5: 1438

Cronin NM, Dawson LW & DeMali KA (2024) Shear stress-stimulated AMPK couples endothelial cell mechanics, metabolism and vasodilation. *J Cell Sci* 137: jcs262232

Davis MJ, Earley S, Li Y-S & Chien S (2023) Vascular mechanotransduction. *Physiol Rev* 103: 1247–1421

De Bock K, Georgiadou M, Schoors S, Kuchnio A, Wong BW, Cantelmo AR, Quaegebeur A, Ghesquière B, Cauwenberghs S, Eelen G, *et al* (2013) Role of PFKFB3-driven glycolysis in vessel sprouting. *Cell* 154: 651–663

Englund DA, Zhang X, Aversa Z & LeBrasseur NK (2021) Skeletal muscle aging, cellular senescence, and senotherapeutics: Current knowledge and future directions. *Mech Ageing Dev* 200: 111595

Fan W & Evans R (2015) PPARs and ERRs: molecular mediators of mitochondrial metabolism. *Curr Opin Cell Biol* 33: 49–54

Ge SX, Jung D & Yao R (2020) ShinyGO: a graphical gene-set enrichment tool for animals and plants. *Bioinforma Oxf Engl* 36: 2628–2629

Große-Segerath L, Follert P, Behnke K, Ettich J, Buschmann T, Kirschner P, Hartwig S, Lehr S, Korf-Klingebiel M, Eberhard D, *et al* (2024) Identification of myeloid-derived growth factor as a mechanically-induced, growth-promoting angiocrine signal for human hepatocytes. *Nat Commun* 15: 1076

van Hall G, Sacchetti M & Rådegran G (2002) Whole body and leg acetate kinetics at rest, during exercise and recovery in humans. *J Physiol* 542: 263–272

Halonen EJ, Gabriel I, Kelahaara MM, Ahtiainen JP & Hulmi JJ (2024) Does Taking a Break Matter-Adaptations in Muscle Strength and Size Between Continuous and Periodic Resistance Training. *Scand J Med Sci Sports* 34: e14739

Handzlik MK & Metallo CM (2023) Sources and Sinks of Serine in Nutrition, Health, and Disease. *Annu Rev Nutr* 43: 123–151

Hinkle ER, Blue RE, Tsai Y-H, Combs M, Davi J, Coffey AR, Boriek AM, Taylor JM, Parker JS & Giudice J (2022) Stretching muscle cells induces transcriptional and splicing transitions and changes in SR proteins. *Commun Biol* 5: 987

Hood DA, Tryon LD, Carter HN, Kim Y & Chen CCW (2016) Unravelling the mechanisms regulating muscle mitochondrial biogenesis. *Biochem J* 473: 2295–2314

Horibata Y, Ando H, Itoh M & Sugimoto H (2013) Enzymatic and transcriptional regulation of the cytoplasmic acetyl-CoA hydrolase ACOT12. *J Lipid Res* 54: 2049–2059

Hulmi JJ, Ahtiainen JP, Kaasalainen T, Pöllänen E, Häkkinen K, Alen M, Selänne H, Kovanen V & Mero AA (2007) Postexercise myostatin and activin IIb mRNA levels: effects of strength training. *Med Sci Sports Exerc* 39: 289–297

Ilaiwy A, Quintana MT, Bain JR, Muehlbauer MJ, Brown DI, Stansfield WE & Willis MS (2016) Cessation of biomechanical stretch model of C2C12 cells models myocyte atrophy and anaplerotic changes in metabolism using non-targeted metabolomics analysis. *Int J Biochem Cell Biol* 79: 80–92

Jing X, Lyu J & Xiong J (2023) Acetate regulates GAPDH acetylation and T helper 1 cell differentiation. *Mol Biol Cell* 34: br10

Jufri NF, Mohamedali A, Avolio A & Baker MS (2015) Mechanical stretch: physiological and pathological implications for human vascular endothelial cells. *Vasc Cell* 7: 8

Kim J, Cross JM & Bamman MM (2005) Impact of resistance loading on myostatin expression and cell cycle regulation in young and older men and women. *Am J Physiol Endocrinol Metab* 288: E1110-1119

Kobialka P & Graupera M (2019) Revisiting PI3-kinase signalling in angiogenesis. *Vasc Biol Bristol Engl* 1: H125–H134

Kong X, Kapustka A, Sullivan B, Schwarz GJ & Leckband DE (2022) Extracellular matrix regulates force transduction at VE-cadherin junctions. *Mol Biol Cell* 33: ar95

Kujala UM, Mäkinen V-P, Heinonen I, Soininen P, Kangas AJ, Leskinen TH, Rahkila P, Würtz P, Kovanen V, Cheng S, *et al* (2013) Long-term leisure-time physical activity and serum metabolome. *Circulation* 127: 340–348

Kwee BJ, Budina E, Najibi AJ & Mooney DJ (2018) CD4 T-cells regulate angiogenesis and myogenesis. *Biomaterials* 178: 109–121

Langmead B & Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat*
Methods 9: 357–359

Lautaoja JH, M O'Connell T, Mäntyselkä S, Peräkylä J, Kainulainen H, Pekkala S,
Permi P & Hulmi JJ (2021) Higher glucose availability augments the metabolic
responses of the C2C12 myotubes to exercise-like electrical pulse stimulation.
Am J Physiol Endocrinol Metab 321: E229–E245

Lautaoja JH, Turner DC, Sharples AP, Kivelä R, Pekkala S, Hulmi JJ & Ylä-Outinen
L (2023) Mimicking exercise in vitro: effects of myotube contractions and
mechanical stretch on omics. *Am J Physiol Cell Physiol* 324: C886–C892

Law CW, Chen Y, Shi W & Smyth GK (2014) voom: Precision weights unlock linear
model analysis tools for RNA-seq read counts. *Genome Biol* 15: R29

Lee G, Wong C, Cho A, West JJ, Crawford AJ, Russo GC, Si BR, Kim J, Hoffner L,
Jang C, *et al* (2024) E-Cadherin Induces Serine Synthesis to Support
Progression and Metastasis of Breast Cancer. *Cancer Res* 84: 2820–2835

Legrand C, Bour JM, Jacob C, Capiaumont J, Martial A, Marc A, Wudtke M,
Kretzmer G, Demangel C & Duval D (1992) Lactate dehydrogenase (LDH)
activity of the cultured eukaryotic cells as marker of the number of dead cells
in the medium [corrected]. *J Biotechnol* 25: 231–243

Li S, Xu Z, Wang Y, Chen L, Wang X, Zhou Y, Lei D, Zang G & Wang G (2024)
Recent advances of mechanosensitive genes in vascular endothelial cells for
the formation and treatment of atherosclerosis. *Genes Dis* 11: 101046

Li W & Sumpio BE (2005) Strain-induced vascular endothelial cell proliferation
requires PI3K-dependent mTOR-4E-BP1 signal pathway. *Am J Physiol Heart*
Circ Physiol 288: H1591-1597

Liao Y, Smyth GK & Shi W (2014) featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinforma Oxf*
Engl 30: 923–930

Liu X, Cooper DE, Cluntun AA, Warmoes MO, Zhao S, Reid MA, Liu J, Lund PJ,
Lopes M, Garcia BA, *et al* (2018) Acetate Production from Glucose and
Coupling to Mitochondrial Metabolism in Mammals. *Cell* 175: 502-513.e13

Lorenz L, Axnick J, Buschmann T, Henning C, Urner S, Fang S, Nurmi H, Eichhorst
N, Holtmeier R, Bódis K, *et al* (2018) Mechanosensing by β1 integrin induces
angiocrine signals for liver growth and survival. *Nature* 562: 128–132

Lunt SY & Vander Heiden MG (2011) Aerobic glycolysis: meeting the metabolic
requirements of cell proliferation. *Annu Rev Cell Dev Biol* 27: 441–464

Manning BD & Toker A (2017) AKT/PKB Signaling: Navigating the Network. *Cell*
169: 381–405

Mäntyselkä S, Kolari K, Baumert P, Ylä-Outinen L, Kuikka L, Lahtonen S, Permi P,
Wackerhage H, Kalenius E, Kivelä R, *et al* (2024) Serine synthesis pathway

enzyme PHGDH is critical for muscle cell biomass, anabolic metabolism, and
mTORC1 signaling. *Am J Physiol Endocrinol Metab* 326: E73–E91

Millard P, Delépine B, Guionnet M, Heuillet M, Bellvert F & Létisse F (2019) IsoCor:
isotope correction for high-resolution MS labeling experiments. *Bioinforma Oxf*
Engl 35: 4484–4487

Miller KD, O'Connor S, Pniewski KA, Kannan T, Acosta R, Mirji G, Papp S, Hulse M,
Mukha D, Hlavaty SI, *et al* (2023) Acetate acts as a metabolic
immunomodulator by bolstering T-cell effector function and potentiating
antitumor immunity in breast cancer. *Nat Cancer* 4: 1491–1507

Nakai N, Kawano F & Nakata K (2015) Mechanical stretch activates mammalian
target of rapamycin and AMP-activated protein kinase pathways in skeletal
muscle cells. *Mol Cell Biochem* 406: 285–292

Nan W, He Y, Wang S & Zhang Y (2023) Molecular mechanism of VE-cadherin in
regulating endothelial cell behaviour during angiogenesis. *Front Physiol* 14:
1234104

Pacold ME, Brimacombe KR, Chan SH, Rohde JM, Lewis CA, Swier LJYM,
Possemato R, Chen WW, Sullivan LB, Fiske BP, *et al* (2016) A PHGDH
inhibitor reveals coordination of serine synthesis and one-carbon unit fate. *Nat*
Chem Biol 12: 452–458

Peng Z, Shu B, Zhang Y & Wang M (2019) Endothelial Response to
Pathophysiological Stress. *Arterioscler Thromb Vasc Biol* 39: e233–e243

Pepe GJ & Albrecht ED (2023) Microvascular Skeletal-Muscle Crosstalk in Health
and Disease. *Int J Mol Sci* 24: 10425

Pillon NJ, Gabriel BM, Dollet L, Smith JAB, Sardón Puig L, Botella J, Bishop DJ,
Krook A & Zierath JR (2020) Transcriptomic profiling of skeletal muscle
adaptations to exercise and inactivity. *Nat Commun* 11: 470

Powers SK, Radak Z, Ji LL & Jackson M (2024) Reactive oxygen species promote
endurance exercise-induced adaptations in skeletal muscles. *J Sport Health*
Sci 13: 780–792

Qasem RJ, Farh IK & Al Essa MA (2017) A novel LC-MS/MS method for the
quantitative measurement of the acetate content in pharmaceutical peptides.
J Pharm Biomed Anal 146: 354–360

Qiu J, Villa M, Sanin DE, Buck MD, O'Sullivan D, Ching R, Matsushita M, Grzes KM,
Winkler F, Chang C-H, *et al* (2019) Acetate Promotes T Cell Effector Function
during Glucose Restriction. *Cell Rep* 27: 2063-2074.e5

Ren D, Song J, Liu R, Zeng X, Yan X, Zhang Q & Yuan X (2021) Molecular and
Biomechanical Adaptations to Mechanical Stretch in Cultured Myotubes. *Front*
Physiol 12: 689492

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W & Smyth GK (2015) limma
powers differential expression analyses for RNA-sequencing and microarray
studies. *Nucleic Acids Res* 43: e47

Robinson MD, McCarthy DJ & Smyth GK (2010) edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinforma*
Oxf Engl 26: 139–140

Robinson MD & Oshlack A (2010) A scaling normalization method for differential
expression analysis of RNA-seq data. *Genome Biol* 11: R25

Ross M, Kargl CK, Ferguson R, Gavin TP & Hellsten Y (2023) Exercise-induced
skeletal muscle angiogenesis: impact of age, sex, angiocrines and cellular
mediators. *Eur J Appl Physiol* 123: 1415–1432

Schiaffino S, Pereira MG, Ciciliot S & Rovere-Querini P (2017) Regulatory T cells
and skeletal muscle regeneration. *FEBS J* 284: 517–524

Schranner D, Kastenmüller G, Schönfelder M, Römisch-Margl W & Wackerhage H
(2020) Metabolite Concentration Changes in Humans After a Bout of
Exercise: a Systematic Review of Exercise Metabolomics Studies. *Sports*
Med - Open 6: 11

Seo H-R, Jeong HE, Joo HJ, Choi S-C, Park C-Y, Kim J-H, Choi J-H, Cui L-H, Hong
SJ, Chung S, *et al* (2016) Intrinsic FGF2 and FGF5 promotes angiogenesis of
human aortic endothelial cells in 3D microfluidic angiogenesis system. *Sci*
Rep 6: 28832

Smith JAB, Gabriel BM, Abdelmoez AM, Savikj M, Wright SC, Koutsilieri S, Barrès
R, Lauschke VM, Krook A, Zierath JR, *et al* (2024) Downregulation of NR4A3
during inactivity alters glucose metabolism and impairs translation in human
skeletal muscle. doi:10.1101/2024.03.11.584422 [PREPRINT]

Smith JAB, Murach KA, Dyar KA & Zierath JR (2023) Exercise metabolism and
adaptation in skeletal muscle. *Nat Rev Mol Cell Biol* 24: 607–632

Sottiurai VS, Sue SL, Breaux JR & Smith LM (1989) Adaptability of endothelial
orientation to blood flow dynamics--a morphologic analysis. *Eur J Vasc Surg*
3: 145–151

Stegen S, Loopmans S, Stockmans I, Moermans K, Carmeliet P & Carmeliet G
(2022) De novo serine synthesis regulates chondrocyte proliferation during
bone development and repair. *Bone Res* 10: 14

Tsugawa H, Kanazawa M, Ogiwara A & Arita M (2014) MRMPROBS suite for
metabolomics using large-scale MRM assays. *Bioinforma Oxf Engl* 30: 2379–
2380

Turner DC, Gorski PP, Seaborne RA, Viggars M, Murphy M, Jarvis JC, Martin NRW,
Stewart CE & Sharples AP (2021) Mechanical loading of bioengineered
skeletal muscle in vitro recapitulates gene expression signatures of resistance
exercise in vivo. *J Cell Physiol* 236: 6534–6547

Vandekeere S, Dubois C, Kalucka J, Sullivan MR, García-Caballero M, Goveia J,
Chen R, Diehl FF, Bar-Lev L, Souffreau J, *et al* (2018) Serine Synthesis via
PHGDH Is Essential for Heme Production in Endothelial Cells. *Cell Metab* 28:
573-587.e13

Verma M, Asakura Y, Murakonda BSR, Pengo T, Latroche C, Chazaud B, McLoon
LK & Asakura A (2018) Muscle Satellite Cell Cross-Talk with a Vascular Niche
maintains quiescence via VEGF and Notch Signaling. *Cell Stem Cell* 23: 530-
543.e9

Wickramasinghe NM, Sachs D, Shewale B, Gonzalez DM, Dhanan-Krishnan P,
Torre D, LaMarca E, Raimo S, Dariolli R, Serasinghe MN, *et al* (2022)
PPARdelta activation induces metabolic and contractile maturation of human
pluripotent stem cell-derived cardiomyocytes. *Cell Stem Cell* 29: 559-576.e7

Yoon SJ, Combs JA, Falzone A, Prieto-Farigua N, Caldwell S, Ackerman HD, Flores
ER & DeNicola GM (2023) Comprehensive Metabolic Tracing Reveals the
Origin and Catabolism of Cysteine in Mammalian Tissues and Tumors.
Cancer Res 83: 1426–1442

Yu G, Wang L-G, Han Y & He Q-Y (2012) clusterProfiler: an R package for
comparing biological themes among gene clusters. *Omics J Integr Biol* 16:
284–287

Yung YC, Chae J, Buehler MJ, Hunter CP & Mooney DJ (2009) Cyclic tensile strain
triggers a sequence of autocrine and paracrine signaling to regulate
angiogenic sprouting in human vascular cells. *Proc Natl Acad Sci U S A* 106:
15279–15284

Zhang D, Li AM, Hu G, Huang M, Yang F, Zhang L, Wellen KE, Xu X, Conn CS, Zou
W, *et al* (2023) PHGDH-mediated endothelial metabolism drives glioblastoma
resistance to chimeric antigen receptor T cell immunotherapy. *Cell Metab* 35:
517-534.e8

Zhang T, Gillies MC, Madigan MC, Shen W, Du J, Grünert U, Zhou F, Yam M & Zhu
L (2018) Disruption of De Novo Serine Synthesis in Müller Cells Induced
Mitochondrial Dysfunction and Aggravated Oxidative Damage. *Mol Neurobiol*
55: 7025–7037

Zhang X, Sun M, Jiao Y, Lin B & Yang Q (2022) PHGDH Inhibitor CBR-5884 Inhibits
Epithelial Ovarian Cancer Progression via ROS/Wnt/β-Catenin Pathway and
Plays a Synergistic Role with PARP Inhibitor Olaparib. *Oxid Med Cell Longev*
2022: 9029544

1131