Stress-induced ribosome degradation in *Bacillus subtilis* is mediated by the RNase Y-

specificity complex*.*

Fabián A. Cornejo[1,*], Kristina Driller[1,2,*], Rina Ahmed-Begrich[1], Katja Schmidt[1], Michael Jahn[1],

Vivekanandan Shanmuganathan[1], Karin Hahnke[1], Florian Kondrot[1], Thomas F. Wulff[1],

Sebastian Rämisch[1], Kathirvel Alagesan[1], Emmanuelle Charpentier[1,3], and Kürşad Turgay[1,2]

[1]Max Planck Unit for the Science of Pathogens, Berlin, Germany

[2]Leibniz Universität Hannover, Institute of Microbiology, Hannover, Germany

[3]Humboldt-Universität zu Berlin, Institute of Biology, Berlin, Germany

* These authors equally contributed to this work

Correspondence to: cornejo@mpusp.mpg.de, turgay@mpusp.mpg.de

Abstract

Limiting the synthesis and activity of ribosomes is crucial for adaptation to stresses, such as

heat or nutrient starvation. In *Bacillus subtilis,* this can be achieved through the coordinated

action of the alarmones (p)ppGpp and the transcription factor Spx. Here, we performed a

genetic screen to uncover novel factors contributing to the heat shock response of *B. subtilis*.

We identified the Y-complex, which confers specificity to the endonuclease RNase Y, as a

critical player during stress conditions, such as heat or transition into stationary phase. This

protein complex is required for processing diverse RNAs, notably the maturation of mRNAs

encoding proteins involved in translation and metabolism. We further demonstrate that the Y-

complex and RNase Y initiate the degradation of rRNAs of mature ribosomes, lowering their

abundance. We propose that the Y-complex is a regulatory hub that modulates gene

expression, adjusts protein synthesis and resource allocation.

Keywords: *Bacillus subtilis*, Ribosome, Stress response, Y-complex, RNase Y, Ribosome

degradation, Heat shock, (p)ppGpp

Introduction

Bacteria can sense, adapt to, and survive stressful environmental changes facilitated by various general and specific stress response systems. Proteotoxic stresses, like high temperatures, significantly disrupt protein folding and can lead to the formation of toxic protein aggregates. Chaperones and proteases are expressed as a response due to the inactivation of the transcriptional repressors HrcA and CtsR in *Bacillus subtilis*[1–4]. The activation of the protein quality control (PQC) allows the refolding of un- or misfolded proteins by different chaperone systems or the degradation of irreparable proteins through AAA+ protease complexes like ClpCP[5,6].

In addition to the repairing or removal of protein aggregates, inhibiting translation is crucial for survival during proteotoxic stresses, as it can alleviate the extra load on the PQC system[7–10]. In *Bacillus subtilis*, the nucleotide second messengers guanosine tetraphosphate and guanosine pentaphosphate, collectively referred to as (p)ppGpp, and the transcription factor Spx independently regulate synthesis and activity of ribosomes during stress[7,11]. Spx is a redox-sensitive heat shock transcription factor that controls the expression of redox chaperones and simultaneously represses transcription of translation-related genes during oxidative and heat shock response[11–14].

The alarmones (p)ppGpp were initially discovered to be synthesized during amino acid starvation, but more recently, it was observed that their synthesis is also induced by protein folding stresses, such as heat or oxidative stress[7]. During heat stress, their accumulation results in increased thermoresistance, inhibition of translation, and a reduction in protein aggregation[7]. (p)ppGpp inhibit translation by binding to specific GTPases (e.g. IF2, RbgA)[15–17]. Furthermore, (p)ppGpp accumulation can lead to decreased transcription of e.g. rRNA and ribosomal protein (r-protein) transcripts by inhibiting GTP biosynthesis and reducing its levels[18,19]. This downregulation of translation is an important component of the response to nutrient scarcity as well as heat shock.

In this study, we aimed to identify new genes or pathways involved in the heat shock response using a genetic screen that reports on fitness. We identified that the Y-complex (RicAFT-complex), which interacts with the endonuclease RNase Y and conveys substrate specificity, plays a role in the heat shock response. We demonstrate further that the Y-complex regulates ribosome levels by initiating ribosome degradation during nutrient depletion or heat shock. Mutants lacking a Y-complex member show increased ribosome levels before and during heat shock, increasing the burden on the PQC system and, in consequence, protein aggregation during heat stress. These findings not only indicate the importance of RNases during stress response but also emphasize that controlling the stability of RNA is an important general mechanism to modify cellular function.

Results

The Y-complex is necessary for survival during heat shock

To identify genes that play a role in the heat shock response, we used the BKE single-mutant library[20], which includes a barcoded deletion cassette for each non-essential gene in *B. subtilis*. All gene deletion mutant strains were pooled, grown at 30°C, and subjected to a mild heat shock at 50°C during the mid-exponential phase, while a control was maintained at 30°C. Relative fitness scores for each gene deletion in the pooled population were calculated by quantifying each barcode at different temperatures (**Fig. 1A, Table S1**). The relative fitness of the heat shocked cultures compared to the cultures grown at 30°C was used to identify genes whose deletion results in a change in fitness during heat shock (**Fig. 1A-B, Table S1**).

We observed that *B. subtilis* strains lacking the genes coding for the heat shock transcriptional repressors HrcA or CtsR exhibit higher fitness at 50°C compared to 30°C (**Fig. S1A**), consistent with their known regulatory role. In contrast, deletion of the genes coding for the DnaK chaperone system, the protease ClpP, and the adaptor protein complex McsA-McsB show greater sensitivity to increased temperatures (**Fig. S1B**), demonstrating that our screening approach can identify genes with already known heat-shock phenotypes[21–23].

The control of transcription initiation by transcription factors has been well established[24]. However, the contribution of RNA stability in stress response is not well understood. Interestingly, six mutants lacking RNases or RNase-regulating genes exhibit differential fitness at 50°C compared to 30°C (**Fig. 1B**). The deletion of RNase Mini-III (*ΔmrnC)* results in lower fitness at 30°C, whereas its fitness remains unaffected at 50°C. Deletions of RNase J2 (*ΔrnjB)*, the nano-RNase A (*ΔnrnA*), and the RNA pyrophosphohydrolase and (p)ppGpp hydrolase NahA (*Δyvcl*) exhibit reduced fitness during increased temperatures (**Fig. 1B**).

One of the mutants with the lowest fitness score at 50°C is *ΔymcA* (**Fig. 1B**). YmcA (RicA) interacts with YaaT (RicT) and YlbF (RicF) to form the Y-complex, which associates with the endonuclease RNase Y for maturation of polycistronic mRNA and riboswitches[25,26]. In addition, we also identified *ΔyaaT* in our screen (**Fig. 1B**). We could confirm the heat sensitivity of all three deletion mutants of the Y-complex genes in single-culture experiments during a severe heat shock at 54°C (**Fig. 1C**).

Identification of Y-complex-dependent cleavages at late exponential and transition phases

We evaluated the growth of all three strains lacking the Y-complex encoding genes. They exhibit a similar growth defect that begins at the end of the exponential phase and continues during transition phase until the stationary phase (**Fig. 2A**). Given that this complex associates

with the endonuclease RNase Y and regulates substrate selection[26], we investigated Y-
complex-dependent cleavages in the late exponential and transition phase (**Fig. 2A**) using the
ISCP (Identification of Specific Cleavage Positions) approach[27]. We compared the coverage
of RNA 3' and 5' ends in the WT and ΔymcA mutant, and the ends significantly reduced in the
ΔymcA were classified as putative Y-complex-dependent cleavages (**Table S2**). It is worth
noting that deleting any member of the Y-complex results in a significant decrease in the
processing of substrate RNA[26]. When a transcript is cleaved, it results in two products: one is
quickly degraded by exonucleases, while the other is stabilized[27]. Consequently, it is not
always possible to detect both 5' and 3' ends after cleavage.

We identified 221 and 258 Y-complex-dependent ends in late exponential and transition
phases, with six also detected in a previous study at the mid-exponential phase[26] (**Table S3**).
In the late exponential phase, 5' and 3' ends are almost equally distributed, yet more 5' ends
are detected in the transition phase (**Fig. 2B**).

It is known that the Y-complex and RNase Y generate fragments with differential stability in the
mid-exponential phase[26]. Here, we evaluated the stability of the resulting fragments in the late
exponential and transition phases by calculating the ratio of coverage upstream and
downstream of the cleavage site (**Table S2**). We observed a trend toward stabilizing
downstream fragments. However, upstream fragments can also be stabilized, but to a lesser
extent (**Fig. S2A**). The ISCP method differentiates processing sites with blunt ends (Unique,
U-type) or when it was further processed by exonucleases[27] (Stepped, S-type). Most S-types
were found at the 3' ends, suggesting further processing by exonucleases, while most U-types
were detected at 5' ends (**Fig. 2C**). This is consistent with the trend to stabilise the downstream
fragment (**Fig. S2A**).

The Y-complex could target a particular sequence motif, thereby helping to recognize specific
substrates. However, a motif search using U-type ends resembles the simple motif described
for RNase Y in *B. subtilis*[28], *S. aureus*[29], and *S. pyogenes*[27], being most likely a G upstream of
the cleavage site (**Fig. S2B**). Given the simple sequence motif, we investigated whether
cleavages occur randomly in longer or more abundant transcripts. Notably, approximately 81%
of targeted transcripts are cleaved only once, about 16% twice, and very few are cleaved
multiple times (**Fig. S2C**). Neither the length nor the abundance of a transcript predicts whether
it is cleaved, suggesting that other factors influence where cleavages occur (**Fig. S2D-E, Table
S4**). Recent studies have shown that the cleavage efficiency is multifactorial, depending on
the sequence and secondary structures surrounding the cleavage site[28,30].

As we observed that RNA processing could result in fragments with different stability (**Fig.
S2A**), we wondered if this could be a consequence of secondary structures protecting them
from exonucleases. We used U-type ends and compared transcripts with the same cleavage

motif but not cleaved as controls. For the 5' ends, a strong decrease in the minimum free
energy ~25 nt downstream of the cleavage site indicated a possible secondary structure, while
the upstream fragment remained similar to the control (**Fig. 2D**). The upstream fragment's
lower stability may be due to its lack of secondary structure, making it accessible to 3'-5'
exonucleases. For the 3' ends, the more stable upstream fragment could form a possible
secondary structure starting 10 nt upstream of the cleavage site (**Fig. 2D**). The generation of
mRNA fragments with different stability might have a regulatory role in gene expression.

Y-complex-mediated transcript cleavages and mRNA maturation

Most detected Y-complex-dependent ends are inside ORFs, mainly in polycistronic mRNAs
(**Fig. 2E-F**), confirming the role of the Y-complex in the maturation and stoichiometry regulation
of co-transcribed genes[26]. An example is the *cggR-gapA* transcript, where the Y-complex
mediates cleavage in the coding sequence of *cggR* near the stop codon[25,26]. Cleavage renders
the *cggR* fragment less stable compared to the *gapA* fragment (**Fig. S3A**), thereby uncoupling
the transcript levels of these co-transcribed genes. Interestingly, we observed other cleavages
generating similar uncoupling of transcripts, such as in the gene encoding the endonuclease
YloC (**Fig. S3B**, **Table S2)**. In addition, cleavages in genes transcribed as monocistronic
mRNAs, such as *guaB,* suggest that the Y-complex could initiate their degradation (**Fig. S3C**,
Table S2).

We observe Y complex-dependent processing in 5', 3', and internal UTRs (**Fig. 2F**). Cleavages
in 5' UTR suggest a possible regulatory role by decreasing the abundance of the mRNA
containing them, as already described for riboswitches[26,31], or by producing a shorter version
of a 5' UTR, which could affect gene expression. Processing of 3' and internal UTR could
influence mRNA stability and polycistronic mRNA maturation. We observed that some 3' UTR
cleavages are located upstream of the Rho-independent terminator, decreasing the stability of
the upstream fragment and leaving an apparent more stable downstream fragment that
includes its terminator structure, as observed for *trxB* (**Fig. S3D**). This phenomenon has been
observed in *E. coli*, where RNase E cleaves upstream of the Rho-independent terminator[32,33].

In polycistronic mRNA maturation, we observed that Y-complex processes internal UTRs,
decoupling the RNA levels of coding fragments (**Fig. S3E-F**). For example, the *bipA-ylaH*
transcript is cleaved 21 nt after the *bipA* stop codon, rendering the upstream fragment less
stable than the *ylaH* fragment (**Fig. S3E**). We also observed a cleavage between the *fusA* (EF-
G) and *tufA* (EF-Tu) genes, both coding for translation elongation factors (**Fig. S3F**).

The Y-complex targets translation and metabolism-related genes in the late exponential phase

We wondered if the Y-complex primarily processes transcripts bearing genes with specific functions required for switching the growth phase. Therefore, we conducted an overrepresentation analysis and observed that the most significantly enriched categories pertain to "Metabolism" and "Information processing" (**Fig. 2G, Fig. S4A, Table S5**). Overall, the Y-complex is predominantly involved in the processing of operons encoding genes for carbon metabolism (e.g., glycolysis and gluconeogenesis), nucleotide metabolism (e.g., pyrimidine and purine biosynthesis), and amino acid biosynthesis and utilization (**Fig. 2G**). This allows adjusting the utilization and biosynthesis of nutrients when they become limited. In addition, all transcripts containing rRNA and ~50% of transcripts carrying genes encoding translation factors and r-proteins are processed by the Y-complex (**Fig. 2G**).

The reduced RNA processing in the *ymcA* deletion strain results in significant changes in the transcriptome (**Fig. S4B, Table S6-7**). Consequently, transcripts of certain pathways related to metabolism and translation, among others, show maturation events that depend on the Y-complex and also demonstrate differential expression upon deletion of *ymcA* (**Fig. 2G**, **Fig. S4B**). This may result from either a direct or indirect effect of decreased RNA maturation events. Our results suggest that the Y-complex and RNase Y are important for transitioning to the stationary phase by processing transcripts related to translation and metabolism, which may impact their stability and/or gene expression.

The Y-complex targets ribosomes for degradation

Several Y complex-dependent cleavages were detected in the rRNA sequence during both growth phases, most of them being in the 23S rRNA during the late exponential phase (**Fig. 3A**). We detected cleavages in specific helices in the 23S and 16S rRNA, mainly in loops of hairpin structures (**Fig. S5A-B**). These helices are accessible and primarily located close to the A-site and the interface between the two ribosomal subunits (**Fig. S6**), which could be recognized and cut by the Y-complex with the endonuclease RNase Y. These cleavages will most likely destabilize the ribosome structure. Generating free 5' and 3' RNA ends, which are substrates for exonucleases such as PNPase, RNase R, and RNase J, will result in further RNA degradation and ribosome disassembly, releasing the ribosomal proteins as well.

We confirmed the observed rRNA degradation via northern blots of 23S or 16S rRNA (**Fig. 3B**, **Fig. S7A**). Deletion of *ymcA* strongly reduces the presence of rRNA degradation intermediates during the late exponential phase. As ribosomes are assembled co-transcriptionally[34], we inhibited transcription by adding rifampicin to cultures at the late exponential phase. We observed that stopping transcription does not affect rRNA degradation of the 23S and even

raises the levels of 16S degradation intermediates, indicating that mature ribosomes are
degraded (**Fig. 3B**, **Fig. S7A**).

Cleavages within mature rRNA can trigger ribosome disassembly and degradation, which,
during the transition phase, coincides with translation downregulation through reduced
synthesis and inhibition of ribosome activity. This suggests that the Y-complex may be involved
in reducing ribosome levels to accommodate the new, slower growth rate. Therefore, we
quantified ribosome content by measuring the abundance of all r-proteins via mass
spectrometry (**Table S8**). When *B. subtilis* shifts from exponential to the transition phase, we
observed a coordinated reduction in r-protein levels (**Fig. 3C**), indicating a decrease in
ribosome levels. This reduction is highly impaired in $\Delta ymcA$ (**Fig. 3C**), where the levels of r-
proteins do not decline to the same degree as in the WT, likely due to the reduced degradation
of ribosomes.

Accumulation of (p)ppGpp leads to inhibition of transcription of rRNA and r-proteins[19,35] upon
shortages of nutrients during the transition to the stationary phase. To test if the deletion of
ymcA is interfering with this transcriptional downregulation, we measured the promoter activity
of the rRNA promoters *rrnJ* and *rrnB* and r-protein promoters of *rplK*, *rplJ*, and *rpsJ*. All these
promoters show similar downregulation in the late exponential and transition phases in both
WT and $\Delta ymcA$ strains (**Fig. 3D-E**, **Fig. S7B-D**), confirming that their transcriptional control
remains unaffected and is not the cause of the increased ribosome amounts in the $\Delta ymcA$
strain.

Notably, we observed that the increased levels of ribosomes in a $\Delta ymcA$ strain led to the
accumulation of hibernating 100S ribosomes during the transition phase (**Fig. 3F**). This 100S
is no longer observed after the deletion of the gene encoding the hibernation factor *hpf*, which
triggers their formation[36,37] (**Fig. S7E**). The presence of hibernating ribosomes at this time point
indicates that functional and superabundant ribosomes in the $\Delta ymcA$ strain are sequestered
away from the actively translating pool through hibernation.

(p)ppGpp and the Y-complex play a crucial role in regulating the levels of ribosomes

Since ribosome levels and translation are lowered by (p)ppGpp under nutrient limitation or
other stresses, we wondered if there is a genetic interaction between the two systems. Deleting
ymcA in a strain that cannot synthesize (p)ppGpp [(p)ppGpp^0] leads to an even stronger growth
defect during the transition phase (**Fig. 4A**). The (p)ppGpp^0 strain, like $\Delta ymcA$, shows impaired
reduction of ribosome levels during the transition phase (**Fig. 4B**). The double mutant
(p)ppGpp^0 $\Delta ymcA$ shows almost no reduction in ribosome quantities during this growth phase
(**Fig. 4B-C**). The same is observed when the r-protein mass fraction is calculated at the
transition phase (**Fig. 4D**). These experiments suggest that the Y-complex and (p)ppGpp

control ribosome levels and translation independently during post-exponential growth. The
absence of regulation by both systems results in a severe growth defect.

Increased ribosome levels lead to protein aggregation during heat shock

Since we discovered the $\Delta ymcA$ strain in a competition experiment during heat shock, we
explored whether the Y-complex is involved in lowering ribosome levels upon proteotoxic
stress. We evaluated the degradation of 23S and 16S rRNA by the Y-complex before and after
heat shock (**Fig. 5A**, **Fig. S8A**). After 15 minutes at 50°C, degradation intermediates are
observed in the WT but not in $\Delta ymcA$, confirming a role of the Y-complex in degrading
ribosomes during heat stress (**Fig. 5A**). Comparing r-protein mass fractions before and during
heat shock shows that both WT and $\Delta ymcA$ downregulate ribosome levels during the
experiment. Overall, there are increased levels of r-proteins in the $\Delta ymcA$ strain already before
the heat shock (**Fig. 5B**), which suggests a role of the Y-complex in maintaining ribosome
levels during normal growth. Notably, the levels and induction of known heat-shock-regulated
proteins are not affected in the *ymcA* deletion (**Fig. 5C, Table S9, Fig. S8B**). This suggests
that chaperone and protease induction during heat stress in the $\Delta ymcA$ strain is intact.

The downregulation of translation is crucial to limit the increased burden of misfolding and
aggregation-prone nascent polypeptide chains on the PQC system. In agreement, a (p)ppGpp^0
strain shows increased aggregate formation during heat shock[7]. We quantified the relative
amount of aggregated proteins before and after 15 minutes of severe heat shock at 54°C and
observed that the $\Delta ymcA$ strain contained significantly higher amounts of aggregated proteins
already prior to heat exposure (**Fig. 5D**). This is consistent with the observed increased
ribosome levels of the $\Delta ymcA$ strain already without heat shock (time point 0 in **Fig. 5B**). After
15 minutes of heat shock, $\Delta ymcA$ shows 20% more aggregated proteins than WT. The
(p)ppGpp^0 $\Delta ymcA$ mutant exhibits even higher protein aggregation than $\Delta ymcA$ or (p)ppGpp^0
strains alone, further supporting that (p)ppGpp and the Y-complex act independently on the
regulation of ribosome levels and translation (**Fig. S8C**). Mutants of the Y-complex are also
strongly affected by other proteotoxic stresses, such as osmotic and oxidative stress, but not
by ethanol (**Fig. S8D**), which further demonstrates the importance of controlling ribosome
levels under proteotoxic stress conditions.

We observed that the Y-complex globally controls the RNase Y-mediated maturation of
transcripts and is involved in adjusting ribosome levels to the growth state, allowing the
endonuclease RNase Y to initiate the processing and degradation of rRNA and r-protein
transcripts. The absence of the Y-complex results in an abundance of idle ribosomes, which
interfere with the downregulation of translation and increase sensitivity to protein folding
stresses.

Discussion

Bacterial adaptation to environmental challenges depends on their ability to quickly change gene expression. Beyond the well-known regulation of transcription initiation, we observed here that post-transcriptional mechanisms, which influence RNA stability or translation efficiency, also contribute to the rapid and tailored cellular response. In this context, the control of RNases plays a critical role, since we show that the fitness of *B. subtilis* during heat shock is critically dependent on certain non-essential RNases and the Y-complex, which controls the substrate selection of the quasi-essential endonuclease RNase Y[26] (**Fig. 1B-C**). Together, our findings demonstrate that the Y-complex – RNase Y system is a regulatory hub that modulates gene expression and the cell's translational capacity during stress response.

Our RNA-seq analyses reveal that the Y-complex may regulate transcripts involved in a wide range of metabolic pathways, developmental processes, stress responses, and translation (**Fig. 2G, Fig. S4A-B, Table S2, Table S6-7**). RNA processing by the Y-complex-guided RNase Y can create new transcript isoforms[26] or modify regulatory elements such as UTRs[26,31](**Fig. 2F**, **Fig. S3**), thereby tailoring gene expression to specific growth conditions. This wide regulatory scope of the Y-complex could explain the known pleiotropic phenotypes in these mutants, like decreased sporulation, competence, and defective biofilm formation[38–40].

Among these diverse functions in RNA processing, we discovered that the Y-complex initiates ribosome degradation at the end of exponential growth by directing RNase Y to cleave 23S and 16S rRNA (**Fig. 3A-B**, **Fig. S7A**), thereby decreasing the levels of ribosomes. This finding is consistent with previous suggestions that RNase Y can degrade rRNA in aging *B. subtilis* spores[41].

The exact nature of how the Y-complex (YmcA, YlbF, and YaaT) associates with RNase Y is an active subject of research. Recent evidence indicates that YmcA and YlbF transfer YaaT to RNase Y, forming a stable, membrane-localized YaaT-RNase Y complex[42]. However, the deletion of any of the Y-complex members leads to a strong decrease in the processing of target mRNA[26], indicating that all Y-complex proteins are equally indispensable for RNA processing. Based on the dynamic localization of YmcA and YlbF between the cell membrane and the cytosol[43], we propose that these Y-proteins may also recognize substrates, including ribosomes, and direct them to RNase Y for cleavage.

The two $[4Fe-4S]^{+2}$ clusters in the Y-complex[44] likely act as sensors for cellular growth, energy, and stress, affecting how the complex controls RNase Y. This regulatory complex is localized at the cell membrane, where RNase Y forms a dynamic degradosome-like network with other RNases, helicases, and glycolytic enzymes[45–47]. This spatial co-localization may enable a

sequential degradation process: the Y-complex-guided RNase Y performs initial cleavage on targets like rRNA, making fragments susceptible to further degradation by exonucleases such as RNase R, PNPase, and RNase J. This suggests complex spatial control of RNA processing and ribosome decay, separate from transcription and translation in the cytoplasm (**Fig. 6**).

Most of the cleavages we observed in rRNA were in the 23S (**Fig. 3A**); however, the Y-complex and RNase Y can also initiate 16S degradation in the late exponential phase and during heat shock (**Fig. 3A**, **Fig. S7A**, **Fig. S8A**). Recently, it was observed that the exoribonuclease RNase R alone can degrade 16S[48], suggesting that endonucleolytic cleavages might not be strictly necessary for 16S degradation. Since RNase R is induced as part of the general stress response[49], it suggests a possible stress-specific and Y-complex-independent degradation pathway for 16S rRNA.

Complete ribosome decay requires the concurrent degradation of r-proteins (**Fig. 6**). These proteins might be passively released from the ribosome into the cytoplasm during the degradation of the rRNA. Since this would also change their interaction with rRNA, they might misfold and aggregate, and can subsequently be degraded by AAA+ protease complexes. Alternatively, these AAA+ protease complexes could be actively recruited to the site of ribosome degradation to simultaneously extract and degrade the r-proteins along with rRNA. AAA+ protease complexes such as ClpCP with McsB[50–52] or ClpXP, which functions in ribosome-associated quality control in *B. subtilis*[53,54], could be involved in the proteolysis of the released r-proteins.

Strikingly, our work reveals a second tier of regulation of ribosome levels. The Y-complex additionally processes transcripts encoding r-proteins and other translation factors (**Fig. 2G**), such as BipA, EF-Tu, and EF-G (**Fig. S3E-F**), potentially regulating their translation or leading to their decay. This is supported by the findings that the Y-complex is necessary to process some r-protein transcripts at the mid-logarithmic phase[26], and RNase Y depletion in this growth phase leads to an increased abundance of transcripts of r-proteins, such as *rplS, rpsO, rpmA, rpsD, rpsB,* and *rplU*[55].

The control of translation is an important component of many stress response mechanisms. During nutrient starvation, downregulating translation allows the reallocation of resources to biosynthetic pathways[56]. During proteotoxic stresses, it could relieve the overwhelmed PQC system by reducing the synthesis of new proteins that could potentially aggregate[57]. The stress transcription factor Spx and the alarmones (p)ppGpp regulate ribosome levels in *B. subtilis*, and the lack of both impairs growth at 50°C[7]. Our work establishes a mechanistic link between the regulation of ribosome levels through synthesis and decay. The severe growth defect observed in cells lacking both (p)ppGpp and the Y-complex indicates that these systems are important for transitioning from exponential to stationary phase. Furthermore, the absence of

their combined regulation of translation results in elevated ribosome levels and increased
accumulation of protein aggregates. (**Fig. S8C**). In times of nutrient limitation, the collective
function of (p)ppGpp and the Y-complex in regulating ribosome levels by inhibiting their
synthesis and initiating degradation allows resource allocation to other biosynthetic pathways
(**Fig. 6**).

These strategies are not exclusive to *B. subtilis*, as different organisms use various
mechanisms to degrade ribosomes. These must specifically recognize the ribosome or its
subunits to disassemble them and target rRNA and r-proteins for degradation. For example,
E. coli cells transitioning into the stationary phase also initiate ribosome degradation by the
membrane-associated endonuclease RNase E[58,59], which is functionally equivalent to RNase
Y in *B. subtilis*[60]. RNase E forms the scaffold of the membrane-associated degradosome
network in *E. coli*[61]. This could allow for the combined activity of endo- and exonucleases,
facilitating the full degradation of the ribosome[60,62]. In addition, it was observed that *E. coli* r-
proteins are degraded by the Lon AAA+ protease during starvation[63]. In eukaryotic cells,
nutrient starvation also triggers ribosome degradation via the ubiquitin-proteasome system or
via specialized autophagy pathways, such as ribophagy[64]. Recent studies have described how
stress- or starvation-induced degradation of the 40S ribosome involves the ubiquitylation of
proteins S3 and S5. This allows RIOK3 binding, leading to the decay of 18S rRNA by unknown
RNases[65–67].

In summary, we identified the Y-complex as a key regulator of ribosome abundance in *B.*
subtilis. The Y-complex regulates the translational activity by triggering ribosome decay and
processing transcripts for new ribosomal components. Our findings contribute to the
understanding of ribosome degradation by identifying a direct pathway and illustrating how this
process is integrated with other stress response mechanisms to control cellular growth and
survival. Further studies are required to understand the molecular signals that trigger ribosome
degradation and how the decay of rRNA and r-proteins is temporarily and spatially coordinated.

Materials and methods

Growth conditions

Unless stated the contrary, strains (**Table S10**) were streaked on LB plates (5 g/l yeast extract, 10 g/l tryptone agar, 10 g/l NaCl, 1.5% agar) containing the required antibiotics for selection (1 μg/ml erythromycin + 25 μg/ml lincomycin, 150 μg/ml spectinomycin or 10 μg/ml kanamycin) and incubated overnight at 37°C. A single colony was inoculated in 3 ml of LB medium and grown overnight at 37°C and 180 rpm, then used as the inoculum for subsequent experiments.

Strain construction

For constructing the strains $\Delta ymcA::erm^R$ and $\Delta yaaT::erm^R$, the loci containing the resistance cassette were PCR-amplified from the respective strains of the BKE library[20] and transformed in *B. subtilis* WT. The deletions of $\Delta rel::kan^R$ and $\Delta ylbF::erm^R$ were constructed as described in Ref.[20]. For promoter activity reporter strains, the *gfp*(mut2) gene was cloned downstream of the promoters of interest, using the integrative plasmid pDR111 as a backbone.

For transformation, 600 μl of an overnight culture of the receptor strains were diluted in 5 ml competence medium (17.5 g/l K_2HPO_4, 7.5 g/l KH_2PO_4, 2.5 g/l $(NH_4)_2SO_4$, 1.25 g/l tri-sodium citrate x2 H_2O, 0.5% glucose, 7 mM $MgSO_4$, 0.1 mg/ml L-tryptophan, 0.02% casamino acids, 0.22 g/ml ammonium iron citrate) and grown at 37°C, 180 rpm, for 3 hours. Subsequently, 5 ml of starvation medium (17.5 g/l K_2HPO_4, 7.5 g/l KH_2PO_4, 2.5 g/l $(NH_4)_2SO_4$, 1.25 g/l tri-sodium citrate x2 H_2O, 0.5% glucose, 7 mM $MgSO_4$) were added and grown for 2 more hours. One ml of this culture was incubated with the DNA to be transformed and grown for 2 hours at 37°C and 180 rpm. Positive clones were selected on LB agar containing the required antibiotics. Genetic modifications were verified using PCR and DNA sequencing.

Genetic screening during heat shock

The BKE library[20] was replicated on LB agar plates containing 1 μg/ml erythromycin. Liquid cultures were prepared in 200 μl LB medium supplemented with 1 μg/ml erythromycin and grown for 42 hours at 30°C 200 rpm. All mutant strains were pooled and mixed for 15 minutes using a magnetic stirrer. The pooled culture was centrifuged at 5000 rpm for 10 minutes at 18°C. The pellet was resuspended in LB medium to an OD_{600} of 65 and mixed with an equal volume of 50% glycerol. The pooled library was aliquoted and kept at -80°C.

For each replicate, a tube of the pooled library was quickly unfrozen in a water bath at 30°C and used to inoculate 75 ml LB medium in 500 ml flasks to a starting of OD_{600} 0.05. The culture was grown in a water bath at 30°C, 180 rpm, until OD_{600} 0.4. Fifty ml of culture were split into two 250 ml flasks. One flask was transferred to 50°C, 180 rpm for heat shock. The other flask was used as a control sample and was kept at 30°C. One ml samples were collected for gDNA

purification at 0 and 2 hours post temperature shift. Samples were pelleted for 2 minutes at 13,000 rpm and washed once with 500 µl PBS. gDNA was purified using the Wizard® Genomic DNA Purification Kit (Promega, A1120) and stored at -20°C until further processing.

The strain-specific barcodes were amplified using indexed primers (**Table S7**) with the Collibri Library Amplification Master Mix (ThermoFisher, A38539050) according to the manufacturer's instructions, with 100 ng of gDNA as input, for 22 cycles. Barcodes were gel-purified, quantified, pooled equimolarly, and sequenced using the Illumina NovaSeq2 single-read method at the Sequencing Core Facility of the Max Planck Institute for Molecular Genetics (Berlin, Germany).

To calculate the fitness score of each deletion mutant from barcode abundance, BEAN-counter v2.6.1[68] was used with the following settings. A fixed amplicon read structure was used with an index at position 0, the common primer sequence GCAGGCGAGAAAGGAGAG at position 9, and the mutant barcode at position 27 of read 1. All other settings were used with default values. The sample before heat shock served as a reference to which all other time points were compared. All comparisons were performed with three biological replicates.

Fitness data was further analyzed using the R statistical computing language (v4.3.3). For annotation, protein-related information was fetched using the Uniprot REST API (29 February 2024). Average fitness per gene, condition, and time point was calculated as the arithmetic mean of the fitness score from three biological replicates. Differential fitness ΔF, e.g. for comparison of treatment *versus* control at a particular time point, was calculated as $\Delta F = F_{treatment} - F_{control}$. A two-sided Student's t-test was performed to evaluate statistical significance for each gene, and the obtained *p*-value was corrected using the Benjamini-Hochberg procedure[69].

Growth curves

Twenty-five ml of LB medium were inoculated with an overnight culture to a starting OD_{600} 0.05 in 250 ml Erlenmeyer flasks. Cultures were grown at 37°C 180 rpm in a water bath. OD_{600} was monitored every 30 minutes. The average and standard deviation of three biological replicates were calculated.

Heat shock survival

Cultures were inoculated in 25 ml LB medium to OD_{600} 0.05 and incubated in a water bath at 37°C 180 rpm. At OD_{600} 0.3 – 0.4, the cultures were exposed to a heat shock at 54°C for 2 hours, and samples for viability measurement were obtained every 30 minutes. Viability was measured by spotting 4 µl of 1:10 serial dilutions in PBS of each sample on LB agar. The plates were incubated overnight at 30°C, and the colony-forming units (CFU) were counted. The

CFU/ml was calculated for each sample, and relative viability was determined using samples
before heat shock as a reference.

<u>Isolation of protein aggregates</u>

Twenty-five ml of culture were harvested before and 15 minutes after heat shock at 54°C by
centrifugation for 5 minutes at 10,000 rpm. The pellets were washed once with buffer A (100
mM HEPES pH 8.0, 150 mM NaCl) and lysed by bead-beating in buffer B (100 mM HEPES
pH 8.0, 150 mM NaCl, cOmplete EDTA-free protease inhibitor) using 0.1 mm glass beads.
Afterwards, samples were centrifuged at 1000 x g for 2 minutes to obtain the cell extract. The
soluble fraction was separated from protein aggregates by centrifugation for 15 minutes at
11,000 x g at 4°C. To remove membrane proteins, the aggregates were resuspended in buffer
B containing 1% Triton X-100 and incubated for 1 hour at 4°C. Protein aggregates were
recovered by centrifugation at 11,000 x g, 4°C for 15 minutes and resuspended and incubated
as described above. Protein aggregates were washed again with buffer B + 0.5% Triton X-
100, and the detergent was removed in a final wash using buffer B. Pellets containing protein
aggregates were resuspended in buffer B + 2% SDS by boiling for 5 minutes at 95°C. Protein
concentration was measured using the Rapid Gold BCA Protein-Assay-Kit (Pierce™, A53225).
The percentage of aggregated protein was calculated to the total protein content in the cell
extract.

<u>Spot test</u>

Overnight cultures were diluted 1:1000 in LB medium without antibiotics and incubated at 37°C
180 rpm for 7-8 hours. The cultures were diluted to OD_{600} 1 in fresh LB medium in a 96-well
plate. Seven serial 1:10 dilutions were prepared in 1x PBS. Four μl of each dilution was spotted
on LB agar containing different stressors (concentration indicated in the figures). Plates were
freshly prepared before use.

<u>RNA purification</u>

Samples for RNA purification were collected at the indicated time points: exponential phase
(OD_{600} ~ 0.3), late exponential phase (OD_{600} ~ 1) and transition phase (100 minutes after late
exponential phase). Rifampicin was added to a final concentration of 100 μg/ml for 10 minutes.

Samples were harvested by mixing with the same volume of cold 1:1 acetone:ethanol and
centrifugated at 9,000 rpm for 5 minutes, 4°C. RNA purification was performed according to
Ref.[27] with slight modifications: The pellet was washed twice in 1 ml TE-sucrose buffer (50 mM
Tris-HCl pH 8.0, 10 mM EDTA, 25% sucrose) and centrifugated at 10,000 rpm for 5 minutes,
4°C. Cells were resuspended in 130 μl lysis buffer (20 mM Tris-HCl pH 8.0, 50 mM EDTA, 20
% sucrose) supplemented with 3.2 mg/ml lysozyme and incubated on ice for 5 minutes. Two
hundred μl of lysis executioner (2% SDS, 1 mg/ml Proteinase K) were added to the samples

and incubated at 95°C for 1.5 minutes. Samples were mixed with 1 ml of TRIzol (Ambion) and

incubated at room temperature. After 5 minutes, 250 µl chloroform was added to each sample

and mixed by vortexing. After 10 minutes at room temperature, the samples were centrifuged

for 10 minutes at 11,000 rpm. The RNA was isopropanol-precipitated at -20°C overnight. RNA

was pelleted at 11,000 rpm for 20 minutes at 4°C, washed with 1 ml 70% ethanol followed by

another centrifugation at 13,000 rpm for 10 minutes at 4°C. The pellet was air-dried and

resuspended in 50 µl nuclease-free water.

ISCP RNA sequencing

RNA sequencing was performed as previously described[70]. In brief, total RNA was treated with

TURBO DNase (Ambion, AM2238), depleted of ribosomal RNA using the riboPOOL rRNA

depletion kit for *B. subtilis* (siTOOLS; two reactions with 4.5 µg each) and purified using the

RNA Clean & Concentrator-5 kit (RCC-5; Zymo Research). RNA 5' triphosphates were

converted to monophosphates using RppH (NEB), and RNA was recovered using

phenol:chloroform:isoamyl alcohol extraction followed by ethanol precipitation. RNA ends were

repaired using T4 Polynucleotide Kinase (Thermo Scientific), before RNA was purified using

the RCC-5 kit and fragmented using an M220 Focused-ultrasonicator (Covaris) as described

by the manufacturer but with a treatment time of 100 s (microTUBE AFA Fiber Snap-Cap). After

purification using the RCC-5 kit, sequencing libraries were prepared from the recovered RNA

using the NEXTflex Small RNA-Seq Kit v3 (Bioo Scientific) according to the manufacturer's

protocol (including step G). Libraries were purified as described in step I of the NEXTflex Rapid

Directional qRNA-Seq Kit (Bioo Scientific) using Agencourt AMPure XP Beads (Beckman

Coulter; 0.9x bead-to-sample ratio). Sequencing was performed on an Illumina NovaSeq 6000

in paired-end mode with a read length of 100 bp at the Sequencing Core Facility of the Max

Planck Institute for Molecular Genetics (Berlin, Germany).

Processing of RNA sequencing data

FastQC (v0.11.5) was used to assess the quality of the data. Reads were filtered with a

minimum quality score of 10 and a length of at least 18 nt, cleaned from adapter sequences

using Cutadapt (v3.5)[71], and mapped against the *B. subtilis* reference genome (NC_000964.3)

using STAR (v2.7.10a_alpha_220314)[72] in 'random best' and 'end to end' modes. Resulting

BAM files were sorted and indexed with Samtools (v1.9)[73], and PCR duplication artefacts were

removed using UMI Tools (v1.1.0)[74]. Gene counts of annotated RefSeq genes were

determined with featureCounts (v2.0.3)[75] and differentially expressed genes (DEG) were

calculated with three replicates per condition and time point using DESeq2 (v1.24.0)[76]. *p*-

values were corrected for multiple testing using the Benjamini-Hochberg method[69]. To call

DEGs, a threshold of adjusted *p*-value < 0.05 and absolute log2 fold change > 1 was used.

Genome coverage profiles (total, 5' and 3' ends) were generated using a custom script utilizing

the HTSeq library (v2.0.2)[77]. We performed the aforementioned analysis steps using a

customized pipeline implemented in Snakemake (v7.14.0)[78].

Identification of Y complex cleavage sites

The Y-complex-dependent cleavage positions were identified following the previously

published procedure[70]. In brief, the genome coverage data was prefiltered with a count per

million (cpm) value ≥0.05, and only the RNA ends displaying a cpm ≥ 5 were further analyzed.

We carried out differential expression analysis of normalized 5′ and 3′ read ends by comparing

WT vs. $\Delta ymcA$ at the late exponential and transition phase. Y-complex-dependent ends were

identified using edgeR (v3.28.0)[79] with absolute log2 fold change ≥ 1 and FDR < 0.05, and kept

only if present in both comparisons. These results were further filtered with additional

parameters, i.e., the "proportion of ends (cleavage ratio)" and the "ratio of WT and $\Delta ymcA$

proportion of ends" as was previously described[70]. Moreover, to reduce the potential set of

false positive processing sites even further, we applied an additional cut-off on the "proportion

of ends" parameter, i.e., we kept only processing sites with a cleavage ratio of the reference

strain (WT) ≥ 95%-percentile of the cleavage ratio of the mutant strain.

Impact of processing sites on transcript and operon expression

To examine the impact of cleavage sites on transcript and operon expression, as well as other

genomic features such as untranslated regions (UTRs) and regulatory elements, we calculated

transcript per million (TPM) values from variance-stabilized data for each annotated RefSeq

gene (transcript level) and for each operon as defined in the BSGatlas (v1.1)[80] using the

DESeq2 package and a custom Python script. Furthermore, all Y-complex-dependent

cleavage sites were mapped to various genomic features, including coding regions, UTRs,

terminator regions, and riboswitches, as defined in the BSGatlas[80] (**Table S2**).

Processing site sequence logo

533 The sequence logos were generated with Logomaker (v0.8)[81], specifically for 5' and 3' end U-

534 type cleavage sites. For this, we extracted 30 nt sequences centered at the cleavage site and

535 calculated the logos with a GC content of 43.5%.

RNA secondary structure prediction

The RNA structure surrounding Y-complex-dependent cleavage sites was estimated by

calculating the MFE (ΔG in kcal/mol) utilizing RNAfold (v2.6.4)[82]. This analysis employed a

sliding window approach with 50 nt sequences, focusing on a 200 nt region centered around

the position of interest. The average MFE at each nucleotide was then calculated. To calculate

and visualize the background folding energy, we randomly draw potential processing sites from

the genome determined with the corresponding 5' or 3' end motif and calculate the average
MFE at each nucleotide as described above. Potential processing sites were drawn 50 times
independently.

<u>Overrepresentation analysis</u>

Annotations of operons were obtained from BSGatlas[80] and gene categories from Subtiwiki
(24 July 2023)[83]. The overrepresentation analysis was performed using the R statistical
computing language (v.4.2.2) and the Bioconductor package Category (v.2.64.0).

<u>Northern blot</u>

Biotinylated RNA probes were prepared using the HighYield T7 Biotin11 RNA Labeling Kit
(UTP-based) (Jena Bioscience, RNT-101-BIOX) from PCR amplified fragments containing the
promoter for the T7 RNA polymerase (**Table S11**). Probes were purified using the Monarch®
Spin RNA Cleanup Kit (NEB, T2040) according to the manufacturer's instructions. Probes were
designed taking into consideration the predicted cleavage sites.

One and a half µg of total RNA was separated on denaturing agarose gels (1% agarose,
20 mM MOPS, 5 mM NaOAc, 1 mM EDTA, 6.6% formaldehyde, pH 7.0) in 1x AGE-NB
Running Buffer (20 mM MOPS, 5 mM NaOAc, 1 mM EDTA, 0.74% formaldehyde, pH 7.0).
The gel was washed 3 times with milli-Q water and once with 10x SSC buffer, and the
nitrocellulose membrane was washed once with milli-Q water and once with 10x SSC. The
RNA was transferred using a vacuum blotter at 300 mbar covered with 10x SSC for 1.5 hours.
After transfer, the membrane was washed for 2 minutes with 6x SSC, once with water, and
dried and crosslinked at 254 nm 0.12 J twice. The membrane was developed with methylene
blue, and the staining was removed by washing with 0.2x SSC containing 1% SDS, followed
by washing with milli-Q water. The hybridization of 16S or 23S rRNA probes was performed
using the North2South Chemiluminescent Hybridization and Detection Kit (Thermo) and
North2South Hybridization Stringency Wash Buffer (2X) according to the manufacturer's
instructions. Blots were developed in a Vilber imaging system.

<u>RT-qPCR</u>

Twenty µg of total RNA were treated with TURBO DNase (Ambion, AM2238) and purified using
the Monarch® Spin RNA Cleanup Kit (NEB, T2040). Subsequently, cDNA was prepared using
the LunaScript RT Mastermix (NEB, E3025) and 2.5 µg of DNase-treated RNA as input.
Quantitative PCR was performed using the Luna Universal qPCR Master Mix (NEB, M3003)
in a QuantStudio 5 Real-Time PCR System (Applied Biosystems). To calculate changes in *gfp*
transcription, samples from the late exponential and transition phases were compared to those

from the exponential phase, with *pcp* serving as a reference gene using the Pfaffl method[84].

Primers used for RT-qPCR are listed in **Table S11**.

Ribosome sedimentation in sucrose gradients

Cultures of WT and $\Delta ymcA$ were grown until the exponential phase (OD_{600} ~ 0.3), late

exponential (OD_{600} ~ 1), and transition phase (100 minutes after the late exponential phase).

Chloramphenicol was added at a concentration of 100 µg/ml 2 minutes prior to sample

collection. Fifty ml of culture were pelleted at 4,500 rpm for 10 minutes at room temperature.

Pellets were transferred to ice and resuspended in 2 ml of lysis buffer (20 mM Tris-HCl pH 8.0,

100 mM NH_4Cl, 10 mM $MgCl_2$, 0.4% Triton X-100, 0.4% NP-40, 1 mM chloramphenicol, 100

U/µl DNase I), and flash-frozen. Cells were lysed under cryogenic conditions using 5 cycles of

mix-milling at 30 Hz for 1 minute. Lysates (240 µg RNA) were laid over a 10-50% sucrose

gradient in 20 mM Tris-HCl pH 8.0, 100 mM NH_4Cl, 10 mM $MgCl_2$, 0.5 mM DTT and resolved

at 35,000 rpm for 3 hours at 4°C. Gradients were analyzed using a piston fractionator (Biocomp

instruments), and absorbance at 260 and 280 nm was monitored.

Sample preparation for MS-based proteomics

Samples were collected at the indicated time points: late exponential phase (OD_{600} ~ 1) and

transition phase (100 minutes after the late exponential phase). For the proteomic changes

during heat shock, cultures were grown at 37°C until OD_{600} ~ 0.3 and shifted to 54°C. Samples

were collected at 0, 15, 30, and 60 minutes after heat shock. Ten ml of culture were collected

by centrifugation at 4,500 rpm for 10 minutes, and the pellet was stored at -80°C until further

processing.

Cell pellets were resuspended in 1 ml of cold PBS and centrifuged at 13,000 rpm for 5 minutes

at 4°C. The pellet was washed once more with PBS, resuspended in 150 µL of lysis buffer

(100 mM HEPES pH 8.0, 2% SDS, 20 mM TCEP, 60 mM CAA), and lysed using bead beating.

Cell debris was discarded by centrifuging at 13,000 rpm for 5 minutes at 4°C. The supernatant

was boiled at 95°C for 5 minutes, cooled down to room temperature, and the protein

concentration was estimated using the Rapid Gold BCA protein assay kit (Pierce). Samples

were flash-frozen in liquid nitrogen and stored at -80°C until further processing.

All samples were subjected to SP3 sample preparation[85] on an Agilent BRAVO liquid handling

robot. Ten µg of a 1:1 mixture of hydrophilic and hydrophobic carboxyl-coated paramagnetic

beads (SeraMag, #24152105050250 and #44152105050250, GE Healtcare) were added for

each µg of protein. Protein binding was induced by the addition of acetonitrile to a final

concentration of 50% (v/v). Samples were incubated at room temperature for 10 minutes. The

tubes were placed on a magnetic rack, and beads were allowed to settle for 3 minutes. The

supernatant was discarded, and beads were rinsed three times with 200 µL of 80% ethanol

without removing the tubes from the rack. Beads were resuspended in digestion buffer containing 50 mM triethylammonium bicarbonate and both Trypsin (Serva) and Lys-C (Wako) in a 1:50 enzyme-to-protein ratio. Protein digestion was performed for 14 hours at 37°C in a PCR cycler. Afterward, the supernatant was recovered and dried down in a vacuum concentrator.

Proteomics using Mass spectrometry

All samples obtained during the late exponential and transition phase were analyzed on an Orbitrap Exploris (Thermo Scientific) equipped with a FAIMS Pro device and coupled to a 3000 RSLC nano UPLC (Thermo Scientific). Samples were loaded on a PepMap trap cartridge (300 µm i.d. x 5 mm, C18, Thermo) with 2% acetonitrile and 0.1% TFA at a flow rate of 20 µl/min. Peptides were separated over a 50 cm analytical column (Picofrit, 360 µm O.D., 75 µm I.D., 10 µm tip opening, non-coated, New Objective) that was packed in-house with Poroshell 120 EC-C18, 2.7 µm (Agilent). Solvent A consists of 0.1% formic acid in water. Elution was carried out at a constant flow rate of 300 nl/min using a 60-minute method: 8-30% solvent B (0.1% formic acid in 80% acetonitrile) within 23 minutes, 30-45% solvent B within 3 minutes, 45-98% buffer B within 0.5 minutes, followed by column washing and equilibration. Data acquisition on the Orbitrap Exploris was carried out using a data-dependent acquisition (**DDA**) method in positive ion mode. MS survey scans were acquired from 375-1500 m/z in profile mode at a resolution of 60,000. AGC target was set to 200% at a maximum injection time of 25 ms. Peptides with charge states 2-6 were isolated within a window of 1.2 m/z and subjected to HCD fragmentation at a normalized collision energy of 27%. The MS2 AGC target was set to 200%, allowing a maximum injection time of 22 ms. Product ions were detected in the Orbitrap at a resolution of 15,000. Precursors were dynamically excluded for 20 s. The cycle time was set to 1 s for each of the two FAIMS voltages of −45 and −65 V.

LC-MS/MS analysis of the heat shock samples was performed on an Ultimate 3000 UPLC coupled to a Lumos hybrid mass spectrometer (Thermo Scientific). Peptides were loaded onto a PepMap trap cartridge (300 µm i.d. x 5 mm, C18, Thermo) with 2% acetonitrile and 0.1% TFA at a 20 µl/min flow rate. Peptides were separated over a 50 cm analytical column (Picofrit, 360 µm O.D., 75 µm I.D., 10 µm tip opening, non-coated, New Objective) that was packed in-house with Poroshell 120 EC-C18, 2.7 µm (Agilent). Solvent A consists of 0.1% formic acid in water. Elution was carried out at a constant flow rate of 250 nl/min using a 60-minute method: 8-30% solvent B (0.1% formic acid in 80% acetonitrile) within 23 minutes, 30-45% solvent B within 3 minutes, 45-98% buffer B within 0.5 minutes, followed by column washing and equilibration.

The mass spectrometer was operated in data-independent acquisition (**DIA**) and positive ionization mode. MS1 full scans (375–1500 m/z) were acquired with a resolution of 60,000, a

normalized automatic gain control target value of 120%, and a maximum injection time set to
55 ms. Peptide fragmentation was performed using higher energy collision-induced
dissociation and a normalized collision energy of 30%. MS2 spectra were acquired with a
resolution of 30,000 (first mass 100 m/z) with 8 m/z DIA windows between 400 and 1200 m/z.
A normalized automatic gain control target value of 100%, and a maximum injection time of
54 ms.

<u>Proteomics data analysis</u>

For samples at the late exponential and transition phase, Raw files were processed with
MSFragger v3.5 using the default settings. Briefly, peak lists were extracted from raw files and
searched using a Uniprot *Bacillus subtilis* database (v220530, taxonomy ID 224308) and a
database containing sequences of common contaminants and decoys. Trypsin/P was set as
enzyme specificity, allowing a maximum of two missed cleavages. Peptide matches were
filtered to 1% FDR using PeptideProphet and Philosopher (v4.4.0).

For samples during heat shock, DIA-NN (DIA-NN v1.8.1) was employed for DIA data analysis.
The main DIA-NN report, "report.txt", was used for all calculations. Fasta file
(Bacillus_subtilis_Strain168_UP000001570.fasta) was digested with a maximum of one
missed cleavage. Peptide length was restricted from 7 to 30 peptides, and the precursor m/z
range was set from 300 to 1800. N-terminal methionine excision was enabled. Cysteine
carbamidomethylation was selected as a fixed modification, and methionine oxidation and N-
terminal acetylation as variable modifications. The maximum number of variable modifications
was set to two. All other parameters were at default settings, including the 1% precursor FDR
and enabled match between runs (MBR). The main DIA-NN report, "report.pg_matrix.tsv," was
used for all calculations.

The R statistical computing language was used for data processing (v4.4.1). The main
comparison of WT, $\Delta ymcA$, (p)ppGpp^{0,} and (p)ppGpp^0 $\Delta ymcA$ mutants was based on samples
from data-dependent acquisition (DDA) experiments. Raw peptide intensity values from label-
free quantitation were imported as tabular output from MSFragger. The MSstats package
(v4.12.1)[86] was used to convert raw intensities to an MSstats experiment. In this process,
peptide quantification data was log$_2$-transformed, median-normalized, and summarized to
protein abundance. All features mapping to a protein were used for quantification, Tukey's
median polish was used as a summarization strategy, and missing values were imputed using
MSstats' default strategy. Feature abundance and inferred protein quantity were visually
inspected using profile plots. Experimental groups were compared using MSstats'
groupComparison function, and the similarity of conditions and replicates was inspected using
principal component analysis (PCA, function *prcomp*). For annotation, protein-related
information was fetched from Uniprot using the Uniprot REST API (14 June 2024). In addition

to the DDA experiments, a time series of WT and Δ*ymcA* mutants was obtained using data-independent acquisition (DIA) proteomics. To quantify relative changes, the $\log_2$ fold change of each time point was calculated using the zero time point as a reference. To quantify absolute protein abundance for the ribosomal proteome sector, the coarse-graining strategy was used as in Ref.[87,88]. Briefly, two protein sectors (groups) were created containing all annotated proteins of ribosomal subunits. Raw MS1 intensities for all proteins of the respective group and time points were summed up and divided by the total sum of protein intensities, yielding the protein mass fraction per sector.

Data availability

Amplicon-seq and RNA-seq raw reads have been deposited at the European Nucleotide Archive (ENA) under accession number PRJEB90661. Raw proteomic data have been deposited at the PRIDE database under the accession number PXD065465 for DDA proteomics of late exponential and transition phase samples (Reviewer access in PRIDE (https://www.ebi.ac.uk/pride/login), Token: ijwEKpkL6sTq). The DIA proteomics raw data from the heat shock experiment can be accessed using the accession number PXD065462 (Reviewer access in PRIDE, Token: 8IRshuji8qRS).

Author contributions

K.T., K.D., and F.A.C.; Conceptualization. K.D., F.A.C., K.T., R.A.-B., T.F.W., and K.A.: Methodology. F.A.C., K.D., K.S., V.S., K.H., and F.K.: Investigation. K.D., F.A.C., R.A.-B., M.J., K.A., S.R., and K.T.: Formal analysis. F.A.C., K.D., R.A.-B., M.J.: Visualization. F.A.C., K.D., and K.T.: Writing – Original Draft. K.T. and E.C.: Funding acquisition. All authors: Writing – Review & Editing.

Acknowledgments

We thank Gert Bange (Philipps-Universität Marburg) and Rainer Nikolay (Max Planck Institute for Molecular Genetics) for critically reading the manuscript and providing valuable feedback. This work was funded by the Deutsche Forschungsgemeinschaft (Tu106/8, SPP1879 for K.D. and K.T., Leibniz Prize for E.C.) and the Max Planck Society (for E.C. and K.T.).

Figure 1. Genetic screening for genes involved in the heat shock response. A) Genetic screening experiment: the pooled library of deletion mutants (BKE) was grown at 30°C; when it reached an OD_{600nm} of 0.4, one fraction was exposed to 50°C while the other was maintained at 30°C. Genomic DNA (gDNA) was extracted from the library, and barcodes were amplified and sequenced using Next-Generation Sequencing. The fitness of each mutant at the mentioned temperatures and time points was calculated using BEANcounter. **B)** Relative fitness of each deletion mutant at 50°C (*y*-axis) and 30°C (*x*-axis) two hours post-temperature shift. Each point represents a deletion mutant, with mutants lacking RNase-related genes highlighted in magenta. The dotted lines indicate the range in which the relative fitness between the conditions differed by less than 2. Data represent the average of three biological replicates. A negative relative fitness value means that the abundance of the deletion mutant, compared to the rest of the population, was negatively affected by the treatment, while a positive value means an increased abundance in this condition. **C)** Changes in the viability (log_{10}CFU/ml) of WT and *ymcA, ylbF,* and *yaaT* deletion mutants to a severe heat shock at 54°C. Values were normalized to the viability at timepoint 0 min. The data represents the average ± standard error of three biological replicates.

Figure 2. Identification of Y-complex-dependent cleavages at late exponential and transition phase. A) Growth curves of WT, *ymcA*, *ylbF*, and *yaaT* deletion mutants in LB at 37°C. The blue bar marks the exponential phase. Arrows indicate sample collection for the ISCP method at the late exponential and transition phases. Three biological replicates are displayed with a smooth curve regression. **B)** Number of RNA 3' or 5' ends identified at the two growth phases. **C)** Fraction of identified 3' or 5' ends displaying U-type (unique) or S-type (stepped) profiles. **D)** Minimum free energy (MFE) was calculated on 50 nt windows for RNAs cleaved by the Y-complex with U-type ends (in magenta). Transcripts that share the same cleavage motif but are not processed were used as a control (in gray). rRNA genes were excluded from the analysis since they are highly structured. **E)** Fractions of ends identified in genes transcribed in polycistronic or monocistronic RNA. **F)** Count of RNA features where

Y-complex-dependent ends are detected at either growth phase. Annotations were obtained from
BSGatlas[80]. **G)** Overrepresentation analysis of pathways whose transcripts are cleaved by the Y-
complex during the late exponential phase. Operons were assigned to be involved in a pathway if they
code for one gene participating in such a pathway. As genes in the same operon can have different
functions, transcripts may be linked to multiple roles. Pathway annotations were retrieved from
SubtiWiki[83]. The color indicates the statistical significance of the enrichment. Biosyn = biosynthesis, acq
= acquisition, excl TCS = excluding two-component systems.

Figure 3. The Y-complex controls ribosome levels by initiating rRNA degradation. A) Number of cleavages detected by ISCP in *B. subtilis* rRNA during the late exponential and transition phase. **B)** Northern blot against 23S of WT and Δ*ymcA* cells at the late exponential phase, before and after treatment with 100 μg/ml rifampicin to stop transcription. The methylene blue-stained membrane is shown as a loading control. This image is representative of three biological replicates. **C)** Volcano plot of protein quantification of WT and Δ*ymcA* strains comparing transition to late exponential phase. The proteins composing the small (30S) and major (50S) ribosome subunits are highlighted in cyan and magenta, respectively. The data shows the average of four biological replicates. Activity of the promoter controlling **D)** *rrnJ* **(**rRNA) or **E)** the *rplK* (r-protein) transcription at late exponential and transition phase compared to the exponential phase in WT and Δ*ymcA*. The promoters were cloned controlling *gfp* transcription, which was measured using RT-qPCR. The *pcp* transcript was used as a reference. Individual values and the average ± standard deviation of three biological replicates and two technical replicates are shown. The statistical significance was tested using student's *t*-test. ns: *p*-value > 0.05. **F)** Ribosome sedimentation profile in 10-50% sucrose gradients of WT and Δ*ymcA* in exponential, late exponential, and transition phases. The 70S and 100S ribosomes are highlighted with a blue or red box, respectively. The data is representative of three biological replicates.

Figure 4. The alarmones (p)ppGpp and the Y-complex are important for controlling ribosome levels. A) Growth curve of (p)ppGpp⁰ and (p)ppGpp⁰ Δ*ymcA* strains in LB at 37°C. The blue bar marks the exponential phase. Three biological replicates are displayed with a smooth curve regression. **B)** Volcano plot of proteomic changes of (p)ppGpp⁰ and (p)ppGpp⁰ Δ*ymcA* cultures comparing transition to exponential phase. The proteins composing the small (30S) and major (50S) ribosome subunits are highlighted in cyan and magenta, respectively. The data shows the average of four biological replicates. **C)** Comparison of r-proteins downregulation in WT, Δ*ymcA,* (p)ppGpp⁰, and (p)ppGpp⁰ Δ*ymcA* during the transition phase compared to the late exponential phase. The boxplot represents the interquartile range (IQR) and the median in the center. Whiskers show the variability outside quartile 1 (Q1) and Q3 and were calculated as Q1-1.5*IQR and Q5+1.5*IQR, respectively. Proteins are highlighted like in **B**. Alternative r-proteins were excluded from the analysis. The statistical significance was tested using paired Wilcoxon test. ****: p-value ≤ 0.0001. **D)** Proteome fraction (in %) of r-proteins mass, measured by mass spectrometry. The r-protein mass fraction was calculated for the indicated strains. Individual values and the average ± standard deviation of three biological replicates are shown. The statistical significance was tested using student's t-test. ns: p-value > 0.05, **: p-value ≤ 0.01, ***: p-value ≤ 0.001.

Figure 5. Control of ribosome levels and protein homeostasis in *ymcA* mutant during heat shock.
A) Northern blot against 23S of WT and Δ*ymcA* cells at exponential phase before or after a heat shock.
The methylene blue-stained membrane is shown as a loading control. This image is representative of
three biological replicates. **B)** Proteome fraction (in %) of r-proteins mass, measured by mass
spectrometry. Individual values and the average ± standard deviation of three biological replicates are
shown. The statistical significance was tested using student's *t*-test. *: *p*-value ≤ 0.05, **: *p*-value ≤ 0.01.
C) Induction of CtsR and HrcA regulons after heat shock (50°C) in WT and Δ*ymcA*. Proteins were
measured by mass spectrometry. The color bar shows the $\log_2$ fold change compared to the control
before heat shock. **D)** Percentage of aggregated proteins before and after a 15 min heat shock at 54°C
in WT and Δ*ymcA*. Individual values and the average ± standard deviation of three biological replicates
are shown. The statistical difference was tested using student's *t*-test. *: *p*-value ≤ 0.05, **: *p*-value ≤
0.01, ***: *p*-value ≤ 0.001.

Figure 6. Control of ribosome levels by (p)ppGpp and the Y-complex. The alarmone (p)ppGpp reduces the biogenesis of new ribosomes by indirectly inhibiting the transcription of rRNA and r-proteins mRNA. (p)ppGpp inhibits GTPases that are involved in new ribosome assembly. Additionally, (p)ppGpp reduces translation by binding to specific GTPases. The Y-complex can initiate the degradation of ribosomes, which need to be transported to the membrane by unknown mechanisms. Exonucleases and AAA+ protease complexes might assist in ribosome degradation for the complete degradation of rRNA and r-proteins, which could partially replenish the nucleotide and amino acid pools during starvation.

References

1. Hecker, M., Schumann, W., and Völker, U. (1996). Heat-shock and general stress response in
Bacillus subtilis. Mol. Microbiol. *19*, 417–428. https://doi.org/10.1046/j.1365-2958.1996.396932.x.

2. Mogk, A., Homuth, G., Scholz, C., Kim, L., Schmid, F.X., and Schumann, W. (1997). The GroE
chaperonin machine is a major modulator of the CIRCE heat shock regulon of *Bacillus subtilis*.
EMBO J. *16*, 4579–4590. https://doi.org/10.1093/emboj/16.15.4579.

3. Krüger, E., and Hecker, M. (1998). The first gene of the *Bacillus subtilis clpC* operon, *ctsR*,
encodes a negative regulator of its own operon and other Class III heat shock genes. J. Bacteriol.
180, 6681–6688. https://doi.org/10.1128/jb.180.24.6681-6688.1998.

4. Elsholz, A.K.W., Michalik, S., Zühlke, D., Hecker, M., and Gerth, U. (2010). CtsR, the Gram-
positive master regulator of protein quality control, feels the heat. EMBO J. *29*, 3621–3629.
https://doi.org/10.1038/emboj.2010.228.

5. Krüger, E., Witt, E., Ohlmeier, S., Hanschke, R., and Hecker, M. (2000). The Clp proteases of
Bacillus subtilis are directly involved in degradation of misfolded proteins. J. Bacteriol. *182*, 3259–
3265. https://doi.org/10.1128/jb.182.11.3259-3265.2000.

6. Alver, R., Hantke, I., Cornejo, F.A., Gunka, K., Rämisch, S., Molière, N., Charpentier, E., and
Turgay, K. (2023). The control of protein arginine phosphorylation facilitates proteostasis by an
AAA+ chaperone protease system. Preprint at bioRxiv, https://doi.org/10.1101/2022.09.15.508104
https://doi.org/10.1101/2022.09.15.508104.

7. Schäfer, H., Beckert, B., Frese, C.K., Steinchen, W., Nuss, A.M., Beckstette, M., Hantke, I.,
Driller, K., Sudzinová, P., Krásný, L., et al. (2020). The alarmones (p)ppGpp are part of the heat
shock response of *Bacillus subtilis*. PLOS Genet. *16*, e1008275.
https://doi.org/10.1371/journal.pgen.1008275.

8. Clay, K.J., Yang, Y., Clark, C., and Petrascheck, M. (2023). Proteostasis is differentially
modulated by inhibition of translation initiation or elongation. eLife *12*, e76465.
https://doi.org/10.7554/eLife.76465.

9. Cherkasov, V., Hofmann, S., Druffel-Augustin, S., Mogk, A., Tyedmers, J., Stoecklin, G., and
Bukau, B. (2013). Coordination of translational control and protein homeostasis during severe
heat stress. Curr. Biol. *23*, 2452–2462. https://doi.org/10.1016/j.cub.2013.09.058.

10. Schramm, F.D., Schroeder, K., and Jonas, K. (2020). Protein aggregation in bacteria. FEMS
Microbiol. Rev. *44*, 54–72. https://doi.org/10.1093/femsre/fuz026.

11. Schäfer, H., Heinz, A., Sudzinová, P., Voß, M., Hantke, I., Krásný, L., and Turgay, K. (2018). Spx,
the central regulator of the heat and oxidative stress response in *B. subtilis* , can repress
transcription of translation-related genes. Mol. Microbiol., mmi.14171.
https://doi.org/10.1111/mmi.14171.

12. Nakano, S., Erwin, K.N., Ralle, M., and Zuber, P. (2005). Redox-sensitive transcriptional control
by a thiol/disulphide switch in the global regulator, Spx. Mol. Microbiol. *55*, 498–510.
https://doi.org/10.1111/j.1365-2958.2004.04395.x.

13. Runde, S., Molière, N., Heinz, A., Maisonneuve, E., Janczikowski, A., Elsholz, A.K.W., Gerth, U.,
Hecker, M., and Turgay, K. (2014). The role of thiol oxidative stress response in heat-induced
protein aggregate formation during thermotolerance in *Bacillus subtilis*. Mol. Microbiol. *91*, 1036–
1052. https://doi.org/10.1111/mmi.12521.

14. Rochat, T., Nicolas, P., Delumeau, O., Rabatinová, A., Korelusová, J., Leduc, A., Bessières, P.,
Dervyn, E., Krásný, L., and Noirot, P. (2012). Genome-wide identification of genes directly
regulated by the pleiotropic transcription factor Spx in *Bacillus subtilis*. Nucleic Acids Res. *40*,
9571–9583. https://doi.org/10.1093/nar/gks755.

15. Diez, S., Ryu, J., Caban, K., Gonzalez, R.L., and Dworkin, J. (2020). The alarmones (p)ppGpp
directly regulate translation initiation during entry into quiescence. Proc. Natl. Acad. Sci. *117*,
15565–15572. https://doi.org/10.1073/pnas.1920013117.

16. Corrigan, R.M., Bellows, L.E., Wood, A., and Gründling, A. (2016). ppGpp negatively impacts
ribosome assembly affecting growth and antimicrobial tolerance in Gram-positive bacteria. Proc.
Natl. Acad. Sci. *113*. https://doi.org/10.1073/pnas.1522179113.

17. Pausch, P., Steinchen, W., Wieland, M., Klaus, T., Freibert, S.-A., Altegoer, F., Wilson, D.N., and
Bange, G. (2018). Structural basis for (p)ppGpp-mediated inhibition of the GTPase RbgA. J. Biol.
Chem. *293*, 19699–19709. https://doi.org/10.1074/jbc.RA118.003070.

18. Krásný, L., Tišerová, H., Jonák, J., Rejman, D., and Šanderová, H. (2008). The identity of the
transcription +1 position is crucial for changes in gene expression in response to amino acid
starvation in *Bacillus subtilis*. Mol. Microbiol. *69*, 42–54. https://doi.org/10.1111/j.1365-
2958.2008.06256.x.

19. Krásný, L., and Gourse, R.L. (2004). An alternative strategy for bacterial ribosome synthesis:
Bacillus subtilis rRNA transcription regulation. EMBO J. *23*, 4473–4483.
https://doi.org/10.1038/sj.emboj.7600423.

20. Koo, B.-M., Kritikos, G., Farelli, J.D., Todor, H., Tong, K., Kimsey, H., Wapinski, I., Galardini, M.,
Cabal, A., Peters, J.M., et al. (2017). Construction and analysis of two genome-scale deletion
libraries for *Bacillus subtilis*. Cell Syst. *4*, 291-305.e7. https://doi.org/10.1016/j.cels.2016.12.013.

21. Msadek, T., Dartois, V., Kunst, F., Herbaud, M.-L., Denizot, F., and Rapoport, G. (1998). ClpP of
Bacillus subtilis is required for competence development, motility, degradative enzyme synthesis,
growth at high temperature and sporulation. Mol. Microbiol. *27*, 899–914.
https://doi.org/10.1046/j.1365-2958.1998.00735.x.

22. Hajdusits, B., Suskiewicz, M.J., Hundt, N., Meinhart, A., Kurzbauer, R., Leodolter, J., Kukura, P.,
and Clausen, T. (2021). McsB forms a gated kinase chamber to mark aberrant bacterial proteins
for degradation. eLife *10*, e63505. https://doi.org/10.7554/eLife.63505.

23. Matavacas, J., Hallgren, J., and von Wachenfeldt, C. (2023). *Bacillus subtilis* forms twisted cells
with cell wall integrity defects upon removal of the molecular chaperones DnaK and trigger factor.
Front. Microbiol. *13*. https://doi.org/10.3389/fmicb.2022.988768.

24. Schumann, W. (2016). Regulation of bacterial heat shock stimulons. Cell Stress Chaperones *21*,
959–968. https://doi.org/10.1007/s12192-016-0727-z.

25. DeLoughery, A., Dengler, V., Chai, Y., and Losick, R. (2016). Biofilm formation by *Bacillus subtilis*
requires an endoribonuclease-containing multisubunit complex that controls mRNA levels for the
matrix gene repressor SinR. Mol. Microbiol. *99*, 425–437. https://doi.org/10.1111/mmi.13240.

26. DeLoughery, A., Lalanne, J.-B., Losick, R., and Li, G.-W. (2018). Maturation of polycistronic
mRNAs by the endoribonuclease RNase Y and its associated Y-complex in *Bacillus subtilis*. Proc.
Natl. Acad. Sci. *115*, E5585–E5594. https://doi.org/10.1073/pnas.1803283115.

27. Broglia, L., Lécrivain, A.-L., Renault, T.T., Hahnke, K., Ahmed-Begrich, R., Le Rhun, A., and
Charpentier, E. (2020). An RNA-seq based comparative approach reveals the transcriptome-wide
interplay between 3′-to-5′ exoRNases and RNase Y. Nat. Commun. *11*, 1587.
https://doi.org/10.1038/s41467-020-15387-6.

28. Taggart, J.C., Dierksheide, K.J., LeBlanc, H.J., Lalanne, J.-B., Durand, S., Braun, F., Condon, C.,
and Li, G.-W. (2025). A high-resolution view of RNA endonuclease cleavage in *Bacillus subtilis*.
Nucleic Acids Res. *53*, gkaf030. https://doi.org/10.1093/nar/gkaf030.

29. Khemici, V., Prados, J., Linder, P., and Redder, P. (2015). Decay-initiating endoribonucleolytic
cleavage by RNase Y is kept under tight control via sequence preference and sub-cellular
localisation. PLOS Genet. *11*, e1005577. https://doi.org/10.1371/journal.pgen.1005577.

30. Le Scornet, A., Jousselin, A., Baumas, K., Kostova, G., Durand, S., Poljak, L., Barriot, R.,
Coutant, E., Pigearias, R., Tejero, G., et al. (2024). Critical factors for precise and efficient RNA
cleavage by RNase Y in *Staphylococcus aureus*. PLOS Genet. *20*, e1011349.
https://doi.org/10.1371/journal.pgen.1011349.

31. Shahbabian, K., Jamalli, A., Zig, L., and Putzer, H. (2009). RNase Y, a novel endoribonuclease,
initiates riboswitch turnover in *Bacillus subtilis*. EMBO J. *28*, 3523–3533.
https://doi.org/10.1038/emboj.2009.283.

32. Chao, Y., Li, L., Girodat, D., Förstner, K.U., Said, N., Corcoran, C., Śmiga, M., Papenfort, K.,
Reinhardt, R., Wieden, H.-J., et al. (2017). In vivo cleavage map illuminates the central role of
RNase E in coding and non-coding RNA pathways. Mol. Cell *65*, 39–51.
https://doi.org/10.1016/j.molcel.2016.11.002.

33. Kime, L., Jourdan, S.S., Stead, J.A., Hidalgo-Sastre, A., and McDowall, K.J. (2010). Rapid
cleavage of RNA by RNase E in the absence of 5′ monophosphate stimulation. Mol. Microbiol. *76*,
590–604. https://doi.org/10.1111/j.1365-2958.2009.06935.x.

34. Davis, J.H., and Williamson, J.R. (2017). Structure and dynamics of bacterial ribosome
biogenesis. Philos. Trans. R. Soc. B Biol. Sci. *372*, 20160181.
https://doi.org/10.1098/rstb.2016.0181.

35. Eymann, C., Homuth, G., Scharf, C., and Hecker, M. (2002). *Bacillus subtilis* functional genomics:
global characterization of the stringent response by proteome and transcriptome analysis. J.
Bacteriol. *184*, 2500–2520. https://doi.org/10.1128/JB.184.9.2500-2520.2002.

36. Tagami, K., Nanamiya, H., Kazo, Y., Maehashi, M., Suzuki, S., Namba, E., Hoshiya, M., Hanai,
R., Tozawa, Y., Morimoto, T., et al. (2012). Expression of a small (p)ppGpp synthetase, YwaC, in
the (p)ppGpp^0 mutant of *Bacillus subtilis* triggers YvyD-dependent dimerization of ribosome.
MicrobiologyOpen *1*, 115–134. https://doi.org/10.1002/mbo3.16.

37. Beckert, B., Abdelshahid, M., Schäfer, H., Steinchen, W., Arenz, S., Berninghausen, O.,
Beckmann, R., Bange, G., Turgay, K., and Wilson, D.N. (2017). Structure of the *Bacillus subtilis*
hibernating 100S ribosome reveals the basis for 70S dimerization. EMBO J. *36*, 2061–2072.
https://doi.org/10.15252/embj.201696189.

38. Carabetta, V.J., Tanner, A.W., Greco, T.M., Defrancesco, M., Cristea, I.M., and Dubnau, D.
(2013). A complex of YlbF, YmcA and YaaT regulates sporulation, competence and biofilm
formation by accelerating the phosphorylation of Spo0A. Mol. Microbiol. *88*, 283–300.
https://doi.org/10.1111/mmi.12186.

39. Tortosa, P., Albano, M., and Dubnau, D. (2000). Characterization of *ylbF*, a new gene involved in
competence development and sporulation in *Bacillus subtilis*. Mol. Microbiol. *35*, 1110–1119.
https://doi.org/10.1046/j.1365-2958.2000.01779.x.

40. Branda, S.S., González-Pastor, J.E., Dervyn, E., Ehrlich, S.D., Losick, R., and Kolter, R. (2004).
Genes involved in formation of structured multicellular communities by *Bacillus subtilis*. J.
Bacteriol. *186*, 3970–3979. https://doi.org/10.1128/jb.186.12.3970-3979.2004.

41. Segev, E., Smith, Y., and Ben-Yehuda, S. (2012). RNA dynamics in aging bacterial spores. Cell
148, 139–149. https://doi.org/10.1016/j.cell.2011.11.059.

42. Dubnau, E., DeSantis, M., and Dubnau, D. (2023). Formation of a stable RNase Y-RicT (YaaT)
complex requires RicA (YmcA) and RicF (YlbF). mBio, e01269-23.
https://doi.org/10.1128/mbio.01269-23.

43. Hinrichs, R., Pozhydaieva, N., Höfer, K., and Graumann, P.L. (2022). Y-complex proteins show
RNA-dependent binding events at the cell membrane and distinct single-molecule dynamics. Cells
11, 933. https://doi.org/10.3390/cells11060933.

44. Adusei-Danso, F., Khaja, F.T., DeSantis, M., Jeffrey, P.D., Dubnau, E., Demeler, B., Neiditch,
M.B., and Dubnau, D. (2019). Structure-function studies of the *Bacillus subtilis* Ric proteins
identify the Fe-S cluster-ligating residues and their roles in development and RNA processing.
mBio *10*, e01841-19, /mbio/10/5/mBio.01841-19.atom. https://doi.org/10.1128/mBio.01841-19.

45. Lehnik-Habrink, M., Pförtner, H., Rempeters, L., Pietack, N., Herzberg, C., and Stülke, J. (2010).
The RNA degradosome in *Bacillus subtilis*: identification of CshA as the major RNA helicase in the
multiprotein complex. Mol. Microbiol. *77*, 958–971. https://doi.org/10.1111/j.1365-
2958.2010.07264.x.

46. Commichau, F.M., Rothe, F.M., Herzberg, C., Wagner, E., Hellwig, D., Lehnik-Habrink, M.,
Hammer, E., Völker, U., and Stülke, J. (2009). Novel activities of glycolytic enzymes in *Bacillus
subtilis*. Mol. Cell. Proteomics *8*, 1350–1360. https://doi.org/10.1074/mcp.M800546-MCP200.

47. Durand, S., and Condon, C. RNases and helicases in Gram-positive bacteria. Microbiol. Spectr. *6*,
10.1128/microbiolspec.rwr-0003–2017. https://doi.org/10.1128/microbiolspec.rwr-0003-2017.

48. Dimitrova-Paternoga, L., Kasvandik, S., Beckert, B., Granneman, S., Tenson, T., Wilson, D.N.,
and Paternoga, H. (2024). Structural basis of ribosomal 30S subunit degradation by RNase R.
Nature *626*, 1133–1140. https://doi.org/10.1038/s41586-024-07027-6.

49. Boylan, S.A., Redfield, A.R., Brody, M.S., and Price, C.W. (1993). Stress-induced activation of the
Sigma B transcription factor of *Bacillus subtilis*. J. Bacteriol. *175*, 7931–7937.
https://doi.org/10.1128/jb.175.24.7931-7937.1993.

50. Trentini, D.B., Suskiewicz, M.J., Heuck, A., Kurzbauer, R., Deszcz, L., Mechtler, K., and Clausen,
T. (2016). Arginine phosphorylation marks proteins for degradation by a Clp protease. Nature *539*,
48–53. https://doi.org/10.1038/nature20122.

51. Schmidt, A., Trentini, D.B., Spiess, S., Fuhrmann, J., Ammerer, G., Mechtler, K., and Clausen, T.
(2014). Quantitative phosphoproteomics reveals the role of protein arginine phosphorylation in the
bacterial stress response. Mol. Cell. Proteomics *13*, 537–550.
https://doi.org/10.1074/mcp.M113.032292.

52. Elsholz, A.K.W., Turgay, K., Michalik, S., Hessling, B., Gronau, K., Oertel, D., Mäder, U.,
Bernhardt, J., Becher, D., Hecker, M., et al. (2012). Global impact of protein arginine
phosphorylation on the physiology of *Bacillus subtilis*. Proc. Natl. Acad. Sci. *109*, 7451–7456.
https://doi.org/10.1073/pnas.1117483109.

53. Lytvynenko, I., Paternoga, H., Thrun, A., Balke, A., Müller, T.A., Chiang, C.H., Nagler, K.,
Tsaprailis, G., Anders, S., Bischofs, I., et al. (2019). Alanine tails signal proteolysis in bacterial
ribosome-associated quality control. Cell *178*, 76-90.e22.
https://doi.org/10.1016/j.cell.2019.05.002.

54. Wiegert, T., and Schumann, W. (2001). SsrA-mediated tagging in *Bacillus subtilis*. J. Bacteriol.
183, 3885–3889. https://doi.org/10.1128/JB.183.13.3885-3889.2001.

55. Lehnik-Habrink, M., Schaffer, M., Mäder, U., Diethmaier, C., Herzberg, C., and Stülke, J. (2011).
RNA processing in *Bacillus subtilis*: identification of targets of the essential RNase Y: Targets of
the essential RNase Y in *Bacillus subtilis*. Mol. Microbiol. *81*, 1459–1473.
https://doi.org/10.1111/j.1365-2958.2011.07777.x.

56. Anderson, B.W., Fung, D.K., and Wang, J.D. (2021). Regulatory themes and variations by the
stress-signaling nucleotide alarmones (p)ppGpp in bacteria. Annu. Rev. Genet. *55*, 115–133.
https://doi.org/10.1146/annurev-genet-021821-025827.

57. Driller, K., Cornejo, F.A., and Turgay, K. (2023). (p)ppGpp – an important player during heat
shock response. microLife *4*, uqad017. https://doi.org/10.1093/femsml/uqad017.

58. Zundel, M.A., Basturea, G.N., and Deutscher, M.P. (2009). Initiation of ribosome degradation
during starvation in *Escherichia coli*. RNA *15*, 977–983. https://doi.org/10.1261/rna.1381309.

59. Sulthana, S., Basturea, G.N., and Deutscher, M.P. (2016). Elucidation of pathways of ribosomal
RNA degradation: an essential role for RNase E. RNA *22*, 1163–1171.
https://doi.org/10.1261/rna.056275.116.

60. Lehnik-Habrink, M., Newman, J., Rothe, F.M., Solovyova, A.S., Rodrigues, C., Herzberg, C.,
Commichau, F.M., Lewis, R.J., and Stulke, J. (2011). RNase Y in *Bacillus subtilis*: a natively
disordered protein that is the functional equivalent of RNase E from *Escherichia coli*. J. Bacteriol.
193, 5431–5441. https://doi.org/10.1128/JB.05500-11.

61. Marcaida, M.J., DePristo, M.A., Chandran, V., Carpousis, A.J., and Luisi, B.F. (2006). The RNA
degradosome: life in the fast lane of adaptive molecular evolution. Trends Biochem. Sci. *31*, 359–
365. https://doi.org/10.1016/j.tibs.2006.05.005.

62. Piir, K., Paier, A., Liiv, A., Tenson, T., and Maiväli, Ü. (2011). Ribosome degradation in growing
bacteria. EMBO Rep. *12*, 458–462. https://doi.org/10.1038/embor.2011.47.

63. Kuroda, A., Nomura, K., Ohtomo, R., Kato, J., Ikeda, T., Takiguchi, N., Ohtake, H., and Kornberg,
A. (2001). Role of inorganic polyphosphate in promoting ribosomal protein degradation by the Lon
protease in *E. coli*. Science *293*, 705–708. https://doi.org/10.1126/science.1061315.

64. An, H., and Harper, J.W. (2020). Ribosome abundance control via the ubiquitin–proteasome
system and autophagy. J. Mol. Biol. *432*, 170–184. https://doi.org/10.1016/j.jmb.2019.06.001.

65. Huang, Z., Diehl, F.F., Wang, M., Li, Y., Song, A., Chen, F.X., Rosa-Mercado, N.A., Beckmann,
R., Green, R., and Cheng, J. (2025). RIOK3 mediates the degradation of 40S ribosomes. Mol.
Cell *85*, 802-814.e12. https://doi.org/10.1016/j.molcel.2025.01.013.

66. Ford, P.W., Garshott, D.M., Narasimhan, M., Ge, X., Jordahl, E.M., Subramanya, S., and Bennett,
E.J. (2025). RNF10 and RIOK3 facilitate 40S ribosomal subunit degradation upon 60S biogenesis
disruption or amino acid starvation. Cell Rep. *44*, 115371.
https://doi.org/10.1016/j.celrep.2025.115371.

67. Coria, A.R., Shah, A., Shafieinouri, M., Taylor, S.J., Orgebin, E., Guiblet, W., Miller, J.T., Sharma,
I.M., and Wu, C.C.-C. (2025). The integrated stress response regulates 18S nonfunctional rRNA
decay in mammals. Mol. Cell *85*, 787-801.e8. https://doi.org/10.1016/j.molcel.2025.01.017.

68. Simpkins, S.W., Deshpande, R., Nelson, J., Li, S.C., Piotrowski, J.S., Ward, H.N., Yashiroda, Y.,
Osada, H., Yoshida, M., Boone, C., et al. (2019). Using BEAN-counter to quantify genetic
interactions from multiplexed barcode sequencing experiments. Nat. Protoc. *14*, 415–440.
https://doi.org/10.1038/s41596-018-0099-1.

69. Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and
powerful approach to multiple testing. J. R. Stat. Soc. Ser. B Methodol. *57*, 289–300.
https://doi.org/10.1111/j.2517-6161.1995.tb02031.x.

70. Lécrivain, A.-L., Le Rhun, A., Renault, T.T., Ahmed-Begrich, R., Hahnke, K., and Charpentier, E.
(2018). In vivo 3′-to-5′ exoribonuclease targetomes of *Streptococcus pyogenes*. Proc. Natl. Acad.
Sci. *115*, 11814–11819. https://doi.org/10.1073/pnas.1809663115.

71. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal *17*, 10. https://doi.org/10.14806/ej.17.1.200.

72. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21.
https://doi.org/10.1093/bioinformatics/bts635.

73. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352.

74. Smith, T., Heger, A., and Sudbery, I. (2017). UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. Genome Res. *27*, 491–499.
https://doi.org/10.1101/gr.209601.116.

75. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics *30*, 923–930.
https://doi.org/10.1093/bioinformatics/btt656.

76. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. *15*, 550. https://doi.org/10.1186/s13059-014-0550-
8.

77. Putri, G.H., Anders, S., Pyl, P.T., Pimanda, J.E., and Zanini, F. (2022). Analysing high-throughput
sequencing data in Python with HTSeq 2.0. Bioinformatics *38*, 2943–2945.
https://doi.org/10.1093/bioinformatics/btac166.

78. Mölder, F., Jablonski, K.P., Letcher, B., Hall, M.B., Tomkins-Tinch, C.H., Sochat, V., Forster, J.,
Lee, S., Twardziok, S.O., Kanitz, A., et al. (2021). Sustainable data analysis with Snakemake.
Preprint at F1000Research, https://doi.org/10.12688/f1000research.29032.2
https://doi.org/10.12688/f1000research.29032.2.

79. McCarthy, D.J., Chen, Y., and Smyth, G.K. (2012). Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. Nucleic Acids Res. *40*, 4288–4297.
https://doi.org/10.1093/nar/gks042.

80. Geissler, A.S., Anthon, C., Alkan, F., González-Tortuero, E., Poulsen, L.D., Kallehauge, T.B.,
Breüner, A., Seemann, S.E., Vinther, J., and Gorodkin, J. (2021). BSGatlas: a unified *Bacillus*
subtilis genome and transcriptome annotation atlas with enhanced information access. Microb.
Genomics *7*, 000524. https://doi.org/10.1099/mgen.0.000524.

81. Tareen, A., and Kinney, J.B. (2020). Logomaker: beautiful sequence logos in Python.
Bioinformatics *36*, 2272–2274. https://doi.org/10.1093/bioinformatics/btz921.

82. Lorenz, R., Bernhart, S.H., Höner zu Siederdissen, C., Tafer, H., Flamm, C., Stadler, P.F., and
Hofacker, I.L. (2011). ViennaRNA Package 2.0. Algorithms Mol. Biol. *6*, 26.
https://doi.org/10.1186/1748-7188-6-26.

83. Elfmann, C., Dumann, V., van den Berg, T., and Stülke, J. (2025). A new framework for SubtiWiki,
the database for the model organism *Bacillus subtilis*. Nucleic Acids Res. *53*, D864–D870.
https://doi.org/10.1093/nar/gkae957.

84. Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic Acids Res. *29*, 45e–445. https://doi.org/10.1093/nar/29.9.e45.

85. Hughes, C.S., Moggridge, S., Müller, T., Sorensen, P.H., Morin, G.B., and Krijgsveld, J. (2019).
Single-pot, solid-phase-enhanced sample preparation for proteomics experiments. Nat. Protoc.
14, 68–85. https://doi.org/10.1038/s41596-018-0082-x.

86. Kohler, D., Staniak, M., Tsai, T.-H., Huang, T., Shulman, N., Bernhardt, O.M., MacLean, B.X.,
Nesvizhskii, A.I., Reiter, L., Sabido, E., et al. (2023). MSstats version 4.0: statistical analyses of
quantitative mass spectrometry-based proteomic experiments with chromatography-based
quantification at scale. J. Proteome Res. *22*, 1466–1482.
https://doi.org/10.1021/acs.jproteome.2c00834.

87. Jahn, M., Vialas, V., Karlsen, J., Maddalo, G., Edfors, F., Forsström, B., Uhlén, M., Käll, L., and
Hudson, E.P. (2018). Growth of cyanobacteria is constrained by the abundance of light and
carbon assimilation proteins. Cell Rep. *25*, 478-486.e8.
https://doi.org/10.1016/j.celrep.2018.09.040.

88. Jahn, M., Crang, N., Janasch, M., Hober, A., Forsström, B., Kimler, K., Mattausch, A., Chen, Q.,
Asplund-Samuelsson, J., and Hudson, E.P. (2021). Protein allocation and utilization in the

versatile chemolithoautotroph *Cupriavidus necator*. eLife *10*, e69019.
https://doi.org/10.7554/eLife.69019.

89. Sohmen, D., Chiba, S., Shimokawa-Chiba, N., Innis, C.A., Berninghausen, O., Beckmann, R., Ito,
K., and Wilson, D.N. (2015). Structure of the *Bacillus subtilis* 70S ribosome reveals the basis for
species-specific stalling. Nat. Commun. *6*, 6941. https://doi.org/10.1038/ncomms7941.

1088

Figure S1. Fitness of selected deletion mutants during heat shock in the genetic screening. Heatmap of the relative fitness of **A)** heat shock transcriptional repressors, and **B)** chaperones, proteases and adaptors. The color represents the average relative fitness of three biological replicates. A negative relative fitness value means that the abundance of the deletion mutant, compared to the rest of the population, was negatively affected by the treatment, while a positive value means an increased abundance on those conditions. h.s. = heat shock.

Figure S2. Recognition motif, length, or expression levels do not determine the Y-complex substrates. A) Stabilization of the resulting fragments after Y-complex processing, the coverage ratio in a window of 5 nt before and after the processing site was calculated and used as a proxy of the stabilization of the resulting fragments. The predicted stability correlates with the end detected in the ISCP method. **B)** Motif search using 5' and U-type ends. The arrow indicates the cleavage site. **C)** Number of Y-complex-dependent cleavages detected per monocistronic or polycistronic transcripts in the late exponential or transition phase. **D)** Number of cleavages detected by operon length of the

transcript. The color indicates the numbers of Y-complex-dependent ends detected per transcript **E)** RNA levels of WT and Δ*ymcA* mutant transcripts at late exponential and transition phases. Transcripts processed by the Y-complex are highlighted in magenta. Transcript annotations were obtained from BSGatlas.

1110

1111

Figure S3. The Y-complex processes mRNA in UTRs and coding regions. RNA end counts per
million (cpm), and RNA-seq coverage of selected transcripts in WT and Δ*ymcA.* The arrow indicates the
Y-complex-dependent processing site. **A)** The *gapA-cggR* transcript was used as a positive control. **B)**
Processing in the coding region of *yloC* in the polycistronic mRNA *yloC-remA-gmk-yloH*. **C)** Processing
in the coding region of the monocistronic transcript of *guaB*. **D)** Processing at the 3' UTR of *trxB,* near
the Rho-independent terminator. Processing in the internal UTR of translation-associated transcripts
like **E)** the *bipA-ylaH* transcript, and **F)** between *fusA-tufA*.

A

B

1120

Figure S4. mRNA processing by the Y-complex and effect on the transcriptome. A)
Overrepresentation analysis of pathways whose transcripts are cleaved by the Y-complex during the

late exponential phase. Operons were assigned to be involved in a pathway if they code for one gene
participating in such a pathway. Pathway annotations were retrieved from SubtiWiki[83]. The color
indicates the statistical significance of the enrichment. **B)** Gene set enrichment analysis (GSEA) of
differentially expressed genes at the late exponential and transition phase when comparing Δ*ymcA* to
WT. The size of the dot represents the statistical significance. The normalized enrichment score is
displayed on the *x*-axis; positive values mean upregulation; meanwhile negative values mean
downregulation. GSEA results were prefiltered for pathways showing and enrichment with an FDR ≤
0.01. Biosyn. = Biosynthesis, acq. = Acquisition.

1131

1132 **Figure S5. Mapping of Y-complex-dependent processing in the secondary structure of rRNA at**
1133 **the late exponential phase.** Cleavages are marked with an arrow and the helix number for **A)** 23S and
1134 **B)** 16S rRNA.

1135

1136

1137 **Figure S6. Mapping of Y-complex dependent processing in the ribosome 3D structure.** The RNA
1138 end detected by the ISCP method is highlighted in magenta. The ribosome structure was retrieved from
1139 PDB (ID: 3J9W)[89]. The A site is shown as a green ellipse.

Figure S7. Y-complex-dependent initiates degradation of rRNA but does not affect transcriptional control of stringent response. A) Northern blot against 16S of WT and Δ*ymcA* cells at the late exponential phase, before and after treatment with 100 µg/ml rifampicin to inhibit transcription. The methylene blue-stained membrane is shown as a loading control. This image is representative of three biological replicates. Activity of the promoter controlling **B)** *rrnB* (rRNA) transcription, **C)** *rplJ*, and **D)** *rpsJ* (r-proteins) genes at late exponential and transition phase compared to the exponential phase in WT and Δ*ymcA*. The promoters were cloned controlling *gfp* transcription, which was measured using RT-qPCR. The *pcp* transcript was used as a reference. Individual values and the average ± standard deviation of three biological replicates and two technical replicates are shown. The statistical significance was tested using student's *t*-test. ns: *p*-value > 0.05, *: *p*-value ≤ 0.05. **E)** Ribosome sedimentation profile in 10-50% sucrose gradients of WT, Δ*hpf*, Δ*ymcA*, and Δ*ymcA* Δ*hpf* in the transition phase. The 70S and 100S ribosomes are highlighted with a blue or red box, respectively. The data is representative of three biological replicates.

1154

1155

1156

1157

1158

1159

Figure S8. Control of ribosome levels and protein homeostasis in *ymcA* is important for responding against proteotoxic stresses. A) Northern blot against 16S of WT and ΔymcA cells at exponential phase before and after heat shock. The methylene blue-stained membrane is shown as a loading control. This image is representative of three biological replicates. **B)** Induction of CtsR and HrcA regulons after heat shock (50°C) ΔymcA compared to the WT. Proteins were measured by mass spectrometry. The color bar shows the $\log_2$ fold change compared to the WT control. **C)** Percentage of aggregated proteins before and after a 15 min heat shock at 54°C in WT and ΔymcA, (p)ppGpp^0 and (p)ppGpp^0 ΔymcA. Individual values and the average ± standard deviation of three biological replicates are shown. The statistical difference was tested using student's *t*-test. ns: *p*-value > 0.05, *: *p*-value ≤ 0.05, **: *p*-value ≤ 0.01, ***: *p*-value ≤ 0.001. **D)** Spot test of deletion mutants of Y-complex members in the displayed stresses. The picture is representative of three biological replicates.