Targeting the translation machinery: The RNase Y specificity factor Y-complex coordinates ribosome degradation

Fabián A. Cornejo[1,*], Kristina Driller[1,2,*], Rina Ahmed-Begrich[1], Katja Schmidt[1], Michael Jahn[1], Vivekanandan Shanmuganathan[1], Karin Hahnke[1], Florian Kondrot[1], Thomas F. Wulff[1], Sebastian Rämisch[1], Kathirvel Alagesan[1], Emmanuelle Charpentier[1,3], and Kürşad Turgay[1,2]

[1]Max Planck Unit for the Science of Pathogens, Berlin, Germany

[2]Leibniz Universität Hannover, Institute of Microbiology, Hannover, Germany

[3]Humboldt-Universität zu Berlin, Institute of Biology, Berlin, Germany

* These authors equally contributed to this work

Correspondence to: cornejo@mpusp.mpg.de, turgay@mpusp.mpg.de

Abstract

Limiting the synthesis and activity of ribosomes is crucial for responding to various stresses, including nutrient starvation or proteotoxic stress. In *Bacillus subtilis,* this can be achieved through the coordinated action of the alarmones (p)ppGpp and the transcription factor Spx. In this study, we performed a genetic screen to identify new factors involved in the heat shock response of *B. subtilis*. We discovered that the RNase Y specificity factor Y-complex plays an important role in the heat shock response and the transition to the stationary growth phase. The Y-complex is required for the processing of various RNAs, including the maturation of mRNAs encoding proteins involved in translation and metabolism. We further demonstrate that the Y-complex and RNase Y initiate the degradation of rRNAs in mature ribosomes, lowering their abundance. We propose the Y-complex as a system by which *B. subtilis* regulates protein synthesis and resource allocation after exponential growth.

Keywords: *Bacillus subtilis*, Ribosome, Stress response, Y-complex, RNase Y, Ribosome degradation, Heat shock, (p)ppGpp

Introduction

Bacteria can sense, adapt to, and survive stressful environmental changes facilitated by various general and specific stress response systems. Proteotoxic stresses, like high temperatures, significantly disrupt protein folding and can lead to the formation of toxic protein aggregates. Chaperones and proteases are expressed as a response due to the inactivation of the transcriptional repressors HrcA and CtsR in *Bacillus subtilis*[1–4]. The activation of the protein quality control (PQC) allows the refolding of un- or misfolded proteins by different chaperone systems or the degradation of irreparable proteins through AAA+ protease complexes like ClpCP[5,6].

In addition to the repairing or removal of protein aggregates, inhibiting translation is crucial for survival during proteotoxic stresses, as it can alleviate the extra load on the PQC system[7–10]. In *Bacillus subtilis*, the nucleotide second messengers guanosine tetraphosphate and guanosine pentaphosphate, collectively referred to as (p)ppGpp, and the transcription factor Spx independently regulate synthesis and activity of ribosomes during stress[7,11]. Spx is a redox-sensitive heat shock transcription factor that controls the expression of redox chaperones and simultaneously represses transcription of translation-related genes during oxidative and heat shock response[11–14].

The alarmones (p)ppGpp were initially discovered to be synthesized during amino acid starvation, but more recently, it was observed that their synthesis is also induced by protein folding stresses, such as heat or oxidative stress[7]. During heat stress, their accumulation results in increased thermoresistance, inhibition of translation, and a reduction in protein aggregation[7]. (p)ppGpp inhibit translation by binding to specific GTPases (e.g. IF2, RbgA)[15–17]. Furthermore, (p)ppGpp accumulation can lead to decreased transcription of e.g. rRNA and ribosomal protein (r-protein) transcripts by inhibiting GTP biosynthesis and reducing its levels[18,19]. This downregulation of translation is an important component of the response to nutrient scarcity as well as heat shock.

In this study, we aimed to identify new genes or pathways involved in the heat shock response using a genetic screen that reports on fitness. We identified that the Y-complex (RicAFT-complex), which interacts with the endonuclease RNase Y and conveys substrate specificity, plays a role in the heat shock response. We demonstrate further that the Y-complex regulates ribosome levels by initiating ribosome degradation during nutrient depletion or heat shock. Mutants lacking a Y-complex member show increased ribosome levels before and during heat shock, increasing the burden on the PQC system and, in consequence, protein aggregation during heat stress. These findings not only indicate the importance of RNases during stress response but also emphasize that controlling the stability of RNA is an important general mechanism to modify cellular function.

Results

The Y-complex is necessary for survival during heat shock

To identify genes that play a role in the heat shock response, we used the BKE single-mutant library[20], which includes a barcoded deletion cassette for each non-essential gene in *B. subtilis*. All gene deletion mutant strains were pooled, grown at 30°C, and subjected to a mild heat shock at 50°C during the mid-exponential phase, while a control was maintained at 30°C. Relative fitness scores for each gene deletion in the pooled population were calculated by quantifying each barcode at different temperatures (**Fig. 1A, Table S1**). The relative fitness of the heat shock sample compared to the control at 30°C was used to identify genes whose deletion results in a change in fitness during heat shock (**Fig. 1A-B, Table S1**).

We observed that *B. subtilis* strains lacking the genes coding for the heat shock transcriptional repressors HrcA or CtsR exhibit higher fitness at 50°C compared to 30°C (**Fig. S1A**), consistent with their known regulatory role. In contrast, deletion of the genes coding for the DnaK chaperone system, the protease ClpP, and the adaptor protein complex McsA-McsB show greater sensitivity to increased temperatures (**Fig. S1B**), demonstrating that our screening approach can identify genes with already known heat-shock phenotypes[21–23].

The control of transcription initiation by transcription factors has been well established[24]. However, the contribution of RNA stability in stress response is not well understood. Interestingly, six mutants lacking RNases or RNase-regulating genes exhibit differential fitness at 50°C compared to 30°C (**Fig. 1B**). The deletion of RNase Mini-III (Δ*mrnC)* results in lower fitness at 30°C, whereas its fitness remains unaffected at 50°C. Deletions of RNase J2 (Δ*rnjB)*, the nano-RNase A (Δ*nrnA*), and the RNA pyrophosphohydrolase and (p)ppGpp hydrolase NahA (Δ*yvcI*) exhibit reduced fitness during increased temperatures (**Fig. 1B**).

One of the mutants with the lowest fitness score at 50°C is Δ*ymcA* (**Fig. 1B**). YmcA (RicA) interacts with YaaT (RicT) and YlbF (RicF) to form the Y-complex, which associates with the endonuclease RNase Y for maturation of polycistronic mRNA and riboswitches[25,26]. In addition, we also identified Δ*yaaT* in our screen (**Fig. 1B**). We could confirm the heat sensitivity of all three deletion mutants of the Y-complex genes in single-culture experiments during a severe heat shock at 54°C (**Fig. 1C**).

Identification of Y-complex-dependent cleavages at late exponential and transition phases

We evaluated the growth of all three strains lacking the Y-complex encoding genes. They exhibit a similar growth defect that begins at the end of the exponential phase and continues during transition phase until the stationary phase (**Fig. 2A**). Given that this complex associates

with the endonuclease RNase Y and regulates substrate selection[26], we investigated Y-

complex-dependent cleavages in the late exponential and transition phase (**Fig. 2A**) using the

ISCP (Identification of Specific Cleavage Positions) approach[27]. We compared the coverage

of RNA 3' and 5' ends in the WT and $\Delta ymcA$ mutant, and the ends significantly reduced in the

$\Delta ymcA$ were classified as putative Y-complex-dependent cleavages (**Table S2**). It is worth

noting that deleting any member of the Y-complex results in a significant decrease in the

processing of substrate RNA[26]. When a transcript is cleaved, it results in two products: one is

quickly degraded by exonucleases, while the other is stabilized[27]. Consequently, it is not

always possible to detect both 5' and 3' ends after cleavage.

We identified 221 and 258 Y-complex-dependent ends in late exponential and transition

phases, with six also detected in a previous study at the mid-exponential phase[26] (**Table S3**).

In the late exponential phase, 5' and 3' ends are almost equally distributed, yet more 5' ends

are detected in the transition phase (**Fig. 2B**).

It is known that the Y-complex and RNase Y generate fragments with differential stability in the

mid-exponential phase[26]. Here, we evaluated the stability of the resulting fragments in the late

exponential and transition phases by calculating the ratio of coverage upstream and

downstream of the cleavage site (**Table S2**). We observed a trend toward stabilizing

downstream fragments. However, upstream fragments can also be stabilized to a lesser extent

(**Fig. S2A**). The ISCP method differentiates processing sites with blunt ends (Unique, U-type)

or when it was further processed by exonucleases[27] (Stepped, S-type). Most S-types were

found at the 3' ends, suggesting further processing by exonucleases, while most U-types were

detected at 5' ends (**Fig. 2C**). This is consistent with the trend to stabilise the downstream

fragment (**Fig. S2A**).

The Y-complex could target a particular sequence motif, thereby helping to recognize specific

substrates. However, a motif search using U-type ends resembles the simple motif described

for RNase Y in *B. subtilis*[28], *S. aureus*[29], and *S. pyogenes*[27], being most likely a G upstream of

the cleavage site (**Fig. S2B**). Given the simple sequence motif, we investigated whether

cleavages occur randomly in longer or more abundant transcripts. Notably, approximately 81%

of targeted transcripts are cleaved only once, about 16% twice, and very few are cleaved

multiple times (**Fig. S2C**). Neither the length nor the abundance of a transcript predicts whether

it is cleaved, suggesting that other factors influence where cleavages occur (**Fig. S2D-E, Table**

S4). Recent studies have shown that the cleavage efficiency is multifactorial, depending on

the sequence and secondary structures surrounding the cleavage site[28,30].

As we observed that RNA processing could result in fragments with different stability (**Fig.**

S2A), we wondered if this could be a consequence of secondary structures protecting them

from exonucleases. We used U-type ends and compared transcripts with the same cleavage

motif but not cleaved as controls. For the 5' ends, a strong decrease in the minimum free

energy ~25 nt downstream of the cleavage site indicated a possible secondary structure, while

the upstream fragment remained similar to the control (**Fig. 2D**). The upstream fragment's

lower stability may be due to its lack of secondary structure, making it accessible to 3'-5'

exonucleases. For the 3' ends, the more stable upstream fragment could form a possible

secondary structure starting 10 nt upstream of the cleavage site (**Fig. 2D**). The generation of

mRNA fragments with different stability might have a regulatory role in gene expression.

Y-complex-mediated transcript cleavages and mRNA maturation

Most detected Y-complex-dependent ends are inside ORFs, mainly in polycistronic mRNAs

(**Fig. 2E-F**), confirming the role of the Y-complex in the maturation and stoichiometry regulation

of co-transcribed genes[26]. An example is the *cggR-gapA* transcript, where the Y-complex

mediates cleavage in the coding sequence of *cggR* near the stop codon[25,26]. Cleavage renders

the *cggR* fragment less stable compared to the *gapA* fragment (**Fig. S3A**), thereby uncoupling

the transcript levels of these co-transcribed genes. Interestingly, we observed other cleavages

generating similar uncoupling of transcripts, such as in the gene encoding the endonuclease

YloC (**Fig. S3B**, **Table S2)**. In addition, cleavages in genes transcribed as monocistronic

mRNAs, such as *guaB,* suggest that the Y-complex could initiate their degradation (**Fig. S3C**,

Table S2).

We observe Y complex-dependent processing in 5', 3', and internal UTRs (**Fig. 2F**). Cleavages

in 5' UTR suggest a possible regulatory role by decreasing the abundance of the mRNA

containing them, as already described for riboswitches[26,31], or by producing a shorter version

of a 5' UTR, which could affect gene expression. Processing of 3' and internal UTR could

influence mRNA stability and polycistronic mRNA maturation. We observed that some 3' UTR

cleavages are located upstream of the Rho-independent terminator, decreasing the stability of

the upstream fragment and leaving an apparent more stable downstream fragment that

includes its terminator structure, as observed for *trxB* (**Fig. S3D**). This phenomenon has been

observed in *E. coli*, where RNase E cleaves upstream of the Rho-independent terminator[32,33].

In polycistronic mRNA maturation, we observed that Y-complex processes internal UTRs,

decoupling the RNA levels of coding fragments (**Fig. S3E-F**). For example, the *bipA-ylaH*

transcript is cleaved 21 nt after the *bipA* stop codon, rendering the upstream fragment less

stable than the *ylaH* fragment (**Fig. S3E**). We also observed a cleavage between the *fusA* (EF-

G) and *tufA* (EF-Tu) genes, both coding for translation elongation factors (**Fig. S3F**).

The Y-complex targets translation and metabolism-related genes in the late exponential phase

We wondered if the Y-complex primarily processes transcripts bearing genes with specific functions required for switching the growth phase. Therefore, we conducted an overrepresentation analysis and observed that the most significantly enriched categories pertain to "Metabolism" and "Information processing" (**Fig. 2G, Fig. S4A, Table S5**). Overall, the Y-complex is predominantly involved in the processing of operons encoding genes for carbon metabolism (e.g., glycolysis and gluconeogenesis), nucleotide metabolism (e.g., pyrimidine and purine biosynthesis), and amino acid biosynthesis and utilization (**Fig. 2G**). This allows adjusting the utilization and biosynthesis of nutrients when they become limited. In addition, all transcripts containing rRNA and ~50% of transcripts carrying genes encoding translation factors and r-proteins are processed by the Y-complex (**Fig. 2G**).

The reduced RNA processing in the *ymcA* deletion strain results in significant changes in the transcriptome (**Fig. S4B, Table S6-7**). Consequently, transcripts of certain pathways related to metabolism and translation, among others, show maturation events that depend on the Y-complex and also demonstrate differential expression upon deletion of *ymcA* (**Fig. 2G**, **Fig. S4B**). This may result from either a direct or indirect effect of decreased RNA maturation events. Our results suggest that the Y-complex and RNase Y are important for transitioning to the stationary phase by processing transcripts related to translation and metabolism, which may impact their stability and/or gene expression.

The Y-complex targets ribosomes for degradation

Several Y complex-dependent cleavages were detected in the rRNA sequence during both growth phases, most of them being in the 23S rRNA during the late exponential phase (**Fig. 3A**). We detected cleavages in specific helices in the 23S and 16S rRNA, mainly in loops of hairpin structures (**Fig. S5A-B**). These helices are accessible and primarily located close to the A-site and the interface between the two ribosomal subunits (**Fig. S6**), which could be recognized by the Y-complex and the endonuclease RNase Y, initiating ribosome degradation.

We confirmed the observed rRNA degradation via northern blots of 23S or 16S rRNA (**Fig. 3B**, **Fig. S7A**). Deletion of *ymcA* strongly reduces the presence of rRNA degradation intermediates during the late exponential phase. As ribosomes are assembled co-transcriptionally[34], we inhibited transcription by adding rifampicin to cultures at the late exponential phase. We observed that stopping transcription does not affect rRNA degradation of the 23S and even raises the levels of 16S degradation intermediates, indicating that mature ribosomes are degraded (**Fig. 3B**, **Fig. S7A**).

Cleavages within mature rRNA can trigger ribosome disassembly and degradation, which,
during the transition phase, coincides with translation downregulation through reduced
synthesis and inhibition of ribosome activity. This suggests that the Y-complex may be involved
in reducing ribosome levels to accommodate the new, slower growth rate. Ribosome
degradation has also been observed in *E. coli,* where it is initiated by the functionally equivalent
endonuclease RNase E, resulting in a decrease in ribosome levels of up to 50%[35].

To understand this process, we measured ribosome content via quantitative mass
spectrometry (**Table S8**), using all r-proteins as a proxy. When *B. subtilis* shifts from
exponential to the transition phase, we observed a coordinated reduction in r-protein levels
(**Fig. 3C**), indicating a decrease in ribosome levels. This reduction is highly impaired in Δ*ymcA*
(**Fig. 3C**), where the levels of r-proteins do not decline to the same degree as in the WT, likely
due to the reduced degradation of ribosomes.

Accumulation of (p)ppGpp leads to inhibition of transcription of rRNA and r-proteins[19,36] upon
shortages of nutrients during the transition to the stationary phase. To test if the deletion of
ymcA is interfering with this transcriptional downregulation, we measured the promoter activity
of the rRNA promoters *rrnJ* and *rrnB* and r-protein promoters of *rplK*, *rplJ*, and *rpsJ*. All these
promoters show similar downregulation in the late exponential and transition phases in both
WT and Δ*ymcA* strains (**Fig. 3D-E**, **Fig. S7B-D**), confirming that their transcriptional control
remains unaffected and is not the cause of the increased ribosome amounts in the Δ*ymcA*
strain.

Notably, we observed that the increased levels of ribosomes in a Δ*ymcA* strain led to the
accumulation of hibernating 100S ribosomes during the transition phase (**Fig. 3F**). This 100S
is no longer observed after the deletion of the gene encoding the hibernation factor *hpf*, which
triggers their formation[37,38] (**Fig. S7E**). The presence of hibernating ribosomes at this time point
indicates that functional and over abundant ribosomes in the Δ*ymcA* strain are sequestered
away from the actively translating pool through hibernation.

(p)ppGpp and the Y-complex play a crucial role in regulating the levels of ribosomes

Since ribosome levels and translation are lowered by (p)ppGpp under nutrient limitation or
other stresses, we wondered if there is a genetic interaction between the two systems. Deleting
ymcA in a strain that cannot synthesize (p)ppGpp [(p)ppGpp^0] leads to an even stronger growth
defect during the transition phase (**Fig. 4A**). The (p)ppGpp^0 strain, like Δ*ymcA*, shows impaired
reduction of ribosome levels during the transition phase (**Fig. 4B**). The double mutant
(p)ppGpp^0 Δ*ymcA* shows almost no reduction in ribosome quantities during this growth phase
(**Fig. 4B-C**). The same is observed when the r-protein mass fraction is calculated at the
transition phase (**Fig. 4D**). These experiments suggest that the Y-complex and (p)ppGpp

control ribosome levels and translation independently during post-exponential growth. The
absence of regulation by both systems results in a severe growth defect.

Increased ribosome levels lead to protein aggregation during heat shock

Since we discovered the $\Delta ymcA$ strain in a competition experiment during heat shock, we
explored whether the Y-complex is involved in lowering ribosome levels upon proteotoxic
stress. We evaluated the degradation of 23S and 16S rRNA by the Y-complex before and after
heat shock (**Fig. 5A**, **Fig. S8A**). After 15 minutes at 50°C, degradation intermediates are
observed in the WT but not in $\Delta ymcA$, confirming a role of the Y-complex in degrading
ribosomes during heat stress (**Fig. 5A**). Comparing r-protein mass fractions before and during
heat shock shows that both WT and $\Delta ymcA$ downregulate ribosome levels during the
experiment. Overall, there are increased levels of r-proteins in the $\Delta ymcA$ strain already before
the heat shock (**Fig. 5B**), which suggests a role of the Y-complex in maintaining ribosome
levels during normal growth. Notably, the levels and induction of known heat-shock-regulated
proteins is not affected in the *ymcA* deletion (**Fig. 5C, Table S9, Fig. S8B**). This suggests that
chaperone and protease induction during heat stress in the $\Delta ymcA$ strain is intact.

The downregulation of translation is crucial to limit the increased burden of misfolding and
aggregation-prone nascent polypeptide chains on the PQC system. In agreement, a $(p)ppGpp^0$
strain shows increased aggregate formation during heat shock[7]. We quantified the relative
amount of aggregated proteins before and after 15 minutes of severe heat shock at 54°C and
observed that the $\Delta ymcA$ strain contained significantly higher amounts of aggregated proteins
already prior to heat exposure (**Fig. 5D**). This is consistent with the observed increased
ribosome levels of the $\Delta ymcA$ strain already without heat shock (time point 0 in **Fig. 5B**). After
15 minutes of heat shock, $\Delta ymcA$ shows 20% more aggregated proteins than WT. The
$(p)ppGpp^0$ $\Delta ymcA$ mutant exhibits even higher protein aggregation than $\Delta ymcA$ or $(p)ppGpp^0$
strains alone, further supporting that (p)ppGpp and the Y-complex act independently on the
regulation of ribosome levels and translation (**Fig. S8C**). Mutants of the Y-complex are also
strongly affected by other proteotoxic stresses, such as osmotic and oxidative stress*,* but not
by ethanol (**Fig. S8D**), which further demonstrates the importance of controlling ribosome
levels under proteotoxic stress conditions.

In summary, we observed that the Y-complex controls ribosome levels by degrading rRNA and
processing r-protein transcripts. The absence of the Y-complex leads to an increase in the
amount of ribosomes, resulting in a higher load on the PQC by nascent polypeptide chains,
which in turn may increase sensitivity to protein folding stresses.

Discussion

Regulation of transcription initiation plays a crucial role in controlling stress response pathways. Gene expression can also be controlled post-transcriptionally, e.g., by modifying mRNA stability or translation efficiency. Due to the short half-life of mRNA, their levels can change rapidly through transcription initiation, which may lead to the assumption that RNases do not play a direct role in regulating gene expression during stress. However, we observed that the fitness of *B. subtilis* during heat shock is affected by the deletion of some non-essential RNases and the Y-complex, which regulates substrate selection of the quasi-essential endonuclease RNase Y (**Fig. 1B-C**). This suggests that controlling RNA maturation and stability can play an important role in stress response and gene expression.

Our results suggest that the Y-complex may regulate transcripts involved in a wide range of metabolic pathways, developmental processes, stress responses as well as translation (**Fig. 2G, Fig. S4A-B, Table S2, Table S6-7**). RNA processing by the Y-complex-guided RNase Y could potentially create new transcript isoforms[26] or modify regulatory elements such as UTRs[26,31] (**Fig. 2F**, **Fig. S3**), thereby tailoring gene expression to specific growth conditions. This could explain some pleiotropic phenotypes already observed in these mutants, such as reduced sporulation, competence, and defective biofilm formation[39–41]. More research is necessary to understand the exact role of the Y-complex in developmental pathways.

We also show that the Y-complex initiates ribosome degradation at the end of exponential growth by directing RNase Y to cleave 23S and 16S rRNA (**Fig. 3A-B**, **Fig. S7A**), thereby decreasing ribosome levels. Consistent with our findings, it has been previously suggested that RNase Y could degrade rRNA in aging spores[42]. In *E. coli*, ribosome degradation is also triggered during nutrient starvation. This process is initiated by the endonuclease RNase E, which is functionally equivalent to RNase Y but does not share sequence homology, exemplifying a case of convergent evolution[43].

Recent evidence indicates that YmcA and YlbF transfer YaaT to RNase Y, forming a stable, membrane-localized YaaT-RNase Y complex[44]. However, the deletion of any of the Y-complex members leads to a strong decrease in the processing of target mRNA[26], indicating that all of them are equally indispensable for RNA processing. Both YmcA and YlbF show a dynamic localization between the cell membrane and the cytosol[45]. It is possible that YmcA and YlbF also recognize substrates, including ribosomes, and direct them to the membrane for cleavage. This represents a sophisticated spatial regulation of RNA processing and ribosome decay, separating them from transcription and translation in the cell.

RNase Y in *B. subtilis* also interacts in the membrane with other RNases, helicases, and glycolytic enzymes to form a degradosome-like network (DLN), which participates in the decay

of RNA[46–48]. This membrane localization would also imply that once rRNA is cleaved by RNase

Y and the Y-complex, exonucleases such as RNase R, PNPase, and RNase J could complete

its degradation. AAA+ protease complexes could degrade the released ribosomal proteins. For

instance, in *E. coli*, r-proteins are degraded by the Lon protease during starvation[49].

Most of the cleavages we observed in rRNA were located in the 23S (**Fig. 3A**); however, the

Y-complex and RNase Y can also initiate 16S degradation in the late stationary phase and

during heat shock (**Fig. 3A**, **Fig. S7**, **Fig. S8**). Recently, it was observed that the

exoribonuclease RNase R alone can degrade 16S[50], suggesting that endonucleolytic

cleavages might not be strictly necessary for 16S degradation. It is worth noting that RNase R

is induced as part of the general stress response, which is activated in the stationary phase or

during heat shock[51], indicating a possible stress-specific degradation of 16S rRNA

independent of the Y-complex. These observations suggest the presence of different possible

pathways for ribosome degradation.

Interestingly, ribosomes have been observed to reach membranes during starvation conditions

in several organisms. For instance, in *E. coli*, the hibernation factors YqjD/ElaB/YgaM can bind

to 70S ribosomes near the peptide exit tunnel and tether them to the inner membrane[52]. In

eukaryotes, during starvation, the proteins NUFIP1 and ZHNIT3 bind ribosomes and act as

cargo receptors in the forming autophagosome for their degradation[53]. Recently, it has also

been shown that, during nutrient scarcity, cytoplasmic ribosomes can hibernate by binding to

the mitochondrial membrane in yeast[54].

The regulation of translation by the Y-complex is not limited to ribosome degradation. The Y-

complex additionally processes transcripts encoding r-proteins and other translation factors

(**Fig. 2G**), such as BipA, EF-Tu, and EF-G (**Fig. S3**), which might regulate their translation or

lead to their decay. Strikingly, the Y-complex is necessary to process some transcripts coding

for r-proteins at the mid-logarithmic phase[26], and the depletion of RNase Y in similar conditions

leads to an increased abundance of transcripts of r-proteins, such as *rplS, rpsO, rpmA, rpsD,*

rpsB, and *rplU*[55].

Translation control is an important component of certain stress response mechanisms. For

example, during nutrient starvation, downregulating translation allows the reallocation of

resources to biosynthetic pathways[56]. During proteotoxic stresses, it could relieve the

overwhelmed PQC system by reducing the synthesis of new proteins that could potentially

aggregate[57]. The stress transcription factor Spx and the alarmones (p)ppGpp regulate

ribosome levels in *B. subtilis*, and the lack of both impairs growth at 50°C [7]. The strong growth

defect of cells lacking both (p)ppGpp and the Y-complex suggests that the two systems are

important for transitioning from the exponential to the stationary phase. Missing their combined

regulation of translation leads to high ribosome levels and increased accumulation of protein

aggregates (**Fig. S8C**). In times of nutrient limitation, the collective function of (p)ppGpp and
the Y-complex in regulating ribosome levels by inhibiting their synthesis and initiating
degradation allows resource allocation to other biosynthetic pathways (**Fig. 6**).

In conclusion, ribosome degradation is crucial during nutrient scarcity. Different organisms
employ various strategies to achieve ribosome degradation. We demonstrate that the Y-
complex and RNase Y are responsible for this function in *B. subtilis*, likely applicable to many
other bacteria that lack RNase E. Further studies are required to fully describe how this process
operates. This includes, for example, how ribosomes reach the membrane, whether and how
the Y-complex is activated during the late exponential phase, and which proteases are involved
in r-protein degradation.

Materials and methods

<u>Growth conditions</u>

Unless stated the contrary, strains (**Table S10**) were streaked on LB plates (5 g/l yeast extract, 10 g/l tryptone agar, 10 g/l NaCl, 1.5% agar) containing the required antibiotics for selection (1 µg/ml erythromycin + 25 µg/ml lincomycin, 150 µg/ml spectinomycin or 10 µg/ml kanamycin) and incubated overnight at 37°C. A single colony was inoculated in 3 ml of LB medium and grown overnight at 37°C and 180 rpm, then used as the inoculum for subsequent experiments.

<u>Strain construction</u>

For constructing the strains $\Delta ymcA::erm^R$ and $\Delta yaaT::erm^R$, the loci containing the resistance cassette were PCR-amplified from the respective strains of the BKE library[20] and transformed in *B. subtilis* WT. The deletions of $\Delta rel::kan^R$ and $\Delta ylbF::erm^R$ were constructed as described in Ref.[20]. For promoter activity reporter strains, the *gfp*(mut2) gene was cloned downstream of the promoters of interest, using the integrative plasmid pDR111 as a backbone.

For transformation, 600 µl of an overnight culture of the receptor strains were diluted in 5 ml competence medium (17.5 g/l K_2HPO_4, 7.5 g/l KH_2PO_4, 2.5 g/l $(NH_4)_2SO_4$, 1.25 g/l tri-sodium citrate x2 H_2O, 0.5% glucose, 7 mM $MgSO_4$, 0.1 mg/ml L-tryptophan, 0.02% casamino acids, 0.22 g/ml ammonium iron citrate) and grown at 37°C, 180 rpm, for 3 hours. Subsequently, 5 ml of starvation medium (17.5 g/l K_2HPO_4, 7.5 g/l KH_2PO_4, 2.5 g/l $(NH_4)_2SO_4$, 1.25 g/l tri-sodium citrate x2 H_2O, 0.5% glucose, 7 mM $MgSO_4$) were added and grown for 2 more hours. One ml of this culture was incubated with the DNA to be transformed and grown for 2 hours at 37°C and 180 rpm. Positive clones were selected on LB agar containing the required antibiotics. Genetic modifications were verified using PCR and DNA sequencing.

<u>Genetic screening during heat shock</u>

The BKE library[20] was replicated on LB agar plates containing 1 µg/ml erythromycin. Liquid cultures were prepared in 200 µl LB medium supplemented with 1 µg/ml erythromycin and grown for 42 hours at 30°C 200 rpm. All mutant strains were pooled and mixed for 15 minutes using a magnetic stirrer. The pooled culture was centrifuged at 5000 rpm for 10 minutes at 18°C. The pellet was resuspended in LB medium to an OD_{600} of 65 and mixed with an equal volume of 50% glycerol. The pooled library was aliquoted and kept at -80°C.

For each replicate, a tube of the pooled library was quickly unfrozen in a water bath at 30°C and used to inoculate 75 ml LB medium in 500 ml flasks to a starting of OD_{600} 0.05. The culture was grown in a water bath at 30°C, 180 rpm, until OD_{600} 0.4. Fifty ml of culture were split into two 250 ml flasks. One flask was transferred to 50°C, 180 rpm for heat shock. The other flask was used as a control sample and was kept at 30°C. One ml samples were collected for gDNA

purification at 0 and 2 hours post temperature shift. Samples were pelleted for 2 minutes at 13,000 rpm and washed once with 500 µl PBS. gDNA was purified using the Wizard® Genomic DNA Purification Kit (Promega, A1120) and stored at -20°C until further processing.

The strain-specific barcodes were amplified using indexed primers (**Table S7**) with the Collibri Library Amplification Master Mix (ThermoFisher, A38539050) according to the manufacturer's instructions, with 100 ng of gDNA as input, for 22 cycles. Barcodes were gel-purified, quantified, pooled equimolarly, and sequenced using the Illumina NovaSeq2 single-read method at the Sequencing Core Facility of the Max Planck Institute for Molecular Genetics (Berlin, Germany).

To calculate the fitness score of each deletion mutant from barcode abundance, BEAN-counter v2.6.1 was used with the following settings. A fixed amplicon read structure was used with an index at position 0, the common primer sequence GCAGGCGAGAAAGGAGAG at position 9, and the mutant barcode at position 27 of read 1. All other settings were used with default values. The sample before heat shock served as a reference to which all other time points were compared. All comparisons were performed with three biological replicates.

Fitness data was further analyzed using the R statistical computing language (v4.3.3). For annotation, protein-related information was fetched using the Uniprot REST API (29 February 2024). Average fitness per gene, condition, and time point was calculated as the arithmetic mean of the fitness score from three biological replicates. Differential fitness ΔF, e.g. for comparison of treatment *versus* control at a particular time point, was calculated as $\Delta F = F_{treatment} - F_{control}$. A two-sided Student's t-test was performed to evaluate statistical significance for each gene, and the obtained *p*-value was corrected using the Benjamini-Hochberg procedure[58].

<u>Growth curves</u>

Twenty-five ml of LB medium were inoculated with an overnight culture to a starting OD_{600} 0.05 in 250 ml Erlenmeyer flasks. Cultures were grown at 37°C 180 rpm in a water bath. OD_{600} was monitored every 30 minutes. The average and standard deviation of three biological replicates were calculated.

<u>Heat shock survival</u>

Cultures were inoculated in 25 ml LB medium to OD_{600} 0.05 and incubated in a water bath at 37°C 180 rpm. At OD_{600} 0.3 – 0.4, the cultures were exposed to a heat shock at 54°C for 2 hours, and samples for viability measurement were obtained every 30 minutes. Viability was measured by spotting 4 µl of 1:10 serial dilutions in PBS of each sample on LB agar. The plates were incubated overnight at 30°C, and the colony-forming units (CFU) were counted. The

CFU/ml was calculated for each sample, and relative viability was determined using samples

before heat shock as a reference.

<u>Isolation of protein aggregates</u>

Twenty-five ml of culture were harvested before and 15 minutes after heat shock at 54°C by

centrifugation for 5 minutes at 10,000 rpm. The pellets were washed once with buffer A (100

mM HEPES pH 8.0, 150 mM NaCl) and lysed by bead-beating in buffer B (100 mM HEPES

pH 8.0, 150 mM NaCl, cOmplete EDTA-free protease inhibitor) using 0.1 mm glass beads.

Afterwards, samples were centrifuged at 1000 x g for 2 minutes to obtain the cell extract. The

soluble fraction was separated from protein aggregates by centrifugation for 15 minutes at

11,000 x g at 4°C. To remove membrane proteins, the aggregates were resuspended in buffer

B containing 1% Triton X-100 and incubated for 1 hour at 4°C. Protein aggregates were

recovered by centrifugation at 11,000 x g, 4°C for 15 minutes and resuspended and incubated

as described above. Protein aggregates were washed again with buffer B + 0.5% Triton X-

100, and the detergent was removed in a final wash using buffer B. Pellets containing protein

aggregates were resuspended in buffer B + 2% SDS by boiling for 5 minutes at 95°C. Protein

concentration was measured using the Rapid Gold BCA Protein-Assay-Kit (Pierce™, A53225).

The percentage of aggregated protein was calculated to the total protein content in the cell

extract.

<u>Spot test</u>

Overnight cultures were diluted 1:1000 in LB medium without antibiotics and incubated at 37°C

180 rpm for 7-8 hours. The cultures were diluted to OD_{600} 1 in fresh LB medium in a 96-well

plate. Seven serial 1:10 dilutions were prepared in 1x PBS. Four µl of each dilution was spotted

on LB agar containing different stressors (concentration indicated in the figures). Plates were

freshly prepared before use.

<u>RNA purification</u>

Samples for RNA purification were collected at the indicated time points: exponential phase

(OD_{600} ~ 0.3), late exponential phase (OD_{600} ~ 1) and transition phase (100 minutes after late

exponential phase). Rifampicin was added to a final concentration of 100 µg/ml for 10 minutes.

Samples were harvested by mixing with the same volume of cold 1:1 acetone:ethanol and

centrifugated at 9,000 rpm for 5 minutes, 4°C. RNA purification was performed according to

Ref.[27] with slight modifications: The pellet was washed twice in 1 ml TE-sucrose buffer (50 mM

Tris-HCl pH 8.0, 10 mM EDTA, 25% sucrose) and centrifugated at 10,000 rpm for 5 minutes,

4°C. Cells were resuspended in 130 µl lysis buffer (20 mM Tris-HCl pH 8.0, 50 mM EDTA, 20

% sucrose) supplemented with 3.2 mg/ml lysozyme and incubated on ice for 5 minutes. Two

hundred µl of lysis executioner (2% SDS, 1 mg/ml Proteinase K) were added to the samples

and incubated at 95°C for 1.5 minutes. Samples were mixed with 1 ml of TRIzol (Ambion) and
incubated at room temperature. After 5 minutes, 250 µl chloroform was added to each sample
and mixed by vortexing. After 10 minutes at room temperature, the samples were centrifuged
for 10 minutes at 11,000 rpm. The RNA was isopropanol-precipitated at -20°C overnight. RNA
was pelleted at 11,000 rpm for 20 minutes at 4°C, washed with 1 ml 70% ethanol followed by
another centrifugation at 13,000 rpm for 10 minutes at 4°C. The pellet was air-dried and
resuspended in 50 µl nuclease-free water.

ISCP RNA sequencing

RNA sequencing was performed as previously described[59]. In brief, total RNA was treated with
TURBO DNase (Ambion, AM2238), depleted of ribosomal RNA using the riboPOOL rRNA
depletion kit for *B. subtilis* (siTOOLS; two reactions with 4.5 µg each) and purified using the
RNA Clean & Concentrator-5 kit (RCC-5; Zymo Research). RNA 5' triphosphates were
converted to monophosphates using RppH (NEB), and RNA was recovered using
phenol:chloroform:isoamyl alcohol extraction followed by ethanol precipitation. RNA ends were
repaired using T4 Polynucleotide Kinase (Thermo Scientific), before RNA was purified using
the RCC-5 kit and fragmented using an M220 Focused-ultrasonicator (Covaris) as described
by the manufacturer but with a treatment time of 100 s (microTUBE AFA Fiber Snap-Cap). After
purification using the RCC-5 kit, sequencing libraries were prepared from the recovered RNA
using the NEXTflex Small RNA-Seq Kit v3 (Bioo Scientific) according to the manufacturer's
protocol (including step G). Libraries were purified as described in step I of the NEXTflex Rapid
Directional qRNA-Seq Kit (Bioo Scientific) using Agencourt AMPure XP Beads (Beckman
Coulter; 0.9x bead-to-sample ratio). Sequencing was performed on an Illumina NovaSeq 6000
in paired-end mode with a read length of 100 bp at the Sequencing Core Facility of the Max
Planck Institute for Molecular Genetics (Berlin, Germany).

Processing of RNA sequencing data

FastQC (v0.11.5) was used to assess the quality of the data. Reads were filtered with a
minimum quality score of 10 and a length of at least 18 nt, cleaned from adapter sequences
using Cutadapt (v3.5)[60], and mapped against the *B. subtilis* reference genome (NC_000964.3)
using STAR (v2.7.10a_alpha_220314)[61] in 'random best' and 'end to end' modes. Resulting
BAM files were sorted and indexed with Samtools (v1.9)[62], and PCR duplication artefacts were
removed using UMI Tools (v1.1.0)[63]. Gene counts of annotated RefSeq genes were
determined with featureCounts (v2.0.3)[64] and differentially expressed genes (DEG) were
calculated with three replicates per condition and time point using DESeq2 (v1.24.0)[65]. *p*-
values were corrected for multiple testing using the Benjamini-Hochberg method[58]. To call
DEGs, a threshold of adjusted *p*-value < 0.05 and absolute log2 fold change > 1 was used.

Genome coverage profiles (total, 5' and 3' ends) were generated using a custom script utilizing

the HTSeq library (v2.0.2)[66]. We performed the aforementioned analysis steps using a

customized pipeline implemented in Snakemake (v7.14.0)[67].

<u>Identification of Y complex cleavage sites</u>

The Y-complex-dependent cleavage positions were identified following the previously

published procedure[59]. In brief, the genome coverage data was prefiltered with a count per

million (cpm) value ≥0.05, and only the RNA ends displaying a cpm ≥ 5 were further analyzed.

We carried out differential expression analysis of normalized 5′ and 3′ read ends by comparing

WT vs. $\Delta ymcA$ at the late exponential and transition phase. Y-complex-dependent ends were

identified using edgeR (v3.28.0)[68] with absolute log2 fold change ≥ 1 and FDR < 0.05, and kept

only if present in both comparisons. These results were further filtered with additional

parameters, i.e., the "proportion of ends (cleavage ratio)" and the "ratio of WT and $\Delta ymcA$

proportion of ends" as was previously described[59]. Moreover, to reduce the potential set of

false positive processing sites even further, we applied an additional cut-off on the "proportion

of ends" parameter, i.e., we kept only processing sites with a cleavage ratio of the reference

strain (WT) ≥ 95%-percentile of the cleavage ratio of the mutant strain.

<u>Impact of processing sites on transcript and operon expression</u>

To examine the impact of cleavage sites on transcript and operon expression, as well as other

genomic features such as untranslated regions (UTRs) and regulatory elements, we calculated

transcript per million (TPM) values from variance-stabilized data for each annotated RefSeq

gene (transcript level) and for each operon as defined in the BSGatlas (v1.1)[69] using the

DESeq2 package and a custom Python script. Furthermore, all Y-complex-dependent

cleavage sites were mapped to various genomic features, including coding regions, UTRs,

terminator regions, and riboswitches, as defined in the BSGatlas[69] (**Table S2**).

<u>Processing site sequence logo</u>

The sequence logos were generated with Logomaker (v0.8)[70], specifically for 5' and 3' end U-

type cleavage sites. For this, we extracted 30 nt sequences centered at the cleavage site and

calculated the logos with a GC content of 43.5%.

<u>RNA secondary structure prediction</u>

The RNA structure surrounding Y-complex-dependent cleavage sites was estimated by

calculating the MFE (ΔG in kcal/mol) utilizing RNAfold (v2.6.4)[71]. This analysis employed a

sliding window approach with 50 nt sequences, focusing on a 200 nt region centered around

the position of interest. The average MFE at each nucleotide was then calculated. To calculate

and visualize the background folding energy, we randomly draw potential processing sites from

the genome determined with the corresponding 5' or 3' end motif and calculate the average MFE at each nucleotide as described above. Potential processing sites were drawn 50 times independently.

<u>Overrepresentation analysis</u>

Annotations of operons were obtained from BSGatlas[69] and gene categories from Subtiwiki (24 July 2023)[72]. The overrepresentation analysis was performed using the R statistical computing language (v.4.2.2) and the Bioconductor package Category (v.2.64.0).

<u>Northern blot</u>

Biotinylated RNA probes were prepared using the HighYield T7 Biotin11 RNA Labeling Kit (UTP-based) (Jena Bioscience, RNT-101-BIOX) from PCR amplified fragments containing the promoter for the T7 RNA polymerase (**Table S11**). Probes were purified using the Monarch® Spin RNA Cleanup Kit (NEB, T2040) according to the manufacturer's instructions. Probes were designed taking into consideration the predicted cleavage sites.

One and a half µg of total RNA was separated on denaturing agarose gels (1% agarose, 20 mM MOPS, 5 mM NaOAc, 1 mM EDTA, 6.6% formaldehyde, pH 7.0) in 1x AGE-NB Running Buffer (20 mM MOPS, 5 mM NaOAc, 1 mM EDTA, 0.74% formaldehyde, pH 7.0). The gel was washed 3 times with milli-Q water and once with 10x SSC buffer, and the nitrocellulose membrane was washed once with milli-Q water and once with 10x SSC. The RNA was transferred using a vacuum blotter at 300 mbar covered with 10x SSC for 1.5 hours. After transfer, the membrane was washed for 2 minutes with 6x SSC, once with water, and dried and crosslinked at 254 nm 0.12 J twice. The membrane was developed with methylene blue, and the staining was removed by washing with 0.2x SSC containing 1% SDS, followed by washing with milli-Q water. The hybridization of 16S or 23S rRNA probes was performed using the North2South Chemiluminescent Hybridization and Detection Kit (Thermo) and North2South Hybridization Stringency Wash Buffer (2X) according to the manufacturer's instructions. Blots were developed in a Vilber imaging system.

<u>RT-qPCR</u>

Twenty µg of total RNA were treated with TURBO DNase (Ambion, AM2238) and purified using the Monarch® Spin RNA Cleanup Kit (NEB, T2040). Subsequently, cDNA was prepared using the LunaScript RT Mastermix (NEB, E3025) and 2.5 µg of DNase-treated RNA as input. Quantitative PCR was performed using the Luna Universal qPCR Master Mix (NEB, M3003) in a QuantStudio 5 Real-Time PCR System (Applied Biosystems). To calculate changes in *gfp* transcription, samples from the late exponential and transition phases were compared to those

from the exponential phase, with *pcp* serving as a reference gene using the Pfaffl method[73].

Primers used for RT-qPCR are listed in **Table S11**.

<u>Ribosome sedimentation in sucrose gradients</u>

Cultures of WT and Δ*ymcA* were grown until the exponential phase (OD_{600} ~ 0.3), late

exponential (OD_{600} ~ 1), and transition phase (100 minutes after the late exponential phase).

Chloramphenicol was added at a concentration of 100 µg/ml 2 minutes prior to sample

collection. Fifty ml of culture were pelleted at 4,500 rpm for 10 minutes at room temperature.

Pellets were transferred to ice and resuspended in 2 ml of lysis buffer (20 mM Tris-HCl pH 8.0,

100 mM NH_4Cl, 10 mM $MgCl_2$, 0.4% Triton X-100, 0.4% NP-40, 1 mM chloramphenicol, 100

U/µl DNase I), and flash-frozen. Cells were lysed under cryogenic conditions using 5 cycles of

mix-milling at 30 Hz for 1 minute. Lysates (240 µg RNA) were laid over a 10-50% sucrose

gradient in 20 mM Tris-HCl pH 8.0, 100 mM NH_4Cl, 10 mM $MgCl_2$, 0.5 mM DTT and resolved

at 35,000 rpm for 3 hours at 4°C. Gradients were analyzed using a piston fractionator (Biocomp

instruments), and absorbance at 260 and 280 nm was monitored.

<u>Sample preparation for MS-based proteomics</u>

Samples were collected at the indicated time points: late exponential phase (OD_{600} ~ 1) and

transition phase (100 minutes after the late exponential phase). For the proteomic changes

during heat shock, cultures were grown at 37°C until OD_{600} ~ 0.3 and shifted to 54°C. Samples

were collected at 0, 15, 30, and 60 minutes after heat shock. Ten ml of culture were collected

by centrifugation at 4,500 rpm for 10 minutes, and the pellet was stored at -80°C until further

processing.

Cell pellets were resuspended in 1 ml of cold PBS and centrifuged at 13,000 rpm for 5 minutes

at 4°C. The pellet was washed once more with PBS, resuspended in 150 µL of lysis buffer

(100 mM HEPES pH 8.0, 2% SDS, 20 mM TCEP, 60 mM CAA), and lysed using bead beating.

Cell debris was discarded by centrifuging at 13,000 rpm for 5 minutes at 4°C. The supernatant

was boiled at 95°C for 5 minutes, cooled down to room temperature, and the protein

concentration was estimated using the Rapid Gold BCA protein assay kit (Pierce). Samples

were flash-frozen in liquid nitrogen and stored at -80°C until further processing.

All samples were subjected to SP3 sample preparation[74] on an Agilent BRAVO liquid handling

robot. Ten µg of a 1:1 mixture of hydrophilic and hydrophobic carboxyl-coated paramagnetic

beads (SeraMag, #24152105050250 and #44152105050250, GE Healtcare) were added for

each µg of protein. Protein binding was induced by the addition of acetonitrile to a final

concentration of 50% (v/v). Samples were incubated at room temperature for 10 minutes. The

tubes were placed on a magnetic rack, and beads were allowed to settle for 3 minutes. The

supernatant was discarded, and beads were rinsed three times with 200 µL of 80% ethanol

without removing the tubes from the rack. Beads were resuspended in digestion buffer

containing 50 mM triethylammonium bicarbonate and both Trypsin (Serva) and Lys-C (Wako)

in a 1:50 enzyme-to-protein ratio. Protein digestion was performed for 14 hours at 37°C in a

PCR cycler. Afterward, the supernatant was recovered and dried down in a vacuum

concentrator.

<u>Proteomics using Mass spectrometry</u>

All samples obtained during the late exponential and transition phase were analyzed on an

Orbitrap Exploris (Thermo Scientific) equipped with a FAIMS Pro device and coupled to a 3000

RSLC nano UPLC (Thermo Scientific). Samples were loaded on a PepMap trap cartridge (300

µm i.d. x 5 mm, C18, Thermo) with 2% acetonitrile and 0.1% TFA at a flow rate of 20 µl/min.

Peptides were separated over a 50 cm analytical column (Picofrit, 360 µm O.D., 75 µm I.D.,

10 µm tip opening, non-coated, New Objective) that was packed in-house with Poroshell 120

EC-C18, 2.7 µm (Agilent). Solvent A consists of 0.1% formic acid in water. Elution was carried

out at a constant flow rate of 300 nl/min using a 60-minute method: 8-30% solvent B (0.1%

formic acid in 80% acetonitrile) within 23 minutes, 30-45% solvent B within 3 minutes, 45-98%

buffer B within 0.5 minutes, followed by column washing and equilibration. Data acquisition on

the Orbitrap Exploris was carried out using a data-dependent acquisition (**DDA**) method in

positive ion mode. MS survey scans were acquired from 375-1500 m/z in profile mode at a

resolution of 60,000. AGC target was set to 200% at a maximum injection time of 25 ms.

Peptides with charge states 2-6 were isolated within a window of 1.2 m/z and subjected to

HCD fragmentation at a normalized collision energy of 27%. The MS2 AGC target was set to

200%, allowing a maximum injection time of 22 ms. Product ions were detected in the Orbitrap

at a resolution of 15,000. Precursors were dynamically excluded for 20 s. The cycle time was

set to 1 s for each of the two FAIMS voltages of −45 and −65 V.

LC-MS/MS analysis of the heat shock samples was performed on an Ultimate 3000 UPLC

coupled to a Lumos hybrid mass spectrometer (Thermo Scientific). Peptides were loaded onto

a PepMap trap cartridge (300 µm i.d. x 5 mm, C18, Thermo) with 2% acetonitrile and 0.1%

TFA at a 20 µl/min flow rate. Peptides were separated over a 50 cm analytical column (Picofrit,

360 µm O.D., 75 µm I.D., 10 µm tip opening, non-coated, New Objective) that was packed in-

house with Poroshell 120 EC-C18, 2.7 µm (Agilent). Solvent A consists of 0.1% formic acid in

water. Elution was carried out at a constant flow rate of 250 nl/min using a 60-minute method:

8-30% solvent B (0.1% formic acid in 80% acetonitrile) within 23 minutes, 30-45% solvent B

within 3 minutes, 45-98% buffer B within 0.5 minutes, followed by column washing and

equilibration.

The mass spectrometer was operated in data-independent acquisition (**DIA**) and positive

ionization mode. MS1 full scans (375–1500 m/z) were acquired with a resolution of 60,000, a

normalized automatic gain control target value of 120%, and a maximum injection time set to 55 ms. Peptide fragmentation was performed using higher energy collision-induced dissociation and a normalized collision energy of 30%. MS2 spectra were acquired with a resolution of 30,000 (first mass 100 m/z) with 8 m/z DIA windows between 400 and 1200 m/z. A normalized automatic gain control target value of 100%, and a maximum injection time of 54 ms.

<u>Proteomics data analysis</u>

For samples at the late exponential and transition phase, Raw files were processed with MSFragger v3.5 using the default settings. Briefly, peak lists were extracted from raw files and searched using a Uniprot *Bacillus subtilis* database (v220530, taxonomy ID 224308) and a database containing sequences of common contaminants and decoys. Trypsin/P was set as enzyme specificity, allowing a maximum of two missed cleavages. Peptide matches were filtered to 1% FDR using PeptideProphet and Philosopher (v4.4.0).

For samples during heat shock, DIA-NN (DIA-NN v1.8.1) was employed for DIA data analysis. The main DIA-NN report, "report.txt", was used for all calculations. Fasta file (Bacillus_subtilis_Strain168_UP000001570.fasta) was digested with a maximum of one missed cleavage. Peptide length was restricted from 7 to 30 peptides, and the precursor m/z range was set from 300 to 1800. N-terminal methionine excision was enabled. Cysteine carbamidomethylation was selected as a fixed modification, and methionine oxidation and N-terminal acetylation as variable modifications. The maximum number of variable modifications was set to two. All other parameters were at default settings, including the 1% precursor FDR and enabled match between runs (MBR). The main DIA-NN report, "report.pg_matrix.tsv," was used for all calculations.

The R statistical computing language was used for data processing (v4.4.1). The main comparison of WT, $\Delta ymcA$, (p)ppGpp^{0,} and (p)ppGpp^{0} $\Delta ymcA$ mutants was based on samples from data-dependent acquisition (DDA) experiments. Raw peptide intensity values from label-free quantitation were imported as tabular output from MSFragger. The MSstats package (v4.12.1)[75] was used to convert raw intensities to an MSstats experiment. In this process, peptide quantification data was $\log_2$-transformed, median-normalized, and summarized to protein abundance. All features mapping to a protein were used for quantification, Tukey's median polish was used as a summarization strategy, and missing values were imputed using MSstats' default strategy. Feature abundance and inferred protein quantity were visually inspected using profile plots. Experimental groups were compared using MSstats' *groupComparison* function, and the similarity of conditions and replicates was inspected using principal component analysis (PCA, function *prcomp*). For annotation, protein-related information was fetched from Uniprot using the Uniprot REST API (14 June 2024). In addition

to the DDA experiments, a time series of WT and Δ*ymcA* mutants was obtained using data-
independent acquisition (DIA) proteomics. To quantify relative changes, the $\log_2$ fold change
of each time point was calculated using the zero time point as a reference. To quantify absolute
protein abundance for the ribosomal proteome sector, the coarse-graining strategy was used
as in Ref.[76,77]. Briefly, two protein sectors (groups) were created containing all annotated
proteins of ribosomal subunits. Raw MS1 intensities for all proteins of the respective group and
time points were summed up and divided by the total sum of protein intensities, yielding the
protein mass fraction per sector.

Data availability

674 Amplicon-seq and RNA-seq raw reads have been deposited at the European Nucleotide
675 Archive (ENA) under accession number PRJEB90661. Raw proteomic data have been
676 deposited at the PRIDE database under the accession number PXD065465 for DDA
677 proteomics of late exponential and transition phase samples. The DIA proteomics raw data
678 from the heat shock experiment can be accessed using the accession number PXD065462.

Author contributions

680 K.T., K.D., and F.A.C.; Conceptualization. K.D., F.A.C., K.T., R.A.-B., T.F.W., and K.A.:
681 Methodology. F.A.C., K.D., K.S., V.S., K.H., and F.K.: Investigation. K.D., F.A.C., R.A.-B., M.J.,
682 K.A., S.R., and K.T.: Formal analysis. F.A.C., K.D., R.A.-B., M.J.: Visualization. F.A.C., K.D.,
683 and K.T.: Writing – Original Draft. K.T. and E.C.: Funding acquisition. All authors: Writing –
684 Review & Editing.

Acknowledgments

686 This work was funded by the Deutsche Forschungsgemeinschaft (Tu106/8, SPP1879 for K.D.
687 and K.T., Leibniz Prize for E.C.) and the Max Planck Society (for E.C. and K.T.).

Figure 1. Genetic screening for genes involved in the heat shock response. A) Genetic screening experiment: the pooled library of deletion mutants (BKE) was grown at 30°C; when it reached an OD_{600nm} of 0.4, one fraction was exposed to 50°C while the other was maintained at 30°C. Genomic DNA (gDNA) was extracted from the library, and barcodes were amplified and sequenced using Next-Generation Sequencing. The fitness of each mutant at the mentioned temperatures and time points was calculated using BEANcounter. **B)** Relative fitness of each deletion mutant at 50°C (*y*-axis) and 30°C (*x*-axis) two hours post-temperature shift. Each point represents a deletion mutant, with mutants lacking RNase-related genes highlighted in magenta. The dotted lines indicate the range in which the relative fitness between the conditions differed by less than 2. Data represent the average of three biological replicates. A negative relative fitness value means that the abundance of the deletion mutant, compared to the rest of the population, was negatively affected by the treatment, while a positive value means an increased abundance in this condition. **C)** Changes in the viability (log_{10}CFU/ml) of WT and *ymcA, ylbF,* and *yaaT* deletion mutants to a severe heat shock at 54°C. Values were normalized to the viability at timepoint 0 min. The data represents the average ± standard error of three biological replicates.

Figure 2. Identification of Y-complex-dependent cleavages at late exponential and transition phase. A) Growth curves of WT, *ymcA*, *ylbF*, and *yaaT* deletion mutants in LB at 37°C. The blue bar marks the exponential phase. Arrows indicate sample collection for the ISCP method at the late exponential and transition phases. Three biological replicates are displayed with a smooth curve regression. **B)** Number of RNA 3' or 5' ends identified at the two growth phases. **C)** Fraction of identified 3' or 5' ends displaying U-type (unique) or S-type (stepped) profiles. **D)** Minimum free energy (MFE) was calculated on 50 nt windows for RNAs cleaved by the Y-complex with U-type ends (in magenta). Transcripts that share the same cleavage motif but are not processed were used as a control (in gray). rRNA genes were excluded from the analysis since they are highly structured. **E)** Fractions of ends identified in genes transcribed in polycistronic or monocistronic RNA. **F)** Count of RNA features where

Y-complex-dependent ends are detected at either growth phase. Annotations were obtained from
BSGatlas[69]. **G)** Overrepresentation analysis of pathways whose transcripts are cleaved by the Y-
complex during the late exponential phase. Operons were assigned to be involved in a pathway if they
code for one gene participating in such a pathway. As genes in the same operon can have different
functions, transcripts may be linked to multiple roles. Pathway annotations were retrieved from
SubtiWiki[72]. The color indicates the statistical significance of the enrichment. Biosyn = biosynthesis, acq
= acquisition, excl TCS = excluding two-component systems.

Figure 3. The Y-complex controls ribosome levels by initiating rRNA degradation. A) Number of
cleavages detected by ISCP in *B. subtilis* rRNA during the late exponential and transition phase. **B)**
Northern blot against 23S of WT and Δ*ymcA* cells at the late exponential phase, before and after
treatment with 100 μg/ml rifampicin to stop transcription. The methylene blue-stained membrane is
shown as a loading control. This image is representative of three biological replicates. **C)** Volcano plot
of protein quantification of WT and Δ*ymcA* strains comparing transition to late exponential phase. The
proteins composing the small (30S) and major (50S) ribosome subunits are highlighted in cyan and
magenta, respectively. The data shows the average of four biological replicates. Activity of the promoter
controlling **D)** *rrnJ* **(**rRNA) or **E)** the *rplK* (r-protein) transcription at late exponential and transition phase
compared to the exponential phase in WT and Δ*ymcA*. The promoters were cloned controlling *gfp*
transcription, which was measured using RT-qPCR. The *pcp* transcript was used as a reference.
Individual values and the average ± standard deviation of three biological replicates and two technical
replicates are shown. The statistical significance was tested using student's *t*-test. ns: *p*-value > 0.05.
F) Ribosome sedimentation profile in 10-50% sucrose gradients of WT and Δ*ymcA* in exponential, late
exponential, and transition phases. The 70S and 100S ribosomes are highlighted with a blue or red box,
respectively. The data is representative of three biological replicates.

Figure 4. The alarmones (p)ppGpp and the Y-complex are important for controlling ribosome levels. A) Growth curve of (p)ppGpp⁰ and (p)ppGpp⁰ Δ*ymcA* strains in LB at 37°C. The blue bar marks the exponential phase. Three biological replicates are displayed with a smooth curve regression. **B)** Volcano plot of proteomic changes of (p)ppGpp⁰ and (p)ppGpp⁰ Δ*ymcA* cultures comparing transition to exponential phase. The proteins composing the small (30S) and major (50S) ribosome subunits are highlighted in cyan and magenta, respectively. The data shows the average of four biological replicates. **C)** Comparison of r-proteins downregulation in WT, Δ*ymcA,* (p)ppGpp⁰, and (p)ppGpp⁰ Δ*ymcA* during the transition phase compared to the late exponential phase. The boxplot represents the interquartile range (IQR) and the median in the center. Whiskers show the variability outside quartile 1 (Q1) and Q3 and were calculated as Q1-1.5*IQR and Q5+1.5*IQR, respectively. Proteins are highlighted like in **B**. Alternative r-proteins were excluded from the analysis. The statistical significance was tested using paired Wilcoxon test. ****: *p*-value ≤ 0.0001. **D)** Proteome fraction (in %) of r-proteins mass, measured by mass spectrometry. The r-protein mass fraction was calculated for the indicated strains. Individual values and the average ± standard deviation of three biological replicates are shown. The statistical significance was tested using student's *t*-test. ns: *p*-value > 0.05, **: *p*-value ≤ 0.01, ***: *p*-value ≤ 0.001.

Figure 5. Control of ribosome levels and protein homeostasis in *ymcA* mutant during heat shock.
A) Northern blot against 23S of WT and Δ*ymcA* cells at exponential phase before or after a heat shock. The methylene blue-stained membrane is shown as a loading control. This image is representative of three biological replicates. **B)** Proteome fraction (in %) of r-proteins mass, measured by mass spectrometry. Individual values and the average ± standard deviation of three biological replicates are shown. The statistical significance was tested using student's *t*-test. *: *p*-value ≤ 0.05, **: *p*-value ≤ 0.01. **C)** Induction of CtsR and HrcA regulons after heat shock (50°C) in WT and Δ*ymcA*. Proteins were measured by mass spectrometry. The color bar shows the log$_2$ fold change compared to the control before heat shock. **D)** Percentage of aggregated proteins before and after a 15 min heat shock at 54°C in WT and Δ*ymcA*. Individual values and the average ± standard deviation of three biological replicates are shown. The statistical difference was tested using student's *t*-test. *: *p*-value ≤ 0.05, **: *p*-value ≤ 0.01, ***: *p*-value ≤ 0.001.

Figure 6. Control of ribosome levels by (p)ppGpp and the Y-complex. The alarmone (p)ppGpp reduces the biogenesis of new ribosomes by indirectly inhibiting the transcription of rRNA and r-proteins mRNA. (p)ppGpp inhibits GTPases that are involved in new ribosome assembly. Additionally, (p)ppGpp reduces translation by binding to specific GTPases. The Y-complex can initiate the degradation of ribosomes, which need to be transported to the membrane by unknown mechanisms. Exonucleases and AAA+ protease complexes might assist in ribosome degradation for the complete degradation of rRNA and r-proteins, which could partially replenish the nucleotide and amino acid pools during starvation.

References

1. Hecker, M., Schumann, W. & Völker, U. Heat-shock and general stress response in

Bacillus subtilis. *Mol. Microbiol.* **19**, 417–428 (1996).

2. Mogk, A. *et al.* The GroE chaperonin machine is a major modulator of the CIRCE heat

shock regulon of *Bacillus subtilis*. *EMBO J.* **16**, 4579–4590 (1997).

3. Krüger, E. & Hecker, M. The first gene of the *Bacillus subtilis clpC* operon, *ctsR*, encodes

a negative regulator of its own operon and other class III heat shock genes. *J. Bacteriol.*

180, 6681–6688 (1998).

4. Elsholz, A. K. W., Michalik, S., Zühlke, D., Hecker, M. & Gerth, U. CtsR, the Gram-

positive master regulator of protein quality control, feels the heat. *EMBO J.* **29**, 3621–

3629 (2010).

5. Krüger, E., Witt, E., Ohlmeier, S., Hanschke, R. & Hecker, M. The Clp proteases of

Bacillus subtilis are directly involved in degradation of misfolded proteins. *J. Bacteriol.*

182, 3259–3265 (2000).

6. Alver, R. *et al.* The control of protein arginine phosphorylation facilitates proteostasis by

an AAA+ chaperone protease system. Preprint at

https://doi.org/10.1101/2022.09.15.508104 (2023).

7. Schäfer, H. *et al.* The alarmones (p)ppGpp are part of the heat shock response of

Bacillus subtilis. *PLOS Genet.* **16**, e1008275 (2020).

8. Clay, K. J., Yang, Y., Clark, C. & Petrascheck, M. Proteostasis is differentially modulated

by inhibition of translation initiation or elongation. *eLife* **12**, e76465 (2023).

9. Cherkasov, V. *et al.* Coordination of translational control and protein homeostasis during

severe heat stress. *Curr. Biol.* **23**, 2452–2462 (2013).

10. Schramm, F. D., Schroeder, K. & Jonas, K. Protein aggregation in bacteria. *FEMS*

Microbiol. Rev. **44**, 54–72 (2020).

11. Schäfer, H. *et al.* Spx, the central regulator of the heat and oxidative stress response in

B. subtilis, can repress transcription of translation-related genes. *Mol. Microbiol.*

mmi.14171 (2018).

12. Nakano, S., Erwin, K. N., Ralle, M. & Zuber, P. Redox-sensitive transcriptional control by

a thiol/disulphide switch in the global regulator, Spx. *Mol. Microbiol.* **55**, 498–510 (2005).

13. Runde, S. *et al.* The role of thiol oxidative stress response in heat-induced protein

aggregate formation during thermotolerance in *Bacillus subtilis*. *Mol. Microbiol.* **91**,

1036–1052 (2014).

14. Rochat, T. *et al.* Genome-wide identification of genes directly regulated by the pleiotropic

transcription factor Spx in *Bacillus subtilis*. *Nucleic Acids Res.* **40**, 9571–9583 (2012).

15. Diez, S., Ryu, J., Caban, K., Gonzalez, R. L. & Dworkin, J. The alarmones (p)ppGpp
directly regulate translation initiation during entry into quiescence. *Proc. Natl. Acad. Sci.*
117, 15565–15572 (2020).
16. Corrigan, R. M., Bellows, L. E., Wood, A. & Gründling, A. ppGpp negatively impacts
ribosome assembly affecting growth and antimicrobial tolerance in Gram-positive
bacteria. *Proc. Natl. Acad. Sci.* **113**, (2016).
17. Pausch, P. *et al.* Structural basis for (p)ppGpp-mediated inhibition of the GTPase RbgA.
J. Biol. Chem. **293**, 19699–19709 (2018).
18. Krásný, L., Tišerová, H., Jonák, J., Rejman, D. & Šanderová, H. The identity of the
transcription +1 position is crucial for changes in gene expression in response to amino
acid starvation in *Bacillus subtilis*. *Mol. Microbiol.* **69**, 42–54 (2008).
19. Krásný, L. & Gourse, R. L. An alternative strategy for bacterial ribosome synthesis:
Bacillus subtilis rRNA transcription regulation. *EMBO J.* **23**, 4473–4483 (2004).
20. Koo, B.-M. *et al.* Construction and analysis of two genome-scale deletion libraries for
Bacillus subtilis. *Cell Syst.* **4**, 291-305.e7 (2017).
21. Msadek, T. *et al.* ClpP of *Bacillus subtilis* is required for competence development,
motility, degradative enzyme synthesis, growth at high temperature and sporulation. *Mol.*
Microbiol. **27**, 899–914 (1998).
22. Hajdusits, B. *et al.* McsB forms a gated kinase chamber to mark aberrant bacterial
proteins for degradation. *eLife* **10**, e63505 (2021).
23. Matavacas, J., Hallgren, J. & von Wachenfeldt, C. *Bacillus subtilis* forms twisted cells
with cell wall integrity defects upon removal of the molecular chaperones DnaK and
trigger factor. *Front. Microbiol.* **13**, (2023).
24. Schumann, W. Regulation of bacterial heat shock stimulons. *Cell Stress Chaperon.* **21**,
959–968 (2016).
25. DeLoughery, A., Dengler, V., Chai, Y. & Losick, R. Biofilm formation by *Bacillus subtilis*
requires an endoribonuclease-containing multisubunit complex that controls mRNA levels
for the matrix gene repressor SinR. *Mol. Microbiol.* **99**, 425–437 (2016).
26. DeLoughery, A., Lalanne, J.-B., Losick, R. & Li, G.-W. Maturation of polycistronic mRNAs
by the endoribonuclease RNase Y and its associated Y-complex in *Bacillus subtilis*. *Proc.*
Natl. Acad. Sci. **115**, E5585–E5594 (2018).
27. Broglia, L. *et al.* An RNA-seq based comparative approach reveals the transcriptome-
wide interplay between 3′-to-5′ exoRNases and RNase Y. *Nat. Commun.* **11**, 1587
(2020).
28. Taggart, J. C. *et al.* A high-resolution view of RNA endonuclease cleavage in *Bacillus*
subtilis. *Nucleic Acids Res.* **53**, gkaf030 (2025).

29. Khemici, V., Prados, J., Linder, P. & Redder, P. Decay-initiating endoribonucleolytic
cleavage by RNase Y is kept under tight control via sequence preference and sub-
cellular localisation. *PLOS Genet.* **11**, e1005577 (2015).
30. Le Scornet, A. *et al.* Critical factors for precise and efficient RNA cleavage by RNase Y in
Staphylococcus aureus. *PLOS Genet.* **20**, e1011349 (2024).
31. Shahbabian, K., Jamalli, A., Zig, L. & Putzer, H. RNase Y, a novel endoribonuclease,
initiates riboswitch turnover in *Bacillus subtilis*. *EMBO J.* **28**, 3523–3533 (2009).
32. Chao, Y. *et al.* In vivo cleavage map illuminates the central role of RNase E in coding
and non-coding RNA pathways. *Mol. Cell* **65**, 39–51 (2017).
33. Kime, L., Jourdan, S. S., Stead, J. A., Hidalgo-Sastre, A. & McDowall, K. J. Rapid
cleavage of RNA by RNase E in the absence of 5′ monophosphate stimulation. *Mol.*
Microbiol. **76**, 590–604 (2010).
34. Davis, J. H. & Williamson, J. R. Structure and dynamics of bacterial ribosome biogenesis.
Philos. Trans. R. Soc. B Biol. Sci. **372**, 20160181 (2017).
35. Piir, K., Paier, A., Liiv, A., Tenson, T. & Maiväli, Ü. Ribosome degradation in growing
bacteria. *EMBO Rep.* **12**, 458–462 (2011).
36. Eymann, C., Homuth, G., Scharf, C. & Hecker, M. *Bacillus subtilis* functional genomics:
global characterization of the stringent response by proteome and transcriptome
analysis. *J. Bacteriol.* **184**, 2500–2520 (2002).
37. Tagami, K. *et al.* Expression of a small (p)ppGpp synthetase, YwaC, in the (p)ppGpp^0
mutant of *Bacillus subtilis* triggers YvyD-dependent dimerization of ribosome.
MicrobiologyOpen **1**, 115–134 (2012).
38. Beckert, B. *et al.* Structure of the *Bacillus subtilis* hibernating 100S ribosome reveals the
basis for 70S dimerization. *EMBO J.* **36**, 2061–2072 (2017).
39. Carabetta, V. J. *et al.* A complex of YlbF, YmcA and YaaT regulates sporulation,
competence and biofilm formation by accelerating the phosphorylation of Spo0A: YmcA-
YlbF-YaaT complex stimulates the phosphorelay. *Mol. Microbiol.* **88**, 283–300 (2013).
40. Tortosa, P., Albano, M. & Dubnau, D. Characterization of *ylbF*, a new gene involved in
competence development and sporulation in *Bacillus subtilis*. *Mol. Microbiol.* **35**, 1110–
1119 (2000).
41. Branda, S. S. *et al.* Genes involved in formation of structured multicellular communities
by *Bacillus subtilis*. *J. Bacteriol.* **186**, 3970–3979 (2004).
42. Segev, E., Smith, Y. & Ben-Yehuda, S. RNA Dynamics in aging bacterial spores. *Cell*
148, 139–149 (2012).
43. Lehnik-Habrink, M. *et al.* RNase Y in *Bacillus subtilis*: a natively disordered protein that Is
the functional equivalent of RNase E from *Escherichia coli*. *J. Bacteriol.* **193**, 5431–5441
(2011).

44. Dubnau, E., DeSantis, M. & Dubnau, D. Formation of a stable RNase Y-RicT (YaaT)
complex requires RicA (YmcA) and RicF (YlbF). *mBio* e01269-23 (2023).

45. Hinrichs, R., Pozhydaieva, N., Höfer, K. & Graumann, P. L. Y-complex proteins show
RNA-dependent binding events at the cell membrane and distinct single-molecule
dynamics. *Cells* **11**, 933 (2022).

46. Lehnik-Habrink, M. *et al.* The RNA degradosome in *Bacillus subtilis*: identification of
CshA as the major RNA helicase in the multiprotein complex. *Mol. Microbiol.* **77**, 958–
971 (2010).

47. Commichau, F. M. *et al.* Novel activities of glycolytic enzymes in *Bacillus subtilis*. *Mol.*
Cell. Proteomics **8**, 1350–1360 (2009).

48. Durand, S. & Condon, C. RNases and Helicases in Gram-positive Bacteria. *Microbiol.*
Spectr. **6**, 10.1128/microbiolspec.rwr-0003–2017 (2018).

49. Kuroda, A. *et al.* Role of inorganic polyphosphate in promoting ribosomal protein
degradation by the Lon protease in *E. coli*. *Science* **293**, 705–708 (2001).

50. Dimitrova-Paternoga, L. *et al.* Structural basis of ribosomal 30S subunit degradation by
RNase R. *Nature* **626**, 1133–1140 (2024).

51. Boylan, S. A., Redfield, A. R., Brody, M. S. & Price, C. W. Stress-induced activation of
the sigma B transcription factor of *Bacillus subtilis*. *J. Bacteriol.* **175**, 7931–7937 (1993).

52. Njenga, R. K. *et al.* Ribosome-inactivation by a class of widely distributed C-tail anchored
membrane proteins. *Structure* **32**, 2259-2275.e6 (2024).

53. Wyant, G. A. *et al.* NUFIP1 is a ribosome receptor for starvation-induced ribophagy.
Science **360**, 751–758 (2018).

54. Gemin, O. *et al.* Ribosomes hibernate on mitochondria during cellular stress. *Nat.*
Commun. **15**, 8666 (2024).

55. Lehnik-Habrink, M. *et al.* RNA processing in *Bacillus subtilis*: identification of targets of
the essential RNase Y: Targets of the essential RNase Y in *Bacillus subtilis*. *Mol.*
Microbiol. **81**, 1459–1473 (2011).

56. Anderson, B. W., Fung, D. K. & Wang, J. D. Regulatory themes and variations by the
stress-signaling nucleotide alarmones (p)ppGpp in Bacteria. *Annu. Rev. Genet.* **55**, 115–
133 (2021).

57. Driller, K., Cornejo, F. A. & Turgay, K. (p)ppGpp – an important player during heat shock
response. *microLife* **4**, uqad017 (2023).

58. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A practical and
powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300
(1995).

59. Lécrivain, A.-L. *et al.* In vivo 3′-to-5′ exoribonuclease targetomes of *Streptococcus*
pyogenes. *Proc. Natl. Acad. Sci.* **115**, 11814–11819 (2018).

60. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17**, 10 (2011).

61. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

62. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

63. Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in Unique
Molecular Identifiers to improve quantification accuracy. *Genome Res.* **27**, 491–499
(2017).

64. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

65. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

66. Putri, G. H., Anders, S., Pyl, P. T., Pimanda, J. E. & Zanini, F. Analysing high-throughput
sequencing data in Python with HTSeq 2.0. *Bioinformatics* **38**, 2943–2945 (2022).

67. Mölder, F. *et al.* Sustainable data analysis with Snakemake. Preprint at
https://doi.org/10.12688/f1000research.29032.2 (2021).

68. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* **40**, 4288–
4297 (2012).

69. Geissler, A. S. *et al.* BSGatlas: a unified *Bacillus subtilis* genome and transcriptome
annotation atlas with enhanced information access. *Microb. Genomics* **7**, 000524 (2021).

70. Tareen, A. & Kinney, J. B. Logomaker: beautiful sequence logos in Python.
Bioinformatics **36**, 2272–2274 (2020).

71. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, 26 (2011).

72. Elfmann, C., Dumann, V., van den Berg, T. & Stülke, J. A new framework for SubtiWiki,
the database for the model organism *Bacillus subtilis*. *Nucleic Acids Res.* **53**, D864–
D870 (2025).

73. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic Acids Res. **29**, 45e–445 (2001).

74. Hughes, C. S. *et al.* Single-pot, solid-phase-enhanced sample preparation for proteomics
experiments. *Nat. Protoc.* **14**, 68–85 (2019).

75. Kohler, D. *et al.* MSstats Version 4.0: Statistical analyses of quantitative Mass
Spectrometry-based proteomic experiments with chromatography-based quantification at
scale. *J. Proteome Res.* **22**, 1466–1482 (2023).

76. Jahn, M. *et al.* Growth of cyanobacteria is constrained by the abundance of light and
carbon assimilation proteins. *Cell Rep.* **25**, 478-486.e8 (2018).

77. Jahn, M. *et al.* Protein allocation and utilization in the versatile chemolithoautotroph

Cupriavidus necator. eLife **10**, e69019 (2021).