1   **The *Drosophila* proventriculus lacks stem cells but compensates for age-related cell loss**
2   **via endoreplication-mediated cell growth**
3

4   **Authors:** Ben Ewen-Campen[1], Weihang Chen[1], Sudhir Gopal Tattikota[1], Ying Liu[1], Yanhui Hu[1],
5   Norbert Perrimon[1,2]
6
7   **Affiliations:**
8
9   1. Department of Genetics, Blavatnik Institute, Harvard Medical School, Boston, MA 02115,
10   USA
11   2. Howard Hughes Medical Institute, Boston, MA 02115, USA
12
13
14 **Corresponding author**: Ben Ewen-Campen and Norbert Perrimon
15 **Email:** bewencampen@genetics.med.harvard.edu, perrimon@genetics.med.harvard.edu

Abstract

The *Drosophila* proventriculus is a bulb-shaped structure at the juncture of the foregut and the midgut, which plays important roles in ingestion, peritrophic membrane synthesis, and the immune response to oral pathogens. A previous study identified a population of cells in the proventriculus which incorporate bromodeoxyuridine (BrdU), a marker of DNA synthesis, and proposed that these cycling cells are multipotent stem cells that replace dying cells elsewhere in the tissue. Here, we re-investigate these cycling cells and find that they do not undergo mitosis, do not generate clonal lineages, and do not proliferate in response to tissue damage, and are therefore not stem cells. Instead, we find that these cells continually endoreplicate throughout the fly's life, increasing their ploidy and cell volume, while at the same time nearby cells are lost into the gut lumen in an age-dependent process. These cells play a critical role in the synthesis of peritrophic membrane components, and we show that when endocycling in these cells is experimentally increased or decreased, nuclear and cellular volume as well as peritrophic membrane synthesis is concomitantly enhanced or reduced. Further, we show that blocking endocycling in these cells leads to functional defects in the peritrophic membrane, ultimately leading to increased susceptibility to orally infectious bacteria. Altogether, we show that continual endocycling of these cells is critical for maintaining tissue size and function over the course of the adult lifespan.

Introduction

The *Drosophila* gut is a highly regionalized organ with distinct functional domains along its length [1–3]. Within each domain, in addition to primary roles such as nutrient absorption and immunity, there are cellular mechanisms to ensure that the gut is repaired and renewed following tissue damage or cell loss. In the midgut, there are multipotent intestinal stem cells (ISCs) that continually divide to replace the differentiated cell types of the gut, both during homeostasis [4,5] and in response to tissue damage [6,7]. Analogous ISCs are present in the mammalian small intestine [8,9], which is functionally comparable to the *Drosophila* midgut [10].

In contrast to the midgut, in the hindgut there is no evidence of multipotent stem cells [11,12]. Instead, hindgut cells respond to tissue damage by undergoing extensive endocycling, a specialized cell cycle in which DNA is replicated without undergoing mitosis, thus increasing the ploidy and nuclear size of hindgut cells [11]. In *Drosophila*, this endocycling response to tissue damage or wounding is conserved in the adult epidermis [11,13,14], and is important to maintain tissue integrity in the absence of stem cell activity.

At the anterior of the gut, where the foregut meets the midgut, there is a specialized bulb-shaped structure called the proventriculus (also known as the cardia, based on its shape). A previous study [15] identified a ring of cells in the proventriculus which incorporate the nucleotide analog BrdU, a marker of S phase, and proposed that these cycling cells are stem cells that replace dying cells elsewhere in the proventriculus.

Here, we re-investigate the function of these previously-identified cycling cells in the
proventriculus. We find no evidence that these cells undergo mitosis, generate clonal lineages,
or respond to tissue damage, either by mitosis or by increasing endocycling. Instead, we find
that these cells continually endocycle throughout the fly's life, at a rate that varies with systemic
nutritional signals. Further, we find that cells from this region of the proventriculus are
progressively lost into the gut lumen as the fly ages, and the remaining nuclei and cells become
progressively larger in volume via endocycling. Lastly, we use functional studies to show that
blocking endocycling in these cells disrupts peritrophic membrane formation, whereas
inappropriately enhancing endocycling leads to increased peritrophic membrane formation. We
conclude that the proventriculus does not contain stem cells, but that it instead compensates for
age-related cell loss by endocycling-mediated cell growth to maintain tissue function.
71
72   **Results and Discussion**
73
The proventriculus contains a ring of cycling cells in Zone 4. The proventriculus consists of
a doubly-folded tubular epithelium and associated muscles and neuronal structures (Figure 1A)
[16–18]. It forms during embryogenesis by the invagination of an anterior, ectodermal portion of
the embryonic gut (the future esophagus) into an adjacent segment of endodermal gut, creating
a bulb structure with an internal lumen [19–21]. In addition to serving as a muscular valve
surrounding the esophagus, the proventriculus plays a critical role in the synthesis of the
peritrophic membrane [16–18]. Studies in other Dipterans have shown that the proventriculus is
capable of producing peritrophic membrane at a rate exceeding six mm/hour at 25ºC,
representing an enormous capacity for synthesis and secretion [22,23]. In addition to peritrophic
membrane synthesis, the proventriculus plays important roles in the immune system [24,25],
and as a site for regulation of the microbiota [26].
85
86   Morphological studies have identified six distinct "zones" arranged linearly along the
87   proventriculus epithelium, based upon cellular morphology and on their proposed role in the
88   synthesis of the peritrophic membrane [16]. These anatomical zones have been shown via
89   single cell RNAseq (scRNAseq) to largely correspond to transcriptionally distinct cell types [25].
90
91   Zone 4 is a ring-shaped structure that encircles the esophagus at the anterior of the
92   proventriculus (Figure 1A). A previous study identified a ring of cells located at the posterior limit
93   of Zone 4 which consistently incorporate BrdU, indicating that these cycles progress through S
94   phase in adult flies [15]. This study proposed that these cycling cells are multipotent stem cells
95   that proliferate and replace dying cells elsewhere in the proventriculus [15].
96
Functionally, Zone 4 cells are highly secretory and play an important role in the synthesis of
chitin and other peritrophic membrane components [16,25,27]. scRNAseq studies indicate it is
enriched for genes involved in chitin synthesis and chitin modifications, including *chitin synthase*
2 (chs2) [25] which can be visualized with *chs2-Gal4 > UAS:2XEGFP* (Figure 1B, left). Studies
in larval flies confirm that mutations in *chs2* lead to defects in peritrophic membrane synthesis
[27], and visualizing the peritrophic membrane with fluorescent HPA lectin [28,29] confirms
previous observations that peritrophic membrane components is highly enriched in the lumen

directly abutting Zone 4 cells (Figure 1B, middle.) Zone 4 cells also express high levels of the
WNT pathway ligand Wingless (Figure 1C, right), as reported by [15].
106
To re-investigate the function of these cycling cells, we fed flies with EdU for two days and
confirmed the presence of a ring of EdU+ cells at the posterior edge of Zone 4 (Figure 1C, left).
EdU+ cells are occasionally observed elsewhere in the proventriculus (see e.g. Figure 1G, 2A),
but here we focus exclusively on the EdU+ population in Zone 4. Previous studies suggested
that, in addition to Wg expression, JAK-STAT signaling is also enriched in this area [15], and
when we co-labeled proventriculi for EdU and 10XSTAT:GFP, a JAK-STAT pathway reporter,
we found that the EdU+ population is directly anterior to STAT+ cells, suggesting that EdU+
cells are enriched at the posterior edge of Zone 4, with 10XSTAT expression representing the
anterior limit of Zone 3 (Figure 1C, right).
116
117   To characterize the dynamics of EdU incorporation over time, we fed flies with EdU for 1, 3, or 5
118   days. We performed this experiment with two standard lab genotypes, w^{1118} and *OreR,* and
119   observed that the number of labeled cells increased linearly with time (Figure 1D), at a rate of
120   between 3.3 (w^{1118}) and 4.1 (*OreR*) EdU+ cells per day, suggesting that cells in this region are
121   continually entering S phase at a relatively constant rate.
122
To test whether individual cells cycle continuously over time or whether they cease cycling after
a discrete number of S phases, we performed an EdU pulse-chase experiment by feeding EdU
for 3 days, followed by a 20 day chase in the absence of EdU (Figure S1). After 20 days, we
observed a marked decrease in EdU+ cells in Zone 4, suggesting that the EdU signal was
diluted by continued cycling (Figure S1). This dilution over time is similar to what is observed for
the continually cycling ISCs (mitotic) and enterocytes (ECs; endocycling) of the posterior midgut
(Figure S1), and contrasts with what is observed in the hindgut, where BrdU signal is retained
for 20 days due to a lack of continual cycling under homeostasis [11].
131
In these initial observations, we noted progressive changes in the nuclear morphology of EdU+
cells. Specifically, after 1d of EdU feeding, Edu+ nuclei tended to be small in volume and
located at the posterior limit of Zone 4 (Figure 1D, left) whereas after 5d of feeding labeled
nuclei were larger and could be seen a few cell diameters anteriorly (Figure 1D, right). This
increase in nuclear size suggests that EdU+ incorporation in these cells may reflect an increase
in genomic size via endocycling, a modified cell cycle in which cells become polyploid by
replicating DNA without undergoing mitosis or cell division, rather than mitotic activity as
previously suggested.
140
It has previously been shown that systemic nutritional signals tightly correlate with endocycle
progression in a variety of cellular contexts in the *Drosophila* larva, whereas mitotic activity is
not clearly linked to nutritional status [30,31]. We tested whether the EdU+ cells in Zone 4 are
responsive to nutrition by varying the amount of dry active yeast, a primary source of protein
and other macronutrients for *Drosophila*, added to a standard minimal lab food (see Methods).
We found that the number of EdU+ cells increased in direct proportion to the amount of yeast

added to the fly's food ([Figure 1E](#)), suggesting that the cycling of these cells is influenced by
systemic nutritional cues.

We further tested this sensitivity to systemic nutrition using a genetic tumor model in the midgut
(*esg-Gal4, tubGal80^{ts} > UAS:yki^{3SA};* note that *esg-Gal4* is not expressed in the proventriculus),
which promotes systemic muscle wasting and cachexia via the upregulation of the systemic
insulin antagonist *Impl2,* the cytokine *upd3,* and other cytokines [32–34]. In cachectic flies, we
observed a significant decrease in the number of Zone 4 cells labeled with EdU+ ([Figure
S2A,B](#)), indicating that cycling in the cells is directly responsive to systemic reduction in insulin
signaling. We then tested whether over-expressing the insulin antagonists *Impl2, upd3*, or both
together, would lead to a similar reduction in EdU+ cells in the proventriculus. Indeed, when we
expressed these factors from other ISCs (*esg-Gal4*) or from muscles (*dMef2-Gal4*), both of
which decrease systemic insulin signaling, we observed a significant decrease in EdU+ cells of
the proventriculus ([Figure S2A,B](#)).

Altogether, these observations confirm that the adult proventriculus contains a ring of cells at
the posterior of Zone 4 which consistently cycle through S phase, and that they do so in
response to systemic nutritional signals. Further, initial observations of their cellular morphology
over time suggests that they may be endoreplicating rather than undergoing mitosis.

A molecular atlas and Gal4 toolkit for functional studies of the proventriculus. In order to
functionally characterize these EdU+ cells *in vivo*, we generated a toolkit of Gal4 lines to label
and manipulate various cell types of the proventriculus. First, we screened the K-Gut database
[35] for Gal4 lines that specifically label Zone 4. We identified four lines that labeled Zone 4,
inclusive of EdU+ cells, (GMR41F06-GAL4, GMR26D04-GAL4, GMR56D02-GAL4, and
GMR71F06-Gal4) and selected a particularly specific and consistent line, 71F06-Gal4, for
further study ([Figure 2A](#)).

Next, to identify molecular markers for other proventriculus cell types and to serve as a
foundation for further molecular study, we created a single cell transcriptional atlas of the
proventriculus using single nucleus RNAseq (snRNAseq). We sequenced a total of 22,332
nuclei from approximately 600 dissected proventriculus tissues, which ultimately resolved into
14 transcriptionally-defined clusters ([Figure 2B,C,D](#)). This atlas is available via an online data
portal (https://www.flyrnai.org/scRNA/).

We mapped the identity of these snRNAseq clusters to *in vivo* anatomy using cluster-specific
marker genes. One cluster was enriched for molecular markers of muscles (e.g. *Mhc, up*) and
two clusters were enriched for pan-neuronal markers (e.g. *nSyb, elav, brp*), one of which was
enriched for *Akh-* and *Impl2*-expressing cells and therefore likely represents the corpora
cardiaca [36–38]. To map the remaining 11 clusters to the *in vivo* proventriculus, we examined
the expression of top marker genes for each cluster using either Gal4 lines or split-intein Gal4
lines crossed to a ubiquitous reporter ([Figure 2C,D,E.](#)) [39].

We were able to identify the remaining clusters as anatomically distinguished epithelial cell
types. One cluster corresponded to the esophagus (marked by *Elovl7-Gal4*), another to Zone 1
(marked by *CG11550-Gal4* and *Cht7-Gal4*). At the clustering resolution we selected, Zone 2
and 3 cells were both found in a single cluster (enriched for *Ppn-Gal4* and *Cyp49a1-Gal4*). We
observed a distinct cluster enriched for *Lgr1* and *Delta* that corresponds to a population of cells
specifically at the anterior border of Zone 3, immediately posterior to Zone 4. We recovered
distinct clusters corresponding to Zone 4 (*Wnt4-Gal4*) and Zone 5 (*Ca-Ma2d-Gal4*), as well as a
cluster that corresponds to a discrete band of cells at the widest diameter of the proventriculus,
at the border of Zone 5 and 6 (marked by *CG10702-Gal4, Notum-Gal4*, *Mco4-Gal4[N-int]*). A
cluster enriched for *SecCl-Gal4* corresponded to Zone 6, and a cluster enriched *mex1-Gal4* and
P5CS-Gal4 corresponded to the anterior midgut. Two additional minor clusters (labeled
"unknown" and "anterior midgut 1" in [Figure 2B]) were enriched for markers corresponding to
multiple cell types and were excluded from further analysis. In a previous scRNAseq atlas of the
Drosophila midgut [40], midgut ISC clusters were enriched for the stem cell marker *esg*, as well
as cell-cycle related genes *PCNA* and *stg.* Notably, we did not observe notable expression of
any of these genes, nor any cluster-specific enrichment for any *cyclin* genes, in our
proventriculus atlas, whether in our Zone 4 cluster or elsewhere.

Our snRNAseq atlas is largely concordant with a recently published scRNAseq atlas of the
proventriculus [25], with minor deviations in the proportions of cells present in each cluster
([Figure S3]). This high degree of concordance suggests that these cell types we identify are
robust to experimental approach, as these two atlases were created using entirely different
single cell protols; Zhu (2024) sequenced single cells from freshly dissected and chemically
dissociated suspensions, while we sequenced isolated nuclei from flash frozen tissues. We also
note that our atlas was generated from two distinct biological samples: proventriculi from control
flies and from cachectic flies, and thus includes cells across a range of nutritional status (see
Methods). This snRNAseq atlas thus serves as a resource for studies of the proventriculus, and
allows us to identify cell type-specific Gal4 driver lines for functional experiments.

No evidence for mitosis in the proventriculus. Stem cells are defined as multipotent
precursor cells that undergo repeated asymmetric mitosis, giving rise to one daughter cell that
retains a stem identity and another daughter cell capable of differentiation. Thus, mitosis is a
critical feature of any stem cell system. A previous study of the proventriculus suggested that
rare mitoses do occur in this tissue, albeit at a rate that is seemingly out of proportion with the
steady numbers of cells passing through S phase [15].

To re-examine whether these cells undergo mitosis, we first stained guts from two different
control genotypes (20 *OreR* wildtype guts, 36 *71F06 > UAS:GFP* guts, total *n* = 56 guts) for
phosphorylated Histone 3 (pH3+), a marker of mitosis. We fed these flies colchicine overnight
prior to staining, which enriches for pH3+ cells by blocking exit from mitosis [11]. We observed
zero pH3+ cells in the epithelia of Zone 4 in these flies, or anywhere else in the proventriculus.

The cell cycle can be experimentally driven forward by over-expressing such oncogenes as
activated Yki or Ras, or by *Notch* loss-of-function, and in cells that are capable of mitosis these

manipulations will strikingly increase M phase. For example, in ISCs of the posterior midgut,
over-expression of *UAS-yki^{3SA}* [41–43] or *UAS-Ras1^A* [44–46]*,* or knock down of *Notch [5,47]* all
lead to dramatic over-proliferation via excess mitosis, leading to massive tumor growth. We
hypothesized that, if Zone 4 cells are competent to undergo mitosis, over-expression of such
cell cycle promoting factors should lead to an increase in mitotic cells.
239
To test whether these factors can drive mitotic proliferation in the proventriculus, we used
71F06-Gal4 to drive *UAS-yki^{3SA}, UAS-Ras1^A,* or *Notch-RNAi* in Zone 4 cells. These
manipulations caused a dramatic enlargement of Zone 4, increased cytoplasmic extrusions into
the gut lumen (Figure 3 and S4A,B), and a expansion of the domain that typically incorporates
EdU to cover the entire Zone 4 rather than a band at its posterior edge (Figure 3A). However,
we did not observe a single pH3+ cell under any of these conditions, which also included
colchicine treatment to enrich for pH3+ (Figure 3B). As a positive control, imaging the posterior
midguts from these same samples showed extensive pH3+ staining (Figure 3B).
248
These results demonstrate that EdU+ cells located in Zone 4 of the proventriculus do not
undergo mitosis either under homeostatic conditions or when experimentally driven through cell
cycle progression via ectopic oncogene expression. Furthermore, we note that despite
morphological deformities caused by over-expressing activated Yki or Ras1, we never saw
these cells travel from Zone 4 into any other portion of proventriculus, which provides further
evidence that EdU+ cells in Zone 4 are not capable of replacing damaged or dead cells in other
regions of the tissue as suggested by [15].
256
No evidence for clonal growth of Zone 4 cells in the adult proventriculus. Despite the
absence of pH3 staining, it is possible that rare mitoses occur. Another hallmark of adult stem
cells is the ability to generate multipotent clones in the adult organism. For example, in the adult
posterior midgut, MARCM-based lineage tracing of ISCs leads to the presence of GFP-labeled
clones throughout the midgut that contain all of the epithelial cell types that can be generated by
ISC differentiation [5,47]. We therefore used MARCM-based lineage tracing to test whether
there is clonal growth in the adult proventriculus.
264
As a positive control for our labeling protocol, we heat-shocked MARCM19A larvae during the
L3 larval stage, when this tissue is still undergoing developmental proliferation. We observed
large GFP+ clones throughout the proventriculus, both in Zone 4 and elsewhere, in every
sample examined, demonstrating that MARCM labeling functions as expected in this tissue
(Figure 4A). As a negative control, we examined GFP+ patterns in both midguts and
proventriculi from non heat-shocked flies, and observed small numbers of scattered GFP+ cells
throughout the proventriculus, as well as in the midgut. We scored proventriculi as having either
no GFP+ clones, between 1-5 small groups of GFP+ cells (typically 1-2 cells each), or having
large GFP+ clones that include >10 cells (Figure 4D).
274
275   To test for clonal growth during the adult stage, we heat-shocked MARCM19A flies several days
276   after eclosion, then reared them for varying lengths of time before scoring for GFP+ clones. We

277 challenged an additional cohort of flies with a 2-day starvation followed by re-feeding, which
278 increases ISC proliferation in the midgut [48,49].
279
As expected, in the posterior midgut we observed GFP+ clones in the midguts of heat-shocked
flies after 14 days, and we also found that these clones tended to be larger in flies that
experienced a 2 day starvation (Figure 4B). In contrast, in the proventriculus of these flies,
observed scattered singlet and doublet GFP+ cells throughout the tissue at roughly similar rates
between negative controls and heat-shocked flies (Figure 4C,D). This was true at 14 days after
heat-shock, 19 days after heat-shock, and 30 days at 18ºC after heat-shock (Figure 4C,D).
286
In the midgut, ISCs divisions increase dramatically upon tissue damage [6,7]. To test whether
there may be damage-responsive proliferation in the proventriculus, we repeated these MARCM
experiments in flies subjected to a variety of damage: bleomycin treatment, infection with
pathogenic *ECC15* bacteria, and 2000 RADs of X-rays [6,7]. Given that we observed that EdU+
incorporation increases with additional yeast content, we also performed MARCM experiments
with 10% yeast supplementation. We did not observe discernable GFP+ clones emanating from
Zone 4 or any other area of the proventriculus in any of these conditions (Figure S5). We note
that, unlike enterocytes of the midgut, Zone 4 cells do not directly contact the contents gut
lumen and therefore likely do not experience damage to the same extent as ECs, but
nonetheless we did not observe GFP+ clonal growth under any of these conditions. Altogether,
we find no evidence for clonal growth of the EdU+ cells in Zone 4 of the proventriculus.
298
Zone 4 cells do not cycle or proliferate in response to apoptotic damage in the
proventriculus. In the midgut, ISCs proliferate at low levels under homeostatic conditions, and
increase proliferation in response to tissue damage and/or dying enterocytes [6,7]. In contrast,
in the hindgut, cells do not mitotically proliferate in response to damage but instead undergo
extensive endoreplication [11,12]. A previous study in the proventriculus suggested that the
EdU+ population of cells in the proventriculus proliferates in order to homeostatically replace
dying cells elsewhere in the tissue [15].
306
To directly test how the EdU+ population of cells responds to nearby tissue damage, we used
the Gal4-UAS system to induce cell death in various cell types of the proventriculus by over-
expressing the proapoptotic factor *UAS-RPR*. We drove cell death in three different cell types of
the proventriculus: in Zone 5 (*Ca-Ma2d-Gal4*), in Zone 1 (*CG11550-Gal4*), and in the
esophagus (*Evolv7-Gal4*) (Figure 4). In these experiments, we silenced *UAS-RPR* throughout
development using *tubGal80^{ts}* and raising flies at 18ºC until several days post-eclosion, and
then activated tissue damage by shifting adult flies to 29ºC for 48 hours. We placed flies on
EdU-containing food for the duration of the tissue damage, and throughout a recovery phase of
2 days or 5 days at 18ºC following damage. If these EdU+ cells in Zone 4 proliferate to
compensate for dying cells, we would expect to see an increase in EdU+ cells in response to
damage, and likely EdU+ cells growing towards the area of damage.
318
We observed pyknotic nuclei, indicative of apoptosis, in the intended cell type in *UAS-RPR* flies,
and not in control flies (Figure 5A, B, C), indicating that this system induces apoptosis as

expected. However, in all three experiments we observed not an increase in EdU+ cells, but
rather a significant decrease (Figure 5.) These results are inconsistent with the hypothesis that
these cells proliferate in response to tissue damage elsewhere in the proventriculus. The
decrease in EdU+ cells in all cases further supports the previous observation that these cells
cycle in response to systemic cues indicative of organismal health.
326
Interestingly, when we drove apoptosis in the esophagus with *Elovl7-Gal4*, while we observed a
decrease in EdU+ cells in Zone 4, we did observe indications of massive endoreplication in the
esophagus itself: unusually large, EdU+ nuclei (Figure 5C, asterisks). This is similar to what has
been reported in the hindgut [11], and we note that both the foregut and hindgut are ectodermal
in origin [50]. In contrast, the remainder of the proventriculus, like the midgut proper, is
endodermal in origin and we did not observe any similar indications of endoreplication in the
damaged tissue (Figure 5A,B). We thus propose that damage-responsive endoreplication may
be a phenomenon primarily found in ectodermally-derived tissues, and note that it is also
observed in the epidermis which is ectodermal in origin[11,13,14].
336
Genetic lineage tracing using a highly specific split-intein Gal4 line that labels EdU+ cells
in Zone 4. As an additional method for lineage tracing independent of MARCM, we wished to
genetically label the descendants of Zone 4 EdU+ cells using a Gal4 line specific to these
cycling cells. However, we did not observe any discernible transcriptional signatures of the Zone
4 EdU+ cells, and were initially unable to identify a Gal4 driver to specifically label these cells
within Zone 4.
343
We thus used the split-intein Gal4 system to generate a genetic driver that specifically labels the
EdU+ population of cells in Zone 4 [39]. We generated a split-intein Gal4$^{N\text{-}int}$ line using the
71F06 enhancer fragment that drives expression in Zone 4 (Figure 5A), and then used our
snRNAseq atlas to identify *Delta* as a candidate gene which is primarily enriched in Zones 1-3
and in Zone 3-4 border cells, with some expression as well in Zone 4 (Figure 6A). When we
examined the intersectional labeling of *71F06-Gal4::Gal4-C-int ∩ Delta$^{Gal4\text{-}N\text{-}int}$ > UAS:2xEGFP*,
we observed strikingly specific expression in a band of cells at the posterior edge of Zone 4
(Figure 6A). We confirmed that these cells contain the large majority of EdU+ cells (Figure 6B).
352
To lineage trace these cells, we used the G-TRACE system combined with *tubGal80^{ts}*, which
labels current expression in RFP and lineage expression in GFP [51]. After development at
18ºC, we placed *71F06-Gal4::Gal4-C-int ∩ Delta$^{Gal4\text{-}N\text{-}int}$, tubGal80^{ts} > G-TRACE* at 29ºC to
initiate lineage tracing and examined the proventriculus after 1 day, 6 days at 10 days (Figure
6C,D). If these cells are proliferating and generating clones, we would expect to see
progressively larger areas of GFP+ / RFP- cells as descendants of these cells grow into the rest
of the tissue. However, while we did observe small numbers of GFP+ / RFP- cells immediately
adjacent to the GFP+/RFP+ domain, the number of these cells did not increase or move over
time, but were constant over a 10 day period (Figure 6C,D). These results are consistent with
leaky and/or minor dynamic changes in the expression pattern of our split-intein Gal4 line, rather
than with proliferation of cells in this domain.

364
365   Taken all together, we find no evidence that the EdU+ cells located in Zone 4 are stem cells.
366   We do not observe mitosis, nor the generation of MARCM clones, nor a proliferative response
367   to nearby tissue damage, nor any evidence of genetic cell lineages growing over time. We thus
368   propose that these cells must be endocycling through a cell cycle that lacks an M phase.
369
370   **Zone 4 cells progressively increase in ploidy over the fly lifespan, as others are lost into**
371   **the lumen.** The consistency with which EdU is incorporated into cells in Zone 4 suggests that
372   these cells are continually endocycling throughout the fly's life and thus increasing in ploidy.
373   Indeed, microscopy-based ploidy analysis [52] of Zone 4 cells indicated a wide range of ploidy
374   states, with the majority between 4N and 16N, and including cells up to 40N (Figure S6).
375
If this cell cycling reflects consistent endocycling throughout the fly's life, these cells should
progressively increase in nuclear volume as flies age [31,53]. To test this directly, we performed
a time-course experiment using confocal microscopy to quantify the number and volume of
Zone 4 nuclei over the course of fly aging. In this experiment, *71F06-Gal4 > UAS:GFP^stinger* flies
(hereafter, *UAS:GFP^NUC*), which express nuclear-localized GFP in Zone 4, were split into vials
either containing dried active yeast powder *ad libitum* or standard food lacking additional yeast
powder, and were examined at six timepoints between 3 days and 35 days after eclosion. At
each time-point, we collected 5 confocal z-stacks and quantified both the number of GFP+
nuclei as well as the 3D volume of these nuclei (Figure S6A).
385
We observed a consistent, monotonic increase in nuclear volume over the course of the fly's
life, supporting the hypothesis that these cells are endocycling and thereby increasing in ploidy
(Figure 7A,B,C). Consistent with the observation that supplemental yeast increases EdU
incorporation, we found that nuclear volume increased more quickly in flies which received dried
active yeast, but that by 28 days the average nuclear volume was not significantly different
between these treatments (Figure 7A,B,C). As an internal comparison, we also quantified the
volume of diploid ISCs from these same guts and using the same volume quantification pipeline.
At 3d after eclosion, many Zone 4 nuclei were approximately the same volume as diploid ISCs
(Figure S6B). Over the course of the fly's life, nuclear volume increased until at 35 days old the
largest nuclei were ~16X larger in volume than diploid ISCs (Figure S6B). Co-staining these
samples with the f-actin marker phalloidin indicated that Zone 4 cells remain mononucleate,
even at the latest time point (35d) when the tissue begins to lose a clearly monolayered
epithelial structure (Figure S6C).
399
Surprisingly, we observed that while the volume of Zone 4 nuclei progressively increased, the
total number of nuclei steadily decreased (Figure 7D). Over the first 14 days, cell numbers
decreased at equivalent rates for flies fed with or without yeast, but beginning at 21 days the
number fell faster for flies without added yeast (Figure 7D). Remarkably, throughout this time-
course, we observed that GFP+ nuclei were being shed into the gut lumen (Figure 7E,F and
S7). We quantified the number of GFP+ nuclei present in the lumen outside the Zone 4
epithelium, representing a snapshot of cell loss at the time of dissection, and found that the
number of cells shed into the lumen first started to increase at 14 days after eclosion, and

subsequently increased until reaching a maximum at 28 days (Figure 7D). The mechanism by
which these cells exit the epithelium is not known, but we observed holes forming in the F-actin-
rich barrier separating Zone 4 cells from the gut lumen, with GFP+ nuclei passing through these
holes into the lumen (Figure S7). We note that the patterns of cell loss correspond neatly with
our total cell counts, with the greatest decreases in cell numbers corresponding to those time-
points when more nuclei were being lost.
414
415   Together, these data suggest that Zone 4 nuclei are endocycling and thus increasing in ploidy
416   and nuclear volume at the same time that cells from this region are being lost into the lumen.
417   One simple explanation for these observations is that genomic copy number is increasing in
418   order to maintain metabolic output and tissue integrity as cells are progressively lost during
419   aging.
420
Disruptions to endocycling in Zone 4 lead to altered peritrophic membrane synthesis and
function. Increasing genomic copy number is a widespread mechanism to enhance synthesis
of proteins and other biological components in non-proliferative tissues [31,53,54]. Given the
role that Zone 4 cells play in the synthesis and secretion of chitin and other peritrophic
membrane components [25,27], we hypothesized that endocycling in this tissue is linked to
peritrophic membrane synthesis. We therefore used genetic manipulations to both increase and
decrease endocycling, and then visualized the peritrophic membrane and tested for functional
defects.
429
To increase endocycling, we overexpressed *UAS:CycE* or *UAS:Myc* in Zone 4 cells using
71F06-Gal4. CycE is a key regulator of both endocycling and the mitotic cell cycle, and drives
entry into S phase [31,55], and Myc is a transcription factor whose overexpression is sufficient
to drive endocycling and increased nuclear volume in many contexts [31,54,56,57]. In Zone 4
cells of the proventriculus, both of these manipulations led to increased EdU incorporation and
the formation of significantly enlarged nuclei after 7-9 days at 29ºC (Figure 8A, B and S9A). We
visualized the peritrophic membrane formation using HPA lectin staining, and observed
enlarged streams of glycosylated material emanating from Zone 4, forming thickened cords that
were not observed in controls. We interpret these to indicate that increased endocycling leads to
enhanced synthesis of peritrophic membrane components in Zone 4.
440
To inhibit endocycling, we noted that the cycling of these cells is highly dependent on nutritional
status. We therefore knocked down the insulin receptor (*InR*) in Zone 4, which mimics starvation
specifically in these cells. Using two independent RNAi constructs, we observed a near total
loss of EdU staining in *InR RNAi* animals after 6 and 15 days at 29ºC (Figure S9B,C) and a
concordant failure to increase in nuclear volume over time (Figure 8B). In control flies (*w^{1118}* or
mCherry-RNAi), Zone 4 nuclei steadily increase in nuclear volume between 6 days and 21 days
at 29ºC, where in *71F06-Gal4 > InR-RNAi* flies, these nuclei did not increase in volume (Figure
8B). The overall size of Zone 4, visualized with cytoplasmic *UAS:2XEGFP,* was dramatically
reduced by *InR RNAi*, indicating that nuclear volume and overall cell size are linked, as
expected (Figure 8C.)
451

Visualization of the peritrophic membrane in *71F06-Gal4 > InR-RNAi* revealed notably less
material emanating from Zone 4, and overall far less visible peritrophic membrane within the
proventriculus (Figure 8D). Consistent with the reduced nuclear volume and tissue size, the
quantity of peritrophic membrane emanating from Zone 4 in these animals appeared to be
reduced and altered (Figure 8D). These alterations in peritrophic membrane morphology are
consistent with the hypothesis that increased genomic copies allow for increased synthesis of
peritrophic membrane components.
459
To assess the functional consequences of this reduction in peritrophic membrane production,
we assayed their resistance to orally-ingested pathogen infection. The peritrophic membrane is
a critical barrier to pathogens [58–60], so we hypothesized that reduced endocycling in Zone 4
cells may lead to functional defects in pathogen resistance. Under standard conditions at 29ºC,
the lifespan *71F06-Gal4 > InR-RNAi* flies was indistinguishable from control flies (Figure 8E).
However, when these flies were challenged with the pathogenic bacteria *Pseudomonas*
aeruginosa (*PA14*), *InR RNAi* flies were far more susceptible to infection than controls (Figure
8F). This is consistent with a compromised peritrophic membrane.
468
These experiments also allowed us to test whether endocycling-based cell growth was
functionally causal for the loss of cells from Zone 4 during aging. If cell loss was driven by the
process of tissue growth via endocycling, then experimentally increasing endocycling via *UAS-*
Myc or *UAS-CycE* should lead to enhanced less loss. We counted the number of GFP+ cells in
these genotypes one day after shifting to 29ºC, and after 12 days (Figure S10). The number of
cells in each genotype decreased after 12 days by between 21% - 31%, but the percent
decrease between genotypes were not significantly different (Figure S10), despite the fact that
nuclear volume is greatly increased in these genotypes compared to controls (Figure 8A,B,
Figure S8). This suggests that the shedding of Zone 4 cells during aging is not driven by
endocycling itself, and that a distinct mechanism must govern the loss of cells.
479
480 Altogether, these experiments demonstrate that endocycling leads to increased nuclear and
481 cellular volume, ultimately increasing the production of peritrophic membrane components.
482 Given the observation that under wildtype conditions, Zone 4 cells are lost over the course of a
483 fly's life, we propose that endocycling is a compensatory mechanism to ensure that continued
484 production of sufficient quantities of peritrophic membrane components.
485
Conclusions. We find no evidence of stem cells in the *Drosophila* proventriculus. The EdU+
cells in Zone 4 do not undergo mitosis, do not produce clonal lineages, and do not cycle in
response to nearby tissue damage. Instead, we find that the cells in Zone 4 endocycle
continuously throughout the fly's life, independently of exogenous damage, while at the same
time cells are shed into the gut lumen as the fly ages. We find that blocking endocycling in Zone
4 causes defects in peritrophic membrane synthesis, and we propose that continual endocycling
ensures high levels of peritrophic membrane synthesis throughout the fly's life.
493
Endoreplication and polyploidy is extremely widespread in *Drosophila* and across animals and
plants, both under healthy, homeostatic conditions and in response to cell or tissue damage

[31,53,54,61,62]. In the *Drosophila* larva, many tissues cease mitotic proliferation during
embryogenesis, and subsequently grow via developmentally programmed endocycling [31,63].
In adult flies, many cell types become polyploid via endoreplication under wildtype conditions,
including the nurse cells and follicle cells of the ovary, the enterocytes of the midgut, and rectal
papillae cells [61], while other cell types endocycle in response to tissue or cellular damage,
including the epidermis in response to wounding [13,62], the hindgut in response to tissue injury
[11,12], and the adult brain in response to oxidative and DNA damage [64]. Mammals also
display widespread examples of both developmentally programmed endocycling as well as
damage-responsive endocycling, including for example in the mammalian liver which displays
high levels of endoreplication in response to damage and injury [31,53,54,61,62].
506
The pattern of endocycling that we observe in the proventriculus appears to be unique in certain
respects compared to other well-characterized examples of endoreplication in *Drosophila* and
other tissues. Unlike in the adult brain [64] or ectodermal gut regions including the hindgut
[11,12] or the esophagus (Figure 5E), endoreplication in the proventriculus does not occur in
response to tissue damage, but is instead reduced by nearby injury (Figure 5). And whereas
many other highly metabolically active cells become highly polyploid via endoreplication,
including the nurse cells or the follicle cells of the ovary, in those cell types endoreplication
appears to proceed for a specific number of cycles and then cease [65], rather than continual
endoreplication over the fly's lifespan. What we observe in the proventriculus appears to be a
continuous increase in ploidy throughout the adult stage, as part of a homeostatic system that
accounts for the steady loss of neighboring cells and also responds to the nutritional input of the
fly. We propose that this system may provide insights into how the endocycle can be regulated
over long time-scales, as well as into the mechanisms that link age-related cell loss to
endoreplication.
521
522 **Methods and Materials**
523
Experimental animals. Genotypes and sources for *Drosophila melanogaster* lines used in this
study are provided in Table S1. Flies were maintained at 18ºC, 25ºC, 29ºC, as indicated in the
text, on standard laboratory cornmeal food, except for EdU treatments, yeast supplementation
experiments, or drug treatments, which were maintained on Formula 4-24 Instant Blue Food
("Instant Blue Food", Carolina Biological Supply) as described below. All experiments were
performed on female flies.
530
EdU feeding and EdU+ cell counting. EdU (Thermo Fisher) was diluted to 0.2mM in water,
then used to hydrate an equal volume of Carolina Instant Blue Food, supplemented with yeast
powder as indicated in the text. Adult flies were transferred to EdU-containing food, and flipped
to fresh EdU-containing food every two days, for any experiment lasting longer than two days.
EdU signal was detected using Click-iT™ EdU Cell Proliferation Kit (Thermo Fisher). Except
where noted in the text, 5% yeast was used in EdU experiments (100 mg dried yeast per 2 mL
of food), and flies were < 2 weeks old. EdU+ cells were manually counted on a compound
epifluorescent microscope, specifically focusing on those EdU+ cells located in Zone 4.
539

Yeast variation experiments. To test the effect of supplemental dietary yeast on EdU incorporation, dried active yeast was added as a percentage by weight of Instant Blue Food.

Colchicine treatment. To enrich mitotic nuclei, flies were fed with 0.2 mg/mL colchicine (EMD Millipore) for 12 hours prior to dissection. Colchicine was diluted in water and used to dilute Instant Blue Food supplemented with 10% yeast.

Gut damage with X-rays, bleomycin, and *ECC15* infection. For X-ray damage, adult flies were irradiated with 0, 1000, or 2000 RADs in a TORREX 120D X-ray Inspection System (ScanRay Corporation). For gut damage with bleomycin, bleomycin (Sigma) was diluted to 25 ng/µL in water and used to rehydrate 2 mL of Instant Blue Food. Flies were fed bleomycin-contained food for two days at 29ºC to increase ingestion, and were flipped to fresh food each day, with water as a control. For *ECC15* treatment, 100mL of LB medium were inoculated with 20 µL of a *ECC15* glycerol stock and grown at 30ºC for 20-24 hours. *ECC15* cultures were then centrifuged for 30 minutes at 3000 rpm and resuspended in 1.4 mL of a 5% sucrose. Flies were starved for two hours at 29ºC, and then flipped to a vial containing only a kimwipe soaked in 700 µL of the *ECC15* solution, where they fed for 24 hours at 29ºC to induce gut damage. After gut damage, flies were maintained on standard food at 25ºC for the indicated amount of time.

Immunohistochemistry and confocal imaging. Tissues were dissected in 1X phospho-buffered saline (PBS), fixed for 25-30 minutes in 4% paraformaldehyde in PBS, permeabilized in PBS with 0.3% Triton-X (PBST), and stained following standard antibody staining procedures using 5% normal goat serum as a blocking agent. Filamentous actin was stained using Alexa 555-coupled phalloidin (Molecular Probes), peritrophic membranes were stained with Alexa647-coupled Lectin HPA (Thermo Fisher), and nuclei were stained using DAPI. Stained tissues were mounted in Vectashield (Vector Labs), except for Alexa647-labeled tissues which were mounted in Prolong Gold (Thermo Fisher Scientific). Rabbit anti-GFP AlexaFluor488 conjugate (Molecular Probes, 1:300) was used to visualize GFP expression, Wg expression was detected using mouse anti-Wg (Developmental Studies Hybridoma Bank 4D4, 1:100) and phospho-Histone 3 was detected using rabbit anti-PH3 (Cell Signaling Technologies 9701S, 1:500). Primary antibodies were detected with Alexa Fluor 488- or 555- coupled secondary antibodies (1:400). Confocal images were captured on a Zeiss LSM 780, LSM 980, or an Olympus IX83 confocal microscope, through the Microscopy Resources of the North Quad (MicRoN) facility at Harvard Medical School.

Single nucleus RNAseq (snRNAseq) atlas. To capture proventriculus transcriptional states in both well-fed flies and flies with reduced systemic insulin signaling, single nuclei suspensions were generated from proventriculus samples from both control flies (*esg-Gal4, tubGal80^ts^, UAS:GFP / +*) and cachexic flies (*esg-Gal4, tubGal80^ts^, UAS:GFP ; UAS-yki^3SA^*), after 8 or 9 days at 29ºC. We verified via confocal analysis that *esg-Gal4* is not expressed in the proventriculus itself, which was also confirmed via snRNA sequencing. Proventriculus samples were manually dissected from adult female flies in cold Schneider's medium and immediately flash frozen on dry ice in eppendorf tubes until a total of approximately 600 samples were collected (~300 of each genotype). Proventriculi were manually severed from the crop and

584 esophagus, and from the midgut just posterior to the proventriculus, anterior to where *esg-Gal4*
585 is detected in midgut ISCs.
586
587 Single nuclei were isolated following the protocol described in [66], derived originally from [67],
588 and stained with DAPI immediately before FACS sorting, performed at the Immunology Flow
589 Cytometry Core at Harvard Medical School. Sorted nuclei were resuspended in 1X PBS
590 supplemented with 0.5% BSA with RNAse inhibitor (Promega) at a concentration of ~2000
591 cells/µl.
592
593 snRNAseq was performed following the 10X Genomics Chromium protocol (Chromium Next
594 GEM 551 Single Cell 3'_v3.1_Rev_D), and sequenced using a Illumina NovaSeq 6000 at the
595 Harvard Medical School Biopolymers Facility. 10,525 nuclei from esg^{ts} flies and 11,801 nuclei
596 from esg^{ts} > *yki* flies were sequenced (total = 22,332 nuclei) and reads were mapped to
597 BDGP6.32.
598
Bioinformatic analysis was performed largely following [68] with minor modifications. Raw data
were processed with Cell Ranger software (10x Genomics, v7.0.0) to generate a single cell
matrix for each sample. The count matrices were imported into Seurat (v4.3.0) and cells with
less than 1300 or greater than 60,000 UMIs were filtered out, and low abundance genes with
less than 10 counts were removed from further analysis. The two biological samples were
bioinformatically pooled for PCA and clustering analysis to generate a single atlas including
cells from both conditions. After a PCA reduction, the first 40 components were used for UMAP
and clustering analysis, at a 0.2 resolution. We removed from further analysis one cluster that
appeared to be an artefact, as it contained markers for multiple different unrelated cell-types as
well as stress and heat shock proteins. Differentially expressed genes were calculated using the
Wilcox Rank Sum Test across each cluster and were then further selected using a Log2FC > X
cut-off and adjusted p value < 0.05. All downstream analysis was performed in R (v4.3).
611
612 To map snRNAseq clusters to proventriculus anatomy, we examined the expression pattern of
613 Gal4 lines for top-ranking marker genes using confocal microscopy.
614
615 **MARCM analysis.** Flies of the genotype *hsFLP, FRT19A, tubGal80 ; tub-Gal4, UAS-GFP* were
616 crossed to *FRT19A* flies (BL1709), and offspring were heat-shocked in a 37ºC water bath either
617 during the L3 stage for during the adult stage. Two one hour heat-shocks were administered,
618 separated by 2-3 hours at 25ºC. MARCM crosses were maintained at either 25ºC or 18ºC as
619 indicated in the text.
620
Tissue damage protocol using *UAS-rpr*. Similarly to [11], we induced ectopic apoptosis by
crossing Gal4 drivers to flies containing *UAS-rpr* ; *tubGal80^{ts}* or, for non-damaged controls, to
w^{1118} flies. These crosses were maintained at 18ºC until several days after eclosion. Adult flies
were then transferred to EdU+ food and placed at 29ºC for 48 hours to induce apoptosis, then
returned to recover at 18ºC for either 2 days or 5 days, on EdU+ food throughout. Thus, all
EdU+ incorporation events that occurred during the damage or during the recovery period could
be visualized and quantified.

Creation of split-intein Gal4 line. To generate *GMR71F06-Gal4::Gal4-C-int*, the 3,547 bp
enhancer fragment from *GMR71F06* FlyLight line [69] was amplified using the following primers:
caccTGCCCCGAGCAGCAAAAGAAGTCCG and TCTCCGAAATCTGCGCCAATTCCAG, and
cloned into the pENTR-dTOPO vector (Thermo Fisher Scientific). This fragment was cloned into
the pBP-Gal4^{C-int} backbone [39] (Addgene 199202) via LR gateway cloning, and was inserted
into the attP40 landing site using standard phiC31 transgenesis.

Time series of nuclear morphology. To quantify nuclear morphology over the course of the
fly's life, age matched flies of the genotype *GMR7106-Gal4 / UAS-Stinger ; MKRS / +* were
collected on eclosion day, and maintained at 25ºC in standard fly vials on standard lab food
(approximately 10-15 adult flies per vial), either with or without dried active yeast sprinkled on
the food's surface. In a pilot experiment, we observed that *UAS-Stinger* is tightly nuclear-
localized in this tissue, whereas other common nuclear-localized fluorophores transgenes
(*UAS:GFP^{NLS}* [BL4776], *UAS-unc84-GFP* [lab stock] and *UAS:RedStinger* [BL8545]) were not.
Flies were flipped onto fresh food every two or three days. On each day of sampling (3d, 7d, 14,
21d, 28d, and 35d), approximately 6-8 cardia were dissected and stained overnight with Alexa
488-coupled anti-GFP, DAPI, and phalloidin as described above. At each time-point, confocal z-
stacks were collected through the entirety of Zone 4 (n=5-6 proventriculus per time point) and
analyzed as described below. We also collected samples for eclosion day and for 1d later, but
found that the Gal4 expression pattern extended beyond Zone 4 at these time points, so they
were excluded from further analysis.

Image quantification. Image quantification was performed with arivis Vision4D software (Zeiss,
v4.1.2). GFP+ nuclei were segmented using the Cellpose-based Segmenter, and segmentation
was then manually corrected for each sample by splitting or removing nuclei which were
erroneously fused and removing artefacts that did not correspond to nuclei. Nuclear volume was
extracted for all objects >10µm^3. For cell count, those nuclei missed by Cellposed-based
segmentation were manually counted using the "add marker" function while viewing the tissue in
3D. To quantify cell loss during aging, cells that had been shed into the lumen were manually
counted in FIJI [70].

Ploidy measurements. Ploidy was quantified as described in [52]. Flies of the genotype *w ;
UAS:GFP^{Stinger} ; 71F06-Gal4* were aged to 17-19 days old at 25ºC, and then squashed to a
cellular monolayer on a charged slide, in parallel with adult testes samples as a haploid
standard. The summed DAPI intensity of GFP+ nuclei were quantified and compared to haploid
nuclei measured under the same conditions.

Lifespan and *Pseudomonas* survival experiments. Lifespan experiments were conducted at
29ºC, with 15 mated females per vial on standard fly food supplemented with dried yeast
powder, and flipped every 1-2 days. Flies of the genotype *w ; tubGal80^{ts}, UAS:GFP ; 71F06-
Gal4* were crossed to control or UAS:RNAi lines, and the offspring were reared at 18ºC until
eclosion. Adults were collected and mated over a 4 day period at 18ºC, then shifted to 29ºC

671   three days later to begin life-span measurements. The number of dead flies was recorded every
672   day.
673
Infection with *Pseudomonas aeruginosa (PA14)* followed a protocol similar to [59]. 5mL of LB
medium was seeded from a glycerol stock of *PA14* and grown overnight in a shaking 37ºC
incubator. The following morning, 1 mL of this culture was diluted in 50 mL of fresh LB and
grown for an additional 6 hours, shaking at 37ºC. This culture was then diluted 1:3 in 5%
sucrose solution. Fly vials containing half of a standard kimwipe, compressed into the bottom of
the vial, were loaded with 1.5 mL of this bacterial suspension. In order to promote feeding on
the bacteria, flies were starved for two hours in empty vials immediately prior to bacterial
treatment.
682
These experiments were performed at 29ºC on flies that had been maintained at 29ºC for 15
days prior to *PA14* infection. After a 24 hour infection, flies were flipped to fresh vials containing
kimwipes saturated with 5% sucrose solution, and flipped to fresh 5% sucrose solution daily
after that. The number of dead flies was recorded approximately every 24 hours. Four vials of 8-
10 flies each were used per genotype, and the entire experiment was performed in biological
duplicate, with similar results in both experiments.
689
690   **Graphing and statistical analysis.** All graphs were created and statistical tests performed
691   using Prism (GraphPad, v10).
692
693   **Data Sharing/ Resource Availability**
694   The snRNA-seq Atlas is available for data mining at https://www.flyrnai.org/scRNA/. Raw RNA-
695   seq reads and processed data files have been deposited in the NCBI Gene Expression
696   Omnibus (GEO) database under accession code GSE301009.
697
698
699
700   **Acknowledgements**
701

We thank Mujeeb Qadiri, Austin Veal, and Aram Comjean for assistance with snRNAseq data
analysis and sharing, and Rich Binari for assistance with fly work. We thank Hannah Dayton
and Liz Lane for reagents and advice with *Pseudomonas* experiments. This work was supported
by the National Institute of Health R24 OD031952. NP is an investigator of the Howard Hughes
Medical Institute. This article is subject to HHMI's Open Access to Publications policy. HHMI lab
heads have previously granted a nonexclusive CC BY 4.0 license to the public and a
sublicensable license to HHMI in their research articles. Pursuant to those licenses, the author-
accepted manuscript of this article can be made freely available under a CC BY 4.0 license
immediately upon publication.

711
712
713

Literature Cited

1. Buchon N, Osman D, David FPA, Fang HY, Boquete J-P, Deplancke B, et al. Morphological and Molecular Characterization of Adult Midgut Compartmentalization in Drosophila. Cell Reports. 2013;3: 1725–1738. doi:10.1016/j.celrep.2013.04.001

2. Buchon N, Osman D. All for one and one for all: Regionalization of the Drosophila intestine. Insect Biochem Molec. 2015;67: 2–8. doi:10.1016/j.ibmb.2015.05.015

3. Marianes A, Spradling AC. Physiological and stem cell compartmentalization within the Drosophila midgut. Elife. 2013;2: e00886. doi:10.7554/elife.00886

4. Ohlstein B, Spradling A. Multipotent Drosophila Intestinal Stem Cells Specify Daughter Cell Fates by Differential Notch Signaling. Science. 2007;315: 988–992. doi:10.1126/science.1136606

5. Micchelli CA, Perrimon N. Evidence that stem cells reside in the adult Drosophila midgut epithelium. Nature. 2005;439: 475–479. doi:10.1038/nature04371

6. Jiang H, Tian A, Jiang J. Intestinal stem cell response to injury: lessons from Drosophila. Cell Mol Life Sci. 2016;73: 3337–3349. doi:10.1007/s00018-016-2235-9

7. Lucchetta EM, Ohlstein B. The Drosophila midgut: a model for stem cell driven tissue regeneration. Wiley Interdiscip Rev: Dev Biol. 2012;1: 781–788. doi:10.1002/wdev.51

8. Cheng H, Leblond CP. Origin, differentiation and renewal of the four main epithelial cell types in the mouse small intestine V. Unitarian theory of the origin of the four epithelial cell types. Am J Anat. 1974;141: 537–561. doi:10.1002/aja.1001410407

9. Barker N, Es JH van, Kuipers J, Kujala P, Born M van den, Cozijnsen M, et al. Identification of stem cells in small intestine and colon by marker gene Lgr5. Nature. 2007;449: 1003–1007. doi:10.1038/nature06196

10. Zwick RK, Ohlstein B, Klein OD. Intestinal renewal across the animal kingdom: comparing stem cell activity in mouse and Drosophila. Am J Physiol-Gastrointest Liver Physiol. 2019;316: G313–G322. doi:10.1152/ajpgi.00353.2018

11. Fox DT, Spradling AC. The Drosophila Hindgut Lacks Constitutively Active Adult Stem Cells but Proliferates in Response to Tissue Damage. Cell Stem Cell. 2009;5: 290–297. doi:10.1016/j.stem.2009.06.003

12. Cohen E, Allen SR, Sawyer JK, Fox DT. Fizzy-Related dictates A cell cycle switch during organ repair and tissue growth responses in the Drosophila hindgut. eLife. 2018;7: e38327. doi:10.7554/elife.38327

13. Losick VP, Fox DT, Spradling AC. Polyploidization and Cell Fusion Contribute to Wound Healing in the Adult Drosophila Epithelium. Curr Biol. 2013;23: 2224–2232. doi:10.1016/j.cub.2013.09.029

14. Losick VP, Jun AS, Spradling AC. Wound-Induced Polyploidization: Regulation by Hippo and JNK Signaling and Conservation in Mammals. PLoS ONE. 2016;11: e0151251. doi:10.1371/journal.pone.0151251

15. Singh SR, Zeng X, Zheng Z, Hou SX. The adult Drosophila gastric and stomach organs are maintained by a multipotent stem cell pool at the foregut/midgut junction in the cardia (proventriculus). Cell Cycle. 2011;10: 1109–1120. doi:10.4161/cc.10.7.14830

16. King DG. Cellular organization and peritrophic membrane formation in the cardia (Proventriculus) of Drosophila melanogaster. J Morphol. 1988;196: 253–282.

757   doi:10.1002/jmor.1051960302

17.   Wigglesworth VB. The Formation of the Peritrophic Membrane in Insects, with Special Reference to the Larvae of Mosquitoes. J Cell Sci. 1930;s2-73: 593–616. doi:10.1242/jcs.s2-73.292.593

18.   Rizki MTM. The secretory activity of the proventriculus of Drosophila melanogaster. J Exp Zoöl. 1956;131: 203–221. doi:10.1002/jez.1401310202

19.   Pankratz MJ, Hoch M. Control of epithelial morphogenesis by cell signaling and integrin molecules in the Drosophila foregut. Development. 1995;121: 1885–1898. doi:10.1242/dev.121.6.1885

20.   Fuß B, Hoch M. Drosophila endoderm development requires a novel homeobox gene which is a target of Wingless and Dpp signalling. Mech Dev. 1998;79: 83–97. doi:10.1016/s0925-4773(98)00172-5

21.   Fuss B, Josten F, Feix M, Hoch M. Cell movements controlled by the Notch signalling cascade during foregut development in Drosophila. Development. 2004;131: 1587–1595. doi:10.1242/dev.01057

22.   Waterhouse D. The Rate of Production of the Peritrophic Membrane in Some Insects. Aust J Biol Sci. 1954;7: 59–72. doi:10.1071/bi9540059

23.   Becker B. Determination of the formation rate of peritrophic membranes in some Diptera. J Insect Physiol. 1978;24: 529–533. doi:10.1016/0022-1910(78)90054-9

24.   Tzou P, Ohresser S, Ferrandon D, Capovilla M, Reichhart J-M, Lemaitre B, et al. Tissue-Specific Inducible Expression of Antimicrobial Peptide Genes in Drosophila Surface Epithelia. Immunity. 2000;13: 737–748. doi:10.1016/s1074-7613(00)00072-8

25.   Zhu H, Ludington WB, Spradling AC. Cellular and molecular organization of the Drosophila foregut. Proc Natl Acad Sci. 2024;121: e2318760121. doi:10.1073/pnas.2318760121

26.   Dodge R, Jones EW, Zhu H, Obadia B, Martinez DJ, Wang C, et al. A symbiotic physical niche in Drosophila melanogaster regulates stable association of a multi-species gut microbiota. Nat Commun. 2023;14: 1557. doi:10.1038/s41467-023-36942-x

27.   Bertran-Mas J, Giorgio ED, Martín N, Llimargas M. Different cellular and molecular mechanisms of chitin deposition contribute to the specificity of the two chitin synthases in D. melanogaster. bioRxiv. 2024; 2024.12.02.626067. doi:10.1101/2024.12.02.626067

28.   Pandey A, Galeone A, Han SY, Story BA, Consonni G, Mueller WF, et al. Gut barrier defects, intestinal immune hyperactivation and enhanced lipid catabolism drive lethality in NGLY1-deficient Drosophila. Nat Commun. 2023;14: 5667. doi:10.1038/s41467-023-40910-w

29.   Zhang L, Turner B, Ribbeck K, Hagen KGT. Loss of the mucosal barrier alters the progenitor cell niche via Janus kinase/signal transducer and activator of transcription (JAK/STAT) signaling. J Biol Chem. 2017;292: 21231–21242. doi:10.1074/jbc.m117.809848

30.   Britton JS, Edgar BA. Environmental control of the cell cycle in Drosophila: nutrition activates mitotic and endoreplicative cells by distinct mechanisms. Development. 1998;125: 2149–2158. doi:10.1242/dev.125.11.2149

31.   Edgar BA, Orr-Weaver TL. Endoreplication Cell Cycles More for Less. Cell. 2001;105: 297–306. doi:10.1016/s0092-8674(01)00334-8

32.   Kwon Y, Song W, Droujinine IA, Hu Y, Asara JM, Perrimon N. Systemic Organ Wasting Induced by Localized Expression of the Secreted Insulin/IGF Antagonist ImpL2. Dev Cell. 2015;33: 36–46. doi:10.1016/j.devcel.2015.02.012

33. Ding G, Xiang X, Hu Y, Xiao G, Chen Y, Binari R, et al. Coordination of tumor growth and
host wasting by tumor-derived Upd3. Cell Rep. 2021;36: 109553.
doi:10.1016/j.celrep.2021.109553

34. Liu Y, Saavedra P, Perrimon N. Cancer cachexia: lessons from Drosophila. Dis Model
Mech. 2022;15: dmm049298. doi:10.1242/dmm.049298

35. Lim SY, You H, Lee J, Lee J, Lee Y, Lee K-A, et al. Identification and characterization of
GAL4 drivers that mark distinct cell types and regions in the Drosophila adult gut. J Neurogenet.
2021;35: 33–44. doi:10.1080/01677063.2020.1853722

36. Ghosh S, Leng W, Wilsch-Bräuninger M, Barrera-Velázquez M, Léopold P, Eaton S. A local
insulin reservoir in Drosophila alpha cell homologs ensures developmental progression under
nutrient shortage. Curr Biol. 2022;32: 1788-1797.e5. doi:10.1016/j.cub.2022.02.068

37. Kim SK, Rulifson EJ. Conserved mechanisms of glucose sensing and regulation by
Drosophila corpora cardiaca cells. Nature. 2004;431: 316–320. doi:10.1038/nature02897

38. Lee G, Park JH. Hemolymph Sugar Homeostasis and Starvation-Induced Hyperactivity
Affected by Genetic Manipulations of the Adipokinetic Hormone-Encoding Gene in Drosophila
melanogaster. Genetics. 2004;167: 311–323. doi:10.1534/genetics.167.1.311

39. Ewen-Campen B, Luan H, Xu J, Singh R, Joshi N, Thakkar T, et al. split-intein Gal4
provides intersectional genetic labeling that is repressible by Gal80. Proc Natl Acad Sci United
States Am. 2023;120: e2304730120. doi:10.1073/pnas.2304730120

40. Hung R-J, Hu Y, Kirchner R, Liu Y, Xu C, Comjean A, et al. A cell atlas of the adult
Drosophila midgut. Proc National Acad Sci. 2020;117: 1514–1523.
doi:10.1073/pnas.1916820117

41. Ren F, Wang B, Yue T, Yun E-Y, Ip YT, Jiang J. Hippo signaling regulates Drosophila
intestine stem cell proliferation through multiple pathways. Proc National Acad Sci. 2010;107:
21064–21069. doi:10.1073/pnas.1012759107

42. Karpowicz P, Perez J, Perrimon N. The Hippo tumor suppressor pathway regulates
intestinal stem cell regeneration. Development (Cambridge, England). 2010;137: 4135–4145.
doi:10.1242/dev.060483

43. Shaw RL, Kohlmaier A, Polesello C, Veelken C, Edgar BA, Tapon N. The Hippo pathway
regulates intestinal stem cell proliferation during Drosophila adult midgut regeneration.
Development. 2010;137: 4147–4158. doi:10.1242/dev.052506

44. Jiang H, Grenley MO, Bravo M-J, Blumhagen RZ, Edgar BA. EGFR/Ras/MAPK Signaling
Mediates Adult Midgut Epithelial Homeostasis and Regeneration in Drosophila. Cell Stem Cell.
2011;8: 84–95. doi:10.1016/j.stem.2010.11.026

45. Xu C, Luo J, He L, Montell C, Perrimon N. Oxidative stress induces stem cell proliferation
via TRPA1/RyR-mediated Ca2+ signaling in the Drosophila midgut. eLife. 2017;6: e22441.
doi:10.7554/elife.22441

46. Leung NY, Xu C, Li JSS, Ganguly A, Meyerhof GT, Regimbald-Dumas Y, et al. Gut tumors
in flies alter the taste valence of an anti-tumorigenic bitter compound. Curr Biol. 2024;34: 2623-
2632.e5. doi:10.1016/j.cub.2024.04.082

47. Ohlstein B, Spradling A. The adult Drosophila posterior midgut is maintained by pluripotent
stem cells. Nature. 2006;439: 470–474. doi:10.1038/nature04333

48. McLeod CJ, Wang L, Wong C, Jones DL. Stem cell dynamics in response to nutrient
availability. Current biology : CB. 2010;20: 2100–2105. doi:10.1016/j.cub.2010.10.038

49. Lucchetta EM, Ohlstein B. Amitosis of Polyploid Cells Regenerates Functional Stem Cells
in the Drosophila Intestine. Stem Cell. 2017;20: 1–19. doi:10.1016/j.stem.2017.02.012
50. Miguel-Aliaga I, Jasper H, Lemaitre B. Anatomy and Physiology of the Digestive Tract of
Drosophila melanogaster. Genetics. 2018;210: 357–396. doi:10.1534/genetics.118.300224
51. Evans CJ, Olson JM, Ngo KT, Kim E, Lee NE, Kuoy E, et al. G-TRACE: rapid Gal4-based
cell lineage analysis in Drosophila. Nat Methods. 2009;6: 603–5. doi:10.1038/nmeth.1356
52. Clay DE, Stormo BM, Fox DT. Measuring Cellular Ploidy In Situ by Light Microscopy.
Methods Mol Biol (Clifton, NJ). 2023;2545: 401–412. doi:10.1007/978-1-0716-2561-3_21
53. Morris JP, Baslan T, Soltis DE, Soltis PS, Fox DT. Integrating the Study of Polyploidy
Across Organisms, Tissues, and Disease. Annu Rev Genet. 2024. doi:10.1146/annurev-genet-
111523-102124
54. Øvrebø JI, Edgar BA. Polyploidy in tissue homeostasis and regeneration. Development.
2018;145: dev156034. doi:10.1242/dev.156034
55. Molano-Fernández M, Hickson ID, Herranz H. Cyclin E overexpression in the Drosophila
accessory gland induces tissue dysplasia. Front Cell Dev Biol. 2023;10: 992253.
doi:10.3389/fcell.2022.992253
56. Pierce SB, Yost C, Britton JS, Loo LWM, Flynn EM, Edgar BA, et al. dMyc is required for
larval growth and endoreplication in Drosophila. Development. 2004;131: 2317–2327.
doi:10.1242/dev.01108
57. Demontis F, Perrimon N. Integration of Insulin receptor/Foxo signaling and dMyc activity
during muscle growth regulates body size in Drosophila. Development (Cambridge, England).
2009;136: 983–993. doi:10.1242/dev.027466
58. Kuraishi T, Binggeli O, Opota O, Buchon N, Lemaitre B. Genetic evidence for a protective
role of the peritrophic matrix against intestinal bacterial infection in Drosophila melanogaster.
Proc Natl Acad Sci. 2011;108: 15966–15971. doi:10.1073/pnas.1105994108
59. Kim K, Hung R-J, Perrimon N. miR-263a Regulates ENaC to Maintain Osmotic and
Intestinal Stem Cell Homeostasis in Drosophila. Dev Cell. 2017;40: 23–36.
doi:10.1016/j.devcel.2016.11.023
60. Kim K, Lane EA, Saftien A, Wang H, Xu Y, Wirtz-Peitz F, et al. Drosophila as a model for
studying cystic fibrosis pathophysiology of the gastrointestinal system. Proc Natl Acad Sci.
2020;117: 10357–10367. doi:10.1073/pnas.1913127117
61. Fox DT, Duronio RJ. Endoreplication and polyploidy: insights into development and
disease. Development. 2012;140: 3–12. doi:10.1242/dev.080531
62. Bailey EC, Kobielski S, Park J, Losick VP. Polyploidy in Tissue Repair and Regeneration.
Cold Spring Harb Perspect Biol. 2021;13: a040881. doi:10.1101/cshperspect.a040881
63. Smith AV, Orr-Weaver TL. The regulation of the cell cycle during Drosophila
embryogenesis: the transition to polyteny. Development. 1991;112: 997–1008.
doi:10.1242/dev.112.4.997
64. Nandakumar S, Grushko O, Buttitta LA. Polyploidy in the adult Drosophila brain. eLife.
2020;9: e54385. doi:10.7554/elife.54385
65. Zielke N, Edgar BA, DePamphilis ML. Endoreplication. Cold Spring Harb Perspect Biol.
2013;5: a012948. doi:10.1101/cshperspect.a012948
66. Liu Y, Dantas E, Ferrer M, Liu Y, Comjean A, Davidson EE, et al. Tumor Cytokine-Induced
Hepatic Gluconeogenesis Contributes to Cancer Cachexia: Insights from Full Body Single

Nuclei Sequencing. bioRxiv. 2023; 2023.05.15.540823. doi:10.1101/2023.05.15.540823

67. Li H, Janssens J, Waegeneer MD, Kolluru SS, Davie K, Gardeux V, et al. Fly Cell Atlas: A

single-nucleus transcriptomic atlas of the adult fruit fly. Science. 2022;375: eabk2432.

doi:10.1126/science.abk2432

68. Saavedra P, Dumesic PA, Hu Y, Filine E, Jouandin P, Binari R, et al. REPTOR and

CREBRF encode key regulators of muscle energy metabolism. Nat Commun. 2023;14: 4943.

doi:10.1038/s41467-023-40595-1

69. Jenett A, Rubin GM, Ngo T-TB, Shepherd D, Murphy C, Dionne H, et al. A GAL4-Driver

Line Resource for Drosophila Neurobiology. Cell Reports. 2012;2: 991–1001.

doi:10.1016/j.celrep.2012.09.011

70. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.

doi:10.1038/nmeth.2019

Main Figures
905
906

907
908
Figure 1. The *Drosophila* proventriculus contains a ring of nutrition-sensitive EdU+ cells
located in Zone 4. (A) Schematic of the single-cell epithelium comprising the *Drosophila*
proventriculus, located at the juncture of the foregut and anterior midgut. Zone 4 cells are
indicated in green. (B) Left panel: *chitin synthase 2* (*Chs2-Gal4*) is specifically expressed in
Zone 4 cells. Middle panel: HPA lectin stains the peritrophic membrane (PM). Left panel: Wg
protein is enriched in Zone 4 cells. Confocal z-slices are shown. (C) Left panel: EdU labeling is
enriched in a ring of cells at the posterior edge of Zone 4. A maximum intensity projection of
EdU staining is overlaid on a single z-slice of DAPI staining. Right panel: EdU+ cells are
immediately anterior to a ring of cells expressing the 10XSTAT:GFP reporter. Single z-slice is
shown. Scale bar is 50 µm. (D) The number of EdU+ cells increases with the duration of EdU
feeding. Maximum intensity projections are shown. (E) Quantification of EdU+ cell numbers at 1,
3, and 5 days of feeding, for two different fly strains, *OregonR* and *w1118*. (F) After a 1 day EdU
pulse, staining is largely restricted to small nuclei at the border of Zone 3 and Zone 4. After 5
days of EdU feeding, staining is also visible in larger nuclei. Confocal z-slices are shown. (G)
The number of EdU+ cells increases as more yeast is added to the food source. Maximum
intensity projections are shown. (H) Quantification of the experiment shown in (G). Anterior is
up.

926
Figure 2. Molecular markers and a single-nuclei atlas for the proventriculus. (A) 71F06-
Gal4 drives specific expression in Zone 4 cells, including EdU+ cells. (B) UMAP projection of a
single snRNAseq dataset created from 22,332 nuclei, containing 14 clusters representing the
major epithelial, muscle, and neuronal cells found in the proventriculus. (C) Schematic of the
major epithelial cell types in the proventriculus, corresponding to the clusters identified in (B).
(D) Dot plot of selected marker genes for the 12 major clusters identified via snRNAseq; note
that two minor clusters ("anterior midgut 1" and "unknown") are excluded from this figure as they
could not be clearly linked to *in vivo* cell types. (E) Expression patterns of marker genes for
each of the 9 major epithelial cell types in the proventriculus. Anterior is up. Confocal z-slices
are shown.
937
938

Figure 3. Mitosis is undetectable in Zone 4 cells. (A) Over-expression of the tumor drivers *yki³ˢᴬ, Ras¹ᴬ,* or *Notch* RNAi in Zone 4 cells causes increased EdU+ incorporation and enlarged and altered morphology of the tissue. (B) No phospho-histone 3 staining was detected in any of the proventriculi examined in any of these genotypes. Flies were fed colchecine for 18 hours prior to dissection to enrich for pH3+ cells. Midgut ISCs from these same guts were included as positive controls, which contained high numbers of pH3+ staining in all samples. Maximum intensity projections are shown. Anterior is up.

Lineage marking with MARCM19A (*hsFLP, FRT19A, tubGal80 / FRT19A ;; tub-Gal4, UAS-GFP*)

14 days at 25ºC after heatshock

Figure 4. Clonal analysis with MARCM does not label large clones in Zone 4. (A) Positive control for clone induction; clones were induced during the L3 larval stage, leading to large GFP+ clones throughout the proventriculus including in Zone 4. (B) In the posterior midgut, clone induction labels ISCs their descendants and generates GFP+ clones throughout the midgut. These clones are generally larger when flies endure a 2-day starvation followed by re-feeding. (C) In the proventriculus, small scattered GFP+ cells are detected throughout the tissue, but no discernable enrichment in Zone 4, where EdU+ cells are located. (D) Quantification of several MARCM experiments, with dissection 14 days after clone induction (at 25ºC), 19 days after clone induction (at 25ºC), or 30days after clone induction (at 18ºC). Maximum intensity projections are shown. Scale bar = 50µm. Anterior is up.

Figure 5. Tissue damage causes a decrease, not an increase, in Zone 4 cell cycling.
Tissue damage was induced by over-expressing the pro-apoptotic gene *rpr* for two days, in 3 distinct cell types of the proventriculus, and EdU+ cells were visualized after two and five days of recovery. (A) *rpr* expression in Zone 5, using *Ca-Ma2d-Gal4*, causes a decrease in EdU+ cells after both two and five days. Blue arrows indicate pyknotic nuclei indicative of cell death. (B) Quantification of experiment shown in (A). (C) *rpr* expression in Zone 1, using *CG11550-Gal4,* causes a decrease in EdU+ cells after both two and five days. (D) Quantification of experiment shown in (C). (E) *rpr* expression in the esophagus, using *ELOVL7-Gal4*, leads to decreased EdU+ cells in Zone 4, and endoreplication in esophageal cells. White bracket indicates EdU+ signal in Zone 4, blue arrowheads and white asterisks indicate large, EdU+ nuclei in the esophagus itself. Scale bar = 50µm. Anterior is up. (F) Quantification of experiment shown in E. Significance values are from t-test, with Welch's correction for any comparisons with unequal variances.

986
987
988

Figure 6. Genetic lineage tracing using a highly-specific split-intein Gal4 that labels Edu+

cells. (A) Expression pattern of *71F06-Gal4::Gal4-C-int*, *Delta^{Gal4-N-int}*, and the intersection of

71F06 ∩ Delta. (B) *71F06 ∩ Delta* expression largely overlaps with EdU+ cells in Zone 4 cells.

(C) Using *71F06 ∩ Delta* to drive the G-TRACE lineage system, with *tubGal80^{ts}* to restrict

labeling to the adult stage for the indicated duration. Colored arrows indicate individual cells with

either solely GFP (past but not current expression) or RFP (very recently initiated expression.)

Anterior is up.(D) Quantification of experiment shown in (C). Each column represents a single

proventriculus sample at the indicated time-point.

997
998

Figure 7. Zone 4 nuclear volume progressively increases over the fly lifespan, while some cells are shed into the gut lumen during aging. (A) Time-course of nuclear morphology of the lifespan of the fly, from three days to 35 days post-eclosion. Nuclear-localized GFP (UAS:GFP^Stinger) was expressed in Zone 4 using 71F06-Gal4, and flies were provided with or without supplemental yeast powder. (B) Nuclei were false-colored by volume, indicating a consistent increase in nuclear volume over the course of a fly's life. A posterior-to-anterior gradient of increasingly large nuclear volume is also apparent. (C) Quantification of nuclear volume of Zone 4 nuclei, as well as diploid ISCs for comparison. Nuclear volume increases throughout the lifespan of a fly, with consistently higher volumes for yeast-supplemented flies until the 35d timepoint. "n.s." = not significantly different; * p = 0.02; *** p = 0.0005; **** p < 0.0001, t-test, with Welch's correction for any comparison with unequal variances. (D) The number of Zone 4 cells steadily decreases throughout this time-course, and decreases faster in the absence of yeast powder. *** p = 0.007, t-test. (E) GFP+ nuclei are consistently observed shed into the lumen adjacent to Zone 4. (F) Quantification of GFP+ cells observed in the lumen at each time point.

1019
1020

Figure 8. Perturbations to endocycling in Zone 4 cells lead to altered nuclear volume, tissue size, and peritrophic membrane synthesis. (A) Ectopic expression of CycE or Myc drives excess endocycling, leading to larger nuclei (top) and increased peritrophic membrane synthesis from Zone 4 cells, visualized with HPA-lectin. Insets show increased magnification of nuclear GFP. Images are maximum intensity projections of confocal z-stacks (B) Quantification of nuclear volume of *UAS:CycE, UAS:Myc* (n = 4 proventriculi per genotype, 231-255 nuclei per genotype)*,* and *InR RNAi* experiments (n = 3-5 proventriculi, 179-346 nuclei per genotype per timepoint). Statistical tests are one-way ANOVA with multiple comparisons; *** $p < 0.001$; **** $p < 0.0001$. (C) Cytoplasmic UAS:EGFP expression reveals that RNAi against *InR* reduces the overall size of Zone 4. Images are single confocal z-slices. (D) The peritrophic membrane, stained with HPA-Lectin, is reduced and disrupted when endocycling is blocked by RNAi against *InR.* Top two row shows maximum intensity projection, middle row shows single z-stack, and bottom row shows maximum intensity projection.(E) Lifespan at 29ºC under standard conditions is not affected by *InR RNAi*. Sample sizes: *w* = 134; *mCherry RNAi* = 95 flies; *InR RNAi 51518* = 128 flies; *InR RNAi 35251* = 58 flies; 15 flies per vial in all cases. (F) RNAi against *InR* in Zone 4 leads to increased susceptibility to *P. aeruginosa* infection. Sample sizes: *w* = 40 flies;

1037   *mCherry RNAi* = 40 flies; *InR RNAi 51518* = 38 flies; *InR RNAi 35251* = 40 flies; 9-10 flies per
1038   vial.
1039
1040   **Supplemental Figures**
1041

1042
1043
1044   **Supplemental Figure 1. EdU+ cells continue cycling throughout adult life.** (A) After 3 days
1045   of EdU feeding, a ring of labeled cells is visible at the posterior edge of Zone 4, as well as in the
1046   ISCs and enterocytes of the midgut. (B) Following a 20 day chase period, a reduction of EdU+
1047   cells is visible in both Zone 4 cells and in the midgut, where continual cycling has diluted out the
1048   original EdU pulse.
1049
1050

Supplemental Figure 2. Systemic reduction in insulin signaling reduces the number of EdU+ cells in the proventriculus. (A) Tumors in midgut ISCs (driven by *esg-Gal4 > yki*[3SA]) cause a reduction in EdU+ cells in the proventriculus. Note that *esg-Gal4* is not expressed in the proventriculus. Anterior is up, scale bar is 50 μm. (B) Quantification of the experiment shown in (A). (C) Over-expression of insulin-antagonists *ImpL2, upd3,* or a combination of both from the ISCs (*esg-Gal4*) leads to a reduction in EdU+ cells. (D) Over-expression of insulin-antagonists *ImpL2, upd3,* or a combination of both from muscles using *dMef2-Gal4* leads to a reduction in EdU+ cells.

Zhu et al. (scRNAseq)

This study (snRNAseq)

1062
1063
1064 **Supplemental Figure 3. Merged UMAP projection of two single cell atlases of the**
1065 **proventriculus.** Data from the snRNAseq atlas presented in this manuscript and scRNAseq
1066 data from [25] were bioinformatically pooled and re-analyzed into a single dataset.
1067
1068

A GMR71F06-Gal4, tubGal80^{ts}, UAS:2XEGFP

w¹¹¹⁸   _UAS:yki^{3SA}_   _UAS:Ras^{1A}_

3 days at 29°C — surface / medial

DAPI
GFP
Phal.

extruded cytoplasm

50 µm

B 6 days at 29°C — surface / medial

1069
1070
1071 **Supplemental Figure 4. Phenotypes of ectopic oncogene expression in Zone 4 cells.**
1072 Ectopic expression of oncogenes _yki^{3SA}_ or _Ras^{1A}_ in Zone 4 cells of the proventriculus for either
1073 (A) three days or (B) six days leads to enlarged, distended tissue morphologies. These enlarged
1074 structures can be seen extruding cytoplasm into the gut lumen, and in no cases were cells
1075 observed to be growing into adjacent zones.
1076
1077

1078
1079
Supplemental Figure 5. MARCM analysis in the proventriculus following gut damage.
Analysis of MARCM clones in the proventriculus and midgut following a number of experimental
perturbations to the gut. Three representative proventriculus images are shown for each
condition, from a total sample size between 10 and 23 guts. (A) X-ray damage to entire flies at
1000 or 2000 RADS, examined seven days after damage. (B) Gut damage from *ECC15*
bacteria, examined 5 days after damage. (C) Gut damage from bleomycin, examined 11 days
after damage. (D) Undamaged guts, fed with food supplemented with 10% yeast. Anterior is up,
scale bar is 50μm.
1088
1089

1090
1091
1092 **Supplemental Figure 6. Ploidy analysis of Zone 4 nuclei.** Histogram of estimated ploidy of
1093 Zone 4 nuclei (n = 230 nuclei) normalized to haploid sperm nuclei.
1094
1095

1096
1097
1098 **Supplemental Figure 7. Time course analysis of nuclear volume and morphology in Zone**
1099 **4 cells.** (A) Representative images of each time-point for the experiment shown in Figure 7. (B)
1100 Histogram of nuclear volume calculated from the data represented in Figure 7.
1101
1102

1103
1104
1105 **Supplemental Figure 8. Additional examples of Zone 4 nuclei loss into the gut lumen.** In
1106 aging flies, GFP+ nuclei can be seen shed into the gut lumen from Zone 4. Multiple examples
1107 are shown of these nuclei appearing to exit Zone 4 via disrupted cell barriers, visible as loss of
1108 phalloidin, as well as blebs on the lumen-facing surface of Zone 4. Cells are observed at various
1109 locations along the anterior-posterior axis of Zone 4. Anterior is up, scale bar is 20μm.
1110
1111

1112
1113
Supplemental Figure 9. Additional characterization of genetic manipulations to
endocycling in Zone 4. (A) Over-expression of CycE (left) or Myc (right) leads to increased
endocycling and enlargement of nuclei, as well as increased overall tissue size. Flies were fed
EdU for two days prior to dissection. (B) RNAi against *InR* in Zone 4 abolishes cycling in these
cells, evidenced by a lack of EdU staining and smaller nuclear size. (C) Quantification of EdU+
cells in *InR* RNAi proventrici after 6 days and 15 days at 29ºC. Letters indicate statistical
significance groups (one-way ANOVA with Dunnett's multiple comparisons test.) All images are
maximum intensity projections of confocal z-stacks.
1122
1123

71F06-Gal4, tubGal80ts, UAS:GFPNUC

Number of GFP cells
(1d vs. 12d at 29ºC)

Percent decrease
(1d vs. 12d at 29ºC)

1124
1125
Supplemental Figure 10. Increased endocycling does not enhance cell loss in Zone 4.
Left: Cell counts of GFP+ Zone 4 cells after 1 day or 12 days at 29ºC for the indicated
genotypes. Right: the average percentage decrease, calculated from the data shown on the left,
was not significantly different between these conditions. Error bars represent standard deviation
propagated through the calculation of percentage decrease, and mean differences were
compared using a t-test.

 Table S1: Genotypes used in this study

Figure	Genotype	Source Notes
1B, left	yw ; + / wg[Sp-1] ; TI{GFP[3xP3.cLa]=CRIMIC.TG4.0}Chs2[CR60212-TG4.0] / P{w[+mC]=UAS-2xEGFP}AH3	Chs2-Gal4: BL93534; UAS:2xEGFP: BL60293
1B, middle	w ; P{w[+mC]=UAS-2xEGFP}AH2 ; P{w[+mW.hs]=GawB}Lk6[NP0101] / P{y[+t7.7] v[+t1.8]=VALIUM20-mCherry.RNAi}attP2	"Cardia-Gal4": Kyoto DGRC 103522; UAS:2xEGFP: BL60292; mCherry RNAi: BL35785
1B, right	w[1118] ; esg-Gal4, UAS:GFP, tubGal80[ts]	Perrimon lab stock
1C, left	w[1118] ; esg-Gal4, UAS:GFP, tubGal80[ts]	Perrimon lab stock
1C right	w[1118]; P{w[+mC]=10XStat92E-GFP}1	Perrimon lab stock
1D	Oregon R	
1F	Oregon R	
1G	w[1118] ; esg-Gal4, UAS:GFP, tubGal80[ts]	Perrimon lab stock
2A	w ; P{UAS-Stinger}2 / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / MKRS	UAS-Stinger: BL90920, GMR71F06-Gal4: BL39596
2E	yw ; wg[Sp-1] / + ; Mi{Trojan-GAL4.0}Elovl7[MI01455-TG4.0] / P{w[+mC]=UAS-2xEGFP}AH3	Elovl7-Gal4: BL67431
2E	yw ; wg[Sp-1] / + ; TI{GFP[3xP3.cLa]=CRIMIC.TG4.2}CG11550[CR02741-TG4.2] / P{w[+mC]=UAS-2xEGFP}AH3	CG11550-Gal4: BL97589
2E	yw ; P{y[+t7.7] w[+mC]=Tub-GAL4.C-int}attP40 / + ; TI{RFP[3xP3.PB]=2A-GAL4(1-20)::N-int}Ppn[G4-N-int] / P{w[+mC]=UAS-2xEGFP}AH3	Ppn-C-int: BL602730; tub-Gal4-C-int: BL602738
2E	yw ; wg[Sp-1] / + ; TI{GFP[3xP3.cLa]=CRIMIC.TG4.2}Lgr1[CR01685-TG4.2] / P{w[+mC]=UAS-2xEGFP}AH3	Lgr1-Gal4: BL86491
2E	yw ; y[1] w[*]; Mi{Trojan-GAL4.2}Wnt4[MI03717-TG4.2] / wg[Sp-1] ; P{w[+mC]=UAS-2xEGFP}AH3	Wnt4-Gal4: BL67449
2E	yw ; TI{GFP[3xP3.cLa]=CRIMIC.TG4.2}Ca-Ma2d[CR01742-TG4.2] / wg[Sp-1] ; P{w[+mC]=UAS-2xEGFP}AH3	Ca-Ma2d-Gal4: BL86499
2E	w[*] TI{RFP[3xP3.PB]=2A-GAL4(1-20)::N-int}Mco4[G4-N-int] ; P{y[+t7.7] w[+mC]=Tub-GAL4.C-int}attP40 / + ; P{w[+mC]=UAS-2xEGFP}AH3 / +	Mco4-N-int: BL605167; tub-Gal4-C-int: BL602738

2E	*yw ; wg[Sp-1] / + ; TI{GFP[3xP3.cLa]=CRIMIC.TG4.2}SecCl[CR01836-TG4.2]/TM3, Sb[1] Ser[1] / P{w[+mC]=UAS-2xEGFP}AH3*	SecCl-Gal4: BL91420
2E	*w ; Mex1-Gal4, UAS-2xEGFP*	Mex1-Gal4 gift of A. Petsakou (originally from Phillips & Thomas, 2006)
3A, left	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2*	GMR71F06-Gal4: BL39596, UAS:2xEGFP: BL60292, tubGal80ts: BL7108
3A, second to left	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / P{y[+t7.7] w[+mC]=UAS-yki.S111A.S168A.S250A.V5}attP2*	yki[3SA]: BL28817
3A, third to left	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / UAS:Ras[1A]*	Ras1A: gift of Chiwei Xu, Perrimon Lab
3A, fourth to left	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2*	
3A, right	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / P{y[+t7.7] v[+t1.8]=TRiP.HMS00001}attP2*	Notch RNAi: BL33611
4	hsFLP, FRT19A, tubGal80 / P{ry[+t7.2]=neoFRT}19A; ry[506] ; ; tub-Gal4, UAS-GFP	FRT19A: BL1709; MARCM19A: Gift of P. Jouandin
5A, top	*w ; P{w[+mC]=UAS-rpr.C}14 / + ; P{w[+mC]=tubP-GAL80[ts]}ncd[GAL80ts-7] / +*	UAS-rpr: BL5824; tubGal80ts: BL7018
5A, bottom	*w ; P{w[+mC]=UAS-rpr.C}14 / TI{GFP[3xP3.cLa]=CRIMIC.TG4.2}Ca-Ma2d[CR01742-TG4.2] ; P{w[+mC]=tubP-GAL80[ts]}ncd[GAL80ts-7] / +*	Elovl7-Gal4: BL67431
5B, top	*w ; P{w[+mC]=UAS-rpr.C}14 / + ; P{w[+mC]=tubP-GAL80[ts]}ncd[GAL80ts-7] / +*	
5B, bottom	*w ; P{w[+mC]=UAS-rpr.C}14 / + ; P{w[+mC]=tubP-GAL80[ts]}ncd[GAL80ts-7] / TI{GFP[3xP3.cLa]=CRIMIC.TG4.2}CG11550[CR02741-TG4.2]*	CG11550-Gal4: BL97589
5C, top	*w ; P{w[+mC]=UAS-rpr.C}14 / + ; P{w[+mC]=tubP-GAL80[ts]}ncd[GAL80ts-7] / +*	
5C, bottom	*w ; P{w[+mC]=UAS-rpr.C}14 / + ; P{w[+mC]=tubP-GAL80[ts]}ncd[GAL80ts-7] / Mi{Trojan-GAL4.0}Elovl7[MI01455-TG4.0]*	Ca-Ma2d-Gal4: BL86499

6A, left	*w ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4.C-int}attP40 / P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=Tub-GAL4.C-int}attP2*	GMR71F06-Gal4-C-int: This manuscript
6A, middle	*w ; P{y[+t7.7] w[+mC]=Tub-GAL4.C-int}attP40 ; TI{RFP[3xP3.PB]=2A-GAL4(1-20)::N-int}Delta[G4-N-int] / P{w[+mC]=UAS-2xEGFP}AH3*	Delta-Gal4-N-int: BL602717
6A, right	*w ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4.C-int}attP40 / P{w[+mC]=UAS-2xEGFP}AH2 ; TI{RFP[3xP3.PB]=2A-GAL4(1-20)::N-int}Delta[G4-N-int]*	
6B	*w ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4.C-int}attP40 / P{w[+mC]=UAS-2xEGFP}AH2 ; TI{RFP[3xP3.PB]=2A-GAL4(1-20)::N-int}Delta[G4-N-int]*	
6C	*w ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4.C-int}attP40 / w ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4.C-int}attP40 / P{w[+mC]=UAS-2xEGFP}AH2 ; TI{RFP[3xP3.PB]=2A-GAL4(1-20)::N-int}Delta[G4-N-int] ; TI{RFP[3xP3.PB]=2A-GAL4(1-20)::N-int}Delta[G4-N-int] / P{w[+mC]=tubP-GAL80[ts]}ncd[GAL80ts-7]*	GTRACE: BL28280
7	*w ; P{UAS-Stinger}2 / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / MKRS*	UAS-Stinger: BL90920, GMR71F06-Gal4: BL39596
8A, left	*w ; P{UAS-Stinger}2, P{w[+mC]=tubP-GAL80[ts]}10] / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / +*	
8A, middle	*w ; P{UAS-Stinger}2, P{w[+mC]=tubP-GAL80[ts]}10] / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / P{w[+mC]=UAS-CycE.L}ML1*	UAS-CycE: BL4781
8A, right	*w ; P{UAS-Stinger}2, P{w[+mC]=tubP-GAL80[ts]}10] / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / M{w[+mC]=UAS-Myc.HA.WT}ZH-86Fb*	UAS-Myc: BL64759
8C, left	*w ; P{w[+mC]=tubP-GAL80[ts]}10], P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / +*	
8C, second to left	*w ; P{w[+mC]=tubP-GAL80[ts]}10], P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / P{y[+t7.7] v[+t1.8]=VALIUM20-mCherry.RNAi}attP2*	mCherry RNAi: BL35785
8C, second to right	*w ; P{w[+mC]=tubP-GAL80[ts]}10], P{w[+mC]=UAS-2xEGFP}AH2 / P{y[+t7.7] v[+t1.8]=TRiP.HMS03166}attP40 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / +*	InR RNAi: BL51518

8C, right	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / P{y[+t7.7] v[+t1.8]=TRiP.GL00139}attP2*	InR RNAi: BL35251
8D	Same as 8C, except UAS-Stinger instead of UAS-2xEGFP	
S1	*Oregon R*	
S2A, left	*w ; esg-Gal4, UAS:GFP, tubGal80[ts] ; +*	
S2A, right	*w ; esg-Gal4, UAS:GFP, tubGal80[ts] ; P{y[+t7.7] w[+mC]=UAS-yki.S111A.S168A.S250A.V5}attP2*	
S2C	*w ; esg-Gal4, UAS:GFP, tubGal80[ts] ; +*	
S2C	*w ; esg-Gal4, UAS:GFP, tubGal80[ts] ; UAS:Impl2*	UAS:ImpL2: Gift of Ying Liu
S2C	*w ; esg-Gal4, UAS:GFP, tubGal80[ts] / UAS:upd3 ; +*	UAS:upd3: Gift of Ying Liu
S2C	*w ; esg-Gal4, UAS:GFP, tubGal80[ts] / UAS:upd3 ; UAS:Impl2*	UAS:Impl2; UAS:upd3; Gift of A. Petsakou
S2D	*w ; dMef2-Gal4 / + ; tubGal80[ts] / +*	dMef2-Gal4: Perrimon Lab stock
S2D	*w ; dMef2-Gal4 / UAS:Impl2 ; tubGal80[ts] / +*	UAS:ImpL2: Gift of Ying Liu
S2D	*w ; dMef2-Gal4 / + ; tubGal80[ts] / UAS:upd3*	UAS:upd3: Gift of Ying Liu
S2D	*w ; dMef2-Gal4 / UAS:upd3 ; tubGal80[ts] / UAS:Impl2*	UAS:Impl2; UAS:upd3; Gift of A. Petsakou
S3	hsFLP, FRT19A, tubGal80 / P{ry[+t7.2]=neoFRT}19A; ry[506] ; ; tub-Gal4, UAS-GFP	FRT19A: BL7109; MARCM19A: Gift of P. Jouandin
S4, left	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2*	GMR71F06-Gal4: BL39596, UAS:2xEGFP: BL60292, tubGal80ts: BL7108
S4, middle	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / P{y[+t7.7] w[+mC]=UAS-yki.S111A.S168A.S250A.V5}attP2*	yki[3SA]: BL28817
S4, right	*w ; P{w[+mC]=tubP-GAL80[ts]}10, P{w[+mC]=UAS-2xEGFP}AH2 ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / Ras[1A]*	Ras1A: gift of Chiwei Xu, Perrimon Lab

S5	*w ; P{UAS-Stinger}2 / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / MKRS*	UAS-Stinger: BL90920, GMR71F06-Gal4: BL39596
S6	*w ; P{UAS-Stinger}2 / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / MKRS*	UAS-Stinger: BL90920, GMR71F06-Gal4: BL39596
S7	*w ; P{UAS-Stinger}2 / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / MKRS*	
S8	*w ; P{UAS-Stinger}2 / + ; P{y[+t7.7] w[+mC]=GMR71F06-GAL4}attP2 / MKRS*	
S9,S10	See Figure 8.	

1135