The role of the L421P mutation in Penicillin-Binding Protein 1 (PBP1) in the evolution of chromosomally mediated penicillin resistance in *Neisseria gonorrhoeae*

Running Title: Evolution of Penicillin Resistance in *N. gonorrhoeae*

Gabriella Gentile[a], Bryan Guzman[a], Adriana Le Van[b,c], Ann E. Jerse[b], Yonatan H. Grad[d], Daniel Dominguez[a,e,f,g,h], Tatum D. Mortimer[i] & Robert A. Nicholas[a,j#]

[a]Department of Pharmacology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[b]Department of Microbiology and Immunology, Uniformed Services University of the Health Sciences, Bethesda, MD 20814, USA
[c]Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc, Bethesda, MD 20817, USA
[d]Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of Public Health, Boston, MA 02115, USA
[e]Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[f]Department of Bioinformatics and Computational Biology Curriculum, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[g]RNA Discovery Center, University of North Carolina, Chapel Hill, NC27599, USA
[h]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC27599, USA
[i]Department of Population Health, College of Veterinary Medicine,University of Georgia, Athens, GA 30602, USA
[j]Department of Microbiology and Immunology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

[#]Corresponding author. Email: robert_nicholas@med.unc.edu

Abstract:

$ponA^{L421P}$ encodes a mutated variant of penicillin-binding protein 1 (PBP1) and is a key resistance determinant that increases the penicillin MIC (MIC_{PEN}) above the clinical breakpoint in *Neisseria gonorrhoeae*. Despite the removal of penicillin from treatment guidelines for gonococcal infections in the 1980s, $ponA^{L421P}$ is present in nearly 50% of current *N. gonorrhoeae* isolates in the PubMLST database. Bioinformatic analysis indicates that $ponA^{L421P}$ is exclusive to *N. gonorrhoeae* isolates, whereas Leu-421 is 100% conserved in other *Neisseria* species. To understand the involvement of $ponA^{L421P}$ in antibiotic resistance, we introduced *ponA* variants encoding 16 different amino acids at position-421 into FA6140, a penicillin-resistant gonococcal isolate that naturally harbors $ponA^{L421P}$. Proline-421 was the only mutation that increased the MIC_{PEN} to the same level as FA6140. We also assessed the fitness of strains with the 16 mutant *ponA* alleles over multiple serial passages, both with and without sub-MIC levels of penicillin. There was no fitness defect attributed to $ponA^{L421P}$ under these experimental conditions; instead, our analyses suggest that the widespread occurrence of $ponA^{L421P}$ is driven by its capacity to increase the MIC_{pen} above the clinical breakpoint. In FA6140 transformed with the mosaic *penA* allele from strain H041, a ceftriaxone-resistant isolate, $ponA^{L421P}$ increased the MIC of ceftriaxone, suggesting that ceftriaxone targets PBP1 in this strain. We conclude that the $ponA^{L421P}$ allele emerged in gonococcal isolates, increasing the MIC_{PEN} above the clinical breakpoint, and has remained in the population even after the removal of penicillin from treatment guidelines.

Importance:

The emergence of antibiotic-resistant *Neisseria gonorrhoeae* threatens effective treatment of gonorrhea, one of the most common sexually transmitted infections worldwide. Understanding the genetic changes that drive and maintain resistance is crucial for anticipating future resistance trends. Here, we investigated the impact of a key resistance mutation in PBP1 (encoded by *ponA*^{L421P}). Although penicillin has not been used to treat gonorrhea for decades, this mutation remains widespread even in recent *N. gonorrhoeae* isolates. *ponA*^{L421P} confers clinically relevant penicillin resistance without imposing an *in vitro* fitness cost. *ponA*^{L421P} also increases resistance to ceftriaxone in strains with *penA* alleles that are associated with ceftriaxone resistance. This work highlights the role of the *ponA*^{L421P} allele in shaping the current antibiotic resistance landscape and supports the need for ongoing surveillance and evolutionary studies of such mutations in the gonococcal population.

Introduction:

Neisseria gonorrhoeae is the etiologic agent of the sexually transmitted infection gonorrhea, which causes ~80 million infections annually (1, 2). The β-lactam antibiotic penicillin was first used to treat gonorrhea in the early 1940s. For the following 40 years, penicillin remained effective, but the minimum inhibitory concentrations of penicillin (MIC_{PEN}) steadily increased, requiring higher doses to cure infections (3). By 1985, over 5% of circulating N. gonorrhoeae strains had MIC_{PEN} values above the breakpoint (≥2 μg/ml), prompting a switch in treatment guidelines to other antibiotics. The rise in penicillin resistance was driven by two mechanisms: plasmid-mediated production of penicillinases (TEM-1 and TEM-135 β-lactamases) and expression of resistance determinants that are variants of chromosomal genes (4–11). In the US, the percentage of penicillinase-producing N. gonorrhoeae isolates (PPNG) decreased by the 1990s, while the percentage of chromosomally mediated penicillin-resistant N. gonorrhoeae isolates (CMRNG) increased (2, 8).

β-lactam antibiotics target penicillin-binding proteins (PBPs), which are involved in the cross-linking of the peptide chains of peptidoglycan. N. gonorrhoeae has four PBPs, two of which (PBP1 and PBP2) are essential. PBP1 is a bifunctional PBP that catalyzes both glycan polymerization (transglycosylation) and peptide cross-linking (transpeptidation), and PBP2 is a monofunctional PBP, catalyzing transpeptidation reactions. Because PBP2 has a much higher rate of acylation than PBP1 for penicillin and other β-lactam antibiotics used to treat N. gonorrhoeae in susceptible strains, PBP2 is considered the lethal target for these antibiotics. CMRNG is facilitated by the step-wise transfer (12–14) of at least four resistance determinants from a penicillin-resistant strain

to an antibiotic-susceptible strain: 1) mutant *penA* alleles, which encode variants of PBP2

that reduce the acylation rate of penicillin by 16-fold (15); 2) *mtr* mutations, which increase

mtrCDE operon expression and result in increased penicillin efflux through the MtrCDE

efflux pump (9, 16); 3) mutations in *porB*, which encodes the major outer membrane porin,

that reduce the influx of penicillin into the periplasm (7, 17); and 4) *ponA*[L421P], encoding a

PBP1 variant with an L421P missense mutation. Mutations in *penC* were thought to

contribute to high-level penicillin resistance (8), but *penC* was later shown to encode the

pore-forming secretion protein, PilQ (18). These *pilQ* mutations compromise its secretin

function and increase the MIC of multiple antibiotics (18, 19), but to date, *pilQ* mutations

have not been observed in clinical isolates, as they disrupt the type IV pili that play a key

role in gonococcal pathogenesis (20, 21). While the first three resistance determinants

are well characterized in the literature, the contribution of *ponA*[L421P] to penicillin resistance

remains less well understood.

Our laboratory has shown that the L421P mutation in PBP1 lowers the acylation

rate by 2- to 3-fold for penicillin G and other β-lactam antibiotics (8), but it is not clear

whether this lower acylation rate explains the role of *ponA*[L421P] in facilitating penicillin

resistance. Transformation of the *penA, mtr,* and *porB* determinants from strain FA6140,

a high-level-penicillin-resistant clinical isolate (MIC$_{PEN}$ = 4 µg/ml), into the antibiotic-

susceptible strain FA19 (MIC$_{PEN}$ = 0.01 µg/ml) results in MIC$_{PEN}$ between 0.5 to 1.0 µg/ml

(12, 13), but further transformation with FA6140 DNA or *ponA*[L421P] does not increase the

MIC$_{PEN}$ (22). All attempts to reach donor levels of resistance in gonococcal isolates by

natural transformation have been unsuccessful (8, 10, 12). By contrast, we showed that

reversion of *ponA*[L421P] in two penicillin-resistant clinical isolates, including FA6140, back

to the wild-type *ponA* allele decreases the MIC$_{PEN}$ 2-fold (22). These data imply that

ponA^{L421P} requires a yet to be determined genetic background that cannot be transferred

between strains by transformation to exert its phenotypic effect on resistance.

We hypothesized that *ponA*^{L421P} arose during the years of penicillin use and has

continued to remain in the population because *ponA*^{L421P} continues to confer a fitness

benefit, even with the current β-lactam antibiotics recommended for use*.* In the present

study, we sought to test this prediction by investigating the origins of *ponA*^{L421P} to

elucidate the relationship between *ponA*^{L421P} and the evolution of β-lactam resistance in

N. gonorrhoeae.

Results:

***ponA*^{L421P} is exclusive to *N. gonorrhoeae* among *Neisseria* species.**

N. gonorrhoeae isolates can acquire resistance-conferring mutations from commensal *Neisseria* species and from *N. meningitidis* (23–25). We examined whether *ponA*^{L421P} arose via transfer of DNA from other *Neisseria* species, or if this mutation is specific to gonococcal isolates. We accessed the sequences of all unique *ponA* alleles across *Neisseria* species from the PubMLST Database, and divided them into three groups: (1) *N. gonorrhoeae* (n = 20,513 isolates, 172 *ponA* alleles), (2) *N. meningitidis* (n = 75,612 isolates, 897 *ponA* alleles), and (3) all other commensal *Neisseria* species (n = 1,604 isolates, 369 *ponA* alleles).

The PBP1 protein sequences across all three groups of *Neisseria* isolates are highly conserved (Fig. S1), particularly the region surrounding residue 421 (Fig. 1). Leu-421 is 100% conserved across all *N. meningitidis* and commensal *Neisseria ponA* alleles (Fig. 1B-C), but for *N. gonorrhoeae* isolates, residue 421 is either a leucine or a proline (Fig. 1A). Moreover, *ponA*^{L421P} was observed in 49.5% of all gonococcal isolates in the PubMLST database and was not found in other *Neisseria* species (Table S5). These data strongly suggest that *ponA*^{L421P} arose in *N. gonorrhoeae* and not through horizontal gene transfer from other *Neisseria* species.

***ponA*^{L421P} is prevalent amongst gonococcal isolates with chromosomally-mediated penicillin resistance.**

To investigate the emergence and distribution of *ponA*^{L421P} in the gonococcal population, we assembled a dataset of sequenced *N. gonorrhoeae* isolates and the associated MIC$_{PEN}$ values (n=16,891). The first appearance of *ponA*^{L421P} in a sequenced

isolate was in 1961, and it rapidly expanded in the population in the 1980s (Fig. 2).

Interestingly, we first observed the insertion of aspartate after position 345 in *penA*

(Asp345a)*,* which is a hallmark of early penicillin-resistant isolates (26, 27), in our dataset

of sequenced isolates in 1956, only a few years prior to the L421P mutation in *ponA* (Fig.

2)*.* We examined the phylogenetic distribution of *ponA* alleles in a representative sample

of *N. gonorrhoeae* isolates without plasmid-mediated resistance (n=5,192) and found

ponA^{L421P} emerged across multiple lineages in the gonococcal population (Fig. 3).

We next examined the relationship between *ponA*^{L421P} and penicillin-resistant

CMRNG strains (Fig. 4, n=8,999). Less than 1% of isolates harboring the wild-type *ponA*

allele included in Fig. 4 have resistant MIC$_{PEN}$ values (≥2 µg/mL), while the majority (90%)

fall within the intermediate MIC$_{PEN}$ range (0.12–1 µg/mL). In contrast, 25% of isolates

harboring *ponA*^{L421P} have resistant MIC$_{PEN}$ (≥2 µg/mL), and 75% of isolates have

intermediate MIC$_{PEN}$. Few isolates, regardless of the *ponA* or *ponA*^{L421P} allele, had

susceptible MIC$_{PEN.}$ Additionally, 91.61% of isolates that harbor mosaic *penA* alleles, which

encode highly remodeled PBP2 variants that confer increased resistance to cephalosporins,

also harbor *ponA*^{L421P} (Fig. 4B).

Identification of other polymorphisms in gonococcal *ponA*.

Amongst *N. gonorrhoeae* isolates in PubMLST*,* there were only two other

mutations observed in gonococcal PBP1 at frequencies above 1%: A375T (7.07%) and

Y537C (2.35%) (Fig. 5). In keeping with these frequencies, a phylogeny of 6,082

representative gonococcal genomes reveals these mutations are present in fewer

lineages, compared to the phylogenetic distribution of the L421P mutation (Fig. S2). We

next examined the relationship between each mutation and penicillin-resistant CMRNG

strains (n=8,999). The majority (95%) of isolates harboring the A375T mutation in *ponA*

have intermediate (0.12–1 µg/mL) to resistant (≥ 2 µg/mL) MIC_{PEN}. This proportion does

not change in isolates with alanine at residue 375 of *ponA*, as 95% of these isolates also

have intermediate to resistant MIC_{PEN}. The same trend is observed with the Y537C

mutation. We do not observe A375T and Y537C in sequenced isolates collected during

the penicillin treatment era, as they first appear in 1998 and 2001, respectively, well after

penicillin was removed as a treatment for gonorrhea infections in the US in 1985. It

remains unclear whether A375T and Y537C are directly involved in facilitating resistance

to penicillin, but, given the fact that isolates with those mutations are not enriched for

penicillin resistance and they are present in fewer gonococcal lineages than L421P, we

did not interrogate them further.

Only proline and serine at PBP1 position 421 increase the MIC_{PEN}.

To understand why only a proline mutation at residue 421 is observed in PBP1*,*

we created an isogenic library of 16 mutant strains in FA6140, each with a different amino

acid at position 421 (Table 1). FA6140 (MIC_{PEN}= 4 µg/mL) naturally harbors $ponA^{L421P}$,

and reversion to $ponA^{421L}$ reduces the MIC_{PEN} to 2 µg/mL (8, 22) (Table 1). In the strain

library, FA6140 $ponA^{L421P}$ was the only strain with an MIC_{PEN} of 4 µg/mL. FA6140

$ponA^{L421S}$, which had a median MIC_{PEN} of 2.5 µg/mL, was the only other mutation that

increased the MIC_{PEN} > 2 µg/mL, with all the other strains having the same MIC_{PEN} as

FA6140 *ponA* (Table 1).

***In vitro* serial passaging of the FA6140 $ponA^{L421X}$ mutant library in the absence or**

presence of sub-MIC concentrations of penicillin.

The 2-fold higher MIC_{PEN} conferred by $ponA^{L421P}$ and the timing of its first appearance

in sequenced isolates suggest that $ponA^{L421P}$ emerged in response to selection by

penicillin. To test this, we performed *in vitro* serial passaging on our set of FA6140

$ponA^{L421X}$ mutants in increasing sub-MIC_{PEN} concentrations (0.25, 0.5, and 1.0 µg/ml).

Illumina sequencing revealed that equal amounts of each FA6140 $ponA^{L421X}$ mutant were

used to inoculate the cultures at hour 0 across all 3 trials (Fig. 6A, Fig. S3). In the absence

of penicillin, the amino acids proline, cystine, asparagine, and glycine at position 421

appear to be more fit in this growth environment, as they have the largest fold increase in

read counts at hour 36 relative to the inoculum (Fig. 6C). The $ponA^{L421P}$ reads consistently

doubled by hour 36 in all three trials, while the $ponA^{421L}$ reads remained similar to, or

decreased from, the number of reads in the inoculum. The proline and serine mutants

were the only strains with an MIC_{PEN} above 2 µg/mL (Table 1), but the read counts for the

serine mutant at hour 36 did not surpass its read counts at hour 0 (Fig. 6C). None of the

strains differed significantly in growth rate from the parental strain (Table S4).

We grew the collection of strains in growth medium with increasing concentrations of

penicillin to examine which strains would grow in the presence of selective pressure.

Optical Density at 600 nm (OD_{600}) values during growth in medium containing 0.25 mg/ml

penicillin were very similar to no antibiotic (Fig. S4), while there was more variability in

the OD_{600} values across trials for the cultures supplemented with 0.5 µg/mL of penicillin

(Fig. 6B). Of the 16 mutants, the two with the highest MIC_{PEN}, FA6140 $ponA^{L421P}$ and

FA6140 $ponA^{L421S}$, increased in read counts when grown in 0.5 µg/mL penicillin, while all

the others decreased in read counts. FA6140 $ponA^{L421P}$ consistently increased in read

counts across all three trials, while read counts for FA6140 $ponA^{L421S}$ increased in trials

1 and 2, but not in trial 3. In trial 1, FA6140 $ponA^{L421P}$ had more read counts at 36 h than

FA6140 $ponA^{L421S}$, and both strains had similar reads counts at 36 h in trial 2 (Fig. 6C).

FA6140 $ponA^{L421P}$ is the only mutant that consistently increased in read counts over 36 h

in cultures either with or without sub-MIC_{PEN} concentrations.

At the highest sub-MIC_{PEN} concentration tested, we observed more variable results.

After 6 h of growth and passaging into fresh medium with 1 μg/ml penicillin, the OD_{600}

increased very little over the next 12 h (Fig. 7A). However, in two of the three trials (1 &

3), there were still live cells present, as later passages showed measurable growth during

the 6 h period. These data suggest that gonococcal cells remained viable but were not

dividing during the prior 6 h growth periods and eventually acquired either phenotypic or

genetic changes that allowed for measurable growth at the later growth periods. In trial 1,

we observed increasing growth during consecutive passages, beginning at 18 h. Illumina

sequencing of the *ponA* region revealed that serine was the predominant mutation at

position 421 (Fig. 7B).

Whole genome sequencing of cells from *in vitro* serial passaging FA6140 *ponA*^{L421X}

mutant library reveals emergence of a mutation in *pilQ*.

To determine if the FA6140 $ponA^{L421X}$ mutants acquired additional mutations during

the 36 hours of *in vitro* growth, we sequenced the genomes of cells collected at 0 h and

36 h. We looked for mutations that were distinct from the inoculum and present in greater

than 20% of the reads, as anything detected below this would likely not be responsible

for any changes in the overall codon counts from the Illumina reads (Figs. 6D, 7B). Across

all 3 trials, most, but not all, mutations above the 20% threshold were either errors in the

reference genome sequence used for alignment (not true genetic changes), also present

in the inoculum, or in genes that would most likely be inconsequential to resistance.

In trial 1, we detected a mutation in *pilQ* at 36 h at 85% frequency in the culture

supplemented with 1.0 µg/mL of penicillin. The mutation was an insertion of two bases

(+CG) at position 459, which would result in a frameshift and early termination. Previous

work from our lab showed that missense and frameshift mutations in *pilQ* increase MIC$_{PEN}$

by inactivating PilQ and decreasing permeation of drug into the gonococcal cell (8, 18,

19, 28), which likely explains the marked increase in codon counts for the serine mutation

at 24 h seen in trial 1 (Fig. 7C). Our data suggest that during trial 1, FA6140 *ponA*^{L421S}

acquired a mutation in *pilQ* between hours 18 and 24, which allowed it to expand in the

culture compared to FA6140 *ponA*^{L421P}.

ponA^{L421P} increases the ceftriaxone MICs of gonococcal isolates that also harbor a

mosaic _penA_ allele.

Our data indicate that *ponA*^{L421P} is involved in penicillin resistance, but penicillin

was removed as a recommended treatment in 1985, and ceftriaxone has been the only

recommended β-lactam antibiotic since 2012 for treating gonococcal infections. Given

that *ponA*^{L421P} appears to confer a modest fitness deficit *in vivo* (29), it might be expected

that *ponA*^{L421P} would decrease in prevalence without selection by penicillin use. However,

ponA^{L421P} is observed in a high percentage of isolates with a mosaic *penA* allele (91.61%,

Fig. 5B), which is the predominant determinant for decreased susceptibility to ceftriaxone

(30, 31). One possibility to explain the prevalence of *ponA*^{L421P} is that *ponA*^{L421P} influences

the MIC of ceftriaxone in the presence of a mosaic *penA* allele.

To test this hypothesis, we replaced the *penA4* allele in FA6140 and FA6140 *ponA*

strains with *penA41*, the mosaic allele from the multi-drug resistant gonococcal isolate,

H041 (32). This allele encodes a PBP2 with a markedly reduced acylation rate for

penicillin, cefixime, and ceftriaxone (31, 33, 34). Reversion of *ponA*^{L421P} to wild-type *ponA*

reduced the MIC$_{PEN}$ in both strain backgrounds, regardless of the *penA* allele (Fig. 8A),

suggesting that PBP1 is the lethal target of penicillin in these strains. While reversion of

ponA^{L421P} to *ponA* in strain FA6140 had no effect on the MIC of ceftriaxone (Fig. 8B),

reversion of *ponA*^{L421P} to *ponA* in strain FA6140 *penA41* caused a 3-fold decrease in the

MIC of ceftriaxone, suggesting again that PBP1 is the lethal target of ceftriaxone in this

strain.

Discussion:

In this study, we sought to understand the origins and role of the PBP1-L421P

mutation (encoded by the $ponA^{L421P}$ allele) in β-lactam antibiotic resistance in *N.*

gonorrhoeae. Our data revealed that $ponA^{L421P}$ is found exclusively in *N. gonorrhoeae*

and its emergence in gonococcal isolates is likely due to it increasing the MIC_{PEN} above

the clinical breakpoint of 2 μg/ml in CMRNG. While $ponA^{L421P}$ has a modest fitness cost

in vivo (29), there is no apparent fitness cost during growth *in vitro*. Our data suggest that

this cost turns into a fitness benefit during growth in sub-MIC_{PEN} concentrations. Moreover,

while penicillin G is no longer used to treat gonococcal infections, $ponA^{L421P}$ also

increases the MIC of ceftriaxone in strains harboring mosaic *penA41* alleles that confer

ceftriaxone resistance. Thus, the increases in MIC_{PEN} (in FA6140) and MIC_{CEF} (in FA6140

penA41) that result from harboring $ponA^{L421P}$ appear to have provided enough of an

evolutionary advantage to remain in the population despite its modest fitness cost.

Mosaic *penA* alleles are defined by a sub-set of mutations that surround the active

site of PBP2 and decrease the conformational dynamics of the PBP, which reduces

acylation rates and increases MICs for various β-lactam antibiotics (15, 30, 31). These

mutations emerged in *N. gonorrhoeae* through interspecies recombination events with

Neisseria commensals. In contrast, $ponA^{L421P}$ appears to have arisen spontaneously in

N. gonorrhoeae, and, unlike resistance-conferring mutations in PBP2, L421P is located

on a hinge that links the oligosaccharide-binding domain to the start of the transpeptidase

domain in PBP1 and is over 20 Å from the active-site serine residue (Fig. 5). How the

L421P mutation decreases acylation is unknown, and compared to mosaic PBP2 variants,

the effects on k_2/K_S are modest (8). The small decrease in acylation, however, could

explain the 2-fold increase in the MIC$_{PEN}$.

While $ponA^{L421P}$ was the only mutation in FA6140 to increase the MIC$_{PEN}$ to 4 μg/ml,

another mutation, $ponA^{L421S}$, increased the MIC$_{PEN}$ to 2.5 μg/mL. However, to date

$ponA^{L421S}$ has not been documented in clinical isolates. This could be due to three

reasons: 1) the L421P mutation requires only a single base change in the Leu-421 codon

in $ponA$ (C$\underline{T}$G → C$\underline{C}$G), whereas the L421S mutation would require at least two base

changes (CTG → $\underline{TCT}$/$\underline{TCC}$/$\underline{TCA}$/$\underline{TC}$G/$\underline{AGT}$/$\underline{AGC}$); 2) strain FA6140 $ponA^{L421S}$ is less fit

in vitro compared to other strains; and 3) the increase in resistance is insufficient to

overcome its fitness cost.

Following the six serial passages of the 16 mutants, we sequenced the genomes of

the 36-hour cultures from all three biological replicates to be sure that the results were

not skewed by the acquisition of random mutations, particularly under antibiotic selective

pressure. No consistent or biologically relevant mutations were detected, with the

exception of a single frameshift mutation in *pilQ* observed in the culture supplemented

with 1.0 μg/mL of penicillin during trial 1. PilQ increases the MICs of a range of

antimicrobials in *N. gonorrhoeae* strains that also have the *mtr* and *porB* resistance

determinants (18, 19), which decrease the concentration of antibiotics in the periplasm by

increasing efflux and decreasing influx of antibiotics, respectively. This implies that

permeation through the PilQ secretin is low and does not impact the MIC unless both *mtr*

and *porB* determinants are present, as they are in FA6140 (3, 8, 18, 22, 35). Based on

the increase in read counts for $ponA^{L421S}$ at 24 hours, we surmise that the PilQ mutation

arose in FA6140 $ponA^{L421S}$ sometime between 18-24 hours of growth, which increased

its MIC$_{PEN}$ above that of FA6140 *ponA*^{L421P} and allowed it to survive in higher penicillin

concentrations. However, this is an outcome that would likely occur only during *in vitro*

growth, as expression of type IV pili (of which PilQ is an essential component) is important

for *N. gonorrhoeae* survival during pathogenesis (36). This is supported by the absence

of inactive PilQ mutants identified in clinical isolates (37, 38).

We recently reported that *N. gonorrhoeae* lineages harboring *ponA*^{L421P} between

1993 and 2013 were associated with a modest fitness disadvantage, compared to the

baseline type (lineages that do not carry any resistance determinants), suggesting that

ponA^{L421P} imposes a fitness cost (29). These data were consistent with results from

competitive co-infections between *ponA*- and *ponA*^{L421P}-containing strains of FA19 and

FA6140 (29). Despite this observed fitness cost, we showed here that *ponA*^{L421P} remains

prevalent in the gonococcal population and is present in over 90% of isolates that also

harbor a mosaic *penA* allele (as classified by the NG-STAR database). Why is *ponA*^{L421P}

found so frequently in strains with a mosaic *penA* allele? There are several possible

explanations for this: 1) mosaic *penA* alleles emerged by transformation almost

exclusively into penicillin-resistant strains that already had *ponA*^{L421P}, as these strains

also contain other resistance determinants and presumably compensatory mutations that

lessen their fitness cost; 2) with penicillin resistance at or near 100% in Asia (39), where

ceftriaxone resistance has been emerging, the link between *ponA*^{L421P} and ceftriaxone

resistance may reflect the gonococcal diversity/population structure in the places

experiencing the most selection for ceftriaxone resistance; and 3) *ponA*^{L421P} confers

additional resistance to ceftriaxone and thus provides a benefit to these strains, given the

widespread use of ceftriaxone to treat current infections. We tested the latter possibility

by determining the MIC$_{CEF}$ values of FA6140 and FA6140 *ponA* also harboring the mosaic

penA41 allele from H041, which confers ceftriaxone resistance to these strains.

Importantly, the MIC$_{CEF}$ drops ~3-fold when *ponA*[L421P] is replaced with *ponA*, indicating

that *ponA*[L421P] confers additional resistance to ceftriaxone and potentially explaining, at

least in part, why *ponA*[L421P] is so prevalent in strains also harboring mosaic *penA* alleles,

particularly with mosaic *penA* alleles that confer the largest increases in MIC$_{CEF}$.

The MIC data we present in this study can be best interpreted by knowing which PBP

is being targeted by penicillin and ceftriaxone (Fig. 9). The acylation constant k_2/K_S for

each antibiotic determines which PBP first becomes acylated as the antibiotic

concentration increases (the higher the k_2/K_S, the lower the concentration of antibiotic

needed to inhibit the PBP). Thus, k_2/K_S and the MIC have an inverse relationship, and the

essential PBP that first becomes acylated as the antibiotic concentration increases is the

killing target for that antibiotic and determines the MIC. For penicillin, the killing target for

FA6140 is likely a combination of PBP2[FA6140] and PBP1[L421P], but when PBP1[L421P] is

replaced by PBP1, the MIC decreases because PBP1 becomes the exclusive killing

target of penicillin. We showed previously that in 35/02, a ceftriaxone-intermediate

resistant (and penicillin-resistant) gonococcal isolate, reversion of *ponA*[L421P] to *ponA* did

not alter MIC$_{CEF}$ values (40). In contrast, expression of PBP2[H041] in FA6140, which has a

much lower k_2/K_s for ceftriaxone than PBP2[35/02], shifts the killing target to PBP1[L421P]. Thus,

exchanging PBP1[L421P] for PBP1 in the FA6140 genetic background decreases MIC$_{CEF}$ by

3-fold, suggesting that when strains acquire a "strong" mosaic PBP2 variant, e.g.

PBP2[H041], that has a sufficiently low acylation rate constant for ceftriaxone, the target

shifts from PBP2 to PBP1 and *ponA*[L421P] confers a selective advantage over *ponA.*

In conclusion, this is the first study to investigate the origins of $ponA^{L421P}$ in the context

of penicillin resistance and address why it is maintained in the *N. gonorrhoeae* population.

During the 40+ years of penicillin use, *N. gonorrhoeae* strains acquired β-lactamases, as

well as mutations in PBP1, PBP2, *mtr*, and *porB*, which eventually increased the MIC_{PEN}

enough to result in its removal as a treatment option. When expanded-spectrum

cephalosporins (ESCs) were introduced to treat gonococcal infections, mosaic PBP2

variants emerged (because PBP2 was the exclusive killing target of ESCs), in existing

penicillin-resistant strains, resulting in the proliferation of mosaic *penA* alleles that

increased resistance to ESCs. Over time, the MICs of ESCs for gonococcal isolates

increased as the mosaic *penA* alleles picked up additional mutations, and the killing target

shifted to PBP1, allowing the L421P mutation to further increase the MIC. These findings

reinforce the need to identify antibiotics that are PBP1-selective and thus can avoid the

highly mutated mosaic PBP2 variants that are compromising the use of ESCs. β-lactam

antibiotics that target both PBPs or a combination of two β-lactam antibiotics that target

PBP2 and PBP1 may provide a new strategy for treating strains with mosaic *penA* alleles.

Methods:

Strains and plasmids.

Strains, plasmids, and primers used in this study are listed in Tables S1 and S2. To

aid in subsequent screening for generating random mutations at codon-421, we

transformed FA6140 with the *ponA* allele from FA19, which naturally has a PstI site

encompassing codons 421 and 422. A portion of the *ponA* gene (bp 956 – 2400) from

FA19 genomic DNA was amplified via PCR and transformed into FA6140 via spot

transformation, performed as described (41)*.* Colonies were screened for *ponA* by PCR

amplification of *ponA* and digestion with PstI. To generate FA6140 and FA6140 *ponA*

harboring the mosaic *penA* allele (*penA41*) from the ceftriaxone-resistant strain H041,

piliated cells were transformed with pUC18us-*penA41* plasmid containing *penA41* starting

at bp 133 and ending 200 bp downstream *penA41* to facilitate recombination.

Transformants that acquired *penA41* were selected on GCB agar plates containing 1.0

μg/mL cefixime. All transformants were verified by PCR and Sanger sequencing.

Descriptions of the cloning plasmids used to create the library of isogenic FA6140

ponA^{L421X} mutants can be found in Table S3. pUC18us-*ponA*^{421X}-Ω contains a portion of

the *ponA* gene (from bp 831 to 54 bp downstream of the stop codon) harboring

randomized codons to introduce amino acid substitutions at codon-421, the *aad1*

resistance cassette (Ω) conferring spectinomycin/streptomycin resistance (42), and 531

bp of sequence downstream of *ponA* to facilitate recombination. We employed overlap-

extension (43) PCR mutagenesis to introduce amino acid substitutions at codon-421 in

ponA fragments, using the primers listed in Table S2. pUC18us-*ponA*^{421X}-Ω plasmids

were created using the NEBuilder HiFi DNA Assembly Master Mix and transformed into

Escherichia coli XL-10 Gold cells, according to manufacturer's protocol. Clones were

screened by PCR amplification of the *ponA* gene followed by Sanger sequencing.

Plasmids containing unique mutations were isolated using a QIAPrep miniprep kit

(Qiagen). Plasmids (Table S3) were transformed into FA6140 *ponA* by allelic exchange,

and transformants were selected on GCB agar plates containing 25 μg/mL spectinomycin.

Individual colonies from each transformation were screened by PCR by amplifying the

ponA gene and digesting with PstI (loss of PstI indicated the presence of the mutation).

All clones were verified by Sanger sequencing.

PCR reaction conditions were as follows; 94°C for 30s (denaturation), 60°C for 30s

(annealing), and 72°C for 60s (elongation) for 35 cycles, and fragments were purified

using the QIAquick PCR Purification kit (Qiagen) as per manufacturers guidelines. All

gonococcal strains were propagated on solid GCB agar containing Kellogg's supplement

I (38) and 12 mM $Fe(NO_3)_3$ for 18 to 20 h at 37°C in a 5% CO_2-enriched humidified

atmosphere. Serial passaging was minimized to reduce the risk of acquiring secondary

mutations.

Bacterial growth.

FA6140 *ponA*[L421X] mutants were grown in liquid culture to verify individual growth

phenotypes (Table S4). Nonpiliated bacterial colonies were swabbed from GCB agar

grown for 16 h and inoculated into GCB (per liter: 15 g protease peptone 3, 4 g K_2HPO_4,

dibasic, 1 g KH_2PO_4, 5 g NaCl) supplemented with Kellogg's supplement I (38) and 12

mM $Fe(NO_3)_3$ at a starting optical density at 600 nm (OD_{600}) of 0.08. Cultures were shaken

at 180 rpm at 37°C in a 5% CO_2-enriched atmosphere, and bacterial growth was

assessed by measuring the OD_{600} at hourly intervals for a total of 8 h. Experiments were

repeated in biological triplicate, and statistical significance between individual strains was

determined by using a repeated-measures 2-way analysis of variance (ANOVA) with

Tukey's multiple comparisons. Differences in growth rate were measured by comparing

the average time to reach an OD_{600} of 0.8 for each strain as previously described (44).

Results were compared using a one-way analysis of variance to determine overall

significance followed by Dunnett's multiple comparison test to determine significance

between individual strains.

***Neisseria* sequence diversity analysis.**

The datasets used to identify polymorphisms and assess overall sequence diversity

of the *ponA* gene across all *Neisseria* species were all exported from the *Neisseria* page

of the PubMLST data base (last accessed September 30, 2024) (45). Allele mining

analysis of the *ponA* gene was performed as described (46). An individual search of the

profiles was made via the locus/sequence definitions tool to identify the locus assigned

to *ponA* (NEIS0414). Data was exported from the two-field breakdown section of the

Neisseria isolate database as a Microsoft Excel file, which provided a list of unique *ponA*

allelic profiles and a count for how many isolates possess that unique allele within each

Neisseria spp. Microsoft Excel was used to calculate the number of isolates that possess

each allele. Mutations identified in the *ponA* alleles from *N. gonorrhoeae* isolates with

frequency greater than 5% were mapped onto the predicted PBP1 structure, which was

generated using Alphafold (47). The PDB file for PBP1 from *N. gonorrhoeae* (UniProt:

O05131) was downloaded from the AlphaFold Protein Structure Database (AFDB) (48,

49) and mutations detected frequencies greater than 5% were mapped to the structure

using Pymol (2.0.7) (50). *ponA* sequences from PubMLST were exported using the script

available in the allele/sequence definition database. Molecular Evolutionary Genetics

Analysis software, version 7 (MEGA7) (51) was used to translate the sequences and

perform a multiple sequence comparison by log-expectation (MUSCLE) (52). Sequence

logo images were created using WebLogo (53). MUSCLE alignments of unique *ponA*

alleles from PubMLST were uploaded to WebLogo, background composition of proteome

was set to equiprobable, and y-axis units were set to bits.

Genomic Analyses

Genome assembly and typing. Publicly available *N. gonorrhoeae* genomic data with

available MICs of penicillin were downloaded from the European Nucleotide Archive (54–

72). Resistance-associated SNPs were determined from variant calls produced by Pilon

v 1.23 (73) after mapping to the NCCP11945 reference genome (NC_011035.1) using

BWA-MEM v 0.7.17 (74). Mosaic alleles and the presence or absence of resistance-

associated accessory genes (e.g. *bla*$_{TEM}$) were identified using blastn v 2.14.0 (75)

searches of *de novo* assemblies generated with SPAdes v 3.12.0 (76). Mosaic *penA*

alleles were typed according to the NG-STAR database (77).

Genomic data was included if the total *de novo* assembly length was within the range

of expected lengths for *N. gonorrhoeae* (1.89 Mb – 2.31 Mb) and that the number of

contigs was less than 250. We also required more than 40X coverage of the reference

genome, that at least 80% of reads mapped to the reference genome, and less than 12%

of positions in the reference genome had ambiguous calls defined as at least 90% of

reads supporting either the reference or alternate nucleotide. Lastly, we removed isolates

with ambiguous calls in genes that contribute to penicillin resistance, including *ponA,*

penA, and the *mtr* operon.

Phylogenetic analysis. An initial alignment and core genome phylogeny was estimated

using ParSNP v 2.0.5 (78) from 10,282 gonococcal *de novo* assemblies. To reduce the

computational resources needed for a recombination-corrected phylogenetic analysis, we

used Treemmer v 0.3 (79) to prune the ParSNP tree to 99.9% of the original tree length,

removing the most closely related pairs of isolates from our dataset. For this

representative subset of 6,082 isolates, pseudogenomes were generated from variant

calls and concatenated to produce an alignment for recombination detection and

phylogenetic analysis with Gubbins v 3.3.3 (80). Phylogenetic trees were visualized using

iTOL (81). We additionally pruned the dataset to remove isolates encoding *blaTEM*,

leaving a representative dataset of 5,192 isolates to investigate the distribution of *ponA*

among CMRNG.

Antimicrobial susceptibility testing.

All MIC values were determined by the agar dilution method exactly as described

previously (34). Briefly, FA6140 transformants were grown overnight on GCB plates,

resuspended in GCB broth at a final concentration of ~1 × 10^7 colony forming units

(CFU)/ml. Aliquots (5 µl) of each strain were spotted onto GCB plates containing

antibiotics at <2-fold dilutions. The MIC was defined as the lowest concentration of

antibiotic on which fewer than 5 colonies appeared after 24 hrs of incubation.

***In vitro* serial passaging.**

Non-piliated colonies from each individual FA6140 *ponA*^{L421X} strain (Table 1) were

harvested and used to prepare a starting inoculum, which consisted of an equal ratio

(verified through Illumina sequencing) of all 16 strains. An equal volume of the starting

inoculum was used to inoculate liquid cultures at 0 h at an OD_{600} of 0.08. Three cultures

were supplemented with a different sub-MIC concentration of penicillin (0.25, 0.5, and 1.0

μg/mL), and one culture remained penicillin-free. Penicillin was added at hour 0 at the

specified concentration and was maintained throughout each of the six 6-hr growth

periods. After inoculation, cultures grew for 6 hours, at which time the OD_{600} of each

culture was measured to calculate the volume of cells needed to inoculate the next culture

at an OD_{600} of 0.08, and an aliquot was removed for analysis. This passaging procedure

was performed every 6 h for a total of 36 h. At every 6-hour timepoint, the cultures were

checked for contamination by dipping a sterile swab in the media and streaking it out on

a GCB agar plate. This experiment was repeated 3 separate times. All liquid cultures

consisted of 40 mL of GCB + supplements and were shaken at 180 rpm at 37°C in a 5%

CO_2-enriched atmosphere.

Genomic DNA extraction.

DNA was extracted from cells collected every 6 h during the *in vitro* serial passaging

experiments using the Promega Wizard genomic DNAs purification kit (Promega,

Madison, WI) according to the manufacturer's protocol. Extracted gDNA is from a mixture

of cells that represent the ratio of all sixteen FA6140 *ponA*^{L421X} mutants growing in the

culture at that time point. DNA concentrations were measured on the Invitrogen Qubit 2.0

Fluorometer. Samples were prepped using the Qubit™ dsDNA Quantification Assay Kits

according to the manufacturer' guidelines (Invitrogen).

PCR and DNA sequencing of FA6140 *ponA*^{L421X} mutants during *in vitro* serial

passaging.

Library preparation. Fragments for the library were prepped in a series of two PCR

reactions. The first PCR reaction (PCR 1; 20 cycles of 94°C for 30 s, 60°C for 30 s, and

72°C for 15 s) used genomic DNA extracted from cells collected during the serial *in vitro*

passaging experiments as a template. Primers were designed to target a region of the

ponA gene (bp 1212-1309), which encompasses codon-421*,* and to attach the

appropriate adaptor sequences to the 5' and 3' ends of each amplified fragment. The

amplicons obtained were then purified and used as templates in the second PCR (PCR

2; 12 cycles of 94°C for 30 s, 60°C for 30 s, and 72°C for 15 s). Primers used in PCR 2

target the 5' and 3' Illumina adaptor sequences from PCR 1. Barcoding was achieved by

adding a unique, 5 bp sequence to the 5' end of fragments to signify the time point it refers

to. A unique index sequence was attached to the 3' ends of the fragments to mark the

penicillin concentration of the culture from which the fragment was obtained. All PCR

fragments were extracted from 3% agarose gels and purified using the QIAquick Gel

Extraction Kit (QIAGEN) according to manufacturers' guidelines. Concentrations were

measured on the Invitrogen Qubit 2.0 Fluorometer.

Illumina sequencing. Purified fragments from PCR 2 were pooled and sequencing was

performed on an Illumina Next-seq 1000. This was done three separate times, once per

in vitro serial passaging trial.

Data analysis. Read files generated from our libraries contained between 10 and 70

million reads. The reads in this library are essentially all genetically identical copies of the

amplified region of *ponA*, except for codon-421*,* meaning each read correlates to a strain

from the FA6140 *ponA*[L421X] library, based on the amino acid at codon-421. Data were

parsed using a UNIX script to count codons and condition specific barcodes. This process

yielded a table for each timepoint, listing all of the codons identified at position-421, and

a count of how many reads have that codon. Read counts across time points were

532 normalized to the total number of reads in the file. Only informative reads, containing the

533 expected codons at the expected positions, were analyzed to limit normalization and

534 counting based on uncertain read calls. After filtering, we retained ~99% of reads that

535 were informative. Bar graphs were generated by normalizing codon counts at 36 hr to the

536 strain's inoculum value (codon count at 0 h). Since this is a zero-sum competitive

537 experiment, any gain in codon counts by one isolate directly corresponds to a loss in

538 codon counts by the others.

539 **Whole genome sequencing and assembly.**

540  Whole genome sequencing was performed by SeqCenter (Pittsburgh, PA) on genomic

541 DNA from the inoculum and the 36 h timepoint. Illumina sequencing libraries were

542 prepared using the tagmentation-based and PCR-based Illumina DNA Prep kit and

543 custom IDT 10 bp unique dual indices (UDI) with a target insert size of 280 bp. No

544 additional DNA fragmentation or size selection steps were performed. Illumina

545 sequencing was performed on an Illumina NovaSeq X Plus sequencer in one or more

546 multiplexed shared-flow-cell runs, producing 2 x 151 bp paired-end reads. Demultiplexing,

547 quality control and adapter trimming was performed with bcl-convert1 (v4.2.4). Variant

548 calling analysis on the reads was also performed by SeqCenter, using their standard

549 methods. Illumina-generated 2 x 151 bp paired-end read data was used as the input for

550 variant calling against the reference genome of gonococcal clinical isolate, FA6140 (82),

551 the parental strain used to create the library of mutants (GCF_001047255.1). Variant

552 calling was carried out by SeqCenter using BreSeq1 under default settings.

553 **Data availability**

This publication made use of the PubMLST website (http://pubmlst.org/) developed by

Keith Jolley (45) and sited at the University of Oxford. The development of that website

was funded by the Welcome Trust. Data was retrieved and is available from

https://pubmlst.org/organisms/neisseria-spp. Data supporting the findings are included in

the manuscript. All metadata and quality metrics for isolates included in the datasets for

our genomic and phylogenetic analysis are available at https://github.com/mortimer-

lab/ponA421. Experimental data, raw FASTQ files, and tables with codons and counts for

each Illumina sequencing trial, as well as the code used to generate these tables are

available upon request.

Supplemental Material

564
TABLE S1
TABLE S2
TABLE S3
TABLE S5
FIG S1
FIG S2
FIG S3
FIG S4
FIG S5

574
Acknowledgements

The contents of this article are solely the responsibility of the authors and do not

necessarily represent the official views of the Uniformed Services University of the Health

Sciences, the Department of Defense, the University of North Carolina, Harvard

University, and the University of Georgia. This work was supported by grants AI153521

(YHG, AEJ, and RAN) and AI164794 (RAN), as well as the Pharmacological sciences

T32 Training Program award, GM135095 (GG), and the National Institute of Allergy and

Infectious Disease (NIAID) STI/HIV T32 Training Program award, T32AI007001 (GG).

The funders had no role in study design, data collection and interpretation, or the decision

to submit the work for publication.

GG and RAN designed and conceptualized the research, GG performed all

experiments, analyzed data, provided opinions, and wrote the manuscript. TDM

performed genome assembly and typing and phylogenetic analysis, reviewed and edited

the article, and provided opinions. BG and DD assisted with Illumina sequencing,

providing insightful advice on experimental design and data analysis. ALV, AEJ, and YHG

provided input and opinions during the course of this study, and critical review of the

manuscript. Supervision, funding acquisition, and article revisions provided by RAN.

References:

592   1.   World Health Organization. 2024. Multi-drug resistant gonorrhoea. https://www.who.int/news-room/fact-sheets/detail/multi-drug-resistant-gonorrhoea.

2.   Wang SA, National Center for HIV, STD, and TB Prevention (U.S.)., Division of STD Prevention. 1998. Sexually Transmitted Disease Surveillance 1998 Supplement: Gonococcal Isolate Surveillance Project (GISP) Annual Report – 1998. Department of Health and Human Services, Public Health Service. Centers for Disease Control and Prevention, Atlanta, Ga.

3.   Unemo M, Del Rio C, Shafer WM. 2016. Antimicrobial Resistance Expressed by *Neisseria gonorrhoeae*: A Major Global Public Health Problem in the 21st Century. Microbiol Spectr 4:4.3.35.

4.   Spratt BG, Balthazar JT, Hagman KE, Pan W, Judd RC, Shafer WM. 1995. Resistance of *Neisseria gonorrhoeae* to antimicrobial hydrophobic agents is modulated by the *mtrRCDE* efflux system. Microbiology 141:611–622.

5.   Spratt BG. 1988. Hybrid penicillin-binding proteins in penicillin-resistant strains of *Neisseria gonorrhoeae*. Nature 332:173–176.

6.   Pan W, Spratt BG. 1994. Regulation of the permeability of the gonococcal cell envelope by the *mtr* system. Mol Microbiol 11:769–775.

7.   Olesky M, Hobbs M, Nicholas RA. 2002. Identification and Analysis of Amino Acid Mutations in Porin IB That Mediate Intermediate-Level Resistance to Penicillin and Tetracycline in *Neisseria gonorrhoeae*. Antimicrob Agents Chemother 46:2811–2820.

8.   Ropp PA, Hu M, Olesky M, Nicholas RA. 2002. Mutations in *ponA*, the Gene Encoding Penicillin-Binding Protein 1, and a Novel Locus, *penC*, Are Required for High-Level Chromosomally Mediated Penicillin Resistance in *Neisseria gonorrhoeae*. Antimicrob Agents Chemother 46:769–777.

9.   Veal WL, Nicholas RA, Shafer WM. 2002. Overexpression of the MtrC-MtrD-MtrE Efflux Pump Due to an *mtrR* Mutation Is Required for Chromosomally Mediated Penicillin Resistance in *Neisseria gonorrhoeae*. J Bacteriol 184:5619–5624.

10.   Cannon JG, Sparling PF. 1984. THE GENETICS OF THE GONOCOCCUS. Annu Rev Microbiol 38:111–133.

11.   Elsener TA, Cehovin A, Philp C, Fortney K, Spinola SM, Maiden MCJ, Tang CM. 2025. Origin, evolution, and success of pbla, the gonococcal beta-lactamase plasmid, and implications for public health. PLoS Pathog 21:e1013151.

12.   Dougherty TJ. 1986. Genetic analysis and penicillin-binding protein alterations in *Neisseria gonorrhoeae* with chromosomally mediated resistance. Antimicrob Agents Chemother 30:649–652.

13. Faruki H, Sparling PF. 1986. Genetics of resistance in a non-beta-lactamase-
producing gonococcus with relatively high-level penicillin resistance. Antimicrob
Agents Chemother 30:856–860.

14. Sparling PF, Sarubbi FA, Blackman E. 1975. Inheritance of low-level resistance to
penicillin, tetracycline, and chloramphenicol in *Neisseria gonorrhoeae*. J Bacteriol
124:740–749.

15. Powell AJ, Tomberg J, Deacon AM, Nicholas RA, Davies C. 2009. Crystal
Structures of Penicillin-binding Protein 2 from Penicillin-susceptible and -resistant
Strains of *Neisseria gonorrhoeae* Reveal an Unexpectedly Subtle Mechanism for
Antibiotic Resistance. J Biol Chem 284:1202–1212.

16. Spratt BG, Balthazar JT, Hagman KE, Pan W, Judd RC, Shafer WM. 1995.
Resistance of *Neisseria gonorrhoeae* to antimicrobial hydrophobic agents is
modulated by the *mtrRCDE* efflux system. Microbiology 141:611–622.

17. Gill MJ, Simjee S, Al-Hattawi K, Robertson BD, Easmon CSF, Ison CA. 1998.
Gonococcal Resistance to β-Lactams and Tetracycline Involves Mutation in Loop 3
of the Porin Encoded at the *penB* Locus. Antimicrob Agents Chemother 42:2799–
2803.

18. Zhao S, Tobiason DM, Hu M, Seifert HS, Nicholas RA. 2005. The *penC* mutation
conferring antibiotic resistance in *Neisseria gonorrhoeae* arises from a mutation in
the PilQ secretin that interferes with multimer stability. Mol Microbiol 57:1238–
1251.

19. Nandi S, Swanson S, Tomberg J, Nicholas RA. 2015. Diffusion of Antibiotics
through the PilQ Secretin in *Neisseria gonorrhoeae* Occurs through the Immature,
Sodium Dodecyl Sulfate-Labile Form. J Bacteriol 197:1308–1321.

20. Swanson J. 1973. Studies on gonococcus infection. IV. Pili: their role in attachment
of gonococci to tissue culture cells. J Exp Med 137:571–589.

21. Hamrick TS, Dempsey JAF, Cohen MS, Cannon JG. 2001. Antigenic variation of
gonococcal pilin expression *in vivo*: analysis of the strain FA1090 pilin repertoire
and identification of the *pilS* gene copies recombining with *pilE* during experimental
human infection. Microbiology (Reading) 147:839–849.

22. Zhao S, Duncan M, Tomberg J, Davies C, Unemo M, Nicholas RA. 2009. Genetics
of Chromosomally Mediated Intermediate Resistance to Ceftriaxone and Cefixime
in *Neisseria gonorrhoeae*. Antimicrob Agents Chemother 53:3744–3751.

23. Bowler LD, Zhang QY, Riou JY, Spratt BG. 1994. Interspecies recombination
between the *penA* genes of *Neisseria meningitidis* and commensal *Neisseria*
species during the emergence of penicillin resistance in *N. meningitidis*: natural
events and laboratory simulation. J Bacteriol 176:333–337.

24. Takahata S, Senju N, Osaki Y, Yoshida T, Ida T. 2006. Amino Acid Substitutions in
Mosaic Penicillin-Binding Protein 2 Associated with Reduced Susceptibility to
Cefixime in Clinical Isolates of *Neisseria gonorrhoeae*. Antimicrob Agents
Chemother 50:3638–3645.

25. Spratt BrianG, Bowler LucasD, Zhang Q-Y, Zhou J, Smith J. 1992. Role of
interspecies transfer of chromosomal genes in the evolution of penicillin resistance
in pathogenic and commensal *Neisseria* species. J Mol Evol 34.

26. Dowson CG, Jephcott AE, Gough KR, Spratt BG. 1989. Penicillin-binding protein 2
genes of non-β-lactamase-producing, penicillin-resistant strains of *Neisseria*
gonorrhoeae. Mol Microbiol 3:35–41.

27. Fedarovich A, Cook E, Tomberg J, Nicholas RA, Davies C. 2014. Structural Effect
of the Asp345a Insertion in Penicillin-Binding Protein 2 from Penicillin-Resistant
Strains of *Neisseria gonorrhoeae*. Biochemistry 53:7596–7603.

28. Folster JP, Dhulipala V, Nicholas RA, Shafer WM. 2007. Differential Regulation of
ponA and *pilMNOPQ* Expression by the MtrR Transcriptional Regulatory Protein in
Neisseria gonorrhoeae. J Bacteriol 189:4569–4577.

29. Helekal D, Mortimer TD, Mukherjee A, Gentile G, Le Van A, Nicholas RA, Jerse
AE, Palace SG, Grad YH. 2025. Quantifying the impact of genetic determinants of
antibiotic resistance on bacterial lineage dynamics. bioRxiv
https://doi.org/10.1101/2025.02.03.636319.
686
30. Tomberg J, Unemo M, Davies C, Nicholas RA. 2010. Molecular and Structural
Analysis of Mosaic Variants of Penicillin-Binding Protein 2 Conferring Decreased
Susceptibility to Expanded-Spectrum Cephalosporins in *Neisseria gonorrhoeae*:
Role of Epistatic Mutations. Biochemistry 49:8062–8070.

31. Tomberg J, Unemo M, Ohnishi M, Davies C, Nicholas RA. 2013. Identification of
Amino Acids Conferring High-Level Resistance to Expanded-Spectrum
Cephalosporins in the *penA* Gene from *Neisseria gonorrhoeae* Strain H041.
Antimicrob Agents Chemother 57:3029–3036.

32. Ohnishi M, Golparian D, Shimuta K, Saika T, Hoshina S, Iwasaku K, Nakayama S,
Kitawaki J, Unemo M. 2011. Is *Neisseria gonorrhoeae* Initiating a Future Era of
Untreatable Gonorrhea?: Detailed Characterization of the First Strain with High-
Level Resistance to Ceftriaxone. Antimicrob Agents Chemother 55:3538–3545.

33. Tomberg J, Unemo M, Davies C, Nicholas RA. 2010. Molecular and Structural
Analysis of Mosaic Variants of Penicillin-Binding Protein 2 Conferring Decreased
Susceptibility to Expanded-Spectrum Cephalosporins in *Neisseria gonorrhoeae*:
Role of Epistatic Mutations. Biochemistry 49:8062–8070.

34. Tomberg J, Fedarovich A, Vincent LR, Jerse AE, Unemo M, Davies C, Nicholas
RA. 2017. Alanine 501 Mutations in Penicillin-Binding Protein 2 from *Neisseria*

gonorrhoeae: Structure, Mechanism, and Effects on Cephalosporin Resistance and
Biological Fitness. Biochemistry 56:1140–1150.

35. Johnson SR, Grad Y, Ganakammal SR, Burroughs M, Frace M, Lipsitch M, Weil R,
    Trees D. 2014. *In Vitro* Selection of *Neisseria gonorrhoeae* Mutants with Elevated
    MIC Values and Increased Resistance to Cephalosporins. Antimicrob Agents
    Chemother 58:6986–6989.

36. Drake SL, Koomey M. 1995. The product of the *pilQ* gene is essential for the
    biogenesis of type IV pili in *Neisseria gonorrhoeae*. Mol Microbiol 18:975–986.

37. Whiley DM, Jacobsson S, Tapsall JW, Nissen MD, Sloots TP, Unemo M. 2010.
    Alterations of the pilQ gene in *Neisseria gonorrhoeae* are unlikely contributors to
    decreased susceptibility to ceftriaxone and cefixime in clinical gonococcal strains. J
    Antimicrob Chemother 65:2543–2547.

38. Kellogg DS, Peacock WL, Deacon WE, Brown L, Pirkle CI. 1963. *NEISSERIA
    GONORRHOEAE* I: Virulence Genetically Linked to Clonal Variation. J Bacteriol
    85:1274–1279.

39. Adamson PC, Van Le H, Le HHL, Le GM, Nguyen TV, Klausner JD. 2020. Trends
    in antimicrobial resistance in *Neisseria gonorrhoeae* in Hanoi, Vietnam, 2017–
    2019. BMC Infect Dis 20:809.

40. Zhao S, Duncan M, Tomberg J, Davies C, Unemo M, Nicholas RA. 2009. Genetics
    of chromosomally mediated intermediate resistance to ceftriaxone and cefixime in
    Neisseria gonorrhoeae. Antimicrob Agents Chemother 53:3744–3751.

41. Dillard JP, Chan JM. 2024. Genetic Manipulation of *Neisseria gonorrhoeae* and
    Commensal *Neisseria* Species. Curr Protoc 4:e70000.

42. Prentki P, Krisch HM. 1984. *In vitro* insertional mutagenesis with a selectable DNA
    fragment. Gene 29:303–313.

43. Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. 1989. Site-directed
    mutagenesis by overlap extension using the polymerase chain reaction. Gene
    77:51–59.

44. Kunz AN, Begum AA, Wu H, D'Ambrozio JA, Robinson JM, Shafer WM, Bash MC,
    Jerse AE. 2012. Impact of Fluoroquinolone Resistance Mutations on Gonococcal
    Fitness and *In Vivo* Selection for Compensatory Mutations. J Infect Dis 205:1821–
    1829.

45. Jolley KA, Bray JE, Maiden MCJ. 2018. Open-access bacterial population
    genomics: BIGSdb software, the PubMLST.org website and their applications.
    Welcome Open Res 3:124.

46. Baarda BI, Zielke RA, Nicholas RA, Sikora AE. 2018. PubMLST for Antigen Allele
Mining to Inform Development of Gonorrhea Protein-Based Vaccines. Front
Microbiol 9:2971.

47. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,
Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl
SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T,
Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M,
Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K,
Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
AlphaFold. Nature 596:583–589.

48. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D,
Stroe O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S,
Jumper J, Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N,
Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar S. 2022. AlphaFold Protein
Structure Database: massively expanding the structural coverage of protein-
sequence space with high-accuracy models. Nucleic Acids Res 50:D439–D444.

49. Varadi M, Bertoni D, Magana P, Paramval U, Pidruchna I, Radhakrishnan M,
Tsenkov M, Nair S, Mirdita M, Yeo J, Kovalevskiy O, Tunyasuvunakool K, Laydon
A, Žídek A, Tomlinson H, Hariharan D, Abrahamson J, Green T, Jumper J, Birney
E, Steinegger M, Hassabis D, Velankar S. 2024. AlphaFold Protein Structure
Database in 2024: providing structure coverage for over 214 million protein
sequences. Nucleic Acids Res 52:D368–D375.

50.   Schrodinger, LLC. 2010. The PyMOL Molecular Graphics System, Version X.X.

51. Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics
Analysis Version 7.0 for Bigger Datasets. Mol Biol and Evol 33:1870–1874.

52. Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced
time and space complexity. BMC Bioinformatics 5:113.

53. Crooks GE, Hon G, Chandonia J-M, Brenner SE. 2004. WebLogo: A Sequence
Logo Generator: Figure 1. Genome Res 14:1188–1190.

54. Alfsnes K, Eldholm V, Olsen AO, Brynildsrud OB, Bohlin J, Steinbakk M, Caugant
DA. 2020. Genomic epidemiology and population structure of *Neisseria
gonorrhoeae* in Norway, 2016–2017. Microbial Genomics
https://doi.org/10.1099/mgen.0.000359.

55. Bristow CC, Mortimer TD, Morris S, Grad Y, Soge OO, Wakatake E, Pascual R,
Murphy SM, Fryling KE, Adamson PC, Dillon J-A, Palmer NR, Le HHL, Van Le H,
Ovalles Ureña RM, Mitchev N, Mlisana K, Wi T, Dickson SP, Klausner JD. 2023.
Whole Genome Sequencing to Predict Antimicrobial Susceptibility Profiles in
Neisseria gonorrhoeae. J Infect Dis 227:917-925.

56. Demczuk W, Lynch T, Martin I, Domselaar GV, Graham M, Bharat A, Allen V,
Hoang L, Lefebvre B, Tyrrell G, Horsman G, Haldane D, Garceau R, Wylie J, Wong
T, Mulvey MR. 2015. Whole-Genome Phylogenomic Heterogeneity of *Neisseria*
gonorrhoeae Isolates with Decreased Cephalosporin Susceptibility Collected in
Canada between 1989 and 2013. J Clin Microbiol 53:191–200.

57. Eyre DW, De Silva D, Cole K, Peters J, Cole MJ, Grad YH, Demczuk W, Martin I,
Mulvey MR, Crook DW, Walker AS, Peto TEA, Paul J. 2017. WGS to predict
antibiotic MICs for *Neisseria gonorrhoeae*. J Antimicrob Chemother 72:1937–1947.

58. Golparian D, Harris SR, Sánchez-Busó L, Hoffmann S, Shafer WM, Bentley SD,
Jensen JS, Unemo M. 2020. Genomic evolution of *Neisseria gonorrhoeae* since
the preantibiotic era (1928–2013): antimicrobial use/misuse selects for resistance
and drives evolution. BMC Genomics 21:116.

59. Golparian D, Bazzo ML, Golfetto L, Gaspar PC, Schörner MA, Schwartz Benzaken
A, Ramos MC, Ferreira WA, Alonso Neto JB, Mendes Pereira GF, Unemo M, the
Brazilian-GASP Network. 2020. Genomic epidemiology of *Neisseria gonorrhoeae*
elucidating the gonococcal antimicrobial resistance and lineages/sublineages
across Brazil, 2015–16. J Antimicrob Chemother 75:3163–3172.

60. Golparian D, Kittiyaowamarn R, Paopang P, Sangprasert P, Sirivongrangson P,
Franceschi F, Jacobsson S, Wi T, Unemo M. 2022. Genomic surveillance and
antimicrobial resistance in *Neisseria gonorrhoeae* isolates in Bangkok, Thailand in
2018. J Antimicrob Chemother 77:2171–2182.

61. Grad YH, Kirkcaldy RD, Trees D, Dordel J, Harris SR, Goldstein E, Weinstock H,
Parkhill J, Hanage WP, Bentley S, Lipsitch M. 2014. Genomic epidemiology of
Neisseria gonorrhoeae with reduced susceptibility to cefixime in the USA: a
retrospective observational study. Lancet Infect Dis 14:220–226.

62. Grad YH, Harris SR, Kirkcaldy RD, Green AG, Marks DS, Bentley SD, Trees D,
Lipsitch M. 2016. Genomic Epidemiology of Gonococcal Resistance to Extended-
Spectrum Cephalosporins, Macrolides, and Fluoroquinolones in the United States,
2000–2013. J Infect Dis 214:1579–1587.

63. Kwong JC, Chow EPF, Stevens K, Stinear TP, Seemann T, Fairley CK, Chen MY,
Howden BP. 2018. Whole-genome sequencing reveals transmission of gonococcal
antibiotic resistance among men who have sex with men: an observational study.
Sex Transm Infect 94:151–157.

64. Lan PT, Golparian D, Ringlander J, Van Hung L, Van Thuong N, Unemo M. 2020.
Genomic analysis and antimicrobial resistance of *Neisseria gonorrhoeae* isolates
from Vietnam in 2011 and 2015-16. J Antimicrob Chemother 75:1432–1438.

65. Lee RS, Seemann T, Heffernan H, Kwong JC, Gonçalves da Silva A, Carter GP,
Woodhouse R, Dyet KH, Bulach DM, Stinear TP, Howden BP, Williamson DA.

2018. Genomic epidemiology and antimicrobial resistance of *Neisseria*
gonorrhoeae in New Zealand. J Antimicrob Chemother 73:353–364.

66. Pinto M, Borges V, Isidro J, Rodrigues JC, Vieira L, Borrego MJ, Gomes JP. 2020.
Neisseria gonorrhoeae clustering to reveal major European whole-genome-
sequencing-based genogroups in association with antimicrobial resistance. Microb
Genom 7:000481.

67. Reimche JL, Chivukula VL, Schmerer MW, Joseph SJ, Pham CD, Schlanger K, St
Cyr SB, Weinstock HS, Raphael BH, Kersh EN, Gernert KM, Antimicrobial
Resistant *Neisseria gonorrhoeae* Working Group. 2021. Genomic analysis of the
predominant strains and antimicrobial resistance determinants within 1479
Neisseria gonorrhoeae isolates from the U.S. Gonococcal Isolate Surveillance
Project in 2018. Sex Transm Dis https://doi.org/10.1097/OLQ.0000000000001471.

68. Town K, Harris S, Sánchez-Busó L, Cole MJ, Pitt R, Fifer H, Mohammed H, Field
N, Hughes G. 2020. Genomic and Phenotypic Variability in *Neisseria gonorrhoeae*
Antimicrobial Susceptibility, England. Emerg Infect Dis 26:505–515.

69. Williamson DA, Chow EPF, Gorrie CL, Seemann T, Ingle DJ, Higgins N, Easton M,
Taiaroa G, Grad YH, Kwong JC, Fairley CK, Chen MY, Howden BP. 2019. Bridging
of *Neisseria gonorrhoeae* lineages across sexual networks in the HIV pre-exposure
prophylaxis era. Nat Commun 10:1–10.

70. Reimche JL, Clemons AA, Chivukula VL, Joseph SJ, Schmerer MW, Pham CD,
Schlanger K, St Cyr SB, Kersh EN, Gernert KM, †Antimicrobial- Resistant Working
Group. 2023. Genomic analysis of 1710 surveillance-based Neisseria *gonorrhoeae*
isolates from the USA in 2019 identifies predominant strain types and
chromosomal antimicrobial-resistance determinants. Microbial Genomics
9:001006.

71. Salmerón P, Moreno-Mingorance A, Trejo J, Amado R, Viñado B, Cornejo-Sanchez
T, Alberny M, Barbera MJ, Arando M, Pumarola T, Hoyos-Mallecot Y, Serra-
Pladevall J, González-López JJ. 2021. Emergence and dissemination of three mild
outbreaks of Neisseria gonorrhoeae with high-level resistance to azithromycin in
Barcelona, 2016–18. J Antimicrob Chemother 76:930–935.

72. Sánchez-Busó L, Golparian D, Corander J, Grad YH, Ohnishi M, Flemming R,
Parkhill J, Bentley SD, Unemo M, Harris SR. 2019. The impact of antimicrobials on
gonococcal evolution. Nat Microbiol 4:1941–1950.

73. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA,
Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: An Integrated Tool for
Comprehensive Microbial Variant Detection and Genome Assembly Improvement.
PLOS ONE 9:e112963.

74. Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. ArXiv:1303.3997 [q-bio.GN].

75. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden
856   TL. 2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421.

857   76. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
858   Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,
859   Alekseyev MA, Pevzner PA. 2012. SPAdes: A New Genome Assembly Algorithm
860   and Its Applications to Single-Cell Sequencing. J Comput Biol 19:455–477.

861   77. Demczuk W, Sidhu S, Unemo M, Whiley DM, Allen VG, Dillon JR, Cole M, Seah C,
862   Trembizki E, Trees DL, Kersh EN, Abrams AJ, Vries HJC de, Dam AP van, Medina
863   I, Bharat A, Mulvey MR, Domselaar GV, Martin I. 2017. *Neisseria gonorrhoeae*
864   Sequence Typing for Antimicrobial Resistance, a Novel Antimicrobial Resistance
865   Multilocus Typing Scheme for Tracking Global Dissemination of *N. gonorrhoeae*
866   Strains. J Clin Microbiol 55:1454–1468.

867   78. Kille B, Nute MG, Huang V, Kim E, Phillippy AM, Treangen TJ. 2024. Parsnp 2.0:
868   Scalable Core-Genome Alignment for Massive Microbial Datasets. bioRxiv
869   https://doi.org/10.1101/2024.01.30.577458.

870   79. Menardo F, Loiseau C, Brites D, Coscolla M, Gygli SM, Rutaihwa LK, Trauner A,
871   Beisel C, Borrell S, Gagneux S. 2018. Treemmer: a tool to reduce large
872   phylogenetic datasets with minimal loss of diversity. BMC Bioinformatics 19:164.

873   80. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J,
874   Harris SR. 2015. Rapid phylogenetic analysis of large samples of recombinant
875   bacterial whole genome sequences using Gubbins. Nucl Acids Res 43:e15–e15.

876   81. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
877   developments. Nucl Acids Res https://doi.org/10.1093/nar/gkz239.

878   82. Abrams AJ, Trees DL, Nicholas RA. 2015. Complete Genome Sequences of Three
879   *Neisseria gonorrhoeae* Laboratory Reference Strains, Determined Using PacBio
880   Single-Molecule Real-Time Technology. Genome Announc 3:e01052-15.

881   83. Clinical and Laboratory Standard Institute (CLSI). 2020. Performance Standards for
882   Anti-Microbial Susceptibility Testing. 30th Edition, M100.
883   https://clsi.org/media/3481/m100ed30_sample.pdf.

Table 1 FA6140 *ponA*[L421X] Isogenic Mutant Library and penicillin MICs.

Strains Name[a]	Amino Acid at Position 421 in *ponA*	Codon	Penicillin MIC (μg/mL)[b]
FA6140[c]	Proline (P)	CCG	4
FA6140 *ponA*[d]	Leucine (L)	CTG	2
FA6140 *ponA*[L421P]	Proline (P)	CCG	4
FA6140 *ponA*[L421L]	Leucine (L)	CTG	2
FA6140 *ponA*[L421A]	Alanine (A)	GCC	2
FA6140 *ponA*[L421R]	Arginine (R)	AGA	2
FA6140 *ponA*[L421C]	Cysteine (C)	TGC	2
FA6140 *ponA*[L421N]	Asparagine (N)	AAC	2
FA6140 *ponA*[L421E]	Glutamic Acid (E)	GAA	2
FA6140 *ponA*[L421G]	Glycine (G)	GGT	2
FA6140 *ponA*[L421H]	Histidine (H)	CAT	2
FA6140 *ponA*[L421Q]	Glutamine (Q)	CAG	2
FA6140 *ponA*[L421W]	Tryptophan (W)	TGG	2
FA6140 *ponA*[L421Y]	Tyrosine (Y)	TAT	2
FA6140 *ponA*[L421S]	Serine (S)	AGT	2.5
FA6140 *ponA*[L421F]	Phenylalanine (F)	TTC	2
FA6140 *ponA*[L421I]	Isoleucine (I)	ATC	2
FA6140 *ponA*[L421T]	Threonine (T)	ACT	2

[a]All strains are created in the FA6140 background, a penicillin-resistant gonococcal isolate, as described in Methods and Materials. Each strain possesses a different amino acid at position 421 in *ponA*.

[b]The MICs reported are the median values determined from a minimum of three independent experiments.

[c]Penicillin-resistant gonococcal isolate, FA6140, which harbors *ponA*[L421P].

[d]FA6140 with *ponA*[L421P] replaced with WT *ponA*.

A). *N. gonorrhoeae*

B). *N. meningitidis*

C). *Neisseria spp.*

FIG 1 Sequence logo diagrams comparing the conservation of positions 415-430 in PBP1 from various *Neisseria* isolates. Diagram generated on WebLogo with MUSCLE-aligned *ponA* sequences. Sequences were downloaded from PubMLST and are a representative sub-set of all of the *ponA* alleles documented for each *Neisseria* species group: *N. gonorrhoeae* (n=172), *N. meningitidis* (n=897), and *Neisseria spp.* (n=369). The X-axis indicates the position of the amino acid. The height of the logo element (letter) represents its log-transformed frequency displayed in bits of information, and the overall height of the stack indicates the sequence conservation at that position.

FIG 2 Historical analysis of *N. gonorrhoeae* isolates (n=16,891). Isolates are represented by the year the isolate was collected. Asp345a refers to the insertion of asparagine at position 345 in *penA* (encodes PBP2), a known penicillin-resistance-conferring mutation. Count represents the number of isolates that harbor *ponA* (blue) or *ponA*^{L421P} (red).

FIG 3 Phylogenetic distribution of *ponA* alleles in a representative sample of *N. gonorrhoeae* isolates without plasmid-mediated resistance (n=5,192). The outer annotation ring indicates the presence of *ponA* (L421, grey) and *ponA*[L421P] (L421P, black). The inner ring indicates the level of penicillin resistance. S is susceptible (≤0.06 µg/mL), I is intermediate (0.25 - 1 µg/mL), and R is resistant (≥2 µg/mL) breakpoints of PenG as defined by the Clinical and Laboratory Standards Institute (CLSI) (83).

FIG 4 Distribution of MIC values for penicillin G (PenG) among *N. gonorrhoeae* isolates (n = 8,999). Strains are separated by harboring either (A) a non-mosaic *penA* allele, or (B) a mosaic *penA* allele. The X-axis shows the PenG MIC (MIC_{PEN}), and dotted lines indicate the susceptible (≤0.06 µg/mL), intermediate (0.25 - 1 µg/mL), and resistant (≥2 µg/mL) breakpoints of PenG as defined by CLSI (83). The Y-axis shows the number of isolated *N. gonorrhoeae* strains for each MIC_{PEN}, value. Red bars show distribution of strains with the *ponA*[L421P] allele and blue bars shows the number of strains with the wild-type *ponA* (L421) allele. The mosaic *penA* alleles in isolates were defined and typed according to the NG-STAR database (56).

FIG 5 Polymorphisms of *N. gonorrhoeae* PBP1 mapped onto the predicted protein structure. Polymorphisms are colored according to their prevalence in the gonococcal population. Only polymorphisms detected at frequencies greater than 1% are shown: L421P (47.15%), A375T (7.07%), and Y537C (2.35%). The Transpeptidase (TPase), Transglycosylase (TGase), and the Oligosaccharide-Binding (OB) domains of PBP1 are labeled, with the TPase active site residues marked in red. MUSCLE-aligned sequences of gonococcal PBP1 were taken from PubMLST isolate database in order to identify polymorphisms and calculate population frequencies, see Materials and Methods. Predicted structure of gonococcal PBP1 is from the AlphaFold Protein Structure Database (AFDB).

FIG 6 Optical densities (OD_{600}) and Illumina read counts from *in vitro* serial passaging of the FA6140 *ponA*[L421X] mutant library in the absence or presence of sub-MIC concentrations of penicillin. (A) Percentage of reads, relative to the total number of reads, for each mutant at hour 0 (inoculum). The inoculum was used to inoculate each culture at 0.08 OD_{600}. (B) The serial passaging scheme included a total of 4 cultures, 3 of which were supplemented with different sub-MIC concentrations of penicillin (0.25, 0.5, and 1.0 µg/mL), and one with no penicillin added. Cultures were passaged every 6 hours, for a total of 36 hours, always with a starting OD_{600} of 0.08. (C) OD_{600} and (D) Illumina read counts of the cultures with no penicillin, compared to cultures supplemented with 0.5 µg/mL penicillin. (D) Reads for each mutant in the inoculum were normalized to 1, as indicated by the dotted line at y=1. Bars represent read counts for each mutant at hour 36, relative to their inoculum value. This experiment was performed 3 separate times

(trials) and each trial represents a separate Illumina sequencing run; (C-D) Trial 1 is shown in red, trial 2 in yellow, and trial 3 in blue. The pie chart in (A) is a representative figure from trial 1, but all 3 trials had similar results. See supplemental for read counts from the cultures supplemented with 0.25 µg/mL of penicillin (Fig. S4) and for the inoculum used in trials 2 and 3 (Fig. S3).

FIG 7 A *pilQ* mutation arose in the culture supplemented with 1.0 ug/ml of penicillin. Optical densities (A) and Illumina read counts (B) for the FA6140 *ponA*[L421X] strains grown in cultures supplemented with 1.0 µg/mL of penicillin. (B) Read counts for each mutant at hour 36, relative to their inoculum (Hr 0) value. Inoculum reads for each mutant were normalized to 1, as indicated by the dotted line at y=1. (C) Read counts over time for FA6140 *ponA*[L421S] (orange) FA6140 *ponA*[L421P] (black), relative to their inoculum value (y=1), during trials 1 and 3. Each trial represents a separate Illumina sequencing run (A-B, trial 1 = red, trial 2 = yellow, trial 3 = blue). Cells growing in the culture from trial 2 appeared to die at hour 18 (A; yellow line), resulting in poor sequencing reads. See supplemental Fig. S5 for trial 2 data.

FIG 8 MICs of penicillin G and ceftriaxone for FA6140 transformants. The MICs reported above each bar are the median values (± the standard deviation) determined from three independent experiments. Data points within each bar represent the MIC values measured during each experiment.

FIG 9. Diagram outlining the drug targets for penicillin and ceftriaxone in gonococcal strains with different *penA* and *ponA* alleles. PBP1 variants are marked by green rectangles and PBP2 variants are marked as purple rectangles. The left axis indicates the acylation rate, and the right axis indicates the MIC of the indicated strain. Arrows point to the killing target for each of the indicated strains. A). Penicillin has a low k_2/K_S for all PBP2 variants, so it kills gonococcal cells by inhibiting PBP1. Replacing PBP1^{L421P} with PBP1^{WT} decreases the MIC. B). Ceftriaxone normally kills by inhibiting PBP2, but PBP2^{H041} has a much lower k_2/K_S and this shifts the killing target to PBP1 (and maybe some PBP2^{H041}). Replacing PBP1^{L421P} with PBP1^{WT} decreases the MIC.