# Large sharp-wave ripples promote hippocampo-cortical memory reactivation and consolidation

Heath L. Robinson[1,3], Ralitsa Todorova[1,3], Gergo A. Nagy[2], Anna Gruzdeva[1], Praveen Paudel[1], Azahara Oliva[1], Antonio Fernandez-Ruiz[1,4]

Affiliations:

[1]Department of Neurobiology and Behavior, Cornell University, Ithaca, NY, USA.

[2]Institute of Experimental Medicine – Hungarian Research Network, Budapest, Hungary

[3]These authors contributed equally to this work.

[4]Lead Contact

*Correspondence to: afr77@cornell.edu

## Summary

During sleep, ensemble activity patterns encoding recent experiences are reactivated in the hippocampus and cortex. This reactivation is coordinated by hippocampal sharp-wave ripples (SWRs) and is believed to support the early stages of memory consolidation. However, only a minority of sleep SWRs are associated with memory reactivation in the hippocampus and its downstream areas. Whether that subset of SWRs have specific physiological characteristics and directly contribute to memory performance is not known. We identified a specific subset of large SWRs linked to memory reactivation in both the hippocampus and prefrontal cortex (PFC) of mice, and found that their occurrence selectively increased during sleep following new learning. Closed-loop optogenetic SWR boosting during sleep was sufficient to enhance ensemble memory reactivation in hippocampus and PFC. This manipulation also improved subsequent memory retrieval and hippocampal-PFC coordination, causally linking both phenomena to SWR-associated ensemble reactivation during sleep.

## Keywords

Hippocampus; Memory; Learning; Sharp-wave ripples; Reactivation; Replay; Cell assemblies; Synchrony; Network dynamics; Oscillations

## Introduction

New memories are consolidated during sleep following learning, a process that leads to the formation of stable, long-term representations [1,2]. The hippocampus plays a key role in the initial memory encoding [3]. According to the influential systems memory consolidation theory, memories are gradually transferred from the hippocampus to the neocortex for permanent storage [4–6]. This process has been hypothesized to be mediated by synchronous network patterns generated in the hippocampus, known as sharp-wave ripples (SWRs) [7–10]. SWRs occur during non-rapid eye movement (NREM) sleep and waking immobility periods. They coordinate the reactivation of hippocampal neurons that encoded recent experiences and broadcast these memory representations to the neocortex and other brain areas [10–14]. Numerous studies including both correlational analyses as well as causal manipulations, have established the role of hippocampal SWRs in memory formation and consolidation [15–25]. In downstream structures such as the prefrontal cortex (PFC), SWRs entrain local circuits driving the coordinated reactivation of experience-related activity patterns [26–30]. Furthermore, the reactivation of neural population activity in both the hippocampus and the prefrontal cortex has been linked to novel experiences and memory demands [26,31–37].

Despite this wealth of evidence, an apparent paradox is that only a small subset of sleep SWRs are linked to the reactivation of recent experiences. Most studies, using a variety of different methods, converge to only ~10 to 30 % of SWRs significantly associated with significant memory reactivation or "replay" [12,15,20,25,34,36,38]. This fraction could be even smaller for memory reactivation in downstream structures involved in memory consolidation, such as the PFC. These observations raise the question of whether the subset of SWRs that successfully elicit memory reactivation in the neocortex, and thus presumably contribute to memory consolidation, have unique physiological characteristics or whether their impact is determined by some additional variables that converge with SWRs. In this study, we aimed to distinguish between these two hypotheses and establish a direct link between SWR features and memory performance. Furthermore, studies in humans have shown that increased functional coordination between hippocampus and cortex during rest following new learning correlates with successful memory recall [39,40]. However, whether sleep SWRs mediate a persistent enhancement of hippocampo-cortical coordination and subsequent memory recall is not known.

To answer these questions, we performed neural ensemble recordings in the hippocampus and PFC of mice during a memory task and sleep, together with closed-loop optogenetic manipulation of SWRs. We found that only a specific subset of large SWRs were associated with memory reactivation in both structures. The rate of these large SWRs selectively increased in sleep after learning and correlated with successful memory performance. To directly test their contribution to memory, we optogenetically boosted sleep SWRs following a learning paradigm otherwise insufficient for mice to consolidate a new experience. SWR boosting enhanced memory reactivation in both the hippocampus and PFC and, remarkably, allowed mice to successfully recall the new experience after sleep. Oscillatory synchrony between hippocampus and PFC during memory recall was also enhanced, linking SWR-mediated sleep memory reactivation to persistent functional connectivity changes, similarly to reports from human memory studies.

## Results

**Large SWRs are associated with stronger memory reactivation in CA1 and PFC and modulated by learning.**

We implanted mice with high-density silicon probes in the dorsal hippocampus (CA1 region) and PFC (pre-limbic region) (Fig. 1A and Fig. S1). We simultaneously recorded large ensembles of cells in both regions (n = 1827 CA1 and 1103 PFC neurons from 21 mice) while mice performed a hippocampal-dependent memory task (object displacement, see Methods) and in sleep before and after the task. To analyze memory reactivation, we employed an unsupervised method to detect the synchronous coactivity of groups of neurons, or cell assemblies, during learning, as in previous work [23,24,41,42]. We then compared whether these task-related cell assemblies in PFC and CA1 reactivated during SWRs in subsequent sleep. Using this approach, we found that 35.4% of SWRs were associated with significant memory reactivation in the hippocampus and 19.7% in PFC (P < 0.05, n = 353.600 SWRs from 21 mice). The probability of PFC reactivation was higher for SWRs associated with significant CA1 reactivation (Fig. S2A).

We asked whether SWRs that elicited ensemble reactivation in downstream areas, such as the PFC, and could therefore promote memory consolidation, have specific defining features. To gain intuition about which features may be more relevant, we first represented all SWRs as points in a high-dimensional space constructed using different features (see Methods) and embedded them in a low-dimensional space using Uniform Manifold Approximation and Projection (UMAP) for visualization. This method revealed that SWR properties such as amplitude, duration or frequency varied along a continuum (Fig. 1B and Fig. S2) [36,43]. Color-coding SWRs based on the reactivation strength of concurrent PFC ensemble activity suggested a correlation between SWR amplitude and duration with PFC reactivation strength (Fig. 1B). Classifying all sleep SWRs in two binary categories, those with and without significant associated memory reactivation in PFC, confirmed that the former had significantly larger duration and amplitude (Fig.1C-E), but did not significantly differ in other features (Fig. S2).

To examine in more detail the relationship between memory reactivation strength and SWR duration and amplitude, we classified SWRs based on these features and calculated peri-SWR histograms of PFC assembly reactivation strength for each of the categories. Only the topmost duration and amplitude SWR quantiles were associated with strong PFC memory reactivation (Fig. 1F-G). We obtained similar results when we performed the same analysis with CA1 assembly reactivation strength (Fig. 1H-I). However, in the case of CA1, reactivation strength increased more gradually with increasing SWR duration and amplitude in contrast to a more abrupt increase in PFC.

SWR duration and amplitude were only moderately correlated (Fig. S2), and our initial observations suggested that only a subset of SWRs with both large amplitude and duration were associated with strong memory reactivation (Fig. 1B). Therefore, we classified SWR into groups of increasing joint amplitude and duration. Memory reactivation strength in both the PFC (Fig. 1J) and CA1 (Fig. 1K) was largest for SWRs in the top duration and amplitude quantile, but weak for SWRs with large amplitude but short duration or long duration but low amplitude.

**Figure 1: Enhanced memory reactivation in hippocampus and cortex during a subset of SWRs**

**A)** Schematic of recording design. **B)** Visualization of SWR features using a 3D UMAP embedding. Each point is a single SWR from an example session, color coded according to different features. **C)** Examples of SWR without (left) and with (right) associated PFC cell assembly reactivation. Top: CA1 LFP traces. Middle: raster plots of neuronal firing, a colored circle indicates assembly activity accompanying the hippocampal SWR; Bottom: assembly reactivation strength. **D)** Average filtered CA1 LFP trace for SWRs associated with (red trace) or without (black trace) significant reactivation of PFC cell assemblies. Data is from the same session as B-C. **E)** SWRs associated with significant reactivation of PFC cell assemblies were longer (P = 0.043, n = 69,522 SWRs, hierarchical bootstrap test) and had higher amplitude (P = 0.025, n = 284,078) than the remaining SWRs. Black dots: mean; black lines: 95% confidence intervals; curves: the resampled mean distributions. **F)** Left: reactivation strength of PFC assemblies around SWRs sorted by duration. Right: PFC assembly reactivation during SWRs classified by duration quantiles (P = 0.25 / 0.20 / 0.0104 for short, mid-length and long ripples respectively, hierarchical bootstrap test). **G)** Same as F but with SWRs sorted by amplitude (P= 0.55 / 0.14 / 0.013 for low, mid and high-amplitude SWRs respectively). **H)** Same as F but for CA1 assemblies (P=1 / 0.65 / 0 for short, mid-length and long SWRs respectively). **I)**

Same as H but with SWRs sorted by SWR amplitude (P=1 / 0.57 / 0 for low, mid and high-amplitude SWRs). **J)** Left: PFC assembly responses to SWR quantile grid split according to both duration (y-axis) and amplitude (x-axis). Note that PFC responses were significant only for SWR quantiles of long-duration (top) and high-amplitude (right) (*P<0.05, Wilcoxon signed-rank tests). Right: Hierarchically bootstrapped estimates of PFC reactivation strength. Note that small and mid-length SWRs showed no significant PFC reactivation (P = 0.42 / 0.34, hierarchical bootstrap test), but large SWRs were followed by significant reactivation (P = 0.0076). **K)** Same as J but for CA1 assemblies (P = 1 / 0.96 / 0 hierarchical bootstrap test for small, mid-size, and large SWRs respectively).

Finding that memory reactivation was associated with a specific subset of larger SWRs prompted us to ask whether the occurrence of these SWRs was correlated with successful memory consolidation. We tested mice in two versions of the object displacement task (Fig. 2A) [44] . In the first one, mice were allowed to explore two identical objects for three 5-minute trials. For the memory recall test, conducted after a 4 hour sleep session in their home cages, one object was displaced to a different location. Mice displayed an increased exploration of the displaced object as quantified by a discrimination index (Fig. 2B and Fig. S3), establishing that they correctly remembered its original location. However, when mice were only allowed to explore the object for one 5-minute trial, they did not show an exploratory bias for the displaced object during the following test (Fig. 2B and Fig. S3). Mice did show an increased exploration of the displaced object after 5-minute exploration when the test was conducted only 1 hour later (Fig. S3). This suggests that the impaired recall at 4 hours was not due to an insufficient encoding of the new experience but that reduced experience was not enough for successful memory consolidation after 4 hours. Sleep structure and the amount of NREM were similar in the rest periods after both versions of the task (Fig. S4).

We next compared SWR features in the sleep sessions before and after the initial object exploration. SWRs before and after the 5-minute exploration sessions were similar in amplitude and duration (Fig. 2C-E). In contrast, SWR following 15-minute exploration sessions were larger on average than in the preceding sleep period (Fig. 2C-E). We thus divided all SWRs into "large" and "small" using a compound measure combining duration and amplitude (see Methods) and calculated their relative fraction in each sleep period. Overall SWR rate (0.43 and 0.47 Hz in pre and post sleep respectively, p = 0.079, Wilcoxon signed-rank test) and the relative fraction of small and large SWRs did not change following 5-minute object exploration sessions (Fig. 2F). Following 15-minute exploration sessions, the overall SWR rate increased (0.46 and 0.52 Hz, p = 6.2e-4, Wilcoxon signed-rank test), specifically due to a larger fraction of large, but not small, SWRs (Fig. 2F). Furthermore, there was a significant positive correlation between the post-sleep fraction of large SWRs and task recall performance after 15-minute but not 5-minute exploration sessions (Fig. S5).

We then examined the temporal dynamics of SWR occurrence. SWR rate and fraction of large SWRs after 15-minute exploration increased sharply at the start of the post-learning NREM sleep and then gradually decayed but remained high for ~1 hour of NREM (Fig. 2G-H). Following the 5-minute exploration, only a transient increase of SWR rate was observed right at the start of NREM sleep, and no specific increase of large SWR occurrence (Fig. 2G-H).

**Figure 2: Increased fraction of large SWRs correlates with memory performance.**
**A)** Animals were trained on the object displacement memory task, with either a single or three 5-minute exploration sessions. **B)** Memory performance (object discrimination index) on the test session was significant for the 15-minute (n = 13 mice, P = 2.4e-4, Wilcoxon signed rank test) but not the 5-minute group (n = 13 mice, P = 0.68, Wilcoxon signed rank test; P = 0.011 for 5 vs 15-minute group, Wilcoxon rank-sum test). **C)** Spectral power within the ripple band (150–250 Hz) increased for SWRs following 15-minute but not 5-minute exploration (n = 38 and 20 sessions, respectively). A line indicates the significant interval (P < 0.05, Monte Carlo test). **D)** Spectral profile of SWRs (within 50 ms of the SWR peak) was unaffected after 5-minute exploration (left), while power in the ripple frequency band was enhanced following 15-minute exploration (P < 0.05, Monte Carlo test, n = 20 / 38 sessions); a line indicates the significant interval **E)** Average sleep SWR wavelet spectrograms for pre (left), post-task sleep (middle) and their difference (right), following either 5-minute (top, n = 20) or 15-minute (bottom, n = 38) exploration sessions. **F)** Fraction of large SWRs increased following 15-minute (bottom), but not 5-minute (top), object exploration sessions (5-minute: P = 0.68, n = 20 and 15-minute: P = 8.9×10⁻⁶, n = 38, Wilcoxon signed-rank test). **G)** Incidence of small SWRs in NREM (one hour of contiguous NREM sleep before and after the task) was similar between 5-minute and 15-minute training conditions (P > 0.05, hierarchical bootstrap test). **H)** Incidence of large SWRs in post-task NREM sleep increased following 15-minute exploration (P < 0.05); a line indicates the significant interval. C-G show mean ± SEM.

## Closed-loop optogenetic SWR boosting enhanced memory reactivation and recall performance

After describing a correlation between large SWRs during sleep and successful memory retrieval, we next wanted to establish whether there is a causal link between large SWRs and memory performance by specifically manipulating them during the post-task sleep period. For this purpose, we expressed channelrhodopsin (ChR2) in CA1 pyramidal cells using viral and transgenic approaches (see Methods).

Mice were implanted with silicon probes (or wire arrays) in the hippocampus and the prefrontal cortex and bilateral optic fibers above CA1 (Fig. 3A). We detected SWRs in real-time during post-task sleep sessions and optogenetically boosted them using a closed-loop approach (Fig. 3B and Fig. S6). A 100 ms low-intensity blue-light pulse delivered upon SWR detection, prolonged the ongoing oscillation and associated spiking activity (Fig. 3C). This manipulation also resulted in a boost of SWR power (Fig. 3D). To test whether this manipulation could enhance memory performance, we applied it during the sleep period following 5-minute object exploration sessions. SWR boosting resulted in an increase in the total fraction of large SWRs during post-task sleep similar to that naturally occurring following 15-minute exploration sessions (Fig. 3E). Remarkably, while after 5-minute exploration sessions without optogenetic stimulation mice did not recall the displaced object, memory recall was successful in sessions with SWR boosting during sleep (Fig. 3F). This result indicates that optogenetic boosting of sleep SWR was able to enhance memory performance.

We hypothesized that the observed differences in memory performance following 5 or 15-minute exploration and the behavioral effects of optogenetic SWR boosting could be explained by differences in post-task neural reactivation in CA1 and PFC. To test that, we assessed sleep memory reactivation using several approaches. First, we measured the correlation of neural firing patterns in pairs of cells between the task and post-sleep SWRs. Pairwise reactivation was significant (relative to baseline sleep) in CA1, but not in the PFC following 5-minute exploration (Fig. 3G-H). In contrast, after both 15-minute or 5-minute exploration followed by SWR boosting, we observed significant memory reactivation in both CA1 and PFC (Fig. 3G-H). Next, we examined the reactivation of CA1 and PFC cell assemblies in sleep SWRs. Cell assemblies in CA1 only showed a weak reactivation following 5-minute exploration, and PFC assemblies displayed no significant reactivation at all (Fig. 3I-L). In contrast, we observed robust significant assembly reactivation in both structures after 15-minute exploration sessions, as well as 5-minute exploration with SWR boosting (Fig. 3I-L). Furthermore, we observed a session-wise positive correlation between the rate of optogenetically boosted SWRs and memory performance (Fig. S5). Optogenetic stimulation did not alter overall sleep structure or the relative fraction of NREM sleep epochs (Fig. S4). These results suggest that the improvement in memory performance due to optogenetic SWR boosting was due to an enhancement of task-related neural reactivation in CA1 and PFC.

**Figure 3: Optogenetic SWR boosting enhances memory reactivation and performance.**
**A)** Schematic of CA1 and PFC recordings combined with optic fibers targeting the hippocampus bilaterally.
**B)** Histological section showing the expression of ChR2-EYFP (green) in pyramidal cells; white arrow: optic fiber track. **C)** Example of optogenetic SWR boosting. Black trace is CA1 LFP, vertical dashed bar shows SWR real-time detection and blue rectangle shows optogenetic stimulation pulse. Raster plot below shows CA1 (in black) and PFC (in blue) units firing. **D)** Average sleep SWR wavelet spectrograms for pre (left) and post-task sleep (middle) with SWR boosting (n = 13 mice; 120 -230 Hz spectral power comparison between post-task and baseline sleep, P < 0.05, Monte Carlo test). **E)** SWR boosting resulted in an increased incidence of large SWRs in post-task sleep (*P = 0.028, n = 13, Wilcoxon signed-rank test). **F)** Memory recall performance (object discrimination index) for no stimulation 5-minute and 15-minute training sessions (same data as Figure 2B) and SWR boosting sessions (n = 13 mice, P = 1.7e-3, Wilcoxon signed rank test. SWR boost vs 5-minute P = 8.9e-3, Wilcoxon rank sum test. SWR boost vs 15-minute P = 0.96). **G)** CA1 pairwise reactivation. Grey: 5-minute exploration (P = 0.039, n = 18,708 cell pairs, hierarchical bootstrap test); green: 15-minute exploration (P = 0, n = 16,215); blue: 5-minute exploration followed by SWR-boosting (P = 0.001, n = 10,963); black dots: mean estimates; black lines: 95% confidence intervals; curves: the resampled mean distributions. **H)** Same as G) but for PFC. 5-minute exploration (grey, P = 0.14, n = 12,739 pairs); 15-minute (green: P = 0, n = 10,729); SWR-boosting (blue: P = 0.020, n = 11,157). **I)** Averaged peri-SWR CA1 assembly reactivation strength following 5-minute exploration (left), 15-minute (middle), and 5-minute followed by SWR-boosting (right); light grey: baseline pre-task assembly activity. Mean ± SEM. **J)** Quantification of I) measured by hierarchical bootstrap estimate of assembly reactivation in post-task sleep relative to baseline sleep: 5-minute exploration (grey: P = 0.032, n = 97 assemblies); 15-minute (green: P = 1.8e-3, n = 189); SWR boosting (blue: P = 0.016, n = 131 assemblies). **K-L)** Same as I-J) but for PFC assemblies. 5-minute (grey: P = 0.35, n = 79 assemblies); 15-minute (green: P = 0.0104, n = 96), SWR-boosting (blue: P = 0.011, n = 115).

**Enhanced CA1-PFC coordination during memory consolidation and recall**

Recent work in humans has shown that functional connectivity between the hippocampus and cortex increases after learning, and that this increase correlates with subsequent memory recall performance [39,40]. We thus hypothesized that the coordination between CA1 and PFC during sleep would increase following 15-minute but not 5-minute exploration sessions, and that a similar effect would be induced by optogenetic SWR boosting. We performed two different analyses to test this hypothesis.

First, we examined the coordination of the reactivation of CA1 and PFC cell assemblies. PFC cell assemblies increased their reactivation rate for SWRs associated with significant CA1 reactivation in sleep following 15-minute exploration or 5-minute exploration followed by sleep SWR boosting, but not after 5-minute exploration without stimulation (Fig. S7A). Moreover, we found evidence for an assembly–assembly-specific coordination as PFC assemblies consistently reactivated with particular CA1 assemblies throughout the sleep sessions that took place following 15-minute training or 5-minute exploration followed by SWR boosting (Fig. S7). To test for CA1-PFC coordination directly, we investigated coordinated reactivation of pairs of CA1 and PFC cells in SWR before and after the task. CA1-PFC pairwise reactivation increased following 15-minute but not 5-minute exploration sessions (Fig. 4A). SWR boosting after 5-minute exploration also resulted in enhanced pairwise cross-regional reactivation (Fig. 4A). Next, we identified cross-regional assemblies, synchronous groups of coactive cells with members from both CA1 and the PFC (Fig. 4B)[29,45], and assessed their reactivation in sleep SWR before and after the task. CA1-PFC cell assemblies showed significant coordinated reactivation after 15-minute exploration or 5-minute exploration followed by sleep SWR boosting, but not after 5-minute exploration without stimulation (Fig. 4C).

Finally, we asked whether the enhanced cross-regional coordination during sleep persisted during waking memory recall. To answer this, we measured CA1-PFC theta frequency LFP coherence, as it has been previously shown to be associated with cross-regional communication and successful memory performance[46,47]. We found that CA1-PFC theta coherence during the test phase was largely increased compared to the initial object exploration in both 15-minute sessions and 5-minute sessions followed by SWR boosting, in contrast to 5-minute exploration sessions (Fig. 4D-E). This observation suggests that extended training and artificial SWR boosting result in an enhanced CA1-PFC coordination that supports successful memory recall.

**Figure 4: Coordination of CA1 and PFC activity during sleep and memory recall**
**A)** CA1-PFC cell pair reactivation: 5-minute exploration (grey: P = 0.32, n = 10,342 cell pairs; hierarchical bootstrap test); 15-minute (green: P = 0, n = 4,573); 5-minute exploration followed by SWR-boosting (blue: P = 0.012, n = 16,334 cell pairs). **B)** Example CA1-PFC assembly. Left: CA1 and PFC units sorted by their contribution to selected assembly (assembly members in bright color at the bottom). Right: two examples of raster plot of unit firing showing assembly reactivation (shaded ovals) during a SWR (CA1 filtered LFP on top). **C)** Quantification of peri-SWR CA1-PFC assembly reactivation strength measured by a hierarchical bootstrap estimate of assembly reactivation in post-task sleep relative to baseline sleep: 5-minute (grey, P = 0.032, n = 29 assemblies); 15-minute (green, P = 1.8e-3, n = 28 assemblies); 5-minute with SWR boosting (blue, P = 0.016, n = 26 assemblies). **D)** CA1-PFC LFP coherence in training and test. Left: CA1-PFC coherence was similar during 5-minute exploration (light grey) and the subsequent test session (black). 15-minute exploration (middle) and SWR boosting following 5-minute exploration (right) resulted in enhanced CA1-PFC coherence in the subsequent test session (green or blue) compared to training (light grey). Mean ± SEM. Dashed vertical lines indicate theta frequency band. **E)** Quantification of D) measured by a hierarchical bootstrap estimate of the increase in hippocampo-prefrontal coherence during test, normalized by coherence during training (5-minute versus 15-minute training, P = 0, n = 9 and 13 sessions respectively; 5-minute versus 5-minute training with SWR boosting, P = 0, n = 9 / 5 sessions; 15-minute versus 5-minute training with SWR boosting P = 0.24, n = 13 / 5 sessions).

## Discussion

Neural ensemble reactivation in the hippocampus and associated cortical areas, such as the PFC, during sleep is regarded as a cellular hallmark of memory consolidation [11–13,15,33]. This process is primarily associated with a minority of SWRs (~10-20%) during NREM sleep. We found that the fraction of SWRs that were associated with the reactivation of a hippocampus-dependent memory task in CA1 and PFC were those with both the largest duration and amplitude. The fraction of these large SWRs selectively increased during sleep following new learning, but only when learning time was sufficient for successful memory retrieval 4 hours later. These findings help explain previous observations on the global increase in SWR

rate in sleep following learning [23,44,48] and the association of SWR duration and amplitude with increased memory load [21,49,50].

To move beyond correlational observations and establish a more direct link between SWRs, ensemble reactivation and memory performance, we deployed a closed-loop optogenetic approach [21]. Brief optogenetic activation of CA1 pyramidal cells triggered by real-time detection of sleep SWRs boosted them, mimicking the effect of extended learning. Optogenetically boosted SWRs resulted in enhanced memory reactivation in both CA1 and PFC, demonstrating a direct link between SWR features and the associated dynamics of neuronal population activity. Remarkably, this manipulation also improved mice memory performance, enabling them to successfully recall a brief prior experience, normally insufficient for successful retrieval. This effect was similar to that of extended learning experience, suggesting that sleep reactivation can compensate for reduced learning time, in line with theoretical proposals on the role of "offline" memory replay for training artificial and biological neural networks [51–53]. Multiple brain-wide phenomena, such as cortical oscillations and brain-state fluctuations, correlate with SWRs and are associated with memory consolidation [36,54–59]. Our results suggest that the contribution of some of these processes to memory could be mediated by their modulation of SWR features that in turn have a direct effect in promoting neural ensemble reactivation. For example, changes in neuromodulatory tone such as a decrease in the level of acetylcholine release in the hippocampus could create permissive substates when large SWRs are preferentially generated and memory reactivation is promoted.

A leading mechanism for the systems consolidation of new memories is the coordination of hippocampal and cortical activity promoted by SWRs [8,9,55,58,60,61]. While most of the evidence supporting this proposal is based on the coupling of hippocampal and cortical network patterns or mesoscopic activity, much less is known about how the millisecond timescale dynamics of cell assemblies across structures changes during consolidation and how sleep processes are related to persistent cross-regional coordination during waking. We found that CA1-PFC spiking coactivity patterns increased their coordinated reactivation following extended learning experience, and correlated with successful memory recall. During waking memory recall the PFC and hippocampus exhibit enhanced coherence in the theta frequency band [46,47,62]. We found similar enhanced theta coherence following extended learning experience. Importantly, optogenetically boosting SWRs in sleep following reduced learning experience produced similar effects to extended training, that is increased CA1-PFC cell assembly reactivation during sleep and enhanced theta coherence during subsequent awake recall. This result provides a missing link between the coordination of hippocampus and cortex during sleep SWRs and persistent changes in cross-regional interactions during waking. This finding offers a potential mechanism to explain observations in humans showing enhanced hippocampo-cortical functional connectivity following learning and its correlation with memory performance.

## Resource availability

### Lead Contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Dr. Antonio Fernandez-Ruiz (afr77@cornell.edu).

### Materials Availability
This study did not generate new unique reagents.

### Data and Code Availability
• Raw electrophysiological and mice behavioral data have been deposited on servers at Cornell University, and are available upon request by contacting the lead contact.
• The original code that supports the findings of this study can be downloaded from Github (https://github.com/ayalab1/neurocode).
• Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

## Acknowledgments
The authors thank members of the Oliva and Fernandez-Ruiz labs for providing useful feedback on the manuscript. This work was supported by NIH grant R00MH122582 and R01MH130367 (AO); NIH grant R01MH136355, DP2MH136496, Sloan Fellowship, Whitehall Research Grant, Klingenstein-Simons Fellowship and Pershing Square Foundation's MIND Prize (AFR); NIH grant F32MH134673 (H.L.R); Eric and Wendy and Schmidt AI in Science Postdoctoral Fellowship (R.T.).

## Author contributions
H.L.R., R.T., G.A.N., A.O. and A.F-R. conceived and designed experiments and analyses and wrote the manuscript. H.L.R., G.A.N., A.G., P.P., and R.T. performed experiments. R.T. and H.L.R. conducted the analysis.

## Declaration of interests
The authors declare no competing interests.

# METHODS

## Experimental model and study participant details
All experiments complied with guidelines established by the National Institutes of Health and approved by the Cornell University Institutional Animal Care and Use Committee (IACUC). All mice were housed in vivariums on a 12-hour light/dark cycle with a maximum of 5 animals per cage with *ad libitum* access to water and food. After surgery, mice were individually housed. Adult C57BL/6J (The Jackson Laboratory #000664, ~25-35g, ~3-6 months) or adult 'hybrid' offspring (~35-45g, ~3-6 months) were used for experiments not involving optogenetic manipulation. Hybrid mice were the first-generation offspring from a cross between C57BL/6J and FVB/NJ (The Jackson Laboratory, #001800)[63]. For optogenetic manipulation, transgenic animals or viral approaches (see Viral Injections) were used to express Channelrhodopsin (ChR2/H134R) in pyramidal neurons. For the transgenic approach, offspring of Ai32 (The Jackson Laboratory, #024109) and CamKIIα-Cre homozygous animals (The Jackson Laboratory, #00539) were used to express ChR2/H134R in pyramidal neurons.

## Method details

### Surgical Procedures
For silicon probes and viral injections, mice were anesthetized with isoflurane, and coordinates were taken following stereotaxic techniques following previously reported methods [23,36,44]. Briefly, fur was removed using Nair and skin was washed three times with alcohol and iodine. A skin incision was then made and craniotomies were performed. For hippocampal virus injections, craniotomies were made at AP -2.3 mm, ML ±1.6 mm, DV -1.1 mm from Bregma, and AP -2.7 mm, ML ±2.5 mm, DV -1.25 mm. For CA1 silicon probe implantation, craniotomies were made at AP -2.5 mm, ML -2.0 mm. For the PFC, probes were inserted in a craniotomy made at AP +1.85 mm, ML ±0.35 mm, DV 1.5 mm. Silicon probes (128-channels probes from Diagnostic Biochips or 64-channel probes from Neuronexus) were attached to micro-drives (Cambridge Neurotech or custom-printed drives). For optogenetics, 200 $\mu$m optical fibers (ThorLabs, part# FP200ERT) were glued to a probe. A stainless-steel screw (JI Morris, 000-120 x 1/16) attached to a thin wire (Phoenix Wire Inc., cat# 36744MHW) was used to connect the ground screw over the cerebellum. Next, custom 3D printed baseplates with copper mesh attached were glued to the skull using metabond (C&B Metabond Quick Adhesive Cement System). Probes were inserted above the target region, and the base of microdrives was secured to the skull using dental cement (UNIFAST Trad Acrylic Resin). Craniotomies were covered with Dura-gel (Cambridge NeuroTech Dura-Gel). Ground wires were connected to probes' ground wires and finally to the shielding copper mesh. Animals were allowed to recover for a week before recording started, and they were given carprofen subcutaneously for the first 3 days. After recovery, probes were slowly moved between 50 $\mu$m to 200 $\mu$m steps per day until the desired location was reached, where LFP physiological landmarks and characteristic LFP patterns were used to identify hippocampal and cortical layers[44,64].

### Viral Injections
Stock AAVs were acquired from Addgene. AAV-CamKIIα-ChR2/H13R-EYFP (3.4 x $10^{12}$ gc/ml, Addgene #26969-AAV5) was used. Specifically, 400nl of AAV-CamKIIα-ChR2/H13R-EYFP full virus titer was injected into the hippocampus bilaterally (see Surgical Procedures). Injections were delivered through a pulled pipette using Nanoject III (Drummond Scientific Company, 3-000-207). Animals were allowed two weeks for recovery and for viral expression to take effect before probe implantation (see Surgical Procedures).

The pAAV-CaMKIIa-hChR2(H134R)-EYFP was a gift from Karl Deisseroth (Addgene plasmid # 26969; http://n2t.net/addgene:26969; RRID: Addgene_26969)[65].

## Recording System and Data Preprocessing

Recordings were conducted with Intan RHD USB interface board or Intan Recording System. The sampling rate was 20,000 Hz, and amplification and digitization were done in the head stage by Intan RHD Electrophysiology Amplifier Chips. Acquisition was done with RHX Data Acquisition Software. LFPs were down-sampled to 1250 Hz before analysis. For video recording, an overhead mounted Basler Ace camera (Basler, acA3100-200uc) was used to record video simultaneously with neural signals. The camera's frame rate (40 Hz) was recorded as digital inputs to Intan recording systems. Mice were tracked with custom DeepLabCut [66] models trained on recorded videos andtracking was synchronized with individual frame timestamps.

## Behavioral Tasks

Before surgery, animals were exposed to an open field environment to decrease neophobia after implantation. After surgery and recovery, animals were handled daily and habituated to the experimenter, room, open field environment, and recording cables for at least a week before the start of behavioral experiments. Recordings of brain signals were done during the object displacement memory task, as previously described [44]. Specifically, ~2 hours of presleep was recorded before a 5-minute open field exploration. After open-field habituation, mice were trained to explore two identical objects for 5-minute periods. Depending on the experiment, animals were allowed a single 5-minute exploration of objects or three 5-minute exposures to objects. After object training, animals were allowed to sleep either 1 or 4 hours post-sleep, depending on experimental conditions. After postsleep, animals were tested in a 5-minute single trial where one of the identical objects was moved to a new location. A discrimination index was used to assess memory performance [67]. Specifically, the discrimination index is calculated as:

Discrimination Index =(timeObject A - timeObject B)/(timeObject A+ timeObject B)x 100

time Object A is the time during the test exploring the unmoved object, and time Object B is the time during the test exploring the moved object. The discrimination index results in a score between 1 and -1, where one corresponds to time spent studying only the moved object, -1 corresponds to exploring the unmoved object. Only sessions where animals explored for at least 5 seconds were used for behavioral experiments [44].

## Optogenetic manipulations and real-time SWR Detection

During the first 2-hours of post-task sleep, real-time SWR detection was conducted as done previously [21,23]. Three LFP channels were used for online SWR detection. An LFP channel with the strongest ripple amplitude was selected for the ripple channel, filtering between 100 and 300 Hz. Another channel with the largest sharp-wave amplitude was selected to detect sharp-wave, filtering between 10 and 40 Hz. A third noise channel was selected by choosing a channel in the neocortex where no SWR activity was present. The noise channel was filtered between 100 and 300 Hz. The root-mean-square (RMS) of the three signals was then computed by a custom-made diode bridge rectifier (1N4001 diodes, 560 kOhm resistor, 68 nF capacitor). The filtered and rectified LFP were sent to a programmable real-time processor (Cambridge Electronic Design, CED) at 20,000 Hz sampling rate. Custom scripts in SpikeDetect2 (CED) were used to manually set the detection thresholds for ripple, sharp-wave, and noise for each animal individually. SWRs were defined as events crossing the ripple and sharp-wave thresholds but not crossing the noise channel threshold.

Once a SWR was detected, a small amplitude, 100 ms, light-pulse was delivered through bilateral optic fibers. The light intensity was manually adjusted for each animal the day of the experiment while the animal

was in its homecage, where light power was increased until a ripple was produced. A trapezoid pulse with a 20 ms initial ramp and 20 ms ramp after the pulse was used [21,68].

**Tissue Processing and Immunohistochemistry**
After experiments, mice were deeply anesthetized under high-concentration (4-5%) isoflurane anesthesia. Mice were then perfused transcardially with 0.9% phosphate buffer saline (PBS) solution, after which 4% paraformaldehyde (PFA) solution was perfused. Brains were kept for 24 hours in PFA before sectioning them into 70 $\mu$m thick slices[69,70]. Slices were then stained with DAPI (Thermo Fisher, #D21490) and incubated for 30 minutes before washing. After washing, slices were mounted on glass slides with a Polyvinyl alcohol mounting medium with DABCO antifading (Millipore Sigma, #10981). Slides were imaged with a confocal microscope (Zeiss LSM 800) to image DAPI and virally expressed EYFP (see Virus Injections).

# Quantification and statistical analyses

**Spike Sorting and Single Unit Classification**
Spike sorting was performed semi-automatically using Kilosort[71] (https://github.com/cortex-lab/KiloSort). After sorting, manual curation was done in Phy (https://github.com/kwikteam/phy-data) using custom-designed plugins to acquire single well-isolated units (https://github.com/petersenpeter/phy1-plugins). Clusters were manually inspected for waveforms, auto-correlograms, and isolation distance metrics. Multi-units, noise clusters, or poorly clustered units were discarded, whereas well-isolated units were classified into putative cell types using the CellExplorer Matlab package (https://cellexplorer.org/)[72]. Three cell types were assigned: pyramidal, narrow-waveform interneurons, and wide-waveform interneurons. Two key metrics separated different putative cell types: the burst index and trough-to-peak latency [73]. The burst index was calculated as the average number of spikes in the 3-5 ms bins of a cell's autocorrelogram divided by the average number of spikes in the 200 to 300 ms bins. The average waveform from the recording site with maximum amplitude was taken to determine the trough-to-peak latency. Only well-isolated neurons with at least 100 spikes were kept for further analysis.

**Sleep and Brain States Classification**
Sleep and brain state scoring was done as previously described[29,44,74,75]. State scoring used broadband LFP, narrow band theta frequency, and estimated electromyogram (EMG) to differentiate brain states. From broadband LFP, spectrograms were calculated using a fast Fourier transform with 10 s epochs and a sliding window of 1 s. Principal Component Analysis was then computed from the Z-transform of the spectrogram from 1 to 100 Hz. The first component reflected the power in the low-frequency range (<20 Hz) with opposite weighted power at higher frequencies (>32 Hz) and distinguished non-rapid eye movement (NREM) from non-NREM periods [74]. For theta epochs, a channel with high theta power was selected, and theta dominance was quantified as the ratio of power in the 5-10 Hz and 2-16 Hz frequency bands [75]. For EMG, the zero-lag correlation of filtered signal between 300-600 Hz (Butterworth filter with filter padding between 275-625 Hz) between recording sites was used to estimate EMG activity. Pairwise correlations between channels of non-adjacent probe shank were calculated, and the mean of all correlations in time bins of 0.5 s was used as an EMG score [76]. Soft sticky thresholds were used on broadband LFP, narrow band theta, and EMG were used to determine states. High LFP in the first principal component of broadband spectrogram and high EMG were considered NREM, high narrow band theta activity and low EMG were considered rapid eye movement (REM), and remaining data (i.e., high EMG) were taken as waking state.

To prevent variations in the total amount of recorded NREM sleep from affecting our results given that SWR rate and reactivation are known to change throughout NREM sleep, we defined "baseline sleep" as the first cumulative hour of NREM sleep in the rest session recorded before training, and "post-task sleep" as the first cumulative hour of NREM sleep in the rest session recorded after training.

**SWR analyses**

For offline SWR detection, the wide-band signal was band-pass filtered (difference-of-Gaussians; zero-lag, linear phase FIR), and instantaneous power was calculated by clipping at 4 SD, rectified, and low-pass filtered. The low-pass filter cut-off was at a frequency corresponding to p cycles of the mean bandpass (for 80-250 Hz band-pass, the low-pass was 55 Hz). One channel around the CA1 pyramidal layer was chosen for ripple detection, and one channel from CA1 str. radiatum was chosen for sharp wave detection. Candidate events were identified as local minima in the sharp-wave band-filtered signal, and the corresponding ripple power was recorded. K-means clustering was used to define clustering of SWR from non-SWR events, and manual curation was used to better define the boundary and remove outliers. The events were then expanded until the (non-clipped) ripple power fell below 1 SD and the sharp-wave band-filtered signal fell below 0.5 SD above the local (±5 s) median; short events (<15 ms) and longer events (>400 ms) were discarded.

LFP signals were extracted around detected ripple events. For each event, a wavelet spectrogram was computed on the detrended LFP segment within ±500 ms of the ripple peak. The mean wavelet spectrogram for SWRs in baseline sleep and SWRs in post-task sleep, and each was corrected by subtracting power at -200 ms. Subsequently, the minimum and maximum power within the 150 Hz frequency band during the ±100 ms window around the event were used to normalize the spectrogram as w = (w-min)/(max-min).

To compare SWR spectral power in the ripple band (e.g. Figure 1E), the power values in the ripple band (150–250 Hz) within ±50 ms around the ripple peak from the baseline sleep spectrogram and the post-task sleep spectrogram for each session were normalized together by subtracting the combined mean of both and dividing by their combined standard deviation. A Monte-Carlo test was then used to compare the normalized baseline sleep power to the normalized post-task sleep power for each task condition.

To compare SWR spectral profiles (e.g. Figure 1F), the mean spectra within ±100 ms around the ripple peak from baseline sleep and post-task sleep for each session were normalized together by subtracting the combined mean of both conditions and dividing by their combined standard deviation. A Monte-Carlo test was then used to compare the normalized baseline sleep spectra to the normalized post-task sleep spectra for each task condition.

SWR start and end are defined as part of the offline SWR detection described above. SWR peak was defined as the moment of maximal ripple amplitude power within the event. SWR asymmetry was defined as how far the ripple peak deviated from the center of the event, computed as the absolute deviation of the peak from the midpoint between ripple start and end: |(peak - start) / (end - start) - 0.5|, with values near 0 indicating a symmetrical event. SWR frequency was defined as the frequency at which the wavelet spectrogram reached its maximum within ±100 ms of the ripple peak, and SWR amplitude was the corresponding power at that frequency. To visualize the SWR features, we included SWR duration, SWR amplitude, SWR frequency, SWR asymmetry, and reactivation strength of PFC cell assemblies (see below), and reduced the dimensions using an unsupervised Uniform Manifold Approximation and Projection (UMAP)[36,77] and plotted the first two dimensions, color-coding by feature.

To select SWRs with respect to their size in terms of both duration and amplitude, we designed a compound measure combining these features. For each session, SWR amplitude and SWR duration were z-scored using the mean and standard deviation of the SWR amplitude and SWR duration values,

respectively, measured from baseline sleep SWRs. SWR size was defined as the minimum of these two values, and events with an SWR size exceeding the mean size of baseline sleep SWRs were classified as "large SWRs,", and other SWRs were classified as "small SWRs".

## Cell assembly analyses

Cell assemblies were detected using an unsupervised statistical framework based on a hybrid principal component analysis (PCA) followed by independent component analysis (ICA) as reported previously[24,41,60]. In brief, for each region (the hippocampus and the prefrontal cortex), spike trains of each neuron were binned in 50-ms intervals for the whole session, the pairwise firing correlation coefficients were computed for all pairs. Next, we calculated the number of assemblies based on those principal components whose eigenvalues exceeded the threshold for random firing correlations (using the Marčenko-Pastur law), ensuring that each of the detected patterns explained more variance of the spike train correlation matrix than other patterns that would result from independently firing neurons. Independent component analysis (fast-ICA algorithm) was then used to determine for each assembly (component) the vector of weights with which each neuron's firing contributes to that assembly. The strength of each assembly $i$'s activation for a given time bin $k$ was computed as follows:

$$S_{ik} = Z_k^T P Z_k$$

where Z is the population activity matrix of the z-scored firing rate of each unit, and $P$ is the outer product of the component $i$'s weights, in which the diagonal has been set to zero so that isolated spikes from individual units do not contribute to S.

The goal of the cell assembly analysis was to study the patterns associated with the learning experience (i.e. object exploration). However, because the task design required the animals to be habituated to the environment in advance, we expected a number of cell assemblies to have formed during this habituation period, reflecting task-related variables (such as the environment layout, distal cues, etc.) that are orthogonal to the learning experience. We therefore discarded cell assemblies with significant reactivation in baseline sleep before object exploration from subsequent analyses.

To detect cross-structural cell assemblies, we followed a procedure described previously [29,45]. Briefly, we combined the recordings in the hippocampus and the prefrontal cortex and performed assembly detection as described above. For a given assembly, the weight vector was normalized to have unit Euclidean norm, and units whose weights exceeded $1/\sqrt{N}$, where N is the number of recorded units, were considered assembly members. Cell assemblies with members in both the hippocampus and the prefrontal cortex were retained as cross-structural cell assemblies.

Assembly responses to SWRs were quantified as the mean reactivation strength within 0–100 ms of SWR onset. To allow pooling across assemblies, responses were z-scored within each assembly. For analyses requiring classification of individual SWR responses (Figure S7), responses exceeding z = 1.96 (P < 0.05) were considered significant.

## Pairwise reactivation

Pairwise reactivation analyses were performed as described previously [24,78]. Briefly, we binned the activity of all units during SWRs in baseline sleep, and used this spike count matrix to compute a pairwise cross-correlation matrix $C_{baseline}$. Similarly, we binned the activity of all units in 100 ms during movement intervals of training of the object task, and used this spike count matrix to compute a pairwise cross-correlation matrix $C_{behavior}$. Finally, we binned the activity of all units during SWRs in post-task sleep, and used this spike count matrix to compute a pairwise cross-correlation matrix $C_{post}$. We used

linear regression between $C_{baseline}$ and $C_{behavior}$ and between $C_{baseline}$ and $C_{post}$ to remove the portion of the variance that can be explained by preexisting patterns. The reactivation of pairwise behavioral patterns in post-task sleep was quantified as the correlation between the resultant residuals $R_{behavior}$ and $R_{post}$.

## Hierarchical bootstrap

Hierarchical bootstrap [79] was performed to analyze data with hierarchical structure. Briefly, bootstrap datasets were created by resampling with replacement following levels of hierarchical order in the order of animals followed by sessions followed by the within-session observations. Each resampled bootstrap dataset was used to calculate an estimate of the mean, in a total of 5000 resampling times.

For the pairwise reactivation analysis, hierarchical bootstrap was used to estimate a distribution of correlation values between $R_{behavior}$ and $R_{post}$ . To this end, $R_{behavior}$ and $R_{post}$ were resampled together according to the hierarchical structure, and the correlation between them was computed for each bootstrap sample.

In all cases (whether estimating a mean or a correlation), the P-value reflects the direct comparison of the distribution of estimates to zero, or comparisons of different distributions to each other, after controlling for the other nesting variables.

## Statistical Analysis

Data analysis was performed using custom routines in Python and MATLAB (MathWorks). No specific analysis was used to estimate minimal population sample or group size, but the number of animals, sessions and recorded cells were larger or similar to those employed in previous related work [20,25,36,44]. Unless otherwise noted, non-parametric Wilcoxon rank-sum or Wilcoxon signed-rank test was used for unpaired and paired data comparisons respectively, and ANOVA and Kruskal-Wallis was used for multiple comparisons. All statistical tests were two-tailed with $p < 0.05$ as the cutoff for statistical significance, which is indicated by asterisks (*$p<0.05$, **$p<0.01$, ***$p<0.001$, and ****$p<0.0001$). Error bars show SEMs, and boxplots show median (central mark), 75th (box), and 90th (whiskers) percentile, unless indicated otherwise. Due to experimental constraints of optogenetic experiments, experimenters were not blind to these manipulations.

# References

1. Diekelmann, S., and Born, J. (2010). The memory function of sleep. Nat. Rev. Neurosci. *11*, 1*14–126*. https://doi.org/10.*1038*/nrn2762.

2. Frankland, P.W., and Bontempi, B. (2005). The organization of recent and remote memories. Nat. Rev. Neurosci. 6, 1*19–130*. https://doi.org/10.1038/nrn1607.

3. Squire, L.R. (1*992*). Memory and the Hippocampuss: A Synthesis From Findings With Rats, Monkeys, and Humans. Psychol. Rev. 99, 1*95–231*. https://doi.org/10.10*37*/0033-295x.99.2.195.

4. McClelland, J.L., McNaughton, B.L., and O'Reilly, R.C. (1995). Why There Are Complementary Learning Systems in the Hippocampus and Neocortex: Insights From the Successes and Failures of Connectionist Models of Learning and Memory. Psychol. Rev. *102*, 419–457. https://doi.org/10.1037/0033-295x.102.3.419.

5. Marr, D. (1971). Simple memory: a theory for archicortex. Philos. Trans. R. Soc. Lond. B, Biol. Sci. *262*, 23–81. https://doi.org/10.*1098*/rstb.1971.0078.

6. Alvarez, P., and Squire, L.R. (1994). Memory consolidation and the medial temporal lobe: a simple network model. Proc. Natl. Acad. Sci. *91*, 7041–7045. https://doi.org/10.1073/pnas.91.*15*.7041.

7. Buzsáki, G. (1989). Two-stage model of memory trace formation: A role for "noisy" brain states. Neuroscience 31, 551–570. https://doi.org/10.*1016*/0306-45*22*(89)*90423*-5.

8. Preston, A.R., and Eichenbaum, H. (2013). Interplay of Hippocampus and Prefrontal Cortex in Memory. Curr. Biol. 23, R764–R773. https://doi.org/10.1016/j.cub.2013.05.041.

9. Klinzing, J.G., Niethard, N., and Born, J. (2019). Mechanisms of systems memory consolidation during sleep. Nat. Neurosci. 22, 1598–1610. https://doi.org/10.1038/s41593-019-0*467*-3.

10. Joo, H.R., and Frank, L.M. (20*18*). The hippocampal sharp wave–ripple in memory retrieval for immediate use and consolidation. Nat. Rev. Neurosci. 19, 744–757. https://doi.org/10.1038/s41583-018-0077-1.

11. Buzsáki, G. (2015). Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. Hippocampus *25*, 1073–1188. https://doi.org/10.*1002*/hipo.22488.

12. Wilson, M.A., and McNaughton, B.L. (1994). Reactivation of Hippocampal Ensemble Memories During Sleep. Science *265*, 676–679. https://doi.org/10.1126/science.80*36517*.

13. Foster, D.J. (2017). Replay Comes of Age. Annu. Rev. Neurosci. *40*, 581–602. https://doi.org/10.1146/annurev-neuro-07*21*16-031538.

14. Diba, K., and Buzsáki, G. (2007). Forward and reverse hippocampal place-cell sequences during ripples. Nat. Neurosci. 10, 1241–1242. https://doi.org/10.1038/nn1*961*.

15. Lee, A.K., and Wilson, M.A. (2002). Memory of Sequential Experience in the Hippocampus during Slow Wave Sleep. Neuron 36, 1183–1194. https://doi.org/10.1016/s0896-6273(02)01096-6.

16. Girardeau, G., Benchenane, K., Wiener, S.I., Buzsáki, G., and Zugaro, M.B. (2009). Selective suppression of hippocampal ripples impairs spatial memory. Nat. Neurosci. 12, 1222–1223. https://doi.org/10.1038/nn.23*84*.

17. Gupta, A.S., Meer, M.A.A. van der, Touretzky, D.S., and Redish, A.D. (2010). Hippocampal Replay Is Not a Simple Function of Experience. Neuron 65, 695–705. https://doi.org/10.1016/j.neuron.2010.01.034.

18. Jadhav, S.P., Kemere, C., German, P.W., and Frank, L.M. (2012). Awake Hippocampal Sharp-Wave

Ripples Support Spatial Memory. Science *336*, 1454–1458. https://doi.org/10.1126/science.1217230.

19. Pfeiffer, B.E., and Foster, D.J. (2013). Hippocampal place-cell sequences depict future paths to remembered goals. Nature *497*, 74–79. https://doi.org/10.1038/nature12112.

20. Farooq, U., Sibille, J., Liu, K., and Dragoi, G. (2019). Strengthened Temporal Coordination within Pre-existing Sequential Cell Assemblies Supports Trajectory Replay. Neuron 103, 719-733.e7. https://doi.org/10.1016/j.neuron.2019.05.040.

21. Fernández-Ruiz, A., Oliva, A., Oliveira, E.F. de, Rocha-Almeida, F., Tingley, D., and Buzsáki, G. (2019). Long-duration hippocampal sharp wave ripples improve memory. Science *364*, 1082–1086. https://doi.org/10.1126/science.aax0758.

22. Gillespie, A.K., Maya, D.A.A., Denovellis, E.L., Liu, D.F., Kastner, D.B., Coulter, M.E., Roumis, D.K., Eden, U.T., and Frank, L.M. (2021). Hippocampal replay reflects specific past experiences rather than a plan for subsequent choice. Neuron 109, 3149-3163.e6. https://doi.org/10.1016/j.neuron.2021.07.029.

23. Oliva, A., Fernández-Ruiz, A., Leroy, F., and Siegelbaum, S.A. (2020). Hippocampal CA2 sharp-wave ripples reactivate and promote social memory. Nature *587*, 264–269. https://doi.org/10.1038/s41586-020-2758-y.

24. Liu, C., Todorova, R., Tang, W., Oliva, A., and Fernandez-Ruiz, A. (2023). Associative and predictive hippocampal codes support memory-guided behaviors. Science *382*, eadi8237. https://doi.org/10.1126/science.adi8237.

25. Giri, B., Kinsky, N., Kaya, U., Maboudi, K., Abel, T., and Diba, K. (2024). Sleep loss diminishes hippocampal reactivation and replay. Nature *630*, 935–942. https://doi.org/10.1038/s41586-024-07538-2.

26. Peyrache, A., Khamassi, M., Benchenane, K., Wiener, S.I., and Battaglia, F.P. (2009). Replay of rule-learning related neural patterns in the prefrontal cortex during sleep. Nat. Neurosci. 12, 919–926. https://doi.org/10.1038/nn.2337.

27. Euston, D.R., Tatsuno, M., and McNaughton, B.L. (2007). Fast-Forward Playback of Recent Memory Sequences in Prefrontal Cortex During Sleep. Science *318*, 1147–1150. https://doi.org/10.1126/science.1148979.

28. Norman, Y., Yeagle, E.M., Khuvis, S., Harel, M., Mehta, A.D., and Malach, R. (2019). Hippocampal sharp-wave ripples linked to visual episodic recollection in humans. Science 365, eaax1030. https://doi.org/10.1126/science.aax1030.

29. Harvey, R.E., Robinson, H.L., Liu, C., Oliva, A., and Fernandez-Ruiz, A. (2023). Hippocampo-cortical circuits for selective memory encoding, routing, and replay. Neuron *111*, 2076-2090.e9. https://doi.org/10.1016/j.neuron.2023.04.015.

30. Tang, W., Shin, J.D., Frank, L.M., and Jadhav, S.P. (2017). Hippocampal-Prefrontal Reactivation during Learning Is Stronger in Awake Compared with Sleep States.  J. Neurosci. 37, 11789–11805. https://doi.org/10.1523/jneurosci.2291-17.2017.

31. Ólafsdóttir, H.F., Carpenter, F., and Barry, C. (2017). Task Demands Predict a Dynamic Switch in the Content of Awake Hippocampal Replay. Neuron 96, 925-935.e6. https://doi.org/10.1016/j.neuron.2017.09.035.

*32.* Gorriz, M.H., Takigawa, M., and Bendor, D. (2023). The role of experience in prioritizing hippocampal replay. Nat. Commun. 14, 8157. https://doi.org/10.1038/s41467-023-43939-z.

33. Dupret, D., O'Neill, J., Pleydell-Bouverie, B., and Csicsvari, J. (2010). The reorganization and

reactivation of hippocampal maps predict spatial memory performance. Nat. Neurosci. 13, 995–1002. https://doi.org/10.1038/nn.2599.

34. Shin, J.D., Tang, W., and Jadhav, S.P. (2019). Dynamics of Awake Hippocampal-Prefrontal Replay for Spatial Learning and Memory-Guided Decision Making. Neuron *104*, 1*110*-1125.e7. https://doi.org/10.1016/j.neuron.2019.09.012.

35. Kaefer, K., Nardin, M., Blahna, K., and Csicsvari, J. (2020). Replay of Behavioral Sequences in the Medial Prefrontal Cortex during Rule Switching. Neuron *106*, 154-165.e6. https://doi.org/10.1016/j.neuron.2020.01.015.

36. Chang, H., Tang, W., Wulf, A.M., Nyasulu, T., Wolf, M.E., Fernandez-Ruiz, A., and Oliva, A. (2025). Sleep microstructure organizes memory replay. Nature *637*, 1161–1169. https://doi.org/10.1038/s41586-024-08340-w.

37. O'Neill, J., Senior, T.J., Allen, K., Huxter, J.R., and Csicsvari, J. (2008). Reactivation of experience-dependent cell assembly patterns in the hippocampus. Nat. Neurosci. 11, 209–215. https://doi.org/10.1038/nn2037.

38. Silva, D., Feng, T., and Foster, D.J. (2015). Trajectory events across hippocampal place cells require previous experience. Nat. Neurosci. 18, 1772–1779. https://doi.org/10.1038/nn.4151.

39. Tambini, A., Ketz, N., and Davachi, L. (2010). Enhanced Brain Correlations during Rest Are Related to Memory for Recent Experiences. Neuron 65, 280–290. https://doi.org/10.1016/j.neuron.2010.01.001.

40. Tambini, A., and Davachi, L. (2013). Persistence of hippocampal multivoxel patterns into postencoding rest is related to memory. Proc. Natl. Acad. Sci. 110, 19591–19596. https://doi.org/10.1073/pnas.1308499110.

41. Lopes-dos-Santos, V., Ribeiro, S., and Tort, A.B.L. (2013). Detecting cell assemblies in large neuronal populations. J. Neurosci. Methods *220*, 149–166. https://doi.org/10.1016/j.jneumeth.2013.04.010.

42. Fernández-Ruiz, A., Oliva, A., Soula, M., Rocha-Almeida, F., Nagy, G.A., Martin-Vazquez, G., and Buzsáki, G. (2021). Gamma rhythm communication between entorhinal cortex and dentate gyrus neuronal assemblies. Science *372*, eabf3119. https://doi.org/10.1126/science.abf3119.

43. Sebastian, E.R., Quintanilla, J.P., Sánchez-Aguilera, A., Esparza, J., Cid, E., and Prida, L.M. de la (2023). Topological analysis of sharp-wave ripple waveforms reveals input mechanisms behind feature variations. Nat. Neurosci. 26, 2171–2181. https://doi.org/10.1038/s41593-023-01471-9.

44. Karaba, L.A., Robinson, H.L., Harvey, R.E., Chen, W., Fernandez-Ruiz, A., and Oliva, A. (2024). A hippocampal circuit mechanism to balance memory reactivation during sleep. Science *385*, 738–743. https://doi.org/10.1126/science.ado5708.

45. Oberto, V.J., Boucly, C.J., Gao, H., Todorova, R., Zugaro, M.B., and Wiener, S.I. (2022). Distributed cell assemblies spanning prefrontal cortex and striatum. Curr. Biol. 32, 1-13.e6. https://doi.org/10.1016/j.cub.2021.10.007.

46. Benchenane, K., Peyrache, A., Khamassi, M., Tierney, P.L., Gioanni, Y., Battaglia, F.P., and Wiener, S.I. (2010). Coherent Theta Oscillations and Reorganization of Spike Timing in the Hippocampal-Prefrontal Network upon Learning. Neuron 66, 921–936. https://doi.org/10.1016/j.neuron.2010.05.013.

47. Nardin, M., Kaefer, K., Stella, F., and Csicsvari, J. (2023). Theta oscillations as a substrate for medial prefrontal-hippocampal assembly interactions. Cell Rep. 42, 113015. https://doi.org/10.1016/j.celrep.2023.113015.

48. Eschenko, O., Ramadan, W., Mölle, M., Born, J., and Sara, S.J. (2008). Sustained increase in

hippocampal sharp-wave ripple activity during slow-wave sleep after learning. Learn. Mem. 15, 222–228. https://doi.org/10.1101/lm.726008.

49. Samanta, A., Aleman-Zapata, A., Agarwal, K., Özsezer, P., Alonso, A., Meij, J. van der, Rayan, A., Navarro-Lobato, I., and Genzel, L. (2023). CBD lengthens sleep but shortens ripples and leads to intact simple but worse cumulative memory. iScience 26, 108327. https://doi.org/10.1016/j.isci.2023.108327.

50. Vancura, B., Geiller, T., Grosmark, A., Zhao, V., and Losonczy, A. (2023). Inhibitory control of sharp-wave ripple duration during learning in hippocampal recurrent networks. Nat. Neurosci. 26, 788–797. https://doi.org/10.1038/s41593-023-01306-7.

51. Mattar, M.G., and Daw, N.D. (2018). Prioritized memory access explains planning and hippocampal replay. Nat. Neurosci. 21, 1609–1617. https://doi.org/10.1038/s41593-018-0232-z.

52. Levenstein, D., Efremov, A., Eyono, R.H., Peyrache, A., and Richards, B. (2024). Sequential predictive learning is a unifying theory for hippocampal representation and replay. bioRxiv, 2024.04.28.591528. https://doi.org/10.1101/2024.04.28.591528.

53. Lengyel, M., and Dayan, P. (2008). Hippocampal Contributions to Control: The Third Way. In Advances in Neural Information Processing Systems (Curran Associates, Inc.).

54. Siapas, A.G., and Wilson, M.A. (1998). Coordinated Interactions between Hippocampal Ripples and Cortical Spindles during Slow-Wave Sleep. Neuron 21, 1123–1128. https://doi.org/10.1016/s0896-6273(00)80629-7.

55. Maingret, N., Girardeau, G., Todorova, R., Goutierre, M., and Zugaro, M. (2016). Hippocampo-cortical coupling mediates memory consolidation during sleep. Nat. Neurosci. 19, 959–964. https://doi.org/10.1038/nn.4304.

56. Sirota, A., Csicsvari, J., Buhl, D., and Buzsáki, G. (2003). Communication between neocortex and hippocampus during sleep in rodents. Proc. Natl. Acad. Sci. 100, 2065–2069. https://doi.org/10.1073/pnas.0437938100.

57. Battaglia, F.P., Sutherland, G.R., and McNaughton, B.L. (2004). Hippocampal sharp wave bursts coincide with neocortical "up-state" transitions. Learn. Mem. 11, 697–704. https://doi.org/10.1101/lm.73504.

58. Staresina, B.P., Bergmann, T.O., Bonnefond, M., Meij, R. van der, Jensen, O., Deuker, L., Elger, C.E., Axmacher, N., and Fell, J. (2015). Hierarchical nesting of slow oscillations, spindles and ripples in the human hippocampus during sleep. Nat. Neurosci. 18, 1679–1686. https://doi.org/10.1038/nn.4119.

59. Latchoumane, C.-F.V., Ngo, H.-V.V., Born, J., and Shin, H.-S. (2017). Thalamic Spindles Promote Memory Formation during Sleep through Triple Phase-Locking of Cortical, Thalamic, and Hippocampal Rhythms. Neuron 95, 424-435.e6. https://doi.org/10.1016/j.neuron.2017.06.025.

60. Todorova, R., and Zugaro, M. (2019). Isolated cortical computations during delta waves support memory consolidation. Science *366*, 377–381. https://doi.org/10.1126/science.aay0616.

61. Todorova, R., and Zugaro, M. (2020). Hippocampal ripples as a mode of communication with cortical and subcortical areas. Hippocampus 30, 39–49. https://doi.org/10.1002/hipo.22997.

62. Siapas, A.G., Lubenov, E.V., and Wilson, M.A. (2005). Prefrontal Phase Locking to Hippocampal Theta Oscillations. Neuron 46, 141–151. https://doi.org/10.1016/j.neuron.2005.02.028.

63. Sloin, H.E., Bikovski, L., Levi, A., Amber-Vitos, O., Katz, T., Spivak, L., Someck, S., Gattegno, R., Sivroni, S., Sjulson, L., et al. (2022). Hybrid Offspring of C57BL/6J Mice Exhibit Improved Properties for Neurobehavioral Research. eNeuro 9, ENEURO.0221-22.2022. https://doi.org/10.1523/eneuro.0221-

22.2022.

64. Senzai, Y., Fernandez-Ruiz, A., and Buzsáki, G. (2019). Layer-Specific Physiological Features and Interlaminar Interactions in the Primary Visual Cortex of the Mouse. Neuron 101, 500-513.e5. https://doi.org/10.1016/j.neuron.2018.12.009.

65. Lee, J.H., Durand, R., Gradinaru, V., Zhang, F., Goshen, I., Kim, D.-S., Fenno, L.E., Ramakrishnan, C., and Deisseroth, K. (2010). Global and local fMRI signals driven by neurons defined optogenetically by type and wiring. Nature *465*, 788–792. https://doi.org/10.1038/nature09108.

66. Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., and Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. Nat. Neurosci. 21, 1281–1289. https://doi.org/10.1038/s41593-018-0209-y.

67. Genzel, L., Schut, E., Schröder, T., Eichler, R., Khamassi, M., Gomez, A., Lobato, I.N., and Battaglia, F. (2019). The object space task shows cumulative memory expression in both mice and rats. PLoS Biol. 17, e3000322. https://doi.org/10.1371/journal.pbio.3000322.

68. Oliva, A., Fernández-Ruiz, A., Oliveira, E.F. de, and Buzsáki, G. (2018). Origin of Gamma Frequency Power during Hippocampal Sharp-Wave Ripples. Cell Rep. 25, 1693-1700.e4. https://doi.org/10.1016/j.celrep.2018.10.066.

69. Tan, Z., Robinson, H.L., Yin, D.-M., Liu, Y., Liu, F., Wang, H., Lin, T.W., Xing, G., Gan, L., Xiong, W.-C., et al. (2018). Dynamic ErbB4 Activity in Hippocampal-Prefrontal Synchrony and Top-Down Attention in Rodents. Neuron 98, 380-393.e4. https://doi.org/10.1016/j.neuron.2018.03.018.

70. Robinson, H.L., Tan, Z., Santiago-Marrero, I., Arzola, E.P., Dong, T.V., Xiong, W.-C., and Mei, L. (2021). Neuregulin 1 and ErbB4 Kinase Actively Regulate Sharp Wave Ripples in the Hippocampus. J. Neurosci. 42, 390–404. https://doi.org/10.1523/jneurosci.1022-21.2021.

71. Pachitariu, Steinmetz, M., Kadir, Carandini, S., and Harris, M. (2016). Fast and accurate spike sorting of high-channel count probes with KiloSort. In Advances in Neural Information Processing Systems., Lee, D. and Sugiyama, M. and Luxburg, U. and Guyon, and I. and Garnett, eds.

72. Petersen, P.C., Siegle, J.H., Steinmetz, N.A., Mahallati, S., and Buzsáki, G. (2021). CellExplorer: A framework for visualizing and characterizing single neurons. Neuron 109, 3594-3608.e2. https://doi.org/10.1016/j.neuron.2021.09.002.

73. Sirota, A., Sirota, A., Montgomery, S.M., Montgomery, S.M., Fujisawa, S., Fujisawa, S., Isomura, Y., Isomura, Y., Zugaro, M., Zugaro, M., et al. (2008). Entrainment of neocortical neurons and gamma oscillations by the hippocampal theta rhythm. Neuron. https://doi.org/10.1016/j.neuron.2008.09.014.

74. Levenstein, D., Buzsáki, G., and Rinzel, J. (2019). NREM sleep in the rodent neocortex and hippocampus reflects excitable dynamics. Nat. Commun. 10, 2478. https://doi.org/10.1038/s41467-019-10327-5.

75. Watson, B.O., Levenstein, D., Greene, J.P., Gelinas, J.N., and Buzsáki, G. (2016). Network Homeostasis and State Dynamics of Neocortical Sleep. Neuron 90, 839–852. https://doi.org/10.1016/j.neuron.2016.03.036.

76. Schomburg, E.W., Fernández-Ruiz, A., Mizuseki, K., Berényi, A., Anastassiou, C.A., Koch, C., and Buzsáki, G. (2014). Theta Phase Segregation of Input-Specific Gamma Patterns in Entorhinal-Hippocampal Networks. Neuron 84, 470–485. https://doi.org/10.1016/j.neuron.2014.08.051.

77. Meehan, C., Ebrahimian, J., Moore, W., and Meehan, S. (2025). Uniform Manifold Approximation and Projection (UMAP). Preprint.

78. Giri, B., Miyawaki, H., Mizuseki, K., Cheng, S., and Diba, K. (2019). Hippocampal Reactivation Extends for Several Hours Following Novel Experience.  J. Neurosci. 39, 866–875. https://doi.org/10.1523/jneurosci.1950-18.2018.

79. Saravanan, V., Berman, G.J., and Sober, S.J. (2020). Application of the hierarchical bootstrap to multi-level data in neuroscience. Neurons, Behav., data Anal. theory 3.