Title

Socio-acoustic co-selection? Vocal encoding of sociability prevails over emotions in sheep bleats

Authors

Avelyne S. Villain [1,2], Alain Boissy [3], Gaetan Bonnafé [4], Christian Durand [4], Sébastien Douls [4],
Paul Renaud-Goud [1,5], Marie-Madeleine Mialon [3], Dominique Hazard [2], Elodie F. Briefer [1]

Affiliations

[1] Behavioural Ecology Group, Section for Ecology and Evolution, Department of Biology,
University of Copenhagen, 2100 Copenhagen Ø, Denmark

[2] GenPhySE, Université de Toulouse, INRAE, ENVT, 31326 Castanet-Tolosan, France

[3] UMR Herbivores, Universit*é* Clermont Auvergne, INRAE, VetAgro Sup, 63122 Saint-Genès-
Champanelle , France

[4] UE Domaine de la Fage, INRAE, 12250 Saint-Jean Saint-Paul, France

[5] IRIT, Université Toulouse III - Paul Sabatier, Université Toulouse Jean Jaurès, 31400 Toulouse,
France

Key words

Acoustic communication, Emotions, Social behaviour, Temperament, *Ovis aries, G*enetic selection.

Abstract

Vocalisations of animals are good indicators of their emotions. Temperament is known to influence the
regulation and expression of emotions. However, how animal temperament affects their vocalisations
and particularly their vocal expressions of emotions remains largely unexplored. Sociability is often
measured as the behavioural reactivity to social separation and is a temperament trait intrinsically

linked to emotional reactivity. Most social species respond to this challenging situation using contact

calls. Here, we investigated whether the acoustic structure of these calls reflect sociability, emotions or

both. We used 42 female lambs from two diverging sheep lines selected for high or low sociability.

High bleats were recorded both in isolation (social challenge) and before receiving a food treat (non

social context) to investigate the link between vocalisations, emotions and heritable sociability. The

acoustic features of isolation bleats differed between the lines, but it was not the case for pre-feeding

bleats. Surprisingly, the genetic selection index and social behaviour were better predictors of the struc-

ture of isolation bleats than the arousal. Last, encoding of individuality in isolation bleats was impaired

by the genetic selection. Our findings suggest a socio-acoustic co-selection: selecting for sociable ani-

mals affects vocal signatures in calls produced during a social separation.

33 *1. Introduction*

Vocalisations encode both static (i.e. relatively stable) information about the sender, such as its sex,

identity and social rank [1,2], as well as more dynamic information about its internal motivation and

emotional states [3]. Indeed, variation in acoustic signaling in term of call types, rate of production of

these types, or spectro-temporal features of vocalisations arising as a function of the sender's emotions

have been discovered in many non-human animals and show construct validity as indicators of emo-

tional states [3–5]. Regulation and expression of dynamic features such as emotions depend on more

stable traits, namely personality, in humans [6,7], with some evidence suggesting that this is also the

case in non-human animals [8]. As a result, a link between temperament or personality and vocal ex-

pressions of emotions is expected, but has rarely been explored in non-human animals.

Temperament traits and personalities are defined as consistent individual differences in response to

species-specific environmental or social stimuli [9]. Temperament traits are heritable, and selective

breeding of divergent lines for opposite temperament traits provide experimental models to study phe-

notypic consequences of heritable components in ecology [9,10], medical [11] or agricultural research

[12]. Boldness-shyness, exploration-avoidance, activity, sociability, aggressiveness or emotional reac-

tivity, are temperament traits commonly assessed in both humans and non-human animals [9]. Most of

these traits are inherently linked to, and measured as, individual differences in physiological and be-

havioural responses to challenging situations [9–11], and are thus intrinsically linked to emotions

[13,14]. In particular, sociability may be defined as "an individual's tendency or propensity to associate

with other individuals, where the association is not driven by reproduction or aggression" [15] and may

be partly measured as "an individual's reaction to the absence of conspecifics" [9], i.e. quantifying the stress response and/or exacerbated behaviours to isolation.

In response to challenging situations, many animal species respond using vocalisations (alarm calls, contact calls) [12]. Yet, studies on the link between the acoustic structure of vocalisations (i.e. variation in duration, frequencies and amplitude) and temperament traits are rare, despite their potential to highlight non-invasive and rapid indicators of temperament. A few studies have incorporated call rate in their behavioural assessment of temperament [13,16–19]. In some cases, call types were produced in a context-specific way [16,20], which may emphasize the functional link between a call type and a situation. Going further into the acoustic structure of vocalisations produced during temperament assessment, Leliveld et al (2007) focused on two common call types emitted by pigs, grunts and squeals, to investigate vocal correlate of emotional reactivity [21]. They showed that grunts could be consistently divided into high and low frequency grunts and that high frequency grunts were indicative of more active, more explorative and less stressful individuals. As vocalisations are strongly affected by emotions [3–5], and the range of emotions experienced by an individual is driven by its temperament, we can predict that the acoustic structure of vocalisations may reflect the temperament of non-human animals in a broader range of species as currently demonstrated [21], similarly to vocal correlates of personality in human voice [22]. Within a social group, acoustic correlates of temperament could be very useful for receivers, as it would allow them to predict the behaviour of others.

Sheep (*Ovis aries)* are highly gregarious and vocal species [23]. Their vocal repertoire contains two main structural call types: high and low bleats, differentiated by the opening of the month during the call (open and closed, respectively). High and low bleats are context specify; namely, high bleats are usually produced in higher arousal states, especially in negative situations such as isolation from conspecifics [23], and low bleats in lower arousal states, such as during short distance mother-lamb communication [24,25]. The acoustic structure of sheep bleats encodes individuality [24], hence enabling vocal recognition, especially between mother and lamb [25,26] as well as between kin and non-kin lambs [27], which is at the basis of the strong social bonds on which the lamb's survival depends at early stages of its live. Several temperament traits, such as 'exploration-avoidance', 'boldness-shyness' or 'sociability', were identified in sheep [28]. Vocal production rate (i.e. the number of vocalisation produced per time unit) was found as indicative of sociability [28]. In addition, selective breeding based on sociability revealed that the rate of calls emitted during isolation from conspecifics was heritable and that highly sociable sheep genetic line produced bleats at a higher rate than less sociable ones

[29]. However, whether the acoustic structure of sheep bleats also encodes sociability remains un-
known.

In the present study, we studied a population of sheep consisting of two divergent lines of female sheep
selected for high or low sociability for over 11 years (at the time of data collection). We evaluated the
consequences of three generations of selection for sociability on the structure of sheep bleats recorded
in two contexts that trigger emotions of different valence (negative/unpleasant versus positive/pleasant)
of moderate to high arousal and vary in whether they are social-related (isolation from conspecifics) or
not social-related (prior to a repeated and announced food treat delivery), to investigate the drivers of
bleat structure and study potential vocal correlates of temperament in sheep. We designed two non-
exclusive hypotheses to explain bleat structure: the emotional hypothesis and the socio-acoustic co-
selection hypothesis, while (partially) controlling for heritability of the structure of the vocal apparatus.
According to the emotional hypothesis, the isolation test should trigger a higher level of arousal for
high-sociability sheep than less sociable ones, which should explain variation in acoustic structure,
while the 'anticipation' of a food treat should not provoke differential arousal states between the two
lines. Considering vocal indicators of negative arousal, which are consistent across species [3], this
should result in bleats of longer duration and with higher frequencies in the isolation bleats of high-
sociability sheep. Based on the socio-acoustic co-selection hypothesis, selection for sociability should
be the main driver of the acoustic structure of bleats, meaning that genetic selection for sociability and
social behaviour should better explain variation in acoustic structure than emotional arousal. We first
tested for an acoustic signature of the two divergent lines of sheep, in the two recorded contexts. We
predicted that if selection for sociability drives the acoustic structure of bleats, a 'line acoustic
signature' should be stronger in calls produced in a social context (isolation) than in a non social
context (food treat 'anticipation'). We then tested the importance of several predictors (biometry,
emotional arousal, selection index and social behaviour) on the acoustic structure of isolation bleats
while partially controlling for pedigree effects to disentangle between the emotional and socio-acoustic
co-selection hypotheses. Last, as the salience of vocal individuality (i.e. how much acoustic structure
differ between individuals) has been shown to be affected by both emotional arousal [30] and valence
[31], we additionally evaluated the potential consequences of the divergent selection for sociability on
individual signature in bleats, a key stone for social communication.

2. Materials and Methods

2.1 Ethical note

This study was approved by the Animal Ethics Committee SCIENCE ET SANTE ANIMALES N°115 of the Veterinary School of Toulouse under the number: SSA_2022_003.

2.2 Animals and management

The experimental animals were Romane meat sheep, reared at the INRAE experimental farm of La Fage (Saint-Jean Saint-Paul, Causse du Larzac, France) exclusively outdoors. The flock comprised about 250 reproductive females reared on 280 ha of rangelands (see [32] for details about the farming system and management characteristics). This population has been established in 2000 and selections started in 2011. Lambs used in this study were born outdoors in the spring (from April 2nd to May 2nd 2022), and remained with their dams until weaning (late June) with limited contact with humans. Lambs had access to additional food concentrate from four weeks of age. At 1 day of age, lambs and mothers were subjected to an isolation test in the same environment as the arena test (presented below) and a human approach test. Lambs were weighed on June 14th 2022. From July 4th to 14th 2022, 152 female lambs were subjected to individual tests, upon which 42 female lambs were selected according to their high (S+) or low (S-) reactivity to social isolation to constitute the two divergent lines (see next section for procedure). Four groups of five female lambs within each line were composed (two lambs, selected as replacement were not introduced into the groups). Three lambs (3S+) died from respiratory disease early after the constitution of the groups, one of which was replaced, leading to groups of three to five lambs. Sheep were individually marked with color spray and brought to pasture fields (approx. 50*50m). Lambs had unlimited access to water and grass and were supplemented in hay due to a very dry season in 2022. Every morning, between 7:00 and 10:00, at random times and for 5 weeks, each group received a 800g portion of barley.

2.3 Divergent selection based on behavioural reactivity to isolation and human presence

2.3.1 Tests procedure

The tests were performed between July 4th and 14th 2022 on 152 females. The day before being tested, 20 to 30 lambs were removed from the rest of the flock, moved to a barn with both indoor (approxi-

141 mately 15m×5 m) and outdoor space (approximately 20m×7 m), and with ad libitum access to water,

142 hay and concentrate food. The following day, lambs were individually tested in an arena test (morning)

143 and a corridor test (afternoon). At the end of each testing day, lambs returned to pasture with the with

144 the rest of the flock (see below for a summary of the tests [33,34] for a detailed description).

Arena test:

The test arena was a wooden pen (7m long and 2m wide), with the entrance for the tested lamb on one

side and an additional pen with three flock-mates separated by a grid and an optional wooden panel on

the other side. In a nutshell, each lamb was placed by an operator in the corridor and was exposed to

three consecutive phases: visual contact with the flock-mates (30s), no visual contact with the flock-

mates (isolation phase, 60s), visual contact with the flock-mates while a human stood between the

tested lamb and the flock-mates (reunion phase, 60s). Several behaviours (live data) were collected

during the test and used for the selection of the animals (number of low and high bleats, displacement).

The test was also video recorded to score the behaviour of each lamb in further details later on during

each phase.

Corridor test:

156 The test corridor was a non-opaque squared corridor (1m large and 25m long). Each lamb was placed

157 there by an operator and exposed to two phases: isolation (30s) and exposure to a moving human (60s).

158 The vocalisations were not used in the present study due too their lower quality, but the first phase of

159 the test was used for selection (see below).

2.3.2 Selection indexes and divergent lines

To summarize, the selection for social traits is based on the reactivity to social separation from con-

specifics, and the main variable used the selection is the number of bleats produced by the lamb during

the isolation and reunion phases of the arena test and during the isolation of the corridor test (see [29]

for a detailed description of the statistical procedure). A PCA was applied on the 3 variables corre-

sponding to the number of vocalisations emitted during the 3 phases of tests to construct a synthetic

variable (i.e. first principal component). For genetic selection of the divergent lines, the Social Selec-

tion Index (hereafter 'SSI'), corresponding to individual estimated breeding values, was calculated us-

ing genetic animal mixed model and considering this synthetic variable and pedigree information. Ex-

tremes individuals were chosen based on the SSI to constitute high and low divergent lines. The 42

most extremes individuals for the SSI were selected: the 21 individuals with the highest SSIs, compos-

ing the S+ line, *i.e* individuals highly reactive to separation from conspecifics, and the 21 individuals

with the lowest SSIs, composing the S- line, *i.e* individuals with low behavioural expression during
separation from conspecifics. Intermediate individuals were not selected.

Similarly, two additional divergent lines were selected for human reactivity (see detailed procedure in
[29]). A symmetrical pattern was used to constitute the H+ and H- lines by selecting individuals with
the highest and lowest Human Selection Indexes (hereafter 'HSI'), respectively expressing the highest
and lowest proximity to the human. The S+ and S- lines were selected for having simultaneously inter-
mediate values for the HSI.

The divergent selections had been carried out for 11 years at the time of the study. Considering the
breeding cycle of the sheep and overlaps between generations, this study is composed of animals after
2 (19%), 3 (53%) or 4 (22%) and 5 (6%) generations. Heritability of sociability and human reactivity
traits have already been quantified and are available in [29], including the behavioural profiles of S+
and S- sheep. Overall, S+ lambs produce significantly more calls, have a higher locomotion and main-
tain closer proximity towards conspecifics than S- lamb [29].

185 **2.4 Recording contexts and analysis of vocalisations structure**

186 *2.4.1 Isolation bleats*

An omnidirectional microphone (Sennheiser MK20), connected to a recorder (Marantz PMD670, fre-
quency sampling 44100 Hz) was placed on the top of the arena at 2.5m from the ground (between 2.3
and 5m from the lamb). Calls, produced during isolation phase of the arena test, when the lamb was
fully separated from the flock mates, were recorded. The context of production of these calls is thus a
negative situation of relatively high arousal, with regards to the gregarious behaviour of sheep. In total,
152 females were tested and their bleats recorded.

193 *2.4.2 Pre-feeding bleats*

Eight groups composed of lambs of each line (17S+ and 20S- lambs) were given a barley reward right
after shaking a metal barley box every morning for a week, resulting in an association between the
sound and reward. Every day of the four following weeks, vocalisations produced before the delivery
of the barley treat were recorded by the experimenter at group level using an omnidirectional micro-
phone (Sennheiser MK20), connected to a recorder (Marantz PMD670), while located at 3 to 15 meters
from the lamb. The identity of the caller was assigned when possible by the experimenter. The context
of production of these calls could be a food anticipation context, which we assume as positive and

lower in arousal than the social isolation [35,36]. However, the valence of this context was not vali-
dated using emotion indicators.

2.4.3 Acoustic analysis of high bleats

Only high bleats, which were consistently produced in both contexts and by the two divergent lines,
were analysed in the present study (hereafter 'bleats'). Isolation and pre-feeding bleats were analysed
separately using the same methodology. All calls of good quality (no overlapping sound) were kept for
acoustic analysis. For the isolation context, 1287 bleats were analysed; 322 bleats from 21 S+ females,
67 bleats from 14 S- females, and 898 bleats from 117 females that were recorded but not selected as
S+ or S- due to their intermediate SSI (hereafter 'non-selected'). Regarding the pre-feeding context,
519 bleats were analysed; 262 bleats from 14 S+ females (including 64 from unknown callers), and 257
bleats from S- females (including 52 from unknown caller). The process for extracting acoustic
parameters was automatically computed using the *extract_chunk_features_dir()* function from
SoundChunk R-package [37], which compiles several low level functions of the Seewave and tuneR R
package [38–40] (details on computation are available as supplementary material (section 'Additional
content > Acoustics')). Sixteen acoustic parameters were extracted to describe the spectro-temporal
features of the bleats: duration, energy quartiles (Q25, Q50, Q75 and IQR), spectral flatness, dominant
frequency over the duration of the call (mean and standard deviation), fundamental frequency over the
duration of the call (mean and standard deviation), mean and relative amplitude of the three first
formants (table S1).

It has to be noted that the environment of the two recording contexts (indoor vs. outdoor) prevented us
from running direct comparison of call structure between contexts.

2.5 Statistics

223 All statistics were performed in R version 4.4.0. [34]

2.5.1 Investigation of acoustic difference between socially selected lines

Two complementary approaches were used: on the one hand, a Linear Discriminant Analysis (LDA) on
bleats from selected S+ and S- sheep was used built an acoustic score describing bleat structure and
extract key potentially discriminant acoustic parameters. Predictions of linear discriminant functions on
non selected animals allowed to test for their intermediate score using a linear mixed effect model
(LMM). On the other hand, a permuted Discriminant Function Analysis (pDFA) was used to test for
correct classification of the genetic line of the sheep based on the acoustic structure of their bleats.

<u>Linear discriminant function analysis and test for intermediate acoustic structure in bleats from non selected females.</u>

A LDA including the 16 acoustic parameters extracted from the bleats and with line as a factor was performed to search for the acoustic parameters more likely to discriminate the calls from S+ and S- lambs (*lda()* function from 'MASS' R package [41], no warning for collinearity between parameters was raised). One Acoustic Score (LD) was computed for the isolation calls (isoLD), and one for the pre-feeding calls (feedLD), by entering the isolation calls and pre-feeding calls in separate LDAs. The LDA ran on the isolation calls from S+ and S- lambs was then used to predict the Acoustic Score of the calls produced by non-selected females (*predict()* function from 'MASS' R package).

Following the LDA, a LMM (*lmer()* function, 'lme4' R package [42]) was built using the isoLD as response variable, in order to test for intermediate acoustic structure of non selected females. Two interacting fixed effects were used: 'group_line', a categorical factor with three levels (the two sheep lines (S+ or S-) and the group predicted by the LDA using the non-selected lambs) in interaction with the weight of the lamb ('Zweight', centered and reduced covariate, i.e. 'scaled'). The identity of the genetic father and identity of the animal were used as a nested random factor (animal ID within father ID):

```
isoLD ~ group_line*Zweight + (1|father_ID/animal_ID), data = all_females
```

<u>Permuted Discriminant Function Analysis</u>

Focusing on selected lambs only, pDFAs were computed within each recording context to test for discrimination of the line respectively in isolation calls and in pre-feeding calls. The statistical approach is described in [43] and the code used here was developed in R by Roger Mundry using the R 'MASS' package (built mainly using functions *lda()* and *predict()*).

To test for acoustic discrimination according to the factor line, nested pDFAs were computed within each recording context, with N=100 random selection and cross classification and N=1000 permutations, to generate p-values for correct classifications in general and within each group (confusion matrix of p-values). Two pDFA per context were ran: the first pDFA tested for correct classification of all calls without taking into account the individual identity of the emitter (so that all calls could be kept for analysis, even from unknown callers in pre-feeding calls and from individuals that had produced less than four calls in isolation). The second pDFA controlled for the individual identity of the lambs (removing calls produced by unknown callers and individuals having produced less than 4 calls). For the pDFA controlling for individual identity, only females that had produced at least four calls were kept to allow selection of three calls (training dataset) and cross classification on the remaining one(s). For

isolation calls, the first pDFA included calls from 21 S+ and 14 S- females, with median number of

calls per female [first;last decile]=9[2;16], while the second pFDA included 21 S+ and eight S- females

with median number of calls per female [first;last decile]=9[2;16]. For pre-feeding calls, the first pDFA

inclluded calls from 12 S+, 10 S- females, among which, 116 calls out of 519 were produced by un-

known callers but known line. The second pDFA included calls from seven S+ and nine S- females that

had produced at least four calls, with median number of calls per female [first;last decile] = 12[4;34].

2.5.2 Study of predictors of acoustic structure of isolation bleats

To study predictors of the acoustic structure of the isolation bleats, we designed a LMM including the

Acoustic Score (isoLD) as a response variable and several predictors as fixed effects. Since the Acous-

tic Score was built to discriminate calls from S+ and S- females, data from these selected sheep were

excluded and the LMM was run on predicted isoLD corresponding to the calls of non-selected females

only. According to our two hypotheses, isolation bleats structure may be driven by 1) the arousal level

expressed during the recording (emotional hypothesis) and/or 2) the selection for sociability (socio-

acoustic co-selection hypothesis). To disentangle between hypotheses, proxies of arousal and social

behaviour were extracted from scoring the videos of the isolation and reunion phases to be tested along

with the SSI. The detailed description of the procedure is available as supplementary material (section

'Behavioural profiles', table S2 and figure S1). In a nutshell, Principal Component Analyses allowed us

to compute behavioural scores (PCs) that described the behaviour of S+ and S- sheep, using behaviour

that could be symmetrically scored in both phases: mobility, interest toward either conspecifics or the

rest of the environment and attraction toward conspecifics. Three behavioural scores showed signifi-

cant difference or a trend between S+ and S- sheep and were kept as potential predictors of bleats struc-

ture: IsoPC1 was a proxy of vigilance and mobility (i.e. bodily activation, arousal), ReuPC1 was a

proxy of interest toward conspecifics and ReuPC3 was a proxy of attraction toward conspecifics (see

loadings of PCs in table S2). As controls, we were first able to partially control for potential heritability

of vocal apparatus using the identity of the father (10 fathers) as random factor in the LMM. Second,

based on the source-filter theory of sound production, bigger lambs are expected to produce calls con-

taining lower frequencies [44], which could be a strong predictor of bleat acoustic structure. Hence, the

weight of the lamb was used as interacting covariate in our LMM. Third, to control for effect of the

diverging selections, the HSI was also included in our LMM, as a negative control for selections.

Therefore, the full model was composed of the following fixed effects: isoPC1 (proxy of the arousal of

the lamb in isolation, emotional hypothesis), reuPC1 and reuPC3 (proxy of social interest and attraction

toward conspecifics, socio-acoustic co-selection hypothesis), the SSI and HSI (proxy of genetic selec-
tion for sociability and selection on human reactivity, socio-acoustic co-selection hypothesis), and the
scaled weight of the lamb 'Zweight' (proxy of the size of the lamb, biometric effects) in interaction
with all other factors. The random factor included the identity of the animal nested within the identity
of the father (partially controlling for heritability of vocal apparatus structure and repetitions of several
calls from the same individual):

```
300  isoLD ~ lmer (ZWeight *(isoPC1 + reuPC1 + reuPC3 + SSI  + HSI) + (1|father_ID/ani-
301  mal_ID), data = (intermediate) non selected females)
```

302 *2.5.3 Impact of genetic selection for sociability on encoding of individuality in*
303 *bleats*

To test the impact of the genetic line on identity coding in isolation and pre-feeding bleats, nested
pDFAs were computed within each context of recording to extract scores of correct classification ac-
cording to the identity of the caller.

For the isolation bleats, the calls of 124 females (selected or not) were used in the analysis, and permu-
tations for testing call classification according to individuals were restricted to within father identity
([min;max] = [1;18] females per father for permutations). For the pre-feeding bleats, the calls of 16
females were used in the analysis, and due to the lower sample size ([min;max] = [1;3] females per
father), no restriction for permutations within father identity was applied. The analysis confirmed that
the set of acoustic parameters used in this study allowed correct classification according to the caller, in
both contexts (supplementary table S2), as previously shown in sheep on a different set of parameters
[24]. The ratio of correctly cross-classified bleats, in each context, was then extracted from the diagonal
of the prediction/reference confusion matrix as a proxy of success of cross classification of calls ac-
cording to the identity of the caller (the higher the ratio, the higher the individuality) and used as re-
sponse variable after square-root transformation in a LMM to test the effect of the genetic line (and
predictors cited above when possible).

The LMM built for the isolation bleats included the same predictors as above (2.5.2) to explain the cor-
rect classification score: the SSI, isoPC1, reuPC1, reuPC3 and the weight of the lamb as interacting
scaled covariate ('Zweight'). Arousal level may decrease identity coding in vocalisations [30]: if iden-
tity coding was affected by the level of arousal expressed during the test, then we expected a negative
relationship between the ratio of correctly classified calls and isoPC1 (emotional hypothesis). If iden-
tity coding was affected by the selection for sociability and social behaviour, then we expected a rela-

tionship between the ratio of correctly classified calls and reuPC1, reuPC3 and/or the Social Selection

Index (socio-acoustic co-selection hypothesis). The father identity was included as a random effect:

```
327 sqrt(ratio_of_correctly_cross_classified_bleats) ~ lmer(ZWeight *(isoPC1 + reuPC1
328 + reuPC3 + SocialSelectionIndex) + (1|father_ID))
```

329 For the LMM built for the pre-feeding bleats, only selected females were recorded and no behavioural

330 data was available, hence we included only the line (S+ vs. S-) in interaction with the weight of the

331 lamb as an interacting scaled covariate ('Zweight'), along with father identity as a random effect:

```
332 sqrt(ratio_of_correctly_cross_classified_bleats) ~ lmer(ZWeight *line + (1|
333 father_ID))
```

334 For both models, the response variables were transformed using a square root function to reach symet-

335 rical distribution prior to modeling.

2.5.4 Model inspection and statistical tests on LMM

337 After visual inspection of the symmetrical distribution of the residuals, a type II Wald Chi-Square test

338 on the full model was computed to test the effect of the predictors (*Anova()* function, 'car' R package).

339 Slope estimates were extracted with *emtrends()* function ('emmeans' R package, [45]). When interac-

340 tions between covariates were significant, pairwise comparisons of slopes were computed, fixing one

341 covariate at mean, mean-se, mean+se to test the variation of the slope of the other covariate.

3 Results

3.1 Differences in acoustic signature of lambs selected for sociability

According to the loadings of acoustic parameters on the first linear discriminant function (LD) ex-

tracted from a LDA maximizing the difference between the two lines, the median frequency (Q50),

mean fundamental frequency (fo_mean), the mean frequency of the second formant and its relative

amplitude (F2_mean and F2_amp) consistently loaded highly on LD in bleats recorded in both contexts

(table 1 and figure 1A, 1B). Other parameters loaded highly on the LD in one of the contexts but not

the other: duration and mean dominant frequency (dfreq_mean) loaded more strongly on isoLD, carried

out on isolation bleats, whereas the spectral flatness (sfm), the energy distribution (Q25, Q75 and IQR)

and the variability of fundamental frequency (fo_sd) loaded more on the feedLD, carried out on pre-

feeding bleats (figure 1B). For isolation bleats, the projection of the acoustic parameters of bleats pro-

duced by non-selected females on the linear discriminant function describing differences between S+

and S- bleats (isoLD) showed that non selected females had intermediate values (LMM, Chisq = 23.94,

p < 0.001, figure 1C) between the bleats from S+ and S- females.

Table 1: Loadings of acoustic parameters on the first linear discriminant function (LD) extracted from the linear discriminant function analysis (LDA) performed on isolation calls, leading to the acoustic score 'isoLD', and on LD extracted from the LDA performed on pre-feeding calls, leading to the acoustic score 'feedLD'. In bold, loading values having absolute relative loading above 0.4, considered as descriptors of the linear discriminant function. Underligned acoustic parameter names illustrate consistently high loadings across both contexts, bold loading value illustrate parameters loadings significantly on LD in one of the context.

Acoustic parameter	Loading on isoLD	Loading on feedLD
duration	**0.515**	-0.174
Q50	**-0.745**	**-0.640**
Q25	0.131	**-0.526**
Q75	0.161	**0.895**
IQR	0.112	**-1.000**
sfm	-0.172	**0.932**
dfreq_mean	**1.000**	-0.036
dfreq_sd	**-0.866**	-0.051
fo_mean	**-0.895**	**0.488**
fo_sd	-0.260	**-0.545**
F1_mean	-0.087	-0.316
F2_mean	**-0.952**	**0.656**
F3_mean	0.129	-0.132
F2_amp	**0.926**	**0.639**
F3_amp	**-0.430**	-0.175

NB: dfreq_mean and dfreq_sd for pre-feeding LDA were square-root transformed

Figure 1: Acoustic differences in isolation bleats (left side of panel) and in pre-feeding bleats (right side of panel) according to the 'line' (S+, S- and intermediate not selected when considering the isolation calls). A: Spectrogram of calls of one representative S+ female and one representative S- female according to the context of recording (FFT wl = 2048 pts, ov= 90%, frequency sampling=44.1 kHz, amplitude range from blue to red color scale: -70 to 0dB). B: Averaged mean-spectrum (and standard deviation in shade) over the total amount of calls analyzed per line and contexts (Number of isolation calls: N=322 from S+, N=67 from S-, N=899 from not selected females; Number of pre-feeding calls: N=262 from S+, N=257 from S- females). C: Acoustic parameters explaining the Acoustic Score computed using a LDA on 'line' within each context of recording (isoLD for isolation and pre-feedLD for pre-feeding). For isolation calls, the predictions of isoLD for calls recorded in non-selected females and comparisons (groups with different letters illustrate statistical differences, see supplementary statistical tables S3 for model estimates and pairwise comparisons).

359 The pDFA on isolation and pre-feeding bleats showed that, without controlling for individual identity,

360 the correct classification according to the line was significant on the training data set, i.e. selected calls

361 (24% and 12% correct classification above percentage expected at random for isolation and pre-feeding

362 calls, respectively; p = 0.001 for both; table 2A). However, correct cross classification (on the validat-

ing dataset) was only significant for isolation bleats and not pre-feeding bleats (31%, p = 0.001 vs. 15%, p = 0.223 correct classification above percentage expected at random for isolation and pre-feeding calls, respectively; table 2A). Since the results could be driven both by the line and individual vocal signatures of the lamb belonging to each line, a second analysis was performed, controlling for repeated measures of the same individuals. The analysis showed a trend for correct classification for isolation bleats according to the line (p = 0.105 and p = 0.084 for correct classification of calls using the training and validating (cross classification) dataset respectively, table 2A). Although the significance 0.05 threshold was not reached for cross classification in general, the confusion matrix of p-values showed significant discrimination of S+ bleats for isolation calls (table 2B). For pre-feeding bleats, no evidence of correct classification of the line was found neither considering classification scores (p = 0.501 and p = 0.435 for correct classification of calls according to the line using the training and validating (cross classification) dataset respectively, table 2A), nor inspecting the confusion matrix of p-values for each line (table 2B).

Table 2: Call classification according to the line (S+/S-) using pDFA in isolation bleats and in pre-feeding bleats. A: result of pDFA, percentage of classified and cross classified calls and p-values computed according to permuted (random) classification. B: confusion matrix of classification p-values for S+ and S- calls in each context.

	Isolation bleats		Pre-feeding bleats	
A. Results of pDFA on discrimination of calls from line S+ and S-				
Control factor (for selection)	X	Individual	X	Individual
Number of levels of control factor selected out of total	X	14/29	X	12/16
Correct classification of selected calls (%)	88.02	93.52	68.41	85.04
Expected correct classification of selected calls (%)	64.23	85.47	56.60	84.36
P value for classification of selected calls	**0.001**	**0.105**	**0.001**	**0.501**
Correct cross classification (%)	80.01	68.91	64.43	56.75
Expected correct cross classification of calls (%)	49.84	58.21	49.69	55.74
P value for cross classification of calls	**0.001**	**0.084**	**0.223**	**0.435**
B. Confusion matrices of P values of cross classification				

Reference \ Prediction	S-	S+	S-	S+	S-	S+	S-	S+
S-	0.476	0.505	0.443	0.913	0.515	0.486	0.370	0.598
S+	1	**0.001**	0.947	**0.039**	0.875	0.126	0.586	0.459

3.2 Drivers of acoustic signature of isolation bleats

A significant positive relationship between the SSI and the acoustic score isoLD was found (LMM: Chisq = 8.15 p = 0.004, slope estimate [95%CI] = 0.36[0.10;0.63], figure 2), without any interaction with the weight of the lamb (LMM: Chisq = 0.31 p = 0. 576). Based on the acoustic loadings (table 1), this suggests that non-selected lambs having SSI closer to S+ sheep produced calls with higher mean dominant frequency, longer duration, and a more pronounced second formant. A significant interaction was found between the weight of the lamb and reuPC3 (LMM: Chisq = 4.29 p= 0.04). Further posthoc tests revealed that there was a positive relationship between reuPC3 and the acoustic score isoLD, especially for heavier lambs (lstrends posthoc tests, slope estimates [95%CI]: 0.27[0.06;0.48] and 0.13[-0.01;0.26] respectively for high and medium), but not for lighter lamb (-0.02[-0.19;0.16], figure 2). According to the loadings of the behavioural PCA (table S2), this suggests that lambs - at least heavy and medium weight ones - exhibiting closer proximity with their conspecifics in the reunion that followed the isolation phase produced bleats with a higher mean dominant frequency, longer duration, and a more pronounced second formant during isolation. There was no evidence for any effect of the behavioural scores isoPC1 - correlated with bodily activation during the isolation - or reuPC1 - correlated with exploration and bodily activation during the reunion - on isoLD (LMM: Chisq < 2.3 p > 0.13, supplementary table S6). There was also no evidence of any effect of the HSI on isoLD (LMM: Chisq = 0.006 p = 0.94, supplementary table S6).

Figure 2: Model estimates and 95% confidence interval depicting the relationship between the Acoustic Score describing isolation bleat structure (isoLD) of non-selected females (hence having intermediate social selection score compared to selected animals) and (A) the Social Selection Index or (B)the behavioural score reuPC3, which positively correlated with proximity. For reuPC3, a significant interaction was found with the weight of the lamb, so slope estimated are showed based on the mean, mean + se and mean – se of the scaled weight used as covariate. Stars (*) show significant slopes. Statistical results of Anova() and model estimates are available as supplementary material tables S6 and S7 repsectively.

3.3 Consequence of genetic divergence for sociability on individuality coding in bleats

For isolation bleats, using 122 females that produced more than four calls during the arena test, the SSI was negatively correlated with the individuality in calls (measured as the ratio of correctly cross-classified calls in the pDFA) (LMM: Chisq = 4.6 p = 0.03); higher SSI had slightly lower ratios of correctly cross-classified calls (slope estimates [95%CI]: -0.04[-0.09;0.002], figure 3). No evidence of any effect of other potential predictors (weight as interacting covariate of isoPC1, reuPC1, reuPC3) were found (LMM: Chisq < 1.59 p > 0.21, table S8). For pre-feeding bleats, using the 16 females that produced more than four calls during the pre-feeding sessions, no evidence of any effect of the line nor the

404 weight was found on the ratio of correctly cross-classified calls (LMM: Chisq < 0.95 p > 0.33, table
405 S8).

Figure 3: Relationship between the ratio of correctly cross classified calls (proxy of individuality in spectro-temporal features of bleats) and the Social Selection Index. Model estimate (line) and 95% confidence interval represented (shade). Each point represents individual estimated value. *: significant slope. Statistical results of Anova() are available as supplementary material tables S8.

Discussion

High bleats of lambs selected for high (S+) or low (S-) sociability (i.e. reactivity to social isolation)
were recorded in emotional contexts, namely isolation and pre-feeding contexts. Our acoustic analysis
showed that several acoustic parameters, such as median frequency, fundamental frequeny, frequency
and relative amplitude of second formant , consistently contributed to the difference between the S+
and S- lines across contexts, while others were more context specific. Correct classification of calls
according to the line was significant only for isolation bleats, particularly S+ bleats, and not pre-feed-
ing bleats. This suggests that a sociability related 'line acoustic signature' may arise in isolation calls
only. The Social Selection Index (SSI) and proximity to conspecifics expressed during the reunion
phase were predictors of the acoustic structure of isolation bleats, which was not the case of emotional
arousal. These results suggest that the temperament trait sociability, more than emotional arousal per
se, influenced sheep vocalisations structure in our study, especially in the context of an isolation test
where such trait is strongly expressed. This may reveal that specific acoustic markers are co-selected
when selecting for sociable animals. Last, the results showed that the SSI negatively impacted the en-
coding of individual's identity in isolation bleats: highly sociable animals tended to produce isolation
bleats with lower probability of being correctly attributed to the caller. Overall, this suggests that tem-

perament traits such as sociability strongly impact on vocalisations of farm animals such as sheep, pre-
vailing in some contexts over the effect of emotional arousal and impacting on vocal individuality.

Context specific divergence of acoustic structure of bleats in sociable and less sociable sheep lines

The results from the acoustic analysis showed that common acoustic parameters were involved in the
discrimination of high bleats produced by female lambs belonging to two divergent genetic lines based
on social traits (S+, highly sociable and S-, less sociable): mean fundamental frequency, median of the
mean spectrum, mean value of second formant and its relative amplitude. Based on the source-filter
theory of vocal production, vocal signals result from a two-stage production, with the glottal wave gen-
erated in the larynx (the source) being subsequently filtered in the supralaryngeal vocal tract (the filter)
[38]. Here, we show that both source (fundamental frequency, median of the mean spectrum) and filter
(formants) parameters may participate in the discrimination of the divergent lines consistently in both
contexts. It is likely that the anatomy of the larynx and vocal tract may be influenced by genetic charac-
teristics (heritability of structures of the vocal apparatus), leading to diverging vocal signature between
sheep of the two lines. If this was the only mechanism underlying the acoustic difference we found
between genetic lines, we would have expected a correct classification of bleats according to the line
regardless of the recording contexts. On the contrary, only isolation bleats could be correctly classified
according to the line and not the pre-feeding bleats. The 'line acoustic signature' is hence context-spe-
cific, suggesting that selection (in general) and heritability of structures of the vocal apparatus cannot,
alone, explain the acoustic difference in bleat structure revealed in our study

This conclusion may find support in the fact that other acoustic parameters discriminated between the
two lines in one context but not the other, showing some flexibility. Indeed, dominant frequency mean
and its standard deviation, relative amplitude of third formant and duration were important parameters
for distinguishing S+ from S- in isolation bleats, but not for pre-feeding bleats. As sheep are highly
gregarious, isolation from conspecifics was assumed to be a negative situation of high arousal (provok-
ing high bodily activation [46]). In several species, stress involves the production of calls with longer
durations and higher frequencies [3–5]. Here, S+ lamb, which expressed a higher behavioral reaction to
isolation (see also [29]), also produced high bleats with higher dominant frequency and longer duration.
Hence, our acoustic results are in line with the hypothesis that S+ lambs may be more stressed in social
isolation than S- lambs. However, specific changes in acoustic structure of bleats as a function of the

emotional valence and arousal of the producer are still lacking for sheep, and this should be further
validated with standard tests [5].

Different parameters played an important role in distinguishing high bleats from S+ and S- lamb in pre-
feeding context: quartiles of the mean spectrum and its the inter-quartile range as well as the spectral
flatness (noisiness), but the classification of these calls according to the line was not above chance
level. The pre-feeding context may represent an emotional state of intermediate arousal (walking or
running), but characterized by a positive valence (pleasant), and not related to social behaviour like in
the case of the social isolation [35,36]. In fact, it is possible that the selection process, based on socia-
bility, leads to a stronger selective pressure on isolation bleats than on pre-feeding bleats, explaining
why the difference emerges in isolation bleats, especially in calls from highly sociable lambs.

Beyond social stress: vocal encoding of heritable sociability in isolation bleats

To investigate whether the above-described 'line acoustic signature' is driven by emotional arousal or
other factors, our model tested predictors related to the social selection process, the social behaviour
expressed during the test and emotional arousal, while controlling for the weight of the animal. This
allowed us to potentially disentangle between two hypothesis that could explain difference in bleat
acoustic structure between our two lines: the emotional hypothesis and the socio-acoustic co-selection
hypothesis. Results showed that, although the emotional arousal during isolation was clearly different
between the two lines, with S+ being more active, there was no evidence of a relationship between our
arousal proxy and the acoustic score describing isolation bleats structure. Instead, the SSI and spatial
proximity to conspecifics during the reunion that immediately followed the isolation were the main
predictors of the acoustic score describing the isolation bleat structure. Our results are therefore in fa-
vor of the socio-acoustic co-selection. In other words, selecting for sociability, based on behavioural
reaction to isolation, likely also selected for individuals more motivated to be closer to conspecifics and
for individuals producing isolation bleats having a typical vocal signature or expressing emotional
arousal/motivation to reunite with conspecifics to a higher extent than predicted by their behaviour.
Indeed, temperament has been shown to affect expression of emotions in other ungulates such as
horses, in which proactive horses appear more stressed based on their behaviour compared to their
physiology [8]. In genetic selections, selection for a trait might co-select another one, with several ex-
amples in poultry reviewed by Jones and Hocking (1999) [47]. We show here a possible vocal encod-

ing of sociability, a heritable temperament trait linked to emotional reactivity during social separation
and to the motivation to associate with others.

Implications of a socio-acoustic co-selection for social cognition

Our results confirmed that an individual vocal signature is encoded in high bleats, in accordance with
previous studies [24,25]. Interestingly, based on the acoustic parameters used here, the classification of
calls to the correct caller was slightly impacted by the SSI: the higher the index, and thus the higher the
sociability, the lower the ratio of isolation bleats correctly classified to the caller. Several hypotheses
have been proposed to explain individual distinctiveness in calls, summarized in [48]. Based on the
'distance communication hypothesis', individually in long distance calls should be higher than in short
range ones, for which identity of the caller may be encoded by other modalities (e.g. visual or olfactory
[49]). In our study species, high bleats may be considered as rather long-distance calls: contrary to low
bleats, they are produced at higher amplitude and when an individual loses contact with conspecifics
(however, short separation distance may also trigger the production of high bleats in experimental set-
tings) [25]. In accordance to the distance communication hypothesis, high bleats encode more individu-
ality than low bleats, at least in adult female sheep [25]. Our results revealed a rather high individuality
in the high bleats, as shown by high cross classification scores in general. In addition, higher scores
were found in isolation bleats than pre-feeding bleats, which may be in line with the distance communi-
cation hypothesis, since during isolation, sheep were at further distance from conspecifics. Secondly,
based on the 'social hypothesis', individuality in call structure should be driven either by the context or
the receiver the call is intended to: calls addressed to a specific individual are predicted to display
higher individuality than calls directed to a group or produced in a general context (food, alarm calls,
distress) [50,51]. In baby cries for example, individual signatures in pain cries are less reliable than in
discomfort cries and it was hypothesized that low individuality in distress calls produced within a so-
cial group function to trigger the attention of the entire group rather than to the caretaker only [30].
Recently, using a database of contact calls from seven ungulate species, Osiecka et al. showed that in-
dividual distinctiveness in calls was impaired in negative contexts compared to positive ones of similar
arousal [31], and hypothesized that calls produced in negative contexts may be addressed to a broader
audience than those produced in affiliative contexts. Our results show here that, within the same nega-
tive context (social isolation) and beyond social stress, encoding of sociability also impairs individual-
ity in high bleats: calls from more sociable animals carry less individuality. Calls produced by more
sociable animals in a context of social separation, where they are highly motivated to reunite with con-

specifics, may thus be perceived by multiple receivers and increase the probability of triggering a response from at least one group member, independently from the identity of the caller. If this hypothesis is true, it could have implications for social cognition, as it might impair individual recognition (mother-lamb or social partners) of more sociable sheep based on their high bleats. More research is needed to investigate the consequences of vocal encoding of sociability for communication and its interplay with vocal perception of emotions.

Conclusion

To our knowledge, our study provides the first evidence of the consequences of genetic selection for sociability traits on vocal communication, using sheep as a model species. Our findings lead us to suggest a socio-acoustic co-selection: selecting for sociable animals leads to exaggerated vocal expressions of emotional arousal during social isolation and impairs vocal signatures at the group and individual level. The phenomenon we quantified opens the question of the long-term consequences of such selection on social cognition. The divergent lines we describe here also provide a model to study the co-evolution between social traits and social and emotional communication in a gregarious species and explore welfare consequences.

Acknowledgments and funding

The authors would like to thank Sara Parisot and Charlotte Allain for the support of INRAE Experimental Unit of La Fage, Monica Padilla de la Torre for participating to data collection and Eric Delval (UMRH) for scoring the videos of the arena test. This study was funded by a grant from Carlsberg Foundation attributed to Elodie F. Briefer (CF20-0538).

Authors' contribution

ASV designed the study, collected the acoustic data, ran the analyses and drafted the manuscript, AB designed the behavioural tests, discussed the design of the study and revised the manuscript, GB, CD and SD carried out the behavioural tests, MMM discussed the design of the study and revised the manuscript, PRG collected the acoustic data, ran the analyses and revised the manuscript, DH designed and computed the genetic selections and revised the manuscript, EFB designed the study, revised the manuscript and funded the study.

540 *References*

1. Charlton BD, Pisanski K, Raine J, Reby D. 2020 Coding of Static Information in Terrestrial Mammal Vocal Signals. In *Coding Strategies in Vertebrate Acoustic Communication* (eds T Aubin, N Mathevon), pp. 115–136. Cham: Springer International Publishing. (doi:10.1007/978-3-030-39200-0_5)

2. Bradbury JW, Vehrencamp SL. 2011 *Principles of Animal Communication*. Second. See http://www.sinauer.com/principles-of-animal-communication.html.

3. Briefer EF. 2020 Coding for 'Dynamic' Information: Vocal Expression of Emotional Arousal and Valence in Non-human Animals. In *Coding Strategies in Vertebrate Acoustic Communication* (eds T Aubin, N Mathevon), pp. 137–162. Cham: Springer International Publishing. (doi:10.1007/978-3-030-39200-0_6)

4. Briefer EF. 2012 Vocal expression of emotions in mammals: mechanisms of production and evidence. *J Zool* **288**, 1–20. (doi:10.1111/j.1469-7998.2012.00920.x)

5. Villain AS, Briefer EF. in production Vocal signals as indicators of emotions. In *Assessment of Animal Welfare - a guide to the valid use of indicators of affective states*, United Kingdom: John Wiley & Sons Ltd, Oxford,.

6. Hughes DJ, Kratsiotis IK, Niven K, Holman D. 2020 Personality traits and emotion regulation: A targeted review and recommendations. *Emotion* **20**, 63–67. (doi:10.1037/emo0000644)

7. Keltner D. 1996 Chapter 21 - Facial Expressions of Emotion and Personality. In *Handbook of Emotion, Adult Development, and Aging* (eds C Magai, SH McFadden), pp. 385–401. San Diego: Academic Press. (doi:10.1016/B978-012464995-8/50022-4)

8. Squibb K, Griffin K, Favier R, Ijichi C. 2018 Poker Face: Discrepancies in behaviour and affective states in horses during stressful handling procedures. *Applied Animal Behaviour Science* **202**, 34–38. (doi:10.1016/j.applanim.2018.02.003)

9. Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. 2007 Integrating animal temperament within ecology and evolution. *Biological Reviews* **82**, 291–318. (doi:10.1111/j.1469-185X.2007.00010.x)

10. Carere C, Welink D, Drent PJ, Koolhaas JM, Groothuis TGG. 2001 Effect of social defeat in a territorial bird (*Parus major*) selected for different coping styles. *Physiology & Behavior* **73**, 427–433. (doi:10.1016/S0031-9384(01)00492-9)

11. Brunelli SA, Hofer MA. 2007 Selective breeding for infant rat separation-induced ultrasonic vocalizations: Developmental precursors of passive and active coping styles. *Behavioural Brain Research* **182**, 193–207. (doi:10.1016/j.bbr.2007.04.014)

12. Faure JM, Mills AD. 2014 Chapter 8 - Improving the Adaptability of Animals by Selection. In *Genetics and the Behavior of Domestic Animals (Second Edition)* (eds T Grandin, MJ Deesing), pp. 291–316. San Diego: Academic Press. (doi:10.1016/B978-0-12-394586-0.00008-1)

13. Forkman B, Boissy A, Meunier-Salaün M-C, Canali E, Jones RB. 2007 A critical review of fear tests used on cattle, pigs, sheep, poultry and horses. *Physiology & Behavior* **92**, 340–374. (doi:10.1016/j.physbeh.2007.03.016)

14. Koolhaas JM, Korte SM, De Boer SF, Van Der Vegt BJ, Van Reenen CG, Hopster H, De Jong IC, Ruis MAW, Blokhuis HJ. 1999 Coping styles in animals: current status in behavior and stress-physiology. *Neuroscience & Biobehavioral Reviews* **23**, 925–935. (doi:10.1016/S0149-7634(99)00026-3)

15. Gartland LA, Firth JA, Laskowski KL, Jeanson R, Ioannou CC. 2022 Sociability as a personality trait in animals: methods, causes and consequences. *Biological Reviews* **97**, 802–816. (doi:10.1111/brv.12823)

16. Šlipogor V, Gunhold-de Oliveira T, Tadić Z, Massen JJM, Bugnyar T. 2016 Consistent inter-individual differences in common marmosets (Callithrix jacchus) in Boldness-Shyness, Stress-Activity, and Exploration-Avoidance. *American Journal of Primatology* **78**, 961–973. (doi:10.1002/ajp.22566)

17. Naguib M, Schmidt R, Sprau P, Roth T, Floercke C, Amrhein V. 2008 The ecology of vocal signaling: male spacing and communication distance of different song traits in nightingales. *Behav. Ecol.* **19**, 1034–1040. (doi:10.1093/beheco/arn065)

18. Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G. 2012 How not to measure boldness: novel object and antipredator responses are not the same in wild baboons. *Animal Behaviour* **84**, 603–609. (doi:10.1016/j.anbehav.2012.06.015)

19. Friel M, Kunc HP, Griffin K, Asher L, Collins LM. 2016 Acoustic signalling reflects personality in a social mammal. *Royal Society Open Science* **3**, 160178. (doi:10.1098/rsos.160178)

20. Nelson XJ, Wilson DR, Evans CS. 2008 Behavioral Syndromes in Stable Social Groups: An Artifact of External Constraints? *Ethology* **114**, 1154–1165. (doi:10.1111/j.1439-0310.2008.01568.x)

21. Leliveld LMC, Düpjan S, Tuchscherer A, Puppe B. 2017 Vocal correlates of emotional reactivity within and across contexts in domestic pigs (Sus scrofa). *Physiology & Behavior* **181**, 117–126. (doi:10.1016/j.physbeh.2017.09.010)

22. Guidi A, Gentili C, Scilingo EP, Vanello N. 2019 Analysis of speech features and personality traits. *Biomedical Signal Processing and Control* **51**, 1–7. (doi:10.1016/j.bspc.2019.01.027)

23. Dwyer CM. 2008 *The welfare of sheep*. Dordrecht: Springer.

24. Sèbe F, Poindron P, Ligout S, Sèbe O, Aubin T. 2018 Amplitude modulation is a major marker of individual signature in lamb bleats. *Bioacoustics* **27**, 359–375. (doi:10.1080/09524622.2017.1357146)

25. Sèbe F, Duboscq J, Aubin T, Ligout S, Poindron P. 2010 Early vocal recognition of mother by lambs: contribution of low- and high-frequency vocalizations. *Animal Behaviour* **79**, 1055–1066. (doi:10.1016/j.anbehav.2010.01.021)

26. Sèbe F, Nowak R, Poindron P, Aubin T. 2007 Establishment of vocal communication and discrimination between ewes and their lamb in the first two days after parturition. *Developmental Psychobiology* **49**, 375–386. (doi:10.1002/dev.20218)

27. Sebe F, Ligout S, Porter R. 2004 Vocal discrimination of kin and non-kin agemates among lambs. *Behaviour* **141**, 355–369. (doi:10.1163/156853904322981905)

28. Atkinson L, Doyle RE, Woodward A, Jongman EC. 2022 Behavioural reactivity testing in sheep indicates the presence of multiple temperament traits. *Behavioural Processes* **201**, 104711. (doi:10.1016/j.beproc.2022.104711)

29. Hazard D *et al.* 2022 118. Divergent genetic selections for social attractiveness or tolerance toward humans in sheep. In *Proceedings of 12th World Congress on Genetics Applied to Livestock Production (WCGALP),* pp. 520–523. Rotterdam, the Netherlands: Wageningen Academic Publishers. (doi:10.3920/978-90-8686-940-4_118)

30. Corvin S, Fauchon C, Patural H, Peyron R, Reby D, Theunissen F, Mathevon N. 2024 Pain cues override identity cues in baby cries. *iScience* **27**. (doi:10.1016/j.isci.2024.110375)

31. Osiecka AN, Lefèvre R, Briefer EF. 2025 Emotional contexts influence vocal individuality in ungulates. (doi:https://doi.org/10.1101/2024.09.18.613506)

32. González-García E, Gozzo de Figuereido V, Foulquie D, Jousserand E, Autran P, Camous S, Tesniere A, Bocquier F, Jouven M. 2014 Circannual body reserve dynamics and metabolic profile changes in Romane ewes grazing on rangelands. *Domestic Animal Endocrinology* **46**, 37–48. (doi:10.1016/j.domaniend.2013.10.002)

33. Boissy A, Bouix J, Orgeur P, Poindron P, Bibé B, Neindre PL. 2005 Genetic analysis of emotional reactivity in sheep: effects of the genotypes of the lambs and of their dams. *Genet. Sel. Evol.* **37**, 381–401. (doi:10.1051/gse:2005007)

34. Ligout S, Foulquié D, Sèbe F, Bouix J, Boissy A. 2011 Assessment of sociability in farm animals: The use of arena test in lambs. *Applied Animal Behaviour Science* **135**, 57–62. (doi:10.1016/j.applanim.2011.09.004)

35. Briefer EF, Tettamanti F, McElligott AG. 2015 Emotions in goats: mapping physiological, behavioural and vocal profiles. *Animal Behaviour* **99**, 131–143. (doi:10.1016/j.anbehav.2014.11.002)

36. Boissy A *et al.* 2007 Assessment of positive emotions in animals to improve their welfare. *Physiology & Behavior* **92**, 375–397. (doi:10.1016/j.physbeh.2007.02.003)

37. Villain AS, Renaud-Goud P. 2023 SoundChunk R Package and example data. (doi:10.5281/zenodo.10796326)

38. Sueur J, Aubin T, Simonis C. 2008 Seewave, a free modular too for sound analysis ans synthesis. *Bioacoustics* **18**, 213–226. (doi:10.1080/09524622.2008.9753600)

39. Sueur J. 2018 *Sound Analysis and Synthesis with R*. Cham: Springer International Publishing. (doi:10.1007/978-3-319-77647-7)

40. Ligges U, Krey S, Mersmann O, Schnackenberg S. 2023 *tuneR: Analysis of Music and Speech*. See https://CRAN.R-project.org/package=tuneR.

41. Ripley B, Venables B, Bates DM, ca 1998) KH (partial port, ca 1998) AG (partial port, polr) DF (support functions for. 2025 MASS: Support Functions and Datasets for Venables and Ripley's MASS.

42. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* **67**, 1–48. (doi:10.18637/jss.v067.i01)

43. Mundry R, Sommer C. 2007 Discriminant function analysis with nonindependent data: consequences and an alternative. *Animal Behaviour* **74**, 965–976. (doi:10.1016/j.anbehav.2006.12.028)

44. Taylor AM, Reby D. 2010 The contribution of source–filter theory to mammal vocal communication research. *Journal of Zoology* **280**, 221–236. (doi:10.1111/j.1469-7998.2009.00661.x)

45. Lenth RV. 2016 Least-Squares Means: The R Package lsmeans. **69**, 1–33. (doi:doi:10.18637/jss.v069.i01)

46. Bradley MM, Codispoti M, Cuthbert BN, Lang PJ. 2001 Emotion and motivation I: defensive and appetitive reactions in picture processing. *Emotion* **1**, 276–298. (doi:10.1037/1528-3542.1.3.276)

47. Jones RB, Hocking PM. 1999 Genetic Selection for Poultry Behaviour: Big Bad Wolf or Friend in Need? *Animal Welfare* **8**, 343–359. (doi:10.1017/S0962728600021977)

48. Linn SN, Schmidt S, Scheumann M. 2021 Individual distinctiveness across call types of the southern white rhinoceros (Ceratotherium simum simum). *Journal of Mammalogy* **102**, 440–456. (doi:10.1093/jmammal/gyab007)

49. Mitani JC, Gros-Louis J, Macedonia JM. 1996 Selection for acoustic individuality within the vocal repertoire of wild chimpanzees. *Int J Primatol* **17**, 569–583. (doi:10.1007/BF02735192)

50. Snowdon CT, Hausberger M. 1997 *Social Influences on Vocal Development*. Cambridge University Press.

51. Lemasson A, Hausberger M. 2011 Acoustic variability and social significance of calls in female Campbell's monkeys (Cercopithecus campbelli campbelli). *The Journal of the Acoustical Society of America* **129**, 3341–3352. (doi:10.1121/1.3569704)

Isolation bleats

Pre-feeding bleats

A

B

Norm.amplitude

C

Individuality in isolation bleats