Exposure to polycyclic aromatic hydrocarbons and phthalate diesters in a threatened freshwater turtle *Emys orbicularis*: effects of age, sex, and temporal variation

Leslie-Anne Merleau*[1,2,3], Olivier Lourdais[2,4], Anthony Olivier[3], Marion Vittecoq[3], Fabrice Alliot[1], Sira Traore[1], Carole Leray[3], and Aurélie Goutte[1]

[1] École Pratique des Hautes Études, PSL Research University, UMR 7619 METIS, Sorbonne University – Paris, France *
[2] Centre d'Études Biologiques de Chizé, CNRS, UMR 7372 – Villiers en Bois, France
[3] La Tour du Valat, Research Institute for the Conservation of Mediterranean Wetlands – Arles, France
[4] School of Life Sciences, Arizona State University – Tempe, Arizona, USA

*Corresponding author
Correspondence: leslie-anne.merleau@ephe.psl.eu

ABSTRACT
Freshwater ecosystems constitute major sinks for organic contaminants, increasing pressure on already threatened environments and jeopardising the unique biodiversity they harbour. In addition to the persistence of legacy compounds, pollutants of various families are being released into the environment, such as polycyclic aromatic hydrocarbons (PAHs) and phthalate diesters (PAEs). Heretofore, these pollutants have received insufficient attention in freshwater vertebrates, particularly in reptiles, considering their potential to trigger detrimental effects on organisms. During two years (2019 and 2020), we quantified plasma levels of 46 compounds from 6 chemical families in two monitored populations of the European pond turtle (*Emys orbicularis*) in the Camargue wetland, France. PAHs and PAEs were often found concomitantly and with similar occurrences and levels in the two populations. We observed similar inter-annual variations in PAHs and PAEs with differences between males and females, highlighting the need for a better understanding of the role of sex in the exposure pathway and the toxicokinetics of contaminants, especially in turtles. The negative relationship between PAH levels and age, as well as the high intra-individual variation in levels of both contaminant families, provides further evidence of limited bioaccumulation of these pollutants in the blood of *E. orbicularis*. This could be explained by the metabolic biotransformation of parent compounds and the need to quantify the levels of PAH metabolites and phthalate monoesters. Finally, our work highlights the need for long-term monitoring and assessment of the potential maternal transfer of these contaminants in order to determine the risks for the conservation of turtle populations that are already exposed to a wide range of contaminants.

Keywords: organic contaminants, plasma, reptile, wetlands

Introduction

Freshwater ecosystems are continuously exposed to a wide range of organic pollutants from agricultural, urban, and industrial sources. These include legacy contaminants such as organochlorine pesticides (OCPs) and polychlorinated biphenyls (PCBs), as well as currently emitted compounds like polycyclic aromatic hydrocarbons (PAHs) and phthalate esters (PAEs) (Malaj et al., 2014; Rasmussen et al., 2015). Most of these compounds are considered Persistent Organic Pollutants (POPs) and are banned by international regulations, such as PCBs and OCPs, but their persistence in aquatic environments still poses a long-term threat to biodiversity (Lohmann et al., 2007; Mateo et al., 2016). Some PAHs, and to a lesser extent PAEs, can also persist, notably in sediments (Manzetti, 2013; Net et al., 2015). However, international regulations targeting industrial and domestic releases of PAHs and PAEs remain limited or nonexistent, leading to steady environmental contamination (Ravindra et al., 2008; Malaj et al., 2014; Net et al., 2015).

Once in the environment, these contaminants can accumulate in aquatic organisms. PCBs and OCPs tend to biomagnify through food webs because of their lipophilicity and slow metabolism (Mackintosh et al., 2004; Walters et al., 2016; Goutte et al., 2020) whereas PAHs and PAEs are more prone to trophic dilution due to higher metabolization and excretion in vertebrate species (Mackintosh et al., 2004; Wan et al., 2007; Walters et al., 2016; Goutte et al., 2020) However, repeated contamination through water and sediments can lead to chronic exposure and the saturation of metabolic capacities for these molecules (Van Der Oost et al., 2003; Tyler et al., 2018).

Exposure to such contaminants can lead to detrimental effects in a wide range of organisms. PCBs, OCPs, PAHs, and PAEs can cause endocrine disruption, particularly in the thyroid and reproductive hormonal systems (Vethaak & Legler, 2013; Matthiessen et al., 2018; Wallace et al., 2020). Impairment of immunological functions, oxidative stress, genotoxicity and embryotoxicity have also been reported in vertebrate species exposed to these contaminants (Reynaud & Deschaux, 2006; Marasco & Costantini, 2016; González-Mille et al., 2019; Wallace et al., 2020). These various sublethal effects can ultimately compromise individual fitness through higher mortality rates or reduced reproductive outputs, potentially leading to population decline (Aulsebrook et al., 2020; Campioni et al., 2024). However, the majority of ecotoxicological studies on organic contaminants still focus on abiotic environmental monitoring, excluding wildlife exposure (Simms et al., 2025).

This knowledge gap is particularly concerning given the current state of freshwater ecosystems. Wetlands are among the most threatened ecosystems worldwide, and pollution constitutes a major issue (Malaj et al., 2014; Reid et al., 2019; Dudgeon, 2019). Vertebrate populations in wetlands are declining faster and sharper than in any other ecosystem (Reid et al., 2019), highlighting the necessity for targeted monitoring of contaminant burdens in wildlife species (Simms et al., 2025). Furthermore, field studies have frequently focused on a specific class of pollutants, thereby neglecting the diversity of contaminants and potential synergistic effects (Molbert et al., 2019).

The distribution of field studies across taxa is also uneven. Among vertebrates, reptiles remain largely overlooked in ecotoxicology despite being one of the most threatened taxa (Sparling et al., 2010; Simms et al., 2025). Nevertheless, some organic contaminants have been quantified in wild reptile species (Yu et al., 2012; Meyer et al., 2016; González-Mille et al., 2019; Sanjuan et al., 2023). Freshwater turtles are key species for monitoring pollutant exposure because of their high trophic positions and slow generational turnover that increase their susceptibility to bioaccumulation (Rowe, 2008). In addition, their lifespans and high site fidelity (Olivier, 2002; Fay et al., 2023) enable them to be exposed to pollutants over decades, providing insights into local contamination patterns (Rowe, 2008; Merleau, Goutte, et al., 2024). The European pond turtle (*Emys orbicularis*) has a large Eurasian distribution with fragmented and declining populations, exposed to a wide range of anthropogenic disturbances. Several studies have investigated levels of metallic trace elements (MTEs), pesticides, and POPs, suggesting that this species is a good

candidate for ecotoxicological studies (Swartz et al., 2003; Namroodi et al., 2017; Guillot et al., 2018; Beau et al., 2019; Burkart et al., 2021; Merleau, Lourdais, et al., 2024; Merleau, Goutte, et al., 2024).

We measured organic contaminants in two neighboring populations of European pond turtles monitored since 1976 (Olivier, 2002; Olivier et al., 2010) in the Camargue wetland (Southern France), a biodiversity hotspot. These two populations are located in the Rhône Delta and exposed to several sources of urban and industrial pollution (Roche et al., 2000; Berny et al., 2002; Ribeiro et al., 2005; Burkart et al., 2021). The Rhône River, one of France's main catchment areas, is particularly exposed to PCBs and PAEs (Paluselli et al., 2018; Dendievel et al., 2020), and a major industrial port zone (ZIP) and petrochemical complex are located a few kilometers downstream. The region is also characterized by intensive rice cultivation, which is historically associated with heavy OCP use (Roche et al., 2000). The two populations of *E. orbicularis* differ in their hydrology in relation to the irrigation and drainage systems of the rice fields, leading to contrasting exposure to pollution (Burkart et al., 2021; Merleau, Lourdais, et al., 2024; Merleau, Goutte, et al., 2024). Furthermore, our long-term monitoring provides insight into the effects of individual characteristics on contamination, such as age and sex. Using a multiresidue method of quantification (Molbert et al., 2019), we measured plasma concentration of 15 PAHs, 7 PAEs, and 24 other organic contaminants (PCBs, OCPs, pyrethroids, DEET, PBDEs) over two years (2019 and 2020) in 162 individuals. We aimed to assess:

1) How spatiotemporal factors influenced organic contaminant burdens in *E. orbicularis* in relation to chemical family
2) How individual characteristics such as sex, body condition, and age shaped the levels of contaminants
3) If contaminant levels exhibited fluctuations at the intra-individual level in a subset of individuals sampled over the two years

Material and methods

Animal captures and marking

Turtles were captured in two populations located in the Camargue wetland (Southern France) in the Natural Reserve of the Tour du Valat (43°30′N, 4°40′E). These two populations of *E. orbicularis* are only a few kilometers apart, but their habitats differ strongly regarding the hydrological characteristics (Olivier et al., 2010; Burkart et al., 2021; Merleau, Lourdais, et al., 2024; Merleau, Goutte, et al., 2024). The population of Esquineau occupies canals and marshes associated with the rice paddies irrigation system, using water coming from the Rhône River. The population of Faïsses is found in the paddy drainage system and the associated canals and marshes. Turtles were collected from the 23rd of April to the 26th of July in 2019 and from the 9th of April to the 7th of August in 2020, either manually or with funnel traps. Turtles are individually recognized by notching their marginal and nuchal scales as part of a long-term capture-mark-recapture survey (1976–2025) (Olivier, 2002; Ficheux et al., 2014). Traps located in 20 and 16 capture stations (Esquineau and Faïsses, respectively) were checked every day, and individuals were released at the precise location of capture. Captures of this protected species were performed under French Departmental Authorities authorization (Permits: DREAL Cerfa 13616-01; N°13-2020-03-27-007).

Blood sampling and biometrics

When collecting and processing biological samples, we were careful to ensure that plastic sampling materials did not contain PAEs, and we did not use polyvinyl chloride (PVC) to avoid contamination of biological samples. We sampled 1.5 mL of blood from 162 individuals using the dorsal coccygeal vein and a previously heparinized syringe and a 25G needle. In order to comply with animal welfare standards, we performed blood sampling on turtles weighing more than 300 g, i.e., sampling volume corresponding to less than 1% body mass. During the study, 56 individuals were sampled both years (2019 and 2020), resulting in a total of 218 blood samples. After centrifugation to separate plasma from erythrocytes, the samples were kept at -18°C until analyses at Sorbonne University's UMR 7619 METIS. An Ethics Committee approved this procedure in accordance with EU Directive 2010/63/EU for animal experiments (permit: APAFIS 17899–201 812 022 345 423).

Turtles were weighed with a precision scale (Mettler Toledo PB3001-S) to the nearest gram and measured with a caliper. We used the dorsal shell length (hereafter "carapace length") as the measure of the size. We estimated body condition using the residuals of a linear regression of the log-transformed mass against the log-transformed carapace length with the addition of sex as a control variable. Thanks to the CMR monitoring, the age of the individual, obtained by counting the number of growth rings on the plastron scutes at first capture before 5 years old (Castanet, 1988; Olivier, 2002), was known for 111 samples (68 from 48 females, range: 5- 32 years; 43 from 35 males, range: 4-25 years).

Multiresidue analytic protocol

Using a method developed by Molbert et al., (2019), we quantified 15 PAHs (acenaphtylene, naphthalene, anthracene, fluorene, phenanthrene, fluoranthene, pyrene, chrysene, benzo[a]anthracene, benzo[a]pyrene, benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[g,h,i]perylene, dibenzo[a,h]anthracene, indeno[1,2,3-cd]pyrene), 7 PAEs (dimethyle phthalate (DMP), diethyle phthalate (DEP), di-iso-butyl phthalate (DIBP), di-n-butyl phthalate (DNBP), n-butyl benzyl phthalate (BBP), di-2-ethylhexyl phthalate (DEHP), di-n-octyl phthalate (DnOP), 7 pyrethroids (bifenthrin, permethrin, phenothrin, cyfluthrin, cypermethrin, fenvarelate, deltamethrin), 4 OCPs (PeCB, HCB, lindane, p,p'-DDE), 7 PCB congeners (28, 52, 101, 118, 138, 153, 180), and 6 polybrominated diphenyl ethers (PBDE) congeners (28, 47, 99, 100, 153, 154). Quantification was carried out by internal calibration using a mix of standards (phenanthrene d10; pyrene d10; benzo(a)anthracene d12; DEP d4; DEHP d4; lindane-d6; cis-permethrin 13C; PCB 30/107, and BDE Standard Mix; see Supplementary Material) that were added to 100 µL of thawed plasma. After being vortexed and stored one night at 4°C, the extraction was performed with hexane/dichloromethane/acetone (1:1:1, v/v/v) with two rounds of centrifugation, then concentrated under a nitrogen stream to approximately 0.5 mL. Purification was performed on Florisil cartridges (Supelco 1 g/6 mL, Sigma-Aldrich) using two cycles of elution with 2x5 mL of hexane/dichloromethane (1:1, v/v) and 2x5 mL of hexane/acetone (1:1, v/v). After concentration to 0.5 mL under nitrogen stream and with the Genevac Concentrator EZ-2 (Biopharma Technologies), the extracted samples were filtered with nylon filter tubes (porosity 0.2 µm, VWR, Fontenay-sous-Bois) and rinsed twice with 70 µL of hexane. Filtrates were obtained by centrifugation (6000 rpm, 2 min) and rinsing (50 µL hexane) of the tubes. The samples were then analysed by gas chromatography coupled with tandem mass spectrometry (GC-MS/MS). Blanks were performed for each series of 12 samples. The method was validated in terms of yield, repeatability, and quantification limits for these compounds and in this matrix (see Supplementary Material).

Statistical analyses

We performed all statistical analyses with R software (4.4.2) and the packages lme4 (Bates et al., 2015), emmeans (Lenth, 2025), DHARMa (Hartig, 2024), and rptR (Stoffel et al., 2017). Non-

significant variables (p > 0.05) were gradually eliminated from the models. The statistical values
acquired in the final stage prior to their removal from the models are supplied for non-significant
variables. Tukey post hoc tests in the emmeans package were used for pairwise comparisons
(Lenth, 2025). Using the DHARMa package, residual analysis was performed to check model
assumptions (Bates et al., 2015).

Variation in contaminants
To account for repeated sampling of individuals, we employed LMMs (Linear Mixed Models)
with individual identity as a random effect. We examined how the sums of plasmatic concentrations
of compounds from the same chemical family were affected by year, month, site, sex, carapace
length, body condition, and the interactions site × year, sex × year, sex × size, and sex × age. To
best suit the model assumptions, the sums of PAEs and PAHs were log-transformed. We also
performed the same LMMs by examining the impact of age on the subset of known-aged
individuals (n = 111), removing size and body condition variables from the models as they were
highly associated with age.

Intra-individual variation over time
We also assessed annual variations at the intra-individual level using repeated measures from
56 individuals sampled in 2019 and 2020. We used t-tests to compare groups of individuals
sampled each year in order to evaluate variations in individuals collected across the two years.
The differences were tested for normality using the Shapiro-Wilk normality test. For the PAH
concentrations, as the Shapiro-Wilk test indicated a statistically significant deviation from
normality, we confirmed the results of the t-test using a Wilcoxon test which confirmed the
statistical result obtained. Using the program rptR, we then conducted individual repeatability tests
to evaluate the intra-individual variation between 2019 and 2020 (Stoffel et al., 2017).
"Repeatability" is defined as the stability of the intra-individual measured contaminants over time.
220
221
222

Results

Contaminant levels in the plasma of *E. orbicularis*

Organic pollutants were detected in 93.1% of the plasma samples. On average, 3 different
chemicals were quantified per sample, and only 15.1% of the contaminated samples contained a
single compound. The highest number of compounds detected in a single sample was 11,
observed in one individual, and 58.3% of the samples contained compounds from at least 2
chemical families.
PAEs were the compound with the highest quantification rates (78.4%), followed by the PAHs
(66.5%). Among PAEs, DIBP and DNPB were the most frequently quantified (49.1% and 43.1%,
respectively) and had the maximum mean concentrations, 380.9 and 187.4 ng/mL of plasma,
respectively. Despite being quantified in substantial proportions, DEP, DEHP, and DMP had lower
plasma concentrations (41.4, 32.6, and 4.8 ng/mL, respectively).
Regarding PAHs, naphthalene was the most quantified (54.1%) with the highest mean
plasmatic concentration (5.2 ng/mL). Two other PAHs were quantified in substantial frequencies:
phenanthrene (23.9%) and fluorene (22.9%). However, their plasmatic levels were lower than
naphthalene: 3.4 and 0.7 ng/mL, respectively. The summary of PAEs and PAHs plasmatic
concentrations, by site and by year, is available in Supplementary Material.
We quantified DEET in 19 samples (8.7%). Among the PCB congeners, we quantified PCB 28,
52, and 138 in 3, 10, and 1 samples, respectively. No pyrethroids, OCPs, or PBDE, were
quantified.

Table 1 - Effects of environmental and individual variables on the levels of PAEs and PAHs in the plasma of *E. orbicularis* in two populations of the Camargue wetland (France). Models were selected using a top-to-bottom approach, based on two general models: Σcontaminants ~ population*year + month + sex*year + sex*scale(CL) + body condition and Σcontaminants ~ population*year + sex*year + sex*age for the subset of known-aged individuals. For each effect tested, we present either (1) if selected in the final model, the observed significant effect (in bold), or (2) if removed, the level before exclusion.

	All (218 samples)			Known-aged (111 samples)		
A. Sum of 5 PAEs	chi²	df	p value	chi²	df	p value
population*year	0.525	1	0.469	1.702	1	0.192
year	**7.739**	**1**	**0.005**	**22.191**	**1**	**< 0.001**
population	0.020	1	0.889	**4.922**	**1**	**0.027**
month	1.070	4	0.899	5.375	4	0.251
sex*year	**5.300**	**1**	**0.021**	1.683	1	0.858
sex*CL(scaled)	0.275	1	0.6	-	-	-
sex*age	-	-	-	0.002	1	0.966
sex	0.012	1	0.912	0.345	1	0.557
CL(scaled)	0.020	1	0.887	-	-	-
body condition (scaled)	0.115	1	0.734	-	-	-
age	-	-	-	**7.765**	**1**	**0.005**
B. Sum of 3 PAHs						
population *year	0.040	1	0.841	0.041	1	0.840
year	**19.084**	**1**	**< 0.001**	**39.081**	**1**	**< 0.001**
population	0.688	1	0.407	0.128	1	0.721
month	**12.698**	**4**	**0.013**	8.936	4	0.063
sex*year	0.156	1	0.693	0.047	1	0.828
sex*CL(scaled)	1.420	1	0.233	-	-	-
sex*age	-	-	-	**11.912**	**1**	**0.001**
sex	0.072	1	0.788	0.607	1	0.436
CL(scaled)	0.437	1	0.509	-	-	-
body condition (scaled)	0.653	1	0.419	-	-	-
age	-	-	-	0.626	1	0.429

Note: Results of linear mixed models (LMM) for the sums of PAEs (phthalate diesters) and PAHs (polycyclic aromatic hydrocarbons) levels. Known-aged: a subset of individuals of known age. CL: carapace length, a measure of the body size. The significant variables retained in the final model are in bold font. Sum of 5 PAEs: DIBP, DNPB, DEP, DEHP, and DMP. Sum of 3 PAHs: naphthalene, phenanthrene, and fluorene.

PAE concentrations

The sum of the PAE concentrations differed between males and females in interaction with the year (Table 1). Males had significantly higher levels compared to females, but only in 2020 (Fig. 1). We did not detect a significant effect of the month of capture (Table 1) nor a difference between the two populations when all the samples were taken into account (Table 1). Regarding individual characteristics, plasmatic levels of PAEs were not affected by individual size and body condition (Table 1). Although the final model for the subset of known-age individuals included the site of capture as an explanatory variable, this effect was nearly non-significant (Table 1). We also found an effect of the year: plasmatic levels of PAEs were lower in 2019 than in 2020 (Table 1, Fig. 2). The levels of PAEs significantly decreased with age (Table 1, Fig. 2).

Figure 1 - Sum (Σ) of the five phthalate diester (PAE) concentrations (ng.mL^{-1}) quantified in the plasma of *E. orbicularis* from two populations in the Camargue (France) sampled in 2019 and 2020, according to the year and the sex of individuals (female: red; male: blue; n_{Female} = 134, n_{Male} = 84). Statistical significance obtained from the LMM model: **: $p < 0.01$, ns: not significant.

Figure 2 - Sum (Σ) of the five phthalate diester (PAE) concentrations (ng.mL^{-1}) quantified in the plasma of *E. orbicularis* from the Camargue (France) sampled in 2019 and 2020, according to the age of the individuals, the year (2019: green; 2020: purple), and the population (dots: data; lines: predictions of the LMM model including the age, the year, and the population; R^2 conditional = 0.34, n_{2019} = 55, n_{2020} = 56).

PAH concentrations
The plasmatic levels of PAHs were lower in 2019 compared to 2020 (Table 1, Fig. 3) and were
variable between months, although post-hoc tests did not show statistical differences between
pairs of months (Fig. S1). We did not detect any other effect on the sum of PAH concentrations
when considering all the samples. However, for samples from known-aged individuals, we also
detected an interaction between sex and age (Table 1). Older females had higher PAH levels,
whereas this correlation was negative for males (Fig. 4). Females reached higher ages in our
dataset, but the interaction between sex and age was still observed when retrieving older females.
However, females' levels were stable when only individuals under 25 years old were considered,
indicating that levels increased later in females.

300
301
302

Figure 3 - Sum (Σ) of the three polycyclic aromatic hydrocarbon (PAH)
concentrations (ng.mL^{-1}) quantified in the plasma of *E. orbicularis* from two
populations in the Camargue (France) sampled in 2019 and 2020, according to the
year and the sex of individuals (2019: green; 2020: purple; n_{2019} = 107; n_{2020} = 111).
Statistical significance obtained from the LMM model: ***: $p < 0.001$.

Intra-individual variation

Using the contamination data of the 56 individuals sampled during two years, we were able to
assess the inter-annual variation at the individual level. Individuals exhibited statistically
significantly higher levels of PAEs and PAHs in 2020 compared to 2019 (paired t-tests: PAEs, t =
-3.017, df = 55, p = 0.004; PAHs, t = -4.047, df = 55, p < 0.001). We found low and non-significant
intra-individual repeatability for both the PAE and the PAH concentrations, with repeatability
coefficients of 0.04 and 0.097, respectively (Fig. 5).
317
318

Figure 4 - Sum (Σ) of the three polycyclic aromatic hydrocarbons (PAH)
concentrations (ng.mL⁻¹) quantified in the plasma of *E. orbicularis* from the
Camargue (France) sampled in 2019 and 2020, according to the age of the
individuals, the sex (female: red; male: blue), and the year (dots: data; lines:
predictions of the LMM model including the age, the sex, and the year; R^2 conditional
= 0.38; n_{Female} = 68; n_{Male} = 43).

Figure 5 - Sum (Σ) of (A) the five phthalate diester (PAE) and (B) the three polycyclic
aromatic hydrocarbon (PAH) concentrations (ng.mL⁻¹) quantified in the plasma of *E.*
orbicularis from the Camargue (France) sampled both in 2019 and 2020, n = 36
(Esquineau: blue; Faïsses: orange; R: coefficient of individual repeatability; p:
significance).

334
335

Discussion

In this study, we investigated the circulating levels of organic compounds from 7 major chemical families in two populations of *E. orbicularis*, a threatened freshwater turtle species, in relation to spatiotemporal characteristics and individual traits. PAHs and PAEs were quantified predominantly, whereas DEET and PCBs exhibited low detection frequencies, and no pyrethroids, OCPs, and PBDEs were quantified. At the spatiotemporal level, the year of sampling emerged as the strongest driver of contaminant concentrations. In contrast, individual traits had limited influence, with age and sex contributing primarily through interactions, either between them or with other variables.

Diversity of contaminants

The contaminant plasmatic burden of the two populations was overwhelmingly due to PAEs and PAHs, and the majority of the individuals were contaminated by both of these chemical families. These results highlight the ubiquity of these two families in the environment in the vicinity of the petrochemical complex of Fos-sur-Mer.

PAHs are characterized by their number of benzene rings and can thus be classified as low (LMW) or high molecular weight (HMW) (Ravindra et al., 2008). Our results showed that the PAHs quantified in the plasma of *E. orbicularis* were almost exclusively two or three-ringed: naphthalene, phenanthrene, and fluorene. These PAHs are known to be dominant in the surface water of the area (Guigue et al., 2014). As LMW PAHs, they exhibit higher water solubility, less sorption on dissolved carbon matters, and thus slower sedimentation than PAHs with 4 or more rings, and thus may be more bioavailable for aquatic organisms (Meador et al., 1995; Rabodonirina et al., 2015). In addition, HMW PAHs appear to be less absorbed and more metabolized in higher vertebrates (Perugini et al., 2007; Frapiccini et al., 2020), which could explain the poor detection in our populations. However, the lower yield rates of some HMW PAHs in our study might provide an additional hypothesis to explain the occurrence differences between HMW and LMW PAHs (see Supplementary Material). The high occurrence of naphthalene, phenanthrene, and fluorene was also observed in sea turtle *Caretta caretta* and *Chelonia mydas* populations (Camacho et al., 2012, 2013; Cocci et al., 2018; Sinaei & Zare, 2019; López-Berenguer et al., 2023). To our knowledge, circulating levels of PAHs in freshwater turtles were only observed in *Actinemys marmorata* populations in California (Meyer et al., 2016). The sums of the plasmatic PAHs in our populations were lower than those found in *A. marmorata*, mainly due to higher levels of pyrene in this species. In detail, phenanthrene concentrations were similar to our populations of *E. orbicularis*, but fluorene was two times higher (Meyer et al., 2016). Only one study has monitored PAH levels in *E. orbicularis* populations in Azerbaijan, quantifying mostly naphthalene and phenanthrene in fat tissues (Swartz et al., 2003), but the difference between the matrices limits the comparison with our results. However, a later study suggested that chromosomal damages in these populations were correlated with environmental levels of three-ring PAHs (Matson et al., 2005). In *C. caretta*, levels of circulating LMW PAHs have been highly correlated with increased DNA methylation in blood cells (Cocci et al., 2018). The genotoxicity of LMW PAHs has also been highlighted in both in vitro and in vivo studies in fish (Holth et al., 2009; Yazdani, 2020). Our results suggest that even though HMW-PAH with high toxicity were not quantified, the LMW-PAH burden may have adverse effects on the two *E. orbicularis* populations.

We quantified 5 PAEs in the plasma of *E. orbicularis*, with higher occurrences for intermediate (DIBP, DNBP) and high (DEHP) molecular weight PAEs. These PAEs are the most abundant in freshwater environments and have various origins, from atmospheric deposition to leaching from plastic debris (Net et al., 2015). Despite the scarcity of PAE monitoring in wildlife and in reptiles in particular, these 3 PAEs exhibit frequent quantifications in vertebrate populations (Savoca et al., 2018; Liu et al., 2019; Blasi et al., 2022; Molbert et al., 2025). To our knowledge, our work is the first to assess PAE burden in the plasma of a freshwater turtle species. In Mediterranean populations of *C. caretta*, plasma levels of DEHP were similar to our populations of *E. orbicularis*,

whereas DIBP and DnBP were respectively 9 and 3 times lower (Blasi et al., 2022). PAEs are usually not acutely toxic to vertebrates and mainly cause endocrine disruption (Oehlmann et al., 2009; Tyler et al., 2018), but long-term toxicity thresholds are unknown for reptile species. In mammal, fish, and bird species, chronic exposure to DEHP, DnBP, or DIBP has been linked to testicular toxicity (Tyler et al., 2018; Alam et al., 2025). In addition, DEHP exposure can lead to ovarian and hepatotoxicity (Alam et al., 2025). In wild populations of *Squalius cephalus*, Molbert et al. (2021) found a negative association between PAE metabolite burden in muscles and telomere length, suggesting an effect on life expectancy. The concentrations measured in *E. orbicularis* in this study might thus have a deleterious impact on the general health of the individuals.

In a previous study on the same populations carried out in 2018, PCB52 and PCB153 were quantified in 45% and 41% of the individuals from Esquineau and Faïsses (Burkart et al., 2021). In the present study, we quantified PCB with low occurrence (6%), and PCB52 was detected in only 10 samples from 2019 and 2020. PCBs are persistent in the environment, and the main explanation for this discrepancy over one or two years in the same populations lies in the extraction protocol used. The method used by Burkart et al. (2021) was specifically designed to extract and purify PCB and OCP, which allowed them to obtain three to 84 times lower LOQ for these compounds than in the multiresidue method used in 2019 and 2020 (Molbert et al., 2019).

Spatiotemporal effects

At the temporal scale, significant inter-annual variations were observed for both PAEs and PAHs, with overall higher concentrations in 2020, except for PAEs in females. These results are consistent with changes in emission patterns and meteorological conditions, particularly for PAHs (Miura et al., 2019; Dron et al., 2021). Surprisingly, plasmatic PAH burdens were higher in 2020. While lockdown periods were associated with temporary reductions in some pollutants (Bakola et al., 2022), bounce-back effects in post-lockdown periods have been observed (Li et al., 2024). Lockdown in France ended shortly after the beginning of the fieldwork, and the higher PAH levels might reflect this rebound in anthropogenic activity. We also highlighted a monthly variation for PAHs that seemed mostly driven by a few individuals with high levels in May and June (see Supplementary Material). Although PAEs and PAHs are continuously released by anthropogenic activities, environmental concentrations exhibit seasonal variations, with PAHs typically peaking in winter, a pattern also observed in the study area (Ravindra et al., 2008; Guigue et al., 2014). Such patterns can be reflected in biota contamination, as shown in two Mediterranean fish species (Perugini et al., 2007; Frapiccini et al., 2020), but generally require longer sampling intervals than the present study.

Globally, plasmatic burdens of PAEs and PAHs were similar between the two populations. These findings suggest that atmospheric deposition might be the primary source for these populations that are geographically close but differ in the hydrology of their habitats. Although the Rhône River is a source of PAE (Paluselli et al., 2018), turtles from the irrigation site (Esquineau) did not exhibit higher levels than the ones from the drainage site (Faïsses). We previously showed that site (and presumably the hydrology) was a significant explanatory factor in both circulating levels of MTEs and pesticides in these two populations (Merleau, Lourdais, et al., 2024; Merleau, Goutte, et al., 2024). In addition, PAEs and PAHs are present in substantial amounts in the atmosphere, and wet and dry depositions in freshwater environments are important factors of contamination (Ravindra et al., 2008; Bergé et al., 2013; Rabodonirina et al., 2015; Paluselli et al., 2018). At the regional scale, the ZIP of Fos-sur-Mer is a major source of PAHs, but lichen biomonitoring showed low disparity in rural and urban sampling points outside of the complex (Ratier et al., 2018). In surface waters, dissolved PAH concentrations exhibited similar concentrations in the region, except for the ZIP, with atmospheric deposition being a major source of contamination (Guigue et al., 2014). Contrasting results have been found in wildlife in the area, with high PAH disparity in eels sampled in the Vaccares Lagoon (Ribeiro et al., 2005) but no difference in little egret eggs from three colonies of the Camargue wetland (Berny et al., 2002). All

together, these results highlight that differences in the ecological characteristics of the species and in the biological matrix used lead to a contrasting pattern of contamination.

Individuals characteristics

The long-term monitoring of the two populations allowed us to assess the effect of age, which is rarely accessible in wild populations. Age was negatively correlated with PAE levels for both sexes and with PAH levels in males only. These results are consistent with the hypothesis that these compounds do not bioaccumulate in vertebrates (Mackintosh et al., 2004; Tyler et al., 2018; Goutte et al., 2020). Metabolization capacity might improve with age, leading to lower parent compounds in older individuals. Few studies have investigated the links between PAEs and age, but a positive relationship between age and PAE metabolites, but not parent compounds, has been shown in *S. cephalus* (Molbert et al., 2020). Changes in diet or habitat use have been suggested to influence PAE levels in *C. caretta* (Blasi et al., 2022) and could be linked to age in *E. orbicularis* (Ottonello et al., 2005). Growth dilution, as highlighted in *C. caretta* for PAHs and for PAE metabolites (Camacho et al., 2012; Sanjuan et al., 2023), is unlikely to play a role in *E. orbicularis*, given the absence of a link between the contaminant burden and the size of individuals in our dataset.

Sex-related differences were linked in contrasting ways to year for PAEs and age for PAHs. Higher levels of PAEs in males in 2020 only are more likely to reflect temporary changes in habitat or diet than biological differences in toxicokinetics. However, the significant decrease of PAH concentrations with age in males might indicate differential uptake/elimination or metabolization rates between sexes, as already suggested in *Lepidochelys kempii* (Muñoz et al., 2021). In the same populations of *E. orbicularis*, results supporting this hypothesis were observed for lead, whose levels increased in older females but not in males (Merleau, Goutte, et al., 2024). Maternal transfer has been identified for PAEs in *Enhydris chinensis* and *C. caretta* (Liu et al., 2019; Savoca et al., 2021) and for PAHs in *Eretmochelys imbricata* (Muñoz & Vermeiren, 2018). The contaminant burden quantified in *E. orbicularis* females during the reproductive season might thus have deleterious consequences for their offspring.

Intra-individual variation across the years

The individual repeatability between years was particularly low for both PAEs and PAHs, indicating that exposure is fluctuating over time for individuals. Even though plasma is a dynamic biological compartment, these results reinforce the conclusion that PAEs and PAHs are efficiently metabolized in *E. orbicularis*, as already shown in numerous vertebrate species (Mackintosh et al., 2004; Wan et al., 2007; Fourgous et al., 2016; Tyler et al., 2018; Goutte et al., 2020). Further studies should also investigate plasmatic levels of PAH metabolites and phthalate monoesters to better assess exposure and toxicokinetics. Temporal variations at the intra-individual levels are contaminant-dependent: mercury, lead, and selenium had high repeatability in the same individuals, whereas bentazone, a widely used pesticide in the area, exhibited strong intra-individual variations over the years, similarly to PAEs and PAHs (Merleau, Lourdais, et al., 2024; Merleau, Goutte, et al., 2024).

Conclusion

Freshwater ecosystems constitute sinks for multiple organic contaminants, yet exposure studies remain scarce for many vertebrate species, particularly reptiles. In this study, we showed that a threatened freshwater turtle, *E. orbicularis*, presented elevated high plasmatic concentrations of organic pollutants. While legacy contaminant occurrences were considerably

low, PAEs and PAHs were frequently quantified at the same time. Our results suggest a global local contamination, which appears to be independent of the hydrology type of the habitats, probably originating from atmospheric deposition and fluctuating over time. Both the relationship of PAE and PAH levels with age and intra-individual variation indicate limited potential for bioaccumulation in this species, in accordance with the toxicokinetics of these compounds in vertebrates. However, the significant differences between males and females underline the need to better understand the sex-specific exposure pathways and contamination levels. In particular, monitoring the potential maternal transfer of these contaminants, which can exhibit endocrine-disrupting properties and embryotoxicity, deserves priority attention. Finally, the detection of PAEs and PAHs in these two populations, in which contamination by pesticides and MTEs has already been identified, raises additional concern for the conservation of this threatened species. Long-term monitoring efforts should be maintained to detect potential effects on individual health and population dynamics.

Acknowledgments

The authors are grateful to Aésane Méric, Lousiane Burkart, and Louis Spinner for their help in collecting data in the field. We thank the French Ministry of Environment for giving permission to capture *E. orbicularis*.

Funding

The present work was funded by the Water Agency Rhône Méditerranée Corse, the Fondation de la Tour du Valat, the Office Français de la Biodiversité, Plan Ecophyto II under the project "CISTOX" OFB.21.09.42, and the École Pratique des Hautes Études for financing L.A.M's temporary teaching and research associate contract.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

Data and script are available online: https://doi.org/10.5281/zenodo.15721641; Merleau et al., 2025.

Supplementary information is available online: https://doi.org/10.5281/zenodo.15756310; Merleau et al., 2025.

References

Alam MS, Maowa Z, Hasan MN (2025) Phthalates toxicity in vivo to rats, mice, birds, and fish: A thematic scoping review. *Heliyon*, **11**, e41277. https://doi.org/10.1016/j.heliyon.2024.e41277

Aulsebrook LC, Bertram MG, Martin JM, Aulsebrook AE, Brodin T, Evans JP, Hall MD, O'Bryan MK, Pask AJ, Tyler CR, Wong BBM (2020) Reproduction in a polluted world: implications for wildlife. *Reproduction*, **160**, R13–R23. https://doi.org/10.1530/REP-20-0154

Bakola M, Hernandez Carballo I, Jelastopulu E, Stuckler D (2022) The impact of COVID-19 lockdown on air pollution in Europe and North America: a systematic review. *European Journal of Public Health*, **32**, 962–968. https://doi.org/10.1093/eurpub/ckac118

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using **lme4**. *Journal of Statistical Software*, **67**. https://doi.org/10.18637/jss.v067.i01

Beau F, Bustamante P, Michaud B, Brischoux F (2019) Environmental causes and reproductive correlates of mercury contamination in European pond turtles (*Emys orbicularis*). *Environmental Research*, **172**, 338–344. https://doi.org/10.1016/j.envres.2019.01.043

Bergé A, Cladière M, Gasperi J, Coursimault A, Tassin B, Moilleron R (2013) Meta-analysis of environmental contamination by phthalates. *Environmental Science and Pollution Research*, **20**, 8057–8076. https://doi.org/10.1007/s11356-013-1982-5

Berny P, Sadoul N, Dol S, Videman B, Kayser Y, Hafner H (2002) Impact of local agricultural and industrial practices on organic contamination of little egret (*Egretta garzetta*) eggs in the rhône Delta, Southern France. *Environmental Toxicology and Chemistry*, **21**, 520–526. https://doi.org/10.1002/etc.5620210308

Blasi MF, Avino P, Notardonato I, Di Fiore C, Mattei D, Gauger MFW, Gelippi M, Cicala D, Hochscheid S, Camedda A, De Lucia GA, Favero G (2022) Phthalate esters (PAEs) concentration pattern reflects dietary habitats (δ13C) in blood of Mediterranean loggerhead turtles (*Caretta caretta*). *Ecotoxicology and Environmental Safety*, **239**, 113619. https://doi.org/10.1016/j.ecoenv.2022.113619

Burkart L, Olivier A, Lourdais O, Vittecoq M, Blouin-Demers G, Alliot F, Le Gac C, Martin N, Goutte A (2021) Determinants of Legacy Persistent Organic Pollutant Levels in the European Pond Turtle (*Emys orbicularis*) in the Camargue Wetland, France. *Environmental Toxicology and Chemistry*, **40**, 2261–2268. https://doi.org/10.1002/etc.5077

Camacho M, Boada LD, Orós J, Calabuig P, Zumbado M, Luzardo OP (2012) Comparative study of polycyclic aromatic hydrocarbons (PAHs) in plasma of Eastern Atlantic juvenile and adult nesting loggerhead sea turtles (*Caretta caretta*). *Marine Pollution Bulletin*, **64**, 1974–1980. https://doi.org/10.1016/j.marpolbul.2012.06.002

Camacho M, Luzardo OP, Boada LD, López Jurado LF, Medina M, Zumbado M, Orós J (2013) Potential adverse health effects of persistent organic pollutants on sea turtles: Evidences from a cross-sectional study on Cape Verde loggerhead sea turtles. *Science of The Total Environment*, **458–460**, 283–289. https://doi.org/10.1016/j.scitotenv.2013.04.043

Campioni L, Oró-Nolla B, Granadeiro JP, Silva MC, Madeiros J, Gjerdrum C, Lacorte S (2024) Exposure of an endangered seabird species to persistent organic pollutants: Assessing levels in blood and link with reproductive parameters. *Science of The Total Environment*, **930**, 172814. https://doi.org/10.1016/j.scitotenv.2024.172814

Castanet J (1988) Les méthodes d'estimation de l'âge chez les chéloniens. *Mésogée*, **48**, 21–28.

Cocci P, Mosconi G, Bracchetti L, Nalocca JM, Frapiccini E, Marini M, Caprioli G, Sagratini G, Palermo FA (2018) Investigating the potential impact of polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) on gene biomarker expression and global DNA methylation in loggerhead sea turtles (*Caretta caretta*) from the Adriatic Sea. *Science of The Total Environment*, **619–620**, 49–57. https://doi.org/10.1016/j.scitotenv.2017.11.118

Dendievel A-M, Mourier B, Dabrin A, Delile H, Coynel A, Gosset A, Liber Y, Berger J-F, Bedell J-P (2020) Metal pollution trajectories and mixture risk assessed by combining dated cores and subsurface sediments along a major European river (Rhône River, France). *Environment International*, **144**, 106032. https://doi.org/10.1016/j.envint.2020.106032

Dron J, Ratier A, Austruy A, Revenko G, Chaspoul F, Wafo E (2021) Effects of meteorological conditions and topography on the bioaccumulation of PAHs and metal elements by native lichen (*Xanthoria parietina*). *Journal of Environmental Sciences*, **109**, 193–205. https://doi.org/10.1016/j.jes.2021.03.045

Dudgeon D (2019) Multiple threats imperil freshwater biodiversity in the Anthropocene. *Current Biology*, **29**, R960–R967. https://doi.org/10.1016/j.cub.2019.08.002

Fay R, Ficheux S, Béchet A, Besnard A, Crochet P, Leblois R, Crivelli A, Wattier R, Olivier A (2023) Direct and indirect estimates of dispersal support strong juvenile philopatry and male-biased dispersal in a freshwater turtle species (*Emys orbicularis*). *Freshwater Biology*, fwb.14171. https://doi.org/10.1111/fwb.14171

Ficheux S, Olivier A, Fay R, Crivelli A, Besnard A, Béchet A (2014) Rapid response of a long-lived species to improved water and grazing management: The case of the European pond turtle (*Emys orbicularis*) in the Camargue, France. *Journal for Nature Conservation*, **22**, 342–348. https://doi.org/10.1016/j.jnc.2014.03.001

Fourgous C, Chevreuil M, Alliot F, Amilhat E, Faliex E, Paris-Palacios S, Teil MJ, Goutte A (2016) Phthalate metabolites in the European eel (*Anguilla anguilla*) from Mediterranean coastal lagoons. *Science of The Total Environment*, **569–570**, 1053–1059. https://doi.org/10.1016/j.scitotenv.2016.06.159

Frapiccini E, Panfili M, Guicciardi S, Santojanni A, Marini M, Truzzi C, Annibaldi A (2020) Effects of biological factors and seasonality on the level of polycyclic aromatic hydrocarbons in red mullet (*Mullus barbatus*). *Environmental Pollution*, **258**, 113742. https://doi.org/10.1016/j.envpol.2019.113742

González-Mille DJ, Ilizaliturri-Hernández CA, Espinosa-Reyes G, Cruz-Santiago O, Cuevas-Díaz MDC, Martín Del Campo CC, Flores-Ramírez R (2019) DNA damage in different wildlife species exposed to persistent organic pollutants (POPs) from the delta of the Coatzacoalcos river, Mexico. *Ecotoxicology and Environmental Safety*, **180**, 403–411. https://doi.org/10.1016/j.ecoenv.2019.05.030

Goutte A, Alliot F, Budzinski H, Simonnet-Laprade C, Santos R, Lachaux V, Maciejewski K, Le Menach K, Labadie P (2020) Trophic Transfer of Micropollutants and Their Metabolites in an Urban Riverine Food Web. *Environmental Science & Technology*, **54**, 8043–8050. https://doi.org/10.1021/acs.est.0c01411

Guigue C, Tedetti M, Ferretto N, Garcia N, Méjanelle L, Goutx M (2014) Spatial and seasonal variabilities of dissolved hydrocarbons in surface waters from the Northwestern Mediterranean Sea: Results from one year intensive sampling. *Science of The Total Environment*, **466–467**, 650–662. https://doi.org/10.1016/j.scitotenv.2013.07.082

Guillot H, Bonnet X, Bustamante P, Churlaud C, Trotignon J, Brischoux F (2018) Trace Element Concentrations in European Pond Turtles (*Emys orbicularis*) from Brenne Natural Park, France. *Bulletin of Environmental Contamination and Toxicology*, **101**, 300–304. https://doi.org/10.1007/s00128-018-2376-7

Hartig F (2024). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. R package version 0.4.7. https://doi.org/10.32614/CRAN.package.DHARMa

Holth TF, Beylich BA, Skarphédinsdóttir H, Liewenborg B, Grung M, Hylland K (2009) Genotoxicity of Environmentally Relevant Concentrations of Water-Soluble Oil Components in Cod (*Gadus morhua*). *Environmental Science & Technology*, **43**, 3329–3334. https://doi.org/10.1021/es803479p

Lenth RV (2025) emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.11.0. https://doi.org/10.32614/CRAN.package.emmeans

Li R, Luo Y, Zhu X, Zhang J, Wang Z, Yang W, Li Y, Li H (2024) Anthropogenic impacts on polycyclic aromatic hydrocarbons in surface water: Evidence from the COVID-19 lockdown. *Water Research*, **262**, 122143. https://doi.org/10.1016/j.watres.2024.122143

Liu Y-E, Tang B, Liu Y, Luo X-J, Mai B-X, Covaci A, Poma G (2019) Occurrence, biomagnification and maternal transfer of legacy and emerging organophosphorus flame retardants and plasticizers in water snake from an e-waste site. *Environment International*, **133**, 105240. https://doi.org/10.1016/j.envint.2019.105240

Lohmann R, Breivik K, Dachs J, Muir D (2007) Global fate of POPs: Current and future research directions. *Environmental Pollution*, **150**, 150–165. https://doi.org/10.1016/j.envpol.2007.06.051

López-Berenguer G, Acosta-Dacal A, Luzardo OP, Peñalver J, Martínez-López E (2023) Assessment of polycyclic aromatic hydrocarbons (PAHs) in mediterranean top marine predators stranded in SE Spain. *Chemosphere*, **336**, 139306. https://doi.org/10.1016/j.chemosphere.2023.139306

Mackintosh CE, Maldonado J, Hongwu J, Hoover N, Chong A, Ikonomou MG, Gobas FAPC (2004) Distribution of Phthalate Esters in a Marine Aquatic Food Web: Comparison to Polychlorinated

Biphenyls. *Environmental Science & Technology*, **38**, 2011–2020. https://doi.org/10.1021/es034745r

Malaj E, von der Ohe PC, Grote M, Kühne R, Mondy CP, Usseglio-Polatera P, Brack W, Schäfer RB (2014) Organic chemicals jeopardize the health of freshwater ecosystems on the continental scale. *Proceedings of the National Academy of Sciences*, **111**, 9549–9554. https://doi.org/10.1073/pnas.1321082111

Manzetti S (2013) Polycyclic Aromatic Hydrocarbons in the Environment: Environmental Fate and Transformation. *Polycyclic Aromatic Compounds*, **33**, 311–330. https://doi.org/10.1080/10406638.2013.781042

Marasco V, Costantini D (2016) Signaling in a Polluted World: Oxidative Stress as an Overlooked Mechanism Linking Contaminants to Animal Communication. *Frontiers in Ecology and Evolution*, **4**. https://doi.org/10.3389/fevo.2016.00095

Mateo R, Lacorte S, Taggart MA (2016) An Overview of Recent Trends in Wildlife Ecotoxicology. In: *Current Trends in Wildlife Research* Wildlife Research Monographs. (eds Mateo R, Arroyo B, Garcia JT), pp. 125–150. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-319-27912-1_6

Matson CW, Palatnikov G, Islamzadeh A, Mcdonald TJ, Autenrieth RL, Donnelly KC, Bickham JW (2005) Chromosomal Damage in Two Species of Aquatic Turtles (*Emys orbicularis* and *Mauremys caspica*) Inhabiting Contaminated Sites in Azerbaijan. *Ecotoxicology*, **14**, 513–525. https://doi.org/10.1007/s10646-005-0001-0

Matthiessen P, Wheeler JR, Weltje L (2018) A review of the evidence for endocrine disrupting effects of current-use chemicals on wildlife populations. *Critical Reviews in Toxicology*, **48**, 195–216. https://doi.org/10.1080/10408444.2017.1397099

Meador JP, Stein JE, Reichert WL, Varanasi U (1995) Bioaccumulation of Polycyclic Aromatic Hydrocarbons by Marine Organisms. In: *Reviews of Environmental Contamination and Toxicology* Reviews of Environmental Contamination and Toxicology. (ed Ware GW), pp. 79–165. Springer New York, New York, NY. https://doi.org/10.1007/978-1-4612-2542-3_4

Merleau L-A, Goutte A, Olivier A, Vittecoq M, Bustamante P, Leray C, Lourdais O (2024) Blood levels of metallic trace elements are influenced by sex, age and habitat in the European pond turtle (*Emys orbicularis*). *Science of The Total Environment*, **957**, 177487. https://doi.org/10.1016/j.scitotenv.2024.177487

Merleau L-A, Lourdais O, Olivier A, Vittecoq M, Blouin-Demers G, Alliot F, Burkart L, Foucault Y, Leray C, Migne E, Goutte A (2024) Pesticide concentrations in a threatened freshwater turtle (*Emys orbicularis*): Seasonal and annual variation in the Camargue wetland, France. *Environmental Pollution*, **341**, 122903. https://doi.org/10.1016/j.envpol.2023.122903

Meyer E, Eskew EA, Chibwe L, Schrlau J, Massey Simonich SL, Todd BD (2016) Organic contaminants in western pond turtles in remote habitat in California. *Chemosphere*, **154**, 326–334. https://doi.org/10.1016/j.chemosphere.2016.03.128

Miura K, Shimada K, Sugiyama T, Sato K, Takami A, Chan CK, Kim IS, Kim YP, Lin N-H, Hatakeyama S (2019) Seasonal and annual changes in PAH concentrations in a remote site in the Pacific Ocean. *Scientific Reports*, **9**, 12591. https://doi.org/10.1038/s41598-019-47409-9

Molbert N, Alliot F, Goutte A, Hansson MC (2025) The dead can talk: Investigating trace element and organic pollutant exposure in mammalian roadkill under contrasting habitats. *Environmental Pollution*, **367**, 125648. https://doi.org/10.1016/j.envpol.2025.125648

Molbert N, Alliot F, Leroux-Coyau M, Médoc V, Biard C, Meylan S, Jacquin L, Santos R, Goutte A (2020) Potential Benefits of Acanthocephalan Parasites for Chub Hosts in Polluted Environments. *Environmental Science & Technology*, **54**, 5540–5549. https://doi.org/10.1021/acs.est.0c00177

Molbert N, Alliot F, Santos R, Chevreuil M, Mouchel J-M, Goutte A (2019) Multiresidue Methods for the Determination of Organic Micropollutants and Their Metabolites in Fish Matrices. *Environmental Toxicology and Chemistry*, **38**, 1866–1878. https://doi.org/10.1002/etc.4500

Molbert N, Angelier F, Alliot F, Ribout C, Goutte A (2021) Fish from urban rivers and with high pollutant levels have shorter telomeres. *Biology Letters*, **17**, 20200819. https://doi.org/10.1098/rsbl.2020.0819

Muñoz CC, Hendriks AJ, Ragas AMJ, Vermeiren P (2021) Internal and Maternal Distribution of Persistent Organic Pollutants in Sea Turtle Tissues: A Meta-Analysis. *Environmental Science & Technology*, **55**, 10012–10024. https://doi.org/10.1021/acs.est.1c02845

Muñoz CC, Vermeiren P (2018) Profiles of environmental contaminants in hawksbill turtle egg yolks reflect local to distant pollution sources among nesting beaches in the Yucatán Peninsula, Mexico. *Marine Environmental Research*, **135**, 43–54. https://doi.org/10.1016/j.marenvres.2018.01.012

Namroodi S, Zaccaroni A, Rezaei H, Hosseini SM (2017) European pond turtle (*Emys orbicularis persica*) as a biomarker of environmental pollution in Golestan and Mazandaran provinces, Iran. *Veterinary Research Forum: An International Quarterly Journal*, **8**, 333–339.

Net S, Sempéré R, Delmont A, Paluselli A, Ouddane B (2015) Occurrence, Fate, Behavior and Ecotoxicological State of Phthalates in Different Environmental Matrices. *Environmental Science & Technology*, **49**, 4019–4035. https://doi.org/10.1021/es505233b

Oehlmann J, Schulte-Oehlmann U, Kloas W, Jagnytsch O, Lutz I, Kusk KO, Wollenberger L, Santos EM, Paull GC, Van Look KJW, Tyler CR (2009) A critical analysis of the biological impacts of plasticizers on wildlife. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **364**, 2047–2062. https://doi.org/10.1098/rstb.2008.0242

Olivier A (2002) Ecologie, traits d'histoire de vie et conservation d'une population de Cistude d'Europe, *Emys orbicularis*, en Camargue.

Olivier A, Barbraud C, Rosecchi E, Germain C, Cheylan M (2010) Assessing spatial and temporal population dynamics of cryptic species: an example with the European pond turtle. *Ecological Applications*, **20**, 993–1004. https://doi.org/10.1890/09-0801.1

Ottonello D, Salvidio S, Rosecchi E (2005) Feeding habits of the European pond terrapin *Emys orbicularis* in Camargue (Rhône delta, Southern France). *Amphibia-Reptilia*, **26**, 562–565. https://doi.org/10.1163/1568538057748062241

Paluselli A, Aminot Y, Galgani F, Net S, Sempéré R (2018) Occurrence of phthalate acid esters (PAEs) in the northwestern Mediterranean Sea and the Rhone River. *Progress in Oceanography*, **163**, 221–231. https://doi.org/10.1016/j.pocean.2017.06.002

Perugini M, Visciano P, Giammarino A, Manera M, Di Nardo W, Amorena M (2007) Polycyclic aromatic hydrocarbons in marine organisms from the Adriatic Sea, Italy. *Chemosphere*, **66**, 1904–1910. https://doi.org/10.1016/j.chemosphere.2006.07.079

Rabodonirina S, Net S, Ouddane B, Merhaby D, Dumoulin D, Popescu T, Ravelonandro P (2015) Distribution of persistent organic pollutants (PAHs, Me-PAHs, PCBs) in dissolved, particulate and sedimentary phases in freshwater systems. *Environmental Pollution*, **206**, 38–48. https://doi.org/10.1016/j.envpol.2015.06.023

Rasmussen SB, Abrahamsen P, Nielsen MH, Holm PE, Hansen S (2015) Effects of Single Rainfall Events on Leaching of Glyphosate and Bentazone on Two Different Soil Types, using the DAISY Model. *Vadose Zone Journal*, **14**, vzj2014.11.0164. https://doi.org/10.2136/vzj2014.11.0164

Ratier A, Dron J, Revenko G, Austruy A, Dauphin C-E, Chaspoul F, Wafo E (2018) Characterization of atmospheric emission sources in lichen from metal and organic contaminant patterns. *Environmental Science and Pollution Research*, **25**, 8364–8376. https://doi.org/10.1007/s11356-017-1173-x

Ravindra K, Sokhi R, Vangrieken R (2008) Atmospheric polycyclic aromatic hydrocarbons: Source attribution, emission factors and regulation. *Atmospheric Environment*, **42**, 2895–2921. https://doi.org/10.1016/j.atmosenv.2007.12.010

Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PTJ, Kidd KA, MacCormack TJ, Olden JD, Ormerod SJ, Smol JP, Taylor WW, Tockner K, Vermaire JC, Dudgeon D, Cooke SJ (2019) Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biological Reviews*, **94**, 849–873. https://doi.org/10.1111/brv.12480

Reynaud S, Deschaux P (2006) The effects of polycyclic aromatic hydrocarbons on the immune system of fish: A review. *Aquatic Toxicology*, **77**, 229–238. https://doi.org/10.1016/j.aquatox.2005.10.018

Ribeiro CAO, Vollaire Y, Sanchez-Chardi A, Roche H (2005) Bioaccumulation and the effects of organochlorine pesticides, PAH and heavy metals in the Eel (*Anguilla anguilla*) at the Camargue Nature Reserve, France. *Aquatic Toxicology*, **74**, 53–69. https://doi.org/10.1016/j.aquatox.2005.04.008

Roche H, Buet A, Jonot O, Ramade F (2000) Organochlorine residues in european eel (*Anguilla anguilla*), crucian carp (*Carassius carassius*) and catfish (*Ictalurus nebulosus*) from Vaccarès lagoon (French National Nature Reserve of Camargue) – effects on some physiological parameters. *Aquatic Toxicology*, **48**, 443–459. https://doi.org/10.1016/S0166-445X(99)00061-2

Rowe CL (2008) "The Calamity of So Long Life": Life Histories, Contaminants, and Potential Emerging Threats to Long-lived Vertebrates. *BioScience*, **58**, 623–631. https://doi.org/10.1641/B580709

Sanjuan ON, Sait STL, Gonzalez SV, Tomás J, Raga JA, Asimakopoulos AG (2023) Phthalate metabolites in loggerhead marine turtles (*Caretta caretta*) from the Mediterranean Sea (East Spain region). *Environmental Chemistry and Ecotoxicology*, **5**, 178–185. https://doi.org/10.1016/j.enceco.2023.08.003

Savoca D, Arculeo M, Barreca S, Buscemi S, Caracappa S, Gentile A, Persichetti MF, Pace A (2018) Chasing phthalates in tissues of marine turtles from the Mediterranean sea. *Marine Pollution Bulletin*, **127**, 165–169. https://doi.org/10.1016/j.marpolbul.2017.11.069

Savoca D, Arculeo M, Vecchioni L, Cambera I, Visconti G, Melfi R, Arizza V, Palumbo Piccionello A, Buscemi S, Pace A (2021) Can phthalates move into the eggs of the loggerhead sea turtle *Caretta caretta*? The case of the nests on the Linosa Island in the Mediterranean Sea. *Marine Pollution Bulletin*, **168**, 112395. https://doi.org/10.1016/j.marpolbul.2021.112395

Simms A, Robert K, Spencer R-J, Treby S, Williams-Kelly K, Sexton C, Korossy-Horwood R, Terry R, Parker A, Van Dyke J (2025) A systematic review of how endocrine-disrupting contaminants are sampled in environmental compartments: wildlife impacts are overshadowed by environmental surveillance. *Environmental Science and Pollution Research*. https://doi.org/10.1007/s11356-025-36211-y

Sinaei M, Zare R (2019) Polycyclic aromatic hydrocarbons (PAHs) and some biomarkers in the green sea turtles (*Chelonia mydas*). *Marine Pollution Bulletin*, **146**, 336–342. https://doi.org/10.1016/j.marpolbul.2019.06.024

Sparling DW, Linder G, Bishop CA, Krest SK (2010) Recent Advancements in Amphibian and Reptile Ecotoxicology. In: *Ecotoxicology of Amphibians and Reptiles* , p. . CRC Press.

Stoffel MA, Nakagawa S, Schielzeth H (2017) rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models (S Goslee, Ed,). *Methods in Ecology and Evolution*, **8**, 1639–1644. https://doi.org/10.1111/2041-210X.12797

Swartz CD, Donnelly KC, Islamzadeh A, Rowe GT, Rogers WJ, Palatnikov GM, Mekhtiev AA, Kasimov R, McDonald TJ, Wickliffe JK, Presley BJ, Bickham JW (2003) Chemical Contaminants and their Effects in Fish and Wildlife from the Industrial Zone of Sumgayit, Republic of Azerbaijan. *Ecotoxicology*, **12**, 509–521. https://doi.org/10.1023/B:ECTX.0000003038.02643.08

Tyler CR, Parsons A, Rogers NJ, Lange A, Brown AR (2018) Plasticisers and Their Impact on Wildlife. In: *Plastics and the Environment* (eds Harrison RM, Hester RE), pp. 106–130. The Royal Society of Chemistry. https://doi.org/10.1039/9781788013314-00106

Van Der Oost R, Beyer J, Vermeulen NPE (2003) Fish bioaccumulation and biomarkers in environmental risk assessment: a review. *Environmental Toxicology and Pharmacology*, **13**, 57–149. https://doi.org/10.1016/S1382-6689(02)00126-6

Vethaak D, Legler J (2013) Endocrine Disruption in Wildlife: Background, Effects, and Implications. In: *Endocrine Disrupters* (ed Matthiessen P), pp. 7–58. Wiley. https://doi.org/10.1002/9781118355961.ch2

Wallace SJ, De Solla SR, Head JA, Hodson PV, Parrott JL, Thomas PJ, Berthiaume A, Langlois VS (2020) Polycyclic aromatic compounds (PACs) in the Canadian environment: Exposure and effects on wildlife. *Environmental Pollution*, **265**, 114863. https://doi.org/10.1016/j.envpol.2020.114863

Walters DM, Jardine TD, Cade BS, Kidd KA, Muir DCG, Leipzig-Scott P (2016) Trophic Magnification of Organic Chemicals: A Global Synthesis. *Environmental Science & Technology*, **50**, 4650–4658. https://doi.org/10.1021/acs.est.6b00201

Wan Y, Jin X, Hu J, Jin F (2007) Trophic Dilution of Polycyclic Aromatic Hydrocarbons (PAHs) in a Marine Food Web from Bohai Bay, North China. *Environmental Science & Technology*, **41**, 3109–3114. https://doi.org/10.1021/es062594x

Yazdani M (2020) Comparative toxicity of selected PAHs in rainbow trout hepatocytes: genotoxicity, oxidative stress and cytotoxicity. *Drug and Chemical Toxicology*, **43**, 71–78. https://doi.org/10.1080/01480545.2018.1497054

Yu S, Halbrook RS, Sparling DW (2012) Accumulation of Polychlorinated Biphenyls (PCBs) and Evaluation of Hematological and Immunological Effects of PCB Exposure on Turtles. *Bulletin of Environmental Contamination and Toxicology*, **88**, 823–827. https://doi.org/10.1007/s00128-012-0590-2