Interaction of Wnt and SHH gradients synchronizes cell cycle exit and differentiation

Authors:

Ruiqi Li[1,2], Yiqun Jiang[3,4*], Sarah Platt[3*], Kevin A. Peterson[5], Hainan Lam[3], Sarah Van[3], Chae Ho Lim[6], Jonathan Levinsohn[7], Eva-Larue Barber[3], David Wang[8], M. Mark Taketo[9], Yuval Kluger[1,2,10,11], Peggy Myung[2,3,4,11,12]

Affiliations:

[1]Computational Biology & Biomedical Informatics Program, Yale University, New Haven, CT 06520, USA

[2]Department of Pathology, Yale University, New Haven, CT 06520, USA

[3]Department of Dermatology, Yale University, New Haven, CT 06520, USA

[4]Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06520, USA

[5]The Jackson Laboratory, Bar Harbor, ME 04609, USA

[6]Ronald O. Perelman Department of Dermatology, and Department of Cell Biology, NYU Grossman School of Medicine, New York, NY, USA

[7]Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, PA 19104.

[8]Veterans Administration, Ann Arbor Healthcare System, Ann Arbor, MI, USA

[9]Colon Cancer Project, Graduate School of Medicine, Kyoto University, Sakyo, Kyoto 606-8501, Japan

[10]Applied Mathematics Program, Yale University, New Haven, CT 06511, USA

[11]Yale Cancer Center, New Haven, CT 06520, USA

[12]Yale Stem Cell Center, New Haven, CT 06520, USA

*These authors contributed jointly

Keywords:

Hair follicle, dermal condensate, scRNA-sequencing, Wnt, Hedgehog, skin, dermis

Summary:

How signals coordinate cellular proliferation with differentiation to regulate cell fate transitions is poorly

understood. Hair follicle dermal condensates (DCs) emerge in developing skin from an acute cell fate transition

in which cell cycle exit and molecular differentiation occur simultaneously. Here, we show that the coincident

levels of Wnt and Hedgehog signals synchronize these two processes and when uncoupled, trigger them

discordantly, resulting in asynchronous arrest and fuzzy cell fate borders. We use an innovative computational

approach to dissect out independent processes from scRNA-sequencing data to show that high Wnt activity

alone elicits cell cycle exit through a Hedgehog mediator, Gli3. Furthermore, Hedgehog induces DC genes in a

Wnt-dependent manner while cell-autonomously accelerating Wnt activity, effectively synchronizing cell cycle

exit with DC gene expression. These results show that the interaction of signal gradients can balance

proliferation with differentiation to regulate cell fate transitions, revealing a tunable logic to tissue patterning.

Introduction:

To make functional organs, stem and progenitor cells must precisely coordinate proliferation with

differentiation, ensuring that distinct cell types assemble into organized populations of the correct size,

location, and relative cell proportions. In many tissues, cell cycle exit is associated with terminal differentiation[1].

In certain scenarios such as in mouse or human embryonic stem cells, G1 phase represents a permissive state

in which cells are more responsive to cell type transcription factors that induce differentiation genes [2-4]. Here,

cells may asynchronously differentiate upon entering G1/G0 phase, and the signals that regulate cell cycle

arrest may be independent of those that induce differentiation. In other scenarios such as during cortical

neurogenesis, cell cycle regulators may have a dual function in regulating cell type differentiation such that

differentiation occurs synchronously with cell cycle phase[5,6]. During development and regeneration, cells can

also progress through intermediate states of differentiation (e.g., transit-amplifying progenitors) before

undergoing cell cycle exit and terminal differentiation[7-11]. Here, intermediate states of differentiation are

coordinated with proliferation. At the same time, failure to coordinate these events can lead to pathological

states such as cancer. Therefore, understanding how these processes are coupled is essential for efforts to

promote proper regeneration and to mitigate uncontrolled and autonomous growth.

A challenge to examining this question during development or regeneration is that different cell types

dynamically undergo a mixture of unique and shared biological processes as they progress along a lineage

trajectory. This makes it challenging to accurately delineate the progression of cell states within a lineage and

the biological processes that govern transitions between cell states. Knowledge of a cell's past and future cell

state has been largely dependent on biased tools that durably label specific cell types or cell states. Live

imaging has given the ability to track individual cells over time in the same tissue allowing for real-time analysis

of cellular differentiation[12-15]. These tools have been instrumental to our understanding of lineage trajectories

and the signals that regulate cell fate decisions. Still, dissecting out the specific mechanisms of how cell

proliferation and differentiation are coordinated remains limited by the lack of tools and biomarkers to track

dynamic changes in these biological processes and to interrogate them individually, especially when they

occur concurrently. With the emergence of single-cell RNA-sequencing (scRNA-seq) technology, the ability to

profile molecular differences between cells has provided a useful tool to infer signals that regulate cell type

differentiation. Still, cell-based trajectory methods often fail to accurately delineate independent processes

when they co-occur in the same group of cells, as they collapse cells into a single pseudotemporal path that

mixes their distinct dynamics. Thus, the molecular programs that coordinate tissue growth and cell type

patterning remain poorly defined in many tissues and poses a major barrier to efforts aimed at recapitulating

organ formation *in vitro* or in adult animals.

In this study, we examine this question using a defined biological system that represents the first differentiated

population in mammalian dermis and heralds the onset of dermal lineage diversification[16-18]. In mouse

embryos, the hair follicle dermal condensate (DC) emerges at around E14.0-E14.5 in association with

overlying epithelial placodes [19-23]. DCs are clusters of specialized quiescent dermal cells that are essential for

hair follicle formation. A major obstacle to delineating the sequence of steps that lead to DC genesis is the fact

that DC cells form through an acute cell fate transition with few observable intermediate states to identify and

track DC progenitors using conventional methods. To address this issue, we previously used scRNA-seq to

infer events and signals that precede DC commitment [24,25]. Combined with *in vivo* lineage tracing, we showed

that DC cells are immediate quiescent progeny of a highly proliferative population (*Dkk1*-expressing) that

divides over a short time frame before DC gene expression and quiescence. We further showed that dermal

Wnt/β-catenin and SHH signaling, together, are required and sufficient to induce DC genesis. In other

biological systems (e.g., hair follicle epithelial placode, neural tube), Wnt and SHH signaling antagonize each

other[13,26,27]. We showed that in the dermis, they unexpectedly cooperate to induce DC genesis. However, it

remains unknown how Wnt and Shh act together intracellularly to coordinate a cell fate transition in which cell

cycle exit is tightly coupled with molecular DC differentiation, especially as Wnt and SHH antagonize each

other in other systems. As such, the DC system provides a unique opportunity to understand how these two

key signals can be differentially combined to regulate diverse cell fate patterns and transitions across tissues.

One problem is that both Wnt and SHH ligands are secreted from a common epidermal source, resulting in

covarying and overlapping Wnt and SHH signaling gradients in which both signals are simultaneously

upregulated across the transition. Furthermore, both cell cycle exit and molecular DC differentiation occur

synchronously across the pre-DC-to-DC transition, making it difficult to separate them apart and to identify their

respective molecular mediators. As neither Wnt nor SHH activation alone is sufficient to induce the transition,

we sought an approach to dissect out how these two signals regulate the pre-DC-to-DC transition. Here, we an

innovative computational method recently developed specifically to deconvolve co-occurring biological

processes from whole-transcriptome scRNA-seq data[28]. In contrast to standard cell trajectory inference

methods, GeneTrajectory (GT), identifies parallel gene trajectories, allowing independent biological processes

to be resolved. Using this method and genetic models that modulate Wnt or SHH activity, we show that while

cell cycle exit and DC gene expression occur concurrently, they are not actually regulated by shared

downstream effectors as is the case in other tissues (e.g. neurogenesis). Rather, they are synchronized by the

mere coupling of upstream Wnt and SHH co-activation, which when combined, result in a sharp cell fate

transition. In particular, we show that Wnt and SHH signaling induce cell cycle exit and molecular differentiation

through distinct downstream effectors, which revealed a previously unknown role for Gli3 in cell cycle exit. By

genetically decoupling these two graded morphogens cell cycle exit and DC differentiation occur

asynchronously, resulting in a gradual transition across time. Notably, we show that the observed overlapping

levels of Wnt and SHH during normal DC formation is caused by their ability to modulate each other, thereby

ensuring their paired activity and the synchronization of cell cycle arrest and DC differentiation. This study

provides a model to understand how other tissues may differentially combine morphogen signals to balance

proliferation with differentiation during patterned tissue growth.

Results:

Cell cycle exit and molecular differentiation are tightly coupled during dermal condensate development

To examine how Wnt and SHH signaling regulate the pre-DC-to-DC transition, we first characterized where

and when quiescence and DC gene expression occur in relationship to Wnt and SHH activity. During mouse

skin development, dermal condensates (DCs) emerge around E14.0-E14.5 underneath hair follicle epithelial

placodes (Figure 1A, S1A). DC cells express cell type markers, including *Sox2*, *Sox18*, and *Gal*. Cells

expressing these markers correspond to cells that are quiescent as shown by a very low rate of EdU

nucleotide incorporation (Figure 1B, S1B). DC marker expression also corresponds to expression of the *in vivo*

G1/G0 Fucci2 reporter (Figure 1C) [25,29], and virtually all *Sox2*+ cells co-express the cell cycle inhibitor,

p21(*Cdkn1a)* with only rare cells expressing *Cdkn1a* but not *Sox2* (Figure 1D). Thus, markers of quiescence

largely colocalize with cells that express DC marker genes. Using genetic lineage tracing and EdU pulse-chase

experiments, we previously showed that while DC cells are quiescent, they are immediate progeny of highly

proliferative *Dkk1+* progenitors that are located in the upper dermis at E13.5 and later surrounding the growing

DC (Figure 1D, S1B) [24]. We showed that these *Dkk1+* proliferative progenitors undergo a short phase of

proliferation early during morphogenesis (between approximately E13.5-E14.5) to give rise to the vast majority

of arrested DC cells. Thus, cell cycle exit and molecular DC differentiation occur synchronously after a phase

of high proliferation (Figure S1C).

We next asked where and when Wnt and SHH activity are expressed during DC formation. As SHH ligand is

exclusively expressed by hair follicle epithelial placodes, SHH is not active at E13.5. However at this time

point, Wnt signaling is activated broadly across the upper dermis (UD) in response to Wnt ligands secreted

from the epidermis. This Wnt activity is required for DC and hair follicle initiation [25,30]. Although Wnt activity

alone is not sufficient for DC gene expression, we asked if Wnt activity alone may promote quiescence

independent of SHH. We used *Lef1* as a canonical marker of Wnt activity, whose expression we previously

showed highly correlates with other known Wnt target genes (e.g., *Axin2*, *Wif1*, *Tcf7*; Figure S1C, S1D)[24,25].

We found that *Lef1* expression gradually decreases over the first four upper dermal cell layers with the most

superficial dermal layer (closest to the epidermis) showing the highest average *Lef1* expression (Figure 1E,

1F). Prior to SHH, we found no significant correlation between the rate of EdU incorporation and *Lef1*

expression levels (Figure 1E). By E14.5, SHH is secreted by placode epithelial cells, resulting in SHH

activation in the placode epithelium and underlying dermis (Figure 1G). Using *Ptch1* as a canonical marker of

SHH activity, we found that SHH activity correlates with Wnt signaling levels at E14.5, and both correlate

spatially with pre-DC-to-DC cell states, with highest levels of *Ptch1* and *Lef1* seen in cells that express DC

markers. At this point, increasing levels of both Wnt and SHH activity correlate with decreasing rates of EdU

incorporation and quiescence. Notably, Wnt signaling is amplified after E13.5, and the highest levels of Wnt

and SHH activity now correspond with cell cycle exit and DC gene expression (Figure 1H). Hence, prior to

SHH (E13.5), neither DC genes nor quiescence appear to be regulated by Wnt signaling levels. When

epithelial placodes upregulate Wnt and SHH ligands at E14.5, both dermal Wnt and SHH activity are

concurrently upregulated during the pre-DC-to-DC transition. During this transition, cell cycle exit and DC gene

expression occur synchronously.

High Wnt activity induces cell cycle exit independent of SHH

As high levels of both Wnt and SHH activity seen at E14.5 correspond to cell cycle exit and DC gene

expression, we asked if high levels of Wnt and/or SHH activity regulate these events. We and others showed

that expression of an activated form of β-catenin in the dermis induces uniformly high Wnt activity [30,31]. This

results in larger DCs where placodes normally form (Figure S2A), suggesting that cells with high levels of Wnt

signaling are primed to undergo DC differentiation upon SHH expression. To assess if increasing Wnt activity

to a level higher than that seen at E13.5 (when there are no detectable DC changes) can promote pre-DC

changes prior to SHH expression, we expressed activated β-catenin in the dermis (*Axin2CreER;βcat*[fl/EX3] or

Actβcat, Figure 2A) by administering tamoxifen at E11.5. To profile molecular changes induced by high Wnt

activation, we performed scRNA-seq analysis of dermal populations obtained from microdissected dorsolateral

skin from E13.5 mutant and paired control embryos. Dermal cells were analyzed after filtering cells based on

known dermal markers (e.g., *Col1a1*a) as previously described. As expected, the mutant dermal population

showed increased expression of Wnt target genes, including *Lef1*, *Axin2* and *Tcf7*, but lacked DC markers

(e.g., *Sox2*) or SHH target genes (Figure 2B). We next conducted differential gene expression analysis

between E13.5 control and Actβcat dermal cells (Supp table 1). Despite upregulation of Wnt target genes in

mutant dermal cells, DC markers such as *Sox2* and *Sox18* were not significantly expressed.

However, *Cdkn1a*, a cell cycle inhibitor expressed by DC cells at E14.5, was among the most significantly

upregulated genes in the Actβcat dermis, suggesting that high Wnt activation alone may induce cell cycle exit

independent of DC gene expression (Figure 2B). By FISH, we verified the lack of *Gli1* or *Sox2* expression in

the Actβcat dermis. However, significant expression of *Cdkn1a* was detected in the mutant but not in the

control dermis (Figure 2B, 2C). These *Cdkn1a*+ cells showed high levels of *Lef1* expression and correlated

with a low rate of EdU incorporation, indicating a low rate of proliferation (Figure 2C). These data demonstrate

that constitutive activation of Wnt signaling is sufficient to induce expression of *Cdkn1a*, independent of SHH

activation (Figure 2D). By E14.5, *Cdkn1a* was highly upregulated in both control and mutant DCs. However, in

contrast to controls in which *Cdkn1a* expression almost perfectly overlaps with *Sox2* expression, many cells

surrounding *Sox2*+ clusters and within the upper dermis also expressed high levels of *Cdkn1a* but not DC

markers in the Actβcat mutant. They also showed a low rate of EdU incorporation (Figure 2E). *Cdkn1a*+ cells

that co-expressed *Sox2* in the Actβcat mutant were spatially located nearest to the placode and corresponded

to cells with both high Wnt and high SHH activity. Notably, cells that expressed *Sox2* in the Actβcat mutant

showed highest *Ptch1* expression, suggesting a threshold level of SHH activity may be required for DC gene

expression even if high levels of Wnt activity are induced. To assess Wnt activity levels caused by expression

of activated β-catenin, we quantified *Lef1* levels by quantitative FISH. As expected, E13.5 Actβcat mutant

dermal cells showed higher levels of *Lef1* expression than control E13.5 dermal cells. By E14.5, higher levels

of *Lef1* are also observed in the mutant dermis compared to controls but were never greater than those seen in

control DC cells, indicating that dermal expression of stabilized β-catenin does not induce levels above those

seen physiologically in forming DCs (Figure S2B). Taken together, these results suggest that increased Wnt

signaling can induce quiescence independent of SHH, while DC gene expression appears to require Wnt

activity as well as high levels of SHH activation (Figure 2F, 2G). Hence, Wnt and SHH interact to induce cell

cycle exit or DC gene expression through different mechanisms during DC genesis.

SHH cell-autonomously augments Wnt signaling to induce cell cycle exit

Based on these results, we next asked how high SHH affects DC gene expression and cell cycle exit in the

absence of high Wnt activity. We induced uniformly high SHH activity in the dermis by expressing a

constitutively activated form of Smoothened (*Axin2CreER;RosaLSL-Smom2YFP* or ActSMO) at E13.5 when

Wnt signaling levels are low (tamoxifen E11.5; Figure 3A). Whole-mount analysis of E14.5 skin showed that

high SHH activation in Wnt-active upper dermal cells induced large DCs that formed autonomously without

placodes (Figure 3A). Interestingly, Sox2+ cells in the center of these dermal clusters lacked EdU

incorporation, indicating that cell cycle exit still occurred. At the same time, many Sox2+ cells in the periphery

were actively proliferating as shown by Sox2+/EdU+ double positive cells (Figure 3A, Figure S3A) [24]. The

dermal clusters in the ActSMO mutant were also diffuse and less tightly aggregated, suggesting that the

coordination of DC gene expression with quiescence may be critical for DC formation. This result suggests that

high SHH can induce both DC genes and cell cycle exit asynchronously.

We asked how quiescence can occur if high SHH was induced in cells with low Wnt activity and in the absence

of placodes that are typically thought to be the source of high Wnt ligands. We first measured Wnt activity in

the E13.5 ActSMO mutant and surprisingly found that *Lef1* levels in the ActSMO mutant were significantly

higher than in E13.5 controls and were similar to that seen in wildtype DC cells (Figure 3B, 3C, Figure S3A).

Further similar to the Actβcat mutant, these cells co-expressed high levels of *Cdkn1a* and corresponded with

lower rates of EdU incorporation (Figure 3D). Notably, only cells that expressed *Ptch1* (i.e., cells expressing

the ActSMO mutation) showed elevated *Lef1* expression while neighboring wildtype *Ptch1*-negative cells (Cre

non-recombined) in the upper dermis showed levels similar to the control upper dermis (Figure 3C). Thus, *Lef1*

expression is amplified cell-autonomously by SHH activation and possibly independent of epidermal Wnt

ligands. Consistent with this, we found that many cells with the highest *Lef1* transcript levels frequently were

not spatially closest to the epidermis where Wnt ligands are expressed (Figure 3B). Collectively, these results

show that SHH activation cell-autonomously accelerates Wnt activity. High Wnt activity, in turn, induces cell

cycle exit, These data also suggest that high levels of dermal Wnt activity seen at E14.5 may be caused by

SHH activation rather than by placode Wnt ligands (Figure 3E).

DC gene expression and cell cycle exit are off sync upon high SHH activation. We next sought to use this

model to dissect out how each process was mediated. Here, we employed an innovative computational

approach, GeneTrajectory (GT), to unbiasedly separate out co-occurring independent biological gene

programs from whole transcriptome scRNA-seq data. Briefly, GT is a computational method developed to

unbiasedly detect independent gene dynamics from whole transcriptome data in which each GT represents a

distinct biological process (Figure S3B). Traditional cell trajectory inference methods first organize cells into

lineages and assign a one-dimensional pseudotime and then gene dynamics are assessed with respect to

those cell lineages. A major limitation of these approaches arises when multiple biological processes coexist

within the same group of cells (e.g., when cell cycle is coupled with differentiation). Here, the inferred cell

pseudotime often conflates the effects of distinct processes, making it difficult to disentangle their respective

gene programs. By contrast, GT does not require cell trajectory inference or predefining an initial and terminal

cell state. It directly constructs gene trajectories and allows each cell to participate in multiple concurrent

biological processes. This is done by computing pairwise optimal transport distances between gene expression

distributions over the cell-cell graph from which gene programs, and their pseudotemporal gene order is

extracted based on their sequence of activation along the inferred biological processes. This separates apart

coexisting biological processes that are entangled at the cell level into independent gene programs with distinct

sequential gene expression patterns (Figure S3B).

We first used GT to examine E14.5 wildtype dermal scRNA-seq data, which identified three GTs that

correspond to cell cycle (CC), DC differentiation and lower dermal (LD) differentiation processes, respectively

(Figure 3F, Figure S3C). Here, the DC GT identifies an order of gene activity in which DC genes are expressed

at the terminus along with *Cdkn1a* (Figure 3F). This method was benchmarked against other cell trajectory

inference methods, showing superior accuracy in delineating the order of genes involved in the DC

differentiation process based on biological validaton[28]. We then defined different cell stages along the DC

differentiation process by partitioning the DC GT into gene bins, with each bin representing marker genes for a

specific stage. For each cell, we computed a gene bin score representing the proportion of genes from a given

bin that were expressed. Based on these scores, cells were exclusively assigned to a stage or left unassigned

(Method - Gene bin and cell stages, Figure 3F). Canonical DC genes, including *Gal*, *Sox18* and *Sox2,* were

largely expressed at the last stage 6 (Figure 3G). Similarly, the CC GT represents an independent process in

which cells are colored according to their expression of genes along a progression from S and G2/M phases

(Figure 3F). As shown, cells expressing genes along the DC GT also simultaneously express genes that

correspond to an independent cell cycle process (proliferation). Measuring the fraction of cells in G1/G0 phase

by DC GT stages showed that cells at stage 6 are quiescent with a G1 fraction near 1.0 (Figure 3H). At the

same time, both *Lef1* and *Ptch1* covary and show increasing levels across DC GT stages with highest levels of

both observed at stage 6 (Figure 3I). Hence as expected, Wnt activity is coupled to SHH activation in the

wildtype condition, and cell cycle exit and molecular DC differentiation occur simultaneously at the terminus

(Figure 3J).

A gene program that corresponds to cell cycle exit is identified by GeneTrajectory

Based on the observation that there were proliferating Sox2+ cells surrounding a quiescent center in the

ActSMO condition, this suggests that cell cycle exit does not always occur simultaneously with DC gene

expression. We asked if we can use GT to separate out the two processes of cell cycle exit and DC gene

expression in the ActSMO condition. We obtained scRNA-seq data from E13.5 ActSMO and paired control skin

to exclude any possible confounding effects of endogenous SHH activation and DC formation. As expected,

the ActSMO mutant showed uniformly high *Ptch1* expression in dermal cells, while Wnt target genes remained

localized to cells that express UD markers (Figure 4A). Interestingly, GT identified four GTs in the ActSMO

mutant: one corresponded to CC (S/G2/M) and another to LD differentiation, which were similar to the control

(Figure 4B, Figure S4A). As expected, one GT contained DC genes as well as Wnt target genes, which was

not detected in the E13.5 control. Additionally, a new GT was detected in the ActSMO mutant that was not

present in the E13.5 control (Figure 4B). To characterize this new gene module, we performed pathway

analysis. We found that genes belonging to this GT were most highly associated with regulation of TP53, which

is involved in tumor suppression and inhibits cell cycle progression (Figure 4C). These genes included cell

cycle regulators such as *Trp53inp1* and *Plk2*, which overlap with *Cdkn1a* expression (Figure 4C, Supp Table

1). Cells expressing this gene program were predominantly Wnt-active and restricted to G1/G0 phase (Figure

4B). Based on these findings, we defined this GT as a cell cycle exit GT (Exit GT). By FISH, both *Cdkn1a* and

Trp53inp1 were significantly upregulated in the E13.5 ActSMO dermis compared to E13.5 control dermis,

which showed negligible expression of these genes (Figure 4D). We first measured Wnt activity in the E13.5

ActSMO mutant by measuring *Lef1* levels in *Trp53inp1+* cells. Consistent with the ability of SHH activation to

cell-autonomously upregulate Lef1 levels (Figure 3C), *Trp53inp1+* cells showed significantly higher *Lef1*

transcript levels than E13.5 control upper dermal cells with some *Trp53inp1+* cells showing levels similar to

E14.5 wildtype DC cells (Figure 4D). In contrast to E14.5 wildtype embryos in which *Trp53inp1* and *Cdkn1a*

are largely restricted to *Sox2+* DC cells, these genes were expressed by both *Sox2+* and surrounding *Sox2-*

cells in the E13.5 ActSMO mutant. This indicates that Exit GT genes are expressed not exclusive in cells that

express terminal DC genes in the ActSMO mutant.

We next asked if this gene program can be detected in the E13.5 Actβcat mutant, which also induced cell cycle

exit but not DC genes. GT method also identified a similar GT in the E13.5 Actβcat dermis, which shared

several genes with the E13.5 ActSMO Exit GT, including *Trp53inp1* along with *Cdkn1a*, which was also

upregulated in cells expressing this GT (Figure S4B, S4C). To further compare these GTs across conditions,

we applied a module projection strategy (i.e., label transfer) to assess if a gene program is preserved in a

subset of cells under different conditions. We calculated the module score of the gene set in the other condition

and visualized local neighborhoods of cells that exhibited a high module score, indicating that the average

expression of the gene program is retained in that cell population. This enables the identification of shared or

condition-specific cell states across samples without requiring sample integration, thus avoiding potential

biases introduced by batch correction methods that may obscure true biological differences. Label transfer of

the E13.5 ActSMO Exit GT onto the E13.5 Actβcat and paired control showed that the two mutants share a

similar gene module and molecular cell state, which was not detected in the control sample (Figure 4E).

Moreover, we obtained scRNA-seq data from E13.5 Wntless cKO (*K14Cre;Wls^fl/fl*) embryos, which ablates

epidermal Wnt ligands, resulting in decreased dermal Wnt signaling and a lack of DCs or hair follicles (Figure

S4C) [30,32]. Here, an Exit gene program could not be detected upon label transfer (Figure 4E), indicating that

this Exit GT is Wnt-dependent. These results show that high Wnt activity alone is sufficient to induce genes

that regulate cell cycle exit independent of DC gene expression, and this Exit gene program is Wnt-dependent.

Furthermore, despite the decreased rate of proliferation seen in the Wntless cKO, an Exit GT was not

detected, indicating that this Exit gene program does not simply reflect prolonged G1 phase caused by slower

rates of proliferation.

Although some Sox2+ cells are proliferating in ActSMO dermal clusters at E14.5, most *Sox2*+ cells are

quiescent by E15.5, similar to wildtype E14.5 DC cells (Figure S4E). It was possible that the proliferating cells

in E14.5 ActSMO dermal clusters represent cells that will immediately exit the cell cycle upon cell division

similar to wildtype transitioning cells but continuously over days. To assess if the Exit GT represents a gene

program expressed by cells that are actively exiting the cell cycle, we performed EdU pulse-chase

experiments. First, we observed that most cells in the ActSMO mutant that express *Trp53inp1* at both E13.5

and E14.5 show a low rate of EdU incorporation similar to quiescent DC cells (Figure 4F). We next tested if

these quiescent *Trp53inp1*+ cells are recent progeny of a cell division by pulsing embryos 24 hours prior to

harvest. At both E13.5 and E14.5, there was a significant proportion of *Trp53inp1*+ cells that retained EdU

nucleotide, similar to wildtype E14.5 DC cells, indicating *Trp53inp1*+ cells are quiescent but many are progeny

of a recent cell division. Interestingly, we did not observe an accumulation of *Trp53inp1*+ cells over days

(Figure 4F), suggesting that its expression may be downregulated after a cell has undergone cell cycle exit

(Figure 4F). Thus, in the ActSMO condition, *Trp53inp1*+ cells represent cells that are undergoing cell cycle exit

continuously (Figure 4F). These data also suggest that the Exit GT reflects a process in which cells enter

quiescence as immediate progeny of a cell division similar to the cell cycle dynamics characteristic of the

normal pre-DC-to-DC transition (Figure 4G, Figure S1B).

Loss of GLI3 chromatin binding regulates cell cycle exit downstream of high Wnt activity

To next examine how elevated Wnt signaling promotes cell cycle exit, we first profiled genes differentially

bound by CTNNB1 in the E13.5 Actβcat mutant versus control skin. Although CTNNB1 was more bound at

Wnt target genes (e.g., *Sp5*, *Axin2*, *Lef1*, *Twist2*, *Tcf7*), we did not detect binding to *Cdkn1a* or any Exit GT

genes, nor did we detect significant binding to genes associated with cell cycle inhibition (Figure S5A, S5B).

These results are in keeping with other work showing that Wnt signaling is essential for upper dermal cell

proliferation and indirectly inhibits *Cdkn1a* expression[30]. Given the paradoxical finding that high Wnt activity

promotes cell cycle exit and that this high Wnt signaling is induced by SHH activation (Figure 3B, 3C) [24], we

asked if this cell cycle exit program is specifically induced in the context of SHH pathway mediators. We

reasoned that high Wnt activity and cell cycle exit normally is coupled to high SHH activity and DC gene

expression during DC formation. Hence, SHH effectors may cooperate with Wnt signaling to induce cell cycle

exit. Yet cell cycle exit is observed in the Actβcat mutant at E13.5 and independent of SHH. As *Gli1* is only

expressed upon SHH expression, Gli1 activator would not likely mediate this effect seen in the Actβcat

condition. By contrast, Wnt signaling is known to promote *Gli3* expression independent of SHH in the neural

tube [26]. Similarly, *Gli3* was also expressed in Wnt-active cells in both control and Actβcat dermal cells at E13.5

prior to SHH (Figure 5A). GLI3 acts as a SHH transcription repressor in the absence of SHH and binds to a

broad range of SHH target genes, including cell cycle regulators in the limb bud [33,34]. To profile genes that GLI3

binds and that may be regulated in the Actβcat condition, we performed CUT&RUN to identify GLI3-bound

genomic regions at E13.5. Surprisingly, we found reproducible global reduction of GLI3 binding to chromatin in

the Actβcat condition, indicated by decreased GLI3-bound read density at transcriptional start sites (Figure

5B). As expected, GLI motifs were the most enriched in the differentially bound regions based on HOMER de

novo motif analysis [35] (Figure S5C). KEGG pathway analysis also showed that those unbound regions are

enriched for genes involved in SHH signaling despite the fact that most canonical SHH target genes are not

significantly expressed (Figure 5C). This is consistent with recent work showing that GLI3 binding prior to SHH

is actually inert at many GLI3-bound regions and does not actually repress them[36]. We also found that loss of

GLI3 binding did not derepress many canonical SHH target genes, including *Ptch1* or *Gli1* (Figure S5B). By

western blot analysis, we confirmed that loss of GLI3 binding in the Actβcat condition was not due to

decreased protein expression (Figure 5D). Additionally, Gli3 remained largely expressed in its cleaved

repressor form in both the control and mutant, suggesting other mechanisms (e.g., other post-translational

modifications or cofactors) likely regulate GLI3 binding. These results show that high Wnt signaling alone

unexpectedly induces loss of GLI3 binding at many genomic regions independent of SHH but without

derepression of many of those unbound genes.

Even though many GLI3-unbound regions were not derepressed in the Actβcat mutant, we hypothesized that a

subset of unbound regions may be sufficient to induce cell cycle exit downstream of high Wnt activity. If so,

loss of Gli3 itself should be sufficient to induce cell cycle exit. We genetically ablated *Gli3* in the dermis

(*Axin2CreER;Gli3^{fl/fl}* or Gli3 cKO; tamoxifen E11.5). At E14.5, DCs of Gli3 cKO embryos were smaller than

control DCs (Figure 5E, Figure S5D). Additionally, *Dkk1+* peri-DC progenitors showed aberrant upregulation of

Cdkn1a with lower rates of EdU incorporation compared to controls in which most *Dkk1+* cells do not express

Cdkn1a and are proliferative (Figure 5E, Figure S5D). Examination of E13.5 Gli3 cKO dermis prior to DC

formation also showed that *Cdkn1a* was highly upregulated similar to E13.5 Actβcat embryos and correlated

with a lower rate of EdU incorporation (Figure 5F). We obtained scRNA-seq data from E13.5 Gli3 cKO

embryos and paired controls. Examination of the Gli3 cKO dermal population showed that *Gli3* gene

expression was diminished, but expression of Wnt target genes was not affected, indicating that *Gli3* acts

downstream of high Wnt activity (Figure S5E, S5F). Also, neither canonical SHH target genes nor DC genes

(e.g., *Sox18*, *Gal*) were induced upon loss of *Gli3*, confirming that GLI3 binding does not repress many SHH

target genes (Figure S5E, S5F). Interestingly, GT identified four GTs in the Gli3 cKO dermal scRNA-seq data

(Figure 5G, Figure S5H). Three GTs corresponded to those found in the paired E13.5 control dermal data (UD,

LD, CC). An additional GT identified in the mutant data and not found in the control revealed that several of the

genes contained in this GT overlapped with those found in the Exit GT of the Actβcat mutant, including

Trp53inp1 (Figure 5G). Similarly, most cells within this GT were confined to G1/G0 phase and aberrantly

expressed *Cdkn1a* (Figure 5G). We next intersected genes that were unbound by GLI3 in the E13.5 Actβcat

mutant to genes differentially expressed in the E13.5 Gli3 cKO versus control. Genes involved in cell cycle

inhibition, including *Mxd4* and *Efna5*, were upregulated and unbound in the Gli3 cKO and contained more

active promoter marks (H3K4me3, Figure 5H). We also found Exit GT genes, *Baiap2* and *Cpt1c*, were

unbound by GLI3 and upregulated.

We also transferred the Exit GT label from the ActSMO mutant to the Gli3 cKO, which identified the same Exit

GT population (Figure 5I), indicating that this gene program is shared among all three mutants (ActSMO,

Actβcat, Gli3 cKO). Similar to the ActSMO condition, *Cdkn1a*-high cells in the Gli3 cKO showed a low rate of

active EdU incorporation, while pulsing embryos with EdU 24 hours prior showed that a significant proportion

of *Cdkn1a*-high cells are progeny of a recent division (Figure 5F). These data show that both β-catenin and

Gli3 function in a common pathway that causes cell cycle exit. As DC genes were not expressed in the Gli3

cKO dermis, these data suggest that high Wnt signaling (downstream of SHH co-activation) induces loss of

GLI3 binding specifically to regulate cell cycle exit (Figure 5J).

DC gene expression correlates with Wnt activity levels upon SHH activation

We next examined the E13.5 ActSMO GT that corresponded to DC genes (Figure 6A, 6B). This DC GT

showed that the gene order correlated with known DC genes, including *Gal*, *Sox18* and *Sox2*, which

interestingly, were sequentially activated over DC GT stages in a manner that correlated with increasing *Lef1*

levels (Figure 6C, 6D). This was in contrast to the E14.5 wildtype DC GT in which DC genes were

simultaneously activated at the last stage when both Wnt and SHH activity are highest. As SHH target genes

are uniformly expressed across the dermis and no longer progresses with DC differentiation, canonical SHH

pathway genes were not detected in the ActSMO DC GT. These data show that high SHH activity can induce

DC genes in cells with varying levels of Wnt activity and not only in cells that have high levels of Wnt signaling

(Figure 6E). However, the level of DC genes expressed corresponds with the level of Wnt activity. This implies

that the synchronization of cell cycle exit with high DC gene expression seen in the wildtype scenario may

occur by restricting high SHH activity to cells with high Wnt signaling. As cell cycle exit occurs in cells that do

not express high DC genes in the ActSMO condition, the level of Wnt signaling necessary for cell cycle exit

would be lower than that required to express high DC genes (Figure 6E).

Coupled Wnt and SHH gradients is sufficient to synchronize cell cycle exit with molecular

differentiation

Based on the above results, cell cycle exit and expression of high DC genes are only seen in cells with coupled

high Wnt and high SHH activity induced by placode SHH ligand. We asked if inducing SHH ligand expression

across the epidermis is sufficient to accelerate Wnt activity and to couple high levels of Wnt signaling with high

SHH activity without placodes. Here, we genetically induced a covarying gradient of Wnt and SHH activation in

the upper dermis by expressing SHH ligand uniformly across the epidermis where Wnt ligands are normally

expressed prior to hair follicle morphogenesis (*K14Cre;Rosa-LSL-SHH* or SHH OE; Figure 7A). ScRNA-seq

analysis of E13.5 SHH overexpressing (SHH OE) mutant and paired control dermal cells showed that both

Lef1 and *Ptch1* were expressed in an overlapping covarying pattern over the UMAP cell embedding in the

mutant (Figure 7B). Cells with the highest levels of *Lef1* and *Ptch1* also expressed DC genes (e.g., *Gal*, *Sox18*,

Sox2). Additionally, *Cdkn1a* was largely localized to cells expressing DC genes in contrast to ActSMO dermal

cells in which many *Cdkn1a+* cells do not express many DC genes (Figure 7B, 7C). Notably, GT analysis

showed that only three GTs were identified that corresponded to similar gene programs identified in E14.5

wildtype dermal cells (i.e., CC, LD, DC; Figure 7D, Figure S7A). A separate Exit GT was not detected. Similar

to the E14.5 wildtype DC GT (Figure 2C), the E13.5 SHH OE DC GT showed a gene order in which Wnt and

SHH levels correlate with each other and with DC genes, which were restricted to the last stage (Figure 7D).

Cdkn1a was also activated together with DC genes (Figure 7D). Additionally, label transfer of the ActSMO Exit

GT did not detect a separate population in the SHH OE or control dermis (Figure 7E). As expected, *Lef1*

expression was accelerated in the SHH OE compared to the E13.5 control, with the highest levels

corresponding to arrest and DC genes (Figure 7F, S7B, S7D). Taken together, these data suggest that

covarying levels of Wnt and SHH activity synchronize cell cycle exit with DC genes and recapitulate the

wildtype pre-DC-to-DC transition.

Quantification of *Lef1* and *Ptch1* levels by spatial dermal layer showed that both genes covary with higher *Lef1*

levels in the first two layers of the dermis in the SHH OE compared to controls, consistent with SHH

accelerating *Lef1* levels (Figure 7F). By FISH, we found DC genes were restricted to the most superficial

dermal layers where both *Lef1* and *Ptch1* levels were highest and where proliferation is low (Figure 7F, Figure

S7B-C). We next performed GLI3 CUT&RUN in the mutant and control and found global loss of GLI3 binding

at genomic regions similar to the Actβcat mutant (Figure 7G, Figure S7D), indicating that loss of GLI3 binding

represents an endogenous molecular event during DC genesis and not simply an artifact of genetically

stabilizing β-catenin. As expected, we found loss of GLI3 binding at SHH target genes (*Ptch1*, *Gli1*) but now

with increased transcriptional activity as marked by increased H3K4Me3 signal at these genes. This suggests

that Gli activator may be required for expression of most SHH target genes and that GLI3 may function to

prime those sites for GLI activator upon SHH expression. We also detected loss of binding at cell cycle

regulators, *Mxd4* and *Efna5* similar to the Actβcat mutant (Figure 7H).

We next asked if we can modulate Wnt and SHH levels to recapitulate the interaction between these signals

and their regulated biological processes by establishing an in vitro dermal culture system. Previous attempts to

culture dermal papilla cells used neonatal dermal cells that contain preformed DC/dermal papilla cells[37,38].

However, they show limited ability to maintain DC/DP markers and immediately lose Wnt, SHH, and DC gene

expression. Further, these cells show poor responses to Wnt or SHH agonists. We found similar results using

neonatal dermal cells with only a mild increase in *Lef1* or *Ptch1* expression upon Wnt agonism with

CHIR99021 (CHIR) or SHH agonism (SAG), respectively (Figure S7E). Consistent with a previous report, we

found HDAC inhibitors incrementally augmented CHIR *Lef1* responses but had no effect on SAG responses

(Figure S7F). We next cultured dermal cells FACS isolated from E13.5 PDGFRαH2BGFP embryos. In contrast

to neonatal fibroblasts, E13.5 dermal cells showed robust dose-dependent target gene responses to either

CHIR or SAG after 48-72 hours of culture (Figure S7E).

Using this model, we asked if SHH activation can augment Wnt signaling by assessing the effect of SAG on

CHIR-induced *Lef1* gene expression. We found that SAG treatment boosted *Lef1* expression in CHIR-treated

cells (Figure 7I). Moreover, the fold increase in *Lef1* expression was significantly more in cells treated with

higher levels of CHIR (5 μM vs 2 μM) upon SAG treatment, suggesting that the amount of Wnt acceleration

depends on the initial levels of Wnt activity (Figure 7I). Consistent with in vivo results, high levels of CHIR

(5 μM) alone are sufficient to induce *Cdkn1a* expression independent of SHH activation (Figure 7J), while co-

treatment with CHIR and SAG induced DC genes such as *Gal* and *Sox18* (Figure 7K). Importantly, consistent

with our in vivo E14.5 Actβcat results (Figure S7G), DC gene induction was SAG-dependent, indicating that

higher levels of SHH activation is essential for DC gene expression. Altogether, these results demonstrate that

levels of Wnt and SHH activation can be tuned to coordinate cell cycle exit and DC gene expression and can

be used to recapitulate these events in vitro.

Discussion:

Our results fit with a scenario in which cell cycle exit is tightly coupled to differentiation. The decision to exit the

cell cycle may be made in G2/M phase or in G1 phase. To explain this coupling, previous work have proposed

a direct model in which cell cycle regulators can have a dual role in regulating cell cycle progression and cell

type differentiation (e.g. activating cell type-specific transcription factors). Here, we reveal an unexpected

mechanism in which these two processes are independently mediated but are coupled to each other due to the

mere coupling of Wnt and SHH activity. In the DC system, Wnt activity levels are mechanistically dependent

Model: Wnt and SHH ligands are secreted from epidermal cells, creating gradients of pathway activation in underlying dermal cells. SHH activation in the dermis further accelerates Wnt signaling, resulting in loss of GLI3 chromatin binding which is sufficient to cause cell cycle arrest. Concurrently, SHH activation in these high Wnt activity cells can induce DC marker genes, thereby coupling DC differentiation and cell cycle exit below the hair placode.

on SHH activation and the reason for the synchronization of cell cycle exit and molecular DC differentiation, yet the processes themselves are separable because the threshold level of Wnt activity required for cell cycle exit is less than that required for expressing maximal DC genes (Figure 6). This contrasts other models in which the processes are coupled to each other by co-dependent downstream mediators. Our results also demonstrate that two covarying gradients do not merely reflect overlapping independent signals induced by a common ligand source. Rather, the spatial gradients seen during DC formation, in part, reflect the dynamic interaction between Wnt and SHH to regulate each other and to coordinate distinct biological processes over a cell fate

transition. Specifically, we show that SHH is sufficient to cell-autonomously amplify Wnt activity, and the high

levels of Lef1 normally seen in DC cells may be caused by SHH co-activation rather than by local extrinsic

placode Wnt ligands. In the wildtype scenario, only a subset of cells that have sufficient pre-existing Wnt

activity (dermal layer 1 or peri-DC cells) will induce cell cycle exit and expression of all DC genes

synchronously once their Wnt activity is accelerated by SHH co-activation (Figure 7L). Therefore, the

coordination of Wnt and SHH signaling levels in pre-DC progenitors is likely due to the spatial proximity of

these cells to epidermal cells that produce both SHH and Wnt ligands. However, the acceleration of Wnt

signaling that initiates the pre-DC-to-DC transition is independent of epidermal Wnt ligands. How SHH boosts

Wnt activity in dermal cells is unclear, but it is likely indirect, as we did not find that GLI transcription factors

bind to Wnt target genes, including *Lef1*. We posit that this model may also help explain how other scenarios

can time proliferation with differentiation. For example, hair follicle epithelial stem cells rely on Wnt and SHH

activation during anagen phase and give rise to transit-amplifying cells that undergo proliferation and

differentiation before they terminally arrest into different spatially organized cell types. It is possible that relative

levels of Wnt and SHH (or other signals) regulate the coordination of these two processes to time when a

progenitor exits the cell cycle and the specific cell type genes it expresses. From a different perspective, when

distinct biological processes of a cell fate transition occur synchronously, there may be few intermediate states

to capture the transition, giving the appearance that a population emerged from "thin air". Determining the

individual biological processes that induce a cell fate transition and why those processes are synchronized can

help address how some patterned structures appear to from without a pre-existing pattern.

In the ActSMO condition, uniformly high SHH activation was induced in cells with varying Wnt-activity, and the

number and expression level of DC genes correlated with Wnt signaling levels. However, in the wildtype

condition, DC marker genes are only expressed in cells that show both high Wnt and high SHH activity. This

suggests that a threshold level of SHH activity, which is induced in the ActSMO mutant, is required for DC

gene expression. It is plausible that high Wnt activity can reciprocally accelerate SHH activity to ensure that

only cells with high Wnt activity activate high SHH signaling to induce DC genes. Consistent with this, we

observed that cells in the peri-DC region of the E14.5 Actβcat mutant exhibited higher levels of Wnt activity

than control peri-DC cells but also higher *Ptch1* levels than in the control, suggesting that Wnt activity levels

may regulate SHH activity. Such an intricate feedback mechanism between Wnt and SHH signaling may

deterministically drive a rapid sequence of cell cycle exit and DC gene expression. Further studies to

temporally regulate Wnt and SHH activity can give insight into the intimate relationship between dermal Wnt

and SHH signaling.

How is cell cycle exit induced upon high Wnt activity? Wnt signaling is known to promote G1-S transition and is

typically associated with stemness and proliferation [7,39-41]. Consistent with this, we and others showed that loss

of dermal Wnt activity prior to SHH results in decreased dermal cell proliferation with concomitant loss of

dermal cell density [25,30]. We found that inducing high Wnt activity, above that seen prior to SHH, results in cell

cycle exit. This high Wnt signaling unexpectedly results in a global reduction in GLI3 binding to chromatin, and

loss of Gli3 is sufficient to mediate cell cycle exit. Interestingly, canonical SHH target genes such as *Ptch1* or

Gli1 were not derepressed despite loss of GLI3 binding at those sites in the Actβcat mutant. These genes

become expressed upon SHH, suggesting GLI activator (e.g., GLI1) is required to activate those GLI sites. It

remains unclear how high Wnt signaling causes GLI3 unbinding. Other transcription co-factors or other GLI3

post-translational modifications may regulate GLI3 binding. Prior to SHH, Wnt signaling may promote GLI3

binding at canonical and non-canonical SHH target genes. It is possible that upon SHH, cell cycle exit is

immediately induced by GLI3 unbinding (after SHH-induced Wnt amplification) while DC and SHH target genes

may be activated soon after upon *Gli1* expression and GLI1 binding to those unbound regions. The global loss

of GLI3 binding upon acceleration of Wnt signaling could serve as mechanism to couple high Wnt activity with

efficient upregulation of SHH activity. This would ensure that DC genes are only expressed coincident with or

just after cell cycle exit. In the ActSMO condition, molecular DC differentiation occurs while cells are still

proliferating. These dermal clusters are diffuse with fuzzy borders and largely fail to contact the epidermis and

do not induce placodes. We hypothesize that the clustering of DC cells and their ability to communicate with

the epidermis relies on their ability to stop dividing and to form stable dense clusters. It will be interesting to

assess if cell-cell adhesion genes or extracellular matrix genes are affected by cell cycle regulators.

This study also underscores the utility of computational methods to give mechanistic insight into complex cell

fate transitions by separating out independent gene programs. This method was useful to identify Gli3 as a

molecular mediator that regulates a specific subpopulation of G1/G0 phase cells, which represent immediate

quiescent progeny of a division. The possibility of distinguishing heterogeneity within G1/G0 phase cells during

lineage specification holds the potential to uncover mechanisms that dictate if and when a cell will undergo cell

cycle arrest as well as how differentiation is balanced with cell divisions (e.g., asymmetric cell divisions in stem

cells and transit amplifying cells). From a translational perspective, this study sheds light on how distinct

biological processes of DC formation can be tuned to grow and differentiate DC cells *in vitro*, a critical step

towards making DC organoids and organotypic hair follicles to model hair growth and disease.

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Peggy Myung (peggy.myung@yale.edu).

Data and Code Availability

Genomic datasets used in this study are deposited in NCBI Gene Expression Omnibus database and are publicly available as of the date of publication (GSE280825, token: qfirggaarvypzet). For other all other scRNA-seq data presented in this study, the data will be deposited in a GEO repository and will be made available with a token upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

No statistical methods were used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment.

Mice

Axin2CreER [42] mice were bred to *Rosa-LSLSmoM2YFP* [43], *β-catenin*[fl(Ex3/+)] [44] and *Gli3*[fl/fl] [45] mice. *K14Cre* [46] mice were bred to *Shh* Overexpressor (Want et al. 2010) mice. *Rosa-Flag-Gli3YFP* [33] mice were bred to *Axin2CreER* and *β-catenin*[fl(Ex3/+)] mice to enable detection of GLI3-Flag using CUT&RUN. *R26Fucci2aR* [47] reporter mice were used to visualize cell cycle phase. A random population of both male and female mice were used for all experiments. All procedures involving animal subjects were performed under the approval of the Institutional Animal Care and Use Committee of the Yale School of Medicine.

Acknowledgements: We thank D. Gay, M. Ito, and R. Atit for review and discussion of this manuscript. This work was supported by National Institutes of Health grants R01AR076420 (P.M.)and LEO Foundation LF-OC-23-001347 (P.M.).

Author Contributions: Conceptualization, P.M., Y.K., R.L., Y.J., S.P.; Methodology, P.M., R.L., Y.K., Y.J., S.P.; Software, R.L., Y.K.; Resources, M.T., D.W., K.P.; Investigation, P.M., R.L., Y.J., S.P. H.L, S.V.; Writing –

578   Original Draft, P.M.; Writing – Review & Editing, P.M., R.L., Y.J., S.P., Y.K., C.L., J.L., E.B.; Supervision, P.M.,

579   Y.K.; Funding Acquisition, P.M.

  Main Figure Titles and Legends:

  Figure 1: Cell cycle exit and molecular differentiation are tightly coupled during dermal condensate

  development

  (A) Hair follicle development over time. **(B)** FISH images of E13.5 and E14.5 flank skin with *Sox2*+ DC with

  lack of EdU copositivity at E14.5. **(C)** Skin whole-mount image of E14.5 Sox2eGFP skin showing GFP

  colocalization with Fucci2 reporter. **(D)** FISH stained for indicated genes of E14.5. **(E)** FISH of E13.5 wildtype

  skin stained for Lef1, Ptch1 or EdU after 1-hour EdU pulse, quantification of *Lef1* and *Ptch1* levels (H-score) by

  dermal cell layer, %EdU+ by *Lef1* bin **(F)** Cartoon depicting graded *Lef1* expression spatially. **(G)** FISH of

  E14.5 skin stained for *Lef1*, *Ptch1* or EdU; quantification of *Lef1* and *Ptch1* levels by indicated spatial regions,

  %EdU+ by *Lef1* bin. **(H)** Cartoon depicting *Lef1* and *Ptch1* expression levels spatially; n=5. Data as mean±

  SEM (n=5 per day); *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, one-way ANOVA; ns, not significant. Scale

  bars, 50 µm. Related to Figure S1.

  Figure 2: High Wnt activity induces cell cycle exit independent of SHH

  (A) Cartoon of Wnt activity levels in E13.5 wildtype or Actβcat mutant (tamoxifen E11.5). **(B)** UMAP colored by

  expression of indicated genes or cell cycle (CC) phase for E13.5 Actβcat mutant or paired control dermal

  scRNA-seq data. **(C)** FISH showing E13.5 *Lef1*, *Cdkn1a*, EdU (left) and quantification of FISH *Cdkn1a* levels

  (H-score) or %EdU of *Cdkn1a* (p21)+ cells. **(D)** Cartoon of cell cycle and DC molecular differentiation dynamics

  in E13.5 Actβcat showing cells exit the cell cycle but do not express DC genes. **(E)** FISH showing E14.5 *Sox2*,

  Cdkn1a, EdU (left) and FISH quantification of %EdU+ cells within indicated control or Actβcat populations. **(F)**

  FISH showing E14.5 *Lef1*, *Ptch1*, *Sox2* (left) and FISH quantification of *Ptch1* level within indicated control or

  Actβcat populations. **(G)** Cartoon of E14.5 Actβcat condition in which quiescence can occur without DC gene

  expression. Related to Figure S2.

  Figure 3: SHH cell-autonomously augments Wnt signaling to induce cell cycle exit

  (A) Cartoon of experimental condition induced by expressing a constitutively activated form of Smo. Whole-

  mount image of E14.5 ActSMO and control and quantification of %EdU+ of Sox2+ cells at E14.5. **(B)** FISH

  showing indicated genes in control or ActSMO mutant (n=4). *Lef1* levels by **(C)** *Ptch1* status or **(D)** %EdU by

Lef1 levels (bin) **(E)** Cartoon showing that SHH activation cell-autonomously boosts Lef1 levels. **(F)** Expression

level of indicated DC genes by DC GT stage and UMAPS with indicated genes. **(G)** DC gene expression by

DC GT stage of the ActSMO mutant and UMAPs. **(H)** *Cdkn1a* expression and G1/G0 fraction by DC GT stage.

(I) *Lef1* and *Ptch1* levels by DC GT stage. **(J)** Cartoon of covarying Wnt and SHH activity across the pre-DC-

to-DC transition that corresponds to synchronous cell cycle exit and DC gene expression. Data as mean±

SEM; *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, one-way ANOVA. Scale bars, 50 µm. Related to Figure

S3.

Figure 4: A Wnt-dependent gene program that corresponds to cell cycle exit is identified by

GeneTrajectory

(**A**) UMAPs of control and ActSMO colored by indicate genes. (B) GT analysis of E13.5 ActSMO and paired

control dermal scRNA-seq data showing four GTs and DC GT of ActSMO mutant which is not present in

control. Exit GT of ActSMO colored by GT stage or indicated genes. (C) Pathway analysis of Exit GT. UMAPs

of control and ActSMO colored by indicate genes. (D) FISH showing indicated genes of E13.5 control or

ActSMO mutant skin; corresponding quantification of *Trp53inp1* or *Cdkn1a* levels. (E) Label transfer of E13.5

ActSMO Exit GT onto indicated E13.5 mutants. **(F)** EdU pulse-chase experiments of E13.5-E14.5 ActSMO

mutant and corresponding quantification of %EdU positive of *Trp53inp1*+ cells. (**G**) Cartoons depict the

interpretation of cell cycle dynamics of *Trp53inp1*+ cells which represent cells that are immediate progeny of a

division. Data as mean ± SEM, one-way ANOVA. Scale bars, 50 µm. Related to Figure S4.

Figure 5: Loss of GLI3 chromatin binding regulates cell cycle exit downstream of high Wnt activity

(A) FISH image of E13.5 wildtype skin showing Gli3 expression prior to SHH. **(B)** Heat map of GLI3 binding at

all transcriptional start sites (TSS) in E13.5 control or Actβcat mutant skin. **(C)** KEGG pathway analysis of GLI3

unbound regions in the Actβcat mutant. (D**)** Western blot of cell lysates from control of Actβcat mutant stained

for anti-Flag (Gli3) or b-actin. (**E**) FISH of E14.5 control and Gli3 cKO for indicated genes or EdU; quantification

of #Sox2+ cells per DC and Cdkn1a transcript levels by skin population. **(F)** FISH images of E13.5 control and

Gli3 cKO with indicated genes or EdU; Quantification of Cdkn1a transcripts or %EdU by population after 1-hour

EdU pulse or after 24-hour pulse-chase. **(G)** GT analysis of E13.5 Gli3 cKO showing an Exit GT. **(H)** Volcano

plot of differentially expressed genes in the Gli3 cKO vs control and those genes that overlap with GLI3

unbound genes in the Actβcat mutant (blue); genome browser tracks showing binding signal of respective

antibodies at the promoter region of indicated genes. **(I)** Label transfer of E14.5 Actβcat Exit GT onto indicated

mutant scRNA-seq data. **(J)** Cartoon showing model of mechanism by which high Wnt activity induces cell

cycle exit. Data as mean± SEM; *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, one-way ANOVA; ns, not

significant; scale bars, 50 µm. Related to Figure S5.

Figure 6: DC gene expression correlates with Wnt activity levels upon SHH activation

(A)GT analysis of E13.5 ActSMO and paired control dermal scRNA-seq data showing four GTs and DC GT

of ActSMO mutant which is not present in control. **(B)** DC gene expression by DC GT stage of the ActSMO

mutant and UMAPs. **(C)** Expression level of indicated DC genes by DC GT stage and UMAPS with indicated

genes. **(D)** Lef1 levels by DC GT stage. **(E)** Cartoon showing that SHH activation cell-autonomously boosts

Lef1 levels. Data as mean± SEM; *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, one-way ANOVA. Scale bars,

50 µm. Related to Figure S6.

Figure 7: Coupled Wnt and SHH gradients is sufficient to synchronize cell cycle exit with molecular

differentiation

(A) Cartoon of wildtype scenario showing covarying gradients and experimental scenario to induce covarying

Wnt and SHH gradients by expressing SHH ligand coordinately with Wnt ligands in the epidermis. (**B**) UMAPs

colored by gene or cell cycle phase of control and E13.5 SHH OE mutant dermal cells. (**C**) Quantification of the

fraction of *Cdkn1a+* cells that co-express DC genes (Gal) in ActSMO vs SHH OE. **(D)** GT analysis of E13.5

SHH OE dermal scRNA-seq data. **(E)** Label transfer of E13.5 ActSMO Exit GT onto SHH OE and control

dermal populations. **(F)** Lef1 levels by DC GT stage; FISH image showing spatially covarying gradients of

Ptch1 and Lef1 that corresponds to dermal layer. **(G)** Heat map of GLI3 binding at all TSS in E13.5 control

versus SHH OE showing global loss of binding. **(H)** Genome browser tracks showing binding signal of

respective antibodies at the promoter region of indicated genes. **(I)** qPCR for *Lef1* of E13.5 PDGFRaH2B

dermal cells after SAG and/or CHIR stimulation for 48 hours (n=4). **(J)** qPCR for *Cdkn1a* after 48 hours. (K)

qPCR for DC genes after 48 hours of culture with CHIR and/or SAG. **(L)** Model of how gradient levels of Wnt

664   and SHH interact dynamically to coordinate DC differentiation gene expression and cell cycle exit and that the

665   covariance of the two signals is responsible for the synchronization of cell cycle exit and DC genes during DC

666   genesis. Data as mean ± SEM, one-way ANOVA; scale bars, 50 μm. Related to Figure S7.

667
668

669 **STAR Methods**

670 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-IgG	EpiCypher	Cat#13-0042; RRID: AB_2923178
Rabbit anti-H3K4Me3	EpiCypher	Cat#13-0041; RRID: AB_3076423
Rabbit anti-H3K27Ac	EpiCypher	Cat#13-0059; RRID: AB_2923489
Chicken anti-GFP	Abcam	Cat#Ab13970; RRID: AB_300798
Rabbit anti-Sox2	Abcam	Cat#Ab97959; RRID: AB_2341193
Mouse anti-Flag	Millipore	Cat#F1804; RRID: AB_262044
Rabbit anti-RFP	Rockland	Cat# 600-401-379; RRID: AB_2209751
Rabbit anti-CTNNB1	Antibodies-online	Cat#ABIN2855042; RRID: AB_10825282
Donkey anti-rabbit Alexa Fluor 488	Thermo Fisher Scientific	Cat#A21206; RRID: AB_2535792
Goat anti-chicken Alexa Fluor 488	Thermo Fisher Scientific	Cat#A11039; RRID: AB_2534096
Donkey anti-rabbit Alexa Fluor 568	Thermo Fisher Scientific	Cat#A10042; RRID: AB_2534017
Chemicals, peptides, and recombinant proteins		
Hoechst	Invitrogen	H3570
DAPI solution	Novus Biologicals	Cat#NBP2-31156
Tamoxifen	Sigma Aldrich	Cat#T5648
EdU (5-ethynyl-2-deoxyuridine)	Thermo Fisher Scientific	Cat#E10187
CHIR99021	Sigma-Aldrich	Cat#SML-1046
SAG	Sigma-Aldrich	Cat#SML-1314
Critical commercial assays		
Click it EdU Imaging Kit 555	Invitrogen	Cat#C10338
Click it EdU Imaging Kit 488	Invitrogen	Cat#A10266
RNAscope Multiplex Fluorescent Detection Kit v2	ACDBio	Cat#323270
Deposited data		
Raw sequence data files	NCBI Gene Expression Omnibus	GEO: GSE280825
Experimental models: Organisms/strains		
Mouse: β−catenin EX3/+	Harada et al., 1999	N/A
Mouse: Axin2CreER	Amerongen et al., 2012	N/A
Mouse: Fucci2	Abe et al., 2013	N/A
Mouse: K14Cre	Dassule et al., 2000	N/A
Mouse: Wls flox/flox	Carpenter et al., 2010	N/A
Mouse: Sox2eGFP	Arnold et al., 2011	N/A
Mouse: R26R-LSL-SmoM2YFP	Jeong et al., 2004	N/A
Mouse: Gli3 flox/flox	Blaess S et al., 2008	N/A
Mouse: $Gli3^{3XFlag}$	Lopez-Rios J , et al., 2014	N/A
Mouse: *lox-EGFP-STOP-lox-Shh*	PI: David Wang, University of Michigan Ann Arbor	N/A
Mouse: PDGFRaH2BGFP	Hamilton et al., 2003	N/A
Oligonucleotides		
Mouse-Cdkn1a	ACDBio	Cat#408551
Mouse-Ptch1	ACDBio	Cat#402811
Mouse-Lef1	ACDBio	Cat#441861
Mouse-Sox2	ACDBio	Cat#401041
Mouse-Trp53inp1	ACDBio	Cat#1161531

Mouse-Gli3	ACDBio	Cat#445511
Mouse-Dkk1	ACDBio	Cat#402521
Mouse-Gal	ACDBio	Cat#400961
Software and algorithms		
Adobe photoshop CC 2024	Adobe Systems	http://www.adobe.com/products/photoshop.html
Adobe Illustrator CC 2024	Adobe Systems	http://www.adobe.com/products/illustrator/features.html
Fiji	National Institutes of Health	https://imagej.net/software/fiji/
R	R Core Team	https://www.r-project.org/
rARPACK	R Package	https://cran.r-project.org/web/packages/rARPACK/index.html
Differential Abundance analysis	Landa et al., 2020	https://arxiv.org/abs/2011.03418
Seurat	Macosko et al., 2015	http://satijalab.org/seurat/
GraphPad Prism 9.1.0	GraphPad software	https://www.graphpad.com/scientific-software/prism/

METHOD DETAILS

Tamoxifen induction of mice

Embryos were staged as days post coitum, with embryonic day E0.5 considered as noon of the day a vaginal

plug was detected after overnight mating. Pregnant dams were given one dose of Tamoxifen dissolved in corn

oil (20mg/ml, Sigma) at 40-60 µg/gm body weight by oral gavage at E11.5 as indicated.

EdU Incorporation Assay

To assess active proliferation, EdU was administered to pregnant mice intraperitoneally (25 µg/gm) and

embryos were harvested after 1.5 hour. For pulse-chase experiments, 25 µg/gm EdU was administered to

pregnant mice at indicated times and embryos were harvested at later times. EdU incorporation was assessed

using the Click-it EdU Imaging kit Alexa 555 or Alexa 488 (Life technologies, c10338) according to

manufacturer's instructions. Briefly, skin explants were treated with a mixture of 1X Click-iT reaction buffer,

$CuSO_4$, Alexa Fluor azide dye and 1X reaction buffer additive, all provided in the Click-it EdU Imaging Kit, for

30 minutes at RT before washing in PBS.

Histology

10% formalin-fixed paraffin embedded (FFPE) whole embryos were used for histological analysis. FFPE

embryos were sectioned at 10 µm thickness for FISH.

Whole mount immunofluorescence

Dorsolateral skin from embryos were microdissected and placed on nucleopore filters (VWR, WHA 800281)

and fixed overnight with 4%PFA at 4° C. Skin explants were blocked for 6 hours in 5% normal donkey serum,

1% bovine serum albumin and 0.5% Triton-X100 at RT. Explants were incubated overnight at 4° C in the

following primary antibodies: rabbit anti-Sox2 (1:400, Abcam, ab97959), chicken anti-GFP (1:500, Abcam,

ab13970), rabbit anti-RFP (1:500, Rockland, 600-401-379). Explants were then washed in 0.2% Tween20/PBS

for 4-6 hours on a rotator and then incubated with the following secondary antibodies: Alexa Fluor 568-donkey

anti-rabbit (1:200, Invitrogen, A10042), Alexa Fluor 488-donkey anti-rabbit (1:200, Invitrogen, A21206), Alexa

Fluor 488-goat anti-chicken (1:200, Invitrogen, A11039) overnight at RT. Washes were carried out for 6 hours

with 0.2% Tween20/PBS. Hoechst (1:750 in PBS) was used for 45 min before mounting on slides with

SlowFade Gold Antifade Mountant (Invitrogen, S36936).

In-situ hybridization

The RNAscope Multiplex Fluorescent Detection Kit v2 (ACDBio, 323110) was used for single-molecule

fluorescence in situ hybridization (FISH) according to the manufacturer's protocol. Briefly, sections were

deparaffinized and permeabilized with hydrogen peroxide followed by antigen retrieval and protease treatment

before probe hybridization. After hybridization, amplification and probe detection were done using the Amp 1–3

reagents. Probe channels were targeted using the provided HRP-C1-3 reagents and tyramide signal

amplification VIVID fluorophores—650 and 570 (ACD biotechne, 323271). EdU staining was done using the

Click-it EdU Imaging Kit Alexa 488 (Life Technologies, c10338) according to the manufacturer's instructions.

Nuclear counter-stain was done using Hoechst 33342 (Invitrogen, H3570) before mounting with SlowFade

Mountant. RNA scope probes used (ACDBio)—Mm-Lef1 (441861), Mm-Ptch1 (402811), Mm-Dkk1 (402521),

Mm-Sox2 (401041), Mm-Gal (400961), Mm-Gli3 (400961) and Mm-Trp53inp1 (1161531).

Microscopy

FISH paraffin-embedded images were acquired using the Leica Stellaris 8 DMi8 confocal microscope with a

×40 oil immersion (Numerical Aperture 1.3) objective lens, scanned at 5 µm thickness, 1, 024 × 1, 024 pixel

width, 400Hz. Whole mount explants were imaged in 3 dimensions using the LaVision TriM Scope II (LaVision

Biotec) microscope equipped with a Chameleon Vision II (Coherent) two-photon laser (810 -1000 nm) to

acquire *z*-stack images ranging from 50-120 µm (2 µm serial optical sections) using a 20X water immersion

lens (NA 1.0; Olympus), scanned with a field of view of 0.3–0.5 mm^2 at 800 Hz.

Single-cell dissociation

Embryonic dorsolateral/flank skin was micro-dissected and pooled after genotyping (3 embryos per condition)

and dissociated into a single-cell suspension using 0.25% trypsin (Gibco, Life Technologies) for 14 minutes at

37° C. Single-cell suspensions were then stained with DAPI (Fisher Scientific, NBP2-31156) just prior to

fluorescence-activated cell sorting.

<u>Fluorescence-activated cell sorting</u>

DAPI-excluded live skin cells were sorted on a BD FACS Aria II (Biosciences) sorter with a 100 µm nozzle.

Cells were sorted in bulk and submitted for 10X Genomics library preparation at 0.75-1.0x10^6/mL concentration

in 4% FCS/PBS solution.

<u>Single-cell RNA sequencing and library preparation</u>

Chromium Single cell 3' Library and Gel Bead Kit v2 Chromium Single Cell 3′ Library & Gel Bead Kit v2 (PN-

120237), Chromium Single Cell 3′ Chip kit v2 (PN-120236) and Chromium i7 Multiplex Kit (PN-120262) were

used according to the manufacturer's instructions in the Chromium Single Cell 3′ Reagents Kits V2 User

Guide. After cDNA libraries were created, they were subjected to Novaseq 6000 (Illumina) sequencing.

<u>10X scRNA-seq data pipeline to matrix</u>

The transcriptomes of live single cells from E13.5 or E14.5 mouse skin samples under different biological

conditions were sequenced. Raw 10X sequencing data was processed into a matrix employing the standard

10X CellRanger pipeline. Briefly, base call files were fastq format which were aligned to the mm10 reference

genome followed by nUMI and barcode counting, constructing the nUMI count matrices. nUMI matrices were

filtered, centered and normalized using Seurat@1. Briefly, for each original run condition (e.g., E13.5 wildtype),

cells with > 1,000 detected genes, mitochondrial content < 25%, and ribosomal content > 5% (calculated via

PercentageFeatureSet in Seurat) were retained. Doublets were identified and removed using scDblFinder@2.

Data were then log-normalized by total UMI count.

<u>Principal Component Calculation and UMAP Visualization</u>

Principal components were computed using Seurat's RunPCA on the top 2,000 variable genes. The number of

PCs retained was determined from the elbow plot@7, and used for downstream analysis, including

dimensionality reduction for visualization and clustering.

<u>Cell type specification</u>

For each original run, UMAP was performed on log-normalized, centered, and scaled UMI count matrices

using the retained PCs. Unsupervised clustering was then conducted using Seurat's FindClusters function with

a resolution of 0.5. Clusters expressing Col1a1 were defined as dermal, while clusters expressing Lef1 or Dkk1

were defined as upper-dermal.

Gene trajectory inference

We applied GeneTrajectory@3 to each sample to identify multiple, independent gene dynamic processes. For

each sample, we followed the recommended guidelines. We selected the top 2,000 highly variable genes

(identified using Seurat's FindVariableFeatures) that were expressed in >1% and <50% of cells as input. The

top 20PCs were used to compute a diffusion map, and the top 10 diffusion components were used to construct

a cell k-nearest neighbor graph (k = 10).

The number of gene trajectories and the time steps for gene inclusion within each trajectory were determined

interactively by inspecting the top three or more gene embeddings. For the samples analyzed in this study, we

identified the following gene trajectories and associated time steps (t):

• E14.5 wild type: 3 trajectories (t = 13, 10, 5)

• E13.5 SmoM2: 4 trajectories (t = 1, 3, 2, 1); control: 3 trajectories (t = 3, 2, 2)

• E13.5 TG: 3 trajectories (t = 6, 3, 3); control: 3 trajectories (t = 5, 3, 3)

• E13.5 EX3: 4 trajectories (t = 3, 3, 2, 2); control: 3 trajectories (t = 1, 3, 3)

• E13.5 Gli3-KO: 4 trajectories (t = 8, 7, 6, 1); control: 3 trajectories (t = 4, 3, 1)

Gene bins and cell stages

Following GeneTrajectory guidelines, each trajectory from mutant samples was divided into 5 or 6 gene bins.

For each bin, a gene bin score was computed for both mutant and control samples to compare gene

expression dynamics across the cell embedding. Cell involved in each trajectory were stratified into the same

number of stages, based on whether they expressed more than 50% of the genes in the corresponding gene

bin. Cells assigned to multiple stages were resolved by assigning them to the stage closer to the terminal end

of the trajectory (i.e., farther from the branching point).

Exit GT Score

Exit GT scores for each sample were calculated using Seurat's AddModuleScore, based on all genes from the

exit gene trajectory identified in E13.5 SmoM2.

<u>Pathway enrichment analysis</u>

Reactome pathway enrichment analysis was performed using ReactomePA@4, with pathways showing p <

0.05 considered significantly enriched.

<u>Cell cycle estimation of scRNA-seq data</u>

We employed Seurat's cell cycle scoring [48]. Briefly, averaged relative expression of those cell cycle related

genes were used to calculate G2/M and S scores, which were used for binning cells into G2/M, S and G1/G0

bins.

<u>Data merge and normalization</u>

For comparative analysis, we merged upper-dermal cell matrices from each pair of conditions separately (i.e.

E13.5 control and Gli3-KO), followed by standard preprocessing and normalization using the workflow provided

by Seurat.

<u>Differential abundance and Differential gene expression analysis</u>

Differential abundant cells were identified using DA-seq@6, with a DA score threshold > 0.8. Differential gene

expression analysis between DA cells from E13.5 Gli3-KO and all other upper-dermal cells from both E13.5

Gli3-KO and control was performed using Seurat's FindMarkers without prefiltering. Significant DEGs for

visualization were defined using a combined threshold of padj < 1e-5 and log2FC > 0.23.

<u>CUT&RUN experiments</u>

CUT&RUN was performed with the Epicypher CUTANA™ ChIC/CUT&RUN Kit (14-1048) according to the

manufacturer's specifications with the following conditions: 500,000 cells per sample were prepared first and

nuclei were extracted according to the Epicypher CUT&RUN Manual Appendix. Nuclei were incubated with

activated ConA beads. The following antibodies (0.5 ug of antibody per reaction) were used:

IgG Control antibody: CUTANA™ Kit Rabbit IgG CUT&RUN Negative Control Antibody, EpiCypher Rabbit anti-

H3K27Ac antibody, Millipore Mouse anti-Flag antibody, Antibodies-online Rabbit anti-CTNNB1 antibody.

Libraries were prepared using the Epicypher CUTANA™ CUT&RUN Library Prep Kit (14-1002) according to

the manufacturer's specifications. Library quality was analyzed using Agilent Tapestation D1000 High

Sensitivity Tapes (#5067-5584). Sequencing was performed with 5 million 150bp paired-end reads per sample

on an Illumina NovaSeq 6000.

<u>CUT&RUN data analysis</u>

CUT&RUN reads were analyzed for quality using FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and adaptors trimmed using Trimmomatic. Reads

were then aligned to mm9 by Bowtie2 version 2.3.1. SAMTools version 1.11 was then used to convert to BAM

format, index, isolate uniquely mapped paired reads, and remove duplicates. MACS2 version 2.2.8 was used

to call peaks with IgG as input control. Read counts across genomic intervals and peak visualization were

performed using deepTools version 3.5 and bigwig files were visualized using the Integrative Genomics

Viewer. Deseq2 was used to identify differentially bound peaks between conditions, DiffBind used to plot

heatmaps of accessible/bound regions and HOMER to identify *de novo* motifs most enriched in the

differentially bound regions and annotate associated genes.

<u>Western blot</u>

Cells were lysed in RIPA buffer (Thermo Fisher Scientific, Cat# 89900) supplemented with protease inhibitors

(Thermo Fisher Scientific, Cat# 87786) for 30 minutes at 4 °C. The lysates were clarified by centrifugation at

$10,000 \times g$ for 20 minutes at 4 °C, and protein concentrations were determined using Bradford Protein Assay

Kit (Bio-Rad, Cat# 5000006). Equal amounts of protein (10 µg) were denatured at 95 °C for 5 minutes and

loaded onto 4–12% precast polyacrylamide gels (Bio-Rad, Cat# 3450125) for electrophoresis. Proteins were

transferred to PVDF membranes (Thermo Fisher Scientific, Cat# IB24001) using an iBlot2 transfer device

(Invitrogen, Cat# IB21001). Membranes were blocked with 5% BSA (vol/vol) in 1× TBST (AmericanBio, Cat#

AB14330-01000) at room temperature, incubated with primary antibodies overnight at 4 °C, washed three

times with TBST for 5 minutes each, incubated with HRP-conjugated secondary antibodies for 1 hour at room

temperature, and washed again three times with TBST.

Membranes were developed using SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Fisher

Scientific, Cat# 34577) and imaged on a ChemiDoc imaging system (Bio-Rad). The following antibodies were

used: mouse anti-Flag (Millipore, Cat# F1804), rabbit anti-Actin (Cell Signaling Technology, Cat# 4970), HRP-

conjugated goat anti-rabbit IgG (Cell Signaling Technology, Cat# 7074), and HRP-conjugated goat anti-mouse

IgG (Bio-Rad, Cat# STAR207P, 1:10,000).

<u>Dermal cell culture</u>

E13.5 PDGFRαH2BGFP skin cells were sorted based on their GFP status. GFP+ cells were collected and

pelleted (300 × g, 10 min, 4°C), resuspended at 2×10^6 cells/mL in complete growth medium (DMEM + 10%

FBS, Penicillin-Streptomycin, and L-Glutamine), and viability was assessed by trypan blue exclusion. Cells

were plated at 3×10^6 cells/well in 6-well plates in 2 mL total volume. For chemical perturbation assays, cells

were mixed 1:1 with a 2× agonist solution. For example, 10 µM CHIR99021 (Sigma, Cat# SML1046) was

prepared in DMSO and diluted to a working concentration of 5 µM in culture. Similarly, SAG (Sigma, Cat#

SML1314) was prepared as a 50 µM stock and diluted as needed. Media with agonist was used fresh or stored

at 4°C for up to two days. Cells were cultured in 48-72 hours.

<u>qPCR</u>

Total RNA was extracted from cultured cells using the RNeasy Mini Kit (Qiagen, Cat# 74134) according to the

manufacturer's protocol. Complementary DNA (cDNA) was synthesized from 500 ng–1 µg of total RNA using

the SuperScript™ III First-Strand Synthesis System (Invitrogen, Cat# 18080051) with random hexamer

primers. Quantitative PCR was performed using the PowerUp™ SYBR™ Green Master Mix (Thermo Fisher

Scientific, Cat# 4364344) on a Viia7 Real-Time PCR System (Applied Biosystems) in 384-well optical plates.

Reactions were run in technical triplicates using gene-specific primers, and threshold cycle (Ct) values were

normalized to Gapdh as a housekeeping gene. Relative expression was calculated using the ΔΔCt method.

QUANTIFICATION AND STATISTICAL ANALYSIS

<u>Image analysis</u>

Raw image stacks acquired from whole-mount explants were imported into Fiji or Adobe Photoshop for

analysis. ggplot2 and cowplot R libraries were used for graphical representation of the scRNA-seq data.

Diffusion maps were generated as described above.

<u>Quantification</u>

For quantification of DC density, 3-dimensional whole-mount tiled mosaics of skin explants with a 1000 x 1000

µm field of view and a z-depth of 60-100 µm were used for n=3-4 embryos. For volumetric quantification of DC

cell number, EdU+ or Tom+ percentage of DC or UD, 3-dimensional whole mount mosaics stained with Sox2

and EdU or RFP were used to manually count positive cells, using ImageJ (Fiji) software using planar (XY) and

orthogonal (YZ, XZ) views. For volumetric quantification of cell cycle phases, 3-dimensional whole mount

mosaics stained with markers for G0/G1 cells (based on Fucci2 reporter) were done manually by counting

nuclear positive cells using ImageJ (Fiji) software. For quantification of upper dermal cells, a distance of 10 um

below the epidermis was considered upper dermis. For quantification of the peri-DC region by whole-mount,

cells two cell layers out from the Sox2+ DC were counted. For quantification based on FISH, cells with 4-5 dots

were considered positive (according to the RNAScope manufacturer's instructions) and sections from a total of

n=4 different embryos were examined. To measure RNA expression levels, H-scores were calculated

according to ACDBio manufacturer's instructions: a cell with 0 dot is scored 0, 1-3 dots score 1, 4-9 dots score

2, 10-15 dots and/or less than 10% clustered dots score 3, and more than 15 dots and/or more than 10%

clustered dots score 4; then the final H-score of a given cell type A is calculated by summing the (% cells

scored B within all cells in A)*B for score B in 0-4. For quantification of upper dermal cells, a distance of 10-20

μm below the epidermis was considered as upper dermis. For quantification of cells in the peri-DC region

Dkk1+Sox2- was considered.

Statistical analysis

All statistical values are expressed as mean ± SEM. An unpaired Student's *t*-test was used to analyze data

sets with two groups and *P <0.05 to, **P<0.01, ***P< 0.001 and ****P<0.00001 indicated a significant

difference. When comparing more than two groups, P values were determined by one-way ANOVA with

Tukey's HSD test performed as the post hoc analysis. Statistical calculations were performed using Prism

software package (GraphPad). For categorical data, a chi-square test was employed to estimate a P value.

References

1.   Soufi, A., and Dalton, S. (2016). Cycling through developmental decisions: how cell cycle dynamics

control pluripotency, differentiation and reprogramming. Development *143*, 4301-4311.

10.1242/dev.142075.

2.   Liu, L., Michowski, W., Kolodziejczyk, A., and Sicinski, P. (2019). The cell cycle in stem cell

proliferation, pluripotency and differentiation. Nat Cell Biol *21*, 1060-1067. 10.1038/s41556-019-0384-

4.

3.   Pauklin, S., and Vallier, L. (2013). The cell-cycle state of stem cells determines cell fate propensity. Cell

155, 135-147. 10.1016/j.cell.2013.08.031.

4.   Calder, A., Roth-Albin, I., Bhatia, S., Pilquil, C., Lee, J.H., Bhatia, M., Levadoux-Martin, M., McNicol,

J., Russell, J., Collins, T., and Draper, J.S. (2013). Lengthened G1 phase indicates differentiation status

in human embryonic stem cells. Stem Cells Dev *22*, 279-295. 10.1089/scd.2012.0168.

5.   Nguyen, L., Besson, A., Roberts, J.M., and Guillemot, F. (2006). Coupling cell cycle exit, neuronal

differentiation and migration in cortical neurogenesis. Cell Cycle *5*, 2314-2318. 10.4161/cc.5.20.3381.

6.   Theilgaard-Monch, K., Pundhir, S., Reckzeh, K., Su, J., Tapia, M., Furtwangler, B., Jendholm, J.,

Jakobsen, J.S., Hasemann, M.S., Knudsen, K.J., et al. (2022). Transcription factor-driven coordination

of cell cycle exit and lineage-specification in vivo during granulocytic differentiation : In memoriam

Professor Niels Borregaard. Nat Commun *13*, 3595. 10.1038/s41467-022-31332-1.

7.   Greco, V., Chen, T., Rendl, M., Schober, M., Pasolli, H.A., Stokes, N., Dela Cruz-Racelis, J., and Fuchs,

E. (2009). A two-step mechanism for stem cell activation during hair regeneration. Cell Stem Cell *4*,

155-169. 10.1016/j.stem.2008.12.009.

8.   Hsu, Y.C., Li, L., and Fuchs, E. (2014). Transit-amplifying cells orchestrate stem cell activity and tissue

regeneration. Cell *157*, 935-949. 10.1016/j.cell.2014.02.057.

9.   Yi, R. (2017). Concise Review: Mechanisms of Quiescent Hair Follicle Stem Cell Regulation. Stem

Cells *35*, 2323-2330. 10.1002/stem.2696.

10.   Sun, Q., Lee, W., Hu, H., Ogawa, T., De Leon, S., Katehis, I., Lim, C.H., Takeo, M., Cammer, M.,

Taketo, M.M., et al. (2023). Dedifferentiation maintains melanocyte stem cells in a dynamic niche.

Nature *616*, 774-782. 10.1038/s41586-023-05960-6.

11.   Cotsarelis, G., Sun, T.T., and Lavker, R.M. (1990). Label-retaining cells reside in the bulge area of

pilosebaceous unit: implications for follicular stem cells, hair cycle, and skin carcinogenesis. Cell *61*,

1329-1337.

12.   Morita, R., Sanzen, N., Sasaki, H., Hayashi, T., Umeda, M., Yoshimura, M., Yamamoto, T., Shibata, T.,

Abe, T., Kiyonari, H., et al. (2021). Tracing the origin of hair follicle stem cells. Nature *594*, 547-552.

10.1038/s41586-021-03638-5.

13.   Ouspenskaia, T., Matos, I., Mertz, A.F., Fiore, V.F., and Fuchs, E. (2016). WNT-SHH Antagonism

Specifies and Expands Stem Cells prior to Niche Formation. Cell *164*, 156-169.

10.1016/j.cell.2015.11.058.

14.   Rompolas, P., Deschene, E.R., Zito, G., Gonzalez, D.G., Saotome, I., Haberman, A.M., and Greco, V.

(2012). Live imaging of stem cell and progeny behaviour in physiological hair-follicle regeneration.

Nature *487*, 496-499. 10.1038/nature11218.

15.   Zhang, C., Wang, D., Wang, J., Wang, L., Qiu, W., Kume, T., Dowell, R., and Yi, R. (2021). Escape of

hair follicle stem cells causes stem cell exhaustion during aging. Nat Aging *1*, 889-903. 10.1038/s43587-

021-00103-w.

16.   Driskell, R.R., Lichtenberger, B.M., Hoste, E., Kretzschmar, K., Simons, B.D., Charalambous, M.,

Ferron, S.R., Herault, Y., Pavlovic, G., Ferguson-Smith, A.C., and Watt, F.M. (2013). Distinct fibroblast

lineages determine dermal architecture in skin development and repair. Nature *504*, 277-281.

10.1038/nature12783.

17.   Myung, P., Andl, T., and Atit, R. (2022). The origins of skin diversity: lessons from dermal fibroblasts.

Development *149*. 10.1242/dev.200298.

18.   Chuong, C.M., Widelitz, R.B., Ting-Berreth, S., and Jiang, T.X. (1996). Early events during avian skin

appendage regeneration: dependence on epithelial-mesenchymal interaction and order of molecular

reappearance. J Invest Dermatol *107*, 639-646.

19.   Hardy, M.H. (1992). The secret life of the hair follicle. Trends Genet *8*, 55-61.

20.   Millar, S.E. (2002). Molecular mechanisms regulating hair follicle development. J Invest Dermatol *118*,

216-225. 10.1046/j.0022-202x.2001.01670.x.

21.   Paus, R., Muller-Rover, S., Van Der Veen, C., Maurer, M., Eichmuller, S., Ling, G., Hofmann, U.,

Foitzik, K., Mecklenburg, L., and Handjiski, B. (1999). A comprehensive guide for the recognition and

classification of distinct stages of hair follicle morphogenesis. J Invest Dermatol *113*, 523-532.

10.1046/j.1523-1747.1999.00740.x.

22.   Saxena, N., Mok, K.W., and Rendl, M. (2019). An updated classification of hair follicle morphogenesis.

Exp Dermatol *28*, 332-344. 10.1111/exd.13913.

23.   Chase, H.B., Rauch, R., and Smith, V.W. (1951). Critical stages of hair development and pigmentation

in the mouse. Physiol Zool *24*, 1-8. 10.1086/physzool.24.1.30152098.

24.   Qu, R., Gupta, K., Dong, D., Jiang, Y., Landa, B., Saez, C., Strickland, G., Levinsohn, J., Weng, P.L.,

Taketo, M.M., et al. (2022). Decomposing a deterministic path to mesenchymal niche formation by two

intersecting morphogen gradients. Dev Cell *57*, 1053-1067 e1055. 10.1016/j.devcel.2022.03.011.

25.   Gupta, K., Levinsohn, J., Linderman, G., Chen, D., Sun, T.Y., Dong, D., Taketo, M.M., Bosenberg, M.,

Kluger, Y., Choate, K., and Myung, P. (2019). Single-Cell Analysis Reveals a Hair Follicle Dermal

Niche Molecular Differentiation Trajectory that Begins Prior to Morphogenesis. Dev Cell *48*, 17-31 e16.

10.1016/j.devcel.2018.11.032.

26.   Alvarez-Medina, R., Cayuso, J., Okubo, T., Takada, S., and Marti, E. (2008). Wnt canonical pathway

restricts graded Shh/Gli patterning activity through the regulation of Gli3 expression. Development *135*,

237-247. 10.1242/dev.012054.

27.   Xu, Z., Wang, W., Jiang, K., Yu, Z., Huang, H., Wang, F., Zhou, B., and Chen, T. (2015). Embryonic

attenuated Wnt/beta-catenin signaling defines niche location and long-term stem cell fate in hair follicle.

Elife *4*, e10567. 10.7554/eLife.10567.

28.   Qu, R., Cheng, X., Sefik, E., Stanley Iii, J.S., Landa, B., Strino, F., Platt, S., Garritano, J., Odell, I.D.,

Coifman, R., et al. (2024). Gene trajectory inference for single-cell data by optimal transport metrics.

Nat Biotechnol. 10.1038/s41587-024-02186-3.

29.   Biggs, L.C., Makela, O.J., Myllymaki, S.M., Das Roy, R., Narhi, K., Pispa, J., Mustonen, T., and

Mikkola, M.L. (2018). Hair follicle dermal condensation forms via Fgf20 primed cell cycle exit, cell

motility, and aggregation. Elife *7*. 10.7554/eLife.36468.

30.   Chen, D., Jarrell, A., Guo, C., Lang, R., and Atit, R. (2012). Dermal beta-catenin activity in response to

epidermal Wnt ligands is required for fibroblast proliferation and hair follicle initiation. Development

139, 1522-1533. 10.1242/dev.076463.

960 31. Zhang, Y., Andl, T., Yang, S.H., Teta, M., Liu, F., Seykora, J.T., Tobias, J.W., Piccolo, S., Schmidt-

961    Ullrich, R., Nagy, A., et al. (2008). Activation of beta-catenin signaling programs embryonic epidermis

962    to hair follicle fate. Development *135*, 2161-2172. 10.1242/dev.017459.

963 32. Fu, J., and Hsu, W. (2013). Epidermal Wnt controls hair follicle induction by orchestrating dynamic

964    signaling crosstalk between the epidermis and dermis. J Invest Dermatol *133*, 890-898.

965    10.1038/jid.2012.407.

966 33. Vokes, S.A., Ji, H., Wong, W.H., and McMahon, A.P. (2008). A genome-scale analysis of the cis-

967    regulatory circuitry underlying sonic hedgehog-mediated patterning of the mammalian limb. Genes Dev

968    *22*, 2651-2663. 10.1101/gad.1693008.

969 34. te Welscher, P., Zuniga, A., Kuijper, S., Drenth, T., Goedemans, H.J., Meijlink, F., and Zeller, R.

970    (2002). Progression of vertebrate limb development through SHH-mediated counteraction of GLI3.

971    Science *298*, 827-830. 10.1126/science.1075620.

972 35. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H.,

973    and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-

974    regulatory elements required for macrophage and B cell identities. Mol Cell *38*, 576-589.

975    10.1016/j.molcel.2010.05.004.

976 36. Lex, R.K., Zhou, W., Ji, Z., Falkenstein, K.N., Schuler, K.E., Windsor, K.E., Kim, J.D., Ji, H., and

977    Vokes, S.A. (2022). GLI transcriptional repression is inert prior to Hedgehog pathway activation. Nat

978    Commun *13*, 808. 10.1038/s41467-022-28485-4.

979 37. Park, M., Jang, S., Chung, J.H., Kwon, O., and Jo, S.J. (2021). Inhibition of class I HDACs preserves

980    hair follicle inductivity in postnatal dermal cells. Sci Rep *11*, 24056. 10.1038/s41598-021-03508-0.

981 38. Kishimoto, J., Burgeson, R.E., and Morgan, B.A. (2000). Wnt signaling maintains the hair-inducing

982    activity of the dermal papilla. Genes Dev *14*, 1181-1185.

983 39. Choi, Y.S., Zhang, Y., Xu, M., Yang, Y., Ito, M., Peng, T., Cui, Z., Nagy, A., Hadjantonakis, A.K.,

984    Lang, R.A., et al. (2013). Distinct functions for Wnt/beta-catenin in hair follicle stem cell proliferation

and survival and interfollicular epidermal homeostasis. Cell Stem Cell *13*, 720-733.

10.1016/j.stem.2013.10.003.

40. Huggins, I.J., Bos, T., Gaylord, O., Jessen, C., Lonquich, B., Puranen, A., Richter, J., Rossdam, C.,

Brafman, D., Gaasterland, T., and Willert, K. (2017). The WNT target SP5 negatively regulates WNT

transcriptional programs in human pluripotent stem cells. Nat Commun *8*, 1034. 10.1038/s41467-017-

01203-1.

41. Myung, P.S., Takeo, M., Ito, M., and Atit, R.P. (2013). Epithelial Wnt ligand secretion is required for

adult hair follicle growth and regeneration. J Invest Dermatol *133*, 31-41. 10.1038/jid.2012.230.

42. van Amerongen, R., Bowman, A.N., and Nusse, R. (2012). Developmental stage and time dictate the

fate of Wnt/beta-catenin-responsive stem cells in the mammary gland. Cell Stem Cell *11*, 387-400.

10.1016/j.stem.2012.05.023.

43. Jeong, J., Mao, J., Tenzen, T., Kottmann, A.H., and McMahon, A.P. (2004). Hedgehog signaling in the

neural crest cells regulates the patterning and growth of facial primordia. Genes Dev *18*, 937-951.

10.1101/gad.1190304.

44. Harada, N., Tamai, Y., Ishikawa, T., Sauer, B., Takaku, K., Oshima, M., and Taketo, M.M. (1999).

Intestinal polyposis in mice with a dominant stable mutation of the beta-catenin gene. EMBO J *18*,

5931-5942. 10.1093/emboj/18.21.5931.

45. Blaess, S., Stephen, D., and Joyner, A.L. (2008). Gli3 coordinates three-dimensional patterning and

growth of the tectum and cerebellum by integrating Shh and Fgf8 signaling. Development *135*, 2093-

2103. 10.1242/dev.015990.

46. Dassule, H.R., Lewis, P., Bei, M., Maas, R., and McMahon, A.P. (2000). Sonic hedgehog regulates

growth and morphogenesis of the tooth. Development *127*, 4775-4785.

47. Abe, T., Sakaue-Sawano, A., Kiyonari, H., Shioi, G., Inoue, K., Horiuchi, T., Nakao, K., Miyawaki, A.,

Aizawa, S., and Fujimori, T. (2013). Visualization of cell cycle in mouse embryos with Fucci2 reporter

directed by Rosa26 promoter. Development *140*, 237-246. 10.1242/dev.084111.

48.   Nestorowa, S., Hamey, F.K., Pijuan Sala, B., Diamanti, E., Shepherd, M., Laurenti, E., Wilson, N.K.,

Kent, D.G., and Gottgens, B. (2016). A single-cell resolution map of mouse hematopoietic stem and

progenitor cell differentiation. Blood *128*, e20-31. 10.1182/blood-2016-05-716480.

Figure 1

Figure 2

Figure 3

Figure 4

A

CC Phase Control *Lef1* *Ptch1* **E13.5 ActSMO Dermis** CC Phase ActSMO *Lef1* *Ptch1*

- G1/G0
- S
- G2/M

B E13.5 ActSMO Gene Trajectories

CC DC LD New GT

Exit GT projected on ActSMO and Control

Stage 1 Stage 2 Stage 3 Stage 4 Stage 5

ActSMO

Control

C Reactome Pathway Analysis

Transcriptional Regulation by TP53
TP53 Regulates Metabolic Genes
TNFR2 non-canonical NF-kB
Programmed cell death
Keratinization

p.adjust

E13.5 ActSMO *Trp53inp1* *Cdkn1a*

ActSMO

Control

D

E13.5 Control E13.5 ActSMO E14.5 Wildtype

Sox2 *Cdkn1a* *Cdkn1a* *Sox2* *Cdkn1a* *Sox2* *Cdkn1a* *Sox2*

Sox2 *Trp53inp1* *Trp53inp1* *Sox2* *Trp53inp1* *Sox2* *Trp53inp1* *Sox2*

E13.5 Gene Expression *Trp53inp1* *Cdkn1a* ****

Lef1 level *Lef1* H-Score UD *Trp53*+ DC E13.5 E14.5 **** ns

E Label transfer ActSMO Exit GT

ActSMO Actβcat Wntless cKO Control

F

EdU 12.5 13.5 14.5 EdU 12.5 13.5 14.5 EdU 12.5 13.5 14.5

ActSMO *Trp53inp1* EdU E13.5 *Trp53inp1* EdU E14.5 *Trp53inp1* EdU E14.5 *Trp53inp1* EdU

EdU pulse (1 hr) vs EdU chase (24 hr)

%EdU EdU 1 hr EdU 24 hr **** ns **** ****

E13.5 *Trp53*+ E14.5 *Trp53*+ E14.5 DC

E13.5 EdU pulse → EdU pulse-chase → E14.5 EdU chase E14.5 EdU pulse

○ *Trp53inp* - ○ EdU -
● *Trp53inp* + ● EdU + (S phase) ● EdU + (retained by daughter cells)

G

G2/M cell → G0 cell (*Trp53inp1^{HI}*) → G0 cell (*Trp53inp1^{LO}*)

Figure 5

Figure 6

A E13.5 ActSMO Gene Trajectories

CC
DC
LD
Exit GT

B E13.5 ActSMO DC GeneTrajectory Gene Order

Runx3 Lef1   Bmp4   Gal   Sox2

Stage 1   Stage 2   Stage 3   Stage 4   Stage 5

ActSMO

Control

C DC gene expression by GT stage

Gal

Sox18

Sox2

Gal

Sox18

Sox2

D *Lef1* level by stage

Control   ActSMO

Lef1 level

mean
1.6
1.2
0.8
0.4
0.0

1 2 3 4 5

ActSMO DC stage

E

E14.5 Wildtype

Synchronized CC exit and DC genes

ActSMO

Uncoupled CC exit and DC genes

DC Genes   ◯ Quiescent Cell

E14.5 Wildtype

Placode

DC

Proliferation   Quiescence

SHH activity

SHH-induced Wnt acceleration

Wnt activity

DC Genes

CC Exit (GLI3 loss)   High DC Genes

CC Exit   DC Genes

Figure 7