Sharp cell-type boundaries emerge from temporal coordination between morphogen

signals

Authors:

Ruiqi Li[1,2,13], Yiqun Jiang[3,4,13], Sarah Platt[3,13], Tianchi Xin[5], Ryan Driskell[6], Kevin A. Peterson[7],

Sarah Van[3], Hainan Lam[3], Shagun Lukkad[3], Eva-LaRue Barber[3], Chae Ho Lim[8], M. Mark

Taketo[9], Yuval Kluger[1,2,10,11], Peggy Myung[2,3,4,12*]

*Corresponding author

Affiliations:

[1]Computational Biology & Biomedical Informatics Program, Yale University, New Haven, CT

06520, USA

[2]Department of Pathology, Yale University, New Haven, CT 06520, USA

[3]Department of Dermatology, Yale University, New Haven, CT 06520, USA

[4]Molecular, Cellular and Developmental Biology, Yale University, New Haven, CT 06520, USA

[5]Department of Genetics, Yale University, New Haven, CT 06520, USA

[6] School of Molecular Biosciences, Washington State University, Pullman, WA, USA

[7]The Jackson Laboratory, Bar Harbor, ME 04609, USA

[8]Ronald O. Perelman Department of Dermatology, and Department of Cell Biology, NYU

Grossman School of Medicine, New York, NY, USA

[9]Colon Cancer Project, Graduate School of Medicine, Kyoto University, Sakyo, Kyoto 606-8501,

Japan

[10]Applied Mathematics Program, Yale University, New Haven, CT 06511, USA

[11]Yale Cancer Center, New Haven, CT 06520, USA

[12]Yale Stem Cell Center, New Haven, CT 06520, USA

[13]These authors contributed equally

Keywords: Morphogen, patterning, development, scRNA-sequencing, Wnt, Hedgehog, skin,

cell cycle, differentiation

Abstract

Classic models of the French flag problem depict sharp cell-type boundaries emerging from

threshold responses to morphogen gradients. Yet, how such boundaries arise during dynamic

cell-state transitions remains unclear. We use hair follicle dermal condensates to study a sharp

cell-type transition in which proliferative progenitors synchronously undergo cell-cycle exit and

molecular differentiation. Using genetic and genomic approaches, we show that Wnt and

Hedgehog signaling interact to coordinate the timing of these processes. When their activities

are temporally aligned, intermediate transitional states are compressed through cell-cycle exit,

producing a sharp cell-type boundary; when misaligned, transitional states expand, yielding

fuzzy borders. Mechanistically, elevated Wnt activity promotes cell-cycle exit by regulating

chromatin binding of the Hedgehog mediator GLI3. Hedgehog signaling induces differentiation

genes in a Wnt-dependent manner and simultaneously elevates Wnt activity, aligning arrest and

differentiation in time. These findings reveal a mechanism wherein morphogen interactions

coordinate transitions to generate precise cell-type boundaries.

Introduction

Developing tissues must position distinct cell types across space and time to assemble

functional organs. How such patterns form was framed by Lewis Wolpert as the "French flag"

problem, in which discrete cell types emerge at defined positions along a morphogen gradient

(Fig. 1a)[1]. In this framework, graded morphogen signals are interpreted through concentration

thresholds that delineate spatial domains of cell fate[2-6]. Yet, a fundamental question remains

unresolved: how sharp cell-type boundaries emerge *in vivo* when the underlying signals and cell

states vary continuously across a tissue. In developing systems, boundaries do not arise from

instantaneous fate switches. Instead, cells progress through dynamic transitions and pass

through intermediate states before committing to a new identity. Moreover, in many mammalian

tissues, these fate transitions unfold while cells continue to proliferate, raising the question of

how ongoing cell-cycle dynamics are coordinated with graded morphogen inputs. Thus,

boundary sharpness may not only reflect where fate thresholds are crossed, but how rapidly

cells traverse the same underlying developmental progression.

Addressing the temporal dimension of the French flag problem has been challenging, as most

approaches to studying cell fate transitions emphasize cell origins or endpoints, rather than

intermediate regulatory programs, or can visualize intermediate cell behaviors without molecular

resolution[7-11]. We address this problem in a mammalian system by examining hair follicle dermal

condensate formation, a uniquely tractable developmental transition in which the principal

morphogen signals are known and cell-state progression occurs over a narrow temporal

window[12-14].

Dermal condensates are spheroid clusters of quiescent dermal cells that form beneath epithelial

hair follicle placodes and are essential for hair follicle development[15-21]. At E13.5, prior to DC

formation, Wnt signaling is broadly activated across the upper dermis, forming a gradient that is

highest in cells closest to the epidermis[16, 22, 23]. Wnt activity is required but not sufficient for DC

formation. Upon placode formation, epithelial cells secrete Sonic Hedgehog (SHH) and

additional Wnt ligands, resulting in localized co-activation of Wnt and SHH in underlying dermal

cells[21, 24-26]. Together, these signals are necessary and sufficient for DC formation, yet how Wnt

and SHH signaling interact to drive the pre-DC-to-DC transition remains unknown, and DCs

appear to emerge abruptly with few detectable intermediates.[8, 27, 28].

Previous work has examined how tissue growth and proliferation affect morphogen gradients by

shaping field size, ligand distribution, or the scaling of positional information[29-33]. In these

frameworks, proliferation is typically considered in terms of how it modifies the signaling

environment cells interpret. Many mammalian tissues, however, undergo fate specification while

cells continue to divide, introducing an additional constraint, as proliferative expansion of

intermediate states becomes a key factor influencing the precision of pattern formation. In this

study, we focus on how interacting morphogen signals coordinate cell-cycle exit with

differentiation to ensure spatially precise patterning in such proliferative contexts.

Here, we ask how interacting morphogen signals regulate the progression of a rapid cell-state

transition to generate a sharp tissue boundary. By combining *in vivo* genetic perturbations,

chromatin profiling, and a single-cell computational framework that disentangles concurrent

biological programs, we are able to capture intermediate transcriptional states that are normally

difficult to detect and to determine how their timing and spatial extent are regulated by these two

pathways *in vivo*[34]. We show that Wnt and SHH differentially control cell-cycle exit and

molecular differentiation, and that the temporal coordination of these processes determines

whether transitional states are compressed or expanded, thereby controlling boundary

sharpness. Mechanistically, we identify cross-regulation of transcription factor occupancy as a

key mechanism linking signaling dynamics to cell-cycle exit.

Results

Cell-cycle exit and molecular differentiation are temporally coordinated during dermal

condensate formation

Prevailing models of the French flag problem posit that distinct cell states are specified by

threshold levels of morphogen concentration that vary with distance from a polarized source,

predicting that sharp cell-type boundaries arise from spatial differences in morphogen signaling

(Fig. 1a). To examine how this model applies to hair follicle dermal condensate (DC) formation,

we asked how dermal cell types are spatially organized before and during DC formation. Prior to

DC formation, dermal cells are proliferative and lack DC-specific markers (Extended Data Fig.

1a). Upon DC formation, DC cells express characteristic differentiation markers, including *Sox2*,

Sox18, and *Gal*, and undergo cell-cycle exit, as indicated by minimal EdU nucleotide

incorporation, while neighboring upper dermal cells that do not express DC genes remain highly

proliferative. (Fig. 1b,c and Extended Data Fig. 1b,c).

Using *in vivo* labeling, we previously showed that DC cells arise from highly proliferative

progenitors located in the upper dermis at E13.5 (Extended Data Fig. 1d)[16, 27]. As DC formation

initiates, proliferating cells that give rise to DCs become localized to a region surrounding the

nascent condensate (peri-DC region) where they express *Dkk1* (Fig. 1c and Extended Data

Fig.1c). During the pre-DC–to–DC transition, *Dkk1*+ peri-DC cells exit the cell cycle while

upregulating DC differentiation genes[27]. Consistent with this, DC marker expression overlapped

with the *in vivo* G1/G0 Fucci2 reporter and with expression of the cell-cycle inhibitor *Cdkn1a*

(*p21*), with only rare cells expressed *Cdkn1a* in the absence of *Sox2* (Fig. 1c and Extended

Data Fig. 1e)[16, 35]. These data indicate that cell-cycle exit and DC differentiation occur

concurrently during DC formation.

Given this temporal coincidence, we next asked how these two processes relate to Wnt and

Sonic Hedgehog (SHH) signaling during DC formation. At E13.5, prior to SHH expression, Wnt

signaling is active in the upper dermis (Extended Data Fig. 1a,b) but is not sufficient to induce

DC gene expression[16, 22, 36]. Using *Lef1* as a canonical readout of Wnt activity, which is

concordant with expression of multiple Wnt target genes (*Axin2*, *Wif1*, *Tcf7*) (Extended Data Fig.

1f)[16, 27], we examined the relationship between Wnt signaling and proliferation. At E13.5, *Lef1*

expression was highest in the most superficial layer of upper dermis and decreased across

deeper dermal cell layers, yet this spatial pattern did not correlate with EdU incorporation, nor

with DC gene expression (Extended Data Fig. 1a).

By E14.5, SHH expression in placode cells resulted in SHH activation in the underlying dermis.

At this stage, dermal SHH activity (assessed by *Ptch1*) covaried with Wnt signaling (*Lef1*)

across dermal populations, with significantly higher *Lef1* levels detected compared to E13.5

(Fig. 1d). The highest levels of both signals were observed in DC cells, intermediate levels in

peri-DC (*Dkk1*+) progenitors, and progressively lower levels in surrounding upper dermal cells.

In contrast to E13.5, elevated Wnt activity at E14.5 correlated with reduced proliferation, with

the lowest EdU incorporation observed in DC cells (Fig. 1d,e). Thus, quiescence and molecular

differentiation coincide spatially and temporally within a narrow domain characterized by

elevated Wnt and SHH signaling.

High Wnt activity induces cell-cycle exit independently of SHH signaling

To dissect the respective contributions of Wnt and SHH signaling to the pre-DC–to-DC

transition, we perturbed each pathway prior to DC formation. Because both pathways are

required for DC differentiation, standard loss-of-function approaches preclude analysis of their

individual roles at this stage. Given that DC cells ultimately exhibit the highest levels of both Wnt

and SHH activity, we asked whether these signals contribute to distinct components of the

transition or act through a shared mechanism. We therefore examined the effects of elevating

Wnt signaling at E13.5, prior to DC formation by expressing a constitutively active form of β-

catenin (*Axin2CreER;βcat$^{fl/EX3}$*, hereafter Actβcat; Fig. 2a, tamoxifen at E11.5).

To unbiasedly assess how increased Wnt activity reshapes dermal transcriptional states before

DC differentiation, we performed single-cell RNA sequencing on E13.5 Actβcat and control

dermis. Dermal cells were filtered for analysis using established markers (e.g., *Col1a1*,

PDGFRα)[16, 27]. Consistent with prior work, Actβcat mutant dermal cells showed increased

expression of Wnt target genes, including *Lef1*, *Axin2* and *Tcf7*, but not DC markers (e.g., *Sox2*)

or SHH target genes (e.g., *Gli1*) (Fig. 2b). Notably, differential gene expression analysis

revealed that *Cdkn1a*, a cell-cycle inhibitor normally expressed in DC cells at E14.5, was among

the most significantly upregulated genes in E13.5 Actβcat dermal cells (Fig. 2b and

Supplementary Table 1). Quantitative FISH confirmed elevated *Lef1* expression in E13.5

Actβcat dermal cells relative to controls and showed that Wnt target gene levels did not exceed

those observed in wildtype DC cells, indicating that activated β-catenin elevates Wnt signaling

but not above physiological levels (Fig. 2c and Extended Data Fig. 2a). FISH analysis further

showed that constitutive Wnt activation was sufficient to induce expression of *Cdkn1a* (*p21*) at

E13.5 in the absence of SHH activation, and *Cdkn1a*+ cells exhibited reduced EdU

incorporation (Fig. 2c).

By E14.5, DCs were present in both control and Actβcat embryos but were enlarged in mutants

(Extended Data Fig. 2b). In control embryos, *Cdkn1a* expression was largely restricted to *Sox2+*

DC cells, whereas in Actβcat mutants, it extended into surrounding *Sox2-* upper dermal cells

(Fig. 2d). Both *Sox2+* and *Sox2−* cells expressing high *Cdkn1a* exhibited low EdU incorporation

(Fig. 2d). *Sox2+* cells in Actβcat mutants were closest to placodes and expressed the highest

levels of *Ptch1*, whereas surrounding mutant *Cdkn1a+Sox2-* cells expressed lower *Ptch1* levels,

indicating that DC gene expression requires SHH signaling above a defined threshold (Fig. 2e).

Thus, elevated Wnt signaling is sufficient to induce cell-cycle exit in the absence of SHH,

whereas molecular DC differentiation requires additional SHH-dependent input (Fig. 2f).

SHH signaling cell-autonomously elevates Wnt activity coincident with cell-cycle exit

Our results show that elevated Wnt activity induces cell-cycle exit without DC differentiation. We

next examined how SHH signaling influences this transition. Specifically, we asked whether

elevated SHH activity could initiate DC gene expression independently of cell-cycle exit. To

address this, we expressed a constitutively activated form of Smoothened

(*Axin2CreER;RosaLSL-Smom2YFP*, hereafter ActSMO) in the dermis prior to DC formation

when Wnt signaling levels are low (tamoxifen E11.5; Fig. 3a). Whole-mount analysis of E14.5

skin showed that high SHH activation in Wnt-active dermal cells induced large dermal clusters

expressing DC genes in the upper dermis (Fig. 3b). These clusters formed autonomously

without placodes, indicating that DC gene expression can occur in the absence of placode-

derived Wnt ligands[27]. Notably, many cells in the outer layers of Sox2+ dermal clusters

remained proliferative, as indicated by %Sox2+/EdU+ double labeling, demonstrating that DC

gene expression can occur independently of cell-cycle arrest. By contrast, Sox2+ cells in the

center of clusters lacked EdU incorporation similar to quiescent DC cells in wildtype embryos.

Given the observed quiescent central Sox2+ dermal cells (Fig. 3b), we asked whether there

were similar molecular changes to those seen in the E13.5 Actβcat mutant. We obtained

scRNA-seq data from E13.5 ActSMO and paired control skin (Fig. 3c). As expected, E13.5

ActSMO dermal cells exhibited uniformly high *Ptch1* expression. *Lef1* expression remained

largely enriched in upper dermal (UD) cells but was significantly elevated compared to E13.5

controls (Fig. 3c,d). FISH analysis confirmed increased *Lef1* expression in E13.5 ActSMO

dermis despite the absence of placodes (Fig. 3d). Importantly, only *Ptch1+* cells (recombined)

showed elevated *Lef1* expression, while neighboring *Ptch1*-negative cells (non-recombined)

retained normal *Lef1* levels comparable to E13.5 control UD cells, indicating a cell-autonomous

effect. ActSMO cells with high *Lef1* expression also exhibited elevated *Cdkn1a* expression and

low rates of EdU incorporation (Fig. 3d,e). Together, these findings show that SHH activation

elevates Wnt signaling cell-autonomously and is associated with induction of cell-cycle exit,

independent of placode Wnt ligands (Fig. 3f).

GeneTrajectory resolves transcriptional programs associated with cell-cycle exit

In the ActSMO condition, DC genes are expressed in cells that remain proliferative (Fig. 3b),

indicating that cell-cycle exit and DC gene expression can be temporally uncoupled. To resolve

the gene programs underlying these processes, we applied GeneTrajectory (GT), a

computational approach that separates co-occurring transcriptional programs from whole

transcriptome scRNA-seq data[34]. Unlike conventional pseudotime methods that assign cells to a

single lineage trajectory, GT directly constructs gene trajectories, allowing individual cells to be

analyzed across multiple overlapping biological processes.

We first applied GT to E14.5 wildtype dermal scRNA-seq data, identifying three gene

trajectories, corresponding to proliferation (S/G2/M), DC differentiation and lower dermal

differentiation (Fig. 4a). The DC trajectory was partitioned into stages using gene bins, and cells

were visualized by gene bin scores (Fig. 4a and Extended Data Fig. 3a)[34]. *Lef1* and *Ptch1*

expression covaried across DC trajectory stages, with maximal expression at the terminal stage

6 (Fig. 4b and Extended Data Fig. 3b). Canonical DC genes, including *Gal*, *Sox18* and *Sox2,*

were similarly enriched at stage 6 (Fig. 4c). Cell-cycle analysis revealed that stage 6 cells were

quiescent with a G1 fraction near 1.0, and elevated *Cdkn1a* expression (Fig. 4c). Thus, in the

wildtype condition, cell-cycle exit and molecular DC differentiation coincide at the terminus of

the DC trajectory.

We next applied GT to E13.5 ActSMO scRNA-seq data in which cell-cycle exit and DC

differentiation are uncoupled. GT identified four trajectories (Fig. 4d and Extended Data Fig. 3c).

Two were similar to the control, corresponding to proliferation (S/G2/M) and LD differentiation.

Two additional trajectories were unique to the ActSMO dermis: one containing DC genes and a

second lacking DC markers. Cells expressing this latter program localized to a subset of Wnt-

active cells and were largely restricted to G1/G0 phase (Fig. 4e and 3c). Pathway analysis

revealed enrichment for TP53-associated cell-cycle inhibitory genes, including *Trp53inp1* and

Plk2, overlapping with *Cdkn1a* expression (Fig. 4f and Supplementary Table 1). FISH analysis

confirmed significant upregulation of both *Cdkn1a* and *Trp53inp1* in the E13.5 ActSMO dermis

relative to controls (Fig. 4g and 3d and Extended Data Fig. 3d)

In ActSMO mutants, SHH-active cells expressing *Trp53inp1* exhibited significantly higher *Lef1*

levels than control upper dermal cells (Fig. 4g), consistent with earlier results showing that SHH

elevates Wnt activity and promotes quiescence (Figs. 3d and 2c). To test whether elevatd Wnt

signaling alone induces this TP53-associated program, we applied GT to the E13.5 Actβcat

dermis, which induces *Cdkn1a* but not DC genes (Extended Data Fig. 3e). GT analysis

identified a similar *Trp53inp1*-containing program in E13.5 Actβcat mutants (Supplementary

Table 1). We applied a module projection (label transfer) method to assess whether the gene

program was shared across different perturbations. Specifically, we transferred the E13.5

ActSMO-specific trajectory onto E13.5 Actβcat and control datasets, which revealed a shared

molecular state in both mutants that was absent in controls (Fig. 4h).

To determine whether this program reflects a general consequence of reduced proliferation, we

analyzed E13.5 Wntless conditional knockout (*K14Cre;Wls*$^{fl/fl}$) embryos in which epidermal Wnt

ligand secretion is disrupted, resulting in loss of dermal Wnt signaling and reduced dermal

proliferation (Extended Data Fig. 3f)[22, 34, 37]. Label transfer of the ActSMO/Actβcat-specific

trajectory was not detected in Wntless cKO dermis (Fig. 4h and Extended Data Fig. 3f),

indicating that this TP53-associated program is Wnt-dependent and specific to DC-associated

cell-cycle exit.

Finally, we examined the cell-cycle dynamics of cells expressing this gene program. At E13.5

and E14.5, most ActSMO *Trp53inp1+* cells showed low EdU incorporation comparable to

quiescent DC cells (Fig. 4i and Extended Data Fig. 4a). EdU pulse-chase experiments

demonstrated that a substantial fraction of *Trp53inp1+* cells retained EdU nucleotide similarly to

wildtype DCs (Fig. 4i and Extended Data Fig.1d)[16, 27]. Although ActSMO dermal clusters

contained proliferating Sox2+ cells at E14.5, most became quiescent by E15.5 (Extended Data

Fig. 4b), indicating that these proliferating cells exit the cell cycle after division. Together, these

data define a shared transcriptional program underlying cell-cycle exit during the pre-DC-to-DC

transition, which we term the *"Exit" gene trajectory*.

Loss of GLI3 chromatin binding induces cell-cycle exit downstream of elevated Wnt

signaling

Although Wnt activity is required for dermal proliferation, our results indicate that elevated Wnt

activity promotes cell-cycle exit during DC formation[16, 22]. To examine the underlying

mechanism, we profiled genes differentially bound by CTNNB1 (β–catenin) in E13.5 Actβcat

and control dermis. CTNNB1 binding was enriched at canonical Wnt target genes (e.g., *Sp5*,

Axin2, *Lef1*), but not at *Cdkn1a,* Exit GT genes, or other loci associated with cell cycle inhibition

(Extended Data Fig. 5a,b)[22]. These findings raised the question of how elevated Wnt activity

promotes cell-cycle exit despite the absence of direct CTNNB1 binding at cell-cycle inhibitory

genes. Given that Wnt activity is elevated upon SHH activation, it was plausible that

components of the SHH pathway mediate Wnt-dependent cell-cycle exit. In other tissues, Wnt

signaling regulates *Gli3* expression independently of SHH[38]. In the absence of SHH ligand, Gli3

is predominantly present in its cleaved repressor form, which binds a broad range of SHH target

genes[39, 40]. FISH analysis showed that *Gli3* was expressed in Wnt-active dermal cells in both

control and Actβcat embryos at E13.5 prior to SHH expression (Fig. 5a). To profile GLI3 binding,

we performed CUT&RUN in E13.5 Actβcat and control dermis using *Gli3Flag* knockin embryos.

Interestingly, GLI3 occupancy was globally reduced in the Actβcat condition as indicated by

decreased GLI3-bound reads at transcriptional start sites (Fig. 5b). De novo motif analysis

confirmed enrichment of GLI motifs at regions showing reduced binding (Extended Data Fig.

5c)[41]. These regions were enriched for genes associated with SHH signaling, despite the fact

that most canonical SHH target genes (e.g., *Ptch1*, *Gli1*) were not expressed and lacked active

histone marks (H3K4me3 or H3K27ac) at E13.5 (Fig. 5c). These observations are consistent

with a recent report indicating that GLI3 binding prior to SHH does not actively repress

transcription (i.e., is inert) and that many targets become activated only after SHH expression[42].

Western Blot analysis confirmed that cleaved Gli3 protein levels were retained in the Actβcat

mutant, showing that reduced chromatin occupancy was not due to loss of Gli3 protein and that

Gli3 remained predominantly in its cleaved repressor form (Fig. 5d). These results show that

elevated Wnt signaling alone reduces GLI3 chromatin occupancy at many genomic regions

independently of SHH.

To test whether loss of GLI3 binding is sufficient to induce cell-cycle exit, we ablated *Gli3* in the

dermis (*Axin2CreER;Gli3[fl/fl]* or Gli3 cKO; tamoxifen E11.5). At E14.5, DCs in Gli3 cKO embryos

were smaller than in controls (Fig. 5e), and *Dkk1+* peri-DC progenitors showed premature

upregulation of *Cdkn1a* and reduced EdU incorporation (Fig. 5e and Extended Data Fig. 5d).

Similarly, at E13.5 prior to DC formation, Gli3 cKO dermal cells exhibited elevated *Cdkn1a/p21*

expression and reduced proliferation, phenocopying the Actβcat condition (Fig. 5f and Extended

Data Fig. 5e).

FISH analysis confirmed reduced *Gli3* expression in the dermis of E13.5 Gli3 cKO embryos,

while Wnt target genes were unaffected, suggesting that Gli3 acts downstream of elevated Wnt

activity (Fig. 5f and Extended Data Fig. 5f). Canonical SHH target genes (e.g. *Ptch1*) were also

not induced, confirming that loss of Gli3 does not broadly derepress SHH targets and that the

lack of SHH target expression in the Actβcat condition is unlikely due to residual GLI3

repression. To assess whether GLI3 binding is associated with chromatin organization, we

performed chromatin capture (Micro-C) in E13.5 wildtype dermis. GLI3-bound loci (e.g., *Ptch1*)

were actually modestly enriched in A (open) rather than B (closed) compartments but rarely

participated in chromatin loops (<1%) and lacked active promoter marks (Extended Data Fig.

5g). By contrast, canonical Wnt target genes (e.g., *Lef1* and *Sp5)* formed chromatin loop

interactions and showed active histone modifications (Extended Data Fig. 5g). Moreover,

snATAC+RNA-seq analysis of E14.5 embryos that lack epidermal *Shh* showed significant GLI

motif accessibility localized to Wnt-active cells even without Gli activator function (Fig. 5g).

Together, these data indicate that GLI3 does not actively repress many bound regions and may

instead prime GLI sites for accessibility prior to SHH.

As expected, scRNA-seq analysis of E13.5 Gli3 cKO dermis showed reduced *Gli3* expression.

Neither DC nor canonical SHH target genes were detected, and Wnt target genes were retained

(Extended Data Fig. 6a,b). GT analysis identified four trajectories in Gli3 cKO dermis, three

corresponding to control populations and a fourth absent in controls (Fig. 5h and Extended Data

Fig.6c). This trajectory overlapped with the Exit GT identified in the Actβcat and ActSMO

mutants and was enriched for cell-cycle inhibitory genes, including *Trp53inp1* (Fig. 5h and

Supplementary Table 1). Additionally, label transfer of the Exit GT from the ActSMO mutant to

the Gli3 cKO dataset identified a corresponding population that expressed this gene program

(Fig. 5i), showing that this Exit gene program is shared across ActSMO, Actβcat, and Gli3 cKO

conditions. Using EdU pulse-chase experiments, we found that *Cdkn1a*-high cells in the Gli3

cKO were largely quiescent, while representing recent progeny of cell divisions (Extended Data

Fig. 5e). Thus, β-catenin and Gli3 function in a common pathway regulating cell-cycle exit.

Cross-referencing genes that lost GLI3 binding in E13.5 Actβcat dermis with those upregulated

in E13.5 Gli3 cKO dermis identified multiple cell-cycle inhibitory genes, including *Mxd4* and

Efna5, which showed increased expression and more active promoter marks (Extended Data

Fig. 6d). In addition, Exit GT genes such as *Baiap2* and *Cpt1c* also exhibited reduced GLI3

binding and increased expression. Together, these data indicate that β-catenin and Gli3 function

within a shared pathway regulating cell-cycle exit in which elevated Wnt signaling causes loss of

GLI3 chromatin occupancy and induction of genes associated with cell-cycle exit (Fig. 5j).

Uniformly high SHH activity induces dermal condensate gene expression across the Wnt

gradient

High SHH activity can induce DC gene expression before cell-cycle exit, indicating that DC

differentiation does not strictly require a preceding quiescent state (Fig. 3b). However, in the

wildtype condition, DC genes are induced abruptly as cells exit the cell cycle, suggesting that

Wnt activity contributes to the timing of DC gene induction. Given that SHH cell-autonomously

elevates Wnt activity, we examined whether Wnt signaling levels modulate how DC genes are

induced in response to high SHH activity.

To address this, we analyzed the DC gene trajectory in the E13.5 ActSMO mutant, in which

SHH activity is uniformly high across cells spanning a range of Wnt activity (Fig. 6a and

Extended Data Fig 7a). In the E14.5 wildtype DC gene trajectory, DC genes are induced

primarily at the terminal stage, coinciding with the highest levels of *Lef1* and *Ptch1* expression

(Fig. 4a,c). By contrast, in the ActSMO condition, DC genes, including *Gal*, *Sox18,* and *Sox2*,

were induced progressively across DC GT stages, with cumulative expression correlating with

increasing *Lef1* levels (Fig. 6b,c,d and Extended Data Fig. 7b). Cells with highest *Lef1* levels

showed the most advanced progression along the trajectory and the strongest DC gene

expression (Fig. 6c,d).

Consistent with the requirement for Wnt signaling in DC differentiation, cells with high SHH

activity but lacking Wnt signaling in the ActSMO condition failed to express DC genes (Fig. 6c

and Extended Data Fig. 7c)[27]. Thus, uniform SHH activation expands the window over which

Wnt-competent cells initiate DC gene expression, resulting in graded rather than abrupt

induction of DC genes. In both E14.5 wildtype and Actβcat (high Wnt) conditions, DC genes

were restricted to Wnt-active cells with high SHH activity, indicating that DC differentiation

requires SHH signaling above a defined threshold (Fig. 6e).

Coupling Wnt and SHH gradients is sufficient to synchronize the timing of cell-cycle exit

with molecular differentiation

Given that cell-cycle exit and DC differentiation coincide where high Wnt activity overlaps with

SHH signaling, we tested whether coupling these gradients is sufficient to restore temporal

coordination between the two processes. In wildtype skin, this overlap occurs in dermal cells

adjacent to placodes. We therefore induced epidermal SHH expression at E13.5 (*K14Cre;Rosa-*

LSL-SHH; SHH OE) to activate SHH signaling in Wnt-active dermal cells prior to placode

formation (Fig. 7a)[43, 44]. FISH analysis revealed that *Lef1* and *Ptch1* covaried with dermal depth

with the highest levels in cells closest to the epidermis (Fig. 7b). Compared to E13.5 controls,

Lef1 levels were increased in the superficial dermis, indicating that SHH ligand expression

enhanced Wnt activity. DC genes (e.g., *Sox2*, *Gal*) were restricted to these superficial layers

and coincided with *Cdkn1a* expression and low EdU incorporation (Fig. 7c and Extended Data

Fig. 8a). *Dkk1* expression localized to a deeper layer beneath cells expressing DC genes.

scRNA-seq analysis of E13.5 SHH OE dermis confirmed overlapping *Lef1* and *Ptch1*

expression, with DC genes and *Cdkn1a* restricted to cells with the highest combined signal

levels (Fig. 7d).

GT analysis identified three trajectories corresponding to proliferation, lower dermal

differentiation and DC differentiation, mirroring E14.5 wildtype dermis (Figs. 4a and 7e and

Extended Data Fig. 8b). Notably, a distinct Exit trajectory was not detected. Within the DC

trajectory, *Lef1* and *Ptch1* increased coordinately, and DC genes and *Cdkn1a* were induced

synchronously at the terminal stage, coincident with quiescence (Fig. 7f and Extended Data Fig.

8c). Thus, coupling Wnt and SHH activity is sufficient to restore the temporal alignment of

differentiation and cell-cycle exit.

Based on our findings in Actβcat mutants, we next asked whether loss of GLI3 chromatin

occupancy also occurs when Wnt and SHH signaling are coupled. GLI3 CUT&RUN analysis in

SHH OE dermis revealed global depletion of GLI3 binding (Fig. 7g and Extended Data Fig. 8d),

indicating that GLI3 displacement is not an artifact of genetic stabilization of β-catenin. Loss of

GLI3 binding was observed at cell cycle regulatory genes (*Mxd4* and *Efna5*) associated with

active histone marks (Extended Data Fig. 8e). In contrast to Actβcat mutants, SHH target genes

(*Ptch1*, *Gli1*) were transcriptionally activated, accompanied by increased H3K4Me3 and

H3K27Ac signal. Because SHH signaling converts cleaved GLI3 repressor into full-length GLI

activator, both detectable by anti-Flag immunoprecipitation, the observed loss of GLI3

occupancy suggests replacement by other GLI activators at SHH target loci (Fig. 7h). These

data indicate that GLI3 eviction both derepresses cell-cycle exit genes and may also prime

SHH-dependent target loci for activation.

To test whether graded Wnt and SHH inputs are sufficient to coordinate these processes

outside of the *in vivo* tissue context, we reconstituted the system *in vitro* by using E13.5

PDGFRαH2BGFP dermal cells competent to respond to both pathways[45]. These cells showed

dose-dependent gene responses to the Wnt agonist (CHIR99021) or the Smoothened agonist

(SAG) (Extended Data Fig. 9a). SAG enhanced *Lef1* expression in CHIR-treated cells in a

manner dependent on baseline Wnt activity (Extended Data Fig. 9b). High doses of CHIR alone

induced *Cdkn1a*, whereas combined CHIR and SAG treatment induced DC genes, including

Gal and *Sox18* (Extended Data Fig. 9c,d).

Finally, to assess how signal coupling affects DC morphogenesis, E13.5 dermal cells were

seeded onto a collagen-based 3D matrix with varying doses of CHIR and/or SAG (Fig. 7i and

Extended Data Fig. 9e). High-dose CHIR (5 μM) combined with SAG (50 nM) produced large

dermal clusters that expressed Robo2, a gene expressed in DC cells that regulates cell

migration and tissue morphogenesis[46, 47]. These clusters showed low proliferation, whereas

other conditions showed higher proportions of Ki67+ cells. Thus, coupling high Wnt and SHH

activity is sufficient to align molecular differentiation with cell-cycle exit *in vitro*.

Collectively, these results indicate that the pre-DC–to-DC transition comprises separable

biological processes whose timing can be differentially coordinated by Wnt and SHH. These

relationships are summarized in 7j.

Discussion

In this study, we identify a mechanism by which interacting morphogen signals generate sharp

cell-type boundaries by coordinating the timing of separable components of a cell-state

transition. During dermal condensate (DC) formation, we show that Sonic Hedgehog (SHH) acts

on Wnt-primed cells to elevate Wnt activity, which is sufficient to induce cell-cycle exit through

eviction of the transcriptional repressor GLI3 from chromatin, while SHH concurrently initiates

the differentiation program. By aligning differentiation with cell-cycle exit, this coordination

prevents proliferative expansion of intermediate states, causing transitional cells to remain

sparse and difficult to detect. In this way, boundary sharpness reflects regulated temporal

compression of a developmental transition, rather than the absence of intermediate cell states.

While sharp transitions can also arise from nonlinear ligand diffusion, our findings reveal a

distinct mechanism in which interactions between morphogen pathways synchronize concurrent

transcriptional programs.

These findings revisit classic interpretations of the French flag problem, which emphasized fixed

concentration thresholds of a polarized morphogen as primary determinants of spatial

patterning[2-6, 48]. Our data instead indicate that during DC specification, morphogen thresholds

function as triggers that initiate a committed developmental progression. In this context, SHH

gates entry into the DC differentiation program, whereas Wnt activity shapes the timing and

synchrony of its execution, establishing a division of labor between permissive and kinetic

morphogen inputs. Abrupt boundaries therefore emerge not from static positional decoding, but

from coordinated progression through a shared cell-state transition, providing a framework for

understanding how gradual signaling changes can be converted into abrupt cell-state

transitions.

Although tight coupling of Wnt and SHH activities generates a sharp and synchronous transition

during DC formation, variation in the timing, strength, or spatial alignment of interacting

morphogen gradients may support alternative modes of tissue organization. In systems

containing transit-amplifying populations, differentiation programs can proceed without

immediate cell-cycle exit, expanding intermediate states and producing graded or continuous

boundaries[7, 49, 50]. Our findings suggest that the duration and apparent abundance of transitional

states are tunable outcomes of morphogen coordination, providing a general framework for how

interacting signals regulate the emergence of distinct tissue boundaries tissues (Fig. 7j).

How SHH signaling elevates Wnt activity is unlikely to involve direct transcriptional activation of

Wnt target genes by GLI factors. Instead, our data indicate that elevated Wnt activity is

associated with loss of GLI3 chromatin occupancy at loci involved in cell-cycle regulation.

Although GLI3 eviction from chromatin does not induce large-scale chromatin architectural

reorganization, it coincides with SHH-dependent activation of differentiation genes, suggesting

that chromatin-level cross-regulation between Wnt and SHH pathways synchronizes cell-cycle

exit with molecular differentiation.

At the tissue level, this SHH-dependent enhancement of Wnt responsiveness shapes how

transitions unfold in space and time. Prior to SHH signaling, dermal cells occupy positions along

a Wnt-defined continuum, reflecting differences in transcriptional state. These differences

encode competence – how rapidly and synchronously cells traverse the transition once SHH is

engaged rather than pre-specified fates. Cells with higher Wnt activity reach cell-cycle exit more

rapidly upon SHH activation, whereas cells with lower Wnt activity transition more slowly,

expanding intermediate states. In this view, SHH synchronizes progression across a pre-

patterned landscape, compressing developmental time locally and producing sharp boundaries

without altering the sequence of states traversed.

Finally, by focusing on an early dermal transition, this work establishes a tractable mammalian

system for investigating how interacting morphogen gradients regulate the timing of concurrent

biological processes. Integrating *in vivo* genetics with trajectory-based analysis enables

resolution of overlapping differentiation programs that are otherwise difficult to detect, revealing

how dynamic signaling interactions shape discrete cell-state transitions[34].

Methods

<u>Tamoxifen induction of mice</u>

Embryos were staged as days post coitum, with embryonic day E0.5 considered as noon of the

day a vaginal plug was detected after overnight mating. Pregnant dams were given one dose of

Tamoxifen dissolved in corn oil (20mg/ml, Sigma) at 40-60 µg/gm body weight by oral gavage at

E11.5 as indicated.

<u>Mice details</u>

Axin2CreER [51] mice were bred to *Rosa-LSLSmoM2YFP* [52], *β-catenin*$^{fl(Ex3/+}$ [53] and *Gli3*$^{fl/fl}$ [54]

mice. *K14Cre* [55] mice were bred to *Shh* Overexpressor (Wang et al. 2010) and *Wntless*$^{fl/fl}$[56]mice.

Rosa-Flag-Gli3YFP [39] mice were bred to *Axin2CreER* and *β-catenin*$^{fl(Ex3/+}$ mice to enable

detection of GLI3-Flag using CUT&RUN. *R26Fucci2aR* [57] reporter mice were used to visualize

cell cycle phase. *PDGFRaH2BGFP*[45] mice were used culture embryonic dermal cells.

Sox2eGFP[58] mice were used to visualize DC cells in embryonic skin whole mounts. A random

population of both male and female mice were used for all experiments. All procedures involving

animal subjects were performed under the approval of the Institutional Animal Care and Use

Committee of the Yale School of Medicine.

<u>EdU Incorporation Assay</u>

To assess active proliferation, EdU was administered to pregnant mice intraperitoneally (25

µg/gm) and embryos were harvested after 1.5 hour. For pulse-chase experiments, 25 µg/gm

EdU was administered to pregnant mice at indicated times and embryos were harvested at later

times. EdU incorporation was assessed using the Click-it EdU Imaging kit Alexa 555 or Alexa

488 (Life technologies, c10338) according to manufacturer's instructions. Briefly, skin explants

were treated with a mixture of 1X Click-iT reaction buffer, $CuSO_4$, Alexa Fluor azide dye and 1X

reaction buffer additive, all provided in the Click-it EdU Imaging Kit, for 30 minutes at RT before

washing in PBS.

<u>Histology</u>

10% formalin-fixed paraffin embedded (FFPE) whole embryos were used for histological

analysis. FFPE embryos were sectioned at 10 µm thickness for FISH.

<u>Whole mount immunofluorescence</u>

Dorsolateral skin from embryos were microdissected and placed on nucleopore filters (VWR,

WHA 800281) and fixed overnight with 4%PFA at 4° C. Skin explants were blocked for 6 hours

in 5% normal donkey serum, 1% bovine serum albumin and 0.5% Triton-X100 at RT. Explants

were incubated overnight at 4° C in the following primary antibodies: rabbit anti-Sox2 (1:400,

Abcam, ab97959), chicken anti-GFP (1:500, Abcam, ab13970), rabbit anti-RFP (1:500,

Rockland, 600-401-379). Explants were then washed in 0.2% Tween20/PBS for 4-6 hours on a

rotator and then incubated with the following secondary antibodies: Alexa Fluor 568-donkey

anti-rabbit (1:200, Invitrogen, A10042), Alexa Fluor 488-donkey anti-rabbit (1:200, Invitrogen,

A21206), Alexa Fluor 488-goat anti-chicken (1:200, Invitrogen, A11039) overnight at RT.

Washes were carried out for 6 hours with 0.2% Tween20/PBS. Hoechst (1:750 in PBS) was

used for 45 min before mounting on slides with SlowFade Gold Antifade Mountant (Invitrogen,

S36936).

<u>*In-situ* hybridization</u>

The RNAscope Multiplex Fluorescent Detection Kit v2 (ACDBio, 323110) was used for single-

molecule fluorescence in situ hybridization (FISH) according to the manufacturer's protocol.

Briefly, sections were deparaffinized and permeabilized with hydrogen peroxide followed by

antigen retrieval and protease treatment before probe hybridization. After hybridization,

amplification and probe detection were done using the Amp 1–3 reagents. Probe channels were

targeted using the provided HRP-C1-3 reagents and tyramide signal amplification VIVID

fluorophores—650 and 570 (ACD biotechne, 323271). EdU staining was done using the Click-it

EdU Imaging Kit Alexa 488 (Life Technologies, c10338) according to the manufacturer's

instructions. Nuclear counter-stain was done using Hoechst 33342 (Invitrogen, H3570) before

mounting with SlowFade Mountant. RNA scope probes used (ACDBio)—Mm-Lef1 (441861),

Mm-Ptch1 (402811), Mm-Dkk1 (402521), Mm-Sox2 (401041), Mm-Gal (400961), Mm-Gli3

(400961) and Mm-Trp53inp1 (1161531).

<u>Microscopy</u>

FISH paraffin-embedded images were acquired using the Leica Stellaris 8 DMi8 confocal

microscope with a ×40 oil immersion (Numerical Aperture 1.3) objective lens, scanned at 5 μm

thickness, 1, 024 × 1, 024 pixel width, 400Hz. Whole mount explants were imaged in 3

dimensions using the LaVision TriM Scope II (LaVision Biotec) microscope equipped with a

Chameleon Vision II (Coherent) two-photon laser (810 -1000 nm) to acquire z-stack images

ranging from 50-120 μm (2 μm serial optical sections) using a 20X water immersion lens (NA

1.0; Olympus), scanned with a field of view of 0.3–0.5 mm^2 at 800 Hz.

Single-cell dissociation

Embryonic dorsolateral/flank skin was micro-dissected and pooled after genotyping (3 embryos

per condition) and dissociated into a single-cell suspension using 0.25% trypsin (Gibco, Life

Technologies) for 14 minutes at 37° C. Single-cell suspensions were then stained with DAPI

(Fisher Scientific, NBP2-31156) just prior to fluorescence-activated cell sorting.

Fluorescence-activated cell sorting

DAPI-excluded live skin cells were sorted on a BD FACS Aria II (Biosciences) sorter with a 100

μm nozzle. Cells were sorted in bulk and submitted for 10X Genomics library preparation at

0.75-1.0x10^6/mL concentration in 4% FCS/PBS solution.

Single-cell RNA sequencing and library preparation

Chromium Single cell 3' Library and Gel Bead Kit v2 Chromium Single Cell 3′ Library & Gel

Bead Kit v2 (PN-120237), Chromium Single Cell 3′ Chip kit v2 (PN-120236) and Chromium i7

Multiplex Kit (PN-120262) were used according to the manufacturer's instructions in the

Chromium Single Cell 3′ Reagents Kits V2 User Guide. After cDNA libraries were created, they

were subjected to Novaseq 6000 (Illumina) sequencing.

10X scRNA-seq data pipeline to matrix

The transcriptomes of live single cells from E13.5 or E14.5 mouse skin samples under different

biological conditions were sequenced. Raw 10X sequencing data was processed into a matrix

employing the standard 10X CellRanger pipeline. Briefly, base call files were fastq format which

were aligned to the mm10 reference genome followed by nUMI and barcode counting,

constructing the nUMI count matrices. nUMI matrices were filtered, centered and normalized

using Seurat@1. Briefly, for each original run condition (e.g., E13.5 wildtype), cells with > 1,000

detected genes, mitochondrial content < 25%, and ribosomal content > 5% (calculated via

PercentageFeatureSet in Seurat) were retained. Doublets were identified and removed using

scDblFinder@2. Data were then log-normalized by total UMI count.

<u>Principal Component Calculation and UMAP Visualization</u>

Principal components were computed using Seurat's RunPCA on the top 2,000 variable genes.

The number of PCs retained was determined from the elbow plot@7, and used for downstream

analysis, including dimensionality reduction for visualization and clustering.

<u>Cell type specification</u>

For each original run, UMAP was performed on log-normalized, centered, and scaled UMI count

matrices using the retained PCs. Unsupervised clustering was then conducted using Seurat's

FindClusters function with a resolution of 0.5. Clusters expressing Col1a1 were defined as

dermal, while clusters expressing Lef1 or Dkk1 were defined as upper-dermal.

<u>Gene trajectory inference</u>

We applied GeneTrajectory@3 to each sample to identify multiple, independent gene dynamic

processes. For each sample, we followed the recommended guidelines. We selected the top

2,000 highly variable genes (identified using Seurat's FindVariableFeatures) that were

expressed in >1% and <50% of cells as input. The top 20PCs were used to compute a diffusion

map, and the top 10 diffusion components were used to construct a cell k-nearest neighbor

graph (k = 10).

The number of gene trajectories and the time steps for gene inclusion within each trajectory

were determined interactively by inspecting the top three or more gene embeddings. For the

samples analyzed in this study, we identified the following gene trajectories and associated time

steps (t):

•   E14.5 wild type: 3 trajectories (t = 13, 10, 5)

•   E13.5 SmoM2: 4 trajectories (t = 1, 3, 2, 1); control: 3 trajectories (t = 3, 2, 2)

•   E13.5 TG: 3 trajectories (t = 6, 3, 3); control: 3 trajectories (t = 5, 3, 3)

•   E13.5 EX3: 4 trajectories (t = 3, 3, 2, 2); control: 3 trajectories (t = 1, 3, 3)

•   E13.5 Gli3-KO: 4 trajectories (t = 8, 7, 6, 1); control: 3 trajectories (t = 4, 3, 1)

<u>Gene bins and cell stages</u>

Following GeneTrajectory guidelines, each trajectory from mutant samples was divided into 5 or

6 gene bins. For each bin, a gene bin score was computed for both mutant and control samples

to compare gene expression dynamics across the cell embedding. Cell involved in each

trajectory were stratified into the same number of stages, based on whether they expressed

more than 50% of the genes in the corresponding gene bin. Cells assigned to multiple stages

were resolved by assigning them to the stage closer to the terminal end of the trajectory (i.e.,

farther from the branching point).

<u>Exit GT Score</u>

Exit GT scores for each sample were calculated using Seurat's AddModuleScore, based on all

genes from the exit gene trajectory identified in E13.5 SmoM2.

<u>Pathway enrichment analysis</u>

Reactome pathway enrichment analysis was performed using ReactomePA@4, with pathways

showing $p < 0.05$ considered significantly enriched.

<u>Cell cycle estimation of scRNA-seq data</u>

We employed Seurat's cell cycle scoring [59]. Briefly, averaged relative expression of those cell

cycle related genes were used to calculate G2/M and S scores, which were used for binning

cells into G2/M, S and G1/G0 bins.

<u>Data merge and normalization</u>

For comparative analysis, we merged upper-dermal cell matrices from each pair of conditions

separately (i.e. E13.5 control and Gli3-KO), followed by standard preprocessing and

normalization using the workflow provided by Seurat.

<u>Differential abundance and Differential gene expression analysis</u>

Differential abundant cells were identified using DA-seq@6, with a DA score threshold > 0.8.

Differential gene expression analysis between DA cells from E13.5 Gli3-KO and all other upper-

dermal cells from both E13.5 Gli3-KO and control was performed using Seurat's FindMarkers

without prefiltering. Significant DEGs for visualization were defined using a combined threshold

of padj < 1e-5 and log2FC > 0.23.

Integrated snATAC+RNA-seq Multi-omic Analysis

Single-cell RNA-seq (scRNA-seq) and single-cell ATAC-seq (scATAC-seq) data from E14.5

K14Cre;SHH$^{fl/fl}$ and paired control embryonic skin were processed and integrated using the

Seurat and Signac packages in R. The scRNA-seq data was normalized using SCTransform.

For the scATAC-seq component, dimensionality reduction was performed using Latent

Semantic Indexing (LSI). To integrate the modalities, we applied Weighted Nearest Neighbor

(WNN) analysis, constructing a joint neighbor graph that represents the weighted combination of

RNA and ATAC modalities. A joint UMAP was generated from the WNN graph for visualization.

Transcription factor motif accessibilities were calculated using chromVAR.

<u>CUT&RUN experiments</u>

CUT&RUN was performed with the Epicypher CUTANA™ ChIC/CUT&RUN Kit (14-1048)

according to the manufacturer's specifications with the following conditions: 500,000 cells per

sample were prepared first and nuclei were extracted according to the Epicypher CUT&RUN

Manual Appendix. Nuclei were incubated with activated ConA beads. The following antibodies

(0.5 ug of antibody per reaction) were used:

IgG Control antibody: CUTANA™ Kit Rabbit IgG CUT&RUN Negative Control Antibody,

EpiCypher Rabbit anti-H3K27Ac antibody, Millipore Mouse anti-Flag antibody, Antibodies-online

Rabbit anti-CTNNB1 antibody.

Libraries were prepared using the Epicypher CUTANA™ CUT&RUN Library Prep Kit (14-1002)

according to the manufacturer's specifications. Library quality was analyzed using Agilent

Tapestation D1000 High Sensitivity Tapes (#5067-5584). Sequencing was performed with 5

million 150bp paired-end reads per sample on an Illumina NovaSeq 6000.

<u>CUT&RUN data analysis</u>

CUT&RUN reads were analyzed for quality using FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and adaptors trimmed using

Trimmomatic. Reads were then aligned to mm9 by Bowtie2 version 2.3.1. SAMTools version

1.11 was then used to convert to BAM format, index, isolate uniquely mapped paired reads, and

remove duplicates. MACS2 version 2.2.8 was used to call peaks with IgG as input control. Read

counts across genomic intervals and peak visualization were performed using deepTools

version 3.5 and bigwig files were visualized using the Integrative Genomics Viewer. Deseq2

was used to identify differentially bound peaks between conditions, DiffBind used to plot

heatmaps of accessible/bound regions and HOMER to identify *de novo* motifs most enriched in

the differentially bound regions and annotate associated genes.

Micro-C Experiments

Micro-C was performed using the Dovetail® Micro-C Kit (Cantata Bio, PN 21006E) according to

the manufacturer's protocol using 1 million cells per reaction. E13.5 mouse dorsolateral dermis

was microdissected, dissociated, and crosslinked with formaldehyde, followed by quenching

and nuclei isolation. Chromatin was digested with micrococcal nuclease, end-repaired, and

proximity ligated to capture nucleosome-level chromatin contacts. Reverse crosslinking and

DNA purification were performed following kit specifications. Libraries were prepared using the

Dovetail® Library Module for Illumina (PN 25004E) and indexed with the manufacturer-provided

i5/i7 primer sets. Library quality and fragment size distribution were assessed using the Agilent

TapeStation High Sensitivity D5000 assay (Agilent #5067-5592, **#**5067-5593, **#**5067-5594).

Final libraries were sequenced on an Illumina NovaSeq 6000 platform to a depth of 300 million

150-bp paired-end reads per biological replicate.

Micro-C data analysis

Micro-C reads were aligned to the *mm10* genome using BWA-MEM with proximity-ligation–

optimized settings, and ligation junctions were identified, sorted, and deduplicated using the

Pairtools pipeline to generate uniquely mapped, non-duplicate valid pairs. Valid pairs were

converted into 5-kb contact matrices using cooler (v0.10.4**),** balanced by iterative correction,

and aggregated into multi-resolution .mcool files for downstream analysis. Chromatin loops

were identified using Mustache (v1.2.0), a multi-scale loop-calling algorithm that applies scale-

space representation to detect interaction peaks across resolutions in high-resolution Micro-C

maps. Loop anchors were intersected with CUT&RUN peaks using Bedtools (v2.27.1) to

annotate interactions. Genome-wide A/B chromatin compartments were computed at 40-kb

resolution using FAN-C (v0.9.28), which generated Pearson correlation matrices and extracted

the first principal component to assign A/B compartment identity. Processed matrices, loops,

and eigenvector tracks were visualized using FAN-C.

<u>Western blot</u>

Cells were lysed in RIPA buffer (Thermo Fisher Scientific, Cat# 89900) supplemented with

protease inhibitors (Thermo Fisher Scientific, Cat# 87786) for 30 minutes at 4 °C. The lysates

were clarified by centrifugation at 10,000 × g for 20 minutes at 4 °C, and protein concentrations

were determined using Bradford Protein Assay Kit (Bio-Rad, Cat# 5000006). Equal amounts of

protein (10 µg) were denatured at 95 °C for 5 minutes and loaded onto 4–12% precast

polyacrylamide gels (Bio-Rad, Cat# 3450125) for electrophoresis. Proteins were transferred to

PVDF membranes (Thermo Fisher Scientific, Cat# IB24001) using an iBlot2 transfer device

(Invitrogen, Cat# IB21001). Membranes were blocked with 5% BSA (vol/vol) in 1× TBST

(AmericanBio, Cat# AB14330-01000) at room temperature, incubated with primary antibodies

overnight at 4 °C, washed three times with TBST for 5 minutes each, incubated with HRP-

conjugated secondary antibodies for 1 hour at room temperature, and washed again three times

with TBST.

Membranes were developed using SuperSignal West Pico PLUS Chemiluminescent Substrate

(Thermo Fisher Scientific, Cat# 34577) and imaged on a ChemiDoc imaging system (Bio-Rad).

The following antibodies were used: mouse anti-Flag (Millipore, Cat# F1804), rabbit anti-Actin

(Cell Signaling Technology, Cat# 4970), HRP-conjugated goat anti-rabbit IgG (Cell Signaling

Technology, Cat# 7074), and HRP-conjugated goat anti-mouse IgG (Bio-Rad, Cat# STAR207P,

1:10,000).

<u>Dermal cell culture</u>

E13.5 PDGFRαH2BGFP skin cells were sorted based on their GFP status. GFP+ cells were

collected and pelleted (300 × g, 10 min, 4°C), resuspended at 2 × 10^6 cells/mL in complete

growth medium (DMEM + 10% FBS, Penicillin-Streptomycin, and L-Glutamine), and viability

was assessed by trypan blue exclusion. Cells were plated at 3×10^6 cells/well in 6-well plates in

2 mL total volume. For chemical perturbation assays, cells were mixed 1:1 with a 2× agonist

solution. For example, 5 µM CHIR99021 (Sigma, Cat# SML1046) was prepared in DMSO and

diluted to a working concentration of 5 µM in culture. Similarly, SAG (Sigma, Cat# SML1314)

was prepared as a 50 µM stock and diluted as needed. Media with agonist was used fresh or

stored at 4°C for up to two days. Cells were cultured in 48-72 hours. For 3D culture, 3×10^5 cells

were seeded onto each well of a µ-slide 8 Well Glass Bottom chamber (ibidi, Cat#80807) which

has been precoated by 1.5mg/mL Collagen Type I (Corning, Cat#354236). Collagen Type I was

neutralized by manufacture instructions and 200uL was added to each well 1 hour prior to

seeding the cells for solidification.

<u>Quantitative PCR</u>

Total RNA was extracted from cultured cells using the RNeasy Mini Kit (Qiagen, Cat# 74134)

according to the manufacturer's protocol. Complementary DNA (cDNA) was synthesized from

500 ng–1 µg of total RNA using the SuperScript™ III First-Strand Synthesis System (Invitrogen,

Cat# 18080051) with random hexamer primers. Quantitative PCR was performed using the

PowerUp™ SYBR™ Green Master Mix (Thermo Fisher Scientific, Cat# 4364344) on a Viia7

Real-Time PCR System (Applied Biosystems) in 384-well optical plates. Reactions were run in

technical triplicates using gene-specific primers, and threshold cycle (Ct) values were

normalized to Gapdh as a housekeeping gene. Relative expression was calculated using the

ΔΔCt method.

<u>3D culture immunofluorescence</u>

Cultured cells with collagen were fixed for 25 minutes with 4%PFA at room temperature. They

were blocked for 2-4 hours in 5% normal donkey serum, 1% bovine serum albumin and 0.2%

Triton-X100 at RT. Cells were incubated overnight at 4° C in the following primary antibodies:

rat anti-Ki67 (1:200, Invitrogen, 14-5698-82), rabbit anti-Robo2 (1:200, Cell signaling, 45568).

Cells were then washed in 1x PBS for 4-6 hours and then incubated with the following

secondary antibodies: Alexa Fluor 568-goat anti-rat (1:200, Invitrogen, A11077) and Alexa Fluor

647-donkey anti-rabbit (1:200, Invitrogen, A31573) overnight at 4° C. Washes were carried out

for 6 hours with 1x PBS. Hoechst (1:750 in PBS) was used for 45 min before washing in 1x

PBS.

<u>Image analysis</u>

Raw image stacks acquired from whole-mount explants were imported into Fiji or Adobe

Photoshop for analysis. ggplot2 and cowplot R libraries were used for graphical representation

of the scRNA-seq data. Diffusion maps were generated as described above.

<u>Quantification</u>

For quantification of DC density, 3-dimensional whole-mount tiled mosaics of skin explants with

a 1000 x 1000 μm field of view and a z-depth of 60-100 μm were used for n=3-4 embryos. For

volumetric quantification of DC cell number, EdU+ or Tom+ percentage of DC or UD, 3-

dimensional whole mount mosaics stained with Sox2 and EdU or RFP were used to manually

count positive cells, using ImageJ (Fiji) software using planar (XY) and orthogonal (YZ, XZ)

views. For volumetric quantification of cell cycle phases, 3-dimensional whole mount mosaics

stained with markers for G0/G1 cells (based on Fucci2 reporter) were done manually by

counting nuclear positive cells using ImageJ (Fiji) software. For quantification of upper dermal

cells, a distance of 10 um below the epidermis was considered upper dermis. For quantification

of the peri-DC region by whole-mount, cells two cell layers out from the Sox2+ DC were

counted. For quantification based on FISH, cells with 4-5 dots were considered positive

(according to the RNAScope manufacturer's instructions) and sections from a total of n=4

different embryos were examined. To measure RNA expression levels, H-scores were

calculated according to ACDBio manufacturer's instructions: a cell with 0 dot is scored 0, 1-3

dots score 1, 4-9 dots score 2, 10-15 dots and/or less than 10% clustered dots score 3, and

more than 15 dots and/or more than 10% clustered dots score 4; then the final H-score of a

given cell type A is calculated by summing the (% cells scored B within all cells in A)*B for score

B in 0-4. For quantification of upper dermal cells, a distance of 10-20 μm below the epidermis

was considered as upper dermis. For quantification of cells in the peri-DC region *Dkk1+Sox2-*

was considered.

<u>Statistical analysis</u>

All statistical values are expressed as mean ± SEM. An unpaired Student's *t*-test was used to

analyze data sets with two groups and *P <0.05 to, **P<0.01, ***P< 0.001 and ****P<0.00001

indicated a significant difference. When comparing more than two groups, P values were

determined by one-way ANOVA with Tukey's HSD test performed as the post hoc analysis.

Statistical calculations were performed using Prism software package (GraphPad). For

categorical data, a chi-square test was employed to estimate a P value.

No statistical methods were used to predetermine sample size. The experiments were not

randomized. The investigators were not blinded to allocation during experiments and outcome

assessment.

Data and Code Availability

Genomic datasets used in this study will be deposited in NCBI Gene Expression Omnibus

database and made publicly available as of the date of publication or with a token upon request.

References

1.   Wolpert, L. Positional information and the spatial pattern of cellular differentiation. *J*

Theor Biol **25**, 1-47 (1969).

2.   Benzinger, D. & Briscoe, J. Investigating morphogen and patterning dynamics with

optogenetic control of morphogen production. *Dev Cell* **60**, 3421-3430 e3426 (2025).

3.   Driever, W. & Nusslein-Volhard, C. The bicoid protein determines position in the

Drosophila embryo in a concentration-dependent manner. *Cell* **54**, 95-104 (1988).

4.   Greenfeld, H., Lin, J. & Mullins, M.C. The BMP signaling gradient is interpreted through

concentration thresholds in dorsal-ventral axial patterning. *PLoS Biol* **19**, e3001059

(2021).

5.   Roelink, H. *et al.* Floor plate and motor neuron induction by different concentrations of

the amino-terminal cleavage product of sonic hedgehog autoproteolysis. *Cell* **81**, 445-

455 (1995).

6.   Teague, S. *et al.* Time-integrated BMP signaling determines fate in a stem cell model for

early human development. *Nat Commun* **15**, 1471 (2024).

7. Morita, R. *et al.* Tracing the origin of hair follicle stem cells. *Nature* **594**, 547-552 (2021).

8. Ouspenskaia, T., Matos, I., Mertz, A.F., Fiore, V.F. & Fuchs, E. WNT-SHH Antagonism

Specifies and Expands Stem Cells prior to Niche Formation. *Cell* **164**, 156-169 (2016).

9. Rompolas, P. *et al.* Live imaging of stem cell and progeny behaviour in physiological

hair-follicle regeneration. *Nature* **487**, 496-499 (2012).

10. Zhang, C. *et al.* Escape of hair follicle stem cells causes stem cell exhaustion during

aging. *Nat Aging* **1**, 889-903 (2021).

11. Leybova, L. *et al.* Radially patterned morphogenesis of murine hair follicle placodes

ensures robust epithelial budding. *Dev Cell* **59**, 3272-3289 e3275 (2024).

12. Driskell, R.R. *et al.* Distinct fibroblast lineages determine dermal architecture in skin

development and repair. *Nature* **504**, 277-281 (2013).

13. Myung, P., Andl, T. & Atit, R. The origins of skin diversity: lessons from dermal

fibroblasts. *Development* **149** (2022).

14. Chuong, C.M., Widelitz, R.B., Ting-Berreth, S. & Jiang, T.X. Early events during avian

skin appendage regeneration: dependence on epithelial-mesenchymal interaction and

order of molecular reappearance. *J Invest Dermatol* **107**, 639-646 (1996).

15. Chase, H.B., Rauch, R. & Smith, V.W. Critical stages of hair development and

pigmentation in the mouse. *Physiol Zool* **24**, 1-8 (1951).

16. Gupta, K. *et al.* Single-Cell Analysis Reveals a Hair Follicle Dermal Niche Molecular

Differentiation Trajectory that Begins Prior to Morphogenesis. *Dev Cell* **48**, 17-31 e16

(2019).

17. Hardy, M.H. The secret life of the hair follicle. *Trends Genet* **8**, 55-61 (1992).

18. Millar, S.E. *et al.* WNT signaling in the control of hair growth and structure. *Dev Biol* **207**,

133-149 (1999).

19. Paus, R. *et al.* A comprehensive guide for the recognition and classification of distinct

stages of hair follicle morphogenesis. *J Invest Dermatol* **113**, 523-532 (1999).

20. Saxena, N., Mok, K.W. & Rendl, M. An updated classification of hair follicle

morphogenesis. *Exp Dermatol* **28**, 332-344 (2019).

21. Woo, W.M., Zhen, H.H. & Oro, A.E. Shh maintains dermal papilla identity and hair

morphogenesis via a Noggin-Shh regulatory loop. *Genes Dev* **26**, 1235-1246 (2012).

22. Chen, D., Jarrell, A., Guo, C., Lang, R. & Atit, R. Dermal beta-catenin activity in

response to epidermal Wnt ligands is required for fibroblast proliferation and hair follicle

initiation. *Development* **139**, 1522-1533 (2012).

23. Noramly, S., Freeman, A. & Morgan, B.A. beta-catenin signaling can initiate feather bud

development. *Development* **126**, 3509-3521 (1999).

24. Chiang, C. *et al.* Essential role for Sonic hedgehog during hair follicle morphogenesis.

Dev Biol **205**, 1-9 (1999).

25. St-Jacques, B. *et al.* Sonic hedgehog signaling is essential for hair development. *Curr*

Biol **8**, 1058-1068 (1998).

26. Gritli-Linde, A. *et al.* Abnormal hair development and apparent follicular transformation to

mammary gland in the absence of hedgehog signaling. *Dev Cell* **12**, 99-112 (2007).

27. Qu, R. *et al.* Decomposing a deterministic path to mesenchymal niche formation by two

intersecting morphogen gradients. *Dev Cell* **57**, 1053-1067 e1055 (2022).

28. Xu, Z. *et al.* Embryonic attenuated Wnt/beta-catenin signaling defines niche location and

long-term stem cell fate in hair follicle. *Elife* **4**, e10567 (2015).

29. Ben-Zvi, D., Shilo, B.Z., Fainsod, A. & Barkai, N. Scaling of the BMP activation gradient

in Xenopus embryos. *Nature* **453**, 1205-1211 (2008).

30. Dekanty, A. & Milan, M. The interplay between morphogens and tissue growth. *EMBO*

Rep **12**, 1003-1010 (2011).

31. Gregor, T., Wieschaus, E.F., McGregor, A.P., Bialek, W. & Tank, D.W. Stability and

nuclear dynamics of the bicoid morphogen gradient. *Cell* **130**, 141-152 (2007).

32. Kicheva, A., Bollenbach, T., Wartlick, O., Julicher, F. & Gonzalez-Gaitan, M.

Investigating the principles of morphogen gradient formation: from tissues to cells. *Curr*

Opin Genet Dev **22**, 527-532 (2012).

33. Lander, A.D., Nie, Q. & Wan, F.Y. Do morphogen gradients arise by diffusion? *Dev Cell*

2, 785-796 (2002).

34. Qu, R. *et al.* Gene trajectory inference for single-cell data by optimal transport metrics. *Nat Biotechnol* (2024).

35. Biggs, L.C. *et al.* Hair follicle dermal condensation forms via Fgf20 primed cell cycle exit, cell motility, and aggregation. *Elife* **7** (2018).

36. Zhang, Y. *et al.* Reciprocal requirements for EDA/EDAR/NF-kappaB and Wnt/beta-catenin signaling pathways in hair follicle induction. *Dev Cell* **17**, 49-61 (2009).

37. Fu, J. & Hsu, W. Epidermal Wnt controls hair follicle induction by orchestrating dynamic signaling crosstalk between the epidermis and dermis. *J Invest Dermatol* **133**, 890-898 (2013).

38. Alvarez-Medina, R., Cayuso, J., Okubo, T., Takada, S. & Marti, E. Wnt canonical pathway restricts graded Shh/Gli patterning activity through the regulation of Gli3 expression. *Development* **135**, 237-247 (2008).

39. Vokes, S.A., Ji, H., Wong, W.H. & McMahon, A.P. A genome-scale analysis of the cis-regulatory circuitry underlying sonic hedgehog-mediated patterning of the mammalian limb. *Genes Dev* **22**, 2651-2663 (2008).

40. te Welscher, P. *et al.* Progression of vertebrate limb development through SHH-mediated counteraction of GLI3. *Science* **298**, 827-830 (2002).

41. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589 (2010).

42. Lex, R.K. *et al.* GLI transcriptional repression is inert prior to Hedgehog pathway activation. *Nat Commun* **13**, 808 (2022).

43. Phan, Q.M. *et al.* Lef1 expression in fibroblasts maintains developmental potential in adult skin to regenerate wounds. *Elife* **9** (2020).

44. Reddy, S. *et al.* Characterization of Wnt gene expression in developing and postnatal hair follicles and identification of Wnt5a as a target of Sonic hedgehog in hair follicle morphogenesis. *Mech Dev* **107**, 69-82 (2001).

45.   Hamilton, T.G., Klinghoffer, R.A., Corrin, P.D. & Soriano, P. Evolutionary divergence of

platelet-derived growth factor alpha receptor signaling mechanisms. *Mol Cell Biol* **23**,

4013-4025 (2003).

46.   Guzman-Palma, P. *et al.* Slit/Robo Signaling Regulates Multiple Stages of the

Development of the Drosophila Motion Detection System. *Front Cell Dev Biol* **9**, 612645

(2021).

47.   Goncalves, A.N., Correia-Pinto, J. & Nogueira-Silva, C. ROBO2 signaling in lung

development regulates SOX2/SOX9 balance, branching morphogenesis and is

dysregulated in nitrofen-induced congenital diaphragmatic hernia. *Respir Res* **21**, 302

(2020).

48.   Reyes, R., Lander, A.D. & Nahmad, M. Dynamic readout of the Hh gradient in the

Drosophila wing disc reveals pattern-specific tradeoffs between robustness and

precision. *Elife* **13** (2024).

49.   Sun, Q. *et al.* Dedifferentiation maintains melanocyte stem cells in a dynamic niche.

Nature **616**, 774-782 (2023).

50.   Myung, P. & Ito, M. Dissecting the bulge in hair regeneration. *J Clin Invest* **122**, 448-454

(2012).

51.   van Amerongen, R., Bowman, A.N. & Nusse, R. Developmental stage and time dictate

the fate of Wnt/beta-catenin-responsive stem cells in the mammary gland. *Cell Stem Cell*

11, 387-400 (2012).

52.   Jeong, J., Mao, J., Tenzen, T., Kottmann, A.H. & McMahon, A.P. Hedgehog signaling in

the neural crest cells regulates the patterning and growth of facial primordia. *Genes Dev*

18, 937-951 (2004).

53.   Harada, N. *et al.* Intestinal polyposis in mice with a dominant stable mutation of the beta-

catenin gene. *EMBO J* **18**, 5931-5942 (1999).

54.   Blaess, S., Stephen, D. & Joyner, A.L. Gli3 coordinates three-dimensional patterning

and growth of the tectum and cerebellum by integrating Shh and Fgf8 signaling.

Development **135**, 2093-2103 (2008).

55.   Dassule, H.R., Lewis, P., Bei, M., Maas, R. & McMahon, A.P. Sonic hedgehog regulates

growth and morphogenesis of the tooth. *Development* **127**, 4775-4785 (2000).

56.   Carpenter, A.C., Rao, S., Wells, J.M., Campbell, K. & Lang, R.A. Generation of mice

with a conditional null allele for Wntless. *Genesis* **48**, 554-558 (2010).

57.   Abe, T. *et al.* Visualization of cell cycle in mouse embryos with Fucci2 reporter directed

by Rosa26 promoter. *Development* **140**, 237-246 (2013).

58.   Arnold, K. *et al.* Sox2(+) adult stem and progenitor cells are important for tissue

regeneration and survival of mice. *Cell Stem Cell* **9**, 317-329 (2011).

59.   Nestorowa, S. *et al.* A single-cell resolution map of mouse hematopoietic stem and

progenitor cell differentiation. *Blood* **128**, e20-31 (2016).

Acknowledgements

We thank D. Gay, M. Ito, J. Levinsohn, and R. Atit for review and discussion of this manuscript.

We thank J. Wang and K. Sumigray for generously sharing their Stellaris equipment and help

with imaging troubleshooting. Inducible SHH ligand transgenic mice were kindly provided by

David Wang. This work was supported by National Institutes of Health grants R01AR076420

(P.M.) and LEO Foundation LF-OC-23-001347 (P.M.).

Author Contributions

Conceptualization, P.M., Y.K., R.L., Y.J., S.P.; Methodology, P.M., R.L., Y.K., Y.J., S.P., E.B.,

S.V., H.L.; Software, R.L., Y.K.; Resources, M.T., K.P.; Investigation, P.M., R.L., Y.J., S.P. H.L,

E.B., S.V.; Writing – Original Draft, P.M.; Writing – Review & Editing, P.M., R.L., Y.J., S.P., T.X.,

R.D., Y.K., C.L., S.L., E.B.; Supervision, P.M., Y.K.; Funding Acquisition, P.M.

Competing Interests

The authors declare no competing interests.

Main Figure Legends:

Fig. 1: Cell-cycle exit and molecular differentiation are tightly coupled during dermal

condensate formation

a, Left, illustration of the French flag problem with stripes representing distinct cell-type readouts

positioned relative to a fixed morphogen source. Classical models propose that threshold

concentrations of morphogen specify different cell-types with sharp borders. Right, schematic

depiction of cellular responses to a linear morphogen gradient, illustrating either linear

responses or non-linear transitions across morphogen levels. This contrasts static spatial

threshold interpretations with a dynamic view in which cells traverse morphogen levels during

fate transitions. **b,** FISH images of E14.5 mouse skin showing *Sox2*+ DCs form by E14.5, which

lack EdU copositivity; quantification of %EdU by cell-type (UD, upper interfollicular dermis)

(n=10). **c,** FISH image showing peri-DC (*Dkk1*+) cells adjacent to *Sox2*+ with *Cdkn1a* enriched

at the DC boundary; (n=10). **d,** E14.5 skin stained for *Lef1* and *Ptch1* with transcripts quantified

by spatial regions (LD, lower dermis) or %EdU+ by *Lef1* bin (n=5). **e,** Schematic summarizing

dermal cell states with respect to spatial position and relative Wnt and SHH activity levels. DCs

arise within regions of overlapping Wnt and SHH activity, where cell-cycle exit and DC gene

expression occur within a narrow temporal window. Data as mean± SEM; *$P<0.05$, **$P<0.01$,

***$P<0.001$, ****$P<0.0001$, one-way ANOVA; ns, not significant. Scale bars=50 µm.

Fig. 2: High Wnt activity induces cell-cycle exit independently of SHH signaling

a, Experimental setup in which β–catenin was constitutively activated in the dermis prior to DC

formation (tamoxifen E11.5). **b,** UMAPs colored by indicated genes or cell cycle (CC) phase for

E13.5 Actβcat mutant or paired control dermal scRNA-seq data (n=2). **c,** FISH showing E13.5

Lef1, *Cdkn1a*, EdU (left) and quantification of UD *Cdkn1a* transcripts (H-score) or %EdU of

Cdkn1a (*p21*)+ or *p21*- cells in Actβcat UD vs. control UD (*p21*-) (n=3). **d,** FISH showing E14.5

Sox2, Cdkn1a, EdU (left) and quantification of %EdU+ cells in control or Actβcat *Cdkn1a*+ cells

(almost all *Cdkn1a*+ cells co-express *Sox2* in control, n=5). **e,** E14.5 *Lef1*, *Ptch1*, *Sox2* FISH-

stained sections (left) and quantification of *Ptch1* levels in Actβcat by cell type. **f,** Illustration

showing that activated βcatenin (elevated Wnt activity) induces cell-cycle exit at E13.5 in the

absence of SHH. At later stages, DC gene expression is restricted to cells experiencing high

SHH activity with high Wnt activity. Data as mean± SEM; *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001, one-way ANOVA; ns, not significant. Scale bars=50 μm.

Fig. 3: SHH signaling cell-autonomously elevates Wnt activity in association with cell-

cycle exit

a, Experimental setup showing expression of constitutively activated Smoothened in dermal

cells prior to DC formation when Wnt activity is low (tamoxifen E11.5). **b,** Whole-mount images

of E14.5 ActSMO and control skin stained with EdU and Sox2 with quantification of %EdU+ of

Sox2+ cells (n=8). **c,** UMAPs of E13.5 control and ActSMO dermal scRNA-seq data with

indicated gene expression; right, quantification of scRNA-seq *Lef1* levels in UD cells. **d,** E13.5

control or ActSMO mutant skin stained for *Lef1*, *Ptch1* and *Cdkn1a* with quantification of

average *Lef1* expression by *Ptch1* expression status (n=8). **e,** E13.5 skin showing ActSMO

boosts *Lef1* gene expression which corresponds with decreased EdU incorporation (%EdU). **f,**

Cartoon illustrating that high SHH activity cell-autonomously elevates Wnt signaling, and

elevated Wnt activity coincides with cell-cycle exit. Data as mean± SEM; *$P<0.05$, **$P<0.01$,

***$P<0.001$, ****$P<0.0001$, one-way ANOVA. Scale bars, 50 µm.

Fig. 4: GeneTrajectory defines transcriptional changes accompanying cell-cycle exit

a, GeneTrajectory diffusion map and gene order of E14.5 wildtype DC GT with dermal UMAPs

colored by DC GT stages. **b,** *Lef1* and *Ptch1* levels by DC GT stage. **c,** DC gene expression by

DC GT stage. Right, G1/G0 fraction by DC GT stage with UMAP of *Cdkn1a* expression or CC

phase. **d,** E13.5 ActSMO GT diffusion map. **e,** ActSMO and control dermal UMAPs colored by

expression of the mutant New GT stages or CC phase. **f,** Pathway analysis of genes from new

mutant GT and UMAPs of control and mutant dermal cells colored by indicated genes. **g,** FISH

images of indicated genes in E13.5 control or ActSMO skin with quantification of *Lef1*

expression in control E13.5 upper dermal (UD) or E14.5 DC cells vs. E13.5 ActSMO *Trp53inp1+*

cells (n=3). **h,** Label transfer of the E13.5 ActSMO new GT onto indicated E13.5 mutant or

control dermal populations showing cells that express this similar gene module. **i,** EdU pulse-

chase experiments of E13.5-E14.5 ActSMO mutant and corresponding quantification of %EdU

of *Trp53inp1+* cells (n=3). Bottom cartoon illustrating *Trp53inp1+* cells undergo cell cycle exit as

immediately after a division. Data as mean ± SEM, one-way ANOVA. Scale bars, 50 µm.

Fig. 5: Loss of GLI3 chromatin binding induces premature cell-cycle exit downstream of elevated Wnt signaling

a, *Gli3* expression in E13.5 control and Actβcat skin prior to SHH. **b,** Heat map of GLI3 binding at all transcriptional start sites (TSS) in E13.5 control or Actβcat skin. **c,** KEGG pathway analysis of GLI3 unbound regions in the Actβcat mutant. Right, Genome browser tracks showing binding signal of respective antibodies at indicated genes. **d,** Western blot of cell lysates from control or Actβcat mutant skin stained with anti-Flag (Gli3) or β-actin (n=3). **e,** FISH of E14.5 control and Gli3 dermal cKO (tamoxifen E11.5) stained for indicated genes; quantification of DC size and %EdU of control or Gli3 cKO by cell type (n=3). **f,** FISH of E13.5 control and Gli3 cKO stained for indicated genes with quantification of transcripts by FISH (n=4). **g,** snATAC+RNA-seq UMAPs of co-embedded E14.5 control and epidermal SHH cKO showing either gene expression or GLI motif accessibility. **h,** GT diffusion map of E13.5 Gli3 cKO dermal cells, showing an Exit GT. UMAPs of control or Gli3 cKO colored by Exit GT bins, indicated genes or cell-cycle phase. **i,** Label transfer of E13.5 ActSMO Exit GT onto Gli3 cKO or control scRNA-seq data showing a shared Exit gene GT. **j,** Elevated Wnt activity reduces GLI3 chromatin occupancy, derepressing genes that mediate cell-cycle exit, but not DC differentiation. Here, canonical SHH target genes and DC differentiation genes remain inactive in the absence of SHH signaling. Upon co-activation of Wnt and SHH, canonical SHH targets and DC genes are induced. Data as mean± SEM; *$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, one-way ANOVA; ns, not significant; scale bars, 50 μm.

Fig. 6: Uniformly high SHH activity induces dermal condensate gene expression across

the Wnt gradient

a, E13.5 ActSMO dermal GT diffusion map. **b,** Gene order (top) of ActSMO DC GT, showing DC

genes expressed across the trajectory; expression of DC GT stages colored onto ActSMO

dermal UMAPs. **c,** Expression levels of DC genes by DC GT stage. **d,** *Lef1* levels by DC GT

stage of ActSMO and control. **e,** Cartoon illustrating that uniform SHH activation permits DC

gene expression across a range of Wnt activity levels, resulting in sequential induction of DC

genes that correlates with *Lef1* levels. Here, DC genes can be expressed in proliferating cells of

lower Wnt activity, expanding intermediate states before coordinated cell-cycle exit is achieved.

The orange/gray bars represent the extent of intermediate DC transcriptional states along the

trajectory and not a direct measure of temporal kinetics.

1027

Fig. 7: Coordinated Wnt and SHH signaling synchronizes cell-cycle exit and dermal condensate differentiation

a, Experimental design to test whether coordinated Wnt and SHH activity is sufficient to synchronize cell cycle exit with DC genes by co-expressing Wnt and SHH ligands across the epidermis at E13.5 (*K14Cre;Lox-STOP-Lox-Shh*; SHH OE). **b,** FISH images of E13.5 control and mutant showing covarying levels of *Ptch1* and *Lef1* across vertical dermal layers in the SHH OE mutant and elevated *Lef1* levels in layers where *Ptch1* is concurrently activated. **c,** E14.5 control and SHH OE with spatial partitioning of cells by cell type genes and *Cdkn1a* coinciding with *Sox2* expression (n=3). **d,** E13.5 control or SHH OE UMAPs with indicated genes or cell-cycle phase. **e,** GT diffusion map of E13.5 SHH OE dermal scRNA-seq data. **f,** DC GT gene order of SHH OE (top) with expression of DC genes and fraction G1/G0 by DC GT stage (bottom) showing coincident induction of DC genes and cell-cycle exit at the terminal stage. Right, covarying *Lef1* and *Ptch1* levels by DC GT stage. **g,** Heat map of GLI3 binding at TSS in E13.5 control versus SHH OE. Genome browser tracks showing reduced GLI3 binding and active histone marks at *Gli1* and *Ptch1*. **h,** Cartoon showing loss of GLI3 binding upon Wnt and SHH coactivation with activation of canonical SHH targets. **i,** Confocal images of 3D dermal cultures stained with Robo2 (DC marker) showing Robo2+ clusters lacking Ki67 in high CHIR+SAG condition. Right, dermal cluster number in 3D cultures by condition. **j,** Model illustrating how coordination between Wnt and SHH signaling regulates the timing of cell-cycle exit and molecular differentiation during the pre-DC-to-DC transition. When morphogen activities are uncoupled, intermediates states proliferate and persist over a broader temporal and/or spatial window. Coordinated morphogen dynamics compress the transitional window, yielding an abrupt cell-state transition. The pink bar in wildtype indicates the spatial field of SHH activation. The dotted line represents the spatial profile of Wnt activity observed at E13.5 prior to SHH expression, modeled here as a baseline gradient. Data as mean± SEM; **$P<0.01$, ***$P<0.001$, ****$P<0.0001$, one-way ANOVA; ns, not significant; scale bars, 50 μm.

Fig. 1

a   **How do sharp cell-type borders form from a smooth morphogen gradient?**

Gradient model for the French Flag Problem

Morphogen concentration ▭ Source

Cell-type Source

Sharp borders by threshold levels of morphogen

Morphogen Concentration C_1 C_2 Source

Cells traverse a range of morphogen levels during transition

Cell-type Transition

······ Spatial morphogen gradient
—— Morphogen activity

Morphogen Activity/Concentration C_1
Gradual cell-type Transition
Fuzzy border

Morphogen Activity/Concentration C_1
Temporally condensed cell-type Transition
Sharp border

b   *Sox2* EdU   E14.5

Proliferation
% EdU+
60
40
20
0
**** ****
UD   Peri DC (*Dkk1*+)   DC (*Sox2*+)

c   *Cdkn1a* *Sox2* *Dkk1*   E14.5
Sox2
Cdkn1a
Dkk1

d   *Lef1* *Ptch1* EdU   E14.5
Lef1
Ptch1
EdU

Lef1 or *Ptch1* Level
400
300
200
100
**** **** **
ns
LD UD Peri DC (*Dkk1*+) DC (*Sox2*+)
Lef1 *Ptch1*

Proliferation
%EdU
*** **** ns ns *
0 1 2 3 4
Lef1 level (bin)

E14.0-E14.5
Wnts   ↑Wnts   Wnts
SHH
Upper Dermis (UD)
Lower Dermis

○ G0 (Queiescent) cell
◌ G1 cell
⬭ Proliferating cell
■ DC cell (*Sox2*+)
▨ Peri-DC cell (*Dkk1*+)
□ Other UD cell

e   **How does a sharp cell-type border form by overlapping Wnt and SHH gradients?**

E14.5 Cell types
UD PeriDC DC
Cell-type Responses:
DC genes
Cell cycle arrest
Source

Peri-DC-to-DC transition

Synchronized temporal progression
Wnt/SHH Activity
← DC genes
Cell-type Transition
Sharp border

Wnt/SHH Activity
← Cell cycle arrest
Cell-type Transition

Fig. 2

Fig. 3

a Experimental spatial activation of signal

Control E13.5 ActSMO

Epidermis
Dermis

Wnts / Wnt / SHH

dermal Smoothened activation

b E14.5

Control / ActSMO

Sox2 / EdU

DC Proliferation
% EdU
CTL ActSMO

c E13.5 ActSMO Dermal scRNA-seq

Lef1 *Ptch1* *Sox2* *Gal* *Cdkn1a*

Control
ActSMO

Lef1 Expression
Expression
ActSMO Control

d E13.5

Control / ActSMO

Lef1 / Ptch1 / Cdkn1a

E13.5 *Lef1* level by *Ptch1* status
Lef1 H-Score
Control / ActSMO
Ptch1- *Ptch1+*

e E13.5

Control / ActSMO

Lef1 / EdU

%EdU by *Lef1* level
%EdU
Lef1 bin
ActSMO

f E13.5 ActSMO
+SHH activation

Wnt Activity

SHH cell-autonomously boosts Wnt activity

E13.5 Wildtype

Wnt activity / SHH activity
Wnt threshold (Exit)
SHH = 0
CC exit
Cell-type transition

E13.5 ActSMO

Wnt activity / SHH activity
Wnt threshold (Exit)
High SHH (ActSMO)
ActSMO
Wlildtype
CC exit
Cell-type transition

ActSMO Wnt activity (High SHH)
Wnt activity (No SHH)

Uniform SH activation increases Wnt output cell-autonomously enabling cells with higher baseline Wnt activity to reach the threshold required for cell-cycle exit

Fig. 4

a Gene Trajectories

GeneTrajectory on E14.5 Wildtype dermis

E14.5 DC Gene Trajectory Gene Order

Grem1 Dkk1 Runx3 Lef1 Ptch1 Bmp4 Cdkn1a DC genes Sox2 Foxd1 Gal Sox18

Stage 1 Stage 2 Stage 3 Stage 4 Stage 5 Stage 6

DC

b *Lef1* level by stage

Ptch1 level by stage

c DC gene expression by GT stage

Gal   *Sox18*   *Sox2*   *Sox2*   G1/G0 fraction by DC stage   *Cdkn1a*   CC phase
G1/G0
S
G2/M

d E13.5 ActSMO Gene Trajectories

S/G2/M
New GT
DC
LD

e New GT projected on ActSMO and Control

Stage 1   Stage 2   Stage 3   Stage 4   Stage 5   CC Phase
ActSMO
Control

f Reactome Pathway Analysis

Transcriptional Regulation by TP53
TP53 Regulates Metabolic Genes
TNFR2 non-canonical NF-kB
Programmed cell death
Keratinization

E13.5 ActSMO
Trp53inp1   *Cdkn1a*
ActSMO
Control

g E13.5 Control

Lef1
Ptch1
Trp53inp1

Trp53inp1
Lef1
Ptch1

E13.5 ActSMO

Lef1
Ptch1
Trp53inp1

Trp53inp1
Lef1
Ptch1

Lef1 level
Control
ActSMO

UD *Trp53*+ DC
E13.5   E14.5

h Label transfer ActSMO
New GT
ActSMO   Actβcat

Wntless cKO   Control

Exit GT Score

i EdU

12.5 13.5 14.5

EdU

12.5 13.5 14.5

E14.5   *Trp53inp1*
EdU

E14.5   *Trp53inp1*
EdU

E14.5
EdU pulse

Trp53inp +
EdU- (G0)

E13.5
EdU pulse

Trp53inp -
EdU+ (S phase)

E13.5 -> E14.5
EdU pulse-chase

E14.5
EdU chase

Trp53inp +
EdU-retained (G0)

EdU pulse (1 hr) vs EdU chase (24 hr)
Control
ActSMO

Duration after EdU
1 hr 24 hr   1 hr 24 hr   1 hr 24 hr
E13.5   E14.5   E14.5
Trp53+   *Trp53*+   DC

Fig. 5

a E13.5 Control
Lef1
Gli3

E13.5 Actβcat
Lef1
Gli3

b E13.5 Control | E13.5 Actβcat

sgGroup
Sites
signal

Gli3 binding signal

c KEGG Pathway analysis on GLI3-bound genes in E13.5 Control vs. Actβcat

Basal cell carcinoma
Hippo pathway
Notch pathway
Hedgehog pathway
Longevity regulating
Stem cell pluripotency
Hepatic carcinoma
TGF-beta signaling
Transcript misregulation
Fatty acid metabolism

EnrichmentScore (-log10(pvalue))

Count
pvalue
0.009
0.006
0.003

GLI3 binding E13.5 Actβcat
Control
Actβcat

GLI3
H3K4 Me3
H3K 27Ac
IgG

Lef1 *Gli1* *Ptch1*

d
kDa CTL Act βcat
α-Flag
250 — ◄ FL GLI3
150 —
100 — ◄ Cleaved GLI3
75 —
50 — β–actin
37 —
25 —

e Control | Gli3 cKO
Cdkn1a/p21
Dkk1
EdU
E14.5
Cdkn1a
Dkk1
EdU

#*Sox2* cells/DC
**
Cont Gli3 cKO

E14.5
EdU%
ns * ns
Control Gli3 cKO
UD *Dkk1*+ DC

f Control | Gli3 cKO
Cdkn1a/p21
Lef1
EdU
E13.5
Lef1
Cdkn1a
EdU

E13.5 Gene expression
H-score
**** ns ns ****
Gli3 *Lef1* *Ptch1* *Cdkn1a*

g E14.5 SHH cKO snATAC+RNA-seq
Control SHH cKO
Gene Expression (GEX)
Lef1
Gli1
Gli3
Motif Accessibility
GLI

h E13.5 Gli3 cKO Gene Trajectories
UD LD
S2/G2/M
Exit

E13.5 Gli3 cKO Exit GT onto cKO and control UMAP
Gli3 cKO
Stage 1 Stage 2 Stage 3 Stage 4 Stage 5 CC phase *Trp53inp1* *Sox18*
Control

CC phase
G1/G0
S
G2/M

i Label transfer of E13.5 ActSMO Exit GT
ActSMO Gli3 cKO
Control
Exit GT Score
0.8
0.6
0.4
0.2
0.0

j
E13.5 Wildtype

GLI3 binding is inert
GLI3 → SHH target genes

Loss of GLI3 binding derepresses
GLI3 → CC Exit Genes

E13.5 Actβcat
Elevated Wnt Activity

GLI3 → SHH target genes

GLI3 → CC Exit Genes

Fig. 6

a E13.5 ActSMO Gene Trajectories

b E13.5 ActSMO DC GeneTrajectory Gene Order

Stage 1 | Stage 2 | Stage 3 | Stage 4 | Stage 5

DC
ActSMO

c DC gene expression by DC GT stage

Gal   *Sox18*   *Sox2*

ActSMO DC GT stage

d *Lef1* level by stage
Control

Lef1 level

ActSMO

Lef1 level

ActSMO DC stage

e

E14.5 Wildtype

Wnts   ↑Wnts   Wnts
SHH

E13.5 ActSMO

Wnts

Wnt activity   SHH activity

Wnt threshold (Exit)   SHH threshold (DC genes)

Cell-type transition

DC genes
Binary

Compressed Transition

Cell Density

Sparse intermediates (Few visible cells)

Wnt activity   SHH activity

High SHH

Wnt threshold (Exit)   SHH threshold (DC genes)

Cell-type transition

DC genes
Graded across Wnt levels

—— Wnt activity (With SHH)
····· Wnt activity (No SHH)

Expanded Transition

Cell Density

Expanded intermediates (More visible cells)

Coupling of Wnt-dependent cell-cycle exit with SHH-driven differentiation prevents proliferative expansion of intermediate states, restricting their accumulation during the transition

Fig. 7

a Experimental spatial activation of signal
E13.5 SHH OE

Temporally aligned DC genes with arrest

b E13.5 Expression by space

Lef1 by Layer

Ptch1 by Layer

c E14.5

d E13.5 Control   E13.5 K14Cre;LSL-SHH (SHH OE)

G1/G0
S
G2/M

e Gene Trajectories
E13.5 SHH OE

f Stage 1 Stage 2 Stage 3 Stage 4 Stage 5

DC Gene Order

DC gene expression by stage

Lef1 by stage   *Ptch1 by stage*

Gal   *Sox18*   *Sox2*   Fraction in G1/G0

DC Stage

g Control   SHH OE

Gli3 binding signal

Control
Gli3-Flag
SHH OE

H3K4Me3

H3K27Ac

IgG

Gli1 *Ptch1*

h

E13.5 Actβcat
High Wnt activity (No SHH)

GLI3

SHH targets

GLI3 binding is inert

E13.5 SHH OE
High Wnt + High SHH activity

GLI3
GLIA
SHH targets

SHH-dependent GLI Activator

i DMSO   SAG

Low CHIR + SAG   High CHIR + SAG

Robo2
Ki67
DAPI

No. clusters per well

CHIR (uM) 0 1 5 0 1 5
SAG (nM)   0 0 0 50 50 50

j E14.5 Wildtype

SHH Gate
Localized SHH field

Wnt Activity

Arrest

Compressed transition

Sharp border
Synchronous neighbors

Wnt activity (with SHH+)
Baseline Wnt activity (E13.5)

Pre-DC ▶ DC (Narrow transition)

Local SHH synchronizes cell-cycle exit + differentiation

ActSMO
Uniform SHH

SHH Gate

Wnt Activity

Arrest

Expanded transition

Fuzzy border
Asynchronous

Pre-DC ▶ DC (Wider transition)

Differentiation without synchronized exit