<u>Title</u>

Parallel circuits in the posterior parietal cortex balance behavioral flexibility and

stability

<u>Authors</u>

Eunji Jung[1,6], Jong-Hoon Lee[2,3,6], Woochul Choi[4], Gil-Hyun Kim[3], Geonhui Ryu[1], Se-Bum

Paik[5], Seung-Hee Lee[1,2,3,*]

<u>Affiliations</u>

[1] Department of Biological Sciences, Korea Advanced Institute of Science and Technology,

Daejeon, 34141, Republic of Korea

[2] KAIST Institute for the BioCentury, Korea Advanced Institute of Science and Technology,

Daejeon, 34141, Republic of Korea

[3] Center for Synaptic Brain Dysfunctions, Institute for Basic Science, Daejeon 34141,

Republic of Korea

[4] Cognitive Science Research Group, Korea Brain Research Institute, Daegu 41062, Republic

of Korea

[5] Department of Brain and Cognitive Sciences, Korea Advanced Institute of Science and

Technology, Daejeon 34141, Republic of Korea

[6] These authors contributed equally.

[*] Corresponding author: shlee1@kaist.ac.kr

<u>Abstract</u>

Rapid behavioral adaptation requires the brain to solve a fundamental computational dilemma:

how to flexibly update learned rules while maintaining stable motor performance. This

flexibility-stability trade-off is central to cognitive function and artificial intelligence, yet its

neural circuit basis is poorly understood. Here, we identify the circuit architecture that

coordinates this balance. Using a within-session auditory reversal learning task combined

with circuit-specific manipulations in mice, we demonstrate that the posterior parietal cortex

(PPC) orchestrates adaptive learning through two parallel, anatomically distinct pathways to

the auditory cortex (AC) and inferior colliculus (IC). These two circuits perform a double

dissociation of function: the PPC-to-AC circuit facilitates flexible rule updating by encoding

dynamic stimulus-outcome contingencies, while the PPC-to-IC circuit preserves stable action

execution across changing rules. A feedforward network model further demonstrates that

segregated stable and flexible top-down projections are necessary and sufficient for rapid

reversal learning. This parallel architecture segregates adaptive and stable computations into

distinct circuits, providing a mechanistic framework for cognitive flexibility and offering

potential insight into conditions where this balance is disrupted, such as obsessive-

compulsive disorder and schizophrenia.

Introduction

Behavioral flexibility, the ability to adapt behavioral responses to changes in environmental contexts even under identical sensory conditions, requires flexible association of sensory information with appropriate motor outputs[1,2]. To achieve rapid adaptation to rule changes, the brain must maintain stable, goal-directed actions while simultaneously updating task rules to support successful behavioral transitions according to new rules[3,4] Disruption of this coordination is a hallmark of several neurological and psychiatric disorders. For instance, patients with schizophrenia often exhibit unconstrained action patterns and impaired stability in goal-directed behaviors[5,6], whereas those with Parkinson's disease or obsessive-compulsive disorder demonstrate maladaptation to new rules, characterized by excessive perseveration[2,7-10]. This flexibility-stability dilemma also poses a significant challenge in computational learning, and modern machine-learning algorithms address this by balancing flexible updates with stable maintenance through functional modularization[11-13]. However, the neural mechanisms by which the brain negotiates this flexibility-stability trade-off during adaptive learning, enabling the acquisition of new behaviors without compromising execution stability, remain poorly understood. It is still debated whether this balance is maintained within a single network through specific representational dynamics or whether it relies on anatomically segregated networks operating in parallel.

Reversal learning tasks, which require updating behavioral responses following reversed stimulus-outcome contingencies, provide a powerful paradigm for interrogating behavioral flexibility[2,14]. While the prefrontal cortex has traditionally been considered central to behavioral flexibility through the encoding of task rules[15-17], recent evidence suggests that a broader network of brain areas dynamically adjusts representations in response to task-relevant contingencies[18-20]. The posterior parietal cortex (PPC), with its extensive

bidirectional connections to both sensory and frontal areas[21,22], has emerged as a critical node

in this process. In flexible learning tasks, the PPC encodes task-relevant choices rather than

raw sensory stimuli[23], is required for rule-based sensory categorization[24], and retains a history

of behavioral contingencies to guide future decisions[25,26]. These properties suggest that the

PPC acts as a flexible hub that can update outcome contingencies during reversal learning,

potentially through adaptive control of inter-areal connectivity and compositional coding[27,28].

However, the underlying circuit architecture that implements these principles while

simultaneously ensuring behavioral stability during rapid transitions is a fundamental

question that has yet to be addressed.

During reversal learning, the frontal cortex exhibits increased activity associated with

rule updating[29,30], while sensory cortical activity is enhanced only after full adaptation to new

contingencies[31]. While sensory representations are known to be modulated to prioritize task-

relevant stimuli[32-34], how stability and flexibility are manifested across distinct sensory areas

during adaptive perceptual performance has yet to be fully elucidated. Recent evidence

indicates that both sensory and context-dependent prior information is distributed across

multiple brain areas[35,36]. Remaining question is how such priors shape sensory

representations across the sensory hierarchy and, more importantly, whether these

representational dynamics are causally linked to flexible behavioral outcomes. In the auditory

system, both the midbrain inferior colliculus (IC) and the auditory cortex (AC) exhibit

significant response modulations driven by task demands[34,37,38] and internal behavioral

states[39-42]. Furthermore, both structures encode non-sensory information, including action-

and reward-related signals[38,43-49], suggesting that top-down influences orchestrate auditory

processing across the auditory pathway. Nevertheless, a systematic comparison of cortical

and subcortical auditory processing during the course of behavioral adaptation and how they

differentially contribute to the stability and flexibility of auditory behavior is currently

lacking.

89   In this study, we demonstrate that parallel top-down projections from the PPC to the

AC and IC enable rapid behavioral adaptation during within-session auditory Go/No-go

reversal learning. We first found that AC neurons flexibly alter their representations during

reversal learning to enhance cue discrimination, whereas IC neurons maintain stable

representations across rules. Consistent with these observations, pharmacological inactivation

of the AC impaired adaptation to reversed contingencies, while silencing the IC disrupted

auditory discriminability regardless of contingency. Using dual retrograde tracing, we

identified anatomically segregated PPC subpopulations projecting to the AC (PPC_{AC}) and IC

(PPC_{IC}). Projection-specific calcium imaging and optogenetic perturbations revealed that

PPC_{AC} neurons flexibly update stimulus-outcome associations, while PPC_{IC} neurons support

stable, goal-directed action execution during reversal learning. Finally, simulations using a

feed-forward network model indicated that this parallel architecture is a computational

necessity for rapid adaptation following contingency reversal. Together, our findings reveal a

circuit-level division of labor within the PPC that balances flexibility and stability to support

rapid and robust behavioral adaptation.

Result

Fast transitions of behavioral responses in mice performing a within-session auditory
reversal learning

To examine behavioral flexibility during rapid contingency changes, we
developed a within-session auditory reversal-learning task for head-fixed mice using a
Go/No-go paradigm (Figure 1A; Methods). Mice were initially trained under the first rule
(R1) to lick in response to a low-frequency tone (Go, 5kHz) and to withhold licking in
response to a high-frequency tone (No-go, 10kHz). Upon reaching expert-level
performance (correct response rate > 0.75 within 200 trials), stimulus-outcome
contingencies were reversed within the same session (R2; Go, 10 kHz; No-go, 5 kHz).

On the first day of reversal training, mice exhibited high levels of false alarm (FA)
rates in R2, continuing to lick in response to the previously rewarded but now incorrect
stimulus (5 kHz), along with reduced hit rates for the newly rewarded stimulus (10 kHz;
Figure 1B). With continued training, mice rapidly adjusted their licking behavior
following rule reversal, achieving accurate performance within a single session (Figure
1C). Reversal-learning efficiency was quantified as the trial-number-to-reversal (TTR),
defined as the number of trials required to reach expert-level performance in R2 (Methods).
Across training sessions, R2 performance progressively improved, accompanied by a
corresponding decrease in TTR, indicating enhanced reversal-learning efficiency over
time (Figure 1D; Figure S1A). In expert mice, the average TTR was 30.82 ± 6.25 trials
(Figure 1E). Lick reaction times were modestly increased in R2 relative to R1 (Figure 1F),
whereas overall lick rates remained stable across rules (Figure 1G).

Differential contributions of AC and IC during auditory reversal learning

To determine the causal contributions of auditory cortical and subcortical regions to rapid adaptation during auditory reversal learning, we inactivated the AC and IC using muscimol (MUS), a $GABA_A$ receptor agonist, in expert mice performing the task (Figure 1H; Figures S1B and Table S1; Methods). Bilateral AC inactivation selectively impaired auditory reversal learning, as evidenced by increased TTRs, reduced Hit rates, and elevated FA rates in R2, while leaving performance in R1 intact (Figures 1I-1L; Figures S1C, S1F and S1G). In contrast, unilateral IC inactivation disrupted auditory discrimination performance in both R1 and R2, significantly reducing hit rates across task phases, including R1 (Figures 1M and 1N; Figures S1D, S1H-S1I). Control PBS injections into either AC or IC had no effect on behavior performance (Figures 1I-1N; Figures S1C, S1D, S1F-S1I).

Given these distinct behavioral effects, we next examined how auditory representations in AC and IC evolve across task rules during reversal learning. We recorded single-unit activity from AC (n = 400 neurons) and IC (n = 179 neurons) using linear silicon probes in mice performing the auditory reversal-learning task (three mice per region, with two recorded in both regions; Figure 1O; Table S2). Behavioral performance, including correct rates, TTRs, reaction times, and lick rates, did not differ between recording groups (Figures S2A-S2E; Supplementary Table 2). Auditory-evoked activity was analyzed across three task phases: R1 (initial rule), TR (transition from R1 to R2 up to TTR), and R2 (reversed rule after TTR). To minimize motor-related confounds, analyses were restricted to trials without licking within ±1 s of stimulus onset. Neurons exhibiting significant responses to either the 5 kHz or 10 kHz tone in each phase were identified using a Wilcoxon signed-rank test (p < 0.01). The fraction of stimulus-responsive neurons

decreased during TR but was comparable between R1 and R2 in both AC and IC (Figures

1P and 1Q).

We first examined whether stimulus-responsive neurons altered their responses to

the same auditory stimuli across task phases (Figures 1R-1U). Changes in response

amplitude were quantified using the auditory gain modulation index (GMI), where

positive values indicate enhanced responses and negative values indicate suppressed

responses relative to R1. AC neurons exhibited significantly enhanced auditory responses

in R2 relative to R1 for both tones (Figures 1R and 1T), indicating an overall amplification

of auditory representations following rule reversal. Notably, these enhanced responses

emerged during the transition phase, suggesting that gain modulation develops early

during behavioral adaptation in the AC (Figures 1R and 1T). In contrast, IC neurons

exhibited stable tone-evoked responses across task phases (Figures 1S and 1U).

We next examined whether tone selectivity changed across task phases. The tone

selectivity index (SI) was quantified for each neuron using receiver operating

characteristic (ROC) analysis[23,31], based on differences in firing rates during the stimulus

epoch between the 5kHz and 10kHz tones (Figures 1V and 1W; Methods). Stimulus

selectivity was significantly enhanced in R2 exclusively in AC neurons, with no

corresponding change in IC neurons (Figure 1X). Consistently, a larger fraction of AC

neurons exhibited changes in their auditory representations across rules compared to IC

neurons (Figure 1Y, Figures S2F and S2G). Together, these results indicate that AC

neurons undergo pronounced remapping of auditory representations following rule

reversal, whereas IC neurons maintain stable auditory encoding across task rules.

Collectively, these findings reveal a functional dissociation between auditory

cortical and subcortical regions during auditory reversal learning. The AC is specifically

required for behavioral flexibility following contingency reversal and exhibits enhanced

representation of reversed Go/No-go stimuli through integration of rule information with

auditory inputs. In contrast, the IC supports stable auditory discrimination performance by

maintaining reliable auditory encoding and consistent execution of learned responses

across task phases.

PPC, projecting to AC and IC, supports auditory reversal learning

To identify candidate top-down brain regions that balance flexibility in the AC and

stability in the IC during auditory reversal learning, we performed dual-color retrograde

tracing by injecting retrograde adeno-associated viruses (AAVrg) expressing GFP and

tdTomato into the AC and IC, respectively (Figure 2A). Whole-brain clearing combined with

light-sheet microscopy revealed distinct anatomical projection patterns to each structure

(Figures 2A-2C; Supplementary Movies 1 and 2). The orbitofrontal cortex (OFC), secondary

motor cortex (M2), and cingulate cortex area 1 (Cg1) preferentially projected to the AC

(Figure 2C$_{i-ii}$), whereas the medial prefrontal cortex, including prelimbic (PrL) and

infralimbic (IL) cortices, as well as cingulate cortex area 2 (Cg2), primarily innervated the IC

(Figure 2c$_{i-ii}$; Supplementary Movie 2). The insular cortex (InC), extending along the

anterior-posterior axis, mainly projected to the IC, whereas the adjacent claustrum (Cl)

exhibited strong projections to the AC (Figure 2C$_{ii-iii}$).

In contrast to these regions, which selectively projected to either the AC or IC, both

the PPC and temporal association cortex (TeA), corresponding to dorsal and ventral

association cortices, respectively, contained two anatomically distinct and spatially

segregated neuronal populations projecting to the AC and IC (Figures 2B and 2C$_{iv}$;

Supplementary Movies 1 and 2). In particular, robust dual labeling in the PPC indicated

prominent top-down projections to both auditory areas (Figures 2B, 2D and 2E). Within the

PPC, neurons projecting to the IC (PPC$_{IC}$) accounted for 19.30 % ± 4.69% of labeled cells,

whereas neurons projecting to the AC (PPC$_{AC}$) comprised 82.03 % ± 4.61% (Figure S3).

Neurons co-labeled with both tracers were rare, constituting only 1.66 % ± 0.31% of PPC$_{AC}$

neurons and 7.85% ± 1.93% of PPC$_{IC}$ neurons. Spatially, PPC$_{AC}$ neurons were distributed

along the mediolateral axis with higher density in lateral PPC (PPC-L), whereas PPC$_{IC}$

neurons were predominantly localized to medial PPC (PPC-M) (Figures 2B, 2C$_{iv,}$ and 2D).

Depth analysis using 4',6-diamidino-2-phenylindole (DAPI)-stained sections revealed that

PPC$_{AC}$ neurons spanned superficial to deep cortical layers, whereas PPC$_{IC}$ neurons were

largely confined to deep layers (Figures 2D and 2E; Methods). These anatomical distinctions

are consistent with previously described intratelencephalic (IT) and pyramidal tract (PT)

cortical neuron subtypes[50,51].

We next examined the functional contribution of the PPC to auditory reversal learning.

Similar to the effects of AC inactivation (Figures 1I-1L; Figures S1C, S1F, and S1G),

bilateral PPC inactivation using muscimol significantly increased TTRs following the rule

switch and reduced hit rates in R2 (Figures 2F-2I; Figures S1E, S1J and S1K). Importantly,

this deficit was specific to reversal learning, as auditory discrimination performance during

R1 remained intact (Figures 2F and 2H; Figures S1E, S1J and S1K). Control PBS injections

had no effect on TTRs or task performance (Figures 2F-2I; Figures S1E, S1J and S1K).

Together, these results demonstrate that PPC activity is required for flexible adaptation to

reversed stimulus-outcome contingencies during auditory reversal learning.

Double dissociation of top-down projections from the PPC during reversal learning

Given that segregated PPC neuronal populations project to the AC and IC, we next

examined the functional roles of these two top-down pathways in auditory reversal

learning. To selectively inhibit each projection during task performance, we expressed the

inhibitory opsin archaerhodopsin (ArchT) in PPC_{AC} or PPC_{IC} neurons via bilateral

injections of AAVrg expressing ArchT under the CaMKIIa promoter into the AC or IC,

respectively (Figures 3A and 3B; Table S3). Green laser illumination was delivered to the

PPC from 1 s before stimulus onset through the response window either selectively during

the transition phase (TR; early R2 up to the TTR; Type I inactivation) or on randomly

selected trials during stable phases (R1 or R2; Type II inactivation).

Type I inactivation of PPC_{AC} neurons induced pronounced perseverative errors,

reflected by decreased Hit rates, increased FA rates, and elevated TTR values (Figures 3C-

3F), indicating a failure to update reversed stimulus-outcome contingencies. In contrast,

Type I inactivation of PPC_{IC} neurons markedly reduced licking following rule reversal and

resulted in a significant decrease in Hit rates, indicating an inability to maintain Go

responses after the contingency switch (Figures 3G-3J). Control mice expressing eYFP in

the PPC showed no behavioral effects under the same stimulation protocol (Figure S4),

confirming the specificity of these effects. By comparison, Type II inactivation of either

pathway had no detectable impact on task performance across rules (Figures 3K-3R),

indicating that neither PPC_{AC} nor PPC_{IC} activity is required once mice have fully adapted

to stable contingencies. Together, these results indicate that both PPC output pathways are

specifically engaged during contingency reversal, with PPC_{AC} supporting contingency

updating and PPC_{IC} supporting stable action execution.

Collectively, projection-specific optogenetic perturbations reveal a double

dissociation between PPC_{AC} and PPC_{IC} function during reversal learning. Inactivation of

PPC_{AC} disrupts contingency updating, leading to perseverative licking responses to

previously rewarded stimuli that have become No-go. In contrast, inactivation of PPC_{IC}

impairs the execution of Go responses, reflecting a failure to sustain goal-directed licking

behavior following the rule change.

PPC_{AC} exhibits flexible reorganization of population activity, whereas PPC_{IC} remains
251
stable during reversal learning
252

We next investigated how the two PPC output pathways differentially support

adaptive behavior during reversal learning at the population level. PPC neurons projecting to

either the AC or IC were selectively labeled by injecting retrograde AAVs expressing Cre

recombinase into each target region in Ai148 mice expressing GCaMP6f (Figures 4A and 4B;

Methods). This approach yielded 283 PPC_{AC} neurons and 111 PPC_{IC} neurons with sufficient

activity for analysis. Behavioral performance, including correct rates, TTRs, lick rates, and

reaction times across rules and trial types, did not differ between the two imaging groups

(Figures S5A-S5E; Table S4).

To visualize population activity across task phases, we applied Rastermap, a scaled k-

means clustering algorithm[52,53], to the mean activity of neurons during Hit and CR trials and

plotted cluster-ordered activity across R1, TR, and post-transition R2 (Figures 4C and 4D).

PPC_{AC} neurons exhibited more robust activity reorganization compared with PPC_{IC} neurons.

To further characterize population dynamics, we performed principal component analysis

(PCA; Methods) and projected activity trajectories for hit and CR trials from R1, TR, and R2

into a shared low-dimensional space (Figures 4E and 4F; Figure S5F). PPC_{AC} population

trajectories showed clear remapping across phases, with divergence reflecting both

contingency-dependent trial types (hit vs. CR) and task rules (R1 vs. R2; Figure 4E). In

contrast, PPC_{IC} population activity was primarily driven by hit trials and remained largely

stable, with minimal trajectory shifts following rule reversal (Figure 4F).

We next quantified population-level changes using subspace overlap analysis (Figure

4G; Methods). Both PPC_{AC} and PPC_{IC} populations exhibited strong within-rule overlap,

indicating consistent activity under stable contingencies. However, PPC_{AC} neurons showed

significantly reduced overlap across task phases (Figure 4H), consistent with dynamic

reorganization during reversal learning. By contrast, PPC_{IC} neurons maintained high subspace

overlap across all phases (Figure 4H), and across-rule overlap was significantly greater in

PPC_{IC} than in PPC_{AC} populations (Figures 4I and 4J). Together, these results demonstrate a

functional dissociation between PPC output pathways: PPC_{AC} neurons flexibly reorganize

their population activity to encode updated stimulus-outcome relationships, whereas PPC_{IC}

neurons preserve stable activity patterns during Go trials, potentially supporting reliable

execution of learned actions across rule reversals.

PPC top-down projections encode distinct task variables during the reversal transition

To determine how PPC_{AC} and PPC_{IC} populations encode task variables (TVs) during

the reversal transition (TR), when stimulus-outcome updating occurs, we fit a generalized

linear model (GLM; Methods) to single-neuron activity across phases. TVs included stimulus

identity (5kHz, 10kHz), licking behavior (bout onset and offset), and trial outcomes (hit, FA,

CR, miss; Figures 5A and 5B). Approximately half of PPC_{AC} and PPC_{IC} neurons encoded at

least one TV across phases, with PPC_{IC} neurons exhibiting overall stronger model fits (Figure

5A; Figures S6A and S6B).

Despite comparable encoding prevalence, the structure of TV representation differed

between pathways. PPC_{AC} neurons encoded a greater number of TVs per neuron and

exhibited more multiplexed co-encoding across variables, whereas PPC_{IC} neurons were more

selectively tuned to licking-related variables and lick-associated outcomes (Figures 5C and

5D). PPC_{AC} neurons showed stronger encoding of auditory stimulus identity and broadly

represented trial outcomes, including both lick-associated (hit and FA) and no-lick-associated

(CR and miss), whereas PPC_{IC} neurons preferentially encoded licking behavior and lick-

associated outcomes (hit and FA; Figures 5C-5H). Both populations increased encoding of

task-relevant outcomes during TR relative to R1 and R2, consistent with engagement during

contingency updating (Figures 5E and 5F; Figures S6D and S6E). However, PPC_{AC} neurons

showed greater instability of task-variable encoding across phases, with fewer neurons

maintaining stable representations through TR compared to PPC_{IC} neurons (Figure S6C).

Temporal analysis of GLM weights revealed further dissociation. PPC_{AC} neurons

exhibited prolonged and distributed encoding of both lick and no-lick outcomes across the

trial, whereas PPC_{IC} neurons encoded lick-related outcomes predominantly within the

response window (Figures 5I and 5J). Hit and FA encoding largely overlapped with lick

encoding in both populations but more strongly in PPC_{IC} neurons, while encoding of no-lick

outcomes showed minimal overlap with lick-related variables (Figures 5K and 5L). Notably,

PPC_{AC} neurons differentiated rewarded and unrewarded lick outcomes, with divergent

activity patterns for Hit versus FA trials, whereas PPC_{IC} neurons showed similar activity

across these conditions (Figures 5M and 5O). Sensory encoding in PPC_{AC} was largely

segregated from motor and outcome representations, indicating parallel processing of

auditory information within this pathway (Figure S6F).

Together, these results demonstrate that PPC_{AC} neurons flexibly and dynamically

encode newly relevant stimulus-outcome relationships during reversal, while PPC_{IC} neurons

maintain stable representations of licking and lick-associated outcomes, supporting consistent

action execution across changing contingencies.

A feedforward network model with parallel PPC pathways for rapid reversal learning

To examine the circuit mechanisms underlying the flexibility-stability trade-off

during reversal learning, we developed a hierarchical feedforward network model linking

sensory input to action decision. The network architecture reflects the biological hierarchy

of auditory processing, with sensory signals transmitted from the IC to the AC, and

convergent projections from both regions to a downstream motor unit that generates lick

or no-lick decisions (Figures 6A and 6B). In the model, the IC provides a stable auditory-

to-action drive, whereas the AC implements rule-dependent plasticity to update lick/no-

lick associations.

These two pathways are independently modulated by parallel PPC control modules,

PPC_{IC} and PPC_{AC}, whose signals evolve dynamically across task phases. During initial

learning (R1), PPC_{AC} contingency error signals decrease as stimulus-outcome associations

are acquired, while PPC_{IC} output stabilizes to support consistent action execution.

Following rule reversal (R2), PPC_{AC} detects a sharp increase in contingency error and

transiently gates elevated plasticity in the AC, enabling rapid overwriting of outdated

associations. In parallel, PPC_{IC} maintains stable feedback to the IC pathway, preserving

effective sensory-to-motor transformation and sustaining task engagement despite

transiently unreliable outcomes.

The functional roles of each pathway were validated through simulated circuit-

specific inactivation (Figure S7). Disabling the PPC_{AC} pathway selectively impaired

contingency updating, reducing behavioral flexibility, whereas disabling the PPC_{IC}

pathway disrupted action stability, reducing behavioral consistency. To test whether

pathway segregation is required, we simulated an integrated architecture in which a single

control module jointly regulated both plasticity and action stability (Figures 6C and 6D).

The integrated model failed to balance flexibility and stability, resulting in slower reversal

learning and reduced synaptic plasticity following contingency reversal (Figures 6B, 6D,

and 6E-6G). Together, these simulations demonstrate that segregation of PPC top-down

control into parallel pathways is necessary to resolve the fundamental dilemma between

behavioral persistence and flexibility, enabling rapid and stable adaptation following

contingency reversal.

Discussion

Behavioral flexibility requires the ability to update internal models in response to

changing rules while maintaining stable, goal-directed actions. Although prefrontal and

parietal regions have long been implicated in reversal learning and executive control[2,54,55], the

circuit mechanisms that balance flexibility and stability remain poorly defined. Here, we

identify a circuit-level organizational principle in the mouse brain in which anatomically

segregated top-down projections from the PPC to AC and IC play complementary roles

during auditory reversal learning (Figure 6A). PPC projections to AC (PPC_{AC}) flexibly

reorganize their activity to encode stimulus-outcome contingencies and support rule updating,

whereas projections to IC (PPC_{IC}) maintain stable representations related to goal-directed

licking execution. This double dissociation reveals how parallel PPC circuits coordinate

flexible updating with stable action control.

This circuit architecture suggests a general top-down motif for resolving the

flexibility-stability trade-off through parallel control. By anatomically segregating these

outputs, the PPC minimizes interference between adaptive and stable computations. In a

unified network, the rapid remapping of task rules could inadvertently disrupt the signals

required for stable motor execution, leading to premature stopping or a collapse of behavioral

drive, a phenomenon we observed during PPC_{IC} inactivation. Thus, parallel gating serves as a

hardware solution to the fundamental challenge in cognitive systems: integrating new

information without overwriting previously acquired execution policies[56]. This buffering

mechanism allows biological systems to maintain high behavioral drive (via PPC_{IC}) even

when the internal model of the environment is in a state of high uncertainty (during PPC_{AC}-

mediated remapping).

Our analysis of PPC_{AC} neural dynamics further reveals a coding strategy that

combines model-based and model-free reinforcement learning. On one hand, PPC_{AC}

populations exhibit systematic shifts in their activity patterns that correspond to the hidden

rule state, a hallmark of model-based representation[57]. This compositional coding allows for a

rapid transition of the population state-space when a rule change is detected. On the other

hand, these same neurons display a prolonged update of stimulus-outcome associations

following contingency changes, characteristic of model-free outcome-based learning[26]. This

hybrid representation suggests that PPC_{AC} does not merely switch between static maps.

Rather, it uses a model-based framework to guide rapid adaptation while simultaneously

utilizing model-free feedback to fine-tune the new contingency. This dual-coding scheme

enables the PPC to bridge the gap between abstract rule inference and the gradual acquisition

of new sensory-motor associations.

Our findings extend the functional roles of the AC and IC beyond traditional sensory

processing. While AC activity was dispensable for stable discrimination under familiar rules,

it was essential for adaptive learning following contingency changes. This result aligns with

evidence suggesting the auditory cortex is selectively recruited during the acquisition of new

sensory-motor associations[32,38]. In contrast, the IC remained necessary for behavioral

responses to cues across all task phases, consistent with its emerging role in integrating

auditory and motor-related signals to support goal-directed behavior[47,48,58,59]. Extending this

framework, our data suggest that PPC_{IC} projections convey action execution signals that

stabilize sensory-to-motor transformation, whereas PPC_{AC} projections enable flexible, rule-

dependent remapping. This division of labor may reflect an evolutionarily conserved strategy

that prioritizes reliable execution while preserving the capacity for rapid adaptation.

Moreover, imbalances within this dual-control system may lead to cognitive dysfunction.

Insufficient flexible updating or impaired stabilization could contribute to pathological

perseveration or instability, respectively, as observed in various neuropsychiatric disorders[5-

10,60-63]. Future studies in disease models will be essential to explore whether selectively

rebalancing these parallel PPC outputs can restore adaptive behavioral control.

Methods

Animals

All experimental procedures were approved by the KAIST Institutional Animal Care

and Use Committee (IACUC-18-100, IACUC-18-235). Animals were housed under

controlled environmental conditions with a 12-hour light/dark cycle (8 am - 8 pm / 8 pm - 8

am) and provided *ad libitum* access to food and water. Throughout the experimental period,

mice were individually housed at a controlled temperature (20 - 22□) and humidity (30 to

50%). We used C57BL/6 wild-type mice and B6.Cg-Igs7tm148.1(tetO-GCaMP6f, CAG-

tTA2)Hze/J mice (Ai148, Jackson Laboratory, stock no. 030328) of both male and female,

aged 2-6 months. Detailed information regarding the number of animals used in each

experimental condition is provided in the corresponding figure legends and supplementary

data tables.

Surgery

Common procedure for animal surgery

Adult mice (P45-P80) were anesthetized with 1.5-2% isoflurane in oxygen, and their

heads were fixed in a stereotaxic apparatus using ear bars. Body temperature was maintained

at 37°C using a feedback-controlled heating pad (CWE Inc.). Following hair removal, the

scalp was sterilized with 70% ethanol and povidone-iodine solution. A midline incision was

made using surgical scissors for virus injection, or a portion of the scalp was removed for

head plate implantation. After removing the periosteum above the skull, the exposed skull

surface was cleaned with 70% ethanol, povidone-iodine solution, and sterile filtered

phosphate-buffered saline.

Virus injection for dual-color retrograde tracing

For dual-color retrograde tracing, we injected ~ 0.5 µl of retrograde adeno-associated

virus (AAVrg) expressing tdTomato (pAAV-CAG-tdTomato, Cat# 59462-AAVrg, Addgene)

and eGFP (pAAV-CAG-GFP, Cat# 37825-AAVrg, Addgene) into the right AC (posterior 2.5

mm and lateral 4.2 mm from bregma; depth 0.7 mm) and the right IC (posterior 5.2 mm and

lateral 0.8 mm from bregma; depth 0.8 mm), respectively, of a wild-type mouse using a KDS

Legato 130 Syringe Pump (RWD) at a rate of 0.3nl/sec. After 14 ~ 15 days post-injection,

brain samples were collected for histological analysis following transcardial perfusion.

Head plate implantation surgery for mouse head-fixation

For head plate implantation, mice were prepared as described above, including scalp

incision and removal of connective tissues. We marked the PPC (posterior + 1.95 mm and

lateral 1.4~1.6 mm from bregma; depth 0.4 mm), AC, and IC for subsequent pharmacological

inactivation experiments, *in vivo* extracellular recordings, and optogenetic manipulations. For

lens implantation, a 1 mm × 1 mm area was demarcated on the skull 0.2 mm away from the

PPC target coordinates. The custom-designed head plate was securely attached to the skull

using a combination of small screws (Small Parts), cyanoacrylate glue (Loctite super glue

401, Henkel), and dental cement (Super-Bond C&B Kit, Sun Medical Co.). Following

surgery, mice were closely monitored for their health status, and behavioral training began

after a minimum recovery period of one week.

Surgical procedures for in vivo Ca^{2+} imaging with a miniature microscope

For circuit-specific Ca^{2+} imaging, ~0.5 µl of AAVrg-hsyn-cre (pENN.AAV.hSyn.Cre.

WPRE.hGH, Cat# 105553-AAVrg, Addgene) was injected into either the AC or IC of the

right hemisphere in Ai148 mice before head-plate mounting. The exposed brain tissue at the

injection site was covered with cured Kwik-Sil (WPI). Following a 7 – 10 day viral

incubation period, mice began water restriction for behavioral training. During training on the

auditory reversal-learning task (see Methods; Behavior), mice had full access to water until

they returned to their baseline weight, ensuring proper health before the subsequent lens

implantation surgery.

Gradient refractive index (GRIN) lens implantation was performed following

established protocols[64]. Mice received subcutaneous injections of carprofen (5 mg/kg) and

dexamethasone (5 mg/kg) at least 1 hour before surgery. Under anesthesia, a 1 mm × 1 mm

craniotomy was made 0.2-0.3 mm anterior to the target field of view (FOV: PPC$_{AC}$, AP -1.65

mm; PPC$_{IC}$, AP -1.45 mm). The craniotomy location accounted for the working distance of

the miniature microscope. A small linear incision was made in the cortical surface using a

surgical knife (Cat# 10055-12, Fine Science Tools,) at the planned lens insertion site. The

prism-attached GRIN lens (Cat# 1050-004624, Inscopix) was slowly inserted at a rate of 0.1

mm/20 s to a depth of 0.9-1.0 mm along the medio-lateral axis. The exposed brain tissue

between the GRIN lens and skull was sealed with cured Kwik-Sil, followed by Super-Bond

application to secure the lens. To protect the GRIN lens surface from debris during the

recovery period, Kwik-Sil was applied to the top of the lens. Post-operative care included

subcutaneous carprofen and dexamethasone injections for three consecutive days to manage

pain and inflammation. After a 2-week recovery, water restriction was reinstated to resume

behavioral training. Once mice regained stable performance (>0.75 Correct rate) in the

auditory reversal task, the FOV for each mouse was examined for clear detection of neuronal

activity. Following confirmation of a suitable imaging quality of FOV, a baseplate (Cat#

1050-004683, Inscopix) was attached to the GRIN lens using dental acrylic (Ortho-Jet, Lang

Dental) and Super-Bond, then secured with a baseplate cover (Cat# 1050-004639, Inscopix).

Surgical procedure for optogenetic manipulation in task-performing mice

For circuit-specific optogenetic manipulations, AAVrg-CaMKIIα-ArchT (pAAV-

CaMKIIα-ArchT-GFP, Cat# 99039-AAVrg, Addgene) was bilaterally injected into the AC or

IC of wild-type mice in the test group. Control animals were injected with AAV2-CaMKIIα-

eYFP (UNC) bilaterally. All viral injections were performed on the same day prior to head-

plate implantation. After a 7 - 10 day period for virus expression, mice began water

restriction for behavioral training. Once mice reached expert performance in the auditory

reversal-learning task (see Methods; Behavior), they were provided unrestricted water access

for at least 2 days to prepare for optic fiber implantation. Under anesthesia with 1.5 ~ 2 %

isoflurane, bilateral craniotomies (~0.5 mm diameter) were made over the PPC using either

manual drilling or a biopsy punch (Cat# 12-460-414, Fisher scientific, manufacturer). Optic

fibers (200 μm core diameter, Cat# FT200UMT, Thorlabs) with ferrules (CFLC230-10,

Thorlabs) were implanted into the PPC at a depth of 0.35mm and secured with Kwik-Sil

(WPI) and Super-Bond. Following a 2-day recovery period, water restriction was resumed for

optogenetic manipulation experiments.

Behavior assay

Hardware setup

Auditory stimuli were digitally generated using custom MATLAB code (sampling rate

192,000 Hz, 16bit) with 5 ms sinusoidal ramps at stimulus onset and offset. A speaker

(Ultrasonic Dynamic Speaker Vifa, Avisoft-Bioacoustics) was positioned 10 ~ 12cm from the

head-fixation apparatus. The sound pressure level (SPL) of both stimuli was calibrated to

75dB using a Reference Signal Generator and an externally polarized condenser microphone

(CM16/CMPA, Avisoft-Bioacoustics), positioned at the same distance as the mouse's left ear

relative to the speaker. A custom-built lickometer was used to detect individual licking events

during the task. Both outcomes (water reward and air puff punishment) were delivered

through separate solenoid valves (EV-2-24, Clippard). All hardware components were

controlled by Presentation (Neurobehavioral Systems) using a USB data acquisition board

(USB-201, Measurement Computing).

Behavior training for the auditory reversal-learning task

Before behavioral training, mice underwent initial water deprivation until their body

weight dropped below 80% of their pre-deprivation weight, typically after three days of

deprivation. Body weight was monitored daily and maintained above 75% of the initial

weight throughout the water restriction period. During behavioral experiments, mice were

kept on ~1 ml of water per day to ensure consistent performance. Task performance was

evaluated based on hit rate, False Alarm (FA) rate, and Correct rate, which were calculated as

follows:

(1) Hit rate $= \frac{\# \, of \, Hit \, trials}{\# \, of \, Go \, trials}$

514   (2) FA rate $= \dfrac{\#\ of\ FA\ trials}{\#\ of\ No-go\ trials}$

515   (3) Correct rate $= \dfrac{\#\ of\ Hit\ trials\ +\ \#\ of\ Correct\ Rejection\ trials}{\#\ of\ Go\ trials\ +\ \#\ of\ No-go\ trials}$

To assess perceptual flexibility in mice, we adapted and modified a reversal-learning task

from previous research[23,65]. The task consisted of five stages:

1. Reward conditioning stage: Water-restricted mice received 1μl of water upon licking

the water port. If mice did not initiate licking behavior within 3 minutes from the

session start, water was manually delivered through the port until voluntary licking

was established. Mice progressed to the next stage if they achieved more than 1,000

licks within 20 minutes.

2. R1 conditioning stage: Mice were trained to discriminate between Go (5kHz) and No-

go (10kHz) pure tones (500 ms duration), presented pseudo-randomly in a 1:1 ratio. A

2-second response window followed a 500 ms delay after stimulus presentation.

During Go trials, licking within the response window (hit) triggered a 3 μl water

reward and a 2-second consumption period. If mice did not lick (miss trials), water

was automatically delivered after the response window to encourage the Go stimulus-

reward association. No-go trials had no consequences with licking (FA) or not licking

(CR). The session included a 4-second inter-trial interval and terminated after 20

consecutive miss trials. Mice advanced to the next stage if they achieved a hit rate

above 0.8.

3. R1 learning stage: Mice were trained to discriminate between 5 kHz and 10 kHz tones

in a Go/No-go paradigm with reward and punishment. The paradigm was similar to

the conditioning stage, except for two critical differences: no water reward for miss

trials and negative outcomes for FA trials (4 - 8s timeout and/or 100 - 800 ms air puff).

4. Reversal-learning task with conditioning stage: Once mice reached a ≥0.75 Correct

rate in the initial rule (R1), they proceeded to the reversal-learning stage, where the

Go/No-go contingencies were reversed within a single session. To facilitate the rule

transition, classical conditioning (CS) trials were introduced immediately following

the reversal, during which a reward was delivered after Go stimuli regardless of

licking behavior. CS trials were progressively reduced in 20-trial increments once

mice achieved ≥0.75 Correct rates for both R1 and R2. When the number of CS trials

reached 40, the delay period was extended from 500 ms to 1,000 ms. Mice exhibiting

persistent cross-rule impulsive licking were returned to R1 training. Task performance

during CS trials was evaluated based on anticipatory licking behavior[65], with licks

during the delay period classified as hits or FAs depending on the stimulus

contingency. All subsequent experiments, including electrophysiology, drug

treatments (MUS and PBS), Ca^{2+} imaging and optogenetics were conducted at this

stage.

5. Reversal-learning expert stage: Mice that demonstrated a ≥0.75 Correct rate for at

least three consecutive days in the R2 stage (without CS trials) were classified as

reversal experts. Subsequent experiments, including drug treatments, Ca^{2+} imaging,

and optogenetics were conducted at this stage.

Quantification of trial-number-to-reversal (TTR) and transition (TR) phase

To quantify the speed of adaptation to reversed contingencies (R2), we defined a

metric termed trial-number-to-reversal (TTR). Following rule reversal, a 10-trial moving

average of binary choice accuracy (Correct = 1, Incorrect = 0) was computed. TTR was

defined as the final trial in the first sequence of three consecutive trials where this moving

average exceeded 0.7. The TR phase was defined as the period between the contingency

reversal and TTR (Figure 1). In sessions that included CS trials, TTR was calculated using

both CS and standard trials, with anticipatory licking responses used to assess response

accuracy.

Pharmacological inactivation

To investigate the role of specific brain regions in auditory reversal learning, we

performed local neural inactivation through bilateral microinjections of either the $GABA_A$

receptor agonist muscimol (MUS, 1 μg/μl; Sigma-Aldrich) or 1X PBS (control) into PPC, AC,

and IC. Each hemisphere received 0.23 μl of solution. Behavioral testing in the auditory

reversal task commenced 20 minutes post-injection to ensure adequate drug diffusion and

effect. To control for individual variability and enable within-subject comparisons, mice were

tested across multiple sessions in the following sequence: pre-injection baseline (no injection;

Pre), muscimol injection (MUS), and PBS injection. Fluoro-ruby (Cat# AG335, Sigma-

Aldrich) was injected with PBS to verify the injection site by histology. For specific

subgroups (AC, N=1; IC, N=3; PPC, N=2), performance measures were averaged across

sessions to generate a single data point per animal.

Optogenetic manipulations

For circuit-specific optogenetic inactivation during the reversal task, AAVrg-

CaMKIIa-ArchT-GFP was bilaterally injected into either the AC (PPC_{AC}) or IC (PPC_{IC}),

followed by bilateral optical fiber implantation in the PPC. To probe the temporal specificity

of PPC inputs during behavioral adaptation, we employed two laser stimulation protocols

(Type I and II inactivation):

1. Type I: Continuous green laser stimulation (532 nm, 10-12 mW) from 1 s before

stimulus onset to the end of the response window (trial epoch).

2. Type II (Control): Trial-epoch stimulation delivered in pseudo-randomly selected trials.

For Type I, the number of stimulated trials was set by rounding up the TTR measured during

matched laser-off sessions (up to a maximum of 80 trials) to ensure coverage of the TR phase.

Type II served as controls to confirm that observed effects were specific to the transition

stage.

In vivo electrophysiological recordings

Neural activity recording during task

Neural recordings from AC and IC in task-performing mice were conducted using a

64-channel probe (P64-12, Diagnostic Biochips. Inc.). A 1 mm diameter craniotomy was

made one day prior to recording and covered with an artificial dura-like material (Duragel,

Cambridge Neurotech) to protect the exposed brain tissue. For each animal, 3 to 5 recording

tracks were made in each target area. Neural data were acquired using an OpenEphys data

acquisition board and software at 30kHz, with an amplifier gain of 192 for each channel.

Signals were bandpass filtered with cutoff frequencies at 300 Hz and 6000 Hz. To confirm the

recording sites and perform post hoc histological analysis, electrodes were coated with

lipophilic dye (1,1'- dioctadecyl-3,3,303'- tetramethylindocarbocyanine perchlorate (DiI),

Invitrogen) before insertion.

Spike sorting for single-unit isolation

Spike sorting was performed using an automated spike-sorting MATLAB toolbox

Kilosort to cluster units from raw data[66]. The resulting spike clusters were manually curated

using Phy (Kwik team). Further quality control was conducted with custom MATLAB codes

to ensure high-quality single-unit data. A total of 579 single units were recorded from AC and

IC in 4 animals. To plot firing rate changes in representative neurons, peri-stimulus time

histograms (PSTHs) of spike trains for 5kHz and 10kHz stimuli in R1 and R2 were generated

separately by averaging trial-by-trial firing rates in 10-ms bins and smoothing with a

Gaussian kernel (σ = 50 ms).

Gain modulation index

For each stimulus-responsive neuron (Wilcoxon signed-rank test, p < 0.01), the

change in firing rate during the stimulus period (0 to 500 ms) relative to the pre-stimulus

baseline (-1500 to -500 ms) was computed on a trial-averaged basis, yielding Δ firing rates.

For each stimulus frequency (5 kHz and 10 kHz, analyzed separately), the gain modulation

index (GMI) was calculated by comparing response magnitudes between task phases (R2 or

TR vs. R1), normalized by the maximum firing rate across these phases using a soft

normalization parameter ($\alpha = 0.01$).

$$(4)\ Gain\ modulation\ index = \frac{\Delta FR_{\mathrm{R2(or\ TR)}} - \Delta FR_{\mathrm{R1}}}{\Delta FR_{\mathrm{R2(or\ TR)}} + \Delta FR_{\mathrm{R1}} + \alpha}$$

Selectivity index

Neuronal selectivity for stimulus identity was quantified using receiver operating characteristic (ROC) analysis, as previously described[23,67]. Analyses were performed separately for R1 and R2 by comparing time-averaged firing-rate changes evoked by the 5kHz and 10kHz stimuli across trials. To capture both increased and decreased responses, firing-rate changes (Δfiring rates) were calculated for each trial as the difference between activity during the stimulus period (0 to 500 ms) and the pre-stimulus baseline (-1500 to -500 ms). To minimize contamination from motor-related activity, trials with licking events occurring near the stimulus period (-500 to +750 ms relative to stimulus onset) were excluded from analysis. The stimulus selectivity index (SI) was then computed for each neuron from the area under the ROC curve (AUC) as follows:

(5) SI = (AUC - 0.5) x 2

An SI value of +1 indicates consistently greater firing-rate changes in response to the 5kHz tone than the 10kHz tone, whereas an SI value of -1 indicates a preference for the 10kHz tone. Statistical significance of SI was assessed using a permutation test across trials (1,000 iterations; $p < 0.01$). To quantify changes in stimulus discriminability across task phases, we calculated ΔSI as follows:

(6) $\Delta\ SI = |SI_{TR\ or\ R2}| - |SI_{R1}|$

Neurons that did not exhibit a significant SI in any task phase were excluded from this analysis.

Outcome selectivity was quantified using the same ROC-based approach by

comparing hit and CR trials during the response window (1500 to 3500 ms). In this case,

an SI value of +1 indicates consistently greater firing-rate changes during hit trials,

whereas −1 indicates greater responses during CR trials.

Circuit-specific Ca^{2+} imaging and data analysis

Data acquisition and processing

We performed *in vivo* Ca^{2+} imaging using a miniature fluorescence microscope and

the nVista 3.0 data acquisition system (Inscopix)[68,69] with LED power of 0.2~ 0.4 mW and a

gain value of 1 ~ 4, collecting data at 20 frames per second.

Data processing was performed using Inscopix Data Processing Software (IDPS). The

preprocessing pipeline included spatial down-sampling (by 4), spatial filtering, and motion

correction to mitigate motion artifacts. We used Constrained Non-negative Matrix

Factorization for micro-endoscopic data (CNMF-E) to identify regions of interest (ROIs)

corresponding to individual cells. ROI detection was configured with a cell diameter between

5 to 7 pixels and standard thresholds for minimum pixel correlation (0.60 to 0.80) and

minimum peak-to-noise ratio (8.00 to 10.00) to ensure Ca^2 signal detection while minimizing

false positives. After automated ROI detection, we manually curated the results to match

fluorescence traces ($\Delta F/F_0$) with ROIs and eliminate contamination from out-of-focus

neuropil fluorescence. To correct for fluorophore photo-bleaching, we applied a correction

method from a previously published protocol[70], using the following equation:

$(7)\ F_{corrected(t)} = F_{subt(t)} - G_{(t)} + <F_{subt(t)}>_t$

This equation calculated the corrected fluorescence signal ($F_{corrected(t)}$) by subtracting

slow baseline fluctuations averaged over a 300-second sliding window ($G_{(t)}$) and adding back

the overall mean fluorescence across the recording session ($<F_{sub(t)}>_t$).

Rastermap visualization

Out of 798 recorded neurons (594 PPC_{AC} neurons and 194 PPC_{IC} neurons), we

selected neurons with an average $\Delta F/F_0$ greater than 0.05 (283 PPC_{AC} neurons and 111 PPC_{IC}

neurons). To visualize task-related Ca^{2+} activity, we applied Rastermap[52,53], a k-means

clustering-based sorting algorithm that highlights temporally correlated patterns in neural

population activity. Among the tunable parameters, we selected a cluster number of 50 for

sorting population activity. Importantly, choosing alternative values (e.g., 25 clusters) did not

alter the overall patterns or conclusions of our observations (Data not shown).

Principal Component Analysis (PCA)

Ca^{2+} activity was averaged for each trial type (hit and CR) and task phase (R1, TR,

and R2), then concatenated to form a $6T \times N$ matrix, where T represents the duration of each

trial segment (2 second pre-stimulus onset and 6 seconds post-stimulus onset) and N is the

number of imaged neurons. We then performed principal component analysis (PCA) using

pca function in Matlab library). Neural trajectories were computed by projecting the data onto

the first three principal components (PCs).

Subspace overlap analysis

Normalized Ca^{2+} traces were trial-averaged in 50-trial segments S_i (i = 1, ... 7) across

R1 (S1-S3; 1 to 150 trials), TR (S4; R2 to TTR + 15 trials) and R2 (S5-S8; 1 to 150 trials post

S4). We then performed PCA using *scikit-learn* (Python library) to compute low-dimensional

neural subspaces for each segment. To quantify subspace overlap, we calculated the cosine of

the principal angle between each pair of subspaces, as previously published[71]. The principal

angle (θ_p) between two subspaces *U* and *V* corresponds to the largest angle between any two

pairs of vectors in *U* and *V*, providing a measure of their alignment. For each segment S_i, we

extracted the first N principal components and organized them into an N × N matrix **R** (S_i),

defining the subspace of that segment. The cosine of the principal angle between subspaces S_i

and S_j was computed using the method described by Knyazev and Argentati[72], as follows:

(8) $cos\theta_p(s_i, s_i) = \sigma_{max}(R(s_i)^T R(s_j))$

where $\sigma_{max}(A)$ denotes the largest singular value of matrix A.

Generalized linear models (GLM)

We used generalized linear models (GLM) to regress recorded Ca^{2+} signals against

task event variables[70,73,74]. To capture contingency-dependent neural dynamics, GLMs were

trained on specific task phase (R1, TR or R2) rather than on the full session. Ca^{2+} responses

for each neuron were modeled as a linear sum of task components and events, including lick-

bout onset and offset, Go stimulus (10kHz) and No-go stimulus (5kHz), hit, FA, CR, and

miss.

The neural response at time t ($\hat{y}_t$) was modeled by the sum of the bias term (β_0) and

the weighted sum of various binary predictors (x_i) as follows:

$$(9)\ \hat{y}_t = \beta_0 + \sum_{i=0}^{15} \beta_i^{Go} x_i^{Go} + \sum_{i=0}^{15} \beta_i^{Nogo} x_i^{Nogo} + \sum_{i=0}^{60} \beta_i^{Hit} x_i^{Hit} + \sum_{i=0}^{60} \beta_i^{FA} x_i^{FA}$$

$$+ \sum_{i=0}^{60} \beta_i^{Miss} x_i^{Miss} + \sum_{i=0}^{60} \beta_i^{CR} x_i^{CR} + \sum_{i=-5}^{10} \beta_i^{lon} x_i^{lon} + \sum_{i=-5}^{10} \beta_i^{loff} x_i^{loff}$$

Predictors for each task variable were convolved with temporal filters to capture the

slow dynamics of Ca^{2+} signals[75]. Predictors for Go (x_i^{Go}) and No-go (x_i^{Nogo}) stimuli indicated

stimulus presence, while lick predictors represented onset (x_i^{lon}) and offset (x_i^{loff}) of lick

bouts, defined as sequences with inter-lick intervals of less than 2 seconds. Predictors for

behavioral outcomes, including hit (x_i^{Hit}), FA (x_i^{FA}), CR (x_i^{CR}), miss (x_i^{Miss}), included time

lags to cover the duration of stimulus presentation, reward consumption, and slow Ca^{2+}

dynamics. Time lags were set to 0-1.5 second for stimulus predictors, 0-6 seconds for

outcome predictors, and -0.5 to 1 s for lick-related predictors, where 0 denotes the stimulus

(for stimulus and outcome) or lick onset time. The full model contained up to 321 coefficients,

including the bias term (β_0).

To avoid overfitting, we used a bottom-up model selection approach. We first fit

single-variable GLMs for each neuron using *ElasticNet* linear regression (scikit-learn, Python

library), which combines L1(lasso) and L2 (ridge) regularization. Model performance was

evaluated using twenty-fold cross-validation, with 30% of trials held out for validation. For

each neuron, the task variable (TV) whose model produced the highest mean cross-validated

R^2 score significantly greater than 0 (Student's t-test) was selected as the initial predictor.

Neurons for which no TV reached significance were classified as not fitted by the GLM.

Additional TVs were then incorporated iteratively. At each step, we compared model fits with

and without the added TV and retained the new TV only if it significantly improved model

performance (two-sample Kolmogorov–Smirnov test). This process was repeated until no

further improvement was detected. The ElasticNet regularization parameter α was optimized

across a broad range (1×10^{-5} to 10) to maximize model performance.

To analyze patterns of neuronal multiplexing (co-encoding of multiple TVs), we

identified all unique combinations of co-encoded variables and ranked multiplexing patterns

by their frequency across neurons. For neurons encoding more than two TVs, all pairwise

combinations were counted separately. For example, a neuron encoding "Hit," "FA," and

"CR" contributed one count each to the pairs "Hit-FA", "Hit-CR", and "FA-CR". This

approach enabled us to quantify how frequently each pair of TVs was co-encoded across the

population. Co-encoding relationships were visualized using a customized circular Sankey

diagram (Matlab File Exchange, ID: 128679)[76]. In these diagrams, curved links connect pairs

of co-encoded variables, and the thickness of each link is proportional to the frequency of co-

occurrence.

Histology and imaging

Fluorescence imaging of brain sections

Histological experiments were conducted to verify anatomical circuit tracings,

confirm targeting sites after pharmacological inactivation, identify electrode tracks following

in vivo recordings, and validate optogenetic manipulation and Ca^{2+} imaging sites. Mice were

deeply anesthetized with isoflurane and intraperitoneal injection of avertin (2,2,2,-

Tribromoethanol; 125-250 mg/kg; Sigma-Aldrich), and transcardially perfused with 10ml of

phosphate-buffered saline (PBS) followed by 4% paraformaldehyde (wt/vol in 1X PBS).

Brain samples were carefully extracted and post-fixed for 4 hours at room temperature, then

washed 3 times in 1X PBS for 10 minutes each. The samples were transferred to a filtered 30%

sucrose solution (wt/vol in 1X PBS) and allowed to sink completely for 1~2 days at 4°C.

Brain samples were embedded in the optimal cutting temperature (OCT) compound (Tissue-

Tek, Sakura Finetek), rapidly frozen at -80°C, and coronally sectioned at 30 μm (for tracing)

or 40 μm (for site verification) using a cryostat (Leica). Sections were washed three times in

1X PBS for 10 minutes to remove residual OCT medium. The sections were mounted on

glass slides with a mounting medium containing 4',6-diamidino-2-phenylindole (DAPI,

Vector Labs) for nuclear staining and tissue visualization. Fluorescent images were acquired

using a slide scanner (Axio Scan Z1, Carl Zeiss).

Tracing analysis

For dual-retrograde tracing analysis, three coronal brain sections (30 μm thick) were

collected from each of three mice. Sections were selected based on stereotaxic coordinates

from the Paxinos atlas (AP -1.70 mm, -1.84 mm, and -2.06 mm from the bregma). Using

IMARIS software (Oxford Instruments), brain section images were aligned to the reference

atlas, and anatomical boundaries were matched to delineate the PPC, including its medial

(PPC-M) and lateral (PPC-L) subdivisions. Labeled cells within these boundaries were

detected using the IMARIS Cell Detection function, with DAPI colocalization used to

confirm accurate cell identification. The spatial coordinates (x, y, z) of verified cells were

exported for custom MATLAB-based analysis. To assess layer-specific distributions, atlas

boundaries were imported into IMARIS to guide selection of DAPI signals corresponding to

pia and ventricular surfaces. Cortical depth was normalized by regression fitting of DAPI

signals along these boundaries, with the pia-to-ventricle distance scaled to a depth of 1. This

normalization allowed direct comparison of relative laminar distributions between PPC_{AC} and

PPC_{IC} populations.

Whole-brain clearing and 3D imaging

Whole-brain clearing was performed using the Binaree Tissue-Clearing Kit (HRTC-

012 and HRMO-006, Binaree, Inc.) following a previously published protocol[77]. After

perfusion and brain extraction, the tissue-clearing process proceeded as follows: The brain

was incubated in a starting solution at 4°C for 24 hours. The brain was then transferred to

tissue-clearing solutions and incubated at 37°C with 50 rpm agitation for 48 hours. This

clearing step was repeated to achieve complete tissue transparency. The brain was rinsed

three times in a washing solution, then incubated in a mounting and storage solution at 37°C

for 24 hours. For imaging, a Lightsheet 7 fluorescence microscope (Carl Zeiss) was used to

acquire z-stack images of the cleared brain. The resulting images were reconstructed in 3D

using IMARIS software.

Model Simulation

Neural network configuration

To simulate an auditory Go/No-go task with within-session rule reversal, we

implemented a hierarchical feed-forward network consisting of sensory (input, IC, AC),

modulatory (PPC_{AC}, PPC_{IC}) and action output modules (Figure 6). Auditory tone information

was propagated from the IC to the AC and integrated to generate action logits corresponding

to lick versus no-lick responses. Two parallel modulatory modules controlled the gain of the

IC and AC layers. Specifically, PPC_{AC} monitored outcome contingency statistics (hit, miss,

FA and CR), computed errors as deviations from the expected contingencies, and generated a

signal that promoted adaptive updating in the AC. PPC_{IC} computed a goal-directed action

signal (the difference between the probabilities of lick and no-lick responses) and modulated

IC gain to stabilize action execution.

Reversal learning

Agents were trained on a Go/No-Go task with a within-session reversal of stimulus-

action contingencies, identical to the behavioral experiment. Actions were sampled from a

softmax policy, and scalar rewards were assigned according to the task rule: r = +1 for licking

on Go trials and r = −1 for licking on No-go trials. Parameters were optimized using a

standard policy-gradient objective, $L = -E[(r - b) \log \pi(a|x)] - \beta H(\pi)$, which included

an entropy bonus ($H(\pi)$: categorical entropy with $\beta = 1 \times 10^{-3}$) to encourage exploration. The

reward baseline b was computed as an exponential moving average of rewards, updated as b

$\leftarrow \lambda b + (1-\lambda)E[r]$ with $\lambda = 0.95$.

Flexible-stable synaptic weight updates

Synaptic weights were updated using a flexibility-weighted meta-update rule[56,78], in

which each parameter's gradient update was gated by its assigned flexibility. Stable pathways

(input–IC, IC–readout) were assigned low flexibility, whereas IC–AC and AC–readout

pathways were assigned higher flexibility, enabling selective adaptation following rule

reversal. Low-flexibility synapses were anchored to their initial values, thereby suppressing

updates as parameters deviated from this anchor.

Specifically, for each synapse, the deviation from the anchor weight was defined as

$\Delta = p - w_0$, and parameters were updated according to

(10) $p \leftarrow p - \eta \, S(\Delta, f) \, g, \; S(\Delta, f) = 1 - \tanh^2 \left(\alpha \frac{1-f}{f} \Delta \right),$

where g denotes the gradient of the loss, $f \in [0,1]$ represents synaptic flexibility,

with $\alpha = 4$ controlling the sharpness of the gating function, chosen to provide a clear

separation between flexible and stable update regimes. Under this rule, high-flexibility

synapses $(f \approx 1)$ undergo near-standard gradient-based updates, whereas low-flexibility

synapses $(f \ll 1)$ rapidly suppress updates as they deviate from the anchor, effectively

preserving previously learned synaptic structure.

Agent performance and inactivation conditions

Circuit-specific inactivation (AC, IC, PPC_{AC} and PPC_{IC}) was simulated by

multiplicative suppression of activity or flexibility signals in the corresponding modules,

matching the experimental manipulations. Behavioral performance was quantified by

stimulus-conditioned lick probabilities, estimated from held-out trials, for hit and FA

responses.

Alternative network architectures for comparison

To evaluate the functional advantage of the parallel circuit motif, we implemented an

integrated PPC model and two control networks with uniform synaptic properties. In the

integrated PPC model, a single population integrated both goal-directed and contingency-

related errors into a global population error signal. Crucially, this global error signal was

jointly coupled to both stability and flexibility modulation, preventing the independent

control characteristic of the parallel model. All simulation scripts are available at

https://github.com/seungheelee1789/PPC_RL_Jung.

Statistical analysis

Data analysis was conducted using Microsoft Excel, custom MATLAB (MathWorks)

and Python scripts. All data are presented as means ± standard error of the mean (SEM)

unless otherwise specified. 'N' denotes the number of mice, while 'n' represents the number of

sessions or neurons. Statistical differences were evaluated through repeated measures two-

way ANOVA, Wilcoxon signed-rank test (non-parametric, paired datasets), Paired t-test

(parametric, paired datasets), Wilcoxon rank-sum test (non-parametric, unpaired datasets),

two-sided chi-square test (non-parametric, frequency counts), Mood's median test (non-

parametric, unpaired datasets), Sign-test (non-parametric, zero median of single population),

and Two-sample Kolmogorov-Smirnov test. All the statistical significances are indicated as *,

, *, **** for $p < 0.05$, $p < 0.01$, $p < 0.001$, and $p < 0.0001$, respectively.

Acknowledgments

This work has been supported by grants to S.H.L. from the Institute for Basic Science (IBS-

R002-A2) and to S.P. from the National Research Foundation of Korea (RS-2002-NR070602).

Author Contributions

E.J. and S.L. conceptualized and initiated the project. E.J., J.L., and S.L. designed

experiments and wrote the manuscript. E.J. and J.L. jointly performed data analysis and

visualization. E.J. performed behavioral training, histology, Ca^{2+} imaging, optogenetic

experiments, and some *in vivo* recordings. J.L. established *in vivo* electrophysiology

recordings and implemented subspace overlap and GLM analyses. W.C. designed the

computational model. S.P. provided advice and conceptual guidance. G.R. conducted some *in*

vivo recordings and behavioral training. G.K. performed brain clearing and cell detection.

Data and code availability

Source data are provided with this paper. All the representative data and code that support the

findings are publicly available on GitHub: https://github.com/

seungheelee1789/PPC_RL_Jung. Further requests for the data generated in this study can be

directed to the corresponding author (shlee1@kaist.ac.kr).

Ethics declarations

Competing interests

The authors declare no competing financial interests.

Figure 1. Distinct functional roles of AC and IC in rapid reversal learning

A. Schematic of the auditory Go/No-go reversal-learning task.

B-C. Example lick raster plots from a naïve (**B**) and expert (**C**) mouse during reversal learning.
Left, 5kHz trials; right, 10kHz trials. Each black dot represents a single lick. Blue and
magenta vertical bands denote R1 and R2 blocks, respectively. Vertical black lines denote
sound onset (0 s); horizontal black lines indicate the rule-reversal point; green line and arrow
mark the TTRs; cyan dots indicate lick trials; blue dots indicate no-lick trials. Naïve mice
exhibited licking after passive water delivery during R2 (**B**), whereas expert mice initiated
licking in response to the 10kHz stimulus following reversal (**C**).

D. Scatter plot of TTR vs. R2 correct rates across training sessions. Sessions for each mouse
were divided into six equal training segments, averaged within each segment, and then

averaged across mice (one circle per segment). Red dashed line indicates linear regression.
Pearson's correlation shows improved performance with training.

E. Average TTRs for naïve (189.82 ± 19.04) and expert (30.82 ± 6.25) mice, demonstrating
faster adaptation after training.

F. Lick reaction times for hit and FA trials. Hit reaction times were significantly longer in R2
than in R1 (p = 0.0275), whereas FA reaction times did not differ. Wilcoxon rank-sum test.

G. Lick rates for hit and FA trials, showing no significant differences across rules.

H. Schematic of MUS or PBS injections into bilateral AC or unilateral IC.

I-L. Behavioral effects of AC inactivation (N=6).

I. Average hit, FA, and correct rates during Pre (no injection; open bars), MUS (red bars), and
PBS (black bars) sessions. Repeated measures two-way ANOVA; hit rate (drug, p = 7.88 x
10^{-7}; rule, p = 3.53 x 10^{-5}; interaction, p = 6.69 x 10^{-6}); FA rate (drug, p = 0.566; rule, p =
0.409; interaction, p = 0.850); correct rate (drug, p = 1.04 x 10^{-5}; rule, p = 9.3 x 10^{-5};
interaction, p = 4.83 x 10^{-6}). Inset shows bilateral AC injection sites aligned to the Paxinos
atlas. Scale bar, 500 μm.

J. Session-averaged performance with AC inactivation. Hit, FA, and correct rates are shown
for MUS (red) and PBS (black). Shading indicates ± SEM. Grayscale bars denote p-values
for significant differences between MUS and PBS (Wilcoxon signed-rank test).

K. Average TTRs for Pre, MUS, and PBS sessions (mean ± SEM). Pre (13.42 ± 5.87) vs.
MUS (148.08 ± 23.38), p = 0.002; MUS vs. Post (13.08 ± 5.70), p = 0.004; paired t-test.

L. Scatter plot of TTRs vs. R2 correct rate across sessions. Each circle represents one session
(open, Pre; red, MUS; black, PBS). Crosses denote group means ± SEM.

M-N. Same as **I-J**, but for IC inactivation (N=7). Repeated measures two-way ANOVA; hit
rate (drug, p = 6.21 x 10^{-11}; rule, p = 0.005; interaction, p = 0.067); FA rate (drug, p = 0.566;
rule, p = 0.409; interaction, p = 0.850); correct rate (drug, p = 2.99 x 10^{-6}; rule, p = 0.044;
interaction, p = 0.164).

O. DiI-labeled brain sections showing multi-channel electrode tracks in the AC (top) and IC
(bottom). AuD, dorsal secondary auditory cortex; Au1, primary auditory cortex; AuV, ventral
secondary auditory cortex; DCIC, dorsal cortex of the inferior colliculus; CIC, central
nucleus of the inferior colliculus; ECIC, external cortex of the inferior colliculus. Scale bar,
500 μm.

P. Fraction of AC neurons (n=400) responsive to auditory stimuli during the stimulus epoch
and task outcomes during the response window across R1, TR, and R2 phases. Black bars
indicate neurons responsive across phases. Light blue (5kHz) and pink (10kHz) bars indicate

stimulus-responsive neurons within each phase. Light gray bars indicate non-responsive
neurons. Right, fraction of neurons responsive to 5 kHz only (light blue), 10 kHz only (pink),
or both stimuli (light yellow) across task rules. Neurons were classified as responsive if they
exhibited a significant stimulus-evoked response in at least one phase. Wilcoxon signed-rank
test, $p < 0.01$.

Q. Same as **P**, but for IC neurons (n=179).

R. Left, example PSTHs from an AC neuron (#48) illustrating the gain modulation index
(GMI) for the 5 kHz stimulus. Solid lines, R1; dotted line, R2; dark gray shading, stimulus
epoch. Right, violin plots of population GMI values for 5 kHz-responsive AC neurons during
TR and R2. Horizontal lines indicate medians across neurons responsive in at least one phase.
Median GMI_{5kHz}: TR = 0.055 (p = 0.044), R2 = 0.126 (p = 0.001); one-sample Wilcoxon
signed-rank test with zeros.

S. Same as **R,** but for IC neurons (#119). Median GMI_{5kHz}: TR = -0.032, R2 = 0.004 (both
n.s.). Between-group comparison: TR, p = 0.048.; R2, p = 0.016; Mood's median test.

T-U. Same as **R-S**, but for GMI_{10kHz}. (**t**) AC median GMI_{10kHz}: TR = 0.118 (p < 0.0001), R2 =
0.128 (p < 0.0001). (**u**) IC median GMI_{10kHz}: TR = -0.022, R2 = -0.019 (both n.s.). Between-
group comparison: TR, p = 0.001; R2, p = 0.0005.

V. Schematic illustrating selectivity index (SI) calculation using ROC analysis.

W. Example PSTHs from an AC neuron (#194) showing stimulus selectivity (5kHz versus
10kHz) during R1 (top) and R2 (bottom). SI values are indicated above each PSTH.

X. Violin plots of changes in absolute stimulus selectivity (Δ |SI|) during R2 (|R2| - |R1|).
Median Δ |SI|: AC = 0.056 (p = 0.026), IC = -0.003 (p = 0.542); Wilcoxon signed-rank test
with zeros. Between-group comparison: p = 0.011; Mood's median test.

Y. Fraction of stable and flexible stimulus-selective neurons in AC and IC. Stable: AC =
0.180 ± 0.002, IC = 0.239 ± 0.002, $p < 6.38 \times 10^{-8}$; Flexible: AC = 0.261 ± 0.003, IC = 0.241
± 0.004, p = 0.001; Wilcoxon rank-sum test.

Figure 2. PPC sends segregated top-down projections to AC and IC and is required for auditory reversal learning.

A-E. Dual-color retrograde tracing reveals anatomically segregated PPC projections to the AC (green) and IC (red).

A. Coronal view of a cleared whole brain showing retrograde injection sites in the AC (AAVrg-CAG-GFP, left) and IC (AAVrg-CAG-tdTomato, right), aligned to the Paxinos atlas (gray lines). Scale bars, 500 μm.

B. Dorsal view of a cleared brain imaged with light-sheet microscopy and reconstructed in 3D. White outlines indicate the right cortical hemisphere and PPC. GFP-labeled PPC$_{AC}$ neurons are positioned laterally, whereas tdTomato-labeled PPC$_{IC}$ neurons are located medially. Scale bar, 1 mm.

C. Coronal sections from cleared brains imaged by light-sheet microscopy and reconstructed in 3D. DAPI (blue) indicates brain architecture; AC-projecting (green) and IC-projecting (red) neurons are visualized across cortical regions. Images aligned to the Paxinos atlas (gray lines). Scale bars, 1mm.

C$_i$, OFC (orbitofrontal cortex; medial, ventral, and lateral parts); Prl (prelimbic cortex); M2 (secondary motor cortex).

C$_{ii}$, M1 (primary motor cortex); M2; S1 (primary somatosensory cortex); S2 (secondary somatosensory cortex); Cg1 (cingulate cortex area 1); Cg2 (cingulate cortex area 2); InC (insular cortex); Cl (claustrum).

C$_{iii}$, M1; M2; S1; S2; InC.

C$_{iv}$, PPC (medial and lateral); S1; AC; TeA (temporal association cortex).

D. Representative coronal sections along the anterior-posterior (AP) axis of the PPC showing segregated PPC$_{AC}$ (green) and PPC$_{IC}$ (red) neuronal populations. Images are aligned to the Paxinos atlas (gray lines). Scale bars, 500 μm.

E. Laminar and spatial distribution of retrogradely labeled neurons in medial PPC (PPC-M, left) and lateral PPC (PPC-L, right). Neuronal fractions were binned by normalized cortical depth (pia = 0, ventricle = 1) using DAPI signals. Green, PPC$_{AC}$ neurons; red, PPC$_{IC}$ neurons; yellow, dual-labeled neurons.

F-I. PPC inactivation impairs auditory reversal learning.

F. Behavioral effects of PPC inactivation (N=7). Average hit, FA, and correct rates during Pre (no injection; open bars), MUS (red), and PBS (black) sessions. Repeated-measures two-way ANOVA; hit rate: drug, $p = 3.84 \times 10^{-9}$; rule, $p = 1.83 \times 10^{-7}$; interaction, $p = 3.45 \times 10^{-8}$; FA rate: drug, $p = 0.666$; rule, $p = 0.423$; interaction, $p = 0.050$; Correct rate: drug, $p = 0.0001$; rule, $p = 0.001$; interaction, $p = 0.016$. Inset, bilateral PPC injection sites labeled with fluoro-ruby and aligned to the Paxinos atlas. Scale bars, 500 μm.

G. Average TTRs for Pre (28.92 ± 9.08), MUS (133.21 ± 32.37), and PBS (42.29 ± 12.55) sessions (mean ± SEM). Paired t-test: Pre vs. MUS, $p = 0.019$; MUS vs. Post; $p = 0.034$.

H. Session-averaged performance with or without PPC inactivation (N=7). hit (top); FA (middle), correct (bottom) rates are shown for MUS (red) and PBS (black) sessions. Shaded

997 regions indicate ± SEM. Gray-scaled bars indicate p-values for significant differences
998 between MUS and PBS sessions; Wilcoxon signed-rank tests.

999 **I.** Scatter plot of TTRs vs. R2 correct rate across individual sessions. Circles, individual
1000 sessions (open, Pre; red, MUS; black, PBS). Crosses denote group means ± SEM.

1001

Figure 3. Double dissociation of PPC top-down projections during auditory reversal learning

A-B. Schematics of projection-specific optogenetic inactivation targeting PPC$_{AC}$ neurons (**A**) and PPC$_{IC}$ neurons (**B**)

C-F. Impaired reversal learning induced by Type I PPC$_{AC}$ inactivation (N=14 mice). Top, Type I inactivation protocol. Green laser light (532 nm) was delivered immediately after rule reversal, from 1 s before stimulus onset to the end of the response window.

C. Time course of hit (top), FA (middle), and correct (bottom) rates aligned to the start of R2 (trial 0). Green, light-on sessions; gray, light-off sessions; shading, ± SEM. Gray-scaled bars indicate p-values comparing light-off and light-on sessions; Wilcoxon signed-rank test.

D. TTRs were significantly increased during light-on (66.79 ± 10.98) compared to light-off (31.29 ± 5.37) sessions; p = 0.009, Wilcoxon signed-rank test.

E. Changes in performance (Δ = light-on - light-off) for hit (R1, -0.015 ± 0.011; R2, -0.148 ± 0.057; p = 0.042), FA (R1 = -0.027 ± 0.032; R2 = 0.096 ± 0.040; p = 0.007), and correct rates (R1, 0.005 ± 0.019; R2, -0.167 ± 0.052; p = 0.001). Wilcoxon signed-rank test.

F. Scatter plot of TTR vs. R2 correct rates across sessions. Filled green circles, light-on; open green circles, light-off. Crosses show group means ± SEM (dark-green, light-on; gray, light-

off).

G-J. Same as **C-D**, but for Type I PPC$_{IC}$ inactivation (N=12).

H. TTR was significantly increased in light-on (104.08 ± 14.95) compared to light-off
sessions (25.08 ± 5.94); p = 0.0005, Wilcoxon signed-rank test.

I. Change in performance for hit (R1, -0.022 ± 0.015; R2, -0.388 ± 0.077; p = 0.0005), FA
(R1 = 0.032 ± 0.045; R2 = -0.069 ± 0.049; p = 0.003), and correct rates (R1, -0.027 ± 0.027;
R2, -0.160 ± 0.038; p = 0.0005). Wilcoxon signed-rank test.

K-N. Same as **C-F** but for Type II PPC$_{AC}$ inactivation (N=6). Top, Type II inactivation
protocol. Laser light was delivered randomly in 50% of trials across all task phases, from 1 s
before stimulus onset to the end of the response window. Note that mice showed intact
reversal learning.

L. TTRs did not differ between light-on (29.33 ± 5.58) and light-off (23.00 ± 6.84) sessions.

M. Change in performance rates for hit (R1, 0.003 ± 0.003; R2, 0.010 ± 0.018; n.s.), FA (R1
= -0.036 ± 0.037; R2 = 0.0.35 ± 0.030; n.s.), and Correct rates (R1, 0.019 ± 0.019; R2, -0.012
± 0.014; n.s.).

O-R. Same as **C-F,** but for Type II PPC$_{IC}$ inactivation (N=6).

P. TTR did not differ between light-on (9.40 ± 2.29) and light-off (12.40 ± 1.74) sessions.

Q. Change in performance for hit (R1, -0.035 ± 0.035; R2, -0.010 ± 0.026; n.s.), FA (R1 =
0.105 ± 0.085; R2 = 0.253 ± 0.138; n.s.), and correct rates (R1, -0.072 ± 0.057; R2, -0.131 ±
0.074; n.s.).

1040

1041 **Figure 4. Distinct population dynamics of PPC_AC and PPC_IC neurons during auditory**
1042 **reversal learning**

1043

A. Projection-specific Ca²⁺ imaging of PPC_AC neurons. Top left, schematic of the imaging

strategy. Top right, representative field of view (FOV) in the PPC. Bottom left, dorsal view
showing all FOV locations. Bottom right, example Ca^{2+} trace from a single neuron during
task performance. Vertical blue and magenta lines indicate 5kHz and 10kHz stimulus onset,
respectively. Scale bar, 1mm.

B. Same as **A**, but for PPC_{IC} neurons.

C. Rastermap visualization (50 clusters) of trial-averaged, z-scored Ca^{2+} activity from PPC_{AC}
neurons (n = 283) during concatenated hit and CR trials across R1 (left), TR (middle), and R2
(right). Neurons were clustered based on activity during R1.

D. Same as **C**, but for PPC_{IC} neurons (n=111).

E-F. Principal component trajectories of trial-averaged population activity from PPC_{AC} (**E**)
and PPC_{IC} (**F**) neurons, aligned to stimulus onset (-2 to +6 s) and projected onto the first three
PCs. Colors indicate trial type and task phase, as shown in the legend.

G. Schematic of the subspace-overlap analysis, computed from angular differences between
PC axes.

H. Subspace-overlap heatmaps for PPC_{AC} (left) and PPC_{IC} (right) populations across task
phases. Color scale: 0 (no overlap) to 1 (complete overlap).

I. Left, difference in subspace overlap (PPC_{AC} - PPC_{IC}), showing greater overlap in PPC_{IC}
neurons, particularly during TR. Right, statistical comparison of overlap values between
PPC_{AC} and PPC_{IC} populations across 20-fold cross-validation (Wilcoxon rank-sum test).

J. Average subspace overlaps (mean ± SEM) within and across task phases for PPC_{AC} (black
bars; within, 0.601 ± 0.036; across, 0.419 ± 0.028) and PPC_{IC} neurons (open bars; within,
0.799 ± 0.019; across, 0.756 ± 0.014). PPC_{AC} neurons exhibited a significantly greater
reduction in overlap across phases compared to within phases (p = 0.0072), whereas PPC_{IC}
neurons showed no within-group difference (Wilcoxon rank-sum test). Between-group
comparisons revealed significantly lower overlaps in PPC_{AC} than PPC_{IC} populations both
within (p = 0.0022) and across phases (p = 3.39 x 10^{-6}; Wilcoxon rank-sum test).

1071

1072

Figure 5. Distinct and multiplexed task-variable encoding by PPC$_{AC}$ and PPC$_{IC}$ neurons

during reversal learning.

1075

A. Left, schematic of the GLM used to quantify encoding of task variables (TVs) by
individual neurons. Right, cumulative distribution of GLM-fitted R^2 values during the TR for
PPC$_{AC}$ (black) and PPC$_{IC}$ (gray) neurons.

B. Example Ca^{2+} traces with corresponding GLM fits and TV weights from representative
PPC$_{AC}$ (top) and PPC$_{IC}$ (bottom) neurons. Solid lines indicate measured Ca^{2+} activity; dotted
lines indicate GLM-predicted activity. Vertical blue and magenta lines indicate onset of the
5kHz and 10kHz stimuli, respectively. TV weights are color-coded as in **A**.

C. Fraction of neurons encoding each TV during TR. Filled bars, PPC$_{AC}$; open bars, PPC$_{IC}$.
Two-sided chi-square tests; 10kHz, $p = 0.004$; 5kHz, $p = 0.026$; CR, $p < 0.0001$; miss, $p =$
0.022; Lick$_{on}$, $p = 0.044$; hit, FA and Lick$_{off}$, not significant.

D. Left, fraction of PPC$_{AC}$ (top) and PPC$_{IC}$ (bottom) neurons grouped by the number of TVs
encoded. Right, circular Sankey diagrams illustrating multiplexed TV co-encoding in PPC$_{AC}$
(top) and PPC$_{IC}$ (bottom) populations. Line thickness indicates frequency of each
combination.

E-F. Time-resolved fractions of neurons encoding each TV across the trial for PPC$_{AC}$ (**E**) and
PPC$_{IC}$ (**F**) neurons. Left, stimulus- and outcome-encoding from 0 to 7 s relative to stimulus
onset. Dark and light gray shading denote stimulus and response epochs, respectively. Right,
fractions of neurons encoding motor variables (lick-bout onset and offset, aligned to 0 s).
Task variables are color-coded as in **a**.

G. Average GLM weights for PPC$_{AC}$ neurons encoding 10kHz (Go, magenta), 5kHz (No-go,
blue), Lick$_{on}$ (black), and Lick$_{off}$ (gray) variables. Positive (top) and negative (bottom)
weights are shown separately; shading indicates ± SEM. Gray-scaled bars indicate

significance (one-sample Wilcoxon signed-rank test).

H. Same as **G**, but for PPC_{IC} neurons. When fewer than five neurons encoded a given
variable, individual traces are plotted as dotted lines.

I-J. Average GLM weights for neurons encoding outcome variables (hit, FA, CR, and miss)
in PPC_{AC} (**I**) and PPC_{IC} (**J**).

K-L. Proportional Euler diagrams showing overlap among PPC_{AC} (**K**) and PPC_{IC} (**L**) neurons
encoding hit, FA, and lick variables (left), and CR, miss, and lick variables (right).

M-N. PCA trajectories of hit - and FA-encoding PPC_{AC} (**M**) and PPC_{IC} (**N**) neurons during
hit and FA trials. Black dots denote stimulus onset; colored dots indicate response-window
onset.

O. Quantification of trajectory separation using subspace-overlap analysis. Wilcoxon signed-
rank test, $p = 6.796 \times 10^{-8}$.

1110

1111

1112 **Figure 6. Computational dissociation of flexibility and stability *via* parallel PPC circuits**

1113

A. Schematic of the parallel PPC top-down model. Auditory input is processed by the IC
layer and relayed to the AC layer. Activity in both layers converges to drive motor output
(lick or no-lick). In this architecture, the PPC$_{AC}$ module regulates contingency-dependent
plasticity in AC, whereas the PPC$_{IC}$ module stabilizes auditory-to-lick output in the IC
through goal-directed signals.

B. Evolution of lick probabilities for the two auditory stimuli across learning in the parallel
top-down circuit. The vertical dashed red line indicates the rule-reversal point.

C. Schematic of an integrated PPC model, in which a single PPC module jointly monitors
contingency and goal errors to regulate both AC plasticity and IC gain.

D. Same as **B**, but for the integrated top-down circuit.

E. Learning curves showing task accuracy for the parallel (black) and integrated (gray) networks. The parallel network exhibits faster recovery following reversal, indicating more rapid adaptation.

F. Comparison of behavioral performance between parallel and integrated networks during R1 (iterations 1-100) and R2 (iterations 300-400). Wilcoxon rank-sum test; R1 hit rate (p = 0.0003); R1 FA rate (p = 0.0003); R2 hit rate (p < 0.00001); R2 FA rate (p < 0.00001).

G. TTR was significantly higher in the integrated network (83.300 ± 2.264) than in the parallel network (416.700 ± 22.263); p < 0.00001, Wilcoxon rank-sum test.

 Extended Data Figures

 Supplementary Figure 1. Task performance during within-session auditory reversal
 learning and effects of MUS inactivation

1137

A. Performance changes in mice during auditory reversal learning. Top, Correct rates across
training stages in mice used for single-unit recordings (N=3, blue) and MUS experiments
(N=13, gray). Left, auditory discrimination (R1 only) with a 500 ms delay of the response
window from the stimulus offset; middle, reversal learning with a 500 ms delay; right,
reversal learning with a 1000 ms delay. Red horizontal lines indicate the expert-level
performance threshold (Correct rate ≥ 0.75). Bottom, Correct rates for R1 and R2 trials
during reversal learning with 500 ms (left) and 1000 ms (right) delays.

B. Baseline performance prior to MUS inactivation. Mice reached Correct rates > 0.75 for

both rules across three consecutive days before MUS injections into AC, IC, or PPC.

C. Lick reaction times (Top) and lick rates (Bottom) for hit and FA trials during AC
inactivation. Circles, individual sessions; open bars, Pre; red bars, MUS; black bars, PBS.
Wilcoxon rank-sum test; R2 hit lick rates, Pre vs. MUS (p = 0.034); Mus vs. PBS (p = 0.024).

D. Same as **C**, but for IC inactivation sessions. Wilcoxon rank-sum test; reaction times of R1
hit trials, MUS vs. PBS (p = 0.053); lick rates of R1 hit trials, Pre vs. MUS (p = 0.008); MUS
vs. PBS (p = 0.010); lick rates of R2 hit trials, Pre vs. MUS (p = 0.004); MUS vs. PBS (p =
0.013).

E. Same as **C**, but for PPC inactivation sessions. Wilcoxon rank-sum test; R2 hit lick rates,
MUS vs. PBS (p = 0.019).

F. Example lick rasters from AC inactivation sessions. Left, MUS; right, PBS.

G. Session-averaged lick histograms for hit and FA trials during Pre (left), MUS (middle),
and PBS (right) sessions targeting bilateral AC. Top bars, gray-scaled p-values relative to
MUS sessions (Wilcoxon signed-rank test).

H-I. Same as **F-G**, but for IC inactivation sessions.

J-K. Same as **F-G**, but for PPC inactivation sessions.

1162

1163 Supplementary Figure 2. Behavioral performance during single unit recordings in AC
1164 and IC

1165

A. Average hit, FA, and Correct rates in mice recorded from AC (N=3, n=8 sessions; black
bars) and IC (N=3, n=9 sessions; open bars). No significant group differences (two-way
ANOVA without repetition).

B. Average TTRs for AC-recorded (24.37 ± 6.95) and IC-recorded (42.66 ± 10.16) sessions.
Circles represent individual sessions.

C. Scatter plot of TTRs vs. R2 Correct rate (AC = 0.87 ± 0.04; IC = 0.93 ± 0.01). Circles,
individual sessions (black, AC; empty, IC); crosses, group means ± SEM.

D-E. Lick reaction times (**D**) and lick rates (**E**) for hit and FA trials. Circles, sessions; black
bars, AC-recording sessions; open bars, IC-recording sessions. No significant group
differences (Wilcoxon rank-sum test).

F-G. Left, pie charts showing fractions of neurons in each SI category in AC (**F**) and IC (**G**).
Right, scatter plots of SI in R1 versus SI in R2 for individual neurons in AC (**F**) and IC (**G**).
Inset, color key for six neuron categories defined by rule-dependent changes in SI: stimulus-
selective (green), contingency-selective (blue), lost selectivity (yellow), acquired selectivity
(red), non-selective (dark gray), and non-responsive (light gray).

Supplementary Figure 3. Quantification of projection-specific PPC neurons

1182

1183 Proportions (%) of PPC$_{AC}$, PPC$_{IC}$, and dual-projecting neurons across individual brain
1184 samples (S1-S3, N=3) and group average (bottom).

1185

1186 **Supplementary Figure 4. Control experiment for optogenetic manipulations of PPC$_{AC}$**
1187 **and PPC$_{IC}$ circuits during auditory reversal learning**

1188

A-D. Control green-light (532 nm) stimulation of PPC during Type I optogenetic inhibition
experiments.

B. Δ Performance (light-on - light-off) for hit (R1, 0.006 ± 0.009; R2, -0.002 ± 0.026; n.s.),
FA (R1 = 0.102 ± 0.057; R2 = 0.021 ± 0.043; n.s.), and Correct rates (R1, -0.049 ± 0.032; R2,
-0.010 ± 0.032; n.s.). Wilcoxon signed-rank test.

C. TTRs during light-off (24.00 ± 3.89) and light-on (18.83 ± 2.77) sessions; p = 0.009.
Wilcoxon signed-rank test.

D. Scatter plot of TTRs vs. R2 Correct rates. Green-open circles, light-off sessions; green-
filled circles, light-on sessions; Crosses indicate group means ± SEM (gray, light-off; dark
green, light-on)

1199

Supplementary Figure 5. Behavioral performance during *in vivo* imaging of PPC$_{AC}$ and PPC$_{IC}$ neurons and robustness of neural population analyses

A. Average hit, FA, and Correct rates for PPC$_{AC}$ (n=6, black bars) and PPC$_{IC}$ (n=12, open bars) imaging groups. No group differences (two-way ANOVA without repetition).

B. Average TTRs for PPC$_{AC}$ (34.50 ± 9.02) and PPC$_{IC}$ (24.58 ± 5.12). Circles, individual sessions.

C. Scatter plot of TTRs vs. R2 Correct rates across all imaging sessions (PPC$_{AC}$, 0.76 ± 0.01; PPC$_{IC}$, 0.77 ± 0.02). Circles, imaging sessions (black, PPC$_{AC}$; empty, PPC$_{IC}$); crosses, group means ± SEM (black, PPC$_{AC}$; gray, PPC$_{IC}$).

D-E. Lick reaction times **(D)** and lick rates **(E)** for hit and FA trials. Circles, individual sessions; black bars, PPC$_{AC}$; open bars, PPC$_{IC}$. No significant group differences (Wilcoxon rank-sum test).

F. Explained variance of Ca^{2+} population activity for PPC$_{AC}$ (black) and PPC$_{IC}$ (gray) neurons across task phases (same dataset as **Figures 4E** and **4F**). The top 20 PCs capture ~80% of total variance, validating the use of 20 PCs.

 Supplementary Figure 6. Temporal dynamics of task variable encoding in projection-
 specific PPC populations across rules

1219

A. Left, cumulative distribution of GLM-fitted R^2 values across rules. Blue line, R1; black
line, TR; magenta line, R2. Right, fraction of neurons encoding each task variable across
rules. Blue bar, R1; black bar, TR; magenta bar, R2. Noted that similar proportions of neurons
were fit across phases.

B. Same as **A**, but for PPC$_{IC}$ neurons.

C. Neurons grouped by the number of TVs encoded during TR (box outlines); filled colors
indicate stability of TV encoding across rules. PPC$_{AC}$ exhibited lower pattern stability (0.57)
than PPC$_{IC}$ (0.40), indicating more stable TV encoding in PPC$_{IC}$ across rule changes.

D. Left, time-resolved fractions of PPC$_{AC}$ neurons encoding each task variable (color-coded)
during R1 (top) and R2 (bottom), aligned to stimulus onset. Right, fractions encoding lick-
bout onset (black) and offset (gray), aligned to their respective motor events.

E. Same as **D**, but for PPC$_{IC}$ neurons.

F. Left, Euler diagrams of PPC$_{AC}$ neurons encoding 10kHz and 5kHz stimuli. Right,
comparison of co-encoding fractions among 10kHz-encoding (magenta outline), 5kHz-

encoding (blue outline), and all TV-encoding (no outline) PPC_{AC} neurons. No significant

difference (two-sided chi-square test).

1236

1237

 Supplementary Figure 7. Inactivation simulations support a parallel top-down control
 model

A. Left, lick probability across learning iterations under simulated AC inactivation across task
rules. Right, average hit and FA rates under control conditions (intact AC; open bars) and AC
inactivation (red bars) across simulations (N = 20). Wilcoxon rank-sum test; R1 hit rate (n.s.);
R1 FA rate (n.s.); R2 hit rate ($p < 0.00001$); R2 FA rate ($p < 0.00001$).

B. Same as **A,** but for IC inactivation. Wilcoxon rank-sum tests; R1 hit rate ($p < 0.00001$); R1
FA rate (n.s.); R2 hit rate ($p < 0.00001$); R2 FA rate ($p < 0.00001$).

C. Left, lick probability across learning iterations under simulated PPC_{AC} pathway
inactivation following contingency reversal. Right, average hit and FA rates under control
(circuit on; open bars) and PPC_{AC} inactivation (circuit off; green bars); N = 20 simulations.
Wilcoxon rank-sum test; R1 hit rate (n.s.); R1 FA rate (n.s.); R2 hit rate ($p < 0.00001$); R2 FA
rate ($p < 0.00001$).

D. Same as **C**, but for PPC_{IC} pathway inactivation. Wilcoxon rank-sum test; R1 hit rate (n.s.);
R1 FA rate (n.s.); R2 hit rate ($p < 0.00001$); R2 FA rate ($p = 0.026$).

1254

1255

 <u>Supplementary Movies</u>

 Supplementary movie 1. 3D view of a whole mouse brain injected with retrograde
 tracers in AC and IC

 360-degree rotation of a cleared whole mouse brain around the mediolateral (ML) axis, with
 a fixed anteroposterior (AP) position.

 Supplementary movie 2. Mapping neurons projecting to AC and IC in whole mouse
 brains

 Series of coronal sections from five cleared whole mouse brain samples, progressing from
 posterior to anterior regions. The central sample corresponds to the one shown in **Fig. 2** and
 Supplementary movie 1, aligned to the Paxinos brain atlas. Scale bars: 1 mm. Note that
 images appear left-right reversed due to microscope orientation. Green fluorescence indicates
 AC-projecting neurons, while red fluorescence marks IC-projecting neurons. Key anatomical
 structures include: central nucleus of inferior colliculus (CIC); external cortex of inferior
 colliculus (ECIC); dorsal cortex of inferior colliculus (DCIC); auditory cortex including
 primary (A1), dorsal (AuD), and ventral (AuV) regions; primary (S1) and secondary (S2)
 somatosensory cortices; primary (M1) and secondary (M2) motor cortices; cingulate cortex
 areas 1 and 2 (Cg1, Cg2); insular cortex (InC); claustrum (Cl); infralimbic cortex (IL);
 prelimbic cortex (PrL); and orbitofrontal cortex (OFC) with medial (m), ventral (v), and
 lateral (l) subdivisions.

 Supplementary Tables

 Supplementary Data Table 1. Behavior performance rates and TTRs of mice used for
 MUS experiments

Target	# of mice	Session	R1						R2							
			Hit rate		FA rate		Correct rate		Hit rate		FA rate		Correct rate		TTR	
			Mean	±	Mean	±	Mean	±	Mean	±	Mean	±	Mean	±	Mean	±
AC	6	Pre	0.94	0.03	0.17	0.04	0.88	0.03	0.93	0.03	0.18	0.06	0.87	0.04	13.42	5.87
		MUS	0.92	0.04	0.15	0.01	0.88	0.03	0.26	0.13	0.29	0.07	0.49	0.04	148.08	23.38
		PBS	0.98	0.01	0.14	0.04	0.92	0.02	0.94	0.04	0.21	0.08	0.86	0.04	13.08	5.70
IC	7	Pre	0.88	0.05	0.23	0.05	0.83	0.04	0.87	0.04	0.32	0.06	0.77	0.03	16.71	6.03
		MUS	0.40	0.13	0.13	0.04	0.64	0.05	0.03	0.01	0.06	0.04	0.48	0.02	187.29	12.71
		PBS	0.93	0.03	0.21	0.03	0.86	0.02	0.76	0.12	0.30	0.05	0.73	0.05	29.50	10.42
PPC	7	Pre	0.99	0.01	0.13	0.03	0.93	0.01	0.96	0.01	0.32	0.08	0.82	0.04	28.93	9.08
		MUS	0.94	0.03	0.35	0.10	0.80	0.06	0.31	0.34	0.20	0.03	0.55	0.04	133.21	32.37
		PBS	0.96	0.02	0.24	0.03	0.86	0.01	0.90	0.05	0.32	0.09	0.79	0.05	42.29	12.55

 Supplementary Data Table 2. Behavior performance rates and TTRs of mice used for
 electrophysiology

Target	mouse #	# of neurons	R#1			R#2			TTR
			Hit rate	FA rate	Correct rate	Hit rate	FA rate	Correct rate	
AC	602	40	1.00	0.11	0.95	0.95	0.06	0.95	21
		42	0.66	0.16	0.75	1.00	0.03	0.99	41
		56	0.68	0.01	0.84	0.91	0.02	0.95	6
	605	37	0.96	0.16	0.90	0.97	0.01	0.98	16
		89	0.93	0.17	0.88	0.80	0.31	0.75	66
	606	64	0.42	0.01	0.71	0.77	0.11	0.83	18
		37	0.91	0.18	0.86	0.86	0.15	0.86	14
		35	0.69	0.25	0.72	0.97	0.07	0.95	13
AC total	N=3	400	0.78 ± 0.03	0.17 ± 0.04	0.83 ± 0.03	0.90 ± 0.03	0.10 ± 0.04	0.90 ± 0.03	24.38 ± 6.95
IC	602	22	0.93	0.26	0.84	0.95	0.42	0.77	45
		58	0.83	0.41	0.71	1.00	0.14	0.93	63
		13	0.97	0.41	0.78	0.98	0.12	0.93	22
		9	0.96	0.07	0.95	0.96	0.05	0.96	51
	606	20	1.00	0.22	0.89	0.85	0.14	0.86	22
		23	1.00	0.19	0.91	0.96	0.03	0.97	5
		24	1.00	0.24	0.88	0.97	0.04	0.97	22
	609	10	0.86	0.44	0.71	0.96	0.10	0.93	88
IC total	N=3	179	0.94 ± 0.02	0.28 ± 0.05	0.83 ± 0.03	0.95 ± 0.02	0.13 ± 0.04	0.91 ± 0.02	39.75 ± 9.61

1289 **Supplementary Data Table 3. Behavior performance rates and TTRs of mice used for**

1290 **optogenetic manipulations**

Group	# of mice	Type	Condition	R1 Hit rate Mean	±	R1 FA rate Mean	±	R1 Correct rate Mean	±	R2 Hit rate Mean	±	R2 FA rate Mean	±	R2 Correct rate Mean	±	TTR Mean	±
PPC$_{AC}$	14	Type I	Light on	0.96	0.02	0.22	0.05	0.87	0.02	0.74	0.06	0.37	0.04	0.68	0.03	66.79	10.98
			Light off	0.98	0.01	0.25	0.05	0.86	0.02	0.88	0.03	0.28	0.05	0.80	0.02	31.29	5.37
	6	Type II	Light on	1.00	0.04	0.12	0.04	0.94	0.02	0.92	0.04	0.34	0.08	0.79	0.03	23.00	6.84
			Light off	1.00	0.00	0.15	0.05	0.92	0.02	0.90	0.05	0.31	0.06	0.80	0.02		
PPC$_{IC}$	14	Type I	Light on	0.95	0.02	0.24	0.05	0.85	0.03	0.51	0.08	0.21	0.04	0.65	0.04	104.08	14.95
			Light off	0.97	0.01	0.22	0.04	0.88	0.02	0.91	0.03	0.29	0.03	0.81	0.02	24.57	5.09
	5	Type II	Light on	0.95	0.04	0.40	0.11	0.77	0.07	0.94	0.02	0.46	0.15	0.74	0.08	12.40	2.14
			Light off	0.98	0.01	0.30	0.07	0.84	0.04	0.95	0.03	0.21	0.08	0.87	0.04		
PPC$_{ctrl}$	6	Type I	Light on	0.98	0.01	0.29	0.07	0.85	0.04	0.93	0.03	0.29	0.05	0.82	0.04	18.83	2.77
			Light off	0.98	0.02	0.19	0.06	0.89	0.03	0.93	0.03	0.27	0.05	0.83	0.03	24.00	3.89

1291

1292

1293 **Supplementary Data Table 4. Behavior performance rates and TTRs of mice used for**

1294 **Ca^{2+} imaging**

Group	# of neurons	R1 Hit rate	R1 FA rate	R1 Correct rate	TR Hit rate	TR FA rate	TR Correct rate	R2 Hit rate	R2 FA rate	R2 Correct rate	TTR
PPC$_{AC}$	149	0.99	0.50	0.75	0.77	0.92	0.42	0.97	0.39	0.79	27
	180	0.98	0.18	0.90	0.60	0.50	0.55	0.98	0.48	0.75	12
	92	1.00	0.27	0.87	0.85	0.82	0.50	0.84	0.18	0.83	43
	54	0.80	0.38	0.71	0.57	0.58	0.49	0.74	0.18	0.78	74
	72	0.97	0.66	0.66	1.00	0.81	0.58	0.98	0.29	0.85	32
	47	0.98	0.22	0.88	0.70	0.88	0.44	0.92	0.28	0.82	19
PPC$_{AC}$ total	594	0.95 ± 0.03	0.37 ± 0.08	0.79 ± 0.04	0.75 ± 0.07	0.75 ± 0.07	0.50 ± 0.02	0.90 ± 0.04	0.30 ± 0.05	0.80 ± 0.01	34.5 ± 9.02
PPC$_{IC}$	57	1.00	0.38	0.81	0.80	0.75	0.50	0.87	0.14	0.86	23
	37	1.00	0.78	0.61	0.73	0.82	0.45	1.00	0.22	0.89	23
	5	1.00	0.39	0.81	1.00	0.67	0.60	0.98	0.19	0.90	6
	4	0.99	0.09	0.95	0.83	0.67	0.58	0.95	0.29	0.83	13
	1	1.00	0.18	0.91	0.83	0.75	0.54	0.92	0.32	0.80	25
	13	0.61	0.65	0.48	0.50	0.64	0.43	0.89	0.51	0.69	29
	14	0.76	0.24	0.76	1.00	0.00	1.00	0.82	0.20	0.81	5
	10	0.94	0.41	0.77	0.89	0.71	0.59	0.78	0.07	0.85	57
	27	0.99	0.34	0.83	0.50	0.70	0.39	0.79	0.50	0.65	19
	11	0.35	0.17	0.59	0.36	0.30	0.52	0.76	0.10	0.83	22
	10	0.88	0.08	0.90	0.67	0.60	0.55	0.82	0.18	0.82	12
	5	0.93	0.43	0.75	0.17	0.23	0.47	0.75	0.22	0.76	61
PPC$_{IC}$ total	194	0.87 ± 0.06	0.34 ± 0.06	0.76 ± 0.04	0.69 ± 0.07	0.57 ± 0.07	0.55 ± 0.05	0.86 ± 0.02	0.25 ± 0.04	0.81 ± 0.02	24.58 ± 5.12

1295

1296

1297

Reference

1. Miller, E.K., and Cohen, J.D. (2001). An integrative theory of prefrontal cortex function. Annu Rev Neurosci *24*, 167-202. DOI 10.1146/annurev.neuro.24.1.167.
2. Uddin, L.Q. (2021). Cognitive and behavioural flexibility: neural mechanisms and clinical considerations. Nat Rev Neurosci *22*, 167-179. 10.1038/s41583-021-00428-w.
3. Perich, M.G., Gallego, J.A., and Miller, L.E. (2018). A Neural Population Mechanism for Rapid Learning. Neuron *100*, 964-976 e967. 10.1016/j.neuron.2018.09.030.
4. Smith, M.A., Ghazizadeh, A., and Shadmehr, R. (2006). Interacting adaptive processes with different timescales underlie short-term motor learning. PLoS Biol *4*, e179. 10.1371/journal.pbio.0040179.
5. Strauss, G.P., Morra, L.F., Sullivan, S.K., and Gold, J.M. (2015). The role of low cognitive effort and negative symptoms in neuropsychological impairment in schizophrenia. Neuropsychology *29*, 282-291. 10.1037/neu0000113.
6. Gold, J.M., Strauss, G.P., Waltz, J.A., Robinson, B.M., Brown, J.K., and Frank, M.J. (2013). Negative symptoms of schizophrenia are associated with abnormal effort-cost computations. Biol Psychiatry *74*, 130-136. 10.1016/j.biopsych.2012.12.022.
7. Swainson, R., Rogers, R.D., Sahakian, B.J., Summers, B.A., Polkey, C.E., and Robbins, T.W. (2000). Probabilistic learning and reversal deficits in patients with Parkinson's disease or frontal or temporal lobe lesions: possible adverse effects of dopaminergic medication. Neuropsychologia *38*, 596-612. 10.1016/s0028-3932(99)00103-7.
8. Valerius, G., Lumpp, A., Kuelz, A.-K., Freyer, T., and Voderholzer, U. (2008). Reversal Learning as a Neuropsychological Indicator for the Neuropathology of Obsessive Compulsive Disorder? A Behavioral Study. The Journal of Neuropsychiatry and Clinical Neurosciences *20*, 210-218. 10.1176/jnp.2008.20.2.210.
9. Remijnse, P.L., Nielen, M.M., van Balkom, A.J., Cath, D.C., van Oppen, P., Uylings, H.B., and Veltman, D.J. (2006). Reduced orbitofrontal-striatal activity on a reversal learning task in obsessive-compulsive disorder. Arch Gen Psychiatry *63*, 1225-1236. 10.1001/archpsyc.63.11.1225.
10. Apergis-Schoute, A., and Samanth Ip, H.Y. (2020). Reversal Learning in Obsessive Compulsive Disorder: Uncertainty, Punishment, Serotonin and Perseveration. Biological Psychiatry *87*, S125-S126. 10.1016/j.biopsych.2020.02.339.
11. Ditzler, G., Roveri, M., Alippi, C., and Polikar, R. (2015). Learning in nonstationary environments: A survey. IEEE Computational intelligence magazine *10*, 12-25.
12. Merel, J., Botvinick, M., and Wayne, G. (2019). Hierarchical motor control in mammals and machines. Nat Commun *10*, 5489. 10.1038/s41467-019-13239-6.
13. Sandbrink, K., and Summerfield, C. (2024). Modelling cognitive flexibility with deep neural networks. Curr Opin Behav Sci *57*, 101361. 10.1016/j.cobeha.2024.101361.
14. Izquierdo, A., Brigman, J.L., Radke, A.K., Rudebeck, P.H., and Holmes, A. (2017). The neural basis of reversal learning: An updated perspective. Neuroscience *345*, 12-26. 10.1016/j.neuroscience.2016.03.021.
15. Thorpe, S.J., Rolls, E.T., and Maddison, S. (1983). The orbitofrontal cortex: neuronal activity in the behaving monkey. Exp Brain Res *49*, 93-115. 10.1007/BF00235545.
16. Hornak, J., O'Doherty, J., Bramham, J., Rolls, E.T., Morris, R.G., Bullock, P.R., and Polkey, C.E. (2004). Reward-related reversal learning after surgical excisions in orbito-frontal or dorsolateral prefrontal cortex in humans. J Cogn Neurosci *16*, 463-478. 10.1162/089892904322926791.

17. Kim, C., Johnson, N.F., Cilles, S.E., and Gold, B.T. (2011). Common and distinct
mechanisms of cognitive flexibility in prefrontal cortex. J Neurosci *31*, 4771-4779.
10.1523/JNEUROSCI.5923-10.2011.
18. Guo, L., Ponvert, N.D., and Jaramillo, S. (2017). The role of sensory cortex in
behavioral flexibility. Neuroscience *345*, 3-11. 10.1016/j.neuroscience.2016.03.067.
19. Chereau, R., Bawa, T., Fodoulian, L., Carleton, A., Pages, S., and Holtmaat, A. (2020).
Dynamic perceptual feature selectivity in primary somatosensory cortex upon reversal
learning. Nat Commun *11*, 3245. 10.1038/s41467-020-17005-x.
20. Okazawa, G., and Kiani, R. (2023). Neural Mechanisms That Make Perceptual
Decisions Flexible. Annu Rev Physiol *85*, 191-215. 10.1146/annurev-physiol-031722-
024731.
21. Zingg, B., Hintiryan, H., Gou, L., Song, M.Y., Bay, M., Bienkowski, M.S., Foster,
N.N., Yamashita, S., Bowman, I., Toga, A.W., and Dong, H.W. (2014). Neural
networks of the mouse neocortex. Cell *156*, 1096-1111. 10.1016/j.cell.2014.02.023.
22. Lewis, J.W., and Van Essen, D.C. (2000). Corticocortical connections of visual,
sensorimotor, and multimodal processing areas in the parietal lobe of the macaque
monkey. J Comp Neurol *428*, 112-137. 10.1002/1096-
9861(20001204)428:1<112::aid-cne8>3.0.co;2-9.
23. Pho, G.N., Goard, M.J., Woodson, J., Crawford, B., and Sur, M. (2018). Task-
dependent representations of stimulus and choice in mouse parietal cortex. Nature
Communications *9*. ARTN 259610.1038/s41467-018-05012-y.

24. Zhong, L., Zhang, Y., Duan, C.A., Deng, J., Pan, J., and Xu, N.L. (2019). Causal
contributions of parietal cortex to perceptual decision-making during stimulus
categorization. Nat Neurosci *22*, 963-973. 10.1038/s41593-019-0383-6.
25. Akrami, A., Kopec, C.D., Diamond, M.E., and Brody, C.D. (2018). Posterior parietal
cortex represents sensory history and mediates its effects on behaviour. Nature *554*,
368-372. 10.1038/nature25510.
26. Hwang, E.J., Dahlen, J.E., Mukundan, M., and Komiyama, T. (2017). History-based
action selection bias in posterior parietal cortex. Nat Commun *8*, 1242.
10.1038/s41467-017-01356-z.
27. Cole, M.W., Reynolds, J.R., Power, J.D., Repovs, G., Anticevic, A., and Braver, T.S.
(2013). Multi-task connectivity reveals flexible hubs for adaptive task control. Nat
Neurosci *16*, 1348-1355. 10.1038/nn.3470.
28. Ito, T., Yang, G.R., Laurent, P., Schultz, D.H., and Cole, M.W. (2022). Constructing
neural network models from brain data reveals representational transformations linked
to adaptive behavior. Nat Commun *13*, 673. 10.1038/s41467-022-28323-7.
29. Hattori, R., Hedrick, N.G., Jain, A., Chen, S., You, H., Hattori, M., Choi, J.H., Lim,
B.K., Yasuda, R., and Komiyama, T. (2023). Meta-reinforcement learning via
orbitofrontal cortex. Nat Neurosci *26*, 2182-2191. 10.1038/s41593-023-01485-3.
30. Wang, B.A., Veismann, M., Banerjee, A., and Pleger, B. (2023). Human orbitofrontal
cortex signals decision outcomes to sensory cortex during behavioral adaptations. Nat
Commun *14*, 3552. 10.1038/s41467-023-38671-7.
31. Banerjee, A., Parente, G., Teutsch, J., Lewis, C., Voigt, F.F., and Helmchen, F. (2020).
Value-guided remapping of sensory cortex by lateral orbitofrontal cortex. Nature *585*,
245-250. 10.1038/s41586-020-2704-z.
32. Jaramillo, S., Borges, K., and Zador, A.M. (2014). Auditory thalamus and auditory
cortex are equally modulated by context during flexible categorization of sounds. J
Neurosci *34*, 5291-5301. 10.1523/JNEUROSCI.4888-13.2014.
33. Birman, D., and Gardner, J.L. (2019). A flexible readout mechanism of human

sensory representations. Nat Commun *10*, 3500. 10.1038/s41467-019-11448-7.

34. Xin, Y., Zhong, L., Zhang, Y., Zhou, T., Pan, J., and Xu, N.L. (2019). Sensory-to-Category Transformation via Dynamic Reorganization of Ensemble Structures in Mouse Auditory Cortex. Neuron *103*, 909-921 e906. 10.1016/j.neuron.2019.06.004.

35. International Brain, L., Angelaki, D., Benson, B., Benson, J., Birman, D., Bonacchi, N., Bougrova, K., Bruijns, S.A., Carandini, M., Catarino, J.A., et al. (2025). A brain-wide map of neural activity during complex behaviour. Nature *645*, 177-191. 10.1038/s41586-025-09235-0.

36. Findling, C., Hubert, F., International Brain, L., Acerbi, L., Benson, B., Benson, J., Birman, D., Bonacchi, N., Buchanan, E.K., Bruijns, S., et al. (2025). Brain-wide representations of prior information in mouse decision-making. Nature *645*, 192-200. 10.1038/s41586-025-09226-1.

37. David, S.V., Fritz, J.B., and Shamma, S.A. (2012). Task reward structure shapes rapid receptive field plasticity in auditory cortex. Proc Natl Acad Sci U S A *109*, 2144-2149. 10.1073/pnas.1117717109.

38. Toth, J., Sidleck, B., Lombardi, O., Hou, T., Eldo, A., Kerlin, M., Zeng, X., Saeed, D., Agarwal, P., Leonard, D., et al. (2024). Dynamic gating of perceptual flexibility by non-classically responsive cortical neurons. Res Sq. 10.21203/rs.3.rs-4650869/v1.

39. Rinne, T., Balk, M.H., Koistinen, S., Autti, T., Alho, K., and Sams, M. (2008). Auditory selective attention modulates activation of human inferior colliculus. J Neurophysiol *100*, 3323-3327. 10.1152/jn.90607.2008.

40. Schneider, D.M., Nelson, A., and Mooney, R. (2014). A synaptic and circuit basis for corollary discharge in the auditory cortex. Nature *513*, 189-194. 10.1038/nature13724.

41. Yang, Y., Lee, J., and Kim, G. (2020). Integration of locomotion and auditory signals in the mouse inferior colliculus. Elife *9*. 10.7554/eLife.52228.

42. Choi, I., and Lee, S.H. (2025). Locomotion-dependent auditory gating to the parietal cortex guides multisensory decisions. Nat Commun *16*, 2308. 10.1038/s41467-025-57347-y.

43. Metzger, R.R., Greene, N.T., Porter, K.K., and Groh, J.M. (2006). Effects of reward and behavioral context on neural activity in the primate inferior colliculus. Journal of Neuroscience *26*, 7468-7476. 10.1523/Jneurosci.5401-05.2006.

44. Znamenskiy, P., and Zador, A.M. (2013). Corticostriatal neurons in auditory cortex drive decisions during auditory discrimination. Nature *497*, 482-485. 10.1038/nature12077.

45. Li, J.C., Liao, X., Zhang, J.X., Wang, M., Yang, N., Zhang, J., Lv, G.H., Li, H.H., Lu, J., Ding, R., et al. (2017). Primary Auditory Cortex is Required for Anticipatory Motor Response. Cereb Cortex *27*, 3254-3271. 10.1093/cercor/bhx079.

46. Guo, L., Weems, J.T., Walker, W.I., Levichev, A., and Jaramillo, S. (2019). Choice-Selective Neurons in the Auditory Cortex and in Its Striatal Target Encode Reward Expectation. J Neurosci *39*, 3687-3697. 10.1523/JNEUROSCI.2585-18.2019.

47. Ford, A.N., Czarny, J.E., Rogalla, M.M., Quass, G.L., and Apostolides, P.F. (2024). Auditory Corticofugal Neurons Transmit Auditory and Non-auditory Information During Behavior. J Neurosci *44*. 10.1523/JNEUROSCI.1190-23.2023.

48. Quass, G.L., Rogalla, M.M., Ford, A.N., and Apostolides, P.F. (2024). Mixed Representations of Sound and Action in the Auditory Midbrain. Journal of Neuroscience *44*. ARTN e183123202410.1523/JNEUROSCI.1831-23.2024.

49. Drieu, C., Zhu, Z., Wang, Z., Fuller, K., Wang, A., Elnozahy, S., and Kuchibhotla, K. (2025). Rapid emergence of latent knowledge in the sensory cortex drives learning. Nature. 10.1038/s41586-025-08730-8.

50. Winnubst, J., Bas, E., Ferreira, T.A., Wu, Z., Economo, M.N., Edson, P., Arthur, B.J.,
Bruns, C., Rokicki, K., Schauder, D., et al. (2019). Reconstruction of 1,000 Projection
Neurons Reveals New Cell Types and Organization of Long-Range Connectivity in
the Mouse Brain. Cell *179*, 268-281 e213. 10.1016/j.cell.2019.07.042.
51. Munoz-Castaneda, R., Zingg, B., Matho, K.S., Chen, X., Wang, Q., Foster, N.N., Li,
A., Narasimhan, A., Hirokawa, K.E., Huo, B., et al. (2021). Cellular anatomy of the
mouse primary motor cortex. Nature *598*, 159-166. 10.1038/s41586-021-03970-w.
52. Stringer, C., Zhong, L., Syeda, A., Du, F., Kesa, M., and Pachitariu, M. (2025).
Rastermap: a discovery method for neural population recordings. Nat Neurosci *28*,
201-212. 10.1038/s41593-024-01783-4.
53. Pachitariu, M., Stringer, C., Dipoppa, M., Schröder, S., Rossi, L.F., Dalgleish, H.,
Carandini M., and, D., H.K. (2016). Suite2p: beyond 10,000 neurons with standard
two-photon microscopy. BioRxiv. https://doi.org/10.1101/061507.
54. Chudasama, Y., and Robbins, T.W. (2003). Dissociable contributions of the
orbitofrontal and infralimbic cortex to pavlovian autoshaping and discrimination
reversal learning: further evidence for the functional heterogeneity of the rodent
frontal cortex. J Neurosci *23*, 8771-8780. 10.1523/JNEUROSCI.23-25-08771.2003.
55. Kesner, R.P., and Churchwell, J.C. (2011). An analysis of rat prefrontal cortex in
mediating executive function. Neurobiol Learn Mem *96*, 417-431.
10.1016/j.nlm.2011.07.002.
56. Cho, S., Lee, H., Baek, S., and Paik, S.B. (2025). Neuromimetic metaplasticity for
adaptive continual learning without catastrophic forgetting. Neural Netw *190*, 107762.
10.1016/j.neunet.2025.107762.
57. Wilson, R.C., Takahashi, Y.K., Schoenbaum, G., and Niv, Y. (2014). Orbitofrontal
Cortex as a Cognitive Map of Task Space. Neuron *81*, 267-279.
10.1016/j.neuron.2013.11.005.
58. Du, X., Xu, H., Song, P., Zhai, Y., Ye, H., Bao, X., Huang, Q., Tanigawa, H., Tu, Z.,
Chen, P., et al. (2025). The multifaceted role of the inferior colliculus in sensory
prediction, reward processing, and decision-making. Elife *13*. 10.7554/eLife.101142.
59. Lee, T.-Y., Weissenberger, Y., King, A.J., and Dahmen, J.C. (2024). Midbrain encodes
sound detection behavior without auditory cortex.   eLife Sciences Publications, Ltd.
60. Carlisi, C.O., Norman, L.J., Lukito, S.S., Radua, J., Mataix-Cols, D., and Rubia, K.
(2017). Comparative Multimodal Meta-analysis of Structural and Functional Brain
Abnormalities in Autism Spectrum Disorder and Obsessive-Compulsive Disorder.
Biol Psychiatry *82*, 83-102. 10.1016/j.biopsych.2016.10.006.
61. Uddin, L.Q., Yeo, B.T.T., and Spreng, R.N. (2019). Towards a Universal Taxonomy of
Macro-scale Functional Human Brain Networks. Brain Topogr *32*, 926-942.
10.1007/s10548-019-00744-6.
62. Koithan, E.M., Demeter, D.V., Ali, S.A., Feigelis, M., and Greene, D.J. (2024).
Cognitive flexibility in neurodevelopmental disorders: insights from neuroimaging
and neuropsychology. Current Opinion in Behavioral Sciences *59*, 101429.
https://doi.org/10.1016/j.cobeha.2024.101429.
63. Grant, J.E., and Chamberlain, S.R. (2023). Impaired cognitive flexibility across
psychiatric disorders. CNS Spectr *28*, 688-692. 10.1017/S1092852923002237.
64. Gulati, S., Cao, V.Y., and Otte, S. (2017). Multi-layer Cortical Ca2+ Imaging in Freely
Moving Mice with Prism Probes and Miniaturized Fluorescence Microscopy. J Vis
Exp. 10.3791/55579.
65. Jeong, H., Kim, D., Song, M., Paik, S.B., and Jung, M.W. (2020). Distinct roles of
parvalbumin- and somatostatin-expressing neurons in flexible representation of task

variables in the prefrontal cortex. Prog Neurobiol *187*, 101773.
10.1016/j.pneurobio.2020.101773.

66. Pachitariu, M., Sridhar, S., Pennington, J., and Stringer, C. (2024). Spike sorting with
Kilosort4. Nat Methods *21*, 914-921. 10.1038/s41592-024-02232-7.

67. Raposo, D., Kaufman, M.T., and Churchland, A.K. (2014). A category-free neural
population supports evolving demands during decision-making. Nat Neurosci *17*,
1784-1792. 10.1038/nn.3865.

68. Ghosh, K.K., Burns, L.D., Cocker, E.D., Nimmerjahn, A., Ziv, Y., Gamal, A.E., and
Schnitzer, M.J. (2011). Miniaturized integration of a fluorescence microscope. Nat
Methods *8*, 871-878. 10.1038/nmeth.1694.

69. Ziv, Y., Burns, L.D., Cocker, E.D., Hamel, E.O., Ghosh, K.K., Kitch, L.J., El Gamal,
A., and Schnitzer, M.J. (2013). Long-term dynamics of CA1 hippocampal place codes.
Nat Neurosci *16*, 264-266. 10.1038/nn.3329.

70. Pinto, L., and Dan, Y. (2015). Cell-Type-Specific Activity in Prefrontal Cortex during
Goal-Directed Behavior. Neuron *87*, 437-450. 10.1016/j.neuron.2015.06.021.

71. Bondanelli, G., Deneux, T., Bathellier, B., and Ostojic, S. (2021). Network dynamics
underlying OFF responses in the auditory cortex. Elife *10*. 10.7554/eLife.53151.

72. Knyazev, A.V., and Argentati, M.E. (2002). Principal angles between subspaces in an
-based scalar product:: Algorithms and perturbation estimates. Siam J Sci Comput *23*,
2008-2040. Pii S1064827500377332 Doi 10.1137/S1064827500377332.

73. Miri, A., Daie, K., Burdine, R.D., Aksay, E., and Tank, D.W. (2011). Regression-
Based Identification of Behavior-Encoding Neurons During Large-Scale Optical
Imaging of Neural Activity at Cellular Resolution. Journal of Neurophysiology *105*,
964-980. 10.1152/jn.00702.2010.

74. Chen, J.L., Voigt, F.F., Javadzadeh, M., Krueppel, R., and Helmchen, F. (2016). Long-
range population dynamics of anatomically defined neocortical networks. Elife *5*.
10.7554/eLife.14679.

75. Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter,
E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent
proteins for imaging neuronal activity. Nature *499*, 295-300. 10.1038/nature12354.

76. slandarer, Z.L. (2025). sankey plot.
https://kr.mathworks.com/matlabcentral/fileexchange/128679-sankey-plot.

77. Yang, S.H., Yang, E., Lee, J., Kim, J.Y., Yoo, H., Park, H.S., Jung, J.T., Lee, D., Chun,
S., Jo, Y.S., et al. (2023). Neural mechanism of acute stress regulation by trace
aminergic signalling in the lateral habenula in male mice. Nat Commun *14*, 2435.
10.1038/s41467-023-38180-7.

78. Lee, H., Choi, W., Park, Y., and Paik, S.B. (2020). Distinct role of flexible and stable
encodings in sequential working memory. Neural Netw *121*, 419-429.
10.1016/j.neunet.2019.09.034.

1531

A Auditory Go/No-go reversal task

	Lick	No lick
Go	Hit	Miss
No-go	FA	CR

Air puff & time out

Water reward

Trials

Lick → Water reward
Air puff + time out

Stim | Delay | Response window

Session: Rule1 (R1) | Rule2 (R2) | Rule1 (R1) Cont. miss ≥ 20trials

Go: 5kHz → 10kHz → 5kHz
No-go: 10kHz → 5kHz → 10kHz

B Naive — 5kHz, 10kHz

C Expert — 5kHz, 10kHz

D R = -0.9975, p = 9.047 × 10⁻⁶, n=16

E TTR, n=16

F Lick reaction times: R1 Hit, R2 Hit, R1 FA, R2 FA *

G Lick rates: R1 Hit, R2 Hit, R1 FA, R2 FA

H Training ≥0.75 over 3 days → Test
Day1 Pre (No injection), Day2 Test (MUS), Day3 Control (PBS w/ Ruby)
AC: MUS or PBS; IC: MUS or PBS

I AC inactivation (bilateral), ****, **, **, ****, **, **, N=6
Hit rate, FA rate, Correct rate (Pre, MUS, PBS)

J Hit, FA, Correct — Mus, PBS, p-value

K AC inactivation: Pre, MUS, PBS *, *

L Pre, Mus, PBS — R2 Correct rate vs TTR

M IC inactivation (unilateral), *, *, ****, **, **, **, *, *, ****, **, N=7
Hit rate, FA rate, Correct rate (Pre, MUS, PBS)

N Hit, FA, Correct — p-value, Trials

O AuD Au1 AuV; DCIC ECIC CIC

P AC (n=400): R1 TR R1 5kHz 10kHz; Across rules None Both 10kHz 5kHz

Q IC (n=179): R1 TR R1 5kHz 10kHz; Across rules None Both 10kHz 5kHz

R AC #48, GMI₅kHz = 0.30; AC *, **

S IC #119, GMI₅kHz = -0.44; IC *, *, n.s., n.s.

T AC #48, GMI₁₀kHz = 0.48; AC ****, ****

U IC #119, GMI₁₀kHz = 0.04; IC n.s., n.s.

V N₁ threshold Nₙ; 10kHz 5kHz Firing rates; ROC True positive, False positive, AUC

W AC #194, SI_R1 = 0.00, SI_R2 = -0.43

X ΔISI: AC *, IC n.s. *

Y Fraction: Stable (AC, IC) ****, Flex. (AC, IC) **

A Parallel model

B

C Integrated model

D

E Accuracy over iteration

F

G TTR

A **SU recording sessions** B C

D Lick reaction times E Lick rates

F AC (n=400) G IC (n=179)

Stimulus
Contingency
Lost
Acquired
non-selective
non-responsive

	PPC_{AC}	PPC_{IC}	Dual
S1	75.74	23.37	1.11
S2	79.32	22.48	1.80
S3	91.10	10.07	1.11
Ave	82.03	19.30	1.13

A Imaging sessions

Performance rate / Hit, FA, Correct across R1, TR, R2

PPC_AC (n=6)
PPC_C (n=12)

B TTR — n.s. — PPC_AC, PPC_C

C R2 Correct rate vs TTR

D Lick reaction times
Reaction times (sec): R1 Hit, R2 Hit, R1 FA, R2 FA

E Lick rates
Lick rates (Hz): R1 Hit, R2 Hit, R1 FA, R2 FA

F Explained variance vs # of PCs
PPC_AC (84.77%)
PPC_C (89.41%)