Plasmin-Mediated Processing of Clumping Factor A Impacts Bacterial, Aggregation and Abscess Formation during *Staphylococcus aureus* Infection

Mary B. Turley[1,2,3], Zhicheng Hu[4], Ross W. Ward[5], Ed C. Lavelle[5], Tao Jin[4,6], Joan A. Geoghegan[1,2,3,*]

[1]Department of Microbiology, Moyne Institute of Preventive Medicine, School of Genetics and Microbiology, Trinity College Dublin, Dublin, Ireland.

[2]Department of Microbes, Infection and Microbiomes, College of Medicine and Health, University of Birmingham, Birmingham, United Kingdom.

[3]Institute of Microbiology and Infection, University of Birmingham, Birmingham, United Kingdom.

[4]Department of Rheumatology and Inflammation Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden.

[5]Adjuvant Research Group, School of Biochemistry and Immunology, Trinity Biomedical Sciences Institute, Trinity College Dublin, Dublin, Ireland.

[6]Department of Rheumatology, Sahlgrenska University Hospital, Gothenburg, Sweden.

*Corresponding author: Joan A. Geoghegan. Mailing address: Department of Microbes, Infections and Microbiomes, College of Medicine and Health, University of Birmingham. E-mail: j.geoghegan@bham.ac.uk.

Abstract

The interface between pathogenic bacteria and the host plays a critical role during the progression of infection. Bacterial surface components, such as the *Staphylococcus aureus* cell wall-anchored protein clumping factor A (ClfA), interact with host factors. ClfA promotes bacterial aggregation in plasma by binding to fibrinogen. Here, we show that the host serine endopeptidase plasmin cleaves ClfA, disrupting its interaction with fibrinogen and promoting bloodstream survival and virulence in *S. aureus.* We found that plasmin cleaves ClfA between residues Arg214 and Ala215. Cleavage occurs in human serum regardless of whether plasmin becomes activated by host tissue-plasminogen activator or by the activity of the *S. aureus* enzyme staphylokinase. Expressing a plasmin-resistant form of ClfA, and a truncated form that mimics the plasmin-cleaved product on the bacterial cell surface, revealed that following cleavage, ClfA cannot mediate aggregation of *S. aureus* in fibrinogen. Truncated ClfA on the cell surface cannot bind to fibrinogen and promotes the establishment of renal abscesses *in vivo.* Collectively, our findings suggest that activation of plasmin during infection leads to site-specific processing of ClfA, and this not only abrogates fibrinogen binding but also facilitates enhanced *S. aureus* persistence in the host. Thus, we demonstrate that proteolytic processing by plasmin is a mechanism for modulating the host-pathogen interaction during infection.

Author Summary

Staphylococcus aureus is a major human bacterial pathogen that binds to host factors during infection using its surface adhesins. Here we describe a novel

mechanism by which host proteolytic activity modulates bacterial interaction with the

plasma protein fibrinogen to influence the course of infection. The surface adhesin

clumping factor A (ClfA), binds to fibrinogen, causing bacterial aggregation in the

bloodstream. In this study, we show that the host enzyme plasmin cleaves ClfA at a

specific site, disrupting its ability to bind fibrinogen. Cleavage eliminates ClfA-

mediated aggregation in plasma fibrinogen and enhances *S. aureus* virulence in

vivo, increasing the formation of kidney abscesses. Our work therefore demonstrates

that specific protease-mediated processing of bacterial adhesins by host factors can

influence the progression of infection.

Introduction

Staphylococcus aureus bacteraemia is among the most common and serious bacterial infections affecting the global population (1). S. aureus was the leading bacterial cause of death in 135 countries in 2019 (2), and more than 300,000 people die each year as a direct result of S. aureus bacteraemia (3). Complications arise when S. aureus evades host immune defences and disseminates to organs and tissues giving rise to metastatic infections such as infective endocarditis, septic arthritis, and infections associated with implanted medical devices. These infections increase morbidity and mortality, often requiring prolonged treatment and, in some cases, surgical intervention.

One of the key virulence factors of S. aureus is clumping factor A (ClfA), a cell wall-anchored protein that plays a central role during bacteraemia, septic arthritis and infective endocarditis through fibrinogen-dependent and independent mechanisms (4-10). ClfA binds to fibrinogen, a glycoprotein present in blood plasma, promoting the aggregation of S. aureus in plasma. However, fibrinogen also functions in host defense by activating neutrophils and macrophages, as well as trapping bacteria in matrices and inhibiting dissemination (11-15). Fibrinogen-deficient mice have reduced ability to clear S. aureus from the peritoneal cavity following infection, and disruption of the S. aureus-fibrinogen interaction inhibits dissemination from the peritoneum during acute infection (16).

ClfA binds to the extreme C-terminus of the γ chain of fibrinogen. The C-terminal residues of the γ chain bind in a trench formed between the separately folded adjacent immunoglobulin G-like domains N2 and N3 domains by the 'dock, lock, and latch' mechanism (17-19). Docking of the peptide induces a conformational

change in the flexible extension at the C-terminus of the ClfA N3 domain, which

redirects to cover the bound fibrinogen peptide locking it in place. In doing so it forms

a β strand which is complementary to a β sheet in the N2 domain. The D domain of

fibrinogen establishes further interactions with another site on the surface of the ClfA

N3 domain (20). The overall stability of the ClfA-fibrinogen complex is enhanced by

shear stress (21). ClfA has also been reported to bind to the host protein factor I,

facilitating cleavage of complement opsonin C3b into iC3b to inhibit

opsonophagocytosis (10, 22). However, no studies have yet detailed the specific

mechanism or binding site involved.

There is growing evidence that host-derived proteases alter the *S. aureus* cell

surface, potentially impacting its interactions with the host during infection (23-25).

The host serine protease plasmin is particularly relevant in the context of infection

due to its activation at sites of tissue damage and inflammation (26, 27). Notably, *S.*

aureus produces staphylokinase (Sak), a secreted protein that activates host

plasminogen to its proteolytically active form, plasmin (28). This interaction suggests

a potential mechanism by which *S. aureus* may exploit the host fibrinolytic system to

remodel its own cell surface. This study set out to determine if ClfA is susceptible to

plasmin and whether plasmin can modulate the biological activity of ClfA.

Materials and Methods

Reagents, Strains, Media, and Growth Conditions

All strains and plasmids used in this study are listed in Table 1. If not specified otherwise, *S. aureus* strains were cultured on tryptic soy agar (TSA) plates for isolation of individual colonies or for liquid cultures in tryptic soy broth (TSB) (Difco) at 37°C with shaking (200 rpm). *Escherichia coli* strains were grown at 37°C on lysogeny Agar (LA) or in lysogeny Broth (LB) with shaking (200 rpm) for liquid cultures. Where appropriate, antibiotics were incorporated into media; chloramphenicol (Cm,10 µg/mL), erythromycin (Em, 10 µg/mL), ampicillin (Amp, 100 µg/mL); anhydrotetracycline (Atc, 1 ng/mL) (Sigma).

Plasmin treatment

Bacteria were grown overnight for 16-20 h, then pelleted by centrifugation (7 min, 8000 x *g*), washed with an equal volume of PBS once and pelleted again. Samples were then resuspended in 1 mL PBS and adjusted to the desired OD_{600} for the assay. Human plasmin (Sigma) was added to the sample to a final concentration of 0.075 U/mL and incubated at 37°C for 1 h in a water bath. Samples were then washed with an equal volume of PBS three times before a final reconstitution to original volume. The protease inhibitor alpha-2-macroglobulin (α2M, Sigma) was added where indicated at a concentration of 10 times that of plasmin used and preincubated with plasmin for 20 minutes to inactivate the protease. Recombinant proteins were typically diluted to a concentration of 5 µM in sterile PBS to which plasmin was added to a final concentration of 0.075 U/mL and incubated at 37°C for 1 h in a water bath.

Preparation of human and murine blood and serum

Fresh blood samples were drawn from healthy human volunteers, into serum clot activator vacuette tubes (Greiner) or EDTA-containing vacuettes (Greiner). Serum was collected by inverting the serum clot activator tubes, then incubating at room temperature for 15 min before centrifuging at 1,500 x g for 16 min. To collect murine serum, whole blood was collected from female C57BL6/JCR mice by tail bleed and was allowed to clot overnight at 4°C. Serum was then harvested from clotted blood by centrifugation at 1100 x g for 11 min. To catalyse the activation of plasminogen to plasmin in human serum, 40% serum was incubated with tissue plasminogen activator (DiaPharma, 25 µg/mL) or *S. aureus* supernatant for 20 min at 37°C before being used in the treatment of recombinant proteins as outlined above.

SDS-PAGE and Western immunoblotting

Cell wall extractions were carried out as described by O'Halloran *et al.*, 2015 (29)**.** Briefly, bacteria grown to exponential phase (OD_{600} = 0.35 - 0.5) or stationary phase (overnight culture) were pelleted and washed with PBS, then adjusted to an OD_{600} = 10. Bacteria were then incubated with or without plasmin (0.075 U/mL) in a 37°C waterbath for 1 h before being centrifuged and resuspended in 250 µL extraction solution (30% w/v Raffinose, 50 mM Tris-HCL, 20 mM MgCl2, pH 7.5, 50 µl/ml complete protease inhibitor [Roche], lysostaphin 10 U/mL) for 8 min at 37°C. Protoplasts were removed by centrifugation at 15000 x g for 5 minutes.

Protein samples were separated by polyacrylamide gel electrophoresis (7.5%, 10% or 12.5%). Gels were either stained using Instant Blue (Expedeon) or Gel code

(ThermoFisher Scientific) or transferred to nitrocellulose (GE Healthcare) or

polyvinylidene difluoride (PVDF, Roche) filters for western immunoblotting.

Filters were blocked with 10% (w/v) skimmed milk (Marvel) and then probed

with polyclonal anti-ClfA N1N2N3 IgG (1:1000, (30)) followed by horseradish

peroxidase (HRP)-conjugated protein A (1:500, Sigma), 7E8 IgG (1:1000 (31))

followed by goat anti-mouse IgG-HRP (1:2500, Dako) or alternatively with Strep-

tag® II IgG-HRP (1:2500, Novagen). Filters were developed using LumiGLO

Reagent and peroxide detection solution (Cell signalling Technology) and imaged

using the ImageQuant TL (GE).

DNA manipulation

Isolation of DNA Genomic DNA was isolated using the PureElute Bacterial

genomic DNA Purification kit (Edge Biosystems), with the addition of lysostaphin

(100 µg/ml) to the cell lysis buffer to digest *S. aureus* cell wall peptidoglycan to

improve DNA extraction. Plasmid DNA was isolated using Wizard® Plus SVMiniprep

kit (Promega) or Qiagen® Plasmid plus Midi-kit.

Strains	Description	Reference or source
Staphylococcus aureus		
BH1CC	MRSA, CC8	(32)
Newman	MSSA, CC8	(33)
Newman *clfA5*	Derivative of Newman deficient in ClfA, frameshift mutation	(34)
S. aureus ClfA$^+$	Newman *clfA5* (pRMC2::*clfA*). ClfA-deficient mutant carrying the entire *clfA* gene on a plasmid	This study
S. aureus ClfA^R	Newman *clfA5* (pRMC2::*clfA^R*). ClfA-deficient mutant carrying the entire *clfA* gene with DNA encoding residues 211-214 exchanged for DNA encoding AlaSerAlaSer on a plasmid	This study
S. aureus ClfA$^{\Delta40\text{-}214}$	Newman *clfA5* (pRMC2::*clfA$_{\Delta40\text{-}214}$*). ClfA-deficient mutant carrying the *clfA* gene lacking DNA encoding residues 40-214 on a plasmid	This study
S. aureus ClfA$^-$	Newman *clfA5* (pRMC2). ClfA-deficient mutant carrying the empty plasmid.	This study
LAC*	MRSA; CC8, Em sensitive	(35)
LAC* *sak*::tn	Derivative of LAC* deficient in *sak*. *sak*::Tn Em^R	(36)
14-2533T	MRSA, CC22	(37)
Escherichia coli		
XL-1-Blue	Cloning host. *recA1 endA1 gyr96 thi-1 hsdR17 suppe44 relA1 lac*	Stratagene

Topp3	Expression host [F'proAB+ lacqZΔM15 Tn10 (Tet□) (Kan□)]	Stratagene
IM08B	Host for plasmid propagation. SA30BΩPN25-*hsdS* (CC30-1)	(38)
Plasmids		
pQE30::*clfA*40-559*	pQE30 carrying DNA encoding ClfA residues 40-559 in-frame with a C-terminal Strep tag II. Amp^R	This study
pQE30::*clfA*^{R}40-559*	pQE30::ClfA40-559* with residues 211-214 of ClfA substituted with AlaSerAlaSer. Amp^R	This study
pRMC2	Tetracycline-inducible expression vector. Cm^R	(39)
pRMC2::*clfA*	pRMC2 containing full-length *clfA* gene from strain Newman. Cm^R	(40)
pRMC2::*clfA*^{R}	pRMC2::*clfA* with DNA encoding residues 211-214 of ClfA substituted for AlaSerAlaSer. Cm^R	This study
pRMC2::*clfA*$^{\Delta40-214}$	pRMC2::*clfA* lacking DNA encoding residues 40-214 of ClfA. Cm^R	This study

170 MRSA = Methicillin resistant *Staphylococcus aureus*; MSSA = Methicillin sensitive

171 *Staphylococcus aureus*; CC = Clonal complex; Tn = Transposon; ^{R} = resistant; Em =

172 erythromycin; Cm = chloramphenicol; Amp = Ampicillin

Plasmid construction

Plasmids were constructed by inverse PCR or via sequence and ligase independent cloning (51). Plasmids pQE30::ClfA$^R_{40-559*}$ and pRMC2::ClfA^R were constructed by sequence and ligase independent cloning (41) where vectors were amplified using Velocity polymerase and the insert was amplified from a region of DNA commercially synthesised as a gBlock (IDT). Primers used in this study are listed in **Table S1**. Plasmids pQE30::ClfA$_{40-559*}$ and pRMC2::ClfA$^{\Delta40-214}$ were generated by inverse PCR. Inverse PCR products were treated with DpnI and blunt-end ligation was carried out using the LigaFast® Rapid RNA ligation system (Promega). Chemically competent *E. coli* (42) XL-1 Blue was transformed with the ligation products. Plasmids were then isolated and used to transform *E. coli* Topp3 made chemically competent for protein purification or *E. coli* IM08B for transformation of *S. aureus* made electrocompetent as described previously (43). Colony PCRs were carried out to screen for transformants using Phire Hot Start DNA Polymerase (Sigma) as per the manufacturer's instructions.

Expression and purification of recombinant proteins

Briefly, *E. coli* Topp3 carrying plasmids was grown to an OD$_{600}$ = 0.4 - 0.8, before isopropyl β-D-1-thiogalactopyranoside (IPTG, 1 mM) was added to induce protein expression. Growth was continued for 3 h before cell sedimentation was carried out. The pellet was resuspended in 25 mL Imidazole Buffer (40 mM Tris-HCL, 1 M NaCl, 0.1 M imidazole, pH 7.9) with an EDTA-free protease inhibitor. Bacteria were then lysed using a French Press at 1500 psi before centrifuging the lysates at 18,000 x g for 20 mins at 4°C. DNase (1 mg/mL) was added to the supernatant before filtration through a 0.45 µm filter (Whatman). Proteins were purified with an N-terminal

6xhexa-histidine (His) affinity tag and a C-terminal Strep-tag II affinity tag for

purification via nickel affinity chromatography (5 mL HiTrap affinity column; GE

Healthcare) and Strep-Tactin affinity chromatography (Strep-Spin Protein Miniprep

kit; Zymo Research) respectively. Protein concentrations were determined by

measuring the absorbance at 280 nm using a Nanodrop spectrophotometer and

calculations according to Beer-Lambert Law.

207   N-terminal sequencing

Samples sent for N-terminal sequencing were first incubated with plasmin (0.075

U/mL) as outlined above prior to purification of the digested product by Strep-Tactin

affinity chromatography. Purified samples were then subjected to SDS-PAGE,

transferred to PVDF membrane and stained with Ponceau S (ThermoScientific). N-

terminal sequencing was performed by Edman degradation at Alta Bioscience,

Redditch, UK.

215   Synthetic peptide cleavage

Synthetic peptides used in this study were commercially synthesised by

AltaBioscience (Redditch, UK). Peptide size and purity was confirmed by mass

spectrometry at AltaBioscience. All peptides were dissolved in molecular grade water

to a concentration of 150 µM. As the peptides were 5/6-FAM labelled they were

protected from light when in use and stored at -20°C. Synthetic peptides were diluted

to a concentration of 37.5 µM in molecular grade water, then incubated with human

plasmin (0.075 U/mL) at 37°C for 30 min in a water bath. Reaction products were

separated by agarose gel electrophoresis (4% w/v) and gel images were captured

under UV light.

Bacterial clumping in fibrinogen

Bacteria were grown to stationary phase, then pelleted by centrifugation (8000 x *g,* 7

min). Cells were washed with PBS and then adjusted to OD_{600} of 1.5 in 1 mL PBS.

Purified human fibrinogen depleted of plasminogen, von Willebrand Factor and

fibronectin (Enzyme Research Labs) or murine fibrinogen (Innovative Research) was

added to the tube to a final concentration of 18.4 μg/mL and left to stand at 37°C.

After incubation for 1 h 45 min, the top 0.5 mL was removed from the tube, diluted

1:2 in PBS for OD_{600} measurements (Genesys 10 UV, Thermo Spectronic) and

calculated as a percentage of the starting OD_{600} where $(100/T0\_OD_{600})*T105\_OD_{600}$,

where T0 is the starting OD_{600} and T105 is the OD_{600} after 1 h 45 min. For each *S.*

aureus strain, there was a fibrinogen-deficient control. In cases where plasmin

treated bacteria were included, on adjustment to an $OD_{600} = 1.5$ the bacteria were

incubated with plasmin (0.075 U/mL) for 1 h at 37°C. Following three washes in PBS,

the bacteria were resuspended to the original volume in BHI and made up to a final

volume of 1 mL with PBS. Fibrinogen (18.4 ug/mL) was then added, and the protocol

followed as outlined above. In each case for plasmin treatment, a no plasmin control

for each strain was included.

Ex vivo **whole blood survival assay**

The quantification of survival of *S. aureus* in human blood was performed according

to the protocol previously described by Zapotoczna *et al.*, 2018 (37). Briefly, *S.*

aureus was grown overnight in RPMI then adjusted to 2×10^4 cfu/mL. 25 μL was

added to 475 μL of freshly drawn EDTA-treated blood obtained from human

volunteers. Tubes were inverted and 100 μL of each sample was immediately added

to 900 µL ice-cold endotoxin free water and serial dilutions were made in water for

plating on agar to establish the CFU/mL of the sample at T0. The tubes were then

incubated at 37°C on a 360° rotation wheel and after 3 h (T3), serial dilutions were

made and bacteria plated on agar to establish the number of CFU per mL. In

parallel, bacteria were incubated in serum derived from the same donor. Percentage

survival was calculated using the following formula (T3 CFU/mL)/(T0 cfu/mL)*100.

Mouse model of *S. aureus* bloodstream infection

Female NMRI, aged 9-10 weeks, were obtained from Envigo (Venray, Netherlands)

and housed at the animal facility at the University of Gothenburg. Mice were

maintained under standard temperature and light conditions, with unrestricted

access to laboratory chow and water. *S. aureus* strains were cultured for 24 hours on

horse blood agar plates and then preserved according to established protocols (44).

Before each experiment, the bacterial solutions were thawed, washed, and adjusted

to the desired concentration. To investigate the clinical relevance of ClfA plasmin

cleavage, we used a well-established *S. aureus* sepsis mouse model (45). NMRI

mice (10 per group) were intravenously injected via the tail vein with 200 µL of PBS

containing 75 ng/mL anhydrotetracycline and the respective *S. aureus* strain

Newman *clfA5* expressing pRMC2 variants: ClfA$^+$, ClfA^R or ClfA$^{\Delta 40\text{-}214}$ (6 ×

10^6 colony-forming units [CFU]/mouse). Expression of ClfA was induced via an

intraperitoneal injection of inducer anhydrotetracycline (5 µg/mouse) on Day 0, 1 h

before infection and 1h post-infection. The mice were monitored individually two

times daily by an observer (Z.H). The blood samples were collected on both day 3

and day 10 for subsequent Interleukin 6 (IL-6) level analysis. Surviving mice were

sacrificed on day 10, the blood, kidneys and livers were aseptically collected.

Quantification of bacterial load in murine kidneys and livers

Kidneys and livers were collected aseptically, and kidney abscess scores were assessed by two investigators (T.J. and Z. H.) in a blinded manner (46). The scoring system used ranged from 0 to 3 (0 indicates healthy kidneys; 1, 1–2 small abscesses in the kidneys without structure changes; 2, >2 abscesses, but <75% kidney tissue involved; and 3, large amounts of abscess with >75% kidney tissue involved). The kidneys and livers were homogenized and plated on horse blood agar plates to quantify the CFU counts.

Quantification of interleukin 6 using enzyme-linked immunosorbent assay

The levels of Interleukin 6 (IL 6) in the blood of NMRI mice infected with *S. aureus* were measured on day 3 and 10 post-infection using a DuoSet ELISA kit (R&D Systems, Abingdon, UK) according to the manufacturer's instructions.

Ethical Approval

Our research complies with all relevant ethical regulations. Informed consent was obtained from all adult participants. Ethical approval for the collection and use of human blood samples from adult volunteers was obtained from the Trinity College Dublin Faculty of Health Sciences Ethics Committee, Ireland (reference 2020111) and through the Human Biomaterials Resource Centre, University of Birmingham, United Kingdom (reference 25/NW/0013). Personal information such as sex and age

were not recorded. All participants were over the age of 18 and provided written

informed consent. Participants did not receive compensation for their participation.

For animal studies conducted in Trinity College Dublin, C57BL/6 mice were bred in-

house in the Comparative Medicine Unit, as per university and The Health Products

Regulatory Authority (HPRA) welfare guidelines. All animal studies were approved

by the TCD Animal Research Ethics Committee (Ethical Approval Number 091210)

and HPRA license AE191364/P079. For mouse work, the study was approved by the

Gothenburg Ethics Committee of Animal Research (Ethical Approval Number 5.8.18-

02443/2021), and all procedures adhered to established guidelines for animal

experimentation.

310   Statistical analysis

Statistical analysis was performed using GraphPad Prism (GraphPad Software, Inc,

USA; version 8.0.2.263 and version 10). Unless stated otherwise, the data presented

in this study show mean ± standard deviations of at least three independent

experiments. Statistically significant differences were calculated by using methods as

indicated in the figure legends. P values of less than 0.05 were considered

significant with *, **, *** and **** representing P values of <0.05, <0.01, <0.001 and

<0.0001 respectively. P values of >0.05 represents non-significant values (ns). For

experiments with blood or serum, samples from different donors were used for each

of the biological replicates.

Results

ClfA is cleaved by human plasmin

To determine if cell wall-anchored ClfA was susceptible to plasmin-mediated proteolytic processing in *S. aureus*, bacteria were grown to stationary phase, washed and incubated with plasmin. Cell wall-associated proteins were solubilised from the cell wall using lysostaphin during protoplast formation. Probing cell wall extracts of wild-type *S. aureus* Newman with anti-ClfA IgG in a western blot revealed a ca. 170 kDa immunoreactive band corresponding to full length ClfA (**Fig. 1A**). Incubation with plasmin resulted in the appearance of a ca. 130 kDa band in addition to the ca. 170 kDa band indicating that cell-wall anchored ClfA had been cleaved by plasmin. The broad-spectrum protease inhibitor alpha-2-macroglobulin prevented the activity of plasmin as only the band corresponding to full-length ClfA was detected (**Fig. 1A**). As a control, cell wall extracts were prepared from a mutant deficient in ClfA (Newman *clfA5*). The absence of immunoreactive bands confirmed the specificity of the anti-ClfA IgG (**Fig. 1A**). These results suggested that ClfA on the surface of *S. aureus* Newman can be cleaved by purified plasmin.

To explore plasmin cleavage further, three methicillin-resistant *S. aureus* (MRSA) isolates were incubated with plasmin and the integrity of ClfA in cell wall extracts was examined. For each of the isolates, ClfA was cleaved by plasmin as indicated by the disappearance of the ca. 170 kDa band corresponding to full-length ClfA and the appearance of a ca. 130 kDa immunoreactive band following plasmin treatment (**Fig. 1B**). A weak immunoreactive band migrating around 130 kDa was detected in cell wall extracts from LAC that had not been incubated with plasmin. This was likely due to the activity of the *S. aureus* extracellular protease aureolysin

that cleaves at the junction of N2 and N3 of ClfA (47). In summary, these findings

demonstrate that plasmin cleaves ClfA on the surface of *S. aureus*.

Recognition by an antibody specific for ClfA domains N1N2N3 (**Fig. 1**)

suggested that ClfA undergoes cleavage towards the N-terminus, leaving a portion

of the N1N2N3 domains intact and anchored to the cell wall. To examine this further,

a recombinant protein corresponding to the N1N2N3 domains of ClfA with an N-

terminal hexahistidine tag and a C-terminal Strep-tagII was purified from *E. coli*

(ClfA$_{40-559*}$). ClfA$_{40-559*}$ migrates aberrantly with an apparent molecular mass of ca. 95

kDa (**Fig. 2A**) due to the presence of the elongated unfolded N1 region (48).

Incubation of ClfA$_{40-559*}$ with plasmin resulted in the conversion of the ca. 95 kDa

band to a single product of ca. 43 kDa. The remainder of the protein could not be

detected (**FIG. 2A, B**) suggesting that it was digested into smaller fragments. ClfA$_{40-}$

$_{559*}$ remained mostly intact when the protease activity of plasmin was inhibited by

alpha-2-macroglobulin (**FIG. 2A, B**) confirming that cleavage of ClfA$_{40-559*}$ was due to

the proteolytic activity of plasmin.

To determine if the single band remaining following plasmin treatment

retained the N-terminal and/or C-terminal tags, the same samples were probed with

anti-His IgG to detect the N-terminal hexa-histidine tag (**FIG. 2C**) or anti-Strep tagII

IgG to detect the C-terminal strep tag (**Fig. 2D**). While a band corresponding to full-

length ClfA$_{40-559*}$ was detected in samples without plasmin or when plasmin was

inhibited with alpha-2-macroglobulin, using anti-His IgG, no band was detected after

the addition of plasmin (**FIG. 2C**). These data suggest that the hexa-histidine tag,

and therefore the N-terminal region of the recombinant protein, is removed by

plasmin cleavage. In contrast, the plasmin-cleaved product reacted with anti-Strep

tag II IgG (**FIG. 2D**) indicating that the C-terminus of the protein was intact. Other

bands detected in lanes 1 and 3 (**FIG. 2B, D**) are likely to be ClfA breakdown

product(s) that were already in the sample or formed during the incubation process.

To elucidate if ClfA can be cleaved by activated plasmin in human serum,

ClfA$_{40-559*}$ was incubated in serum with or without added tissue plasminogen

activator (tPA) which catalyses activation of the zymogen plasminogen (**Fig. 2E**) or

with concentrated culture supernatants containing the secreted *S. aureus*

plasminogen activating-enzyme Sak (**Fig. 2F**). Full length ClfA$_{40-559*}$ was detected

following incubation in serum, but only a band corresponding to cleaved ClfA was

detected when plasmin was activated (**Fig. 2E**). The addition of alpha-2-

macroglobulin protected full-length ClfA from cleavage. The presence of some

truncated protein in the absence of tPA or when plasmin activity was inhibited by

alpha-2-macroglobulin indicates that this species was present in the protein

preparation at the onset of the experiment or developed independently of plasmin

activation. No immunoreactive bands were detected in serum only controls

confirming the specificity of the polyclonal antibody for ClfA. Full length ClfA$_{40-559*}$

was detected following incubation in serum treated with concentrated supernatants

from an *S. aureus* mutant deficient in *sak* (LAC* *sak*::Tn) but not when ClfA was

incubated with supernatants from wild-type LAC* (**Fig. 2F**), indicating that plasmin

was activated by Sak and cleaved ClfA (**Fig. 2F**). Taken together, these findings

indicate that purified human plasmin, or plasmin activated by the host or bacterium

during infection, is capable of cleaving ClfA.

Plasmin cleaves ClfA between residues 214 and 215

394   To identify the plasmin cleavage site, ClfA$_{40-559*}$ was incubated with plasmin and the

395   cleaved protein was purified by Strep-Tactin affinity chromatography. The 7 most N-

terminal amino-acids of the cleaved product were identified using Edman degradation, revealing that plasmin cleaves the peptide bond after $Arg_{214,}$ towards the end of the N1 domain (**Fig. 3A**). To verify the plasmin cleavage site experimentally, an 11-mer fluorescein labelled peptide comprising ClfA residues 211-221, was incubated with plasmin and analysed by agarose gel electrophoresis. The 11-mer peptide was chosen based on the evidence from N-terminal sequencing that plasmin would cleave between residues 214 and 215 resulting in a clear size shift from a fluorescein-labelled 11-mer peptide to a fluorescein-labelled 4-mer. In agreement with this, the 11-mer peptide (WT) underwent complete conversion to a smaller truncate following plasmin treatment (**Fig. 3B**). Next a series of peptides where amino-acids 211-216 were individually substituted with serine were tested for susceptibility to plasmin cleavage (**Fig. 3B**). Only the R214S substitution affected cleavage with this peptide undergoing partial truncation following plasmin treatment underscoring the importance of residue Arg214 for plasmin recognition and/or cleavage. A scrambled version of the 11-mer peptide (S□) and peptide where the entire plasmin-cleavage site and surrounding residues were substituted (amino-acids 211-214, WT□) remained intact post-incubation with plasmin. Similarly, recombinant $ClfA_{40-559^*}$ harbouring amino-acid substitutions of residues 211-214 ($ClfA^R_{40-559^*}$) was resistant to plasmin cleavage (**Fig. 3C**). Overall, these data indicate that substitution of residues 211-214 prevents the cleavage of ClfA by plasmin and that R214 is particularly important for optimal cleavage by plasmin. Failure of plasmin to cleave the scrambled peptide indicates that the order of the residues is critical for plasmin to cleave this sequence.

Plasmin cleavage disrupts the fibrinogen binding activity of ClfA

ClfA binds fibrinogen in human plasma which facilitates clumping of *S. aureus* by way of the formation of fibrinogen bridges between bacteria (4, 49). The ability of ClfA to interact with fibrinogen was studied by incubating *S. aureus* in a solution of fibrinogen and monitoring the optical density at 600 nm of the suspension as the bacteria clump. A suspension of *S. aureus* expressing full-length ClfA (*S. aureus* ClfA$^+$) treated with plasmin to promote processing of ClfA prior to incubation with fibrinogen, underwent significantly less clumping than untreated bacteria (**Fig. 4A**). Conversely, bacteria expressing ClfA carrying amino-acid substitutions in residues 211-214 that render the protein resistant to plasmin cleavage (*S. aureus* ClfA^R) did not change clumping phenotype following plasmin treatment (**Fig. 4B**). Together these results demonstrate that the cleavage of ClfA by plasmin reduces the protein's ability to interact with soluble fibrinogen and promote clumping.

Next the ability of *S. aureus* expressing a ClfA truncate that mimics the plasmin-cleaved state, (*S. aureus* ClfA$^{\Delta 40-214}$), was tested for its ability to clump in a solution of fibrinogen (**Fig. 4C**). *S. aureus* ClfA$^{\Delta 40-214}$ lost the ability to clump (**Fig. 4C**) while *S. aureus* ClfA$^+$ (**Fig. 4D**) and *S. aureus* ClfA^R (**Fig. 4E**) incubated under the same conditions clumped in a solution of fibrinogen. *S. aureus* ClfA$^-$ did not clump demonstrating that clumping activity is mediated solely by ClfA under the conditions used here (**Fig. 4F**). Together these results suggest that removal of residues 40-214 either by treatment with plasminogen or genetically, renders ClfA incapable of binding to soluble fibrinogen and promoting clumping of *S. aureus*.

Truncation of ClfA enhances ClfA-mediated survival in human blood.

To determine whether truncation of ClfA, and the resulting loss of fibrinogen binding, influences bacterial killing by the host in blood, the survival of *S. aureus* was monitored over three hours in freshly drawn human blood (**Fig. 5**). *S. aureus* ClfA$^{\Delta40\text{-}214}$ displayed a significantly increased survival in whole human blood compared to both *S. aureus* ClfA^{+} and the cleavage resistant variant ClfA^{R}. There was no significant difference in the survival rates of *S. aureus* ClfA^{+} and ClfA^{R}. Importantly, all strains grew similarly in serum (**Fig, S1**), indicating that the better survival of ClfA$^{\Delta40\text{-}214}$ compared to ClfA^{+} and ClfA^{R} is not due to a difference in the ability to grow in the liquid component of blood, but rather a difference in resistance to phagocytic killing. Together these results suggest that removal of the N1 domain increases ClfA-mediated survival in blood.

Truncation of ClfA increases kidney abscess formation by *S. aureus*

ClfA promotes renal abscess formation in murine infection models (4, 50-52). Having shown that the truncated form of ClfA interacts poorly with soluble fibrinogen (**Fig. 4**) but enhances whole blood survival (**Fig. 5**), we sought to determine how this might impact *S. aureus* infection in a mouse model. ClfA was not susceptible to cleavage by mouse plasmin in serum (**Fig. S2**) so the effect of truncation on the activity of ClfA *in vivo* was examined using the *S. aureus* expressing full-length ClfA and its variants. In agreement with our findings with human fibrinogen (**Fig. 4**), *S. aureus* ClfA$^{\Delta40\text{-}214}$ was impaired in its ability to induce clumping in a solution of mouse fibrinogen (**Fig.**

S2). A sub-lethal intravenous dose of *S. aureus* ClfA$^{\Delta40\text{-}214}$ resulted in more severe

abscess formation (**Fig**. **6A**) and a higher bacterial load (**Fig. 6B**) in the infected

mouse kidneys compared to *S. aureus* ClfA^{+} or *S. aureus* ClfA^{R} indicating that the

ability of *S. aureus* to disseminate to the kidney and establish renal abscesses is

enhanced when residues 40-214 are missing from ClfA. There was no difference in

bacterial load in the liver following infection with any of the strains (**Fig. 6C**).

Comparing the survival of mice challenged with *S. aureus* expressing ClfA^{+}, ClfA^{R}

and ClfA$^{\Delta40\text{-}214}$ indicated that there were no significant differences in the survival rates

of mice infected with any of the three strains over the 10 days of the experiment (**Fig.**

6D). The concentrations of IL-6 mice infected by all three strains was similar (**Fig.**

S3). Together these data indicate that removal of residues 40-214 and therefore

inactivation of the fibrinogen binding function of ClfA enhances the ability of *S.*

aureus to form abscesses in mouse kidneys.

Discussion

In this study we show that cleavage of ClfA by plasmin results in a pronounced loss of bacterial clumping in fibrinogen, reduced survival of *S. aureu*s in human blood and increased abscess formation in the kidney. We pinpoint the plasmin cleavage site to between residues R214 and A215 in ClfA (**Fig. 2**) and demonstrate that the activity of plasmin leads to the formation of a stable truncate of ClfA on the cell surface lacking residues 40-214.

S. aureus actively recruits plasminogen to its surface and converts it to plasmin using Sak (28), thus providing a mechanism to inactivate the fibrinogen-binding activity of ClfA on the cell surface, even in the absence of host-activated plasmin. Existing evidence supports fibrinogen binding to bacteria conferring protective benefits to the host (11, 13, 15, 51, 53-56) while conversely, other studies find it can drive pathogenesis (4, 12, 16, 49, 50, 52, 57-61). Such differences in outcome likely reflect infection model or strain-dependent factors. Here we used a sub-lethal intravenous challenge with *S. aureus* and followed survival of the infected animals and dissemination to the kidneys and liver. This showed that loss of fibrinogen binding in the truncate resulted in impaired bacterial killing, enhanced dissemination and increased abscess formation. This agrees with previous findings showing that mice expressing a fibrinogen truncate lacking the ClfA binding site demonstrate delayed bacterial killing and increased dissemination from the peritoneal cavity (16).

The proteolytic cleavage of cell wall-anchored proteins in *S. aureus* has been described previously. In some instances, surface-exposed cell wall-anchored proteins are completely degraded by proteases, such as FnBPA and FnBPB (62, 63). However, in other cases, cleavage at specific sites by bacterial or host

proteases (24, 64, 65) serves as a post-translational modification that modulates

protein function. For example, ClfB loses its ability to bind fibrinogen when cleaved

by aureolysin (47) and SasG no longer inhibits bacterial clumping when processing

by trypsin (25).

In this study, we demonstrate that removal of residues 40-214 from the

mature ClfA protein abolishes its ability to interact with soluble fibrinogen. This

finding was unexpected, as the canonical fibrinogen-binding domain of ClfA is

thought to reside entirely within the N2 and N3 domains (residues 229-545) (18, 20).

Interestingly, the loss of fibrinogen binding in the ClfA$^{\Delta 40-214}$ variant mirrors the

ligand-binding behaviour of ClfB, where cleavage by aureolysin outside of the

minimal fibrinogen-binding domain results in a complete loss of function (47). These

observations suggest that the ClfA-fibrinogen interaction is more complex than

previously appreciated. While residues 229-545 are sufficient to support recombinant

protein binding to a peptide derived from fibrinogen, they are clearly not sufficient to

promote bacterial interactions with fibrinogen and fibrin in the bloodstream.

Elucidating the structure of the entire ClfA protein in complex with fibrinogen

would reveal the full binding interface and uncover how ClfA engages

fibrinogen/fibrin. This new finding highlights the drawbacks of studying host-

pathogen interactions using recombinant bacterial adhesin domains *in vitro* and

indicates that the interaction between ClfA and fibrinogen, and likely other

MSCRAMMs and their ligands, is far more complex than the dock, lock and latch

mechanism encapsulates and best studied using whole cells expressing the protein

of interest.

Plasmin cleaves ClfA at the carboxyl terminus of arginine residues, consistent

with its known substrate specificity (66). Proteolytic processing of ClfA results in the

release of residues N-terminal of A215, yet the corresponding fragment (the majority

of N1) is undetectable by SDS-PAGE suggesting it undergoes rapid secondary

degradation. Recombinant N1N2N3 domains of MSCRAMMs often resolve into a

single proteolytically stable fragment corresponding to N2N3, even in the absence of

added protease (67). Degradation of the N1 domain of MSCRAMMs is likely driven

by intrinsic disorder within N1 and trace protease contamination during recombinant

protein preparation (25, 47, 67). Our *in vivo* experiments were conducted exclusively

with a truncated ClfA variant as we found that murine plasmin was inefficient at

cleaving ClfA (Fig. S2). This confirming that the observed phenotypes stem from loss

of ClfA function rather than activity of the released fragment. However future studies

should investigate if the released portion, or peptides derived from it, have biological

activity.

545   Funding information

M.B.T. was supported by a Postgraduate Ussher Fellowship Award from Trinity

College Dublin. J.A.G. is supported by a Royal Society Wolfson Fellowship (Award

193002). Z.H. and T.J. are supported by Swedish Medical Research Council 2024-

02667 and the Swedish state under the agreement between the Swedish

Government and the county councils ALFGBG-1005076. E.C.L. and R.W.W.

received support from Science Foundation Ireland Frontiers for the Future award

19/FFP/6484. R.W.W. received support from the Irish Research Council

GOIPG/2019/4236. The funders were not involved with the design of this study,

collection or analysis of data, decision to publish or preparation of the manuscript.

556   Acknowledgements

We gratefully acknowledge the contribution to this study made by the University of

Birmingham's Human Biomaterials Resource Centre which has been supported

through Birmingham Science City - Experimental Medicine Network of Excellence

project. We thank Deirdre Muldowney and Kevin Lyons for technical assistance.

Conflict of Interest

The authors declare there are no conflicts of interest of a financial or any other

nature.

FIG. 1. Cell wall-associated ClfA is cleaved by human plasmin. Stationary phase

cultures of *S. aureus* strains Newman, Newman *clfA5*, BH1CC, LAC and 14-2533T

were adjusted to an OD_{600} = 10 and incubated at 37°C for 1 h with and without

plasmin (± P, 0.075 U/mL) and alpha-2 macroglobulin (+ α2M, 1 mg/mL) before cell

wall proteins were extracted and separated by SDS-PAGE and probed with anti-ClfA

A domain polyclonal IgG (1:1000) in a western blot. Bound primary antibodies was

detected with protein A peroxidase (1:500). Blots presented are representative of two

independent experiments.

FIG. 2. The ClfA N1N2N3 domains are cleaved by human plasmin. $ClfA_{40-559*}$ (5

μM) was incubated in PBS with or without added human plasmin (± P, 0.07 U/mL)

and alpha-2 macroglobulin (+ α2M, 1 mg/mL), where indicated, at 37°C for 1 h (A-D).

Alternatively, $ClfA_{40-559*}$ (5 μM) was incubated in human serum (40%) that was

pretreated with either E) tissue plasminogen activator (± tPA, 25 μg/mL) or F)

concentrated supernatants (s/n) from *S. aureus* strain LAC* or its *sak*-deficient

mutant (*sak*::Tn) for 20 min at 37°C. Serum only controls did not include $ClfA_{40-559*}$.

Samples were separated by SDS-PAGE and either stained with SimplyBlue

Coomassie stain (A) or probed in a western immunoblot with anti-ClfA IgG (1:1000;

B, E, F), anti-His 7E8 IgG (1:1000; C) or anti-Strep Tag II IgG-HRP (1:2500; D).

Bound anti-ClfA IgG was detected with protein A peroxidase (1:500) and bound anti-

His IgG was detected with HRP-conjugated rabbit anti-mouse IgG (1:2500). Blots

represent two independent experiments.

FIG. 3. Plasmin cleaves ClfA between residues 214 and 215. A) Domain organisation of ClfA. The amino acid coordinates of the signal peptide (S) and N1N2N3 domains are indicated. The SD repeat region and wall (W) and membranes-spanning (M) regions are shown. The 7 most N-terminal amino-acids of the plasmin-cleaved product identified using Edman degradation are indicated. Sequence identity is indicated in green with red arrows highlighting matches to the ClfA sequence. Arg214 predicted to be cleaved by plasmin is underlined (R). B) A fluorescently-labelled 11-mer peptide (WT) encompassing the region around the proposed cleavage site at Arg214 (residues 211-221) or variants P211S, R212S, M213S, R214S, A215S and R216S harbouring single amino-acid substitutions with serine (37.5 μM) were incubated in the absence (-) or presence (+) of plasmin (0.075 U/mL) for 30 min at 37°C. Peptide S^c was a scrambled version of the WT peptide and WT^R a peptide with four substitutions (P211A, R212S, M213A, R214S). Samples were analysed by agarose (4% w/v) gel electrophoresis. Gels represent three independent experiments. C) Recombinant $ClfA_{40-559^*}$ and $ClfA_{40-559^*}$ (5 μM)

609   were incubated with or without plasmin (± P, 0.075 U/mL) at 37°C for 1 h and

610   analysed by SDS-PAGE. Gels were stained with Instant Blue Coomassie stain. The

611   gel image presented is representative of two independent experiments.

FIG. 4. Plasmin cleavage disrupts ClfA-mediated clumping in fibrinogen. Stationary phase cultures of *S. aureus* ClfA$^+$(A) and ClfA☐ (B) were adjusted to OD_{600} = 1.5 in PBS and incubated in the absence and presence of plasmin (± P, 0.075 U/mL) for 1 h at 37°C. Bacteria were washed and incubated statically at 37°C for 105 min in the presence of fibrinogen (18.4 µg/mL). *S. aureus* ClfA$^{Δ40-214}$ (C) ClfA$^+$ (D) ClfA☐ (E) and ClfA$^-$ (F) were incubated in PBS with (+) and without (-) fibrinogen (18.4 µg/mL) for 105 min at 37°C. The OD_{600} after 105 min is expressed as a percentage of the original input OD_{600} at 0 h. Each symbol represents the % of input OD_{600} from a single matched experiment and the bars represent the mean value for each condition from four independent experiments. Error bars represent standard deviation, and statistical significance was determined by an unpaired student t-test. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ns, non-significant.

FIG. 5. Truncation of ClfA enhances ClfA-mediated survival in human blood.

Stationary phase cultures of *S. aureus* ClfA⁺, ClfA□ and ClfA$^{\Delta40\text{-}214}$ were inoculated

(ca. 2.5 x 10□ CFU/mL) into freshly drawn human blood and incubated for 3 h at

37°C. Viable counts after 3 h are expressed as a percentage of the original input

CFU at 0 h. Each symbol represents the % of input CFU recovered from a single

experiment and the bars represent the mean value for each condition from three

independent experiments. Error bars represent standard deviation, and statistical

significance was determined by one-way Anova following Tukey's multi-comparison

test. Where significant P is represented: ns. $P > 0.05$; *, $P < 0.05$; **, $P < 0.01$.

642
FIG. 6. Truncation of ClfA increases kidney abscess formation by *S. aureus*. *S.*

aureus ClfA[+], ClfA[R] and ClfA$^{\Delta40-214}$ were intravenously administered to NMRI mice (10

per group) via the tail vein with 200 µL of PBS containing 75 ng/mL

anhydrotetracycline. Expression of ClfA was induced via an intraperitoneal injection

of inducer anhydrotetracycline (5 µg/mouse) on Day 0, 1 h before infection and 1h

post-infection. Kidneys and livers were collected aseptically, and kidney abscess

scores (A) were assessed in a blinded manner; 0 represented healthy kidneys; 1

indicated one to two small abscesses without structural changes; 2 denoted more

than two abscesses affecting less than 75% of the kidney tissue; and 3 indicated

extensive abscess formation involving more than 75% of the tissue. Bacterial load in

the kidney (B) and liver (C) were quantified by homogenising the organs and

culturing on horse blood agar. (D) Mouse survival after infection. Percent survival of

mice (n = 20 per group) after infection with 6 x 10^6 CFU/mouse. The dashed line

colour pairs to symbol colour for strains in panel A-C. Error bars represent 95%

confidence intervals and statistical significance was determined by one-way Anova

multiple comparisons with the Kruskal-Wallis post-test. *P < 0.05; **, P < 0.01. No

symbol P > 0.05.

References

1. Tong SYC, Davis JS, Eichenberger E, Holland TL, Fowler VG. *Staphylococcus aureus* Infections: Epidemiology, Pathophysiology, Clinical Manifestations, and Management. Clinical microbiology reviews. 2015;28(3):603-61.

2. Naghavi M, Vollset SE, Ikuta KS, Swetschinski LR, Gray AP, Wool EE, et al. Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. The Lancet. 2024;404(10459):1199-226.

3. Bai AD, Lo CKL, Komorowski AS, Suresh M, Guo K, Garg A, et al. *Staphylococcus aureus* bacteraemia mortality: a systematic review and meta-analysis. Clinical Microbiology and Infection. 2022;28(8):1076-84.

4. McAdow M, Kim HK, DeDent AC, Hendrickx APA, Schneewind O, Missiakas DM. Preventing Staphylococcus aureus Sepsis through the Inhibition of Its Agglutination in Blood. PLoS pathogens. 2011;7(10):e1002307.

5. Vilacosta I, Graupner C, San Román JA, Sarriá C, Ronderos R, Fernández C, et al. Risk of embolization after institution of antibiotic therapy for infective endocarditis. J Am Coll Cardiol. 2002;39(9):1489-95.

6. Nappi F, Spadaccio C, Moon MR. A management framework for left sided endocarditis: a narrative review. Ann Transl Med. 2020;8(23):1627.

7. Avtaar Singh SS, Costantino MF, D'Addeo G, Cardinale D, Fiorilli R, Nappi F. A narrative review of diagnosis of infective endocarditis-imaging methods and comparison. Ann Transl Med. 2020;8(23):1621.

8. Nappi F, Martuscelli G, Bellomo F, Avtaar Singh SS, Moon MR. Infective Endocarditis in High-Income Countries. Metabolites. 2022;12(8).

9.   Moreillon P, Entenza JM, Francioli P, McDevitt D, Foster TJ, François P, et al.

Role of *Staphylococcus aureus* coagulase and clumping factor in pathogenesis of

experimental endocarditis. Infection and immunity. 1995;63(12):4738-43.

10.   Hair PS, Echague CG, Sholl AM, Watkins JA, Geoghegan JA, Foster TJ, et al.

Clumping factor A interaction with complement factor I increases C3b cleavage on

the bacterial surface of *Staphylococcus aureus* and decreases complement-

mediated phagocytosis. Infection and immunity. 2010;78(4):1717-27.

11.   Macrae FL, Duval C, Papareddy P, Baker SR, Yuldasheva N, Kearney KJ, et

al. A fibrin biofilm covers blood clots and protects from microbial invasion. The

Journal of Clinical Investigation. 2018;128(8):3356-68.

12.   Negrón O, Flick MJ. Does fibrinogen serve the host or the microbe in

Staphylococcus infection? Current Opinion in Hematology. 2019;26(1531-7048

(Electronic)):343-8.

13.   Forsyth CB, Solovjov DA, Ugarova TP, Plow EF. Integrin αMβ2-Mediated Cell

Migration to Fibrinogen and Its Recognition Peptides. Journal of Experimental

Medicine. 2001;193(10):1123-34.

14.   Rubel C, Fernández GC, Rosa FA, Gómez S, Bompadre MB, Coso OA, et al.

Soluble Fibrinogen Modulates Neutrophil Functionality Through the Activation of an

Extracellular Signal-Regulated Kinase-Dependent Pathway1. The Journal of

Immunology. 2002;168(7):3527-35.

15.   Smiley ST, King JA, Hancock WW. Fibrinogen Stimulates Macrophage

Chemokine Secretion Through Toll-Like Receptor 41. The Journal of Immunology.

2001;167(5):2887-94.

16.   Negrón O, Hur WS, Prasad J, Paul DS, Rowe SE, Degen JL, et al.

Fibrin(ogen) engagement of *S. aureus* promotes the host antimicrobial response and

suppression of microbe dissemination following peritoneal infection. PLoS
pathogens. 2022;18(1):e1010227.

17.   McDevitt D, Nanavaty T, House-Pompeo K, Bell E, Turner N, McIntire L, et al.
Characterization of the interaction between the *Staphylococcus aureus* clumping
factor (ClfA) and fibrinogen. Eur J Biochem. 1997;247(1):416-24.

18.   Ganesh VK, Rivera JJ, Smeds E, Ko Y-P, Bowden MG, Wann ER, et al. A
structural model of the *Staphylococcus aureus* ClfA-fibrinogen interaction opens new
avenues for the design of anti-staphylococcal therapeutics. PLoS pathogens.
2008;4(11):e1000226-e.

19.   Ponnuraj K, Bowden MG, Davis S, Gurusiddappa S, Moore D, Choe D, et al.
A "dock, lock, and latch" Structural Model for a Staphylococcal Adhesin Binding to
Fibrinogen. Cell. 2003;115(2):217-28.

20.   Ganesh VK, Liang X, Geoghegan JA, Cohen ALV, Venugopalan N, Foster TJ,
et al. Lessons from the Crystal Structure of the *S. aureus* Surface Protein Clumping
Factor A in Complex With Tefibazumab, an Inhibiting Monoclonal Antibody.
EBioMedicine. 2016;13:328-38.

21.   Herman-Bausier P, Labate C, Towell AM, Derclaye S, Geoghegan JA,
Dufrêne YF. *Staphylococcus aureus* clumping factor A is a force-sensitive molecular
switch that activates bacterial adhesion. Proceedings of the National Academy of
Sciences of the United States of America. 2018;115(21):5564-9.

22.   Hair PS, Ward MD, Semmes OJ, Foster TJ, Cunnion KM. *Staphylococcus
aureus* clumping factor A binds to complement regulator factor I and increases factor
I cleavage of C3b. The Journal of infectious diseases. 2008;198(1):125-33.

23.   Geoghegan JA, Monk IR, O'Gara JP, Foster TJ. Subdomains N2N3 of
fibronectin binding protein A mediate *Staphylococcus aureus* biofilm formation and

adherence to fibrinogen using distinct mechanisms. Journal of bacteriology.

2013;195(11):2675-83.

24.   Hildén P, Savolainen K, Tyynelä J, Vuento M, Kuusela P. Purification and

characterisation of a plasmin-sensitive surface protein of *Staphylococcus aureus*.

Eur J Biochem. 1996;236(3):904-10.

25.   Crosby Heidi A, Keim K, Kwiecinski Jakub M, Langouët-Astrié Christophe J,

Oshima K, LaRivière Wells B, et al. Host-derived protease promotes aggregation of

Staphylococcus   aureus   by   cleaving   the   surface   protein   SasG.   mBio.

2024;15(4):e03483-23.

26.   Baker SK, Strickland S. A critical role for plasminogen in inflammation.

Journal of Experimental Medicine. 2020;217(4).

27.   Vago JP, Zaidan I, Perucci LO, Brito LF, Teixeira LCR, Silva CMS, et al.

Plasmin   and   plasminogen   prevent   sepsis   severity   by   reducing   neutrophil

extracellular traps and systemic inflammation. JCI Insight. 2023;8(8).

28.   Bokarewa MI, Jin T, Tarkowski A. Staphylococcus aureus: Staphylokinase.

The International Journal of Biochemistry & Cell Biology. 2006;38(4):504-9.

29.   O'Halloran Dara P, Wynne K, Geoghegan Joan A. Protein A Is Released into

the *Staphylococcus aureus* Culture Supernatant with an Unprocessed Sorting

Signal. Infection and immunity. 2015;83(4):1598-609.

30.   Josefsson E, Hartford O, O'Brien L, Patti JM, Foster T. Protection against

Experimental *Staphylococcus aureus* Arthritis by Vaccination with Clumping Factor

A,   a   Novel   Virulence   Determinant.   The   Journal   of   infectious   diseases.

2001;184(12):1572-80.

31.   Visai L, Xu Y Fau - Casolini F, Casolini F Fau - Rindi S, Rindi S Fau - Höök M,

Höök M Fau - Speziale P, Speziale P. Monoclonal antibodies to CNA, a collagen-

binding microbial surface component recognizing adhesive matrix molecules, detach

Staphylococcus aureus from a collagen substrate. 2000(0021-9258 (Print)).

32.   O'Neill E, Pozzi C, Houston P, Humphreys H, Robinson DA, Loughman A, et

al. A novel *Staphylococcus aureus* biofilm phenotype mediated by the fibronectin-

binding proteins, FnBPA and FnBPB. Journal of bacteriology. 2008;190(11):3835-50.

33.   Duthie Es Fau - Lorenz LL, Lorenz LL. Staphylococcal coagulase; mode of

action and antigenicity. (0022-1287 (Print)).

34.   Fitzgerald JR, Loughman A, Keane F, Brennan M, Knobel M, Higgins J, et al.

Fibronectin-binding proteins of *Staphylococcus aureus* mediate activation of human

platelets via fibrinogen and fibronectin bridges to integrin GPIIb/IIIa and IgG binding

to the FcγRIIa receptor. Molecular microbiology. 2006;59(1):212-30.

35.   Boles BR, Thoendel M, Roth AJ, Horswill AR. Identification of Genes Involved

in Polysaccharide-Independent *Staphylococcus aureus* Biofilm Formation. PloS one.

2010;5(4):e10146.

36.   Pietrocola G, Nobile G, Rindi S, Speziale P. *Staphylococcus aureus*

Manipulates Innate Immunity through Own and Host-Expressed Proteases. Frontiers

in cellular and infection microbiology. 2017;7:166.

37.   Zapotoczna M, Riboldi GP, Moustafa AM, Dickson E, Narechania A,

Morrissey JA, et al. Mobile-Genetic-Element-Encoded Hypertolerance to Copper

Protects *Staphylococcus aureus* from Killing by Host Phagocytes. mBio. 2018;9(5).

38.   Monk IR, Tree JJ, Howden BP, Stinear TP, Foster TJ. Complete Bypass of

Restriction Systems for Major *Staphylococcus aureus* Lineages. mBio.

2015;6(3):e00308-15.

39.   Corrigan RM, Miajlovic H, Foster TJ. Surface proteins that promote adherence

of *Staphylococcus aureus* to human desquamated nasal epithelial cells. BMC

microbiology. 2009;9:22.

40.   McCormack N, Foster TJ, Geoghegan JA. A short sequence within

subdomain N1 of region A of the *Staphylococcus aureus* MSCRAMM clumping factor

A is required for export and surface display. Microbiology (Reading, England).

2014;160(4):659-70.

41.   Li MZ, Elledge SJ. SLIC: a method for sequence- and ligation-independent

cloning. Methods Mol Biol. 2012;852:51-9.

42.   Dagert M, Ehrlich SD. Prolonged incubation in calcium chloride improves the

competence of *Escherichia coli* cells. Gene. 1979;6(1):23-8.

43.   Löfblom J, Kronqvist N, Uhlén M, Ståhl S, Wernérus H. Optimization of

electroporation-mediated transformation: *Staphylococcus carnosus* as model

organism. Journal of applied microbiology. 2007;102(3):736-47.

44.   Mohammad M, Nguyen M-T, Engdahl C, Na M, Jarneborn A, Hu Z, et al. The

YIN and YANG of lipoproteins in developing and preventing infectious arthritis by

Staphylococcus aureus. PLoS pathogens. 2019;15(6):e1007877.

45.   Jin T. Exploring the role of bacterial virulence factors and host elements in

septic arthritis: insights from animal models for innovative therapies. Frontiers in

microbiology. 2024;Volume 15 - 2024.

46.   Ali A, Welin A, Schwarze J-C, Svensson MND, Na M, Jarneborn A, et al.

CTLA4 Immunoglobulin but Not Anti–Tumor Necrosis Factor Therapy Promotes

Staphylococcal Septic Arthritis in Mice. The Journal of infectious diseases.

2015;212(8):1308-16.

47.   McAleese FM, Walsh EJ, Sieprawska M, Potempa J, Foster TJ. Loss of

Clumping Factor B Fibrinogen Binding Activity by *Staphylococcus aureus* Involves

Cessation of Transcription, Shedding and Cleavage by Metalloprotease*. Journal of

Biological Chemistry. 2001;276(32):29969-78.

48.   Perkins S, Walsh EJ, Deivanayagam CCS, Narayana SVL, Foster TJ, Höök

M. Structural Organization of the Fibrinogen-binding Region of the Clumping Factor

B MSCRAMM of *Staphylococcus aureus* *. Journal of Biological Chemistry.

2001;276(48):44721-8.

49.   Higgins J, Loughman A, Van Kessel KPM, Van Strijp JAG, Foster TJ.

Clumping factor A of *Staphylococcus aureus* inhibits phagocytosis by human

polymorphonuclear leucocytes. FEMS microbiology letters. 2006;258(2):290-6.

50.   Flick MJ, Du X, Prasad JM, Raghu H, Palumbo JS, Smeds E, et al. Genetic

elimination of the binding motif on fibrinogen for the *S. aureus* virulence factor ClfA

improves host survival in septicemia. Blood. 2013;121(10):1783-94.

51.   Malachowa N, Kobayashi SD, Porter AR, Braughton KR, Scott DP, Gardner

DJ, et al. Contribution of *Staphylococcus aureus* Coagulases and Clumping Factor A

to Abscess Formation in a Rabbit Model of Skin and Soft Tissue Infection. PloS one.

2016;11(1932-6203 (Electronic)).

52.   Cheng AG, Kim Hk Fau - Burts ML, Burts Ml Fau - Krausz T, Krausz T Fau -

Schneewind O, Schneewind O Fau - Missiakas DM, Missiakas DM. Genetic

requirements for *Staphylococcus aureus* abscess formation and persistence in host

tissues. PNAS. 2009;115(1530-6860 (Electronic)):5564-9.

53.   Sully EK, Malachowa N, Elmore BO, Alexander SM, Femling JK, Gray BM, et

al. Selective Chemical Inhibition of agr Quorum Sensing in *Staphylococcus aureus*

Promotes Host Defense with Minimal Impact on Resistance. PLoS pathogens. 2014;10(6):e1004174.

54.   Josefsson E, Higgins J, Foster TJ, Tarkowski A. Fibrinogen Binding Sites P336 and Y338 of Clumping Factor A Are Crucial for *Staphylococcus aureus* Virulence. PloS one. 2008;3(5):e2206.

55.   Ko YP, Flick MJ. Fibrinogen Is at the Interface of Host Defense and Pathogen Virulence in *Staphylococcus aureus* Infection. 2016(1098-9064 (Electronic)).

56.   Prasad JM, Negrón O, Du X, Mullins ES, Palumbo JS, Gilbertie JM, et al. Host fibrinogen drives antimicrobial function in *Staphylococcus aureus* peritonitis through bacterial-mediated prothrombin activation. Proceedings of the National Academy of Sciences. 2021;118(1):e2009837118.

57.   Peetermans M, Vanassche T, Liesenborghs L, Claes J, Vande Velde G, Kwiecinksi J, et al. Plasminogen activation by staphylokinase enhances local spreading of *S. aureus* in skin infections. BMC microbiology. 2014;14:310-.

58.   Kwiecinski J, Jacobsson G, Karlsson M, Zhu X, Wang W, Bremell T, et al. Staphylokinase Promotes the Establishment of Staphylococcus aureus Skin Infections While Decreasing Disease Severity. The Journal of infectious diseases. 2013;208(6):990-9.

59.   Loughman A, Fitzgerald JR, Brennan MP, Higgins J, Downer R, Cox D, et al. Roles for fibrinogen, immunoglobulin and complement in platelet activation promoted by *Staphylococcus aureus* clumping factor A. Molecular microbiology. 2005;57(3):804-18.

60.   Panizzi P, Nahrendorf M, Figueiredo J-L, Panizzi J, Marinelli B, Iwamoto Y, et al. In vivo detection of *Staphylococcus aureus* endocarditis by targeting pathogen-specific prothrombin activation. Nature Medicine. 2011;17(9):1142-6.

61. Palmqvist N, Josefsson E, Tarkowski A. Clumping factor A-mediated virulence

during *Staphylococcus aureus* infection is retained despite fibrinogen depletion.

Microbes Infect. 2004;6(2):196-201.

62. McGavin MJ, Zahradka C, Rice K, Scott JE. Modification of the

Staphylococcus aureus fibronectin binding phenotype by V8 protease. Infection and

immunity. 1997;65(7):2621-8.

63. Karlsson A, Saravia-Otten P, Tegmark K, Morfeldt E, Arvidson S. Decreased

Amounts of Cell Wall-Associated Protein A and Fibronectin-Binding Proteins in

Staphylococcus aureus sarA Mutants due to Up-Regulation of Extracellular

Proteases. Infection and immunity. 2001;69(8):4742-8.

64. Geoghegan JA, Monk IR, Gara JP, Foster TJ. Subdomains N2N3 of

Fibronectin Binding Protein A Mediate *Staphylococcus aureus* Biofilm Formation and

Adherence to Fibrinogen Using Distinct Mechanisms. Journal of bacteriology.

2013;195(11):2675.

65. Savolainen K, Paulin L, Westerlund-Wikström B, Foster TJ, Korhonen TK,

Kuusela P. Expression of pls, a gene closely associated with the mecA gene of

methicillin-resistant Staphylococcus aureus, prevents bacterial adhesion in vitro.

Infection and immunity. 2001;69(5):3013-20.

66. Hervio LS, Coombs GS, Bergstrom RC, Trivedi K, Corey DR, Madison EL.

Negative selectivity and the evolution of protease cascades: the specificity of

plasmin for peptide and protein substrates. Chemistry & biology. 2000;7(6):443-52.

67. Barbu EM, Ganesh VK, Gurusiddappa S, Mackenzie RC, Foster TJ, Sudhof

TC, et al. β-Neurexin Is a Ligand for the *Staphylococcus aureus* MSCRAMM SdrC.

PLoS pathogens. 2010;6(1):e1000726.