Distinct functional potential of bacterial extracellular vesicles across biogeographic provinces of the South Pacific Ocean

Eduard Fadeev[1,*], Neza Orel[2], Tinkara Tinta[2], Leila Afjehi-Sadat[3], Haoran Liu[4], Thomas J. Browning[4], Zhongwei Yuan[4], Eric P. Achterberg[4], Steven J. Biller[5], Daniel J. Sher[6], and Gerhard J. Herndl[1,3,7]

[1] Department of Functional and Evolutionary Ecology, Bio-Oceanography and Marine Biology Unit, University of Vienna, Vienna, Austria

[2] Marine Biology Station Piran, National Institute of Biology, Piran, Slovenia

[3] Mass Spectrometry Unit, Research Support Facilities, University of Vienna, Vienna, Austria

[4] GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany

[5] Department of Biological Sciences, Wellesley College, Wellesley, USA

[6] Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Haifa, Israel

[7] NIOZ, Department of Marine Microbiology and Biogeochemistry, Royal Netherlands Institute for Sea Research, Den Burg, The Netherlands

*Corresponding author

Keywords: bacterioplankton, membrane vesicles, horizontal gene transfer, siderophores, extracellular enzymes

Abstract

Bacterial extracellular vesicles (BEVs) are spherical membrane-bound structures released by bacterial cells that can carry various biological compounds. Experimental investigation of BEVs in marine bacteria, largely limited to laboratory model systems, have suggested putative roles in bacterial ecophysiology. Yet the functions of BEVs in their ecological context has never been explored directly in the ocean. Here we report the results a large-scale oceanographic survey of marine BEVs in surface waters, conducted over more than 5,000 nautical miles in the South Pacific, linking their protein cargoes to the bacterial communities producing them. The presence of marine BEVs was consistently observed across a range of biogeochemical conditions, with an overall abundance comparable to that of bacterial cells (ca. 10^8 BEVs L^{-1}). The protein cargoes of marine BEVs differed significantly between environmental conditions, being enriched in carbohydrate transporters and hydrolytic enzymes under phytoplankton bloom conditions, and enriched in iron uptake-related proteins in nutrient-limited waters, suggesting key extracellular functions they could perform in marine bacterial communities. Our observations highlight the ubiquity of marine BEVs and their potential functions across oceanic scales, with implications for our general understanding of key processes in marine microbial ecology.

Introduction

Bacterial extracellular vesicles (BEVs) are spherical membranous structures typically ranging between 20 and 250 nm in diameter that can carry various biological cargoes (e.g., nucleic acids and proteins) over long distances in a protected and locally concentrated manner [1]. Since their discovery in human-associated *Escherichia coli* in the 1960s [2], most research on BEVs has focused on pathogens and other host-associated bacteria [3]. Although it remains unclear to what extent the release of BEVs is an active cellular process or simply a by-product of cell growth or death (i.e., lysis), a variety of BEV functions have been described to date. One of the best-documented functions of BEVs is the intercellular transfer of DNA molecules [4,5], which can facilitate the exchange of antibiotic resistance genes between different bacterial lineages [6], as well as the transfer of entire plasmids [7,8], recently termed 'vesiduction' [9]. In addition to DNA transfer, BEVs can also play a role in bacterial quorum sensing [10,11], biofilm formation [12], and contribute to microbial "warfare", where they deliver compounds that kill competing bacteria [13,14].

Because BEVs contain fragments of the releasing bacterial cell, as well as cell membrane-derived enzymes, receptors and transporters, it has been proposed that BEVs may have a variety of functions in the exocellular milieu [15,16] in addition to intercellular interactions, which may be particularly relevant for bacteria living in aqueous environments [3,17]. The ocean - the largest aquatic ecosystem on Earth where bacteria are among the most abundant organisms [18] - is a highly dynamic environment both physically and biochemically. The surface waters of the ocean are characterised by a depletion of bioavailable nutrients required for life in some areas (e.g., oceanic gyres) [19] and a high availability in others (e.g., coastal upwelling regions) [20,21]. To survive in such dynamic ecosystems, marine bacteria have developed a wide range of ecophysiological capabilities [22–24], and implement highly diverse nutrient and energy utilisation strategies [25–27]. In the interaction between marine bacteria and their nutrient depleted environment, high cellular surface area to volume ratios provide optimal conditions for nutrient uptake [28,29]. In this sense, marine BEVs can be conceptualised as an expansion of the cell surface [1] that diffuses through seawater and randomly encounters biotic and abiotic components. Existing laboratory-based observations of several isolates of globally abundant marine bacteria [30–35] suggest that BEVs may mediate specific extracellular functions in the ecophysiology of marine bacteria, such as binding specific elements (e.g., iron or phosphate) and delivering them in concentrated form to bacterial cells. However, despite the apparent ubiquity of BEVs in the marine environment [36,37] and their potential ability to impact bacterial functions, how the contents and functional potential of these nanoparticles might vary across broad oceanographic contexts is unknown.

In this study we determined the abundance and protein cargo diversity of BEVs across the surface waters of the South Pacific Ocean. By combining molecular characterisation of marine BEVs with their local microbial and biogeochemical context, we show that BEV contents vary among ocean regions and may contribute to acquisition of limiting nutrients by the local microbial community. Our observations across a large oceanic basin reinforce the hypothesis that important extracellular bacterial functions may be mediated by BEVs in the global ocean.

Results and Discussion

Distinct biogeochemical conditions across the South Pacific Ocean

To investigate the potential ecophysiological functions of BEVs in marine bacterial communities we sampled across diverse environmental conditions, varying in nutrient limitations and phytoplankton bloom conditions. We collected surface water (5-6 m depth) at 20 stations along a transect that spanned the South Pacific Ocean (Fig. 1A; Table S1). Detailed physicochemical characterization across this transect [38], showed that the field sampling was carried out across four oceanic provinces. Briefly, the easternmost region was the Chilean coastal upwelling zone (UP), characterised by elevated nitrate and phosphate concentrations, while depleted in dissolved iron. In this area there were elevated chlorophyll-a concentrations (ca. 3 mg m^{-3}; Fig. S1), with the phytoplankton community mostly composed of diatoms (Fig. 1B). The transition zone between the upwelling and the subtropical gyre (TRAN) was

characterised by elevated concentrations of phosphate and dissolved iron, but depleted in nitrate. This
area also had relatively high chlorophyll-a concentrations (0.04-0.26 mg m^{-3}), with mixed
phytoplankton communities. The centre of the subtropical gyre (GYRE) was characterised by very low
nitrate concentrations and a declining spatial gradient from east to west of phosphate concentrations,
with very low chlorophyll-a concentrations (< 0.03 mg m^{-3}). The westernmost region (WEST) was
defined as the area of the subtropical gyre with the lowest phosphate concentrations (< 10 nM) and
higher (compared to GYRE) chlorophyll-a concentrations (0.03-0.14 mg m^{-3}) of predominantly
cyanobacterial origins.

Fig. 1 – Diverse biogeochemical regimes encountered across the South Pacific Ocean. (A) Spatial grouping of
the sampling stations according to the four oceanic provinces: 'UP' - Chilean coastal upwelling zone; 'TRAN' -
transition zone between the upwelling and the subtropical gyre; 'GYRE' - South Pacific subtropical gyre; 'WEST'
- westernmost region. Chl. a concentration is overlaid on the cruise transect, with the grey scale background
representing the bathymetry.(B) The composition of the surface water phytoplankton at each station, estimated
on the basis of diagnostic pigments.

We found that bacterial abundance (analysed using flow cytometry) varied between the biogeochemical
provinces, and the taxonomic composition of the bacterial communities (based on metagenomic
characterisation) also differed significantly between them (Fig. 2; PERMANOVA test; $F_{3,14}$=21.16,
R^2=0.82, p=0.001). To investigate the physiological and metabolic processes expressed within the
bacterial communities, we characterised the proteome of the bacterial cells, identifying between 4504-

6532 cellular proteins at each station (Table S1). The composition of bacterial cellular proteins was significantly different between the four regions (Fig. S2; PERMANOVA test; $F_{3,16}=2.85$, $R^2=0.35$, p=0.001). To test for the differences between the physiological and metabolic processes in bacterial communities in the different provinces, we performed differential protein abundance analyses between each pair of adjacent provinces (Table S2). The bacterial abundance was strongly elevated in the upwelling zone ($9\text{-}26\times10^8$ cells L^{-1}), with the bacterial community showing a strong heterotrophic response to the availability of organic carbon as a result of the diatom bloom. The dominant bacteria were *Flavobacteriales* and *Cellvibrionales*, which are typically active during phytoplankton blooms [39,40]. In the transition zone, bacterial abundance reached $0.4\text{-}3.5\times10^8$ cells L^{-1}, with an elevated photosynthetic activity of the cyanobacterial order *Synechococcales* (*Prochlorococcaceae* and *Synechococcaceae*), which was probably stimulated by a nutrient influx from a returning lateral current [38]. The subtropical gyre exhibited the lowest bacterial abundances ($0.2\text{-}0.9\times10^8$ cells L^{-1}), characterised by the dominance of the oligotrophic order *Pelagibacterales* [41], which has previously been shown to numerically dominate the bacterial communities of the ultra-oligotrophic South Pacific subtropical gyre [42,43]. In the westernmost region, bacterial abundance was significantly higher than in the subtropical gyre ($0.8\text{-}1.3\times10^8$ cells L^{-1}; Student's t-test, p=0.002), characterised by increased activity of *Prochlorococcaceae*, which may have been partly facilitated by micronutrient input from a large volcanic eruption in the region two months prior to sampling [44].

Fig. 2 – Bacterial community composition based on taxonomic classification of metagenomic sequences. *'Other classes < 0.5%' – refers to taxonomic classes that comprised less than 0.5% of the sequences; 'unclassified' – refers to bacterial sequences with unknown classification on a class level. All samples contained $2\text{-}5\times10^8$ classified sequences.*

Abundance and characteristics of marine BEVs

To investigate the abundance and diversity of marine BEVs, we sampled the extracellular size fraction corresponding to BEV-like structures (100 kDa-0.22 µm) across all stations using size fractionated filtration. To some extent, marine viruses are in the same size fraction as BEVs, and they can be a major obstacle to studying marine BEVs due to our limited ability to mechanically separate them [45–47]. Therefore, we used density gradient separation to further enrich each sample for BEVs in the laboratory (see Materials and Methods). At least 50% of the purified BEVs were in the 70 to 130 nm diameter range, matching the size distributions reported for cultures of marine bacteria (Fig. S3)[33,35]. The abundance of BEVs was in the same order of magnitude as bacterial cells (Fig. 3). However, because the density gradient separation resulted in a reduced yield of 6-52% of all nanoparticles (Fig. S4), these are likely conservative estimates of the BEVs concentration in the seawater. We then characterised the

protein cargoes of the BEV-like structures and identified 2029-3580 BEV-associated proteins at each
station (Table S1). Of these proteins only 56-121 were taxonomically or functionally (e.g., capsid or
tail sheath) related to viruses, suggesting a strong enrichment of BEVs in our samples. We found that
78-96% of proteins observed in the BEVs were also found in the corresponding cellular fraction at each
station, but the composition of the proteins significantly varied between the two fractions
(PERMANOVA test; $F_{1,38}$=15.00, R^2=0.28, p =0.001). This indicates that the BEVs represent a subset
that differs from the full bacterial proteome.

146
Fig. 3 – Bacterial cell and EVs abundances across different provinces. *The abundances were*
determined by flow cytometry and nanoparticle tracking analysis (NTA), respectively. The N represents
the number of stations in each province.

Gram-negative marine bacteria can release BEVs through blebbing of the cellular membrane, forming
particles bounded by fragments of cellular membranes and containing periplasm of the releasing
bacterial cell [1,48]. In addition to biases in packaging relating to subcellular localization, external factors
have also been shown to have an impact on the production of BEVs and their contents [49–52]. Other BEVs
may result from bacterial cell lysis due to bacteriophages, and thus represent a pseudo-random
assemblage of fragments from different subcellular origins [3,53]. To differentiate between these different
origins of BEVs, we assessed whether there is an enrichment in non-cytoplasmic proteins in the BEV
fraction. Overall, we identified 1,121 proteins that were significantly enriched in BEVs compared to
the cellular fraction across the dataset (log_2 fold change >1, adjusted p-value < 0.1; Table S3). Prediction
of sub-cellular origins revealed that half of the BEV-enriched proteins were non-cytoplasmic,
comprising mostly outer membrane and extracellular proteins, compared to ca. 25% of the enriched
proteins in the cellular fraction (Fig. 4). Although we cannot specifically identify their production
process in our dataset, our results suggest that the packaging of marine BEVs is strongly biased towards
proteins of non-cytoplasmic origin.

*Fig. 4 – **Proportions of significantly enriched proteins between BEV and cellular fractions.** Proportion of enriched proteins in each fraction according to their predicted sub-cellular origins (A) and taxonomic affiliation (B). The total number of enriched proteins in the BEVs and the cellular fractions was 1121 and 2075, respectively. Taxonomic affiliations of enriched proteins in the cellular and BEVs fractions according to different oceanic provinces. The taxonomic class of each order is given in parentheses: 'A' – Alphaproteobacterial, 'C' – Cytophagia, 'CY' – Cyanophyceae, 'F' – Flavobacteriia, 'G' – Gammaproteobacteria.*

The enrichment analyses revealed that proteins associated with the cyanobacterial order *Synechococcales* (families *Prochlorococcaceae* and *Synechococcaceae*) made up the largest number of enriched proteins in the BEVs and cellular fractions, proportionally accounting for 17% and 22% of proteins, respectively (Fig. 4). The second largest taxonomic group was the order *Pelagibacterales* (the SAR11 clade of Alphaproteobacteria), which accounted for 9% of BEV-enriched proteins while comprising 18% of enriched proteins in the cellular fraction. In contrast, enriched proteins associated with other bacterial orders, such as *Flavobacteriales* (mostly the family *Flavobacteriaceae*), *Alteromonadales* (mostly the families *Alteromonadaceae* and *Pseudoalteromonadaceae*), and *Cellvibrionales* (mostly the families *Halieaceae* and *Cellvibrionaceae*), accounted for more than twice the relative amount of BEV-enriched proteins compared to the cellular fraction. A previous comparison of isolates from the major classes of marine bacteria suggested that there may be a relationship between growth and BEVs production rates [35]. Based on the assumption that the number of enriched proteins

reflects the abundance of BEVs to some extent, our results support the previously raised hypothesis that
fast-growing bacteria (e.g. *Flavobacteria* and *Alteromonas*) contribute more to the BEVs population [35].

It has been postulated that the production of BEVs is influenced by environmental factors that affect
their potential ecological impact [16,35]. Therefore, to account for the differences in microbial communities
and the biogeochemical conditions, we further examined each oceanic province separately. To infer the
potential functions of BEVs in the interaction with the extracellular environment, we examined 2,700
outer membrane and extracellular proteins identified in our dataset for enrichment in BEVs. We found
a total of 608 of these proteins (Table S4), which were associated with 51 different bacterial lineages
(Fig. S5), that were significantly enriched in BEVs compared to bacterial cells. The BEV-enriched
proteins were found to be associated with a variety of functional groups, including solute-binding
proteins, porins and various ribosomal proteins, as well as numerous proteins whose function is
unknown. Different proteins were found enriched in BEVs in different oceanic provinces, suggesting
that BEVs may play a role in multiple ecophysiological processes, the main ones of which are discussed
below.

Marine BEVs as vectors for iron scavenging
Iron is a micronutrient that occurs at low concentrations in surface seawater but is vital for life [54]. Iron
limitation can lead to enhanced BEVs production in non-marine bacteria [55]. Preferential packaging of
iron-related transporters into BEVs has been proposed to facilitate iron accumulation [52,56,57], and
subsequent uptake of such BEVs by bacterial cells has previously been demonstrated in a culture of a
pathogenic Gammaproteobacteria [58]. To scavenge for iron in the environment, some marine bacteria
release siderophores that strongly bind iron, and are subsequently acquired by bacterial cells through
specialised TonB-dependent transport systems [59–61]. The largest group of proteins that were significantly
enriched in marine BEVs consisted of proteins related to the TonB-dependent transport system (Fig. 5),
which accounted for 34-47% of all BEV-enriched outer membrane proteins in each region, compared
to 22-27% of outer membrane proteins enriched in the cellular fraction (Table S3). The majority of
these proteins were TonB-dependent receptors, which are outer membrane proteins that perform high-
affinity binding and energy-dependent transport of specific substrates across the membrane [62]. Although
the exact substrate for most of these receptors is unknown, we were able to identify 7 siderophore-
specific TonB-dependent receptors that were enriched in the BEVs across the transect. To actively
transport the siderophore across the outer membrane, the TonB-dependent receptor requires proton
motive force from the inner membrane [62]. While a proton motive force is likely lacking in BEVs, it has
recently been demonstrated in a non-marine bacterial isolate that TonB-dependent receptors are capable
of binding the siderophores even in the absence of such an energy source [58]. Therefore, it is possible
that iron or iron-rich proteins are trapped on the BEVs even if they are not internalized.

While production of siderophores in marine bacterial communities is mostly restricted to the class
Gammaproteobacteria, the capacity to acquire them from the environment (i.e., synthesizing specific
TonB-dependent receptors) is a taxonomically widespread functional trait [22]. Nonetheless, the vast
majority of the BEV-enriched TonB-dependent receptors we observed was also assigned to members
of the class Gammaproteobacteria (orders *Alteromonadales* and *Cellvibrionales*, as well as other
unclassified lineages), which represented 20-28% of all BEV-enriched outer membrane proteins,
compared to 12-16% of outer membrane proteins enriched in the cellular fraction. Enrichment of TonB-
dependent receptors on BEVs has been previously observed in cultures of marine Gammaproteobacteria
[31,63], among which are also several isolates of the order *Alteromonadales* [32]. Members of
Alteromonadales are copiotrophs that thrive under high availability of organic matter [64] and are
considered to be major siderophore producers and key mediators of iron acquisition in marine bacterial
communities [65,66]. We propose that during episodes of high metabolic activity (e.g., associated with
increased primary production), *Alteromonadales* not only release siderophores that bind dissolved iron,
but also produce BEVs that 'concentrate' them in the environment.

Fig. 5 – Enrichment of selected groups of proteins in marine BEVs. The upper panel shows the enrichment in BEVs and the lower panel shows the enrichment in the cellular fraction. 'Porins' – includes the cyanobacterial iron-specific porins and the putative carbohydrate-selective porin OprB. 'SusC/D' - includes both the SusC- and SusD-like proteins of the flavobacterial carbohydrate transport system. 'TBDR' – includes various TonB-dependent receptors. The total number of enriched proteins is indicated by the size of each circle and the number next to it.

Along with nitrogen, iron was the limiting nutrient of surface water primary production across most of the oceanic transect [38]. In the transition zone, where there was an elevated cyanobacterial activity without iron limitation, we did not identify any iron-selective porins among the BEV-enriched cyanobacterial proteins, compared to 6 enriched porins in the cellular fraction (Table S4). In contrast, in the westernmost region, where a cyanobacterial activity was observed and iron was identified as one of the limiting nutrients, 14 of the 37 BEV-enriched proteins associated with cyanobacteria (mostly *Prochlorococcaceae*) were putative iron-selective porins (Fig. 5). We propose that these iron-specific porins form channels that allow passive diffusion of inorganic iron into the BEVs, as they would do in intact cells [67]. The inorganic iron could then be bound by periplasmic iron-binding proteins [68], trapping it inside the BEVs.

In ocean regions with limited iron availability, BEVs with the characteristics described above could be considered as nanoscale "hotspots" of bioavailable iron that could potentially be available to marine bacteria in a relatively concentrated form. The producers of these BEVs could be either pelagic (i.e., free-living)[34,35], particle-associated [32,63], or associated with phytoplankton cells [30]. The BEVs produced by particle- or phytoplankton-associated bacteria are (to some extent) likely to be taken up by the same type of cell, since particle-associated bacteria may live in "microcolonies" [69,70]. By contrast, the cells found in the vicinity of a pelagic bacterium are likely to be distinct from the bacterium itself [71]. Although we still lack a thorough understanding of iron-related interactions among pelagic marine bacteria [72], we propose two possible mechanisms for the role of BEVs in iron acquisition by marine bacterial communities. One possible scenario is that, similar to siderophores that can be taken up by different

taxa [22], and these iron-accumulating marine BEVs could be considered to some extent as 'public goods'
that facilitate the iron needs of the entire bacterial community. However, it has also been shown that in
some cases the uptake of BEVs can be selective towards closely related bacterial lineages [56]. Although
the direct recognition mechanism between bacterial cells and BEVs [73], and the indirect effect of
characteristics such as lipid composition and surface charge on BEV uptake remain poorly understood,
the available evidence suggests that biases may exist within cell-vesicle interactions. Therefore, the
release of iron-accumulating BEVs could also be a means of competing for iron in the marine
environment, potentially favouring only closely related phylogenetic populations.

Marine BEVs as mediators of extracellular carbohydrate turnover
Some "selfish" marine bacteria have developed a strategy to prevent the loss of BEVs through
dispersion by producing them in chains that remain partially attached to the cell [30,31]. Such chains were
first observed in an isolate of *Formosa* spp. (class Flavobacteriia), and also in an isolate of marine
Pseudoalteromonas (class Gammaproteobacteria), where they were proposed to play a role in
carbohydrate utilisation based on the transporters and hydrolytic enzyme content of the BEVs.
Flavobacteria are predominantly active during elevated availability of organic matter (e.g., during
phytoplankton blooms), efficiently utilising various carbohydrates using their unique SusC/D transport
system [74,75]. Such BEV chains are fragile and were likely detached and disrupted by our filtration steps
[30], ending up in the bulk BEVs fraction. In our dataset, we identified 57 different integral SusC-like
TonB-dependent transporters that were enriched in the BEVs and 79 SusC-like proteins enriched in the
cellular fraction, with an absolute $\log_2$ fold change of around 4 (Fig. 5). In contrast, the 25 different
extracellular glycan-binding SusD-like proteins, which sit on top of the SusC and bind the substrate [76],
comprised more than twice the number of proteins in the BEV fraction than in the cellular fraction, with
an absolute $\log_2$ fold change of 4-5 compared to 2-4. Taken together with other BEV-enriched
flavobacterial proteins (Table S3), our observations are highly consistent with the previously conducted
culture-based study of *Formosa* spp. [30], confirming that flavobacterial BEVs with similar protein
content also occur in the natural marine environment.

The SusC/D transport system requires energy from the inner membrane in order to transport substrate
across the outer membrane [76]. In *Formosa* spp., based on protein characterization, it has been suggested
that the BEV-chains contain to some extent also an inner membrane (indications for which we also
observed here, such as the inner membrane SecDF protein and the ExbB proton pump; Table S3), and
so it is conceivable that BEV chains may have some ability to take up carbohydrate molecules, which
are then transported into the cell [30]. However, the stronger enrichment of SusD-like lipoproteins in the
BEV fraction compared with SusC suggests a similar mechanism to that proposed above for TonB-
dependent receptors, in which such flavobacterial BEVs are more likely to only be able to bind
carbohydrate molecules. The trapping of carbohydrate molecules on the cell membrane has been shown
to facilitate their efficient digestion by hydrolytic enzymes [26]. Recent observations have shown that gut
bacteria can actively alter the enzyme content of BEVs to optimise carbohydrate utilisation [77], an
analogous process may occur also in marine bacteria. While we lack such a mechanistic understanding
for marine BEVs, the presence of hydrolytic enzymes associated with BEVs and their activity has been
observed in several isolates of heterotrophic marine bacteria [31,32,78,79]. We were also able to identify a
single carbohydrate active enzyme (CAZyme - glycoside hydrolase family 16), taxonomically assigned
to *Flavobacteriales* that was enriched in BEVs in the westernmost region (Table S4). We also identified
an additional glycoside hydrolase taxonomically affiliated with the Alteromonadales and significantly
enriched in BEVs across the entire transect. Conceptually, the association of enzymes with marine
BEVs is advantageous because it allows them to be delivered to the substrate in a more protected manner
compared to direct excretion, and may also allow for more efficient degradation due to the higher
concentration of enzymes that ultimately reach the target molecule [1]. While it is widely accepted that
marine bacteria release hydrolytic enzymes into the extracellular milieu to facilitate carbohydrate
degradation [80], the evidence above suggests that part of this secretion occurs via marine BEVs.

Conclusions

Here we have characterised the abundance and diversity of marine BEVs on an extensive oceanic scale, integrating their ecological and biogeochemical contexts. We found that marine BEVs are ubiquitous in seawater and possess a substantial degree of protein diversity. A relationship was identified between the enrichment of certain cargoes of BEVs and the ecological conditions experienced by the local bacterial community, thus confirming their potential role in the scavenging of extracellular nutrients and energy. The next stage of the research requires methodological investigation into the mechanisms underlying BEV uptake, as this is a major gap in our understanding of the relationship between BEVs and bacterial cells. It also requires quantitative experiments on the contribution of BEVs to extracellular enzymatic activity and iron accumulation in marine bacterial communities. We argue that future research on extracellular processes in marine bacterial communities should consider the possibility that characterisations of bulk seawater may conflate activity mediated by 'truly' dissolved molecules (e.g. siderophores and enzymes) with activity associated with BEVs, the products of which may not be equally available to all cells.

Materials and Methods

Sample collection

The oceanographic sampling was conducted onboard RV *Sonne* (cruise SO289 representing section GP21 of the GEOTRACES program) from 18 February to 8 April 2022 (Fig. 1). Water samples with a total volume of 60 L were taken from a depth of ca. 6 m at each sampling station using the onboard water pump. To remove particulate matter and eukaryotic cells, water samples were pre-filtered at low pressure using a 3 μm pore-size membrane filter (Merck Millipore, MA, USA). From the pre-filtered water sample, 1 mL was preserved with 4% glutaraldehyde (final concentration) and stored at -80°C until flow cytometric analysis. The pre-filtered seawater sample was then filtered through a 142 mm diameter polycarbonate filter with a pore-size of 0.22 μm (Merck Millipore, MA, USA), which was then stored at -80°C and later used for molecular analyses of the bacterial communities. The filtrate (< 0.22 μm) was concentrated ca. 60 times using a Vivaflow tangential flow filtration system (Sartorius, Germany) with a 100 kDa cut-off to 1 L and stored at -20°C until further BEVs purification in the laboratory.

Phytoplankton pigments analysis

Seawater samples (2-4 L) for diagnostic phytoplankton pigments were filtered onto MF300 glass fibre filters (ThermoFisher Scientific, MA, USA) and stored at -80°C until analysis. Pigment extraction and analysis followed the procedure described by [19]. Briefly, upon return to a land-based laboratory, pigments were extracted in 90% acetone by homogenization of the filters using glass beads in a cell mill, centrifuged (10 minutes, 5200 rpm, 4 °C). The supernatant was filtered through 0.2 μm PTFE filters (VWR, PA, USA) and subsequently analysed by reverse-phase high-performance liquid chromatography (HPLC) Dionex UltiMate 3000 LC system (ThermoFisher Scientific, MA, USA)[81]. Diagnostic pigments were converted to estimate contributions of different phytoplankton types using CHEMTAX [82], with initial pigment ratios according to [83]. Total chlorophyll-a derived from HPLC include both chlorophyll-a and divinyl chlorophyll-a, and is referred to as chlorophyll-a throughout this study.

Bacterial cell counts

Bacterial cells were stained with 1×SYBR Green I (Thermo Fisher Scientific, MA, USA) for 15 min and then quantified using a flow cytometer Accuri™ C6 (BD Biosciences, NJ, USA). Counts were processed using R package 'flowWorkspace' v4.18.0 [84] and stained cells were quantified using the 'gate_flowclust_2d' function in R package 'openCyto' [85].

BEVs isolation and purification

A sub-sample of 100 mL of concentrated seawater was further concentrated about 30 times to a final volume of 3 mL using Vivaspin centrifugal filter units (Sartorius, Germany) with a 100 kDa cut-off. The entire concentrate was then loaded onto an iodixanol density gradient following a published protocol [86], and centrifuged at 100,000×g at 4°C for 6 hours. In the 15-30% and 35-40% fractions, the iodixanol solution was exchanged twice with 1×PBS using Vivaspin centrifugal filter units with a 100 kDa cut-off at 2000×g, and resuspended in 1 mL 1×PBS. From each purified fraction, 800 µL was used for molecular analyses and 200 µL for quantification of BEVs.

Nanoparticle tracking analysis

The abundance of BEVs was measured by nanoparticle tracking analysis (NTA) using a NanoSight NS300 instrument (Malvern Panalytycal, UK) equipped with a 488 nm laser. Each sample was injected using a syringe pump with a flow speed of 100 µL min^{-1}, and five videos of 60 sec were recorded. Videos were analysed using NanoSight NTA software v3.2. All particles in the size up to 300 nm in both the 15-30% and 35-40% fractions were included in the final abundance estimates. The flow cell was thoroughly flushed with Milli-Q water (Merck Millipore) between samples, and visually examined to exclude carry over. The performance of the instrument was routinely checked by measuring the concentration and size distribution of standardized 100 nm silica beads.

Metagenomics workflow

One third of the polycarbonate filters with a pore size of 0.22 µm were used for the extraction of genomic DNA using the DNeasy PowerWater Kit (QIAGEN N.V., Germany) according to the manufacturer's recommendations. The extracted DNA was subject to enzymatic shearing and then shotgun sequencing, performed on two lanes of the NovaSeq S4 platform (Illumina, Inc., CA, USA) at the Vienna Biocenter Core Facilities (https://www.viennabiocenter.org/vbcf/_next-generation-sequencing/).

The sequences from all samples were subject to the snakemake [87] metagenomic workflow in Anvi'o v8 [88], with recommended parameters. Briefly, the raw FASTQ files were subject to quality control by 'illumina-utils' v2.13 [89] and taxonomic classification using KrakenUniq v1.0.4 [90] and analysed using R package 'pavian' v1.2.1 [91]. Then the reads were assembled using MEGAHIT v1.2.9 [92] and the gene calling was performed using Prodigal v2.6.3 [93]. To construct the database for downstream metaproteomic analysis, protein-coding genes were clustered at 99% similarity using CD-HIT v4.6.881 [94].

Taxonomic affiliation of the gene calls was estimated using Kaiju v1.9.2 [95] against the NCBI RefSeq database (date: 2024-08-14), followed by classification of sequences with unknown taxonomic affiliation against the NCBI nr database (date: 2024-08-25). Functional annotation of the genes was carried out against InterPro v98 [96], NCBIfam v17.0 [97], Pfam v37 [98] databases using InterProScan v5.66 [99]. The unidentified protein sequences with no hits, were assigned taxonomy and function using BLASTp [100] best-hit approach, against NCBI RefSeq database (date: 2025-01-27). Protein sub-cellular localization was estimated using a combination of PSORTb v3.0.6 [101] and Phobius v1.01 [102].

Proteomics workflow

We used the remaining two thirds of the polycarbonate filters with a pore size of 0.22 µm for extraction of cellular proteins. Protein extraction from the filter was performed as described in detail by [103]. Briefly, the filters were ground into small pieces with a sterile metal spatula after submerging the tubes with the filters into liquid nitrogen. Filter pieces were resuspended in lysis buffer (100mM Tris-HCl pH 7.4, 1% SDS, 150mM NaCl, 1mM DTT, 10mM EDTA) and cells were lysed with five freeze-and-thaw cycles. After centrifugation (20,000x g at 4°C for 25 min) the supernatant was transferred into a tube and proteins were co-precipitated with 0.015% deoxycholate and 7% trichloroacetic acid (TCA) on ice for 1h and washed twice with ice-cold acetone. Dried protein pellets were resuspended in 50 mM TEAB buffer (Millipore Sigma, Burlington, MA, USA) and cysteines were reduced and alkylated with 10 mM DTT and 55 mM iodoacetamide (IAA), respectively. To extract proteins associated with the BEVs, we added the sample reducing agent NuPAGE (Invitrogen, Waltham, MA, USA) to 600 µL of purified BEVs at 1X final concentration.

All samples were re-precipitated using 9 times the sample volume of 96% EtOH at -20°C overnight.
Pellets were resuspended in 50 mM TEAB, followed by overnight in-solution trypsin (Roche, Basel,
Switzerland) digestion (1:100, w/w) at 37°C. TFA was added to the samples at 1% final
concentration to terminate trypsin digestion. Samples were desalted using Pierce C18 Tips
(ThermoFisher Scientific, MA, USA) according to the manufacturer's protocol. Prior the LC-MS
analysis, digested peptides were dissolved in 0.1% formic acid and 2% acetonitrile.

Purified tryptic peptides were dissolved in 0.1% formic acid (FA) and measured using a nano-reversed-
phase high-performance liquid chromatography (RP-HPLC) system Dionex Ultimate 3000
(ThermoFisher Scientific, MA, USA), coupled with a benchtop Quadrupole Orbitrap mass spectrometer
Q-Exactive Plus (ThermoFisher Scientific, MA, USA). Peptides were separated on a PepMap RSLC
C18 column (ThermoFisher Scientific, MA, USA) at 55°C with a flow rate of 300 nL/min. The two-
hour segmented LC gradient ranged from 5% to 80% buffer B (79.9% acetonitrile, 0.1% FA). Mass
spectra were acquired in positive ion mode using a top-15 data-dependent acquisition method. Full MS
scans were performed at a resolution of 70,000 (m/z 200), followed by MS/MS scans at a resolution of
17,500. High-energy collisional dissociation (HCD) fragmentation was applied with a normalized
collision energy (NCE) of 30%. Dynamic exclusion was set to 60 seconds.

The MS raw files of proteins extracted from the bacterial cells and the BEVs were searched using
SEQUEST-HT against the bacterial protein coding genes from the co-assembled metagenome.
Search parameters were as follows: enzyme - trypsin, fragment mass tolerance - 0.8 Da, max. missed
cleavages - 2, fixed modifications - carbamidomethyl (Cys), optional modifications - oxidation
(Met). Percolator parameters were as follows: max. delta Cn: 0.6, max. rank: 0, validation based on
q-value, false discovery rate (calculated by automatic decoy searches) 0.05. Only proteins with more
than two peptides and at least one unique peptide were included in the statistical analyses.

Statistical analyses
428 Statistical analyses were done in R v4.4.2 (R Core Team 2022) using RStudio v2024.09.1 (RStudio
429 Team 2019). Protein enrichment analysis was performed using the R package 'DEqMS' v1.22.0 [104] on
430 a log2-transformed and median-centred protein abundance matrix. In the enrichment test, only those
431 proteins were included that were observed in at least two samples in each fraction. All figures were
432 generated using the R package 'ggplot2' v3.5.1 [105].

Data availability
The metagenomic raw sequences have been deposited in the European Nucleotide Archive (ENA) at
EMBL-EBI under Project accession number PRJEB88044. The mass spectrometry proteomics data
have been deposited to the ProteomeXchange Consortium [106] via the PRIDE [107] partner repository with
the dataset identifier PXD062682.

Scripts for data processing and statistical analyses, as well as raw flow cytometry and NTA files, can
be accessed via Zenodo (https://www.zenodo.org/) under doi: 10.5281/zenodo.15772523 and
10.5281/zenodo.15772578, respectively.

Author contributions

442 EF performed fieldwork, the bioinformatic and statistical analyses, and wrote the manuscript. EF and
443 GJH designed the project. EF, NO, TT performed the protein extraction and LAS carried out the LC-
444 MS measurements. HL, ZY and TJB performed the phytoplankton pigment sampling and analyses. EPA
445 was the chief scientist on board and coordinated the sampling. DS and SJB contributed to data
446 interpretation. All authors contributed to the manuscript and approved its submitted version.

Acknowledgments

448 We thank the captain, the crew and the scientific team of the RV SONNE SO289 GEOTRACES
449 research expedition ('S Pacific GEOTRACES', 03G0289NA, funded by the Bundesministerium für
450 Bildung und Forschung (BMBF)). We thank A. Mutzberg, D. Jasinski, F. Evers, T. Schott, K.

Nachtigall and T. Klüver for technical support. We thank Z. Steiner for his support during the expedition. M. Aubert for assistance with laboratory sample processing. EF acknowledges funding by the Austrian Science Fund (FWF) M2797-B. This work was also supported by the Austrian Science Fund (FWF) project I4978-B to GJH and by the Slovenian Research Agency (ARIS) project J7-2599 to TT.

References

1. Kulp, A. & Kuehn, M. J. Biological functions and biogenesis of secreted bacterial outer membrane vesicles. *Annu. Rev. Microbiol.* **64**, 163–184 (2010).

2. Knox, K. W., Vesk, M. & Work, E. Relation between excreted lipopolysaccharide complexes and surface structures of a lysine-limited culture of Escherichia coli. *J. Bacteriol.* **92**, 1206–1217 (1966).

3. Toyofuku, M., Schild, S., Kaparakis-Liaskos, M. & Eberl, L. Composition and functions of bacterial membrane vesicles. *Nat. Rev. Microbiol.* (2023) doi:10.1038/s41579-023-00875-5.

4. Bitto, N. J. *et al.* Bacterial membrane vesicles transport their DNA cargo into host cells. *Sci. Rep.* **7**, 7072 (2017).

5. Tran, F. & Boedicker, J. Q. Genetic cargo and bacterial species set the rate of vesicle-mediated horizontal gene transfer. *Sci. Rep.* **7**, 8813 (2017).

6. Yaron, S., Kolling, G. L., Simon, L. & Matthews, K. R. Vesicle-mediated transfer of virulence genes from Escherichia coli O157:H7 to other enteric bacteria. *Appl. Environ. Microbiol.* **66**, 4414–4420 (2000).

7. Renelli, M., Matias, V., Lo, R. Y. & Beveridge, T. J. DNA-containing membrane vesicles of Pseudomonas aeruginosa PAO1 and their genetic transformation potential. *Microbiology* **150**, 2161–2169 (2004).

8. Johnston, E. L. *et al.* Planktonic and biofilm-derived Pseudomonas aeruginosa outer membrane vesicles facilitate horizontal gene transfer of Plasmid DNA. *Microbiol. Spectr.* e0517922 (2023).

9. Soler, N. & Forterre, P. Vesiduction: the fourth way of HGT. *Environ. Microbiol.* **22**, 2457–2460 (2020).

10. Toyofuku, M. *et al.* Membrane vesicle-mediated bacterial communication. *ISME J.* **11**, 1504–1509 (2017).

11.   Mashburn, L. M. & Whiteley, M. Membrane vesicles traffic signals and facilitate group activities

in a prokaryote. *Nature* **437**, 422–425 (2005).

12.   Yonezawa, H. *et al.* Outer membrane vesicles of Helicobacter pylori TK1402 are involved in

biofilm formation. *BMC Microbiol.* **9**, 197 (2009).

13.   Li, Z., Clarke, A. J. & Beveridge, T. J. Gram-negative bacteria produce membrane vesicles which

are capable of killing other bacteria. *J. Bacteriol.* **180**, 5478–5483 (1998).

14.   Kadurugamuwa, J. L. & Beveridge, T. J. Bacteriolytic effect of membrane vesicles from

Pseudomonas aeruginosa on other bacteria including pathogens: conceptually new antibiotics. *J.*

Bacteriol. **178**, 2767–2774 (1996).

15.   Zlatkov, N., Nadeem, A., Uhlin, B. E. & Wai, S. N. Eco-evolutionary feedbacks mediated by

bacterial membrane vesicles. *FEMS Microbiol. Rev.* (2020) doi:10.1093/femsre/fuaa047.

16.   Orench-Rivera, N. & Kuehn, M. J. Environmentally controlled bacterial vesicle-mediated export.

Cell. Microbiol. **18**, 1525–1536 (2016).

17.   Toyofuku, M., Tashiro, Y., Hasegawa, Y., Kurosawa, M. & Nomura, N. Bacterial membrane

vesicles, an overlooked environmental colloid: Biology, environmental perspectives and

applications. *Adv. Colloid Interface Sci.* **226**, 65–77 (2015).

18.   Bar-On, Y. M. & Milo, R. The Biomass Composition of the Oceans: A Blueprint of Our Blue

Planet. *Cell* **179**, 1451–1454 (2019).

19.   Browning, T. J. *et al.* Nutrient co-limitation at the boundary of an oceanic gyre. *Nature* **551**, 242–

246 (2017).

20.   Capone, D. G. & Hutchins, D. A. Microbial biogeochemistry of coastal upwelling regimes in a

changing ocean. *Nat. Geosci.* **6**, 711–717 (2013).

21.   Moore, C. M. *et al.* Processes and patterns of oceanic nutrient limitation. *Nat. Geosci.* **6**, 701–710

(2013).

22.   Zoccarato, L., Sher, D., Miki, T., Segrè, D. & Grossart, H.-P. A comparative whole-genome

approach identifies bacterial traits for marine microbial interactions. *Commun Biol* **5**, 276 (2022).

23.   Stocker, R. Marine microbes see a sea of gradients. *Science* **338**, 628–633 (2012).

24. Worden, A. Z. *et al.* Environmental science. Rethinking the marine carbon cycle: factoring in the
multifarious lifestyles of microbes. *Science* **347**, 1257594 (2015).

25. Kirchman, D. L. The uptake of inorganic nutrients by heterotrophic bacteria. *Microb. Ecol.* **28**,
255–271 (1994).

26. Reintjes, G., Arnosti, C., Fuchs, B. M. & Amann, R. An alternative polysaccharide uptake
mechanism of marine bacteria. *ISME J.* **11**, 1640–1650 (2017).

27. Tortell, P. D., Maldonado, M. T., Granger, J. & Price, N. M. Marine bacteria and biogeochemical
cycling of iron in the oceans. *FEMS Microbiol. Ecol.* **29**, 1–11 (1999).

28. Koch, A. L. What size should a bacterium be? A question of scale. *Annu. Rev. Microbiol.* **50**, 317–
348 (1996).

29. Gasol, J. M., del Giorgio, P. A., Massana, R. & Duarte, C. M. Active versus inactive bacteria:size-
dependence in a coastal marine plankton community. *Mar. Ecol. Prog. Ser.* **128**, 91–97 (1995).

30. Fischer, T. *et al.* Biopearling of Interconnected Outer Membrane Vesicle Chains by a Marine
Flavobacterium. *Appl. Environ. Microbiol.* **85**, (2019).

31. Dürwald, A. *et al.* Reaching out in anticipation: bacterial membrane extensions represent a
permanent investment in polysaccharide sensing and utilization. *Environ. Microbiol.* **23**, 3149–
3163 (2021).

32. Fadeev, E. *et al.* Characterization of membrane vesicles in Alteromonas macleodii indicates
potential roles in their copiotrophic lifestyle. *Microlife* **4**, 2022.09.27.509651 (2023).

33. Biller, S. J. *et al.* Bacterial vesicles in marine ecosystems. *Science* **343**, 183–186 (2014).

34. Biller, S. J. *et al.* Prochlorococcus extracellular vesicles: molecular composition and adsorption to
diverse microbes. *Environ. Microbiol.* (2021) doi:10.1111/1462-2920.15834.

35. Biller, S. J. *et al.* Environmental and Taxonomic Drivers of Bacterial Extracellular Vesicle
Production in Marine Ecosystems. *Appl. Environ. Microbiol.* e0059423 (2023).

36. Linney, M. D. *et al.* Microbial Sources of Exocellular DNA in the Ocean. *Appl. Environ.*
Microbiol. **88**, e0209321 (2022).

37. Lücking, D., Mercier, C., Alarcón-Schumacher, T. & Erdmann, S. Extracellular vesicles are the main contributor to the non-viral protected extracellular sequence space. *ISME Commun* **3**, 112 (2023).

38. Liu, H. *et al.* Patterns of (micro)nutrient limitation across the South Pacific Ocean. *Commun. Earth Environ.* **5**, 1–9 (2024).

39. Teeling, H. *et al.* Recurring patterns in bacterioplankton dynamics during coastal spring algae blooms. *Elife* **5**, e11888 (2016).

40. Pontiller, B. *et al.* Rapid bacterioplankton transcription cascades regulate organic matter utilization during phytoplankton bloom progression in a coastal upwelling system. *ISME J.* **16**, 2360–2372 (2022).

41. Morris, R. M. *et al.* SAR11 clade dominates ocean surface bacterioplankton communities. *Nature* **420**, 806–810 (2002).

42. Reintjes, G. *et al.* On-site analysis of bacterial communities of the ultraoligotrophic South Pacific Gyre. *Appl. Environ. Microbiol.* **85**, (2019).

43. Oggerin, M. *et al.* Niche differentiation within bacterial key-taxa in stratified surface waters of the Southern Pacific Gyre. *ISME J.* **18**, (2024).

44. Zhang, Z. *et al.* Substantial trace metal input from the 2022 Hunga Tonga-Hunga Ha'apai eruption into the South Pacific. *Nat. Commun.* **15**, 8986 (2024).

45. Biller, S. J. *et al.* Membrane vesicles in sea water: heterogeneous DNA content and implications for viral abundance estimates. *ISME J.* **11**, 394–404 (2017).

46. Soler, N., Krupovic, M., Marguet, E. & Forterre, P. Membrane vesicles in natural environments: a major challenge in viral ecology. *The ISME journal* vol. 9 793–796 (2015).

47. Nekrouf, N. A., Maestre-Carballa, L., Lluesma-Gomez, M., Martinez-Hernandez, F. & Martínez-García, M. Annual dynamics and metagenomics of marine vesicles: One more layer of complexity in the dissolved organic fraction. *Limnol. Oceanogr.* (2025) doi:10.1002/lno.70031.

48. Schwechheimer, C., Sullivan, C. J. & Kuehn, M. J. Envelope control of outer membrane vesicle production in Gram-negative bacteria. *Biochemistry* **52**, 3031–3040 (2013).

49.   Minhas, T. & Greenman, J. Production of cell-bound and vesicle-associated trypsin-like protease,

alkaline phosphatase and N-acetyl-beta-glucosaminidase by Bacteroides gingivalis strain W50. *J.*

Gen. Microbiol. **135**, 557–564 (1989).

50.   Sabra, W., Lünsdorf, H. & Zeng, A.-P. Alterations in the formation of lipopolysaccharide and

membrane vesicles on the surface of Pseudomonas aeruginosa PAO1 under oxygen stress

conditions. *Microbiology* **149**, 2789–2795 (2003).

51.   Gerritzen, M. J. H. *et al.* High dissolved oxygen tension triggers outer membrane vesicle formation

by Neisseria meningitidis. *Microb. Cell Fact.* **17**, 157 (2018).

52.   Orench-Rivera, N. & Kuehn, M. J. Differential Packaging Into Outer Membrane Vesicles Upon

Oxidative Stress Reveals a General Mechanism for Cargo Selectivity. *Front. Microbiol.* **12**,

561863 (2021).

53.   Pérez-Cruz, C., Delgado, L., López-Iglesias, C. & Mercade, E. Outer-inner membrane vesicles

naturally secreted by gram-negative pathogenic bacteria. *PLoS One* **10**, e0116896 (2015).

54.   Morel, F. M. M. & Price, N. M. The biogeochemical cycles of trace metals in the oceans. *Science*

300, 944–947 (2003).

55.   Prados-Rosales, R. *et al.* Role for Mycobacterium tuberculosis membrane vesicles in iron

acquisition. *J. Bacteriol.* **196**, 1250–1256 (2014).

56.   Wang, M., Nie, Y. & Wu, X.-L. Membrane vesicles from a Dietzia bacterium containing multiple

cargoes and their roles in iron delivery. *Environ. Microbiol.* **23**, 1009–1019 (2021).

57.   Lin, J. *et al.* A Pseudomonas T6SS effector recruits PQS-containing outer membrane vesicles for

iron acquisition. *Nat. Commun.* **8**, 14888 (2017).

58.   Dhurve, G., Madikonda, A. K., Jagannadham, M. V. & Siddavattam, D. Outer Membrane Vesicles

of Acinetobacter baumannii DS002 Are Selectively Enriched with TonB-Dependent Transporters

and Play a Key Role in Iron Acquisition. *Microbiol Spectr* **10**, e0029322 (2022).

59.   Hopkinson, B. M. & Barbeau, K. A. Iron transporters in marine prokaryotic genomes and

metagenomes. *Environ. Microbiol.* **14**, 114–128 (2012).

60.   Armstrong, E., Granger, J., Mann, E. L. & Price, N. M. Outer-membrane siderophore receptors of

heterotrophic oceanic bacteria. *Limnol. Oceanogr.* **49**, 579–587 (2004).

61. Hogle, S. L. *et al.* Siderophores as an iron source for picocyanobacteria in deep chlorophyll
maximum layers of the oligotrophic ocean. *ISME J.* **16**, 1636–1646 (2022).

62. Noinaj, N., Guillier, M., Barnard, T. J. & Buchanan, S. K. TonB-dependent transporters:
regulation, structure, and function. *Annu. Rev. Microbiol.* **64**, 43–60 (2010).

63. Frias, A., Manresa, A., de Oliveira, E., López-Iglesias, C. & Mercade, E. Membrane vesicles: a
common feature in the extracellular matter of cold-adapted antarctic bacteria. *Microb. Ecol.* **59**,
476–486 (2010).

64. Wietz, M., López-Pérez, M., Sher, D., Biller, S. J. & Rodriguez-Valera, F. Microbe Profile:
Alteromonas macleodii - a widespread, fast-responding, "interactive" marine bacterium.
Microbiology **168**, (2022).

65. Manck, L. E. *et al.* Petrobactin, a siderophore produced by Alteromonas, mediates community iron
acquisition in the global ocean. *ISME J.* **16**, 358–369 (2022).

66. Manck, L. E., Espinoza, J. L., Dupont, C. L. & Barbeau, K. A. Transcriptomic Study of Substrate-
Specific Transport Mechanisms for Iron and Carbon in the Marine Copiotroph Alteromonas
macleodii. *mSystems* **5**, (2020).

67. Qiu, G.-W. *et al.* A unique porin meditates iron-selective transport through cyanobacterial outer
membranes. *Environ. Microbiol.* **23**, 376–390 (2021).

68. Qiu, G.-W., Koedooder, C., Qiu, B.-S., Shaked, Y. & Keren, N. Iron transport in cyanobacteria -
from molecules to communities. *Trends Microbiol.* **30**, 229–240 (2022).

69. Grossart, H.-P., Kiørboe, T., Tang, K. & Ploug, H. Bacterial colonization of particles: growth and
interactions. *Appl. Environ. Microbiol.* **69**, 3500–3509 (2003).

70. Seymour, J. R., Amin, S. A., Raina, J.-B. & Stocker, R. Zooming in on the phycosphere: the
ecological interface for phytoplankton–bacteria relationships. *Nature Microbiology* **2**, 1–12
(2017).

71. Kashtan, N. *et al.* Single-cell genomics reveals hundreds of coexisting subpopulations in wild
Prochlorococcus. *Science* **344**, 416–420 (2014).

72.   Pham, A. L.-D., Aumont, O., Ratnarajah, L. & Tagliabue, A. Examining the interaction between

615   free-living bacteria and iron in the global ocean. *Global Biogeochem. Cycles* **36**, e2021GB007194

616   (2022).

617   73.   Knoke, L. R. *et al.* Agrobacterium tumefaciens Small Lipoprotein Atu8019 Is Involved in

618   Selective Outer Membrane Vesicle (OMV) Docking to Bacterial Cells. *Front. Microbiol.* **11**, 1228

619   (2020).

620   74.   Unfried, F. *et al.* Adaptive mechanisms that provide competitive advantages to marine

621   bacteroidetes during microalgal blooms. *ISME J.* **12**, 2894–2906 (2018).

622   75.   Kappelmann, L. *et al.* Polysaccharide utilization loci of North Sea Flavobacteriia as basis for using

623   SusC/D-protein expression for predicting major phytoplankton glycans. *ISME J.* **13**, 76–91 (2019).

624   76.   Glenwright, A. J. *et al.* Structural basis for nutrient acquisition by dominant members of the human

625   gut microbiota. *Nature* **541**, 407–411 (2017).

626   77.   Sartorio, M. G., Pardue, E. J., Scott, N. E. & Feldman, M. F. Human gut bacteria tailor extracellular

627   vesicle cargo for the breakdown of diet- and host-derived glycans. *Proc. Natl. Acad. Sci. U. S. A.*

628   **120**, e2306314120 (2023).

629   78.   Naval, P. & Chandra, T. S. Characterization of membrane vesicles secreted by seaweed associated

630   bacterium Alteromonas macleodii KS62. *Biochem. Biophys. Res. Commun.* **514**, 422–427 (2019).

631   79.   Li, J., Azam, F. & Zhang, S. Outer membrane vesicles containing signalling molecules and active

632   hydrolytic enzymes released by a coral pathogen Vibrio shilonii AK1. *Environ. Microbiol.* **18**,

633   3850–3866 (2016).

634   80.   Arnosti, C. *et al.* The Biogeochemistry of Marine Polysaccharides: Sources, Inventories, and

635   Bacterial Drivers of the Carbohydrate Cycle. *Ann. Rev. Mar. Sci.* **13**, 81–108 (2021).

636   81.   Van Heukelem, L. & Thomas, C. S. Computer-assisted high-performance liquid chromatography

637   method development with applications to the isolation and analysis of phytoplankton pigments. *J.*

638   *Chromatogr. A* **910**, 31–49 (2001).

639   82.   Mackey, M. D., Mackey, D. J., Higgins, H. W. & Wright, S. W. CHEMTAX - a program for

640   estimating class abundances from chemical markers:application to HPLC measurements of

641   phytoplankton. *Mar. Ecol. Prog. Ser.* **144**, 265–283 (1996).

83.   DiTullio, G. R. *et al.* Phytoplankton assemblage structure and primary productivity along 170°W
in the South Pacific Ocean. *Mar. Ecol. Prog. Ser.* **255**, 55–80 (2003).

84.   Finak, G. & Jiang, M. *FlowWorkspace: Infrastructure for Representing and Interacting with Gated*
and Ungated Cytometry Data Sets. (2022).

85.   Finak, G. *et al.* OpenCyto: an open source infrastructure for scalable, robust, reproducible, and
automated, end-to-end flow cytometry data analysis. *PLoS Comput. Biol.* **10**, e1003806 (2014).

86.   Biller, S. J. *et al.* Isolation and Characterization of Cyanobacterial Extracellular Vesicles. *J. Vis.*
Exp. (2022) doi:10.3791/63481.

87.   Köster, J. & Rahmann, S. Snakemake--a scalable bioinformatics workflow engine. *Bioinformatics*
28, 2520–2522 (2012).

88.   Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat.*
Microbiol. **6**, 3–6 (2021).

89.   Eren, A. M., Vineis, J. H., Morrison, H. G. & Sogin, M. L. A filtering method to generate high
quality short reads using illumina paired-end technology. *PLoS One* **8**, e66643 (2013).

90.   Breitwieser, F. P., Baker, D. N. & Salzberg, S. L. KrakenUniq: confident and fast metagenomics
classification using unique k-mer counts. *Genome Biol.* **19**, 198 (2018).

91.   Breitwieser, F. P. & Salzberg, S. L. Pavian: interactive analysis of metagenomics data for
microbiome studies and pathogen identification. *Bioinformatics* **36**, 1303–1304 (2020).

92.   Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics **31**, 1674–1676 (2015).

93.   Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification.
BMC Bioinformatics **11**, 119 (2010).

94.   Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

95.   Menzel, P., Ng, K. L. & Krogh, A. Fast and sensitive taxonomic classification for metagenomics
with Kaiju. *Nat. Commun.* **7**, 11257 (2016).

96. Blum, M. *et al.* InterPro: the protein sequence classification resource in 2025. *Nucleic Acids Res.*
53, D444–D456 (2025).

97. Li, W. *et al.* RefSeq: expanding the Prokaryotic Genome Annotation Pipeline reach with protein
family model curation. *Nucleic Acids Res.* **49**, D1020–D1028 (2021).

98. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, D412–D419
(2021).

99. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**,
1236–1240 (2014).

100. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

101. Lau, W. Y. V. *et al.* PSORTdb 4.0: expanded and redesigned bacterial and archaeal protein
subcellular localization database incorporating new secondary localizations. *Nucleic Acids Res.*
49, D803–D808 (2021).

102. Flores-Uribe, J. Identification of proteins containing transmembrane domains using Phobius v1.
(2018) doi:10.17504/protocols.io.pdwdi7e.

103. Tinta, T. *et al.* Microbial Processing of Jellyfish Detritus in the Ocean. *Front. Microbiol.* **11**,
590995 (2020).

104. Zhu, Y. *DEqMS*. (Bioconductor, 2018). doi:10.18129/B9.BIOC.DEQMS.

105. Villanueva, R. A. M. & Chen, Z. J. ggplot2: Elegant Graphics for Data Analysis (2nd ed.).
Measurement (Mahwah NJ) **17**, 160–167 (2019).

106. Deutsch, E. W. *et al.* The ProteomeXchange consortium at 10 years: 2023 update. *Nucleic Acids*
Res. **51**, D1539–D1548 (2023).

107. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass spectrometry-based
proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2022).