# Bacterial extracellular vesicles exhibit distinct functional potential across the biogeographic provinces of the South Pacific Ocean

Eduard Fadeev[1,*], Neza Orel[2], Tinkara Tinta[2], Leila Afjehi-Sadat[3], Haoran Liu[4], Thomas J. Browning[4], Zhongwei Yuan[4], Eric P. Achterberg[4], Steven J. Biller[5], Daniel J. Sher[6], and Gerhard J. Herndl[1,3,7]

[1] Department of Functional and Evolutionary Ecology, Bio-Oceanography and Marine Biology Unit, University of Vienna, Vienna, Austria

[2] Marine Biology Station Piran, National Institute of Biology, Piran, Slovenia

[3] Mass Spectrometry Unit, Research Support Facilities, University of Vienna, Vienna, Austria

[4] GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany

[5] Department of Biological Sciences, Wellesley College, Wellesley, USA

[6] Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Haifa, Israel

[7] NIOZ, Department of Marine Microbiology and Biogeochemistry, Royal Netherlands Institute for Sea Research, Den Burg, The Netherlands

*Corresponding author

**Keywords: bacterioplankton, membrane vesicles, horizontal gene transfer, siderophores, extracellular enzymes**

## Abstract

Bacterial extracellular vesicles (BEVs) are nanoscale membranous structures released by diverse types of bacteria. Laboratory model systems indicate that these nanoparticles may play several roles in the ecophysiology of marine bacteria. However, their actual functionality in the environment remains unclear. Here we describe the proteomic composition of marine BEVs over more than 5,000 nautical miles of surface waters in the South Pacific, linking BEV cargoes to the bacterial communities producing them. BEVs were consistently present across a range of biogeochemical conditions, with an overall abundance comparable to that of bacterial cells. However, the protein content of the BEVs varied significantly between different ocean regions. The BEVs were enriched in carbohydrate transporters under phytoplankton bloom conditions, and contained iron and phosphate uptake-related proteins in nutrient-limited waters. This suggests that BEVs could enable cells to perform key extracellular functions in the marine environment. Our observations further highlight the prevalence of BEVs and the biogeographic patterns of their functional potential across oceanic scales.

## Introduction

Bacterial extracellular vesicles (BEVs) are nanoscale, membrane-bound structures typically ranging in
diameter from 20 to 250 nm, which are released by bacterial cells. They can result from 'blebbing'
mechanisms in intact cells or be formed as a by-product of cell lysis. This can influence their contents
and, consequently, their functional potential (Kulp & Kuehn, 2010). Since their discovery in human-
associated *Escherichia coli* in the 1960s (Knox et al., 1966), most studies of BEVs have examined their
roles within laboratory model organisms, particularly host-associated bacteria in animal and plant
systems. BEVs can serve as a mechanism for removing harmful compounds from the cell, but can also
transport various biological cargoes (e.g., nucleic acids and proteins) over long distances in a protected
and locally concentrated manner (McMillan & Kuehn, 2021). One of the best-documented functions of
BEVs in intercellular interactions is the transfer of DNA molecules (Bitto et al., 2017; Soler & Forterre,
2020), which can facilitate the exchange of antibiotic resistance genes between different bacterial
lineages (Yaron et al., 2000), as well as the transfer of entire plasmids (Johnston et al., 2023; Renelli et
al., 2004), recently termed 'vesiduction' (Soler & Forterre, 2020). BEVs can also play roles in bacterial
quorum sensing (Mashburn & Whiteley, 2005; Toyofuku et al., 2017), biofilm formation (Yonezawa et
al., 2009), and contribute to microbial "warfare" by delivering compounds that kill competing bacteria
(Kadurugamuwa & Beveridge, 1996; Z. Li et al., 1998). As BEVs can contain a variety of components
released by bacterial cells, including enzymes, receptors and transporters derived from cell membranes,
they may interact with organic and inorganic chemicals in the environment (Orench-Rivera & Kuehn,
2016; Zlatkov et al., 2020). Such extracellular activities, including the enzymatic digestion or binding
of various environmental compounds, are particularly important for bacteria living in aqueous
environments (Toyofuku et al., 2023, 2015).

The ocean, where bacteria are among the most abundant organisms (Bar-On & Milo, 2019), is a highly
dynamic environment both physically and biochemically. The surface ocean is characterised by a
depletion of bioavailable nutrients required for life in some vast areas (e.g., oligotrophic oceanic gyres)
(Browning et al., 2017) and a high availability in others (e.g., coastal upwelling regions) (Capone &
Hutchins, 2013; Moore et al., 2013). To survive in such ecosystems, marine bacteria have developed a
wide range of ecophysiological capabilities (Stocker, 2012; Worden et al., 2015; Zoccarato et al., 2022),
and implement highly diverse nutrient and energy utilisation strategies (Kirchman, 1994; Reintjes et
al., 2017; Tortell et al., 1999). In the interaction between marine bacteria and their oligotrophic
environment, high cellular surface area to volume ratios facilitate efficient nutrient uptake (Gasol et al.,
1995; Koch, 1996). In this sense, the release of marine BEVs by bacterial cells can be conceptualised
as an expansion of the cell surface which disperses through the aqueous medium (Kulp & Kuehn, 2010),
randomly encountering various components of the marine environment. Laboratory observations of
several isolates of globally abundant marine bacteria suggest that BEVs may mediate specific
extracellular functions in the ecophysiology of these bacteria, such as binding nutrients (e.g., iron or
phosphate) and delivering them in concentrated form to bacterial cells (Biller et al., 2023; Biller,
Lundeen, et al., 2022; Biller et al., 2014; Dürwald et al., 2021; Fadeev et al., 2023; Fischer et al., 2019).
However, despite the potential importance of BEVs for marine bacteria, and ubiquity in the marine
environment (Linney et al., 2022; Lücking et al., 2023), our understanding of their biogeographic
diversity across large environmental gradients is unknown.

To address this knowledge gap, we conducted a spatially extensive sampling to examine the abundance
and protein cargo diversity of BEVs across the South Pacific Ocean. We determined how the molecular
cargo of marine BEV populations varies in context with their local microbial and biogeochemical
environment, finding that BEVs may play distinct functions for different microbial populations. Our
observations across large oceanic regions reinforce the emerging picture that important extracellular
bacterial functions may be mediated by BEVs and can inform future efforts to test specific hypotheses
about their ecological roles.

## Results and Discussion

**Distinct biogeochemical conditions across the South Pacific Ocean**

To investigate the potential ecophysiological functions of BEVs in marine bacterial communities we sampled across diverse environmental conditions, varying in nutrient limitations and phytoplankton growth conditions. We collected surface water (5-6 m depth) at 20 stations along a basin-scale transect that spanned the South Pacific Ocean (Fig. 1; Table S1). A detailed physicochemical characterisation of this transect showed that field sampling was carried out across four oceanic provinces (Liu et al., 2024; Yuan et al., 2025). Briefly, the easternmost region was the Chilean coastal upwelling zone (UP), characterized by elevated nitrate and phosphate concentrations, but depleted in dissolved iron. In this area there were elevated chlorophyll-a concentrations (ca. 3 mg m$^{-3}$; Fig. S1), with the phytoplankton community mostly composed of diatoms (Fig. 1). The transition zone between the upwelling and the subtropical gyre (TRAN) was characterized by elevated concentrations of phosphate and dissolved iron, but was depleted in nitrate (Fig. S1). This area also had relatively elevated chlorophyll-a concentrations (0.04-0.26 mg m$^{-3}$), with mixed phytoplankton communities. The centre of the subtropical gyre (GYRE) was characterized by very low nitrate concentrations and a declining spatial gradient from east to west of phosphate concentrations, with strongly depleted chlorophyll-a concentrations (< 0.03 mg m$^{-3}$; Fig. S1). The westernmost region (WEST) was defined as the area of the subtropical gyre with the lowest phosphate concentrations and higher (compared to GYRE) chlorophyll-a concentrations (0.03-0.14 mg m$^{-3}$; Fig. S1), with the phytoplankton comprising predominantly cyanobacteria.

***Fig. 1 – Diverse biogeochemical regimes encountered across the South Pacific Ocean.*** *(A) Spatial grouping of the sampling stations according to four oceanic provinces: 'UP' - Chilean coast upwelling zone; 'TRAN' -*

We found that bacterial abundance varied across the biogeochemical provinces (Table S1), and the
taxonomic composition of the bacterial communities (Fig. 2) also differed significantly between them
(PERMANOVA test; $F_{3,14}$=21.16, $R^2$=0.82, p=0.001). To investigate the physiological and metabolic
processes expressed within the bacterial communities, we characterized the metaproteome of the
bacterial cells, identifying between 4281-6200 cellular proteins at each station (Table S2). The
composition of bacterial cellular proteins significantly varied between the four regions (PERMANOVA
test; $F_{3,16}$=3.06, $R^2$=0.36, p=0.001; Fig. S2) and further differed between each pair of adjacent provinces
(Table S3), indicating that these represented distinct functional ecosystems. Bacterial cell abundance
was the highest in the upwelling zone (9-26 $\times 10^8$ cells $L^{-1}$), with the bacterial community composition
showing a strong heterotrophic response to a diatom bloom (Fig. 2). This response was attributed mainly
to members of the classes Gammaproteobacteria and Flavobacteria, which are typically active during
elevated availability of organic carbon (Pontiller et al., 2022; Teeling et al., 2016). In the transition
zone, bacterial abundance reached 0.7-3.5 $\times 10^8$ cells $L^{-1}$, with the cyanobacteria *Prochlorococcus* and
*Synechococcus* comprising more than 50% of the bacterial community (Table S1). The elevated
photosynthetic cyanobacterial activity in this zone was potentially stimulated by a nutrient influx from
a returning lateral current (Liu et al., 2024). The subtropical gyre had the lowest bacterial cell
abundances (0.2-0.9 $\times 10^8$ cells $L^{-1}$), with cyanobacteria comprising up to 50% of the bacterial
community (Table S1). Metagenomic characterization of the bacterial community composition further
suggested dominance of the oligotrophic Alphaproteobacterial order *Pelagibacterales* (Fig. 2)(R. M.
Morris et al., 2002). This lineage has previously been shown to numerically dominate the bacterial
communities of the ultra-oligotrophic South Pacific subtropical gyre (Oggerin et al., 2024; Reintjes et
al., 2019). In the westernmost region, cell abundance was significantly higher than in the subtropical
gyre (0.8-1.3 $\times 10^8$ cells $L^{-1}$; Student's t-test, p=0.002), with a predominance of *Prochlorococcus*. The
photosynthetic cyanobacterial activity in this region may have been partly facilitated by input from a
large volcanic eruption in the region two months prior to sampling (Zhang et al., 2024).

**Fig. 2 – Bacterial community composition based on taxonomic classification of metagenomic sequences.** *The taxonomic class of each order is given in parentheses: 'AC' – Actinomycetes, 'A' – Alphaproteobacteria, 'B' – Bacilli, 'C' – Clostridia, 'CY' – Cyanophyceae, 'F' – Flavobacteriia, 'G' – Gammaproteobacteria, 'E' – Epsilonproteobacteria. 'Other orders < 1%' – refers to taxonomic orders that comprised less than 1% of the sequences; 'unclassified' – refers to bacterial sequences with unknown classification on an order level. All samples contained $2$-$5 \times 10^8$ classified sequences.*

**Abundance, characteristics, and main producers of marine BEVs**

To investigate the abundance and diversity of marine BEVs, we sampled the extracellular size fraction corresponding to BEV-like structures (100 kDa-0.22 μm) across all stations using size fractionated filtration. The size distribution of marine viruses overlaps with that of BEVs (Biller et al., 2017; Nekrouf et al., 2025; Soler et al., 2015), so we used density gradient separation to further enrich our samples for BEVs in the laboratory  (see Materials and Methods). At least 50% of the purified BEVs were between 70-130 nm in diameter (Fig. 3), matching the size distributions previously observed in cultures of marine bacteria (Biller et al., 2023, 2014). Their abundance was overall comparable to the abundance of bacterial cells (Fig. 3), although this is likely a conservative estimate, since the density gradient separation led to a reduced yield of 6-52% of all nanoparticles (Fig. S3), and the storage of the samples on board at -20°C may also have resulted in a loss of BEV-like structures (Ahmadian et al., 2024).

We then characterised the bulk protein content in the BEV-designated density fraction (see Materials and Methods) and identified 1940-3394 proteins at each station (Table S1). Of these proteins only ca. 3% (58-123 proteins) were taxonomically or functionally (e.g., capsid or tail sheath) related to viruses, suggesting a strong enrichment of BEVs over other types of protein-containing marine nanoparticles in our samples. We found that 78-96% of proteins observed in the BEVs were also found in the corresponding cellular fraction at each station, but the composition of the proteins significantly varied between the two fractions (PERMANOVA test; $F_{1,38}$=15.01, $R^2$=0.28, p=0.001). This indicated that the BEV-associated proteins represented a distinct subset of the bacterial cellular proteome.

**Fig. 3 – Physical characteristics of marine BEV-like structures.** *(A) Estimated diameter distribution of the BEV-like structures. The error bars represent variation between NTA technical replicates. (B) Bacterial cell and BEVs abundances, estimated using flow cytometry and nanoparticle tracking analysis (NTA), respectively.*

Marine bacteria, which are mostly Gram-negative, can produce BEVs through blebbing of the outer membrane, leading to preferential incorporation of outer membrane and periplasmic material from the releasing cell (Kulp & Kuehn, 2010; Schwechheimer et al., 2013). External conditions have been shown to have an impact on the release of BEVs, potentially leading to some degree of preferential protein packaging (Gerritzen et al., 2018; Minhas & Greenman, 1989; Orench-Rivera & Kuehn, 2021; Sabra et al., 2003). Other BEVs may result from bacterial cell lysis, including due to bacteriophages, and thus represent a random assemblage of proteins from different subcellular origins (Pérez-Cruz et al., 2015; Toyofuku et al., 2023). To differentiate between these different origins of BEVs, we determined whether there is an enrichment in non-cytoplasmic proteins in the BEV fraction. We carried out a protein enrichment analysis on all samples in the BEV fraction as compared with the cellular fraction, and identified a total of 1065 proteins that were significantly enriched in the BEVs (absolute $\log_2$ fold difference > 1, adjusted p-value < 0.1), relative to their abundance in the cells. By comparison, 2017 proteins were significantly enriched in the cellular fraction (Table S4). Prediction of sub-cellular origins revealed that half of the BEV-enriched proteins were non-cytoplasmic, comprising mostly outer membrane and extracellular proteins, compared to ca. 25% of such proteins enriched in the cellular fraction (Fig. 4). Although we cannot exclude the presence of lysis-derived BEVs in our samples, the

enrichment of mostly non-cytoplasmic proteins suggests that the marine BEVs in our samples may have derived primarily from membrane blebbing of bacterial cells (Toyofuku et al., 2023).

**Localization**
- Cytoplasmic
- Cytoplasmic Membrane
- Periplasmic
- Outer Membrane
- Extracellular

**Order**
- Hyphomicrobiales(A)
- Pelagibacterales(A)
- Puniceispirillales(A)
- Rhodobacterales(A)
- Rhodospirillales(A)
- Sphingomonadales(A)
- Unknown order(A)
- Cytophagales(C)
- Synechococcales(CY)
- Flavobacteriales(F)
- Alteromonadales(G)
- Cellvibrionales(G)
- Lysobacterales(G)
- Moraxellales(G)
- Oceanospirillales(G)
- Pseudomonadales(G)
- Unknown order(G)
- Other taxa < 1%

*Fig. 4 – Overview of significantly enriched proteins in BEVs and cellular fractions. The bar plots represent the composition of all significantly enriched proteins according to their estimated sub-cellular localization (A) and taxonomic origin (B). The total number of enriched proteins in the BEVs and the cellular fractions was 1121 and 2075, respectively. Taxonomic composition of enriched proteins in the cellular and BEVs fractions according to different oceanic provinces. The taxonomic class of each order is given in parentheses: 'A' – Alphaproteobacteria, 'C' – Cytophagia, 'CY' – Cyanophyceae, 'F' – Flavobacteriia, 'G' – Gammaproteobacteria.*

Studies of laboratory cultures indicate that the major classes of marine bacteria exhibit significant variations in BEV production rates (Biller et al., 2023). Our protein enrichment analyses from field samples revealed that the two taxonomic lineages with the largest number of enriched proteins in both the BEVs and in the cellular fractions were the orders *Synechococcales* (including *Prochlorococcus* and *Synechococcus*) and the order *Pelagibacterales* (class Alphaproteobacteria). The ratio of enriched proteins between the BEVs and cellular fractions of these lineages showed a clear tendency towards the latter (Fig. 5). In contrast, enriched proteins associated with other bacterial orders, such as the *Alteromonadales* and *Cellvibrionales* (class Gammaproteobacteria), and the order *Flavobacteriales* (class Flavobacteriia), accounted for more than twice BEV-enriched proteins compared to the cellular fraction (Fig. 5). These observations could indicate that BEVs from these lineages contain more protein on average than those from Alphaproteobacteria and cyanobacteria. Alternatively, based on the

198 assumption that the number of enriched proteins reflects to some extent the abundance of BEVs, these
199 results are consistent with the previously raised hypothesis that fast-growing heterotrophic marine
200 bacteria contribute more to the BEV population than slow growing ones (Biller et al., 2023).

*Fig. 5 – Ratio of significantly enriched proteins in BEVs and cellular fraction in each taxonomic order. The*
*dashed line represents a theoretical ratio of 1:1 between the number of proteins enriched in each fraction. 'Other*
*taxa' represents classes with less than 3 enriched proteins in each of the fractions.*

The distinct biogeochemical conditions sampled in this study shape bacterial communities (Fig. 2),
which could potentially affect BEV populations and their functions (Biller et al., 2023; Orench-Rivera
& Kuehn, 2016). Therefore, in order to infer the potential functions of BEVs while accounting for these
differences, we examined the proteins enriched in BEVs separately for each oceanic province. As only
two stations sampled in the upwelling region, this region was excluded from further analysis. We found
a total of 1295 proteins (Table S5), which were associated with 77 different bacterial orders (Fig. S4),
that were significantly enriched in BEVs compared to bacterial cells. Different functional groups of
proteins were found enriched in BEVs in different oceanic provinces, suggesting that they may play a
role in multiple ecophysiological processes, the main ones of which are discussed below.

**Marine BEVs as mediators for extracellular electron transfer?**
The surface waters of the South Pacific receive high solar irradiance. High light levels with associated
enhanced UV radiation can induce oxidative stress in photosynthetic microorganisms (Mella-Flores et
al., 2012). Marine cyanobacteria (e.g., *Prochlorococcaceae* and *Synechococcaceae*) have developed

various strategies to cope with this environmental stressor, one of which is the transfer of excess
electrons to ferredoxins and flavoproteins (Scanlan et al., 2009). In our dataset, we observed that 10-
30% of all enriched cyanobacterial proteins in the BEV fraction were outer-membrane porins, compared
to 0.5-5% in the cellular fraction (Table S5). We also found ferredoxins and flavoproteins, which are
typically soluble, cytoplasmic proteins associated with the thylakoid membranes (Pierella Karlusich &
Carrillo, 2017), in cyanobacterial BEVs across all regions, with the strongest enrichment in the centre
of the subtropical gyre (Fig. 6). This enrichment under the nutrient limiting conditions may indicate that
such reduced electron carriers are exported in cyanobacterial BEVs as a cellular defence mechanism
against oxidative stress (Orench-Rivera & Kuehn, 2021).

Increasing light intensity, which can lead to oxidative stress in cyanobacteria, has been shown to
increase vesiculation rates in *Prochlorococcaceae* cultures (Biller et al., 2023; Biller, Lundeen, et al.,
2022). It has been also shown that marine heterotrophic bacteria, such as those belonging to the
*Alteromonadales*, support the growth of *Prochlorococcaceae* by scavenging excreted compounds
produced as a result of oxidative stress by the latter (J. J. Morris et al., 2011). Furthermore,
*Alteromonadales* are capable of taking up BEVs produced by *Prochlorococcaceae*, suggesting that
BEVs can serve as nutrient source for heterotrophic bacteria (Biller et al., 2014). Alternatively, based
on the demonstrated capability of various non-marine bacteria with anaerobic capabilities to acquire
energy from secreted flavoproteins (Marsili et al., 2008; Okamoto et al., 2014), we speculate that
cyanobacterial BEVs enriched with electron acceptors could also serve as a direct energy source for
marine heterotrophic bacteria such as the *Alteromonadales*. Given the recent observation in
*Prochlorococcaceae* and *Synechococcaceae* of intercellular extensions (nanotubes), containing vesicle-
like structures (Angulo-Cánovas et al., 2024), it is also possible that such extracellular electron transfer
occurs through direct contact between cells. In the ocean, such extracellular electron transfer
interactions are more likely to take place among particle-attached bacteria in a spatially confined
environment, which is often characterized by anaerobic conditions (Bianchi et al., 2018). Although this
study focuses on free-living marine bacteria, it is likely that our pre-filtration step, which aimed to
remove the particulate size fraction, disrupted any such structures present in the sample. Parts of these
structures may have ended up in the bulk BEVs fraction.

An alternative mechanism cyanobacteria use to cope with oxidative stress is photoprotection (Mella-
Flores et al., 2012). Interestingly, while in the regions with elevated cyanobacterial activity we observed
phycobilisomes enriched in the cellular fraction, in the centre of the subtropical gyre they were strongly
enriched in the BEVs fraction (Fig. 6). One possible explanation is that the BEV-enriched
phycobilisomes are damaged proteins that were removed from the cell (Schwechheimer & Kuehn,
2015). However, the identification of protein modifications resulting from oxidation revealed that,
while cell-enriched phycobilisomes exhibited up to 5 modifications, BEV-enriched phycobilisomes
exhibited only 0-1 modifications (Table S5). Therefore, it is conceivable that the release of
phycobilisomes from cells into BEVs in response to light and UV stress could be a photoprotective
mechanism, reducing the amount of light energy absorbed by the cell by removing light-harvesting
compounds. Furthermore, given the large size of cyanobacterial phycobilisomes (6.2 MDa; Domínguez-
Martín et al., 2022) and the assumption that BEVs remain attached to the cell, these phycobilisome-
enriched BEVs could provide additional extracellular shading, reducing the amount of light that reaches
the cyanobacterial cell.

**Fig. 6 – *Cyanobacterial proteins significantly enriched in BEVs and cellular fractions in different regions*.** *The mean log₂ fold difference was calculated for each functional protein group in each fraction. Positive log₂ fold difference represents significant enrichment in the BEVs fraction, negative mean log₂ fold difference represents significant enrichment in the cellular fraction. The total number of enriched proteins in the BEVs and the cellular fractions is provided in brackets. The proteins were grouped according to their proposed functions in BEVs (porins, oxidative stress proteins (OS) and other proteins). For visualisation purposes, only functional groups containing at least three enriched proteins in at least one region were included.*

### Marine BEVs as extracellular reserves of bio-available phosphate?

The surface waters of the South Pacific subtropical gyre are one of the most oligotrophic regions in the global ocean (Morel et al., 2010). Among the bacterial lineages that successfully inhabit such limiting environment are members of the order *Pelagibacterales* (also known as SAR11 clade). Their small cell size and large periplasm, which contains many substrate-binding proteins with exceptionally high affinities (Clifton et al., 2024; Sowell et al., 2009), which facilitates their uptake of soluble substrates (e.g., phosphate, sulphate, amino acids, and sugars) from the environment (Bosdriesz et al., 2015). This enables SAR11 to dominate the bacterial communities in the ocean (R. M. Morris et al., 2002) and thrive in nutrient-depleted marine ecosystems (Giovannoni, 2017). We observed that in all regions around one third of all proteins enriched in the cellular fraction were periplasmic substrate-binding proteins annotated to bind amino acids, compared to around 10% in the BEVs fraction, which consisted of other periplasmic binding proteins, mostly with unknown substrates (Table S5). Against this backdrop, in the centre of the subtropical gyre we found significant enrichment of five different homologues of the SAR11 phosphate binding protein PstS (Fig. 7). This periplasmic binding protein has a unique structure that has been suggested to have high affinity for phosphate, as well as being capable of binding larger organic phosphorus molecules (W.-J. Zhu et al., 2025). We propose that outer-membrane porins, which were also observed among the SAR11-derived BEV-enriched proteins (Fig. 7), form channels that allow phosphate molecules to diffuse passively into the BEVs, as they would in intact cells. They become trapped there by binding to high-affinity PstS proteins, making the BEVs a potential phosphate reservoir. Moreover, the accumulation of phosphate in the periplasm is enhanced in the presence of a proton motive force (Kamennaya et al., 2020). Therefore, the observation of proteorhodopsins enriched in BEVs of SAR11 (Fig. 7) raises the possibility that light energy could, in some individual particles, further enhance the accumulation of phosphate inside BEVs in surface waters.

## Marine BEVs as passive scavengers of iron?

Unlike the highly streamlined ecophysiology of SAR11, which relies mostly on passive diffusion from the environment through outer-membrane porins (Giovannoni, 2017), other heterotrophic marine bacteria actively acquire elements using specialised transport systems (Hopkinson & Barbeau, 2012; R. M. Morris et al., 2010; Tang et al., 2012). Our analysis revealed that one of the largest group of proteins significantly enriched in marine BEVs consisted of proteins related to the outer-membrane TonB-dependent transport system (Table S5). The TonB-dependent transport system is a highly diverse functional group of proteins that are involved in cellular uptake of a wide range of substrates, such as iron and iron-containing molecules (e.g., siderophores), as well as vitamins and various organic compounds (Braun, 2024), all of which were observed enriched on BEVs in our dataset.

For BEVs produced by 'blebbing' mechanisms, wherein pieces of the outermost cellular membrane detach from the cell to form the vesicle, their proteome is expected to be biased towards outer-membrane proteins (Kulp & Kuehn, 2010). Therefore, in order to determine whether TonB-related proteins are specifically enriched in BEVs, we separately tested the 3,700 outer-membrane proteins identified in our dataset for enrichment. We identified a total of 301 TonB-related proteins that were enriched in BEVs in one or more regions across the transect, compared to 229 TonB-related proteins enriched in the cellular fraction (out of a total 690 and 623 enriched outer-membrane proteins, respectively; Table S6). The majority of the BEV-enriched TonB-dependent proteins belonged to the one-component system of the outer-membrane receptor (OMR) family that perform both binding and transport of the substrate through the outer membrane. The enriched OMRs were primarily affiliated with the orders *Sphingomonadales* (class Alphaproteobacteria), *Alteromonadales* and *Cellvibrionales* (both class Gammaproteobacteria), and more than 20% of them were linked to the utilization of iron (Fig. 8 and Supplementary Text). Along with nitrate, iron was the main limiting micronutrient for surface water primary production across most of the transect (Liu et al., 2024; Yuan et al., 2025). TonB-dependent receptors are highly expressed in heterotrophic bacterial communities in the upper water column (< 100 m depth) (Bergauer et al., 2018). Their strong enrichment on BEVs has previously been

observed in cultures of related marine bacteria, including several isolates of *Alteromonadales* (Fadeev et al., 2023), an isolate of endosymbiotic marine *Cellvibrionales* (Gasser et al., 2024), and other marine Gammaproteobacteria (Dürwald et al., 2021; Frias et al., 2010). It has been also observed that iron limitation can lead to enhanced BEVs production in non-marine bacteria (Prados-Rosales et al., 2014) and preferential packaging of iron-related transporters into BEVs has been proposed to facilitate iron accumulation (Lin et al., 2017; Orench-Rivera & Kuehn, 2021; Wang et al., 2021). Taken together, these findings suggest that BEVs might play a role in extracellular scavenging of iron by marine bacteria.

To actively transport iron-containing molecules across the outer membrane, such TonB-dependent receptors require proton motive force from the inner membrane (Noinaj et al., 2010). Although BEVs are unlikely to have a proton motive force, these TonB-dependent receptors can still bind the substrates as was recently demonstrated in a culture of a pathogenic Gammaproteobacterium (Dhurve et al., 2022). Taking this into account, we propose that, in conditions of limited availability such as those observed in this study, marine BEVs may 'collect' iron in the environment. They could therefore be considered nanoscale 'hotspots' of bioavailable iron which could potentially provide a relatively concentrated source for marine bacteria.

*The mean log₂ fold difference was calculated for each protein group in each fraction. Positive log₂ fold difference represents significant enrichment in the BEVs fraction, negative log₂ fold difference represents significant enrichment in the cellular fraction. The total number of enriched proteins in the BEVs and the cellular fractions is provided in brackets. The proteins were grouped according to their predicated ligands. The pie charts show the breakdown of protein taxonomic affiliations on order level within each group. The taxonomic class of each order is given in parentheses: 'A' – Alphaproteobacteria, 'B' – Betaproteobacteria, 'C' – Cytophagia, 'CY' – Cyanophyceae, 'F' – Flavobacteriia, 'G' – Gammaproteobacteria.*

**Marine BEVs as mediators of carbohydrate utilisation?**

The second largest functional group of TonB-dependent receptors enriched in BEVs was related to the unique SusC-like proteins family (Fig. 8), which play an important role in carbohydrate utilisation by marine Flavobacteria (i.e., order *Flavobacteriales*). Flavobacteria efficiently utilise various sugars using a two component SusCD system, where SusC-like proteins are integral outer-membrane transporters and SusD-like lipoproteins facilitate extracellular high-affinity binding of the sugars (Glenwright et al., 2017). We found a total of 30 SusC-like proteins enriched in BEVs, compared to 55 SusC-like proteins enriched in the cellular fraction. In contrast, there were 27 SusD-like lipoproteins enriched in BEVs, compared to 15 such proteins in the cellular fraction (Table S6). The stronger enrichment of SusD-like lipoproteins in the BEV fraction compared to SusC suggests that flavobacterial BEVs may only be able to bind to carbohydrate molecules, without the ability to transport them. This phenomenon was previously observed in an isolate of marine *Flavobacteriales* (*Formosa* spp.; (Fischer et al., 2019). Trapping carbohydrate molecules on the BEV membrane (or BEV getting attached to a large carbohydrate) may facilitate their efficient digestion by hydrolytic enzymes, in a manner similar to that observed for cellular outer membranes (Reintjes et al., 2017). While we lack such a mechanistic understanding for marine BEVs, the presence of hydrolytic enzymes associated with BEVs and their activity has been observed in several isolates of heterotrophic marine bacteria (Dürwald et al., 2021; Fadeev et al., 2023; J. Li et al., 2016; Naval & Chandra, 2019).

Recent observations have shown that gut bacteria can actively alter the enzyme content of BEVs to optimise carbohydrate utilisation (Sartorio et al., 2023), an analogous process may occur also in marine bacteria. In our dataset, we identified an enrichment of a single glycoside hydrolase (family GH16) in BEVs in the western region (log₂ fold difference of approximately 4), which was taxonomically affiliated with *Flavobacteriales* (Table S6). The GH16 family contains a diversity of glycoside hydrolases that are active on a wide variety of marine carbohydrate molecules (Viborg et al., 2019). Conceptually, the association of enzymes with marine BEVs is advantageous because it allows them to be delivered to the substrate in a more protected manner compared to being freely dispersed in the water column, and may also allow for more efficient degradation in combination with the SusC-like and Sus-D like proteins (Du Toit, 2023). When considered alongside other BEV-enriched proteins (Table S6), our observations on marine BEVs are in line with previous findings regarding the composition of BEVs in isolates of marine *Flavobacteriales* and *Alteromonadales* (Dürwald et al., 2021; Fischer et al., 2019). In these studies, it was suggested that BEVs play a role in carbohydrate utilisation, based on their transporter and hydrolytic enzyme content. Previous work has shown that marine bacteria release hydrolytic enzymes into the extracellular milieu to facilitate carbohydrate utilisation during periods of increased phytoplankton activity (Arnosti et al., 2021) - a phenomenon observed in the transitional zone and the western regions of the transect. This suggests that some of the secretion occurs via marine BEVs, potentially affecting degradation rates or access to the products.

**Is the production of marine BEVs a 'selfish' trait or a 'public good'?**

The organisms our data indicate to produce most of the BEVs can be found both planktonic (i.e., free-living) and associated with particles (including phytoplankton)(Ivars-Martinez et al., 2008; Seymour et al., 2017). In contrast to the relatively confined particle-associated environments (e.g., biofilms), it is unlikely that BEVs released by a planktonic cell directly into seawater will be encountered again by the

same, or a highly similar, bacterial cell (Kashtan et al., 2014; Słomka et al., 2023). This led to the initial
hypothesis that marine BEVs likely serve to dispose harmful elements or act as a defence mechanism
against viral infection (Biller et al., 2014). However, it has been shown that in some cases the uptake of
BEVs only occurs between closely related bacterial lineages (Wang et al., 2021), suggesting that there
may be mechanisms that bias interactions between BEVs and recipient cells (Knoke et al., 2020). Thus,
BEV production could represent a mechanism through which cells compete for nutrients in the marine
environment, potentially favouring only closely related phylogenetic populations.

The environmental observations of natural bacterial communities in this study, alongside previous
observations of marine bacterial cultures, may also suggest other scenarios. One alternative explanation
is that marine BEVs fulfil their intended function while remaining within the diffusive boundary layer
(Stocker & Seymour, 2012). Another possible scenario is that the BEVs remain attached to the cell.
Some 'selfish' marine bacteria appear to have developed means of producing BEVs that are found in
chains attached to the cell. This trait was previously observed in isolates of marine *Alteromonadales*
and *Flavobacteriales* (Dürwald et al., 2021; Fischer et al., 2019). Chains of such outer membrane-
attached BEVs significantly increase the surface area through which the cell interacts with its
environment, facilitating the accumulation of elements necessary for life in the vicinity of the cell and
their subsequent uptake. Taking into account the potential stability of marine BEVs for at least a number
of days (Biller et al., 2014), they may continue to perform their function (e.g. accumulating phosphate
or iron) even after detachment from the cell. Therefore, whether separated from cells or released directly
into the environment, marine BEVs could also be considered a 'public good' that meets the various
needs of the bacterial community.

## Conclusions

Here we have explored BEVs contents within highly diverse natural marine bacterial communities on
an extensive oceanic scale in the South Pacific. We found that marine BEVs are ubiquitous in surface
seawater and possess a substantial degree of protein diversity. The enrichment of functionally specific
proteins in BEVs, which appears to be linked to the ecological conditions experienced by the local
bacterial community, which suggests BEVs may contribute to marine biogeochemistry and microbial
community metabolism. These findings reinforce the emerging hypothesis that BEVs play a role in
nutrient and energy scavenging in the marine environment. Our study provides initial observations on
the novel functional potential of marine BEVs, such as their ability to perform local enrichment of
molecules necessary for life, which motivate future investigations into the underlying mechanisms. To
develop a better understanding of the role of BEVs in the ecophysiology of marine bacteria, research
into their uptake dynamics is urgently needed. This is a significant knowledge gap regarding the
relationship between BEVs and bacterial cells, which could help us understand whether BEVs are a
'public good' or a competitive feature of marine bacterial communities. Based on the extensive
oceanographic observations presented in this study, the next stage of the research requires the
quantification of the contribution of BEVs to the distribution of essential life-sustaining elements (e.g.
iron and phosphate) in the marine environment. This will shed light on their importance for microbial
processes in the ocean.

## Materials and methods

**Sample collection**
The oceanographic sampling was conducted onboard RV Sonne (cruise SO289 representing section
GP21 of the GEOTRACES programme) from 18 February to 8 April 2022 (Fig. 1A). Water samples
with a total volume of 60 L were taken from a depth of ca. 6 m at each sampling station using the
onboard water pump. To remove particulate matter and eukaryotic cells, water samples were pre-filtered
at low pressure using a 3 μm pore size membrane filter (Merck Millipore, MA, USA). From the pre-
filtered water sample, 1 mL was preserved with 4% glutaraldehyde (final concentration) and stored at -

80°C until flow cytometric analysis. The pre-filtered seawater sample was then filtered through a 142
mm diameter polycarbonate filter with a pore size of 0.22 μm (Merck Millipore, MA, USA), which
were then stored at -80°C and later used for molecular analyses of the bacterial communities. The filtrate
(< 0.22 μm) was concentrated ca. 60 times using a Vivaflow tangential flow filtration system (Sartorius,
Germany) with a 100 kDa cut-off to 1 L and stored at -20°C until further BEVs purification in the
laboratory.

**Phytoplankton pigments analysis**

Seawater samples (2-4 L) for diagnostic phytoplankton pigments were filtered onto MF300 glass fibre
filters (ThermoFisher Scientific, MA, USA) and stored at -80°C until analysis. Pigments extraction and
analysis followed the procedure described by (Browning et al., 2017). Briefly, upon return to a land-
based laboratory, pigments were extracted in 90% acetone by homogenization of the filters using glass
beads in a cell mill, centrifuged (10 min, 5200 rpm, 4 °C). The supernatant was filtered through 0.2 μm
PTFE filters (VWR, PA, USA) and subsequently analysed by reverse-phase high performance liquid
chromatography (HPLC) Dionex UltiMate 3000 LC system (ThermoFisher Scientific, MA, USA)(Van
Heukelem & Thomas, 2001). Diagnostic pigments were converted to estimated contributions of
different phytoplankton types using CHEMTAX (Mackey et al., 1996), with initial pigment ratios
according to (DiTullio et al., 2003). Total chlorophyll-a derived from HPLC include both chlorophyll-
a and divinyl chlorophyll-a, and is referred to as chlorophyll-a throughout this study.

**Bacterial cell counts**

A subsample of 100 μL of seawater from each sampling stations was stained with 1×SYBR Green I
(Thermo Fisher Scientific, MA, USA) for 15 min at room temperature and then quantified using
FACSAria Flow Cytometer (BD Biosciences, NJ, USA). Counts were processed using R package
'flowWorkspace' v4.18.0 (Finak & Jiang, 2022). Quality control of the FCS files was carried out using
'flow_auto_qc' function in R package 'flowAI' (Monaco et al., 2016). The stained cells were subject
to logicle transformation and quantified using the 'gate_flowclust_2d' function in R package
'openCyto' (Finak et al., 2014), applied on the entire dataset (Fig. S5).

Cell abundances of *Prochlorococcus* and *Synechococcus* were reported by (Yuan et al., 2025) at
locations different from those sampled in this study but from the same biogeochemical provinces.
Therefore, the abundance of these cells was used to roughly estimate the contribution of cyanobacteria
to the total bacterial abundance.

**BEVs isolation and purification**

A sub-sample of 100 mL of concentrated seawater was further concentrated about 30 times to a final
volume of 3 mL using Vivaspin centrifugal filter units (Sartorius, Germany) with a 100 kDa cut-off.
The entire concentrate was then loaded onto an iodixanol density gradient following a published
protocol (Biller, Muñoz-Marín, et al., 2022), and centrifuged at 100,000×g at 4°C for 6 h. In the 15-
30% and 35-40% fractions, the iodixanol solution was exchanged twice with 1×PBS using Vivaspin
centrifugal filter units with a 100 kDa cut-off at 2000×g, and resuspended in 1 mL 1×PBS. From each
purified fraction, 800 μL was used for molecular analyses and 200 μL for quantification of BEVs.

**Nanoparticle tracking analysis**

The abundance of BEVs was measured by nanoparticle tracking analysis (NTA) using a NanoSight
NS300 instrument (Malvern Panalytycal, UK) equipped with a 488 nm laser. Each sample was injected
using a syringe pump with a flow speed of 100 μL min$^{-1}$, and five videos of 60 sec were recorded.
Videos were analysed using NanoSight NTA software v3.2. All particles in the size up to 300 nm in
both the 15-30% and 35-40% fractions were included in the final abundance estimates. The flow cell
was thoroughly flushed with Milli-Q water (Merck Millipore) between samples, and visually examined
to exclude carry over. The performance of the instrument was routinely checked by measuring the
concentration and size distribution of standardized 100 nm silica beads. To estimate the background,
we quantified nanoparticles in Optiprep media that was subject to the same washing steps with 1×PBS
as the BEVs samples, yielding concentrations of ca. $10^5$ particles mL$^{-1}$, with most particles being below
50 nm in size.

**Metagenomics workflow**

One third of the polycarbonate filters with a pore size of 0.22 μm were used for the extraction of genomic DNA using the DNeasy PowerWater Kit (QIAGEN N.V., Germany) according to the manufacturer's recommendations. The extracted DNA was subject to enzymatic shearing and then shotgun sequencing, performed on two lanes of the NovaSeq S4 platform (Illumina, Inc., CA, USA) at the Vienna Biocenter Core Facilities (https://www.viennabiocenter.org/vbcf/ next-generation-sequencing/).

The sequences from all samples were subject to the snakemake (Köster & Rahmann, 2012) metagenomic workflow in Anvi'o v8 (Eren et al., 2021), with recommended parameters. Briefly, the raw FASTQ files were subject to quality control by 'illumina-utils' v2.13 (Eren et al., 2013) and taxonomic classification using KrakenUniq v1.0.4 (F. P. Breitwieser et al., 2018) and analysed using R package 'pavian' v1.2.1 (Florian P. Breitwieser & Salzberg, 2020). Then the reads were assembled using MEGAHIT v1.2.9 (D. Li et al., 2015) and the gene calling was performed using Prodigal v2.6.3 (Hyatt et al., 2010).

Taxonomic affiliation of the protein coding genes was estimated using Kaiju v1.9.2 (Menzel et al., 2016) against the NCBI RefSeq database (date: 2024-08-14), followed by classification of sequences with unknown taxonomic affiliation against the NCBI nr database (date: 2024-08-25). Functional annotation of the protein coding genes was carried out against InterPro v98 (Blum et al., 2025), NCBIfam v17.0 (Wenjun Li et al., 2021), and Pfam v37 (Mistry et al., 2021) databases using InterProScan v5.66 (Jones et al., 2014). The unidentified proteins sequences with no hits, were assigned to their taxonomy and function using BLASTp (Camacho et al., 2009) best-hit approach against NCBI RefSeq database (date: 2025-01-27). Protein sub-cellular localization predictions were established using DeepLoc v2.0 (Thumuluri et al., 2022). Porins and TonB-dependent receptors were further classified using PSIBLAST (Camacho et al., 2009) against the Transporter Classification Database (TCDB) (Saier et al., 2006), best hit was defined based on identity > 30%, e-value < 0.001 and bit score > 50.

To construct the database for downstream metaproteomic analysis, bacterial protein coding genes were clustered at 99% similarity using CD-HIT v4.6.881 (Weizhong Li & Godzik, 2006).

**Proteomics workflow**

We used the remaining two thirds of the 0.22 μm pore-size polycarbonate filters for extraction of cellular proteins. Protein extraction from the filter was performed as described elsewhere (Tinta et al., 2020). Briefly, the filters were ground into small pieces with a sterile metal spatula after submerging the tubes with the filters into liquid nitrogen. Filter pieces were resuspended in lysis buffer (100mM Tris-HCl pH 7.4, 1% SDS, 150mM NaCl, 1mM DTT, 10mM EDTA) and cells were lysed with five freeze-and-thaw cycles. After centrifugation (20,000x $g$ at 4°C for 25 min) the supernatant was transferred into a tube and proteins were co-precipitated with 0.015% deoxycholate and 7% trichloroacetic acid (TCA) on ice for 1h and washed twice with ice-cold acetone. Dried protein pellets were resuspended in 50 mM TEAB buffer (Millipore Sigma, Burlington, MA, USA) and cysteines were reduced and alkylated with 10 mM DTT and 55 mM iodoacetamide (IAA), respectively. To extract proteins associated with the BEVs, we added the sample reducing agent NuPAGE (Invitrogen, Waltham, MA, USA) to 600 μL of purified BEVs at 1X final concentration.

All samples were re-precipitated using 9 times the sample volume of 96% EtOH at -20°C overnight. Pellets were resuspended in 50 mM TEAB, followed by overnight in-solution trypsin (Roche, Basel, Switzerland) digestion (1:100, w/w) at 37°C. TFA was added to the samples at 1% final concentration to terminate trypsin digestion. Samples were desalted using Pierce C18 Tips (ThermoFisher Scientific, MA, USA) according to the manufacturer's protocol. Prior to the LC-MS analysis, digested peptides were dissolved in 0.1% formic acid and 2% acetonitrile.

Purified tryptic peptides were dissolved in 0.1% formic acid (FA) and measured using a nano-reversed-phase high-performance liquid chromatography (RP-HPLC) system Dionex Ultimate 3000 (ThermoFisher Scientific, MA, USA), coupled with a benchtop Quadrupole Orbitrap mass spectrometer

Q-Exactive Plus (ThermoFisher Scientific, MA, USA). Peptides were separated on a PepMap RSLC
C18 column (ThermoFisher Scientific, MA, USA) at 55°C with a flow rate of 300 nL/min. The two-
hour segmented LC gradient ranged from 5% to 80% buffer B (79.9% acetonitrile, 0.1% FA). Mass
spectra were acquired in positive ion mode using a top-15 data-dependent acquisition method. Full MS
scans were performed at a resolution of 70,000 ($m/z$ 200), followed by MS/MS scans at a resolution of
17,500. High-energy collisional dissociation (HCD) fragmentation was applied with a normalized
collision energy (NCE) of 30%. Dynamic exclusion was set to 60 s.

The MS raw files of proteins extracted from the bacterial cells and the BEVs were processed using
the Spectrum Files RC node in Proteome Discoverer v2.2.0.338 (ThermoFisher Scientific, MA, USA)
and searched using SEQUEST-HT against the bacterial protein coding genes from the co-assembled
metagenome. Search parameters were as follows: enzyme - trypsin, fragment mass tolerance - 0.8
Da, max. missed cleavages - 2, fixed modifications - carbamidomethyl (Cys), optional modifications
- oxidation (Met). Percolator parameters were as follows: max. delta Cn: 0.6, max. rank: 0, validation
based on q-value, false discovery rate (calculated by automatic decoy searches) 0.05. Abundance of
each protein was estimated by summing the abundance of unique and razor peptides with at least 2
Peptide Spectrum Matches (PSMs). Only proteins with at least one unique peptide and at least two
peptides in total (unique and/or razor peptides) were included in the statistical analyses.

**Statistical analyses**
Statistical analyses were done in R v4.4.2 (R Core Team 2022) using RStudio v2024.09.1 (RStudio
Team 2019). Prior to statistical analyses, the protein abundance matrix was $log_2$-transformed and
median-centred. Protein enrichment analysis was performed using the R package 'DEqMS' v1.22.0 (Y.
Zhu, 2018). Only those proteins were included that were observed in at least two samples in each
fraction. The criteria for a significantly enriched protein were an absolute $log_2$ fold change > 1 and
adjusted p-value < 0.1. All figures were generated using the R package 'ggplot2' v3.5.1 (Villanueva &
Chen, 2019).

**Data availability**
The metagenomic raw sequences have been deposited in the European Nucleotide Archive (ENA) at
EMBL-EBI under Project accession number PRJEB88044. The mass spectrometry proteomics data
have been deposited to the ProteomeXchange Consortium (Deutsch et al., 2023) via the PRIDE (Perez-
Riverol et al., 2022) partner repository with the dataset identifier PXD062682.

Scripts for data processing and statistical analyses, as well as raw flow cytometry and NTA files, can
be accessed via Zenodo (https://www.zenodo.org/) under doi: 10.5281/zenodo.15772578.

# Author contributions

572 EF performed fieldwork, the bioinformatic and statistical analyses, and wrote the manuscript. EF and
573 GJH designed the project. EF, NO, TT performed the protein extraction and LAS carried out the LC-
574 MS measurements and wrote the corresponding methods section. HL, ZY and TJB performed the
575 phytoplankton pigment sampling and analyses. DS and SJB contributed to data interpretation and
576 manuscript writing. All authors approved the submitted version of the article.

# Acknowledgments

We thank the captain, the crew and the scientific team of the SO289 research expedition ('S Pacific
GEOTRACES', funded by the Bundesministerium für Bildung und Forschung (BMBF),
Förderkennzeichen 03G0289NA). We thank A. Mutzberg, D. Jasinski, F. Evers, T. Schott, K.
Nachtigall and T. Klüver for technical support. We thank Z. Steiner for his support during the
expedition. M. Aubert for assistance with laboratory sample processing. EF acknowledges funding by
the Austrian Science Fund (FWF) M2797-B. This work was also supported by the Austrian Science
Fund (FWF) project I4978-B to GJH and by the Slovenian Research Agency (ARIS) project J7-2599
to TT.

## References

Ahmadian, S., Jafari, N., Tamadon, A., Ghaffarzadeh, A., Rahbarghazi, R., & Mahdipour, M. (2024). Different storage and freezing protocols for extracellular vesicles: a systematic review. *Stem Cell Research & Therapy*, *15*(1), 453.

Angulo-Cánovas, E., Bartual, A., López-Igual, R., Luque, I., Radzinski, N. P., Shilova, I., Anjur-Dietrich, M., García-Jurado, G., Úbeda, B., González-Reyes, J. A., Díez, J., Chisholm, S. W., García-Fernández, J. M., & Del Carmen Muñoz-Marín, M. (2024). Direct interaction between marine cyanobacteria mediated by nanotubes. *Science Advances*, *10*(21), eadj1539.

Arnosti, C., Wietz, M., Brinkhoff, T., Hehemann, J.-H., Probandt, D., Zeugner, L., & Amann, R. (2021). The Biogeochemistry of Marine Polysaccharides: Sources, Inventories, and Bacterial Drivers of the Carbohydrate Cycle. *Annual Review of Marine Science*, *13*, 81–108.

Bar-On, Y. M., & Milo, R. (2019). The Biomass Composition of the Oceans: A Blueprint of Our Blue Planet. *Cell*, *179*(7), 1451–1454.

Bergauer, K., Fernandez-Guerra, A., Garcia, J. A. L., Sprenger, R. R., Stepanauskas, R., Pachiadaki, M. G., Jensen, O. N., & Herndl, G. J. (2018). Organic matter processing by microbial communities throughout the Atlantic water column as revealed by metaproteomics. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(3), E400–E408.

Bianchi, D., Weber, T. S., Kiko, R., & Deutsch, C. (2018). Global niche of marine anaerobic metabolisms expanded by particle microenvironments. *Nature Geoscience*, *11*(4), 263–268.

Biller, S. J., Coe, A., Arellano, A. A., Dooley, K., Silvestri, S. M., Gong, J. S., Yeager, E. A., Becker, J. W., & Chisholm, S. W. (2023). Environmental and Taxonomic Drivers of Bacterial Extracellular Vesicle Production in Marine Ecosystems. *Applied and Environmental Microbiology*, e0059423.

Biller, S. J., Lundeen, R. A., Hmelo, L. R., Becker, K. W., Arellano, A. A., Dooley, K., Heal, K. R., Carlson, L. T., Van Mooy, B. A. S., Ingalls, A. E., & Chisholm, S. W. (2022). Prochlorococcus extracellular vesicles: molecular composition and adsorption to diverse microbes. *Environmental Microbiology*, *24*(1), 420–435.

Biller, S. J., McDaniel, L. D., Breitbart, M., Rogers, E., Paul, J. H., & Chisholm, S. W. (2017).

Membrane vesicles in sea water: heterogeneous DNA content and implications for viral

abundance estimates. *The ISME Journal*, *11*(2), 394–404.

Biller, S. J., Muñoz-Marín, M. D. C., Lima, S., Matinha-Cardoso, J., Tamagnini, P., & Oliveira, P.

(2022). Isolation and Characterization of Cyanobacterial Extracellular Vesicles. *Journal of*

*Visualized Experiments: JoVE*, *180*. https://doi.org/10.3791/63481

Biller, S. J., Schubotz, F., Roggensack, S. E., Thompson, A. W., Summons, R. E., & Chisholm, S. W.

(2014). Bacterial vesicles in marine ecosystems. *Science*, *343*(6167), 183–186.

Bitto, N. J., Chapman, R., Pidot, S., Costin, A., Lo, C., Choi, J., D'Cruze, T., Reynolds, E. C.,

Dashper, S. G., Turnbull, L., Whitchurch, C. B., Stinear, T. P., Stacey, K. J., & Ferrero, R. L.

(2017). Bacterial membrane vesicles transport their DNA cargo into host cells. *Scientific*

*Reports*, *7*(1), 7072.

Blum, M., Andreeva, A., Florentino, L. C., Chuguransky, S. R., Grego, T., Hobbs, E., Pinto, B. L.,

Orr, A., Paysan-Lafosse, T., Ponamareva, I., Salazar, G. A., Bordin, N., Bork, P., Bridge, A.,

Colwell, L., Gough, J., Haft, D. H., Letunic, I., Llinares-López, F., … Bateman, A. (2025).

InterPro: the protein sequence classification resource in 2025. *Nucleic Acids Research*,

*53*(D1), D444–D456.

Bosdriesz, E., Magnúsdóttir, S., Bruggeman, F. J., Teusink, B., & Molenaar, D. (2015). Binding

proteins enhance specific uptake rate by increasing the substrate-transporter encounter rate.

*The FEBS Journal*, *282*(12), 2394–2407.

Braun, V. (2024). Substrate uptake by TonB-dependent outer membrane transporters. *Molecular*

*Microbiology*, *122*(6), 929–947.

Breitwieser, F. P., Baker, D. N., & Salzberg, S. L. (2018). KrakenUniq: confident and fast

metagenomics classification using unique k-mer counts. *Genome Biology*, *19*(1), 198.

Breitwieser, Florian P., & Salzberg, S. L. (2020). Pavian: interactive analysis of metagenomics data

for microbiome studies and pathogen identification. *Bioinformatics (Oxford, England)*, *36*(4),

1303–1304.

Browning, T. J., Achterberg, E. P., Rapp, I., Engel, A., Bertrand, E. M., Tagliabue, A., & Moore, C.

M. (2017). Nutrient co-limitation at the boundary of an oceanic gyre. *Nature*, *551*(7679),

242–246.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L.

(2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*(1), 421.

Capone, D. G., & Hutchins, D. A. (2013). Microbial biogeochemistry of coastal upwelling regimes in

a changing ocean. *Nature Geoscience*, *6*(9), 711–717.

Clifton, B. E., Alcolombri, U., Uechi, G.-I., Jackson, C. J., & Laurino, P. (2024). The ultra-high

affinity transport proteins of ubiquitous marine bacteria. *Nature*, *634*(8034), 721–728.

Deutsch, E. W., Bandeira, N., Perez-Riverol, Y., Sharma, V., Carver, J. J., Mendoza, L., Kundu, D. J.,

Wang, S., Bandla, C., Kamatchinathan, S., Hewapathirana, S., Pullman, B. S., Wertz, J., Sun,

Z., Kawano, S., Okuda, S., Watanabe, Y., MacLean, B., MacCoss, M. J., … Vizcaíno, J. A.

(2023). The ProteomeXchange consortium at 10 years: 2023 update. *Nucleic Acids Research*,

*51*(D1), D1539–D1548.

Dhurve, G., Madikonda, A. K., Jagannadham, M. V., & Siddavattam, D. (2022). Outer Membrane

Vesicles of Acinetobacter baumannii DS002 Are Selectively Enriched with TonB-Dependent

Transporters and Play a Key Role in Iron Acquisition. *Microbiology Spectrum*, *10*(2),

e0029322.

DiTullio, G. R., Geesey, M. E., Jones, D. R., Daly, K. L., Campbell, L., & Smith, W. O., Jr. (2003).

Phytoplankton assemblage structure and primary productivity along 170°W in the South

Pacific Ocean. *Marine Ecology Progress Series*, *255*, 55–80.

Domínguez-Martín, M. A., Sauer, P. V., Kirst, H., Sutter, M., Bína, D., Greber, B. J., Nogales, E.,

Polívka, T., & Kerfeld, C. A. (2022). Structures of a phycobilisome in light-harvesting and

photoprotected states. *Nature*, *609*(7928), 835–845.

Du Toit, A. (2023). Capturing glycans. *Nature Reviews. Microbiology*, *21*(8), 485.

Dürwald, A., Zühlke, M.-K., Schlüter, R., Gebbe, R., Bartosik, D., Unfried, F., Becher, D., &

Schweder, T. (2021). Reaching out in anticipation: bacterial membrane extensions represent a

permanent investment in polysaccharide sensing and utilization. *Environmental*

*Microbiology*, *23*(6), 3149–3163.

Eren, A. M., Kiefl, E., Shaiber, A., Veseli, I., Miller, S. E., Schechter, M. S., Fink, I., Pan, J. N.,

Yousef, M., Fogarty, E. C., Trigodet, F., Watson, A. R., Esen, Ö. C., Moore, R. M., Clayssen,

Q., Lee, M. D., Kivenson, V., Graham, E. D., Merrill, B. D., … Willis, A. D. (2021).

Community-led, integrated, reproducible multi-omics with anvi'o. *Nature Microbiology*, *6*(1),

3–6.

Eren, A. M., Vineis, J. H., Morrison, H. G., & Sogin, M. L. (2013). A filtering method to generate

high quality short reads using illumina paired-end technology. *PloS One*, *8*(6), e66643.

Fadeev, E., Carpaneto Bastos, C., Hennenfeind, J. H., Biller, S. J., Sher, D., Wietz, M., & Herndl, G.

J. (2023). Characterization of membrane vesicles in Alteromonas macleodii indicates

potential roles in their copiotrophic lifestyle. *MicroLife*, *4*, 2022.09.27.509651.

Finak, G., Frelinger, J., Jiang, W., Newell, E. W., Ramey, J., Davis, M. M., Kalams, S. A., De Rosa,

S. C., & Gottardo, R. (2014). OpenCyto: an open source infrastructure for scalable, robust,

reproducible, and automated, end-to-end flow cytometry data analysis. *PLoS Computational*

*Biology*, *10*(8), e1003806.

Finak, G., & Jiang, M. (2022). *flowWorkspace: Infrastructure for representing and interacting with*

*gated and ungated cytometry data sets* (4.8.0) [Computer software].

https://www.bioconductor.org/packages/release/bioc/html/flowWorkspace.html

Fischer, T., Schorb, M., Reintjes, G., Kolovou, A., Santarella-Mellwig, R., Markert, S., Rhiel, E.,

Littmann, S., Becher, D., Schweder, T., & Harder, J. (2019). Biopearling of Interconnected

Outer Membrane Vesicle Chains by a Marine Flavobacterium. *Applied and Environmental*

*Microbiology*, *85*(19). https://doi.org/10.1128/AEM.00829-19

Frias, A., Manresa, A., de Oliveira, E., López-Iglesias, C., & Mercade, E. (2010). Membrane vesicles:

a common feature in the extracellular matter of cold-adapted antarctic bacteria. *Microbial*

*Ecology*, *59*(3), 476–486.

Gasol, J. M., del Giorgio, P. A., Massana, R., & Duarte, C. M. (1995). Active versus inactive

bacteria:size-dependence in a coastal marine plankton community. *Marine Ecology Progress*

*Series*, *128*, 91–97.

Gasser, M. T., Liu, A., Altamia, M. A., Brensinger, B. R., Brewer, S. L., Flatau, R., Hancock, E. R.,

Preheim, S. P., Filone, C. M., & Distel, D. L. (2024). Membrane vesicles can contribute to

cellulose degradation by Teredinibacter turnerae, a cultivable intracellular Endosymbiont of

shipworms. *Microbial Biotechnology*, *17*(12), e70064.

Gerritzen, M. J. H., Maas, R. H. W., van den Ijssel, J., van Keulen, L., Martens, D. E., Wijffels, R. H.,

& Stork, M. (2018). High dissolved oxygen tension triggers outer membrane vesicle

formation by Neisseria meningitidis. *Microbial Cell Factories*, *17*(1), 157.

Giovannoni, S. J. (2017). SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Annual*

*Review of Marine Science*, *9*, 231–255.

Glenwright, A. J., Pothula, K. R., Bhamidimarri, S. P., Chorev, D. S., Baslé, A., Firbank, S. J., Zheng,

H., Robinson, C. V., Winterhalter, M., Kleinekathöfer, U., Bolam, D. N., & van den Berg, B.

(2017). Structural basis for nutrient acquisition by dominant members of the human gut

microbiota. *Nature*, *541*(7637), 407–411.

Hopkinson, B. M., & Barbeau, K. A. (2012). Iron transporters in marine prokaryotic genomes and

metagenomes. *Environmental Microbiology*, *14*(1), 114–128.

Hyatt, D., Chen, G.-L., Locascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010). Prodigal:

prokaryotic gene recognition and translation initiation site identification. *BMC*

*Bioinformatics*, *11*(1), 119.

Ivars-Martinez, E., Martin-Cuadrado, A.-B., D'auria, G., Mira, A., Ferriera, S., Johnson, J., Friedman,

R., & Rodriguez-Valera, F. (2008). Comparative genomics of two ecotypes of the marine

planktonic copiotroph Alteromonas macleodii suggests alternative lifestyles associated with

different kinds of particulate organic matter. *The ISME Journal*, *2*(12), 1194–1212.

Johnston, E. L., Zavan, L., Bitto, N. J., Petrovski, S., Hill, A. F., & Kaparakis-Liaskos, M. (2023).

Planktonic and biofilm-derived Pseudomonas aeruginosa outer membrane vesicles facilitate

horizontal gene transfer of Plasmid DNA. *Microbiology Spectrum*, e0517922.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,

Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M.,

Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: genome-scale protein function

classification. *Bioinformatics (Oxford, England)*, *30*(9), 1236–1240.

Kadurugamuwa, J. L., & Beveridge, T. J. (1996). Bacteriolytic effect of membrane vesicles from

Pseudomonas aeruginosa on other bacteria including pathogens: conceptually new antibiotics.

*Journal of Bacteriology*, *178*(10), 2767–2774.

Kamennaya, N. A., Geraki, K., Scanlan, D. J., & Zubkov, M. V. (2020). Accumulation of ambient

phosphate into the periplasm of marine bacteria is proton motive force dependent. *Nature*

*Communications*, *11*(1), 2642.

Kashtan, N., Roggensack, S. E., Rodrigue, S., Thompson, J. W., Biller, S. J., Coe, A., Ding, H.,

Marttinen, P., Malmstrom, R. R., Stocker, R., Follows, M. J., Stepanauskas, R., & Chisholm,

S. W. (2014). Single-cell genomics reveals hundreds of coexisting subpopulations in wild

Prochlorococcus. *Science (New York, N.Y.)*, *344*(6182), 416–420.

Kirchman, D. L. (1994). The uptake of inorganic nutrients by heterotrophic bacteria. *Microbial*

*Ecology*, *28*(2), 255–271.

Knoke, L. R., Abad Herrera, S., Götz, K., Justesen, B. H., Günther Pomorski, T., Fritz, C.,

Schäkermann, S., Bandow, J. E., & Aktas, M. (2020). Agrobacterium tumefaciens Small

Lipoprotein Atu8019 Is Involved in Selective Outer Membrane Vesicle (OMV) Docking to

Bacterial Cells. *Frontiers in Microbiology*, *11*, 1228.

Knox, K. W., Vesk, M., & Work, E. (1966). Relation between excreted lipopolysaccharide complexes

and surface structures of a lysine-limited culture of Escherichia coli. *Journal of Bacteriology*,

*92*(4), 1206–1217.

Koch, A. L. (1996). What size should a bacterium be? A question of scale. *Annual Review of*

*Microbiology*, *50*(1), 317–348.

Köster, J., & Rahmann, S. (2012). Snakemake--a scalable bioinformatics workflow engine.

*Bioinformatics (Oxford, England)*, *28*(19), 2520–2522.

Kulp, A., & Kuehn, M. J. (2010). Biological functions and biogenesis of secreted bacterial outer

membrane vesicles. *Annual Review of Microbiology*, *64*, 163–184.

Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

*Bioinformatics (Oxford, England)*, *31*(10), 1674–1676.

Li, J., Azam, F., & Zhang, S. (2016). Outer membrane vesicles containing signalling molecules and

active hydrolytic enzymes released by a coral pathogen Vibrio shilonii AK1. *Environmental*

*Microbiology*, *18*(11), 3850–3866.

Li, Weizhong, & Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics (Oxford, England)*, *22*(13), 1658–1659.

Li, Wenjun, O'Neill, K. R., Haft, D. H., DiCuccio, M., Chetvernin, V., Badretdin, A., Coulouris, G.,

Chitsaz, F., Derbyshire, M. K., Durkin, A. S., Gonzales, N. R., Gwadz, M., Lanczycki, C. J.,

Song, J. S., Thanki, N., Wang, J., Yamashita, R. A., Yang, M., Zheng, C., … Thibaud-Nissen,

F. (2021). RefSeq: expanding the Prokaryotic Genome Annotation Pipeline reach with protein

family model curation. *Nucleic Acids Research*, *49*(D1), D1020–D1028.

Li, Z., Clarke, A. J., & Beveridge, T. J. (1998). Gram-negative bacteria produce membrane vesicles

which are capable of killing other bacteria. *Journal of Bacteriology*, *180*(20), 5478–5483.

Lin, J., Zhang, W., Cheng, J., Yang, X., Zhu, K., Wang, Y., Wei, G., Qian, P.-Y., Luo, Z.-Q., & Shen,

X. (2017). A Pseudomonas T6SS effector recruits PQS-containing outer membrane vesicles

for iron acquisition. *Nature Communications*, *8*(1), 14888.

Linney, M. D., Eppley, J. M., Romano, A. E., Luo, E., DeLong, E. F., & Karl, D. M. (2022).

Microbial Sources of Exocellular DNA in the Ocean. *Applied and Environmental*

*Microbiology*, *88*(7), e0209321.

Liu, H., Yuan, Z., Gosnell, K. J., Liu, T., Tammen, J. K., Wen, Z., Engel, A., Liu, X., Huang, B., Kao,

S.-J., Achterberg, E. P., & Browning, T. J. (2024). Patterns of (micro)nutrient limitation

across the South Pacific Ocean. *Communications Earth & Environment*, *5*(1), 1–9.

Lücking, D., Mercier, C., Alarcón-Schumacher, T., & Erdmann, S. (2023). Extracellular vesicles are

the main contributor to the non-viral protected extracellular sequence space. *ISME*

*Communications*, *3*(1), 112.

Mackey, M. D., Mackey, D. J., Higgins, H. W., & Wright, S. W. (1996). CHEMTAX - a program for

estimating class abundances from chemical markers:application to HPLC measurements of

phytoplankton. *Marine Ecology Progress Series*, *144*, 265–283.

Marsili, E., Baron, D. B., Shikhare, I. D., Coursolle, D., Gralnick, J. A., & Bond, D. R. (2008).

Shewanella secretes flavins that mediate extracellular electron transfer. *Proceedings of the*

*National Academy of Sciences of the United States of America*, *105*(10), 3968–3973.

Mashburn, L. M., & Whiteley, M. (2005). Membrane vesicles traffic signals and facilitate group

activities in a prokaryote. *Nature*, *437*(7057), 422–425.

McMillan, H. M., & Kuehn, M. J. (2021). The extracellular vesicle generation paradox: a bacterial

point of view. *The EMBO Journal*, e108174.

Mella-Flores, D., Six, C., Ratin, M., Partensky, F., Boutte, C., Le Corguillé, G., Marie, D., Blot, N.,

Gourvil, P., Kolowrat, C., & Garczarek, L. (2012). Prochlorococcus and Synechococcus have

evolved different adaptive mechanisms to cope with light and UV stress. *Frontiers in*

*Microbiology*, *3*, 285.

Menzel, P., Ng, K. L., & Krogh, A. (2016). Fast and sensitive taxonomic classification for

metagenomics with Kaiju. *Nature Communications*, *7*(1), 11257.

Minhas, T., & Greenman, J. (1989). Production of cell-bound and vesicle-associated trypsin-like

protease, alkaline phosphatase and N-acetyl-beta-glucosaminidase by Bacteroides gingivalis

strain W50. *Journal of General Microbiology*, *135*(3), 557–564.

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L.,

Tosatto, S. C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., & Bateman, A. (2021).

Pfam: The protein families database in 2021. *Nucleic Acids Research*, *49*(D1), D412–D419.

Monaco, G., Chen, H., Poidinger, M., Chen, J., de Magalhães, J. P., & Larbi, A. (2016). flowAI:

automatic and interactive anomaly discerning tools for flow cytometry data. *Bioinformatics*

*(Oxford, England)*, *32*(16), 2473–2480.

Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., … Ulloa, O. (2013). Processes and patterns of oceanic nutrient limitation. *Nature Geoscience*, *6*(9), 701–710.

Morel, A., Claustre, H., & Gentili, B. (2010). The most oligotrophic subtropical zones of the global ocean: similarities and differences in terms of chlorophyll and yellow substance. *Biogeosciences*, *7*(10), 3139–3151.

Morris, J. J., Johnson, Z. I., Szul, M. J., Keller, M., & Zinser, E. R. (2011). Dependence of the cyanobacterium Prochlorococcus on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PloS One*, *6*(2), e16805.

Morris, R. M., Nunn, B. L., Frazar, C., Goodlett, D. R., Ting, Y. S., & Rocap, G. (2010). Comparative metaproteomics reveals ocean-scale shifts in microbial nutrient utilization and energy transduction. *The ISME Journal*, *4*(5), 673–685.

Morris, R. M., Rappé, M. S., Connon, S. A., Vergin, K. L., Siebold, W. A., Carlson, C. A., & Giovannoni, S. J. (2002). SAR11 clade dominates ocean surface bacterioplankton communities. *Nature*, *420*(6917), 806–810.

Naval, P., & Chandra, T. S. (2019). Characterization of membrane vesicles secreted by seaweed associated bacterium Alteromonas macleodii KS62. *Biochemical and Biophysical Research Communications*, *514*(2), 422–427.

Nekrouf, N. A., Maestre-Carballa, L., Lluesma-Gomez, M., Martinez-Hernandez, F., & Martínez-García, M. (2025). Annual dynamics and metagenomics of marine vesicles: One more layer of complexity in the dissolved organic fraction. *Limnology and Oceanography*. https://doi.org/10.1002/lno.70031

Noinaj, N., Guillier, M., Barnard, T. J., & Buchanan, S. K. (2010). TonB-dependent transporters: regulation, structure, and function. *Annual Review of Microbiology*, *64*(1), 43–60.

Oggerin, M., Viver, T., Brüwer, J., Voß, D., García-Llorca, M., Zielinski, O., Orellana, L. H., & Fuchs, B. M. (2024). Niche differentiation within bacterial key-taxa in stratified surface

waters of the Southern Pacific Gyre. *The ISME Journal*, *18*(1).

https://doi.org/10.1093/ismejo/wrae155

Okamoto, A., Saito, K., Inoue, K., Nealson, K. H., Hashimoto, K., & Nakamura, R. (2014). Uptake of

self-secreted flavins as bound cofactors for extracellular electron transfer in Geobacter

species. *Energy & Environmental Science*, *7*(4), 1357–1361.

Orench-Rivera, N., & Kuehn, M. J. (2016). Environmentally controlled bacterial vesicle-mediated

export. *Cellular Microbiology*, *18*(11), 1525–1536.

Orench-Rivera, N., & Kuehn, M. J. (2021). Differential Packaging Into Outer Membrane Vesicles

Upon Oxidative Stress Reveals a General Mechanism for Cargo Selectivity. *Frontiers in*

*Microbiology*, *12*, 561863.

Pérez-Cruz, C., Delgado, L., López-Iglesias, C., & Mercade, E. (2015). Outer-inner membrane

vesicles naturally secreted by gram-negative pathogenic bacteria. *PloS One*, *10*(1), e0116896.

Perez-Riverol, Y., Bai, J., Bandla, C., García-Seisdedos, D., Hewapathirana, S., Kamatchinathan, S.,

Kundu, D. J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., Walzer, M., Wang, S.,

Brazma, A., & Vizcaíno, J. A. (2022). The PRIDE database resources in 2022: a hub for mass

spectrometry-based proteomics evidences. *Nucleic Acids Research*, *50*(D1), D543–D552.

Pierella Karlusich, J. J., & Carrillo, N. (2017). Evolution of the acceptor side of photosystem I:

ferredoxin, flavodoxin, and ferredoxin-NADP+ oxidoreductase. *Photosynthesis Research*,

*134*(3), 235–250.

Pontiller, B., Martínez-García, S., Joglar, V., Amnebrink, D., Pérez-Martínez, C., González, J. M.,

Lundin, D., Fernández, E., Teira, E., & Pinhassi, J. (2022). Rapid bacterioplankton

transcription cascades regulate organic matter utilization during phytoplankton bloom

progression in a coastal upwelling system. *The ISME Journal*, *16*(10), 2360–2372.

Prados-Rosales, R., Weinrick, B. C., Piqué, D. G., Jacobs, W. R., Jr, Casadevall, A., & Rodriguez, G.

M. (2014). Role for Mycobacterium tuberculosis membrane vesicles in iron acquisition.

*Journal of Bacteriology*, *196*(6), 1250–1256.

Reintjes, G., Arnosti, C., Fuchs, B. M., & Amann, R. (2017). An alternative polysaccharide uptake

mechanism of marine bacteria. *The ISME Journal*, *11*(7), 1640–1650.

Reintjes, G., Tegetmeyer, H. E., Bürgisser, M., Orlić, S., Tews, I., Zubkov, M., Voß, D., Zielinski, O.,

Quast, C., Glöckner, F. O., Amann, R., Ferdelman, T. G., & Fuchs, B. M. (2019). On-site

analysis of bacterial communities of the ultraoligotrophic South Pacific Gyre. *Applied and*

*Environmental Microbiology*, *85*(14). https://doi.org/10.1128/AEM.00184-19

Renelli, M., Matias, V., Lo, R. Y., & Beveridge, T. J. (2004). DNA-containing membrane vesicles of

Pseudomonas aeruginosa PAO1 and their genetic transformation potential. *Microbiology*

*(Reading, England)*, *150*(Pt 7), 2161–2169.

Sabra, W., Lünsdorf, H., & Zeng, A.-P. (2003). Alterations in the formation of lipopolysaccharide and

membrane vesicles on the surface of Pseudomonas aeruginosa PAO1 under oxygen stress

conditions. *Microbiology*, *149*(Pt 10), 2789–2795.

Saier, M. H., Jr, Tran, C. V., & Barabote, R. D. (2006). TCDB: the Transporter Classification

Database for membrane transport protein analyses and information. *Nucleic Acids Research*,

*34*(Database issue), D181-6.

Sartorio, M. G., Pardue, E. J., Scott, N. E., & Feldman, M. F. (2023). Human gut bacteria tailor

extracellular vesicle cargo for the breakdown of diet- and host-derived glycans. *Proceedings*

*of the National Academy of Sciences of the United States of America*, *120*(27), e2306314120.

Scanlan, D. J., Ostrowski, M., Mazard, S., Dufresne, A., Garczarek, L., Hess, W. R., Post, A. F.,

Hagemann, M., Paulsen, I., & Partensky, F. (2009). Ecological genomics of marine

picocyanobacteria. *Microbiology and Molecular Biology Reviews: MMBR*, *73*(2), 249–299.

Schwechheimer, C., & Kuehn, M. J. (2015). Outer-membrane vesicles from Gram-negative bacteria:

biogenesis and functions. *Nature Reviews. Microbiology*, *13*(10), 605–619.

Schwechheimer, C., Sullivan, C. J., & Kuehn, M. J. (2013). Envelope control of outer membrane

vesicle production in Gram-negative bacteria. *Biochemistry*, *52*(18), 3031–3040.

Seymour, J. R., Amin, S. A., Raina, J.-B., & Stocker, R. (2017). Zooming in on the phycosphere: the

ecological interface for phytoplankton–bacteria relationships. *Nature Microbiology*, *2*(7), 1–

12.

Słomka, J., Alcolombri, U., Carrara, F., Foffi, R., Peaudecerf, F. J., Zbinden, M., & Stocker, R. (2023). Encounter rates prime interactions between microorganisms. *Interface Focus*, *13*(2), 20220059.

Soler, N., & Forterre, P. (2020). Vesiduction: the fourth way of HGT. *Environmental Microbiology*, *22*(7), 2457–2460.

Soler, N., Krupovic, M., Marguet, E., & Forterre, P. (2015). Membrane vesicles in natural environments: a major challenge in viral ecology [Review of *Membrane vesicles in natural environments: a major challenge in viral ecology*]. *The ISME Journal*, *9*(4), 793–796. nature.com.

Sowell, S. M., Wilhelm, L. J., Norbeck, A. D., Lipton, M. S., Nicora, C. D., Barofsky, D. F., Carlson, C. A., Smith, R. D., & Giovanonni, S. J. (2009). Transport functions dominate the SAR11 metaproteome at low-nutrient extremes in the Sargasso Sea. *The ISME Journal*, *3*(1), 93–105.

Stocker, R. (2012). Marine microbes see a sea of gradients. *Science*, *338*(6107), 628–633.

Stocker, R., & Seymour, J. R. (2012). Ecology and physics of bacterial chemotaxis in the ocean. *Microbiology and Molecular Biology Reviews*, *76*(4), 792–812.

Tang, K., Jiao, N., Liu, K., Zhang, Y., & Li, S. (2012). Distribution and functions of TonB-dependent transporters in marine bacteria and environments: implications for dissolved organic matter utilization. *PloS One*, *7*(7), e41204.

Teeling, H., Fuchs, B. M., Bennke, C. M., Krüger, K., Chafee, M., Kappelmann, L., Reintjes, G., Waldmann, J., Quast, C., Glöckner, F. O., Lucas, J., Wichels, A., Gerdts, G., Wiltshire, K. H., & Amann, R. I. (2016). Recurring patterns in bacterioplankton dynamics during coastal spring algae blooms. *ELife*, *5*, e11888.

Thumuluri, V., Almagro Armenteros, J. J., Johansen, A. R., Nielsen, H., & Winther, O. (2022). DeepLoc 2.0: multi-label subcellular localization prediction using protein language models. *Nucleic Acids Research*, *50*(W1), W228–W234.

Tinta, T., Zhao, Z., Escobar, A., Klun, K., Bayer, B., Amano, C., Bamonti, L., & Herndl, G. J. (2020). Microbial Processing of Jellyfish Detritus in the Ocean. *Frontiers in Microbiology*, *11*, 590995.

Tortell, P. D., Maldonado, M. T., Granger, J., & Price, N. M. (1999). Marine bacteria and

biogeochemical cycling of iron in the oceans. *FEMS Microbiology Ecology*, *29*(1), 1–11.

Toyofuku, M., Morinaga, K., Hashimoto, Y., Uhl, J., Shimamura, H., Inaba, H., Schmitt-Kopplin, P.,

Eberl, L., & Nomura, N. (2017). Membrane vesicle-mediated bacterial communication. *The*

*ISME Journal*, *11*(6), 1504–1509.

Toyofuku, M., Schild, S., Kaparakis-Liaskos, M., & Eberl, L. (2023). Composition and functions of

bacterial membrane vesicles. *Nature Reviews. Microbiology*. https://doi.org/10.1038/s41579-

023-00875-5

Toyofuku, M., Tashiro, Y., Hasegawa, Y., Kurosawa, M., & Nomura, N. (2015). Bacterial membrane

vesicles, an overlooked environmental colloid: Biology, environmental perspectives and

applications. *Advances in Colloid and Interface Science*, *226*(Pt A), 65–77.

Van Heukelem, L., & Thomas, C. S. (2001). Computer-assisted high-performance liquid

chromatography method development with applications to the isolation and analysis of

phytoplankton pigments. *Journal of Chromatography A*, *910*(1), 31–49.

Viborg, A. H., Terrapon, N., Lombard, V., Michel, G., Czjzek, M., Henrissat, B., & Brumer, H.

(2019). A subfamily roadmap of the evolutionarily diverse glycoside hydrolase family 16

(GH16). *The Journal of Biological Chemistry*, *294*(44), 15973–15986.

Villanueva, R. A. M., & Chen, Z. J. (2019). ggplot2: Elegant Graphics for Data Analysis (2nd ed.).

*Measurement: Interdisciplinary Research and Perspectives*, *17*(3), 160–167.

Wang, M., Nie, Y., & Wu, X.-L. (2021). Membrane vesicles from a Dietzia bacterium containing

multiple cargoes and their roles in iron delivery. *Environmental Microbiology*, *23*(2), 1009–

1019.

Worden, A. Z., Follows, M. J., Giovannoni, S. J., Wilken, S., Zimmerman, A. E., & Keeling, P. J.

(2015). Environmental science. Rethinking the marine carbon cycle: factoring in the

multifarious lifestyles of microbes. *Science (New York, N.Y.)*, *347*(6223), 1257594.

Yaron, S., Kolling, G. L., Simon, L., & Matthews, K. R. (2000). Vesicle-mediated transfer of

virulence genes from Escherichia coli O157:H7 to other enteric bacteria. *Applied and*

*Environmental Microbiology*, *66*(10), 4414–4420.

Yonezawa, H., Osaki, T., Kurata, S., Fukuda, M., Kawakami, H., Ochiai, K., Hanawa, T., & Kamiya,

942           S. (2009). Outer membrane vesicles of Helicobacter pylori TK1402 are involved in biofilm

943           formation. *BMC Microbiology*, *9*, 197.

944    Yuan, Z., Liu, H., Gosnell, K. J., Wen, Z., Engel, A., Dai, M., Achterberg, E. P., & Browning, T. J.

945           (2025). Localized nutrient colimitation of phytoplankton growth rates across the subtropical

946           South Pacific Ocean. *Proceedings of the National Academy of Sciences of the United States of*

947           *America*, *122*(50), e2526930122.

948    Zhang, Z., Xu, A., Hathorne, E., Gutjahr, M., Browning, T., Gosnell, K. J., Liu, T., Steiner, Z., Kiko,

949           R., Yuan, Z.-X., Liu, H., Achterberg, E. P., & Frank, M. (2024). Substantial trace metal input

950           from the 2022 Hunga Tonga-Hunga Ha'apai eruption into the South Pacific. *Nature*

951           *Communications*, *15*(1), 8986.

952    Zhu, W.-J., Wang, C., Liu, L., Li, J.-X., Wang, H.-Q., Wang, M.-Q., Cao, H.-Y., Chen, X.-L., Qin,

953           Q.-L., Zhang, Y.-Z., Sun, M.-L., & Wang, P. (2025). Structural and molecular basis for

954           phosphate recognition by SAR11 bacteria. *MBio*, *16*(9), e0165425.

955    Zhu, Y. (2018). *DEqMS*. Bioconductor. https://doi.org/10.18129/B9.BIOC.DEQMS

956    Zlatkov, N., Nadeem, A., Uhlin, B. E., & Wai, S. N. (2020). Eco-evolutionary feedbacks mediated by

957           bacterial membrane vesicles. *FEMS Microbiology Reviews*.

958           https://doi.org/10.1093/femsre/fuaa047

959    Zoccarato, L., Sher, D., Miki, T., Segrè, D., & Grossart, H.-P. (2022). A comparative whole-genome

960           approach identifies bacterial traits for marine microbial interactions. *Communications*

961           *Biology*, *5*(1), 276.