1 **One-step induction of human GABAergic neurons promotes presynaptic development**
2 **& synapse maturation**
3

Torben W. van Voorst[1,+], Maaike A. van Boven [1,+], Kevin I. Marinus[2,+], Jennifer M.
Colón-Mercado[3], Jule Schretzmeir[1], Carolin Haag[1,4], Ruud F. Toonen[1,4], Frank
Koopmans[2], Michael E. Ward[3], August B. Smit[2], Ronald E. van Kesteren[2], Matthijs
Verhage[1,4,5], L. Niels Cornelisse[1,5,#]

[1]Department of Functional Genomics, Center for Neurogenomics and Cognitive Research (CNCR),
VU University Amsterdam, de Boelelaan 1100, 1081 HZ Amsterdam, The Netherlands.
[2]Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive
Research (CNCR), VU University Amsterdam, de Boelelaan 1100, 1081 HZ Amsterdam, The
Netherlands.
[3]National Institute of Neurological Disorders and Stroke, National Institutes of Health, 9000
Rockville Pike, Bethesda, MD, USA
[4]Neurospector, de Boelelaan 1100, 1081 HZ Amsterdam, The Netherlands.
[5]Department of Human Genetics, Amsterdam UMC – location VU University Amsterdam, de
Boelelaan 1100, 1081 HV Amsterdam, The Netherlands.
18
[+] Authors contributed equally
[#] Correspondence: l.n.cornelisse@vu.nl

Abstract (174 words)
Human induced pluripotent stem cells (iPSCs) present a powerful approach to study human
brain physiology and disease, yet robust, pure GABAergic induction has remained difficult.
Here we present improved, single-step, transposon-based GABAergic induction with
Ascl1/Dlx2, which yields pure GABAergic neurons, in contrast to lentiviral approaches, and
was tested across three independent iPSC lines. Proteomic and electrophysiological
characterization at different developmental time points showed that these neurons gain a
proteomic profile that maps to different cortical interneuron subtypes, particularly VIP+
interneurons, and display typical GABAergic synaptic properties, producing large,
synchronous and picrotoxin-sensitive currents. During early development synaptic strength
increased threefold, which was accompanied by enhanced expression of multiple GABA-
specific presynaptic gene sets, but few changes in postsynaptic gene sets. Synaptic strength
continued to improve during late development but with only minor proteomic changes. Co-

seeding with NGN2 neurons created stable networks of predefined excitation/inhibition
ratios, with corresponding synapse ratios. Taken together, transposon-based GABAergic
induction yields pure, mature GABAergic neurons suitable for studying gene sets involved in
synaptic maturation and to build excitation/inhibition networks for disease modelling.
39

Keywords (max 10)

41 Induced pluripotent stem cells, induced GABAergic neurons, Transposon, Synaptic maturation,
42 Ascl1, Dlx2, Autaptic neurons, Proteomics, Patch-clamp, Excitation/inhibition
43

Introduction

Human induced pluripotent stem cell (iPSC)-derived models provide a promising platform
for studying human brain physiology and pathology. iPSCs can be generated from patient-
derived tissue or blood to better reflect the human genetics and (patho)physiology compared
to commonly employed rodent models (Breschi *et al*, 2017; Gharib & Robinson-Rechavi,
2011; Hoffman *et al*, 2019). GABAergic dysfunction and excitation/inhibition (E/I) disbalance
have been implicated in many neurodevelopmental and neurodegenerative disorders
(Busche *et al*, 2008; Foss-Feig *et al*, 2017; Nelson & Valakh, 2015; Palop *et al*, 2007a;
Rubenstein & Merzenich, 2003; Sohal & Rubenstein, 2019). In recent years, various
strategies have been developed for generating iPSC-derived GABAergic neurons, but robust,
pure GABAergic induction remains challenging. Directed differentiation, which employs
signalling molecules to mimic neurodevelopment (Liu *et al*, 2013; Maroof *et al*, 2013;
Nicholas *et al*, 2013), yields both glutamatergic and GABAergic neurons, but these type of
inductions often generate inconsistent E/I ratios, tend to yield heterogeneous and immature
neuronal populations and are costly and time-consuming (Engle *et al*, 2018). Rapid
differentiation approaches rely on overexpression of key transcription factors, such as Ascl1
and Dlx2 for GABAergic differentiation (Mossink *et al*, 2022; Yang *et al*, 2017) and often rely
on lentiviral delivery of transgenes. However, such approaches have several limitations: (1)
they require multiple vectors when delivering multiple transgenes due to the size cap of ~8
kb, (2) they pose a risk of disrupting host genes (Baum, 2007), and (3) they are susceptible
to epigenetic silencing during prolonged iPSC culturing (Cabrera *et al*, 2022; Wolf & Goff,
2009; Yao *et al*, 2004). To overcome these challenges, TALEN- and CRISPR-based gene
editing (Miller *et al*, 2011) have been used to target the transgenes to the *AAVS1* safe harbour
locus (Peitz *et al*, 2020; Schmid *et al*, 2021). While this ensures stable transgene expression,
allows for delivery of multiple genes in a single construct, and minimizes off-target effects,

it is time and resource-consuming due to the need for clonal expansion and extensive
validation of every iPSC line used in a study (Lombardo *et al*, 2011).

The PiggyBac transposon system has emerged as a gene-delivery tool that overcomes
most of these limitations. Unlike lentiviral systems, transposons enable stable genomic
integration of large genetic constructs up to 200 kb (Li *et al*, 2013) through a "cut-and-paste"
mechanism (Wilber *et al*, 2007), which enables delivery of multiple transgenes in a single
construct. Additionally, transposon-delivered transgenes are less prone to silencing than
lentivirally delivered transgenes (Ng *et al*, 2021) and the increased packing capacity permits
the inclusion of epigenetic regulators to further diminish silencing. Compared to safe-
harbour insertion of transgenes, the delivery of a transposon cassette is a simpler, single-
step process applicable to multiple iPSC lines in parallel and allows the use of a polyclonal
population for most use cases, eliminating the need for thorough validation, thereby
accelerating experimental timelines.

Here, we characterize PiggyBac transposon-based overexpression of *Ascl1* and *Dlx*2 (PB-
AD) for GABAergic neuronal induction and benchmark this against established GABAergic
induction methods. PB-AD-based induction results in pure GABAergic differentiation across
different genetic backgrounds, with proteomic signatures characteristic of several human
cortical interneuron types, with the strongest resemblance to VIP[+] interneurons, unlike
lentiviral-based AD expression, which induces both GABAergic and glutamatergic
differentiation. We demonstrate that PB-AD neurons integrate into synaptically connected
E/I networks when co-cultured with excitatory NGN2 neurons with synaptic projections
proportional to cellular ratios. Using proteomics, patch-clamp electrophysiology, and
immunocytochemistry we show that functional synapse maturation correlates with
increasing levels of proteins in presynaptic gene sets, resulting in reliable, plastic synapses
that synchronously release GABA, whereas postsynaptic function and protein composition
are stable over time. We propose that this improved GABAergic induction protocol, which
reliably generates pure GABAergic neurons, enables the study of human GABAergic
functioning in pure GABAergic networks, in single-neuron (autaptic) cultures, and stable E/I-
networks.

Results

GABAergic differentiation of human iPSCs using PiggyBac based delivery of Ascl1 and Dlx2
Overexpression of the transcription factors Ascl1 and Dlx2 drives GABAergic lineage
specification of induced pluripotent stem cells (iPSCs) (Yang *et al.*, 2017). Previous approaches

employed separate lentivectors to express both genes in an inducible manner. We aimed to
develop a simplified GABAergic differentiation protocol using a single PiggyBac transposon
construct containing both transcription factors (Figure 1A). This vector contains (1) a
doxycycline (dox)-inducible expression cassette for *Ascl1* and *Dlx2* from a bidirectional
promoter, avoiding variation in expression levels resulting from position effects in multi-
cistronic vectors; (2) reverse tetracycline-controlled transactivator (rtTA), required for dox-
inducible expression; (3) the puromycin resistance gene (pac), allowing for antibiotic
selection, as well as (4) mTagBFP2 (Subach *et al*, 2011), giving transfected cells a fluorescent
marker; and (5) Ubiquitous Chromatin Opening Elements (UCOE) (Müller-Kuller *et al*, 2015)
and 5'HS chicken β-globin (cHS4) insulator elements (Chung *et al*, 1997). Firstly, we tested
whether our newly designed construct could induce GABAergic differentiation (Figure 1B). To
this end, we co-transfected this PiggyBac vector with a transposase vector into BIONi010-C
iPSCs (Rasmussen *et al*, 2014). After puromycin-selection, all cells expressed mTagBFP2
(Figure 1C), confirming successful transgene integration. Upon dox-induction, cells first
clustered together (Figure 1D) and after releasing and reseeding at 3 days post-induction
(DPI), cells developed a neuronal morphology, which became more distinct within one week
(Figure 1E). At 10 weeks post-induction (WPI), neurons displayed MAP2+ dendrites and VGAT[+]
puncta, indicative of GABAergic synapses, whereas glutamatergic VGLUT1/2[+] puncta were
absent (Figure 1F,G).

GABAergic synaptic transmission in PiggyBac Ascl1/Dlx2 neurons
To confirm the GABAergic identity of PiggyBac-induced neurons we performed voltage-clamp
recordings in autaptic cultures (Js *et al*, 2019) between 6 and 10 WPI (Figure 1H). Rapid
depolarization to +30 mV induced action potential evoked postsynaptic currents (ePSCs),
which displayed slow decay times (τ) characteristic of GABAergic currents (Jones &
Westbrook, 1995), as determined by mono-exponential fits (geometric mean (GM) = 104.2 ms,
geometric SD factor (GM_{SDf}) = 1.42; Figure 1I,J). Furthermore, virtually all synaptic signalling
was abolished after applying the GABA receptor antagonist picrotoxin (PTX), confirming
GABAergic synaptic identity (Figure 1I,K).

PiggyBac Ascl1/Dlx2 yields a pure GABAergic neuron population in contrast to lentiviral methods
To benchmark the PiggyBac approach, we compared it with two established lentiviral
methods for GABAergic induction (Mossink *et al.*, 2022; Yang *et al.*, 2017) and an NGN2-based
protocol for glutamatergic induction (Meijer *et al*, 2019; Schmid *et al.*, 2021) (Figure 1L).
Autaptic neurons were generated using all four protocols and ePSC decay time constants
were quantified using mono-exponential fits. As expected, NGN2-derived neurons exhibited

fast ePSCs decay kinetics typical of glutamatergic transmission (GM = 2.43 ms, GM_{SDf} = 1.39,
Figure 1M,N), confirmed pharmacologically and by the presence of VGLUT1/2$^+$ puncta (Figure
S1). Surprisingly, decay time constants from neurons generated with the lentiviral GABAergic
protocols showed a bimodal distribution which could be fitted with a sum of two gaussians.
More than half of the cells in these cultures exhibited ePSC kinetics comparable to those in
the NGN2 cultures (Lenti AF: GM = 2.82 ms, GM_{SDf} = 0.58, 72.8% of the fit; Lenti AD: GM = 4.62
ms, GM_{SDf} = 0.42, 55.3% of the fit). The remaining cells had more than tenfold larger time
constants typical for slow GABAergic ePSCs (Lenti AF: GM = 86.3 ms, GM_{SDf} = 1.56 for 27,2%
of the fit; Lenti AD: GM = 60.5 ms, GM_{SDf} = 1.26 for 44.7% of the fit). In contrast, the PB-AD
method produced a unimodal population with slowly decaying ePSCs (GM = 78.9 ms, GM_{SDf}
= 1.35). Since NGN2 and PB-AD neurons produced decay time constants with unimodal fits,
these values were used to determine cut-off thresholds for synaptic identity (glutamatergic:
$\tau \leqslant 6$ ms; GABAergic: $\tau \geqslant 30$ ms; mixed: $6 < \tau < 30$ ms). This revealed that 68% of PB-AD
neurons exhibited GABAergic currents, while the remaining 32% were synaptically silent
(Figure 1O). In contrast, the lentiviral approaches generated a substantial population of
glutamatergic (Lenti AF: 43%, Lenti AD: 39%) and mixed (Lenti AF: 9% Lenti AD: 15%)
identities. The consistent generation of functional GABAergic neurons and the absence of
glutamatergic neurotransmission underscore the reliability of the PB-AD protocol compared
to previously described methods.

PiggyBac Ascl1/Dlx2 protocol induces GABAergic neurons across genetic backgrounds
To test for generalizability, we applied the PB-AD protocol to generate neurons from two
additional iPSC lines, KOLF2.1J (Pantazis *et al*, 2022) and GM23973 (Nadadhur *et al*, 2019),
and characterized their functional properties at 7 and 6 WPI, respectively. In both lines the
PB-AD approach produced neurons with GABAergic synaptic projections and slow-decaying
synaptic transmission, indicative of GABAergic specification (Figure S2), as was found for the
BIONi010-C line. These findings demonstrate the protocol's applicability across genetic
backgrounds.

PiggyBac Ascl1/Dlx2 neurons integrate into controlled excitatory/inhibitory networks
To test whether PB-AD neurons integrate into E/I networks alongside glutamatergic neurons,
we co-cultured 3 DPI PB-AD neurons with 3 DPI glutamatergic NGN2 neurons in networks
containing 100%, 30% (as found in the human cortex (Bakken *et al*, 2021; Fang *et al*, 2022;
Loomba *et al*, 2022)) and 0% inhibitory neurons (Figure 2A-C). Immunocytochemistry at 10
WPI (Figure 2B,C) revealed comparable neuronal cell densities across all conditions, as
measured by NeuN staining, indicating similar viability of PB-AD and NGN2 neurons. (Figure

2D). Total dendrite length per field of view increased as the proportion of PB-AD neurons
decreased, consistent with the distinct morphological characteristics of GABAergic neurons
(Chang *et al*, 2014) (Figure 2E). Pure PB-AD networks contained VGAT$^+$ (GABAergic) synaptic
puncta, and NGN2 networks contained VGLUT1/2$^+$ (glutamatergic) synaptic puncta, with in
both cases virtually no puncta of the other type present (Figure 2F,G). In contrast, the mixed
(30% PB-AD) networks contained both GABAergic and glutamatergic puncta on the same
dendrite proportional to cellular ratios, demonstrating that PB-AD neurons are viable
components for assembling E/I networks when mixed with NGN2 neurons.

PiggyBac Ascl1/Dlx2 neurons acquire a GABAergic molecular identity over time
Next, we assessed differentiation of PB-AD and NGN2 neurons over time using mass
spectrometry-based proteomics. PB-AD co-cultures with rat astrocytes were compared to
NGN2 co-cultures at 4, 6, 8, and 10 WPI (Figure 3A). We detected 99,154 human peptide
precursors mapping to 7,775 unique proteins across 60 samples that passed quality control.
Variability within groups was low-to-moderate with coefficients of variation between 14.6%
and 32.2%, reflecting consistent detection across replicates (Figure S3A-B). To remove rat
astrocytic signal, subsequent analyses focused on neuronally enriched proteins (1690
proteins). Principal component analysis (PCA) on the neuronally-enriched protein
abundances (Figure 3B, S3C) revealed clear separation between PB-AD and NGN2 cultures
along the first two components (PC1: 71.5% and PC2: 6.1%), as well as the separation of
different time points within each culture. More specifically, time points from both PB-AD and
NGN2 cultures showed clear separation between 4, 6, and 8 WPI, while cultures at 8 and 10
WPI clustered more closely together, suggesting that proteomic maturation plateaus after 8
weeks.

To assess lineage specification of PB-AD and NGN2 inductions, we annotated cell
types based on differentially expressed proteins using cell type specificity scores (CTSS),
computed using single-cell RNA expression data from human cortex (Jorstad *et al*, 2023)
(Figure S4A). Across all timepoints, canonical GABAergic proteins, including SLC32A1 (VGAT),
GAD1, GAD2, were enriched in PB-AD cultures. In addition, other predicted GABAergic
proteins, such as CRHBP, FBN2, CRH, and RELN, were enriched in PB-AD neurons. In contrast,
only a few predicted glutamatergic proteins (e.g., BDNF, PCP4, CPZ, and ADAMTSL1) were
moderately enriched in PB-AD cultures. The glutamatergic marker SLC17A6 (VGLUT2) was
enriched in NGN2 cultures, while SLC17A7 (VGLUT1) was not detected (Figure 3C-F).
Surprisingly, NGN2 neurons also showed strong enrichment of several proteins typically
associated with GABAergic neurons at all timepoints, including SCGN, CALB2, and PVALB.

Immunocytochemistry confirmed PVALB expression in the majority of NGN2 neurons (Figure
S4B).

To quantitatively assess cell type specification, we performed expression-weighted
cell type enrichment (EWCE) analysis (Skene & Grant, 2016) on the top 5% upregulated
neuronal proteins for each induction for each timepoint. At all timepoints, PB-AD-specific
neuronal proteins showed exclusive and significant enrichment for GABAergic neurons, with
enrichment levels increasing until 8 WPI (Figure 2G). GABAergic subclass analysis revealed
strongest enrichment for VIP$^+$ interneurons from 6-10 WPI (Figure S4C). In contrast, NGN2-
specific proteins showed significant enrichment for glutamatergic neurons only at 6 and 10
WPI, while also displaying unexpected GABAergic enrichment, reaching significance at 4 and
10 WPI (Figure 3H). Subclass enrichment analysis revealed that this unexpected GABAergic
enrichment was most likely accounted for by PVALB$^+$ and VIP$^+$ interneuron-like profiles (S4C,
S4D). Taken together, PB-AD neurons acquire a stable GABAergic identity over time,
characterized by consistent expression of canonical and subtype-specific GABAergic proteins.
By contrast, NGN2-induced neurons exhibit a mixed molecular identity, with evidence for
partial GABAergic specification in addition to expected glutamatergic features.

Gradual morphological development of PiggyBac Ascl1/Dlx2 neurons
To examine morphological maturation of PB-AD neurons, we performed immunostaining for
MAP2, VGAT and VGLUT1/2 in autaptic cultures (Figure 4A). Similar to networks, PB-AD
neurons produced only VGAT$^+$ puncta at all timepoints (Figure 4A-C). A progressive increase
in dendrite length was observed during maturation (Figure 4D), accompanied by an increase
in the total number of VGAT$^+$ puncta per neuron and puncta densities per dendritic length
(Figure 4E-F). Additionally, the intensities of VGAT puncta increased over time (Figure 4G),
suggesting increased synaptic vesicles per synaptic terminal. Sholl analysis revealed a rise
in the number of arborizations during maturation (Figure 4H,I). Interestingly, growth of
dendritic length and complexity plateaued by 6 WPI, whereas synaptic projections continued
to develop until 8 and 10 WPI. Together, these results demonstrate that PB-AD neurons
develop complex neuronal morphologies accompanied by an increase in presynaptic
GABAergic terminals during maturation.

Synaptic strengthening during maturation of PiggyBac Ascl1/Dlx2 neurons
Next, we assessed functional maturation in autaptic neurons with voltage-clamp
electrophysiology. Firstly, action-potential independent spontaneous release was recorded.
Both miniature postsynaptic current (mPSC) frequency and amplitude increased during
maturation (Figure 5A-C). Next, neurons were depolarized to induce ePSCs. Across all time

points, 70-80% of neurons were synaptically active, producing a charge of at least 1.5 pC
within 70 ms after stimulation (Figure 5D). In active cells, both the amplitude and charge of
ePSCs increased progressively between 4 and 10 WPI (Figure 5E-G), with the most
pronounced change occurring from 8 to 10 WPI. Decay constants remained consistently
above 30 ms – characteristic of GABAergic currents (Figure 5H). Interestingly, decay kinetics
subtly slowed after 4 WPI, suggesting potential changes in postsynaptic receptor subunits
(Dixon *et al*, 2014; Zhang *et al*, 1997).

To evaluate the impact of maturation on short-term synaptic plasticity, we applied
paired-pulse stimulation at 50 ms, 200 ms, and 1000 ms intervals (Figure 5I-L). This revealed
modest synaptic depression which was stable throughout maturation. Only at a 1000 ms
interval, synaptic depression increased as neurons matured. Importantly, maturation
increased the size of the readily releasable pool (RRP), as determined by back-extrapolation
of the charge evoked by 20 Hz stimulus trains, with the biggest change occurring from 8 to
10 WPI (Figure 5M-O). Taken together, these findings demonstrate that PB-AD GABAergic
neurons develop functional, plastic synapses that strengthen during maturation.

Functional maturation correlates with upregulation of proteins in presynaptic gene sets
To investigate the molecular basis of synaptic maturation, we looked at the differential
expression of SYNGO-annotated synaptic proteins between different timepoints of
development (4 to 6 WPI, 6 to 8 WPI, and 8 to 10 WPI) (Koopmans *et al*, 2019a) (Figure 6A).
Of all 1605 SYNGO-annotated synaptic genes, 929 unique proteins were detected in PB-AD
cultures, of which 478 (51%) were differentially expressed in at least one time contrast
(Figure 6B, D-F). In NGN2 neurons, 967 unique SYNGO-annotated proteins were identified of
which 644 (67%) were differentially expressed in at least one time contrast (Figure 6C, Figure
S5A-C). In both cultures, SYNGO-annotated proteins were abundantly regulated between 4
and 8 WPI, while only few proteins showed altered expression between 8 and 10 WPI (Figures
6B-F, Supplementary table 1). This indicates that proteomic synaptic maturation plateaus
after 8 WPI, consistent with what was observed for neuronal-enriched protein expression.

To examine specifically how pre- and postsynapses develop over time, we used GOAT
(Koopmans, 2024a) to determine which SYNGO gene sets were enriched for developmentally
regulated proteins, where the contribution of each protein was weighted by its $\log_2$ (fold
change) (Koopmans, 2024a). Strongest enrichment was observed between 4-6 WPI and 6-8
WPI, which occurred exclusively in presynaptic gene sets, including *synaptic vesicle* (*SV*),
presynaptic active zone (AZ), and *presynaptic membrane* (Figure 6G-L). Between 8-10 WPI only
SV membrane proteins and its mother term *presynaptic* proteins were enriched. In contrast,

no postsynaptic gene sets were significantly enriched in PB-AD cultures, across all time contrasts. To exclude that these results were biased by a few proteins with large effect sizes, additional SYNGO gene set overrepresentation analysis was performed based on the number of upregulated proteins per gene set. This confirmed the exclusive enrichment of presynaptic terms in PB-AD cultures (Figure S5J–L). Upregulated proteins in NGN2 cultures over time similarly showed a preferential enrichment of presynaptic gene sets using GOAT and SYNGO overrepresentation analyses, with no enrichment for postsynaptic gene sets between 6-10 WPI. Thus, synaptic strengthening correlates with changes specifically in presynaptic protein levels, while postsynaptic protein levels remain relatively stable over time.

In order to test whether our findings on asymmetrical pre- and postsynaptic maturation translate to a more physiological model, we assessed postsynaptic responses to externally applied GABA in E/I networks, as excitation might alter maturation profiles (Paraskevopoulou *et al*, 2019) (Figure 6M). Furthermore, we used gramicidin-perforated patch to allow measurement of postsynaptic responses without influencing the reversal potential for the GABA-receptor (Watanabe & Fukuda, 2015). GABA puffs were applied at varying holding voltages (Figure 6N). The slope of the I/V curve voltages was unaltered by maturation ($p = 0.305$), indicating that the GABAergic postsynaptic response does not change during maturation under these conditions. Thus, increased synaptic strength during synaptic maturation correlates with presynaptic proteomic changes, but not with postsynaptic development also in E/I-networks.

PiggyBac Ascl1/Dlx2 specific synaptic proteins map to cortical interneuron subtypes

Finally, we investigated whether changes in PB-AD and NGN2 SYNGO-annotated proteomes between 4 and 10 WPI are associated with cortical GABAergic and glutamatergic cell types (Jorstad *et al.*, 2023) (Figure 7A). PB-AD cultures showed upregulation of predicted GABAergic and medial ganglionic eminence (MGE) and caudal ganglionic eminence (CGE)-specific proteins (Figure 7B). NGN2 cultures showed upregulation of predicted glutamatergic proteins (Figure 7C). Of note, PB-AD and NGN2 cultures showed upregulation of both GABAergic and glutamatergic receptor subunits (Figure S6A,B), although only input of one of the neurotransmitters was present per culture. Canonical subtype markers for both MGE- and CGE-derived subtypes were also detected in PB-AD neurons (S6C,D).

To analyze lineage-specific synaptic development in more detail, we next contrasted the developing (WPI 4 to WPI 10) PB-AD and NGN2 SYNGO-annotated proteomes with each other to find proteins that were specifically regulated in each induction. To this end, for each protein we calculated the difference between PB-AD and NGN2 in differential expression

between 4-10 WPI (i.e., $\log_2 (FC)_{AD}$- $\log_2 (FC)_{NGN2}$), and selected the 5th and 95th percentile to
define induction-specific synaptic proteins (Figure 7A). We then assessed whether these
proteins were enriched in either cortical glutamatergic or cortical GABAergic reference
expression profiles (Jorstad *et al.*, 2023). Overall, this produced lineage-specific enrichments
in most gene sets. PB-AD specific synaptic proteins showed the highest GABAergic
enrichment in the gene sets *dense core vesicle* (*DCV*), *SV*, *synaptic signalling*, *metabolism* and
presynaptic process (Figure 7D-E). Conversely, NGN2-specific synaptic proteins that belonged
to *synaptic signalling, SV, organization* and *specialization* showed the highest glutamatergic
enrichment (Figure 7F-G).

As synaptic gene expression profiles have been shown to reflect neuronal subtypes
(Roig Adam *et al*, 2023), we queried whether synaptic proteins that specifically increase in
PB-AD or NGN2 cultures show enrichment for neuronal (sub)types. In EWCE analysis, the PB-
AD-specific synaptic proteins were enriched in GABAergic neurons (Figure 7H), specifically
VIP+, SST+, PAX6+ and LAMP5+ interneurons, but not PVALB+ neurons (Figure 7I). NGN2-specific
synaptic proteins were enriched in glutamatergic subtypes (IT, L4 IT, L5 ET, L6 CT, and L6b)
but also in GABAergic subtypes (Figure 7J), particularly VIP+ interneurons (Figure 7K).
Together, these data indicate that PB-AD-based induction leads to GABAergic synaptic
specialization with molecular profiles resembling CGE- and MGE-derived interneurons,
whereas NGN2 induction leads to induction of neurons that also contain typical GABAergic
proteins.

Discussion

Summary
In this work, we present a single-step, transposon-based GABAergic differentiation method,
which uses a single vector to overexpress Ascl1/Dlx2, and we provide a detailed
characterization of these GABAergic neurons at different developmental timepoints using
mass spectrometry proteomics, immunocytochemistry, and patch-clamp electrophysiology.
This method is applicable across different genetic backgrounds and produces exclusively
GABAergic neurons with a proteomic profile that maps to different cortical interneuron
subtypes, particularly VIP+ neurons. Synapses display GABAergic properties such as
picrotoxin-sensitive, slowly decaying, evoked postsynaptic currents and short-term
depression, similar to primary GABAergic neurons (Ikeda *et al*, 2008; Jiang *et al*, 2000; Lambert
& Wilson, 1994; Pearce *et al*, 1995). During development, synapses gradually increase in
synaptic strength, which at earlier timepoints correlates with enhanced expression of specific

presynaptic gene sets associated with the active zone, synaptic vesicle, and presynaptic
membrane. In contrast, little functional and proteomic changes occur in the postsynapse.
Late in development synaptic strength continues to increase, with only minor changes at the
proteomic level. PB-AD neurons can be combined with NGN2 neurons to form networks of
predefined E/I ratios with corresponding synapse ratios. Taken together, transposon-based
GABAergic induction yields pure GABAergic neurons with improved synaptic properties that
can be used to build balanced excitation/inhibition networks, for instance for disease
modelling in those (many) cases where E/I disbalance has been proposed as a disease
mechanism.

PiggyBac Ascl1/Dlx2 neuronal induction produces exclusively GABAergic neurons
By employing autaptic neurons we were able to detect synaptic output on a single neuron
scale, which allowed us to accurately quantify lineage specification and demonstrate how
pure GABAergic transposon-based induction is compared to lentiviral strategies. Previous
studies using a lentiviral approach also detected non-GABAergic currents (Gonzalez-Ramos
et al, 2021) and reported expression of glutamatergic genes (Barretto *et al*, 2020; Song *et al*,
2021), while others focused on measurements in E/I-networks (Mossink *et al.*, 2022; Yang *et
al.*, 2017) or recorded under conditions that blocked glutamatergic transmission (Marcó de la
Cruz *et al*, 2024; Sanchez-Priego *et al*, 2022), making it difficult to compare with our results.
As Ascl1 expression, by itself, induces glutamatergic differentiation (Chanda *et al*, 2014; Wu
et al, 2024), the balance of Ascl1 and Dlx2 expression is likely important in steering
differentiation towards a GABAergic fate. The design of our transposon-based induction
incorporates multiple features to facilitate this and thereby avoids unwanted glutamatergic
specification. Firstly, both factors are present in one cassette and transcribed from a
bidirectional promotor. Secondly, a shared integration site could prevent position effects
(Gierman *et al*, 2007). Lastly, insulator elements may reduce gene silencing, which is observed
for lentiviral cargo (Cabrera *et al.*, 2022; Gödecke *et al*, 2017; He *et al*, 2005; Kwak *et al*, 2024;
Liew *et al*, 2007; Wolf & Goff, 2009; Yao *et al.*, 2004), and less so for transposon delivered
transgenes (Ng *et al.*, 2021).

Proteomic profiles of PiggyBac Ascl1/Dlx2 neurons map to cortical interneuron subtypes
Cell type enrichment analysis revealed that neuronal proteins that are higher expressed in
PB-AD cultures are significantly enriched for GABAergic neurons at all timepoints, with
specific enrichment for VIP$^+$ interneurons from 6 WPI onwards. Zooming in on synaptic
proteins that are upregulated specifically in PB-AD cultures between 4-10 WPI showed
significant enrichment for all GABAergic interneuron subtypes except PVALB$^+$, suggesting

specification towards both MGE- and CGE-derived neurons. In a previous single-cell RNA-sequencing study of lenti-AD induced neurons, predominantly SST$^+$ neurons were detected at 4 WPI, but later time points were not investigated (Allison *et al*, 2021). Perhaps CGE markers only appear further along the maturation process, or transcriptomics do not accurately recapitulate the molecular identity of these neurons (Lindhout *et al*, 2020; Moritz *et al*, 2019; Titlow *et al*, 2023).

NGN2-induction exhibited a mixed glutamatergic and GABAergic molecular identity, although virtually no VGAT puncta were identified and only glutamatergic ePSCs were recorded, consistent with previous reports (Bullmann *et al*, 2024; Fenske *et al*, 2019; Lin *et al*, 2021; Meijer *et al.*, 2019; Nehme *et al*, 2018; Rhee *et al*, 2019; van Berkel *et al*, 2024). Thus, its proteomic profile does not translate to mixed functional identity, making it a good model for pure glutamatergic neurons.

Enhanced expression of proteins in presynaptic gene sets during synaptic maturation

Our multilevel analysis of GABAergic cultures at different time points between 4 and 10 WPI, combining patch-clamp electrophysiology, immunocytochemistry, and mass spec proteomics, revealed the following insights about synapse maturation: Firstly, synaptic strength increased approximately 3-fold during this period. A comparable increase in mPSC frequency and RRP suggests that this is due to an increase in the strength and/or number of presynaptic projections, which is supported by VGAT immunocytochemistry. Similar development has been described for directed-differentiation and safe-harbor AD GABAergic autaptic neurons (Meijer *et al.*, 2019; Rhee *et al.*, 2019).

Secondly, we observed a stark difference between pre- and postsynaptic development. Strong upregulation of *AZ-, SV-* and *presynaptic membrane* proteins occurred, while expression changes in proteins in postsynaptic gene sets were minimal. Current responses to GABA puffs remained constant over time, indicating that functional postsynapses already exist at 4 WPI and the expression of GABA receptors remains unaltered at subsequent weeks. This fits with the notion that expression of postsynaptic neurotransmitter receptors precedes presynaptic innervation (Friedman *et al*, 2000; Kwon & Sabatini, 2011; Mammen *et al*, 1997; Petralia, 2012; Richards *et al*, 2005). Release from presynaptic projections subsequently induces clustering of postsynaptic molecules, initiating postsynaptic maturation (Andreae & Burrone, 2015; Friedman *et al.*, 2000; Harms *et al*, 2005; Okabe *et al*, 2001; Sanes & Lichtman, 1999). Indeed, alignment of the pre- and postsynapse and/or clustering of postsynaptic receptors could explain the 1.5-fold increase in mPSC amplitude with the total number of GABA receptors remaining constant. A similar asymmetric proteomic development of the pre- and

postsynapse was observed in NGN2 neurons and has previously also been described for
directed differentiation of iPSC-derived neurons (Nadadhur *et al*, 2017; Togo *et al*, 2021).

Thirdly, functional development follows molecular development with a delay.
Proteomic changes abruptly stop after 8 WPI, while the number of synapses and synaptic
strength continues to increase beyond this point. Possibly, the bulk of *AZ, SV* and *presynaptic*
membrane proteins that are upregulated until 8 WPI are needed to create functional
presynapses. Enhancement of GABAergic synaptic strength after this timepoint could rely on
cell intrinsic and/or astrocyte-induced (Chung *et al*, 2015; Elmariah *et al*, 2004; Liu *et al*, 2022)
structural rearrangement, as well as posttranslational modifications of synaptic proteins,
which can occur via various mechanisms. For example, posttranslational modification,
mediated by e.g. ERK, PKC, PKA, CaMKII and/or RhoA/ROCK signaling, affects many pre- and
postsynaptic proteins, including gephyrin (Flores *et al*, 2015; Ghosh *et al*, 2016; Lorenz-
Guertin & Jacob, 2018; Tyagarajan *et al*, 2013; Tyagarajan *et al*, 2011), GABA-receptors (Bright
& Smart, 2013; Farzana *et al*, 2016; Hao *et al*, 2023; Hausrat *et al*, 2015; Houston *et al*, 2009;
Lorenz-Guertin & Jacob, 2018; Lützenkirchen *et al*, 2024; Nakamura *et al*, 2015), SV-anchored
synapsins (Cui *et al*, 2008; Song & Augustine, 2023) and microtubules along with their
associated proteins (Heisler *et al*, 2011; Jacob *et al*, 2008; Lorenz-Guertin & Jacob, 2018; Prag
et al, 2007; Velasco *et al*, 2023; Waites *et al*, 2021). These modifications influence the
localization and functional properties of these proteins, thereby modulating synaptic
strength. Additionally, synaptic strength can be regulated by (trans-synaptic) alignment of
functionally related proteins in nanodomains and nanocolumns, enhancing their cooperative
interactions (Biederer *et al*, 2017; Olah *et al*, 2023; Sanderson *et al*, 2020; Scimemi &
Diamond, 2012; Südhof, 2018, 2021; Tang *et al*, 2016). Of the few proteins that were
upregulated – or for the first time detected – in PB-AD neurons at 10 WPI (Table S1), we
found several cell adhesion molecules (CAMs) (CNTNAP4, SLITRK1, CDH1, EPHA7, ITGB3,
NPTX2, ITGA2, ELFN1, IGSF8 and CNTNAP1) (Beuter *et al*, 2016; Fiederling *et al*, 2011;
Karayannis *et al*, 2014; Li *et al*, 2020; Park & Goda, 2016; Ray & Treloar, 2012; Sia *et al*, 2007;
Stachniak *et al*, 2019; Um *et al*, 2014; Wiera *et al*, 2022), extra cellular matrix (ECM) proteins
(ITGB3, ITGA2, TNC, GPR158 and RELN) (Condomitti *et al*, 2018; Park & Goda, 2016;
Quattrocchi *et al*, 2002; Tucić *et al*, 2021; Wiera *et al*., 2022), and proteins involved in receptor
clustering (NPTX2, SLITRK1, PRRT1, ELFN1) (Martin *et al*, 2021; Matsunaga & Aruga, 2021;
O'Brien *et al*, 1999; Yim *et al*, 2013). Some of which are known to strengthen GABAergic
transmission (Beuter *et al*., 2016; Karayannis *et al*., 2014). Altogether, this suggests that
maturation and synaptic strengthening in iPSC-derived neuronal synapses depends on

presynaptic development driven by enhanced expression of multiple presynaptic gene sets
and alignment/reorganisation of pre- and postsynapse driven by synapse-organising genes.

A versatile in vitro model for studying the role of inhibition in brain disease
Dysfunction of inhibitory neurons and E/I disbalance are common aspect of
neurodevelopmental and neurodegenerative disorders (Gabitto *et al*, 2023; Goel *et al*, 2018;
Hijazi *et al*, 2020; Malwade *et al*, 2022; Mossner *et al*, 2020; Palop *et al*, 2007b; Zott *et al*,
2018). Given the purity of the GABAergic cultures produced with the PB-AD protocol, neurons
generated with this method provide a versatile model to study the role of inhibition and E/I
disbalance in these disorders using a human, and patient-own, genetic background.
Compared to a safe harbour approach, our method is directly applicable to patient derived
iPSCs, which makes it time and cost efficient, and prevents the introduction of clonal gene
editing effects. The reliable production of GABAergic autaptic cultures allows for a detailed
assessment of interneuron synaptic properties, such as priming, fusion and synaptic plasticity
in patient cell lines (Østergaard *et al*, 2025; Öttl *et al*, 2024; van Boven *et al*, 2024). Application
to disease-lines will be important steps to substantiate the full potential of this method. To
this end, PB-AD neurons can be mixed with NGN2 neurons at defined ratios, enabling
accurate modelling of E/I networks at any ratio, including those found in the human brain
(Bakken *et al.*, 2021; Fang *et al.*, 2022; Loomba *et al.*, 2022).
461

Acknowledgements
The authors thank Robbert Zalm and Ingrid Saarloos for their help with cloning and plasmid
production, Lisa Laan and Desiree Schut for preparation of glia feeder plates, Judith Huijgen
and Solange Lopes Cardozo for preparation and maintenance of iPSCs, and Berna Özer and
Remco Klaassen for performing mass-spectrometry proteomics.
This work was supported by the following grants (ZonMW-TOP BECAUSE (91216064),
to M.V and L.N.C. ZonMW-PSIDER Brainmodel (10250022110003) to M.V and L.N.C., NWO-
Dutch Research Agenda (NWA) NewTDec (1160.18.200) to M.V and L.N.C.), NWO Gravitation
program grant BRAINSCAPES (NWO 024.004.012) to M.V. and ZonMW-OPEN
(9120232310098) from the Netherlands Organisation for Scientific Research (NWO) to M.V.
and L.N.C.

Declaration of interest
The authors declare no competing interests.
476

Author contributions

M.E.W., and J.M.C.M. designed and produced the PiggyBac construct

T.W.v.V, M.A.v.B., K.I.M., R.F.T., A.B.S., R.E.v.K, M.V. and L.N.C. designed the experiments

T.W.v.V, M.A.v.B. and J.S. performed and maintained neuronal inductions and cultures

T.W.v.V M.A.v.B. performed the patch-clamp experiments with contributions of J.S. to Fig. 1

and C.F. to Fig. S2

T.W.v.V, M.A.v.B.,. performed the immunocytochemistry experiments with contributions of J.S

to Fig. 1

K.I.M., and F.K. performed the proteomics experiments and analysis

T.W.V.V, M.A.v.B., K.I.M., R.E.v.K, M.V. and L.N.C. designed figures and wrote the manuscript with

input from all authors.

Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
MAP2	Merck Millipore	(Millipore Cat# AB5543, RRID:AB_571049)
VGAT	Synaptic Systems	(Synaptic Systems Cat# 131 002, RRID:AB_887871)
VGLUT1	Antibodies Inc.	(Antibodies Incorporated Cat# 75-066, RRID:AB_2187693)
VGLUT2	Antibodies Inc.	(Antibodies Incorporated Cat# 75-067, RRID:AB_2239153)
NeuN	Merck Millipore	(Millipore Cat# ABN78, RRID:AB_10807945)
Parvalbumin	Synaptic Systems	(Synaptic Systems Cat# 195 004, RRID:AB_2156476)
Bacterial and virus strains		
FUW-TetO-*Ascl1*-T2A-puro	Addgene	RRID:Addgene_973 29

FUW-TetO-*Dlx2*-IRES-hygro	Addgene	RRID:Addgene_973 30
FUW-M2rtTA	Addgene	RRID:Addgene_203 42
Deposited data		
PRIDE dataset " One-step induction of human GABAergic neurons promotes presynaptic development & synapse maturation"	This study	PXD064163
Transcriptomic cytoarchitecture reveals principles of human neocortex organization	Jorstad, et al. (Jorstad *et al.*, 2023)	RRID:SCR_016152
Experimental models: Cell lines		
BiONI010-C-10	EBiSC	RRID:CVCL_RM84
BioNi010-C-13	EBiSC	RRID:CVCL_RF90
Kolf2.1J	Jackson Laboratory	RRID:CVCL_B5P3
GM23973	Coriell	RRID:CVCL_BW42
Recombinant DNA		
PB-TO-ASCL1-DLX2	This study	RRID:Addgene_182 307
Transposase	This study	N/A
Software and algorithms		
DIA-NN 1.8.1	Demichev, et al (2020,2022) (Demichev *et al*, 2020; Demichev *et al*, 2022)	
MS-DAP 1.0.5	Koopmans, et al. 2023 (Koopmans *et al*, 2023)	
DEqMS	Zhu, et al, 2020 (Zhu *et al*, 2020)	
GOAT	Miedema, et al. 2022 (Koopmans, 2024b)	
SynGO	Koopmans, et al. 2019 (Koopmans *et al*, 2019b)	

iPSCs origin and culture

Unless stated otherwise, the BiONi010-C-10 stem cell line (EBISC, RRID:CVCL_RM84) was
used for all experiments in this study. For glutamatergic induction, a gene-edited variant of
the same line containing an integrated NGN2-casette was used (EBISC, RRID:CVCL_RF90). As
a genetically independent control the KOLF2.1J (JAX, JiPSC1000, RRID:CVCL_B5P3) and
GM23973 (Coriell, RRID:CVCL_BW42) lines were used. All iPSCs were maintained on Matrigel
(Cornig) or Geltrex (Fisher Scientific)-coated dishes in Essential 8 (Gibco) or Essential 8 Flex
(Gibco) medium and passaged with EDTA and replated in the culture medium supplemented
with ROCK inhibitor Y27632 (5 μM, Tetu bio).

Lentiviral infection

Ultrahigh titre lentiviral particles were generated by Alstem (Richmond, CA, USA) from FUW-
M2rtTA (Addgene, #20342, RRID:Addgene_20342), FUW-TetO-*Ascl1*-T2A-puro (Addgene
#97329, RRID:Addgene_97329), and FUW-TetO-*Dlx2* -IRES-hygro (Addgene #97330,
RRID:Addgene_97330). For *Ascl1*+Forskolin, iPSCs were infected with only FUW-M2rtTA and
TetO-*Ascl1*-Puro, for *Ascl1* + *Dlx2* induction, iPSCs were infected with all three lentiviruses.
For both lentiviral induction methods, 24 hours after dox induction, puromycin (1 μg/ml)
selection was performed. For lentiviral *Ascl1* + *Dlx2* induction, additional hygromycin (75
μg/ml) selection was performed. For both selection regimes an uninfected control well was
included to confirm successful selection before proceeding towards induction.

PiggyBac transfection

To prepare for PiggyBac transfection, iPSCs were released with accutase and plated as single
cells on a Matrigel-coated 6-well plate in E8 + ROCK Inhibitor Y27632 at a density of 1500k
per well. Cells were placed in the incubator to adhere for 1 to 2 hours. A total of 3 μg of DNA
containing the *Ascl1* + *Dlx2* PiggyBac transposon (Presented here; available on Addgene,
182307, RRID:Addgene_182307) and transposase constructs (extended data 1) at a 1:3 molar
ratio was prepared in 100 μL of opti-MEM (GIBCO, 31985062). To another 100 μL op opti-
MEM 5 μL of Lipofectamine STEM (Invitrogen, STEM00001) was added. Opti-MEM
preparations were quickly mixed by pipetting, after which the mixture was incubated at RT
for 15 to 30 minutes during which cells received fresh E8 + ROCK Inhibitor Y27632 to wash
away non-adherent cells. The opti-MEM solution was then added to the cells in a dropwise
fashion. 24 hours post transfection, selection with puromycin (1 μg/ml) was initiated to select
only for cells containing the transposon, and selection was repeated every 3-4 days.

PB-AD GABAergic neuronal induction

PB-AD positive iPSCs were released with Accutase (Merck) on day post induction (DPI) 0 and
replated with at an approximate density of 3.500 cells/cm^2 in N2 medium (DMEM/F12
medium, Life technologies) containing Glutamax (200 mM, Life Technologies), dextrose (0.3%,
Life Technologies) N2 supplement B (StemCell Technologies), Pen/Strep (0.1%, Gibco), ROCK
inhibitor Y27632 (10 µM, Tetubio) and doxycycline hyclate (2 µg/mL) onto Matrigel (Cornig)
or Geltrex (Fisher Scientific) coated dishes. On DPI 1, medium was refreshed using the same
composition without ROCK inhibitor Y27632. On DPI 2, the medium was refreshed once more
with the same composition as on DPI 1, with the addition of 5-Fluoro-2'-deoxyuridine (FUDR)
(10 µM, Sigma-Aldrich) or Cytosine β-D-arabinofuranoside (AraC, 20 µM, Sigma-Aldrich) to
remove proliferative cells. On DPI 3, neurons were released with accutase, pelleted, and
resuspended and plated in Neurobasal medium (Life Technologies) supplemented with
Glutamax (200 mM), B27 (0.1%) Pen-Strep (0.1%), fetal bovine serum (0.5%), BDNF (10 ng/ml,
Peprotech), CNTF (10ng/ml, Peprotech), GDNF (10 ng/ml, Peprotech) and doxycycline hyclate
(2 µg/mL; Sigma-Aldrich). Neurons were plated to the required respective preparations (see
Autaptic culture and *Network culture*).

Glutamatergic neuronal induction
For glutamatergic induction, Bioni010-C-13 iPSCs were released with Accutase (Merck) on
DPI 0 and replated with at an approximate density of 3.500 cells/cm^2 in N2 medium
(DMEM/F12 medium, Life technologies) containing Glutamax (200 mM, Life Technologies),
dextrose (0.3%, Life Technologies) N2 supplement B (StemCell Technologies), Pen/Strep
(0.1%, Gibco) and doxycycline hyclate (2 µg/mL; Sigma-Aldrich) combined with dual-SMAD
inhibitors LDN193189 (100 nM, Stemgent), SB431542 (10 µM, Tocris Bioscience), and XAV939
(2 µM, Stemgent) NT3 (10 ng/ml, Peprotech) and doxycycline hyclate (2 µg/mL) onto Matrigel
(Cornig) or Geltrex (Fisher Scientific) coated dishes. On DPI 1, medium was refreshed using
the same composition without ROCK inhibitor Y27632. On DPI 2, the medium was refreshed
once more with the same composition as on day 2, with the addition of 5-Fluoro-2'-
deoxyuridine (FUDR) (10 µM, Sigma-Aldrich) or Cytosine β-D-arabinofuranoside (AraC, 20 µM,
Sigma-Aldrich) to remove proliferative cells. On DPI 3, neurons were released with accutase,
pelleted, resuspended and plated in Neurobasal medium (Life Technologies) supplemented
with Glutamax (200 mM), B27 (0.1%) Pen-Strep (0.1%), foetal bovine serum (0.5%), BDNF (10
ng/ml, Peprotech), CNTF (10ng/ml, Peprotech), GDNF (10 ng/ml, Peprotech) and doxycycline
hyclate (2 µg/mL; Sigma-Aldrich). Neurons were plated to the required respective
preparations (see *Autaptic culture* and *Network culture*).

Lentiviral GABAergic induction

In general, lentiviral GABAergic induction protocols followed a similar workflow as PB-AD
induction. To better adhere to established protocols (Mossink *et al.*, 2022; Yang *et al.*, 2017),
the medium composition differed at certain steps (Figure 1H). For Ascl1 + Forskolin induction,
from DPI 0 onwards, Forskolin (10 μM, Merck) was added to the medium up to DPI 14.
Furthermore, from DPI0 onwards non-essential amino acids (1:100, Sigma) were added up to
day DPI 14. BDNF (10 ng/ml) and NT3 (10 ng/ml) were present from DPI1 onwards. For
lentiviral *Ascl1* + *Dlx2* induction BDNF (10 ng/ml) and NT3 (10 ng/ml) were added from DPI
0 onward, as well. Furthermore, laminin (1μg/ml, Sigma-Aldrich), cyclic AMP (1 μM, Sigma-
Aldrich), and Ascorbic Acid (200 μM, Sigma-Aldrich), were added to the medium at all
subsequent refreshments to improve viability.

Autaptic culture
Neurons were plated at a density of ~ 400 cells / cm^2 on micro-islands prepared from rat
glial cells plated on agarose-coated coverslips stamped with a custom stamp in Poly-D-
Lysine (0.1 mg/ml, Sigma), rat tail collagen (0.7 mg/ml, Corning) and Fibronectin (100 μg/ml,
Corning) in acetic acid (10 mM, sigma). Glial islands were prepared approximately 7 days
before plating neurons and were treated with AraC (20 μM, Sigma-Aldrich) before plating
neurons. Autaptic cultures were refreshed every 7 days with a ~50% media refreshment, with
the same medium as used for plating neurons, but omitting doxycycline hyclate (except for
Figure 1E-G, where doxycycline hyclate was added up to DPI 14).

Network culture
Neurons were plated at a density of ~ 7000 cells / cm^2 on coverslips covered with a glial-
monolayer prepared from rat astrocytes a week in advance and treated with Cytosine β-D-
arabinofuranoside (20 μM, Sigma-Aldrich) 24-48 hours before plating neurons. Networks
cultures were refreshed with a ~50% media refreshment every 3-4 days, with the same
medium as used for plating neurons, but omitting doxycycline hyclate.

Electrophysiology
Whole-cell voltage-clamp recordings ($V_m = -70$ mV) were performed on autaptic neurons,
with borosilicate glass pipettes (2–5MΩ) filled with (in mM) 136 KCl, 17.8 HEPES, 1 EGTA,
0.6 MgCl$_2$, 4 ATP-Mg, 0.3 GTP-Na, 12 phosphocreatine-K$_2$ and 50 units/mL phosphocreatine
kinase (pH = 7.30, 290 mOsmol). The external solution contained in mM: 10 HEPES, 10
glucose, 140 NaCl, 2.4 KCl, 4 MgCl$_2$, and 4 CaCl$_2$ (pH = 7.30, 290 mOsmol). Experiments were
performed at room temperature (~22℃ using a MultiClamp700B amplifier (Axon
Instruments) and Digidata 1440A digitizer under control of Clampex 10 software (Molecular

devices). Only cells with an access resistance < 15MΩ (80% compensated) and leak current
of <300 pA were included. Synaptic responses were elicited by 0.5 ms depolarization to
30 mV.

For GABA puffs, perforated patch recordings were produced by filling the patch pipette
(in mM) 140 KCl, 10 HEPES, gramicidin (100 µg/ml) and allowing cells to perforate for 15-30
minutes after reaching a GΩ seal to < 100 MΩ series resistance. Puff pipettes filled with 100
µM GABA in ACSF were positioned near the soma, at approximately 25 µm distance. 100 ms
puffs were applied at -110, -90, -70, -50, -30 and -10 mV holding voltages to induce GABAergic
currents.

Pharmacology
To Block GABA-currents, picrotoxin (100 µM, Hello Bio) was used. A 100 mM stock was
prepared in DMSO and was added to the bath to reach the final concentration. To block
glutamatergic currents, AP-V (100 µM, R&D systems) and DNQX (40 µM, Tocris) were used.
For DNQX, A 1000x of stock of each was prepared in DMSO. For AP-V, the stock was prepared
in water.

Analysis electrophysiological data.
Offline analysis was performed using custom-written programs (Huson & Cornelisse, 2019)
in Matlab (Mathworks). Experimental conditions were blinded during analysis. In all figures,
stimulation artefacts have been removed. For evoked release during 20 Hz stimulation, back-
extrapolation of linear fits from the last 20 pulses of the cumulative charge was used to
estimate the RRP (y-intercept) (Neher, 2015).

Immunocytochemistry and morphological analysis
Neurons were fixed in ice cold 100% methanol (VWR) for 15 minutes at 4°C, blocked with
2.5% normal goat serum (NGS, Life Technologies) and 0.1% Triton X-100 in PBS for
45 minutes and incubated with primary antibodies in PBS with 2.5% NGS and 0.1% Triton X-
100 overnight at 4°C. The following primary antibodies were used: chicken anti-MAP2
(1:1000, Merck AB5543, RRID:AB_571049); rabbit anti-VGAT (1:2000, SySy #131 002,
RRID:AB_887871), mouse anti-VGlut1 (1:250, NeuroMab clone N28/9, Antibodies inc. 75-066,
RRID:AB_2187693), mouse anti-VGlut2 (1:250, NeuroMab clone N29/29, Antibodies inc. 75-
067, RRID:AB_2239153), Rabbit anti-NeuN (1:500, Merck ABN78, RRID:AB_10807945) and
anti-Parvalbumin (1:500, SySy #195 004, RRID:AB_2156476). After washing with PBS the next
day, cells were incubated for 1 h at room temperature with secondary antibodies
conjugated to Alexa dyes (1:1000, Invitrogen Molecular Probes), washed again and mounted

with DABCO-Mowiol (Invitrogen). Images were acquired at 1024x1024 pixels on a Nikon
Eclipse Ti confocal laser scanning microscope (40× objective; NA 1.3) with NIS-Elements
software. Z-stacks were produced at a 0.5 μm step-size until no more puncta and/or dendrites
were in focus. Confocal settings were kept constant for all scans within an experimental
week. Analysis of neuronal morphology was performed on maximum-intensity projections
using the automated image analysis routine SynD (Schmitz *et al*, 2011) for autaptic neurons
and a custom ImageJ workflows for network analysis, in either case, for VGAT and VGLUT
puncta counts, only puncta on dendrites were counted.

Statistics of electrophysiological and morphological data
Statistical testing was performed in MATLAB and GraphPad Prism For descriptive statistics
(Figure 1) the likelihood of normality vs lognormality was first calculated; for normally
distributed data, the arithmetic mean and standard deviation were calculated. For
lognormally distributed data, the geometric mean and geometric SD factor were calculated,
instead. In autaptic development timelines (Figure 4&5), to test general developmental
trajectories, nonparametric Spearman correlations were computed with a two-tailed test for
significance and a 95% confidence interval. For timepoint comparisons, Kruskal-Wallis
ANOVAs were performed with Dunn's multiple comparisons test. Pairwise comparisons were
made between 4 and 6, 6 and 8, and 8 and 10 WPI. #, ##, ###, #### and *ns* represent results
of spearman correlation results, *, **, ***, **** and ns represent Kruskal-Wallis ANOVA results.
ns and ns represent non-significant results. Single, double, triple and quadruple marks
represent $p < 0.05$, $p < 0.01$, $p < 0.001$ and $p < 0.0001$, respectively. GABA-puff responses (figure
6O) were fitted using a first-order polynomial linear fit and an extra sum-of-squares F-test
was used to determine whether GABA responses were identical throughout development. In
plots, unless stated otherwise, the lines in boxplots represent the median value. Crosses
represent the mean value. Boxes represent 50% off the values. Circular markers represent
individual observations.

Cell pellet harvesting for MS
Cells were washed three times with 1 ml sterile Dulbecco's PBS (Gibco), whereof first twice
at 37°C and finally once at 4 °C. Cultures were kept on ice, scraped off using VWR cell lifters,
and harvested in 1.5 ml protein LoBind Eppendorf tubes (VWR). Pellets were obtained by
centrifugation at 1,000 × g for 5 minutes at 4 °C. Supernatant was removed and pellets were
kept at -80 °C before subsequent sample preparation.

MS sample preparation

MS sample preparation was performed following the DNA micro spin column suspension
trapping protocol (Thanou *et al*, 2023), with some minor adaptations. Cell pellets were
extracted and reduced in 6% SDS buffer, containing 50 mM Tris-HCl (pH 8) and 5 mM tris (2-
carboxyethyl)phosphine (TCEP), by incubation at 55°C for 15 minutes in a thermomixer set
to 1400 RPM. Free sulfhydryl groups were alkylated by incubation with 20 mM methyl
methanethiosulfonate (MMTS; 200 mM stock solution) for 15 minutes at RT. Insoluble debris
was cleared by centrifugation at 10000g for 1 minute at RT and the supernatant was
transferred to a new tube (Eppendorf). Next, samples were acidified by addition of 1.1%
phosphoric acid (12% stock solution), mixed with six volumes of binding/washing buffer
(90% methanol and 100mM Tris-HCl, pH 8) and loaded onto a plasmid DNA micro column
(HiPure from Magen Biotechnology). The protein particulate was retained on the column
upon centrifugation at 1400x g for 1 minute and the columns were washed four times with
binding/washing buffer. Columns were transferred to a new tube (Eppendorf), supplemented
with 500 ng Trypsin/Lys-C (Promega) in 50 mM NH4HCO3 and incubated overnight at 37°C
in a humidified incubator. Tryptic peptides were eluted and pooled by subsequent addition
of 50 mM NH4HCO3, 0.1% formic acid and 0.1% formic acid in acetonitrile. Collected peptides
were dried by SpeedVac and stored at -80°C.

LC-MS Analysis
Each sample of tryptic digest was redissolved in 0.1% formic acid and the peptide
concentration was determined by tryptophan-fluorescence assay (Wiśniewski & Gaugaz,
2015); 75 ng of peptide was loaded onto an Evotip Pure (Evosep). Peptide samples were
separated by standardized 30 samples per day method on the Evosep One liquid
chromatography system, using a 15 cm × 150 μm reverse-phase column packed with 1.5 μm
C18-beads (EV1137 from Evosep) connected to a 20 μm ID ZDV emitter (Bruker Daltonics).
Peptides were electro-sprayed into the timsTOF Pro 2 mass spectrometer (Bruker Daltonics)
equipped with CaptiveSpray source and measured with the following settings: Scan range
100-1700 m/z, ion mobility 0.65 to 1.5 Vs/cm2, ramp time 100 ms, accumulation time 100
ms, and collision energy decreasing linearly with inverse ion mobility from 59 eV at 1.6
Vs/cm2 to 20 eV at 0.6 Vs/cm2. Operating in dia-PASEF mode, each cycle took 1.38 s and
consisted of 1 MS1 full scan and 12 dia-PASEF scans. Each dia-PASEF scan contained two
isolation windows, in total covering 300-1200 m/z and ion mobility 0.65 to 1.50 Vs/cm2. Dia-
PASEF window placement was optimized using the py-diAID tool (Skowronek *et al*, 2022). Ion
mobility was auto calibrated at the start of each sample (calibrant m/z, 1/K0: 622.029, 0.992
Vs/cm2; 922.010, 1.199 Vs/cm2; 1221.991, 1.393 Vs/cm2).

MS data Analysis

DIA-PASEF raw data were processed with DIA-NN 1.8.1 (Demichev *et al.*, 2020; Demichev *et al.*, 2022). An in-silico spectral library was generated from the uniprot human reference proteome (SwissProt and TrEMBL, canonical and additional isoforms, release 2023-2) using Trypsin/P digestion and at most 1 missed cleavage. Fixed modification was set to beta-methylthiolation (C) and variable modifications were oxidation (M) and N-term M excision (at

most 1 per peptide). Peptide length was set to 7-30, precursor charge range was set to 2-4, precursor m/z was limited to 280–1220, both MS1 and MS2 mass accuracy were set to 15 ppm, scan window was set to 9, double-pass-mode and match-between-runs were enabled. Protein identifiers (isoforms) were used for protein inference. All other settings were left as default.

MS-DAP 1.0.5 (Koopmans *et al.*, 2023) was used for downstream differential abundance analyses using the DIA-NN results. Filtering and normalization was applied to respective samples per statistical contrast. Peptide-level filtering was configured to retain only peptides identified in at least 4 samples in both experimental conditions (within current contrast) and only proteins with 2 valid peptides were retained. Next, peptide abundance values were normalized by minimizing the mode $\log_2$ (FC) between sample pairs (MWMB algorithm), followed by protein-level mode-between normalization. The DEqMS (Zhu *et al.*, 2020) algorithm was used for statistical analysis and resulting p-values were adjusted for multiple testing using the Benjamini–Hochberg False Discovery Rate (FDR) procedure. To investigate proteins with strong differences in detection rates between conditions, e.g. consistently detected in one condition but (near) absent in another, the MS-DAP differential detection test was applied to all proteins that were observed with at least 2 peptides in at least 4 samples in either condition (i.e. confidently detected proteins). Using the default MS-DAP workflow, we merged differential detection (DD) results into the DEqMS result table for the subset of proteins that had both an absolute DD z-score of at least 6 and a smaller adjusted p-value in DD (or no p-value from DEqMS due to a lack of data points in either condition).

Expression-weighted cell type enrichment analysis

Cell type enrichment analysis gives insight into the cell types that potentially drive differential expression. EWCE uses single-cell RNA-sequencing data to generate a probability distribution associated with a gene list having an average level of expression within a cell (Skene & Grant, 2016). Although it's based on RNA expression levels, it can also be applied

to proteomics data (Miedema *et al*, 2022). We used published single nucleus RNA-sequencing
data (49,495 nuclei) from human cortical brain regions – including middle temporal gyrus,
anterior cingulate cortex, primary visual cortex, primary motor cortex, primary somatosensory
cortex and primary auditory cortex – published by Jorstad et al (Jorstad *et al.*, 2023). For cell
type annotations we used the cell type annotations: class (e.g., GABAergic neurons) and
subclass (e.g. SST[+] interneurons) for first and second level EWCE analysis, respectively.
Bootstrap enrichment tests were performed using 10,000 reps to look whether gene sets
(i.e., foreground) were significantly enriched for a cell class or subclass relative to the
background (i.e., all detected proteins included in that contrast). Since we focused on
neuronally-enriched proteins or SYNGO-annotated synaptic proteins, we excluded non-
neuronal enrichment scores.

EWCE calculates cell type-specificity scores (CTSS) for all genes, which is the
percentage of mean expression per cell type relative to the sum of the mean expressions of
all cell types. We used the CTSS to predict for each protein the cell type the protein probably
is derived from. We only included CTSS from genes that passed mRNA detection threshold
(mean expression >4 in at least one cell class). Cell type predictions included neuronal
(neuronal CTSS > 70%), GABAergic neurons (GABA CTSS > 0.8 and glutamatergic CTSS < 0.1),
glutamatergic neurons (glutamatergic CTSS > 0.8 and GABAergic CTSS < 0.1). Cell type
predictions were labelled in volcano plots. We also used the EWCE-calculated mean mRNA
expression for each cell type to estimate the differential expression of genes in GABAergic
vs glutamatergic neurons, which was used in the SYNGO manual colour plots (Figure 6H)).

Heatmaps and line plots
We used the mean $\log_2$-transformed normalized protein intensities per group (i.e., PBAD 4
WPI) and plotted these values using line plots in RStudio (Team, 2020) using ggplot
(Wickham, 2016). The relative normalized protein intensities ($\log_2$ (FC)) was used in
heatmaps, which were produced using Morpheus. To map these proteins to predicted cell
types we also show EWCE-calculated cell type specificity scores. All heatmaps visualizations
are with fixed colour scaling (i.e., not relative colour scaling per gene), as indicated by
heatmap legend.

Gene-set enrichment analysis using GOAT and over-representation analysis using SYNGO
We used the output of the differential abundance analysis using MSDAP (PB-AD 10 vs 4 WPI
and NGN2 10 vs 4 WPI) as input for gene-set enrichment analysis using GOAT. The GOAT
online v1.0.2 (Koopmans, 2024b) was used to perform gene set enrichment analyses using
gene sets from the SynGO knowledgebase v1.2 (Koopmans *et al.*, 2019b). Effect size-derived

gene scores (i.e., log$_2$ (FC)) were used to test for enriched gene sets that contained at least
10 and at most 1500 genes (or 50% of the gene list, whichever was smaller) that overlapped
with the input gene list. Multiple testing correction was independently applied per gene set
"source" (i.e. SYNGO_CC, SYNGO_BP) using Bonferroni adjustment and all p-values were again
adjusted using Bonferroni correction to account for 2 separate tests across "sources". The
significance threshold for adjusted p-values was set to 0.05. SYNGO over-representation
analysis was performed using the same MSDAP output. Foreground was defined as
upregulated proteins per contrast (p-adjusted < 0.05 and log$_2$ (FC) > 0.1), and the background
contained all tested genes for that contrast. SynGO release is dataset version: 20231201, and
default settings were used. Ontologies are significantly enriched at 1% FDR, while including
terms containing at least three matching input genes.

Material availability
PiggyBac *Ascl1-Dlx2* construct is available from Addgene #182307. The mass spectrometry
proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE
partner repository (Perez-Riverol *et al*, 2025) with the dataset identifier PXD064163.
775

References
Allison T, Langerman J, Sabri S, Otero-Garcia M, Lund A, Huang J, Wei X, Samarasinghe RA,
Polioudakis D, Mody I *et al* (2021) Defining the nature of human pluripotent stem cell-derived
interneurons via single-cell analysis. *Stem Cell Reports* 16: 2548-2564

Andreae LC, Burrone J (2015) Spontaneous Neurotransmitter Release Shapes Dendritic Arbors
via Long-Range Activation of NMDA Receptors. *Cell Rep* 10: 873-882

Bakken TE, Jorstad NL, Hu Q, Lake BB, Tian W, Kalmbach BE, Crow M, Hodge RD, Krienen FM,
Sorensen SA *et al* (2021) Comparative cellular analysis of motor cortex in human, marmoset
and mouse. *Nature* 598: 111-119

Barretto N, Zhang H, Powell SK, Fernando MB, Zhang S, Flaherty EK, Ho SM, Slesinger PA,
Duan J, Brennand KJ (2020) ASCL1- and DLX2-induced GABAergic neurons from hiPSC-derived
NPCs. *J Neurosci Methods* 334: 108548

Baum C (2007) Insertional mutagenesis in gene therapy and stem cell biology. *Current opinion*
in hematology 14: 337-342

Beuter S, Ardi Z, Horovitz O, Wuchter J, Keller S, Saha R, Tripathi K, Anunu R, Kehat O, Kriebel
M *et al* (2016) Receptor tyrosine kinase EphA7 is required for interneuron connectivity at
specific subcellular compartments of granule cells. *Scientific Reports* 6: 29710

Biederer T, Kaeser PS, Blanpied TA (2017) Transcellular Nanoalignment of Synaptic Function.
Neuron 96: 680-696

Breschi A, Gingeras TR, Guigó R (2017) Comparative transcriptomics in human and mouse.
Nature Reviews Genetics 18: 425-440

Bright DP, Smart TG (2013) Protein kinase C regulates tonic GABAA receptor-mediated
inhibition in the hippocampus and thalamus. *European Journal of Neuroscience* 38: 3408-3423

Bullmann T, Kaas T, Ritzau-Jost A, Wöhner A, Kirmann T, Rizalar FS, Holzer M, Nerlich J, Puchkov
D, Geis C *et al* (2024) Human iPSC-Derived Neurons with Reliable Synapses and Large
Presynaptic Action Potentials. *J Neurosci* 44

Busche MA, Eichhoff G, Adelsberger H, Abramowski D, Wiederhold K-H, Haass C, Staufenbiel
M, Konnerth A, Garaschuk O (2008) Clusters of Hyperactive Neurons Near Amyloid Plaques
in a Mouse Model of Alzheimer's Disease. *Science* 321: 1686-1689

Cabrera A, Edelstein HI, Glykofrydis F, Love KS, Palacios S, Tycko J, Zhang M, Lensch S, Shields
CE, Livingston M *et al* (2022) The sound of silence: Transgene silencing in mammalian cell
engineering. *Cell Systems* 13: 950-973

Chanda S, Ang Cheen E, Davila J, Pak C, Mall M, Lee Qian Y, Ahlenius H, Jung Seung W, Südhof
Thomas C, Wernig M (2014) Generation of Induced Neuronal Cells by the Single
Reprogramming Factor ASCL1. *Stem Cell Reports* 3: 282-296

Chang C-L, Trimbuch T, Chao H-T, Jordan J-C, Herman MA, Rosenmund C (2014) Investigation
of Synapse Formation and Function in a Glutamatergic-GABAergic Two-Neuron Microcircuit.
The Journal of Neuroscience 34: 855-868

Chung JH, Bell AC, Felsenfeld G (1997) Characterization of the chicken beta-globin insulator.
Proc Natl Acad Sci U S A 94: 575-580

Chung WS, Allen NJ, Eroglu C (2015) Astrocytes Control Synapse Formation, Function, and
Elimination. *Cold Spring Harb Perspect Biol* 7: a020370

Condomitti G, Wierda KD, Schroeder A, Rubio SE, Vennekens KM, Orlandi C, Martemyanov KA,
Gounko NV, Savas JN, de Wit J (2018) An Input-Specific Orphan Receptor GPR158-HSPG
Interaction Organizes Hippocampal Mossy Fiber-CA3 Synapses. *Neuron* 100: 201-215.e209

Cui Y, Costa RM, Murphy GG, Elgersma Y, Zhu Y, Gutmann DH, Parada LF, Mody I, Silva AJ (2008)
Neurofibromin regulation of ERK signaling modulates GABA release and learning. *Cell* 135:
549-560

Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M (2020) DIA-NN: neural networks and
interference correction enable deep proteome coverage in high throughput. *Nat Methods* 17:
41-44

Demichev V, Szyrwiel L, Yu F, Teo GC, Rosenberger G, Niewienda A, Ludwig D, Decker J, Kaspar-
Schoenefeld S, Lilley KS *et al* (2022) dia-PASEF data analysis using FragPipe and DIA-NN for
deep proteomics of low sample amounts. *Nat Commun* 13: 3944

Dixon C, Sah P, Lynch JW, Keramidas A (2014) GABA$_A$ Receptor α and
β Subunits Shape Synaptic Currents via Different Mechanisms *. *Journal of Biological*
Chemistry 289: 5399-5411

Elmariah SB, Hughes EG, Oh EJ, Balice-Gordon RJ (2004) Neurotrophin signaling among
neurons and glia during formation of tripartite synapses. *Neuron Glia Biol* 1: 1-11

Engle SJ, Blaha L, Kleiman RJ (2018) Best Practices for Translational Disease Modeling Using
Human iPSC-Derived Neurons. *Neuron* 100: 783-797

Fang R, Xia C, Close JL, Zhang M, He J, Huang Z, Halpern AR, Long B, Miller JA, Lein ES *et al*
(2022) Conservation and divergence of cortical cell organization in human and mouse
revealed by MERFISH. *Science* 377: 56-62

Farzana F, Zalm R, Chen N, Li KW, Grant SG, Smit AB, Toonen RF, Verhage M (2016)
Neurobeachin Regulates Glutamate- and GABA-Receptor Targeting to Synapses via Distinct
Pathways. *Mol Neurobiol* 53: 2112-2123

Fenske P, Grauel MK, Brockmann MM, Dorrn AL, Trimbuch T, Rosenmund C (2019) Autaptic
cultures of human induced neurons as a versatile platform for studying synaptic function
and neuronal morphology. *Scientific Reports* 9: 4890

Fiederling A, Ewert R, Andreyeva A, Jüngling K, Gottmann K (2011) E-cadherin is required at
GABAergic synapses in cultured cortical neurons. *Neuroscience Letters* 501: 167-172

Flores CE, Nikonenko I, Mendez P, Fritschy JM, Tyagarajan SK, Muller D (2015) Activity-
dependent inhibitory synapse remodeling through gephyrin phosphorylation. *Proc Natl Acad*
Sci U S A 112: E65-72

Foss-Feig JH, Adkinson BD, Ji JL, Yang G, Srihari VH, McPartland JC, Krystal JH, Murray JD,
Anticevic A (2017) Searching for Cross-Diagnostic Convergence: Neural Mechanisms
Governing Excitation and Inhibition Balance in Schizophrenia and Autism Spectrum
Disorders. *Biol Psychiatry* 81: 848-861

Friedman HV, Bresler T, Garner CC, Ziv NE (2000) Assembly of New Individual Excitatory
Synapses: Time Course and Temporal Order of Synaptic Molecule Recruitment. *Neuron* 27:
57-69

Gabitto MI, Travaglini KJ, Rachleff VM, Kaplan ES, Long B, Ariza J, Ding Y, Mahoney JT, Dee N,
Goldy J *et al* (2023) Integrated multimodal cell atlas of Alzheimer's disease. *Res Sq*

Gharib WH, Robinson-Rechavi M (2011) When orthologs diverge between human and mouse.
Briefings in Bioinformatics 12: 436-441

Ghosh H, Auguadri L, Battaglia S, Simone Thirouin Z, Zemoura K, Messner S, Acuña MA,
Wildner H, Yévenes GE, Dieter A *et al* (2016) Several posttranslational modifications act in
concert to regulate gephyrin scaffolding and GABAergic transmission. *Nature Communications*
7: 13365

Gierman HJ, Indemans MH, Koster J, Goetze S, Seppen J, Geerts D, van Driel R, Versteeg R
(2007) Domain-wide regulation of gene expression in the human genome. *Genome Res* 17:
1286-1295

Gödecke N, Zha L, Spencer S, Behme S, Riemer P, Rehli M, Hauser H, Wirth D (2017) Controlled
re-activation of epigenetically silenced Tet promoter-driven transgene expression by
targeted demethylation. *Nucleic Acids Research* 45: e147-e147

Goel A, Cantu DA, Guilfoyle J, Chaudhari GR, Newadkar A, Todisco B, de Alba D, Kourdougli N,
Schmitt LM, Pedapati E *et al* (2018) Impaired perceptual learning in a mouse model of Fragile
X syndrome is mediated by parvalbumin neuron dysfunction and is reversible. *Nat Neurosci*
21: 1404-1411

Gonzalez-Ramos A, Waloschková E, Mikroulis A, Kokaia Z, Bengzon J, Ledri M, Andersson M,
Kokaia M (2021) Human stem cell-derived GABAergic neurons functionally integrate into
human neuronal networks. *Sci Rep* 11: 22050

Hao Y, Liu H, Zeng X-T, Wang Y, Zeng W-X, Qian K-Y, Li L, Chi M-X, Gao S, Hu Z *et al* (2023) UNC-
43/CaMKII-triggered anterograde signals recruit GABAARs to mediate inhibitory synaptic
transmission and plasticity at C. elegans NMJs. *Nature Communications* 14: 1436

Harms KJ, Tovar KR, Craig AM (2005) Synapse-specific regulation of AMPA receptor subunit
composition by activity. *J Neurosci* 25: 6379-6388

Hausrat TJ, Muhia M, Gerrow K, Thomas P, Hirdes W, Tsukita S, Heisler FF, Herich L, Dubroqua
S, Breiden P *et al* (2015) Radixin regulates synaptic GABAA receptor density and is essential
for reversal learning and short-term memory. *Nat Commun* 6: 6872

He J, Yang Q, Chang L-J (2005) Dynamic DNA Methylation and Histone Modifications
Contribute to Lentiviral Transgene Silencing in Murine Embryonic Carcinoma Cells. *Journal of*
Virology 79: 13497-13508

Heisler FF, Loebrich S, Pechmann Y, Maier N, Zivkovic AR, Tokito M, Hausrat TJ, Schweizer M,
Bähring R, Holzbaur EL *et al* (2011) Muskelin regulates actin filament- and microtubule-based
GABA (A) receptor transport in neurons. *Neuron* 70: 66-81

Hijazi S, Heistek TS, van der Loo R, Mansvelder HD, Smit AB, van Kesteren RE (2020)
Hyperexcitable Parvalbumin Interneurons Render Hippocampal Circuitry Vulnerable to
Amyloid Beta. *iScience* 23: 101271

Hoffman GE, Schrode N, Flaherty E, Brennand KJ (2019) New considerations for hiPSC-based
models of neuropsychiatric disorders. *Molecular Psychiatry* 24: 49-66

Houston CM, He Q, Smart TG (2009) CaMKII phosphorylation of the GABA (A) receptor:
receptor subtype- and synapse-specific modulation. *J Physiol* 587: 2115-2125

Huson V, Cornelisse LN, 2019. viewEPSC, 85b8899 ed.

Ikeda K, Yanagawa Y, Bekkers JM (2008) Distinctive Quantal Properties of Neurotransmission
at Excitatory and Inhibitory Autapses Revealed Using Variance−Mean Analysis. *The Journal of*
Neuroscience 28: 13563-13573

Jacob TC, Moss SJ, Jurd R (2008) GABA (A) receptor trafficking and its role in the dynamic
modulation of neuronal inhibition. *Nat Rev Neurosci* 9: 331-343

Jiang L, Sun S, Nedergaard M, Kang J (2000) Paired-pulse modulation at individual GABAergic
synapses in rat hippocampus. *J Physiol* 523 Pt 2: 425-439

Jones MV, Westbrook GL (1995) Desensitized states prolong GABAA channel responses to
brief agonist pulses. *Neuron* 15: 181-191

Jorstad NL, Close J, Johansen N, Yanny AM, Barkan ER, Travaglini KJ, Bertagnolli D, Campos J,
Casper T, Crichton K *et al* (2023) Transcriptomic cytoarchitecture reveals principles of human
neocortex organization. *Science* 382

Js, J. RH, Shaib AH, Rehbach K, Lee C, Seif P, Thomas C, Gideons E, Guenther A, Krutenko T *et al*
(2019) An Autaptic Culture System for Standardized Analyses of iPSC-Derived Human
Neurons. *Cell reports* 27

Karayannis T, Au E, Patel JC, Kruglikov I, Markx S, Delorme R, Héron D, Salomon D, Glessner J,
Restituito S *et al* (2014) Cntnap4 differentially contributes to GABAergic and dopaminergic
synaptic transmission. *Nature* 511: 236-240

Koopmans F (2024a) GOAT: efficient and robust identification of gene set enrichment.
Communications Biology 7: 744

Koopmans F (2024b) GOAT: efficient and robust identification of gene set enrichment.
Commun Biol 7: 744

Koopmans F, Li KW, Klaassen RV, Smit AB (2023) MS-DAP Platform for Downstream Data
Analysis of Label-Free Proteomics Uncovers Optimal Workflows in Benchmark Data Sets and
Increased Sensitivity in Analysis of Alzheimer's Biomarker Data. *J Proteome Res* 22: 374-386

Koopmans F, Van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, Cornelisse LN,
Farrell RJ, Goldschmidt HL, Howrigan DP *et al* (2019a) SynGO: An Evidence-Based, Expert-
Curated Knowledge Base for the Synapse. *Neuron* 103: 217-234.e214

Koopmans F, van Nierop P, Andres-Alonso M, Byrnes A, Cijsouw T, Coba MP, Cornelisse LN,
Farrell RJ, Goldschmidt HL, Howrigan DP *et al* (2019b) SynGO: An Evidence-Based, Expert-
Curated Knowledge Base for the Synapse. *Neuron* 103: 217-234.e214

Kwak CS, Oflaz FE, Qiu J, Wang X (2024) Human stem cell-specific epigenetic signatures
control transgene expression. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory*
Mechanisms 1867: 195063

Kwon HB, Sabatini BL (2011) Glutamate induces de novo growth of functional spines in
developing cortex. *Nature* 474: 100-104

Lambert NA, Wilson WA (1994) Temporally distinct mechanisms of use-dependent depression
at inhibitory synapses in the rat hippocampus in vitro. *Journal of neurophysiology* 72: 121-130

Li MA, Pettitt SJ, Eckert S, Ning Z, Rice S, Cadiñanos J, Yusa K, Conte N, Bradley A (2013) The
piggyBac Transposon Displays Local and Distant Reintegration Preferences and Can Cause
Mutations at Noncanonical Integration Sites. *Molecular and Cellular Biology* 33: 1317-1330

Li W, Yang L, Tang C, Liu K, Lu Y, Wang H, Yan K, Qiu Z, Zhou W (2020) Mutations of CNTNAP1
led to defects in neuronal development. *JCI Insight* 5

Liew C-G, Draper JS, Walsh J, Moore H, Andrews PW (2007) Transient and Stable Transgene
Expression in Human Embryonic Stem Cells. *Stem Cells* 25: 1521-1528

Lin H-C, He Z, Ebert S, Schörnig M, Santel M, Nikolova MT, Weigert A, Hevers W, Kasri NN,
Taverna E *et al* (2021) NGN2 induces diverse neuron types from human pluripotency. *Stem*
cell reports 16: 2118-2127

Lindhout FW, Kooistra R, Portegies S, Herstel LJ, Stucchi R, Snoek BL, Altelaar AFM, MacGillavry
HD, Wierenga CJ, Hoogenraad CC (2020) Quantitative mapping of transcriptome and
proteome dynamics during polarization of human iPSC-derived neurons. *eLife* 9: e58124

Liu J, Feng X, Wang Y, Xia X, Zheng JC (2022) Astrocytes: GABAceptive and GABAergic Cells in
the Brain. *Front Cell Neurosci* 16: 892497

Liu Y, Liu H, Sauvey C, Yao L, Zarnowska ED, Zhang S-C (2013) Directed differentiation of
forebrain GABA interneurons from human pluripotent stem cells. *Nature protocols* 8: 1670-
1679

Lombardo A, Cesana D, Genovese P, Di Stefano B, Provasi E, Colombo DF, Neri M, Magnani Z,
Cantore A, Lo Riso P *et al* (2011) Site-specific integration and tailoring of cassette design for
sustainable gene transfer. *Nature Methods* 8: 861-869

Loomba S, Straehle J, Gangadharan V, Heike N, Khalifa A, Motta A, Ju N, Sievers M, Gempt J,
Meyer HS *et al* (2022) Connectomic comparison of mouse and human cortex. *Science* 377

Lorenz-Guertin JM, Jacob TC (2018) GABA type a receptor trafficking and the architecture of
synaptic inhibition. *Developmental Neurobiology* 78: 238-270

Lützenkirchen FP, Zhu Y, Maric HM, Boeck DS, Gromova KV, Kneussel M (2024) Neurobeachin
regulates receptor downscaling at GABAergic inhibitory synapses in a protein kinase A-
dependent manner. *Commun Biol* 7: 1635

Malwade S, Gasthaus J, Bellardita C, Andelic M, Moric B, Korshunova I, Kiehn O, Vasistha NA,
Khodosevich K (2022) Identification of Vulnerable Interneuron Subtypes in 15q13.3
Microdeletion Syndrome Using Single-Cell Transcriptomics. *Biol Psychiatry* 91: 727-739

Mammen AL, Huganir RL, O'Brien RJ (1997) Redistribution and Stabilization of Cell Surface
Glutamate Receptors during Synapse Formation. *The Journal of Neuroscience* 17: 7351-7358

Marcó de la Cruz B, Campos J, Molinaro A, Xie X, Jin G, Wei Z, Acuna C, Sterky FH (2024) Liprin-
α proteins are master regulators of human presynapse assembly. *Nature Neuroscience* 27:
629-642

Maroof Asif M, Keros S, Tyson Jennifer A, Ying S-W, Ganat Yosif M, Merkle Florian T, Liu B,
Goulburn A, Stanley Edouard G, Elefanty Andrew G *et al* (2013) Directed Differentiation and
Functional Maturation of Cortical Interneurons from Human Embryonic Stem Cells. *Cell Stem*
Cell 12: 559-572

Martin EE, Wleklinski E, Hoang HTM, Ahmad M (2021) Interaction and Subcellular Association
of PRRT1/SynDIG4 With AMPA Receptors. *Frontiers in Synaptic Neuroscience* Volume 13 - 2021

Matsunaga H, Aruga J (2021) Trans-Synaptic Regulation of Metabotropic Glutamate Receptors
by Elfn Proteins in Health and Disease. *Frontiers in Neural Circuits* Volume 15 - 2021

Meijer M, Rehbach K, Brunner JW, Classen JA, Lammertse HCA, van Linge LA, Schut D, Krutenko
T, Hebisch M, Cornelisse LN *et al* (2019) A Single-Cell Model for Synaptic Transmission and
Plasticity in Human iPSC-Derived Neurons. *Cell Rep* 27: 2199-2211 e2196

Miedema SSM, Mol MO, Koopmans FTW, Hondius DC, van Nierop P, Menden K, de Veij
Mestdagh CF, van Rooij J, Ganz AB, Paliukhovich I *et al* (2022) Distinct cell type-specific
protein signatures in GRN and MAPT genetic subtypes of frontotemporal dementia. *Acta
Neuropathologica Communications* 10: 100

Miller JC, Tan S, Qiao G, Barlow KA, Wang J, Xia DF, Meng X, Paschon DE, Leung E, Hinkley SJ
et al (2011) A TALE nuclease architecture for efficient genome editing. *Nature Biotechnology*
29: 143-148

Moritz CP, Mühlhaus T, Tenzer S, Schulenborg T, Friauf E (2019) Poor transcript-protein
correlation in the brain: negatively correlating gene products reveal neuronal polarity as a
potential cause. *Journal of Neurochemistry* 149: 582-604

Mossink B, van Rhijn J-R, Wang S, Linda K, Vitale MR, Zöller JEM, van Hugte EJH, Bak J, Verboven
AHA, Selten M *et al* (2022) Cadherin-13 is a critical regulator of GABAergic modulation in
human stem-cell-derived neuronal networks. *Molecular Psychiatry* 27: 1-18

Mossner JM, Batista-Brito R, Pant R, Cardin JA (2020) Developmental loss of MeCP2 from VIP
interneurons impairs cortical function and behavior. *eLife* 9: e55639

Müller-Kuller U, Ackermann M, Kolodziej S, Brendel C, Fritsch J, Lachmann N, Kunkel H, Lausen
J, Schambach A, Moritz T *et al* (2015) A minimal ubiquitous chromatin opening element
(UCOE) effectively prevents silencing of juxtaposed heterologous promoters by epigenetic
remodeling in multipotent and pluripotent stem cells. *Nucleic Acids Res* 43: 1577-1592

Nadadhur AG, Alsaqati M, Gasparotto L, Cornelissen-Steijger P, van Hugte E, Dooves S,
Harwood AJ, Heine VM (2019) Neuron-Glia Interactions Increase Neuronal Phenotypes in
Tuberous Sclerosis Complex Patient iPSC-Derived Models. *Stem Cell Reports* 12: 42-56

Nadadhur AG, Emperador Melero J, Meijer M, Schut D, Jacobs G, Li KW, Hjorth JJJ, Meredith RM,
Toonen RF, Van Kesteren RE *et al* (2017) Multi-level characterization of balanced inhibitory-
excitatory cortical neuron network derived from human pluripotent stem cells. *PLOS ONE* 12:
e0178533

Nakamura Y, Darnieder LM, Deeb TZ, Moss SJ (2015) Regulation of GABAARs by
phosphorylation. *Adv Pharmacol* 72: 97-146

Neher E (2015) Merits and limitations of vesicle pool models in view of heterogeneous populations of synaptic vesicles. *Neuron* 87: 1131-1142

Nehme R, Zuccaro E, Ghosh SD, Li C, Sherwood JL, Pietilainen O, Barrett LE, Limone F, Worringer KA, Kommineni S *et al* (2018) Combining NGN2 Programming with Developmental Patterning Generates Human Excitatory Neurons with NMDAR-Mediated Synaptic Transmission. *Cell Reports* 23: 2509-2523

Nelson S, Valakh V (2015) Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. *Neuron* 87: 684-698

Ng YH, Chanda S, Janas JA, Yang N, Kokubu Y, Südhof TC, Wernig M (2021) Efficient generation of dopaminergic induced neuronal cells with midbrain characteristics. *Stem Cell Reports* 16: 1763-1776

Nicholas Cory R, Chen J, Tang Y, Southwell Derek G, Chalmers N, Vogt D, Arnold Christine M, Chen Y-Jiun J, Stanley Edouard G, Elefanty Andrew G *et al* (2013) Functional Maturation of hPSC-Derived Forebrain Interneurons Requires an Extended Timeline and Mimics Human Neural Development. *Cell Stem Cell* 12: 573-586

O'Brien RJ, Xu D, Petralia RS, Steward O, Huganir RL, Worley P (1999) Synaptic clustering of AMPA receptors by the extracellular immediate-early gene product Narp. *Neuron* 23: 309-323

Okabe S, Miwa A, Okado H (2001) Spine formation and correlated assembly of presynaptic and postsynaptic molecules. *J Neurosci* 21: 6105-6114

Olah SS, Kareemo DJ, Buchta WC, Sinnen BL, Miller CN, Actor-Engel HS, Gookin SE, Winborn CS, Kleinjan MS, Crosby KC *et al* (2023) Acute reorganization of postsynaptic GABA$_A$ receptors reveals the functional impact of molecular nanoarchitecture at inhibitory synapses. *Cell Reports* 42

Østergaard M, Barbagallo P, Frederiksen HRS, Chung WK, Møller RS, Larsen MR, Freude K, Verhage M, Sørensen JB (2025) SNAP25 variant I67N: synaptic phenotypes, drug response and proteome changes in human neurons. *Brain*

Öttl M, Toonen RF, Verhage M (2024) Reduced synaptic depression in human neurons carrying homozygous disease-causing STXBP1 variant L446F. *Hum Mol Genet* 33: 991-1000

Palop JJ, Chin J, Roberson ED, Wang J, Thwin MT, Bien-Ly N, Yoo J, Ho KO, Yu G-Q, Kreitzer A *et al* (2007a) Aberrant Excitatory Neuronal Activity and Compensatory Remodeling of Inhibitory Hippocampal Circuits in Mouse Models of Alzheimer's Disease. *Neuron* 55: 697-711

Palop JJ, Chin J, Roberson ED, Wang J, Thwin MT, Bien-Ly N, Yoo J, Ho KO, Yu GQ, Kreitzer A *et
al* (2007b) Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory
hippocampal circuits in mouse models of Alzheimer's disease. *Neuron* 55: 697-711

Pantazis CB, Yang A, Lara E, McDonough JA, Blauwendraat C, Peng L, Oguro H, Kanaujiya J, Zou
J, Sebesta D *et al* (2022) A reference human induced pluripotent stem cell line for large-scale
collaborative studies. *Cell Stem Cell* 29: 1685-1702.e1622

Paraskevopoulou F, Herman MA, Rosenmund C (2019) Glutamatergic Innervation onto
Striatal Neurons Potentiates GABAergic Synaptic Output. *J Neurosci* 39: 4448-4460

Park YK, Goda Y (2016) Integrins in synapse regulation. *Nature Reviews Neuroscience* 17: 745-
756

Pearce RA, Grunder SD, Faucher LD (1995) Different mechanisms for use-dependent
depression of two GABAA-mediated IPSCs in rat hippocampus. *J Physiol* 484 (Pt 2): 425-435

Peitz M, Krutenko T, Brüstle O (2020) Protocol for the Standardized Generation of Forward
Programmed Cryopreservable Excitatory and Inhibitory Forebrain Neurons. *STAR Protocols* 1:
100038

Perez-Riverol Y, Bandla C, Kundu DJ, Kamatchinathan S, Bai J, Hewapathirana S, John NS,
Prakash A, Walzer M, Wang S *et al* (2025) The PRIDE database at 20 years: 2025 update.
Nucleic Acids Res 53: D543-d553

Petralia RS (2012) Distribution of Extrasynaptic NMDA Receptors on Neurons. *The Scientific
World Journal* 2012: 267120

Prag S, De Arcangelis A, Georges-Labouesse E, Adams JC (2007) Regulation of post-
translational modifications of muskelin by protein kinase C. *The International Journal of
Biochemistry & Cell Biology* 39: 366-378

Quattrocchi CC, Wannenes F, Persico AM, Ciafré SA, D'Arcangelo G, Farace MG, Keller F (2002)
Reelin is a serine protease of the extracellular matrix. *J Biol Chem* 277: 303-309

Rasmussen Mikkel A, Holst B, Tümer Z, Johnsen Mads G, Zhou S, Stummann Tina C, Hyttel P,
Clausen C (2014) Transient p53 Suppression Increases Reprogramming of Human Fibroblasts
without Affecting Apoptosis and DNA Damage. *Stem Cell Reports* 3: 404-413

Ray A, Treloar HB (2012) IgSF8: a developmentally and functionally regulated cell adhesion
molecule in olfactory sensory neuron axons and synapses. *Mol Cell Neurosci* 50: 238-249

Rhee HJ, Shaib AH, Rehbach K, Lee C, Seif P, Thomas C, Gideons E, Guenther A, Krutenko T,
Hebisch M *et al* (2019) An Autaptic Culture System for Standardized Analyses of iPSC-Derived
Human Neurons. *Cell Reports* 27: 2212-2228.e2217

Richards DA, Mateos JM, Hugel S, de Paola V, Caroni P, Gähwiler BH, McKinney RA (2005)
Glutamate induces the rapid formation of spine head protrusions in hippocampal slice
cultures. *Proc Natl Acad Sci U S A* 102: 6166-6171

Roig Adam A, Martínez-López JA, Van Der Spek SJF, Achsel T, Andres-Alonso M, Bagni C, Bayés
À, Biederer T, Brose N, Chua JJE *et al* (2023) Transcriptional diversity in specific synaptic gene
sets discriminates cortical neuronal identity. *Biology Direct* 18

Rubenstein JLR, Merzenich MM (2003) Model of autism: increased ratio of
excitation/inhibition in key neural systems. *Genes, Brain and Behavior* 2: 255-267

Sanchez-Priego C, Hu R, Boshans LL, Lalli M, Janas JA, Williams SE, Dong Z, Yang N (2022)
Mapping cis-regulatory elements in human neurons links psychiatric disease heritability and
activity-regulated transcriptional programs. *Cell Reports* 39: 110877

Sanderson TM, Georgiou J, Collingridge GL (2020) Illuminating Relationships Between the
Pre- and Post-synapse. *Front Neural Circuits* 14: 9

Sanes JR, Lichtman JW (1999) Development of the vertebrate neuromuscular junction. *Annu
Rev Neurosci* 22: 389-442

Schmid B, Holst B, Poulsen U, Jørring I, Clausen C, Rasmussen M, Mau-Holzmann UA, Steeg R,
Nuthall H, Ebneth A *et al* (2021) Generation of two gene edited iPSC-lines carrying a DOX-
inducible NGN2 expression cassette with and without GFP in the AAVS1 locus. *Stem Cell
Research* 52: 102240

Schmitz SK, Hjorth JJ, Joemai RM, Wijntjes R, Eijgenraam S, de Bruijn P, Georgiou C, de Jong
AP, van Ooyen A, Verhage M (2011) Automated analysis of neuronal morphology, synapse
number and synaptic recruitment. *Journal of neuroscience methods* 195: 185-193

Scimemi A, Diamond JS (2012) The number and organization of Ca2+ channels in the active
zone shapes neurotransmitter release from Schaffer collateral synapses. *J Neurosci* 32:
18157-18176

Sia G-M, Béïque J-C, Rumbaugh G, Cho R, Worley PF, Huganir RL (2007) Interaction of the N-
Terminal Domain of the AMPA Receptor GluR4 Subunit with the Neuronal Pentraxin NP1
Mediates GluR4 Synaptic Recruitment. *Neuron* 55: 87-102

Skene NG, Grant SG (2016) Identification of Vulnerable Cell Types in Major Brain Disorders
Using Single Cell Transcriptomes and Expression Weighted Cell Type Enrichment. *Front*
Neurosci 10: 16

Skowronek P, Thielert M, Voytik E, Tanzer MC, Hansen FM, Willems S, Karayel O, Brunner AD,
Meier F, Mann M (2022) Rapid and In-Depth Coverage of the (Phospho-)Proteome With Deep
Libraries and Optimal Window Design for dia-PASEF. *Mol Cell Proteomics* 21: 100279

Sohal VS, Rubenstein JLR (2019) Excitation-inhibition balance as a framework for
investigating mechanisms in neuropsychiatric disorders. *Molecular Psychiatry* 24: 1248-1257

Song S-H, Augustine GJ (2023) Different mechanisms of synapsin-induced vesicle clustering
at inhibitory and excitatory synapses. *Cell Reports* 42

Song S, Ashok A, Williams D, Kaufman M, Duff K, Sproul A (2021) Efficient Derivation of
Excitatory and Inhibitory Neurons from Human Pluripotent Stem Cells Stably Expressing
Direct Reprogramming Factors. *Curr Protoc* 1: e141

Stachniak TJ, Sylwestrak EL, Scheiffele P, Hall BJ, Ghosh A (2019) Elfn1-Induced Constitutive
Activation of mGluR7 Determines Frequency-Dependent Recruitment of Somatostatin
Interneurons. *J Neurosci* 39: 4461-4474

Subach OM, Cranfill PJ, Davidson MW, Verkhusha VV (2011) An enhanced monomeric blue
fluorescent protein with the high chemical stability of the chromophore. *PLoS One* 6: e28674

Südhof TC (2018) Towards an Understanding of Synapse Formation. *Neuron* 100: 276-293

Südhof TC (2021) The cell biology of synapse formation. *J Cell Biol* 220

Tang AH, Chen H, Li TP, Metzbower SR, MacGillavry HD, Blanpied TA (2016) A trans-synaptic
nanocolumn aligns neurotransmitter release to receptors. *Nature* 536: 210-214

Team R, 2020. RStudio: Integrated Development for R. PBC, Boston, MA.

Thanou E, Koopmans F, Pita-Illobre D, Klaassen RV, Özer B, Charalampopoulos I, Smit AB, Li
KW (2023) Suspension TRAPping Filter (sTRAP) Sample Preparation for Quantitative
Proteomics in the Low µg Input Range Using a Plasmid DNA Micro-Spin Column: Analysis of
the Hippocampus from the 5xFAD Alzheimer's Disease Mouse Model. *Cells* 12

Titlow JS, Kiourlappou M, Palanca A, Lee JY, Gala DS, Ennis D, Yu JJS, Young FL, Susano Pinto
DM, Garforth S *et al* (2023) Systematic analysis of YFP traps reveals common mRNA/protein
discordance in neural tissues. *Journal of Cell Biology* 222: e202205129

Togo K, Fukusumi H, Shofuda T, Ohnishi H, Yamazaki H, Hayashi MK, Kawasaki N, Takei N,
Nakazawa T, Saito Y *et al* (2021) Postsynaptic structure formation of human iPS cell-derived
neurons takes longer than presynaptic formation during neural differentiation in vitro.
Molecular Brain 14: 149

Tucić M, Stamenković V, Andjus P (2021) The Extracellular Matrix Glycoprotein Tenascin C and
Adult Neurogenesis. *Front Cell Dev Biol* 9: 674199

Tyagarajan SK, Ghosh H, Yévenes GE, Imanishi SY, Zeilhofer HU, Gerrits B, Fritschy JM (2013)
Extracellular signal-regulated kinase and glycogen synthase kinase 3 β regulate gephyrin
postsynaptic aggregation and GABAergic synaptic function in a calpain-dependent
mechanism. *J Biol Chem* 288: 9634-9647

Tyagarajan SK, Ghosh H, Yévenes GE, Nikonenko I, Ebeling C, Schwerdel C, Sidler C, Zeilhofer
HU, Gerrits B, Muller D *et al* (2011) Regulation of GABAergic synapse formation and plasticity
by GSK3beta-dependent phosphorylation of gephyrin. *Proc Natl Acad Sci U S A* 108: 379-384

Um JW, Kim KH, Park BS, Choi Y, Kim D, Kim CY, Kim SJ, Kim M, Ko JS, Lee S-G *et al* (2014)
Structural basis for LAR-RPTP/Slitrk complex-mediated synaptic adhesion. *Nature*
Communications 5: 5423

van Berkel AA, Lammertse HCA, Öttl M, Koopmans F, Misra-Isrie M, Meijer M, Dilena R, van
Hasselt PM, Engelen M, van Haelst M *et al* (2024) Reduced MUNC18-1 Levels, Synaptic
Proteome Changes, and Altered Network Activity in STXBP1-Related Disorder Patient
Neurons. *Biol Psychiatry Glob Open Sci* 4: 284-298

van Boven MA, Mestroni M, Zwijnenburg PJG, Verhage M, Cornelisse LN (2024) A de novo
missense mutation in synaptotagmin-1 associated with neurodevelopmental disorder
desynchronizes neurotransmitter release. *Mol Psychiatry* 29: 1798-1809

Velasco CD, Santarella-Mellwig R, Schorb M, Gao L, Thorn-Seshold O, Llobet A (2023)
Microtubule depolymerization contributes to spontaneous neurotransmitter release in vitro.
Communications Biology 6: 488

Waites C, Qu X, Bartolini F (2021) The synaptic life of microtubules. *Curr Opin Neurobiol* 69:
113-123

Watanabe M, Fukuda A (2015) Development and regulation of chloride homeostasis in the
central nervous system. *Front Cell Neurosci* 9: 371

Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis*. Springer International
Publishing

Wiera G, Brzdąk P, Lech AM, Lebida K, Jabłońska J, Gmerek P, Mozrzymas JW (2022) Integrins
Bidirectionally Regulate the Efficacy of Inhibitory Synaptic Transmission and Control
GABAergic Plasticity. *J Neurosci* 42: 5830-5842

Wilber A, Linehan JL, Tian X, Woll PS, Morris JK, Belur LR, McIvor RS, Kaufman DS (2007)
Efficient and stable transgene expression in human embryonic stem cells using transposon-
mediated gene transfer. *Stem cells* 25: 2919-2927

Wiśniewski JR, Gaugaz FZ (2015) Fast and sensitive total protein and Peptide assays for
proteomic analysis. *Anal Chem* 87: 4110-4116

Wolf D, Goff SP (2009) Embryonic stem cells use ZFP809 to silence retroviral DNAs. *Nature*
458: 1201-1204

Wu J, Ren J, Cui H, Xie Y, Tang Y (2024) Rapid and high-purity differentiation of human medium
spiny neurons reveals LMNB1 hypofunction and subtype necessity in modeling Huntington's
disease. *Inflamm Regen* 44: 7

Yang N, Chanda S, Marro S, Ng Y-H, Janas JA, Haag D, Ang CE, Tang Y, Flores Q, Mall M (2017)
Generation of pure GABAergic neurons by transcription factor programming. *Nature methods*
14: 621-628

Yao S, Sukonnik T, Kean T, Bharadwaj RR, Pasceri P, Ellis J (2004) Retrovirus Silencing,
Variegation, Extinction, and Memory Are Controlled by a Dynamic Interplay of Multiple
Epigenetic Modifications. *Molecular Therapy* 10: 27-36

Yim YS, Kwon Y, Nam J, Yoon HI, Lee K, Kim DG, Kim E, Kim CH, Ko J (2013) Slitrks control
excitatory and inhibitory synapse formation with LAR receptor protein tyrosine phosphatases.
Proc Natl Acad Sci U S A 110: 4057-4062

Zhang SJ, Huguenard JR, Prince DA (1997) GABAA Receptor-Mediated Cl− Currents in Rat
Thalamic Reticular and Relay Neurons. *Journal of Neurophysiology* 78: 2280-2286

Zhu Y, Orre LM, Zhou Tran Y, Mermelekas G, Johansson HJ, Malyutina A, Anders S, Lehtiö J
(2020) DEqMS: A Method for Accurate Variance Estimation in Differential Protein Expression
Analysis. *Mol Cell Proteomics* 19: 1047-1057

Zott B, Busche MA, Sperling RA, Konnerth A (2018) What Happens with the Circuit in
Alzheimer's Disease in Mice and Humans? *Annu Rev Neurosci* 41: 277-297
1197
1198

1199

Figure 1. PiggyBac Asc1/Dlx2 efficiently generates pure GABAergic neurons,

A Map of the all-in-one PiggyBac (PB) vector, expressing murine *Ascl1* and *Dlx2* from a bi-directional TRE-responsive promoter, rtTA from the CAG promoter and *pac*-T2A-NLS-mTagBFP from the EF1α promoter. **B** Overview of the protocol for *Ascl1/Dlx2* induction. **C**

Before induction, all iPSCs show nuclear BFP signals, demonstrating successful integration
of the PB construct **D** Dox-induced expression of *Dlx2* and *Ascl1* triggers neuronal
morphology to arise within one day (DPI1) **E** A week after induction (WPI 1), neural
morphology is clearly visible **F** After 10 weeks of culture, PB *Ascl1/Dlx2* induced neurons
show mature neuronal morphology with GABAergic synapses (VGAT+ puncta) and no
glutamatergic synapses (VGLUT+ puncta) Scale bar: 100 µm **G.** Zoom-in on the dendrite
highlighted in F. scale bar = 1 µm. **H** Autaptic cultures of PB *Ascl1/Dlx2* induced neurons were
assessed using patch clamp electrophysiology. **I** Evoked postsynaptic currents (PSCs) were
fitted using a mono-exponential decay function (red) and their currents were diminished by
the application of PTX. **J** Dot-plot of the decay time constants of ePSCs recorded in PB
Ascl1/Dlx2 induced neurons. **K** Dot-plot showing the distribution of cells with 0-10% of the
naïve charge remaining in PTX **L** Overview of the four induction methods used, PB *Ascl1/Dlx2*
induction was compared to the lentiviral *Ascl1/Dlx2* induction, *Ascl1*+Forskolin induction and
NGN2 glutamatergic induction. **M** Example ePSC traces with either fast or slow decay, as
measured in the inductions tested here. **N** Histogram of log-transformed decay time
constants (decay τ) measured in neurons induced with the four different induction strategies,
together with a 'sum of two gaussians' fit (grey line). **O** Pie-charts showing the percentages
of neurons with a slow ($\tau \geqslant 30$ ms), intermediate ($6 > \tau < 30$ ms) and fast ($\tau \leqslant 6$ ms) decay, as
well as the percentage of synaptically silent neurons, of neurons generated with the four
different induction strategies.
1224

1225

1226

Figure 2. PiggyBac Ascl1/Dlx2 neurons integrate into well-defined excitatory/inhibitory networks upon co-culturing with NGN2 induced neurons

A Schematic of co-culturing protocol in which PiggyBac (PB) Ascl1/Dlx2 (AD) neurons were co-cultured with glutamatergic NGN2 induced neurons to generate excitatory/inhibitory (E/I) co-cultures **B** Representative images of E/I co-cultures at different ratios in which dendrites and neuronal nuclei are visualised using MAP2 and NeuN staining, respectively, and PB-AD neurons visualised by nuclear mTagBFP signals. The E/I-ratio is indicated above each image. Scale bar = 100 μm. **C** Representative images of E/I co-cultures at the same ratios as depicted in (B), stained for MAP2 as dendritic marker and VGAT and VGLUT as markers for GABAergic and glutamatergic synapses, respectively. Scale bars = 100 μm, scale bar zoom in = 10 μm **D** The network density (number of nuclei/field of view) is stable across E/I co-culture ratios. **E** The total dendrite length per field of view decreases as the percentage of plated PB-AD

1239 neurons increases. **F** The density of glutamatergic (VGLUT$^+$ puncta) and GABAergic (VGAT$^+$)

1240 puncta scales with the percentage of NGN2 and PB-AD neurons plated. **G** Synapse counts

1241 shown in F expressed as the percentage of all synapses counted.

1242

Figure panels A–H.

1244

Figure 3. PiggyBac *Ascl1/Dlx2* neurons have a cortical GABAergic proteomic profile.

A. Schematic overview of the proteomics analysis in which PiggyBac (PB) Ascl1/Dlx2 (AD) rat astrocyte co-cultures were compared to NGN2 co-cultures per timepoint. Double-sided arrows indicate the contrasts used for analysis. **B**. PCA analysis on neuronal protein abundances reveals separation of culture types as well as time points along the first two components (PC1: 61.7% and PC2: 8.0%). **C-F**. Volcano plots showing regulation of neuronal proteins in PB-AD cultures and in NGN2 cultures at 4 WPI (**C**), 6 WPI (**D**), 8 WPI (**E**) and 10 WPI (**F**). The top 5% significantly upregulated neuronal proteins in PB-AD are shown as blue dots and those in NGN2 as red dots. Proteins are considered significantly regulated if their adjusted p-value is < 0.05 (significance threshold overlaps with x-axis at this scale and is not shown for clarity of the figure) and their $\log_2$ (FC) lies above the 95th or below the 5th percentile (vertical dashed lines). Label colours indicate the predicted cell types based on

cell type specificity scores (CTSS): regulated GABAergic (purple) and glutamatergic proteins (orange). Cell type specificity was defined using the following CTSS cutoffs: neuronal proteins (neuronal CTSS>70%), GABAergic proteins (GABAergic CTSS > 80% and glutamatergic CTSS < 10%), glutamatergic proteins (glutamatergic CTSS > 80% and GABAergic CTSS < 10%). CTSS can be found in Figure S6. **G**. EWCE analysis of top 5% upregulated neuronal proteins in PB-AD cultures reveals specific enrichment for GABAergic neurons, which increased up to 8 WPI. **H**. EWCE analysis of the top 5% upregulated neuronal proteins in NGN2 cultures reveals significant enrichment for both glutamatergic and GABAergic neurons depending on the timepoint.

1269

1270

Figure 4. Morphological development of PiggyBac Ascl1/Dlx2 induced neurons

A Representative images of an autaptic PiggyBac (PB) *Ascl1/Dlx2* (AD) induced neuron at week-post-induction 10, stained for MAP2 as dendritic marker, and VGAT and VGLUT1/2 as

GABAergic and glutamatergic markers, respectively. Scale bar: 100 µm **B** Cut-out of dendrite
marked in A to illustrate individual synaptic puncta. Scale bar: 10 µm **C** Typical examples of
PB-AD autaptic neurons at WPI 4 (i), 6 (ii), 8 (iii) and 10 (iv). Scale bar: 100 µm **D** Total dendrite
length quantified. **E** number of GABAergic synapses per neuron. **F** Synaptic density. **G** Average
VGAT puncta intensity in arbitrary units (AU). Scale bars: 100 µm. Boxplot whiskers extend
the entire data range. **H** Sholl analysis of PB-AD autaptic neurons at WPI 4, 6, 8 and 10.
Shaded areas represent the SEM. **I** Quantification of the area under the curve (AUC) of the
Sholl histogram shown in (**H**).
1282

1283

1284

Figure 5. Functional development of PiggyBac Ascl1/Dlx2 induced neurons

A Representative traces of mPSCs recorded from PiggyBac (PB) *Ascl1/Dlx2* (*AD*) neurons (PB-AD) neurons at 4, 6, 8 and 10 weeks post induction (WPI). **B** mPSC frequency and **(C)**

amplitude of PB-AD neurons at 4, 6, 8 and 10 WPI. **D** Boxplots of the average fractions of synaptically active cells at 4, 6, 8 and 10 WPI. Markers represent averages of individual inductions. **E** Representative ePSC traces of at 4, 6, 8 and 10 WPI. **F** Amplitude charge (**G)** and decay time (**H**) constant of ePSCs recorded from PB-AD neurons at 4, 6, 8 and 10 WPI. **I** Representative traces of paired-pulse ePSCs at a 50, 200 and 1000 ms interval at 10 WPI. **J** Quantified paired-pulse amplitude ratios (PPR) at a 50 ms, 200 ms (**K**) and 100 ms (**L**) interval at 4, 6, 8 and 10 WPI. **M** Representative traces of iPSCs evoked by 20Hz stimulus trains from PB-AD neurons at 4, 6, 8 and 10 WPI with zoom in on first five pulses. **N** Plot of cumulative iPSC charge evoked by 20 Hz stimulus trains, with back-extrapolated linear fits displayed in red. Note, WPI 4 and WPI 6 data overlap. **O** RRP estimate obtained from back-extrapolation of linear fits displayed in N.

Figure 6. Pre- and postsynaptic maturation of PiggyBac Ascl1/Dlx2 neurons

A. Schematic overview of the proteomics analysis with double-sided arrows indicating the contrasts used for analysis. **B-C**. Stacked bar plots showing the number of up- and

downregulated SYNGO annotated proteins over time, distinguishing synaptic (black),
exclusively presynaptic (green) and exclusively postsynaptic (magenta) proteins for PB-AD
(**B**) and NGN2 (**C**) neurons. Classifications were assigned based on SynGO annotations,
synaptic classification contains both pansynaptic and ambiguously localized synaptic
proteins. Regulated proteins have a $\log_2$ (FC) > 0.1, or < -0.1, and adjusted p-value < 0.05. **D-**
F. Volcano plots showing regulation of synaptic proteins between 6 and 4 (**D**), 8 and 6 (**E**),
and 10 and 8 WPI (**F**). Only SYNGO-annotated proteins are plotted and colour coding refer to
significantly regulated synaptic (black) exclusively presynaptic (green) or postsynaptic (pink)
annotations. Proteins were regarded significantly regulated when adjusted p-value < 0.05
(dashed horizontal line) and $\log_2$ (FC) > 0.1 or < -0.1 (dashed vertical lines). Top 5% proteins
regulated proteins were labelled based on 97.5^{th} and 2.5^{th} percentiles marking genes that
drive subsequently detected enrichments. **G-I**. SYNGO plots showing enrichment scores (z-
scores) derived from gene-set enrichment analysis with GOAT on SYNGO cellular
compartment gene sets, using differential expression output of 6 vs 4 (G), 8 vs 6 (H) and 10
vs 8 WPI (I),. Negative z-scores reflect enrichments primarily driven by downregulated
proteins and vice versa for positive z-scores. **J-L.** Lollipop charts showing significant
ontologies with their corresponding adjusted p-values and the number of genes part of the
gene set as analysed by GOAT for 6 vs 4 (J), 8 vs 6 (K) and 8 vs 10 WPI (L) **M.** Schematic
overview of the GABA puff experiment, in which perforated-patch was performed on PB-AD
neurons in E/I-networks and GABA was applied to measure the postsynaptic current it
produced. **N.** Typical example of the postsynaptic current produced by GABA puff at different
holding voltages. **O.** Linear fit of PSC amplitudes as measured in response to GABA puff at 4,
6, 8 and 10 WPI. Error bars represent the SD.
1328
1329

1330

Figure 7. Specialization of PiggyBac Ascl1/Dlx2 and NGN2 neurons into cortical subtypes

A. Schematic overview of the proteomics analysis pipeline. The differential expression of synaptic proteins over time (10 vs 4 WPI) was first quantified for PB-AD and NGN2 culture separately. The difference in expression over time between PB-AD and NGN2 (delta- $\log_2$ (FC)) was next calculated, enabling identification of proteins with PB-AD- or NGN2-specific temporal profiles. Density plot showing 5^{th} and 95^{th} percentiles (dashed vertical lines) selections of proteins that have strong differential temporal regulation (i.e., high or low

delta- $\log_2$ (FC)) in PB-AD compared to NGN2, and vice versa. **B-C**. Volcano plots showing
regulation of synaptic proteins between 10 and 4 WPI. Proteins are considered regulated
when the $\log_2$ (FC) > 0.1 or < -0.1 (dashed vertical lines) and adjusted p-value < 0.05 (dashed
horizontal line). Only SYNGO-annotated proteins are plotted, and dot colors refer to
significantly regulated synaptic (black) exclusively presynaptic (green) or postsynaptic (pink)
annotations. Label colours indicate for predicted cortical GABAergic- (purple) or
glutamatergic (orange) regulated proteins. **D-G** SYNGO sunburst plots showing the
GABAergic and glutamatergic specification of PB-AD and NGN cultures. The SYNGO sunburst
inputs are the proteins that are part of the 5^{th} (**F-G**) and 95^{th} percentiles (**D-E**) from the delta-
$\log_2$ (FC) distribution (**A**), which represent the top 5% exclusively PB-AD and NGN2 regulated
proteins, respectively. GABAergic or glutamatergic annotation for these SYNGO terms is
estimated based on the differential RNA expression ($\log_2$ (FC)) between cortical GABAergic
and glutamatergic neurons of corresponding genes (Jorstad *et al.*, 2023). Both cellular
compartment (**D-F**) and biological process (**E-G**) gene sets were used. **H-K**. EWCE analysis of
PB-AD-specific (**H-I**) or NGN2-specific proteins (**J-K**) show enrichment for GABAergic and
glutamatergic cell sub (classes), respectively. IT and ET refer to intra- and
extratelenchephalic-projecting neurons and NP refers to near-projecting neurons (Jorstad *et*
al., 2023).
1358
1359
1360

1361

Figure S1. NGN2 neurons produce glutamatergic transmission

A Voltage clamp recordings of 6 WPI NGN2 autaptic neurons after stimulation, in normal

ACSF (Naive) and in presence of glutamatergic receptor blockers (APV + CNQX). Synaptic

transmission was fully abolished by blockers, illustrating that NGN2 neurons are

glutamatergic. **B-C** Examples of immunocytochemistry for MAP2, VGAT and VGLUT1+2

performed on NGN2 autaptic neurons, with zoom in on a length of dendrite (**C**). Scale bar =

100 µm, zoom in scale bar = 10 µm

1369

Figure S2. PiggyBac Ascl1/Dlx2 induction reliably generates GABAergic neurons in the KOLF2.1J and GM23973 line

A Representative images of PiggyBac (PB) Ascl1/Dlx2 (AD) neurons induced from KOLF1.2J iPSCs at 7 weeks post induction (WPI), stained for MAP2 as dendritic marker, and VGAT and VGLUT1/2 as GABAergic and glutamatergic markers, respectively**.** Scale bar: 100 μm **B** Zoom-in on dendrite to illustrate punctate VGAT staining. Scale bar: 10 μm **C.** Total dendrite length and (**D**) number of GABAergic synapses per neuron. **E** Synaptic density and intensity (**F**). **G** Representative mPSC trace and (**H**) representative ePSC trace from PB-AD neurons induced from the KOLF2.1J line. **I** Frequency and (**J**) amplitude of miPSCs. **K** Amplitude, (**L**) charge and (**M**) decay time constant of iPSCs in PB-AD neurons generated from the KOLF2.1J line. **N** Representative trace of iPSCs evoked by 20Hz stimulation trains with zoom-in on first five pulses in KOLF2.1J. **O** Plot of cumulative iPSC amplitudes evoked by 20 Hz stimulus trains, with back-extrapolated linear fits displayed in red. **P** RRP estimate obtained from back-extrapolation of linear fits displayed in (**M**). **Q**. representative ePSC traces from PB-AD neurons induced from the GM23973 line. **R.** Representative trace of iPSCs evoked by 20Hz stimulation trains with zoom-in on first five pulses in GM23973. **S** Amplitude, (**T**) charge and (**U**) decay time constant of iPSCs in PB-AD neurons generated from the GM23973.1J line. **V** Plot of cumulative iPSC amplitudes evoked by 20 Hz stimulus trains, with back-extrapolated linear fits displayed in red. **W** RRP estimate obtained from back-extrapolation of linear fits displayed in (M). Boxplot whiskers extend the entire data range.

1393

Figure S3. Peptide and protein detection count and variation is similar between cultures and timepoints.

A-B Boxplots showing variation of peptide (**A**) and protein (**B**) detection for all timepoints per culture, indicating median, 1st and 3rd quartiles, and minimum and maximum values. Number of detected peptides and proteins are indicated in the base of the graph. Median coefficient of variation for peptide and protein detection across groups ranged from 14.6% to 32.2%, reflecting consistent detection performance across replicates. **C** PCA plots showing potential batch effects due to induction. PC1 and PC2 show even spread of different inductions but PC3 (3.9%) shows slight separation based on induction and we therefore corrected for induction effect when performing differential abundance analysis.

1404

1405

Figure S4. Cell type specificity scores for detected GABAergic proteins

1406

A. Cell type specificity scores (CTSS) scores for detected GABAergic and glutamatergic

proteins. CTSS are calculated with the expression weighted cell type enrichment tool (Skene

& Grant, 2016) using single-cell RNA-sequencing data from Jorstad et al. (Jorstad *et al.*, 2023).

MGE and CGE refer to medial and caudal ganglionic eminences. **B.** Representative images of

PB-AD and NGN2 neurons stained for MAP2 and PVALB. Scale bar = 100 µm. **C-D** Expression-

weighted cell type enrichment (EWCE) analysis showing enrichment of proteins higher

expressed in PB-AD or NGN2 for GABAergic **(C)** and glutamatergic subtypes **(D)**. PBAD-

enriched proteins showed strongest enrichment for VIP$^+$ interneurons at most timepoints,

whereas NGN2-enriched proteins map at different timepoints to both GABAergic (PVALB$^+$

and VIP$^+$ interneurons) and glutamatergic subtypes. PB-AD-enriched proteins showed

unexpected enrichment for L5ET neurons at 4 WPI, which reduced below significance

threshold at later timepoints. IT and ET refer to intra- and extratelenchephalic-projecting

neurons, and NP refers to near-projecting neurons – cell subclass annotations consistent

with Jorstad et al. (2023) (Jorstad *et al.*, 2023).

1421

Figure S5. Differential expression of synaptic genes in NGN2 and PB-AD neurons

A-C. Volcano plots showing regulation of synaptic proteins in NGN2 cultures between 6 and 4 (**A**), 8 and 6 (**B**), and 10 and 8 WPI (**C**). Only SYNGO-annotated proteins are plotted and colour coding refer to significantly regulated synaptic (black) exclusively presynaptic (green) or postsynaptic (pink) annotations. Proteins were regarded significantly regulated when adjusted p-value < 0.05 (dashed horizontal line) and $\log_2$ (FC) > 0.1 or < -0.1 (dashed vertical lines). **D-F**. SYNGO plots showing enrichment scores (z-scores) derived from gene-set enrichment analysis with GOAT using SYNGO cellular compartment gene sets for 4 to 6 (**D**), 6 to 8 (**E**) and 8 to 10 WPI (**F**). Negative z-scores (blue) reflect enrichments primarily driven by downregulated proteins and vice versa for positive z-scores (red). **G-I.** Lollipop charts showing significant SYNGO terms with their corresponding adjusted p-values and the number of genes part of the gene set. **J-O.** SYNGO overrepresentation analysis of upregulated proteins in PB-AD (**J-L**) and NGN2 (**M-O**) per timepoint. Foreground are significantly upregulated proteins and background are all tested genes in corresponding contrast.

1436

Figure S6. PiggyBac Ascl1/Dlx2 and NGN2 neurons express markers of diverse GABAergic neuron subtypes

A. Contrasts used to determine cell type specification. **B** and **C** Heatmaps that show temporal expression profiles relative to 4 WPI (log$_2$ (FC)) and cell type specificity scores (CTSS). **B** Typical proteins required for neurotransmitter (NT) synthesis, vesicle loading, and receptors. Undetected proteins are visualized for NT biosynthesis and vesicle loading but not for receptors. **C** Canonical GABAergic subtype markers and proteins with high cell type specificity for GABAergic subtypes and for medial or caudal ganglionic eminences (MGE or CGE). **D.** Line plots show mean expression of canonical GABAergic subtype markers (top) and other

1446   proteins with GABAergic cell type specificity. Error bars represent SD and dot size indicate

1447   detection count.

1448

1449 **Supplemental table 1: SYNGO Proteins with significant expression changes from 8 to 10 WPI**

1450 **in PB-AD neurons**

Protein	new at WPI 10	log$_2$fc	FDR-adjusted p-value
CNTNAP4	yes	1.05	1.50E-26
SLITRK1	yes	0.97	7.75E-23
CDH1	yes	0.91	4.81E-20
EPHA7	yes	0.90	1.26E-19
ITGB3	no	0.80	1.72E-15
PRRT1	yes	0.74	1.35E-13
NPTX2	no	0.73	4.68E-13
ITGA2	no	0.67	3.46E-11
SLC17A5	yes	0.67	5.09E-11
KCNQ2	yes	0.65	1.77E-10
ELFN1	no	0.65	2.09E-10
CYTH1	no	0.65	2.21E-10
TNC	no	0.64	3.74E-10
NRGN	yes	0.63	6.71E-10
NGDN	yes	0.63	6.73E-10
GPR158	no	0.62	1.43E-09
RELN	yes	0.57	3.53E-08
SCG2	no	0.78	2.18E-03
TMEM163	no	0.62	7.64E-03
PTPRN	no	0.59	2.24E-02
NEFH	no	0.40	2.24E-02
IGSF8	no	0.40	2.24E-02
CNTNAP1	no	0.34	3.78E-02
ATP2B4	no	0.30	3.78E-02

1451