1 **One-step induction of human GABAergic neurons promotes presynaptic development**
2 **& synapse maturation**
3

Torben W. van Voorst[1,+], Maaike A. van Boven [1,+], Kevin I. Marinus[2,+], Jennifer M.
Colón-Mercado[3], Jule Schretzmeir[1], Carolin Haag[1,4], Ruud F. Toonen[1,4], Frank
Koopmans[2], Michael E. Ward[3], August B. Smit[2], Ronald E. van Kesteren[2], Matthijs
Verhage[1,4,5], L. Niels Cornelisse[1,5,#]

[1]Department of Functional Genomics, Center for Neurogenomics and Cognitive Research (CNCR),
VU University Amsterdam, de Boelelaan 1100, 1081 HZ Amsterdam, The Netherlands.
[2]Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive
Research (CNCR), VU University Amsterdam, de Boelelaan 1100, 1081 HZ Amsterdam, The
Netherlands.
[3]National Institute of Neurological Disorders and Stroke, National Institutes of Health, 9000
Rockville Pike, Bethesda, MD, USA
[4]Neurospector, de Boelelaan 1100, 1081 HZ Amsterdam, The Netherlands.
[5]Department of Human Genetics, Amsterdam UMC – location VU University Amsterdam, de
Boelelaan 1100, 1081 HV Amsterdam, The Netherlands.
18
19 *[+] Authors contributed equally*
20 *[#] Correspondence: l.n.cornelisse@vu.nl*
21

Abstract (169 words)

Human induced pluripotent stem cells (iPSCs) present a powerful approach to study human
brain physiology and disease, yet robust, pure GABAergic induction has remained difficult.
Here we present improved, single-step, transposon-based GABAergic induction with
Ascl1/Dlx2, which yields, unlike lentiviral approaches, exclusively GABAergic neurons and
was validated across three independent iPSC lines. Co-seeding with Ngn2-induced excitatory
neurons created stable networks of predefined excitation/inhibition ratios, with
corresponding synapse ratios. Proteomic and electrophysiological characterization at
different developmental time points showed that the single-step induced GABAergic neurons
gain a proteomic profile that maps to different cortical interneuron subtypes and display
typical GABAergic synaptic properties, producing large, synchronous and picrotoxin-sensitive
currents. During early development, synaptic strength increased threefold, which was
accompanied by an increase in expression of proteins exclusively enriched for presynaptic

SYNGO terms. Synaptic strength continued to increase during late development but with
only minor proteomic changes. Taken together, transposon-based GABAergic induction yields
exclusively mature GABAergic neurons suitable for studying genes involved in synaptic
maturation and to build excitation/inhibition networks for disease modelling.
39

40 **Keywords (max 10)**
41 Induced pluripotent stem cells, induced GABAergic neurons, Transposon, Synaptic maturation,
42 Ascl1, Dlx2, Autaptic neurons, Proteomics, Patch-clamp, Excitation/inhibition
43

Introduction
Human induced pluripotent stem cell (iPSC)-derived models provide a promising platform
for studying human brain physiology and pathology. iPSCs can be generated from patient-
derived tissue or blood to better reflect the human genetics and (patho)physiology compared
to commonly employed rodent models.[1-3] GABAergic dysfunction and excitation/inhibition
(E/I) disbalance have been implicated in many neurodevelopmental and neurodegenerative
disorders.[4-9] In recent years, various strategies have been developed for generating iPSC-
derived GABAergic neurons, but robust, pure GABAergic induction remains challenging.
Directed differentiation, which employs signalling molecules to mimic neurodevelopment.[10-
12], yields both glutamatergic and GABAergic neurons, but these type of inductions often
generate inconsistent E/I ratios, tend to yield heterogeneous and immature neuronal
populations and are costly and time-consuming.[13] Rapid differentiation approaches rely on
overexpression of key transcription factors, such as Ascl1 and Dlx2 for GABAergic
differentiation.[14,15] and often rely on lentiviral delivery of transgenes. However, such
approaches have several limitations: (1) they require multiple vectors when delivering
multiple transgenes due to the size cap of ~8 kb, (2) they pose a risk of disrupting host
genes[16], and (3) they are susceptible to epigenetic silencing during prolonged iPSC culturing
[17-19] To overcome these challenges, TALEN- and CRISPR-based gene editing.[20] have been
used to target the transgenes to the *AAVS1* safe harbor locus[21,22] While this ensures stable
transgene expression, allows for delivery of multiple genes in a single construct, and
minimizes off-target effects, it is time and resource-consuming due to the need for clonal
expansion and extensive validation of every iPSC line used in a study.[23]
The PiggyBac transposon system has emerged as a gene-delivery tool that overcomes
most of these limitations. Unlike lentiviral systems, transposons enable stable genomic
integration of large genetic constructs up to 200 kb.[24] through a "cut-and-paste"

mechanism[25], which enables delivery of multiple transgenes in a single construct.
Additionally, transposon-delivered transgenes are less prone to silencing than lentivirally
delivered transgenes[26] and the increased packing capacity permits the inclusion of
epigenetic regulators to further diminish silencing. Compared to safe harbor insertion of
transgenes, the delivery of a transposon cassette is a simpler, single-step process applicable
to multiple iPSC lines in parallel and allows the use of a polyclonal population for most use
cases, eliminating the need for thorough validation, thereby accelerating experimental
timelines.

Here, we characterize PiggyBac transposon-based overexpression of *Ascl1* and *Dlx*2 (PB-
AD) for GABAergic neuronal induction and benchmark this against established GABAergic
induction methods. PB-AD-based induction results in exclusively GABAergic differentiation
across different genetic backgrounds, unlike lentiviral approaches which induce both
GABAergic and glutamatergic differentiation. We demonstrate that PB-AD neurons integrate
into synaptically connected E/I networks when co-cultured with excitatory NGN2 neurons
with synaptic projections proportional to cellular ratios. Using proteomics, patch-clamp
electrophysiology, and immunocytochemistry we show that PB-AD induction yields neurons
with reliable, plastic synapses that synchronously release GABA and strengthen during
prolonged culture, with pronounced presynaptic proteomic changes. We propose that this
improved GABAergic induction protocol, which reliably generates pure GABAergic neurons,
enables the study of human GABAergic functioning in pure GABAergic networks, in single-
neuron (autaptic) cultures, and stable E/I-networks.

Results

GABAergic differentiation of human iPSCs using PiggyBac based delivery of Ascl1 and Dlx2
Overexpression of the transcription factors Ascl1 and Dlx2 drives GABAergic lineage
specification of induced pluripotent stem cells (iPSCs)[15] Previous approaches employed
separate lentivectors to express both genes in an inducible manner. We aimed to develop a
simplified GABAergic differentiation protocol using a single PiggyBac transposon construct
containing both transcription factors (RRID: Addgene_182307, Figure 1A). This vector contains
(1) a doxycycline (dox)-inducible expression cassette for *Ascl1* and *Dlx2* from a bidirectional
promoter, avoiding variation in expression levels resulting from position effects in multi-
cistronic vectors; (2) reverse tetracycline-controlled transactivator (rtTA), required for dox-
inducible expression; (3) the puromycin resistance gene (pac), allowing for antibiotic
selection, as well as (4) mTagBFP2[27], giving transfected cells a fluorescent marker; and (5)

Ubiquitous Chromatin Opening Elements (UCOE)[28] and 5'HS chicken β-globin (cHS4)
insulator elements.[29] Firstly, we tested whether our newly designed construct could induce
GABAergic differentiation (Figure 1B). To this end, we co-transfected this PiggyBac vector
with a transposase vector into BIONi010-C iPSCs.[30] After puromycin-selection, all cells
expressed mTagBFP2 (Figure 1C), confirming successful transgene integration. Upon dox-
induction, cells first clustered together (Figure 1D) and after releasing and reseeding at 3
days post-induction (DPI), cells developed a neuronal morphology, which became more
distinct within one week (Figure 1E). At 10 weeks post-induction (WPI), neurons displayed
MAP2+ dendrites and VGAT$^+$ puncta, indicative of GABAergic synapses, whereas glutamatergic
VGLUT1/2$^+$ puncta were absent (Figure 1F,G).

GABAergic synaptic transmission in PiggyBac Ascl1/Dlx2 neurons
To confirm the GABAergic identity of PiggyBac-induced neurons we performed voltage-clamp
recordings in autaptic cultures.[31] between 6 and 10 WPI (Figure 1H). Rapid depolarization to
+30 mV induced action potential evoked postsynaptic currents (ePSCs), which displayed slow
decay times (τ) characteristic of GABAergic currents[32], as determined by mono-exponential
fits (geometric mean (GM) = 104.2 ms, geometric SD factor (GM$_{SDf}$) = 1.42; Figure 1I,J).
Furthermore, virtually all synaptic signalling was abolished after applying the GABA receptor
antagonist picrotoxin (PTX), confirming GABAergic synaptic identity (Figure 1I,K).

PiggyBac Ascl1/Dlx2 yields a pure GABAergic neuron population in contrast to lentiviral methods
To benchmark the PiggyBac approach, we compared it with two established lentiviral
methods for GABAergic induction[14,15], and an NGN2-based protocol for glutamatergic
induction[22,33] (Figure 1L). Autaptic neurons were generated using all four protocols and ePSC
decay time constants were quantified using mono-exponential fits. As expected, NGN2-
derived neurons exhibited fast ePSCs decay kinetics typical of glutamatergic transmission
(GM = 2.43 ms, GM$_{SDf}$ = 1.39, Figure 1M,N), confirmed pharmacologically and by the presence
of VGLUT1/2$^+$ puncta (Figure S1). Surprisingly, decay time constants from neurons generated
with the lentiviral GABAergic protocols showed a bimodal distribution which could be fitted
with a sum of two gaussians. More than half of the cells in these cultures exhibited ePSC
kinetics comparable to those in the NGN2 cultures (Lenti AF: GM = 2.82 ms, GM$_{SDf}$ = 0.58,
72.8% of the fit; Lenti AD: GM = 4.62 ms, GM$_{SDf}$ = 0.42, 55.3% of the fit). The remaining cells
had more than tenfold larger time constants typical for slow GABAergic ePSCs (Lenti AF: GM
= 86.3 ms, GM$_{SDf}$ = 1.56 for 27,2% of the fit; Lenti AD: GM = 60.5 ms, GM$_{SDf}$ = 1.26 for 44.7% of
the fit). In contrast, the PB-AD method produced a unimodal population with slowly decaying
ePSCs (GM = 78.9 ms, GM$_{SDf}$ = 1.35). Since NGN2 and PB-AD neurons produced decay time

constants with unimodal fits, these values were used to determine cut-off thresholds for
synaptic identity (glutamatergic: τ ≤ 6 ms; GABAergic: τ ≥ 30 ms; mixed: 6 < τ < 30 ms). This
revealed that 68% of PB-AD neurons exhibited GABAergic currents, while the remaining 32%
were synaptically silent (Figure 1O). In contrast, the lentiviral approaches generated a
substantial population of glutamatergic (Lenti AF: 43%, Lenti AD: 39%) and mixed (Lenti AF:
9% Lenti AD: 15%) identities. The consistent generation of functional GABAergic neurons and
the absence of glutamatergic neurotransmission underscore the reliability of the PB-AD
protocol compared to previously described methods.

PiggyBac Ascl1/Dlx2 protocol induces GABAergic neurons across genetic backgrounds
To test for generalizability, we applied the PB-AD protocol to generate neurons from two
additional iPSC lines, KOLF2.1J[34] and GM23973[35], and characterized their functional
properties at 7 and 6 WPI, respectively. In both lines the PB-AD approach produced neurons
with GABAergic synaptic projections and slow-decaying synaptic transmission, indicative of
GABAergic specification (Figure S2), as was found for the BIONi010-C line. These findings
demonstrate the protocol's applicability across genetic backgrounds.

PiggyBac Ascl1/Dlx2 neurons integrate into controlled excitatory/inhibitory networks
To test whether PB-AD neurons integrate into E/I networks alongside glutamatergic neurons,
we co-cultured 3 DPI PB-AD neurons with 3 DPI glutamatergic NGN2 neurons in networks
containing 100%, 30% (as found in the human cortex[36-38]) and 0% inhibitory neurons (Figure
2A-C). Immunocytochemistry at 10 WPI (Figure 2B,C) revealed comparable neuronal cell
densities across all conditions, as measured by NeuN staining, indicating similar viability of
PB-AD and NGN2 neurons. (Figure 2D). Total dendrite length per field of view increased as
the proportion of PB-AD neurons decreased, consistent with the distinct morphological
characteristics of GABAergic neurons[39] (Figure 2E). Pure PB-AD networks contained VGAT+
(GABAergic) synaptic puncta, and NGN2 networks contained VGLUT1/2+ (glutamatergic)
synaptic puncta, with in both cases virtually no puncta of the other type present (Figure 2F,G).
In contrast, the mixed (30% PB-AD) networks contained both GABAergic and glutamatergic
puncta on the same dendrite proportional to cellular ratios, demonstrating that PB-AD
neurons are viable components for assembling E/I networks when mixed with NGN2 neurons.

PiggyBac Ascl1/Dlx2 neurons acquire a GABAergic molecular identity over time
Next, we assessed differentiation of PB-AD and NGN2 neurons over time using mass
spectrometry-based proteomics. PB-AD co-cultures with rat astrocytes were compared to
NGN2 co-cultures at 4, 6, 8, and 10 WPI (Figure 3A). We detected 99,154 human peptide

precursors mapping to 7,775 unique proteins across 60 samples that passed quality control. Variability within groups was low-to-moderate with coefficients of variation between 14.6% and 32.2%, reflecting consistent detection across replicates (Figure S3A-B). To remove rat astrocytic signal, subsequent analyses focused on neuronally enriched proteins (1690 proteins), which show >70% neuron-specific expression according to single-cell transcriptomics data of the human cortex (Jostad). Principal component analysis (PCA) on the neuronally-enriched protein abundances (Figure 3B, S3C) revealed clear separation between PB-AD and NGN2 cultures along the first two components (PC1: 71.5% and PC2: 6.1%), as well as the separation of different time points within each culture. More specifically, time points from both PB-AD and NGN2 cultures showed clear separation between 4, 6, and 8 WPI, while cultures at 8 and 10 WPI clustered more closely together, suggesting that proteomic maturation plateaus after 8 weeks.

To assess lineage specification of PB-AD and NGN2 inductions, we annotated cell types based on differentially expressed proteins using cell type specificity scores (CTSS), computed using single-cell RNA expression data from human cortex[40] (Figure S4A). Across all timepoints, canonical GABAergic proteins, including SLC32A1 (VGAT), GAD1, GAD2, were enriched in PB-AD cultures. In addition, other predicted GABAergic proteins, such as CRHBP, FBN2, CRH, and RELN, were enriched in PB-AD neurons. In contrast, only a few predicted glutamatergic proteins (e.g., BDNF, PCP4, CPZ, and ADAMTSL1) were moderately enriched in PB-AD cultures. The glutamatergic marker SLC17A6 (VGLUT2) was enriched in NGN2 cultures, while SLC17A7 (VGLUT1) was not detected (Figure 3C-F). Surprisingly, NGN2 neurons also showed strong enrichment of several proteins typically associated with GABAergic neurons at all timepoints, including SCGN, CALB2, and PVALB. Immunocytochemistry confirmed PVALB expression in the majority of NGN2 neurons (Figure S4B).

To quantitatively assess cell type specification, we performed expression-weighted cell type enrichment (EWCE) analysis[41] on the top 5% upregulated neuronal proteins for each induction for each timepoint. At all timepoints, PB-AD-specific neuronal proteins showed exclusive and significant enrichment for GABAergic neurons, with enrichment levels increasing until 8 WPI (Figure 3G). GABAergic subclass analysis revealed strongest enrichment for VIP[+] interneurons from 6-10 WPI (Figure S4C). In contrast, NGN2-specific proteins showed significant enrichment for glutamatergic neurons only at 6 and 10 WPI, while also displaying unexpected GABAergic enrichment, reaching significance at 4 and 10 WPI (Figure 3H). Subclass enrichment analysis revealed that this unexpected GABAergic enrichment was most likely accounted for by PVALB[+] and VIP[+] interneuron-like profiles (S4C,

S4D). Taken together, PB-AD neurons acquire a stable GABAergic identity over time,
characterized by consistent expression of canonical and subtype-specific GABAergic proteins.
By contrast, NGN2-induced neurons exhibit a mixed molecular identity, with evidence for
partial GABAergic specification in addition to expected glutamatergic features.

Gradual morphological development of PiggyBac Ascl1/Dlx2 neurons
To examine morphological maturation of PB-AD neurons, we performed immunostaining for
MAP2, VGAT and VGLUT1/2 in autaptic cultures (Figure S5A,B). Similar to networks, PB-AD
neurons produced only VGAT$^+$ puncta at all timepoints (Figure S5A,B, Figure 4A-C). A
progressive increase in dendrite length was observed during maturation (Figure 4B),
accompanied by an increase in the total number of VGAT$^+$ puncta per neuron and puncta
densities per dendritic length (Figure 4C,D). Additionally, the intensities of VGAT puncta
increased over time (Figure 4E), suggesting increased synaptic vesicles per synaptic terminal.
Sholl analysis revealed a rise in the number of arborizations during maturation (Figure
S5C,D,I). Interestingly, growth of dendritic length and complexity plateaued by 6 WPI,
whereas synaptic projections continued to develop until 8 and 10 WPI. Together, these results
demonstrate that PB-AD neurons develop complex neuronal morphologies accompanied by
an increase in presynaptic GABAergic terminals during maturation.

Synaptic strengthening during maturation of PiggyBac Ascl1/Dlx2 neurons
Next, we assessed functional maturation in autaptic neurons with voltage-clamp
electrophysiology. Firstly, action-potential independent spontaneous release was recorded.
Both miniature postsynaptic current (mPSC) frequency and amplitude increased during
maturation (Figure 4F-H). Next, neurons were depolarized to induce ePSCs. Across all time
points, 70-80% of neurons were synaptically active, producing a charge of at least 1.5 pC
within 70 ms after stimulation (Figure 4I). In active cells, both the amplitude and charge of
ePSCs increased progressively between 4 and 10 WPI (Figure 4J-L), with the most pronounced
change occurring from 8 to 10 WPI. Decay constants remained consistently above 30 ms –
characteristic of GABAergic currents (Figure 4M). Interestingly, decay kinetics subtly slowed
after 4 WPI, suggesting potential changes in postsynaptic receptor subunits.[42,43]

To evaluate the impact of maturation on short-term synaptic plasticity, we applied
paired-pulse stimulation at 50 ms, 200 ms, and 1000 ms intervals (Figure S5E-H). This
revealed modest synaptic depression which was stable throughout maturation. Only at a
1000 ms interval, synaptic depression increased as neurons matured. Importantly, maturation
increased the size of the readily releasable pool (RRP), as determined by back-extrapolation
of the charge evoked by 20 Hz stimulus trains, with the biggest change occurring from 8 to

10 WPI (Figure S5I-K). Taken together, these findings demonstrate that PB-AD GABAergic
neurons develop functional, plastic synapses that strengthen during maturation.

Functional maturation correlates with upregulation of proteins in presynaptic gene sets

To investigate the molecular basis of synaptic maturation, we looked at the differential
expression of SYNGO-annotated synaptic proteins[44] between different timepoints of
development (4 to 6 WPI, 6 to 8 WPI, and 8 to 10 WPI) (Figure 5A). Of all 1605 SYNGO-
annotated synaptic genes, 929 unique proteins were detected in PB-AD cultures, of which
478 (51%) were differentially expressed in at least one time contrast (Figure 5B, D-F). In
NGN2 neurons, 967 unique SYNGO-annotated proteins were identified of which 644 (67%)
were differentially expressed in at least one time contrast (Figure 5C, Figure S6A-C). In both
cultures, SYNGO-annotated proteins were abundantly regulated between 4 and 8 WPI, while
only few proteins showed altered expression between 8 and 10 WPI (Figures 5B-F,
Supplementary table 1). This indicates that proteomic synaptic maturation plateaus after 8
WPI, consistent with what was observed for neuronal-enriched protein expression.

To examine specifically how pre- and postsynapses develop over time, we used GOAT[45]
to determine which SYNGO gene sets were enriched for developmentally regulated proteins,
where the contribution of each protein was weighted by its $\log_2$(fold change).[45] Strongest
enrichment was observed between 4-6 WPI and 6-8 WPI, which occurred exclusively in
presynaptic gene sets, including *synaptic vesicle (SV)*, *presynaptic active zone (AZ)*, and
presynaptic membrane (Figure 5G-L). Between 8-10 WPI only *SV membrane* proteins and its
mother term *presynaptic* proteins were enriched. In contrast, no postsynaptic gene sets were
significantly enriched in PB-AD cultures, across all time contrasts. To exclude that these
results were biased by a few proteins with large effect sizes, additional SYNGO gene set
overrepresentation analysis was performed based on the number of upregulated proteins
per gene set. This confirmed the exclusive enrichment of presynaptic terms in PB-AD cultures
(Figure S6J–L). Upregulated proteins in NGN2 cultures over time similarly showed a
preferential enrichment of presynaptic gene sets using GOAT and SYNGO overrepresentation
analyses, with no enrichment for postsynaptic gene sets between 6-10 WPI. Thus, synaptic
strengthening correlates with changes specifically in presynaptic protein levels, while
postsynaptic protein levels remain relatively stable over time.

In order to test whether our findings on asymmetrical pre- and postsynaptic
maturation translate to a more physiological model, we assessed postsynaptic responses to
externally applied GABA in E/I networks, as excitation might alter maturation profiles.[46]
(Figure 5M). Furthermore, we used gramicidin-perforated patch to allow measurement of

postsynaptic responses without influencing the reversal potential for the GABA-receptor[47]
GABA puffs were applied at varying holding voltages (Figure 5N). The slope of the I/V curve
voltages was unaltered by maturation (p = 0.305), indicating that the GABAergic postsynaptic
response does not change during maturation under these conditions. Thus, increased
synaptic strength during synaptic maturation correlates with presynaptic proteomic changes,
but not with postsynaptic development also in E/I-networks.

PiggyBac Ascl1/Dlx2 specific synaptic proteins map to cortical interneuron subtypes
Finally, we investigated whether changes in PB-AD and NGN2 SYNGO-annotated proteomes
between 4 and 10 WPI are associated with cortical GABAergic and glutamatergic cell types.[40]
(Figure 6A). PB-AD cultures showed upregulation of predicted GABAergic and medial
ganglionic eminence (MGE) and caudal ganglionic eminence (CGE)-specific proteins (Figure
6B). NGN2 cultures showed upregulation of predicted glutamatergic proteins (Figure 6C). Of
note, PB-AD and NGN2 cultures showed upregulation of both GABAergic and glutamatergic
receptor subunits (Figure S7A,B), although only input of one of the neurotransmitters was
present per culture. Canonical subtype markers for both MGE- and CGE-derived subtypes
were also detected in PB-AD neurons (S7C,D).
To analyze lineage-specific synaptic development in more detail, we next contrasted
the developing (WPI 4 to WPI 10) PB-AD and NGN2 SYNGO-annotated proteomes with each
other to find proteins that were specifically regulated in each induction. To this end, for each
protein we calculated the difference between PB-AD and NGN2 in differential expression
between 4-10 WPI (i.e., $\log_2 (FC)_{AD}$- $\log_2 (FC)_{NGN2}$), and selected the 5th and 95th percentile to
define induction-specific synaptic proteins (Figure 6A). We then assessed whether these
proteins were enriched in either cortical glutamatergic or cortical GABAergic reference
expression profiles[40] Overall, this produced lineage-specific enrichments in most gene sets.
PB-AD specific synaptic proteins showed the highest GABAergic enrichment in the gene sets
dense core vesicle (*DCV*), *SV*, *synaptic signalling*, *metabolism* and *presynaptic process* (Figure 6D-
E). Conversely, NGN2-specific synaptic proteins that belonged to *synaptic signalling, SV,*
organization and *specialization* showed the highest glutamatergic enrichment (Figure 6F-G).
As synaptic gene expression profiles have been shown to reflect neuronal subtypes[48],
we queried whether synaptic proteins that specifically increase in PB-AD or NGN2 cultures
show enrichment for neuronal (sub)types. In EWCE analysis, the PB-AD-specific synaptic
proteins were enriched in GABAergic neurons (Figure 6H), specifically VIP+, SST+, PAX6+ and
LAMP5+ interneurons, but not PVALB+ neurons (Figure 6I). NGN2-specific synaptic proteins
were enriched in glutamatergic subtypes (IT, L4 IT, L5 ET, L6 CT, and L6b) but also in

GABAergic subtypes (Figure 6J), particularly VIP[+] interneurons (Figure 6K). Together, these data indicate that PB-AD-based induction leads to GABAergic synaptic specialization with molecular profiles resembling CGE- and MGE-derived interneurons, whereas NGN2 induction leads to induction of neurons that also contain typical GABAergic proteins.

Discussion

Summary

In this work, we present a single-step, transposon-based GABAergic differentiation method, which uses a single vector to overexpress Ascl1/Dlx2, and we provide a detailed characterization of these GABAergic neurons at different developmental timepoints using mass spectrometry proteomics, immunocytochemistry, and patch-clamp electrophysiology. This method is applicable across different genetic backgrounds and produces exclusively GABAergic neurons with a proteomic profile that maps to different cortical interneuron subtypes, particularly VIP[+] neurons. Synapses display GABAergic properties such as picrotoxin-sensitive, slowly decaying, evoked postsynaptic currents and short-term depression, similar to primary GABAergic neurons.[49-53] During development, synapses gradually increase in synaptic strength, which at earlier timepoints correlates with increased expression of presynaptic proteins associated with active zone, synaptic vesicle, and presynaptic membrane gene sets. In contrast, little functional and proteomic changes occur in the postsynapse. Late in development, synaptic strength continues to increase, with only minor proteomic changes. PB-AD neurons can be combined with NGN2 neurons to form networks of predefined E/I ratios with corresponding synapse ratios. Taken together, transposon-based GABAergic induction yields GABAergic neurons with improved synaptic properties that can be used to build balanced excitation/inhibition networks, for instance to model diseases in which E/I disbalance has been proposed as a disease mechanism.

PiggyBac Ascl1/Dlx2 neuronal induction produces exclusively GABAergic neurons

By employing autaptic neurons we were able to detect synaptic output on a single neuron scale, which allowed us to accurately quantify lineage specification and demonstrate how GABAergic transposon-based induction compares to lentiviral strategies. Previous studies using a lentiviral approach also detected non-GABAergic currents[54] and reported expression of glutamatergic genes[55,56], while others focused on measurements in E/I-networks[14,15] or recorded under conditions that blocked glutamatergic transmission[57,58], making it difficult to compare with our results. As Ascl1 expression, by itself, induces glutamatergic differentiation[59,60], the balance of Ascl1 and Dlx2 expression is likely important in steering

differentiation towards a GABAergic fate. The design of our transposon-based induction
incorporates multiple features to facilitate this and thereby avoids unwanted glutamatergic
specification. Firstly, both factors are present in one cassette and transcribed from a
bidirectional promotor. Secondly, a shared integration site could prevent position effects.[61]
Lastly, insulator elements may reduce gene silencing, which is observed for lentiviral cargo[17-
19,62-65], and less so for transposon delivered transgenes.[26]

Proteomic profiles of PiggyBac Ascl1/Dlx2 neurons map to cortical interneuron subtypes
Cell type enrichment analysis revealed that neuronal proteins that are higher expressed in
PB-AD cultures are significantly enriched for GABAergic neurons at all timepoints, with
specific enrichment for VIP[+] interneurons from 6 WPI onwards. Zooming in on synaptic
proteins that are upregulated specifically in PB-AD cultures between 4-10 WPI, we showed
significant enrichment for all GABAergic interneuron subtypes except PVALB[+], suggesting
specification towards both MGE- and CGE-derived neurons. Assessment of the neuronal
lineages produced by PB-AD induction would require single-cell sequencing approaches. A
previous single-cell RNA-sequencing study of lenti AD neurons predominantly found SST[+]
neurons at 4 WPI, but later time points were not investigated.[66] Perhaps CGE markers only
appear further along the maturation process, or transcriptomics do not accurately reflect the
molecular identity of these neurons.[67-69] To further restrict and direct subtype specification
during PB-AD induction, additional patterning with small molecules, which was recently
developed for safe harbor-AD induction, could potentially be employed.[70]

Ngn2-induction produced cultures with a mixed glutamatergic and GABAergic
molecular identity, including canonical GABAergic markers such as SCGN, PVALB, CALB2.
Although these markers are sometimes found in certain glutamatergic neurons in the brain[71-
76], it is more likely that their expression is a result of nonspecific differentiation. Nevertheless,
virtually no VGAT puncta were identified and only glutamatergic ePSCs were recorded,
consistent with previous reports.[21,77-82]

Enhanced expression of presynaptic proteins during synaptic maturation
Our multilevel analysis of GABAergic cultures at different time points between 4 and 10 WPI,
combining patch-clamp electrophysiology, immunocytochemistry, and mass spec proteomics,
revealed the following insights about synapse maturation. Firstly, synaptic strength increased
approximately 3-fold during this period. A comparable increase in mPSC frequency and RRP
suggests that this is due to an increase in the strength and/or number of presynaptic
projections, which is further supported by VGAT immunocytochemistry. This developmental

trajectory is similar to what has been described for directed-differentiation and safe harbor
AD GABAergic autaptic neurons.[33,79]

Secondly, we observed a stark difference between pre- and postsynaptic molecular
differentiation. Strong upregulation of *AZ-, SV-* and *presynaptic membrane* proteins occurred,
while expression changes in proteins in postsynaptic gene sets were minimal. Asymmetric
proteomic development of the pre- and postsynapse was also observed in NGN2 neurons
and has been described for iPSC-derived neurons obtained through directed
differentiation.[83,84] In PB-AD neurons, current responses to GABA puffs remained constant
over time, indicating that functional postsynapses already exist at 4 WPI and the expression
of GABA receptors remains unaltered at subsequent weeks. This fits with the notion that
expression of postsynaptic neurotransmitter receptors precedes presynaptic innervation.[85-89]
Release from presynaptic projections subsequently induces clustering of postsynaptic
molecules, initiating postsynaptic maturation.[85,90-93] Indeed, postsynaptic receptor clustering
was previously predicted *in silico* to increase mPSC amplitude 1.5-fold, similar to what we
found *in vitro*[94].

Thirdly, although proteomic changes plateau after 8 WPI, both synapse number and
synaptic strength continue to increase. Possibly, the *AZ, SV* and *presynaptic membrane* proteins
that are upregulated until 8 WPI are sufficient to support synapse formation thereafter.
Structural rearrangement and posttranslational modification of synaptic proteins, mediated
by e.g. ERK, PKC, PKA, CaMKII and/or RhoA/ROCK signalling[95-109], could enhance synaptic
strength without proteomic changes. Such processes modulate many pre- and postsynaptic
proteins, including gephyrin[98-100,107,110], GABA-receptors[101-105,108-110], synapsins[96,106] and
microtubules and their associated proteins.[110-115] Additionally, synaptic strength could be
regulated by (trans-synaptic) alignment of functionally related proteins in nanodomains and
nanocolumns, enhancing their cooperative interactions.[94,116-122] Of the few proteins that were
upregulated – or for the first time detected – in PB-AD neurons at 10 WPI (Supplementary
table 1), we found several cell adhesion molecules (CAMs) (CNTNAP4, SLITRK1, CDH1, EPHA7,
ITGB3, NPTX2, ITGA2, ELFN1, IGSF8 and CNTNAP1).[3,123-131], extra cellular matrix (ECM)
proteins (TNC and RELN)[132,133], and proteins involved in receptor clustering (NPTX2, SLITRK1,
PRRT1, ELFN1)[134-137] Some of which are known to strengthen GABAergic transmission.[123,126]
To summarize, during early development, synaptic maturation and strengthening in iPSC-
derived neurons coincides with increased expression of presynaptic proteins compared to
their postsynaptic counterparts. At later stages, further strengthening may result from
alignment and reorganization of the pre- and postsynaptic compartments.

A versatile in vitro model for studying the role of inhibition in brain disease
Dysfunction of inhibitory neurons and E/I disbalance are common aspect of
neurodevelopmental and neurodegenerative disorders.[5,138-143] Given the pure GABAergic
nature of cultures produced with the PB-AD protocol, neurons generated with this method
provide a versatile model to study the role of inhibition and E/I disbalance in these disorders
using a human, and patient-own, genetic background. Compared to a safe harbor approach,
our method is directly applicable to patient derived iPSCs, which makes it time and cost
efficient, and prevents the introduction of clonal gene editing effects. The reliable production
of GABAergic autaptic cultures allows for a detailed assessment of interneuron synaptic
properties, such as priming, fusion and synaptic plasticity in patient cell lines.[144-146]
Application to disease-lines will be important steps to substantiate the full potential of this
method. To this end, PB-AD neurons can be mixed with NGN2 neurons at defined ratios,
enabling accurate modelling of E/I networks at any ratio, including those found in the human
brain.[36-38]
421
422 **Methods**
423
Extended methods are available in the supplementary material.

IPSC culture and delivery of transgenes
IPSC's were maintained on Matrigel- or Geltrex coated-dishes in Essential 8 medium. For viral
induction, viral particles encoding rTTa, Ascl1, and Dlx2 were added to IPSCs and antibiotic
selection was performed 24 hours after transfection. For PiggyBac based strategies, IPSCs
were seeded as single cells after which transfection with PiggyBac transposon and
transposase plasmids was performed using Lipofectamine STEM. Selection was performed
24 hours post-transfection.

Neuronal induction
Transgene-selected IPSCs were seeded in N2 medium with doxycycline to induce transgene
expression. After 2 days of transgene expression, selection for postmitotic cells was
performed with antimitotic agents FUDR or AraC, and cells were collected and replated on
rat astrocytes in neurobasal medium supplemented with neurotrophic factors, in which
neurons were maintained.

Electrophysiology

Whole-cell voltage clamp was performed to measure synaptic release. For GABA puffs,
perforated patch was performed, maintaining cell intrinsic chloride levels. GABAergic and
glutamatergic currents were blocked with picrotoxin and DP-V + DNQX, respectively.

Immunocytochemistry and morphological analysis
Methanol- or PFA-fixed neurons were stained for MAP2, VGAT, VGLUT1/2, NEUN and
Parvalbumin. After confocal microscopy analysis was performed in SynD[147] and ImageJ.

Statistics of electrophysiological and morphological data
Statistical testing was performed in Matlab and GraphPad Prism. Developmental trajectories
were tested using nonparametric Spearman correlations. For timepoint comparisons,
Kruskal-Wallis ANOVAs with Dunn's correction were performed. GABA puff responses were
fitted using a first-order polynomial linear fit and an extra sum-of-squares F-test was
employed.

Mass spectrometry
Cells were scraped, pelleted, and stored at -80°C. Sample were prepared using adapted DNA
micro spin column suspension trapping protocol.[148] Pellets were extracted in SDS buffer and
proteins were isolated and trypsin-digested. Peptides were eluted from the column and
samples were analysed using liquid chromatography MS.

Spectra were mapped to *in silico* spectral library based on the uniprot human
reference proteome peptides using DIA-NN 1.8.1.[149] Protein abundances were quantified
using differential abundance analysis (DAA) of MS-DAP 1.0.5.[150] Proteins (nearly) exclusive to
one group were analysed using MS-DAP differential detection (DD) test. The DEqMS[151]
algorithm was used for statistical analysis and resulting p-values were adjusted using
Benjamini–Hochberg False Discovery Rate (FDR) adjustment. DAA and DD results were
merged. Gene set enrichment analyses were performed using GOAT online v1.0.2.[152] and
SynGO knowledgebase v1.2.[153]

Cell type-associated proteomic changes were identified by expression weighted cell
type enrichment (EWCE) analysis[41], based on snRNA-sequencing data derived from human
cortex.[40] For cell population classifications we used cell class (e.g., GABAergic neurons) and
subclass (e.g. SST[+] interneurons). Cell type-specificity scores (CTSS) for each gene were used
to assign cell type labels.
469
470 **Acknowledgements**

The authors thank Robbert Zalm and Ingrid Saarloos for their help with cloning and plasmid production, Lisa Laan and Desiree Schut for preparation of glia feeder plates, Judith Huijgen and Solange Lopes Cardozo for preparation and maintenance of iPSCs, and Berna Özer and Remco Klaassen for performing mass-spectrometry proteomics.

This work was supported by the following grants (ZonMW-TOP BECAUSE (91216064), to M.V and L.N.C. ZonMW-PSIDER Brainmodel (10250022110003) to M.V and L.N.C., NWO-Dutch Research Agenda (NWA) NewTDec (1160.18.200) to M.V and L.N.C.), NWO Gravitation program grant BRAINSCAPES (NWO 024.004.012) to M.V. and ZonMW-OPEN (9120232310098) from the Netherlands Organisation for Scientific Research (NWO) to M.V. and L.N.C.

Declaration of interest

The authors declare no competing interests.

Author contributions

M.E.W., and J.M.C.M. designed and produced the PiggyBac construct

T.W.v.V, M.A.v.B., K.I.M., R.F.T., A.B.S., R.E.v.K, M.V. and L.N.C. designed the experiments

T.W.v.V, M.A.v.B. and J.S. performed and maintained neuronal inductions and cultures

T.W.v.V M.A.v.B. performed the patch-clamp experiments with contributions of J.S. to Fig. 1 and C.F. to Fig. S2

T.W.v.V, M.A.v.B.,. performed the immunocytochemistry experiments with contributions of J.S to Fig. 1

K.I.M., and F.K. performed the proteomics experiments and analysis

T.W.V.V, M.A.v.B., K.I.M., R.E.v.K, M.V. and L.N.C. designed figures and wrote the manuscript with input from all authors.

Supplementary information

Supplementary information will be made available with the manuscript.

Extended Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		

MAP2	Merck Millipore	(Millipore Cat# AB5543, RRID:AB_571049)
VGAT	Synaptic Systems	(Synaptic Systems Cat# 131 002, RRID:AB_887871)
VGLUT1	Antibodies Inc.	(Antibodies Incorporated Cat# 75-066, RRID:AB_2187693)
VGLUT2	Antibodies Inc.	(Antibodies Incorporated Cat# 75-067, RRID:AB_2239153)
NeuN	Merck Millipore	(Millipore Cat# ABN78, RRID:AB_10807945)
Parvalbumin	Synaptic Systems	(Synaptic Systems Cat# 195 004, RRID:AB_2156476)
Bacterial and virus strains		
FUW-TetO-*Ascl1*-T2A-puro	Addgene	RRID:Addgene_97329
FUW-TetO-*Dlx2*-IRES-hygro	Addgene	RRID:Addgene_97330
FUW-M2rtTA	Addgene	RRID:Addgene_20342
Deposited data		
PRIDE dataset " One-step induction of human GABAergic neurons promotes presynaptic development & synapse maturation"	This study	PXD064163
Transcriptomic cytoarchitecture reveals principles of human neocortex organization	Jorstad, et al.[40]	RRID:SCR_016152
Experimental models: Cell lines		
BiONI010-C-10	EBiSC	RRID:CVCL_RM84
BioNi010-C-13	EBiSC	RRID:CVCL_RF90
Kolf2.1J	Jackson Laboratory	RRID:CVCL_B5P3
GM23973	Coriell	RRID:CVCL_BW42
Recombinant DNA		
PB-TO-ASCL1-DLX2	This study	RRID:Addgene_182307
Transposase	This study	N/A
Software and algorithms		

DIA-NN 1.8.1	Demichev, et al (2020,2022)[154,155]	
MS-DAP 1.0.5	Koopmans, et al. 2023[150]	
DEqMS	Zhu, et al, 2020[151]	
GOAT	Miedema, et al. 2022[152]	
SynGO	Koopmans, et al. 2019[153]	

iPSCs origin and culture

Unless stated otherwise, the BiONi010-C-10 stem cell line (EBISC, RRID:CVCL_RM84) was

used for all experiments in this study. For glutamatergic induction, a gene-edited variant of

the same line containing an integrated NGN2-casette was used (EBISC, RRID:CVCL_RF90). As

a genetically independent control the KOLF2.1J (JAX, JiPSC1000, RRID:CVCL_B5P3) and

GM23973 (Coriell, RRID:CVCL_BW42) lines were used. All iPSCs were maintained on Matrigel

(Cornig) or Geltrex (Fisher Scientific)-coated dishes in Essential 8 (Gibco) or Essential 8 Flex

(Gibco) medium and passaged with EDTA and replated in the culture medium supplemented

with ROCK inhibitor Y27632 (5 μM, Tetu bio).

Lentiviral infection

Ultrahigh titre lentiviral particles were generated by Alstem (Richmond, CA, USA) from FUW-

M2rtTA (Addgene, #20342, RRID:Addgene_20342), FUW-TetO-*Ascl1*-T2A-puro (Addgene

#97329, RRID:Addgene_97329), and FUW-TetO-*Dlx2* -IRES-hygro (Addgene #97330,

RRID:Addgene_97330). For *Ascl1*+Forskolin, iPSCs were infected with only FUW-M2rtTA and

TetO-*Ascl1*-Puro, for *Ascl1* + *Dlx2* induction, iPSCs were infected with all three lentiviruses.

For both lentiviral induction methods, 24 hours after infection, puromycin (1 μg/ml) selection

was performed. For lentiviral *Ascl1* + *Dlx2* induction, additional hygromycin (75 μg/ml)

selection was performed. For both selection regimes an uninfected control well was included

to confirm successful selection before proceeding towards induction.

PiggyBac transfection

A more detailed protocol is found at the end of this document, see protocol 1. To prepare for

PiggyBac transfection, iPSCs were released with accutase and plated as single cells on a

Matrigel-coated 6-well plate in E8 + ROCK Inhibitor Y27632 at a density of 1500k per well.

Cells were placed in the incubator to adhere for 1 to 2 hours. A total of 3 μg of DNA containing

the *Ascl1 + Dlx2* PiggyBac transposon (Presented here; available on Addgene, 182307,
RRID:Addgene_182307) and transposase constructs (extended data 1) at a 1:3 molar ratio
was prepared in 100 µL of opti-MEM (GIBCO, 31985062). To another 100 µL op opti-MEM 5
µL of Lipofectamine STEM (Invitrogen, STEM00001) was added. Opti-MEM preparations were
quickly mixed by pipetting, after which the mixture was incubated at RT for 15 to 30 minutes
during which cells received fresh E8 + ROCK Inhibitor Y27632 to wash away non-adherent
cells. The opti-MEM solution was then added to the cells in a dropwise fashion. 24 hours
post transfection, selection with puromycin (1 µg/ml) was initiated to select only for cells
containing the transposon, and selection was repeated every 3-7 days.

PB-AD GABAergic neuronal induction
A more detailed protocol is found at the end of this document, see protocol 2. PB-AD positive
iPSCs were released with Accutase (Merck) on day post induction (DPI) 0 and replated with
at an approximate density of 3.500 cells/cm^2 in N2 medium (DMEM/F12 medium, Life
technologies) containing Glutamax (200 mM, Life Technologies), dextrose (0.3%, Life
Technologies) N2 supplement B (StemCell Technologies), Pen/Strep (0.1%, Gibco), ROCK
inhibitor Y27632 (10 µM, Tetubio) and doxycycline hyclate (2 µg/mL) onto Matrigel (Cornig)
or Geltrex (Fisher Scientific) coated dishes. On DPI 1, medium was refreshed using the same
composition without ROCK inhibitor Y27632. On DPI 2, the medium was refreshed once more
with the same composition as on DPI 1, with the addition of 5-Fluoro-2'-deoxyuridine (FUDR)
(10 µM, Sigma-Aldrich) or Cytosine β-D-arabinofuranoside (AraC, 20 µM, Sigma-Aldrich) to
remove proliferative cells. On DPI 3, neurons were released with accutase, pelleted, and
resuspended and plated in Neurobasal medium (Life Technologies) supplemented with
Glutamax (200 mM), B27 (0.1%) Pen-Strep (0.1%), fetal bovine serum (0.5%), BDNF (10 ng/ml,
Peprotech), CNTF (10ng/ml, Peprotech), GDNF (10 ng/ml, Peprotech) and doxycycline hyclate
(2 µg/mL; Sigma-Aldrich). Neurons were plated to the required respective preparations (see
Autaptic culture and *Network culture*).

Glutamatergic neuronal induction
For glutamatergic induction, Bioni010-C-13 iPSCs were released with Accutase (Merck) on
DPI 0 and replated with at an approximate density of 3.500 cells/cm^2 in N2 medium
(DMEM/F12 medium, Life technologies) containing Glutamax (200 mM, Life Technologies),
dextrose (0.3%, Life Technologies) N2 supplement B (StemCell Technologies), Pen/Strep
(0.1%, Gibco) and doxycycline hyclate (2 µg/mL; Sigma-Aldrich) combined with dual-SMAD
inhibitors LDN193189 (100 nM, Stemgent), SB431542 (10 µM, Tocris Bioscience), and XAV939
(2 µM, Stemgent) NT3 (10 ng/ml, Peprotech) and doxycycline hyclate (2 µg/mL) onto Matrigel

(Cornig) or Geltrex (Fisher Scientific) coated dishes. On DPI 1, medium was refreshed using
the same composition without ROCK inhibitor Y27632. On DPI 2, the medium was refreshed
once more with the same composition as on day 2, with the addition of 5-Fluoro-2'-
deoxyuridine (FUDR) (10 μM, Sigma-Aldrich) or Cytosine β-D-arabinofuranoside (AraC, 20 μM,
Sigma-Aldrich) to remove proliferative cells. On DPI 3, neurons were released with accutase,
pelleted, resuspended and plated in Neurobasal medium (Life Technologies) supplemented
with Glutamax (200 mM), B27 (0.1%) Pen-Strep (0.1%), foetal bovine serum (0.5%), BDNF (10
ng/ml, Peprotech), CNTF (10ng/ml, Peprotech), GDNF (10 ng/ml, Peprotech) and doxycycline
hyclate (2 μg/mL; Sigma-Aldrich). Neurons were plated to the required respective
preparations (see *Autaptic culture* and *Network culture*).

Lentiviral GABAergic induction
In general, lentiviral GABAergic induction protocols followed a similar workflow as PB-AD
induction. To better adhere to established protocols.[14,15], the medium composition differed at
certain steps (Figure 1H). For Ascl1 + Forskolin induction, from DPI 0 onwards, Forskolin (10
μM, Merck) was added to the medium up to DPI 14. Furthermore, from DPI0 onwards non-
essential amino acids (1:100, Sigma) were added up to day DPI 14. BDNF (10 ng/ml) and NT3
(10 ng/ml) were present from DPI1 onwards. For lentiviral *Ascl1 + Dlx2* induction BDNF (10
ng/ml) and NT3 (10 ng/ml) were added from DPI 0 onward, as well. Furthermore, laminin
(1μg/ml, Sigma-Aldrich), cyclic AMP (1 μM, Sigma-Aldrich), and Ascorbic Acid (200 μM, Sigma-
Aldrich), were added to the medium at all subsequent refreshments to improve viability.

Autaptic culture
Neurons were plated at a density of ~ 400 cells / cm^2 on micro-islands prepared from rat
glial cells plated on agarose-coated coverslips stamped with a custom stamp in Poly-D-
Lysine (0.1 mg/ml, Sigma), rat tail collagen (0.7 mg/ml, Corning) and Fibronectin (100 μg/ml,
Corning) in acetic acid (10 mM, sigma). Glial islands were prepared approximately 7 days
before plating neurons and were treated with AraC (20 μM, Sigma-Aldrich) before plating
neurons. Autaptic cultures were refreshed every 7 days with a ~50% media refreshment, with
the same medium as used for plating neurons, but omitting doxycycline hyclate (except for
Figure 1E-G, where doxycycline hyclate was added up to DPI 14).

Network culture
Neurons were plated at a density of ~ 7000 cells / cm^2 on coverslips covered with a glial-
monolayer prepared from rat astrocytes a week in advance and treated with Cytosine β-D-
arabinofuranoside (20 μM, Sigma-Aldrich) 24-48 hours before plating neurons. Networks

cultures were refreshed with a ~50% media refreshment every 3-4 days, with the same
medium as used for plating neurons, but omitting doxycycline hyclate.

Electrophysiology
Whole-cell voltage-clamp recordings ($V_m = -70$ mV) were performed on autaptic neurons,
with borosilicate glass pipettes ($2-5M\Omega$) filled with (in mM) 136 KCl, 17.8 HEPES, 1 EGTA,
0.6 $MgCl_2$, 4 ATP-Mg, 0.3 GTP-Na, 12 phosphocreatine-K_2 and 50 units/mL phosphocreatine
kinase (pH = 7.30, 290 mOsmol). The external solution contained in mM: 10 HEPES, 10
glucose, 140 NaCl, 2.4 KCl, 4 $MgCl_2$, and 4 $CaCl_2$ (pH = 7.30, 290 mOsmol). Experiments were
performed at room temperature (~22°C using a MultiClamp700B amplifier (Axon
Instruments) and Digidata 1440A digitizer under control of Clampex 10 software (Molecular
devices). Only cells with an access resistance < 15MΩ (80% compensated) and leak current
of <300 pA were included. Synaptic responses were elicited by 0.5 ms depolarization to
30 mV.
For GABA puffs, perforated patch recordings were produced by filling the patch pipette
(in mM) 140 KCl, 10 HEPES, gramicidin (100 µg/ml) and allowing cells to perforate for 15-30
minutes after reaching a GΩ seal to < 100 MΩ series resistance. Puff pipettes filled with 100
µM GABA in ACSF were positioned near the soma, at approximately 25 µm distance. 100 ms
puffs were applied at -110, -90, -70, -50, -30 and -10 mV holding voltages to induce GABAergic
currents. Cells were discarded if a sudden swift decrease in resistance was observed during
perforation, or if the observed reversal potential of Cl$^-$ was ± 0 mV, which indicates exchange
of the patch-pipette solution with the intracellular solution.

Electrophysiology; pharmacology
To Block GABA-currents, picrotoxin (100 µM, Hello Bio) was used. A 100 mM stock was
prepared in DMSO and was added to the bath to reach the final concentration. To block
glutamatergic currents, AP-V (100 µM, R&D systems) and DNQX (40 µM, Tocris) were used.
For DNQX, A 1000x of stock of each was prepared in DMSO. For AP-V, the stock was prepared
in water.

Electrophysiology; analysis electrophysiological data.
Offline analysis was performed using custom-written programs[156] in Matlab (Mathworks).
Experimental conditions were blinded during analysis. In all figures, stimulation artefacts
have been removed. For evoked release during 20 Hz stimulation, back-extrapolation of
linear fits from the last 20 pulses of the cumulative charge was used to estimate the RRP (y-
intercept).[157]

 Immunocytochemistry and morphological analysis

Neurons were fixed in ice cold 100% methanol (VWR) for 15 minutes at 4°C, blocked with

2.5% normal goat serum (NGS, Life Technologies) and 0.1% Triton X-100 in PBS for

45 minutes and incubated with primary antibodies in PBS with 2.5% NGS and 0.1% Triton X-

100 overnight at 4°C. The following primary antibodies were used: chicken anti-MAP2

(1:1000, Merck AB5543, RRID:AB_571049); rabbit anti-VGAT (1:2000, SySy #131 002,

RRID:AB_887871), mouse anti-VGlut1 (1:250, NeuroMab clone N28/9, Antibodies inc. 75-066,

RRID:AB_2187693), mouse anti-VGlut2 (1:250, NeuroMab clone N29/29, Antibodies inc. 75-

067, RRID:AB_2239153). For Rabbit anti-NeuN (1:500, Merck ABN78, RRID:AB_10807945) and

anti-Parvalbumin (1:500, SySy #195 004, RRID:AB_2156476) staining neurons were fixed

with 3.7% PFA for 15 minutes at room temperature, instead of methanol fixation. After

washing with PBS the next day, cells were incubated for 1 h at room temperature with

secondary antibodies conjugated to Alexa dyes (1:1000, Invitrogen Molecular Probes),

washed again and mounted with DABCO-Mowiol (Invitrogen). Images were acquired at

1024x1024 pixels on a Nikon Eclipse Ti confocal laser scanning microscope (40× objective;

NA 1.3) with NIS-Elements software. Z-stacks were produced at a 0.5 µm step-size until no

more puncta and/or dendrites were in focus. Confocal settings were kept constant for all

scans within an experimental week. Analysis of neuronal morphology was performed on

maximum-intensity projections using the automated image analysis routine SynD.[147] for

autaptic neurons and a custom ImageJ workflows for network analysis, in either case, for

VGAT and VGLUT puncta counts, only puncta on dendrites were counted.

Statistics of electrophysiological and morphological data

Statistical testing was performed in MATLAB and GraphPad Prism For descriptive statistics

(Figure 1) the likelihood of normality vs lognormality was calculated, which revealed that

the data was lognormally disturbed. Therefore the geometric mean and geometric SD factor

were calculated and reported. In autaptic development timelines (Figure 4&5), no data was

normally distributed. Therefore, to test general developmental trajectories, nonparametric

Spearman correlations were computed with a two-tailed test for significance and a 95%

confidence interval. For timepoint comparisons, Kruskal-Wallis ANOVAs were performed with

Dunn's multiple comparisons test. Pairwise comparisons were made between 4 and 6, 6 and

8, and 8 and 10 WPI. #, ##, ###, #### and *ns* represent results of spearman correlation results,

*, **, ***, **** and ns represent Kruskal-Wallis ANOVA results. *ns* and ns represent non-

significant results. Single, double, triple and quadruple marks represent (corrected) $p < 0.05$,

$p < 0.01$, $p < 0.001$ and $p < 0.0001$, respectively. GABA-puff responses (figure 6O) were fitted

using a first-order polynomial linear fit and an extra sum-of-squares F-test was used to
determine whether GABA responses were identical throughout development. In plots, unless
stated otherwise, the lines in boxplots represent the median value. Crosses represent the
mean value. Boxes represent 50% off the values. Circular markers represent individual
observations. Quantifications are available in the supplementary data, and each figure
contains n (number of cells) and N (number of inductions) values per parameter in either the
relevant graph or typical example.

Mass spectrometry; Cell pellet harvesting for MS
Cells were washed three times with 1 ml sterile Dulbecco's PBS (Gibco), whereof first twice
at 37°C and finally once at 4 °C. Cultures were kept on ice, scraped off using VWR cell lifters,
and harvested in 1.5 ml protein LoBind Eppendorf tubes (VWR). Pellets were obtained by
centrifugation at 1,000 × g for 5 minutes at 4 °C. Supernatant was removed and pellets were
kept at -80 °C before subsequent sample preparation.

Mass spectrometry; MS sample preparation
MS sample preparation was performed following the DNA micro spin column suspension
trapping protocol[148], with some minor adaptations. Cell pellets were extracted and reduced
in 6% SDS buffer, containing 50 mM Tris-HCl (pH 8) and 5 mM tris (2-carboxyethyl)phosphine
(TCEP), by incubation at 55°C for 15 minutes in a thermomixer set to 1400 RPM. Free
sulfhydryl groups were alkylated by incubation with 20 mM methyl methanethiosulfonate
(MMTS; 200 mM stock solution) for 15 minutes at RT. Insoluble debris was cleared by
centrifugation at 10000g for 1 minute at RT and the supernatant was transferred to a new
tube (Eppendorf). Next, samples were acidified by addition of 1.1% phosphoric acid (12%
stock solution), mixed with six volumes of binding/washing buffer (90% methanol and
100mM Tris-HCl, pH 8) and loaded onto a plasmid DNA micro column (HiPure from Magen
Biotechnology). The protein particulate was retained on the column upon centrifugation at
1400x g for 1 minute and the columns were washed four times with binding/washing buffer.
Columns were transferred to a new tube (Eppendorf), supplemented with 500 ng Trypsin/Lys-
C (Promega) in 50 mM NH4HCO3 and incubated overnight at 37°C in a humidified incubator.
Tryptic peptides were eluted and pooled by subsequent addition of 50 mM NH4HCO3, 0.1%
formic acid and 0.1% formic acid in acetonitrile. Collected peptides were dried by SpeedVac
and stored at -80°C.

Mass spectrometry; LC-MS Analysis

Each sample of tryptic digest was redissolved in 0.1% formic acid and the peptide
concentration was determined by tryptophan-fluorescence assay[158]; 75 ng of peptide was
loaded onto an Evotip Pure (Evosep). Peptide samples were separated by standardized 30
samples per day method on the Evosep One liquid chromatography system, using a 15 cm ×
150 μm reverse-phase column packed with 1.5 μm C18-beads (EV1137 from Evosep)
connected to a 20 μm ID ZDV emitter (Bruker Daltonics). Peptides were electro-sprayed into
the timsTOF Pro 2 mass spectrometer (Bruker Daltonics) equipped with CaptiveSpray source
and measured with the following settings: Scan range 100-1700 m/z, ion mobility 0.65 to
1.5 Vs/cm2, ramp time 100 ms, accumulation time 100 ms, and collision energy decreasing
linearly with inverse ion mobility from 59 eV at 1.6 Vs/cm2 to 20 eV at 0.6 Vs/cm2. Operating
in dia-PASEF mode, each cycle took 1.38 s and consisted of 1 MS1 full scan and 12 dia-PASEF
scans. Each dia-PASEF scan contained two isolation windows, in total covering 300-1200 m/z
and ion mobility 0.65 to 1.50 Vs/cm2. Dia-PASEF window placement was optimized using the
py-diAID tool[159] Ion mobility was auto calibrated at the start of each sample (calibrant m/z,
1/K0: 622.029, 0.992 Vs/cm2; 922.010, 1.199 Vs/cm2; 1221.991, 1.393 Vs/cm2).

Mass spectrometry; MS data Analysis
DIA-PASEF raw data were processed with DIA-NN 1.8.1.[154,155] An in-silico spectral library was
generated from the uniprot human reference proteome (SwissProt and TrEMBL, canonical
and additional isoforms, release 2023-2) using Trypsin/P digestion and at most 1 missed
cleavage. Fixed modification was set to beta-methylthiolation (C) and variable modifications
were oxidation (M) and N-term M excision (at
most 1 per peptide). Peptide length was set to 7-30, precursor charge range was set to 2-4,
precursor m/z was limited to 280–1220, both MS1 and MS2 mass accuracy were set to 15
ppm, scan window was set to 9, double-pass-mode and match-between-runs were enabled.
Protein identifiers (isoforms) were used for protein inference. All other settings were left as
default.
MS-DAP 1.0.5.[150] was used for downstream differential abundance analyses using the
DIA-NN results. Filtering and normalization was applied to respective samples per statistical
contrast. Peptide-level filtering was configured to retain only peptides identified in at least
4 samples in both experimental conditions (within current contrast) and only proteins with
2 valid peptides were retained. Next, peptide abundance values were normalized by
minimizing the mode $\log_2$ (FC) between sample pairs (MWMB algorithm), followed by
protein-level mode-between normalization. The DEqMS[151] algorithm was used for statistical
analysis and resulting p-values were adjusted for multiple testing using the Benjamini–

Hochberg False Discovery Rate (FDR) procedure. To investigate proteins with strong
differences in detection rates between conditions, e.g. consistently detected in one condition
but (near) absent in another, the MS-DAP differential detection test was applied to all
proteins that were observed with at least 2 peptides in at least 4 samples in either condition
(i.e. confidently detected proteins). Using the default MS-DAP workflow, we merged
differential detection (DD) results into the DEqMS result table for the subset of proteins that
had both an absolute DD z-score of at least 6 and a smaller adjusted p-value in DD (or no p-
value from DEqMS due to a lack of data points in either condition).

Mass spectrometry; Expression-weighted cell type enrichment analysis
Cell type enrichment analysis gives insight into the cell types that potentially drive
differential expression. EWCE uses single-cell RNA-sequencing data to generate a probability
distribution associated with a gene list having an average level of expression within a cell[41]
Although it's based on RNA expression levels, it can also be applied to proteomics data.[160]
We used published single nucleus RNA-sequencing data (49,495 nuclei) from human cortical
brain regions – including middle temporal gyrus, anterior cingulate cortex, primary visual
cortex, primary motor cortex, primary somatosensory cortex and primary auditory cortex –
published by Jorstad et al.[40] For cell type annotations we used the cell type annotations: class
(e.g., GABAergic neurons) and subclass (e.g. SST[+] interneurons) for first and second level
EWCE analysis, respectively. Bootstrap enrichment tests were performed using 10,000 reps
to look whether gene sets (i.e., foreground) were significantly enriched for a cell class or
subclass relative to the background (i.e., all detected proteins included in that contrast). Since
we focused on neuronally-enriched proteins or SYNGO-annotated synaptic proteins, we
excluded non-neuronal enrichment scores.

EWCE calculates cell type-specificity scores (CTSS) for all genes, which is the
percentage of mean expression per cell type relative to the sum of the mean expressions of
all cell types. We used the CTSS to predict for each protein the cell type the protein probably
is derived from. We only included CTSS from genes that passed mRNA detection threshold
(mean expression >4 in at least one cell class). Cell type predictions included neuronal
(neuronal CTSS > 70%), GABAergic neurons (GABA CTSS > 0.8 and glutamatergic CTSS < 0.1),
glutamatergic neurons (glutamatergic CTSS > 0.8 and GABAergic CTSS < 0.1). Cell type
predictions were labelled in volcano plots. We also used the EWCE-calculated mean mRNA
expression for each cell type to estimate the differential expression of genes in GABAergic
vs glutamatergic neurons, which was used in the SYNGO manual colour plots (Figure 6H)).

Mass spectrometry; Heatmaps and line plots

We used the mean $\log_2$-transformed normalized protein intensities per group (i.e., PBAD 4
WPI) and plotted these values using line plots in RStudio.[161] using ggplot[162] The relative
normalized protein intensities ($\log_2$ (FC)) was used in heatmaps, which were produced using
Morpheus. To map these proteins to predicted cell types we also show EWCE-calculated cell
type specificity scores. All heatmaps visualizations are with fixed colour scaling (i.e., not
relative colour scaling per gene), as indicated by heatmap legend.

Mass spectrometry; Gene-set enrichment analysis using GOAT and over-representation analysis
using SYNGO
We used the output of the differential abundance analysis using MSDAP (PB-AD 10 vs 4 WPI
and NGN2 10 vs 4 WPI) as input for gene-set enrichment analysis using GOAT. The GOAT
online v1.0.2.[152] was used to perform gene set enrichment analyses using gene sets from the
SynGO knowledgebase v1.2[153] Effect size-derived gene scores (i.e., $\log_2$ (FC)) were used to
test for enriched gene sets that contained at least 10 and at most 1500 genes (or 50% of the
gene list, whichever was smaller) that overlapped with the input gene list. Multiple testing
correction was independently applied per gene set "source" (i.e. SYNGO_CC, SYNGO_BP) using
Bonferroni adjustment and all p-values were again adjusted using Bonferroni correction to
account for 2 separate tests across "sources". The significance threshold for adjusted p-values
was set to 0.05. SYNGO over-representation analysis was performed using the same MSDAP
output. Foreground was defined as upregulated proteins per contrast (p-adjusted < 0.05 and
$\log_2$ (FC) > 0.1), and the background contained all tested genes for that contrast. SynGO
release is dataset version: 20231201, and default settings were used. Ontologies are
significantly enriched at 1% FDR, while including terms containing at least three matching
input genes.

Protocol 1: PiggyBac transfection
Introduction
In this protocol it is described how IPSCs are transfected with the transposon and
transposase to allow for neuronal induction of PB-AD neurons.
Materials
• Geltrex (Cornig) or Matrigel (Fisher Scientific)
• Essential 8 (Gibco) or Essential 8 Flex (Gibco)
• Accutase (Millipore; SCR005)
• DMEM-F12 (Life technologies)
• opti-MEM (GIBCO, 31985062)
• Lipofectamine STEM (Invitrogen, STEM00001)

- 794 • Rock Inhibitor Y27632 (Tetu-Bio, T1725) or CEPT cocktail (chroman 1 (Mechem
- 795 Expresss, HY-15392), emricasan (Selleckchem, S7775), polyamine (Sigma, P8483),
- 796 and trans-ISRIB (Tocris, 5284))
- 797 • Doxycycline hyclate 2 µg/mL (Sigma, D9891)
- 798 • Puromycin (Sanbio, 13884-25)
- 799 • Piggybac transposase (see supplementary plasmid map)
- 800   • *Distribution of the transposase through Addgene is not possible. We advise*
- 801   *synthesis inhouse or commercially using the supplied plasmid map.*
- 802 • *Ascl1 + Dlx2* PiggyBac transposon (Addgene, 182307, RRID:Addgene_182307)

803 *Protocol*

- 804 • Coat your plates with Geltrex or Matrigel according to manufacturer instructions.
- 805 Culture IPSCs in 1x6w until 70-90% confluent.
- 806 • Prepare a fresh coated 1x6w for plating infected cells. Prewarm Essential 8 or
- 807 Essential 8 Flex + Rock Inhibitor Y27632 (RI, 10 µm) or CEPT (50 nM, 5 µM, 1x, 07.
- 808 µM respectively).
- 809   • *Some IPSC lines are more easily transfected then others. If severe cell-death or*
- 810   *bad transfection efficiency are observed, CEPT cocktail can improve this.*
- 811 • Aspirate the medium of the IPSCs and wash 1x with PBS.
- 812 • Aspirate medium and add 350 µl accutase to 1x6w IPSCs. Incubate at 37 °C For
- 813 approx. 5-10 minutes or until cells detach.
- 814 • Meanwhile, aspirate the coating medium from a new plate and add Essential 8 or
- 815 Essential 8 Flex + RI or CEPT.
- 816 • Once cells have detached, collect cells with 4.5 ml DMEM + RI and spin down 5
- 817 minutes at 1000 RPM / 180 RCF.
- 818 • Aspirate the supernatant and resuspend cells in 1 ml of Essential 8 or Essential 8
- 819 Flex + RI or CEPT cocktail.
- 820 • Count cells and plate 1000 to 1500k cells per 1x6w well, 500 to 750k cells per
- 821 1x12w or 200 to 300k per 1x24w.
- 822 • Let cells adhere for 1 to 2 hours at 37 °C/5% CO_2.
- 823 • Per transfection, prepare the mixes as stated in table 1. Give the Opti-MEM a few
- 824 minutes to warm to RT before adding the other components to each mix.

825 *Table: Mixing ratio for transfection of piggyBac constructs*

6 well	Mix 1	Mix 2	12 well	Mix 1	Mix 2	24 well	Mix 1	Mix 2
Opti-MEM	100 µl	100 µl	Opti-MEM	50 µl	50 µl	Opti-MEM	20 µl	20 µl
DNA	3 µg		DNA	1.5 µg		DNA	0.6 µl	
Lipofectamine stem		5 µl	Lipofectamine stem		2.5 µl	Lipofectamine stem		1 µl

- For DNA, mix the transposase and transposon at a molar ratio of 1:3.
- Combine mix 1 and 2 and mix well by pipetting up and down. Incubate transfection mix for 15 to 30 minutes at RT.
- Replace Essential 8 or Essential 8 Flex + RI or CEPT on the cells during this incubation step, to wash away any non-adherent cells and debris.
- Add the mixture to the cells in a dropwise fashion, evenly distributed throughout the well. Immediately swirl the plate to ensure even distribution, and return plate to 37 °C/5% CO_2.
- Over the following 24 to 72 hours, BFP expression should appear. Culture cells to 70-90% confluency or for 1 week before splitting if confluency isn't reached.
- Once cells are 50-70% confluent, add puromycin for selection. This selection regime can be started 48h after transfection, if cells divide quickly, or later. The required concentration can vary between cell-lines. For most lines, 1 µg/ml puromycin for a minimum of two days is recommended.
    - Take along a well of untransfected IPSCs of the same genotype as a control and only stop selection once no cells survive in this well.
- Expand selected cells and use as needed. Repeated antibiotic selection should be performed weekly to ensure stable integration of the transposon is maintained.

846   *Protocol 2: PB-AD GABAergic neuronal induction*

847   *Materials*

- Matrigel (Cornig), Geltrex (Fisher Scientific)
- Accutase (Millipore; SCR005)
- Rock Inhibitor Y27632 10 µM (Tetu-Bio, T1725) (RI)
- Doxycycline hyclate 2 µg/mL (Sigma, D9891) (Dox)
- FUDR (Sigma, F0503) 10 µM **or** AraC (Sigma, C1768) 20 µM
  - AraC more efficiently selects only postmitotic cells and is recommended, but if lentiviral infection after induction is needed FUDR selection allows for higher infection efficiency after selection.
- Neurotrophic Factors
  - BDNF (Stemcell Technologies; 78005.3) - 10 ng/ml
  - CNTF (Peprotech; 450-13) - 10 ng/ml
  - GDNF (Stemcell Technologies; 78058.3) - 10 ng/ml
- DMEM (VWR; 392-0415P)
  - foetal bovine serum (2%, Life Tech; 10270106)
- N2 Medium
  - DMEM/F12 medium (Life Technologies)
  - Glutamax (200 mM, Life Technologies)
  - dextrose (0.3%, Life Technologies)
  - N2 supplement B (StemCell Technologies)
  - Pen/Strep (0.1%, Gibco)
- Neurobasal Medium
  - Neurobasal medium (Life Technologies)
  - Dextrose (Life Tech; A2494001) – 0.3% final concentration – use 1:67
  - Glutamax (200 mM)
  - NEAA (Life Tech; 11350912) - 1:200
  - B27 (0.1%)
  - Pen-Strep (0.1%)
  - foetal bovine serum (Life Tech; 10270106, 0.5%)

Protocol

Day 0: Start induction

1. Per 1x6w IPSCs coat 1x 10cm dish with Geltrex or Matrigel according to manufacturer instructions. Prewarm 10 ml of N2 + RI + Dox, 5 ml of DMEM + 2% FBS per 1x6w to induce, and 5 ml PBS.

2. Remove media from IPSCs and wash 1x with warm PBS.

3. Add 300 µl accutase per 6-well.

4. Incubate for 5 minutes at 37°C.

5. Add 3 ml DMEM and pipet up and down to collect cells.

6. Pellet for 5 minutes at 1000 RPM / 180 RCF.

7. Resuspend pellet in 1 ml N2 medium with RI + Dox.

8. Plate cells as single-cells in a 10 cm dish with 9 ml N2 medium with RI + Dox.

Day 1: Repeat Doxycycline and remove RI
9. The following day, replace 100% of the medium with N2 + Dox

Day 2: Repeat Doxycycline and select postmitotic cells
10. The following day, replace 100% of the medium with N2 + Dox + FUDR **or** N2 + Dox
+ AraC

Day 3: Replate neurons to target vessel
11. Prepare Neurobasal medium + DOX + CNTF + GDNF + BDNF (if using FUDR, + FUDR)

12. When media is warm, transfer it to prepared plates. Induced GABAergic neurons
thrive in co-culture with glia. Neurons can be replated to coverslips pre-seeded with
glia, or be mixed with glia and plated simultaneously.

13. Wash induced neurons 1x with PBS.

14. Add accutase to your induced neurons (1.5 ml in 10 cm dish).

15. Incubate for 5 minutes at 37°C or until cells are released.

16. Add 3 ml of DMEM + 2% FBS and pipette up and down to collect cells.

17. For 1x10 cm dish, use 13 ml of medium for collection.

18. Pellet cells for 5 min at 1000 RPM / 180 RCF.

19. Resuspend the pellet in 1 ml NBM + Dox + BDNF + CNTF + GDNF and count cells.

20. Plate neurons at your required density.

Feeding
21. Feeding rates depend on experimental conditions. For the experiments as described
here, low density (autaptic) cultures are fed once a week. Higher density cultures are

fed twice a week. In all cases, a ~50% medium refreshment is performed. Media
composition is the same as the glia-transfer medium, but AraC/FUDR is omitted.

Material availability

PiggyBac *Ascl1-Dlx2* construct is available from Addgene [#182307](#). The mass spectrometry
proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE
partner repository.[163] with the dataset identifier PXD064163.

References

1.   Breschi, A., Gingeras, T.R., and Guigó, R. (2017). Comparative transcriptomics in human
and mouse. Nature Reviews Genetics *18*, 425-440. 10.1038/nrg.2017.19.

2.   Gharib, W.H., and Robinson-Rechavi, M. (2011). When orthologs diverge between human
and mouse. Briefings in Bioinformatics *12*, 436-441. 10.1093/bib/bbr031.

3.   Stachniak, T.J., Sylwestrak, E.L., Scheiffele, P., Hall, B.J., and Ghosh, A. (2019). Elfn1-
Induced Constitutive Activation of mGluR7 Determines Frequency-Dependent
Recruitment of Somatostatin Interneurons. J Neurosci *39*, 4461-4474.
10.1523/jneurosci.2276-18.2019.

4.   Rubenstein, J.L.R., and Merzenich, M.M. (2003). Model of autism: increased ratio of
excitation/inhibition in key neural systems. Genes, Brain and Behavior *2*, 255-267.
10.1034/j.1601-183x.2003.00037.x.

5.   Palop, J.J., Chin, J., Roberson, E.D., Wang, J., Thwin, M.T., Bien-Ly, N., Yoo, J., Ho, K.O., Yu,
G.Q., Kreitzer, A., et al. (2007). Aberrant excitatory neuronal activity and compensatory
remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease.
Neuron *55*, 697-711. 10.1016/j.neuron.2007.07.025.

6.   Busche, M.A., Eichhoff, G., Adelsberger, H., Abramowski, D., Wiederhold, K.-H., Haass, C.,
Staufenbiel, M., Konnerth, A., and Garaschuk, O. (2008). Clusters of Hyperactive Neurons
Near Amyloid Plaques in a Mouse Model of Alzheimer's Disease. Science *321*, 1686-
1689. doi:10.1126/science.1162844.

7.   Nelson, S., and Valakh, V. (2015). Excitatory/Inhibitory Balance and Circuit Homeostasis
in Autism Spectrum Disorders. Neuron *87*, 684-698. 10.1016/j.neuron.2015.07.033.

8.   Foss-Feig, J.H., Adkinson, B.D., Ji, J.L., Yang, G., Srihari, V.H., McPartland, J.C., Krystal, J.H.,
Murray, J.D., and Anticevic, A. (2017). Searching for Cross-Diagnostic Convergence: Neural
Mechanisms Governing Excitation and Inhibition Balance in Schizophrenia and Autism
Spectrum Disorders. Biol Psychiatry *81*, 848-861. 10.1016/j.biopsych.2017.03.005.

9. Sohal, V.S., and Rubenstein, J.L.R. (2019). Excitation-inhibition balance as a framework
for investigating mechanisms in neuropsychiatric disorders. Molecular Psychiatry *24*,
1248-1257. 10.1038/s41380-019-0426-0.

10. Nicholas, C.R., Chen, J., Tang, Y., Southwell, D.G., Chalmers, N., Vogt, D., Arnold, C.M., Chen,
Y.-J.J., Stanley, E.G., and Elefanty, A.G. (2013). Functional maturation of hPSC-derived
forebrain interneurons requires an extended timeline and mimics human neural
development. Cell stem cell *12*, 573-586.

11. Liu, Y., Liu, H., Sauvey, C., Yao, L., Zarnowska, E.D., and Zhang, S.-C. (2013). Directed
differentiation of forebrain GABA interneurons from human pluripotent stem cells.
Nature protocols *8*, 1670-1679.

12. Maroof, A.M., Keros, S., Tyson, J.A., Ying, S.-W., Ganat, Y.M., Merkle, F.T., Liu, B., Goulburn,
A., Stanley, E.G., and Elefanty, A.G. (2013). Directed differentiation and functional
maturation of cortical interneurons from human embryonic stem cells. Cell stem cell *12*,
559-572.

13. Engle, S.J., Blaha, L., and Kleiman, R.J. (2018). Best Practices for Translational Disease
Modeling Using Human iPSC-Derived Neurons. Neuron *100*, 783-797.
10.1016/j.neuron.2018.10.033.

14. Mossink, B., van Rhijn, J.-R., Wang, S., Linda, K., Vitale, M.R., Zöller, J.E.M., van Hugte, E.J.H.,
Bak, J., Verboven, A.H.A., Selten, M., et al. (2022). Cadherin-13 is a critical regulator of
GABAergic modulation in human stem-cell-derived neuronal networks. Molecular
Psychiatry *27*, 1-18. 10.1038/s41380-021-01117-x.

15. Yang, N., Chanda, S., Marro, S., Ng, Y.H., Janas, J.A., Haag, D., Ang, C.E., Tang, Y., Flores, Q.,
Mall, M., et al. (2017). Generation of pure GABAergic neurons by transcription factor
programming. Nat Methods *14*, 621-628. 10.1038/nmeth.4291.

16. Baum, C. (2007). Insertional mutagenesis in gene therapy and stem cell biology. Current
opinion in hematology *14*, 337-342.

17. Cabrera, A., Edelstein, H.I., Glykofrydis, F., Love, K.S., Palacios, S., Tycko, J., Zhang, M.,
Lensch, S., Shields, C.E., Livingston, M., et al. (2022). The sound of silence: Transgene
silencing in mammalian cell engineering. Cell Systems *13*, 950-973.
https://doi.org/10.1016/j.cels.2022.11.005.

18. Wolf, D., and Goff, S.P. (2009). Embryonic stem cells use ZFP809 to silence retroviral
DNAs. Nature *458*, 1201-1204. 10.1038/nature07844.

19. Yao, S., Sukonnik, T., Kean, T., Bharadwaj, R.R., Pasceri, P., and Ellis, J. (2004). Retrovirus
Silencing, Variegation, Extinction, and Memory Are Controlled by a Dynamic Interplay of

Multiple Epigenetic Modifications. Molecular Therapy *10*, 27-36.
10.1016/j.ymthe.2004.04.007.

20.   Miller, J.C., Tan, S., Qiao, G., Barlow, K.A., Wang, J., Xia, D.F., Meng, X., Paschon, D.E., Leung,
981   E., Hinkley, S.J., et al. (2011). A TALE nuclease architecture for efficient genome editing.
982   Nature Biotechnology *29*, 143-148. 10.1038/nbt.1755.

983   21.   Peitz, M., Krutenko, T., and Brüstle, O. (2020). Protocol for the Standardized Generation
984   of Forward Programmed Cryopreservable Excitatory and Inhibitory Forebrain Neurons.
985   STAR Protoc *1*, 100038. 10.1016/j.xpro.2020.100038.

986   22.   Schmid, B., Holst, B., Poulsen, U., Jørring, I., Clausen, C., Rasmussen, M., Mau-Holzmann,
987   U.A., Steeg, R., Nuthall, H., Ebneth, A., and Cabrera-Socorro, A. (2021). Generation of two
988   gene edited iPSC-lines carrying a DOX-inducible NGN2 expression cassette with and
989   without GFP in the AAVS1 locus. Stem Cell Research *52*, 102240.
990   https://doi.org/10.1016/j.scr.2021.102240.

991   23.   Lombardo, A., Cesana, D., Genovese, P., Di Stefano, B., Provasi, E., Colombo, D.F., Neri, M.,
992   Magnani, Z., Cantore, A., Lo Riso, P., et al. (2011). Site-specific integration and tailoring
993   of cassette design for sustainable gene transfer. Nature Methods *8*, 861-869.
994   10.1038/nmeth.1674.

995   24.   Pi, H.-J., Hangya, B., Kvitsiani, D., Sanders, J.I., Huang, Z.J., and Kepecs, A. (2013). Cortical
996   interneurons that specialize in disinhibitory control. Nature *503*, 521-524.
997   10.1038/nature12676.

998   25.   Wilber, A., Linehan, J.L., Tian, X., Woll, P.S., Morris, J.K., Belur, L.R., McIvor, R.S., and
999   Kaufman, D.S. (2007). Efficient and stable transgene expression in human embryonic
stem cells using transposon-mediated gene transfer. Stem Cells *25*, 2919-2927.
10.1634/stemcells.2007-0026.

26.   Ng, Y.H., Chanda, S., Janas, J.A., Yang, N., Kokubu, Y., Südhof, T.C., and Wernig, M. (2021).
Efficient generation of dopaminergic induced neuronal cells with midbrain
characteristics. Stem Cell Reports *16*, 1763-1776. 10.1016/j.stemcr.2021.05.017.

27.   Subach, O.M., Cranfill, P.J., Davidson, M.W., and Verkhusha, V.V. (2011). An enhanced
monomeric blue fluorescent protein with the high chemical stability of the
chromophore. PLoS One *6*, e28674. 10.1371/journal.pone.0028674.

28.   Müller-Kuller, U., Ackermann, M., Kolodziej, S., Brendel, C., Fritsch, J., Lachmann, N.,
Kunkel, H., Lausen, J., Schambach, A., Moritz, T., and Grez, M. (2015). A minimal ubiquitous
chromatin opening element (UCOE) effectively prevents silencing of juxtaposed

heterologous promoters by epigenetic remodeling in multipotent and pluripotent stem
cells. Nucleic Acids Res *43*, 1577-1592. 10.1093/nar/gkv019.

29.   Chung, J.H., Bell, A.C., and Felsenfeld, G. (1997). Characterization of the chicken beta-
globin insulator. Proc Natl Acad Sci U S A *94*, 575-580. 10.1073/pnas.94.2.575.

30.   Rasmussen, Mikkel A., Holst, B., Tümer, Z., Johnsen, Mads G., Zhou, S., Stummann, Tina C.,
Hyttel, P., and Clausen, C. (2014). Transient p53 Suppression Increases Reprogramming
of Human Fibroblasts without Affecting Apoptosis and DNA Damage. Stem Cell Reports
3, 404-413. 10.1016/j.stemcr.2014.07.006.

31.   Js, J., R.H., Shaib, A.H., Rehbach, K., Lee, C., Seif, P., Thomas, C., Gideons, E., Guenther, A.,
Krutenko, T., et al. (2019). An Autaptic Culture System for Standardized Analyses of iPSC-
Derived Human Neurons. Cell reports *27*. 10.1016/j.celrep.2019.04.059.

32.   Jones, M.V., and Westbrook, G.L. (1995). Desensitized states prolong GABAA channel
responses to brief agonist pulses. Neuron *15*, 181-191. 10.1016/0896-6273(95)90075-
6.

33.   M., M., Rehbach, K., Brunner, J.W., Classen, J.A., Lammertse, H.C.A., van Linge, L.A., Schut,
D., Krutenko, T., Hebisch, M., Cornelisse, L.N., et al. (2019). A Single-Cell Model for
Synaptic Transmission and Plasticity in Human iPSC-Derived Neurons. Cell reports *27*.
10.1016/j.celrep.2019.04.058.

34.   Pantazis, C.B., Yang, A., Lara, E., McDonough, J.A., Blauwendraat, C., Peng, L., Oguro, H.,
Kanaujiya, J., Zou, J., Sebesta, D., et al. (2022). A reference human induced pluripotent
stem cell line for large-scale collaborative studies. Cell Stem Cell *29*, 1685-1702.e1622.
10.1016/j.stem.2022.11.004.

35.   Nadadhur, A.G., Alsaqati, M., Gasparotto, L., Cornelissen-Steijger, P., van Hugte, E., Dooves,
S., Harwood, A.J., and Heine, V.M. (2019). Neuron-Glia Interactions Increase Neuronal
Phenotypes in Tuberous Sclerosis Complex Patient iPSC-Derived Models. Stem Cell
Reports *12*, 42-56. 10.1016/j.stemcr.2018.11.019.

36.   Loomba, S., Straehle, J., Gangadharan, V., Heike, N., Khalifa, A., Motta, A., Ju, N., Sievers,
M., Gempt, J., Meyer, H.S., and Helmstaedter, M. (2022). Connectomic comparison of
mouse and human cortex. Science *377*. 10.1126/science.abo0924.

37.   Bakken, T.E., Jorstad, N.L., Hu, Q., Lake, B.B., Tian, W., Kalmbach, B.E., Crow, M., Hodge, R.D.,
Krienen, F.M., Sorensen, S.A., et al. (2021). Comparative cellular analysis of motor cortex
in human, marmoset and mouse. Nature *598*, 111-119. 10.1038/s41586-021-03465-8.

38.   Fang, R., Xia, C., Close, J.L., Zhang, M., He, J., Huang, Z., Halpern, A.R., Long, B., Miller, J.A.,
Lein, E.S., and Zhuang, X. (2022). Conservation and divergence of cortical cell

organization in human and mouse revealed by MERFISH. Science *377*, 56-62.
10.1126/science.abm1741.

39. Chanda, S., Ang, Cheen E., Davila, J., Pak, C., Mall, M., Lee, Qian Y., Ahlenius, H., Jung,
Seung W., Südhof, Thomas C., and Wernig, M. (2014). Generation of Induced Neuronal
Cells by the Single Reprogramming Factor ASCL1. Stem Cell Reports *3*, 282-296.
10.1016/j.stemcr.2014.05.020.

40. Jorstad, N.L., Close, J., Johansen, N., Yanny, A.M., Barkan, E.R., Travaglini, K.J., Bertagnolli,
D., Campos, J., Casper, T., Crichton, K., et al. (2023). Transcriptomic cytoarchitecture
reveals principles of human neocortex organization. Science *382*.
10.1126/science.adf6812.

41. Skene, N.G., and Grant, S.G. (2016). Identification of Vulnerable Cell Types in Major Brain
Disorders Using Single Cell Transcriptomes and Expression Weighted Cell Type
Enrichment. Front Neurosci *10*, 16. 10.3389/fnins.2016.00016.

42. Dixon, C., Sah, P., Lynch, J.W., and Keramidas, A. (2014). $GABA_A$ Receptor
α and β Subunits Shape Synaptic Currents via Different Mechanisms *.
Journal of Biological Chemistry *289*, 5399-5411. 10.1074/jbc.M113.514695.

43. Zhang, SJ., Huguenard, J.R., and Prince, D.A. (1997). GABAA Receptor-Mediated Cl–
Currents in Rat Thalamic Reticular and Relay Neurons. Journal of Neurophysiology *78*,
2280-2286. 10.1152/jn.1997.78.5.2280.

44. Koopmans, F., Van Nierop, P., Andres-Alonso, M., Byrnes, A., Cijsouw, T., Coba, M.P.,
Cornelisse, L.N., Farrell, R.J., Goldschmidt, H.L., Howrigan, D.P., et al. (2019). SynGO: An
Evidence-Based, Expert-Curated Knowledge Base for the Synapse. Neuron *103*, 217-
234.e214. 10.1016/j.neuron.2019.05.002.

45. Koopmans, F. (2024). GOAT: efficient and robust identification of gene set enrichment.
Communications Biology *7*, 744. 10.1038/s42003-024-06454-5.

46. Paraskevopoulou, F., Herman, M.A., and Rosenmund, C. (2019). Glutamatergic Innervation
onto Striatal Neurons Potentiates GABAergic Synaptic Output. J Neurosci *39*, 4448-4460.
10.1523/jneurosci.2630-18.2019.

47. Watanabe, M., and Fukuda, A. (2015). Development and regulation of chloride
homeostasis in the central nervous system. Front Cell Neurosci *9*, 371.
10.3389/fncel.2015.00371.

48. Roig Adam, A., Martínez-López, J.A., Van Der Spek, S.J.F., Achsel, T., Andres-Alonso, M.,
Bagni, C., Bayés, À., Biederer, T., Brose, N., Chua, J.J.E., et al. (2023). Transcriptional

diversity in specific synaptic gene sets discriminates cortical neuronal identity. Biology
Direct *18*. 10.1186/s13062-023-00372-y.

49. Ikeda, K., Yanagawa, Y., and Bekkers, J.M. (2008). Distinctive Quantal Properties of
Neurotransmission at Excitatory and Inhibitory Autapses Revealed Using Variance –
Mean Analysis. The Journal of Neuroscience *28*, 13563-13573. 10.1523/jneurosci.3350-
08.2008.

50. Jiang, L., Sun, S., Nedergaard, M., and Kang, J. (2000). Paired-pulse modulation at
individual GABAergic synapses in rat hippocampus. J Physiol *523 Pt 2*, 425-439.
10.1111/j.1469-7793.2000.t01-1-00425.x.

51. Lambert, N.A., and Wilson, W.A. (1994). Temporally distinct mechanisms of use-
dependent depression at inhibitory synapses in the rat hippocampus in vitro. Journal of
neurophysiology *72*, 121-130.

52. Pearce, R.A., Grunder, S.D., and Faucher, L.D. (1995). Different mechanisms for use-
dependent depression of two GABAA-mediated IPSCs in rat hippocampus. J Physiol *484*
(Pt 2), 425-435. 10.1113/jphysiol.1995.sp020675.

53. Rebollo, B., Abrahamyan, A., Thomale, U.-W., Kaindl, A.M., Herman, M.A., and Rosenmund,
C. (2025). GABAergic synaptic components are largely preserved across human and
mouse neuronal models. Frontiers in Cellular Neuroscience *Volume 19 - 2025*.
10.3389/fncel.2025.1588894.

54. Gonzalez-Ramos, A., Waloschková, E., Mikroulis, A., Kokaia, Z., Bengzon, J., Ledri, M.,
Andersson, M., and Kokaia, M. (2021). Human stem cell-derived GABAergic neurons
functionally integrate into human neuronal networks. Sci Rep *11*, 22050.
10.1038/s41598-021-01270-x.

55. Song, S., Ashok, A., Williams, D., Kaufman, M., Duff, K., and Sproul, A. (2021). Efficient
Derivation of Excitatory and Inhibitory Neurons from Human Pluripotent Stem Cells
Stably Expressing Direct Reprogramming Factors. Curr Protoc *1*, e141.
10.1002/cpz1.141.

56. Barretto, N., Zhang, H., Powell, S.K., Fernando, M.B., Zhang, S., Flaherty, E.K., Ho, S.M.,
Slesinger, P.A., Duan, J., and Brennand, K.J. (2020). ASCL1- and DLX2-induced GABAergic
neurons from hiPSC-derived NPCs. J Neurosci Methods *334*, 108548.
10.1016/j.jneumeth.2019.108548.

57. Marcó de la Cruz, B., Campos, J., Molinaro, A., Xie, X., Jin, G., Wei, Z., Acuna, C., and Sterky,
F.H. (2024). Liprin-α proteins are master regulators of human presynapse assembly.
Nature Neuroscience *27*, 629-642. 10.1038/s41593-024-01592-9.

58. Sanchez-Priego, C., Hu, R., Boshans, L.L., Lalli, M., Janas, J.A., Williams, S.E., Dong, Z., and
Yang, N. (2022). Mapping cis-regulatory elements in human neurons links psychiatric
disease heritability and activity-regulated transcriptional programs. Cell Reports *39*,
110877. https://doi.org/10.1016/j.celrep.2022.110877.

59. Clifford, M.A., Athar, W., Leonard, C.E., Russo, A., Sampognaro, P.J., Van der Goes, M.S.,
Burton, D.A., Zhao, X., Lalchandani, R.R., Sahin, M., et al. (2014). EphA7 signaling guides
cortical dendritic development and spine maturation. Proc Natl Acad Sci U S A *111*, 4994-
4999. 10.1073/pnas.1323793111.

60. Wu, J., Ren, J., Cui, H., Xie, Y., and Tang, Y. (2024). Rapid and high-purity differentiation of
human medium spiny neurons reveals LMNB1 hypofunction and subtype necessity in
modeling Huntington's disease. Inflamm Regen *44*, 7. 10.1186/s41232-024-00320-x.

61. Gierman, H.J., Indemans, M.H., Koster, J., Goetze, S., Seppen, J., Geerts, D., van Driel, R.,
and Versteeg, R. (2007). Domain-wide regulation of gene expression in the human
genome. Genome Res *17*, 1286-1295. 10.1101/gr.6276007.

62. Liew, C.-G., Draper, J.S., Walsh, J., Moore, H., and Andrews, P.W. (2007). Transient and Stable
Transgene Expression in Human Embryonic Stem Cells. Stem Cells *25*, 1521-1528.
10.1634/stemcells.2006-0634.

63. Kwak, C.S., Oflaz, F.E., Qiu, J., and Wang, X. (2024). Human stem cell-specific epigenetic
signatures control transgene expression. Biochimica et Biophysica Acta (BBA) - Gene
Regulatory Mechanisms *1867*, 195063. https://doi.org/10.1016/j.bbagrm.2024.195063.

64. He, J., Yang, Q., and Chang, L.-J. (2005). Dynamic DNA Methylation and Histone
Modifications Contribute to Lentiviral Transgene Silencing in Murine Embryonic
Carcinoma Cells. Journal of Virology *79*, 13497-13508. doi:10.1128/jvi.79.21.13497-
13508.2005.

65. Gödecke, N., Zha, L., Spencer, S., Behme, S., Riemer, P., Rehli, M., Hauser, H., and Wirth, D.
(2017). Controlled re-activation of epigenetically silenced Tet promoter-driven
transgene expression by targeted demethylation. Nucleic Acids Research *45*, e147-e147.
10.1093/nar/gkx601.

66. Allison, T., Langerman, J., Sabri, S., Otero-Garcia, M., Lund, A., Huang, J., Wei, X.,
Samarasinghe, R.A., Polioudakis, D., Mody, I., et al. (2021). Defining the nature of human
pluripotent stem cell-derived interneurons via single-cell analysis. Stem Cell Reports
16, 2548-2564. https://doi.org/10.1016/j.stemcr.2021.08.006.

67. Lindhout, F.W., Kooistra, R., Portegies, S., Herstel, L.J., Stucchi, R., Snoek, B.L., Altelaar,
A.F.M., MacGillavry, H.D., Wierenga, C.J., and Hoogenraad, C.C. (2020). Quantitative

mapping of transcriptome and proteome dynamics during polarization of human iPSC-
derived neurons. eLife *9*, e58124. 10.7554/eLife.58124.

68. Titlow, J.S., Kiourlappou, M., Palanca, A., Lee, J.Y., Gala, D.S., Ennis, D., Yu, J.J.S., Young, F.L.,
Susano Pinto, D.M., Garforth, S., et al. (2023). Systematic analysis of YFP traps reveals
common mRNA/protein discordance in neural tissues. Journal of Cell Biology *222*,
e202205129. 10.1083/jcb.202205129.

69. Moritz, C.P., Mühlhaus, T., Tenzer, S., Schulenborg, T., and Friauf, E. (2019). Poor transcript-
protein correlation in the brain: negatively correlating gene products reveal neuronal
polarity as a potential cause. Journal of Neurochemistry *149*, 582-604.
https://doi.org/10.1111/jnc.14664.

70. Lin, H.-C., Janssens, J., Eisinger, B., Hornauer, P., Kroell, A.-S., Santel, M., Pascual-Garcia, M.,
Okamoto, R., Karava, K., He, Z., et al. (2025). Human neuron subtype programming via
single-cell transcriptome-coupled patterning screens. Science *389*, eadn6121.
doi:10.1126/science.adn6121.

71. Müller, M., Felmy, F., Schwaller, B., and Schneggenburger, R. (2007). Parvalbumin is a
mobile presynaptic Ca2+ buffer in the calyx of Held that accelerates the decay of Ca2+
and short-term facilitation. J Neurosci *27*, 2261-2271. 10.1523/jneurosci.5582-06.2007.

72. Pór, A., Pocsai, K., Rusznák, Z., and Szucs, G. (2005). Presence and distribution of three
calcium binding proteins in projection neurons of the adult rat cochlear nucleus. Brain
Res *1039*, 63-74. 10.1016/j.brainres.2005.01.057.

73. Laing, B.T., Anderson, M.S., Bonaventura, J., Jayan, A., Sarsfield, S., Gajendiran, A.,
Michaelides, M., and Aponte, Y. (2023). Anterior hypothalamic parvalbumin neurons are
glutamatergic and promote escape behavior. Curr Biol *33*, 3215-3228.e3217.
10.1016/j.cub.2023.06.070.

74. Gartner, W., Lang, W., Leutmetzer, F., Domanovits, H., Waldhäusl, W., and Wagner, L. (2001).
Cerebral Expression and Serum Detectability of Secretagogin, a Recently Cloned EF-
hand Ca2+ -binding Protein. Cerebral Cortex *11*, 1161-1169. 10.1093/cercor/11.12.1161.

75. Tu, Y., Qin, J., Zhang, Q.M., Tang, T.S., Wang, L., and Yao, J. (2023). Secretagogin regulates
asynchronous and spontaneous glutamate release in hippocampal neurons through
interaction with Doc2α. Life Med *2*, lnad041. 10.1093/lifemedi/lnad041.

76. Hanics, J., Tretiakov, E.O., Romanov, R.A., Gáspárdy, A., Hevesi, Z., Schnell, R., Harkany, T.,
and Alpár, A. (2025). Neuronal activity modulates the expression of secretagogin, a
Ca(2+) sensor protein, during mammalian forebrain development. Acta Physiol (Oxf) *241*,
e70031. 10.1111/apha.70031.

77. Lin, H.-C., He, Z., Ebert, S., Schörnig, M., Santel, M., Nikolova, M.T., Weigert, A., Hevers, W.,
Kasri, N.N., Taverna, E., et al. (2021). NGN2 induces diverse neuron types from human
pluripotency. Stem cell reports *16*, 2118-2127. 10.1016/j.stemcr.2021.07.006.

78. Nehme, R., Zuccaro, E., Ghosh, S.D., Li, C., Sherwood, J.L., Pietilainen, O., Barrett, L.E.,
Limone, F., Worringer, K.A., Kommineni, S., et al. (2018). Combining NGN2 Programming
with Developmental Patterning Generates Human Excitatory Neurons with NMDAR-
Mediated Synaptic Transmission. Cell Reports *23*, 2509-2523.
10.1016/j.celrep.2018.04.066.

79. Rhee, H.J., Shaib, A.H., Rehbach, K., Lee, C., Seif, P., Thomas, C., Gideons, E., Guenther, A.,
Krutenko, T., Hebisch, M., et al. (2019). An Autaptic Culture System for Standardized
Analyses of iPSC-Derived Human Neurons. Cell Reports *27*, 2212-2228.e2217.
https://doi.org/10.1016/j.celrep.2019.04.059.

80. Bullmann, T., Kaas, T., Ritzau-Jost, A., Wöhner, A., Kirmann, T., Rizalar, F.S., Holzer, M.,
Nerlich, J., Puchkov, D., Geis, C., et al. (2024). Human iPSC-Derived Neurons with Reliable
Synapses and Large Presynaptic Action Potentials. J Neurosci *44*.
10.1523/jneurosci.0971-23.2024.

81. van Berkel, A.A., Lammertse, H.C.A., Öttl, M., Koopmans, F., Misra-Isrie, M., Meijer, M.,
Dilena, R., van Hasselt, P.M., Engelen, M., van Haelst, M., et al. (2024). Reduced MUNC18-
1 Levels, Synaptic Proteome Changes, and Altered Network Activity in STXBP1-Related
Disorder Patient Neurons. Biol Psychiatry Glob Open Sci *4*, 284-298.
10.1016/j.bpsgos.2023.05.004.

82. Fenske, P., Grauel, M.K., Brockmann, M.M., Dorrn, A.L., Trimbuch, T., and Rosenmund, C.
(2019). Autaptic cultures of human induced neurons as a versatile platform for studying
synaptic function and neuronal morphology. Scientific Reports *9*, 4890. 10.1038/s41598-
019-41259-1.

83. Togo, K., Fukusumi, H., Shofuda, T., Ohnishi, H., Yamazaki, H., Hayashi, M.K., Kawasaki, N.,
Takei, N., Nakazawa, T., Saito, Y., et al. (2021). Postsynaptic structure formation of human
iPS cell-derived neurons takes longer than presynaptic formation during neural
differentiation in vitro. Molecular Brain *14*, 149. 10.1186/s13041-021-00851-1.

84. Nadadhur, A.G., Emperador Melero, J., Meijer, M., Schut, D., Jacobs, G., Li, K.W., Hjorth, J.J.J.,
Meredith, R.M., Toonen, R.F., Van Kesteren, R.E., et al. (2017). Multi-level characterization
of balanced inhibitory-excitatory cortical neuron network derived from human
pluripotent stem cells. PLOS ONE *12*, e0178533. 10.1371/journal.pone.0178533.

85. Friedman, H.V., Bresler, T., Garner, C.C., and Ziv, N.E. (2000). Assembly of New Individual
Excitatory Synapses: Time Course and Temporal Order of Synaptic Molecule
Recruitment. Neuron *27*, 57-69. 10.1016/S0896-6273(00)00009-X.

86. Kwon, H.B., and Sabatini, B.L. (2011). Glutamate induces de novo growth of functional
spines in developing cortex. Nature *474*, 100-104. 10.1038/nature09986.

87. Richards, D.A., Mateos, J.M., Hugel, S., de Paola, V., Caroni, P., Gähwiler, B.H., and McKinney,
R.A. (2005). Glutamate induces the rapid formation of spine head protrusions in
hippocampal slice cultures. Proc Natl Acad Sci U S A *102*, 6166-6171.
10.1073/pnas.0501881102.

88. Mammen, A.L., Huganir, R.L., and O'Brien, R.J. (1997). Redistribution and Stabilization of
Cell Surface Glutamate Receptors during Synapse Formation. The Journal of
Neuroscience *17*, 7351-7358. 10.1523/jneurosci.17-19-07351.1997.

89. Petralia, R.S. (2012). Distribution of Extrasynaptic NMDA Receptors on Neurons. The
Scientific World Journal *2012*, 267120. https://doi.org/10.1100/2012/267120.

90. Okabe, S., Miwa, A., and Okado, H. (2001). Spine formation and correlated assembly of
presynaptic and postsynaptic molecules. J Neurosci *21*, 6105-6114.
10.1523/jneurosci.21-16-06105.2001.

91. Andreae, L.C., and Burrone, J. (2015). Spontaneous Neurotransmitter Release Shapes
Dendritic Arbors via Long-Range Activation of NMDA Receptors. Cell Rep *10*, 873-882.
10.1016/j.celrep.2015.01.032.

92. Sanes, J.R., and Lichtman, J.W. (1999). Development of the vertebrate neuromuscular
junction. Annu Rev Neurosci *22*, 389-442. 10.1146/annurev.neuro.22.1.389.

93. Harms, K.J., Tovar, K.R., and Craig, A.M. (2005). Synapse-specific regulation of AMPA
receptor subunit composition by activity. J Neurosci *25*, 6379-6388.
10.1523/jneurosci.0302-05.2005.

94. MacGillavry, H.D., Song, Y., Raghavachari, S., and Blanpied, T.A. (2013). Nanoscale
scaffolding domains within the postsynaptic density concentrate synaptic AMPA
receptors. Neuron *78*, 615-622. 10.1016/j.neuron.2013.03.009.

95. Vaughan, P.F., Walker, J.H., and Peers, C. (1998). The regulation of neurotransmitter
secretion by protein kinase C. Mol Neurobiol *18*, 125-155. 10.1007/bf02914269.

96. Cui, Y., Costa, R.M., Murphy, G.G., Elgersma, Y., Zhu, Y., Gutmann, D.H., Parada, L.F., Mody,
I., and Silva, A.J. (2008). Neurofibromin regulation of ERK signaling modulates GABA
release and learning. Cell *135*, 549-560. 10.1016/j.cell.2008.09.060.

97. Lützenkirchen, F.P., Zhu, Y., Maric, H.M., Boeck, D.S., Gromova, K.V., and Kneussel, M.
(2024). Neurobeachin regulates receptor downscaling at GABAergic inhibitory synapses
in a protein kinase A-dependent manner. Communications Biology *7*, 1635.
10.1038/s42003-024-07294-z.

98. Ghosh, H., Auguadri, L., Battaglia, S., Simone Thirouin, Z., Zemoura, K., Messner, S., Acuña,
M.A., Wildner, H., Yévenes, G.E., Dieter, A., et al. (2016). Several posttranslational
modifications act in concert to regulate gephyrin scaffolding and GABAergic
transmission. Nature Communications *7*, 13365. 10.1038/ncomms13365.

99. Tyagarajan, S.K., Ghosh, H., Yévenes, G.E., Imanishi, S.Y., Zeilhofer, H.U., Gerrits, B., and
Fritschy, J.M. (2013). Extracellular signal-regulated kinase and glycogen synthase kinase
3β regulate gephyrin postsynaptic aggregation and GABAergic synaptic function in a
calpain-dependent mechanism. J Biol Chem *288*, 9634-9647. 10.1074/jbc.M112.442616.

100. Tyagarajan, S.K., Ghosh, H., Yévenes, G.E., Nikonenko, I., Ebeling, C., Schwerdel, C., Sidler,
C., Zeilhofer, H.U., Gerrits, B., Muller, D., and Fritschy, J.M. (2011). Regulation of GABAergic
synapse formation and plasticity by GSK3beta-dependent phosphorylation of gephyrin.
Proc Natl Acad Sci U S A *108*, 379-384. 10.1073/pnas.1011824108.

101. Bright, D.P., and Smart, T.G. (2013). Protein kinase C regulates tonic GABAA receptor-
mediated inhibition in the hippocampus and thalamus. European Journal of
Neuroscience *38*, 3408-3423. https://doi.org/10.1111/ejn.12352.

102. Lützenkirchen, F.P., Zhu, Y., Maric, H.M., Boeck, D.S., Gromova, K.V., and Kneussel, M.
(2024). Neurobeachin regulates receptor downscaling at GABAergic inhibitory synapses
in a protein kinase A-dependent manner. Commun Biol *7*, 1635. 10.1038/s42003-024-
07294-z.

103. Farzana, F., Zalm, R., Chen, N., Li, K.W., Grant, S.G., Smit, A.B., Toonen, R.F., and Verhage,
M. (2016). Neurobeachin Regulates Glutamate- and GABA-Receptor Targeting to
Synapses via Distinct Pathways. Mol Neurobiol *53*, 2112-2123. 10.1007/s12035-015-
9164-8.

104. Hausrat, T.J., Muhia, M., Gerrow, K., Thomas, P., Hirdes, W., Tsukita, S., Heisler, F.F., Herich,
L., Dubroqua, S., Breiden, P., et al. (2015). Radixin regulates synaptic GABAA receptor
density and is essential for reversal learning and short-term memory. Nat Commun *6*,
6872. 10.1038/ncomms7872.

105. Nakamura, Y., Darnieder, L.M., Deeb, T.Z., and Moss, S.J. (2015). Regulation of GABAARs by
phosphorylation. Adv Pharmacol *72*, 97-146. 10.1016/bs.apha.2014.11.008.

106. Song, S.-H., and Augustine, G.J. (2023). Different mechanisms of synapsin-induced vesicle
clustering at inhibitory and excitatory synapses. Cell Reports *42*.
10.1016/j.celrep.2023.113004.

107. Flores, C.E., Nikonenko, I., Mendez, P., Fritschy, J.M., Tyagarajan, S.K., and Muller, D. (2015).
Activity-dependent inhibitory synapse remodeling through gephyrin phosphorylation.
Proc Natl Acad Sci U S A *112*, E65-72. 10.1073/pnas.1411170112.

108. Hao, Y., Liu, H., Zeng, X.-T., Wang, Y., Zeng, W.-X., Qian, K.-Y., Li, L., Chi, M.-X., Gao, S., Hu, Z.,
and Tong, X.-J. (2023). UNC-43/CaMKII-triggered anterograde signals recruit GABAAARs to
mediate inhibitory synaptic transmission and plasticity at C. elegans NMJs. Nature
Communications *14*, 1436. 10.1038/s41467-023-37137-0.

109. Houston, C.M., He, Q., and Smart, T.G. (2009). CaMKII phosphorylation of the GABA(A)
receptor: receptor subtype- and synapse-specific modulation. J Physiol *587*, 2115-2125.
10.1113/jphysiol.2009.171603.

110. Lorenz-Guertin, J.M., and Jacob, T.C. (2018). GABA type a receptor trafficking and the
architecture of synaptic inhibition. Developmental Neurobiology *78*, 238-270.
https://doi.org/10.1002/dneu.22536.

111. Heisler, F.F., Loebrich, S., Pechmann, Y., Maier, N., Zivkovic, A.R., Tokito, M., Hausrat, T.J.,
Schweizer, M., Bähring, R., Holzbaur, E.L., et al. (2011). Muskelin regulates actin filament-
and microtubule-based GABA(A) receptor transport in neurons. Neuron *70*, 66-81.
10.1016/j.neuron.2011.03.008.

112. Velasco, C.D., Santarella-Mellwig, R., Schorb, M., Gao, L., Thorn-Seshold, O., and Llobet,
A. (2023). Microtubule depolymerization contributes to spontaneous neurotransmitter
release in vitro. Communications Biology *6*, 488. 10.1038/s42003-023-04779-1.

113. Waites, C., Qu, X., and Bartolini, F. (2021). The synaptic life of microtubules. Curr Opin
Neurobiol *69*, 113-123. 10.1016/j.conb.2021.03.004.

114. Jacob, T.C., Moss, S.J., and Jurd, R. (2008). GABA(A) receptor trafficking and its role in the
dynamic modulation of neuronal inhibition. Nat Rev Neurosci *9*, 331-343.
10.1038/nrn2370.

115. Prag, S., De Arcangelis, A., Georges-Labouesse, E., and Adams, J.C. (2007). Regulation of
post-translational modifications of muskelin by protein kinase C. The International
Journal of Biochemistry & Cell Biology *39*, 366-378.
https://doi.org/10.1016/j.biocel.2006.09.003.

116. Biederer, T., Kaeser, P.S., and Blanpied, T.A. (2017). Transcellular Nanoalignment of
Synaptic Function. Neuron *96*, 680-696. 10.1016/j.neuron.2017.10.006.

117. Tang, A.H., Chen, H., Li, T.P., Metzbower, S.R., MacGillavry, H.D., and Blanpied, T.A. (2016).
A trans-synaptic nanocolumn aligns neurotransmitter release to receptors. Nature *536*,
210-214. 10.1038/nature19058.

118. Scimemi, A., and Diamond, J.S. (2012). The number and organization of Ca2+ channels in
the active zone shapes neurotransmitter release from Schaffer collateral synapses. J
Neurosci *32*, 18157-18176. 10.1523/jneurosci.3827-12.2012.

119. Sanderson, T.M., Georgiou, J., and Collingridge, G.L. (2020). Illuminating Relationships
Between the Pre- and Post-synapse. Front Neural Circuits *14*, 9.
10.3389/fncir.2020.00009.

120. Südhof, T.C. (2021). The cell biology of synapse formation. J Cell Biol *220*.
10.1083/jcb.202103052.

121. Südhof, T.C. (2018). Towards an Understanding of Synapse Formation. Neuron *100*, 276-
293. 10.1016/j.neuron.2018.09.040.

122. Olah, S.S., Kareemo, D.J., Buchta, W.C., Sinnen, B.L., Miller, C.N., Actor-Engel, H.S., Gookin,
S.E., Winborn, C.S., Kleinjan, M.S., Crosby, K.C., et al. (2023). Acute reorganization of
postsynaptic GABA$_A$ receptors reveals the functional impact of molecular
nanoarchitecture at inhibitory synapses. Cell Reports *42*. 10.1016/j.celrep.2023.113331.

123. Karayannis, T., Au, E., Patel, J.C., Kruglikov, I., Markx, S., Delorme, R., Héron, D., Salomon,
D., Glessner, J., Restituito, S., et al. (2014). Cntnap4 differentially contributes to GABAergic
and dopaminergic synaptic transmission. Nature *511*, 236-240. 10.1038/nature13248.

124. Um, J.W., Kim, K.H., Park, B.S., Choi, Y., Kim, D., Kim, C.Y., Kim, S.J., Kim, M., Ko, J.S., Lee, S.-
G., et al. (2014). Structural basis for LAR-RPTP/Slitrk complex-mediated synaptic
adhesion. Nature Communications *5*, 5423. 10.1038/ncomms6423.

125. Fiederling, A., Ewert, R., Andreyeva, A., Jüngling, K., and Gottmann, K. (2011). E-cadherin
is required at GABAergic synapses in cultured cortical neurons. Neuroscience Letters
501, 167-172. https://doi.org/10.1016/j.neulet.2011.07.009.

126. Beuter, S., Ardi, Z., Horovitz, O., Wuchter, J., Keller, S., Saha, R., Tripathi, K., Anunu, R., Kehat,
O., Kriebel, M., et al. (2016). Receptor tyrosine kinase EphA7 is required for interneuron
connectivity at specific subcellular compartments of granule cells. Scientific Reports *6*,
29710. 10.1038/srep29710.

127. Park, Y.K., and Goda, Y. (2016). Integrins in synapse regulation. Nature Reviews
Neuroscience *17*, 745-756. 10.1038/nrn.2016.138.

128. Sia, G.-M., Béïque, J.-C., Rumbaugh, G., Cho, R., Worley, P.F., and Huganir, R.L. (2007).
Interaction of the N-Terminal Domain of the AMPA Receptor GluR4 Subunit with the
Neuronal Pentraxin NP1 Mediates GluR4 Synaptic Recruitment. Neuron *55*, 87-102.
10.1016/j.neuron.2007.06.020.

129. Ray, A., and Treloar, H.B. (2012). IgSF8: a developmentally and functionally regulated cell
adhesion molecule in olfactory sensory neuron axons and synapses. Mol Cell Neurosci
50, 238-249. 10.1016/j.mcn.2012.05.007.

130. Li, W., Yang, L., Tang, C., Liu, K., Lu, Y., Wang, H., Yan, K., Qiu, Z., and Zhou, W. (2020).
Mutations of CNTNAP1 led to defects in neuronal development. JCI Insight *5*.
10.1172/jci.insight.135697.

131. Wiera, G., Brzdąk, P., Lech, A.M., Lebida, K., Jabłońska, J., Gmerek, P., and Mozrzymas, J.W.
(2022). Integrins Bidirectionally Regulate the Efficacy of Inhibitory Synaptic
Transmission and Control GABAergic Plasticity. J Neurosci *42*, 5830-5842.
10.1523/jneurosci.1458-21.2022.

132. Tucić, M., Stamenković, V., and Andjus, P. (2021). The Extracellular Matrix Glycoprotein
Tenascin C and Adult Neurogenesis. Front Cell Dev Biol *9*, 674199.
10.3389/fcell.2021.674199.

133. Quattrocchi, C.C., Wannenes, F., Persico, A.M., Ciafré, S.A., D'Arcangelo, G., Farace, M.G.,
and Keller, F. (2002). Reelin is a serine protease of the extracellular matrix. J Biol Chem
277, 303-309. 10.1074/jbc.M106996200.

134. O'Brien, R.J., Xu, D., Petralia, R.S., Steward, O., Huganir, R.L., and Worley, P. (1999). Synaptic
clustering of AMPA receptors by the extracellular immediate-early gene product Narp.
Neuron *23*, 309-323. 10.1016/s0896-6273(00)80782-5.

135. Yim, Y.S., Kwon, Y., Nam, J., Yoon, H.I., Lee, K., Kim, D.G., Kim, E., Kim, C.H., and Ko, J. (2013).
Slitrks control excitatory and inhibitory synapse formation with LAR receptor protein
tyrosine phosphatases. Proc Natl Acad Sci U S A *110*, 4057-4062.
10.1073/pnas.1209881110.

136. Martin, E.E., Wleklinski, E., Hoang, H.T.M., and Ahmad, M. (2021). Interaction and
Subcellular Association of PRRT1/SynDIG4 With AMPA Receptors. Frontiers in Synaptic
Neuroscience *Volume 13 - 2021*. 10.3389/fnsyn.2021.705664.

137. Matsunaga, H., and Aruga, J. (2021). Trans-Synaptic Regulation of Metabotropic
Glutamate Receptors by Elfn Proteins in Health and Disease. Frontiers in Neural Circuits
Volume 15 - 2021. 10.3389/fncir.2021.634875.

138. Hijazi, S., Heistek, T.S., van der Loo, R., Mansvelder, H.D., Smit, A.B., and van Kesteren, R.E.
(2020). Hyperexcitable Parvalbumin Interneurons Render Hippocampal Circuitry
Vulnerable to Amyloid Beta. iScience *23*, 101271. 10.1016/j.isci.2020.101271.

139. Mossner, J.M., Batista-Brito, R., Pant, R., and Cardin, J.A. (2020). Developmental loss of
MeCP2 from VIP interneurons impairs cortical function and behavior. eLife *9*, e55639.
10.7554/eLife.55639.

140. Malwade, S., Gasthaus, J., Bellardita, C., Andelic, M., Moric, B., Korshunova, I., Kiehn, O.,
Vasistha, N.A., and Khodosevich, K. (2022). Identification of Vulnerable Interneuron
Subtypes in 15q13.3 Microdeletion Syndrome Using Single-Cell Transcriptomics. Biol
Psychiatry *91*, 727-739. 10.1016/j.biopsych.2021.09.012.

141. Goel, A., Cantu, D.A., Guilfoyle, J., Chaudhari, G.R., Newadkar, A., Todisco, B., de Alba, D.,
Kourdougli, N., Schmitt, L.M., Pedapati, E., et al. (2018). Impaired perceptual learning in
a mouse model of Fragile X syndrome is mediated by parvalbumin neuron dysfunction
and is reversible. Nat Neurosci *21*, 1404-1411. 10.1038/s41593-018-0231-0.

142. Gabitto, M.I., Travaglini, K.J., Rachleff, V.M., Kaplan, E.S., Long, B., Ariza, J., Ding, Y.,
Mahoney, J.T., Dee, N., Goldy, J., et al. (2023). Integrated multimodal cell atlas of
Alzheimer's disease. Res Sq. 10.21203/rs.3.rs-2921860/v1.

143. Zott, B., Busche, M.A., Sperling, R.A., and Konnerth, A. (2018). What Happens with the
Circuit in Alzheimer's Disease in Mice and Humans? Annu Rev Neurosci *41*, 277-297.
10.1146/annurev-neuro-080317-061725.

144. van Boven, M.A., Mestroni, M., Zwijnenburg, P.J.G., Verhage, M., and Cornelisse, L.N. (2024).
A de novo missense mutation in synaptotagmin-1 associated with neurodevelopmental
disorder desynchronizes neurotransmitter release. Mol Psychiatry *29*, 1798-1809.
10.1038/s41380-024-02444-5.

145. Østergaard, M., Barbagallo, P., Frederiksen, H.R.S., Chung, W.K., Møller, R.S., Larsen, M.R.,
Freude, K., Verhage, M., and Sørensen, J.B. (2025). SNAP25 variant I67N: synaptic
phenotypes, drug response and proteome changes in human neurons. Brain.
10.1093/brain/awaf119.

146. Öttl, M., Toonen, R.F., and Verhage, M. (2024). Reduced synaptic depression in human
neurons carrying homozygous disease-causing STXBP1 variant L446F. Hum Mol Genet
33, 991-1000. 10.1093/hmg/ddae035.

147. Schmitz, S.K., Hjorth, J.J., Joemai, R.M., Wijntjes, R., Eijgenraam, S., de Bruijn, P., Georgiou,
C., de Jong, A.P., van Ooyen, A., and Verhage, M. (2011). Automated analysis of neuronal

morphology, synapse number and synaptic recruitment. Journal of neuroscience methods *195*, 185-193.

148. Thanou, E., Koopmans, F., Pita-Illobre, D., Klaassen, R.V., Özer, B., Charalampopoulos, I., Smit, A.B., and Li, K.W. (2023). Suspension TRAPping Filter (sTRAP) Sample Preparation for Quantitative Proteomics in the Low µg Input Range Using a Plasmid DNA Micro-Spin Column: Analysis of the Hippocampus from the 5xFAD Alzheimer's Disease Mouse Model. Cells *12*. 10.3390/cells12091242.

149. Demichev, V., Messner, C.B., Vernardis, S.I., Lilley, K.S., and Ralser, M. (2020). DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. Nature Methods *17*, 41-44. 10.1038/s41592-019-0638-x.

150. Koopmans, F., Li, K.W., Klaassen, R.V., and Smit, A.B. (2023). MS-DAP Platform for Downstream Data Analysis of Label-Free Proteomics Uncovers Optimal Workflows in Benchmark Data Sets and Increased Sensitivity in Analysis of Alzheimer's Biomarker Data. J Proteome Res *22*, 374-386. 10.1021/acs.jproteome.2c00513.

151. Zhu, Y., Orre, L.M., Zhou Tran, Y., Mermelekas, G., Johansson, H.J., Malyutina, A., Anders, S., and Lehtiö, J. (2020). DEqMS: A Method for Accurate Variance Estimation in Differential Protein Expression Analysis. Mol Cell Proteomics *19*, 1047-1057. 10.1074/mcp.TIR119.001646.

152. Koopmans, F. (2024). GOAT: efficient and robust identification of gene set enrichment. Commun Biol *7*, 744. 10.1038/s42003-024-06454-5.

153. Koopmans, F., van Nierop, P., Andres-Alonso, M., Byrnes, A., Cijsouw, T., Coba, M.P., Cornelisse, L.N., Farrell, R.J., Goldschmidt, H.L., Howrigan, D.P., et al. (2019). SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse. Neuron *103*, 217-234.e214. 10.1016/j.neuron.2019.05.002.

154. Demichev, V., Szyrwiel, L., Yu, F., Teo, G.C., Rosenberger, G., Niewienda, A., Ludwig, D., Decker, J., Kaspar-Schoenefeld, S., Lilley, K.S., et al. (2022). dia-PASEF data analysis using FragPipe and DIA-NN for deep proteomics of low sample amounts. Nat Commun *13*, 3944. 10.1038/s41467-022-31492-0.

155. Demichev, V., Messner, C.B., Vernardis, S.I., Lilley, K.S., and Ralser, M. (2020). DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. Nat Methods *17*, 41-44. 10.1038/s41592-019-0638-x.

156. Huson, V., and Cornelisse, L.N. (2019). viewEPSC.

157. Neher, E. (2015). Merits and limitations of vesicle pool models in view of heterogeneous populations of synaptic vesicles. Neuron *87*, 1131-1142.

158. Wiśniewski, J.R., and Gaugaz, F.Z. (2015). Fast and sensitive total protein and Peptide
assays for proteomic analysis. Anal Chem *87*, 4110-4116. 10.1021/ac504689z.

159. Skowronek, P., Thielert, M., Voytik, E., Tanzer, M.C., Hansen, F.M., Willems, S., Karayel, O.,
Brunner, A.D., Meier, F., and Mann, M. (2022). Rapid and In-Depth Coverage of the
(Phospho-)Proteome With Deep Libraries and Optimal Window Design for dia-PASEF.
Mol Cell Proteomics *21*, 100279. 10.1016/j.mcpro.2022.100279.

160. Miedema, S.S.M., Mol, M.O., Koopmans, F.T.W., Hondius, D.C., van Nierop, P., Menden, K.,
de Veij Mestdagh, C.F., van Rooij, J., Ganz, A.B., Paliukhovich, I., et al. (2022). Distinct cell
type-specific protein signatures in GRN and MAPT genetic subtypes of frontotemporal
dementia. Acta Neuropathologica Communications *10*, 100. 10.1186/s40478-022-
01387-8.

161. Team, R. (2020). RStudio: Integrated Development for R (PBC, Boston, MA).

162. Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (Springer International
Publishing).

163. Perez-Riverol, Y., Bandla, C., Kundu, D.J., Kamatchinathan, S., Bai, J., Hewapathirana, S.,
John, N.S., Prakash, A., Walzer, M., Wang, S., and Vizcaíno, J.A. (2025). The PRIDE database
at 20 years: 2025 update. Nucleic Acids Res *53*, D543-d553. 10.1093/nar/gkae1011.
1462

1463   **Figures**

1464
Figure 1. PiggyBac Asc1/Dlx2 efficiently generates pure GABAergic neurons,
A Map of the all-in-one PiggyBac (PB) vector, expressing murine *Ascl1* and *Dlx2* from a bi-
directional TRE-responsive promoter, rtTA from the CAG promoter and *pac*-T2A-NLS-
mTagBFP from the EF1α promoter. **B** Overview of the protocol for *Ascl1/Dlx2* induction. **C**

Before induction, all iPSCs show nuclear BFP signals, demonstrating successful integration
of the PB construct **D** Dox-induced expression of *Dlx2* and *Ascl1* triggers neuronal
morphology to arise within one day (DPI1) **E** A week after induction (WPI 1), neural
morphology is clearly visible **F** After 10 weeks of culture, PB *Ascl1/Dlx2* induced neurons
show mature neuronal morphology with GABAergic synapses (VGAT+ puncta) and no
glutamatergic synapses (VGLUT+ puncta) Scale bar: 100 µm **G.** Zoom-in on the dendrite
highlighted in F. scale bar = 1 µm. **H** Autaptic cultures of PB *Ascl1/Dlx2* induced neurons were
assessed using patch clamp electrophysiology. **I** Evoked postsynaptic currents (PSCs) were
fitted using a mono-exponential decay function (red) and their currents were diminished by
the application of PTX. **J** Dot-plot of the decay time constants of ePSCs recorded in PB
Ascl1/Dlx2 induced neurons. **K** Dot-plot showing the distribution of cells with 0-10% of the
naïve charge remaining in PTX **L** Overview of the four induction methods used, PB *Ascl1/Dlx2*
induction was compared to the lentiviral *Ascl1/Dlx2* induction, *Ascl1*+Forskolin induction and
NGN2 glutamatergic induction. **M** Example ePSC traces with either fast or slow decay, as
measured in the inductions tested here. **N** Histogram of log-transformed decay time
constants (decay τ) measured in neurons induced with the four different induction strategies,
together with a 'sum of two gaussians' fit (grey line). **O** Pie-charts showing the percentages
of neurons with a slow (τ ≥ 30 ms), intermediate (6 > τ < 30 ms) and fast (τ ≤ 6 ms) decay, as
well as the percentage of synaptically silent neurons, of neurons generated with the four
different induction strategies.
1489

1490

Figure 2. PiggyBac Ascl1/Dlx2 neurons integrate into well-defined excitatory/inhibitory networks upon co-culturing with NGN2 induced neurons

A Schematic of co-culturing protocol in which PiggyBac (PB) Ascl1/Dlx2 (AD) neurons were co-cultured with glutamatergic NGN2 induced neurons to generate excitatory/inhibitory (E/I) co-cultures **B** Representative images of E/I co-cultures at different ratios in which dendrites and neuronal nuclei are visualised using MAP2 and NeuN staining, respectively, and PB-AD neurons visualised by nuclear mTagBFP signals. The E/I-ratio is indicated above each image. Scale bar = 100 μm. **C** Representative images of E/I co-cultures at the same ratios as depicted in (B), stained for MAP2 as dendritic marker and VGAT and VGLUT as markers for GABAergic and glutamatergic synapses, respectively. Scae bars = 100 μm, scale bar zoom in = 10 μm **D** The network density (number of nuclei/field of view) is stable across E/I co-culture ratios. **E** The total dendrite length per field of view decreases as the percentage of plated PB-AD neurons increases. **F** The density of glutamatergic (VGLUT⁺ puncta) and GABAergic (VGAT⁺)

1504   puncta scales with the percentage of NGN2 and PB-AD neurons plated. **G** Synapse counts

1505   shown in F expressed as the percentage of all synapses counted.

1506

Figure 3. *PiggyBac Ascl1/Dlx2 neurons have a cortical GABAergic proteomic profile.*

A. Schematic overview of the proteomics analysis in which PiggyBac (PB) Ascl1/Dlx2 (AD) rat astrocyte co-cultures were compared to NGN2 co-cultures per timepoint. Double-sided arrows indicate the contrasts used for analysis. **B**. PCA analysis on neuronal protein abundances reveals separation of culture types as well as time points along the first two components (PC1: 61.7% and PC2: 8.0%). **C-F**. Volcano plots showing regulation of neuronal proteins in PB-AD cultures and in NGN2 cultures at 4 WPI (**C**), 6 WPI (**D**), 8 WPI (**E**) and 10 WPI (**F**). The top 5% significantly upregulated neuronal proteins in PB-AD are shown as blue dots and those in NGN2 as red dots. Proteins are considered significantly regulated if their adjusted p-value is < 0.05 (significance threshold overlaps with x-axis at this scale and is not shown for clarity of the figure) and their log$_2$ (FC) lies above the 95th or below the 5th percentile (vertical dashed lines). Label colours indicate the predicted cell types based on

cell type specificity scores (CTSS): regulated GABAergic (purple) and glutamatergic proteins

(orange). Cell type specificity was defined using the following CTSS cutoffs: neuronal proteins

(neuronal CTSS>70%), GABAergic proteins (GABAergic CTSS > 80% and glutamatergic CTSS

< 10%), glutamatergic proteins (glutamatergic CTSS > 80% and GABAergic CTSS < 10%). CTSS

can be found in Figure S6. **G**. EWCE analysis of top 5% upregulated neuronal proteins in PB-

AD cultures reveals specific enrichment for GABAergic neurons, which increased up to 8 WPI.

H. EWCE analysis of the top 5% upregulated neuronal proteins in NGN2 cultures reveals

significant enrichment for both glutamatergic and GABAergic neurons depending on the

timepoint.

1530

1531

1532

Figure 4. Morphological and functional development of PiggyBac Ascl1/Dlx2 induced neurons

A Typical examples of PB-AD autaptic neurons at WPI 4 (i), 6 (ii), 8 (iii) and 10 (iv). Scale bar:

100 μm **B** Total dendrite length quantified. **C** number of GABAergic synapses per neuron. **D**

Synaptic density. **E** Average VGAT puncta intensity in arbitrary units (AU). Scale bars: 100 μm.

Boxplot whiskers extend the entire data range. **F** Representative traces of mPSCs recorded

from PiggyBac (PB) *Ascl1/Dlx2* (*AD*) neurons (PB-AD) neurons at 4, 6, 8 and 10 weeks post

induction (WPI). **G** mPSC frequency and **(H)** amplitude of PB-AD neurons at 4, 6, 8 and 10

WPI. **I** Boxplots of the average fractions of synaptically active cells at 4, 6, 8 and 10 WPI.

Markers represent averages of individual inductions. **J** Representative ePSC traces of at 4, 6,

8 and 10 WPI. **K** Amplitude, charge (**L)** and decay time (**M**) constant of ePSCs recorded from

PB-AD neurons at 4, 6, 8 and 10 WPI.

Figure 5. Pre- and postsynaptic maturation of PiggyBac Ascl1/Dlx2 neurons

A. Schematic overview of the proteomics analysis with double-sided arrows indicating the

contrasts used for analysis. **B-C**. Stacked bar plots showing the number of up- and

downregulated SYNGO annotated proteins over time, distinguishing synaptic (black), exclusively presynaptic (green) and exclusively postsynaptic (magenta) proteins for PB-AD (**B**) and NGN2 (**C**) neurons. Classifications were assigned based on SynGO annotations, synaptic classification contains both pansynaptic and ambiguously localized synaptic proteins. Regulated proteins have a $\log_2$ (FC) > 0.1, or < -0.1, and adjusted p-value < 0.05. **D-F.** Volcano plots showing regulation of synaptic proteins between 6 and 4 (**D**), 8 and 6 (**E**), and 10 and 8 WPI (**F**). Only SYNGO-annotated proteins are plotted and colour coding refer to significantly regulated synaptic (black) exclusively presynaptic (green) or postsynaptic (pink) annotations. Proteins were regarded significantly regulated when adjusted p-value < 0.05 (dashed horizontal line) and $\log_2$ (FC) > 0.1 or < -0.1 (dashed vertical lines). Top 5% proteins regulated proteins were labelled based on 97.5^{th} and 2.5^{th} percentiles marking genes that drive subsequently detected enrichments. **G-I.** SYNGO plots showing enrichment scores (z-scores) derived from gene-set enrichment analysis with GOAT on SYNGO cellular compartment gene sets, using differential expression output of 6 vs 4 (G), 8 vs 6 (H) and 10 vs 8 WPI (I),. Negative z-scores reflect enrichments primarily driven by downregulated proteins and vice versa for positive z-scores. **J-L.** Lollipop charts showing significant ontologies with their corresponding adjusted p-values and the number of genes part of the gene set as analysed by GOAT for 6 vs 4 (J), 8 vs 6 (K) and 8 vs 10 WPI (L) **M.** Schematic overview of the GABA puff experiment, in which perforated-patch was performed on PB-AD neurons in E/I-networks and GABA was applied to measure the postsynaptic current it produced. **N.** Typical example of the postsynaptic current produced by GABA puff at different holding voltages. **O.** Linear fit of PSC amplitudes as measured in response to GABA puff at 4, 6, 8 and 10 WPI. Error bars represent the SD.

1576

1577

Figure 6. Specialization of PiggyBac Ascl1/Dlx2 and NGN2 neurons into cortical subtypes

A. Schematic overview of the proteomics analysis pipeline. The differential expression of synaptic proteins over time (10 vs 4 WPI) was first quantified for PB-AD and NGN2 culture separately. The difference in expression over time between PB-AD and NGN2 (delta- $\log_2$ (FC)) was next calculated, enabling identification of proteins with PB-AD- or NGN2-specific temporal profiles. Density plot showing 5th and 95th percentiles (dashed vertical lines) selections of proteins that have strong differential temporal regulation (i.e., high or low

delta- $\log_2$ (FC)) in PB-AD compared to NGN2, and vice versa. **B-C**. Volcano plots showing

regulation of synaptic proteins between 10 and 4 WPI. Proteins are considered regulated

when the $\log_2$ (FC) > 0.1 or < -0.1 (dashed vertical lines) and adjusted p-value < 0.05 (dashed

horizontal line). Only SYNGO-annotated proteins are plotted, and dot colors refer to

significantly regulated synaptic (black) exclusively presynaptic (green) or postsynaptic (pink)

annotations. Label colours indicate for predicted cortical GABAergic- (purple) or

glutamatergic (orange) regulated proteins. **D-G** SYNGO sunburst plots showing the

GABAergic and glutamatergic specification of PB-AD and NGN cultures. The SYNGO sunburst

inputs are the proteins that are part of the 5th (**F-G**) and 95th percentiles (**D-E**) from the delta-

$\log_2$ (FC) distribution (**A**), which represent the top 5% exclusively PB-AD and NGN2 regulated

proteins, respectively. GABAergic or glutamatergic annotation for these SYNGO terms is

estimated based on the differential RNA expression ($\log_2$ (FC)) between cortical GABAergic

and glutamatergic neurons of corresponding genes[40] Both cellular compartment (**D-F**) and

biological process (**E-G**) gene sets were used. **H-K**. EWCE analysis of PB-AD-specific (**H-I**) or

NGN2-specific proteins (**J-K**) show enrichment for GABAergic and glutamatergic cell sub

(classes), respectively. IT and ET refer to intra- and extratelenchephalic-projecting neurons

and NP refers to near-projecting neurons.[40]

1602

1603 **Supplementary figures and table**

1604

Figure S1. NGN2 neurons produce glutamatergic transmission

A Voltage clamp recordings of 6 WPI NGN2 autaptic neurons after stimulation, in normal
ACSF (Naive) and in presence of glutamatergic receptor blockers (APV + CNQX). Synaptic
transmission was fully abolished by blockers, illustrating that NGN2 neurons are
glutamatergic. **B-C** Examples of immunocytochemistry for MAP2, VGAT and VGLUT1+2
performed on NGN2 autaptic neurons, with zoom in on a length of dendrite (**C**). Scale bar =
100 µm, zoom in scale bar = 10 µm

1612

1613

Figure S2. PiggyBac Ascl1/Dlx2 induction reliably generates GABAergic neurons in the KOLF2.1J and GM23973 line

A Representative images of PiggyBac (PB) Ascl1/Dlx2 (AD) neurons induced from KOLF1.2J

iPSCs at 7 weeks post induction (WPI), stained for MAP2 as dendritic marker, and VGAT and

VGLUT1/2 as GABAergic and glutamatergic markers, respectively. Scale bar: 100 µm **B** Zoom-

in on dendrite to illustrate punctate VGAT staining. Scale bar: 10 μm **C.** Total dendrite length
and (**D**) number of GABAergic synapses per neuron. **E** Synaptic density and intensity (**F**). **G**
Representative mPSC trace and (**H**) representative ePSC trace from PB-AD neurons induced
from the KOLF2.1J line. **I** Frequency and (**J**) amplitude of miPSCs. **K** Amplitude, (**L**) charge and
(**M**) decay time constant of iPSCs in PB-AD neurons generated from the KOLF2.1J line. **N**
Representative trace of iPSCs evoked by 20Hz stimulation trains with zoom-in on first five
pulses in KOLF2.1J. **O** Plot of cumulative iPSC amplitudes evoked by 20 Hz stimulus trains,
with back-extrapolated linear fits displayed in red. **P** RRP estimate obtained from back-
extrapolation of linear fits displayed in (**M**). **Q**. representative ePSC traces from PB-AD
neurons induced from the GM23973 line. **R.** Representative trace of iPSCs evoked by 20Hz
stimulation trains with zoom-in on first five pulses in GM23973. **S** Amplitude, (**T**) charge and
(**U**) decay time constant of iPSCs in PB-AD neurons generated from the GM23973.1J line. **V**
Plot of cumulative iPSC amplitudes evoked by 20 Hz stimulus trains, with back-extrapolated
linear fits displayed in red. **W** RRP estimate obtained from back-extrapolation of linear fits
displayed in (M). Boxplot whiskers extend the entire data range.
1633

1634

Figure S3. Peptide and protein detection count and variation is similar between cultures and timepoints.

A-B Boxplots showing variation of peptide (**A**) and protein (**B**) detection for all timepoints per culture, indicating median, 1st and 3rd quartiles, and minimum and maximum values. Number of detected peptides and proteins are indicated in the base of the graph. Median coefficient of variation for peptide and protein detection across groups ranged from 14.6% to 32.2%, reflecting consistent detection performance across replicates. **C** PCA plots showing potential batch effects due to induction. PC1 and PC2 show even spread of different inductions but PC3 (3.9%) shows slight separation based on induction and we therefore corrected for induction effect when performing differential abundance analysis.

1644

1645

Figure S4. Cell type specificity scores for detected GABAergic proteins

A. Cell type specificity scores (CTSS) scores for detected GABAergic and glutamatergic

proteins. CTSS are calculated with the expression weighted cell type enrichment tool.[41]

using single-cell RNA-sequencing data from Jorstad et al. [40] MGE and CGE refer to medial

and caudal ganglionic eminences. **B.** Representative images of PB-AD and NGN2 neurons

stained for MAP2 and PVALB. Scale bar = 100 µm. **C-D** Expression-weighted cell type

enrichment (EWCE) analysis showing enrichment of proteins higher expressed in PB-AD or

NGN2 for GABAergic **(C)** and glutamatergic subtypes **(D)**. PBAD-enriched proteins showed

strongest enrichment for VIP+ interneurons at most timepoints, whereas NGN2-enriched

proteins map at different timepoints to both GABAergic (PVALB+ and VIP+ interneurons) and

glutamatergic subtypes. PB-AD-enriched proteins showed unexpected enrichment for L5ET

neurons at 4 WPI, which reduced below significance threshold at later timepoints. IT and ET

refer to intra- and extratelenchephalic-projecting neurons, and NP refers to near-projecting

neurons – cell subclass annotations consistent with Jorstad et al. (2023).[40]

1660
Figure S5. Extended morphological and functional development of PiggyBac Ascl1/Dlx2 induced neurons

A Representative images of an autaptic PiggyBac (PB) *Ascl1/Dlx2* (AD) induced neuron at
week-post-induction 10, stained for MAP2 as dendritic marker, and VGAT and VGLUT1/2 as
GABAergic and glutamatergic markers, respectively. Scale bar: 100 μm **B** Cut-out of dendrite

marked in A to illustrate individual synaptic puncta. Scale bar: 10 μm **C** Sholl analysis of PB-
AD autaptic neurons at WPI 4, 6, 8 and 10. Shaded areas represent the SEM. **D** Quantification
of the area under the curve (AUC) of the Sholl histogram shown in (**C**) **E** Representative traces
of paired-pulse ePSCs at a 50, 200 and 1000 ms interval at 10 WPI. **F** Quantified paired-pulse
amplitude ratios (PPR) at a 50 ms, 200 ms (**G**) and 100 ms (**H**) interval at 4, 6, 8 and 10 WPI. **I**
Representative traces of iPSCs evoked by 20Hz stimulus trains from PB-AD neurons at 4, 6,
8 and 10 WPI with zoom in on first five pulses. **J** Plot of cumulative iPSC charge evoked by
20 Hz stimulus trains, with back-extrapolated linear fits displayed in red. Note, WPI 4 and
WPI 6 data overlap. **K** RRP estimate obtained from back-extrapolation of linear fits displayed
in (**J**).

1675

Figure S6. Differential expression of synaptic genes in NGN2 and PB-AD neurons

A-C. Volcano plots showing regulation of synaptic proteins in NGN2 cultures between 6 and

4 (**A**), 8 and 6 (**B**), and 10 and 8 WPI (**C**). Only SYNGO-annotated proteins are plotted and

colour coding refer to significantly regulated synaptic (black) exclusively presynaptic (green)

or postsynaptic (pink) annotations. Proteins were regarded significantly regulated when

adjusted p-value < 0.05 (dashed horizontal line) and $\log_2$ (FC) > 0.1 or < -0.1 (dashed vertical

lines). **D-F**. SYNGO plots showing enrichment scores (z-scores) derived from gene-set

enrichment analysis with GOAT using SYNGO cellular compartment gene sets for 4 to 6 (**D**),

6 to 8 (**E**) and 8 to 10 WPI (**F**). Negative z-scores (blue) reflect enrichments primarily driven

by downregulated proteins and vice versa for positive z-scores (red). **G-I.** Lollipop charts

showing significant SYNGO terms with their corresponding adjusted p-values and the

number of genes part of the gene set. **J-O.** SYNGO overrepresentation analysis of upregulated

proteins in PB-AD (**J-L**) and NGN2 (**M-O**) per timepoint. Foreground are significantly

upregulated proteins and background are all tested genes in corresponding contrast.

1690

Figure S7. PiggyBac Ascl1/Dlx2 and NGN2 neurons express markers of diverse GABAergic neuron subtypes

A. Contrasts used to determine cell type specification. **B** and **C** Heatmaps that show temporal

expression profiles relative to 4 WPI (log$_2$ (FC)) and cell type specificity scores (CTSS). **B**

Typical proteins required for neurotransmitter (NT) synthesis, vesicle loading, and receptors.

Undetected proteins are visualized for NT biosynthesis and vesicle loading but not for

receptors. **C** Canonical GABAergic subtype markers and proteins with high cell type specificity

for GABAergic subtypes and for medial or caudal ganglionic eminences (MGE or CGE). **D.** Line

plots show sum expression of canonical GABAergic subtype markers (top) and other proteins

with GABAergic cell type specificity. Error bars represent SD and dot size indicate detection

count.

Supplemental table 1: SYNGO Proteins with significant expression changes from 8 to 10 WPI in PB-AD neurons

Protein	new at WPI 10	log$_2$fc	FDR-adjusted p-value
CNTNAP4	yes	1.05	1.50E-26
SLITRK1	yes	0.97	7.75E-23
CDH1	yes	0.91	4.81E-20
EPHA7	yes	0.90	1.26E-19
ITGB3	no	0.80	1.72E-15
PRRT1	yes	0.74	1.35E-13
NPTX2	no	0.73	4.68E-13
ITGA2	no	0.67	3.46E-11
SLC17A5	yes	0.67	5.09E-11
KCNQ2	yes	0.65	1.77E-10
ELFN1	no	0.65	2.09E-10
CYTH1	no	0.65	2.21E-10
TNC	no	0.64	3.74E-10
NRGN	yes	0.63	6.71E-10
NGDN	yes	0.63	6.73E-10
GPR158	no	0.62	1.43E-09
RELN	yes	0.57	3.53E-08
SCG2	no	0.78	2.18E-03
TMEM163	no	0.62	7.64E-03
PTPRN	no	0.59	2.24E-02
NEFH	no	0.40	2.24E-02
IGSF8	no	0.40	2.24E-02
CNTNAP1	no	0.34	3.78E-02
ATP2B4	no	0.30	3.78E-02

1702

1703