A genome-wide MAGIC kit for recombinase-independent mosaic analysis in *Drosophila*

Yifan Shen[1], Ann T. Yeung[1,2], Payton Ditchfield[3], Elizabeth Korn, Rhiannon Clements, Xinchen

Chen[4], Bei Wang, Michael Sheen[5], Parker A. Jarman[6], and Chun Han*

Department of Molecular Biology and Genetics, Cornell University, Ithaca, New York, USA

Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, New York, USA

[1]These authors contributed equally.

[2]Current address: Harvard Medical School, Boston, MA 02115, USA

[3]Current address: Center for Community Independence, Revere, MA 02151, USA

[4]Current address: Department of Neurobiology, School of Biological Sciences, University of

California, San Diego, La Jolla, CA 92093, USA.

[5]Current address: Warren Alpert Medical School of Brown University, Providence, RI 02903,

USA

[6]Current address: Yale School of Medicine, New Haven, CT 06510, USA

*For correspondence: chun.han@cornell.edu (CH)

ABSTRACT

Mosaic analysis has been instrumental in advancing developmental and cell biology. Most current mosaic techniques rely on exogenous site-specific recombination sequences that need to be introduced into the genome, limiting their application. Mosaic analysis by gRNA-induced crossing-over (MAGIC) was recently developed in *Drosophila* to eliminate this requirement by inducing somatic recombination through CRISPR/Cas9-generated DNA double-strand breaks. However, MAGIC has not been widely adopted because gRNA-markers, a required component for this technique, are not yet available for most chromosomes. Here, we present a complete, genome-wide gRNA-marker kit that incorporates optimized designs for enhanced clone induction and more effective clone labeling in both positive MAGIC (pMAGIC) and negative MAGIC (nMAGIC). With this kit, we demonstrate clonal analysis in a broad range of *Drosophila* tissues, including cell types that have been difficult to analyze using recombinase-based systems. Notably, MAGIC enables clonal analysis of pericentromeric genes and deficiency chromosomes and in interspecific hybrid animals, opening new avenues for gene function study, rapid gene discovery, and understanding cellular basis of speciation. This MAGIC kit complements existing systems and makes mosaic analysis accessible to address a wider range of biological questions.

INTRODUCTION

Mosaic animals containing genetically distinct populations of cells in the same organism are useful for *in vivo* studies of complex biological processes. For this reason, techniques that can generate genetically labeled mosaic clones have been utilized in both vertebrates and invertebrates to study tissue-specific functions of pleiotropic genes, developmental timing, cell lineages, cell proliferation, neural wiring, and many other biological phenomena [1-3]. The most popular mosaic techniques rely on site-specific recombination systems, such as FRT/Flp [4, 5] and LoxP/Cre [6, 7], to induce somatic recombination between homologous chromosomes. Such techniques require the introduction of recombination sites to specific locations in the genome and thus cannot be applied to unmodified chromosomes. To overcome this limitation, we recently developed a recombinase-independent mosaic technique called mosaic analysis by gRNA-induced crossing-over (MAGIC) [8]. In MAGIC, the CRISPR/Cas9 system generates double-strand breaks (DSBs) at a predefined genome location to induce homologous recombination in precursor cells during S/G2 phase. Subsequent chromosomal segregation during mitosis can result in clones homozygous for the chromosomal segments distal to the crossover site. Two variants of this technique in *Drosophila*, positive MAGIC (pMAGIC) and negative MAGIC (nMAGIC), label the resulting homozygous clones by the presence and absence of fluorescent markers, respectively [8]. Like FRT/Flp-based techniques, MAGIC enables characterization of homozygous clones of lethal mutations in otherwise heterozygous animals [8].

However, unlike FRT/Flp-based techniques, MAGIC does not require prior genetic modification of the test chromosome. Thus, it can potentially be used on any chromosome and have much wider applications. Foremost, mutations of diverse natures have been established for most *Drosophila* genes [9-14], and thousands of deficiency strains harbor deletions that collectively uncover 98.4% of the *Drosophila* genome [15]. However, most of these mutant chromosomes cannot be analyzed by traditional mosaic techniques due to the lack of FRT sites or incompatibility with the FRT/Flp system. In contrast, MAGIC can theoretically be applied to these chromosomes as they are, allowing convenient genome-wide mosaic screens. In addition, genes located more proximal to centromeres than existing FRT sites cannot be analyzed by FRT/Flp techniques. In comparison, MAGIC can potentially be used to study these genes because the crossover site in MAGIC can be flexibly defined by users. Lastly, given that MAGIC is compatible with wild-derived chromosomes [8], it may be able to generate

homozygous clones of a chromosome derived from a single species in an interspecific hybrid

animal, allowing the study of species-related cell-cell interactions.

Despite these potentials, MAGIC has not been widely adopted by the *Drosophila*

community. A major barrier is the lack of gRNA-marker transgenes on most chromosomal arms,

which are necessary for DSB induction and fluorescent clone labeling [8]. In addition, the

existing gRNA-markers suffer from several limitations, including low frequency of clone

induction, weak labeling of pMAGIC clones, and suboptimal visualization of nMAGIC clones.

Because of these reasons, MAGIC has only been successfully applied to a few genes in a limited

number of *Drosophila* tissues [8, 16].

To overcome these limitations, here we first optimized gRNA-marker designs to improve

clonal induction, the brightness of pMAGIC clones, and visualization of nMAGIC clones. Then

we generated pMAGIC and nMAGIC gRNA-markers for all chromosomal arms and

characterized their ability to generate clones. Using this kit, we demonstrate mosaic analysis of

centromere-proximal genes, deficiency chromosomes, chromosomes derived from different

Drosophila species in interspecific hybrid animals. This kit allows optimal clone induction in

diverse cell and tissue types and should be useful for studying a wide range of biological

processes.

RESULTS

New gRNA-marker designs improve pMAGIC and nMAGIC

MAGIC relies on a gRNA-marker dual-transgene (Figure 1A) inserted in a specific

chromosomal arm to both induce and visualize clones homozygous for this arm [8]. The gRNA

part of the construct expresses two gRNAs ubiquitously to target a pericentromeric location on

this arm. Two gRNAs targeting two sequences that are close to each other, instead of a single

gRNA, are used to increase the probability of DSBs and thus the clone frequency. The marker

part in pMAGIC utilizes a ubiquitously expressed Gal80 to prevent Gal4-dependent labeling of

heterozygous and homozygous cells for the gRNA-marker, while allowing labeling of

homozygous cells that lose the gRNA-marker. In contrast, nMAGIC uses ubiquitously expressed

BFP, which results in brighter labeling of gRNA-marker homozygous cells, intermediate

labeling of gRNA-marker heterozygous cells, and lack of labeling of gRNA-marker-negative

homozygous cells.

We have previously developed gRNA-marker vectors for both pMAGIC and nMAGIC.

However, gRNA-markers made with these vectors exhibit some limitations. First, the clone

frequency can be low for certain gRNAs [8]. Second, the *ubi-Gal80* in pMAGIC gRNA-markers

does not completely suppress Gal4 activity in certain tissues, such as hemocytes. Third,

pMAGIC clones are sometimes too dim to visualize cell morphology, such as thin dendrite and

axon projections of neurons. Lastly, nMAGIC utilizes a nuclear BFP marker (nlsBFP), which

does not show the cell shape and sometimes cannot mark clones effectively. Thus, to optimize

MAGIC, we first sought to improve the gRNA-marker designs.

Since clone induction in MAGIC depends on gRNA-induced DNA DSBs, we asked

whether a more efficient gRNA design enhances clone frequency in somatic tissues. The

previous gRNA-marker vectors used a tgFE design, which contains the "F+E" gRNA scaffold

and a tRNA^{Gly} before each gRNA targeting sequence (Figure 1A) [8]. However, we have shown

that an improved Qtg2.1 design, which contains the gRNA2.1 scaffold and a single tRNA^{Gln}

spacer between the two gRNAs (Figure 1A), is much more mutagenic than tgFE in somatic

tissues [17]. We thus compared two pMAGIC gRNA-markers that are based on these two gRNA

designs but target the same genomic sequences at cytological band 40D2 to assess their ability to

induce clones in peripheral sensory neurons. When used with the same neuronal/epidermal

precursor Cas9, *zk-Cas9* [8], Qtg2.1 resulted in three times more neuronal clones as compared to

tgFE (Figure 1B), confirming a positive correlation between gRNA efficiency and clone

frequency.

To ensure more complete suppression of Gal4 activity by Gal80 in pMAGIC, we

replaced the *ubi* enhancer driving Gal80 expression with an *αTub84B* (*tub*) enhancer that was

used in *tub-Gal80* in the MARCM system [18]. When combined with the larval hemocyte

marker *pxn-Gal4 UAS-CD4-tdTom* [19] (Figure 1C), *ubi-Gal80* did not completely suppress the

labeling of hemocytes (Figure 1D). In contrast, no labeled hemocytes could be detected in the

presence of *tub-Gal80* (Figure 1E), suggesting that *tub-Gal80* is a better marker for pMAGIC.

To improve clone brightness in pMAGIC, we sought to destabilize Gal80 and reduce its

expression, reasoning that dim clones are due to prolonged Gal80 activity after clone induction.

In addition to replacing the *ubi* enhancer with the *tub* enhancer, we also introduced protein and

mRNA destabilization sequences (DE) [20, 21] at the 3' end of the Gal80 coding sequence, and

replaced the *His2Av* polyA by *SV40* polyA (Figure 1F), as the latter reduces transgene

expression [22]. By measuring the brightness of epidermal (Figure 1G) and neuronal (Figure 1H)

clones induced by *gRNA-40D2* and *gRNA-42A4*, we found that each of the three changes

improved clone brightness.

Lastly, to visualize cell shape in nMAGIC, we replaced nlsBFP with cytosolic BFP

tagged by 3X Hemagglutinin (HA), in addition to utilizing the *tub* enhancer. This new design

allowed us to better discern clone shapes in both the wing imaginal disc and the epidermis

(Figure 1J, 1J', and 1L), in contrast to the previous *ubi-nlsBFP* design (Figure 1I and 1K).

Thus, by altering the designs of the gRNAs and the Gal80 and BFP markers, we created

new vectors optimized for more robust applications of nMAGIC and pMAGIC.

A gRNA-marker kit is established for all four chromosomes of *Drosophila*

To enhance the utility of MAGIC in *Drosophila*, we generated complete sets of pMAGIC and

nMAGIC gRNA-markers for all four chromosomes (Figure 2A). To identify suitable gRNA

target sites, we analyzed the pericentromeric sequences of X, 2L, 2R, 3L, 3R, and 4, based on

three criteria [8]: conserved in closely related *Drosophila* species to minimize the chance of

single nucleotide polymorphism, (2) located away from functionally critical regions to avoid

disrupting essential processes, and (3) unique within the genome to minimize off-target effects.

For each MAGIC construct, we selected a pair of non-repetitive gRNA target sequences in

intergenic regions to maximize the chances of DSBs. These two sequences are closely linked to

minimize the risk of large deletions. Given the variable efficiency of gRNA target sequences, we

selected three pairs of gRNAs targeting three chromosomal locations for each chromosomal arm

and named them according to the corresponding cytoband (Table 1).

To evaluate the clone-induction properties of these gRNAs, we combined the pMAGIC

set with *zk-Cas9* and counted the number of neuronal clones in A1-A7 larval hemi-segments

(Figure 2B). As expected, the clone frequency varied from gRNA to gRNA, but we were able to

identify efficient gRNAs for every chromosomal arm. We previously found that different gRNAs

follow the same trend of relative efficiency in different tissues [8]. Here we additionally tested

nMAGIC gRNAs for 2R in the wing disc (Figure 2C) and noticed a similar trend of clone

induction to that of their pMAGIC counterparts in sensory neurons (Figure 2B), suggesting that

the results in sensory neurons are transferrable to other tissues.

Certain gRNAs (e.g. gRNA-40E1) exhibited very low clone frequency. Such gRNAs can

be useful for inducing sparse clones in highly packed issues such as the brain. For example,

using the same *ey-Cas9* [23], the highly efficient gRNA-40D2 induced too many neuronal clones

in the adult brain for morphological analysis (Figure 2D), while gRNA-40E1 gave rise to few

clones, whose projection patterns were much easier to analyze (Figure 2E).

Together, the pMAGIC and nMAGIC gRNA-marker lines constitute a complete kit

(Table 1) for genome-wide MAGIC applications in *Drosophila*.

MAGIC allows clonal analysis in diverse tissues and cell types

To determine if MAGIC can be applied to diverse tissues in *Drosophila*, we conducted clonal

analysis in the larva using *gRNA-40D2(Gal80)* and several tissue-specific Cas9s. With the

ubiquitous *vas-cas9* [24] and *tub-Gal4 UAS-mCD8-GFP*, we readily detected clones in the larval

brain (Figure 3A), proliferating tissues like eye and leg discs (Figure 3B-C), and polyploid

tissues like the fat body, gut, and trachea (Figure 3D-F). Clone induction in polyploid tissues

suggests that crossing-over events occurred before the last cell division. Using *zk-cas9* and

R38F11-Gal4>UAS-tdTomato (tdTom), we observed frequent epidermal clones (Figure 3G).

Using the glia precursor *gcm-Cas9* and *repo-Gal4>UAS-mCD8-GFP*, we detected individual

glial clones in the brain (Figure 3H). Lastly, the new pMAGIC gRNA-marker design allowed us

to reliably induce hemocyte clones (Figure 3I). These results collectively demonstrate MAGIC's

efficacy in generating clones in diverse *Drosophila* tissues, indicating its value in studying gene

function and cell lineage in various tissue contexts.

The neuromuscular junction (NMJ) in *Drosophila* larvae has been a valuable model for

elucidating synaptic biology, owing to its simplicity, accessibility, and conserved features shared

with mammalian excitatory synapses [25-27]. Despite its popularity, the NMJ has rarely been

analyzed by the MARCM technique in the literature, likely due to the difficulty of inducing

clones in motor neurons. To test the effectiveness of MAGIC in analyzing gene function at the

NMJ, we selected two genes crucial for synaptic function and construction: *Vesicular glutamate*

transporter (*VGlut*) [28] and *bruchpilot* (*brp*) [29, 30], null mutations of each of which exhibit

recessive lethality in larvae. Combined with appropriate gRNA(Gal80) lines (42A4 for *vGluT* on

2R and 40D2 for *brp* on 2L) and *tub-Gal4 UAS-CD8-GFP*, *zk-Cas9* induced frequent clones in

type Ib boutons (Figure 4A-B"). The loss of vGluT or Brp specifically in GFP-labeled NMJs was

confirmed by immunostaining. These results exemplify the value of pMAGIC for dissecting

gene function in NMJ biology at the single cell level.

MAGIC enables clonal analysis of pericentromeric genes, 4th chromosome-associated

mutations, and in interspecific hybrid animals

Because all FRT sites in the existing FRT/Flp mosaic system are located at some distances away

from centromeres, it has not been possible to study genes located between the FRT sites and the

corresponding centromeres by clonal analysis. In contrast, the gRNA target site in a MAGIC

experiment can be user-selected to enable clonal analysis of any given gene. To illustrate this

potential, we used *gRNA-41F9* to induce homozygous mutant clones of *Ecdysone receptor*

(*EcR*), which is located at 42A10 and is inaccessible to all FRT sites on 2R. EcR is a

transcription factor required for neuronal remodeling during metamorphosis [31]. In peripheral

sensory neurons, EcR activation at the beginning of pupariation causes apoptosis of some

dendritic arborization (da) neurons while triggering dendrite pruning of other da neurons [32]. As

expected, a wildtype (WT) pMAGIC clone of class IV da (C4da) neuron exhibited complete

dendrite pruning at 16 h after puparium formation (APF), accompanied by dendrite debris

phagocytosed into epidermal cells [19] (Figure 5A). In contrast, *EcR* mutant da neuron clones

still maintained larval dendritic arbors at 16 h APF (Figures 5B-D), instead of dying (in the case

of C3da) or undergoing dendrite pruning (in the cases of C1da and C4da).

202   Mosaic analysis of genes located on the 4th chromosome had not been possible until

recent introduction of FRT sites onto this chromosome by CRISPR-mediated knock-in [33].

Despite these advances, existing mutations on FRT-lacking 4th chromosomes still cannot be

analyzed by the FRT/Flp system. A valuable gene-disruption resource in *Drosophila* is the

deficiency library consisting of strains harboring chromosomal deletions. MAGIC can

potentially be used in conjunction with the deficiency library to screen for genes important in a

particular biological process. To test the potential of MAGIC for analyzing mutations on the 4th

chromosome and for gene discovery with deficiencies, we generated pMAGIC C4da clones of

Df(4)ED6380, a deficiency that deletes a segment between cytological bands 102B7 and 102D5.

These clones show dramatically reduced dendrites compared to the WT controls (Figure 4E-4G),

indicating the existence of genes important for dendrite growth in this region.

When examining nMAGIC gRNA-markers for the 4th chromosome, we noticed uneven

expression of *tub-3xHA-BFP* in epidermal cells and some imaginal tissues, exemplified by cells

lacking detectable BFP signal (Figures S1A,S1B). This variegated expression is likely due to

transgenes residing in heterochromatin and is common for transgenes located on the 4th

chromosome [34]. While this uneven expression limits the usefulness of our gRNA-markers in

the corresponding epithelial tissues, Gal80 in pMAGIC gRNA-markers efficiently suppressed

Gal4 activities in most neurons (Figure S1C), conforming their effectiveness in neuronal

MAGIC analysis.

Lastly, we wondered if the MAGIC system can generate clones in interspecific hybrid

animals derived from *D. melanogaster* and *D. simulans* parents, given that the two species show

a large degree of synteny [35]. The clones in these animals would contain homozygous

chromosomal arms derived from a single species. To test this idea, we crossed *D. melanogaster*

females containing *gRNA-42A4(BFP); hh-cas9* to *D. simulans* males carrying a loss of function

mutation of *Lethal hybrid rescue mutation* (*Lhr*), which ensures the viability of the hybrid male

progeny [36]. Indeed, we observed twin spots consisting of dark clones, containing only *D.*

simulans 2R, and brighter clones, containing only *D. melanogaster* 2R, in wing discs of both

female and male progeny (Figure 5G and 5H), demonstrating the feasibility of studying species-

specific alleles in cell-cell interactions in interspecific hybrids.

DISCUSSION

Conceptually, MAGIC is a simpler and more convenient mosaic technique compared to

traditional recombinase-dependent methods. However, the lack of gRNA-marker transgenes has

prevented its wide application in *Drosophila*. In this study, we present a complete gRNA-marker

kit that enables genome-wide MAGIC in *Drosophila*, removing this bottleneck. The new gRNA-

markers incorporate optimized designs that improve clone frequency and labeling in both

pMAGIC and nMAGIC. We further demonstrate the compatibility of MAGIC with broad tissues

and cell types. More importantly, MAGIC enables mosaic analyses that could not be

accomplished by existing FRT/Flp systems, such as those of pericentromeric genes, deficiency

chromosomes, and interspecific homologous chromosomes. Thus, this MAGIC kit provides

Drosophila researchers with greater flexibility for conducting mosaic analyses of diverse

purposes.

To implement MAGIC, one needs to first choose a proper Cas9 that is expressed in the

precursor cells of the targeted cell population. We show that ubiquitously expressed Cas9 lines,

such as *vas-Cas9* [24] and *Act5C-Cas9* [37], are sufficient to induce clones in broad tissues.

Alternatively, heat shock (HS)-induced Cas9 [38] can offer temporal control of clone induction

in most tissues, akin to the HS-Flp commonly used in FRT-based mosaic analysis. The third

option is a Cas9 driven by a tissue-specific enhancer specifically in the precursor cells of the

target tissues. Examples shown in this study include *ey-Cas9* expressed in many neuronal

lineages [23], *gcm-Cas9* expressed in glial progenitor cells [39], *ZK-Cas9* expressed in precursor

cells of epidermal cells, motor neurons, and somatosensory neurons [17], and *hh-Cas9* expressed

in imaginal tissues [40]. These Cas9 lines have the advantages of being more specific and

efficient, and more convenient to use than ubiquitous or inducible ones. Multiple strategies,

including enhancer fusion [37, 40], Gal4-to-Cas9 conversion [17, 41], and insertion of Cas9 in

specific gene loci [39], have been developed to ease the generation of such tissue-specific Cas9

lines. As an ongoing effort, we have been converting Gal4 lines known to be expressed in

progenitor cells into Cas9 (available at

https://bdsc.indiana.edu/stocks/genome_editing/crispr_cas9.html), in the hope of making

MAGIC accessible to more *Drosophila* tissues.

The second component of MAGIC is a gRNA-marker line that resides on the appropriate

chromosome arm and targets Cas9 to cut a pericentromeric site on the same arm. In the gRNA-

marker kit, we generated three lines targeting different sites for each chromosome arm. These

lines exhibit a broad range of clone frequencies in larval sensory neurons. As different gRNA-

markers maintain similar relative efficiencies of clone induction across tissues [8], one can

choose a gRNA-marker more appropriate for their applications based on the target gene location

and the desired clone frequency. Notably, a higher clone frequency may not always be

beneficial, such as when studying the projection patterns of individual neurons in the brain. A

low efficiency gRNA-marker in this kit may be more desirable in such applications.

Besides commonly conducted mosaic analysis in *Drosophila*, MAGIC also enables many

novel analyses that are difficult or impossible to accomplish with traditional systems. One

example is to analyze interactions among species-specific cells in an interspecific hybrid animal

for understanding the cell biological basis of hybrid incompatibility. Interspecific crosses

between WT *D. melanogaster* and WT *D. simulans* result in sex-specific lethality of F1 progeny

[36]. The cellular basis of this lethality is poorly understood. With MAGIC, one may generate
mixed cell populations that are homozygous or heterozygous for species-specific alleles in the
same hybrid embryo or larva, in which the relative fitness of the three cell populations can be
compared. Similarly, MAGIC could complement genome-wide association studies (GWAS)
using wild-derived isogenic strains and provide much more mechanistic detail. Such strains as
those in the *Drosophila* Genetic Reference Panel (DGRP) [42] have been very useful for
discovering loci that are responsible for phenotypic variations. MAGIC can be combined with
these strains to analyze the effects of homozygosity of specific variants at the cell biology level
[8]. In addition, MAGIC can be combined with the *Drosophila* deficiency kit for rapid genome-
wide mosaic screens. A deficiency exhibiting a desired phenotype in such screens can be further
dissected with smaller deficiencies within the deleted region or existing mutations in candidate
genes. In this way, the convenience of MAGIC could significantly accelerate phenotype-based
gene discovery. Lastly, because the crossover site in MAGIC can be user-defined, one can easily
generate gRNA-markers that induce somatic recombination at specific genome locations. Such
flexibility could be useful for mapping undefined genetic loci that are responsible for certain
phenotypes, especially in non-traditional model organisms.

MATERIALS AND METHODS:

Fly Stocks and Husbandry

See the Key Resource Table (Table S1) for details of fly stocks used in this study. Most fly lines
were either generated in the Han lab or obtained from the Bloomington *Drosophila* Stock Center.
lethal hybrid rescue (*lhr*) mutant *D. Simulans* was a gift from Dr. Dan Barbash. All flies were
grown on standard yeast-glucose medium, in a 12:12 light/dark cycle, at 25°C unless otherwise
noted. Virgin males and females for mating experiments were aged for 3-5 days.

To generate and label pMAGIC clones in larval peripheral sensory neurons, we used
either *RabX4-Gal4 UAS-MApHS* (for gRNA-markers on chromosomes X, II, and IV) or *21-7-*
Gal4 UAS-MApHS (for gRNA-markers on chromosome III) combined with *zk-cas9*. To count
peripheral sensory neuronal clones on the larval body wall, third instar larvae were mounted
laterally on slides and then counted in segment A1-A7 under a Nikon SMZ18 stereomicroscope.
pMAGIC clones were induced and labeled by *RabX4-Gal4 UAS-MApHS* combined with *ey-cas9*
in the fly adult brain, by *tub-Gal4 UAS-mCD8-GFP* combined with *vas-cas9* in larval brains,

imaginal discs, fat bodies, guts, and trachea, by *repo-Gal4 UAS-mCD8-GFP* combined with

gcm-cas9 in glia, by *pxn-Gal4 UAS-tdTom* combined with *Act-cas9* in hemocytes, by *tub-Gal4*

UAS-mCD8-GFP combined with *zk-cas9* in larval motor neurons, by *R38F11-Gal4 UAS-tdTom*

combined with *zk-cas9* in the larval epidermis. To induce nMAGIC clones in wing imaginal

discs of interspecific hybrid animals, we use *gRNA-42A4(BFP)*; *hh-cas9* virgin females of *D.*

melanogaster to cross with *Lhr[1]* [43] *D. simulans* males.

Molecular Cloning

MAGIC gRNA cloning vectors: pMAGIC gRNA-marker vectors constructed in this study include

pAC-U63-gRNA2.1-ubiGal80(DE)-His2Av, pAC-U63-gRNA2.1-tubGal80(DE)-His2Av and

pAC-U63-gRNA2.1-tubGal80(DE)-SV40. To make pAC-U63-gRNA2.1-ubiGal80(DE)-His2Av,

a fragment containing gRNA2.1 scaffold (gRNA2.1) and U6 3' flanking sequence (U63fl) was

first used to replace gRNA2.1-QtRNA-Gal80(TS)-gRNA2.1-U63fl in pAC-U63-QtgRNA2.1-8R

(Addgene 170514) [17]. Then a destabilization sequence (DE) containing a GS linker, amino

acids (AAs) 422-461 of mouse ornithine decarboxylase [21], 2X RNA destabilizing nonamer

(TTATTTATTgatccTTATTTATT) [21] was added to the C-terminus of Gal80 in frame. To

make pAC-U63-gRNA2.1-tubGal80(DE)-His2Av, a 2.6 kb *tub* enhancer was amplified from

pENTR221-tubP [16] by PCR and used to replace the *ubi* enhancer in pAC-U63-gRNA2.1-

ubiGal80(DE)-His2Av. To make pAC-U63-gRNA2.1-tubGal80(DE)-SV40, a SV40 polyA

sequence was amplified from pAPIC-PHCS [22] and used to replace the His2Av polyA in pAC-

U63-gRNA2.1-ubiGal80(DE)-His2Av. To make the nMAGIC gRNA-marker vector pAC-U63-

gRNA2.1-tubBFP(HA), the gRNA2.1-U63fl fragment was first used to replace gRNA2.1-

QtRNA-BFP(TS)-gRNA2.1-U63fl in pAC-U63-QtgRNA2.1-BR (Addgene 170513) [17]. The

tub enhancer was then used to replace the *ubi* enhancer. Lastly, a synthetic DNA fragment

(GenScript) encoding 3X HA was used to replace the nuclear localization signal at the N-

terminus of mTagBFP. Cloning was carried out by ligation with T4 ligase or NEBuilder DNA

Assembly reactions (New England Biolabs Inc.).

MAGIC gRNA expression vectors: 33 gRNA expression vectors were constructed with the

corresponding gRNA cloning vectors as listed in Table 1 according to published protocols [17].

Briefly, for each expression vector, two primers containing appropriate gRNA target sequences

(Table S2) were used to amplify a fragment consisting of 3' end of U6:3 promotor, the first

target sequence (TS1), gRNA2.1, tRNA^Q, the second target sequence (TS2), and the beginning
sequence of gRNA2.1 with pAC-U63-QtgRNA2.1-BR as the PCR template. The PCR product
was then assembled with SapI-digested gRNA cloning vectors using NEBuilder DNA Assembly.

Injections were carried out by Rainbow Transgenic Flies (Camarillo, CA 93012 USA) or
Genetivision (Stafford, TX 77477) to transform flies through φC31 integrase-mediated
integration into attP docker sites. The 3xP3-RFP selection marker in gRNA-markers inserted at
the attP[ZH-102D] site were removed by crossing to Cre.

Live Imaging

Live imaging of larval epidermal cells, sensory neurons, and hemocytes was performed as
previously described [44]. Animals were collected at 96 (for late third larvae) or 120 hours AEL
(for wandering third instar larvae) hours after egg-laying (AEL) and mounted in glycerol on a
slide with vacuum grease as a spacer. Animals were imaged using a Leica SP8 confocal
microscope with a 40X NA1.3 oil objective, pinhole size 2 airy units, and a z-step size of 1 μm.
For epidermis, images were taken at the dorsal midline of A2 and A3 segments. For dendritic
arborization neurons, images were taken from A1 to A7 hemi-segments.

For imaging sensory neurons in pupae, newly formed pupae were collected and incubated
at 25 °C. After 16 hours, the pupal cases were carefully removed, and the pupae were mounted
dorsal side up on slides with halocarbon oil beneath a coverslip. Double-sided tape was used as a
spacer. The mounted pupae were imaged using a Leica SP8 confocal microscope with a 40X
NA1.3 oil objective.

Imaging of wing discs and other larval tissues

Larval dissections were performed as described previously [40]. Briefly, wandering third instar
larvae were dissected in a small petri dish filled with cold phosphate-buffered saline (PBS). The
anterior half of the larva was inverted. To prepare imaginal discs, trachea and gut were removed.
Samples were then transferred to 4% formaldehyde in PBS and fixed for 20 minutes at room
temperature. After washing with PBS, the tissues were stained in DAPI (1:1000) in 0.2% PBST
(PBS with 0.2% Triton X-100) for 5 minutes. The tissues were washed again in PBST and
mounted in SlowFade Diamond Antifade Mountant (Thermo Fisher Scientific) on a glass slide.

A coverslip was lightly pressed on top. Imaginal discs were imaged using a Leica SP8 confocal

microscope with a 20X NA0.8 oil objective.

Adult brain imaging

365   Flies were aged for 1 day after eclosion. Brains were dissected in PBS at room temperature and

366   then fixed in 4% paraformaldehyde in PBS with constant circular rotation for 20 minutes at room

367   temperature. The brains were subsequently washed in 0.2% PBST and mounted on a glass slide

368   under a glass coverslip. Vacuum grease was used as a spacer between the coverslip and the slide.

369   Brains were imaged using a Leica SP8 confocal microscope with a 40X NA1.3 oil objective.

Larval fillet preparation

Larval fillet dissection was performed on a petri dish half-filled with PMDS gel. Wandering third

instar larvae were pinned on the dish in PBS dorsal-side up and then dissected to expand the

body wall. PBS was then removed and 4% formaldehyde in PBS was added to fix larvae for 15

minutes at room temperature. For VGlut staining, the fillets were fixed in Bouin's solution for 5

minutes at room temperature. Fillets were rinsed and then washed at room temperature in PBS

for 20 minutes or until the yellow color from Bouin's solution faded. After immunostaining, the

head and tail of fillets were removed, and the remaining fillets were placed in SlowFade

Diamond Antifade Mountant on a glass slide. A coverslip was lightly pressed on top. Larval

fillets were imaged using a Leica SP8 confocal microscope with a 40X NA1.3 oil objective.

Immunohistochemistry

Larval brains and larval fillets were rinsed and washed at room temperature in 0.2% PBST after

fixation. The samples were then blocked in PBST with 5% normal donkey serum (NDS) for 1

hour before incubating with appropriate primary antibodies in the blocking solution. Brains were

stained for 2 hours at room temperature, and fillets were stained overnight at 4°C. After

additional rinsing and washing, the samples were incubated with secondary antibodies for 2

hours at room temperature. The samples were then rinsed and washed again before mounting and

imaging. Primary antibodies used in this study are mouse anti-Repo antibody 8D12 (1:50

dilution), mouse anti-Brp antibody nc82 (1:100 dilution), rabbit anti-VGlut (1:200 dilution) [39],

mouse anti-HA antibody (12CA5, 1:100), and goat anti-HRP conjugated with Cy3 (1:200).

Secondary antibodies include donkey anti-mouse antibody conjugated with Cy5 (1:400) and donkey anti-rabbit antibody conjugated with Cy5 (1:400).

Image Analysis and Quantification

ImageJ analyses were conducted in Fiji/ImageJ. To compare brightness of clones in sensory neurons and epidermal cells, clones were detected based on thresholds to generate masks. The clone brightness within the masks was then measured as mean pixel intensity. To quantify clones in wing discs, 2-3 slices of optical sections in the middle of the disc were projected into a 2-dimensional (2D) image. Clones were detected based on a fixed threshold to generate masks, and the total area of clones in the wing pouch of each disc was measured.

The tracing and measurement of da neuron dendrites were done as previously described in detail [44]. Briefly, dendrites were segmented using local thresholding. The segments were then converted into single-pixel-width skeletons. The total length of skeletons was calculated based on pixel distance. Normalized dendrite length was calculated as dendritic length (μm)/segment width (μm).

Statistical Analysis

One-way analysis of variance (ANOVA) with Tukey's honest significant difference (HSD) test was used when dependent variable was normally distributed and there was approximately equal variance across groups. Student's t-test was used when two groups are compared. For additional information on the number of samples, see figure legends. R studio was used for all statistical analyses.

ACKNOWLEDGMENTS

We thank Dion Dickman, Marcus Smolka, and Developmental Studies Hybridoma Bank (DSHB) for antibodies; Dan Barbash, Ethan Bier, and Bloomington *Drosophila* Stock Center for fly stocks; Dan Barbash and Tianzhu Xiong for advice on interspecific crosses; Claire Ho and Christina Breneman for helping to establish and test transgenes; Dan Barbash and Mariana Wolfner for feedback on the manuscript.

AUTHOR CONTRIBUTIONS

Conceptualization: Chun Han, Yifan Shen, Ann Yeung

Data curation: Yifan Shen, Ann Yeung, Elizabeth Korn, Rhiannon Clements, Parker Jarman

Formal analysis: Yifan Shen, Ann Yeung, Rhiannon Clements

Funding acquisition: Chun Han

Investigation: Yifan Shen, Ann Yeung, Elizabeth Korn, Rhiannon Clements, Xinchen Chen,
Michael Sheen, Parker Jarman,

Methodology: Chun Han, Yifan Shen, Ann Yeung

Supervision: Chun Han, Yifan Shen, Ann Yeung, Rhiannon Clements, Elizabeth Korn

Resources: Yifan Shen, Payton Ditchfield, Bei Wang

Validation: Yifan Shen

Visualization: Yifan Shen, Ann Yeung, Rhiannon Clements

Writing – original draft: Chun Han, Yifan Shen

Writing – review & editing: Chun Han, Yifan Shen

REFERENCES
1.   Germani F, Bergantinos C, Johnston LA. Mosaic Analysis in Drosophila. Genetics.
2018;208:4733-490. doi: 10.1534/genetics.117.300256.

2.   Xu T, Rubin GM. The effort to make mosaic analysis a household tool. Development.
2012;139(24):4501-3. doi: 10.1242/dev.085183. PubMed PMID: 23172911; PubMed Central
PMCID: PMCPMC3509720.

3.   Griffin R, Binari R, Perrimon N. Genetic odyssey to generate marked clones in
Drosophila mosaics. Proc Natl Acad Sci U S A. 2014;111(13):4756-63. Epub 20140312. doi:
10.1073/pnas.1403218111. PubMed PMID: 24623854; PubMed Central PMCID:
PMCPMC3977257.

4.   Golic KG, Lindquist S. The FLP recombinase of yeast catalyzes site-specific
recombination in the Drosophila genome. Cell. 1989;59(3):499-509. doi: 10.1016/0092-
8674(89)90033-0. PubMed PMID: 2509077.

5.   Xu T, Rubin GM. Analysis of genetic mosaics in developing and adult Drosophila tissues.
Development. 1993;117(4):1223-37. doi: 10.1242/dev.117.4.1223. PubMed PMID: 8404527.

6.   Henner A, Ventura PB, Jiang Y, Zong H. MADM-ML, a mouse genetic mosaic system
with increased clonal efficiency. PLoS One. 2013;8(10):e77672. Epub 20131015. doi:
10.1371/journal.pone.0077672. PubMed PMID: 24143253; PubMed Central PMCID:
PMCPMC3797059.

7.   Zong H, Espinosa JS, Su HH, Muzumdar MD, Luo L. Mosaic Analysis with Double
Markers in Mice. Cell. 2005;121(3):479-92.

8.   Allen SE, Koreman GT, Sarkar A, Wang B, Wolfner MF, Han C. Versatile CRISPR/Cas9-
mediated mosaic analysis by gRNA-induced crossing-over for unmodified genomes. PLoS Biol.
2021;19(1):e3001061. Epub 20210114. doi: 10.1371/journal.pbio.3001061. PubMed PMID:
33444322; PubMed Central PMCID: PMCPMC7837743.

9.   Thibault ST, Singer MA, Miyazaki WY, Milash B, Dompe NA, Singh CM, et al. A
complementary transposon tool kit for Drosophila melanogaster using P and piggyBac. Nat
Genet. 2004;36(3):283-7. Epub 2004/02/26. doi: 10.1038/ng1314. PubMed PMID: 14981521.

10.   Hacker U, Nystedt S, Barmchi MP, Horn C, Wimmer EA. piggyBac-based insertional
mutagenesis in the presence of stably integrated P elements in Drosophila. Proc Natl Acad Sci U
S A. 2003;100(13):7720-5. Epub 2003/06/13. doi: 10.1073/pnas.1230526100. PubMed PMID:
12802016; PubMed Central PMCID: PMCPMC164654.

11.   Bellen HJ, Levis RW, He Y, Carlson JW, Evans-Holm M, Bae E, et al. The Drosophila
gene disruption project: progress using transposons with distinctive site specificities. Genetics.
2011;188(3):731-43. Epub 2011/04/26. doi: 10.1534/genetics.111.126995. PubMed PMID:
21515576; PubMed Central PMCID: PMCPMC3176542.

12.   Staudt N, Molitor A, Somogyi K, Mata J, Curado S, Eulenberg K, et al. Gain-of-function
screen for genes that affect Drosophila muscle pattern formation. PLoS Genet. 2005;1(4):e55.
Epub 2005/10/29. doi: 10.1371/journal.pgen.0010055. PubMed PMID: 16254604; PubMed
Central PMCID: PMCPMC1270011.

13.   Bellen HJ, Levis RW, Liao G, He Y, Carlson JW, Tsang G, et al. The BDGP gene
disruption project: single transposon insertions associated with 40% of Drosophila genes.
Genetics. 2004;167(2):761-81. Epub 2004/07/09. doi: 10.1534/genetics.104.026427. PubMed
PMID: 15238527; PubMed Central PMCID: PMCPMC1470905.

14.   Yamamoto S, Jaiswal M, Charng WL, Gambin T, Karaca E, Mirzaa G, et al. A drosophila
genetic resource of mutants to study mechanisms underlying human genetic diseases. Cell.
2014;159(1):200-14. Epub 2014/09/27. doi: 10.1016/j.cell.2014.09.002. PubMed PMID:
25259927; PubMed Central PMCID: PMCPMC4298142.

15.   Cook RK, Christensen SJ, Deal JA, Coburn RA, Deal ME, Gresens JM, et al. The
generation of chromosomal deletions to provide extensive coverage and subdivision of the
Drosophila melanogaster genome. Genome Biol. 2012;13(3):R21. Epub 2012/03/27. doi:
10.1186/gb-2012-13-3-r21. PubMed PMID: 22445104; PubMed Central PMCID:
PMCPMC3439972.

16.   Chen X, Wang B, Sarkar A, Huang Z, Ruiz NV, Yeung AT, et al. Phagocytosis-driven
neurodegeneration through opposing roles of an ABC transporter in neurons and phagocytes. Sci
Adv. 2025;11(11):eadr5448. Epub 20250312. doi: 10.1126/sciadv.adr5448. PubMed PMID:
40073145; PubMed Central PMCID: PMCPMC11900885.

17.   Koreman GT, Xu Y, Hu Q, Zhang Z, Allen SE, Wolfner MF, et al. Upgraded
CRISPR/Cas9 tools for tissue-specific mutagenesis in Drosophila. Proc Natl Acad Sci U S A.
2021;118(14). Epub 2021/03/31. doi: 10.1073/pnas.2014255118. PubMed PMID: 33782117;
PubMed Central PMCID: PMCPMC8040800.

18.   Lee T, Luo L. Mosaic analysis with a repressible cell marker for studies of gene function
in neuronal morphogenesis. Neuron. 1999;22(3):451-61. doi: 10.1016/s0896-6273(00)80701-1.
PubMed PMID: 10197526.

19.   Han C, Song Y, Xiao H, Wang D, Franc NC, Jan LY, et al. Epidermal cells are the
primary phagocytes in the fragmentation and clearance of degenerating dendrites in Drosophila.
Neuron. 2014;81(3):544-60. Epub 2014/01/15. doi: 10.1016/j.neuron.2013.11.021. PubMed
PMID: 24412417; PubMed Central PMCID: PMCPMC3995171.

20.   Li X, Zhao X, Fang Y, Jiang X, Duong T, Fan C, et al. Generation of destabilized green
fluorescent protein as a transcription reporter. J Biol Chem. 1998;273(52):34970-5. doi:
10.1074/jbc.273.52.34970. PubMed PMID: 9857028.

21.   Zubiaga AM, Belasco JG, Greenberg ME. The nonamer UUAUUUAUU is the key AU-
rich sequence motif that mediates mRNA degradation. Mol Cell Biol. 1995;15(4):2219-30. doi:
10.1128/MCB.15.4.2219. PubMed PMID: 7891716; PubMed Central PMCID:
PMCPMC230450.

22.   Han C, Jan LY, Jan YN. Enhancer-driven membrane markers for analysis of
nonautonomous mechanisms reveal neuron-glia interactions in Drosophila. Proc Natl Acad Sci U
S A. 2011;108(23):9673-8. Epub 2011/05/25. doi: 10.1073/pnas.1106386108. PubMed PMID:
21606367; PubMed Central PMCID: PMCPMC3111288.

23.   Ji H, Sapar ML, Sarkar A, Wang B, Han C. Phagocytosis and self-destruction break down
dendrites of Drosophila sensory neurons at distinct steps of Wallerian degeneration. Proc Natl
Acad Sci U S A. 2022;119(4). Epub 2022/01/22. doi: 10.1073/pnas.2111818119. PubMed PMID:
35058357; PubMed Central PMCID: PMCPMC8795528.

24.   Lopez Del Amo V, Juste SS, Gantz VM. A nickase Cas9 gene-drive system promotes
super-Mendelian inheritance in Drosophila. Cell Rep. 2022;39(8):110843. doi:
10.1016/j.celrep.2022.110843. PubMed PMID: 35613590; PubMed Central PMCID:
PMCPMC9190248.

25.   Charng WL, Yamamoto S, Bellen HJ. Shared mechanisms between Drosophila peripheral
nervous system development and human neurodegenerative diseases. Curr Opin Neurobiol.
2014;27:158-64. Epub 20140422. doi: 10.1016/j.conb.2014.03.001. PubMed PMID: 24762652;
PubMed Central PMCID: PMCPMC4122633.

26.   Menon KP, Carrillo RA, Zinn K. Development and plasticity of the Drosophila larval
neuromuscular junction. Wiley Interdiscip Rev Dev Biol. 2013;2(5):647-70. Epub 20130205.
doi: 10.1002/wdev.108. PubMed PMID: 24014452; PubMed Central PMCID:
PMCPMC3767937.

27.   Chou VT, Johnson SA, Van Vactor D. Synapse development and maturation at the
drosophila neuromuscular junction. Neural Dev. 2020;15(1):11. Epub 20200802. doi:
10.1186/s13064-020-00147-5. PubMed PMID: 32741370; PubMed Central PMCID:
PMCPMC7397595.

28.   Daniels RW, Collins CA, Gelfand MV, Dant J, Brooks ES, Krantz DE, et al. Increased
expression of the Drosophila vesicular glutamate transporter leads to excess glutamate release
and a compensatory decrease in quantal content. J Neurosci. 2004;24(46):10466-74. doi:

10.1523/JNEUROSCI.3001-04.2004. PubMed PMID: 15548661; PubMed Central PMCID: PMCPMC6730318.

29.   Kittel RJ, Wichmann C, Rasse TM, Fouquet W, Schmidt M, Schmid A, et al. Bruchpilot promotes active zone assembly, Ca2+ channel clustering, and vesicle release. Science. 2006;312(5776):1051-4. Epub 20060413. doi: 10.1126/science.1126308. PubMed PMID: 16614170.

30.   Wagh DA, Rasse TM, Asan E, Hofbauer A, Schwenkert I, Durrbeck H, et al. Bruchpilot, a protein with homology to ELKS/CAST, is required for structural integrity and function of synaptic active zones in Drosophila. Neuron. 2006;49(6):833-44. doi: 10.1016/j.neuron.2006.02.008. PubMed PMID: 16543132.

31.   Brown HL, Cherbas L, Cherbas P, Truman JW. Use of time-lapse imaging and dominant negative receptors to dissect the steroid receptor control of neuronal remodeling in Drosophila. Development. 2006;133(2):275-85. Epub 20051214. doi: 10.1242/dev.02191. PubMed PMID: 16354717.

32.   Williams DW, Truman JW. Cellular mechanisms of dendrite pruning in Drosophila: insights from in vivo time-lapse of remodeling dendritic arborizing sensory neurons. Development. 2005;132(16):3631-42. Epub 20050720. doi: 10.1242/dev.01928. PubMed PMID: 16033801.

33.   Goldsmith SL, Shimell M, Tauscher P, Daly SM, Shimmi O, O'Connor MB, et al. New resources for the Drosophila 4th chromosome: FRT101F enabled mitotic clones and Bloom syndrome helicase enabled meiotic recombination. G3 (Bethesda). 2022;12(4). doi: 10.1093/g3journal/jkac019. PubMed PMID: 35084488; PubMed Central PMCID: PMCPMC8982423.

34.   Riddle NC, Minoda A, Kharchenko PV, Alekseyenko AA, Schwartz YB, Tolstorukov MY, et al. Plasticity in patterns of histone modifications and chromosomal proteins in Drosophila heterochromatin. Genome Res. 2011;21(2):147-63. Epub 20101222. doi: 10.1101/gr.110098.110. PubMed PMID: 21177972; PubMed Central PMCID: PMCPMC3032919.

35.   Chakraborty M, Chang CH, Khost DE, Vedanayagam J, Adrion JR, Liao Y, et al. Evolution of genome structure in the Drosophila simulans species complex. Genome Res. 2021;31(3):380-96. Epub 20210209. doi: 10.1101/gr.263442.120. PubMed PMID: 33563718; PubMed Central PMCID: PMCPMC7919458.

36.   Barbash DA. Ninety years of Drosophila melanogaster hybrids. Genetics. 2010;186(1):1-8. doi: 10.1534/genetics.110.121459. PubMed PMID: 20855573; PubMed Central PMCID: PMCPMC2940278.

37.   Port F, Chen HM, Lee T, Bullock SL. Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in Drosophila. PNAS. 2014;111(29):E2967-E76. doi: 10.1073/pnas.1405500111.

38.   Garcia-Marques J, Espinosa-Medina I, Ku K-Y, Yang C-P, Koyama M, Yu H-H, et al. A programmable sequence of reporters for lineage analysis. Nature Neuroscience. 2020. doi: 10.1038/s41593-020-0676-9.

39.   Chen X, Perry S, Fan Z, Wang B, Loxterkamp E, Wang S, et al. Tissue-specific knockout in the Drosophila neuromuscular system reveals ESCRT's role in formation of synapse-derived

extracellular vesicles. PLoS Genet. 2024;20(10):e1011438. Epub 20241010. doi: 10.1371/journal.pgen.1011438. PubMed PMID: 39388480.

40.   Poe AR, Wang B, Sapar ML, Ji H, Li K, Onabajo T, et al. Robust CRISPR/Cas9-Mediated Tissue-Specific Mutagenesis Reveals Gene Redundancy and Perdurance in Drosophila. Genetics. 2019;211(2):459-72. Epub 2018/12/07. doi: 10.1534/genetics.118.301736. PubMed PMID: 30504366; PubMed Central PMCID: PMCPMC6366929.

41.   Chen HM, Yao X, Ren Q, Chang CC, Liu LY, Miyares RL, et al. Enhanced Golic+: highly effective CRISPR gene targeting and transgene HACKing in Drosophila. Development. 2020;147(11). Epub 20200611. doi: 10.1242/dev.181974. PubMed PMID: 32467238.

42.   MacKay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The Drosophila melanogaster Genetic Reference Panel. Nature. 2012;482(7384):173-8. doi: 10.1038/nature10811.

43.   Brideau NJ, Flores HA, Wang J, Maheshwari S, Wang X, Barbash DA. Two Dobzhansky-Muller genes interact to cause hybrid lethality in Drosophila. Science. 2006;314(5803):1292-5. doi: 10.1126/science.1133953. PubMed PMID: 17124320.

44.   Poe AR, Tang L, Wang B, Li Y, Sapar ML, Han C. Dendritic space-filling requires a neuronal type-specific extracellular permissive signal in Drosophila. PNAS. 2017:E8062-E71. doi: 10.1073/pnas.1707467114.

Figure 1. New gRNA-marker designs improve pMAGIC and nMAGIC.

(A) Original and new designs of gRNA-markers for pMAGIC and nMAGIC. (B) Comparison of
clone frequency in larval sensory neurons between two gRNA designs. The number represents

clones between A1 and A7 segments on one side of each larva. n = larvae number: tgFE (n=10),
Qtg2.1 (n=10). (C-E) Labeling of hemocytes in whole 3rd instar larvae by *pxn*[Gal4]*>CD4-tdTom*
alone (C) or together with *ubi-Gal80* (D) or *tub-Gal80* (E). The panels on the right show
enlarged views of the boxed regions. (F) Designs of Gal80 variants tested in pMAGIC gRNA-
markers. (G) The brightness of epidermal clones labeled by pMAGIC gRNA-markers. n = image
numbers: gRNA-40D2-uH (n = 32), gRNA-40D2-uDEH (n = 31), gRNA-42A4-uDEH (n = 52),
gRNA-42A4-tDEH (n = 39), gRNA-42A4-tDES (n = 38). (H) The brightness of neuronal clones
labeled by pMAGIC gRNA-markers. n = neuron numbers: gRNA-40D2-uH (n = 16), gRNA-
40D2-uDEH (n = 16), gRNA-42A4-uDEH (n = 16), gRNA-42A4-tDEH (n = 15), gRNA-42A4-
tDES (n = 16). (I) Portion of a larval wing disc containing nMAGIC clones visualized by
nlsBFP. (J and J') Portion of a wing disc containing nMAGIC clones labeled by cytosolic BFP
(J) and HA staining (J'). (K) Epidermal clones on the larva body wall labeled by nlsBFP. (L)
Epidermal clones visualized by cytosolic BFP. In all plots, black bar, mean; red bar, SD; AU,
arbitrary unit. Student's t-test in (B); one-way analysis of variance (ANOVA) and Tukey's honest
significant difference (HSD) test in (G) and (H). *p≤0.05, **p≤0.01, ***p≤0.001, ns, not
significance. For (C-E), scale bar, 300 μm. For (I-M), scale bar, 100 μm.

615
616 **Figure 2. A genome-wide gRNA-marker kit suits diverse needs of clone frequency.**

(A) Scheme of gRNA-marker insertion sites and target sites on *Drosophila* chromosomes. (B) Comparison of clone frequencies of all pMAGIC gRNA-markers in larval sensory neurons, clones are labeled using *RabX4-Gal4 UAS-MApHs* (for Chromosome X, II and IV) or *21-7-Gal4 UAS-MApHs* (for Chromosome III). n = larvae number: X2 (n = 10), 20F2 (n = 10), 20F1(n = 10), 40D2 (n = 20), 40D4 (n = 10), 40E1 (n = 10), 41F9 (n = 20), 41F11 (n = 10), 42A4 (n = 10), 80C1 (n = 20), 80C2 (n = 14), 80F5 (n = 15), 81F (n = 10), 82A4 (n = 10), 82C3 (n = 10), 101F1a (n = 10), 101F1b (n = 10), 101F1c (n = 10). (C) Comparison of clone areas in larval wing discs labeled by nMAGIC gRNA-markers on 2R. n = wing disc number: 41F9 (n = 14), 41F11 (n = 16), 42A4 (n = 15). (D and E) Neuronal clones in the central part of the adult brain induced by pMAGIC gRNA-markers *gRNA-40D2* (D) and *gRNA-40E1* (E). In all plots, black bar, mean; red bar, SD. One-way ANOVA and Tukey's HSD test. *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ns, not significance. For (D) and (E), scale bar 100 µm.

Figure 3. MAGIC allows clonal analysis in diverse tissues and cell types.
(A-F) pMAGIC clones induced in different tissues by *vas-Cas9 gRNA-40D2(Gal80)* and labeled by *tub-Gal4 UAS-CD8-GFP* (green). DAPI staining (white) shows all nuclei. (G) A pMAGIC epidermal clone on the larval body wall induced by *zk-Cas9 gRNA-40D2(Gal80)* and labeled by

R38F11-Gal4 UAS-tdTom (green). Epidermal junctions are labeled by *α-Catenin-GFP* (white).
(H) pMAGIC glia clones in the larval brain induced by *gcm-Cas9 gRNA-40D2(Gal80)* and
labeled by *repo-Gal4 UAS-CD8-GFP* (green). Glial nuclei are labeled by Repo staining (white).
(I) pMAGIC hemocyte clones induced by *Act-Cas9 gRNA-40D2(Gal80)* and labeled by *pxn-*
Gal4 UAS-tdTom. For figure A, D-F, H, scale bar 100 μm. For figure B-C, G, scale bar 50 μm.
For Figure I, scale bar 25 μm.

640
Figure 4. MAGIC facilitates clonal analysis at the NMJ.
(A-A") pMAGIC clones of *VGlut¹* mutation in motor neurons at the neuromuscular junction.
Clones were induced by *zk-Cas9 gRNA-40D2(Gal80)* and labeled by *tub-Gal4 UAS-CD8-GFP*. ,
The loss of VGlut is confirmed by VGlut staining. The mutant clones are outlined in (A"). (B-
B") A pMAGIC clone of *brp^d09839* mutation in a motor neuron at the neuromuscular junction.
Clones were induced by *zk-Cas9 gRNA-42A4(Gal80)* and labeled by *tub-Gal4 UAS-CD8-GFP*.
The loss of Brp is confirmed by Brp staining. The mutant clone is outlined in (B"). In both
experiments, HRP staining shows all axons. Scale bars, 10 μm.

Figure 5. MAGIC enables clonal analysis of pericentromeric genes, 4th chromosome-associated mutations, and in interspecific hybrid animals.

(A) A WT pMAGIC class IV da neuron clone exhibiting complete dendrite pruning at 16 hours APF. (B-D) pMAGIC clones of EcR^{M554fs} mutation in da neurons imaged at 16 hours APF, exhibiting the lack of pruning (B and D) or apoptosis (C). In (A-D), the clones were induced by *zk-cas9* with *gRNA-41F9(Gal80)* and labeled by *RabX4-Gal4 UAS-MApHS*. Neuronal cell bodies are indicated by arrows. MApHS contains pHluorin and tdTom [19], but only tdTom signals are shown. The signals in epidermal cells (A) were due to engulfment of pruned dendrites by epidermal cells [19]. (E and F) WT (E) and *Df(4)ED6380* (F) pMAGIC clones in C4da neurons induced by *zk-cas9 gRNA-101Fc(Gal80)* and labeled by *RabX4-Gal4 UAS-MApHS*. Only tdTom signals are shown. (G) Normalized dendrite length of WT clones and deficiency clones. Black bar, mean; red bar, SD. Student's t-test. ***p≤0.001. (H) Scheme for interspecific crosses between *D. melanogaster* (*D.m*) and *D. simulans* (*D.s*). (I and J) Wing discs from male (I) and female (J) progeny carrying clones. Scale bars, 50 μm.

Figure S1. Transgenic markers on the 4th chromosome show uneven expression.

(A-B) Representative epidermal (A) and wing disc (B) images showing uneven expression of

gRNA-101F1c(BFP) inserted at *attP¹⁰²ᴰ* on the 4th chromosome. Yellow arrowheads point to cells

lacking BFP expression. Scale bars, 50 µm. (C) Frequency of labeled neurons by indicated

gRNA(Gal80) in the absence and presence of Cas9. The *zk-Cas9* dataset is the same as that for

chromosome 4 in Figure 2B. Black bar, mean; red bar, SD. One-way ANOVA and HSD test.

***p≤0.001.

Table 1. gRNA-marker collection

Chr Arm	target site	gRNA location	pMAGIC vector	nMAGIC vector	clone frequency
2L	40D2	attP[VK00037]	*pAC-U63-gRNA2.1-ubiGal80(DE)-His2Av*	*pAC-U63-tgRNA-nlsBFP**	31
2L	40D4	attP[VK00037]	*pAC-U63-tgRNA-Gal80**	*pAC-U63-tgRNA-nlsBFP**	13
2L	40E1	attP[VK00037]	*pAC-U63-tgRNA-Gal80**	*pAC-U63-tgRNA-nlsBFP**	4
2R	41F9	attP[VK00018]	*pAC-U63-gRNA2.1-ubiGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	10
2R	41F11	attP[VK00018]	*pAC-U63-gRNA2.1-ubiGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	14
2R	42A4	attP[VK00018]	*pAC-U63-gRNA2.1-tubGal80(DE)-SV40*	*pAC-U63-gRNA2.1-tubBFP(HA)*	14
3L	80F5	attP2	*pAC-U63-gRNA2.1-tubGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	11
3L	80C2	attP2	*pAC-U63-gRNA2.1-tubGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	7
3L	80C1	attP2	*pAC-U63-gRNA2.1-tubGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	4
3R	81F	attP[VK00027]	*pAC-U63-gRNA2.1-tubGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	4
3R	82A4	attP[VK00027]	*pAC-U63-gRNA2.1-tubGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	10

3R	82C3	attP[VK00027]	*pAC-U63-gRNA2.1-tubGal80(DE)-His2Av*	*pAC-U63-gRNA2.1-tubBFP(HA)*	16
X	X2	attP18	*pAC-U63-gRNA2.1-tubGal80(DE)-SV40*	*pAC-U63-gRNA2.1-tubBFP(HA)*	19
X	20F1	attP18	*pAC-U63-gRNA2.1-tubGal80(DE)-SV40*	*pAC-U63-gRNA2.1-tubBFP(HA)*	18
X	20F2	attP18	*pAC-U63-gRNA2.1-tubGal80(DE)-SV40*	*pAC-U63-gRNA2.1-tubBFP(HA)*	19
IV	101F1-a	attP[ZH-102D]**	*pAC-U63-gRNA2.1-tubGal80(DE)-SV40*	*pAC-U63-gRNA2.1-tubBFP(HA)*	4
IV	101F1-b	attP[ZH-102D]**	*pAC-U63-gRNA2.1-tubGal80(DE)-SV40*	*pAC-U63-gRNA2.1-tubBFP(HA)*	25
IV	101F1-c	attP[ZH-102D]**	*pAC-U63-gRNA2.1-tubGal80(DE)-SV40*	*pAC-U63-gRNA2.1-tubBFP(HA)*	28
Clone frequencies are measured by averaging clone numbers in 10 laterally mounted 3rd instar larvae with pMAGIC gRNA-markers. Only clones in A1 to A7 segments of one side of each larva were counted. *Previously published gRNA-markers. **3xP3-RFP has been removed from these lines.					

673