Sensitivity of genome-wide tests for mitonuclear genetic incompatibilities

Kuster, Shady A.[1,2], Schumer, Molly[3,4], Havird, Justin C.[5], Sloan, Daniel B.[2]

- 1: Cell and Molecular Biology, Colorado State University
- 2: Department of Biology, Colorado State University
- 3: Department of Biology, Stanford University
- 4: Freeman Hrabowski Fellow, Howard Hughes Medical Institute
- 5: Department of Integrative Biology, University of Texas at Austin

Author for correspondence: Shady A. Kuster, Shady.Kuster@colostate.edu

<u>ABSTRACT</u>

Mismatches between interacting mitochondrial and nuclear gene products in hybrids have been

proposed to disproportionately contribute to early species boundaries. Under this model, genetic

incompatibilities emerge when mitochondrial haplotypes are in a cellular context without their

coevolved nuclear-encoded mitochondrial (n-mt) proteins. Some case studies have shown that

such disruptions in mitonuclear coevolution can contribute to reproductive isolation, but whether

mitonuclear incompatibilities generate selection that impacts multiple n-mt loci and/or causes

broad, genome-wide contributions to speciation is unclear. Here, we leverage a system with

several hybridizing species pairs (*Xiphophorus* fishes) that have known mitonuclear

incompatibilities of large effect. We divided nuclear-encoded genes into three classes based on

level of interaction with mitochondrial gene products. We found only inconsistent statistical

support for a difference between these classes in the degree of positive covariation in

mitonuclear ancestry. We discuss evidence that these analyses are sensitive to the amount of

non-synonymous divergence between parent species in interacting n-mt genes or the age of the

hybridization event. Overall, our results imply that genome-wide scans focused on enrichment of

broad functional gene classes may often be insufficient for detecting a history of mitonuclear

coevolution, even when strong selection is acting on mitonuclear incompatibilities at multiple

loci.

INTRODUCTION

Since its endosymbiotic origin, the mitochondrion has evolved into an integral component of the eukaryotic cell, functioning in such processes as oxidative phosphorylation (OXPHOS), Ca^{2+} acquisition and signaling, and biosynthesis of heme and iron-sulfur clusters. Bilaterian metazoan mitochondrial genomes (mitogenomes) have been reduced to encode only 13 core protein subunits of the electron transport chain, 2 rRNAs, and 22 tRNAs, making mitochondria dependent on the import of nuclear-encoded, but mitochondrially targeted, (n-mt) proteins for all functions [1–4]. Due to the importance of mitochondrial function to fitness, maintaining these intimate interactions among coevolved partners may be essential.

Mitogenomes evolve at a faster rate than nuclear genomes in most bilaterian metazoan systems, with an average rate of 19× faster in vertebrates [5–8]. These elevated substitution rates may result from a higher underlying mutation rate in combination with inefficient purging of deleterious mutations due to smaller effective population size, lack of recombination, and uniparental inheritance [9–12], although explanations based on reduced efficacy of selection in mitogenomes have been recently challenged [13–15]. Regardless of the cause, the consequences of high mitochondrial DNA (mtDNA) substitution rates are often deleterious [6]. As such, substitutions in mitochondrially encoded proteins may create selection for compensatory changes in interacting n-mt proteins [16–18], and signatures of coevolution have indeed been found between the mitochondrial and nuclear genomes [19–22]. Other forms of mitonuclear coevolution (i.e., reciprocal selection for compatible genotypes) are also possible [19,23–25]. Regardless of the exact process, the outcome is a set of interacting proteins that are functionally dependent on sufficient "matching" of lineage-specific substitutions in mitochondrial and n-mt proteins [23].

As a result, disrupting the association between a mitochondrial haplotype and its coevolved nuclear counterparts (e.g., via secondary contact following speciation) is predicted to uncover genetic incompatibilities that made early contributions to reproductive isolation [5,26]. Given the rate at which mitonuclear genes coevolve and the importance of mitochondrial function to cellular metabolism, mitonuclear incompatibilities are proposed to have a disproportionate effect on the accumulation of reproductive barriers during lineage divergence [23,27–30]. These incompatibilities could potentially be driven by only a few genes with major

effects on fitness or alternatively by many genes of small effect. Although studies have found

evidence of mitonuclear incompatibilities (e.g., [31,32]) in some systems, it is still unclear how

far-reaching these effects are and how many genes are involved. For example, the immediate

effects of selection on mitonuclear incompatibilities could be restricted to a subset of n-mt

proteins involved in direct physical interactions with mitochondrial gene products. On the other

hand, because mitochondria are hubs of various biological processes, selection may be more

widespread and affect nuclear genes that have no direct connection to mitochondrial function.

We hypothesize that, relative to other nuclear genes, n-mt genes are more strongly affected by

selection acting to maintain associations between matched lineage-specific mitonuclear

genotypes. Therefore, examining signatures of mtDNA "match" in n-mt genes vs. nuclear-

encoded genes that lack mitochondrial interactions should reveal a history of mitonuclear

coevolution across the genome at species boundaries. While partitioning genomic data based on

n-mt gene function has been used to investigate mitonuclear incompatibilities in many systems

(Table S1), the effectiveness of this approach has not been tested in a system with previously

known mitonuclear incompatibilities.

Species of the swordtail fish genus *Xiphophorus* have recently begun hybridizing in

multiple river systems in Mexico, providing a model of repeated hybridization and speciation

dynamics [33–42]. Recent work on hybrid zones between sister species *X. birchmanni* and *X.*

malinche has identified severe mitonuclear incompatibilities involving the *ndufs5* and *ndufa13* n-

mt genes [43]. Mitonuclear ancestry mismatches at these loci have severe fitness effects,

including death during embryonic development. Subsequent investigations have identified that at

least nine nuclear genomic regions (which contain a total of 10 known n-mt genes across seven

loci) are involved in mitonuclear incompatibilities in these hybrids [44]. Some of these

incompatibilities, including the *ndufs5* and *ndufa13* n-mt genes, were also found in *X.*

birchmanni and *X. cortezi* hybrids [42]. We therefore leveraged this system to investigate

evidence for mitonuclear matching during hybridization on a genome-wide scale and the ability

to detect these interactions with commonly used approaches that partition nuclear genes into

classes based on their role in mitochondrial function. We categorized nuclear genes into three

groups based on whether the encoded protein (1) interacts with mitogenomic gene products

(interacting n-mt), (2) functions in mitochondria but does not physically interact with

mitogenomic gene products (non-interacting n-mt), or (3) does not function in mitochondria

(non-n-mt). This *a priori* partitioning allows for an unbiased test of whether selection for

matched mitonuclear genotypes leaves genome-wide signatures. Overall, we found inconsistent

statistical evidence for differences among these categories based on analyses of mitonuclear

interactions in *Xiphophorus* hybrids. However, we found that interacting n-mt genes with larger

amounts of non-synonymous substitutions may have more severe consequences for mitonuclear

incompatibilities. Therefore, the role of n-mt genes may not be generally detectable on a

genome-wide scale, and the sensitivity of these tests should be taken into consideration when

interpreting results.

METHODS

Genomic data from *Xiphophorus* hybrid populations

Three datasets resulting from previous local ancestry inference in hybrid *Xiphophorus* fishes

were acquired as described previously [37,40,43]. Each dataset includes samples from a

population of hybrid *Xiphophorus* individuals inhabiting sections of rivers in the states of

Hidalgo or San Luis Potosí, Mexico. The first (Calnali Low, or CALL) is on the Calnali River

and consists of *X. birchmanni* and *X. malinche* hybrids. The second is at Chahuaco Falls (CHAF)

on the Calnali River. It is separated from the CALL population by a waterfall (~50m tall) [43]

and consists of hybrids of the same pair of species as CALL. The third represents two

populations (Huextetitla and Santa Cruz, or HUEX-STAC) downstream of the same fork on the

Santa Cruz River and derived from a shared hybridization event between *X. birchmanni* and *X.*

cortezi followed by independent evolution [37,38].

These datasets were originally analyzed using the *ancestryinfer* pipeline [45] and resulted

in posterior probabilities of species ancestry at between 628,881 and 689,982 ancestry

informative markers (AIMs) across the nuclear and mitochondrial genomes (Table 1), as

described previously [36,39,45]. Informative sites were defined as those with at least a 98%

frequency difference between the parental species, as previously described [33,44]. We used a

published annotation of the *X. birchmanni* reference genome [46] to identify genes containing

each AIM. We then transformed AIM probabilities into categorical genotype calls based on a

posterior probability threshold. If a probability for a parent species was greater than or equal to

0.9 and the other parent species' probability of ancestry was less than or equal to 0.1, ancestry

was assigned to the major probability. If instead the sum of both parents' ancestries was less than

or equal to 0.2, the AIM was assigned as heterozygous. AIMs that did not meet these criteria

were filtered out of the dataset. Allele counts were then averaged across all AIMs that passed the

prior filtering to calculate an average major parent ancestry proportion per individual. These

values were then averaged for nuclear and mitochondrial AIMs to obtain population-level

averages (Table 1).

Identification of *Xiphophorus* n-mt genes

The gffread [47] program was used to obtain predicted protein sequences, using the -V option to

remove any coding sequence with an in-frame stop codon. To annotate n-mt proteins, we

compared the *Xiphophorus* translated proteome to UniProt human (UP000005640,

GCA_000001405.27) and mouse (UP000000589, GCA_000001635.8) protein sequences. N-mt

proteins have been empirically identified in humans and mice as part of the MitoCarta 3.0 project

[4]. Because the UniProt human and mouse datasets differed in nomenclature from the MitoCarta

dataset, we assigned protein names by searching for MitoCarta sequences in the UniProt human

sequences with BLASTP v2.15.0+. A custom Perl script was then used to parse the BLASTP

output, and identified proteins were manually confirmed to be the correct protein if percent

coverage and sequence identity were not 100%. OrthoFinder v2.5.4 [48] was then run with the

primary translated transcript for each *Xiphophorus* sequence and the full sets of human and

mouse protein sequences.

MitoCarta protein classifications were used to differentiate non-n-mt from n-mt

orthogroups. Interacting and non-interacting orthogroups were distinguished based on a

previously published [49] list, updated here, of interacting n-mt proteins that includes OXPHOS

subunits, ribosomal subunits, aminoacyl-tRNA synthetases, and DNA/RNA polymerases (Table

S2).

Linkage disequilibrium calculation

We used mitonuclear linkage disequilibrium (LD) as a metric for the association between

mitochondrial and nuclear alleles in the CALL and CHAF hybrid populations (see below for

analysis methods for the HUEX-STAC population, which differed because their mitogenome is

fixed for the haplotype from the *X. cortezi* parental species [38]). Calculations were performed

using the D' and r^2 statistics [50]. In the equations that follow, A and B represent nuclear and

mitochondrial loci, respectively, and pA and pB represent the corresponding allele frequencies.

pAB is the frequency of the haplotype AB across the two sites. The A and B alleles were assigned

to be from the same parental population so that positive and negative values of LD indicate a

higher frequency of parental or recombinant allele combinations (i.e., matched and mismatched),

respectively.

$$D = p_{AB} - p_A p_B$$

$$D' = \frac{D}{D_{max}}$$

where

$$D_{max} = \begin{cases} \min[p_A(1 - p_B), (1 - p_A)p_B] \; if \; D > 0 \\ \min[p_A p_B, (1 - p_A)(1 - p_B)] \; if \; D < 0 \end{cases}$$

and

$$r^2 = \frac{D^2}{p_A * (1 - p_A) * p_B * (1 - p_B)}$$

Because all mitochondrial genes are linked on the same non-recombining mitogenome, a

single AIM was used (CALL and CHAF's AIM was in *COX2* and HUEX-STAC's AIM was in

ATP6, positions 45 and 846 of GenBank accession CM071424.1, respectively) to identify the

mitochondrial haplotype for each fish and to represent the mitogenome in all LD analyses. This

approach was appropriate for our data, as all mtDNA AIMs within an individual had

approximately the same posterior probability for ancestry assignment. Posterior probabilities

were summed across each nuclear AIM as a proxy for frequency of the two alleles. The two LD

statistics were calculated for each gene class separately in a custom R script. LD values were

visualized using ggplot2 v3.4.0 and ggridges v0.5.4 in RStudio v2023.03.1+446.

LD values were then averaged across AIMs within each nuclear gene to limit pseudo-

replication. A one-way ANOVA from the R stats v4.2.2 package (commands *lm* and *anova*) was

used to determine whether gene class had a statistically significant impact on LD values.

Summary statistics were collected using commands *n, mean*, and *sd* from the dplyr v1.1.0, base

v4.2.2, and stats v4.2.2 packages, respectively.

In the case of selection on a small number of n-mt genes, shifts in genome-wide averages

may not be detectable but could still result in individual loci with the strongest association with

the mitogenome being enriched for interacting n-mt genes. We therefore also identified the AIMs

and genes with the top 1% of LD values with a custom R script.

Partial correlation calculation

High LD values can result from population structure (i.e., proportion of parent species ancestry

in an individual genome), and the populations used here vary in the proportion of background

ancestry (Table 1). We therefore performed a partial correlation (pcor) analysis, which calculated

the correlation between mitochondrial and nuclear ancestry while accounting for the correlation

expected given the variation in genome-wide ancestry among individuals. AIMs were filtered as

described above, and individual ancestry genotype calls for mitochondrial and nuclear AIMs

were used to estimate a correlation coefficient with the *pcor.test* function from the R ppcor v1.1

package. This estimate was then averaged across all AIMs in a gene, and a one-way ANOVA

was run to determine the effect of gene class on the correlation estimate.

HUEX-STAC allele frequency calculation

The HUEX-STAC populations are both fixed for the *X. cortezi* mitochondrial haplotype, so we

expected *X. cortezi* ancestry to be disproportionately represented at n-mt loci. However, due to

the lack of mtDNA variation, the average allele frequency for each nuclear gene was calculated

and visualized (with a custom R script) instead of mitonuclear LD to infer how mtDNA

background affects selection on n-mt genes in this system. A one-way ANOVA and summary

statistics were computed as described above.

A permutation test was then performed to determine whether significant results were

robust to violations of normality assumptions in ANOVA. Allele frequency values were

permutated by randomly reassigning them across genes 9999 times using base R v4.2.2 functions

in a custom R script. Permuted datasets were then used to calculate null expectations for

differences in allele frequencies between interacting n-mt and non-n-mt gene classes. The

observed difference was compared to the permutations to determine the probability of the pattern

arising by chance.

Non-synonymous substitution analysis

We further filtered the three datasets to include only genes that had substitutions in the coding

sequence. We quantified the number of non-synonymous substitutions normalized to the length

of the coding sequence. This filtered dataset was analyzed using a linear model with mitonuclear

LD as the response variable and an interaction between gene class and our normalized count of

non-synonymous substitutions as the explanatory variables. A type III ANOVA was then run

(via *Anova* from the R car v3.1-1 package) on the model to determine statistical significance of

our predictor variables followed by use of the *summary* function (from base R) with non-n-mts

as the reference group. An influential point analysis was performed on the interacting n-mt gene

class using DFFIT with the plotting command *ols_plot_dffits* from the R olsrr v0.6.1 package.

Identified points were then individually removed from the model with all gene classes to

determine their effects on the relationship between mitonuclear LD and non-synonymous

substitutions.

Data and code availability

All code has been deposited in GitHub at https://github.com/sakuster/sensitivity-of-genome-

wide-mtnuc. Data can be accessed on figshare (https://doi.org/10.6084/m9.figshare.c.7902011).

RESULTS

Identification of *Xiphophorus* n-mt sequences

Of the 24,269 *Xiphophorus* annotated genes, 86.5% (20,988) were assigned to an orthogroup by

OrthoFinder. We found that 75.5% (13,264) of the 17,566 identified orthogroups contained at

least one *Xiphophorus* gene, and 69.4% (12,198) included at least one gene from all three species

(*X. birchmanni*, mouse, and human). Gene and/or genome duplications during the evolutionary

history of these three species resulted in some orthogroups with multiple *X. birchmanni* genes

(24.1% of orthogroups, or 4,234, contained 2 or more).

These orthogroups were used to classify *Xiphophorus* genes into our three functional

categories for each population (Table S3). MitoCarta 3.0 consists of 1,136 human and 1,140

mouse n-mt genes. All but three of the mouse n-mt genes (*Gm4984*, *Htd2*, and *Nsun4*) were

identified, and only one human n-mt gene (*NDUFB1*) was not identified in *X. birchmanni*.

Overall, 159 orthogroups were found to contain interacting n-mts, 822 to contain non-interacting

n-mts, and 15,545 to contain non-n-mts. These orthogroups had 184, 1,042, and 23,043 *X.*

birchmanni genes, respectively.

Mitonuclear linkage disequilibrium does not differ among gene classes in hybrid

populations with segregating mitochondrial haplotypes

We sought to determine whether n-mt genes maintained higher mitonuclear LD in *Xiphophorus*

hybrids due to selection against mitonuclear incompatibilities. On average, mitonuclear LD

values were high across the entire nuclear genome in the CALL and CHAF hybrid populations

(Figure 1, Supplemental Figure 1), meaning nuclear ancestry matched the mitochondrial

haplotype (positive D' values). This result is also illustrated by differences in average nuclear

ancestry between individuals with mitochondrial haplotypes from the two possible parent species

(Supplemental Figure 2). These strong associations are likely driven by population structure in

these two populations [33,42]. Accordingly, a partial correlation (pcor) test that controls for

population structure resulted in lower mitonuclear association values (Supplemental Figure 1).

Under a model of mitonuclear coevolution, we predicted that the interacting n-mt gene

class would have higher mitonuclear LD values compared to the other gene classes. However,

we did not find a consistent significant difference between the gene classes for CALL (D': $p =$

0.44; r^2: $p = 0.44$, pcor: $p = 0.04$) or CHAF (D': $p = 0.86$; r^2: $p = 0.29$, pcor: $p = 0.14$; Figure 1,

Supplementary Figure 1, summary statistics in Table S4). In fact, the pcor values in CALL

instead showed higher mitonuclear LD in the non-n-mt gene class than in either of the two n-mt

gene classes. Therefore, it does not appear that mitochondrial haplotypes have preferentially

maintained associations with n-mt loci of the same ancestry at a genome-wide scale in these

hybridizing populations.

We additionally tested whether interacting n-mt genes were overrepresented in the top

1% of all genes based on our association metrics (D', r^2, pcor). However, again, significant

enrichment for interacting n-mt genes was inconsistent (Table S5).

Known mitonuclear incompatibility genes show positive associations with mitochondrial haplotype

Nine of the 10 previously identified mitonuclear incompatibility genes [44] were included in our CALL analysis, and only two (*NDUFS5* and *SMIM8*) were outliers for mitonuclear LD (with a Z-score of >2 above the mean D' value) (Table S6, Figure S3). Therefore, although the top 1% genome-wide LD values did not show an overrepresentation of n-mt genes, we do find that some genes of known biological importance exhibit signatures of elevated mitonuclear LD. Five other genes (*NDUFA13*, *MTERF4*, *ATP5MG*, *LYRM2* and *RMDN3*) were also above the mean but not outliers by this definition, and the final two genes (*MMUT* and *UQCRC2*) were very close to the mean. Repeating these analyses with r^2 and pcor values instead of D' values resulted in slight changes, but the results were generally similar (Table S6, Figures S4-5).

To see whether these genetic incompatibilities showed similar signatures across populations with varying mtDNA frequencies and different parental source species, we also investigated mitonuclear associations for CHAF and HUEX-STAC (Table S6, Figures S6-9. The CALL and CHAF populations, which are hybrids between the same parental species, exhibit a significant correlation in mitonuclear LD (D') across this set of incompatibility genes (Pearson correlation; $r = 0.87$; $p = 0.0051$). Comparisons with allele frequencies in HUEX-STAC, which originates from a different pair of parental species, also show a positive trend, but the relationships are weaker and/or non-significant with both CALL ($r = 0.34$; $p = 0.41$) and CHAF ($r = 0.70$; $p = 0.051$).

Interacting n-mt genes show enriched mitochondrial match in an older hybrid population fixed for a mitochondrial haplotype

The HUEX-STAC populations are fixed for the *X. cortezi* mitochondrial haplotype, so we expected to see *X. cortezi* ancestry disproportionately represented at n-mt loci. HUEX-STAC individuals generally have a large percentage of *X. cortezi* ancestry across their entire nuclear genome with a mean allele frequency of 88.7% ± a standard error of 0.011% (Figure 2B). Non-interacting n-mt and non-n-mt genes showed similar mean *X. cortezi* allele frequencies at 82.2 ± 0.02% and 83.1 ± 0.001%, respectively. Interacting n-mt genes had a higher mean allele

frequency (88.0 ± 0.097%, Figure 2A, Table S4). Thus, unlike in the CALL and CHAF

populations, individuals collected from the HUEX-STAC population did demonstrate a

difference in allele frequencies among gene classes (one way ANOVA: $p = 0.005$) and supported

the predicted higher degree of matched mitonuclear ancestries for interacting n-mt genes (Figure

2A). This result was also supported by a non-parametric permutation test (Figure S10).

Interacting n-mt genes with more non-synonymous substitutions may show higher

mitonuclear LD

Because many nuclear-encoded proteins may show few or no fixed differences in amino acid

sequence between these closely related species, we asked whether the presence of non-

synonymous substitutions between species affected how gene class related to mitonuclear LD.

There were inconsistent effects of gene class, non-synonymous substitutions, and the interaction

between these two across metrics and populations (Tables S7-8). There was, however, some

evidence that interacting n-mt genes with more non-synonymous substitutions have higher

mitonuclear LD compared to non-n-mt genes (Figure 3, Figures S11-12, Table S9). The

significant interactions in CALL and CHAF were found to be the result of a few influential genes

(Table S10). For example, *NDUFS5*, a gene previously shown to underly a lethal mitonuclear

incompatibility, consistently appeared as influential across association metrics and populations

where a significant interaction was found. When *NDUFS5* was removed from the dataset, the

difference between the interacting n-mt and non-n-mt gene classes was eliminated. Other known

incompatibility genes were not consistently influential, although *NDUFA13* and *ATP5MG* were

detected in some analyses in CHAF.

DISCUSSION

We tested the prediction that the interacting n-mt gene class would show the strongest "matched"

association with mitochondrial haplotype. Despite the known role of n-mt genes in hybrid

incompatibilities between *X. birchmanni* and *X. malinche*, the CALL and CHAF populations did

not support this prediction. This negative result likely speaks to the statistical power issues

inherent in identifying mitonuclear interactions. The sensitivity of genome-wide tests for

signatures of selection acting preferentially on n-mt genes will likely be affected by the structure
of the incompatibility and the genomic background of the genes under selection, factors that are
also important for detecting loci involved in nuclear-nuclear hybrid incompatibilities [51].
Specifically, the number of genes under selection, where those genes are distributed in the
genome, and/or whether only n-mt genes are involved could affect the sensitivity of these tests.

N-mt genes make up a small portion of the genome, and only a subset of those genes may
be participating in mitonuclear epistasis. Incompatibilities can potentially involve a few loci with
large effect size or many loci of small effect (or some combination of these extremes). The
analyses performed by Robles et al. [44] identified 10 n-mt genes among seven genomic regions
contributing to mitonuclear incompatibilities in *Xiphophorus* species hybrid populations, one of
which was CALL. These genes represent only 6.8% of all interacting n-mt genes and 1.0% of all
n-mt genes. Selection on these few genes would have to be exceptionally strong and have had
enough time to act to detect a significant shift in the average for the entire interacting n-mt gene
class. In contrast, systems such as *Tigriopus* copepods appear to have many genes involved in
mitonuclear incompatibilities and have proven to be powerful in advancing the field of
mitonuclear genetics and genomics [20,31]. Therefore, even in cases where n-mt genes are
contributing to negative mitonuclear epistasis, the number of loci may be too small to detect
genome-wide effects with the approach implemented here, which is inherently designed to detect
polygenic contributions to incompatibilities.

Elevations in background LD (e.g., due to population structure) and the distribution of n-
mt genes across the genome will also affect the ability to detect a difference in mtDNA ancestry
matching between n-mt and non-n-mt genes. Any signal of this difference arising from selection
on an individual n-mt target could be diluted by hitchhiking at linked non-n-mt loci [52]. This
effect would be particularly problematic for populations that are the result of recent admixture
events and, therefore, have not had many generations for recombination to break up haplotype
blocks in the nuclear genome [41,53], such as in CALL and CHAF (whose initial admixture was
estimated to be ~46 generations ago [35]). In some cases, clustering of n-mt genes in specific
regions of the genome could create strong targets of selection that overcome this dilution effect.
For example, genomic regions with a high density of n-mt genes have been identified in the
long-tailed finch [54] and the eastern yellow robin [55], both of which have found evidence that
n-mt genes disproportionately share ancestry with mtDNA.

Additionally, not all mitonuclear incompatibilities are necessarily mediated by n-mt

genes. Instead, some incompatibilities may involve nuclear-encoded proteins that are not

targeted to mitochondria but nonetheless create genetic interactions with mtDNA through co-

functionality in biochemical and signaling pathways that span multiple cellular compartments

[56]. For example, our definition of an n-mt gene is a gene whose product is targeted for import

into mitochondria. However, mitochondria are a part of a complex cellular network, and any

number of cellular processes could affect mitochondrial function. Genes involved in these

processes are, by our definition, non-n-mt genes. Therefore, any contributions of non-n-mt genes

to mitonuclear incompatibilities would directly undermine search strategies based on partitioning

the genome into n-mt and non-n-mt categories.

However, these challenges were somewhat mitigated in our analyses of CALL and

CHAF by including the number of non-synonymous substitutions for each gene in the model. By

looking only at genes with substitutions between species, we uncovered a pattern in which

interacting n-mt genes tended to show a stronger relationship between mitonuclear association

and the number of non-synonymous substitutions than non-n-mt genes (Figure 3, Figures S11-

12). This stronger, positive relationship could imply that putative functional divergence results in

stronger selection to match ancestry to the mitochondrial haplotype if the divergence occurs in an

interacting n-mt gene. However, this relationship was found to be disproportionately driven by a

few genes, namely *NDUFS5*, a known incompatibility gene [43]. Nevertheless, this result

suggests that including protein sequence divergence could increase the power to detect the

effects of mitonuclear incompatibilities, even when using genome-wide tests in a system with

few genes of large effect. We speculate that these effects will be easiest to detect in interacting n-

mt genes but that this result does not rule out the possibility of other genes contributing to

mitonuclear incompatibilities.

Despite the challenges outlined above, one of the three hybrid populations we

investigated (HUEX-STAC) supported our prediction of enrichment in matched ancestry for n-

mt genes. Notably, this hybrid population is older than CALL and CHAF, estimated to have

formed ~250 generations ago [33,35], and it is the only population we investigated that is fixed

for a mitochondrial haplotype from just one of the parental species. Its older age suggests there

has been more time for incompatible genotype combinations to be purged from the population

and for recombination to have broken up haplotype blocks in the nuclear genome, thereby

limiting the effects of hitchhiking. We therefore speculate that the age of HUEX-STAC and the
fixation of a mitochondrial haplotype may have made selection on n-mt loci stronger and/or
more effective. In such populations, half of the possible mitonuclear combinations have been
removed, so selection for matched ancestry would consistently favor the same n-mt allele. In
contrast, when mitochondrial haplotypes are still segregating, the fitness effects of an n-mt allele
may change depending on an individual's mtDNA. Because mitochondrial and nuclear loci are
not physically linked, very strong selection would be needed to preserve mitonuclear LD in
segregating populations [57].

Despite the potential importance of the age and fixed mitochondrial haplotype of the
HUEX-STAC population, previous studies have found examples of hybrid populations with both
fixed and segregating mitochondrial haplotypes that show a disproportionate effect of selection
on n-mt genes during hybridization (Table S1). Therefore, expanding the availability of genomic
datasets from hybrid systems and performing a formal meta-analysis would be important future
directions to assess relevant variables, such as whether mitochondrial haplotypes are segregating
in the population, the age of the hybrid population, the number of loci involved and their effect
sizes, and the genetic structure of the population. Reanalysis of publicly available genome-level
data from hybrid populations may also be a valuable approach to control for method of ancestry
inference and metric of association between mtDNA and nuclear loci.

Another variable that differentiates the HUEX-STAC population is the parental makeup
of the hybrids. Although geographically distinct, CHAF and CALL share parental species (*X.*
birchmanni and *X. malinche*). In contrast, HUEX-STAC was formed by hybridization between
X. birchmanni and *X. cortezi*. These three species demonstrate mitonuclear discordance [58] and
a history of introgression [42], both of which may impact mitonuclear incompatibilities in these
hybrids, potentially resulting in stronger selection for matched mitonuclear ancestry. Given these
different demographic histories, each of the hybrid populations we analyze here is expected to
have numerous unique features, so caution is warranted in attributing any of our results to a
single feature (such as the fixed mitochondrial haplotype in HUEX-STAC). Nevertheless, we
view the potential role of fixed vs. segregating mitochondrial haplotypes to be an important area
of future research given the small number of studies conducted to date.

Overall, our findings suggest that splitting up the genome into n-mt and non-n-mt gene
classes may be an underpowered method to detect effects of mitonuclear epistasis in recently

admixed populations, even when there are clear biological effects of said epistasis. Although

such tests for genome-wide signatures may offer a relatively unbiased approach, they may be

susceptible to false negatives. Therefore, important contributions of mitonuclear interactions to

speciation may remain unrecognized in many systems.

<u>ACKNOWLEDGEMENTS</u>

SAK was supported by an NIH T32 fellowship (GM132057) and an NSF GRFP (006784). This

work was also supported by NIH grants to DBS (R35GM148134) and JCH (R35GM142836) and

an EDGE - NSF grant (IOS-2421661) to JCH and MS. We thank three anonymous reviewers for

their insightful comments.

TABLES

Table 1. Summary of Xiphophorus hybrid populations used in this study.

Site Abbreviation	CALL (Calnali Low)	CHAF (Chahuaco Falls)	HUEX-STAC (Huextetitla and Santa Cruz)
Population Site	Calnali River	Calnali River	Santa Cruz River
Major Parent Species	*X. malinche*	*X. malinche*	*X. cortezi*
Minor Parent Species	*X. birchmanni*	*X. birchmanni*	*X. birchmanni*
Number of Individuals	281	250	255
Mitochondrial Haplotype Frequency[a]	63.7% *X. birchmanni*	93.5% *X. malinche*	97.8% *X. cortezi*
Average Major Parent Nuclear Ancestry	52.0%	71.3%	88.7%
Nuclear Ancestry Informative Sites	629,430	628,788	689,966
Mitochondrial Ancestry Informative Sites	93	93	16
Estimated Generations Since Initial Admixture	46 ± 1 [40]	46 ± 1 [40]	263 [33]

[a]Haplotype frequencies for each population were calculated following filtering of mitochondrial AIMs with low-confidence genotypes (posterior probability between 0.2 and 0.8). Filtering resulted in sample sizes slightly lower than the Number of Individuals row: CALL n = 278, CHAF n = 249, HUEX-STAC n = 243.

FIGURES

Figure 1. Distribution of mitonuclear D′ *values for each gene class in the CALL (A) and CHAF (B) populations. One-way ANOVA testing if D′ value differs among gene classes: CALL p = 0.44 (A); CHAF p = 0.86 (B). Summary statistics are reported in Table S4, and other mitonuclear LD metrics are shown in Figure S1.*

Figure 2. HUEX-STAC analyses for A) allele frequencies for each gene class and B) nuclear genome-wide ancestry for the HUEX-STAC population, a hybrid population fixed for the X. cortezi *mitogenome. A one-way ANOVA found gene class to significantly affect allele frequency (p = 0.006); see Table S4 for summary statistics showing differences in means Figure S10 for permutation test demonstrating the robustness of this result. The stacked bar histogram in B was made with 12 bins.*

Figure 3. Regression analysis of mitonuclear association metric by non-synonymous substitutions. Partial correlation values for CALL do not show a significant interaction (type III ANOVA p = 0.073) but interacting n-mts have a stronger linear relationship than the non-n-mt relationship (p = 0.034). Partial correlation values for CHAF does not have a significant interaction (p = 0.67) or difference in slopes (p = 0.50). X. cortezi allele frequencies in HUEX-STAC also do not show a significant interaction (p = 0.43) or a difference in slopes (p = 0.85). Other metrics of mitonuclear LD can be found in Figures S10-11. Statistics can be found in Tables S7-10.

REFERENCES

1. Gray MW, Burger G, Lang BF. 1999 Mitochondrial Evolution. *Science (1979)* **283**, 1476–1481.
2. Karlberg O, Canback B, Kurland CG, Andersson SGE. 2000 The dual origin of the yeast mitochondrial proteome. *Yeast* **17**, 170–187.
3. Gray MW. 2015 Mosaic nature of the mitochondrial proteome: Implications for the origin and evolution of mitochondria. *Proc Natl Acad Sci U S A* **112**, 10133–10138. (doi:10.1073/pnas.1421379112)
4. Rath S *et al.* 2021 MitoCarta3.0: An updated mitochondrial proteome now with sub-organelle localization and pathway annotations. *Nucleic Acids Res* **49**, D1541–D1547. (doi:10.1093/nar/gkaa1011)
5. Burton RS. 2022 The role of mitonuclear incompatibilities in allopatric speciation. *Cellular and Molecular Life Sciences* **79**. (doi:10.1007/s00018-021-04059-3)
6. Lynch M, Blanchard JL. 1998 Deleterious mutation accumulation in organelle genomes. *Felsenstein* **102**, 29–39.
7. Nabholz B, Glémin S, Galtier N. 2008 Strong variations of mitochondrial mutation rate across mammals - The longevity hypothesis. *Mol Biol Evol* **25**, 120–130. (doi:10.1093/molbev/msm248)
8. Allio R, Donega S, Galtier N, Nabholz B. 2017 Large variation in the ratio of mitochondrial to nuclear mutation rate across animals: Implications for genetic diversity and the use of mitochondrial DNA as a molecular marker. *Mol Biol Evol* **34**, 2762–2772. (doi:10.1093/molbev/msx197)
9. Lynch M. 1996 Mutation Accumulation in Transfer RNAs: Molecular Evidence for Muller's Ratchet in Mitochondrial Genomes. *Mol. Biol. Evol.* **13**, 209–220.
10. Melvin RG, Ballard JWO. 2017 Cellular and population level processes influence the rate, accumulation and observed frequency of inherited and somatic mtDNA mutations. *Mutagenesis* **32**, 323–334. (doi:10.1093/mutage/gex004)
11. Neiman M, Taylor DR. 2009 The causes of mutation accumulation in mitochondrial genomes. *Proceedings of the Royal Society B: Biological Sciences* **276**, 1201–1209. (doi:10.1098/rspb.2008.1758)
12. Sanchez-Contreras M *et al.* 2021 A replication-linked mutational gradient drives somatic mutation accumulation and influences germline polymorphisms and genome composition in mitochondrial DNA. *Nucleic Acids Res* **49**, 11103–11118. (doi:10.1093/nar/gkab901)
13. Radzvilavicius AL, Kokko H, Christie JR. 2017 Mitigating mitochondrial genome erosion without recombination. *Genetics* **207**, 1079–1088. (doi:10.1534/genetics.117.300273)
14. Cooper BS, Burrus CR, Ji C, Hahn MW, Montooth KL. 2015 Similar efficacies of selection shape mitochondrial and nuclear genes in both Drosophila melanogaster and Homo sapiens. *G3: Genes, Genomes, Genetics* **5**, 2165–2176. (doi:10.1534/g3.114.016493)
15. Edwards DM, Røyrvik EC, Chustecki JM, Giannakis K, Glastad RC, Radzvilavicius AL, Johnston IG. 2021 Avoiding organelle mutational meltdown across eukaryotes with or without a germline bottleneck. *PLoS Biol* **19**. (doi:10.1371/journal.pbio.3001153)
16. Rand DM, Haney RA, Fry AJ. 2004 Cytonuclear coevolution: The genomics of cooperation. *Trends Ecol Evol* **19**, 645–653. (doi:10.1016/j.tree.2004.10.003)

17. Schmidt TR, Wu W, Goodman M, Grossman LI. 2001 Evolution of Nuclear-and Mitochondrial-Encoded Subunit Interaction in Cytochrome c Oxidase. *Mol. Biol. Evol* **18**, 563–569.

18. Hill GE. 2020 Mitonuclear Compensatory Coevolution. *Trends in Genetics* **36**, 403–414. (doi:10.1016/j.tig.2020.03.002)

19. Weaver RJ, Rabinowitz S, Thueson K, Havird JC. 2022 Genomic Signatures of Mitonuclear Coevolution in Mammals. *Mol Biol Evol* **39**. (doi:10.1093/molbev/msac233)

20. Barreto FS, Watson ET, Lima TG, Willett CS, Edmands S, Li W, Burton RS. 2018 Genomic signatures of mitonuclear coevolution across populations of Tigriopus californicus. *Nat Ecol Evol* **2**, 1250–1257. (doi:10.1038/s41559-018-0588-1)

21. Havird JC, Whitehill NS, Snow CD, Sloan DB. 2015 Conservative and compensatory evolution in oxidative phosphorylation complexes of angiosperms with highly divergent rates of mitochondrial genome evolution. *Evolution (N Y)* **69**, 3069–3081. (doi:10.1111/evo.12808)

22. Osada N, Akashi H. 2012 Mitochondrial-nuclear interactions and accelerated compensatory evolution: Evidence from the primate cytochrome c oxidase complex. *Mol Biol Evol* **29**, 337–346. (doi:10.1093/molbev/msr211)

23. Sloan DB, Havird JC, Sharbrough J. 2017 The on-again, off-again relationship between mitochondrial genomes and species boundaries. *Mol Ecol* **26**, 2212–2236. (doi:10.1111/mec.13959)

24. Piccinini G, Iannello M, Puccio G, Plazzi F, Havird JC, Ghiselli F. 2021 Mitonuclear coevolution, but not nuclear compensation, drives evolution of OXPHOS complexes in bivalves. *Mol Biol Evol* **38**, 2597–2614. (doi:10.1093/molbev/msab054)

25. Zavala-Paez M, Sutara B, Keller S, Holliday J, Fitzpatrick MC, Hamilton J. 2025 The role of cytonuclear interactions to plant adaptation across a *Populus* hybrid zone. *bioRxiv* (doi:10.1101/2025.05.13.653687)

26. Ellison CK, Niehuis O, Gadau J. 2008 Hybrid breakdown and mitochondrial dysfunction in hybrids of Nasonia parasitoid wasps. *J Evol Biol* **21**, 1844–1851. (doi:10.1111/j.1420-9101.2008.01608.x)

27. Hill GE. 2016 Mitonuclear coevolution as the genesis of speciation and the mitochondrial DNA barcode gap. *Ecol Evol* **6**, 5831–5842. (doi:10.1002/ece3.2338)

28. Burton RS, Barreto FS. 2012 A disproportionate role for mtDNA in Dobzhansky-Muller incompatibilities? *Mol Ecol* **21**, 4942–4957. (doi:10.1111/mec.12006)

29. Coyne JA. 1992 Genetics and speciation. *Nature* **355**.

30. Gershoni M, Templeton AR, Mishmar D. 2009 Mitochondrial bioenergetics as a major motive force of speciation. *BioEssays* **31**, 642–650. (doi:10.1002/bies.200800139)

31. Lima TG, Burton RS, Willett CS. 2019 Genomic scans reveal multiple mito-nuclear incompatibilities in population crosses of the copepod Tigriopus californicus. *Evolution (N Y)* **73**, 609–620. (doi:10.1111/evo.13690)

32. Forsythe ES, Nelson ADL, Beilstein MA. 2020 Biased gene retention in the face of introgression obscures species relationships. *Genome Biol Evol* **12**, 1646–1663. (doi:10.1093/GBE/EVAA149)

33. Langdon QK *et al.* 2024 Swordtail fish hybrids reveal that genome evolution is surprisingly predictable after initial hybridization. *PLoS Biol* (doi:10.1101/2023.12.21.572897)

34. Schumer M, Cui R, Powell DL, Dresner R, Rosenthal GG, Andolfatto P. 2014 High-resolution mapping reveals hundreds of genetic incompatibilities in hybridizing fish species. *Elife* **2014**. (doi:10.7554/eLife.02535)

35. Powell DL, Rose AD, Rosenthal GG. 2022 A widely-used pollutant causes reversal of conspecific mate preference in a freshwater fish. *bioRxiv* , 2022.09.07.507014. (doi:10.1101/2022.09.07.507014)

36. Powell DL, Payne C, Banerjee SM, Keegan M, Bashkirova E, Cui R, Andolfatto P, Rosenthal GG, Schumer M. 2021 The Genetic Architecture of Variation in the Sexually Selected Sword Ornament and Its Evolution in Hybrid Populations. *Current Biology* **31**, 923-935.e11. (doi:https://doi.org/10.1016/j.cub.2020.12.049)

37. Langdon QK, Powell DL, Kim B, Banerjee SM, Payne C, Dodge TO, Moran B, Fascinetto-Zago P, Schumer M. 2022 Predictability and parallelism in the contemporary evolution of hybrid genomes. *PLoS Genet* **18**. (doi:10.1371/journal.pgen.1009914)

38. Powell DL, Moran BM, Kim BY, Banerjee SM, Aguillon SM, Fascinetto-Zago P, Langdon QK, Schumer M. 2021 Two new hybrid populations expand the swordtail hybridization model system. *Evolution (N Y)* **75**, 2524–2539. (doi:10.1111/evo.14337)

39. Banerjee SM, Powell DL, Moran BM, Ramírez-Duarte WF, Langdon QK, Gunn TR, Vazquez G, Rochman C, Schumer M. 2023 Complex hybridization between deeply diverged fish species in a disturbed ecosystem. *Evolution (N Y)* **77**, 995–1005. (doi:10.1093/evolut/qpad019)

40. Powell DL *et al.* 2020 Natural hybridization reveals incompatible alleles that cause melanoma in swordtail fish. *Science (1979)* **368**, 731–736.

41. Schumer M *et al.* 2018 Natural selection interacts with recombination to shape the evolution of hybrid genomes. *Science (1979)* , 656–660.

42. Aguillon SM *et al.* 2025 Pervasive gene flow despite strong and varied reproductive barriers in swordtails. *Nat Ecol Evol* **9**, 867–878. (doi:https://doi.org/10.1038/s41559-025-02669-9)

43. Moran BM *et al.* 2024 A Lethal Genetic Incompatibility between Naturally Hybridizing Species in Mitochondrial Complex I. *Nature* (doi:https://doi.org/10.1038/s41586-023-06895-8)

44. Robles N V. *et al.* 2025 Admixture mapping reveals evidence for multiple mitonuclear incompatibilities in swordtail fish hybrids. *Mol Ecol* **34**. (doi:https://doi.org/10.1111/mec.70106)

45. Schumer M, Powell DL, Corbett-Detig R. 2020 Versatile simulations of admixture and accurate local ancestry inference with mixnmatch and ancestryinfer. *Mol Ecol Resour* **20**, 1141–1151. (doi:https://doi.org/10.1111/1755-0998.13175)

46. Dodge TO *et al.* 2024 Structural genomic variation and behavioral interactions underpin a balanced sexual mimicry polymorphism. *Current Biology* **34**, 4662-4676.e9. (doi:10.1016/j.cub.2024.08.053)

47. Pertea G, Pertea M. 2020 GFF Utilities: GffRead and GffCompare. *F1000Res* **9**, 304. (doi:10.12688/f1000research.23297.2)

48. Emms DM, Kelly S. 2019 OrthoFinder: Phylogenetic orthology inference for comparative genomics. *Genome Biol* **20**. (doi:10.1186/s13059-019-1832-y)

49. Sloan DB, Fields PD, Havird JC. 2015 Mitonuclear linkage disequilibrium in human populations. *Proceedings of the Royal Society B: Biological Sciences* **282**. (doi:10.1098/rspb.2015.1704)

50. Zhao H, Nettleton D, Soller M, Dekkers JCM. 2005 Evaluation of linkage disequilibrium measures between multi-allelic markers as predictors of linkage disequilibrium between markers and QTL. *Genet Res* **86**, 77–87. (doi:10.1017/S001667230500769X)

51. Schumer M, Brandvain Y. 2016 Determining epistatic selection in admixed populations. *Mol Ecol* **25**, 2577–2591. (doi:10.1111/mec.13641)

52. Barton NH. 2000 Genetic hitchhiking. *Philosophical Transactions of the Royal Society B: Biological Sciences* **355**, 1553–1562. (doi:10.1098/rstb.2000.0716)

53. Veller C, Edelman NB, Muralidhar P, Nowak MA. 2023 Recombination and selection against introgressed DNA. *Evolution (N Y)* **77**, 1131–1144. (doi:10.1093/evolut/qpad021)

54. Lopez KA, McDiarmid CS, Griffith SC, Lovette IJ, Hooper DM. 2021 Evaluating evidence of mitonuclear incompatibilities with the sex chromosomes in an avian hybrid zone. *Evolution (N Y)* **75**, 1395–1414. (doi:10.1111/evo.14243)

55. Low GW *et al.* 2024 Accelerated differentiation of neo-W nuclear-encoded mitochondrial genes between two climate-associated bird lineages signals potential co-evolution with mitogenomes. *Heredity (Edinb)* (doi:10.1038/s41437-024-00718-w)

56. Forsythe ES, Williams AM, Sloan DB. 2021 Genome-wide signatures of plastid-nuclear coevolution point to repeated perturbations of plastid proteostasis systems across angiosperms. *Plant Cell* **33**, 980–997. (doi:10.1093/plcell/koab021)

57. Eyre-Walker A. 2017 Mitochondrial replacement therapy: Are mito-nuclear interactions likely to be a problem? *Genetics* **205**, 1365–1372. (doi:10.1534/genetics.116.196436)

58. Cui R, Schumer M, Kruesi K, Walter R, Andolfatto P, Rosenthal GG. 2013 Phylogenomics Reveals Extensive Reticulate Evolution In Xiphophorus Fishes. *Evolution (N Y)* **67**, 2166–2179. (doi:10.1111/evo.12099)

59. Fields PD *et al.* 2024 Genome Evolution and Introgression in the New Zealand mud Snails Potamopyrgus estuarinus and Potamopyrgus kaitunuparaoa. *Genome Biol Evol* **16**. (doi:10.1093/gbe/evae091)

60. Kato S, Arakaki S, Nagano AJ, Kikuchi K, Hirase S. 2023 Genomic landscape of introgression from the ghost lineage in a gobiid fish uncovers the generality of forces shaping hybrid genomes. *Mol Ecol* (doi:10.1111/mec.17216)

61. Kwon T, Kim K, Caetano-Anolles K, Sung S, Cho S, Jeong C, Hanotte O, Kim H. 2022 Mitonuclear incompatibility as a hidden driver behind the genome ancestry of African admixed cattle. *BMC Biol* **20**. (doi:10.1186/s12915-021-01206-x)

62. Shi X *et al.* 2025 Selection increases mitonuclear DNA discordance but reconciles incompatibility in African cattle. *Mol Biol Evol* (doi:10.1093/molbev/msaf039/8005707)

63. Jensen A *et al.* 2023 Complex Evolutionary History With Extensive Ancestral Gene Flow in an African Primate Radiation. *Mol Biol Evol* **40**. (doi:10.1093/molbev/msad247)

64. Zhu J, Evans BJ. 2023 Mitonuclear Interactions and the Origin of Macaque Societies. *Genome Biol Evol* **15**. (doi:10.1093/gbe/evad010)

65. Evans BJ, Peter BM, Melnick DJ, Andayani N, Supriatna J, Zhu J, Tosi AJ. 2021 Mitonuclear interactions and introgression genomics of macaque monkeys (Macaca) highlight the influence of behaviour on genome evolution. *Proceedings of the Royal Society B: Biological Sciences* **288**. (doi:10.1098/rspb.2021.1756)

66. Bailey NP, Stevison LS. 2021 Mitonuclear conflict in a macaque species exhibiting phylogenomic discordance. *J Evol Biol* **34**, 1568–1579. (doi:10.1111/jeb.13914)

67. Sharbrough J, Havird JC, Noe GR, Warren JM, Sloan DB. 2017 The mitonuclear dimension of Neanderthal and Denisovan ancestry in modern human genomes. *Genome Biol Evol* **9**, 1567–1581. (doi:10.1093/gbe/evx114)

68. Zaidi AA, Makova KD. 2019 Investigating mitonuclear interactions in human admixed populations. *Nat Ecol Evol* **3**, 213–222. (doi:10.1038/s41559-018-0766-1)

69. Morales HE, Pavlova A, Amos N, Major R, Kilian A, Greening C, Sunnucks P. 2018 Concordant divergence of mitogenomes and a mitonuclear gene cluster in bird lineages inhabiting different climates. *Nat Ecol Evol* **2**, 1258–1267. (doi:10.1038/s41559-018-0606-3)

70. Rancilhac L *et al.* 2024 Introgression across narrow contact zones shapes the genomic landscape of phylogenetic variation in an African bird clade. (doi:10.1101/2024.08.13.607717)

71. Musher LJ, Del-Rio G, Marcondes RS, Brumfield RT, Bravo GA, Thom G. 2024 Geogenomic Predictors of Genetree Heterogeneity Explain Phylogeographic and Introgression History: A Case Study in an Amazonian Bird (Thamnophilus aethiops). *Syst Biol* **73**, 36–52. (doi:10.1093/sysbio/syad061)

433 **Supplementary Materials for the manuscript:**

435 Sensitivity of genome-wide tests for mitonuclear genetic
436 incompatibilities

438 Kuster, Shady A.[1,2], Schumer, Molly[3,4], Havird, Justin C.[5], Sloan, Daniel B.[2]

439
440   -   1: Cell and Molecular Biology, Colorado State University
441   -   2: Department of Biology, Colorado State University
442   -   3: Department of Biology, Stanford University
443   -   4: Freeman Hrabowski Fellow, Howard Hughes Medical Institute
444   -   5: Department of Integrative Biology, University of Texas at Austin

445
446 Author for correspondence: Shady A. Kuster, Shady.Kuster@colostate.edu

SUPPLEMENTAL FIGURES

Figure S1. Other metrics of mitonuclear association values for each gene class in the CALL (A and C) and CHAF (B and D) populations. Association is calculated as r² in panels A and B and as a partial correlation that accounts for genome-wide ancestry in panels C and D. Note the change of scale on the x-axis across populations and LD statistics. P-value for a one-way ANOVA testing if gene class affects LD value: CALL r² 0.44 (A), CHAF r² 0.29 (B), CALL partial correlation 0.04 (C), CHAF partial correlation 0.14 (D). Summary statistics are reported in Table S4.

Figure S2. Stacked bar histogram for the proportion of the nuclear genome derived from the major parent for each fish in A) CALL and B) CHAF. Proportion is calculated as the sum of all major parent AIMs divided by number of AIMs. Each fish is colored by which of the two segregating mitochondrial haplotypes it has (blue = X. birchmanni, fuchsia = X. malinche). Both stacked bar histograms were made with 12 bins. Note that y-axis upper limits differ between the panels.

Figure S3. Location of previously identified incompatibility genes [1] in the CALL population D' *values. Dashed line indicates genome-wide mean* D'.

Figure S4. Location of previously identified incompatibility genes [1] in the CALL population r^2 *values. Dashed line indicates genome-wide mean* r^2.

Figure S5. Location of previously identified incompatibility genes [1]in the CALL population partial correlation values. Dashed line indicates genome-wide mean partial correlation score.

Figure S6. Location of previously identified incompatibility genes [1] in the CHAF population D' *values. Dashed line indicates genome-wide mean* D'.

Figure S7. Location of previously identified incompatibility genes [1]in the CHAF population r^2*values. Dashed line indicates genome-wide mean* r^2.

Figure S8. Location of previously identified incompatibility genes [1]in the CHAF population partial correlation values. Dashed line indicates genome-wide mean partial correlation score.

Figure S9. Location of previously identified incompatibility genes [1] in the HUEX-STAC population X. cortezi *allele frequency values. Dashed line indicates genome-wide mean allele frequency.*

Figure S10. Histogram demonstrating permuted differences in mean allele frequency between all interacting n-mt genes and the mean of all other genes. Red line represents the observed difference in interacting n-mt gene class and all other classes. In this permutation test, we found the interacting n-mt gene class to be different from the other two gene classes ($p = 9 \times 10^{-4}$), supporting the parametric analysis (ANOVA) in this population.

Figure S11. Non-synonymous substitution model for the CALL population for the A) D' *and B)* r² *metrics. In a model that accounts for an interaction between gene class and non-synonymous substitutions, interacting n-mt genes have a stronger relationship to the* r² *(p = 0.019) but not the* D' *(p = 0.092) metric (Table S9), but see points driving this relationship in Table S10 and summary statistics in Table S7.*

Figure S12. Non-synonymous substitution model for the CHAF. population for the A) D' *and B)* r² *metrics. In a model that accounts for an interaction between gene class and non-synonymous substitutions, interacting n-mt genes have a stronger relationship to the* r² *(p = 0.027) but not the* D' *(p = 0.37) metric (Table S9), but see points driving this relationship in Table S10 and summary statistics in Table S7.*

SUPPLEMENTAL TABLES

Table S1. Studies investigating differential effects of n-mt and non-n-mt gene groups or genomic windows.

Citation	Admixture category[a]	Hybridizing groups	Segregating mtDNA?[b]	Disproportionate n-mt gene effect?[c]
Mollusks				
Fields et al. 2024 [2]	Historical introgression	*Potamopyrgus estuarinus* and *P. kaitunuparaoa*	No	Yes
Fish				
Kato et al. 2023 [3]	Historical introgression	*Chaenogobius annularis* and a ghost lineage	No	No
Mammals				
Kwon et al. 2022 [4]	Interspecific hybridization	*Bos taurus* and *B. indicus*	No	Yes
Shi et al. 2025 [5]	Interspecific hybridization	*B. taurus* and *B. indicus*	No	No[d]
Jensen et al. 2023 [6]	Historical introgression	*Allochrocebus* and *Chlorocebus* lineages	No	Yes
Zhu & Evans 2023 [7]	Historical introgression	*Macaca arctoides* and *M. fascicularis aurea*	No	No
Evans et al. 2021 [8]	Interspecies individual hybrid	*M. tonkeana* and *M. maura*	This individual has *M. tonkeana* mtDNA	N/A[f]
	Interspecies individual hybrid	*M. tonkeana* and *M. hecki*	This individual has *M. hecki* mtDNA	N/A[f]
Bailey & Stevison 2021 [9]	Interspecific hybridization	*M. arctoides* and *M. sinica* or *M. fascicularis*	No	No
Sloan et al. 2015 [10]	Intraspecific admixture between populations	*Homo sapiens*	Yes	No
Sharbrough et al. 2017 [11]	Historical introgression	*H. sapiens* and *H. neanderthalensis*	No	Yes

	Historical introgression	*H. sapiens* and Denisovans	No	No
	Intraspecific admixture between populations	*H. sapiens* African American and European	Yes	Yes
	Intraspecific admixture between populations	*H. sapiens* Puerto Rican and European	Yes	Yes
Zaidi & Makova 2019 [12]	Intraspecific admixture between populations	*H. sapiens* Caribbean and European	Yes	No
	Intraspecific admixture between populations	*H. sapiens* Columbian and European	Yes	No
	Intraspecific admixture between populations	*H. sapiens* Mexican and European	Yes	No
	Intraspecific admixture between populations	*H. sapiens* Peruvian and European	Yes	No

Birds

Lopez et al. 2021 [13]	Intraspecific admixture between populations	*Poephila acuticauda acuticauda* and *P. a. hecki*	Yes	Yes[e]
Morales et al. 2018 [14]	Intraspecific admixture between populations	northern and southern populations of *Eopsaltria australis*	Yes	Yes, but see [15]
Low et al. 2024 [15]	Intraspecific admixture between populations	inland and coastal mitochondrial lineages of *E. australis*	Yes	Yes[e]
Rancilhac et al. 2024 [16]	Interspecific hybridization	*Pogoniulus pusillus* and *P. chrysoconus*	Yes	No
Musher et al. 2024 [17]	Intraspecific admixture	*Thamnophilus aethiops*	Yes	No

These papers were collected, using two groups of search terms to identify published articles that included similar comparisons between n-mt and non-n-mt genes in the context of hybridization. The first search terms were "n-mt" AND ("reproductive isolation" OR "speciation") AND "hybrid" on Google Scholar in October 2024. Inclusion criteria for the resulting hits were that the work (1) categorized genes into n-mt and non-n-mt classes, (2) measured association between n-mt genetic variants and mitochondrial haplotypes, and (3) was focused on current or ancient admixture of genomes (via hybridization or introgression). Papers were excluded if the only aim was to use phylogenetic analyses to study mutation rates or phylogenetic discordance between mtDNA and non-n-mt loci. This search (conducted on 18 October 2024) identified 193 possible articles, but only 10 met our inclusion criteria. We then performed a separate search using the terms "mito*nuclear" AND ("reproductive isolation" OR "speciation."), returning 57 articles, but none that met our inclusion criteria. We supplemented this list with six additional papers that were not returned as hits in the above searches but that we encountered as citations in other studies or during unrelated literature searches. The combination of search terms and narrow inclusion criteria likely excluded some relevant studies from this table, but the goal was not to capture all studies pertinent to genomic analysis of mitonuclear interactions.

[a]Hybridization systems were categorized as being either intraspecific admixture between populations, interspecific hybridization, or historical introgression to indicate timing of hybridization and how removed from initial admixture a population may be.
[b]Indication of whether the mtDNA from the hybrid source populations or species are still segregating in the hybrid system investigated.
[c]Investigations that found n-mt gene groups or genomic windows to share ancestry more than non-n-mt groups or windows were given a 'yes' to indicate whole-genome based methods successfully identified effects of n-mt selection.
[d]This study instead found evidence of enrichment for mismatch of n-mt and mtDNA ancestry.
[e]These studies find positive relationships between n-mt and mtDNA ancestry, but these relationships also include matching of sex chromosome ancestry or that n-mt genes are located in regions of a chromosome associated with sex determination.
[f]Because these studies only determined n-mt disproportionality within a single individual, they are omitted from our investigation into whether fixing a mitochondrial haplotype affects the sensitivity of genome-wide tests of shared n-mt and mtDNA ancestry.

Table S2. List of human interacting n-mt genes used in this study, modified from [10].

Uniprot ID	Entrez symbol	Gene symbol	Gene Name
P82663	64432	MRPS25	mitochondrial ribosomal protein S25
O15239	4694	NDUFA1	NADH:ubiquinone oxidoreductase subunit A1
Q9Y3B7	65003	MRPL11	mitochondrial ribosomal protein L11
O43678	4695	NDUFA2	NADH:ubiquinone oxidoreductase subunit A2
Q96EL2	64951	MRPS24	mitochondrial ribosomal protein S24
O95167	4696	NDUFA3	NADH:ubiquinone oxidoreductase subunit A3
Q9BYD2	65005	MRPL9	mitochondrial ribosomal protein L9
Q9BYD6	65008	MRPL1	mitochondrial ribosomal protein L1
Q9Y6G3	28977	MRPL42	mitochondrial ribosomal protein L42
P14854	1340	COX6B1	cytochrome c oxidase subunit 6B1
Q9Y2R9	51081	MRPS7	mitochondrial ribosomal protein S7
Q9NVS2	55168	MRPS18A	mitochondrial ribosomal protein S18A
P12074	1337	COX6A1	cytochrome c oxidase subunit 6A1
O00217	4728	NDUFS8	NADH:ubiquinone oxidoreductase core subunit S8
Q9BYN8	64949	MRPS26	mitochondrial ribosomal protein S26
P19404	4729	NDUFV2	NADH:ubiquinone oxidoreductase core subunit V2
Q86Y39	126328	NDUFA11	NADH:ubiquinone oxidoreductase subunit A11
Q9Y676	28973	MRPS18B	mitochondrial ribosomal protein S18B
O14949	27089	UQCRQ	ubiquinol-cytochrome c reductase complex III subunit VII
P30049	513	ATP5F1D	ATP synthase F1 subunit delta
P56381	514	ATP5F1E	ATP synthase F1 subunit epsilon
Q9H9J2	65080	MRPL44	mitochondrial ribosomal protein L44
P24539	515	ATP5PB	ATP synthase peripheral stalk-membrane subunit b
O75306	4720	NDUFS2	NADH:ubiquinone oxidoreductase core subunit S2
Q9UI09	55967	NDUFA12	NADH:ubiquinone oxidoreductase subunit A12
Q96GC5	51642	MRPL48	mitochondrial ribosomal protein L48
P05496	516	ATP5MC1	ATP synthase membrane subunit c locus 1
Q06055	517	ATP5MC2	ATP synthase membrane subunit c locus 2
O75489	4722	NDUFS3	NADH:ubiquinone oxidoreductase core subunit S3
P09001	11222	MRPL3	mitochondrial ribosomal protein L3
P48201	518	ATP5MC3	ATP synthase membrane subunit c locus 3
P49821	4723	NDUFV1	NADH:ubiquinone oxidoreductase core subunit V1
Q5JTZ9	57505	AARS2	alanyl-tRNA synthetase 2, mitochondrial
O43181	4724	NDUFS4	NADH:ubiquinone oxidoreductase subunit S4
O43920	4725	NDUFS5	NADH:ubiquinone oxidoreductase subunit S5
Q9Y3D9	51649	MRPS23	mitochondrial ribosomal protein S23
Q02221	1339	COX6A2	cytochrome c oxidase subunit 6A2
O75380	4726	NDUFS6	NADH:ubiquinone oxidoreductase subunit S6
P82914	64960	MRPS15	mitochondrial ribosomal protein S15
Q9NRX2	63875	MRPL17	mitochondrial ribosomal protein L17
P82912	64963	MRPS11	mitochondrial ribosomal protein S11
P15954	1350	COX7C	cytochrome c oxidase subunit 7C
P82933	64965	MRPS9	mitochondrial ribosomal protein S9
P82932	64968	MRPS6	mitochondrial ribosomal protein S6
Q9BYC9	55052	MRPL20	mitochondrial ribosomal protein L20
P82664	55173	MRPS10	mitochondrial ribosomal protein S10
P82675	64969	MRPS5	mitochondrial ribosomal protein S5
P24311	1349	COX7B	cytochrome c oxidase subunit 7B
O95363	10667	FARS2	phenylalanyl-tRNA synthetase 2, mitochondrial
P14406	1347	COX7A2	cytochrome c oxidase subunit 7A2
Q9Y2R5	51373	MRPS17	mitochondrial ribosomal protein S17
P24310	1346	COX7A1	cytochrome c oxidase subunit 7A1
P56385	521	ATP5ME	ATP synthase membrane subunit e
P09669	1345	COX6C	cytochrome c oxidase subunit 6C
Q9Y291	51650	MRPS33	mitochondrial ribosomal protein S33
P18859	522	ATP5PF	ATP synthase peripheral stalk subunit F6
Q9BZE1	51253	MRPL37	mitochondrial ribosomal protein L37
P82673	60488	MRPS35	mitochondrial ribosomal protein S35

Q9NSE4	55699	IARS2	isoleucyl-tRNA synthetase 2, mitochondrial
Q9UDW1	29796	UQCR10	ubiquinol-cytochrome c reductase, complex III subunit X
Q4U2R6	51258	MRPL51	mitochondrial ribosomal protein L51
O00483	4697	NDUFA4	NDUFA4 mitochondrial complex associated
Q9UHN1	11232	POLG2	DNA polymerase gamma 2, accessory subunit
Q16718	4698	NDUFA5	NADH:ubiquinone oxidoreductase subunit A5
P56181	4731	NDUFV3	NADH:ubiquinone oxidoreductase subunit V3
Q9BRJ2	84311	MRPL45	mitochondrial ribosomal protein L45
Q7Z2W9	219927	MRPL21	mitochondrial ribosomal protein L21
P54098	5428	POLG	DNA polymerase gamma, catalytic subunit
P82650	56945	MRPS22	mitochondrial ribosomal protein S22
O00411	5442	POLRMT	RNA polymerase mitochondrial
Q8IXM3	64975	MRPL41	mitochondrial ribosomal protein L41
Q9NQ50	64976	MRPL40	mitochondrial ribosomal protein L40
Q9P0J6	64979	MRPL36	mitochondrial ribosomal protein L36
Q96DV4	64978	MRPL38	mitochondrial ribosomal protein L38
Q9Y2Q9	28957	MRPS28	mitochondrial ribosomal protein S28
Q99766	27109	DMAC2L	distal membrane arm assembly component 2 like
Q9HD33	57129	MRPL47	mitochondrial ribosomal protein L47
Q8TCC3	51263	MRPL30	mitochondrial ribosomal protein L30
Q9Y3D3	51021	MRPS16	mitochondrial ribosomal protein S16
Q9P015	29088	MRPL15	mitochondrial ribosomal protein L15
P25705	498	ATP5F1A	ATP synthase F1 subunit alpha
Q8N5N7	54534	MRPL50	mitochondrial ribosomal protein L50
Q9Y3D5	51023	MRPS18C	mitochondrial ribosomal protein S18C
Q9P0M9	51264	MRPL27	mitochondrial ribosomal protein L27
Q9UGM6	10352	WARS2	tryptophanyl tRNA synthetase 2, mitochondrial
Q9BQC6	78988	MRPL57	mitochondrial ribosomal protein L57
P48047	539	ATP5PO	ATP synthase peripheral stalk subunit OSCP
O75251	374291	NDUFS7	NADH:ubiquinone oxidoreductase core subunit S7
Q9NP81	54938	SARS2	seryl-tRNA synthetase 2, mitochondrial
O75947	10476	ATP5PD	ATP synthase peripheral stalk subunit d
Q9NX14	54539	NDUFB11	NADH:ubiquinone oxidoreductase subunit B11
O75964	10632	ATP5MG	ATP synthase membrane subunit g
P49590	23438	HARS2	histidyl-tRNA synthetase 2, mitochondrial
P20674	9377	COX5A	cytochrome c oxidase subunit 5A
Q9BQ48	64981	MRPL34	mitochondrial ribosomal protein L34
Q92665	10240	MRPS31	mitochondrial ribosomal protein S31
Q9BYC8	64983	MRPL32	mitochondrial ribosomal protein L32
Q7L3T8	25973	PARS2	prolyl-tRNA synthetase 2, mitochondrial
O60783	63931	MRPS14	mitochondrial ribosomal protein S14
Q6YFQ2	125965	COX6B2	cytochrome c oxidase subunit 6B2
Q7Z7H8	124995	MRPL10	mitochondrial ribosomal protein L10
Q9NYK5	54148	MRPL39	mitochondrial ribosomal protein L39
Q15031	23395	LARS2	leucyl-tRNA synthetase 2, mitochondrial
Q7Z4L0	341947	COX8C	cytochrome c oxidase subunit 8C
Q9NWU5	29093	MRPL22	mitochondrial ribosomal protein L22
P49406	9801	MRPL19	mitochondrial ribosomal protein L19
Q9NP92	10884	MRPS30	mitochondrial ribosomal protein S30
P41250	2617	GARS1	glycyl-tRNA synthetase 1
Q9NX20	54948	MRPL16	mitochondrial ribosomal protein L16
Q9NZE8	51318	MRPL35	mitochondrial ribosomal protein L35
Q16540	6150	MRPL23	mitochondrial ribosomal protein L23
Q9Y2S7	26073	POLDIP2	DNA polymerase delta interacting protein 2
O14957	10975	UQCR11	ubiquinol-cytochrome c reductase, complex III subunit XI
Q96I59	79731	NARS2	asparaginyl-tRNA synthetase 2, mitochondrial
Q96GW9	92935	MARS2	methionyl-tRNA synthetase 2, mitochondrial
Q13084	10573	MRPL28	mitochondrial ribosomal protein L28
Q7Z7F7	128308	MRPL55	mitochondrial ribosomal protein L55
P08574	1537	CYC1	cytochrome c1
O75394	9553	MRPL33	mitochondrial ribosomal protein L33
P56134	9551	ATP5MF	ATP synthase membrane subunit f

Q5JPH6	124454	EARS2	glutamyl-tRNA synthetase 2, mitochondrial
Q5T160	57038	RARS2	arginyl-tRNA synthetase 2, mitochondrial
Q9H0U6	29074	MRPL18	mitochondrial ribosomal protein L18
P22695	7385	UQCRC2	ubiquinol-cytochrome c reductase core protein 2
P31930	7384	UQCRC1	ubiquinol-cytochrome c reductase core protein 1
P47985	7386	UQCRFS1	ubiquinol-cytochrome c reductase, Rieske iron-sulfur polypeptide 1
P07919	7388	UQCRH	ubiquinol-cytochrome c reductase hinge protein
Q9BYD1	28998	MRPL13	mitochondrial ribosomal protein L13
Q96A35	79590	MRPL24	mitochondrial ribosomal protein L24
Q92552	23107	MRPS27	mitochondrial ribosomal protein S27
O95299	4705	NDUFA10	NADH:ubiquinone oxidoreductase subunit A10
Q6P1L8	64928	MRPL14	mitochondrial ribosomal protein L14
Q9BW92	80222	TARS2	threonyl-tRNA synthetase 2, mitochondrial
O14561	4706	NDUFAB1	NADH:ubiquinone oxidoreductase subunit AB1
Q96KJ9	84701	COX4I2	cytochrome c oxidase subunit 4I2
O95178	4708	NDUFB2	NADH:ubiquinone oxidoreductase subunit B2
O43676	4709	NDUFB3	NADH:ubiquinone oxidoreductase subunit B3
Q9Y2Z4	51067	YARS2	tyrosyl-tRNA synthetase 2
Q5T653	51069	MRPL2	mitochondrial ribosomal protein L2
Q8N983	84545	MRPL43	mitochondrial ribosomal protein L43
Q9Y375	51103	NDUFAF1	NADH:ubiquinone oxidoreductase complex assembly factor 1
P56556	4700	NDUFA6	NADH:ubiquinone oxidoreductase subunit A6
O95182	4701	NDUFA7	NADH:ubiquinone oxidoreductase subunit A7
P51970	4702	NDUFA8	NADH:ubiquinone oxidoreductase subunit A8
Q15046	3735	KARS1	lysyl-tRNA synthetase 1
Q86TS9	122704	MRPL52	mitochondrial ribosomal protein L52
Q16795	4704	NDUFA9	NADH:ubiquinone oxidoreductase subunit A9
P82930	65993	MRPS34	mitochondrial ribosomal protein S34
Q5ST30	57176	VARS2	valyl-tRNA synthetase 2, mitochondrial
Q9H2W6	26589	MRPL46	mitochondrial ribosomal protein L46
P52815	6182	MRPL12	mitochondrial ribosomal protein L12
P82921	54460	MRPS21	mitochondrial ribosomal protein S21
O15235	6183	MRPS12	mitochondrial ribosomal protein S12
P13073	1327	COX4I1	cytochrome c oxidase subunit 4I1
O96000	4716	NDUFB10	NADH:ubiquinone oxidoreductase subunit B10
Q6PI48	55157	DARS2	aspartyl-tRNA synthetase 2, mitochondrial
O43677	4717	NDUFC1	NADH:ubiquinone oxidoreductase subunit C1
Q9BYD3	51073	MRPL4	mitochondrial ribosomal protein L4
Q96EL3	116540	MRPL53	mitochondrial ribosomal protein L53
O95298	4718	NDUFC2	NADH:ubiquinone oxidoreductase subunit C2
P28331	4719	NDUFS1	NADH:ubiquinone oxidoreductase core subunit S1
Q13405	740	MRPL49	mitochondrial ribosomal protein L49
Q6P161	116541	MRPL54	mitochondrial ribosomal protein L54
Q9P0J0	51079	NDUFA13	NADH:ubiquinone oxidoreductase subunit A13
O95168	4710	NDUFB4	NADH:ubiquinone oxidoreductase subunit B4
O43674	4711	NDUFB5	NADH:ubiquinone oxidoreductase subunit B5
Q9Y399	51116	MRPS2	mitochondrial ribosomal protein S2
P06576	506	ATP5F1B	ATP synthase F1 subunit beta
O95139	4712	NDUFB6	NADH:ubiquinone oxidoreductase subunit B6
P17568	4713	NDUFB7	NADH:ubiquinone oxidoreductase subunit B7
P10606	1329	COX5B	cytochrome c oxidase subunit 5B
O95169	4714	NDUFB8	NADH:ubiquinone oxidoreductase subunit B8
P36542	509	ATP5F1C	ATP synthase F1 subunit gamma
Q9Y6M9	4715	NDUFB9	NADH:ubiquinone oxidoreductase subunit B9

Table S3. Pre-filtered AIMs and genes (in parentheses) belonging to each gene class. The orthogroup row indicates how many orthogroups containing X. birchmanni *protein sequences were found for each gene class.*

Population	Interacting n-mt	Non-interacting n-mt	Non-n-mt
Orthogroups	159	822	15,545
CALL AIMs (unique genes)	954 (148)	8,580 (883)	287,834 (19,312)
CHAF AIMs (unique genes)	952 (148)	8,575 (883)	287,592 (19,307)
HUEX-STAC AIMs (unique genes)	997 (162)	9,399 (947)	319,679 (20,544)

Table S4. Summary statistics of mitonuclear association values for each hybrid population.

Population	Statistic	Gene Class	Count	Mean	Standard Deviation
CALL	*D'*	Interacting n-mt	148	0.405	0.116
		Non-interacting n-mt	883	0.392	0.105
		Non-n-mt	19,312	0.393	0.114
	r^2	Interacting n-mt	148	0.094	0.064
		Non-interacting n-mt	883	0.094	0.059
		Non-n-mt	19,312	0.097	0.063
	Partial correlation	Interacting n-mt	148	0.008	0.084
		Non-interacting n-mt	881	0.005	0.084
		Non-n-mt	19,293	0.013	0.087
CHAF	*D'*	Interacting n-mt	147	0.428	0.176
		Non-interacting n-mt	883	0.421	0.189
		Non-n-mt	19,296	0.420	0.198
	r^2	Interacting n-mt	147	0.033	0.024
		Non-interacting n-mt	883	0.030	0.024
		Non-n-mt	19,296	0.0315	0.025
	Partial correlation	Interacting n-mt	148	0.021	0.078
		Non-interacting n-mt	882	0.007	0.077
		Non-n-mt	19,297	0.009	0.080
HUEX-STAC	Allele Frequency	Interacting n-mt	162	0.880	0.156
		Non-interacting n-mt	947	0.822	0.222
		Non-n-mt	20543	0.831	0.211

Table S5. Results of Fisher's exact test for enrichment of interacting n-mt genes in top 1% of genome-wide association statistics.

Population	Test	Result
CALL	One-sided Fisher's Exact Test for Count Data on D' values	$p = 0.33$
	One-sided Fisher's Exact Test for Count Data r^2 values	$p = 0.76$
	One-sided Fisher's Exact Test for Count Data partial correlation metric	$p = 0.03$
CHAF	One-sided Fisher's Exact Test for Count Data on D' values	$p = 0.31$
	One-sided Fisher's Exact Test for Count Data r^2 values	$p = 0.42$
	One-sided Fisher's Exact Test for Count Data partial correlation metric	$p = 0.25$

Table S6. Measures of mitonuclear association for incompatible n-mt genes previously identified by Robles et al. [1].

Gene[a]	Incompatible mtDNA[b]	CALL[c]			CHAF[c]			HUEX-STAC[c]
		D'	r^2	Partial correlation	D'	r^2	Partial correlation	Allele frequency
NDUFA13 (g3279.t1)	*X. malinche*	0.58 (1.61)	0.20 (1.70)	0.13 (1.30)	0.52 (0.50)	0.13 (4.02)	0.14 (1.57)	0.99 (0.74)
MTERF4 (g3322.t1)	*X. malinche*	0.60 (1.79)	0.14 (0.76)	0.15 (1.54)	0.63 (1.04)	0.19 (6.24)	0.25 (3.00)	0.97 (0.66)
ATP5MG (g3595.t1)	*X. malinche*	0.57 (1.52)	0.19 (1.54)	0.12 (1.24)	0.45 (0.13)	0.05 (0.90)	-0.02 (-0.32)	0.95 (0.57)
NDUFS5 (g8054.t1)	bidirectional	0.68 (2.54)	0.42 (5.17)	0.35 (3.87)	0.76 (1.71)	0.17 (5.47)	0.21 (2.46)	0.98 (0.70)
MMUT (g15308.t1)	bidirectional	0.48 (0.78)	0.04 (-0.85)	0.17 (1.82)	0.04 (-1.94)	0.00 (-1.24)	0.06 (0.69)	0.51 (-1.51)
SMIM8 (g15060.t1)	*X. birchmanni*	0.64 (2.21)	0.31 (3.39)	0.18 (1.93)	0.69 (1.39)	0.06 (0.96)	0.04 (0.38)	ND
LYRM2 (g15051.t1)	*X. birchmanni*	0.55 (1.36)	0.25 (2.43)	0.17 (1.83)	0.60 (0.92)	0.04 (0.52)	0.10 (1.16)	0.92 (0.42)
RMDN3 (g15056.t1)	*X. birchmanni*	0.62 (1.99)	0.29 (3.09)	0.17 (1.79)	0.61 (0.94)	0.04 (0.38)	0.04 (0.42)	0.92 (0.41)
UQCRC2 (g11575.t1)	*X. birchmanni*	0.34 (-0.49)	0.05 (-0.78)	-0.03 (-0.47)	0.20 (-1.60)	0.00 (-1.20)	-0.04 (-0.59)	0.93 (0.44)

[a]Human gene names are given with the X. birchmanni annotation ID indicated in parentheses. UQCR11 has also been identified as a gene involved in mitonuclear incompatibilities [41], but it was not included in the annotation used in this analysis.

[b]Species listed indicate the mtDNA that is incompatible with the other species' n-mt allele. For example, X. malinche means the X. malinche mtDNA is incompatible with X. birchmanni ancestry at NDUFA13. Bidirectional means the incompatibility occurs for both mtDNAs..

[c]Values are reported for each metric of mitonuclear association (D', r^2, partial correlation, and allele frequency), and the corresponding Z-score for each value is indicated in parentheses, which was calculated by subtracting the mean value and dividing by the standard deviation.

Table S7. Summary statistics for the non-synonymous substitution model. NS is the count of non-synonymous substitutions per coding sequence divided by the coding sequence length.

	Gene class	Number of genes	Mean NS ± standard deviation
CALL	Interacting n-mt	63	0.00085 ± 0.0012
	Non-interacting n-mt	473	0.00051 ± 0.0007
	Non-n-mt	11330	0.00053 ± 0.0007
CHAF	Interacting n-mt	63	0.00085 ± 0.0012
	Non-interacting n-mt	473	0.00051 ± 0.0007
	Non-n-mt	11324	0.00053 ± 0.0007
HUEX-STAC	Interacting n-mt	60	0.00089 ± 0.0011
	Non-interacting n-mt	461	0.00046 ± 0.0007
	Non-n-mt	10587	0.00041 ± 0.0006

Table S8. Type III ANOVA results for non-synonymous substitution models

	Predictor variables	p-value
D'		
CALL	**Intercept**	**< 2e-16**
	Gene class	0.40
	Non-synonymous substitutions / CDS (NS)	0.16
	Gene class * NS	0.11
CHAF	**Intercept**	**< 2e-16**
	Gene class	0.93
	Non-synonymous substitutions / CDS (NS)	0.54
	Gene class * NS	0.35
r²		
CALL	**Intercept**	**< 2e-16**
	Gene class	**0.042**
	Non-synonymous substitutions / CDS (NS)	0.42
	Gene class * NS	**0.011**
CHAF	**Intercept**	**< 2e-16**
	Gene class	0.80
	Non-synonymous substitutions / CDS (NS)	0.12
	Gene class * NS	0.085
Partial correlation		
CALL	**Intercept**	**< 2e-16**
	Gene class	0.062
	Non-synonymous substitutions / CDS (NS)	0.23
	Gene class * NS	0.073
CHAF	**Intercept**	**< 2e-16**
	Gene class	0.67
	Non-synonymous substitutions / CDS (NS)	**0.0028**
	Gene class * NS	0.67
Allele Frequency		
HUEX-STAC	**Intercept**	**< 2e-16**
	Gene class	0.53
	Non-synonymous substitutions / CDS (NS)	**0.0034**
	Gene class * NS	0.83

Table S9. Results of summary() *function testing the difference between interacting n-mt genes' relationship between non-synonymous substitutions and mitonuclear LD and the non-n-mt genes' relationship. Please see attached excel.*

Table S10. Influential points of non-synonymous substitution analysis

Mitonuclear LD metric	Influential genes	Mitonuclear LD value	Normalized non-synonymous substitution count	Removed significance of interaction?[a]	Removed significance of interacting n-mt interaction?[b]
CALL					
r^2	*NDUFS5*	0.42	0.0031	Y	Y
Partial correlation	*NDUFS5*	0.35	0.0031	Y	Y
CHAF					
r^2	*NDUFS5*	0.17	0.0031	Y	Y
	NDUFA13	0.13	0.0023	Y	Y
	ATP5MG	0.45	0.0064	Y	N
	LOC102221627[c]	0.003	0.0031	N	N

The above points were identified as influential (i.e., significantly shaping the slope of the regression) in a model of mitonuclear LD vs an interaction between gene class and non-synonymous substitutions per coding sequence length.

[a]When this gene was removed from the model, did the interaction between gene class and normalized non-synonymous substitution count become non-significant?

[b]When this gene was removed from the model, did the interacting n-mt gene class no longer have a significantly sharper slope than the non-n-mt gene class?

[c]LOC102221627 is annotated as "cytochrome b-c1 complex subunit 8"

REFERENCES

1. Robles N V. *et al.* 2025 Admixture mapping reveals evidence for multiple mitonuclear incompatibilities in swordtail fish hybrids. *Mol Ecol* **34**. (doi:https://doi.org/10.1111/mec.70106)
2. Fields PD *et al.* 2024 Genome Evolution and Introgression in the New Zealand mud Snails Potamopyrgus estuarinus and Potamopyrgus kaitunuparaoa. *Genome Biol Evol* **16**. (doi:10.1093/gbe/evae091)
3. Kato S, Arakaki S, Nagano AJ, Kikuchi K, Hirase S. 2023 Genomic landscape of introgression from the ghost lineage in a gobiid fish uncovers the generality of forces shaping hybrid genomes. *Mol Ecol* (doi:10.1111/mec.17216)
4. Kwon T, Kim K, Caetano-Anolles K, Sung S, Cho S, Jeong C, Hanotte O, Kim H. 2022 Mitonuclear incompatibility as a hidden driver behind the genome ancestry of African admixed cattle. *BMC Biol* **20**. (doi:10.1186/s12915-021-01206-x)
5. Shi X *et al.* 2025 Selection increases mitonuclear DNA discordance but reconciles incompatibility in African cattle. *Mol Biol Evol* (doi:10.1093/molbev/msaf039/8005707)
6. Jensen A *et al.* 2023 Complex Evolutionary History With Extensive Ancestral Gene Flow in an African Primate Radiation. *Mol Biol Evol* **40**. (doi:10.1093/molbev/msad247)
7. Zhu J, Evans BJ. 2023 Mitonuclear Interactions and the Origin of Macaque Societies. *Genome Biol Evol* **15**. (doi:10.1093/gbe/evad010)
8. Evans BJ, Peter BM, Melnick DJ, Andayani N, Supriatna J, Zhu J, Tosi AJ. 2021 Mitonuclear interactions and introgression genomics of macaque monkeys (Macaca) highlight the influence of behaviour on genome evolution. *Proceedings of the Royal Society B: Biological Sciences* **288**. (doi:10.1098/rspb.2021.1756)
9. Bailey NP, Stevison LS. 2021 Mitonuclear conflict in a macaque species exhibiting phylogenomic discordance. *J Evol Biol* **34**, 1568–1579. (doi:10.1111/jeb.13914)
10. Sloan DB, Fields PD, Havird JC. 2015 Mitonuclear linkage disequilibrium in human populations. *Proceedings of the Royal Society B: Biological Sciences* **282**. (doi:10.1098/rspb.2015.1704)
11. Sharbrough J, Havird JC, Noe GR, Warren JM, Sloan DB. 2017 The mitonuclear dimension of Neanderthal and Denisovan ancestry in modern human genomes. *Genome Biol Evol* **9**, 1567–1581. (doi:10.1093/gbe/evx114)
12. Zaidi AA, Makova KD. 2019 Investigating mitonuclear interactions in human admixed populations. *Nat Ecol Evol* **3**, 213–222. (doi:10.1038/s41559-018-0766-1)
13. Lopez KA, McDiarmid CS, Griffith SC, Lovette IJ, Hooper DM. 2021 Evaluating evidence of mitonuclear incompatibilities with the sex chromosomes in an avian hybrid zone. *Evolution (N Y)* **75**, 1395–1414. (doi:10.1111/evo.14243)
14. Morales HE, Pavlova A, Amos N, Major R, Kilian A, Greening C, Sunnucks P. 2018 Concordant divergence of mitogenomes and a mitonuclear gene cluster in bird lineages inhabiting different climates. *Nat Ecol Evol* **2**, 1258–1267. (doi:10.1038/s41559-018-0606-3)
15. Low GW *et al.* 2024 Accelerated differentiation of neo-W nuclear-encoded mitochondrial genes between two climate-associated bird lineages signals potential co-evolution with mitogenomes. *Heredity (Edinb)* (doi:10.1038/s41437-024-00718-w)
16. Rancilhac L *et al.* 2024 Introgression across narrow contact zones shapes the genomic landscape of phylogenetic variation in an African bird clade. (doi:10.1101/2024.08.13.607717)
17. Musher LJ, Del-Rio G, Marcondes RS, Brumfield RT, Bravo GA, Thom G. 2024 Geogenomic Predictors of Genetree Heterogeneity Explain Phylogeographic and Introgression History: A

Case Study in an Amazonian Bird (Thamnophilus aethiops). *Syst Biol* **73**, 36–52. (doi:10.1093/sysbio/syad061)