A reference panel for linkage disequilibrium and genotype imputation using whole-genome sequencing data from 2,680 participants across India

Zheng Li[1,8], Wei Zhao[2,3,8*], Xiang Zhou[1], Yuk Yee Leung[4], Gerard D. Schellenberg[4], Li-San Wang[4], Sharmistha Dey[5], Jinkook Lee[6], Jennifer A. Smith[2,3,9*], Aparajit B. Dey[7,9], Sharon L.R. Kardia[3,9]

1) Department of Biostatistics, School of Public Health, University of Michigan, Ann Arbor, Michigan, United States of America
2) Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, Michigan, United States of America
3) Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, Michigan, United States of America
4) Penn Neurodegeneration Genomics Center, Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania
5) Department of Biophysics, All India Institute of Medical Sciences, New Delhi, India.
6) Department of Economics, University of Southern California, Los Angeles, California, USA.
7) Venu Geriatric Institute, New Delhi, India.
8) These authors contribute equally to this manuscript
9) Senior author

*Correspondence to Wei Zhao (zhaowei@umich.edu) and Jennifer A. Smith (smjenn@umich.edu)

Keywords: whole-genome sequencing, South Asian, Indian, reference panel, linkage disequilibrium, LD blocks, genotype imputation.

Abstract

India is the most populous country globally, yet genetic studies involving Indian individuals remain limited. The Indian population is composed of many founder groups and has a mixed genetic ancestry, including an ancestral component not observed anywhere outside of India. This presents a unique opportunity to uncover novel disease variants and develop more tailored medical interventions. To facilitate genetic research in India, a crucial first step is to create a foundational resource that serves as a benchmark for future population studies and methods development. To this end, we have constructed the largest and most nationally representative linkage disequilibrium and genotype imputation reference panels in India to date, using high-coverage whole-genome sequencing data of 2,680 Indian participants from the Longitudinal Aging Study in India-Harmonized Diagnostic Assessment of Dementia (LASI-DAD). As an LD reference panel, LASI-DAD includes 69.5 million variants, representing 170% and 213% increases relative to the 1000 Genomes Project (1000G) and TOP-LD panels, respectively. Besides serving as an LD lookup panel, LASI-DAD facilitates various statistical analyses that rely on precise LD estimates. In a polygenic risk score (PRS) analysis, LASI-DAD improved the predictive performance of PRS by 2.1% to 35.1% across traits and studies. As an imputation reference panel, LASI-DAD improved the imputation accuracy by 3% to 101% (mean = 38%) compared to the TOPMed panel (Version R3) and by 3% to 73% (mean = 27%) compared to the Genome Asia Pilot (GAsP) panel across different minor allele frequencies. The LASI-DAD reference panel is publicly available to benefit future studies.

Introduction

India, with a population of over 1.4 billion, is home to more than 4,500 anthropologically defined groups that are characterized by a rich diversity in languages, religions, and cultures.[1] The Indian population largely descends from a mixture of two genetically divergent populations: Ancestral North Indians (ANI), who are related to Central Asians, Middle Easterners, Caucasians, and Europeans; and Ancestral South Indians (ASI), who are not related to any groups outside the subcontinent and are only distantly related to the indigenous Andamanese Islanders.[2, 3] Due to this unique ASI component, the Indian population has a distinctive genetic structure and harbors many unique variants not present elsewhere. Thus, genetic variants identified from populations in other areas of the world may not have the same effects in Indians, and many variants specific to the Indian population could play significant roles in predisposing this population to certain diseases. Furthermore, due to consanguineous and endogamous marriages, the ancestral Indian population experienced founder events that are more extreme than Ashkenazi Jews and Finnish populations, which led to many genetically diverse subgroups.[4] As a result, the frequency of deleterious or protective variants may increase substantially, leading to increased power for gene mapping. All of these highlight the critical need and potential for new discovery of variants (otherwise rare outside India) from genetic studies in this population. Despite the need and opportunity, India has been largely underrepresented in genetic studies, making up less than 2% of human study participants in the catalog of genome-wide association studies (GWAS catalog).[5, 6] This lack of representation hinders the discovery of novel disease variants and the development of more tailored medical interventions or treatments towards the Indian population.

To facilitate genetic research in India, a crucial first step is to create an Indian-representative reference panel that can serve as a resource for future studies in India as well as methods development. This resource may serve as a reference panel for linkage disequilibrium (LD) and genotype imputation for the Indian population. LD refers to the non-random co-occurrence of alleles at different genomic loci.[7] A robust and representative estimate of genome-wide LD statistics for a given population is critical for a wide range of genetic analyses,[8-12] such as statistical fine-mapping and polygenic risk score (PRS) construction. Genotype imputation, on the other hand, estimates unobserved genotypes in samples with sparse microarray data and has been widely used in GWAS to boost power, fine-map candidate causal variants, and facilitate

meta-analysis across studies.[13] Extensive efforts have been made by several initiatives such as

the 1000 Genomes Project (1000G), Trans-Omics for Precision Medicine (TOPMed), and

Genome Asia Pilot (GAsP) project to establish reference panels.[14-17] However, these reference

panels have included a limited number of Indian samples and/or primarily focused on expatriate

communities outside of India. For example, 1000G includes 489 unrelated individuals from

South Asia, which includes only 103 Gujarati Indians in Houston, Texas and 102 Indian Telugu

individuals in the United Kingdom. Likewise, the only South Asian representation in TOPMed

comes from the Pakistani Risk of Myocardial Infarction Study (PROMIS) with a sample size of

644. While GAsP consists of a broad range of populations across Asia with a total sample size of

1,739, only 598 individuals are from India. This limited sample size and representation has

restricted the utility of these resources to serve as reference panels for the Indian population.

In the present study, we have constructed the largest and most nationally representative LD and

genotype imputation reference panel in India to date, using high-coverage (30x) whole genome

sequencing (WGS) data from 2,680 Indian participants from the Longitudinal Aging Study in

India-Harmonized Diagnostic Assessment of Dementia (LASI-DAD). For the LD reference

panel, we characterized LD patterns in LASI-DAD by identifying LD blocks and evaluating

regional differences in LD between LASI-DAD and four super-populations from 1000G. Given

the genetic heterogeneity in this population, we further characterized and compared LD patterns

by sub-populations within LASI-DAD and evaluated how such differences manifest in PRS

performance across sub-populations. We further assessed the utility of this LD reference panel in

facilitating PRS construction in South Asians. The LASI-DAD LD reference panel covers a

substantially higher number of genetic variants in India than previous panels and facilitates

various statistical analysis that rely on precise LD estimates. To evaluate LASI-DAD as an

imputation reference panel, we performed imputation using phased LASI-DAD WGS data or

other widely-used reference panels, including TOPMed and GAsP, and compared the imputation

accuracy across the spectrum of minor allele frequencies. As an imputation reference panel,

LASI-DAD enables the imputation of many Indian-specific variants that are not present in the

other panels with improved imputation accuracy.

Material and methods

The Longitudinal Aging Study in India - Harmonized Diagnostic Assessment of Dementia (LASI-DAD)

LASI is a nationally representative survey of more than 70,000 adults aged 45 years and older in India that aims to understand the health, economic, and social well-being of the older adults in India. LASI-DAD is an ancillary study of LASI that subsampled adults who were at least 60 years of age from 18 states and union territories across India to specifically focus on late-life cognition and dementia. Participants were selected based on a two-stage stratified sampling approach that oversampled individuals with a high risk of cognitive impairment.[18] Briefly, participants in the larger LASI cohort were first classified into those at high risk or low risk of cognitive impairment based on cognitive testing or on the proxy report for those who did not complete the cognitive tests. Finally, an approximately equal number of participants were randomly sampled from the high risk and low risk strata with a target sample size for each state/territory proportional to the population size.

A total of 2,762 LASI-DAD participants who consented to blood sample collection were carried forward for whole-genome sequencing (WGS) at MedGenome, Inc. (Bangalore, India) at an average read depth of 30x. Genotype calling and quality control (QC) of the raw WGS data were conducted at the Genome Center for Alzheimer's Disease (GCAD) at the University of Pennsylvania. Specifically, samples were excluded due to low sequencing coverage, sample contamination, sex mismatch, discordance with previous genotyping array data,[19] unexpected duplications, or serving as technical controls. A total of 2,680 samples were retained in the analysis after sample-level QC. The number of samples by state/territory is reported in Table S1, with states/territories grouped by geographic regions. For genotype-level QC, each genotype was examined and set to missing if its read depth (DP) was less than 10 or genotype quality (GQ) score was less than 20. For variant-level QC, a variant was excluded if it was monomorphic, fell in the 99.8% Variant Quality Score Recalibration (VQSR) Tranche or higher, had a call rate less than or equal to 80%, or had an average depth greater than 500 reads. We further excluded variants in low-complexity regions as identified by mdust.[20] After variant-level QC, a total of 71,109,961 biallelic variants that included 66,204,161 autosomal single nucleotide polymorphisms (SNPs) and 4,905,800 indels were retained in the analysis. We phased the

genotype data with Eagle2.4[21, 22] without a reference panel following the user manual of

Eagle2.4 (https://alkesgroup.broadinstitute.org/Eagle/#x1-130004). Finally, we estimated genetic

principal components (PCs) using PC-AiR in both LASI-DAD and the combined samples of

LASI-DAD and 1000G as described previously.[23, 24]

Characterizing and comparing linkage disequilibrium patterns between LASI-DAD and

1000 Genomes Project super-populations

We characterized LD patterns by identifying LD blocks and evaluating the variation in LD

(varLD) scores. For both LD block identification and varLD score calculation, we focused on a

total of 2,607 unrelated samples based on a kinship coefficient threshold of $2^{-3.5}$ (or 0.088)

obtained from PC-Relate.[25] LD blocks represent segments of the genome where genetic variants

are often inherited together. The identification of LD blocks can be useful in many ways for

genetic studies. For example, some polygenic risk score methods estimate SNP effects one block

at a time to achieve a good balance between modeling accuracy and computational efficiency.[26-

28] As another example, local genetic correlation analysis quantifies the genetic similarity of

complex traits within individual LD blocks.[29-31]

We applied LDetect[32] to identify approximately independent LD blocks in LASI-DAD, and

compared with those identified in four super-populations from 1000G: African (AFR), East

Asian (EAS), European (EUR), and South Asian (SAS). LDetect first partitions each

chromosome into smaller chunks to facilitate parallel processing. For each chunk, it computes a

covariance matrix between all pairs of SNPs using the shrinkage estimator from Wen and

Stephens.[33] Notably, both steps require the input of a recombination map to refine the boundary

of chunks and to adjust the level of shrinkage. Following MacDonald et al.,[34] we used the

recombination map inferred for Gujarati Indians in Houston (n = 113) in the 1000G phase 3

data.[35] Recombination rates were linearly interpolated to all SNPs in the LASI-DAD data.

Afterward, LDetect converts the covariance matrix into a vector with each element representing

the sum of elements along a corresponding antidiagonal of the matrix. Boundaries of LD blocks

appear as "dips" in the vector and can be identified by first smoothing the vector and then

searching for local minima. In the analysis, we focused on SNPs with MAF $\geq$ 1% and set the

average block size to 7,000 SNPs following MacDonald et al.[34] To compare LASI-DAD and the

1000G super-populations, we directly obtained LD blocks from MacDonald et al.[34] Notably, the

American (AMR) super-population was not included in MacDonald et al.,[34] likely due to its

extensive genetic admixture.

While LDetect is effective in capturing large LD blocks, it is not well-suited for identifying finer

LD structures within each block, which can provide additional insights into local LD variation.

To further identify LD blocks at a finer scale, we applied Big-LD.[36] Big-LD proceeds by first

identifying LD clusters through a modified clique-based clustering (CLQ) algorithm. Each

cluster consists of SNPs that are in strong LD with each other yet are not necessarily physically

consecutive. Big-LD then expands each LD cluster to form a consecutive genomic segment that

includes all SNPs inside the region. Finally, by merging mutually overlapping segments based on

interval graph modelling, Big-LD effectively partitions the genome into a set of finer LD blocks.

Big-LD was implemented in the gpart R package (version 1.17.0). We set the |r| threshold to be

its default value of 0.5 in the CLQ algorithm, where |r| represents the absolute correlation

coefficient between the dosage values of a SNP pair. For the comparison of LD blocks at the fine

scale, we applied Big-LD to the same four super-populations in 1000G. Specifically, we obtained

the genotype data for 3,202 samples from the 1000G 30x dataset on GRCh38

(https://www.internationalgenome.org/data-portal/data-collection/30x-grch38) and restricted the

samples to 2,504 unrelated individuals as reported by 1000G for our analysis.[15] Following

MacDonald et al.,[34] we excluded the African Caribbean in Barbados (ACB; n = 96) and African

Ancestry in Southwest US (ASW; n = 61) from the AFR population due to genetic admixture, as

well as the Finnish in Finland (FIN; n = 99) from the EUR population due to genetic isolation.

The total number of samples included were 504 for AFR, 504 for EAS, 404 for EUR, and 489

for SAS.

In addition to LD block identification, we compared LD patterns by evaluating the varLD score

between LASI-DAD and the four 1000G super-populations.[37] varLD examines regional patterns

of LD between two populations by comparing the signed r^2 values across all pairs of SNPs in a

genomic region with a MAF $\geq$ 1% that are not strand-ambiguous in both populations, where r^2

represents the squared correlation coefficient between the dosage values of a SNP pair.

Specifically, for a particular genomic region, we first calculated the LD matrix for the two

populations, defined as a SNP-by-SNP correlation matrix with each matrix element representing

the signed r^2 value between a SNP pair. We then obtained the eigenvalues of the LD matrix from

each population and calculated the varLD score as the sum of the absolute differences between

the corresponding eigenvalues derived from the two populations. We employed a sliding-

window approach and calculated varLD scores within consecutive windows of 50 SNPs. Each

consecutive window was obtained by shifting the current window by one SNP in the direction of

the forward strand.[25]

Characterizing LD patterns by LASI-DAD sub-population

In addition to comparisons with 1000G super-populations, we evaluated LD patterns across three

LASI-DAD sub-populations, defined according to the population structure in India. The

population structure in India is correlated with geography, with majority of the individuals lying

along a North/South cline with varying proportions of Ancestral North Indian (ANI) and

Ancestral South Indian (ASI) ancestries. ANI is genetically related to West Eurasians while ASI

is related (distantly) to indigenous Andaman Islanders.[2] In contrast, individuals in the east of

India are considered out-of-cline due to admixture with additional ancestral populations such as

East Asians. Following Kerdoncuff et al.,[38] we clustered individuals into those that lie along the

North/South cline and those that are out of the cline. For individuals that lie along the

North/South cline, we estimated the proportion of ANI ancestry (%ANI) using the f_4-statistic

implemented in the ADMIXTOOLS R package (version 1.3.0).[39] The f_4-statistic measures the

excess of West Eurasian ancestry in a test individual relative to the Onge population based on an

Indian population model described by Moorjani et al.[3] Finally, we grouped individuals from

LASI-DAD into three sub-populations based on their respective cline and %ANI values to

facilitate Indian sub-population analysis. The three sub-populations included 945 individuals in

the high %ANI group ($0.540 \leq$ %ANI < 0.824), 957 in the low %ANI group (%ANI ≤ 0.540),

and 778 that were out of the North/South cline. We conducted this analysis on the set of

unrelated samples, including 935, 941, and 731 individuals in the respective groups. For

simplicity, we focused on the evaluation of varLD scores in the sub-population analysis.

Transferability of polygenic risk scores

A polygenic risk score (PRS) captures an individual's genetic predisposition to a complex trait or disease and is often calculated as the sum of SNP genotypes weighted by their effect sizes estimated from GWAS. However, most GWASs have been conducted in individuals of European ancestry[40] and PRS derived from these GWASs may be substantially less predictive in other genetic ancestries.[11] We evaluated the transferability of the PRS derived from GWASs of primarily European ancestry to LASI-DAD individuals. Specifically, we obtained PRS weights for height and body mass index (BMI) from the Genetic Investigation of Anthropometric Traits (GIANT) consortium. For height, we downloaded PRS weights that were generated by applying SBayesC to a cross-ancestry GWAS meta-analysis[41] (https://portals.broadinstitute.org/collaboration/giant/ index.php/GIANT_consortium_data_files). The meta-analysis included 5,380,080 individuals across 281 studies from the GIANT consortium and 23andMe, with 75.8% of the individuals being primarily of European ancestry and 1.4% being primarily of South Asian ancestry. For BMI, we estimated PRS weights by applying PRS-CS[26] to a GWAS meta-analysis of GIANT consortium studies and UK Biobank.[42] The meta-analysis included ~700,000 individuals of primarily European ancestry. After obtaining the PRS weights, we evaluated the performance of PRS in the three LASI-DAD sub-populations (high %ANI, low %ANI, and out-of-cline) using incremental R^2. Following Yengo et al.,[41] we calculated the incremental R^2 for height by first centering and standardizing height to have a mean of zero and a variance of one within each sex. The incremental R^2 was then calculated as the difference in the coefficient of determination between two linear regression models: one that includes age and the top 20 genotype PCs as covariates, and the other that additionally includes the height PRS. For BMI, we followed a similar procedure to calculate the incremental R^2.

Providing LASI-DAD as a reference panel for linkage disequilibrium

A robust and representative estimate of genome-wide LD statistics for a given population plays an essential role for a wide range of genetic analyses.[8-12] As the largest and most nationally representative WGS study of the Indian population, LASI-DAD has the potential to provide more accurate estimates of LD with higher coverage of genetic variants, serving as a valuable resource for LD in this population. We evaluated the utility of LASI-DAD WGS data as an LD

reference panel to facilitate both LD lookup and various statistical analyses that rely on precise

LD estimates. We constructed an LD reference panel using 2,607 unrelated individuals from the

LASI-DAD WGS data and provided it as a public resource to facilitate future research in the

community. Specifically, for all pairs of genetic variants within 1 million base pairs (Mb) of each

other, we computed the squared Pearson correlation coefficient (r^2) between the phased

haplotypes, D-prime (D'), and the direction of the Pearson correlation coefficient. We reported

LD statistics for pairs of variants that have an r^2 statistic $\geq$ 0.2. Of note, we did not use a

minimum minor allele count to exclude any variants. We compared these LD statistics with those

obtained from the South Asian populations in 1000G and TOP-LD.[16] TOP-LD estimated LD

statistics for 239 selected South Asian individuals from TOPMed WGS data.[14] We obtained the

LD statistics for TOP-LD directly from its website (http://topld.genetics.unc.edu/about.php). For

1000G, we computed the LD statistics following procedures similar to those used for LASI-DAD

and TOP-LD. Of note, an LD panel for the Genome Asia Pilot (GAsP) is not publically available,

thus was not included in this comparison.

In addition to serving as an LD lookup panel, LASI-DAD also facilitates various statistical

analyses that rely on precise LD estimates. To simplify data sharing while ensuring privacy

protection, many statistical methods can directly make use of GWAS summary statistics as

input.[43-46] These summary statistics typically include marginal z-scores for each SNP and a SNP-

SNP correlation matrix. The SNP-SNP correlation matrix is commonly referred to as the LD

matrix and can be estimated from a reference panel consisting of individuals with the same

genetic ancestry as the GWAS. In principle, the closer the LD estimates from the reference panel

match those in the GWAS population, the better the results are likely to be in the statistical

analysis. This often requires the reference panel to be sufficiently large and representative of the

GWAS population. To demonstrate the utility of LASI-DAD as an LD reference panel, we

conducted a PRS analysis where we compared the predictive performance of PRS constructed

with either the LASI-DAD or 1000G SAS LD reference panel. We estimated PRS weights using

PRS-CS, which takes GWAS summary statistics and an external LD reference panel as inputs.

We obtained summary statistics for height and BMI from four studies of South Asian ancestry,

including a GWAS of height[41] and an exome-wide association study (ExWAS) of BMI[47] from

the GIANT consortium, and GWASs of both height and BMI from the Genes & Health.[48] The

GIANT data included 77,890 individuals of primarily South Asian ancestry for height and
29,398 for BMI. The Genes & Health data included 36,317 British Pakistani and British
Bangladeshi individuals for height and 34,408 for BMI. The summary statistics included
marginal association results for 1,339,059 and 246,158 SNPs for height and BMI from the
GIANT consortium, and 24,241,756 and 23,922,099 biallelic variants for height and BMI from
Genes & Health. We constructed the LASI-DAD LD reference panel following a procedure
similar to that of Ge et al.[26] Specifically, we focused on the 2,607 unrelated samples and filtered
out SNPs that have MAF < 1% or are strand-ambiguous. We restricted the SNPs to those in the
HapMap 3 panel to achieve a good balance between computational efficiency and prediction
accuracy. For each LD block as identified by LDetect, we computed a SNP-SNP correlation
matrix, with each element in the matrix representing the Pearson correlation coefficient between
the dosage values of a SNP pair. For each study and LD reference panel, we applied PRS-CS to
estimate PRS weights. We evaluated the performance of PRS in the full 2,680 LASI-DAD
samples using incremental R^2, calculated following the same procedure as described above.

Providing LASI-DAD as a reference panel for genotype imputation
Given the unique population structure of India, LASI-DAD has the potential to serve as a
valuable resource for genotype imputation to enhance the analysis of array-genotyped samples,
especially those collected from the Indian populations. We evaluated the utility of LASI-DAD
WGS data as an imputation reference panel and compared it with the GAsP and TOPMed
imputation panels. Specifically, we randomly selected 180 samples, 10 from each Indian state or
union territory, from the LASI-DAD dataset and selected 530,845 genetic markers that were
present on the Illumina Infinium Global Screening Array-24 v2.0 BeadChip. These 180 samples
served as a test dataset to evaluate the imputation accuracy. Following Yu et al.,[49] we evaluated
the imputation accuracy by calculating the aggregated squared correlation coefficient (r^2),
computed using aggRSquare (https://github.com/yukt/aggRSquare) which groups variants within
a specified MAF range, stacks their imputed dosages and genotype calls from the sequencing
data into two separate vectors, and then calculates the aggregated r^2 as the squared correlation
coefficient between the two vectors. We imputed unobserved genotypes with Minimac4 (v1.0.2)
using either LASI-DAD (n = 2,500 after excluding the 180 samples from the test dataset), GAsP
(n = 1,654), or TOPMed (n = 97,256) as the reference panel. For GAsP and TOPMed, we carried

out the imputation directly using the Michigan Imputation Server and the TOPMed Imputation

Sever,[50] respectively. For LASI-DAD, we followed the same analysis pipeline described on the

TOPMed Imputation Server for data quality control and imputation. Briefly, we split each

chromosome into chunks of 10 Mb, filtered out chunks with fewer than 3 variants, and excluded

monomorphic variants. We then conducted genotype imputation using Minimac4 with the same

parameter settings as those used by the imputation sever. Besides the imputation with a single

reference panel, we also performed meta-imputation by combining imputation results from

LASI-DAD with those from GAsP, TOPMed, or both using MetaMinimac2.[49] We evaluated the

imputation accuracy across different ranges of MAF, where the MAFs were calculated among all

2,680 LASI-DAD samples. To facilitate future research in the community, the LASI-DAD

reference panel is now available on the Michigan Imputation sever.

Results

Characterizing LD patterns in the Indian population from LASI-DAD

We characterized LD patterns in LASI-DAD by identifying LD blocks and evaluating the variation in LD (varLD) scores.[37] In total, we identified 1,262 blocks in LASI-DAD, which is comparable to the 1,291 blocks identified in the SAS population from 1000G (Figure 1A). For the other 1000G populations, the number of identified blocks was 1,605, 1,360, and 1,143 for AFR, EUR, and EAS, respectively (Figure 1A). As expected, we observed the largest number of LD blocks in the AFR population as Africans are the oldest population ancestrally and there has been more time for recombination to break down the LD structure.[51] On average, the length of LD blocks was 2.2 million bases (Mb) in LASI-DAD and 1.7 Mb, 2.1 Mb, 2.2 Mb, and 2.5 Mb in the AFR, EUR, SAS, and EAS populations from 1000G, respectively (Figure 1B).

While the LD blocks identified by LDetect have been widely used by various statistical methods, it is important to note that these blocks represent LD structures at a broad scale, with local LD structures still existing at a finer scale. Indeed, in an evaluation where we compared the average squared Pearson correlation (r^2) between all pairs of variants within an LD block to the average r^2 between all pairs of variants in an LD block and its adjacent two blocks, we found that although the average r^2 was substantially higher within each LD block than with its neighboring two blocks, it remained relatively low, with an average value of 0.015 versus 0.0010 in LASI-DAD (Figure 1C). To better understand the LD structure at a finer scale, we further applied Big-LD.[36] As expected, the number of LD blocks identified by Big-LD was much larger and ranged from 96,415 to 143,383 across the five populations (Table S2). For these LD blocks, we observed a considerably higher average r^2 both within and between blocks, with average values of 0.32 and 0.098, respectively, in LASI-DAD (Figure S1). Finally, we visualized an example LD block from LDetect (Figure S2). While this region corresponds to a single LD block at a broad scale, we observed extensive variation both in the local LD structure in this region (Figure S2A) and across different populations (Figure S2B).

In addition to LD block identification, we evaluated regional differences in LD between pairs of populations with varLD scores. As expected, we observed the smallest difference in LD between

365   LASI-DAD and the SAS population from 1000G, with an average varLD score of 1.67 (Figure
366   1D). This was followed by the EUR (6.21), EAS (8.34), and AFR (18.5) populations.
367

Indian population structure and transferability of PRS

In addition to comparisons with 1000G super-populations, we evaluated LD patterns across the
three LASI-DAD sub-populations (Figures 2A-B and Table S1). We found that the overall
differences in LD across LASI-DAD sub-populations were modest compared to those observed
between LASI-DAD and the 1000G EUR, EAS, and AFR populations, but comparable to those
between LASI-DAD and the 1000G SAS population (Figure 2C versus Figure 1D). The average
varLD scores were 2.10, 1.81, and 1.69 between the out-of-cline and high %ANI groups,
high %ANI and low %ANI groups, and the out-of-cline and low %ANI groups, respectively
(Figure 2C). This level of difference in LD is consistent with the relative genetic distances in the
principal component space, where larger genetic distance corresponds to greater difference in LD
(Figures 2B-C).
379

Despite overall modest differences in LD across LASI-DAD sub-populations, such differences
can still impact the performance of various statistical analyses, such as the PRS. We evaluated
the transferability of a height and body mass index (BMI) PRS derived from GWASs of
primarily European ancestry to LASI-DAD individuals. We found that the predictive
performance of PRS varied across LASI-DAD sub-populations (Figure 2D), with relative
performance consistent with their genetic relatedness to the European population (Figures 2A-B).
As expected, PRS performed the best in the high %ANI group due to its closer relatedness to the
European population, achieving a prediction accuracy of 17% for height and 6% for BMI, as
measured by incremental R^2. In contrast, the prediction accuracy was 14% and 3% in the
low %ANI group and 10% and 2% in the out-of-cline group for height and BMI, respectively.
Despite the variation across sub-populations, the overall prediction accuracy in LASI-DAD
remained substantially lower than that in the European population, where it reached as high as
44.7% for height[41] and 10.2% for BMI.[42] The limited transferability of PRS in LASI-DAD
hinders its clinical utility, especially for South and East Indian populations who are more
distantly related to the European population and benefit less from its large sample size. These

findings suggest that more efforts need to be made toward genetic research in this unique

population to reduce potential health disparities.[40]

Providing LASI-DAD WGS data as an LD reference panel

We evaluated the utility of LASI-DAD WGS data as an LD reference panel to facilitate both LD

lookup and various statistical analyses that rely on precise LD estimates. In the LD lookup panel,

LASI-DAD included a total of 69.5 million variants, representing a 170% (43.8 million) and 213%

(47.3 million) increase in variant coverage relative to the 1000G and TOP-LD panels,

respectively (Figure 3A). To be included in the panels, these variants have at least one tagged

variant with $r^2 \geq 0.2$. As expected, most variants in LASI-DAD that are not present in the other

two panels have a relatively low MAF (e.g., < 1%) (Figure 3A). Interestingly, we found that the

number of variants with a MAF $\geq$ 1% is smaller in the LASI-DAD panel (8.7 million) than in the

1000G (11.6 million) and TOP-LD (9.4 million) panels (Figure 3A). Presumably, this is because

LASI-DAD represents a highly heterogenous yet nationally representative population that

contains individuals with diverse histories of founder events.[38] In contrast, the South Asian

population in 1000G consists of individuals from only five sub-population groups, including 86

Bengali in Bangladesh; 103 Gujarati Indians in Houston, Texas; 102 Indian Telugu in the UK; 96

Punjabi in Lahore, Pakistan; and 102 Sri Lankan Tamil in the UK. Similarly, the only South

Asian representation in TOPMed and its derived LD panel TOP-LD (n = 239) comes from the

Pakistani Risk of Myocardial Infarction Study (PROMIS).[14, 52, 53] The lack of genetic diversity in

1000G and TOP-LD for the South Asian population likely contributed to the larger number of

variants with a relatively high MAF. The coverage of genetic variants remained the highest for

the LASI-DAD panel regardless of the cutoff in r^2 (Figure 3B). In addition, most variants in the

LD lookup panel have at least one tagged variant with $r^2 \geq 0.8$, with proportions of variants being

82%, 76%, and 89% in the LASI-DAD, 1000G, and TOP-LD panels, respectively (Figure 3B).

Notably, the number of variant pairs remained largely similar across different LD lookup panels

regardless of the cutoff in r^2 (Figure 3C). Again, this is likely due to the lack of genetic diversity

in 1000G and TOP-LD, resulting in each variant having more tagged variants in these two panels.

Finally, we evaluated the consistency of LD estimates across the three lookup panels by

examining the common set of variant pairs on chromosome 1. The Pearson correlation of r^2

statistics was 0.94 between LASI-DAD and 1000G, 0.92 between LASI-DAD and TOP-LD, and

0.91 between 1000G and TOP-LD, indicating an overall high consistency across the three panels.

To demonstrate the advantage of LASI-DAD as an LD reference in statistical analyses, we

conducted a PRS analysis where we compared the predictive performance of PRS constructed

with either the LASI-DAD or 1000G SAS LD reference panel. To obtain PRS weights, we

applied PRS-CS[26] to four studies of South Asian ancestries, including a GWAS of height[41] and

an ExWAS of BMI[47] from the GIANT consortium, and GWASs of both height and BMI from

Genes & Health.[48] We evaluated the prediction accuracy of PRS in LASI-DAD individuals. We

found that the predictive performance of PRS was higher with the LASI-DAD panel than with

the 1000G SAS panel for all four studies (Figure 4). Specifically, the LASI-DAD panel

improved the prediction accuracy by 19.5% for height and 35.1% for BMI from the GIANT

consortium, and by 2.2% for height and 2.1% for BMI from the Genes & Health. Notably, the

improvement in PRS performance when using LASI-DAD as the reference panel was smaller for

the Genes & Health data than for the GIANT data. This is likely because the LD patterns in

LASI-DAD match more closely to those in GIANT than Genes & Health, as Genes & Health

consists of British Pakistani and British Bangladeshi individuals, whereas a large proportion of

individuals in GIANT are of Indian origin. These results highlight the benefits of LASI-DAD in

serving as an LD reference panel.

Providing LASI-DAD WGS data as an imputation reference panel

We evaluated the utility of LASI-DAD WGS data as an imputation reference panel and

compared it with the GAsP and TOPMed imputation panels. Specifically, we found that the

LASI-DAD panel enabled the imputation of many Indian-specific variants that are not present in

the other panels, especially those with a low MAF (Figure 5A). For rare variants with a MAF ≤

1%, the LASI-DAD panel was able to impute a total of 61,843,011 variants, representing 182%

(21,948,408) and 940% (5,943,775) increases relative to the TOPMed and GAsP panels,

respectively. Similarly, for common variants with a MAF > 1%, the LASI-DAD panel was able

to impute 8,783,797 variants, representing 4% (363,996) and 20% (1,449,768) increases relative

to the TOPMed and GAsP panels. To ensure a fair comparison, we evaluated the imputation

accuracy using the common set of variants that could be imputed by all three panels (Figure 5A).

We found that the imputation with the LASI-DAD panel achieved substantially higher accuracy

than with the other two panels for variants across most MAF ranges, except for singletons in the

range of (0, 0.02]% (Figure 5B). For non-singletons, the LASI-DAD panel improved the

imputation accuracy by 3% to 101% (mean = 38%) compared to the TOPMed panel and by 3%

to 73% (mean = 27%) compared to the GAsP panel across different MAF ranges. The imputation

performance is consistent with the sample size of South Asian individuals in each panel, with

1,654 samples in the GAsP panel and 644 in the TOPMed panel, both considerably smaller than

that of the LASI-DAD panel. As expected, none of the panels can accurately impute singletons.

By combining imputation results from different reference panels, meta-imputation further

improved the imputation accuracy, especially for variants with a low MAF (Figure 5B). Notably,

meta-imputation using all three panels achieved the highest imputation accuracy across all MAF

ranges. For variants with a relatively small MAF ($\leq$ 0.1%), meta-imputation increased the

accuracy by 14% to 107% (mean = 64%) compared to the best-performing reference panel. On

the other hand, for variants with a relatively large MAF (> 0.1%), incorporating additional

reference panels beyond LASI-DAD offered only limited benefits, with imputation accuracy

improving by 0.2% to 2% (mean = 0.8%) compared to the LASI-DAD panel. These results

highlight the advantages of meta-imputation, in complementing the LASI-DAD panel, for

imputing relatively rare variants in the Indian population.

Finally, we found that the imputation accuracy varied for samples from different states or

territories of India, with the LASI-DAD panel exhibiting the smallest variation (Figure S3). This

variation in imputation accuracy can be explained to some extent by the sample composition of

the reference panels. For example, the TOPMed panel, which consists nearly 50% European

samples, exhibits a strong correlation between the imputation accuracy and the average %ANI

for each state (R^2 = 0.77; Figure S4). This is consistent with the fact that samples with

higher %ANI have closer relatedness to West Eurasians. In contrast, the correlation is much

weaker with the LASI-DAD (R^2 = 0.42) or GAsP (R^2 = 0.15) panel. These results highlight the

importance of having a large and nationally representative dataset for the Indian population to

achieve the best imputation performance due to its complex population structure.

Discussion

As the largest and most nationally representative WGS study of the Indian population, LASI-DAD provides a unique opportunity to serve as a valuable resource for both LD and genotype imputation in this population. In the present study, we characterized the LD patterns in LASI-DAD by identifying LD blocks and evaluating regional differences in LD between the Indian population from LASI-DAD and four super-populations from the 1000G. Besides super-populations, we explored the population structure of India by evaluating LD patterns across Indian sub-populations. We evaluated the utility of LASI-DAD WGS data as an LD reference panel to facilitate both LD lookup and various statistical analyses that rely on precise LD estimates. Finally, we evaluated the utility of LASI-DAD as an imputation reference panel. As an LD reference panel, LASI-DAD provides a more comprehensive representation of genetic variation/structure in India and facilitates downstream LD-dependent analyses. As an imputation panel, it improves imputation accuracy compared to existing panels and demonstrates more robust performance across different Indian states or territories.

The comparison of LD patterns between LASI-DAD and four super-populations from 1000G confirms that the African population has the largest number of LD blocks and smallest overall block sizes. In comparison, the European, East Asian and South Asian populations, which all migrated from Africa, have a comparable number of blocks with similar block sizes. As expected, LASI-DAD has a similar LD structure as the 1000G South Asian population, in terms of both the number and size of LD blocks, according to either the broad-scale LD blocks identified by LDetect or the fine-scale LD blocks identified by BigLD. However, our analysis still reveals some regions with considerable LD differences between LASI-DAD and 1000G SAS. This heterogeneity is further highlighted by global genetic principal components, showing that LASI-DAD significantly broadens the genetic boundaries typically seen in the 1000G South Asian panel (Figure S5). Moreover, due to a significant increase in sample size, LASI-DAD interrogates significantly more (170%) variants, likely specific to the Indian/South Asian population, that are not observed in the 1000G SAS panel. Given the improved representation and coverage, it is not surprising that the LASI-DAD panel, in comparison to 1000G SAS panel, improves polygenic score construction and prediction in South Asians. We expect similar improvement in other LD-based analyses that target Indian and South Asian populations.

Unfortunately, the sample sizes of South Asian GWASs are limited; thus, the overall prediction

accuracy of the corresponding PRS is much lower than that based on the European GWASs

(Figure 2D versus Figure 4). The next important step is to expand GWAS efforts in South Asian

populations, for which a good imputation reference panel is needed to generate high quality

genotype data. LASI-DAD can well serve that purpose as it improves imputation accuracy over

existing popular reference panels for South Asians, including TOPMed and GAsP. Importantly,

it has more robust performance across all Indian states or territories, likely due to its unbiased

representation of the Indian population. Our analysis further highlighted the potential advantage

of leveraging multiple reference panels via meta-imputation to further improve imputation

accuracy. All of these demonstrate that LASI-DAD is a valuable genetic reference panel for

Indian and South Asian populations.

The Indian population is diverse and shaped by multiple ancestral influences. Specifically, the

majority of Indians are part of a North/South cline and are admixed with varying degrees of ANI

(related to European, Central Asia and Middle Eastern populations) and ASI (related to

endogenous ancestry). Meanwhile, a small proportion of Indians who are in the East have an

additional ancestral influence from East Asians. Our analysis by LASI-DAD sub-population

confirms that this population structure leads to LD differences among sub-populations, with the

largest difference observed between North Indians (high %ANI group) and East Indians (out-of-

cline group), while South Indians (low %ANI group) are positioned in between. Since North

Indians are most closely related to Europeans, the predictive performance of European-based

PRSs shows a similar gradient across these sub-populations, with highest performance in North

Indians, followed by South Indians, and then East Indians. This is consistent with many studies,

showing that the transferability of PRS decreases as the genetic distance between target

population and discovery population increases.[54] The heterogeneity could also affect imputation

accuracy across sub-populations when the reference panel is biased. This heterogeneity and

potential consequence on health inequity underscore the importance of including diverse groups

and having population-representative Indian samples, as exemplified by LASI-DAD.

While LASI-DAD has proven to be a valuable resource for serving as a reference panel, several

opportunities are possible to further improve its utility. First, while LASI-DAD represents the

largest and most nationally representative WGS study, it remains insufficient to fully capture the
extensive genetic diversity of the Indian population, which includes over 4,500 anthropologically
defined groups. Substantial efforts to collect more samples are still needed to achieve a more
comprehensive representation. Second, most existing genetic studies analyze South Asian
populations collectively as a single group. However, South Asia is a region that spans eight
countries, making up about 25% of the world's population and having a rich genetic diversity.[52]
As a result, LD patterns observed in one South Asian population may not accurately reflect those
in another. This partly explains why the improvement in PRS performance when using LASI-
DAD as the reference panel was smaller for the Genes & Health data than for the GIANT data,
as the Genes & Health data consists of British Pakistani and British Bangladeshi, whereas a large
proportion of individuals in the GIANT data are of Indian origin. To achieve the best results in
statistical analyses with GWAS summary statistics, identifying the most appropriate reference
panel with LD patterns that best match the GWAS population is critical. In line with this, an
adaptive approach that combines multiple reference panels, akin to the meta-imputation approach,
may provide an alternative way for improving the LD reference panel. However, this would
require the development of new statistical frameworks or analytical procedures.

In summary, we have presented LASI-DAD, the largest and most nationally representative WGS
data from India, as a reference panel for LD and genotype imputation. This valuable resource
fills a critical gap in the field of genetics and could greatly facilitate various genetic analyses in
Indians and South Asians. Meanwhile, the genetic diversity demonstrated in this study also
underscores a critical need to expand data collection and enhance representativeness in this
population.

Acknowledgements

We thank the LASI-DAD participants for being part of the research project. We thank Dr. Priya Moorjani at the University of California, Berkeley for her guidance on defining the LASI-DAD sub-populations. This study was supported by the National Institute on Aging (R01 AG051125, U01 AG064948, and U54-AG-052427). The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions

W.Z., J.S., and S.K. conceived the idea. J.L. and S.K. provided funding support. S.D., A.B.D., and J.L. led data collection in LASI-DAD. Y.Y.L, G.D.S and L.-S.W. processed and cleaned the whole genome sequencing data. Z.L. and W.Z. designed the experiments. Z.L. conducted the data analyses. Z.L., W.Z., and J.S. wrote the manuscript with input from X.Z., Y.Y.L, and G.D.S. All authors read and approved the final manuscript.

Declaration of interests

The authors declare no competing interests.

Institutional review board statement

Investigations were conducted in accordance with the principles outlined in the Declaration of Helsinki (1975, revised in 2013). The study was approved by Institutional Review Boards at the University of Southern California (UP-15-00684) and the University of Michigan (HUM00166956).

Informed consent statement

Written informed consent was obtained from all participants.

Data availability statement

Whole genome sequencing data for the Diagnostic Assessment of Dementia for the Longitudinal Aging Study of India (LASI-DAD) is available from the National Institute on Aging Genetics of Alzheimer's Disease Data Storage Site (NIAGADS), accession number: NG00067 – ADSP Umbrella. Phenotype data is available at the Gateway to Global Aging website,

604 https://g2aging.org/. The LASI-DAD LD reference panel is available at
605 https://github.com/zhengli09/LASI_DAD-LD-Reference-Panel. We are currently working with
606 the Michigan Imputation Server (https://imputationserver.sph.umich.edu) to host the LASI-DAD
607 imputation panel; however, it is not yet available. Researchers wishing to access the imputation
608 panel should contact the corresponding authors.
609

Figures

Figure 1. **Linkage disequilibrium (LD) patterns in LASI-DAD compared to four super-populations from the 1000 Genomes Project (1000G)**. The four super-populations include African (AFR), European (EUR), South Asian (SAS), and East Asian (EAS). (**A**) Number of LD blocks identified by LDetect. (**B**) Distribution of LD block lengths in millions of base pairs (Mb), with LD blocks grouped into different length ranges for clear visualization. (**C**) Boxplots showing the distribution of average LD on chromosome 1, calculated either between all pairs of SNPs within each LD block or between SNPs in each LD block and its adjacent two blocks. LD is evaluated as the squared correlation coefficient (r^2) of genotypes between a SNP pair. (**D**) Distribution of variation in LD (varLD) scores evaluated between LASI-DAD and each of the four super-populations from 1000G.

Figure 2. **Linkage disequilibrium patterns in Indian sub-populations from LASI-DAD**.
Individuals from LASI-DAD were grouped into three sub-populations based on their proportion
of Ancestral North Indian ancestry (%ANI) and presence on the North/South cline of India to
facilitate Indian sub-population analysis. The three sub-populations included the high %ANI
group ($0.540 \le \%\text{ANI} < 0.824$; n = 935), low %ANI group ($\%\text{ANI} \le 0.540$; n = 941), and the out
of the North/South cline group (n = 731). (**A**) Principal component analysis (PCA) plot
illustrating the genetic relationship among Indian, European, and East Asian populations. PCA
was conducted using Indian samples from LASI-DAD along with European, South Asian, and
East Asian samples from 1000G. The proportion of variance explained by each PC is indicated in
the parentheses. LASI-DAD individuals are colored by their geographic region of birth in India
to highlight the relationship between population structure and geography in India. (**B**) The same
PCA plot with LASI-DAD individuals colored by their sub-population groups. (**C**) Distribution
of variation in LD (varLD) scores evaluated between LASI-DAD sub-populations. (**D**) Predictive
performance of polygenic risk scores (PRS) derived from GWAS of primarily European ancestry
to the Indian sub-populations in LASI-DAD. Evaluated traits include height and body mass
index (BMI), with GWAS summary statistics obtained from the GIANT consortium.

640
Figure 3. Number of autosomal variants included in different linkage disequilibrium (LD)
reference panels. Three LD reference panels were compared, including Indian samples from
LASI-DAD (N=2,607), South Asian samples from 1000G (N=489), and South Asian samples
from TOPMed (N=239). (**A**) Number of variants stratified by the range of minor allele frequency
(MAF). (**B**) Number of variants with at least one other variant in LD under different r^2 cutoffs,
where r^2 represents the squared Pearson correlation coefficient between the phased haplotypes of
a variant pair. (**C**) Number of variant pairs stratified by different r^2 cutoffs.
648

649
Figure 4. **PRS analyses demonstrating the utility of LASI-DAD as an LD reference panel**.
The predictive performance of PRS constructed with the LASI-DAD and 1000G SAS LD
reference panels were compared. Summary statistics for height and body mass index (BMI) were
obtained from four studies of South Asian ancestry, including a GWAS of height and an ExWAS
of BMI from the GIANT consortium, and GWASs of both height and BMI from Genes & Health.
PRS weights were estimated by PRS-CS.

Figure 5. **Evaluating LASI-DAD as a reference panel for genotype imputation**. Compared reference panels include LASI-DAD, TOPMed, and GAsP. (**A**) Bar plots showing the number of genetic variants in LASI-DAD that can be imputed by each reference panel and by all three panels. (**B**) Imputation accuracy was evaluated for each reference panel, including meta-imputation to integrate the imputation results from either LASI-DAD and TOPMed, LASI-DAD and GAsP, or all three panels, across different minor allele frequency (MAF) ranges. The MAF was calculated within the full 2,680 LASI-DAD samples. To ensure a fair comparison, the imputation accuracy was evaluated using the common set of variants that can be imputed by all three panels. Aggregated r^2 was used as the evaluation metric, calculated by grouping variants within a specified MAF range, stacking their imputed dosages and genotype calls from the sequencing data into two separate vectors, and then computing the squared correlation coefficient between the two vectors.

References

1. Mastana, S.S. (2014). Unity in diversity: an overview of the genomic anthropology of India. Ann Hum Biol 41, 287-299.

2. Reich, D., Thangaraj, K., Patterson, N., Price, A.L., and Singh, L. (2009). Reconstructing Indian population history. Nature 461, 489-494.

3. Moorjani, P., Thangaraj, K., Patterson, N., Lipson, M., Loh, P.R., Govindaraj, P., Berger, B., Reich, D., and Singh, L. (2013). Genetic evidence for recent population mixture in India. Am J Hum Genet 93, 422-438.

4. Nakatsuka, N., Moorjani, P., Rai, N., Sarkar, B., Tandon, A., Patterson, N., Bhavani, G.S., Girisha, K.M., Mustak, M.S., Srinivasan, S., et al. (2017). The promise of discovering population-specific disease-associated genes in South Asia. Nat Genet 49, 1403-1407.

5. Morales, J., Welter, D., Bowler, E.H., Cerezo, M., Harris, L.W., McMahon, A.C., Hall, P., Junkins, H.A., Milano, A., Hastings, E., et al. (2018). A standardized framework for representation of ancestry data in genomics studies, with application to the NHGRI-EBI GWAS Catalog. Genome Biol 19, 21.

6. Sirugo, G., Williams, S.M., and Tishkoff, S.A. (2019). The Missing Diversity in Human Genetic Studies. Cell 177, 26-31.

7. Wall, J.D., and Pritchard, J.K. (2003). Haplotype blocks and linkage disequilibrium in the human genome. Nat Rev Genet 4, 587-597.

8. Uffelmann, E., Huang, Q.Q., Munung, N.S., de Vries, J., Okada, Y., Martin, A.R., Martin, H.C., Lappalainen, T., and Posthuma, D. (2021). Genome-wide association studies. Nat Rev Method Prime 1.

9. Schaid, D.J., Chen, W., and Larson, N.B. (2018). From genome-wide associations to candidate causal variants by statistical fine-mapping. Nat Rev Genet 19, 491-504.

10. Li, Z., Zhao, W., Shang, L., Mosley, T.H., Kardia, S.L.R., Smith, J.A., and Zhou, X. (2022). METRO: Multi-ancestry transcriptome-wide association studies for powerful gene-trait association detection. Am J Hum Genet 109, 783-801.

11. Kachuri, L., Chatterjee, N., Hirbo, J., Schaid, D.J., Martin, I., Kullo, I.J., Kenny, E.E., Pasaniuc, B., Polygenic Risk Methods in Diverse Populations Consortium Methods Working, G., Witte, J.S., et al. (2024). Principles and methods for transferring polygenic risk scores across global populations. Nat Rev Genet 25, 8-25.

12. Gao, B., and Zhou, X. (2024). MESuSiE enables scalable and powerful multi-ancestry fine-mapping of causal variants in genome-wide association studies. Nat Genet 56, 170-179.

13. Marchini, J., and Howie, B. (2010). Genotype imputation for genome-wide association studies. Nat Rev Genet 11, 499-511.

14. Taliun, D., Harris, D.N., Kessler, M.D., Carlson, J., Szpiech, Z.A., Torres, R., Taliun, S.A.G., Corvelo, A., Gogarten, S.M., Kang, H.M., et al. (2021). Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. Nature 590, 290-299.

15. Byrska-Bishop, M., Evani, U.S., Zhao, X., Basile, A.O., Abel, H.J., Regier, A.A., Corvelo, A., Clarke, W.E., Musunuri, R., Nagulapalli, K., et al. (2022). High-coverage whole-genome sequencing of the expanded 1000 Genomes Project cohort including 602 trios. Cell 185, 3426-3440 e3419.

16. Huang, L., Rosen, J.D., Sun, Q., Chen, J., Wheeler, M.M., Zhou, Y., Min, Y.I., Kooperberg, C., Conomos, M.P., Stilp, A.M., et al. (2022). TOP-LD: A tool to explore linkage disequilibrium with TOPMed whole-genome sequence data. Am J Hum Genet 109, 1175-1181.

17. GenomeAsia, K.C. (2019). The GenomeAsia 100K Project enables genetic discoveries across
Asia. Nature 576, 106-111.
18. Lee, J., Khobragade, P.Y., Banerjee, J., Chien, S., Angrisani, M., Perianayagam, A., Bloom,
D.E., and Dey, A.B. (2020). Design and Methodology of the Longitudinal Aging Study in
India-Diagnostic Assessment of Dementia (LASI-DAD). J Am Geriatr Soc 68 Suppl 3,
S5-S10.
19. Zhao, W., Smith, J.A., Wang, Y.Z., Chintalapati, M., Ammous, F., Yu, M., Moorjani, P.,
Ganna, A., Gross, A., Dey, S., et al. (2023). Polygenic Risk Scores for Alzheimer's
Disease and General Cognitive Function Are Associated With Measures of Cognition in
Older South Asians. J Gerontol A Biol Sci Med Sci 78, 743-752.
20. Morgulis, A., Gertz, E.M., Schaffer, A.A., and Agarwala, R. (2006). A fast and symmetric
DUST implementation to mask low-complexity DNA sequences. J Comput Biol 13,
1028-1040.
21. Loh, P.R., Danecek, P., Palamara, P.F., Fuchsberger, C., Y, A.R., H, K.F., Schoenherr, S.,
Forer, L., McCarthy, S., Abecasis, G.R., et al. (2016). Reference-based phasing using the
Haplotype Reference Consortium panel. Nat Genet 48, 1443-1448.
22. Loh, P.R., Palamara, P.F., and Price, A.L. (2016). Fast and accurate long-range phasing in a
UK Biobank cohort. Nat Genet 48, 811-816.
23. Conomos, M.P., Miller, M.B., and Thornton, T.A. (2015). Robust inference of population
structure for ancestry prediction and correction of stratification in the presence of
relatedness. Genet Epidemiol 39, 276-293.
24. Wang, Y.Z., Zhao, W., Moorjani, P., Gross, A.L., Zhou, X., Dey, A.B., Lee, J., Smith, J.A.,
and Kardia, S.L.R. (2024). Effect of apolipoprotein E epsilon4 and its modification by
sociodemographic characteristics on cognitive measures in South Asians from LASI-
DAD. Alzheimers Dement 20, 4854-4867.
25. Conomos, M.P., Reiner, A.P., Weir, B.S., and Thornton, T.A. (2016). Model-free Estimation
of Recent Genetic Relatedness. Am J Hum Genet 98, 127-148.
26. Ge, T., Chen, C.Y., Ni, Y., Feng, Y.A., and Smoller, J.W. (2019). Polygenic prediction via
Bayesian regression and continuous shrinkage priors. Nat Commun 10, 1776.
27. Zhang, J., Zhan, J., Jin, J., Ma, C., Zhao, R., O'Connell, J., Jiang, Y., andMe Research, T.,
Koelsch, B.L., Zhang, H., et al. (2024). An ensemble penalized regression method for
multi-ancestry polygenic risk prediction. Nat Commun 15, 3238.
28. Zhang, H., Zhan, J., Jin, J., Zhang, J., Lu, W., Zhao, R., Ahearn, T.U., Yu, Z., O'Connell, J.,
Jiang, Y., et al. (2023). A new method for multiancestry polygenic prediction improves
performance across diverse populations. Nat Genet 55, 1757-1768.
29. Werme, J., van der Sluis, S., Posthuma, D., and de Leeuw, C.A. (2022). An integrated
framework for local genetic correlation analysis. Nat Genet 54, 274-282.
30. Zhang, Y., Lu, Q., Ye, Y., Huang, K., Liu, W., Wu, Y., Zhong, X., Li, B., Yu, Z., Travers,
B.G., et al. (2021). SUPERGNOVA: local genetic correlation analysis reveals
heterogeneous etiologic sharing of complex traits. Genome Biol 22, 262.
31. Li, Y., Pawitan, Y., and Shen, X. (2025). An enhanced framework for local genetic
correlation analysis. Nat Genet.
32. Berisa, T., and Pickrell, J.K. (2016). Approximately independent linkage disequilibrium
blocks in human populations. Bioinformatics 32, 283-285.
33. Wen, X., and Stephens, M. (2010). Using Linear Predictors to Impute Allele Frequencies
from Summary or Pooled Genotype Data. Ann Appl Stat 4, 1158-1182.

34. MacDonald, J.W., Harrison, T.A., Bammler, T.K., Mancuso, N., and Lindström, S. (2023).
Ancestry-specific maps of GRCh38 linkage disequilibrium blocks for human genome
research. bioRxiv, 2022.2003.2004.483057.
35. Spence, J.P., and Song, Y.S. (2019). Inference and analysis of population-specific fine-scale
recombination maps across 26 diverse human populations. Sci Adv 5, eaaw9206.
36. Kim, S.A., Cho, C.S., Kim, S.R., Bull, S.B., and Yoo, Y.J. (2018). A new haplotype block
detection method for dense genome sequencing data based on interval graph modeling of
clusters of highly correlated SNPs. Bioinformatics 34, 388-397.
37. Teo, Y.Y., Fry, A.E., Bhattacharya, K., Small, K.S., Kwiatkowski, D.P., and Clark, T.G.
(2009). Genome-wide comparisons of variation in linkage disequilibrium. Genome Res
19, 1849-1860.
38. Kerdoncuff, E., Skov, L., Patterson, N., Zhao, W., Lueng, Y.Y., Schellenberg, G.D., Smith,
J.A., Dey, S., Ganna, A., Dey, A., et al. (2024). 50,000 years of Evolutionary History of
India: Insights from ∼2,700 Whole Genome Sequences. bioRxiv,
2024.2002.2015.580575.
39. Maier, R., Flegontov, P., Flegontova, O., Isildak, U., Changmai, P., and Reich, D. (2023). On
the limits of fitting complex models of population history to f-statistics. Elife 12.
40. Martin, A.R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B.M., and Daly, M.J. (2019).
Clinical use of current polygenic risk scores may exacerbate health disparities. Nat Genet
51, 584-591.
41. Yengo, L., Vedantam, S., Marouli, E., Sidorenko, J., Bartell, E., Sakaue, S., Graff, M.,
Eliasen, A.U., Jiang, Y., Raghavan, S., et al. (2022). A saturated map of common genetic
variants associated with human height. Nature 610, 704-712.
42. Yengo, L., Sidorenko, J., Kemper, K.E., Zheng, Z., Wood, A.R., Weedon, M.N., Frayling,
T.M., Hirschhorn, J., Yang, J., Visscher, P.M., et al. (2018). Meta-analysis of genome-
wide association studies for height and body mass index in approximately 700000
individuals of European ancestry. Hum Mol Genet 27, 3641-3649.
43. Zou, Y., Carbonetto, P., Wang, G., and Stephens, M. (2022). Fine-mapping from summary
data with the "Sum of Single Effects" model. PLoS Genet 18, e1010299.
44. Zhou, X. (2017). A Unified Framework for Variance Component Estimation with Summary
Statistics in Genome-Wide Association Studies. Ann Appl Stat 11, 2027-2051.
45. Mancuso, N., Freund, M.K., Johnson, R., Shi, H., Kichaev, G., Gusev, A., and Pasaniuc, B.
(2019). Probabilistic fine-mapping of transcriptome-wide association studies. Nat Genet
51, 675-682.
46. Li, Z., Gao, B., and Zhou, X. (2025). An alternative framework for transcriptome-wide
association studies to detect and decipher gene-trait associations. bioRxiv,
2025.2003.2014.643391.
47. Turcot, V., Lu, Y., Highland, H.M., Schurmann, C., Justice, A.E., Fine, R.S., Bradfield, J.P.,
Esko, T., Giri, A., Graff, M., et al. (2018). Protein-altering variants associated with body
mass index implicate pathways that control energy intake and expenditure in obesity. Nat
Genet 50, 26-41.
48. Finer, S., Martin, H.C., Khan, A., Hunt, K.A., MacLaughlin, B., Ahmed, Z., Ashcroft, R.,
Durham, C., MacArthur, D.G., McCarthy, M.I., et al. (2020). Cohort Profile: East
London Genes & Health (ELGH), a community-based population genomics and health
study in British Bangladeshi and British Pakistani people. Int J Epidemiol 49, 20-21i.

49. Yu, K., Das, S., LeFaive, J., Kwong, A., Pleiness, J., Forer, L., Schonherr, S., Fuchsberger,
C., Smith, A.V., and Abecasis, G.R. (2022). Meta-imputation: An efficient method to
combine genotype data after imputation with multiple reference panels. Am J Hum Genet
109, 1007-1015.
50. Das, S., Forer, L., Schonherr, S., Sidore, C., Locke, A.E., Kwong, A., Vrieze, S.I., Chew,
E.Y., Levy, S., McGue, M., et al. (2016). Next-generation genotype imputation service
and methods. Nat Genet 48, 1284-1287.
51. Campbell, M.C., and Tishkoff, S.A. (2008). African genetic diversity: implications for
human demographic history, modern human origins, and complex disease mapping.
Annu Rev Genomics Hum Genet 9, 403-433.
52. Dokuru, D.R., Horwitz, T.B., Freis, S.M., Stallings, M.C., and Ehringer, M.A. (2024). South
Asia: The Missing Diverse in Diversity. Behav Genet 54, 51-62.
53. Saleheen, D., Zaidi, M., Rasheed, A., Ahmad, U., Hakeem, A., Murtaza, M., Kayani, W.,
Faruqui, A., Kundi, A., Zaman, K.S., et al. (2009). The Pakistan Risk of Myocardial
Infarction Study: a resource for the study of genetic, lifestyle and other determinants of
myocardial infarction in South Asia. Eur J Epidemiol 24, 329-338.
54. Ziyatdinov, A., Torres, J., Alegre-Diaz, J., Backman, J., Mbatchou, J., Turner, M., Gaynor,
S.M., Joseph, T., Zou, Y., Liu, D., et al. (2023). Genotyping, sequencing and analysis of
140,000 adults from Mexico City. Nature 622, 784-793.
827