Pulsatile flow dynamics maintain pulmonary arterial architecture

Stephen Spurgin,[1,2] Lauren Thai,[2] Tina Wan,[3,4] Christopher P. Chaney,[5] Mitzy A. Cowdin,[1]

Suren V. Reddy,[2] M. Tarique Hussain,[2,6] Munes Fares,[2] M. Luisa Iruela-Arispe,[7] Thomas

Carroll,[5] Andrew D. Spearman,[3] and Ondine Cleaver[1]*

[1]Department of Molecular Biology, University of Texas Southwestern Medical Center, 5323

Harry Hines Blvd., Dallas, Texas, USA 75390.

[2]Department of Pediatrics, Division of Cardiology, University of Texas Southwestern Medical

Center, 5323 Harry Hines Blvd., Dallas, Texas, USA 75390.

[3]Department of Pediatrics, Division of Cardiology, Medical College of Wisconsin, Children's

Wisconsin, Herma Heart Institute, Milwaukee, Wisconsin, USA.

[4]Cardiovascular Center, Medical College of Wisconsin, Milwaukee, Wisconsin, USA.

[5]Department of Internal Medicine and Division of Nephrology, University of Texas

Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, Texas, USA 75390.

[6]Department of Radiology, University of Texas Southwestern Medical Center, 5323 Harry

Hines Blvd., Dallas, Texas, USA 75390.

[7]Department of Cell and Developmental Biology, Northwestern Fienberg School of Medicine,

Chicago, Illinois, USA 60611

*Corresponding author:

Ondine Cleaver

Department of Molecular Biology, University of Texas Southwestern Medical Center

5323 Harry Hines Blvd., NA8.300 Dallas, Texas 75390-9148, USA.

Phone: (214) 648-1647 Fax: (214) 648-1196

E-mail: ondine.cleaver@utsouthwestern.edu

Running title: **Pulsatile stretch drives mural thickness**

Word Count: 11,233 (all text inclusive)

28
Keywords: PDGFB, endothelial cell, mural cell, biomechanical force, hemodynamics,

pulsatility, Glenn, single ventricle, congenital heart disease, rat, mouse

Non-standard Abbreviations and Acronyms

AV - arteriovenous

GSEA - Gene Set Enrichment Analysis

HAECs - human aortic endothelial cells

HPAECs - human pulmonary arterial endothelial cells

IF - immunofluorescence

IHC - immunohistochemistry

LPA - left pulmonary artery

L-SVC - left superior vena cava

NHLFs - normal human lung fibroblasts

PA - pulmonary artery

PASMCs - human pulmonary artery smooth muscle cells

PAVMs - microvascular pulmonary arteriovenous malformations

PDGFB - platelet derived growth factor B

PDGFR β - platelet derived growth receptor β

RPA - right pulmonary artery

SVC - superior vena cava

SV-CHD - Single ventricle congenital heart disease

vSMC - vascular smooth muscle cell

<u>Abstract</u>

Single ventricle congenital heart disease (SV-CHD) is a uniformly lethal condition requiring

the Glenn surgery, which as a side effect eliminates arterial pulsatility and contributes to

pulmonary vascular complications. In Glenn patients, we quantified pulsatility loss in each

dimension of force (flow, pressure, and stretch) using cardiac catheterization and MRI. To

model and investigate the individual impact of each dimension of pulsatility loss on the

pulmonary vasculature, we applied isolated pulsatile and non-pulsatile mechanical stimuli

to pulmonary arterial endothelial cells (ECs) in vitro. We found that each dimension of

force triggered distinct transcriptional responses, revealing force-specific regulation of

structural and signaling pathways. Pulsatile stretch uniquely stimulated EC secretion of

PDGFB, a key driver of vascular smooth muscle cell (vSMC) recruitment. In a rat Glenn

model, loss of pulsatility led to vascular wall thinning, confirming *in vivo* relevance. Our

findings uncover a mechanistic link between endothelial stretch sensing and PDGFB-

mediated EC-vSMC crosstalk, essential for maintaining pulmonary artery architecture.

Clinically, these insights suggest that restoring or mimicking pulsatile forces may help

preserve vascular integrity and prevent remodeling in SV-CHD patients.

Introduction

Globally, 1 in every 2000 babies is born with a heart that has a single functional ventricle (1). Surviving this congenital abnormality requires early and significant medical intervention. A single ventricle cannot perform the work of two for very long, nor can it direct an equal amount of blood flow to the lungs (low vascular resistance) and the body (high vascular resistance). For children with single ventricle congenital heart disease, survival depends on a series of remarkable surgeries: the Glenn and the Fontan. In the circulation created by these procedures, venous blood flows directly to the lungs without the support of a pumping ventricle, while the single functional ventricle supplies pulsatile flow to the rest of the body (2).

A frequent consequence of surgical replumbing of the vasculature in single ventricle patients is the development of a wide variety of vascular malformations: A) diffuse microvascular pulmonary arteriovenous malformations (PAVMs) in the Glenn stage and B) larger, tortuous collateral vessels in both Glenn and Fontan stages. It has been shown that the microvascular lesions that arise in the Glenn are reversible with restoration of the direct flow of liver blood to the lungs in the Fontan (albeit still lacking cardiac pulsatile flow). This has led to the "hepatic factor" hypothesis (3-5). However, interestingly, the proximal macrovascular collateral vessels in the Fontan do not regress (6, 7). Among the abundant potential pathogenic factors—chronic hypoxemia, chronic inflammation, and loss of direct hepatopulmonary blood flow (3, 5, 8)—a role for pulsatility has long been suggested (9-11), but never investigated at the mechanistic level.

Pulsatility, for its part, is a critical characteristic of arterial blood flow. Arteries—whether systemic, pulmonary, or umbilical—are *defined* by the presence of pulsatile blood flow, not by their color or oxygen content. The structure of the arterial wall—with thick

smooth muscle layer and elastic laminae—reflects the need to withstand the high

pulsations of arterial pressure and stretch. Veins, on the other hand, have less

hemodynamic pulsatility and a thinner smooth muscle layer (media) (12). Critically, the

histopathology of vascular malformations often reveals absent or patchy coverage of the

smooth muscle layer (13, 14). This mural insulation must be broken down to form new

connections between vessels. EC proliferation is a then a critical step in the pathogenesis

of vascular malformations (13). Once a connection is formed, the transmission of pulsatile,

arterial blood flow to venous vessels has been shown to induce "arterialization" (medial

thickening, intimal hyperplasia) of the venous vessels (15, 16), further highlighting the

relationship between arterial hemodynamics and the architecture of the vascular wall.

Although arteriovenous differences are most visually evident by their mural

coverage, the endothelial cells of arteries and veins are themselves distinct. They have

unique transcriptional and proteomic signatures (17), and unique developmental origins

(18, 19). The impact of cardiac pulsatility on the arterial side of the vasculature has been

appreciated in the early stages of vascularization (20)—however, the role of *pulsatility* in

maintaining arterial and venous endothelial identity and function *after* birth has received

very limited attention. Scientific investigation of EC mechanobiology has focused on other

physical forces, such as the cellular impact of low versus high shear stress, or of laminar

versus disturbed flow (21). Despite the clinical centrality of pulsatility, it remains an under-

investigated aspect of vascular biology.

ECs are remarkably sensitive to the physical forces transmitted by blood flow.

Endothelial mechanosensitive components include cell-cell adhesion molecules (integrins,

gap-junctions, cadherins), extracellular matrix (ECM) sensing components, ion channels,

and more (21, 22). The forces sensed by these molecular components are applied to ECs

in three dimensions: shear stress , pressure, and stretch (23). Each dimension of force

therefore exerts distinct stresses and deformations to ECs. Laminar shear stress (LSS) is

the frictional drag force that occurs along the inner (apical) surface of the vessel

endothelium (24). Stretch is the effect downstream of each heartbeat, as vessels widen

radially to accommodate an ejected volume of blood and their circumference expands (25).

Hydrostatic pressure is generated by ventricular contraction, and the driving force behind

stretch (26). Notably, ECs align parallel to the direction of LSS (blood flow) but

perpendicular to the direction of stretch (27)—further highlighting the need to consider

each dimension's impact separately. All three biomechanical forces have all been shown

to alter vascular ECM (28), as well as endothelial and smooth muscle transcriptional

programs (29, 30). However, the unique impact of pulsatility within each dimension of force

on vascular homeostasis remains a critical knowledge gap.

The endothelium and vascular smooth muscle communicate closely. Critically,

cross-talk between ECs and vSMCs is mediated by EC-secretion of platelet derived growth

factor B (PDGFB), which binds to PDGF receptor β (PDGFR β) on vSMCs, where it drives

proliferation and dictates mural cell coverage from large vessels down to the capillary level

(31, 32). Loss of Notch signaling in pericytes downregulates PDGFRB in vSMCs in mouse

models, leading to AVMs (33). Conversely, deleting PDGFB specifically from ECs has

shown remarkable dilation and arteriovenous shunting in the lungs of mice (34), and loss

of PDGFB has been observed in human vascular malformations (35). The question arises

as to whether different hemodynamic forces coordinate the maintenance of vascular

structure via EC responses to these forces.

Here, we investigate the biomechanical forces experienced by pulmonary ECs in

Glenn patients and identify pulsatility of blood flow as critical to maintenance of the lung

vasculature. We first use combined cardiac catheterization (cath) and cardiac magnetic

resonance imaging (MRI) to quantify and compare the pulsatility of force that is applied to

pulmonary arterial ECs in children with normal cardiopulmonary anatomy and in those with

Glenn circulation. After noting a clear loss of pulsatility in the Glenn pulmonary arteries, we

model those changes using cultured human pulmonary arterial endothelial cells (HPAECs)

to define the specific transcriptional programs driven by each individual dimension of force.

We demonstrate that pulsatile stretch in vitro, which mimics expansion of the arterial

vessel intima during cardiac ventricular systole, significantly impacts the transcriptional

landscape of ECs. Specifically, we show that pulsatile stretch stimulates ECs to secrete

PDGFB. In addition, using our previously published in vivo rat model of the Glenn surgery,

where pulmonary arteries (PAs) experience loss of pulsatile stretch, we demonstrate

significant reduction of PDGFB in PA ECs. Finally, we build on our recent work in a rat

model of the Glenn surgery(36), which showed early and progressive arteriovenous

shunting, to demonstrate significant thinning of pulmonary arterial vascular walls.

Together, our study identifies pulsatile stretch as a critical and underappreciated

dimension of arterial force that drives EC-secretion of PDGFB, promoting vascular smooth

muscle endowment and providing needed structural support for arterial vessels. Gaining

insight into the impact of *pulsatility* of hemodynamic force is a critical and novel approach

to understanding clinically relevant endothelial cell biology.

Results

The Glenn shunt directs venous blood flow to pulmonary arteries

Arterial vessels normally experience blood flow directly from a pumping cardiac ventricle, adding pulsatility to the magnitude of hemodynamic forces (**Figure 1A**). By reviewing the known characteristics of three distinct pairings of arteries and veins (systemic, pulmonary, and umbilical), we note that pulsatile flow is a shared, definitional quality of arteries (**Supplemental Figure 1**). Pulsatility is thus an inherent quality of blood flow in arterial vessels. After a Glenn surgery, however, the pulmonary arteries no longer receive blood flow directly from a pumping heart ventricle. Instead, pulmonary blood flow is provided by the Glenn shunt: direct venous flow through the superior vena cava (SVC), which is anastomosed to the pulmonary arteries (PA) (**Figure 1A'**). ECs that make up the inner lining of pulmonary arteries downstream of normal cardiac vessel anatomy are subjected to force in three dimensions: flow/sheer stress, pressure, and circumferential stretch (**Figure 1B**) (23). Here, we assess these forces in the Glenn lung arteries.

Pulsatility loss in three dimensions in Glenn patients

To quantitatively determine the loss of pulmonary arterial pulsatility in patients with Glenn anatomy, we used combined cardiac catheterization and cardiac MRI (cath/MRI) in 20 Glenn patients to obtain simultaneous data in all three dimensions of force: pressure, flow, and circumferential stretch (**Figure 1B**). These data were compared to data obtained in 20 age- and gender-matched patients with normal cardiopulmonary connections (see patient demographics in **Supplemental Table 1**). First, as has long been observed clinically, we observed pulsatility loss in the pulmonary arterial *pressure* of Glenn patients

(**Figure 1C, C'**). Similarly, we noted that the Glenn shunt leads to non-pulsatile shear

stress in the pulmonary arteries, as seen by a comparison of the pulmonary artery velocity

profiles in normal (**Figure 1D**) and Glenn (**Figure 1D'**) patients throughout the cardiac

cycle. Finally, we found that the circumference of proximal right pulmonary artery does not

expand repeatedly with each heartbeat in Glenn patients, demonstrating loss of pulsatile

circumferential stretch (**Figure 1E,E'**).

Unique transcriptome for each dimension of pulsatile force

ECs are highly responsive to mechanical force, and can react to differences in both

magnitude and direction of force.(37) Given the observed loss of pulsatility in Glenn

patients, we next investigated the relative impact of pulsatility within each dimension of

force on ECs.

Using our patient cath/MRI pulsatility data, we designed in vitro experiments that

model the pulsatility loss defined in Glenn patients. Each isolated, single dimension of

force was applied to cultured primary human pulmonary artery endothelial cells (HPAECs)

in a pulsatile and non-pulsatile manner. For pulsatile shear stress, we applied 0-15

dyn/cm^2 shear stress at 1 Hz, which mimics the frequency of a resting heart rate. For

pulsatile pressure, we modified the Ibidi equipment to expose HPAECs to columns of

media positioned to alternately provide 5 or 25 mmHg pressure, at 1 Hz, in the absence of

flow or stretch. For stretch, we used a Flexcell system to stretch HPAEC by 10% of their

baseline in a single axis (again at a frequency of 1 Hz). To assess stable transcriptional

changes between non-pulsatile and pulsatile conditions and thus improve in vivo relevance

of our findings, HPAECs were exposed to 48 hours of force or pulsatile force (**Figure 2A**).

To best mimic in vivo conditions in the years of the Glenn stage, pulsatile stretch was

compared to no stretch rather than continuous stretch, which does not happen in vivo. We

observed similar physical changes as have been reported previously: ECs align parallel to

the axis of flow and perpendicular to the axis of stretch and exhibit no alignment

preference under continuous or pulsatile pressure (**Figure 2B**).

The unique changes in cell shape in response to each axis of force highlight the

need to examine them individually. We first investigated the overall pathways that were

regulated by pulsatility. By Gene Set Enrichment Analysis (GSEA) of bulk RNAseq data,

we noted that each dimension of force drove changes in unique signaling pathways, and

that the impact of shear stress was the most distinct. Overall, in HPAECs, laminar flow

affected 19 unique pathways, stretch 10 unique pathways, and pressure 4 unique

pathways (**Figure 2C**). A wide range of pathway types were regulated, many related to

changes in structural cellular components, growth, or interaction of the cell with ECM

(**Figure 2D**). We note that mitogen-activated protein kinase (*Mapk*) and epidermal growth

factor receptor (*ErbB*, also known as *HER*) signaling were only induced by pulsatility of

flow, while Hedgehog and Notch signaling were only induced by pulsatility of stretch.

Pulsatile pressure did not uniquely regulate any obviously EC-related pathways, but did

upregulate the hypertrophic cardiomyopathy pathway.

In addition to these dimension-specific pathways, we find other pathways that are

influenced across multiple dimensions of force. We found 10 pathways upregulated by all 3

dimensions of force, and 17 pathways induced by two of the three dimensions. We note

that shear stress, stretch, and pressure all drove changes in the pathway "Regulation of

Actin Cytoskeleton." However, an analysis of the leading-edge subset (the genes within

the pathway that are most significantly changed and thus responsible for the overall

significance of the pathway) revealed upregulation of different genes by each dimension of

force. For example, within the "Regulation of Actin Cytoskeleton" pathway, pulsatile

pressure shows *Actn1* and pulsatile flow yields *Map2k1*, while pulsatile stretch promotes

Pdgfb, *Arghef6,* and others (**Figure 2E**). Different dimensions of force therefore drive

different transcriptional changes that converge on similar pathways.

Pulsatility does not alter arteriovenous genes in vitro

To test the impact of each dimension of force on ECs, we next analyzed gene

transcription. Using the culture platforms shown in **Figure 3A**, we assessed bulk RNAseq

after application of single dimensions of pulsatile and non-pulsatile force. Strikingly, we

found very little overlap between genes enriched in ECs under different dimensions of

force, suggesting distinct responses to distinct forces. Importantly, we did not find

significant changes in baseline expression of common EC genes between the three

experimental set-ups (**Supplemental Figure 2**), indicating no basic alteration of

endothelial fate. Of the genes significantly regulated by pulsatility, 93.4% (1205/1290) were

driven by a single, unique dimension of force. Pulsatility of stretch accounted for over two-

thirds (888 genes, 68.9%) of these genes (**Figure 3B**). Only 59 genes were uniquely

induced by pulsatility of laminar flow, while 343 were uniquely induced by pulsatility of

pressure. We found less overlap in individual genes regulated by pulsatility than there was

in overall pathways. Twenty-two genes were regulated by pulsatility of both LSS and

stretch, 56 genes by stretch and pressure, and 5 by pressure and LSS.

Finally, we hypothesized that application of pulsatile force on cultured ECs would

induce greater expression of genes previously shown to be specific to arteries (17). We

asked whether the classical markers of arteriovenous (AV) EC differentiation were found in

these groups of genes. In cultured HPAECs, we found no dimension of pulsatility that

significantly affected the expression of a majority of developmental arterial (*Efnb2*, *Eng*,

Sox17, *Notch1, Jag1, Nrp1, Bmx)* or venous (*Ephb4*, *Emcn*, *Nr2f1*, *Nr2f2*/*Coup-TFII, Clu*)

driver genes (**Figure 3C**). Similarly, we found no significant changes arterial or venous

standard marker genes in rat Glenn arterial ECs compared to sham operated rats (data not

shown).

We next asked if there were genes specifically induced by pulsatile LSS, pressure

or stretch, when compared to the non-pulsative forces. After applying a p-value cut-off of

0.05 and fold change cut-off of 2, we found that pulsatile stretch drove a change in 44

genes, compared to 20 genes by pulsatility of flow, and 16 genes by pulsatility of pressure

(**Figure 3D-F**). Expression of these hits in data from other dimensions of force is shown in

Supplemental Figure 3. Genes upregulated under pulsatile LSS included many factors

involved in cell proliferation and immunity (red highlighted genes in **Figure 3D**), and genes

upregulated under pulsatile pressure included factors associated with the cytoskeleton and

RNA processing (red highlighted genes in **Figure 3E**). Gene expression was particularly

impacted by application of pulsatile stretch, with many factors involved in tissue

remodeling and inflammation being upregulated (red highlighted genes in **Figure 3F**) and

factors driving cell division and cell survival pathways (blue highlighted genes in **Figure**

3F).

Identification of *Pdgfb* as a stretch-induced gene in pulmonary arterial endothelium

As stretch accounted for the majority of pulsatility-regulated genes, we utilized

published single cell RNA sequencing of human lung tissue (LungMAP(38, 39)) to assess

the cell type-specific expression pattern of candidate genes. We reasoned that narrowing

the investigation to EC-specific genes might identify endothelial factors that sense flow and

normally respond by bolstering matrix or cellular components of the arterial media that

would increase resistance in proportion to hemodynamic load. *Mgp*, *Cx3cl1*, and *Pdgfb*

were highly expressed in a good proportion of ECs under stretch, but we noted high EC

specificity of the latter two genes (**Figure 4A**). Analysis of the expression of these ligands

and their receptors (*Cx3cl1R* and *Pdgfrb*, respectively) underscored the pulsatility-induced

crosstalk between ECs and immune cells (for *Cx3cl1*) and between ECs and perivascular

cells (for *Pdgfb*) (**Figure 4B-D**). As PDGFB is i) well known to promote smooth muscle cell

proliferation and recruitment (31), ii) a key component of the GSEA for ECs under stretch

(**Figure 2**), and iii) a significant hit in their differential gene expression (**Figure 3**), we

investigated further its role in pulmonary vessels.

Pulsatile stretch uniquely drives EC PDGFB secretion

We next asked if stretch-induction of PDGFB expression was specific to ECs or also

present in other cellular components of the vessel wall (vascular smooth muscle or

fibroblasts). We applied cyclic 10% stretch to HPAECs from two different patients, human

aortic endothelial cells (HAECs), human pulmonary artery smooth muscle cells (PASMCs),

and normal human lung fibroblasts (NHLFs). We found that pulsatile stretch robustly

induced secretion of PDGFB exclusively from endothelial cell lines (**Figure 4E**). As

PDGFB is known to be induced by flow (40), we compared PDGFB secretion from ECs

under continuous flow (mimicking venous condition) to pulsatile stretch (mimicking arterial

condition). Induction of PDGFB by pulsatile stretch was 5x higher than the induction by

laminar, continuous flow (**Figure 4F**).

We next assessed whether we could detect EC secretion of PDGFB in human

patients following similar pulsatile stretch-induction. We anticipated that we would not be

able to detect a change in the circulating blood, as PDGFB is secreted basally to the

perivascular space, not apically into the blood stream, as shown previously (41). To do

this, we tested the blood of patients with either normal or Glenn anatomy. As expected,

analysis of PDGFB expression in plasma obtained from the pulmonary artery and

pulmonary vein revealed significant variation in local PDGFB expression in both normal

and Glenn patients that could not be correlated with pulmonary arterial pulsatility (**Figure**

4G).

Decreased PDGFB expression in pulmonary arteriole ECs of the Glenn rat model

Given the finding that stretch induces PDGFB secretion from pulmonary ECs, we

utilized our animal model of the Glenn surgery to test whether arteries receiving non-

pulsatile flow exhibited decreased endothelial PDGFB synthesis in vivo. We performed an

end-to-end anastomosis of the left SVC (L-SVC) and the left pulmonary artery (LPA)

(**Figure 5A, A'**). This mimics the Glenn surgery, directing only non-pulsatile venous blood

flow from the head and upper extremity into the pulmonary artery. As previously described,

these rats develop hypoxemia secondary to intrapulmonary arteriovenous shunting.(36)

The hypoxemia develops rapidly (after two weeks) and is progressive (to six months (36)).

To test whether arteries receiving non-pulsatile flow exhibited decreased PDGFB

synthesis in their ECs, we focused on the distal lung arterioles—in the vicinity of where

AVMs form in our patients, and where EC-specific expression of PDGFB is highest by

single cell RNAseq (**Figure 4A**). To localize these arterioles and the PDGFB expression

within the ECs, we used immunofluorescent staining with antibodies recognizing the

endothelial junctional marker VE-cadherin (CDH5), the vascular smooth muscle marker

alpha smooth muscle actin (aSMA) and the PDGFB protein (42). We found significant

reduction of endothelial PDGFB in arterioles of the Glenn rat (**Figure 5B**). Critically, we

also noted loss of smooth muscle surrounding these distal arterial vessels in the Glenn.

Quantification of PDGFB IF signal from 5 arterioles each from 3 sham and 3 Glenn-

operated left lungs revealed a significant decrease in endothelial PDGFB expression in the

Glenn arterioles, but not in alveolar capillary ECs (**Figure 5C**).

Rat Glenn displays reduced pulmonary arterial smooth muscle

We hypothesized that a further consequence of reduced PDGFB in pulmonary ECs

under the non-pulsatile Glenn hemodynamics would be loss of the smooth muscle layer.

We analyzed sagittal sections of the left lung of 9 Glenn rats and 5 sham-operated rats

and performed immunohistochemistry (IHC) for smooth muscle actin (SMA). Strikingly, we

found a significant decrease in arterial smooth muscle thickness (SMA+ cells) (**Figure 6A,**

B), while venous smooth muscle thickness remained unchanged (**Figure 6C**). We

observed this reduction in vascular wall thickness in both the largest proximal arteries

(**Figure 6A-C**), as well as the distal arterioles, less than 0.1 mm in diameter (**Figure 6D-**

G).

Discussion

In this study, we identify hemodynamic pulsatility as critical to maintenance of the pulmonary arterial architecture (**Figure 7**). We show that in patients with Glenn circulation the shunted pulmonary blood flow—from the SVC directly to the PA—exerts no pulsatile stretch on the pulmonary artery. Using transcriptomic analysis, we show that cultured pulmonary arterial ECs exposed to each dimension of force elicit distinct molecular signatures. We find that pulsatile stretch induces significant endothelial secretion of PDGFB, known to mediate recruitment of vSMCs to blood vessels (31, 32). As lung biopsies from living patients were not obtainable, our rat model of the Glenn circulation allowed us to investigate the effect of pulsatility loss in vivo. Using this animal model, we show that non-pulsatile, venous flow within an arterial vessel, results in loss of PDGFB protein from ECs and concomitant thinning of the vascular mural wall. Our complementary in vitro and in vivo findings identify a new target signaling pathway for novel therapeutic approaches to treat the diverse vascular malformations that occur in Glenn patients.

Despite clinical appreciation of the fact that Glenn patients experience venous-type, non-pulsatile blood flow in pulmonary arteries, and even though pulsatility loss was the first hypothesized mechanism for Glenn PAVMs (43, 44), pulsatility loss has never been formally documented in three dimensions. Until now, studies comparing surgical outcomes of Glenn patients have only occasionally been able to report some dimensions of pulsatility loss (9, 45), although most studies report none (46-52). This clinical knowledge gap has limited our understanding of the impact of pulsatility within these patients. Our use of combined cardiac catheterization and cardiac MRI (cath/MRI) ensures that these

interdependent forces are measured simultaneously, providing with high temporal

accuracy the first complete 3D view of pulsatility loss in the Glenn pulmonary arteries.

Our pulsatility metrics merit two points of discussion. First, we utilize flow velocities

as a surrogate for wall shear stress (WSS)—the direct hemodynamic force applied to the

endothelium. As WSS depends directly on the velocity gradient, velocity is a reasonable

approximation for this investigation. Given the limited availability and accuracy of WSS

measurements with cardiac MRI, use of velocity instead of the highly regionally variable

WSS supports an easier and more rapid clinical application. Second, the relationship

between pressure and stretch (vascular compliance) is not directly tested in our study. This

will be a critical area of investigation in the future, to answer the question of how much

pulsatility (of flow, pressure, or stretch) is present in the distal lung.

Indeed, estimating the magnitude of pulsatile forces across the pulmonary vascular

tree is difficult. Arterial pulsatility in normal patients is likely significantly diminished as it

reaches distal capillaries. Dynamic measurement of distal pulsatility of stretch in sub-

millimeter vessels in vivo is technically challenging and could not be directly assessed by

our methods. In a normal compliant vessel, proximal stretch acts as a capacitor, absorbing

and lessening distal pulsatility—a phenomenon termed the 'windkessel effect' (53). This

model suggests a reduction of pulsatility in the normal distal microvasculature. Our

proposed mechanism—that pulsatile stretch of ECs stimulates PDGFB followed by smooth

muscle coverage—is supported by the observation that normal arterial smooth muscle

thins as vessels branch further into the lung (12). PDGFB was originally described as flow

responsive (40), and later as stretch responsive (54), although in our in vitro work stretch

elicited 5-fold more PDGFB secretion than flow. The opportunity now arises to characterize

the relative contribution of shear and stretch to PDGFB secretion in these distal vessels.

We also recognize that the presence of respiratory-driven alveolar stretch (at a much lower

frequency) may also contribute to distal pulsatility (55, 56). In all, these studies underscore

the importance of pulsatility throughout the entire arterial network and bring up new

questions about the impact of biomechanical cues on the lung vasculature.

How do ECs sense hemodynamic forces in lung vessels? We previously

hypothesized that endothelial cilia could be involved in sensing pulsatility of flow (57) but

disproved this possibility in the mammalian lung as pulmonary ECs are devoid of primary

cilia (58). Endothelial mechanosensitive components have been well described and are

present on lung ECs. Notch, Cdh5, and many other cell-surface proteins, ECM, and

cytoskeletal elements are all required for proper sensing of flow or stretch (23). In

particular, the protein complex of PECAM1, CDH5, VEGFR2/3, and PLXND1 plays a

critical role in force-sensing, while proteins like LPHN2 and others coordinate downstream

responses (59). In our data, we see significant upregulation of *Plxnd1*, as well as near two-

fold increases in *Cdh5, Kdr,* and *Lphn2* (data not shown) under pulsatile pressure

conditions, but not pulsatile stretch or LSS —suggesting a possible role for pulsatile

pressure in priming the endothelium to respond to force. Our study thus highlights the

need to consider each dimension of force separately, as we continue to evaluate the role

that pulsatility plays in the maintenance of the normal vascular architecture.

One key question is how loss of pulsatility impacts arteriovenous fate in Glenn

vessels. Our studies show that the Glenn shunt supplies venous-type hemodynamic forces

to an artery, as blood flow pulsatility is lost following surgical intervention. The observed

outcome of this change in biomechanical forces is that the artery becomes structurally

more like a vein, in that its vascular wall is significantly diminished in thickness. While

enhancement of arterial EC markers by pulsatility of LSS has been observed previously in

vitro (20), in our studies, we did not find enhancement of classic markers of arterial EC

identity under pulsatile conditions (in any dimension of force), nor did we find shifts in

arteriovenous identity in our rat Glenn model (data not shown). It is possible—if unlikely—

that this difference is due to the highly laminar flow provided by the Ibidi system in our

work, compared to the more turbulent flow in the cone-disc viscometer used in the prior

study. Separately, microvessels exposed to arterial anastomosis have been observed to

increase vSMC coverage (60), however changes in EC arteriovenous identity were not

noted. These observations support our conclusion that pulsatility drives the arterial

vascular structure as a whole (EC + surrounding mural cells), rather than the shifting the

expression of developmentally-regulated markers of arteriovenous EC identity.

Growing evidence has pointed to a potential venous origin of AVMs (61, 62), while

our study focused on upstream impacts within lung arteries and upregulation of endothelial

PDGFB. Endothelial deletion of *Pdgfb* (using Cdh5-CreERT2) in mouse leads to vascular

malformations in mouse lungs, while arterial-specific ablation (using Bmx-Cre) does not

(34). However, we note that in these studies, ablation was induced over a short timeframe

(1 week) and only retinal arteries were assessed. It will be important to examine how

arterial loss of *Pdgfb* over a longer timeframe, like that experienced by patients and Glenn

rats, impacts downstream PAVMs. Hence, the mechanisms and time course of how

upstream biomechanical cues impact arterial ECs and the downstream endothelium of

capillaries and veins, and whether there are organotypic effects, remains to be seen.

Other cells of the vascular wall, such as vSMCs, also certainly sense biomechanical

forces. VSMCs experience stretch similar to ECs, but not LSS as they are not in direct

contact with blood flow. Interestingly, vSMCs respond differently to stretch than ECs,

aligning perpendicular to stretch in vivo, rather than parallel like ECs (63). In our study and

in publicly available single cell RNA sequencing from human lung, we also find that smooth

muscle cells do not secrete PDGFB. We note that vSMCs from rat Glenn do not show

changes in standard vSMC genes such as *myh11, tagln, myocd, cnn1*, and *pdgfrb* (data

not shown). This does not rule out, however, the possible contribution of vSMCs to the

ECM, especially regarding the alterations in pulmonary vascular resistance and vessel

stiffening that so critically drive patient outcomes. Future studies will be necessary to

investigate the responses of vascular smooth muscle to pulsatile stretch.

While we focused here on *Pdgfb*, we also note that under pulsatile flow there is a

significant upregulation of *Col1a2*, a major component of vascular wall. Loss of *Col1a2* is

associated with vasculopathy and aortic dilation (64), suggesting that pulsatility of stretch

likely drives other EC contributions to the structure of the arterial wall—and arterial

vascular identity—beyond the recruitment of vSMCs. Regulation of the ECM is a logical

complimentary function of pulsatility, as the scaffold for the vSMCs and the internal elastic

lamina of the largest arteries are unique features of overall arterial structure. Our work thus

highlights a distinction between the traditional markers of arteriovenous endothelial identity

(EC-centric) and arteriovenous vascular identity (the overall structure of the vessels

cellular layers and ECM).

In addition, we also noted the upregulation of CX3CL1 in pulsatile-stretched

HPAECs. CX3CL1 is a well-known chemokine that increases leukocyte recruitment and

adhesion to ECs (65, 66). Multiple other chemokines (CCL2, CCL14) are regulated by

pulsatility in our in vitro system. The role of the immune system in the vascular

complications of Glenn circulation is a wholly unappreciated factor; however, previous

work in hereditary AVMs identified accumulation of leukocytes in brain AVMs and skin

telangiectasias (67-69). Collectively, these findings suggest that pulsatile stretch may

impact EC-leukocyte crosstalk in pathogenesis of vascular malformations.

Translation of our findings will depend on the relative role of pulsatility in the

pathogenesis of the different vascular malformations that arise in the Glenn lung. They are

highly diverse, and include both large "collateral" vessel formation proximally, as well as

microvascular arteriovenous malformations distally (7). As patients can present with

primarily proximal, primarily distal, or both types of malformations, it is highly likely that the

pathogenesis of these lesions is multifactorial, with both common and distinct elements. As

previously noted, there is 50 years of evidence that a "hepatic factor" is the primary

pathogenic factor for distal microvascular PAVMs. However, we suggest it is not likely the

only factor. An argument for the hepatic factor as the primary pathogenic factor is the

observation that distal PAVMs are reversible in the Fontan stage, which restores direct

hepatopulmonary venous blood flow but not normal pulsatility. However, the proximal

collateral vessels do not resolve in the Fontan, pointing to a critical role for pulsatile flow in

vascular architecture. In our study, we observe thinning of vSMCs both proximally and

distally in the animal model and therefore suggest that pulsatility may be a primary factor in

the generation of proximal collateral vessels and an exacerbating factor in the

development of distal PAVMs.

The relative contribution of pulsatility and hepatic factor to the pathogenesis of distal

AVMs can be assessed by comparing the time of onset in the Glenn and other clinical

scenarios. In Glenn patients (no hepatic factor, no pulsatility), AVMs can develop in

months (7, 70-72). However, in a rare subset of heterotaxy patients (no hepatic factor,

normal pulsatile flow) (73-76), AVMs form more slowly. In these latter patients, who have

interrupted inferior vena cava (strongly suggesting left isomerism heterotaxy syndrome

(77)), the hepatic veins can (rarely) drain directly to the left atrium in an otherwise normal

biventricular heart with intact atrial and ventricular septa. In this scenario (pulsatile

pulmonary blood flow, diversion of hepatic venous blood from the lungs), distal PAVMs can

take years or even decades to appear (73-76). This suggests that loss of pulsatility

critically accelerates the pathogenesis of distal pulmonary AVMs but is not sufficient to

induce them.

Thus, while loss of pulsatility may not be the defining factor that leads to distal

pulmonary AVMs in the Glenn stage, they are correlated in our rat model. Our

immunohistochemical and immunofluorescent evaluation in the Glenn rat model shows

dramatic changes of both proximal and distal pulmonary arterial mural cell recruitment

Currently, descriptions of dilated distal vessels in the Glenn population exist (78-80);

however, there has been no histological analysis of proximal pulmonary arteries. Work in

Fontan patients, however, has shown thinning of the proximal pulmonary arterial media

(81, 82). The persistence of thin media (proximally, where collaterals form) even as distal

PAVMs resolve in the Fontan stage suggests a critical role for pulsatility in maintenance of

proximal vascular architecture. We suggest that pulsatility loss is therefore likely a

common pathogenic factor in the development of both proximal collateral vessels and

distal AVMs.

In summary, we show that the pulsatility of hemodynamic forces in blood vessels is

sensed by the endothelium uniquely in each dimension of force. Our data support a model

whereby pulsatile stretch induces endothelium to secrete PDGFB, signaling for support

from vascular smooth muscle. Indeed, the architecture and cellular composition of the

vessel wall change significantly upon loss of pulsatility in the Glenn, with thinning of the

vascular wall, thereby shifting the arterial vascular wall structure toward that of a vein,

likely impacting downstream endothelium. Together, this work underscores the critical

importance of proper blood flow dynamics for the stability of vascular structure and

function. In addition, it points to biomechanical signaling pathways as potential novel

targets for therapeutic intervention in patients with pulsatility loss, such as in the Glenn

circulation. We seek to better understand and address the mechanistic underpinnings of

vascular defects that occur during single ventricle palliation and anticipate that our data on

pulsatility will provide the foundation for improving clinical outcomes for children that

undergo these necessary surgeries.

Methods

Data Availability

All sequencing data that support the findings of this study have been deposited in the National Center for Biotechnology Information Gene Expression Omnibus (GEO) and are accessible through the GEO Series accession number GSE298790. All other relevant data are available from the corresponding author on request.

Combined cardiac catheterization and cardiac MRI

Combined cardiac catheterization and MRI was performed on a Philips XMR setup (Phillips Ingenia Evolution, Best, Netherlands).(83) This consists of a 1.5T MRI-scanner and a BV Pulsera or Allura Clarity or Ingenia (Philips) cardiac X-Ray unit. An appropriate size balloon wedge catheter (Arrow Intl., Reading, PA) and receiver coil were used depending on the weight of the patient.

All studies were conducted under general anesthesia based on clinical need. Patients were recruited and data was recorded in sequence of the patient's referral for the procedure under University of Texas Southwestern IRB-approved study (STU 032016-009) in 2023 - 2024. All Glenn patients with a single functional ventricle were considered for inclusion in the study. Patients were excluded if they had pulmonary blood flow in addition to the flow provided by the Glenn anastomosis. Both male and female patients were included in the study (see Supplemental Table 1).

Pulsatility was assessed by pulse difference, which was defined as the difference between

maximum and minimum value for each parameter of force. Pulsatility index was not

selected due to the potential for obscuring differences between patients (see Supplemental

Table 2). For each 2D cross-section of the right pulmonary artery obtained, we obtained a

mean velocity (averaged over the cross section of the vessel). This data is obtained over 1

to 2 minutes 'free-breathing' using phase contrast velocity-encoded cine MRI. Sequence

parameters included 40 cardiac phases, TE/TR = 2.7/4.4 ms (TE = echo time, TR =

repetition time), with two signal averages, 1.5-2mm × 1.5-2mm × 6-8 mm resolution,

SENSE acceleration factor = 2, with the velocity encoding gradient set to 25% above the

expected maximum velocity in each vessel. Vendor-provided background phase correction

was used. Artificial intelligence-assisted segmentation using Circle (Circle Cardiovascular

Imaging, CVI42, v6.1.2) was performed to obtain the maximum and minimum area, from

which an idealized circumference was calculated. Pressure tracings were obtained on a

standard Siemens Sensis Hemodynamic recording system (Siemens Healthineers,

Munich, Germany), using a 5- or 6-French Arrow, Balloon Wedge-Pressure Catheter

(Teleflex Medical Headquarter International, Ireland) located in the proximal right branch

pulmonary artery.

In vitro cell-based assays of pulsatile force

Pulsatile force was applied to confluent monolayers of human pulmonary artery endothelial

cells (HPAECs, Lonza) for 48 hours under each condition. Pulsatile (1 Hz, 15 dyn/cm^2) and

continuous (15 dyn/cm2) shear was applied using the Ibidi system (Ibidi USA, Wisconsin).

Pulsatile (25-5 mm Hg) and continuous (25 mm Hg) pressure was applied using a custom

modification to the Ibidi system whereby cells were alternately (1 Hz) exposed to columns

of culture media to deliver the desired hydrostatic pressure. Pressure readings were tested

and confirmed the using the Siemens Sensis Hemodynamic recording system listed

above. Pulsatile stretch was applied using Uniflex plates (1Hz, 1 hr of 3% stretch for

acclimation followed by 47 hrs of 10% stretch) as a part of the Flexcell FX-6000 system

(Flexcell International, Burlington, NC). Ibidi channels were pre-coated with the proprietary

Ibi-treat to promote cell adhesion, and Flexcell uniflex plates were coated with 0.1% gelatin

to promote cell adhesion.

All experiments utilized at least three biological replicates, using separate aliquots of

HPAECs from two different donors. RNA was extracted using RNeasy Plus kit (Qiagen)

prior to bulk RNA sequencing. Library generation and bulk RNA sequencing was carried

out by the McDermott Next Generation Sequencing core facility at UT Southwestern

Medical Center, using an Illumina NextSeq 2000. Processed sequencing data (using

iGenomes annotations) was provided by the core, and a TPM cutoff of 10 was applied (if

any test condition resulted in a gene's TPM >10, the gene was included in our analysis).

PCA plots of each condition are shown in **Supplementary Figure 4**.

For pathway analysis, we performed gene set enrichment analysis was performed for sets

curated in MsigDb (PMID 26771021) using with fgsea (BioRxiv

https://doi.org/10.1101/060012) Bioconductor package. To simultaneously account for both

the magnitude and significance of the measured effects, genes were ranked by the

quotient of log fold-change and adjusted P-value.

Human Lung Single Cell RNA sequencing

The LungMAP Human Lung CellRef atlas was utilized via the online portal at

www.lungmap.net. (38, 39) This dataset includes 148 normal human lung samples from

104 donors, including child, adolescent, and adult tissue. Tissue types include

parenchyma, trachea, bronchi, bronchus, and small airways. 347,970 cells from the

dataset were included in our analysis. Both ShinyCell and cell x gene were utilized for

visualization of cell-specific gene expression. Dataset ID: LMEX0000004396.

Measurement of PDGFB secretion in vitro

Cultured HPAECs, human arterial endothelial cells (HAECs, Lonza), human pulmonary

artery smooth muscle cells (PASMCs, Lonza), and normal human lung fibroblasts (NHLFs,

Lonza were plated onto Flexcell uniflex plates (pre-coated with collagen I). Each cell line

was grown in the recommended cell culture media, obtained from Lonza: HPAECs and

HAECs were grown in EGM-2, PASMCs in SmGM-2, and NHLFs in FGM-2. Cells were

grown to confluency then subjected to pulsatile stretch for 48 hours as above. Media was

collected and PDGF-BB was measured by ELISA (R+D systems DY220). For comparison

to endothelial cells under flow, media was obtained from confluent HPAECs in 6 well

plates under continuous flow at 15 dyn/cm^2 shear stress in a rotational flow apparatus

(Thermo Scientific Model No: 88881101).

For measurement of transpulmonary gradient of PDGF-BB (here termed PDGFB) in

human plasma, samples were obtained directly from Glenn and heart transplant patients in

the course of regularly indicated cardiac catheterization. Patients were consented for the

IRB approved study (University of Texas Southwestern IRB: UTSW-2020-0047). Samples

were obtained from the pre-capillary blood (right pulmonary artery) and the post-capillary

blood (right pulmonary wedge position, with the catheter in the distal pulmonary artery with

balloon inflated and the sample drawn through the end hole) in each patient. Nature of

each sample as pre- or post-capillary was then confirmed by prompt measurement of

oxygen saturation, and only fully oxygenated samples were used for the analysis. All

samples were obtained in K2-EDTA purple-top tubes, placed on ice, and promptly spun

down for plasma isolation. Samples were stored at -80°C up to 2 years maximum prior to

measurement by ELISA as above.

Surgical Rat Model of Glenn Circulation

The rat surgical model of Glenn circulation is described in detail previously.(36) Briefly, we

performed an end-to-end cavopulmonary anastomosis between the left superior vena cava

(L-SVC) and the left pulmonary artery (LPA). This was previously shown to result in

progressive impairment (beginning by two weeks) in oxygenation in tandem with the

development of intrapulmonary arteriovenous shunts as detected by both bubble

echocardiography and microsphere injection.

Adult Sprague–Dawley rats (male and female, 6–12 weeks of age) were used for all

experiments (Taconic). All rats were housed in the Biomedical Research Center at the

Medical College of Wisconsin with access to standard chow diet (PicoLab Lab Diet, 5L0D),

water ad libitum, and maintained in a 12-hour light/dark cycle. All surgical procedures were

performed under isoflurane anesthesia (1%–3%). All experimental protocols were

approved by the Medical College of Wisconsin Institutional Animal Care and Use

Committee prior to initiation of experimental protocols (Animal Use Agreement #7731).

Tissue Preparation and Immunohistochemistry (IHC) /Immunofluorescence (IF)

Lungs were harvested at 2 and 6 months after surgery for these experiments. Lungs were inflated to 10 mmHg for 10 minutes with 10% neutral buffered formalin to promote normal alveolar architecture in sections, followed by overnight fixation in 10% neutral buffered formalin. Formalin-fixed, paraffin-embedded sections were prepared using the left lung of the Glenn and sham operated animals. IHC staining was done for SMA (Abcam, ab7817) using the Leica Bond Rx automated staining platform with Leica BOND Polymer Refine Detection kit (Leica, DS9800) per manufacturer instructions, using standard labeled-streptavidin-biotin detection with secondary antibody (Jackson Immuno Research Labs, Biotinylated donkey anti-mouse 1:500, #715-066-151), streptavidin HRP (Vector Laboratories, #SA-5004), and DAB chromogen (BioCare, BDB2004).

Slides were scanned using Hamamatsu HT whole slide scanner (Hamamatsu, USA) and analyzed with QuPath (Edinburgh, V0.5.1). The file names were randomized and blinded prior to analysis. Due to the possibility of changes in vessel size between sham and Glenn-operated rates, measurements were taken from the most proximal pulmonary arteries and veins to ensure a valid comparison between samples. Arteries were differentiated from veins in consultation with a trained pathologist, by their classic histological characteristics and anatomical proximity to the mainstem bronchi. Slides were not utilized for measurement if no major bronchi or large vessels could be identified. Measurements were taken only in areas where the thickness was uniform and the endothelial cells formed a clear monolayer, to avoid artifactual error from the angle of the slice. 10 measurements of each large vessel (artery or vein, Glenn or sham-operated) were obtained, and the mean measurement value was used for analysis.

We used IF staining and confocal microscopy to identify distal vessels for

quantification of endothelial PDGFB protein and perivascular smooth muscle. Lung

sections were prepared as above. Sections were permeabilized for 10 min in 0.2% Triton

X-100 in PBS, then subjected to antigen retrieval with R-buffer B in a 2100 Retriever

(Electron Microscopy Sciences). Slides were blocked with CAS-block (Invitrogen), and

primary antibody incubations were carried out at 4°C overnight. Primary antibodies were

as follows: VE-cadherin/CDH5 (R+D systems, AF1002, 1:100), PDGFB (Santa Cruz, sc-

365805, 1:100), and SMA-APC (Abcam, ab225143, 1:400). Imaging was performed on a

Nikon CSU-W1 dual-camera inverted spinning-disc confocal microscope. Vessels were

chosen as <50 µm and >20 µm in diameter, with a clear lumen and apical VE-cadherin

staining visible. Images were analyzed in Fiji as follows: the percentage of the

circumference of each vessel covered by SMA staining was calculated, and a mask was

made of the pixels positive for SMA staining in the sub-endothelial region and the mean

pixel intensity of those pixels was calculated.

Statistical Analysis

Sample sizes were based on patient availability and standards in the field. Data were both

analyzed and plotted in Graphpad PRISM 10.2.3. Descriptive statistics (mean, standard

deviation, median, min, and max) were employed for summarizing demographic,

hemodynamic, and clinical variables. Statistical tests (indicated in the relevant methods

sections and figure legends) utilized were the parametric unpaired Student's *t*-test and the

one-way ANOVA with Tukey's multiple comparisons test. Statistical significance is noted

as follows: ns = $p > 0.05$; * = $0.01 < p < 0.05$; ** = $0.001 < p < 0.01$; *** = $0.0001 < p < 0.001$;

**** = $p < 0.0001$. Superplots were made with individual colors representing biological

replicates (individual rats), and the mean of the averages of each biological replicate is

shown (84). Cell images were analyzed in FIJI. Figures and models were made using

Microsoft PowerPoint, data were partially analyzed in Microsoft Excel, and text was written

in Microsoft Word.

<u>ACKNOWLEDGEMENTS</u>

We are grateful to the entire Cleaver Lab for useful discussions and reading of the

manuscript. We thank Denise Marciano for the use of her lab's Flexcell apparatus. We

thank the UT Southwestern Medical Center (UTSW) Quantitative Light Microscopy Core

Facility (Marcel Mettlen, Ph.D.) for training/assistance. The results here are in part based

upon data generated by the LungMAP Consortium and downloaded from

(www.lungmap.net), on March 15, 2025. The LungMAP consortium, the Human Tissue

Core (U01-HL144861), and the LungMAP Data Coordinating Center (U24-HL148865) are

funded by the National Heart, Lung, and Blood Institute (NHLBI).

Author contributions

Conceptualization: S.S., T.H., M.F., M.L.I.A., O.C.; Methodology: S.S., T.H., M.F., S.R.,

A.D.S., O.C.; Investigation: S.S., L.T., T.W., S.R., O.C.; Resources: S.S., T.W., A.D.S.,

T.C., O.C.; Data curation: S.S., M.A.C., L.T., T.H.; Writing - original draft: S.S., O.C.;

Writing - review & editing: S.S., T.H., M.L.I.A., A.D.S., O.C.; Visualization: M.A.C., S.S.;

Supervision: S.S., O.C.; Project administration: S.S., O.C.; Funding acquisition: S.S., T.C.,

A.D.S., O.C.

Funding

This research was supported in part by grants from the American Thoracic

Society/Alveolar Capillary Dysplasia (ACDMPV) Research Grant (23-24PACDA12 to S.S.);

National Heart Lung and Blood Institute (1R01HL-175575 to M.L.I.A., K08HL157510 to

ADS, HL113498 and HL126518 to O.C.); the National Academy of Science, Engineering,

and Medicine Ford Foundation Dissertation Fellowship (M.A.C.); National Institutes of

Health (R01DK127634, RC2 DK125960 to T.C.) and Cancer Prevention Research Institute

of Texas (RP220201 to T.C.); and the Foundation Leducq grant (21CVD03 to M.L.I.A.,

O.C.). Open Access funding provided by University of Texas Southwestern Medical

Center. Deposited in PMC for immediate release.

REFERENCES

1. Liu Y, Chen S, Zuhlke L, Black GC, Choy MK, Li N, et al. Global birth prevalence of congenital heart defects 1970-2017: updated systematic review and meta-analysis of 260 studies. *Int J Epidemiol.* 2019;48(2):455-63.

2. Rychik J, Atz AM, Celermajer DS, Deal BJ, Gatzoulis MA, Gewillig MH, et al. Evaluation and Management of the Child and Adult With Fontan Circulation: A Scientific Statement From the American Heart Association. *Circulation.* 2019:CIR0000000000000696.

3. Srivastava D, Preminger T, Lock JE, Mandell V, Keane JF, Mayer JE, Jr., et al. Hepatic venous blood and the development of pulmonary arteriovenous malformations in congenital heart disease. *Circulation.* 1995;92(5):1217-22.

4. Pike NA, Vricella LA, Feinstein JA, Black MD, and Reitz BA. Regression of severe pulmonary arteriovenous malformations after Fontan revision and "hepatic factor" rerouting. *Ann Thorac Surg.* 2004;78(2):697-9.

5. McElhinney DB, Kreutzer J, Lang P, Mayer JE, Jr., del Nido PJ, and Lock JE. Incorporation of the hepatic veins into the cavopulmonary circulation in patients with heterotaxy and pulmonary arteriovenous malformations after a Kawashima procedure. *Ann Thorac Surg.* 2005;80(5):1597-603.

6. Kavarana MN, Jones JA, Stroud RE, Bradley SM, Ikonomidis JS, and Mukherjee R. Pulmonary arteriovenous malformations after the superior cavopulmonary shunt: mechanisms and clinical implications. *Expert Rev Cardiovasc Ther.* 2014;12(6):703-13.

7. Spearman AD, and Ginde S. Pulmonary Vascular Sequelae of Palliated Single Ventricle Circulation: Arteriovenous Malformations and Aortopulmonary Collaterals. *Journal of Cardiovascular Development and Disease.* 2022;9(9):309.

8. Shah MJ, Rychik J, Fogel MA, Murphy JD, and Jacobs ML. Pulmonary AV Malformations After Superior Cavopulmonary Connection: Resolution After Inclusion of Hepatic Veins in the Pulmonary Circulation. *The Annals of Thoracic Surgery.* 1997;63(4):960-3.

9. Kurotobi S, Sano T, Kogaki S, Matsushita T, Miwatani T, Takeuchi M, et al. Bidirectional cavopulmonary shunt with right ventricular outflow patency: The impact of pulsatility on pulmonary endothelial function. *The Journal of thoracic and cardiovascular surgery.* 2001;121(6):1161-8.

10. Henaine R, Vergnat M, Mercier O, Serraf A, De Montpreville V, Ninet J, et al. Hemodynamics and arteriovenous malformations in cavopulmonary anastomosis: the case for residual antegrade pulsatile flow. *The Journal of thoracic and cardiovascular surgery.* 2013;146(6):1359-65.

11. Sakaki M, Taenaka Y, Tatsumi E, Nakatani T, Kinoshita M, Akagi H, et al. Pulmonary function in a non-pulsatile pulmonary circulation. *ASAIO J.* 1992;38(3):M366-9.

12. Townsley MI. Structure and composition of pulmonary arteries, capillaries, and veins. *Compr Physiol.* 2012;2(1):675-709.

13. Wetzel-Strong SE, Detter MR, and Marchuk DA. The pathobiology of vascular malformations: insights from human and model organism genetics. *J Pathol.* 2017;241(2):281-93.

14. Chen W, Guo Y, Walker EJ, Shen F, Jun K, Oh SP, et al. Reduced mural cell coverage and impaired vessel integrity after angiogenic stimulation in the Alk1-deficient brain. *Arterioscler Thromb Vasc Biol.* 2013;33(2):305-10.

15. Brahmbhatt A, Remuzzi A, Franzoni M, and Misra S. The molecular mechanisms of hemodialysis vascular access failure. *Kidney Int.* 2016;89(2):303-16.

880   16.   Li Y, He Y, Falzon I, Fairbourn B, Tingey S, Imrey PB, et al. Dynamic Remodeling of
881   Human Arteriovenous Fistula Wall Obtained From Magnetic Resonance Imaging
882   During the First 6 Months After Creation. *Kidney Int Rep.* 2022;7(8):1905-9.

883   17.   Corada M, Morini MF, and Dejana E. Signaling pathways in the specification of arteries
884   and veins. *Arterioscler Thromb Vasc Biol.* 2014;34(11):2372-7.

885   18.   Peng T, Tian Y, Boogerd CJ, Lu MM, Kadzik RS, Stewart KM, et al. Coordination of heart
886   and lung co-development by a multipotent cardiopulmonary progenitor. *Nature.*
887   2013;500(7464):589-92.

888   19.   Schwarz MA, and Cleaver OB. *The Pulmonary Endothelium*. 2009:1-24.

889   20.   Buschmann I, Pries A, Styp-Rekowska B, Hillmeister P, Loufrani L, Henrion D, et al.
890   Pulsatile shear and Gja5 modulate arterial identity and remodeling events during flow-
891   driven arteriogenesis. *Development.* 2010;137(13):2187-96.

892   21.   Dessalles CA, Leclech C, Castagnino A, and Barakat AI. Integration of substrate- and
893   flow-derived stresses in endothelial cell mechanobiology. *Commun Biol.* 2021;4(1):764.

894   22.   McEvoy E, Sneh T, Moeendarbary E, Javanmardi Y, Efimova N, Yang C, et al. Feedback
895   between mechanosensitive signaling and active forces governs endothelial junction
896   integrity. *Nat Commun.* 2022;13(1):7089.

897   23.   Kutys ML, and Chen CS. Forces and mechanotransduction in 3D vascular biology. *Curr*
898   *Opin Cell Biol.* 2016;42:73-9.

899   24.   Baeyens N, Bandyopadhyay C, Coon BG, Yun S, and Schwartz MA. Endothelial fluid
900   shear stress sensing in vascular health and disease. *J Clin Invest.* 2016;126(3):821-8.

901   25.   Jufri NF, Mohamedali A, Avolio A, and Baker MS. Mechanical stretch: physiological and
902   pathological implications for human vascular endothelial cells. *Vasc Cell.* 2015;7:8.

903   26.   Mierke CT. Mechanosensory entities and functionality of endothelial cells. *Front Cell*
904   *Dev Biol.* 2024;12:1446452.

905   27.   Zhao S, Suciu A, Ziegler T, Moore JE, Jr., Burki E, Meister JJ, et al. Synergistic effects of
906   fluid shear stress and cyclic circumferential stretch on vascular endothelial cell
907   morphology and cytoskeleton. *Arterioscler Thromb Vasc Biol.* 1995;15(10):1781-6.

908   28.   Russo TA, Banuth AMM, Nader HB, and Dreyfuss JL. Altered shear stress on endothelial
909   cells leads to remodeling of extracellular matrix and induction of angiogenesis. *PLoS*
910   *One.* 2020;15(11):e0241040.

911   29.   Wasserman SM, and Topper JN. Adaptation of the endothelium to fluid flow: in vitro
912   analyses of gene expression and in vivo implications. *Vasc Med.* 2004;9(1):35-45.

913   30.   Wang Y, Cao W, Cui J, Yu Y, Zhao Y, Shi J, et al. Arterial Wall Stress Induces Phenotypic
914   Switching of Arterial Smooth Muscle Cells in Vascular Remodeling by Activating the
915   YAP/TAZ Signaling Pathway. *Cell Physiol Biochem.* 2018;51(2):842-53.

916   31.   Lindahl P, Johansson BR, Leveen P, and Betsholtz C. Pericyte loss and microaneurysm
917   formation in PDGF-B-deficient mice. *Science.* 1997;277(5323):242-5.

918   32.   Hellstrom M, Kalen M, Lindahl P, Abramsson A, and Betsholtz C. Role of PDGF-B and
919   PDGFR-beta in recruitment of vascular smooth muscle cells and pericytes during
920   embryonic blood vessel formation in the mouse. *Development.* 1999;126(14):3047-55.

921   33.   Nadeem T, Bogue W, Bigit B, and Cuervo H. Deficiency of Notch signaling in pericytes
922   results in arteriovenous malformations. *JCI Insight.* 2020;5(21).

923   34.   Lin Y, Gahn J, Banerjee K, Dobreva G, Singhal M, Dubrac A, et al. Role of endothelial
924   PDGFB in arterio-venous malformations pathogenesis. *Angiogenesis.* 2024;27(2):193-
925   209.

35.   Uebelhoer M, Natynki M, Kangas J, Mendola A, Nguyen HL, Soblet J, et al. Venous
malformation-causative TIE2 mutations mediate an AKT-dependent decrease in
PDGFB. *Hum Mol Genet.* 2013;22(17):3438-48.
36.   Wan TC, Rousseau H, Mattern C, Tabor M, Hodges MR, Ramchandran R, et al. Glenn
circulation causes early and progressive shunting in a surgical model of pulmonary
arteriovenous malformations. *Physiol Rep.* 2024;12(22):e70123.
37.   Deng H, Eichmann A, and Schwartz MA. Fluid Shear Stress-Regulated Vascular
Remodeling: Past, Present, and Future. *Arterioscler Thromb Vasc Biol.* 2025.
38.   Gaddis N, Fortriede J, Guo M, Bardes EE, Kouril M, Tabar S, et al. LungMAP Portal
Ecosystem: Systems-level Exploration of the Lung. *Am J Respir Cell Mol Biol.*
2024;70(2):129-39.
39.   Guo M, Morley MP, Jiang C, Wu Y, Li G, Du Y, et al. Guided construction of single cell
reference for human and mouse lung. *Nat Commun.* 2023;14(1):4566.
40.   Resnick N, Collins T, Atkinson W, Bonthron DT, Dewey CF, Jr., and Gimbrone MA, Jr.
Platelet-derived growth factor B chain promoter contains a cis-acting fluid shear-
stress-responsive element. *Proc Natl Acad Sci U S A.* 1993;90(10):4591-5.
41.   Zerwes HG, and Risau W. Polarized secretion of a platelet-derived growth factor-like
chemotactic factor by endothelial cells in vitro. *J Cell Biol.* 1987;105(5):2037-41.
42.   Yuliana ME, Huang ZH, Chou HC, and Chen CM. Effects of uteroplacental insufficiency
on growth-restricted rats with altered lung development: A metabolomic analysis.
Front Pediatr. 2022;10:952313.
43.   McFaul RC, Tajik AJ, Mair DD, Danielson GK, and Seward JB. Development of pulmonary
arteriovenous shunt after superior vena cava-right pulmonary artery (Glenn)
anastomosis. Report of four cases. *Circulation.* 1977;55(1):212-6.
44.   Cloutier A, Ash JM, Smallhorn JF, Williams WG, Trusler GA, Rowe RD, et al. Abnormal
distribution of pulmonary blood flow after the Glenn shunt or Fontan procedure: risk
of development of arteriovenous fistulae. *Circulation.* 1985;72(3):471-9.
45.   Gray RG, Altmann K, Mosca RS, Prakash A, Williams IA, Quaegebeur JM, et al. Persistent
antegrade pulmonary blood flow post-glenn does not alter early post-Fontan outcomes
in single-ventricle patients. *Ann Thorac Surg.* 2007;84(3):888-93; discussion 93.
46.   Chen Q, Tulloh R, Caputo M, Stoica S, Kia M, and Parry AJ. Does the persistence of
pulsatile antegrade pulmonary blood flow following bidirectional Glenn procedure
affect long term outcome? *Eur J Cardiothorac Surg.* 2015;47(1):154-8; discussion 8.
47.   Kowatari R, Suzuki Y, Daitoku K, and Fukuda I. Long-term results of additional
pulmonary blood flow with bidirectional cavopulmonary shunt. *J Cardiothorac Surg.*
2020;15(1):279.
48.   Berdat PA, Belli E, Lacour-Gayet F, Planche C, and Serraf A. Additional pulmonary blood
flow has no adverse effect on outcome after bidirectional cavopulmonary anastomosis.
Ann Thorac Surg. 2005;79(1):29-36; discussion -7.
49.   van de Wal HJCM, Ouknine R, Tamisier D, Lévy M, Vouhé PR, and Leca F. Bi-directional
cavopulmonary shunt: is accessory pulsatile flow, good or bad? *European Journal of*
Cardio-Thoracic Surgery. 1999;16(2):104-10.
50.   Ferns SJ, El Zein C, Multani K, Sajan I, Subramanian S, Polimenakos AC, et al. Is
additional pulsatile pulmonary blood flow beneficial to patients with bidirectional
Glenn? *The Journal of thoracic and cardiovascular surgery.* 2013;145(2):451-4.

51.   Uemura H, Yagihara T, Kawashima Y, Okada K, Kamiya T, and Anderson RH. Use of the
bidirectional Glenn procedure in the presence of forward flow from the ventricles to
the pulmonary arteries. *Circulation.* 1995;92(9 Suppl):II228-32.
52.   Januszewska K, Nawrocki P, Bruenen A, Schnabel A, and Malec E. Significance of
additional pulmonary blood flow between second and third stage in Fontan pathway.
BMC pediatrics. 2024;24(1):705.
53.   Lammers S, Scott D, Hunter K, Tan W, Shandas R, and Stenmark KR. *Comprehensive*
Physiology. 2012:295-319.
54.   Sumpio BE, Du W, Galagher G, Wang X, Khachigian LM, Collins T, et al. Regulation of
PDGF-B in endothelial cells exposed to cyclic strain. *Arterioscler Thromb Vasc Biol.*
1998;18(3):349-55.
55.   Kolpakov V, Rekhter MD, Gordon D, Wang WH, and Kulik TJ. Effect of mechanical forces
on growth and matrix protein synthesis in the in vitro pulmonary artery. Analysis of
the role of individual cell types. *Circ Res.* 1995;77(4):823-31.
56.   Lin C, Zheng X, Lin S, Zhang Y, Wu J, and Li Y. Mechanotransduction Regulates the
Interplays Between Alveolar Epithelial and Vascular Endothelial Cells in Lung. *Front*
Physiol. 2022;13:818394.
57.   R RF, Fukui H, Chow R, Vilfan A, and Vermot J. The cilium as a force sensor-myth versus
reality. *J Cell Sci.* 2019;132(14).
58.   Spurgin S, Nguimtsop AM, Chaudhry FN, Michki SN, Salvador J, Iruela-Arispe ML, et al.
Dynamics of primary cilia in endothelial and mesenchymal cells throughout mouse
lung development. *Dev Dyn.* 2025;254(11):1187-202.
59.   Tanaka K, Chen M, Prendergast A, Zhuang Z, Nasiri A, Joshi D, et al. Latrophilin-2
mediates fluid shear stress mechanotransduction at endothelial junctions. *EMBO J.*
2024;43(15):3175-91.
60.   Van Gieson EJ, Murfee WL, Skalak TC, and Price RJ. Enhanced Smooth Muscle Cell
Coverage of Microvessels Exposed to Increased Hemodynamic Stresses In Vivo.
Circulation Research. 2003;92(8):929-36.
61.   Singh E, Redgrave RE, Phillips HM, and Arthur HM. Arterial endoglin does not protect
against arteriovenous malformations. *Angiogenesis.* 2020;23(4):559-66.
62.   Lee HW, Xu Y, He L, Choi W, Gonzalez D, Jin SW, et al. Role of Venous Endothelial Cells
in Developmental and Pathologic Angiogenesis. *Circulation.* 2021;144(16):1308-22.
63.   Wang JH, Goldschmidt-Clermont P, Wille J, and Yin FC. Specificity of endothelial cell
reorientation in response to cyclic mechanical stretching. *J Biomech.*
2001;34(12):1563-72.
64.   Takeda R, Yamaguchi T, Hayashi S, Sano S, Kawame H, Kanki S, et al. Clinical and
molecular features of patients with COL1-related disorders: Implications for the wider
spectrum and the risk of vascular complications. *Am J Med Genet A.* 2022;188(9):2560-
75.
65.   Lesnik P, Haskell CA, and Charo IF. Decreased atherosclerosis in CX3CR1–/– mice
reveals a role for fractalkine in atherogenesis. *Journal of Clinical Investigation.*
2003;111(3):333-40.
66.   Bazan JF, Bacon KB, Hardiman G, Wang W, Soo K, Rossi D, et al. A new class of
membrane-bound chemokine with a CX3C motif. *Nature.* 1997;385(6617):640-4.
67.   Chen Y, Zhu W, Bollen AW, Lawton MT, Barbaro NM, Dowd CF, et al. Evidence of
Inflammatory Cell Involvement in Brain Arteriovenous Malformations. *Neurosurgery.*
2008;62(6):1340-50.

68.   Zhang R, Han Z, Degos V, Shen F, Choi EJ, Sun Z, et al. Persistent infiltration and pro-
1019   inflammatory differentiation of monocytes cause unresolved inflammation in brain
1020   arteriovenous malformation. *Angiogenesis.* 2016;19(4):451-61.
1021   69.   Braverman IM, Keh A, and Jacobson BS. Ultrastructure and three-dimensional
1022   organization of the telangiectases of hereditary hemorrhagic telangiectasia. *J Invest*
1023   *Dermatol.* 1990;95(4):422-7.
1024   70.   Kim SJ, Bae EJ, Cho DJ, Park IS, Kim YM, Kim W-H, et al. Development of pulmonary
1025   arteriovenous fistulas after bidirectional cavopulmonary shunt. *The Annals of Thoracic*
1026   *Surgery.* 2000;70(6):1918-22.
1027   71.   Vettukattil JJ, Slavik Z, Lamb RK, Monro JL, Keeton BR, Tsang VT, et al. Intrapulmonary
1028   arteriovenous shunting may be a universal phenomenon in patients with the superior
1029   cavopulmonary anastomosis: a radionuclide study. *Heart (British Cardiac Society).*
1030   2000;83(4):425-8.
1031   72.   Feinstein JA, Moore P, Rosenthal DN, Puchalski M, and Brook MM. Comparison of
1032   contrast echocardiography versus cardiac catheterization for detection of pulmonary
1033   arteriovenous malformations. *The American journal of cardiology.* 2002;89(3):281-5.
1034   73.   Stoller JKH, Robert M.
1035   White, Richard D., and Mee RB. Anomalous Hepatic Venous Drainage into the Left Atrium: An
1036   Unusual Cause of Hypoxemia. *Respiratory Care.* 2003;48(1):58-62.
1037   74.   Guardado FJ, Byrd TM, and Petersen WG. Azygous continuation of the inferior vena
1038   cava with anomalous hepatic vein drainage. *Am J Med Sci.* 2012;343(3):259-61.
1039   75.   Lee J, Menkis AH, and Rosenberg HC. Reversal of pulmonary arteriovenous
1040   malformation after diversion of anomalous hepatic drainage. *Ann Thorac Surg.*
1041   1998;65(3):848-9.
1042   76.   Kloppenburg GT, Post MC, Mager HJ, and Schepens MA. Rerouting anomalous hepatic
1043   venous connection to the left atrium. *Ann Thorac Surg.* 2010;90(2):638-40.
1044   77.   Yim D, Nagata H, Lam CZ, Grosse-Wortmann L, Seed M, Jaeggi E, et al. Disharmonious
1045   Patterns of Heterotaxy and Isomerism: How Often Are the Classic Patterns Breached?
1046   *Circ Cardiovasc Imaging.* 2018;11(2):e006917.
1047   78.   Duncan BW, Kneebone JM, Chi EY, Hraska V, Isik FF, Rosenthal GL, et al. A detailed
1048   histologic analysis of pulmonary arteriovenous malformations in children with
1049   cyanotic congenital heart disease. *The Journal of thoracic and cardiovascular surgery.*
1050   1999;117(5):931-8.
1051   79.   Starnes SL, Duncan BW, Kneebone JM, Fraga CH, States S, Rosenthal GL, et al.
1052   Pulmonary microvessel density is a marker of angiogenesis in children after
1053   cavopulmonary anastomosis. *The Journal of thoracic and cardiovascular surgery.*
1054   2000;120(5):902-8.
1055   80.   Starnes SL, Duncan BW, Fraga CH, Desai SY, Jones TK, Mathur SK, et al. Rat model of
1056   pulmonary arteriovenous malformations after right superior cavopulmonary
1057   anastomosis. *Am J Physiol Heart Circ Physiol.* 2002;283(5):H2151-6.
1058   81.   Hays BS, Baker M, Laib A, Tan W, Udholm S, Goldstein BH, et al. Histopathological
1059   abnormalities in the central arteries and veins of Fontan subjects. *Heart (British*
1060   *Cardiac Society).* 2018;104(4):324-31.
1061   82.   Ridderbos FJ, Wolff D, Timmer A, van Melle JP, Ebels T, Dickinson MG, et al. Adverse
1062   pulmonary vascular remodeling in the Fontan circulation. *J Heart Lung Transplant.*
1063   2015;34(3):404-13.

1064   83.   Velasco Forte MN, Roujol S, Ruijsink B, Valverde I, Duong P, Byrne N, et al. MRI for
1065   Guided Right and Left Heart Cardiac Catheterization: A Prospective Study in Congenital
1066   Heart Disease. *J Magn Reson Imaging.* 2021;53(5):1446-57.
1067   84.   Lord SJ, Velle KB, Mullins RD, and Fritz-Laylin LK. SuperPlots: Communicating
1068   reproducibility and variability in cell biology. *J Cell Biol.* 2020;219(6).
1069
1070

Figures

Figure 1. Loss of pulsatility in three dimensions within Glenn pulmonary arteries.

(A, A') Diagrams of representative normal and Glenn cardiopulmonary anatomy. Red

box highlights location of force pulsatility measurements. **(B)** Hemodynamic forces act

on ECs in three dimensions: shear/flow (blue), pressure (orange), and circumferential

stretch (green). (Schematic adapted from graphic on Ibidi.com). **(C)** Outline of data

collection method for each dimension of hemodynamic force. **(D, D')** Example pressure

waveforms from the right pulmonary artery of normal and Glenn patients.

Electrocardiogram shown in red for correlation to cardiac cycle. RPA, right pulmonary

artery. **(E, E')** Variation in flow velocity of the proximal right pulmonary artery measured

by cardiac MRI throughout a cardiac cycle from systole (start) to end-diastole (end).

Variability in length of time for each patient reflects the variable heart rate within each

subject. **(F, F')** Cardiac MRI velocity encoded phase contrast images with color showing

velocity of blood flow out of the plane of the image slice. Dotted orange line shows RPA

border. RPA diameter increases during systole in anormal patients but does not

increase during systole in Glenn patient. **(G)** Quantification of pulse difference of each

dimension of force, obtained during combined cardiac catheterization and cardiac MRI

of normal (n=20) and Glenn (n=20) patients, demonstrating loss of pulsatility in the

Glenn. Circumference % change is calculated from baseline in diastole.

748
749

A Applying pulsatility to ECs in isolated dimensions of force

HPAEC → Non-pulsatile → Bulk RNAseq
48 hours
HPAEC → Pulsatile → Bulk RNAseq

	Non-Pulsatile	Pulsatile
Flow	15 dyn/cm^2	0-15 dyn/cm^2 1 Hz
Pressure	25 mm Hg	5-25 mm Hg 1 Hz
Stretch	0%	10% 1 Hz

B

Flow (LSS)	Pressure	Stretch

Non-Pulsatile / Pulsatile

DAPI CDH5

C Regulated Pathways: Pulsatile [Force] vs Non-pulsatile [Force]

Stretch 10 — 9 — Flow 19
2 — 10 — 6
Pressure 4

D

E Contributions to KEGG pathway: regulation of actin cytoskeleton

Pulsatile Force	Selected Contributors
Flow (LSS)	*Rac2, Map2k1, Myl12a, Pdgfb*
Pressure	*Itga6, Rac3, Cdc42, Actn1*
Stretch	*Actn4, Itga3, Arghef6, Pdgfb*

Figure 2. **Pulsatility drives both similar and distinct signaling pathways within**

each dimension of force. (A) Experimental conditions: HPAECs were exposed to 48

hours of a single dimension of either non-pulsatile or pulsatile force, and bulk RNA

sequencing was performed. **(B)** GSEA of bulk RNAseq data shows overlapping and

unique pathways activated by each dimension of force. **(C)** Representative images of

the cell-shape adjustments that occur in response to the different forces are shown. **(D)**

Normalized enrichment score (NES) and adjusted p-value (adj p-val) shown for select

signaling pathways shows greater impact of pulsatility within flow and stretch relative to

pulsatility of pressure. **(E)** Key genes with the actin cytoskeleton pathway that are

upregulated by each dimension of force. **(F)** *Pdgfb* was downregulated in 11 of

dysregulated pathways identified in E.

Figure 3. Impact of pulsatile stretch is greater than pulsatile pressure or flow but does not change classical arteriovenous differentiation. (A) Diagram of cell culture

setup to apply isolated dimensions of force to ECs. **(B)** Venn diagram of the number of

genes significantly up- or down-regulated by pulsatility of flow, pressure, or stretch. **(C)**

Fold change (pulsatile/non-pulsatile) of any classic EC arteriovenous differentiation

markers by bulk RNAseq. Color scale denotes fold change, exact number in each box.

Significant changes (p-values < 0.05) are underlined. **(D-F)** Volcano plots of significantly

regulated genes within each dimension of pulsatile force: **(D)** LSS; **(E)** pressure; and **(F)**

stretch.

A Lung cell-specific expression of stretch- induced genes

B Cell type population key

C

D

E Isolated lung cells under pulsatile 10% stretch

F PDGFB induction in HPAECs

G Plasma PDGFB

Figure 4. **PDGFB expression in human lungs is endothelial-specific and stretch-**

induced. (A) Genes upregulated by pulsatile stretch in ECs were analyzed within the

LungMAP single cell RNA sequencing data of human lung tissue (347,970 cells).

Expression within ECs (green box) was compared to other cell types. CX3CL1 and

PDGFB (black boxes) were uniquely expressed by ECs. **(B)** Cell population groupings

as annotated by LungMAP cell x gene portal. **(C)** Gene co-expression analysis shows

expression of *Cx3cl1* in ECs with its receptor *Cx3cr1* in immune cells. **(D)** Expression of

Pdgfb is high in ECs, with its receptor *Pdgfrb* found in pericytes and vascular smooth

muscle. **(E)** Different cell lines comprising the primary components of a blood vessel

(endothelium, smooth muscle, fibroblasts) were exposed to pulsatile uniaxial 10%

stretch at 1 Hz and only ECs secreted PDGFB into the media (as measured by ELISA).

(F) Pulsatile stretch of cultured HPAECs induces 5x more PDGFB secretion than

continuous flow. Error bar shows 1 s.d. **(G)** PDGFB measured by ELISA from patient

plasma does not show a significant arteriovenous concentration gradient or difference

between normal and Glenn patients. Error bar shows 1 s.d.

A — Normal

Phrenic Nerve
Aorta
Left SVC
Left PA
Azygous Vein

A' — Glenn

Left SVC
Aorta
Left PA
Azygous Vein

C Endothelial PDGFB after rat Glenn and Sham procedures

B Sham   Glenn

MERGE
DAPI
CDH5
PDGFB
aSMA

Figure 5. **Loss of PDGFB in pulmonary endothelium of rats after Glenn surgery. (A)**
Schematic of normal rat vascular anatomy and (**A'**) rat Glenn surgery (adapted from Wan
2024 with permission). **(B)** Representative immunofluorescent (IF) staining of pulmonary
arterioles from left lung of sham and Glenn-operated rats. Pulmonary arterioles were
identified by presence of VECAD (cyan) and SMA (pink), near lung periphery, and less
than 50 μm in diameter. Scale bar is 5 μm. **(C)** Quantification of PDGFB staining in
pulmonary arteriolar and capillary ECs in sham (N=91 arteriolar ECs, N=74 capillary ECs)
and Glenn-operated (N=89 arteriolar ECs, N=62 capillary ECs) rats demonstrates
significant loss of EC-derived PDGFB in the Glenn lung. 5 vessels each from the left lung
of 3 sham and 3 Glenn rats were analyzed.
804
805

Proximal

A aSMA Sham — Glenn

B Rat #1 25.2 μm — Rat #4 22.2 μm
Rat #2 30.3 μm — Rat #5 13.6 μm
aSMA Rat #3 36.1 μm — Rat #6 19.3 μm

Proximal

C

Distal

D

E Sham — Glenn
CDH5
aSMA
DAPI

Distal

F

Distal

G

Figure 6. **Loss of smooth muscle layer in pulmonary arteries of rats after Glenn**

surgery. (A) Example sagittal sections of lung tissue from sham and Glenn operated

rats, stained by IHC for smooth muscle actin (SMA). Black boxes show typical perihilar

location of analysis of the most proximal portions of pulmonary arteries examined. Scale

bar = 2 mm. **(B)** Left panels - Example images of SMA+ media thickness from 3

different sham operated rats. Scale bar = 20 μm. Right panels - Example images of

SMA+ media thickness from 3 different Glenn rats. Scale bar = 20 μm. **(C)**

Quantification of SMA+ tunica media from proximal arteries versus veins. **(D)** Schematic

showing approximate location of pulmonary vessels assayed in A, B and E. **(E)**

Examples of images of SMA+ media thickness in distal arterioles from 3 different sham

(left panels) and Glenn (right panels). Distal lung vessels < 0.1 mm were identified using

IF staining for CDH5 (green) in Sham (n=3 rats, n=42 vessels) and Glenn (n=3 rats,

n=56 vessels) rat lungs, and staining of SMA (pink) was used to quantify the

circumferential coverage **(F)** and intensity **(G) of** SMA surrounding these arterioles.

Figure 7. Outline of primary findings. **(A)** Overview of experimental results. **(B)**

Comparison of findings between pulsatile and non-pulsatile conditions. Red boxes

indicate region of pulmonary arteries analyzed in patients with either normal lung

vascular architecture or Glenn anatomy. Blue box area shows increased PDGFB

secretion from ECs following blood circulation-induced stretch. EC, endothelial cell

(green); vSMC, vascular smooth muscle cell (brown); BM, basement membrane, and

ECM; extracellular matrix (beige).